

-

SUPPORT THEORY FOR EXTENDED DRINFELD DOUBLES

ERIC M. FRIEDLANDER

ABSTRACT. Following earlier work with Cris Negron on the cohomology of Drinfeld doubles $D(\mathbb{G}_{(r)})$, we develop a “geometric theory” of support varieties for “extended Drinfeld doubles” $\tilde{D}(\mathbb{G}_{(r)})$ of Frobenius kernels $\mathbb{G}_{(r)}$ of smooth linear algebraic groups \mathbb{G} over a field k of characteristic $p > 0$. To a $\tilde{D}(\mathbb{G}_{(r)})$ -module M we associate the space $\Pi(\tilde{D}(\mathbb{G}_{(r)}))_M$ of equivalence classes of “pairs of π -points” and prove most of the desired properties of $M \mapsto \Pi(\tilde{D}(\mathbb{G}_{(r)}))_M$. Namely, this association satisfies the “tensor product property” and admits a natural continuous map $\Psi_{\tilde{D}}$ to cohomological support theory. Moreover, for M finite dimensional and with suitable conditions on $\mathbb{G}_{(r)}$, this association provides a “projectivity test”, $\Psi_{\tilde{D}}$ is a homeomorphism, and identifies $\Pi(\tilde{D}(\mathbb{G}_{(r)}))_M$ with the cohomological support variety of M for various classes of $\tilde{D}(\mathbb{G}_{(r)})$ -modules M .

0. INTRODUCTION

Cohomological support varieties for kG -modules for a finite group G constitute natural extensions of D. Quillen’s geometric description of $\text{Spec } H^\bullet(G, k)$ [19], [20]. As introduced by J. Carlson in [3], the cohomological support variety $|G|_M$ of a finite dimensional kG -module M is the closed subvariety of $\text{Spec } H^\bullet(G, k)$ associated to the (radical of) the kernel of the natural map $H^\bullet(G, k) \rightarrow \text{Ext}_G^*(M, M)$ of k -algebras. To make this more geometric and more computable in the special case that G is an elementary abelian p -group, J. Carlson conjectured and G. Avrunin and L. Scott proved (in [1]) that $|E|_M$ can be identified with the “rank variety” $V(E)_M$ of cyclic shifted subgroups along which the pull-backs of M are not projective. This “geometric interpretation” of cohomological support varieties for modules for an elementary abelian p -group was successively extended by B. Parshall and the author to modules for restricted Lie algebras [6], by A. Suslin, C. Bendel, and the author in [23] to modules for infinitesimal group schemes, and to modules for arbitrary finite group schemes by J. Pevstova and the author [8]. The interweaving of the two approaches (one, primarily cohomological; the second, more geometric) has led to various constructions and invariants.

In this paper, we adapt the theory of π -points for finite group schemes to a theory of π -point pairs for a special class of finite dimensional Hopf algebras over k which are neither commutative nor cocommutative. Namely, beginning with a smooth linear algebraic group \mathbb{G} over k , we consider the Frobenius kernel $\mathbb{G}_{(r)}$ (i.e., the kernel of the r -th iterate of the Frobenius map, $F^r : \mathbb{G} \rightarrow \mathbb{G}^{(r)}$ for some $r > 0$) and take the smash product algebra $\tilde{D}(\mathbb{G}_{(r)}) \equiv k[\mathbb{G}_{(r+1)}] \# k\mathbb{G}_{(r)}$ which maps surjectively to the Drinfeld double $D(\mathbb{G}_{(r)})$ of $\mathbb{G}_{(r)}$.

Date: November 13, 2021.

2010 Mathematics Subject Classification. 16G99, 16S40, 16T05.

Key words and phrases. Drinfeld doubles, Frobenius kernels, Support varieties.

Partially supported by the Simons Foundation.

What enables our approach for $\tilde{D}(\mathbb{G}_{(r)})$ -modules (and does not work for modules for the Drinfeld double $D(\mathbb{G}_{(r)})$) is the existence of a sub Hopf algebra $O(\mathbb{G}_{(r)}) \subset \tilde{D}(\mathbb{G}_{(r)})$ which essentially detects the cohomology of $\tilde{D}(\mathbb{G}_{(r)})$ and which is isomorphic as an algebra to the “group algebra” for a finite group scheme. Indeed, almost by definition, the π -point pair space $\Pi(\tilde{D}(\mathbb{G}_{(r)}))$ maps bijectively to the π -point pair space $\Pi(O(\mathbb{G}_{(r)}))$ which is homeomorphic to the π -point space of the associated finite group scheme. To establish the desired tensor product property $\Pi(\tilde{D}(\mathbb{G}_{(r)}))_{M \otimes M'} = \Pi(\tilde{D}(\mathbb{G}_{(r)}))_M \cap \Pi(\tilde{D}(\mathbb{G}_{(r)}))_N$ in Theorem 3.4, we utilize this tensor product property for finite group schemes. The key step requires the verification that this property remains valid upon changing coproduct structure on this group algebra to the coproduct structure on $O(\mathbb{G}_{(r)})$. This invariance of coproduct of supports of a tensor product is the content of Proposition 3.2. The tensor product property enables a natural topology on $\Pi(\tilde{D}(\mathbb{G}_{(r)}))$.

In Section 4, we extend the arguments of C. Negron and the author in [5] for the cohomology of Drinfeld doubles to the cohomology of the extended doubles $\tilde{D}(\mathbb{G}_{(r)})$ which allows a comparison in Corollary 4.3 of the cohomology of $\tilde{D}(\mathbb{G}_{(r)})$ with that of the simpler subalgebra $O(\mathbb{G}_{(r)})$. In Proposition 4.6, we establish a commutative square of continuous maps relating π -point pair spaces for $O(\mathbb{G}_{(r)})$ and $\tilde{D}(\mathbb{G}_{(r)})$ with (projectivized) prime ideal spectra of their cohomology.

In order to proceed further, we require the determination of the cohomology algebra $H^\bullet(\tilde{D}(\mathbb{G}_{(r)}), k)$, the commutative subalgebra of the Ext-algebra $Ext_{\tilde{D}(\mathbb{G}_{(r)}}^*(k, k)$ of classes of even degree for $p > 2$ (and the full Ext-algebra if $p = 2$). This is achieved in Theorem 5.4 provided that \mathbb{G} admits a quasilogarithm (see Example 5.2) and that $p^{r+1} > 2dim(\mathbb{G})$, again using techniques of [5]. This computation allows us to show that $\Psi_{\tilde{D}} : \Pi(\tilde{D}(\mathbb{G}_{(r)})) \rightarrow \mathbb{P}(H^\bullet(\tilde{D}(\mathbb{G}_{(r)}), k))$ is a homeomorphism. As a consequence, we conclude in Theorem 5.6 that a finite dimensional $\tilde{D}(\mathbb{G}_{(r)})$ -module is projective if and only if $\Pi(\tilde{D}(\mathbb{G}_{(r)}))_M$ is empty.

The fact that $\Psi_{\tilde{D}}$ is a homeomorphism mapping $\Pi(\tilde{D}(\mathbb{G}_{(r)}))_M$ to $\mathbb{P}(H^\bullet(\tilde{D}(\mathbb{G}_{(r)}), k))_M$ does not a priori identify $\Pi(\tilde{D}(\mathbb{G}_{(r)}))_M$ with $\mathbb{P}(H^\bullet(\tilde{D}(\mathbb{G}_{(r)}), k))_M$. In Section 6, we achieve this identification for various special classes of $\tilde{D}(\mathbb{G}_{(r)})$ -modules, including Carlson’s L_ζ -modules, modules whose action factors through the quotient $\tilde{D}(\mathbb{G}_{(r)}) \twoheadrightarrow k\mathbb{G}_{(r)}$ and modules whose restriction to $k[\mathbb{G}_{(r+1)}]$ are projective.

In Section 7, we observe that the same techniques we use to investigate $\tilde{D}(\mathbb{G}_{(r)})$ -modules apply to modules for closely related Hopf algebras. We also mention a couple of questions worthy of further investigation.

We thank both Cris Negron and Julia Pevtsova for their contributions to the foundations of this work, and the Institute for Advanced Study for its (virtual) hospitality.

1. $D(\mathbb{G}_{(r)}), \tilde{D}(\mathbb{G}_{(r)}), O(\mathbb{G}_{(r)}), \mathbb{G}_{(r)} \times (\underline{\mathfrak{g}}^{(r)})_{(1)}$

Throughout this work, \mathbb{G} will denote a connected linear algebraic group smooth over k . If A is a k -algebra and K/k is a field extension, then we denote by A_K the base change $K \otimes A$; for an A -module M , we use the notation M_K for the A_K -module $K \otimes M$. For a finite dimensional k -algebra A , we denote by $H^\bullet(A, k) \subset Ext_A^*(k, k)$ the commutative subalgebra of even dimensional classes if $p > 2$ and the full algebra $Ext_A^*(k, k)$ if $p = 2$.

For any $r > 0$, we denote by $F^r : \mathbb{G} \rightarrow \mathbb{G}^{(r)}$ the r -th iterate of the Frobenius map $F : \mathbb{G} \rightarrow \mathbb{G}^{(1)}$ given by the k -linear map

$$F^* : k[\mathbb{G}^{(1)}] \cong k \otimes_{\phi} k[\mathbb{G}] \rightarrow k[\mathbb{G}], \quad \lambda \otimes f \mapsto \lambda f^p$$

of coordinate algebras (where $k \otimes_{\phi} -$ denotes the base change along the p -th power map $\phi : k \rightarrow k$). The r -th Frobenius kernel of \mathbb{G} is the finite group scheme

$$(1.0.1) \quad \mathbb{G}_{(r)} \equiv \ker\{F^r : \mathbb{G} \rightarrow \mathbb{G}^{(r)}\} \hookrightarrow \mathbb{G},$$

whose coordinate algebra $k[\mathbb{G}_{(r)}]$ is the local commutative algebra defined as the quotient of the localization $(k[\mathbb{G}])_{(e)}$ of $k[\mathbb{G}]$ at the identity $e \in \mathbb{G}$ modulo the ideal generated by elements $x^{p^r}, x \in \mathfrak{m}_e$. For example, $k[GL_n(r)] = k[x_{i,j}]/(x_{i,j}^{p^r} - \delta_{i,j})$.

The coordinate algebra $k[\mathbb{G}]$ of \mathbb{G} is a Hopf algebra with comultiplication determined by the multiplication of \mathbb{G} . For a finite group scheme G such as $\mathbb{G}_{(r)}$, we consider the k -linear dual kG of $k[G]$ which is a finite dimensional, co-commutative Hopf algebra (over k); we refer to kG as the group algebra of G . For such a finite group scheme G , a (rational) G -module is a k -vector space equipped with a k -linear module structure, $kG \times M \rightarrow M$. We shall frequently refer to this structure as the ‘‘action of G on M ’’.

We recall that the Drinfeld double of a Hopf algebra H is the double crossed product of H^{*cop} and H , where H^{*cop} has the same algebra structure as H^* and coproduct given by the opposite of the coproduct of H^* . If H is finite dimensional and cocommutative, then $D(H)$ is isomorphic to the smash product $H^{*op} \# H$ (see [18, 10.3.10]).

In Hopf algebra standard notation, the coadjoint action of an element $h \in H$ on an element $f \in H^*$ is given by $\sum f(h_1 \rightharpoonup f \leftarrow s^{-1}(h_2))$ where s is the antipode of H , Δ is the coproduct of H^* and $\Delta(h)$ is written as $\sum h_1 \otimes h_2$. By a result of D. Radford [21, 10.3.5], the product structure of $D(H)$ is given for finite dimensional H by

$$(f \# h)(f' \# h') = \sum f(h_1 \rightharpoonup f' \leftarrow s^{-1}(h_2)) \# h_3 h'.$$

If H is both finite dimensional and cocommutative, this multiplication is that of the smash product algebra $H^* \# H$:

$$(f \# h)(f' \# h') = \sum f h_1(f) \# h_2 h';$$

the coproduct of $D(H)$ is the tensor product of the opposite coproduct of H^* and the coproduct of H .

Definition 1.1. As discussed above, the Drinfeld double of the finite dimensional cocommutative Hopf algebra $k\mathbb{G}_{(r)}$ has algebra structure given by the smash product

$$(1.1.1) \quad D(\mathbb{G}_{(r)}) \equiv k[\mathbb{G}_{(r)}] \# k\mathbb{G}_{(r)}$$

with respect to the right coadjoint action of $\mathbb{G}_{(r)}$ on $k[\mathbb{G}_{(r)}]$ and coalgebra structure given by the tensor product of the co-opposite coproduct on $k[\mathbb{G}_{(r)}]$ (i.e., the opposite product structure of $k\mathbb{G}_{(r)}$) and the coproduct of $k\mathbb{G}_{(r)}$.

The extended Drinfeld double $\tilde{D}(\mathbb{G}_{(r)})$ is the analogous smash product with respect to the restriction to $\mathbb{G}_{(r)}$ of the coadjoint action of $\mathbb{G}_{(r+1)}$ on $k[\mathbb{G}_{(r+1)}]$:

$$(1.1.2) \quad \tilde{D}(\mathbb{G}_{(r)}) \equiv k[\mathbb{G}_{(r+1)}] \# k\mathbb{G}_{(r)} \hookrightarrow D(\mathbb{G}_{(r+1)}).$$

To verify that (1.1.2) is a well defined inclusion of Hopf algebras, we observe that the restriction of the coadjoint action of $\mathbb{G}_{(r+1)}$ on $k[\mathbb{G}_{(r+1)}]$ along $\mathbb{G}_{(r)} \hookrightarrow \mathbb{G}_{(r+1)}$ is the action used to define $\tilde{D}(\mathbb{G}_{(r)})$, then check directly that the multiplication and comultiplication of $D(\mathbb{G}_{(r+1)})$ restricts via the evident inclusion of k -vector spaces $k[\mathbb{G}_{(r+1)}] \otimes k\mathbb{G}_{(r)} \subset k[\mathbb{G}_{(r+1)}] \otimes k\mathbb{G}_{(r+1)}$ to the multiplication and comultiplication of $\tilde{D}(\mathbb{G}_{(r)})$.

We shall be interested in (left) modules for the extended Drinfeld double $\tilde{D}(\mathbb{G}_{(r)})$ which can be viewed as k -vector spaces N with an action $\tilde{D}(\mathbb{G}_{(r)}) \times N \rightarrow N$. The condition on such a pairing to be a left module structure can be expressed as the restriction to elements in $\tilde{D}(\mathbb{G}_{(r)}) \subset D(\mathbb{G}_{(r+1)})$ of the condition that a pairing $D(\mathbb{G}_{(r+1)}) \times N \rightarrow N$ be a $D(\mathbb{G}_{(r+1)})$ -module structure. This latter condition is made explicit in the following proposition (with $H = k\mathbb{G}_{(r+1)}$).

We first observe that for any Hopf algebra H there are natural algebra embeddings of H, H^* in $D(H)$: $h \in H \mapsto 1\#h$; $f \in H^* \mapsto f\#1$.

Proposition 1.2. [18, 10.6.16], [16] *Let H be a finite dimensional Hopf Algebra. The data of a $D(H)$ -module structure on a vector space N consists of module structures on N for both H^* and H such that*

$$h \cdot (f \cdot n) = \sum (h_1 \rightharpoonup f_2) \cdot ((h_2 \leftarrow f_1) \cdot n) \in N.$$

This is equivalent to a left/right Yetter-Drinfeld module structure on N which consists of the data of a $k\mathbb{G}_{(r)}$ -module structure and a right $k\mathbb{G}_{(r)}$ -comodule structure on N satisfying the Yetter-Drinfeld condition:

$$\sum h_1 \cdot n_o \otimes h_2 n_1 = \sum (h_1 n)_0 \otimes (h_2 \cdot n)_1 h_1 \in N \otimes k\mathbb{G}_{(r)}.$$

Here, we have written the coaction $N \rightarrow N \otimes H$ as $n \mapsto \sum n_o \otimes h_1$.

We relate the extended Drinfeld double $\tilde{D}(\mathbb{G}_{(r)})$ to various other structures.

Definition 1.3. We define

$$(1.3.1) \quad O(\mathbb{G}_{(r)}) \equiv k[(\mathbb{G}^{(r)})_{(1)}] \otimes k\mathbb{G}_{(r)}.$$

Using the natural identification of $(\mathbb{G}^{(r)})_{(1)}$ with $\mathbb{G}_{(r+1)}/\mathbb{G}_{(r)}$, we obtain the natural embedding

$$(1.3.2) \quad i_O : O(\mathbb{G}_{(r)}) \hookrightarrow \tilde{D}(\mathbb{G}_{(r)}).$$

Proposition 1.4. *The embedding $\mathbb{G}_{(r)} \subset \mathbb{G}_{(r+1)}$ determines a $\mathbb{G}_{(r)}$ -equivariant quotient map $k[\mathbb{G}_{(r+1)}] \twoheadrightarrow k[\mathbb{G}_{(r)}]$ of Hopf algebras and thus a quotient map of Hopf algebras*

$$(1.4.1) \quad q : \tilde{D}(\mathbb{G}_{(r)}) \twoheadrightarrow D(\mathbb{G}_{(r)}).$$

Similarly, the quotient map $\mathbb{G}_{(r+1)} \twoheadrightarrow (\mathbb{G}^{(r)})_{(1)}$ determining i_O of (1.3.2) is an embedding of Hopf algebras

The coproduct structure inherited by $O(\mathbb{G}_{(r)})$ is the tensor product of the opposite of the coproduct on $k[(\mathbb{G}^{(r)})_{(1)}]$ (in other words the dual of the opposite multiplication of $k[(\mathbb{G}^{(r)})_{(1)}]$) and the coproduct of $k\mathbb{G}_{(r)}$ (the dual of the multiplication of $k[\mathbb{G}_{(r)}]$). Thus, $O(\mathbb{G}_{(r)})$ is co-commutative if and only if $\mathfrak{g} = \text{Lie}(\mathbb{G})$ is commutative.

Proof. Since each $\mathbb{G}_{(r)}$ is normal in \mathbb{G} , the embedding $\mathbb{G}_{(r)} \subset \mathbb{G}_{(r+1)}$ is \mathbb{G} -equivariant so that $k[\mathbb{G}_{(r+1)}] \twoheadrightarrow k[\mathbb{G}_{(r+1)}/\mathbb{G}_{(r)}]$ is a \mathbb{G} -equivariant quotient map of Hopf algebras; thus, this $\mathbb{G}_{(r)}$ -equivariant quotient determines the map of Hopf algebras $q : \tilde{D}(\mathbb{G}_{(r)}) \twoheadrightarrow D(\mathbb{G}_{(r)})$.

Similarly, the quotient $\mathbb{G}_{(r+1)} \twoheadrightarrow \mathbb{G}^{(r)}_{(1)}$ is also \mathbb{G} -equivariant, determining the \mathbb{G} -equivariant embedding $k[\mathbb{G}^{(r)}_{(1)}] \hookrightarrow k[\mathbb{G}_{(r+1)}]$ of Hopf algebras. Observe that $\mathbb{G}_{(r)} \subset \mathbb{G}$ acts trivially on $k[\mathbb{G}^{(r)}_{(1)}]$, so that we obtain the embedding (as algebras) of the tensor product $k[\mathbb{G}^{(r)}_{(1)}] \otimes k\mathbb{G}_{(r)} \hookrightarrow \tilde{D}(\mathbb{G}_{(r)})$. \square

Proposition 1.5. *The subalgebra $k[\mathbb{G}^{(r)}_{(1)}] \simeq k[\mathbb{G}_{(r+1)}/\mathbb{G}_{(r)}]$ of the commutative algebra $k[\mathbb{G}_{(r+1)}]$ gives $k[\mathbb{G}_{(r+1)}]$ the structure of a free $k[\mathbb{G}^{(r)}_{(1)}]$ -module with basis $\{\tilde{f}_i\}$ given by lifting along $k[\mathbb{G}_{(r+1)}] \rightarrow k[\mathbb{G}_{(r)}]$ a k -basis $\{f_i\}$ for $k[\mathbb{G}_{(r)}]$.*

This structure determines a free (right) $O(\mathbb{G}_{(r)})$ -module structure on $\tilde{D}(\mathbb{G}_{(r)})$ with basis $\{\tilde{f}_i\#1\}$.

Proof. We choose a k -linear section $\sigma : k[\mathbb{G}_{(r)}] \rightarrow k[\mathbb{G}_{(r+1)}]$ of the quotient map $k[\mathbb{G}_{(r+1)}] \twoheadrightarrow k[\mathbb{G}_{(r)}]$ and consider the $k[\mathbb{G}_{(r+1)}/\mathbb{G}_{(r)}]$ -bilinear map

$$(1.5.1) \quad k[\mathbb{G}_{(r)}] \otimes k[\mathbb{G}_{(r+1)}/\mathbb{G}_{(r)}] \rightarrow k[\mathbb{G}_{(r+1)}], \quad f_i \otimes h \mapsto \sigma(f_i) \cdot h.$$

We readily check that this induces a k -linear isomorphism, for example by showing that tensors of elements of k -bases for $k[\mathbb{G}_{(r)}]$ and $k[\mathbb{G}_{(r+1)}/\mathbb{G}_{(r)}]$ determine a k -linearly independent set in $k[\mathbb{G}_{(r+1)}]$.

Using the fact that $k[\mathbb{G}_{(r+1)}/\mathbb{G}_{(r)}] \otimes 1 \subset \tilde{D}(\mathbb{G}_{(r)})$ is central, we readily check that

$$k[\mathbb{G}_{(r)}] \otimes O(\mathbb{G}_{(r)}) \rightarrow \tilde{D}(\mathbb{G}_{(r)}), \quad f \otimes (f' \otimes h) \mapsto (\sigma(f)\#1) \cdot (f' \#h)$$

is an $O(\mathbb{G}_{(r)})$ -linear isomorphism \square

The following observation about $O(\mathbb{G}_{(r)})$ will prove central in our investigation of $\tilde{D}(\mathbb{G}_{(r)})$ -modules. The reader should not confuse the finite group scheme $(\underline{\mathfrak{g}}^{(r)})_{(1)}$ occurring in the proposition below with $(\mathbb{G}^{(r)})_{(1)}$.

Proposition 1.6. *Let $\underline{\mathfrak{g}}$ denote the vector group scheme $\text{Spec } S^\bullet(\mathfrak{g}^*)$ (ignoring the Lie algebra structure on \mathfrak{g}). Then the group algebra of the finite group scheme $(\underline{\mathfrak{g}}^{(r)})_{(1)} \times \mathbb{G}_{(r)}$ is isomorphic as an algebra to $\mathcal{O}(\mathbb{G}_{(r)})$.*

On the other hand, the dual of the Hopf algebra $O(\mathbb{G}_{(r)})$ is not commutative (unlike the coordinate algebra of $(\underline{\mathfrak{g}}^{(r)})_{(1)} \times \mathbb{G}_{(r)}$).

Proof. Essentially by definition, elements $X^{(r)} \in \text{Lie}(\mathbb{G}^{(r)})$ are distributions at the identity of $\mathbb{G}^{(r)}$ of $k[\mathbb{G}^{(r)}]$ of order 1 (see [13, I.7.10]), This identification determines an embedding of $\mathfrak{g}^{(r)*}$ into the maximal ideal of $k[\mathbb{G}_{(r+1)}/\mathbb{G}_{(r)}]$ which necessarily sends each $(X^{(r)})^p \in S^p(\mathfrak{g}^{(r)*})$ to 0. The resulting map

$$\nu : S^\bullet(\mathfrak{g}^{(r)*}) / ((X^{(r)})^p, X \in \mathfrak{g}^*) \rightarrow k[\mathbb{G}_{(r+1)}/\mathbb{G}_{(r)}]$$

is injective, and thus surjective by dimension reasons. The proof is completed by observing that the group algebra of $(\underline{\mathfrak{g}}^{(r)})_{(1)}$ equals $S^\bullet(\mathfrak{g}^{(r)*}) / ((X^{(r)})^p, X \in \mathfrak{g}^*)$. \square

We conclude this section with some remarks about π -point spaces $\Pi(G_1 \times G_2)$ of a product of finite group schemes which we shall use for $G_1 \times G_2 = (\underline{\mathfrak{g}}^{(r)})_{(1)} \times \mathbb{G}_{(r)}$. Recall that a π -point $\alpha_K : K[t]/t^p \rightarrow kG$ of a finite group scheme G over k is a flat

map of algebras which factors through the group algebra of some unipotent abelian subgroup scheme $C_K \subset G_K$, where K is some field extension of k . Thus, a π -point of $G_1 \times G_2$ is a pair of π -points $\alpha_{K,1} : K[t]/t^p \rightarrow KG_1$, $\alpha_{K,2} : K[t]/t^p \rightarrow KG_2$ corresponding to a flat map

$$\alpha_{K,1} + \alpha_2 : K[t]/t^p \rightarrow KC_1 \otimes KC_2, \quad t \mapsto \alpha_{K,1}(t) \otimes 1 + 1 \otimes \alpha_{K,1}(t)$$

with $C_{K,i} \subset G_{K,i}$ unipotent abelian subgroup schemes for $i = 1, 2$.

It is important here for $\alpha_{K,1}(t)$, $\alpha_{K,2}(t)$ to commute in order that $\alpha_{K,1} + \alpha_{K,2}$ is well defined.

By [8, Thm 3.6], two π -points $\alpha_K : K[t]/t^p \rightarrow KG$, $\beta_L : L[t]/t^p \rightarrow LG$ are equivalent if and only if the two ideals of $H^\bullet(G, k)$

$$\ker\{\alpha_K^* : H^\bullet(G, K) \rightarrow H^*(K[t]/t^p, K)\} \cap H^\bullet(G, k),$$

$$\ker\{\beta_L^* : H^\bullet(G, L) \rightarrow H^*(L[t]/t^p, L)\} \cap H^\bullet(G, k)$$

are equal. In other words, this gives a natural bijection of sets (which is sharpened to an isomorphism of schemes in [8])

$$(1.6.1) \quad \Pi_G : \Pi(G) \xrightarrow{\sim} \mathbb{P}H^\bullet(G, k).$$

Observe that the flat map $(\alpha_{K,1}, \alpha_{K,2}) : K[t]/t^p \rightarrow KG_1 \times KG_2$ induces the map

$$(\alpha_{K,1}^* \otimes 1) + (1 \otimes \alpha_{K,2}^*) : H^*(C_1, K) \otimes_K H^*(C_2, K) \rightarrow H^*(K[t]/t^p, K).$$

We denote the composition of this map with the restriction map

$$(\alpha_{K,1} + \alpha_{K,2})^* : H^*(G_1 \times G_2, K) \rightarrow H^*(K[t]/t^p, K).$$

Thus, we have proved the following proposition.

Proposition 1.7. *Two π -points $(\alpha_{K,1}, \alpha_{K,2})$ and $(\beta_{L,1}, \beta_{L,2})$ of the finite group scheme $G_1 \times G_2$ are equivalent if and only if the two ideals of $H^\bullet(G_1 \times G_2, k)$*

$$\ker\{(\alpha_{K,1} + \alpha_{K,2})^* : H^\bullet(G_1 \times G_2, K) \rightarrow H^*(K[t]/t^p, K)\} \cap H^\bullet(G_1 \times G_2, k),$$

$$\ker\{(\beta_{L,1} + \beta_{L,2})^* : H^\bullet(G_1 \times G_2, L) \rightarrow H^*(L[t]/t^p, L)\} \cap H^\bullet(G_1 \times G_2, k)$$

are equal.

In particular, if $\alpha_{K,i}$ is equivalent to $\beta_{L,i}$ for $i = 1, 2$, then $(\alpha_{K,1}, \alpha_{K,2})$ and $(\beta_{L,1}, \beta_{L,2})$ are equivalent.

2. THE SPACE $\Pi(\tilde{D}(\mathbb{G}_{(r)}))$ OF π -POINT PAIRS

In this section we introduce an extension of “rank varieties” which applies to $\tilde{D}(\mathbb{G}_{(r)})$ -modules. Our construction is a modification of the “ π -point construction” of [8]. The strategy we follow is relatively straight-forward. In view of the homological computations of [5], it seems natural to use pairs of π -points (α_K, β_K) with $\alpha_K : K[t]/t^p \rightarrow K[\mathbb{G}_{(r+1)}]$ and $\beta_K : K[t]/t^p \rightarrow K\mathbb{G}_{(r)}$. Yet the most natural way to do this (which would also apply to $D(\mathbb{G}_{(r)})$) appears to fail because $\alpha_K(t)$, $\beta_K(t)$ need not commute. We work around this difficulty by restricting $\tilde{D}(\mathbb{G}_{(r)})$ -modules to $O(\mathbb{G}_{(r)})$.

Definition 2.1. A π -point pair (α_K, β_K) of $\tilde{D}(\mathbb{G}_{(r)})$ is a pair of maps of K -algebras for some field extension K/k ,

$$\alpha_K : K[t]/t^p \rightarrow K[\mathbb{G}_{(r+1)}/\mathbb{G}_{(r)}] \hookrightarrow K[\mathbb{G}_{(r+1)}], \quad \beta_K : K[t]/t^p \rightarrow K\mathbb{G}_{(r)},$$

such that β_K either sends t to 0 or is a π -point of $\mathbb{G}_{(r)}$, α_K either sends t to 0 or is flat, and at least one of α_K, β_K is flat.

The following lemma provides an essential property of a π -point pair (α_K, β_K) .

Lemma 2.2. *A π -point pair (α_K, β_K) of $\tilde{D}(\mathbb{G}_{(r)})$ determines a flat map*

$$\alpha_K + \beta_K : K[t]/t^p \rightarrow O(\mathbb{G}_{(r)})_K, \quad t \mapsto \alpha_K(t)\#1 + 1\#\beta_K(t)$$

whose composition with $i_O : O(\mathbb{G}_{(r),K}) \hookrightarrow \tilde{D}(\mathbb{G}_{(r),K})$ is also flat.

Proof. We consider the composition

$$K[t] \xrightarrow{\Delta} K[t]/p^{\otimes 2} \xrightarrow{\alpha_K \boxtimes \beta_K} K[\mathbb{G}_{(r+1)}/\mathbb{G}_{(r)}] \otimes K\mathbb{G}_{(r)} = O(\mathbb{G}_{(r)})_K, \quad t \mapsto \alpha_K(t) \otimes \beta_K(t).$$

Since $\alpha_K(t), \beta_K(t)$ commute in $O(\mathbb{G}_{(r)})_K$, this composition factors through $\alpha_K + \beta_K : K[t]/t^p \rightarrow O(\mathbb{G}_{(r)})_K$. Since $\Delta : K[t]/p^{\otimes 2} \xrightarrow{\Delta} K[t]/p \otimes K[t]/t^p$ is flat, since both α_K and β_K are either flat or “trivial”, since $O(\mathbb{G}_{(r)})$ is flat over each of its tensor factors, and since $\alpha_K \boxtimes \beta_K$ is flat if $\alpha_K(t) \neq 0 \neq \beta_K(t)$, we conclude that $\alpha_K + \beta_K : K[t]/t^p \rightarrow O(\mathbb{G}_{(r)})_K$ is flat. By Proposition 1.5, this implies the composition $i_O \circ (\alpha_K + \beta_K) : K[t]/t^p \rightarrow \tilde{D}(\mathbb{G}_{(r),K})$ is also flat. \square

The following definition of equivalence for π -point pairs $(\alpha_K, \beta_K), (\alpha'_L, \beta'_L)$ is motivated by Proposition 1.7.

Definition 2.3. Two π -point pairs $(\alpha_K, \beta_K), (\alpha'_L, \beta'_L)$ of $\tilde{D}(\mathbb{G}_{(r)})$ are said to be equivalent (respectively, $O(\mathbb{G}_{(r)})$ -equivalent) if there exists some common field extension Ω of both K and L such that for all finite dimensional $\tilde{D}(\mathbb{G}_{(r)})$ -modules M (resp., all finite dimensional $O(\mathbb{G}_{(r)})$ -modules M) $(\alpha_\Omega + \beta_\Omega)^*(M_\Omega)$ is projective as an $\Omega[t]/t^p$ -module if and only if $(\alpha'_\Omega + \beta'_\Omega)^*(M_\Omega)$ is projective.

We denote by $\Pi(\tilde{D}(\mathbb{G}_{(r)}))$ the set of equivalence classes of π -point pairs of $\tilde{D}(\mathbb{G}_{(r)})$, and we denote by $\Pi(O(\mathbb{G}_{(r)}))$ the set of $O(\mathbb{G}_{(r)})$ -equivalence classes of π -point pairs of $\tilde{D}(\mathbb{G}_{(r)})$. For any $\tilde{D}(\mathbb{G}_{(r)})$ -module M , we define the Π -pair set

$$\Pi(\tilde{D}(\mathbb{G}_{(r)}))_M \subset \Pi(\tilde{D}(\mathbb{G}_{(r)}))$$

to be the subset of equivalence classes of π -point pairs of $\tilde{D}(\mathbb{G}_{(r)})$ such that $(\alpha_K + \beta_K)^*(M)$ is not projective. Similarly, for any $O(\mathbb{G}_{(r)})$ -module M' , we define $\Pi(O(\mathbb{G}_{(r)}))_{M'} \subset \Pi(O(\mathbb{G}_{(r)}))$ to be the subset of equivalence classes of π -point pairs of $\tilde{D}(\mathbb{G}_{(r)})$ such that $(\alpha_K + \beta_K)^*(M')$ is not projective.

Proposition 2.4. *The natural surjection $\Pi_{\tilde{D}} : \Pi(O(\mathbb{G}_{(r)})) \twoheadrightarrow \Pi(\tilde{D}(\mathbb{G}_{(r)}))$ is a bijection.*

Moreover, for any finite dimensional $\tilde{D}(\mathbb{G}_{(r)})$ -module M , $\Pi_{\tilde{D}}$ restricts to a bijection

$$\Pi_{\tilde{D},M} : \Pi(O(\mathbb{G}_{(r)}))_M \xrightarrow{\sim} \Pi(\tilde{D}(\mathbb{G}_{(r)}))_M.$$

Proof. If $(\alpha_K, \beta_K), (\alpha'_L, \beta'_L)$ are $O(\mathbb{G}_{(r)})$ -equivalent, then they are necessarily equivalent since equivalence involves the same condition as $O(\mathbb{G}_{(r)})$ -equivalence except that it only requires consideration of finite dimensional $\tilde{D}(\mathbb{G}_{(r)})$ -modules. Thus, we have well defined, surjective map $\Pi(O(\mathbb{G}_{(r)})) \twoheadrightarrow \Pi(\tilde{D}(\mathbb{G}_{(r)}))$.

To prove injectivity, let $(\alpha_K, \beta_K), (\alpha'_L, \beta'_L)$ represent distinct points in $\Pi(\tilde{D}(\mathbb{G}_{(r)}))$ and choose some finite dimensional $O(\mathbb{G}_{(r)})$ -module N such that $(\alpha_\Omega + \beta_\Omega)^*(N_\Omega)$ is projective whereas $(\alpha'_\Omega + \beta'_\Omega)^*(N_\Omega)$ is not projective for some some common field extension Ω of both K and L . Consider the $\tilde{D}(\mathbb{G}_{(r)})$ -module $M \equiv \tilde{D}(\mathbb{G}_{(r)}) \otimes_{O(\mathbb{G}_{(r)})} N$ and set $N' \equiv i_O^*(M)$. By Proposition 1.5, N' is isomorphic to a direct sum of copies of N (indexed by a k -basis of $k[\mathbb{G}_{(r)}]$). Thus, $(\alpha_\Omega + \beta_\Omega)^*(N'_\Omega)$ is projective whereas $(\alpha'_\Omega + \beta'_\Omega)^*(N'_\Omega)$ is not projective. Since N' is the restriction of the $D(\mathbb{G}_{(r)})$ -module M , we conclude that the images of $(\alpha_K, \beta_K), (\alpha'_L, \beta'_L)$ in $\Pi(\tilde{D}(\mathbb{G}_{(r)}))$ are distinct.

Finally, the bijectivity of $\Pi_{\tilde{D}, M}$ follows immediately from the definitions and the bijectivity of $\Pi_{\tilde{D}}$. \square

We next observe that the condition of equivalence of π -point pairs is equivalent to the seemingly stronger equivalence relation obtained by dropping the finite dimensionality condition on M in Definition 2.3.

Proposition 2.5. *Two π -point pairs $(\alpha_K, \beta_K), (\alpha'_L, \beta'_L)$ of $\tilde{D}(\mathbb{G}_{(r)})$ (respectively $O(\mathbb{G}_{(r)})$) are equivalent if and only if there exists some common field extension Ω of both K and L such that for all $\tilde{D}(\mathbb{G}_{(r)})$ -modules M (resp., all $O(\mathbb{G}_{(r)})$ -modules M) $(\alpha_\Omega + \beta_\Omega)^*(M_\Omega)$ is projective as an $\Omega[t]/t^p$ -module if and only if $(\alpha'_\Omega + \beta'_\Omega)^*(M_\Omega)$ is projective.*

Proof. Denote by $\tilde{\Pi}(\tilde{D}(\mathbb{G}_{(r)}))$ (respectively, $\tilde{\Pi}(O(\mathbb{G}_{(r)}))$) the set of equivalence classes of π -point pairs of $\tilde{D}(\mathbb{G}_{(r)})$ (resp., $O(\mathbb{G}_{(r)})$) using the equivalence relations of the statement of this proposition. We readily verify that we have a commutative square of sets

$$(2.5.1) \quad \begin{array}{ccc} \tilde{\Pi}(O(\mathbb{G}_{(r)})) & \xrightarrow{\Pi_{\tilde{D}}} & \tilde{\Pi}(\tilde{D}(\mathbb{G}_{(r)})) \\ \tilde{\Pi}_O \downarrow & & \downarrow \tilde{\Pi}_D \\ \Pi(O(\mathbb{G}_{(r)})) & \xrightarrow{\Pi_D} & \Pi(\tilde{D}(\mathbb{G}_{(r)})). \end{array}$$

The map Π_D is a bijection as seen in Proposition 2.4; the proof of that proposition also proves that $\Pi_{\tilde{D}}$ is a bijection. The fact that $\tilde{\Pi}_O$ is a bijection follows from [8, Thm 4.6] applied to the finite group scheme $(\mathfrak{g}^{(r)})_{(1)} \times \mathbb{G}_{(r)}$ (see Proposition 2.7 below). Thus, the fact that $\tilde{\Pi}_D$ is a bijection follows from the surjectivity of $\Pi_{\tilde{D}}$ and a simple diagram chase. \square

The following properties of $M \mapsto \Pi(\tilde{D}(\mathbb{G}_{(r)}))_M$ are each verified by restricting along flat maps $(\alpha_\Omega + \beta_\Omega)^*$ associated to π -point pairs (α_K, β_K) , thereby reducing each assertion to the corresponding assertion about $K[t]/t^p$ -modules.

Proposition 2.6. *Let M_1, M_2, M_3 be $\tilde{D}(\mathbb{G}_{(r)})$ -modules.*

- (1) $\Pi(\tilde{D}(\mathbb{G}_{(r)}))_k = \Pi(\tilde{D}(\mathbb{G}_{(r)}))$.
- (2) *If M_1 is a projective $\tilde{D}(\mathbb{G}_{(r)})$ -module, then $\Pi(\tilde{D}(\mathbb{G}_{(r)}))_{M_1} = \emptyset$.*
- (3) *If $0 \rightarrow M_1 \rightarrow M \rightarrow M_2 \rightarrow$ is exact and if σ is a permutation of $\{1, 2, 3\}$, then*

$$(4) \quad \begin{aligned} \Pi(\tilde{D}(\mathbb{G}_{(r)}))_{M_{\sigma(1)}} &\subset \Pi(\tilde{D}(\mathbb{G}_{(r)}))_{M_{\sigma(2)}} \cup \Pi(\tilde{D}(\mathbb{G}_{(r)}))_{M_{\sigma(3)}}. \\ \Pi(\tilde{D}(\mathbb{G}_{(r)}))_{M_1 \oplus M_3} &= \Pi(\tilde{D}(\mathbb{G}_{(r)}))_{M_1} \cup \Pi(\tilde{D}(\mathbb{G}_{(r)}))_{M_3}. \end{aligned}$$

We proceed to show that $\Pi(O(\mathbb{G}_{(r)}))$ can be “identified” with the set of Zariski points of the π -point scheme of the finite group scheme $(\underline{\mathfrak{g}}^{(r)})_{(1)} \times \mathbb{G}_{(r)}$ introduced below.

Proposition 2.7. *Let $\underline{\mathfrak{g}} \simeq (\mathbb{G}_a)^{\times \dim(\mathfrak{g})}$ denote the vector group scheme associated to the underlying vector space of $\mathfrak{g} = \text{Lie}(\mathbb{G})$. As an algebra $O(\mathbb{G}_{(r)})$ is isomorphic to the group algebra of the infinitesimal group scheme $(\underline{\mathfrak{g}}^{(r)})_{(1)} \times \mathbb{G}_{(r)}$.*

Choose an isomorphism of k -algebras $O(\mathbb{G}_{(r)}) \xrightarrow{\sim} k((\underline{\mathfrak{g}}^{(r)})_{(1)} \times \mathbb{G}_{(r)})$. Then sending a π -point pair (α_K, β_K) of $\tilde{D}(\mathbb{G}_{(r)})$ to

$$\alpha_K + \beta_K : K[t]/t^p \rightarrow O(\mathbb{G}_{(r),K}) \xrightarrow{\sim} K((\underline{\mathfrak{g}}^{(r)})_{(1)} \times \mathbb{G}_{(r)})$$

determines a bijection

$$(2.7.1) \quad \Pi_O : \Pi(O(\mathbb{G}_{(r)})) \xrightarrow{\sim} \Pi((\underline{\mathfrak{g}}^{(r)})_{(1)} \times \mathbb{G}_{(r)}),$$

where we abuse notation by interpreting the right hand side as the set of (Zariski) points of the indicated Π -point scheme.

Moreover, for every $O(\mathbb{G}_{(r)})$ -module M , Π_O restricts to a bijection

$$(2.7.2) \quad \Pi_{O,M} : \Pi(O(\mathbb{G}_{(r)}))_M \simeq \Pi((\underline{\mathfrak{g}}^{(r)})_{(1)} \times \mathbb{G}_{(r)})_M.$$

Proof. Since $(\underline{\mathfrak{g}}^{(r)})_{(1)}$ is an abelian unipotent finite group scheme, since β_K is a π -point of $\mathbb{G}_{(r)}$ and thus factors through an abelian unipotent subgroup scheme, and since $\alpha_K + \beta_K : K[t]/t^p \rightarrow KG$ is flat by Lemma 2.2, we conclude that $\alpha_K + \beta_K$ is a π -point of the finite group scheme $(\underline{\mathfrak{g}}^{(r)})_{(1)} \times \mathbb{G}_{(r)}$. Moreover, the discussion preceding shows that every point of $\Pi((\underline{\mathfrak{g}}^{(r)})_{(1)} \times \mathbb{G}_{(r)})$ (i.e., every equivalence class of π -points of $(\underline{\mathfrak{g}}^{(r)})_{(1)} \times \mathbb{G}_{(r)}$) is represented by such a flat map $\alpha_K + \beta_K$ associated to a π -point pair of $\tilde{D}(\mathbb{G}_{(r)})$.

Thus, to prove that Π_O of (2.7.1) is well defined and injective (as well as surjective), it suffices to show that two π -point pairs $(\alpha_K, \beta_K), (\alpha'_K, \beta'_K)$ of $\tilde{D}(\mathbb{G}_{(r)})$ are $O(\mathbb{G}_{(r)})$ -equivalent if and only if $\alpha_K + \beta_K, \alpha'_K + \beta'_K$ are equivalent π -points of $(\underline{\mathfrak{g}}^{(r)})_{(1)} \times \mathbb{G}_{(r)}$. This follows immediately from Definition 2.3, Proposition 1.7, and the bijection $\Psi_{(\underline{\mathfrak{g}}^{(r)})_{(1)} \times \mathbb{G}_{(r)}}$ of (1.6.1).

The fact that (2.7.1) restricts to a bijection $\Pi(O(\mathbb{G}_{(r)}))_M \xrightarrow{\sim} \Pi((\underline{\mathfrak{g}}^{(r)})_{(1)} \times \mathbb{G}_{(r)})_M$ follows immediately from the definitions (i.e., from Definition 2.3 plus Proposition 2.4 and [7, Defn 3.2]). \square

3. THE TENSOR PRODUCT PROPERTY FOR $\tilde{D}(\mathbb{G}_{(r)})$ -MODULES

Propositions 2.4 and 2.7 identify $M \mapsto \Pi(\tilde{D}(\mathbb{G}_{(r)}))_M$ with $M \mapsto \Pi((\underline{\mathfrak{g}}^{(r)})_{(1)} \times \mathbb{G}_{(r)})_M$. This immediately tells us that $M \mapsto \Pi(\tilde{D}(\mathbb{G}_{(r)}))_M$ satisfies most of the properties required for a good theory of support varieties of $\tilde{D}(\mathbb{G}_{(r)})$ -modules. However, this identification does not imply that $M \mapsto \Pi(\tilde{D}(\mathbb{G}_{(r)}))_M$ satisfies the “tensor product property” (namely, that $\Pi(\tilde{D}(\mathbb{G}_{(r)}))_{M \otimes M'} = \Pi(\tilde{D}(\mathbb{G}_{(r)}))_M \cap \Pi(\tilde{D}(\mathbb{G}_{(r)}))_{M'}$) because $O(\mathbb{G}_{(r),K}) \xrightarrow{\sim} K((\underline{\mathfrak{g}}^{(r)})_{(1)} \times \mathbb{G}_{(r)})$ is not a map of Hopf algebras. In this section, we prove this tensor product property.

The following proposition of [7] is the key to our proof of Theorem 3.4 establishing the tensor product property for $M \mapsto \Pi(\tilde{D}(\mathbb{G}_{(r)}))_M$.

Proposition 3.1. [7, Prop 2.2] *Let V be a k -vector space and α, β, γ be pairwise commuting V -endomorphisms. Assume further α, β are p -nilpotent and γ is p^r -nilpotent in $\text{End}_k(V)$ for some $r \geq 1$. Then V is projective as a $k[u]/u^p$ -module where the action of u is given by α if and only if it is projective as a $k[v]/v^p$ module where the action of v is given by $\alpha + \beta\gamma$.*

We apply Proposition 3.1 to prove the following comparison of pull-backs along a given π -point of tensor products of A -modules with two different A -module structures determined by different coproducts on A .

Proposition 3.2. *Let A be a finite dimensional, local, commutative k -algebra with maximal ideal \mathfrak{m} satisfying the property that $a^p = 0$ for all $a \in \mathfrak{m}$. Assume that A is equipped with two A -linear coproducts $\Delta, \Delta' : A \rightarrow A \otimes A$ with the property that*

$$(3.2.1) \quad (\Delta(a) - a \otimes 1 - 1 \otimes a), (\Delta'(a) - a \otimes 1 - 1 \otimes a) \in \mathfrak{m} \otimes \mathfrak{m}, \quad \forall a \in \mathfrak{m}.$$

Let M, M' be A -modules and let $M \boxtimes M'$ be the $A \otimes A$ module given as the external tensor product. Then for any flat map $\alpha : k[t]/t^p \rightarrow A$, the pull-back of $M \boxtimes M'$ via $\Delta \circ \alpha$ is a projective $k[t]/t^p$ -module if and only if the pull-back of $M \boxtimes M'$ via $\Delta' \circ \alpha$ is a projective $k[t]/t^p$ -module.

More generally, let H be finite dimensional, local, commutative Hopf algebra with maximal ideal \mathfrak{m}_H such that $c^p = 0$ for all $c \in \mathfrak{m}_H$ and equip $A \otimes H$ with the coproducts $\Delta \otimes \Delta_H, \Delta' \otimes \Delta_H$. Consider $(A \otimes H)$ -modules M, M' , and let $M \boxtimes M'$ denote the $(A \otimes H) \otimes (A \otimes H)$ module given as the external tensor product. Then for any flat map $\alpha : k[t]/t^p \rightarrow A \otimes H$, the pull-back of $M \boxtimes M'$ via $(\Delta \otimes \Delta_H) \circ \alpha$ is a projective $k[t]/t^p$ module if and only if the pull-back of $M \boxtimes M'$ via $(\Delta' \otimes \Delta_H) \circ \alpha$ is a projective $k[t]/t^p$ -module.

Proof. We apply Proposition 3.1 to the k -vector space $V = M \otimes M'$ and consider triples of p -nilpotent elements in the image of $A \otimes A \rightarrow \text{End}_k(V)$ given by viewing V as the external tensor product $M \boxtimes M'$ (and thus an $A \otimes A$ module for the commutative k -algebra $A \otimes A$).

Consider an element $a \in \mathfrak{m} \subset A$ and write $\Delta(a) - \Delta'(a) = \sum_{i=1}^t b_i \otimes b'_i \in \mathfrak{m} \otimes \mathfrak{m}$. Observe that $\Delta(a) - \Delta'(a) \in A \otimes A$ acts on $V = M \boxtimes M'$ as the sum of the actions commuting elements $(b_i \otimes 1) \cdot (1 \otimes b'_i), 1 \leq i \leq t$ each of which is p -nilpotent.. We apply Proposition 3.1 with $V = M \boxtimes M'$ successively to the triples

$$\alpha = ((\Delta'(a) + \sum_{s=1}^i b_s \otimes b'_s), \quad \beta = (b_{i+1} \otimes 1), \quad \gamma = (1 \otimes b'_{i+1}))$$

with $i+1 \leq t$ to conclude that V is projective for $k[u]/u^p$ with u acting as $\Delta'(a) + \sum_{s=1}^i b_s \otimes b'_s$ if and only if V is projective for $k[v]/t^p$ with v acting as $\Delta'(a) + \sum_{s=1}^{i+1} b_s \otimes b'_s$. This establishes the first assertion.

We extend this argument to $A \otimes H$, a commutative algebra with coproducts $\Delta \otimes \Delta_H, \Delta' \otimes \Delta_H$. We denote by $\mathfrak{n} \subset A \otimes H$ the maximal ideal of $A \otimes H$ generated by $\mathfrak{m} \otimes 1, 1 \otimes \mathfrak{m}_H$, and observe that every element in \mathfrak{n} has p -th power 0. Moreover, $(\Delta \otimes \Delta_H) - (\Delta' \otimes \Delta_H) = (\Delta - \Delta') \otimes \Delta_H$ applied to $a \otimes 1 \in \mathfrak{n}$ lies in $\mathfrak{n} \otimes \mathfrak{n}$ whereas $(\Delta - \Delta') \otimes \Delta_H$ applied to $1 \otimes h \in 1 \otimes \mathfrak{m}_H$ is 0. Consequently,

$$((\Delta \otimes \Delta_H) - (\Delta' \otimes \Delta_H))(b) \in \mathfrak{n} \otimes \mathfrak{n}, \quad \forall b \in \mathfrak{n}.$$

Thus we may repeat the preceding argument with (A, \mathfrak{m}) replaced by $(A \otimes H, \mathfrak{n})$ to complete the proof. \square

Definition 3.3. Consider $O(\mathbb{G}_{(r)})$ -modules M, M' .

We denote by $M \otimes M'$ the $O(\mathbb{G}_{(r)})$ -module determined by the coproduct of $O(\mathbb{G}_{(r)})$ defined as by restriction of the coproduct of $\tilde{D}(\mathbb{G}_{(r)})$ (see Proposition 1.4).

We denote by $M \otimes_G M'$ the k -vector space $M \otimes M'$ equipped with the $O(\mathbb{G}_{(r)}) \simeq (k(\underline{\mathfrak{g}}^{(r)})_{(1)} \times \mathbb{G}_{(r)})$ -module structure determined by the coproduct for the group algebra $k(\underline{\mathfrak{g}}^{(r)})_{(1)} \times \mathbb{G}_{(r)}$ of the finite group scheme $(\underline{\mathfrak{g}}^{(r)})_{(1)} \times \mathbb{G}_{(r)}$.

In the following theorem, we prove the “tensor product property” for $\tilde{D}(\mathbb{G}_{(r)})$ -modules; namely, $M \mapsto \Pi(\tilde{D}(\mathbb{G}_{(r)}))_M$ sends tensor product of modules to intersection of the supports of its tensor factors.

Theorem 3.4. *Let (α_K, β_K) be a π -point pair of $\tilde{D}(\mathbb{G}_{(r)})$ as in Definition 2.1 and let M, M' be $O(\mathbb{G}_{(r)})$ -modules. Then $(\alpha_K + \beta_K)^*(M_K \otimes M'_K)$ is a free $K[t]/t^p$ -module if and only if $(\alpha_K + \beta_K)^*(M_K \otimes_G M'_K)$ is a free $K[t]/t^p$ -module. Thus,*

$$(3.4.1) \quad \Pi(\tilde{D}(\mathbb{G}_{(r)}))_{M \otimes M'} = \Pi(\tilde{D}(\mathbb{G}_{(r)}))_M \cap \Pi(\tilde{D}(\mathbb{G}_{(r)}))_{M'}.$$

Proof. We recall that any $\beta_K : K[t]/t^p \rightarrow K\mathbb{G}_{(r)}$ is equivalent to a map which factors through a map of group algebras induced by a map of finite group schemes $\mathbb{G}_{a(r),L} \rightarrow \mathbb{G}_{(r),L}$ [7, Prop 4.2]. By replacing K if necessary by a field extension K'/K , we may assume $K = L$. Thus when comparing restrictions along a given π -point pair, we may restrict to $K[\mathbb{G}_{(r+1)}/\mathbb{G}_{(r)}] \otimes K\mathbb{G}_{a(r)} \hookrightarrow O(\mathbb{G}_{(r)}) \otimes K$ and replace M, M' by their restrictions to $K[\mathbb{G}_{(r+1)}/\mathbb{G}_{(r)}] \otimes K\mathbb{G}_{a(r)}$.

We apply Proposition 3.2 with A equal to $K[\mathbb{G}_{(r+1)}/\mathbb{G}_{(r)}]$ and $H = K\mathbb{G}_{a(r)}$, and equip $A \otimes K\mathbb{G}_{a(r)}$ with the two coproduct structures obtained by restriction of the two coproduct structures on $O(\mathbb{G}_{(r)})$ discussed in Definition 3.3. As shown in [13, I.2.4], Δ, Δ' satisfy condition (3.2.1) of Proposition 3.2. In our application of Proposition 3.2, we take $\alpha : K[t]/t^p \rightarrow K[\mathbb{G}_{(r+1)}/\mathbb{G}_{(r)}] \otimes K\mathbb{G}_{a(r)}$ to be the composition of the coproduct $K[t]/t^p \rightarrow K[t]/t^p \otimes K[t]/t^p$, $t \mapsto t \otimes 1 + 1 \otimes t$ and the external tensor product $\alpha_K \otimes \beta_K : K[t]/t^p \rightarrow K[t]/t^p \otimes K[t]/t^p \rightarrow K[\mathbb{G}_{(r+1)}/\mathbb{G}_{(r)}] \otimes K\mathbb{G}_{a(r)}$.

By applying Proposition 3.2 with these values for A, H, α , we conclude that it suffices to prove the equality (3.4.1) after replacing the tensor product for $K[\mathbb{G}_{(r+1)}/\mathbb{G}_{(r)}] \otimes K\mathbb{G}_{a(r)}$ -modules (given by the restriction of the coproduct of $O(\mathbb{G}_{(r)})$) by the tensor product given by the restriction of the tensor product of $(\underline{\mathfrak{g}}^{(r)})_{(1)} \times \mathbb{G}_{(r)}$ -modules. To conclude the proof of equality (3.4.1), we apply the tensor product property for modules for the (infinitesimal) group scheme $G = (\underline{\mathfrak{g}}^{(r)})_{(1)} \times \mathbb{G}_{(r)}$ [8, Prop 3.2]; in other words, for $M \otimes_G M'$ in the notation of Definition 3.3. \square

Theorem 3.4 enables topological structures on the sets $\Pi(\tilde{D}(\mathbb{G}_{(r)}))$ and $\Pi(O(\mathbb{G}_{(r)}))$.

Corollary 3.5. *Defining a subset of $\Pi(\tilde{D}(\mathbb{G}_{(r)}))$ to be closed if and only if it is of the form $\Pi(\tilde{D}(\mathbb{G}_{(r)}))_M \subset \Pi(\tilde{D}(\mathbb{G}_{(r)}))$ for some finite dimensional $\tilde{D}(\mathbb{G}_{(r)})$ -module M defines a topology on $\Pi(\tilde{D}(\mathbb{G}_{(r)}))$. Similarly, defining a subset of $\Pi(O(\mathbb{G}_{(r)}))$ to be closed if and only if it is of the form $\Pi(O(\mathbb{G}_{(r)}))_M \subset \Pi(O(\mathbb{G}_{(r)}))$ for some finite dimensional $O(\mathbb{G}_{(r)})$ -module M defines a topology on $\Pi(O(\mathbb{G}_{(r)}))$.*

With these topologies, the bijection $\Pi_{\tilde{D}} : \Pi(\tilde{D}(\mathbb{G}_{(r)})) \rightarrow \Pi(O(\mathbb{G}_{(r)}))$ of Proposition 2.4 is a homeomorphism.

Proof. The asserted formulation of a topology on $\Pi(\tilde{D}(\mathbb{G}_{(r)}))$ is justified by Proposition 2.6(4) and Theorem 3.4. This applies as well to justify the topology on $\Pi(O(\mathbb{G}_{(r)}))$, since $i_O : O(\mathbb{G}_{(r)}) \rightarrow \tilde{D}(\mathbb{G}_{(r)})$ is a map of Hopf algebras. \square

$$4. \Psi_{\tilde{D}} : \Pi(\tilde{D}(\mathbb{G}_{(r)})) \rightarrow \mathbb{P}(H^\bullet(\tilde{D}(\mathbb{G}_{(r)}), k))$$

Cohomological support varieties can be readily defined for any finite dimension Hopf algebra H whose cohomology is finitely generated (i.e., such that $H^*(H, k)$ is a finitely generated k -algebra and that $H^*(H, M)$ is a finite $H^*(H, k)$ -module for any finite dimensional H -module). In this section, we exhibit in Proposition 4.6 a continuous map $\Psi_{\tilde{D}}$ from the π -point pair space $\Pi(\tilde{D}(\mathbb{G}_{(r)}))$ introduced in Section 2 to the (projectivized) cohomology support variety $\mathbb{P}(H^\bullet(\tilde{D}(\mathbb{G}_{(r)}), k))$. To do this, we utilize the subalgebra $O(\mathbb{G}_{(r)}) \subset \tilde{D}(\mathbb{G}_{(r)})$ and the related finite group scheme $\mathbb{G}_{(r)} \times (\underline{\mathfrak{g}}^{(r)})_{(1)}$, relying upon the isomorphism $\Pi(\mathbb{G}_{(r)} \times (\underline{\mathfrak{g}}^{(r)})_{(1)}) \xrightarrow{\sim} \mathbb{P}(H^\bullet(\mathbb{G}_{(r)} \times (\underline{\mathfrak{g}}^{(r)})_{(1)}, k)$.

Associated to a finite dimensional $\tilde{D}(\mathbb{G}_{(r)})$ -module M , we consider the annihilator ideal $I_M \subset H^\bullet(\tilde{D}(\mathbb{G}_{(r)}), k)$ of the $H^\bullet(\tilde{D}(\mathbb{G}_{(r)}), k)$ -module $Ext_{\tilde{D}(\mathbb{G}_{(r)}}^*(M, M)$ which equals the annihilator of $1 \in Ext_{\tilde{D}(\mathbb{G}_{(r)}}^*(M, M)$ which in turn equals the kernel of the graded map of k -algebras $H^\bullet(\tilde{D}(\mathbb{G}_{(r)}), k) \rightarrow Ext_{\tilde{D}(\mathbb{G}_{(r)}}^*(M, M)$. We shall consider the projective scheme $\mathbb{P}(H^\bullet(\tilde{D}(\mathbb{G}_{(r)}), k))$ whose (Zariski) points are (non-trivial) homogenous prime ideals of $H^\bullet(\tilde{D}(\mathbb{G}_{(r)}), k)$ and the reduced projective scheme $\mathbb{P}(H^\bullet(\tilde{D}(\mathbb{G}_{(r)}), k))_M$ whose points are (non-trivial) homogenous prime ideals of $H^\bullet(\tilde{D}(\mathbb{G}_{(r)}), k)$ containing I_M .

Consider the quotient map $\mathbb{G}_{(r+2)} \twoheadrightarrow (\mathbb{G}^{(r+1)})_{(1)}$, the image of $F^{r+1} : \mathbb{G}_{(r+2)} \rightarrow (\mathbb{G}^{(r+1)})_{(r+2)}$ whose kernel is $\mathbb{G}_{(r+1)}$. This quotient map determines the flat map of k -algebras

$$k[(\mathbb{G}^{(r+1)})_{(1)}] \simeq k[\mathbb{G}_{(r+2)}/\mathbb{G}_{(r+1)}] \rightarrow k[\mathbb{G}_{(r+2)}]$$

with the property that $k[\mathbb{G}_{(r+1)}] \simeq k[\mathbb{G}_{(r+2)}] \otimes_{k[\mathbb{G}_{(r+2)}/\mathbb{G}_{(r+1)}]} k$.

The following proposition is a summary of a central technique of [5].

Proposition 4.1. *The deformation $k[\mathbb{G}_{(r+2)}/\mathbb{G}_{(r+1)}] \rightarrow k[\mathbb{G}_{(r+2)}]$ of $k[\mathbb{G}_{(r+1)}]$ (parametrized by $k[\mathbb{G}_{(r+2)}/\mathbb{G}_{(r+1)}] \simeq k[(\mathbb{G}^{(r+1)})_{(1)}]$) determines a \mathbb{G} -equivariant map*

$$(4.1.1) \quad \mathfrak{g}^{(r+1)} \rightarrow H^2(k[\mathbb{G}_{(r+1)}], k).$$

The deformation $k[\mathbb{G}_{(r+2)}/\mathbb{G}_{(r+1)}] \# k\mathbb{G}_{(r)} \rightarrow k[\mathbb{G}_{(r+2)}] \# k\mathbb{G}_{(r)}$ (also parametrized by $k[\mathbb{G}_{(r+2)}/\mathbb{G}_{(r+1)}]$) provides a $\mathbb{G}_{(r)}$ -equivariant lifting of (4.1.1)

$$(4.1.2) \quad \sigma_{\tilde{D}} : \mathfrak{g}^{(r+1)} \rightarrow H^2(\tilde{D}(\mathbb{G}_{(r)}), k).$$

Consequently, we obtain a map of k -algebras

$$(4.1.3) \quad \tilde{\theta}_r : H^*(\mathbb{G}_{(r)}, k) \otimes S^\bullet(\mathfrak{g}^{(r+1)}[2]) \rightarrow H^*(\tilde{D}(\mathbb{G}_{(r)}), k)$$

defined as the product of the inflation map on $H^(\mathbb{G}_{(r)}, k)$ and the algebra extension of $\sigma_{\tilde{D}}$ on $S^\bullet(\mathfrak{g}^{(r+1)}[2])$.*

Proof. Each element of the tangent space at the identity of $(\mathbb{G}^{(r+1)})_{(1)}$ (naturally identified with $\mathfrak{g}^{(r+1)}$) corresponds to a map $k[(\mathbb{G}^{(r+1)})_{(1)}] \rightarrow k[\epsilon]/\epsilon^2$ of k -algebras. Extending the deformation $k[\mathbb{G}_{(r+2)}/\mathbb{G}_{(r+1)}] \rightarrow k[\mathbb{G}_{(r+2)}]$ along such a map determines a deformation of $k[\mathbb{G}_{(r+1)}]$ parametrized by $k[\epsilon]/\epsilon^2$. Thus, the map (4.1.1) is a consequence of the naturality of Gerstenhaber's bijection [12] between deformations of $k[\mathbb{G}_{(r+1)}]$ parametrized by $k[\epsilon]/\epsilon^2$ and the second Hochschild cohomology group $HH^2(k[\mathbb{G}_{(r+1)}])$ of $k[\mathbb{G}_{(r+1)}]$ together with the natural map $HH^2(k[\mathbb{G}_{(r+1)}]) \rightarrow H^2(k[\mathbb{G}_{(r+1)}], k)$ (see [5, Prop 3.4]). The map $\sigma_{\tilde{D}}$ of (4.1.2) follows exactly the same way. Moreover, the naturality of Gerstenhaber's bijection implies that $\sigma_{\tilde{D}}$ lifts (4.1.1).

The construction of $\tilde{\theta}_r : H^*(\mathbb{G}_{(r)}, k) \otimes S^\bullet(\mathfrak{g}^{(r+1)}[2]) \rightarrow H^*(\tilde{D}(\mathbb{G}_{(r)}), k)$ is implicit in the last statement of this proposition. \square

Finite generation of the cohomology algebra $H^*(D(\mathbb{G}_{(r)}), k)$ and the finiteness of $H^*(D(\mathbb{G}_{(r)}), M)$ as an $H^*(D(\mathbb{G}_{(r)}), k)$ -module (for a finite dimensional $D(\mathbb{G}_{(r)})$ -module M) was established in [5, Thm 5.3]. Using Proposition 4.1, we extend this finite generation property to $\tilde{D}(\mathbb{G}_{(r)})$ and $O(\mathbb{G}_{(r)})$ as stated below; the proof is a straight-forward adaptation of that of [5, Thm 5.3] which in turn is an adaptation of [11, Thm 1.1].

Theorem 4.2. *Consider the map $\tilde{\theta}_r : H^*(\mathbb{G}_{(r)}, k) \otimes S^\bullet(\mathfrak{g}^{(r+1)}[2]) \rightarrow H^*(\tilde{D}(\mathbb{G}_{(r)}), k)$ of (4.1.3).*

(1) *The composition*

$$(4.2.1) \quad \begin{aligned} i_O^* \circ \tilde{\theta}_r : H^*(\mathbb{G}_{(r)}, k) \otimes S^\bullet(\mathfrak{g}^{(r+1)}[2]) &\rightarrow H^*(\tilde{D}(\mathbb{G}_{(r)}), k) \\ &\rightarrow H^*(O(\mathbb{G}_{(r)}), k) \simeq H^\bullet(\mathbb{G}_{(r)}, k) \otimes H^*(k[(\mathbb{G}^{(r)})_{(1)}], k) \end{aligned}$$

is a split injection with $H^(\mathbb{G}_{(r)}, k) \otimes S^\bullet(\mathfrak{g}^{(r+1)}[2])$ -module complement*

$$H^*(\mathbb{G}_{(r)}, k) \otimes \Lambda^{>0}(\mathfrak{g}[1]) \otimes S^\bullet(\mathfrak{g}^{(r+1)}[2])$$

which consists entirely of nilpotent elements

(2) *For any finite dimensional $\tilde{D}(\mathbb{G}_{(r)})$ -module M , $H^*(\tilde{D}(\mathbb{G}_{(r)}), M)$ is a finitely generated $H^\bullet(\mathbb{G}_{(r)}, k) \otimes S^\bullet(\mathfrak{g}^{(r+1)}[2])$ -module (with action given by the restriction along $\tilde{\theta}_r$ of the natural action of $H^*(\tilde{D}(\mathbb{G}_{(r)}), k)$ on $H^*(\tilde{D}(\mathbb{G}_{(r)}), M)$).*

(3) *For any finite dimensional $O(\mathbb{G}_{(r)})$ -module M , $H^*(O(\mathbb{G}_{(r)}), M)$ is also a finitely generated $H^\bullet(\mathbb{G}_{(r)}, k) \otimes S^\bullet(\mathfrak{g}^{(r+1)}[2])$ -module.*

Proof. We utilize the spectral sequence

$$(4.2.2) \quad E_2^{s,t} = H^s(\mathbb{G}_{(r)}, H^t(k[G_{(r+1)}], k)) \Rightarrow H^{s+t}(\tilde{D}(\mathbb{G}_{(r)}), k),$$

which can be derived as a special case of Grothendieck's spectral sequence for the derived functors of a composition of left exact functors exactly as in [5, Prop 5.2].

The composition $i_O^* \circ \tilde{\theta}_r$ restricts to the identity on the tensor factor $H^*(\mathbb{G}_{(r)}, k)$ by the definition of $\tilde{\theta}_r$ and the fact that the composition $k\mathbb{G}_{(r)} \rightarrow O(\mathbb{G}_{(r)}) \rightarrow \tilde{D}(\mathbb{G}_{(r)})$ is the evident inclusion. The deformation construction leading to the definition of $\tilde{\theta}_r$ is designed to insure that the composition $S^\bullet(\mathfrak{g}^{(r+1)}[2]) \rightarrow H^*(\tilde{D}(\mathbb{G}_{(r)}), k) \rightarrow H^*(k[\mathbb{G}_{(r+1)}/\mathbb{G}_{(r)}], k)$ sends elements of $\mathfrak{g}^{(r+1)}[2]$ to elements of $H^2(k[\mathbb{G}_{(r+1)}/\mathbb{G}_{(r)}], k)$ which generate $H^*(k[\mathbb{G}_{(r+1)}/\mathbb{G}_{(r)}], k)$ modulo nilpotent elements. The identification of the restriction of the map $i_O^* \circ \tilde{\theta}_r$ to $1 \otimes S^\bullet(\mathfrak{g}^{(r+1)}[2])$ and the computation

of $H^*(k[(\mathbb{G}^{(r)})_{(1)}], k)$ are given in [5, §4], thereby establishing the asserted split surjection.

The fact that $H^*(\tilde{D}(\mathbb{G}_{(r)}), M)$ is a finitely generated $H^\bullet(\mathbb{G}_{(r)}, k) \otimes S^\bullet(\mathfrak{g}^{(r+1)}[2])$ -module for any finite dimensional $\tilde{D}(\mathbb{G}_{(r)})$ -module M arises from the natural pairing of (4.2.2) with the corresponding spectral sequence

$$(4.2.3) \quad E_2^{s,t}(M) = H^s(\mathbb{G}_{(r)}, H^t(k[G_{(r+1)}], M)) \Rightarrow H^{s+t}(\tilde{D}(\mathbb{G}_{(r)}), M)$$

with M coefficients (as in the proof of [5, Prop 5.2]). The same argument with $\tilde{D}(\mathbb{G}_{(r)})$ replaced by $O(\mathbb{G}_{(r)})$ implies that $H^*(O(\mathbb{G}_{(r)}), M)$ is a finitely generated $H^\bullet(\mathbb{G}_{(r)}, k) \otimes S^\bullet(\mathfrak{g}^{(r+1)}[2])$ -module. \square

Corollary 4.3. *The map of commutative k -algebras $\tilde{\theta}_r$ induces a finite, surjective map*

$$\tilde{\Theta}_r : \text{Spec } H^\bullet(\tilde{D}(\mathbb{G}_{(r)}), k) \rightarrow \text{Spec } H^\bullet(\mathbb{G}_{(r)}, k) \times \underline{\mathfrak{g}}^{*(r+1)}.$$

Consequently,

$$\dim(\text{Spec } H^\bullet(\tilde{D}(\mathbb{G}_{(r)}), k)) = \dim(\text{Spec } H^\bullet(\mathbb{G}_{(r)}, k) + \dim(\mathfrak{g})).$$

Proof. By Theorem 4.2, the composition $i^* \circ \tilde{\theta}_r$ is injective, so that $\tilde{\theta}_r$ is injective; this theorem also tells us that $\tilde{\theta}_r$ is a finite map. Consequently, $\tilde{\theta}_r$ is an integral extension which implies that $\tilde{\Theta}_r$ is finite and surjective (see [17, Thm 9.3]).

The determination of $\dim(\text{Spec } H^\bullet(\tilde{D}(\mathbb{G}_{(r)}), k))$ follows from the observation that $\dim(\text{Spec } H^\bullet(\mathbb{G}_{(r)}, k) \times \underline{\mathfrak{g}}^{*(r+1)}) = \dim(\text{Spec } H^\bullet(\mathbb{G}_{(r)}, k) + \dim(\mathfrak{g}))$. \square

We establish the following notation.

Notation 4.4. We denote by $\mathbb{P}(H^\bullet(\tilde{D}(\mathbb{G}_{(r)}), k))$ (respectively, $\mathbb{P}(H^\bullet(O(\mathbb{G}_{(r)}), k))$); respectively, $\mathbb{P}(H^\bullet(\mathbb{G}_{(r)}, k))$) the projective scheme associated to the commutative graded k -algebra $H^\bullet(\tilde{D}(\mathbb{G}_{(r)}), k)$ (resp., $H^\bullet(O(\mathbb{G}_{(r)}), k)$; resp., $H^\bullet(\mathbb{G}_{(r)}, k)$). Thus, the points of $\mathbb{P}(H^\bullet(\tilde{D}(\mathbb{G}_{(r)}), k))$ (resp., $\mathbb{P}(H^\bullet(O(\mathbb{G}_{(r)}), k)$; resp., $\mathbb{P}(H^\bullet(\mathbb{G}_{(r)}, k))$) are the non-trivial homogeneous prime ideals of $H^\bullet(\tilde{D}(\mathbb{G}_{(r)}), k)$ (resp., $H^\bullet(O(\mathbb{G}_{(r)}), k)$; resp., $H^\bullet(\mathbb{G}_{(r)}, k)$). For any finite dimensional $\tilde{D}(\mathbb{G}_{(r)})$ -module M , we denote by

$$\mathbb{P}(H^\bullet(\tilde{D}(\mathbb{G}_{(r)}), k))_M \subset \mathbb{P}(H^\bullet(\tilde{D}(\mathbb{G}_{(r)}), k)),$$

$$\mathbb{P}(H^\bullet(O(\mathbb{G}_{(r)}), k))_M \subset \mathbb{P}(H^\bullet(O(\mathbb{G}_{(r)}), k)),$$

$$\mathbb{P}(H^\bullet(\mathbb{G}_{(r)}, k))_M \subset \mathbb{P}(H^\bullet(\mathbb{G}_{(r)}, k))$$

the reduced closed subscheme given by the radicals of the ideals

$$\ker\{H^\bullet(\tilde{D}(\mathbb{G}_{(r)}), k) \rightarrow \text{Ext}_{\tilde{D}(\mathbb{G}_{(r)}}^*(M, M)\},$$

$$\ker\{H^\bullet(O(\mathbb{G}_{(r)}), k) \rightarrow \text{Ext}_{O(\mathbb{G}_{(r)})}^*(M, M)\},$$

$$\ker\{H^\bullet(\mathbb{G}_{(r)}, k) \rightarrow \text{Ext}_{\mathbb{G}_{(r)}}^*(M, M)\}$$

respectively.

The equivalence relation on π -point pairs (α_K, β_K) of $\tilde{D}(\mathbb{G}_{(r)})$ (and thus on $O(\mathbb{G}_{(r)})$) has been designed to enable the validity of the following proposition.

Proposition 4.5. Define $\Psi_O : \Pi(O(\mathbb{G}_{(r)})) \rightarrow \mathbb{P}(H^*(O(\mathbb{G}_{(r)}), k))$ to be the map of sets sending the equivalence class of $(\alpha_K, \beta_K) \in \Pi(O(\mathbb{G}_{(r)}))$ to

$$\ker\{(\alpha_K + \beta_K)^* : H^\bullet(O(\mathbb{G}_{(r)})_K, K) \rightarrow H^\bullet(K[t]/t^p, K)\} \cap H^*(O(\mathbb{G}_{(r)}), k)$$

in $\mathbb{P}(H^*(O(\mathbb{G}_{(r)}), k))$. A choice of isomorphism of k algebras $j : O(\mathbb{G}_{(r)}) \xrightarrow{\sim} k(\underline{\mathfrak{g}}^{(r)})_{(1)} \times \mathbb{G}_{(r)}$ as in Proposition 1.6 determines the vertical maps in the following commutative square of homeomorphisms

$$(4.5.1) \quad \begin{array}{ccc} \Pi(O(\mathbb{G}_{(r)})) & \xrightarrow{\Psi_O} & \mathbb{P}(H^\bullet(O(\mathbb{G}_{(r)}), k)) \\ \Pi_O \downarrow & & \downarrow \mathbb{P}_O \\ \Pi(\underline{\mathfrak{g}}^{(r)})_{(1)} \times \mathbb{G}_{(r)} & \xrightarrow{\Psi_G} & \mathbb{P}(H^\bullet(\underline{\mathfrak{g}}^{(r)})_{(1)} \times \mathbb{G}_{(r)}, k) \end{array}$$

whose lower horizontal map is the natural homeomorphism of C for G the finite group scheme $(\underline{\mathfrak{g}}^{(r)})_{(1)} \times \mathbb{G}_{(r)}$.

Moreover, for every finite dimensional $O(\mathbb{G}_{(r)})$ -module M , Ψ_O restricts to a bijection $\Psi_{O,M} : \Pi(O(\mathbb{G}_{(r)}))_M \xrightarrow{\sim} \mathbb{P}(H^\bullet(O(\mathbb{G}_{(r)}), k))_M$.

Proof. The definition of Ψ_O is compatible with the definition of $\Psi_G : \Pi(\underline{\mathfrak{g}}^{(r)})_{(1)} \times \mathbb{G}_{(r)} \rightarrow \mathbb{P}(H^\bullet(\underline{\mathfrak{g}}^{(r)})_{(1)} \times \mathbb{G}_{(r)}, k)$ (see $(\underline{\mathfrak{g}}^{(r)})_{(1)} \times \mathbb{G}_{(r)}$ and [7, Prop 3.5]), so that (4.5.1) commutes. The fact that Ψ_O is well defined follows from the fact that Ψ_G is well defined and that the maps Π_O, \mathbb{P}_O are bijections. The topology on $\Pi(\tilde{D}(\mathbb{G}_{(r)})) = \Pi(O(\mathbb{G}_{(r)}))$ introduced in Theorem 3.4 is equal to that induced by the bijection Π_O , so that this bijection is a homeomorphism. The bijection \mathbb{P}_O is a homeomorphism since the isomorphism j induces an isomorphism between the category of finite dimensional $O(\mathbb{G}_{(r)})$ -modules and the category of finite dimensional $k(\underline{\mathfrak{g}}^{(r)})_{(1)} \times \mathbb{G}_{(r)}$ -modules.

The fact that $\Psi_{O,M}$ is a bijection for any finite dimensional $O(\mathbb{G}_{(r)})$ -module follows from the fact that Ψ_O restricts to a bijection $\Psi_{O,M} : \Pi(\underline{\mathfrak{g}}^{(r)})_{(1)} \times \mathbb{G}_{(r)} \xrightarrow{\sim} \mathbb{P}(H^\bullet(\underline{\mathfrak{g}}^{(r)})_{(1)} \times \mathbb{G}_{(r)}, k)_M$. \square

The following proposition complements Proposition 4.5 while introducing $\Psi_{\tilde{D}}$.

Proposition 4.6. The composition $\Psi_{\tilde{D}} \equiv \mathbb{P}_{\tilde{D}} \circ \Psi_O \circ \Pi_{\tilde{D}}^{-1}$:

$$\Pi(\tilde{D}(\mathbb{G}_{(r)})) \xrightarrow{\sim} \Pi(O(\mathbb{G}_{(r)})) \xrightarrow{\sim} \mathbb{P}(H^\bullet(O(\mathbb{G}_{(r)}), k)) \rightarrow \mathbb{P}(H^\bullet(\tilde{D}(\mathbb{G}_{(r)}), k))$$

is continuous, fitting in the commutative square

$$(4.6.1) \quad \begin{array}{ccc} \Pi(\tilde{D}(\mathbb{G}_{(r)})) & \xrightarrow{\Psi_{\tilde{D}}} & \mathbb{P}(H^\bullet(\tilde{D}(\mathbb{G}_{(r)}), k)) \\ \simeq \uparrow \Pi_{\tilde{D}} & & \uparrow \mathbb{P}_{\tilde{D}} \\ \Pi(O(\mathbb{G}_{(r)})) & \xrightarrow{\sim} & \mathbb{P}(H^\bullet(O(\mathbb{G}_{(r)}), k)) \end{array}$$

with the properties that $\Pi_{\tilde{D}}$ and Ψ_O are homeomorphisms, and that the map $\mathbb{P}_{\tilde{D}}$ (induced by $i_O : O(\mathbb{G}_{(r)}) \rightarrow \tilde{D}(\mathbb{G}_{(r)})$) is a closed immersion.

For any finite dimensional $\tilde{D}(\mathbb{G}_{(r)})$ -module M , $\Psi_{\tilde{D}}$ restricts to

$$(4.6.2) \quad \Psi_{\tilde{D},M} : \Pi(\tilde{D})_M \hookrightarrow \mathbb{P}(H^\bullet(\tilde{D}(\mathbb{G}_{(r)}), k))_M.$$

Proof. By Proposition 2.4, $\Pi_{\tilde{D}}$ is a homeomorphism, whereas Ψ_O is a homeomorphism by Proposition 4.5. Theorem 4.2(1) implies that i_O^* is a finite map and is surjective modulo nilpotents which implies that \mathbb{P}_O is a closed embedding. Thus, $\Psi_{\tilde{D}}$ is also a closed embedding and thus is continuous.

To prove that $\Psi_{\tilde{D}}(\Pi(\tilde{D}(\mathbb{G}_{(r)}))_M)$ is contained in $\mathbb{P}(H^\bullet(\tilde{D}(\mathbb{G}_{(r)}), k))_M$, we consider some (α_K, β_K) representing a point of $\Pi(\tilde{D}(\mathbb{G}_{(r)}))_M$; in other words, assume that $(\alpha_K + \beta_K)^*(M_K)$ is not projective. Consider the following commutative square (4.6.3)

$$\begin{array}{ccc} H^\bullet(\tilde{D}(\mathbb{G}_{(r)}), k) & \xrightarrow{(\alpha_K + \beta_K)^*} & H^\bullet(K[t]/t^p, K) \\ \downarrow & & \downarrow \\ Ext_{\tilde{D}(\mathbb{G}_{(r)})}^*(M, M) & \longrightarrow & Ext_{K[t]/t^p}^*((\alpha_K + \beta_K)^*(M_K), (\alpha_K + \beta_K)^*(M_K)) \end{array}$$

whose horizontal maps are induced by restriction along $\alpha_K + \beta_K$. Since $(\alpha_K + \beta_K)^*(M_K)$ is not projective, the right vertical map is injective. A simple diagram chase then implies that $\ker\{(\alpha_K + \beta_K)^* : H^\bullet(\tilde{D}(\mathbb{G}_{(r)}), k) \rightarrow H^\bullet(K[t]/t^p, K)\}$ contains $\ker\{H^\bullet(\tilde{D}(\mathbb{G}_{(r)}), k) \rightarrow Ext_{K[t]/t^p}^*((\alpha_K, \beta_K)^*(M_K), (\alpha_K, \beta_K)^*(M_K))\}$, so that $\Psi_{\tilde{D}}$ applied to the class of (α_K, β_K) lies in $\mathbb{P}(H^\bullet(\tilde{D}(\mathbb{G}_{(r)}), k))_M$. \square

5. $\Pi(\tilde{D}(\mathbb{G}_{(r)}))$ FOR \mathbb{G} ADMITTING A QUASILOGARITHM

One missing aspect of our understanding of $\Psi_{\tilde{D}}$ in (4.6.1) is whether or not it is a homeomorphism. By Proposition 4.6, this is equivalent to whether or not $\mathbb{P}_{\tilde{D}}$ is surjective. With the added hypothesis that \mathbb{G} admits a quasilogarithm and that $p^{r+1} > 2\dim(\mathbb{G})$, we prove this surjectivity of $\mathbb{P}_{\tilde{D}}$ in Theorem 5.4. This enables us to conclude in Theorem 5.6 that $M \mapsto \Pi(\tilde{D}(\mathbb{G}_{(r)}))_M$ detects projectivity: a finite dimensional $\tilde{D}(\mathbb{G}_{(r)})$ -module M is projective if and only if $\Pi(\tilde{D}(\mathbb{G}_{(r)}))_M$ is empty.

We first recall the definition of a quasilogarithm for \mathbb{G} .

Definition 5.1. [14], [5, Defn 6.2] Let \mathbb{G} be a linear algebraic group over k . A quasilogarithm for \mathbb{G} is a \mathbb{G} -equivariant map of k -schemes

$$(5.1.1) \quad L : \mathbb{G} \rightarrow \text{Spec } S^\bullet(\mathfrak{g}^*) \cong \underline{\mathfrak{g}}$$

such that $L(e_{\mathbb{G}}) = 0$ and such that the differential $d_{e_{\mathbb{G}}}L : T_{e_{\mathbb{G}}}\mathbb{G} \rightarrow T_0\underline{\mathfrak{g}} = \mathfrak{g}$ is the identity.

We denote by $I_r \subset S^\bullet(\mathfrak{g}^*)$ the \mathbb{G} -stable ideal generated by $\{f^{p^r}, f \in \mathfrak{g}^*\}$.

Example 5.2. [2, Lem C3], [5, 6.4] If \mathbb{G} is a simple algebraic group for which p is very good, then \mathbb{G} admits a quasilogarithm. Furthermore any Borel subgroup of \mathbb{G} also admits a quasilogarithm. If \mathbb{U} is a unipotent subgroup of a semi-simple algebraic group \mathbb{G} which is normalized by a maximal torus and if p is greater than the nilpotent class of \mathbb{U} , then \mathbb{U} admits a quasilogarithm.

For us, a critical consequence of the condition that \mathbb{G} admits a quasilogarithm $L : \mathbb{G} \rightarrow \underline{\mathfrak{g}}$ is the following result, a summary of the discussion prior to and including Lemma 6.6 of [5].

Proposition 5.3. *Assume that \mathbb{G} is a linear algebraic group smooth over k which admits a quasilogarithm. The \mathbb{G} -equivariant map on coordinate algebras associated to the composition $L \circ i : \mathbb{G}_{(r)} \subset \mathbb{G} \rightarrow \underline{\mathfrak{g}}$ has the form*

$$S^\bullet(\mathfrak{g}^*) \rightarrow S^\bullet(\mathfrak{g}^*)/I_r \xrightarrow{\sim} k[\mathbb{G}_{(r)}].$$

Moreover, the isomorphism $S^\bullet(\mathfrak{g}^*)/I_r \xrightarrow{\sim} k[\mathbb{G}_{(r)}]$ induces an isomorphism of k -algebras

$$S^\bullet(\mathfrak{g}^*)/I_r \# k\mathbb{G}_{(r)} \xrightarrow{\sim} k[\mathbb{G}_{(r)}] \# k\mathbb{G}_{(r)} \simeq D(\mathbb{G}_{(r)}).$$

Similarly, L determines the isomorphism of k -algebras

$$S^\bullet(\mathfrak{g}^*)/I_{r+1} \# k\mathbb{G}_{(r)} \xrightarrow{\sim} k[\mathbb{G}_{(r+1)}] \# k\mathbb{G}_{(r)} \simeq \tilde{D}(\mathbb{G}_{(r)}).$$

These smash products algebras $S^\bullet(\mathfrak{g}^*)/I_r \# k\mathbb{G}_{(r)}$, $S^\bullet(\mathfrak{g}^*)/I_{r+1} \# k\mathbb{G}_{(r)}$ are associated to the coadjoint action of $\mathbb{G}_{(r)}$ on \mathfrak{g}^* . They both admit a $\mathbb{G}_{(r)}$ -equivariant grading, with \mathfrak{g}^* homogenous of degree 1 and $k\mathbb{G}_{(r)}$ homogeneous of degree 0.

Using the structure of a quasilogarithm on \mathbb{G} , we obtain the following strengthening of Corollary 4.3. The proof of this theorem is essentially that of [5, Thm 6.9] which gives the analogous result for $D(\mathbb{G}_{(r)})$.

Theorem 5.4. *Assume that \mathbb{G} is a linear algebraic group smooth over k which admits a quasilogarithm. Further assume that $p^{r+1} > 2\dim(\mathbb{G})$. Then the map*

$$\tilde{\theta}_r : H^*(\mathbb{G}_{(r)}, k) \otimes S^\bullet(\mathfrak{g}^{(r+1)}[2]) \rightarrow H^*(\tilde{D}(\mathbb{G}_{(r)}), k)$$

of Theorem 4.2 is a split injection with $H^*(\mathbb{G}_{(r)}, k) \otimes S^\bullet(\mathfrak{g}^{(r+1)}[2])$ -module complement consisting entirely of nilpotent elements.

Similarly, the map $i_O^* : H^*(\tilde{D}(\mathbb{G}_{(r)}), k) \rightarrow H^*(O(\mathbb{G}_{(r)}), k)$ has nilpotent kernel in $H^*(\tilde{D}(\mathbb{G}_{(r)}), k)$; and the image in $H^*(O(\mathbb{G}_{(r)}), k)$ has $H^*(\mathbb{G}_{(r)}, k) \otimes S^\bullet(\mathfrak{g}^{(r+1)}[2])$ -module complement consisting entirely of nilpotent elements.

Thus, $\mathbb{P}_O : \mathbb{P}(H^*(O(\mathbb{G}_{(r)}), k)) \rightarrow \mathbb{P}(H^*(\tilde{D}(\mathbb{G}_{(r)}), k))$ is a homeomorphism. Consequently, $\Psi_{\tilde{D}} : \mathbb{P}(\tilde{D}(\mathbb{G}_{(r)})) \rightarrow \mathbb{P}(H^*(\tilde{D}(\mathbb{G}_{(r)}), k))$ is also a homeomorphism

Proof. The statement that $\tilde{\theta}_r$ is a split injection is that of Theorem 4.2(1).

The gradings on $S^\bullet(\mathfrak{g}^*)/I_r \# k\mathbb{G}_{(r)}$, $S^\bullet(\mathfrak{g}^*)/I_{r+1} \# k\mathbb{G}_{(r)}$ (and thus also on $O(\mathbb{G}_{(r)})$) determine internal gradings on the spectral sequences of Theorem 4.2; we view these gradings as \mathbb{Z}/p^{r+1} -gradings. As in [5, Lem 6.7], the internal grading on $\mathfrak{g}^{(r+1)}[2] \subset H^2(k[\mathbb{G}_{(r+1)}], k)$ is p^{r+1} , hence $\equiv 0 \pmod{p^{r+1}}$ and the internal grading of $\mathfrak{g}[1] \subset H^1(k[\mathbb{G}_{(r+1)}], k)$ is 1.

We conclude that the map $\tilde{\theta}_r$ of Theorem 4.2 is an isomorphism onto the degree $\equiv 0$ portion of the cohomology of $\tilde{D}(\mathbb{G}_{(r)})$. As argued in the proof of [5, Thm 6.9], the hypothesis that $p^{r+1} > 2\dim(\mathbb{G})$ implies that this image has an $H^*(\mathbb{G}_{(r)}, k) \otimes S^\bullet(\mathfrak{g}^{(r+1)}[2])$ -module complement in $H^*(\tilde{D}(\mathbb{G}_{(r)}), k)$ which is contained in the nilradical.

We may repeat this argument with the spectral sequence (4.2.2) for $\tilde{D}(\mathbb{G}_{(r)})$ mapping to the spectral sequence

$$(5.4.1) \quad E_2^{s,t} = H^s(\mathbb{G}_{(r)}, H^t(k[G_{(r+1)}/\mathbb{G}_{(r)}], k)) \Rightarrow H^{s+t}(O(\mathbb{G}_{(r)}), k).$$

Thus, the composition $H^*(\mathbb{G}_{(r)}, k) \otimes S^\bullet(\mathfrak{g}^{(r+1)}[2]) \rightarrow H^*(\tilde{D}(\mathbb{G}_{(r)}), k) \rightarrow H^*(O(\mathbb{G}_{(r)}), k)$ is also a split injection with $H^*(\mathbb{G}_{(r)}, k) \otimes S^\bullet(\mathfrak{g}^{(r+1)}[2])$ -module complement consisting entirely of nilpotent elements. This readily implies the second statement

concerning $i_O^* : H^*(\tilde{D}(\mathbb{G}_{(r)}), k) \rightarrow H^*(O(\mathbb{G}_{(r)}), k)$. Consequently, \mathbb{P}_O is a bijection as well as a closed embedding, and thus is a homeomorphism. This, together with Proposition 4.6 implies that $\Psi_{\tilde{D}}$ is also a homeomorphism. \square

We “recall” the following projectivity criterion for a finite dimensional $\tilde{D}(\mathbb{G}_{(r)})$ -module in terms of cohomological support varieties.

Proposition 5.5. (see [6, Prop 1.5]) *Let M be a finitely generated $\tilde{D}(\mathbb{G}_{(r)})$ -module. Then M is projective if and only if $\mathbb{P}(H^\bullet(\tilde{D}(\mathbb{G}_{(r)}), k))_M = \emptyset$.*

Similarly, if M is a finitely generated $O(\mathbb{G}_{(r)})$ -module, then M is projective if and only if $\mathbb{P}(H^\bullet(O(\mathbb{G}_{(r)}), k))_M = \emptyset$.

Proof. As shown in [15] for any finite dimensional Hopf algebra with finitely generated cohomology, $\tilde{D}(\mathbb{G}_{(r)})$ and $O(\mathbb{G}_{(r)})$ are Frobenius algebras. This enables the proof of [6, Prop 1.5] to apply, even though that result was stated only for Hopf algebras which are restricted enveloping algebras. \square

Combining Theorem 5.4, with Proposition 5.5, we conclude that the support theory $M \mapsto \Pi(\tilde{D}(\mathbb{G}_{(r)}))_M$ provides a test for projectivity.

Theorem 5.6. *Assume the hypotheses on \mathbb{G} of Theorem 5.4 and consider a finite dimensional $\tilde{D}(\mathbb{G}_{(r)})$ -module M . Then M is projective as a $\tilde{D}(\mathbb{G}_{(r)})$ -module if and only if $\Pi(\tilde{D}(\mathbb{G}_{(r)}))_M = \emptyset$.*

Proof. Assume that M is projective as a $\tilde{D}(\mathbb{G}_{(r)})$ -module. Then the pull-back along $(\alpha_K + \beta_K)^*$ of M is projective (and thus free) as a $K[t]/t^p$ -module for any π -point pair (α_K, β_K) , so that $\Pi(\tilde{D}(\mathbb{G}_{(r)}))_M = \emptyset$.

Conversely, assume that $\Pi(\tilde{D}(\mathbb{G}_{(r)}))_M = \emptyset$. Since all the maps of the commutative square (4.6.1) are bijective (in fact, homeomorphisms), we conclude that $\mathbb{P}H^\bullet((\tilde{D}(\mathbb{G}_{(r)}), k))_M = \emptyset$. Thus, by Proposition 5.5, M is a projective $\tilde{D}(\mathbb{G}_{(r)})$ -module. \square

The arguments of Sections 4 and 5 apply to $\tilde{D}(G') \equiv k[\mathbb{G}_{(r+1)}] \# G' \subset \tilde{D}(\mathbb{G}_{(r)})$ with merely a change of notation, as we state explicitly in the following corollary (and use in the next section).

Corollary 5.7. *Let $G' \hookrightarrow \mathbb{G}_{(r)}$ be a subgroup scheme of $\mathbb{G}_{(r)}$. Theorems 4.2 and 5.4 remain valid upon replacing $i_O : O(\mathbb{G}_{(r)}) \rightarrow \tilde{D}(\mathbb{G}_{(r)})$ by*

$$i_{O'} : k[\mathbb{G}_{(r+1)}/\mathbb{G}_{(r)}] \# kG' \hookrightarrow k[\mathbb{G}_{(r+1)}] \# kG'.$$

Consequently, as in Proposition 5.5 and Theorem 5.6, a finite dimensional $k[\mathbb{G}_{(r+1)}] \# kG'$ -module M is projective if and only if its restriction along $i_{O'}$ is a projective $k[\mathbb{G}_{(r+1)}/\mathbb{G}_{(r)}] \# kG'$ -module.

6. RELATING $\Pi(\tilde{D}(\mathbb{G}_{(r)}))_M$ TO $\mathbb{P}(H^\bullet(\tilde{D}(\mathbb{G}_{(r)}), k))_M$

In this section, we verify for certain finite dimensional $\tilde{D}(\mathbb{G}_{(r)})$ -modules M that the natural inclusion $\Psi_{\tilde{D}, M} : \Pi(\tilde{D}(\mathbb{G}_{(r)}))_M \hookrightarrow \mathbb{P}(H^\bullet(\tilde{D}(\mathbb{G}_{(r)}), k))_M$ is a bijection and thus a homeomorphism.

We begin with an important class of examples (due to J. Carlson; see [3]) of modules constructed to have given cohomological support. Let $P_* \rightarrow k$ be a minimal

resolution of k as a $\tilde{D}(\mathbb{G}_{(r)})$ -module and define $\Omega_{\tilde{D}(\mathbb{G}_{(r)})}^n(k)$ for $n \geq 1$ to be the kernel of $P_{n-1} \rightarrow P_{n-2}$ (where P_{-1} is set equal to k). Recall that $\zeta \in H^n(\tilde{D}(\mathbb{G}_{(r)}), k)$ naturally corresponds to a map $\tilde{\zeta} : \Omega_{\tilde{D}(\mathbb{G}_{(r)})}^n(k) \rightarrow k$ in the stable module category $stmod(\tilde{D}(\mathbb{G}_{(r)}))$. The $\tilde{D}(\mathbb{G}_{(r)})$ -module L_ζ is defined by the short exact sequence of $\tilde{D}(\mathbb{G}_{(r)})$ -modules

$$(6.0.1) \quad 0 \rightarrow L_\zeta \rightarrow \Omega_{\tilde{D}(\mathbb{G}_{(r)})}^n(k) \xrightarrow{\tilde{\zeta}} k \rightarrow 0.$$

For $\xi \in H^n(O(\mathbb{G}_{(r)}), k)$ with corresponding map $\tilde{\zeta} : \Omega_{O(\mathbb{G}_{(r)})}^n(k) \rightarrow k$ in the stable module category $stmod(O(\mathbb{G}_{(r)}))$, we similarly define the $O(\mathbb{G}_{(r)})$ -module L_ξ by the short exact sequence of $O(\mathbb{G}_{(r)})$ -modules

$$0 \rightarrow L_\xi \rightarrow \Omega_{O(\mathbb{G}_{(r)})}^n(k) \xrightarrow{\tilde{\zeta}} k \rightarrow 0.$$

We restrict attention to $n = 2d > 0$. For such $n = 2d$, $\Omega_{k[t]/t^p}^{2d}(k)$ is stably isomorphic to k .

Proposition 6.1. *Consider the $\tilde{D}(\mathbb{G}_{(r)})$ -module L_ζ associated to a cohomology class $\zeta \in H^{2d}(\tilde{D}(\mathbb{G}_{(r)}), k)$. Then a π -point pair (α_K, β_K) of $\tilde{D}(\mathbb{G}_{(r)})$ satisfies the condition $(\alpha_K + \beta_K)^*(L_\zeta)$ is projective as a $K[t]/t^p$ -module if and only if $(\alpha_K + \beta_K)^*(\zeta) \neq 0 \in H^{2d}(K[t]/t^p, K)$.*

Consequently, $\Pi(\tilde{D}(\mathbb{G}_{(r)}))_{L_\zeta}$ consists of equivalence classes of π -point pairs (α_K, β_K) such that $(\alpha_K + \beta_K)^(\zeta) = 0 \in H^{2d}(K[t]/t^p, K)$.*

Proof. Carlson's identification of the support variety of L_ζ in the context of a group algebra of a finite group applies to $\tilde{D}(\mathbb{G}_{(r)})$, for his argument merely requires that the algebra $\tilde{D}(\mathbb{G}_{(r)})$ be a Frobenius algebra and that $\alpha_K + \beta_K : K[t]/t^p \rightarrow \tilde{D}(\mathbb{G}_{(r)})$ be flat. The key observation of Carlson's proof is that the pullback along $\alpha_K + \beta_K$ of (6.0.1) is split if and only if $(\alpha_K + \beta_K)^*(\zeta) \neq 0 \in H^{2d}(K[t]/t^p, K)$. \square

Using Proposition 4.6 and Theorem 5.4, we verify that $\Psi_{\tilde{D}, L_\zeta}$ is a homeomorphism.

Proposition 6.2. *Assume the hypotheses on \mathbb{G} of Theorem 5.4. Consider $\zeta \in H^\bullet(\tilde{D}(\mathbb{G}_{(r)}), k)$ and denote by ζ_O the restriction of ζ to $H^\bullet(O(\mathbb{G}_{(r)}), k)$. Then*

- (1) *The restriction of the $\tilde{D}(\mathbb{G}_{(r)})$ -module L_ζ to $O(\mathbb{G}_{(r)})$ is stably equivalent to L_{ζ_O} .*
- (2) *$\mathbb{P}i_O^*$ restricts to a bijection*

$$\mathbb{P}_O : \mathbb{P}(H^\bullet(O(\mathbb{G}_{(r)}), k))_{L_{\zeta_O}} \xrightarrow{\sim} \mathbb{P}(H^\bullet(\tilde{D}(\mathbb{G}_{(r)}), k))_{L_\zeta}.$$

- (3) *$\Psi_{\tilde{D}}$ restricts to a homeomorphism*

$$\Psi_{\tilde{D}, L_\zeta} : \Pi(\tilde{D}(\mathbb{G}_{(r)}))_{L_\zeta} \xrightarrow{\sim} \mathbb{P}(H^\bullet(\tilde{D}(\mathbb{G}_{(r)}), k))_{L_\zeta}.$$

Proof. To prove the first assertion, observe that the $O(\mathbb{G}_{(r)})$ -module $\Omega_{O(\mathbb{G}_{(r)})}^{2d}(k)$ is stably equivalent to the restriction of $\Omega_{\tilde{D}(\mathbb{G}_{(r)})}^{2d}(k)$ because the restriction of $\tilde{D}(\mathbb{G}_{(r)})$ -projectives to $O(\mathbb{G}_{(r)})$ are projective $O(\mathbb{G}_{(r)})$ -modules. Thus, restricting the short

exact sequence (6.0.1) of $\tilde{D}(\mathbb{G}_{(r)})$ -modules along $i_O : O(\mathbb{G}_{(r)}) \rightarrow \tilde{D}(\mathbb{G}_{(r)})$ takes the form of a distinguished triangle

$$L_{\zeta_O} \rightarrow \Omega_{O(\mathbb{G}_{(r)})}^{2d}(k) \xrightarrow{\zeta_O} k$$

in the stable module category $stmod(O(\mathbb{G}_{(r)}))$.

Using Proposition 6.1, we see that $\mathbb{P}(H^\bullet(O(\mathbb{G}_{(r)}), k))_{L_{\zeta_O}} \subset \mathbb{P}(H^\bullet(O(\mathbb{G}_{(r)}), k))$ is the zero locus of the homogeneous “function” ζ_O on $\mathbb{P}(H^\bullet(O(\mathbb{G}_{(r)}), k))$, and similarly that $\mathbb{P}(H^\bullet(\tilde{D}(\mathbb{G}_{(r)}), k))_{L_\zeta} \subset \mathbb{P}(H^\bullet(\tilde{D}(\mathbb{G}_{(r)}), k))$ is the zero locus of the homogeneous “function” ζ on $\mathbb{P}(H^\bullet(\tilde{D}(\mathbb{G}_{(r)}), k))$. The second assertion thus follows from the fact that $\zeta_O = i_O^*(\zeta)$ and the fact that \mathbb{P}_O is a homeomorphism by Theorem 5.4.

The last assertion now follows from the commutative square (4.6.1) of Proposition 4.6 and the fact that Ψ_O restricts to a bijection $\Psi_{O, L_{\zeta_O}} : \Psi(\tilde{O}(\mathbb{G}_{(r)}))_{L_{\zeta_O}} \xrightarrow{\sim} \mathbb{P}(H^\bullet(\tilde{O}(\mathbb{G}_{(r)}), k))_{L_{\zeta_O}}$ by Proposition 4.5. \square

The following proposition gives some insight into the topology on $\Pi(\tilde{D}(\mathbb{G}_{(r)}))$ beyond knowing that it is the topology inherited from $\mathbb{P}(H^\bullet(\tilde{D}(\mathbb{G}_{(r)}), k))$.

Proposition 6.3. *Assume the hypotheses on \mathbb{G} of Theorem 5.4 so that the maps of the commutative square (4.6.1) are homeomorphisms. Then*

$$\{\Pi(\tilde{D}(\mathbb{G}_{(r)}))_{L_\zeta}, \zeta \in H^\bullet(\tilde{D}(\mathbb{G}_{(r)}), k) \text{ homogeneous}\}$$

is a closed base for the topological space $\Pi(\tilde{D}(\mathbb{G}_{(r)}))$.

Proof. Observe that

$$\{\mathbb{P}(H^\bullet(O(\mathbb{G}_{(r)}), k))_{L_\xi}, \xi \in H^\bullet(O(\mathbb{G}_{(r)}), k) \text{ homogeneous}\}$$

is a closed base for the topology of $\mathbb{P}(H^\bullet(O(\mathbb{G}_{(r)}), k))$. Thus, the fact that $\mathbb{P}_O : \mathbb{P}(H^\bullet(O(\mathbb{G}_{(r)}), k)) \rightarrow \mathbb{P}(H^\bullet(\tilde{D}(\mathbb{G}_{(r)}), k))$ is a homeomorphism (by Theorem 5.4) together with Proposition 6.2(2) implies that

$$\{\mathbb{P}(H^\bullet(\tilde{D}(\mathbb{G}_{(r)}), k))_{L_\zeta}, \zeta \in H^\bullet(\tilde{D}(\mathbb{G}_{(r)}), k) \text{ homogeneous}\}$$

is a closed base for the topology of $\mathbb{P}(H^\bullet(\tilde{D}(\mathbb{G}_{(r)}), k))$. Since $\Psi_{\tilde{D}}$ is a homeomorphism, Proposition 6.2(3) implies that

$$\{\Pi(\tilde{D}(\mathbb{G}_{(r)}))_{L_\zeta}, \zeta \in H^\bullet(\tilde{D}(\mathbb{G}_{(r)}), k) \text{ homogeneous}\}$$

is a closed base for the topological space $\Pi(\tilde{D}(\mathbb{G}_{(r)}))$. \square

We next consider the class of $\tilde{D}(\mathbb{G}_{(r)})$ -modules given as the inflation along $\tilde{D}(\mathbb{G}_{(r)}) \rightarrow \mathbb{G}_{(r)}$ of finite dimensional $\mathbb{G}_{(r)}$ -modules (equivalently, as finite dimensional $\tilde{D}(\mathbb{G}_{(r)})$ -module whose restriction to $k[\mathbb{G}_{(r+1)}] \subset \tilde{D}(\mathbb{G}_{(r)})$ is trivial).

Proposition 6.4. *Assume the hypotheses on \mathbb{G} of Theorem 5.4. Let M be a finite dimensional $\tilde{D}(\mathbb{G}_{(r)})$ -module whose restriction to $k[\mathbb{G}_{(r+1)}] \subset \tilde{D}(\mathbb{G}_{(r)})$ is trivial. Then*

- (1) $\mathbb{P}_{O, M} : \mathbb{P}(H^\bullet(O(\mathbb{G}_{(r)}), k))_M \rightarrow \mathbb{P}(H^\bullet(\tilde{D}(\mathbb{G}_{(r)}), k))_M$ is a homeomorphism.
- (2) $\mathbb{P}(H^\bullet(O(\mathbb{G}_{(r)}), k))_M$ can be identified as the “geometric join” of $\text{Proj}(S^\bullet(\mathfrak{g}^{(r+1)}))$ and $\mathbb{P}(H^\bullet(\mathbb{G}_{(r)}), k)_M$.
- (3) $\Psi_{\tilde{D}, M} : \Pi(\tilde{D}(\mathbb{G}_{(r)}))_M \rightarrow \mathbb{P}(H^\bullet(\tilde{D}(\mathbb{G}_{(r)}), k))_M$ is also a homeomorphism.

Proof. We consider the commutative square

$$(6.4.1) \quad \begin{array}{ccc} H^\bullet(\tilde{D}(\mathbb{G}_{(r)}), k) & \xrightarrow{i_{\tilde{O}}^*} & H^\bullet(O(\mathbb{G}_{(r)}), k) \\ \downarrow & & \downarrow \\ H^\bullet(\tilde{D}(\mathbb{G}_{(r)}), M^* \otimes M) & \xrightarrow{i_{\tilde{O}}^*} & H^\bullet(O(\mathbb{G}_{(r)}), M^* \otimes M). \end{array}$$

We observe that there are multiplicative spectral sequences for $M^* \otimes M$ replacing k corresponding to the spectral sequences (4.2.2) and (5.4.1). Equipping $M^* \otimes M$ with internal grading degree 0, (6.4.1) determines a commutative square of multiplicative spectral sequences with internal grading whose E_2 -page has the form (6.4.2)

$$(6.4.2) \quad \begin{array}{ccc} H^s(\mathbb{G}_{(r)}, H^t(k[\mathbb{G}_{(r+1)}], k)) & \longrightarrow & H^s(\mathbb{G}_{(r)}, H^t(k[\mathbb{G}_{(r+1)}/\mathbb{G}_{(r)}], k)) \\ \downarrow & & \downarrow \\ H^s(\mathbb{G}_{(r)}, H^t(k[\mathbb{G}_{(r+1)}], k) \otimes M^* \otimes M) & \longrightarrow & H^s(\mathbb{G}_{(r)}, H^t(k[\mathbb{G}_{(r+1)}/\mathbb{G}_{(r)}], k) \otimes M^* \otimes M). \end{array}$$

The restriction map $H^*(k[\mathbb{G}_{(r+1)}], k) \rightarrow H^*(k[\mathbb{G}_{(r+1)}/\mathbb{G}_{(r)}], k)$ has nilpotent kernel and cokernel which are nilpotent, $\mathbb{G}_{(r)}$ -summands of $H^*(k[\mathbb{G}_{(r+1)}/\mathbb{G}_{(r)}], k)$. Thus, arguing as in the proof of Theorem 5.4, classes at the E_2 -level with internal grading not congruent to 0 modulo p^{r+1} are torsion; moreover, the lower horizontal arrow of (6.4.2) restricted to internal degrees congruent to 0 modulo p^{r+1} has nilpotent kernel and cokernel which are nilpotent, $\mathbb{G}_{(r)}$ -summands.

Since \mathbb{P}_O is a homeomorphism, to prove (1) it suffices to verify that if $\mathfrak{p} \subset H^\bullet(O(\mathbb{G}_{(r)}), k)$ is a homogeneous prime ideal whose inverse image $\mathfrak{q} \equiv i_{\tilde{O}}^{-1}(\mathfrak{p})$ contains the kernel of the left vertical map of (6.4.1), then \mathfrak{p} contains the kernel of the right vertical map. The validity of this statement is equivalent to the same statement of the commutative square obtained from (6.4.1) by replacing the lower horizontal arrow by its associated map on reduced algebras (i.e., by dividing out be the nilradicals of the domain and range of this map). By the preceding discussion, this latter map is an isomorphism. Thus, having made this replacement, the required statement about $i_{\tilde{O}}^{-1}(\mathfrak{p})$ is immediate.

Assertion (2) follows from the familiar interpretation of the projectivization of the product of two affine varieties.

Assertion (3) follows from the commutativity of (4.6.1) and assertion (1). \square

We consider a third class of finite dimensional $\tilde{D}(\mathbb{G}_{(r)})$ -modules, those with the property that their restrictions to $k[\mathbb{G}_{(r+1)}]$ are projective. We first recall the following detection theorem of A. Suslin whose formulation we quote from [7, Thm 4.10].

Theorem 6.5. [22] *Let G be a finite group scheme, Λ a unital associative G -algebra, and $\zeta \in H^\bullet(G, k)$ be a homogeneous cohomology class of even degree. Then ζ is nilpotent if and only if ζ_K restricts to a nilpotent class in the cohomology of every quasi-elementary subgroup scheme $\mathcal{E}_K \subset G_K$ for any field extension K/k .*

Let M be a finite dimensional $\tilde{D}(\mathbb{G}_{(r)})$ -module with the property that the restriction of M to $k[\mathbb{G}_{(r+1)}]$ is projective. If M satisfies this property, then the spectral sequences (4.2.2) and (4.2.3) degenerate so that the edge homomorphisms

provide the isomorphisms of algebras

$$(6.5.1) \quad \begin{aligned} H^*(\mathbb{G}_{(r)}, H^0(k[\mathbb{G}_{(r+1)}], M^* \otimes M)) &\xrightarrow{\sim} H^*(\tilde{D}(\mathbb{G}_{(r)}), M^* \otimes M), \\ H^*(\mathbb{G}_{(r)}, H^0(k[\mathbb{G}_{(r+1)}/\mathbb{G}_{(r)}], M^* \otimes M)) &\xrightarrow{\sim} H^*(O(\mathbb{G}_{(r)}), M^* \otimes M). \end{aligned}$$

We use (6.5.1) in conjunction with Theorem 6.5 to prove that

$$\Psi_{\tilde{D}, M} : \Pi(\tilde{D}(\mathbb{G}_{(r)}))_M \rightarrow \mathbb{P}(H^\bullet(\tilde{D}(\mathbb{G}_{(r)}), k))_M$$

is a homeomorphism for these modules.

Proposition 6.6. *Assume the hypotheses on \mathbb{G} of Theorem 5.4. Let M be a finite dimensional $\tilde{D}(\mathbb{G}_{(r)})$ -module with the property that the restriction of M to $k[\mathbb{G}_{(r+1)}]$ is projective. Then the restriction of $\Psi_{\tilde{D}}$ is a homeomorphism*

$$\Psi_{\tilde{D}, M} : \Pi(\tilde{D}(\mathbb{G}_{(r)}))_M \xrightarrow{\sim} \mathbb{P}(H^\bullet(\tilde{D}(\mathbb{G}_{(r)}), k))_M.$$

Proof. Let $\Lambda(M)$ denote unital associative $\mathbb{G}_{(r)}$ -algebra $H^0(k[\mathbb{G}_{(r+1)}], M^* \otimes M)$ and let $\Lambda_O(M)$ denote $H^0(k[\mathbb{G}_{(r+1)}/\mathbb{G}_{(r)}], M^* \otimes M)$. Using the isomorphisms (6.5.1), we rewrite (6.4.1) as

$$(6.6.1) \quad \begin{array}{ccc} H^\bullet(\tilde{D}(\mathbb{G}_{(r)}), k) & \xrightarrow{i_O^*} & H^\bullet(O(\mathbb{G}_{(r)}), k) \\ \downarrow & & \downarrow \\ H^\bullet(\mathbb{G}_{(r)}, \Lambda(M)) & \xrightarrow{i_O^*} & H^\bullet(\mathbb{G}_{(r)}, \Lambda_O(M)). \end{array}$$

We set I_M to be the kernel of the left vertical map and J_M the kernel of the right vertical map of (6.6.1).

Since we know that $\Psi_{\tilde{D}, M}$ is well defined and injective and that $\Psi_{\tilde{D}}$ is bijective, to prove the proposition we must show for any homogeneous prime ideal $\mathfrak{q} \subset H^\bullet(O(\mathbb{G}_{(r)}), k)$ with inverse image $\mathfrak{p} \subset H^\bullet(\tilde{D}(\mathbb{G}_{(r)}), k)$ that $J_M \subset \mathfrak{q}$ (i.e., \mathfrak{p} is a point of $\mathbb{P}(H^\bullet(O(\mathbb{G}_{(r)}), k))_M$) whenever $I_M \subset \mathfrak{p}$. Assume that $\zeta \in \mathfrak{p} \subset H^\bullet(\tilde{D}(\mathbb{G}_{(r)}), k)$ is a homogeneous element which is not in I_M , so that its image $\bar{\zeta} \in H^\bullet(\mathbb{G}_{(r)}, \Lambda(M))$ is not nilpotent. Using Theorem 6.5, let $\gamma : \mathbb{G}_{a(r), K} \rightarrow \mathbb{G}_{(r), K}$ be a 1-parameter subgroup with the property that $\gamma^*(\bar{\zeta}) \in H^\bullet(\mathbb{G}_{a(r), K}, \gamma^*(\Lambda(M_K)))$ is not nilpotent.

In other words, $\gamma^*(\zeta) \in H^\bullet(K[\mathbb{G}_{(r+1)}] \#_{\mathbb{G}_{a(r)}} k)$ has image $\overline{\gamma^*(\zeta)} \in Ext_{K[\mathbb{G}_{(r+1)}] \#_{\mathbb{G}_{a(r), K}}}^*(\gamma^*(M_K), \gamma^*(M_K))$ which is not nilpotent. By Corollary 5.7, the image of $\overline{\gamma^*(\zeta)}$ in $Ext_{K[\mathbb{G}_{(r+1)}/\mathbb{G}_{(r)}] \#_{\mathbb{G}_{a(r), K}}}^*(\gamma^*(M_K), \gamma^*(M_K))$ is not nilpotent as well. Consequently, $i_O^*(\zeta)$ does not lie in J_M . Thus, we conclude if \mathfrak{p} contains I_M then \mathfrak{q} must contain J_M . □

7. FURTHER REMARKS

We begin this final section with a natural question.

Question 7.1. *Assume that \mathbb{G} satisfies the conditions of Theorem 5.4. Can one extend the results of Section 6 to prove that*

$$\Psi_{\tilde{D}(\mathbb{G}_{(r)}), M} : \Pi(\tilde{D}(\mathbb{G}_{(r)}))_M \rightarrow \mathbb{P}(H^\bullet(\tilde{D}(\mathbb{G}_{(r)}), k))_M$$

is a homeomorphism for all finite dimensional $\tilde{D}(\mathbb{G}_{(r)})$ -modules?

We proceed to briefly introduce Hopf subalgebras $\tilde{D}^{(s)}(\mathbb{G}_{(r)}) \subset \tilde{D}(\mathbb{G}_{(r)})$ (generalizing $O(\mathbb{G}_{(r)}) \subset \tilde{D}(\mathbb{G}_{(r)})$) and quotient maps of Hopf algebras $k[\mathbb{G}_{r+s}] \twoheadrightarrow D(\mathbb{G}_{(r)})$ (generalizing $\tilde{D}(\mathbb{G}_{(r)}) \twoheadrightarrow D(\mathbb{G}_{(r)})$). These generalizations behave much as the algebras we have consider in previous sections.

Proposition 7.2. *The \mathbb{G} -equivariant quotient map $\mathbb{G}_{(r+1)} \twoheadrightarrow \mathbb{G}_{(r+1)}/\mathbb{G}_{(1)} \simeq (\mathbb{G}^{(1)})_{(r)}$ determines a Hopf algebra embedding of smash products*

$$(7.2.1) \quad i_D : \tilde{D}^{(1)}(\mathbb{G}_{(r)}) \equiv k[(\mathbb{G}^{(1)})_{(r)}] \# k\mathbb{G}_{(r)} \hookrightarrow k[\mathbb{G}_{(r+1)}] \# k\mathbb{G}_{(r)} \equiv \tilde{D}(\mathbb{G}_{(r)}).$$

This is a map of free (right) $O(\mathbb{G}_{(r)})$ -modules

The composition

$$\tilde{D}^{(1)}(\mathbb{G}_{(r)}) \hookrightarrow \tilde{D}(\mathbb{G}_{(r)}) \twoheadrightarrow D(\mathbb{G}_{(r)})$$

is the map $F \# id : k[(\mathbb{G}^{(1)})_{(r)}] \# k\mathbb{G}_{(r)} \rightarrow k[\mathbb{G}_{(r)}] \# k\mathbb{G}_{(r)}$, where F is the Frobenius map.

Proof. As in the justification of (1.1.2), the map of (7.2.1) is an embedding of Hopf algebras. We choose a k -linear section $\sigma : k[\mathbb{G}_{(r)}/\mathbb{G}_{(1)}] \rightarrow k[\mathbb{G}_{(r+1)}/\mathbb{G}_{(1)}]$ of the quotient map induced by the embedding $\mathbb{G}_{(r)}/\mathbb{G}_{(1)} \subset \mathbb{G}_{(r+1)}/\mathbb{G}_{(1)}$ and consider the

$$k[(\mathbb{G}_{(r+1)}/\mathbb{G}_{(1)})/(\mathbb{G}_{(r)}/\mathbb{G}_{(1)})] \simeq k[\mathbb{G}_{(r+1)}/\mathbb{G}_{(r)}] \simeq k[(\mathbb{G}^{(r)})_{(1)}]$$

bilinear map

$$k[(\mathbb{G}^{(1)})_{(r-1)}] \otimes k[(\mathbb{G}^{(r)})_{(1)}] \rightarrow k[(\mathbb{G}^{(1)})_{(r)}]$$

analogous to (1.5.1). Since $k[(\mathbb{G}^{(1)})_{(r-1)}] \otimes 1$ is central in $\tilde{D}^{(1)}(\mathbb{G}_{(r)})$, we conclude as in the proof of Proposition 1.5 that this provides $\tilde{D}^{(1)}(\mathbb{G}_{(r)})$ with the structure of a free (right) $O(\mathbb{G}_{(r)})$ -module.

To verify that the asserted composition equals $F \# id$, we use the identification $\mathbb{G}_{(r+1)}/\mathbb{G}_{(1)} \simeq (\mathbb{G}_{(r)})^{(1)}$. \square

We can view $\tilde{D}^{(1)}(\mathbb{G}_{(r)})$ as a twisted form of $D(\mathbb{G}_{(r)})$ for which the action of $\mathbb{G}_{(r)}$ on $k[(\mathbb{G}^{(1)})_{(r)}]$ equals the Frobenius twist of the coadjoint action of $\mathbb{G}_{(r)}$ on $k[\mathbb{G}_{(r)}]$.

The proof of the following theorem is a basically identical to the proofs given in previous sections of the corresponding properties for $\tilde{D}(\mathbb{G}_{(r)})$ -modules.

Proposition 7.3. *For any $\tilde{D}^{(1)}(\mathbb{G}_{(r)})$ -module M , we can define $M \mapsto \Pi(\tilde{D}^{(1)}(\mathbb{G}_{(r)}))_M$ in strict analogy with $M \mapsto \Pi(\tilde{D}(\mathbb{G}_{(r)}))_M$. So defined, π -point pairs of $\tilde{D}^{(1)}(\mathbb{G}_{(r)})$ are π -point pairs of $O(\mathbb{G}_{(r)})$ with the equivalence relation determined by consideration of finite dimensional $\tilde{D}^{(1)}(\mathbb{G}_{(r)})$ -modules. For any finite dimensional $\tilde{D}^{(1)}(\mathbb{G}_{(r)})$ -module M , we define $\Pi(\tilde{D}^{(1)}(\mathbb{G}_{(r)}))_M \subset \Pi(\tilde{D}^{(1)}(\mathbb{G}_{(r)}))$ to consist of those equivalence classes of π -point pairs (α_K, η_K) with the property that $(\alpha_K + \beta_K)^*(M_K)$ is not projective.*

There is a natural continuous map $\Psi_{\tilde{D}^{(1)}(\mathbb{G}_{(r)})} : \Pi(\tilde{D}^{(1)}(\mathbb{G}_{(r)})) \rightarrow \mathbb{P}(H^\bullet(\tilde{D}^{(1)}(\mathbb{G}_{(r)}), k))$. Assuming that \mathbb{G} admits a quasilogarithm and $p^{r+1} > 2\dim(\mathbb{G})$, this map is a homeomorphism and restricts to a homeomorphism

$$\Psi_{\tilde{D}^{(1)}(\mathbb{G}_{(r)}, M) : \Pi(\tilde{D}^{(1)}(\mathbb{G}_{(r)}))_M \xrightarrow{\sim} \mathbb{P}(H^\bullet(\tilde{D}^{(1)}(\mathbb{G}_{(r)}), k)_M$$

provided that M is in one of the classes analogous to those classes of $\tilde{D}(\mathbb{G}_{(r)})$ -modules considered in Section 6.

Remark 7.4. The triple of Hopf algebras over k

$$O(\mathbb{G}_{(r)}) \hookrightarrow \tilde{D}^{(1)}(\mathbb{G}_{(r)}) \hookrightarrow \tilde{D}(\mathbb{G}_{(r)})$$

can be refined to a sequence of embeddings of Hopf algebras which are free over $O(\mathbb{G}_{(r)})$

$$O(\mathbb{G}_{(r)}) \equiv D^{(r)}(\mathbb{G}_{(r)}) \hookrightarrow \cdots \hookrightarrow \tilde{D}^{(1)}(\mathbb{G}_{(r)}) \hookrightarrow \tilde{D}^{(0)}(\mathbb{G}_{(r)}) \equiv \tilde{D}(\mathbb{G}_{(r)}),$$

where

$$\tilde{D}^{(s)}(\mathbb{G}_{(r)}) \equiv k[(\mathbb{G}_{(r+1-s)}^{(s)})] \# k\mathbb{G}_{(r)}, \quad 0 \leq s \leq r.$$

One readily checks that replacing $\tilde{D}^{(1)}(\mathbb{G}_{(r)})$ in Proposition 7.3 by $\tilde{D}^{(s)}(\mathbb{G}_{(r)})$ gives analogous properties for $\tilde{D}^{(s)}(\mathbb{G}_{(r)})$ -modules.

Remark 7.5. For any $s \geq 1$, we define the Hopf subalgebra

$$k[\mathbb{G}_{(r+s)}] \# k\mathbb{G}_{(r)} \subset k[\mathbb{G}_{(r+s)}] \# k\mathbb{G}_{(r+s)} \equiv D(\mathbb{G}_{(r+s)}).$$

This Hopf algebra admits a quotient map of Hopf algebras

$$k[\mathbb{G}_{(r+s)}] \# k\mathbb{G}_{(r)} \twoheadrightarrow D(\mathbb{G}_{(r)})$$

determined by the quotient map $k[\mathbb{G}_{(r+s)}] \twoheadrightarrow k[\mathbb{G}_{(r+s)}] \twoheadrightarrow k[\mathbb{G}_{(r)}]$ as in (1.4.1). We obtain a sequence of quotient Hopf algebras

$$\cdots \twoheadrightarrow k[\mathbb{G}_{(r+s)}] \# k\mathbb{G}_{(r)} \twoheadrightarrow k[\mathbb{G}_{(r+s-1)}] \# k\mathbb{G}_{(r)} \twoheadrightarrow \cdots \twoheadrightarrow \tilde{D}(\mathbb{G}_{(r)}) \twoheadrightarrow D(\mathbb{G}_{(r)}).$$

We leave to the reader to check that our results for $\tilde{D}(\mathbb{G}_{(r)})$ -modules extend to $k[\mathbb{G}_{(r+s)}] \# k\mathbb{G}_{(r)}$ for $s \geq 1$.

Remark 7.6. We recall that a theorem of Pevtsova, Suslin, and the author [10, Thm 4.10] asserts that if M is a finite dimensional G -module for a finite group scheme G over k and if $\alpha_K : K[t]/t^p \rightarrow G_K$ is a π -point of G at which the Jordan type $\alpha_K^*(M_K)$ is maximal among all Jordan types $\beta_L^*(M_L)$ as $\beta_L : L[t]/t^p \rightarrow G_L$ varies among all π -points of G , then the Jordan type of $\alpha_K^*(M_K)$ equals the Jordan type of $\alpha'_K^*(M_K)$ whenever the π -point α'_K is equivalent to α_K .

This immediately applies to $O(\mathbb{G}_{(r)})$ -modules, since $O(\mathbb{G}_{(r)})$ is isomorphic as a k -algebra to the group algebra of a finite group scheme as seen in Proposition 1.6. By the definition of the equivalence relation on π -point pairs of $\tilde{D}(\mathbb{G}_{(r)})$, this remains valid for a $\tilde{D}(\mathbb{G}_{(r)})$ -module M whose maximal Jordan type is that of a projective module.

Question 7.7. *Does this “independence of representative of equivalence class” remain valid for any finite dimensional $\tilde{D}(\mathbb{G}_{(r)})$ -module M ? Namely, if the equivalence class of a π -point pair of $\tilde{D}(\mathbb{G}_{(r)})$ contains some representative (α_K, β_K) at which $(\alpha_K + \beta_K)^*(M_K)$ has maximal Jordan type, then does $(\alpha'_L + \beta_L)^*(M_L)$ have this same Jordan type for any $(\alpha'_K + \beta'_K) \sim (\alpha_K + \beta_K)$?*

REFERENCES

- [1] G. Avrunin and L. Scott, *Quillen stratification for modules*, Invent. Math. **66** (1982), 277 – 286.
- [2] R. Bezrukavnikov, D. Kazhdan, and Yarshavsky, *On the depth r Bernstein projector*, Selecta Math **22** (2016), 2271 – 2311.
- [3] J. Carlson, *The varieties and the cohomology ring of a module*, J. Algebra **85** (1983), 101-143.

- [4] J. Carlson, E. Friedlander and J. Pevtsova, *Modules of constant Jordan type*, Journal für die reine und angewandte Mathematik **614** (2008), 191 - 234.
- [5] E. Friedlander and C. Negron, *Cohomology of Drinfeld doubles of some infinitesimal group schemes*, Algebra Number Theory **12** (2018), 1281 - 1309.
- [6] E. Friedlander and B. Parshall, *Geometry of p -Unipotent Lie Algebras*, J. Algebra **109** (1987), 25 - 45.
- [7] E. Friedlander and J. Pevtsova, *Representation-theoretic support spaces for finite group schemes*, Amer. J. Math. **127** (2005), 379 - 420.
- [8] E. Friedlander and J. Pevtsova, *Π -supports for modules for finite group schemes*, Duke. Math. J. **139** (2007), 317-368.
- [9] E. Friedlander and J. Pevtsova, *Generalized support varieties for finite group schemes*, Documenta - Extra volume Suslin (2010), 197 - 222.
- [10] E. Friedlander, J. Pevtsova, and A. Suslin, *Generic and maximal Jordan types*, Invent. Math **168** (2007), 485 - 522.
- [11] E. Friedlander and A. Suslin, *Cohomology of finite group schemes over a field*, Invent. Math **127** (1997), 209 - 270.
- [12] M. Gerstenhaber, *On the deformation of rings and algebras*, Ann. of Math **79** (1964), 59 - 103.
- [13] J.C. Jantzen, *Representations of Algebraic groups. Second edition*, Math Surveys and Monographs **107**, AMS (2003).
- [14] D. Kazhdan and Y. Varshavsky, *Endoscopic decomposition of certain depth zero representations*, Studies in Lie theory, Progr. Math. **243**, 223 - 301.
- [15] R. Larson and M. Sweedler, *An associative orthogonal bilinear form for Hopf algebras*, Amer J. Math. **91** (1969), 75 - 94.
- [16] S. Majid, *Doubles of quasitriangular Hopf algebras*, Comm. Alg **19** (1991), 3061 -3073.
- [17] H. Matsumura, *Commutative Algebra*, Benjamin, N.Y (1970).
- [18] S. Montgomery, *Hopf algebras and their actions on rings*, CBMS **82** (1993).
- [19] D. Quillen, *The spectrum of an equivariant cohomology ring: I*, Ann. of Math **94** (1971), 549-572.
- [20] D. Quillen, *The spectrum of an equivariant cohomology ring: II*, Ann. of Math **94** (1971), 573-602.
- [21] D.E. Radford, *Minimal quasi-triangular Hopf algebras*, J. Algebra **157** (1991), 283 - 315.
- [22] A. Suslin, *Detection theorem for finite group schemes*, J.P.A.A. **206** (2006), 189 - 221.
- [23] A. Suslin, E. Friedlander, and C. Bendel *Support varieties for infinitesimal group schemes*, J. Amer. Math. Soc. **10** (1997), 729-759.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF SOUTHERN CALIFORNIA, LOS ANGELES, CA
Email address: ericmf@usc.edu