

Robust Principal Component Analysis: A Median of Means Approach

Debolina Paul¹, Saptarshi Chakraborty², and Swagatam Das^{*3}

¹Indian Statistical Institute, Kolkata, India

²Department of Statistics, University of California, Berkeley

³Electronics and Communication Sciences Unit, Indian Statistical Institute, Kolkata, India

Abstract

Principal Component Analysis (PCA) is a fundamental tool for data visualization, denoising, and dimensionality reduction. It is widely popular in Statistics, Machine Learning, Computer Vision, and related fields. However, PCA is well known to fall prey to the presence of outliers and often fails to detect the true underlying low-dimensional structure within the dataset. Recent supervised learning methods, following the Median of Means (MoM) philosophy, have shown great success in dealing with outlying observations without much compromise to their large sample theoretical properties. In this paper, we propose a PCA procedure based on the MoM principle. Called the **Median of Means Principal Component Analysis (MoMPCA)**, the proposed method is not only computationally appealing, but also achieves optimal convergence rates under minimal assumptions. In particular, we explore the non-asymptotic error bounds of the obtained solution via the aid of Vapnik-Chervonenkis theory and Rademacher complexity, while granting absolutely no assumption on the outlying observations. The efficacy of the proposal is also thoroughly showcased through simulations and real data applications.

1 Introduction

Principal component analysis (PCA) (Pearson, 1901; Wold et al., 1987) is the process of computing a small number of orthonormal basis vectors, which characterize most of the variability within the data cloud. PCA has been one of the most popular approaches for linear dimensionality reduction (Hastie et al., 2009). There has been many modifications to the original versions such as Probabilistic PCA (Tipping and Bishop, 1999), Kernel PCA (Schölkopf et al., 1997), Sparse PCA (Zou et al., 2006), Robust PCA (Candès et al., 2011; Zhang and Tong, 2019), PEA (Paul et al., 2020b) and so on. Among many disadvantages of PCA like information loss and poor interpretation, one of the major concerns is that it is well known to fail in the presence of a single outlying observation. Its fragility against severely corrupted data points often puts its reliability at risk. Popular approaches for making the PCA more robust to these outlying and corrupted observations attempt to represent the original data matrix \mathbf{X} as a sum of a low-rank matrix \mathbf{L} and a sparse matrix \mathbf{S} (Wright et al., 2009; Candès et al., 2011; Chandrasekaran et al., 2011; Kang et al., 2015; Chiang et al., 2016). However, many of the aforementioned methods only allow recovery guarantees under very restrictive assumptions. For example Candès et al. restricts the structure of \mathbf{S} to a Bernoulli model.

Recent advances in the Median of Means (MoM) literature allows us to consider an attractive framework to make an estimator outlier-robust. Opposed to the classical Vapnik-Chervonenkis Empirical Risk Minimization (ERM) (Vapnik, 2013), the MoM philosophy provides a more robust framework of efficiently finding estimates of the true underlying parameter. Although MoM estimator has been in the literature for quite a long time, it has recently been introduced to the Machine learning community (Lugosi et al., 2019; Lecué et al., 2020; Bartlett et al., 2002). MoM estimators are not only insensitive to the presence of observations far away from the bulk of the data cloud, but they are also equipped with finite sample error

*Correspondence to: swagatam.das@isical.ac.in




Figure 1: The first principal component found out by PCA and MoM PCA for the motivating example. Even in the presence of only 1% outlying observations, the classical PCA can render spurious results while MoM PCA finds the direction of maximum variation efficiently.

bounds and large sample theoretical guarantees such as consistency and convergence rates. Moreover, it has been observed in practice that MoM estimators often converge faster than the classical ERM estimators (Lecué et al., 2020).

In this paper, we propose a Principal Component Analysis (PCA) in the framework of the MoM principle. The proposal is not only computationally efficient but also theoretically appealing. It is well known that the theoretical understanding of many classical Empirical Risk Minimization (ERM) (Vapnik, 2013) such as PCA hinges on the assumptions that the data should be independently and identically distributed (i.i.d.) and have sub-gaussian behavior. However, real data that may be corrupted with outliers do not offer us the luxury to make such simplifying assumptions. Towards our theoretical investigation, we assume that the dataset can be split into two categories: the set of inliers (\mathcal{I}) and the set of outliers (\mathcal{O}), i.e. $\{1, \dots, N\} = \mathcal{I} \cup \mathcal{O}$. The data points in \mathcal{I} are assumed to be independently distributed according to the distribution P , which has a finite fourth moment. We make no assumptions on the points in \mathcal{O} ; thus, allowing them to be dependent, unboundedly large, having distributions P_i that are completely dissimilar to P , allowing them to be heavy-tailed etc. Our theoretical analysis hinges on the application of the Rademacher complexity (Bartlett and Mendelson, 2002) and symmetrization arguments (Vapnik, 2013; Devroye et al., 2013).

The summary of our contributions is as follows:

- We propose an efficient robust PCA based on MoM to tackle outlying observations within a dataset.
- Under some minimal and interpretable assumptions, we derive finite-sample uniform concentration bounds for MoMPCA. The analysis is carried out without making any distributional assumptions on the outlying observations. The consistency guarantees and error rates are derived subsequently.
- Through detailed experimental analysis, we show the efficacy of MoMPCA compared to the state-of-the-art.

As a motivating example, in Fig. 1, we show the results of classical PCA and MoMPCA on a toy dataset with 1000 inlying and 10 outlying observations. From Fig. 1, it is clear that a mere 1% outlying observations are enough to render PCA ineffective. However, MoMPCA correctly identifies the direction of maximum variation of the bulk of the data cloud.

The rest of the paper is organized as follows. In section 2, we formulate the MoMPCA, followed by detailed theoretical analysis under minimal and interpretable assumptions in section 3. The experimental results is discussed in section 4, followed by concluding remarks in section 5.

Notations Before we proceed further, we discuss a few notations used in this paper. $\langle A, B \rangle = A^\top B$ denotes the Frobenius inner product between two matrices. $\|A\| = \sqrt{\langle A, A \rangle}$ denotes the Frobenius norm of the matrix A . $\mathbb{P}(E)$ denotes the probability of the event E and $\mathbb{E}_{\mathbf{Z}}(\cdot)$ denotes the expectation with respect to the random variable \mathbf{Z} . A random variable σ is said to be Rademacher if it takes values in $\{+1, -1\}$ with equal probability. $[N]$ denotes the set $\{1, \dots, N\}$, for any $N \geq 0$ and 2^A denotes the power set of the A .

For any two sets A and B , A^B denotes the set of all functions from B to A . $\mu_k = \int \|\mathbf{x}\|_2^k dP$ denote the k -th moment of P .

2 PCA with Median of Means

2.1 Formulation

Suppose $\mathcal{X} = \{\mathbf{X}_1, \dots, \mathbf{X}_N\} \subset \mathbb{R}^p$ be a collection of N (mean-centered) data points in the p -dimensional real vector space. To find a d -dimensional ($d \leq p$) representation of \mathcal{X} , one projects these N points on to a d -dimensional subspace as $\{Q\mathbf{X}_1, \dots, Q\mathbf{X}_N\}$. where, Q is a $p \times p$ projection matrix of rank d . Denoting the set of all $p \times p$ projection matrix of rank d as \mathcal{Q}_d , the PCA minimizes the following objective:

$$\frac{1}{N} \sum_{i=1}^N \|\mathbf{X}_i - Q\mathbf{X}_i\|^2 = \frac{1}{N} \sum_{i=1}^N \mathbf{X}_i^\top (I - Q)\mathbf{X}_i, \quad (1)$$

where $Q \in \mathcal{Q}_d$. The aforementioned objective can be written as $\int f_Q(\mathbf{x})P_n(d\mathbf{x})$, with $f_Q(\mathbf{x}) = \mathbf{x}^\top (I - Q)\mathbf{x}$. Here $P_n = \frac{1}{N} \sum_{i=1}^N \delta_{\mathbf{X}_i}$ denotes the empirical distribution function. However, if the data is corrupted with outliers, minimizing the empirical risk (1), might give spurious estimates of Q as the empirical risk is heavily influenced by the presence of even a single outlying observation.

In the Median of Means philosophy, instead of directly minimizing the empirical risk, one first divides the data into L many partitions, B_1, \dots, B_L (i.e. $B_\ell \subset 2^{[N]}$, $\cup_{\ell=1}^L B_\ell = [N]$ and $B_\ell \cap B_{\ell'} = \emptyset$ for $\ell \neq \ell'$), such that each partition contains exactly B many elements (we can always throw out some observations if N is not divisible by L). Let P_{B_ℓ} denote the empirical distribution of $\{\mathbf{X}_i\}_{i \in B_\ell}$. For notational simplicity, we write, $\mu f = \int f d\mu$. We formulate the **Median of Means Principal Components Analysis** (MoMPCA) as the solution to the minimization problem of the following objective function,

$$\text{MoM}_L^N(f_Q) = \text{Median}(\{P_{B_\ell} f_Q : \ell = 1, \dots, L\}), \quad (2)$$

subject to $Q \in \mathcal{Q}_d$. Here, we will assume that the data is centered not by to the mean (as it is not outlier robust) but by some robust location estimate.

2.2 Optimization

The optimization of (2) is carried out by a gradient descent algorithm. We can write $Q = VV^\top$, where V is a $p \times d$ real column-orthonormal matrix. Thus, making a variable transform, we can write, (2) as

$$\text{MoM}_L^N(g_V) = \text{Median}(\{P_{B_\ell} g_V : \ell = 1, \dots, L\}), \quad (3)$$

where $g_V(\mathbf{x}) = \mathbf{x}^\top (I - VV^\top)\mathbf{x}$. At the t -th iteration, let the median class be $\ell_{\text{med}}^{(t)}$. According to [Lecué et al. \(2020\)](#), $\nabla \text{MoM}_L^N(g_{V^{(t)}}) = \frac{2}{B} \sum_{i \in B_{\ell_{\text{med}}^{(t)}}} \mathbf{X}_i \mathbf{X}_i^\top V^{(t)}$. However, after taking a gradient step, $V^{(t)} - \eta \frac{1}{B} \sum_{i \in B_{\ell_{\text{med}}^{(t)}}} \mathbf{X}_i \mathbf{X}_i^\top V^{(t)}$, the resultant matrix may not be column orthonormal. Following [Arora et al. \(2012\)](#), we apply a Gram-Schmidt algorithm to make the columns of this resultant matrix orthonormal. Let $\mathcal{P}_{\text{orth}}(V)$ denote the resultant matrix after orthonormalizing the columns of V . One can thus use the following gradient descent update:

$$V^{(t+1)} \leftarrow \mathcal{P}_{\text{orth}} \left(V^{(t)} - \eta \frac{1}{B} \sum_{i \in B_{\ell_{\text{med}}^{(t)}}} \mathbf{X}_i \mathbf{X}_i^\top V^{(t)} \right).$$

Algorithm 1 gives a formal description of the proposed gradient descent updates to optimize (2).

Algorithm 1 Median of Means Principal Component Analysis (MoMPCA)

Input: The dataset $\mathcal{X} = \{\mathbf{X}_1, \dots, \mathbf{X}_n\}$, L , d .

Output: $\widehat{Q}_{N,L}$.

Initialization: Randomly partition $[N]$ into L many partitions of equal length.

Compute the feature-wise median $\boldsymbol{\mu}$ of \mathcal{X} . Replace \mathbf{X}_i by $\mathbf{X}_i - \boldsymbol{\mu}$.

Compute the first d eigenvectors of $\sum_{i=1}^n \mathbf{X}_i \mathbf{X}_i^\top$ and stack them as columns of $V^{(0)}$.

repeat

 Compute $v_\ell \leftarrow \sum_{i \in B_\ell} \mathbf{X}_i^\top (I - V^{(t)}(V^{(t)})^\top) \mathbf{X}_i$

 Compute the median class ℓ_{med} , such that $v_{\ell_{\text{med}}} = \text{median}(\{v_\ell\}_{\ell=1}^L)$

$$V^{(t+1)} \leftarrow \mathcal{P}_{\text{orth}} \left(V^{(t)} - \eta \frac{1}{B} \sum_{i \in B_{\ell_{\text{med}}^{(t)}}} \mathbf{X}_i \mathbf{X}_i^\top V^{(t)} \right).$$

until objective $V^{(t)}$ converges

Compute $\widehat{Q}_{N,L} \leftarrow V^{(t)}(V^{(t)})^\top$.

3 Theoretical Properties

To explore the theoretical properties of MoMPCA, we assume that the dataset can be split into two categories: the set of inliers (\mathcal{I}) and the set of outliers (\mathcal{O}), i.e. $[N] = \mathcal{I} \cup \mathcal{O}$. We will assume that

Assumption 1. $\{\mathbf{X}_i\}_{i \in \mathcal{I}}$ are independently and identically distributed (i.i.d) according to the distribution P .

Assumption 2. $\mu_4 = \int \|\mathbf{x}\|^4 P(d\mathbf{x}) < \infty$.

Assumption 3. $\exists \eta > 0$, such that $L > (2 + \eta)|\mathcal{O}|$.

We emphasize that we do not make any assumptions on the outlying observations. They are even allowed to be dependent, and may even come from some heavy-tailed distributions. Assumptions 1 and 2 state that the inlying observations are independently and identically generated from some distribution with a finite fourth moment. To see the significance of Assumption 3, one should note that by pigeon hole principle, (strictly) less than half of the partitions may contain an outlying observation. More than half of the partitions, thus, do not contain an outlier. Now since the median is driven by the majority half of the partitions, naturally by Assumption 3, since the median is only affected by the objective functional value of the majority half of these partitions, the MoM estimates can be expected to only be based on the inlying observations.

For our theoretical understanding of MoMPCA, we need to compute the Rademacher complexity of the function class $\{f_Q : Q \in \mathcal{Q}_d\}$. We first recall the definition of Rademacher complexity. Let $\mathbf{Y}_1, \dots, \mathbf{Y}_m$ $\overset{i.i.d.}{\sim} P$ be random variables in some space \mathcal{S} . Suppose $\mathcal{F} \subseteq \mathbb{R}^{\mathcal{S}}$ be a class of functions from \mathcal{S} to \mathbb{R} and let $\sigma_1, \dots, \sigma_m$ be i.i.d. Rademacher random variables. The empirical Rademacher complexity, based on $\mathcal{Y} = \{\mathbf{Y}_1, \dots, \mathbf{Y}_m\}$ is defined as:

$$R_{\mathcal{Y}}(\mathcal{F}) = \frac{1}{m} \mathbb{E} \left(\sup_{f \in \mathcal{F}} \sum_{i=1}^m \sigma_i f(\mathbf{Y}_i) \middle| \mathcal{Y} \right).$$

Similarly, the population Rademacher complexity is defined as

$$R_m(\mathcal{F}) = \mathbb{E}(R_{\mathcal{Y}}(\mathcal{F})) = \frac{1}{m} \mathbb{E} \sup_{f \in \mathcal{F}} \sum_{i=1}^m \sigma_i f(\mathbf{Y}_i).$$

We now compute the Rademacher complexity of the function class $\{f_Q : Q \in \mathcal{Q}_d\}$ in Theorem 3.1.

Theorem 3.1. *Suppose $\mathcal{F}_d = \{f_Q : Q \in \mathcal{Q}_d\}$. Let $\mathbf{Y}_1, \dots, \mathbf{Y}_m \overset{i.i.d.}{\sim} P$, with P satisfying Assumption 2. Then, $R_m(\mathcal{F}_d) \leq \sqrt{\frac{(p-d)\mu_4}{m}}$.*

Proof. Suppose $\mathbf{Y}_1, \dots, \mathbf{Y}_m \stackrel{\text{i.i.d.}}{\sim} P$ and let $\mathcal{Y} = \{\mathbf{Y}_1, \dots, \mathbf{Y}_m\}$. The empirical Rademacher complexity is thus, given by,

$$\begin{aligned}
\widehat{R}_{\mathcal{Y}}(Q) &= \frac{1}{m} \mathbb{E}_{\boldsymbol{\sigma}} \sup_{Q \in \mathcal{Q}_d} \sum_{i=1}^m \sigma_i \mathbf{Y}_i^{\top} (I - Q) \mathbf{Y}_i \\
&= \frac{1}{m} \mathbb{E}_{\boldsymbol{\sigma}} \sup_{Q \in \mathcal{Q}_d} \sum_{i=1}^m \sigma_i \text{trace} \left(\mathbf{Y}_i^{\top} (I - Q) \mathbf{Y}_i \right) \\
&= \frac{1}{m} \mathbb{E}_{\boldsymbol{\sigma}} \sup_{Q \in \mathcal{Q}_d} \sum_{i=1}^m \sigma_i \text{trace} \left(\mathbf{Y}_i \mathbf{Y}_i^{\top} (I - Q) \right) \\
&= \frac{1}{m} \mathbb{E}_{\boldsymbol{\sigma}} \sup_{Q \in \mathcal{Q}_d} \sum_{i=1}^m \sigma_i \langle \mathbf{Y}_i \mathbf{Y}_i^{\top}, (I - Q) \rangle \\
&= \frac{1}{m} \mathbb{E}_{\boldsymbol{\sigma}} \sup_{Q \in \mathcal{Q}_d} \left\langle \sum_{i=1}^m \sigma_i \mathbf{Y}_i \mathbf{Y}_i^{\top}, (I - Q) \right\rangle \\
&\leq \frac{1}{m} \mathbb{E}_{\boldsymbol{\sigma}} \sup_{Q \in \mathcal{Q}_d} \left\| \sum_{i=1}^m \sigma_i \mathbf{Y}_i \mathbf{Y}_i^{\top} \right\| \| (I - Q) \| \\
&= \frac{1}{m} \mathbb{E}_{\boldsymbol{\sigma}} \sup_{Q \in \mathcal{Q}_d} \left\| \sum_{i=1}^m \sigma_i \mathbf{Y}_i \mathbf{Y}_i^{\top} \right\| \sqrt{\text{trace}((I - Q)^2)} \\
&= \frac{1}{m} \mathbb{E}_{\boldsymbol{\sigma}} \sup_{Q \in \mathcal{Q}_d} \left\| \sum_{i=1}^m \sigma_i \mathbf{Y}_i \mathbf{Y}_i^{\top} \right\| \sqrt{\text{trace}(I - Q)} \\
&= \frac{1}{m} \sqrt{p - d} \mathbb{E}_{\boldsymbol{\sigma}} \left[\sqrt{\text{trace} \left(\left(\sum_{i=1}^m \sigma_i \mathbf{Y}_i \mathbf{Y}_i^{\top} \right)^2 \right)} \right] \\
&\leq \frac{1}{m} \sqrt{p - d} \sqrt{\mathbb{E}_{\boldsymbol{\sigma}} \left[\text{trace} \left(\left(\sum_{i=1}^m \sigma_i \mathbf{Y}_i \mathbf{Y}_i^{\top} \right)^2 \right) \right]} \\
&= \frac{1}{m} \sqrt{p - d} \sqrt{\text{trace} \left(\sum_{i=1}^m \left(\mathbf{Y}_i \mathbf{Y}_i^{\top} \right)^2 \right)} \\
&= \frac{1}{m} \sqrt{p - d} \sqrt{\sum_{i=1}^m \|\mathbf{Y}_i\|^4}.
\end{aligned}$$

Thus,

$$\begin{aligned}
R_m(Q) = \mathbb{E}_{\mathcal{Y}} \widehat{R}_{\mathcal{X}}(Q) &= \frac{1}{m} \sqrt{p - d} \mathbb{E} \left[\sqrt{\sum_{i=1}^m \|\mathbf{Y}_i\|^4} \right] \\
&\leq \frac{1}{m} \sqrt{p - d} \sqrt{\sum_{i=1}^m \mathbb{E} \|\mathbf{Y}_i\|^4} \\
&= \sqrt{\frac{(p - d) \mu_4}{m}}.
\end{aligned}$$

□

We are now ready to state and prove the uniform concentration bound result in Theorem 3.2. Theorem 3.2 asserts that with a high probability, $\sup_{Q \in \mathcal{Q}_d} |\text{MoM}_L^N(f_Q) - Pf_Q|$ decreases with at least a rate of $\max \left\{ \sqrt{\frac{L}{N}}, k \frac{\sqrt{|I|}}{N} \right\}$.

Theorem 3.2. *Under Assumptions 1-3, and $N > L$, with probability at least, $1 - 2e^{-2L(\frac{2}{4+\eta} - \frac{|\mathcal{O}|}{L})^2}$,*

$$\sup_{Q \in \mathcal{Q}_d} |\text{MoM}_L^N(f_Q) - Pf_Q| \leq C \max \left\{ \sqrt{\frac{L}{N}}, \frac{\sqrt{|I|}}{N} \right\},$$

with $C = 2 \max \left\{ \sqrt{\frac{8(4+\eta)C(P)}{\eta}}, \frac{16\sqrt{(p-d)\mu_4(4+\eta)}}{\eta} \right\}$ and $C(P) = (\mu_4 + 2\mu_2^2)(p-d) + (\mu_2^2 - 1)(p-d)^2$.

Proof. Suppose $\epsilon > 0$. We will first bound the probability of $\sup_{Q \in \mathcal{Q}_d} |\text{MoM}_L^N(f_Q) - Pf_Q| > \epsilon$. To do so, we will individually bound the probabilities of the events, $\sup_{Q \in \mathcal{Q}_d} (\text{MoM}_L^N(f_Q) - Pf_Q) > \epsilon$ and $\sup_{Q \in \mathcal{Q}_d} (Pf_Q - \text{MoM}_L^N(f_Q)) > \epsilon$. We note that if $\sup_{Q \in \mathcal{Q}_d} \sum_{\ell=1}^L \mathbb{1}\{(P - P_{B_\ell})f_Q > \epsilon\} > \frac{L}{2}$, then, $\sup_{Q \in \mathcal{Q}_d} (Pf_Q - \text{MoM}_L^N(f_Q)) > \epsilon$. Here $\mathbb{1}\{\cdot\}$ denote the indicator function. Let $\varphi(t) = (t-1)\mathbb{1}\{1 \leq t \leq 2\} + \mathbb{1}\{t > 2\}$. Clearly,

$$\mathbb{1}\{t \geq 2\} \leq \varphi(t) \leq \mathbb{1}\{t \geq 1\}. \quad (4)$$

We observe that,

$$\begin{aligned} & \sup_{Q \in \mathcal{Q}_d} \sum_{\ell=1}^L \mathbb{1}\{(P - P_{B_\ell})f_Q > \epsilon\} \\ & \leq \sup_{Q \in \mathcal{Q}_d} \sum_{\ell \in \mathcal{L}} \mathbb{1}\{(P - P_{B_\ell})f_Q > \epsilon\} + |\mathcal{O}| \\ & \leq \sup_{Q \in \mathcal{Q}_d} \sum_{\ell \in \mathcal{L}} \varphi \left(\frac{2(P - P_{B_\ell})f_Q}{\epsilon} \right) + |\mathcal{O}| \\ & \leq \sup_{Q \in \mathcal{Q}_d} \sum_{\ell \in \mathcal{L}} \mathbb{E} \varphi \left(\frac{2(P - P_{B_\ell})f_Q}{\epsilon} \right) + |\mathcal{O}| + \sup_{Q \in \mathcal{Q}_d} \sum_{\ell \in \mathcal{L}} \left[\varphi \left(\frac{2(P - P_{B_\ell})f_Q}{\epsilon} \right) - \mathbb{E} \varphi \left(\frac{2(P - P_{B_\ell})f_Q}{\epsilon} \right) \right]. \end{aligned} \quad (5)$$

To bound $\sup_{Q \in \mathcal{Q}_d} \sum_{\ell=1}^L \mathbb{1}\{(P - P_{B_\ell})f_Q > \epsilon\}$, we will first bound the quantity, $\mathbb{E} \varphi \left(\frac{2(P - P_{B_\ell})f_Q}{\epsilon} \right)$. We observe that,

$$\begin{aligned} \mathbb{E} \varphi \left(\frac{2(P - P_{B_\ell})f_Q}{\epsilon} \right) & \leq \mathbb{E} \left[\mathbb{1} \left\{ \frac{2(P - P_{B_\ell})f_Q}{\epsilon} > 1 \right\} \right] \\ & = \mathbb{P} \left[(P - P_{B_\ell})f_Q > \frac{\epsilon}{2} \right] \\ & \leq \frac{4}{\epsilon^2} \text{Var}((P - P_{B_\ell})f_Q) \end{aligned} \quad (6)$$

$$\begin{aligned} & = \frac{4}{\epsilon^2} \text{Var}(P_{B_\ell}f_Q) \\ & = \frac{4}{B\epsilon^2} \text{Var}(\mathbf{X}^\top (I - Q)\mathbf{X}) \\ & = \frac{4}{B\epsilon^2} \left[(\mu_4 - 3\mu_2^2) \sum_{i=1}^p (1 - Q_{ii})^2 + (\mu_2^2 - 1)(\text{tr}(I - Q))^2 + 2\mu_2^2 \text{tr}((I - Q)^2) \right] \end{aligned} \quad (7)$$

$$\begin{aligned} & \leq \frac{4}{B\epsilon^2} \left[\mu_4 \text{tr}((I - Q)^2) + (\mu_2^2 - 1)(p-d)^2 + 2\mu_2^2(p-d) \right] \\ & = \frac{4}{B\epsilon^2} \left[\mu_4(p-d) + (\mu_2^2 - 1)(p-d)^2 + 2\mu_2^2(p-d) \right] \\ & = \frac{4}{B\epsilon^2} \left[(\mu_4 + 2\mu_2^2)(p-d) + (\mu_2^2 - 1)(p-d)^2 \right]. \end{aligned} \quad (8)$$

Here equation (6) follows from Chebyshev's inequality and equation (7) follows from the formula of variance of quadratic form of random vectors. We now concentrate on bounding the term $\sup_{Q \in \mathcal{Q}_d} \sum_{\ell \in \mathcal{L}} \left[\varphi \left(\frac{2(P - P_{B_\ell})f_Q}{\epsilon} \right) - \mathbb{E} \varphi \left(\frac{2(P - P_{B_\ell})f_Q}{\epsilon} \right) \right]$. Appealing to Theorem 26.5 of (Shalev-Shwartz and Ben-David, 2014) we observe that, with probability at least $1 - e^{-2L\delta^2}$, $\forall Q \in \mathcal{Q}_d$,

$$\begin{aligned} & \frac{1}{L} \sum_{\ell \in \mathcal{L}} \varphi \left(\frac{2(P - P_{B_\ell})f_Q}{\epsilon} \right) \\ & \leq \mathbb{E} \left[\frac{1}{L} \sum_{\ell \in \mathcal{L}} \varphi \left(\frac{2(P - P_{B_\ell})f_Q}{\epsilon} \right) \right] + 2\mathbb{E} \left[\sup_{Q \in \mathcal{Q}_d} \frac{1}{L} \sum_{\ell \in \mathcal{L}} \sigma_\ell \varphi \left(\frac{2(P - P_{B_\ell})f_Q}{\epsilon} \right) \right] + \delta. \end{aligned} \quad (9)$$

Here $\{\sigma_\ell\}_{\ell \in \mathcal{L}}$ are i.i.d. Rademacher random variables. Let $\{\xi_i\}_{i=1}^n$ be i.i.d. Rademacher random variables, independent from $\{\sigma_\ell\}_{\ell \in \mathcal{L}}$. From equation (9), we get,

$$\begin{aligned} & \frac{1}{L} \sup_{Q \in \mathcal{Q}_d} \sum_{\ell \in \mathcal{L}} \left[\varphi \left(\frac{2(P - P_{B_\ell})f_Q}{\epsilon} \right) - \mathbb{E} \varphi \left(\frac{2(P - P_{B_\ell})f_Q}{\epsilon} \right) \right] \\ & \leq 2\mathbb{E} \left[\sup_{Q \in \mathcal{Q}_d} \frac{1}{L} \sum_{\ell \in \mathcal{L}} \sigma_\ell \varphi \left(\frac{2(P - P_{B_\ell})f_Q}{\epsilon} \right) \right] + \delta \\ & \leq \frac{4}{L\epsilon} \mathbb{E} \left[\sup_{Q \in \mathcal{Q}_d} \sum_{\ell \in \mathcal{L}} \sigma_\ell (P - P_{B_\ell})f_Q \right] + \delta. \end{aligned} \quad (10)$$

Equation (10) follows from the fact that $\varphi(\cdot)$ is 1-Lipschitz and appealing to Lemma 26.9 of Shalev-Shwartz and Ben-David (2014). We now introduce a ghost sample $\mathcal{X}' = \{\mathbf{X}'_1, \dots, \mathbf{X}'_n\}$, which are i.i.d. and follows the law P . Thus, equation (10) further equals the following quantity.

$$\begin{aligned} & = \frac{4}{L\epsilon} \mathbb{E} \left[\sup_{Q \in \mathcal{Q}_d} \sum_{\ell \in \mathcal{L}} \sigma_\ell \mathbb{E}_{\mathcal{X}'} \left((P'_{B_\ell} - P_{B_\ell})f_Q \right) \right] + \delta \\ & \leq \frac{4}{L\epsilon} \mathbb{E} \left[\sup_{Q \in \mathcal{Q}_d} \sum_{\ell \in \mathcal{L}} \sigma_\ell (P'_{B_\ell} - P_{B_\ell})f_Q \right] + \delta \\ & = \frac{4}{L\epsilon} \mathbb{E} \left[\sup_{Q \in \mathcal{Q}_d} \sum_{\ell \in \mathcal{L}} \sigma_\ell \frac{1}{B} \sum_{i \in B_\ell} (f_Q(\mathbf{X}'_i) - f_Q(\mathbf{X}_i)) \right] + \delta \\ & = \frac{4}{BL\epsilon} \mathbb{E} \left[\sup_{Q \in \mathcal{Q}_d} \sum_{\ell \in \mathcal{L}} \sigma_\ell \sum_{i \in B_\ell} \xi_i (f_Q(\mathbf{X}'_i) - f_Q(\mathbf{X}_i)) \right] + \delta \end{aligned} \quad (11)$$

$$\begin{aligned} & = \frac{4}{N\epsilon} \mathbb{E} \left[\sup_{Q \in \mathcal{Q}_d} \sum_{\ell \in \mathcal{L}} \sum_{i \in B_\ell} \sigma_\ell \xi_i (f_Q(\mathbf{X}'_i) - f_Q(\mathbf{X}_i)) \right] + \delta \\ & \leq \frac{4}{N\epsilon} \mathbb{E} \left[\sup_{Q \in \mathcal{Q}_d} \sum_{\ell \in \mathcal{L}} \sum_{i \in B_\ell} \sigma_\ell \xi_i (f_Q(\mathbf{X}'_i) + f_Q(\mathbf{X}_i)) \right] + \delta \\ & = \frac{4}{N\epsilon} \mathbb{E} \left[\sup_{Q \in \mathcal{Q}_d} \sum_{i \in \mathcal{J}} \gamma_i (f_Q(\mathbf{X}'_i) + f_Q(\mathbf{X}_i)) \right] \end{aligned} \quad (12)$$

$$= \frac{8}{N\epsilon} \mathbb{E} \left[\sup_{Q \in \mathcal{Q}_d} \sum_{i \in \mathcal{J}} \gamma_i f_Q(\mathbf{X}_i) \right] + \delta \quad (13)$$

$$\leq \frac{8}{N\epsilon} \sqrt{(p-d)\mu_4|\mathcal{J}|} + \delta$$

$$\leq \frac{8}{N\epsilon} \sqrt{(p-d)\mu_4|\mathcal{I}|} + \delta. \quad (14)$$

Equation (11) follows from observing that $(f_Q(\mathbf{X}'_i) - f_Q(\mathbf{X}_i)) \stackrel{d}{=} \xi_i(f_Q(\mathbf{X}'_i) - f_Q(\mathbf{X}_i))$. In equation (12), $\{\gamma_i\}_{i \in \mathcal{J}}$ are independent Rademacher random variables due to their construction. Equation (13) follows from appealing to Theorem 3.1. Thus, combining equations (9), (10), and (14), we conclude that, with probability of at least $1 - e^{-2L\delta^2}$,

$$\begin{aligned} & \sup_{Q \in \mathcal{Q}_d} \sum_{\ell=1}^L \mathbb{1}\{(P - P_{B_\ell})f_Q > \epsilon\} \\ & \leq L \left(\frac{4}{B\epsilon^2} [(\mu_4 + 2\mu_2^2)(p-d) + (\mu_2^2 - 1)(p-d)^2] + \frac{|\mathcal{O}|}{L} + \frac{8}{N\epsilon} \sqrt{(p-d)\mu_4|\mathcal{I}|} + \delta \right). \end{aligned} \quad (15)$$

For notational simplicity, we write, $C(P) = (\mu_4 + 2\mu_2^2)(p-d) + (\mu_2^2 - 1)(p-d)^2$. We choose $\delta = \frac{2}{4+\eta} - \frac{|\mathcal{O}|}{L}$ and $\epsilon = 2 \max \left\{ \sqrt{\frac{8(4+\eta)C(P)}{\eta}} \sqrt{\frac{L}{N}}, \frac{16\sqrt{(p-d)\mu_4(4+\eta)}}{\eta} \frac{\sqrt{|\mathcal{I}|}}{N} \right\}$. This makes the right hand side of (15) strictly smaller than $\frac{L}{2}$.

Thus, we have shown that

$$\mathbb{P} \left(\sup_{Q \in \mathcal{Q}_d} (Pf_Q - \text{MoM}_L^N(f_Q)) > \epsilon \right) \leq e^{-2L\delta^2}.$$

Similarly, we can show that,

$$\mathbb{P} \left(\sup_{Q \in \mathcal{Q}_d} (\text{MoM}_L^N(f_Q) - Pf_Q) > \epsilon \right) \leq e^{-2L\delta^2}.$$

Combining the above two inequalities, we get,

$$\mathbb{P} \left(\sup_{Q \in \mathcal{Q}_d} |\text{MoM}_L^N(f_Q) - Pf_Q| > \epsilon \right) \leq 2e^{-2L\delta^2}.$$

In other words, with at least probability $1 - 2e^{-2L\delta^2}$,

$$\begin{aligned} & \sup_{Q \in \mathcal{Q}_d} |\text{MoM}_L^N(f_Q) - Pf_Q| \\ & \leq 2 \max \left\{ \sqrt{\frac{16(2+\eta)C(P)}{\eta}} \sqrt{\frac{L}{N}}, \frac{32\sqrt{(p-d)\mu_4(2+\eta)}}{\eta} \frac{\sqrt{|\mathcal{I}|}}{N} \right\} \leq C \max \left\{ \sqrt{\frac{L}{N}}, \frac{\sqrt{|\mathcal{I}|}}{N} \right\}. \end{aligned}$$

□

Let $\widehat{Q}_{N,L} = \text{argmin}_{Q \in \mathcal{Q}_d} \text{MoM}_L^N(f_Q)$ and $Q^* = \text{argmin}_{Q \in \mathcal{Q}_d} Pf_Q$. From Theorem 3.2, we can say that for any any $Q \in \mathcal{Q}_d$, $\text{MoM}_L^N(f_Q)$ and Pf_Q are close to each other with a high probability. Thus, one can expect that their corresponding minimum values be close enough with a high probability. Theorem 3.3 affirms this claim.

Theorem 3.3. *Under Assumptions 1-3, and $N > L$, with probability at least, $1 - 2e^{-2L(\frac{2}{4+\eta} - \frac{|\mathcal{O}|}{L})^2}$,*

$$|Pf_{\widehat{Q}_{N,L}} - Pf_{Q^*}| \leq 2C \max \left\{ \sqrt{\frac{L}{N}}, \frac{\sqrt{|\mathcal{I}|}}{N} \right\},$$

where C is defined as in Theorem 3.2.

Proof. Appealing to the fact that $Pf_{Q^*} \leq Pf_Q$ and $\text{MoM}_L^N(f_{Q_{N,L}}) \leq \text{MoM}_L^N(f_Q)$, for all $Q \in \mathcal{Q}_d$, we observe that,

$$\begin{aligned}
& |Pf_{\widehat{Q}_{N,L}} - Pf_{Q^*}| \\
&= Pf_{Q^*} - Pf_{\widehat{Q}_{N,L}} \\
&= (Pf_{Q^*} - \text{MoM}_L^N(f_{Q^*})) + (\text{MoM}_L^N(f_{\widehat{Q}_{N,L}}) - \text{MoM}_L^N(f_{Q^*})) + (\text{MoM}_L^N(f_{\widehat{Q}_{N,L}}) - Pf_{\widehat{Q}_{N,L}}) \\
&= (Pf_{Q^*} - \text{MoM}_L^N(f_{Q^*})) + (\text{MoM}_L^N(f_{\widehat{Q}_{N,L}}) - Pf_{\widehat{Q}_{N,L}}) \\
&\leq 2 \sup_{Q \in B(M)^k} |\text{MoM}_L^N(f_Q) - Pf_Q| \\
&\leq 2C \max \left\{ \sqrt{\frac{L}{N}}, \frac{\sqrt{|I|}}{N} \right\}. \tag{16}
\end{aligned}$$

From Theorem 3.2, equation (16) holds with probability at least $1 - 2e^{-2L(\frac{2}{4+\eta} - \frac{|\mathcal{O}|}{L})^2}$, proving the desired result. \square

We observe that $\frac{2}{4+\eta} - \frac{|\mathcal{O}|}{L} \geq \frac{2}{4+\eta} - \frac{1}{2+\eta} = \frac{\eta}{(2+\eta)(4+\eta)} > 0$. Thus, if $L \rightarrow \infty$, $1 - 2e^{-2L(\frac{2}{4+\eta} - \frac{|\mathcal{O}|}{L})^2} \rightarrow 1$. Moreover, if we assume that $\frac{L}{N} \rightarrow 0$,

$$\max \left\{ \sqrt{\frac{L}{N}}, \frac{\sqrt{|I|}}{N} \right\} \leq \max \left\{ \sqrt{\frac{L}{N}}, \frac{1}{\sqrt{N}} \right\} \rightarrow 0.$$

Hence, for any prefixed tolerance ϵ ,

$$\lim_{N \rightarrow \infty} \mathbb{P} \left(|Pf_{\widehat{Q}_{N,L}} - Pf_{Q^*}| > \epsilon \right) = 0.$$

In other words, $Pf_{\widehat{Q}_{N,L}} \xrightarrow{P} Pf_{Q^*}$. To formally state this result, we make the following to assumptions.

Assumption 4. $L \rightarrow \infty$, as $N \rightarrow \infty$.

Assumption 5. $L = o(N)$.

Thus, we can immediately infer the following corollary.

Corollary 3.1. *Under assumptions 1-5, $Pf_{\widehat{Q}_{N,L}} \xrightarrow{P} Pf_{Q^*}$.*

To conclude that $\widehat{Q}_{N,L} \xrightarrow{P} Q^*$, we need to ensure that Q^* is identifiable. Towards ensuring that, we make the following identifiability assumption on Q^* . This type of identifiability conditions are especially popular in clustering literature (Pollard, 1981; Chakraborty and Das, 2019; Chakraborty et al., 2020; Chakraborty and Das, 2020; Paul et al., 2020a).

Assumption 6. For all $\epsilon > 0$, there exists $\delta > 0$, such that, if $\|Q - Q^*\| > \epsilon$, $Pf_Q \geq Pf_{Q^*} + \delta$.

We will say that $Q_n \rightarrow Q$ if $\|Q_n - Q\| \rightarrow 0$. With this notion of convergence in the Frobenious sense, we are now ready to prove that $\widehat{Q}_{N,L}$ is consistent for Q^* .

Theorem 3.4. *Under assumptions 1-6, $\widehat{Q}_{N,L} \xrightarrow{P} Q^*$.*

Proof. We fix $\epsilon > 0$. By assumption 6, there exists $\delta > 0$, such that, if $\|Q - Q^*\| > \epsilon$, $Pf_Q \geq Pf_{Q^*} + \delta$. Thus,

$$\mathbb{P}(\|\widehat{Q}_{N,L} - Q^*\| > \epsilon) \leq \mathbb{P} \left(Pf_{\widehat{Q}_{N,L}} \geq Pf_{Q^*} + \delta \right) \leq \mathbb{P} \left(|Pf_{\widehat{Q}_{N,L}} - Pf_{Q^*}| \geq \delta \right) \rightarrow 0,$$

as $N \rightarrow \infty$, by appealing to corollary 3.1. Thus, $\|\widehat{Q}_{N,L} - Q^*\| \xrightarrow{P} 0 \iff \widehat{Q}_{N,L} \xrightarrow{P} Q^*$. \square

n	p	$\text{rank}(X_0)$	PCP	PCPF	MFRPCA	RWL-AN	MoM-PCA
500	500	10	1.6×10^{-3}	1.5×10^{-3}	2.3×10^{-3}	8.7×10^{-3}	1.5×10^{-3}
1000	500	10	7.3×10^{-4}	2.8×10^{-4}	1.8×10^{-3}	2.9×10^{-4}	2.1×10^{-4}
2000	500	10	8.8×10^{-4}	1.1×10^{-4}	8.3×10^{-4}	3.1×10^{-4}	5.6×10^{-5}
5000	500	10	2.1×10^{-4}	9.6×10^{-5}	5.7×10^{-4}	3.3×10^{-4}	7.2×10^{-6}
10000	500	10	7.8×10^{-5}	8.2×10^{-5}	2.9×10^{-4}	9.5×10^{-5}	3.8×10^{-7}

Table 1: Performance of different peer algorithms for low-rank matrix representation in terms of relative reconstruction error.




Figure 2: Background modeling through MoMPCA. The background is modeled as a low-dimensional approximation, calculated through MoMPCA as described in section 4.2.

4 Experimental Results

In this section, we demonstrate the efficacy of MoMPCA on synthetic and real datasets. We apply MoMPCA to perform different tasks including low-rank matrix reconstruction in the presence of outliers, background modeling for video data and anomaly detection for real data benchmarks. All the codes are available at <https://github.com/DebolinaPaul/MoMPCA>.

4.1 A Simulation Study on Recovering Low-rank Matrices

To empirically validate the efficacy of MoMPCA, we performed this dimensionality reduction procedure on various simulated datasets. We have generated the matrix X_0 as a product of two matrices $X_{1,0}$ and $X_{2,0}$ of orders $n \times r$ and $r \times p$ respectively where each entry is simulated from the Gaussian distribution, where r is the rank of X_0 and $r < \min\{n, p\}$. To add outliers, we add random noise generated from $Unif(-500, 500)$ to randomly selected \sqrt{n} many rows of X_0 .

We then run the MoMPCA algorithm along with the peer algorithms on X_0 . We calculate the efficacy of each of the relative reconstruction error as $\frac{\|X - X_0\|}{\|X_0\|}$, where the norm is taken over those rows that do not contain an outlier. Here X is the projected matrix in the lower d -dimensional affine space. Clearly, a lower score represents a better representation in lower-dimensional space. Since the true structure of the matrix lies in a lower-dimensional space, the MoMPCA method successfully plots the dataset in a lower-dimensional space in spite of having \sqrt{n} many outliers due to its robustness property.

We compare our method with the baselines as well as state-of-the-art methods such as Principal Component Pursuit (PCP) (Candès et al., 2011), Principal Component Pursuit with Features (PCPF) (Chiang et al., 2016), Matrix-Factorization Based Robust Principal Component Analysis (MFRPCA) (Chen and Zhou, 2018) and Robust Weight Learning with Adaptive Neighbors (RWL-AN) (Zhang and Tong, 2019). The standard protocols as was used by each original papers of the competing algorithms have been implemented in our experiments. The algorithms providing non-deterministic output have been run 20 times and the average values obtained have been reported. The results demonstrated in table 1 clearly indicates that out of all the methods, the proposed MoMPCA algorithm works best in terms of the reconstruction error.

Data	# Dimensions	# Instances	Anomaly ratio
KDDCUP	120	494,021	0.2
Thyroid	6	3,772	0.025
Arrhythmia	274	452	0.15

Table 2: Statistics of different anomaly detection benchmarks

Method	KDDCUP			Thyroid			Arrhythmia			Average
	Precision	Recall	F_1	Precision	Recall	F_1	Precision	Recall	F_1	rank (F_1)
OC-SVM	0.7457	0.8523	0.7954	0.3639	0.4239	0.3887	0.5397	0.4082	0.4581	6.33
DSEBM-r	0.8521	0.6472	0.7328	0.0404	0.0403	0.0403	0.1515	0.1513	0.1510	11.33
DSEBM-e	0.8619	0.6446	0.7399	0.1319	0.1319	0.1319	0.4667	0.4565	0.4601	7.67
DCN	0.7696	0.7829	0.7762	0.3319	0.3196	0.3251	0.3758	0.3907	0.3815	8
GMM-EN	0.1932	0.1967	0.1949	0.0213	0.0227	0.0220	0.3000	0.2792	0.2886	12.33
PAE	0.7276	0.7397	0.7336	0.1894	0.2062	0.1971	0.4393	0.4437	0.4403	8.67
E2E-AE	0.0024	0.0025	0.0024	0.1064	0.1316	0.1176	0.4667	0.4538	0.4591	10.33
PAE-GMM-EM	0.7183	0.7311	0.7246	0.4745	0.4538	0.4635	0.3970	0.4168	0.4056	8.67
PAE-GMM	0.7251	0.7384	0.7317	0.4532	0.4881	0.4688	0.4575	0.4823	0.4684	6.33
DAGMM-p	0.7579	0.7710	0.7644	0.4723	0.4725	0.4713	0.4909	0.4679	0.4787	4.33
DAGMM-NVI	0.9290	0.9447	0.9368	0.4383	0.4587	0.4470	0.5091	0.4892	0.4981	3.33
DAGMM	0.9297	0.9442	0.9369	0.4766	0.4834	0.4782	0.4909	0.5078	0.4983	2
MoMPCA	0.8966	0.9106	0.9035	0.6974	0.5699	0.6272	0.5469	0.5303	0.5385	1.67

Table 3: Performance of different peer algorithms for anomaly detection. The average ranks on the basis of the F_1 scores are also reported. The best performances for each data w.r.t. each metric are boldfaced.

4.2 Background Modeling in Video

Suppose, we are given a video with f many frames. The objective here is to isolate the moving objects in the video such as moving cars, pedestrians etc. For simplicity, we only consider black and white videos (and thus, there is only a single channel) and let each frame be of size $m \times n$. We construct $\mathcal{X} = \{\mathbf{X}_1, \dots, \mathbf{X}_{mn}\} \subset \mathbb{R}^f$ by computing a $f \times 1$ vector whose features represent the value of a particular pixel for each frame. Since consecutive frames are highly correlated, one can expect that the data cloud \mathcal{X} lies near to a low-dimensional affine space in \mathbb{R}^f . Thus, the background of the video can be modeled as the projection of \mathcal{X} on this low-dimensional affine space. To demonstrate the efficacy of modeling the background through MoMPCA, we take the standard video, called ‘‘highway’’ (available in the github repository), which captures moving cars on a busy highway. The video contains 1700 frames and each frame is of size 240×320 . We construct a 76800×1700 data matrix, whose rows denote the data points. We run the MoMPCA with $L = 40$ and $d = 5$. The background is thus given by $\{Q\mathbf{X}_1, \dots, Q\mathbf{X}_{76800}\}$, where Q is the projection matrix, given by Algorithm 1. The output background for frame no. 863 is shown in Fig. 2 along with the original frame. The object is constructed by taking $\{\|\mathbf{X}_i - Q\mathbf{X}_i\|_1\}_{i=1}^{76800}$. Clearly the object conforms to the moving cars in the original frame and the background only consists of the highway and the motionless trees.

4.3 An application in Anomaly Detection

Anomaly detection or outlier detection is a key problem in machine learning and computer vision. In this section, we will focus on how to detect outliers through MoMPCA. Suppose $\mathcal{X} = \{\mathbf{X}_1, \dots, \mathbf{X}_n\} \subset \mathbb{R}^p$ be n data points in the p -dimensional real vector space. To detect outliers within the data, we first project \mathcal{X} to a d -dimensional affine space via MoMPCA. We then compute the squared Euclidean distance between the original and projected data points $v_i = \|\mathbf{X}_i - \widehat{Q}_{N,L}\mathbf{X}_i\|_2^2$ and sort v_i ’s in ascending order. We call the i -th point an outlier if v_i belongs to the largest $o\%$ of the $\{v_i\}_{i=1}^n$, where o is known beforehand.

For our comparative analysis, we take the KDDCUP, Thyroid, and Arrhythmia datasets, which are available from the UCI machine learning repository (Dua and Graff, 2017) and ODDS library (Rayana, 2016). The details of these datasets are reported in Table 2. We compared our method with different state-of-the-art techniques for outlier detection including OC-SVM (Chen et al., 2001); DSEBM-e & DSEBM-r (Zhang and Woodland, 2017); DCN (Yang et al., 2017); GMM-EN, PAE, E2E-AE, PAE-GMM-EM, PAE-GMM, DAGMM-p, DAGMM-NVI and DAGMM (Zong et al., 2018). Many of the competing algorithms

employ a deep neural network to detect anomalies within the data. To measure the performance of the peer algorithms, we take the average precision, recall, and F_1 score between the ground truth and the obtained labeling (inlier/outlier) of the data points. These measures of accuracy for all the three datasets are reported in Table 3. The performance indicator values for the peer algorithms are quoted from Zong et al. (2018). It is observed from Table 3 that the MoMPCA is quite competitive against the state-of-the-art anomaly detection methods, including even the ones based on deep neural networks.

5 Conclusions

Despite the efficacy, computational simplicity, and ease of visualization of the classical PCA, it often fails to successfully represent the true low-dimensional structure of a dataset in the presence of even a small number of outliers. To circumvent this problem, we propose an alternative PCA method based on the Median of Means (MoM) estimator. The proposed MoMPCA, equipped with a computationally simple gradient descent based optimization procedure, exhibits significant robustness to the presence of outliers.

Under minimal and interpretable assumptions, we establish the consistency of MoMPCA and find the rate of convergence by using uniform concentration bounds. The theoretical analysis of the paper is carried out with the aid of symmetrization arguments and Rademacher complexities, which although extensively used in a supervised learning setting, seldom find application in an unsupervised learning scenario. Through experimental applications in computer vision, MoMPCA is shown to be effective relative to even some of the recent deep learning models. A possible future extension of our work could be to make the optimization faster by using adaptive gradient-based optimizers and prove their convergence properties.

References

- Arora, R., Cotter, A., Livescu, K., and Srebro, N. (2012). Stochastic optimization for pca and pls. In *2012 50th Annual Allerton Conference on Communication, Control, and Computing (Allerton)*, pages 861–868. IEEE.
- Bartlett, P. L., Boucheron, S., and Lugosi, G. (2002). Model selection and error estimation. *Machine Learning*, 48(1):85–113.
- Bartlett, P. L. and Mendelson, S. (2002). Rademacher and gaussian complexities: Risk bounds and structural results. *Journal of Machine Learning Research*, 3(Nov):463–482.
- Candès, E. J., Li, X., Ma, Y., and Wright, J. (2011). Robust principal component analysis? *Journal of the ACM (JACM)*, 58(3):1–37.
- Chakraborty, S. and Das, S. (2019). On the strong consistency of feature-weighted k-means clustering in a nearmetric space. *Stat*, 8(1):e227.
- Chakraborty, S. and Das, S. (2020). Detecting meaningful clusters from high-dimensional data: A strongly consistent sparse center-based clustering approach. *IEEE Transactions on Pattern Analysis and Machine Intelligence*.
- Chakraborty, S., Paul, D., Das, S., and Xu, J. (2020). Entropy weighted power k-means clustering. In *International Conference on Artificial Intelligence and Statistics*, pages 691–701. PMLR.
- Chandrasekaran, V., Sanghavi, S., Parrilo, P. A., and Willsky, A. S. (2011). Rank-sparsity incoherence for matrix decomposition. *SIAM Journal on Optimization*, 21(2):572–596.
- Chen, Y., Zhou, X. S., and Huang, T. S. (2001). One-class svm for learning in image retrieval. In *Proceedings 2001 International Conference on Image Processing (Cat. No. 01CH37205)*, volume 1, pages 34–37. IEEE.
- Chen, Y. and Zhou, Y. (2018). Robust principal component analysis with matrix factorization. In *2018 IEEE International Conference on Acoustics, Speech and Signal Processing (ICASSP)*, pages 2411–2415. IEEE.

- Chiang, K.-Y., Hsieh, C.-J., and Dhillon, I. (2016). Robust principal component analysis with side information. In *Proceedings of The 33rd International Conference on Machine Learning*, volume 48 of *Proceedings of Machine Learning Research*, pages 2291–2299, New York, New York, USA. PMLR.
- Devroye, L., Györfi, L., and Lugosi, G. (2013). *A probabilistic theory of pattern recognition*, volume 31. Springer Science & Business Media.
- Dua, D. and Graff, C. (2017). UCI machine learning repository.
- Hastie, T., Tibshirani, R., and Friedman, J. (2009). *The elements of statistical learning: data mining, inference, and prediction*. Springer Science & Business Media.
- Kang, Z., Peng, C., and Cheng, Q. (2015). Robust pca via nonconvex rank approximation. In *2015 IEEE International Conference on Data Mining*, pages 211–220. IEEE.
- Lecué, G., Lerasle, M., et al. (2020). Robust machine learning by median-of-means: theory and practice. *Annals of Statistics*, 48(2):906–931.
- Lugosi, G., Mendelson, S., et al. (2019). Regularization, sparse recovery, and median-of-means tournaments. *Bernoulli*, 25(3):2075–2106.
- Paul, D., Chakraborty, S., Das, S., and Xu, J. (2020a). Kernel k-means, by all means: Algorithms and strong consistency. *arXiv preprint arXiv:2011.06461*.
- Paul, D., Chakraborty, S., Li, D., and Dunson, D. (2020b). Principal ellipsoid analysis (pea): Efficient non-linear dimension reduction & clustering. *arXiv preprint arXiv:2008.07110*.
- Pearson, K. (1901). Liii. on lines and planes of closest fit to systems of points in space. *The London, Edinburgh, and Dublin Philosophical Magazine and Journal of Science*, 2(11):559–572.
- Pollard, D. (1981). Strong consistency of k -means clustering. *Ann. Statist.*, 9(1):135–140.
- Rayana, S. (2016). ODDS library.
- Schölkopf, B., Smola, A., and Müller, K.-R. (1997). Kernel principal component analysis. In *International conference on artificial neural networks*, pages 583–588. Springer.
- Shalev-Shwartz, S. and Ben-David, S. (2014). *Understanding machine learning: From theory to algorithms*. Cambridge university press.
- Tipping, M. E. and Bishop, C. M. (1999). Probabilistic principal component analysis. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 61(3):611–622.
- Vapnik, V. (2013). *The nature of statistical learning theory*. Springer science & business media.
- Wold, S., Esbensen, K., and Geladi, P. (1987). Principal component analysis. *Chemometrics and intelligent laboratory systems*, 2(1-3):37–52.
- Wright, J., Ganesh, A., Rao, S., Peng, Y., and Ma, Y. (2009). Robust principal component analysis: Exact recovery of corrupted low-rank matrices via convex optimization. In *Advances in Neural Information Processing Systems*, volume 22, pages 2080–2088.
- Yang, B., Fu, X., Sidiropoulos, N. D., and Hong, M. (2017). Towards k-means-friendly spaces: Simultaneous deep learning and clustering. In *international conference on machine learning*, pages 3861–3870. PMLR.
- Zhang, C. and Woodland, P. C. (2017). Joint optimisation of tandem systems using gaussian mixture density neural network discriminative sequence training. In *2017 IEEE International Conference on Acoustics, Speech and Signal Processing (ICASSP)*, pages 5015–5019. IEEE.
- Zhang, R. and Tong, H. (2019). Robust principal component analysis with adaptive neighbors. In *Advances in Neural Information Processing Systems*, volume 32, pages 6961–6969.

Zong, B., Song, Q., Min, M. R., Cheng, W., Lumezanu, C., Cho, D., and Chen, H. (2018). Deep autoencoding gaussian mixture model for unsupervised anomaly detection. In *International Conference on Learning Representations*.

Zou, H., Hastie, T., and Tibshirani, R. (2006). Sparse principal component analysis. *Journal of computational and graphical statistics*, 15(2):265–286.