

EDGEWORTH APPROXIMATIONS FOR DISTRIBUTIONS OF SYMMETRIC STATISTICS

FRIEDRICH GÖTZE¹ AND MINDAUGAS BLOZNELIS¹

*In memoriam Willem Rutger van Zwet *March 31, 1934 †July 2, 2020*

ABSTRACT. We study the distribution of a general class of asymptotically linear statistics which are symmetric functions of N independent observations. The distribution functions of these statistics are approximated by an Edgeworth expansion with a remainder of order $o(N^{-1})$. The Edgeworth expansion is based on Hoeffding's decomposition which provides a stochastic expansion into a linear part, a quadratic part as well as smaller higher order parts. The validity of this Edgeworth expansion is proved under Cramér's condition on the linear part, moment assumptions for all parts of the statistic and an optimal dimensionality requirement for the non linear part.

1991 *Mathematics Subject Classification.* Primary 62E20; Secondary 60F05

Key words and phrases. Edgeworth expansion, Littlewood-Offord problem, concentration in Banach spaces, symmetric statistic, U -statistic, Hoeffding decomposition

1. INTRODUCTION AND RESULTS

1.1. By the classical central limit theorem the distributions of sums $X_1 + \dots + X_N$ of independent and identically distributed random variables can be approximated by the normal distribution. The accuracy of the normal approximation is of order $O(N^{-1/2})$ by the well-known Berry-Esseen theorem.

A function of observations X_1, \dots, X_N is called *linear statistic* if it can be represented by a sum of functions depending on a single observation only. Many important statistics are non linear, but can be approximated by a linear statistic. We call these statistics *asymptotically linear*. The central limit theorem and the normal approximation with rate $O(N^{-1/2})$ extend to the class of asymptotically linear statistics as well. For comparisons of performance of statistical procedures beyond efficiency of first order, that is $o(N^{-1/2})$, at the level of Hodges and Lehmann deficiency, that is $o(N^{-1})$, more precise approximations beyond the normal approximation are required. Such a refinement is provided by Edgeworth expansions of the distribution function. The one-term Edgeworth expansion adds a correction term of order $O(N^{-1/2})$ to the standard normal distribution function $\Phi(x) = \int_{-\infty}^x (2\pi)^{-1/2} e^{-u^2/2} du$ and provides an approximation with the error $o(N^{-1/2})$. Similarly, the two-term Edgeworth expansion includes correction terms of orders $O(N^{-1/2})$ and $O(N^{-1})$ as well with an approximation error of order $o(N^{-1})$.

Normal approximation theory including Edgeworth expansions is well studied for distribution function $F_N(x) = \mathbf{P}\{X_1 + \dots + X_N \leq N\mathbf{E}X_1 + N^{1/2}\sigma x\}$, $\sigma^2 = \mathbf{Var}X_1$ of a sum of independent random variables, see Petrov (1975) [22]. Sums of independent random vectors are considered in Bhattacharya and Rao (1986) [4]. One distinguishes two cases. For summands taking values in an arithmetic progression (lattice case) the distribution function of the sum has jumps of order $O(N^{-1/2})$. Corresponding asymptotic expansions are discontinuous functions designed to capture these jumps see the seminal results by Esseen (1945) [15]. For sums of non-lattice

Date: September 1, 2005.

¹Research supported by CRC 701.

random variables (non-lattice case) the one-term Edgeworth expansion

$$\Phi(x) - \frac{1}{\sqrt{N}} \frac{\kappa_3}{6\sigma^3} (x^2 - 1)\Phi'(x)$$

is a differentiable function. The correcting term reflects the skewness of the distribution of a summand, $\kappa_3 = \mathbf{E}(X_1 - \mathbf{E}X_1)^3$. More generally, a k -term Edgeworth expansion is a differentiable function with all derivatives bounded (it is a sum of Hermite polynomials of increasing order with scalar weights involve cumulants of order at least three of X_1 which vanish for X_1 being Gaussian). Therefore, in order to establish the validity of such an expansion, that is to prove the bound $o(N^{-k/2})$ for the remainder, one should assume that the underlying distribution is sufficiently smooth (see Bickel and Robinson 1982 [9]). A convenient condition to ensure that is Cramér's condition (C):

$$\limsup_{|t| \rightarrow \infty} |\mathbf{E} \exp\{itX_1\}| < 1. \quad (C)$$

In this paper we establish the two-term Edgeworth expansion for a general asymptotically linear statistic $T = T(X_1, \dots, X_N)$ with non-lattice distribution.

There is a rich literature devoted to normal approximation and Edgeworth expansions for various classes of asymptotically linear statistics (see e.g. Babu and Bai (1993) [1], Bai and Rao (1991) [2], Bentkus, Götze and van Zwet (1997) [3], Bhattacharya and Ghosh (1978, 1980) [5, 6], Bhattacharya and Rao (1986) [4], Bickel (1974) [7], Bickel, Götze and van Zwet (1986) [8], Callaert, Janssen and Veraverbeke (1980) [11], Chibisov (1980) [12], Hall (1987) [18], Helmers (1982) [19], Petrov (1975) [22], Pfanzagl (1985) [23], Serfling (1980) [24], etc.

A wide class of statistics can be represented as functions of sample means of vector variables. Edgeworth expansions of such statistics can be obtained by applying the multivariate expansion to corresponding functions, see Bhattacharya and Ghosh (1978, 1980) [5, 6]. In their work the crucial Cramér condition (C) is assumed on the joint distribution of all the components of a vector which may be too restrictive in cases where some components have a negligible influence on the statistic. More often only one or a few of the components satisfy a conditional version of condition (C). Bai and Rao (1991) [2], Babu and Bai (1993) [1] established Edgeworth expansions for functions of sample means under such a conditional Cramér condition. This approach exploits the smoothness of the distribution of vector as well as the smoothness of the function defining the statistic. In particular this approach needs a class of statistics which are smooth functions of observations or can be approximated by such functions via Taylor's expansion, see also Chibisov (1980) [12].

Let us note that the smoothness of the distribution function of a statistic, say $T = \phi(X_1, \dots, X_N)$ may have little to do with the smoothness of the kernel ϕ . Just take Gini's mean difference $\sum_{i < j} |X_i - X_j|$ with absolutely continuous X_i for example. The aim of this paper is to establish the validity of the two-term Edgeworth expansion for general asymptotically linear symmetric statistics in the following setup. We require a minimal Cramér type condition for the linear part of the statistic T (given by its L^2 projection to the space of linear statistics of X_1, \dots, X_N), and require instead of smoothness of ϕ of these observations that the L^2 -complementary non linear part of T does not lie in a linear subspace in L^2 generated by the linear part.

Remark. *Determining in this setup the actual influence of the non linear terms of T on the approximation error represents a considerable challenge. The crucial problem, which required new techniques, was to control the quasi periodic behavior of upper bounds for the characteristic function of T at frequencies $t \sim N$. These upper bounds involve bounds on conditional characteristic functions of the non linear part of T , say $g(t, \mathbf{Y})^m$, given a subset $\mathbf{Y} := \mathbf{X}_I$ of size $N - m$ of the observations. Understanding the separation of large maxima of $(t, \mathbf{Y}) \rightarrow g(t, \mathbf{Y})^m$ in terms of both arguments in a function space setup was finally achieved by a combinatorial argument of Kleitman on symmetric partitions (see Section 4) controlling the concentration of*

sums like in the Littlewood-Offord problem in Banach spaces. The separation of large maxima of $g(t, \mathbf{Y})^m$ then allowed to prove the desired bounds when averaging over t and \mathbf{Y} . Note however, that more standard analytical arguments for concentration bounds would not work in the required generality here.

2.1. Let X, X_1, X_2, \dots, X_N be independent and identically distributed random variables taking values in a measurable space $(\mathcal{X}, \mathcal{B})$. Let P_X denotes the distribution of X on $(\mathcal{X}, \mathcal{B})$. We assume that $\mathbb{T}(X_1, \dots, X_N)$ is a symmetric function of its arguments (symmetric statistic, for short). Furthermore, we assume that the moments $\mathbf{E}\mathbb{T}$ and $\sigma_{\mathbb{T}}^2 := \mathbf{Var}\mathbb{T}$ are finite.

Our approach is based on Hoeffding's decomposition of \mathbb{T} , see Hoeffding (1948) [20], Efron and Stein (1981) [14] and van Zwet (1984) [26]. Hoeffding's decomposition expands \mathbb{T} into the series of centered and mutually uncorrelated U -statistics of increasing order

$$\begin{aligned} \mathbb{T} = \mathbf{E}\mathbb{T} &+ \frac{1}{N^{1/2}} \sum_{1 \leq i \leq N} g(X_i) + \frac{1}{N^{3/2}} \sum_{1 \leq i < j \leq N} \psi(X_i, X_j) \\ &+ \frac{1}{N^{5/2}} \sum_{1 \leq i < j < k \leq N} \chi(X_i, X_j, X_k) + \dots \end{aligned}$$

Let L, Q and K denote the first, the second and the third sum. We call L the linear part, Q the quadratic part and K the cubic part of the decomposition.

We shall assume that the linear part does not vanish, that is, $\mathbf{Var}L^2 > 0$. If, for large N , the linear part dominates the statistic we call \mathbb{T} asymptotically linear. The distribution of an asymptotically linear statistic can be approximated by the normal distribution, via the central limit theorem. An improvement over the normal approximation is obtained by using Edgeworth expansions for the distribution function $\mathbb{F}(x) = \mathbf{P}\{\mathbb{T} - \mathbf{E}\mathbb{T} \leq \sigma_{\mathbb{T}}x\}$. For this purpose we write Hoeffding's decomposition in the form

$$(1) \quad \mathbb{T} - \mathbf{E}\mathbb{T} = L + Q + K + R,$$

where R denotes the remainder. For a number of important examples of asymptotically linear statistics we have $R/\sigma_{\mathbb{T}} = o_P(N^{-1})$ (in probability) as $N \rightarrow \infty$. Therefore, the U -statistic $\sigma_{\mathbb{T}}^{-1}(L+Q+K)$ can be viewed as a stochastic expansion of $(\mathbb{T} - \mathbf{E}\mathbb{T})/\sigma_{\mathbb{T}}$ up to the order $o_P(N^{-1})$. Furthermore, an Edgeworth expansion of $\sigma_{\mathbb{T}}^{-1}(L + Q + K)$ can be used to approximate $\mathbb{F}(x)$. Introduce the following two term Edgeworth expansion of the distribution function of $\sigma_{\mathbb{T}}^{-1}(L + Q + K)$

$$(2) \quad \begin{aligned} G(x) &= \Phi(x) - \frac{1}{\sqrt{N}} \frac{\kappa_3}{6} (x^2 - 1)\Phi'(x) \\ &- \frac{1}{N} \left(\frac{\kappa_3^2}{72} (x^5 - 10x^3 + 15x)\Phi'(x) + \frac{\kappa_4}{24} (x^3 - 3x)\Phi'(x) \right). \end{aligned}$$

Here Φ respectively Φ' denote the standard normal distribution function and its derivative. Furthermore, we introduce $\sigma^2 = \mathbf{E}g^2(X_1)$ and

$$\begin{aligned} \kappa_3 &= \sigma^{-3} \left(\mathbf{E}g^3(X_1) + 3\mathbf{E}g(X_1)g(X_2)\psi(X_1, X_2) \right), \\ \kappa_4 &= \sigma^{-4} \left(\mathbf{E}g^4(X_1) - 3\sigma^4 + 12\mathbf{E}g^2(X_1)g(X_2)\psi(X_1, X_2) \right. \\ &\quad \left. + 12\mathbf{E}g(X_1)g(X_2)\psi(X_1, X_3)\psi(X_2, X_3) \right. \\ &\quad \left. + 4\mathbf{E}g(X_1)g(X_2)g(X_3)\chi(X_1, X_2, X_3) \right). \end{aligned}$$

Our main result, Theorem 1 below, establishes a bound $o(N^{-1})$ for the Kolmogorov distance

$$\Delta = \sup_{x \in \mathbb{R}} |\mathbb{F}(x) - G(x)|.$$

We shall consider a general situation where the kernel $\mathbb{T} = \mathbb{T}^{(N)}$, the space $(\mathcal{X}, \mathcal{B}) = (\mathcal{X}^{(N)}, \mathcal{B}^{(N)})$ and the distribution $P_X = P_X^{(N)}$ all depend on N as $N \rightarrow \infty$. In order to keep the notation simple we drop the subscript N in what follows.

3.1. Let us introduce the conditions we need in order to prove the bound $\Delta = o(N^{-1})$.

(i) *Moment conditions.* Assume that, for some absolute constants $0 < A_* < 1$ and $M_* > 0$ and numbers $r > 4$ and $s > 2$, we have

$$(3) \quad \begin{aligned} \mathbf{E}g^2(X_1) &> A_*\sigma_{\mathbb{T}}^2, & \mathbf{E}|g(X_1)|^r &< M_*\sigma_{\mathbb{T}}^r, \\ \mathbf{E}|\psi(X_1, X_2)|^r &< M_*\sigma_{\mathbb{T}}^r, & \mathbf{E}|\chi(X_1, X_2, X_3)|^s &< M_*\sigma_{\mathbb{T}}^s. \end{aligned}$$

These moment conditions refer to the linear, the quadratic and the cubic part of \mathbb{T} . In order to control the remainder R of the approximation (1) we use moments of differences introduced in Bentkus, Götze and van Zwet (1997) [3], see also van Zwet (1984) [26]. Define, for $1 \leq i \leq N$,

$$D_i\mathbb{T} = \mathbb{T} - \mathbf{E}_i\mathbb{T}, \quad \mathbf{E}_i\mathbb{T} := \mathbf{E}(\mathbb{T}|X_1, \dots, X_{i-1}, X_{i+1}, \dots, X_N).$$

A subsequent application of difference operations D_i, D_j, \dots , (the indices i, j, \dots , are all distinct) produce higher order differences, like

$$D_iD_j\mathbb{T} := D_i(D_j\mathbb{T}) = \mathbb{T} - \mathbf{E}_i\mathbb{T} - \mathbf{E}_j\mathbb{T} + \mathbf{E}_i\mathbf{E}_j\mathbb{T}.$$

For $m = 1, 2, 3, 4$ write $\Delta_m^2 = \mathbf{E}|N^{m-1/2}D_1D_2 \dots D_m\mathbb{T}|^2$.

We shall assume that for some absolute constant $D_* > 0$ and number $\nu_1 \in (0, 1/2)$ we have

$$(4) \quad \Delta_4^2/\sigma_{\mathbb{T}}^2 \leq N^{1-2\nu_1}D_*$$

For a number of important examples of asymptotically linear statistics the moments Δ_m^2 are evaluated or estimated in [3]. Typically we have $\Delta_m^2/\sigma_{\mathbb{T}}^2 = O(1)$ for some m . Therefore, assuming that (4) holds uniformly in N as $N \rightarrow \infty$, we obtain from the inequality $\mathbf{E}R^2 \leq N^{-3}\Delta_4^2$, see (162) (see Appendix), that $R/\sigma_{\mathbb{T}} = O_P(N^{-1-\nu_1})$. Furthermore, assuming that (3), (4) hold uniformly in N as $N \rightarrow \infty$, we obtain from (162), (161), see Appendix, that $\sigma^2/\sigma_{\mathbb{T}}^2 = (1 - O(N^{-1}))$.

(ii) *Cramér type smoothness condition.* Introduce the function

$$\rho(a, b) = 1 - \sup\{|\mathbf{E} \exp\{itg(X_1)/\sigma\}| : a \leq |t| \leq b\}.$$

We shall assume that, for some $\delta > 0$ and $\nu_2 > 0$, we have

$$(5) \quad \rho(\beta_3^{-1}, N^{\nu_2+1/2}) \geq \delta.$$

Here $\beta_3 = \sigma^{-3}\mathbf{E}|g(X_1)|^3$.

It was shown in Götze and van Zwet (1992) [17], see as well Theorem 1.4 of [3], that moment conditions (like (3) and (4)) together with Cramér's condition (on the summand $g(X_i)$ of the linear part) do not suffice to establish the bound $\Delta = o(N^{-1})$. For convenience we state this result in Example 1 below.

Example 1. Let X_1, X_2, \dots be independent random variables uniformly distributed on the interval $(-1/2, 1/2)$. Define $T_N = (W_N + N^{-1/2}V_N)(1 - N^{-1/2}V_N)$, where $V_N = N^{-1/2} \sum\{N^{1/2}X_j\}$ and $W_N = N^{-1} \sum[N^{1/2}X_j]$. Here $[x]$ denotes the nearest integer to x and $\{x\} = x - [x]$.

Assume that $N = m^2$, where m is odd. We have, by the local limit theorem,

$$\mathbf{P}\{W_N = 1\} \geq cN^{-1} \quad \text{and} \quad \mathbf{P}\{|V_N| < \delta\} > c\delta, \quad 0 < \delta < 1,$$

where $c > 0$ is an absolute constant. From these inequalities it follows by the independence of W_N and V_N , that $\mathbf{P}\{1 - \delta^2N^{-1} \leq T_N \leq 1\} \geq c^2\delta N^{-1}$.

The example defines a sequence of U -statistics \mathbb{T}_N whose distribution functions \mathbb{F}_N have $O(N^{-1})$ sized increments in a particular interval of length $o(N^{-1})$. These fluctuations of magnitude $O(N^{-1})$ appear as a result of a nearly lattice structure induced by the interplay between the

(smooth) linear part and the quadratic part. In order to avoid examples with such a (conditional) lattice structure a simple moment condition was introduced in Götze and van Zwet (1992) [17] which, separates (in L^2 distance) the random variable $\psi(X_1, X_2)$ from any random variable of the form $\psi_h(X_1, X_2) = h(X_1)g(X_2) + g(X_1)h(X_2)$, h -measurable. Note that the L^2 distance $\mathbf{E}(\psi(X_1, X_2) - \psi_h(X_1, X_2))^2$ is minimized by $h(x) = b(x)$, where

$$b(x) = \sigma^{-2} \mathbf{E}(\psi(X_1, X_2)g(X_2)|X_1 = x) - (\kappa/2\sigma^4)g(x).$$

Here $\kappa = \mathbf{E}\psi(X_1, X_2)g(X_1)g(X_2)$.

Therefore, we assume that, for some absolute constant $\delta_* > 0$, we have

$$(6) \quad \mathbf{E}\left(\psi(X_1, X_2) - (b(X_1)g(X_2) + b(X_2)g(X_1))\right)^2 \geq \delta_*^2 \sigma_{\mathbb{T}}^2.$$

Define $\nu = 600^{-1} \min\{\nu_1, \nu_2, s - 2, r - 4, \}$.

Theorem 1. *Let $N \geq 4$. Assume that for some absolute constants $A_*, M_*, D_* > 0$ and numbers $r > 4, s > 2, \nu_1, \nu_2 > 0$ and $\delta, \delta_* > 0$, the conditions (3), (4), (5), (6) hold. Then there exists a constant $C_* > 0$ depending only on $A_*, M_*, D_*, r, s, \nu_1, \nu_2, \delta, \delta_*$ such that*

$$\Delta \leq C_* N^{-1-\nu} (1 + \delta_*^{-1} N^{-\nu}).$$

In particular case of U statistics of degree three (the case where $R \equiv 0$ in (1)) the proof of Theorem 1 has been outlined in an unpublished paper by Götze and van Zwet (1992) [17]. We provide a complete and more readable version of the arguments sketched in that preprint and extend them to a general class of symmetric statistics.

Remark 1. Condition (6) can be relaxed. Assume that for some absolute constant G_* we have

$$(7) \quad \mathbf{E}\left(\psi(X_1, X_2) - (b(X_1)g(X_2) + b(X_2)g(X_1))\right)^2 \geq N^{-2\nu} G_* \sigma_{\mathbb{T}}^2.$$

The bound of Theorem 1 holds if we replace (6) by this weaker condition. In this case we have $\Delta \leq C_* N^{-1-\nu}$, where the constant C_* depends on $A_*, D_*, G_*, M_*, r, s, \nu_1, \nu_2, \delta$.

Remark 2. Consider a sequence of statistics $T^{(N)} = T^{(N)}(X_{N1}, \dots, X_{NN})$ based on independent observations X_{N1}, \dots, X_{NN} taking values in $(\mathcal{X}^{(N)}, \mathcal{B}^{(N)})$ and with the common distribution $P_X^{(N)}$. Assume that conditions (3), (4) and (6) (or (7)) hold uniformly in $N = N_0, N_0 + 1, \dots$, for some N_0 . Theorem 1 implies the bound $\Delta = o(N^{-1})$ as $N \rightarrow \infty$.

Remark 3. The value of $\nu = 600^{-1} \min\{\nu_1, \nu_2, s - 2, r - 4, \}$ is far from being optimal. Furthermore, the moment conditions (3) and (4) are not the weakest possible that would ensure the approximation of order $o(N^{-1})$. The condition (3) can likely be reduced to the moment conditions that are necessary to define Edgeworth expansion terms κ_3 and κ_4 , similarly, (4) can be reduced to $\Delta_4^2/\sigma_{\mathbb{T}}^2 = o(N^{-1})$. No effort was made to obtain the result under the optimal conditions. This would increase the complexity of the proof which is already rather involved.

4.1. In order to compare Theorem 1 with earlier results of similar nature let us consider the case of U -statistics of degree two

$$(8) \quad \mathbb{U} = \frac{\sqrt{N}}{2} \binom{N}{2}^{-1} \sum_{1 \leq i < j \leq N} h(X_i, X_j),$$

where $h(\cdot, \cdot)$ denotes a (fixed) symmetric kernel. Assume for simplicity of notation and without loss of generality that $\mathbf{E}h(X_1, X_2) = 0$. Write $h_1(x) = \mathbf{E}(h(X_1, X_2)|X_1 = x)$ and assume that $\sigma_h^2 > 0$, where $\sigma_h^2 = \mathbf{E}h_1^2(X_1)$. In this case Hoeffding's decomposition (1) reduces to $\mathbb{U} = L + Q$, where, by the assumption $\sigma_h^2 > 0$, we have $\mathbf{Var}L > 0$. Since the cubic part vanishes we may remove the moment $\mathbf{E}g(X_1)g(X_2)g(X_3)\chi(X_1, X_2, X_3)$ from the expression for κ_4 . In this way we

obtain the two term Edgeworth expansion (2) for the distribution function $\mathbb{F}_U(x) = \mathbf{P}\{\mathbb{U} \leq \sigma_{\mathbb{U}}x\}$ with $\sigma_{\mathbb{U}}^2 := \mathbf{Var}\mathbb{U}$.

We call a kernel h reducible if for some measurable functions $u, v : \mathcal{X} \rightarrow R$ we have $h(x, y) - \mathbf{E}h(X_1, X_2) = v(x)u(y) + v(y)u(x)$ for $P_X \times P_X$ almost sure $(x, y) \in \mathcal{X} \times \mathcal{X}$. A simple calculation shows that for a sequence of U -statistics (8) with a fixed non-reducible kernel the condition (6) is satisfied, for some $\delta_* > 0$, uniformly in N . A straightforward consequence of Theorem 1 is the following corollary. Write $\tilde{\nu} = 600^{-1} \min\{\nu_2, r - 4, 1\}$.

Corollary 1. *Let $N \geq 4$. Assume that for some $r > 4$*

$$(9) \quad \mathbf{E}|h(X_1, X_2)|^r < \infty.$$

Assume that $\sigma_h^2 > 0$ and the kernel h is non-reducible and that for some $\delta > 0$

$$(10) \quad \sup\{|\mathbf{E}e^{it\sigma_h^{-1}h_1(X_1)}| : |t| \geq \beta_3^{-1}\} \leq 1 - \delta.$$

Then there exist a constant $C_ > 0$ such that*

$$\sup_{x \in \mathbb{R}} |\mathbb{F}_U(x) - G(x)| \leq C_* N^{-1-\tilde{\nu}}$$

For U -statistics with fixed kernel h the validity of the Edgeworth expansion (2) up to the order $o(N^{-1})$ was established by Callaert, Janssen and Veraverbeke (1980) [11] and Bickel, Götze and van Zwet (1986) [8]. In addition to the moment conditions (like (9)) and Cramér's condition (like (10)) they imposed the following rather implicit conditions which ensure the regularity of $\mathbb{F}_U(x)$. Callaert, Janssen and Veraverbeke (1980) [11] assumed that for some $0 < c < 1$ and $0 < \alpha < 1/8$ the event

$$(11) \quad \left| \mathbf{E}(\exp\{it\sigma_{\mathbb{U}}^{-1} \sum_{j=m+1}^N h(X_1, X_j)\} | X_{m+1}, \dots, X_N) \right| \leq c$$

has probability $1 - o(1/N \log N)$ uniformly for all $t \in [N^{3/4}/\log N, N \log N]$. Here $m \approx N^\alpha$.

Bickel, Götze and van Zwet (1986) [8] more explicitly required that the linear operator, $f(\cdot) \rightarrow \mathbf{E}\psi(X, \cdot)f(X)$ defined by ψ has sufficiently large number of non-zero eigenvalues (depending on the existing moments).

Both of these conditions correspond to some techniques used in the parts of our proof. Recall that in the first step of the proof, one reduces the problem of bounding $|\mathbb{F}_U(x) - G(x)|$ by means of Berry-Esseen's smoothing inequality, to that of bounding the difference between the corresponding Fourier transforms $|\hat{\mathbb{F}}_U(t) - \hat{G}(t)|$ in the region $|t| < N\varepsilon_N^{-1}$, for some $\varepsilon_N \downarrow 0$. For $|t| \approx N$ one writes $|\hat{\mathbb{F}}_U(t) - \hat{G}(t)| \leq |\hat{\mathbb{F}}_U(t)| + |\hat{G}(t)|$ and bounds every summand separately. The condition (11) applies more or less directly and shows exponential decay of $|\hat{\mathbb{F}}_U(t)|$ as $N \rightarrow \infty$, for $|t| \approx N$. The eigenvalue condition achieves the same goal, but refers to a more sophisticated approach based on a symmetrization technique introduced by Götze (1979) [16].

The condition (6) provides an alternative to (11) and the eigenvalue condition. It is aimed to exclude the situations, where the interplay between the linear and the quadratic part produces a nearly lattice structure of \mathbb{U} , which in turn results in fluctuations of $\mathbb{F}_U(x)$ of magnitude $O(N^{-1})$. The proof of Theorem 1 uses a result of Kleitman which provides a solution to multidimensional Littlewood-Offord problem. This result establishes bounds for probabilities of the concentration of sums of random variables with values in multidimensional spaces.

Remark 4. The U -statistic (8) with a kernel $h(x, y) = v(x)u(y) + v(y)u(x)$, where $\mathbf{E}u(X_1) = 0$, violates (6). Let us note that in this case one can establish the validity of an Edgeworth expansion with the remainder $o(N^{-1})$, under the following bivariate Cramér's condition, which is essentially more restrictive than condition (ii),

$$1 - \sup\{|\mathbf{E}\exp\{i\frac{t}{\sigma}u(X_1) + i\frac{s}{\sigma}v(X_1)\}| : \beta_3^{-1} < |t| \leq N^{\nu+1/2}, |s| \leq N^\nu\} > \delta,$$

for some $\delta, \nu > 0$. Note that from this condition we immediately obtain the desired exponential decay of $\hat{\mathbb{F}}(t)$ as $N \rightarrow \infty$ for $|t| \approx N$.

The remaining parts of the paper (Sections 2–5) contain the proof of Theorem 1. Auxiliary results are placed in the Appendix.

2. PROOF OF THEOREM 1

The proof combines various techniques developed in earlier papers by Callaert, Janssen and Veraverbeke (1980) [11], Bickel, Götze and van Zwet (1986) [8]. It is based on the manuscript of Götze and van Zwet (1992) [17]. The later paper introduces the condition (6), provides the crucial counter example, see Example 1 above, and contains an outline of the proof in the particular case of U -statistics of degree three ($\mathbb{T} - \mathbf{E}\mathbb{T} = L + Q + K$). In order to extend these arguments to general symmetric statistics we combine stochastic expansions by means of Hoeffding's decomposition and bounds for various parts of the decomposition. This approach was introduced in van Zwet (1984) [26] and further developed in [3].

2.1. Let us start with an outline of the proof. Firstly, using the linear structure induced by Hoeffding's decomposition we replace $\mathbb{T}/\sigma_{\mathbb{T}}$ by the statistic $\tilde{\mathbb{T}}$ which is conditionally linear given X_{m+1}, \dots, X_N . Secondly, invoking a smoothing inequality we pass from distribution functions to Fourier transforms. In the remaining steps we bound the difference $\delta(t) = \mathbf{E}e^{it\tilde{\mathbb{T}}} - \hat{G}(t)$, for $|t| \leq N^{1+\nu}$. For "small frequencies" $|t| \leq CN^{1/2}$, we expand the characteristic function $\mathbf{E}e^{it\tilde{\mathbb{T}}}$ in order to show that $\delta(t) = o(N^{-1})$. For remaining range of frequencies, that is $CN^{1/2} \leq |t| \leq N^{1+\nu}$, we bound the summands $\mathbf{E}e^{it\tilde{\mathbb{T}}}$ and $\hat{G}(t)$ separately. The cases of "large frequencies", that is $N^{1-\nu} \leq |t| \leq N^{1+\nu}$, and "medium frequencies", that is $C\sqrt{N} \leq |t| \leq N^{1-\nu}$, are treated in a different manner. For medium frequencies the Cramér type condition (5) ensures an exponential decay of $|\mathbf{E}e^{it\tilde{\mathbb{T}}}|$ as $N \rightarrow \infty$. For large frequencies we combine conditions (5) and (6). Here we apply a *combinatorial* concentration bound due to Kleitman as described in the introduction.

2.2. Before starting the proof we introduce some notation. By c_* we shall denote a positive constant which may depend only on $A_*, D_*, M_*, r, s, \nu_1, \nu_2, \delta$, but it does not depend on N . In different places the values of c_* may be different.

It is convenient to write the decomposition in the form

$$(12) \quad \mathbb{T} = \mathbf{E}\mathbb{T} + \sum_{1 \leq k \leq N} \mathbb{U}_k, \quad \mathbb{U}_k = \sum_{1 \leq i_1 < \dots < i_k \leq N} g_k(X_{i_1}, \dots, X_{i_k}),$$

where, for every k , the symmetric kernel g_k is centered, i.e., $\mathbf{E}g_k(X_1, \dots, X_k) = 0$, and satisfies, see, e.g., [3],

$$(13) \quad \mathbf{E}(g_k(X_1, \dots, X_k) | X_2, \dots, X_k) = 0 \quad \text{almost surely.}$$

Here we write $g_1 := N^{-1/2}g$, $g_2 := N^{-3/2}\psi$ and $g_3 := N^{-5/2}\chi$. Furthermore, for an integer $k > 0$ write $\Omega_k = \{1, \dots, k\}$. Given a subset $A = \{i_1, \dots, i_k\} \subset \Omega_N$ we write, for short, $T_A := g_k(X_{i_1}, \dots, X_{i_k})$. Put $T_\emptyset := \mathbf{E}\mathbb{T}$. Now the decomposition (12) can be written as follows

$$\mathbb{T} = \mathbf{E}\mathbb{T} + \sum_{1 \leq k \leq N} \mathbb{U}_k = \sum_{A \subset \Omega_N} T_A, \quad \mathbb{U}_k = \sum_{|A|=k, A \subset \Omega_N} T_A.$$

2.3. Proof of Theorem 1. Throughout the proof we assume without loss of generality that

$$(14) \quad 4 < r \leq 5, \quad 2 < s \leq 3 \quad \text{and} \quad \mathbf{E}\mathbb{T} = 0, \quad \sigma_{\mathbb{T}}^2 = 1.$$

Denote, for $t > 0$,

$$\beta_t = \sigma^{-t} \mathbf{E}|g(X_1)|^t, \quad \gamma_t = \mathbf{E}|\psi(X_1, X_2)|^t, \quad \zeta_t = \mathbf{E}|\chi(X_1, X_2, X_3)|^t.$$

The linearization step. Choose number $\nu > 0$ and integer m such that

$$(15) \quad \nu = 600^{-1} \min\{\nu_1, \nu_2, s-2, r-4\}, \quad m \approx N^{100\nu}.$$

Split

$$\mathbb{T} = \mathbb{T}_{[m]} + \mathbb{W}, \quad \mathbb{T}_{[m]} = \sum_{A: A \cap \Omega_m \neq \emptyset} T_A, \quad \mathbb{W} = \sum_{A: A \cap \Omega_m = \emptyset} T_A.$$

Furthermore, write

$$\begin{aligned} \mathbb{T}_{[m]} &= \mathbb{U}_1^* + \mathbb{U}_2^* + \Lambda, & \Lambda &= \Lambda_1 + \Lambda_2 + \Lambda_3 + \Lambda_4 + \Lambda_5, \\ \mathbb{U}_1^* &= \sum_{i=1}^m T_{\{i\}}, & \mathbb{U}_2^* &= \sum_{i=1}^m \sum_{j=m+1}^N T_{\{i,j\}}, \\ \Lambda_1 &= \sum_{1 \leq i < j \leq m} T_{\{i,j\}}, & \Lambda_2 &= \sum_{|A| \geq 3, |A \cap \Omega_m| = 2} T_A, \\ \Lambda_3 &= \sum_{A: |A \cap \Omega_m| \geq 3} T_A, & \Lambda_4 &= \sum_{|A|=3, |A \cap \Omega_m|=1} T_A, \\ \Lambda_5 &= \sum_{i=1}^m \eta_i, & \eta_i &= \sum_{|A| \geq 4, A \cap \Omega_m = \{i\}} T_A. \end{aligned}$$

Before applying a smoothing inequality we replace $\mathbb{F}(x)$ by

$$\tilde{\mathbb{F}}(x) := \mathbf{P}\{\tilde{\mathbb{T}} \leq x\}, \quad \text{where} \quad \tilde{\mathbb{T}} = \mathbb{U}_1^* + \mathbb{U}_2^* + \mathbb{W} = \mathbb{T} - \Lambda.$$

In order to show that Λ can be neglected we apply a simple Slutsky type argument. Given $\varepsilon > 0$, we have

$$(16) \quad \Delta \leq \sup_{x \in \mathbb{R}} |\tilde{\mathbb{F}}(x) - G(x)| + \varepsilon \sup_{x \in \mathbb{R}} |G'(x)| + \mathbf{P}\{|\Lambda| > \varepsilon\}.$$

From Lemma 5 we obtain via Chebyshev's inequality, for $\varepsilon = N^{-1-\nu}$,

$$\begin{aligned} \mathbf{P}\{|\Lambda| > \varepsilon\} &\leq \sum_{i=1}^5 \mathbf{P}\{|\Lambda_i| > \frac{\varepsilon}{5}\} \\ &\leq \left(\frac{5}{\varepsilon}\right)^3 \mathbf{E}|\Lambda_1|^3 + \left(\frac{5}{\varepsilon}\right)^2 (\mathbf{E}\Lambda_2^2 + \mathbf{E}\Lambda_3^2 + \mathbf{E}\Lambda_5^2) + \left(\frac{5}{\varepsilon}\right)^s \mathbf{E}|\Lambda_4|^s \\ &\leq c_* N^{-1-\nu}. \end{aligned}$$

In the last step we used conditions (3), (4) and the inequality (163). Furthermore, using (3) and (4) one can show that

$$(17) \quad \sup_{x \in \mathbb{R}} |G'(x)| \leq c_*.$$

Therefore, (16) implies

$$\Delta \leq \tilde{\Delta} + c_* N^{-1-\nu}, \quad \tilde{\Delta} := \sup_{x \in \mathbb{R}} |\tilde{\mathbb{F}}(x) - G(x)|.$$

It remains to show that $\tilde{\Delta} \leq c_* N^{-1-\nu}$.

A smoothing inequality. Given $a > 0$ and even integer $k \geq 2$ consider the probability density function, see (10.7) in Bhattacharya and Rao (1986) [4],

$$(18) \quad x \rightarrow g_{a,k}(x) = a c(k) (ax)^{-k} \sin^k(ax),$$

where $c(k)$ is the normalizing constant. Its characteristic function

$$\hat{g}_{a,k}(t) = \int_{-\infty}^{+\infty} e^{itx} g_{a,k}(x) dx = 2\pi a c(k) u_{[-a,a]}^{*k}(t)$$

vanishes outside the interval $|t| \leq ka$. Here $u_{[-a,a]}^{*k}(t)$ denotes the probability density function of the sum of k independent random variables each uniformly distributed in $[-a, a]$. It is easy to show that the function $t \rightarrow \hat{g}_{a,k}(t)$ is unimodal and symmetric around $t = 0$.

Let μ be the probability distribution with the density $g_{a,2}$, where a is chosen to satisfy $\mu([-1, 1]) = 3/4$. Given $T > 1$ define $\mu_T(\mathcal{A}) = \mu(T\mathcal{A})$, for $\mathcal{A} \subset \mathbb{R}$ -measurable. Let $\hat{\mu}_T$ denote the characteristic function corresponding to μ_T .

We apply Lemma 12.1 of Bhattacharya and Rao (1986) [4]. It follows from (17) and the identity $\mu_T([-T^{-1}, T^{-1}]) = 3/4$ that

$$(19) \quad \tilde{\Delta} \leq 2 \sup_{x \in \mathbb{R}} |(\tilde{\mathcal{F}} - \mathcal{G}) * \mu_T(-\infty, x]| + c_* T^{-1}.$$

Here $\tilde{\mathcal{F}}$ respectively \mathcal{G} denote the probability distribution of $\tilde{\mathbb{T}}$ respectively the signed measure with density function $G'(x)$. Furthermore, $*$ denotes the convolution operation. Proceeding as in the proof of Lemma 12.2 ibidem we obtain

$$(20) \quad (\tilde{\mathcal{F}} - \mathcal{G}) * \mu_T(-\infty, x] = \frac{1}{2\pi} \int_{-\infty}^{+\infty} e^{-itx} \left(\mathbf{E} e^{it\tilde{\mathbb{T}}} - \hat{G}(t) \right) \frac{\hat{\mu}_T(t)}{-it} dt,$$

where \hat{G} denotes the Fourier transform of $G(x)$. Note that $\hat{\mu}_T(t)$ vanishes outside the interval $|t| \leq 2aT$. Finally, we obtain from (19) and (20) that

$$(21) \quad \tilde{\Delta} \leq \frac{1}{\pi} \sup_{x \in \mathbb{R}} |I(x)| + c_* \frac{2a}{T}, \quad I(x) := \int_{-T}^T e^{-itx} \left(\mathbf{E} e^{it\tilde{\mathbb{T}}} - \hat{G}(t) \right) \frac{\hat{\mu}_{T'}(t)}{-it} dt,$$

where $T' = T/2a$. Here we use the fact that $\hat{\mu}_{T'}(t) = 0$ for $|t| > T$.

Choose $T = N^{1+\nu}$ and denote $K_N(t) = \hat{\mu}_{T'}(t)$. Note that $|K_N(t)| \leq 1$ (since $\mu_{T'}$ is a probability measure). Write

$$\begin{aligned} |I(x)| &\leq c I_1 + c I_2 + |I_3| + |I_4|, \\ I_1 &= \int_{|t| \leq t_1} \left| \mathbf{E} e^{it\tilde{\mathbb{T}}} - \hat{G}(t) \right| \frac{dt}{|t|}, \quad I_2 = \int_{t_1 < |t| < T} \left| \hat{G}(t) \right| \frac{dt}{|t|}, \\ I_3 &= \int_{t_1 < |t| < t_2} e^{-itx} \mathbf{E} e^{it\tilde{\mathbb{T}}} \frac{K_N(t)}{-it} dt, \quad I_4 = \int_{t_2 < |t| < T} e^{-itx} \mathbf{E} e^{it\tilde{\mathbb{T}}} \frac{K_N(t)}{-it} dt. \end{aligned}$$

Here we denote $t_1 = N^{1/2} 10^{-3} / \beta_3$ and $t_2 = N^{1-\nu}$.

In view of (21) the bound $\tilde{\Delta} \leq c_* N^{-1-\nu}$ follows from the bounds

$$(22) \quad |I_k| \leq c_* N^{-1-\nu}, \quad k = 1, 2, 3, \quad \text{and} \quad |I_4| \leq c_* N^{-1-\nu} (1 + \delta_*^{-1} N^{-\nu}).$$

The bound $I_2 \leq c_* N^{-1-\nu}$ is a consequence of the exponential decay of $|\hat{G}(t)|$ as $|t| \rightarrow \infty$. For $k = 3, 4$ the bound (22) is shown in Section 3. The proof of (22), for $k = 1$ is based on careful expansions and is given Section 5.

3. LARGE FREQUENCIES

Here we prove the bounds (22) for I_3 and I_4 . The proof of the bound $|I_3| \leq c_* N^{-1-\nu}$ is relatively simple and it is deferred to the end of the section.

Let us show that

$$(23) \quad \left| \int_{N^{1-\nu} < |t| < N^{1+\nu}} e^{-itx} \mathbf{E} e^{it\tilde{\mathbb{T}}} \frac{K_N(t)}{-it} dt \right| \leq c_* \frac{1 + \delta_*^{-1}}{N^{1+2\nu}}.$$

In what follows we assume that N is sufficiently large, say $N > C_*$, where C_* depends only on $A_*, D_*, M_*, r, s, \nu_1, \nu_2, \delta$. We use this inequality in several places below, where the constant C_* can be easily specified. Note that for small N such that $N \leq C_*$ the inequality (23) becomes trivial.

3.1. Notation. Let us first introduce some notation. Introduce the number

$$\alpha = 3/(r+2).$$

For $r \in (4, 5]$ and ν , defined by (15), we have

$$2/r < \alpha < 1/2 \quad \text{and} \quad 80\nu < \min\{r\alpha - 2, 1 - 2\alpha\}.$$

Given N introduce the integers

$$(24) \quad n \approx N^{50\nu}, \quad M = \lfloor (N - m)/n \rfloor.$$

We have $N - m = Mn + s$, where the integer $0 \leq s < n$. Observe, that the inequalities $\nu < 600^{-1}$ and $m < N^{1/2}$, see (15), imply $M > n$. Therefore, $s < M$. Split the index set

$$(25) \quad \begin{aligned} \{m+1, \dots, N\} &= O_1 \cup O_2 \cup \dots \cup O_n, \\ O_i &= \{j : m + (i-1)M < j \leq m + iM\}, \quad 1 \leq i \leq n-1, \\ O_n &= \{j : m + (n-1)M < j \leq N\} \end{aligned}$$

Clearly, O_1, \dots, O_{n-1} are of equal size ($=M$) and $|O_n| = M + s < 2M$.

We shall assume that the random variable $X : \Omega \rightarrow \mathcal{X}$ is defined on the probability space (Ω, P) and P_X is the probability distribution on \mathcal{X} induced by X . Given $p \geq 1$ let $L^p = L^p(\mathcal{X}, P_X)$ denote the space of real functions $f : \mathcal{X} \rightarrow \mathbb{R}$ with $\mathbf{E}|f(X)|^p < \infty$. Denote $\|f\|_p = (\mathbf{E}|f(X)|^p)^{1/p}$. With a random variable $g(X)$ we associate an element $g = g(\cdot)$ of L^p , $p \leq r$. Let $p_g : L^2 \rightarrow L^2$ denotes the projection onto the subspace orthogonal to the vector $g(\cdot)$ in L^2 . Given $h \in L^2$, decompose

$$(26) \quad h = a_h g + h^*, \quad a_h = \langle h, g \rangle \|g\|_2^{-2}, \quad h^* := p_g(h),$$

where $\langle h, g \rangle = \int h(x)g(x)P_X(dx)$. For $h \in L^r$ we have

$$(27) \quad \|h\|_r \geq \|h\|_2 \geq \|h^*\|_2.$$

Furthermore, for $r^{-1} + v^{-1} = 1$ (here $r \geq 2 \geq v > 1$) we have

$$|\langle h, g \rangle| \leq \|h\|_r \|g\|_v \leq \|h\|_r \|g\|_2.$$

In particular,

$$(28) \quad |a_h| \leq \|h\|_r / \|g\|_2.$$

It follows from the decomposition (26) and (28) that

$$(29) \quad \|h^*\|_r \leq \|h\|_r + |a_h| \|g\|_r \leq \|h\|_r (1 + \|g\|_r / \|g\|_2)$$

$$(30) \quad = c_g \|h\|_r, \quad c_g := 1 + \|g\|_r / \|g\|_2.$$

Note that $c_g \leq c_g^* := 1 + M_*^{1/r} A_*^{-1/2}$. Introduce the numbers

$$(31) \quad a_1 = \frac{1}{4} \min \left\{ \frac{1}{12c_g^*}, \frac{(c_r A_* / 2^r M_*)^{1/(r-2)}}{1 + 4A_*^{-1/2}} \right\}, \quad c_r = \frac{7}{24} \frac{1}{2^{r-1}}.$$

We shall show that there exist $\delta', \delta'' > 0$ depending on A_*, M_*, δ only such that (uniformly in N) Cramér's characteristic ρ , see (5), satisfies

$$(32) \quad \rho(a_1, 2N^{-\nu+1/2}) \geq \delta', \quad \rho((2\beta_3)^{-1}, N^{\nu_2+1/2}) \geq \delta''.$$

We shall prove the first inequality only. In view of (5) it suffices to show that $\rho(a_1, \beta_3^{-1}) \geq \delta'$. Invoking the simple inequality, see, e.g., proof of (187) below,

$$|\mathbf{E}e^{it\sigma^{-1}g(X_1)}| \leq 1 - 2^{-1}t^2(1 - 3^{-1}|t|\beta_3)$$

we obtain, for $|t| \leq \beta_3^{-1}$,

$$|\mathbf{E}e^{it\sigma^{-1}g(X_1)}| \leq 1 - t^2/3.$$

Therefore, $\rho(a_1, \beta_3^{-1}) \geq a_1^2/3$ and we can choose $\delta' = \min\{\delta, a_1^2/3\}$ in (32).

Introduce the constant (depending only on A_*, M_*, δ)

$$(33) \quad \delta_1 = \delta'/(10c_g^*).$$

Note that $0 < \delta_1 < 1/10$.

Given $f \in L^r$ and $T_0 \in \mathbb{R}$ such that

$$(34) \quad N^{-\nu+1/2} \leq |T_0| \leq N^{\nu+1/2},$$

denote

$$(35) \quad \begin{aligned} I(T_0) &= [T_0, T_0 + \delta_1 N^{-\nu+1/2}], \\ \tau(f) &= 1 - v^2(f), \quad v(f) = \sup_{t \in I(T_0)} |u_t(f)|, \end{aligned}$$

$$u_t(f) = \int \exp\{it(g(x) + N^{-1/2}f(x))\} P_X(dx).$$

Given a random variable η with values in L^r and number $0 < s < 1$ define

$$(36) \quad d_s(\eta, I(T_0)) = \mathbb{I}_{\{v^2(\eta) > 1-s^2\}} \mathbb{I}_{\{\|\eta\|_r \leq N^\nu\}}, \quad \delta_s(\eta, I(T_0)) = \mathbf{E}d_s(\eta, I(T_0)).$$

Introduce the function

$$(37) \quad \psi^{**}(x, y) = \psi(x, y) - b(x)g(y) - b(y)g(x)$$

and the number

$$\delta_3^2 = \mathbf{E}|\psi^{**}(X_1, X_2)|^2.$$

It follows from (6) and our assumption that $\sigma_{\mathbb{T}}^2 = 1$, see (14), that $\delta_3^2 \geq \delta_*^2$.

3.2. Proof of (23). The proof of (23) is rather technical and therefore will be illustrated by an outline. In the first step we truncate random variables X_{m+1}, \dots, X_N in a special way using conditioning and replace them by corresponding "truncated" random variables Y_{m+1}, \dots, Y_N . Correspondingly the statistic $\tilde{\mathbb{T}}$ is, then, replaced by T' , see (43). In the second step we split the interval of frequencies $N^{1-\nu} \leq |t| \leq N^{1+\nu}$ into non overlapping intervals $\cup_p J_p$ of sizes $\approx N^{1-\nu}$ so that the integral (45) splits into the sum (46). Conditionally, given Y_{m+1}, \dots, Y_N , the statistic T' is linear in observations X_1, \dots, X_m , since in $\tilde{\mathbb{T}}$ we have removed the higher order terms (in X_1, \dots, X_m) from \mathbb{T} . Let $\mathbf{E}_{\mathbb{Y}}$ denote the conditional expectation given Y_{m+1}, \dots, Y_N . The conditional characteristic function $\mathbf{E}_{\mathbb{Y}} \exp\{itT'\} = \alpha_t^m \exp\{itW'\}$ contains the multiplicative component α_t^m , where

$$\alpha_t = \mathbf{E}_{\mathbb{Y}} \exp\{itN^{-1/2}g(X_1) + itN^{-3/2} \sum_{j=m+1}^N \psi(X_1, Y_j)\}$$

and where the real valued statistic W' is obtained from W replacing X_j by Y_j , for $m+1 \leq j \leq N$. In order to bound $|\mathbf{E}_{\mathbb{Y}} e^{itT'}|$ one would like to show exponential decay (in m) of the product $|\alpha_t^m|$ using a Cramér type condition like (5) above. For $|t| = o(N)$ (the case of medium frequencies) the size of the quadratic part $N^{-3/2} \sum_{j=m+1}^N \psi(X_1, Y_j)$ can be neglected and Cramér's condition implies $|\alpha_t| \leq 1 - v'$ for some $v' > 0$. Thus we obtain $|\alpha_t^m| \leq e^{-mv'}$. For large frequencies $|t| \approx N$, the contribution of the quadratic part becomes significant and we introduce an extra moment

condition (6). Using (6), we show that, for a large set of values $t \in J_p$, Cramér's condition (5) yields the desired decay of $|\alpha_t^m|$. Furthermore, the measure of the remaining t is small with high probability.

Step 1. Truncation. Recall that the random variable $X : \Omega \rightarrow \mathcal{X}$ is defined on the probability space (Ω, P) . Let X' be an independent copy so that (X, X') is defined on $(\Omega \times \Omega', P \times P)$, where $\Omega' = \Omega$. It follows from $\mathbf{E}|\psi(X, X')|^r < \infty$, by Fubini, that for P almost all $\omega' \in \Omega'$ the function $\psi(\cdot, X'(\omega')) = \{x \rightarrow \psi(x, X'(\omega')), x \in \mathcal{X}\}$ is an element of L^r . Furthermore, one can define an L^r -valued random variable $Z' : \Omega' \rightarrow L^r$ such that $Z'(\omega') = \psi(\cdot, X'(\omega'))$, for P almost all ω' . Consider the event $\tilde{\Omega} = \{\|Z'\|_r \leq N^\alpha\} \subset \Omega'$ and denote $q_N = P(\tilde{\Omega})$. Here $\|Z'\|_r = (\int |\psi(x, X'(\omega'))|^r P_X(dx))^{1/r}$ denotes the L^r norm of the random vector Z' . Let $Y : \tilde{\Omega} \rightarrow \mathcal{X}$ denote the random variable X' conditioned on the event $\tilde{\Omega}$. Therefore Y is defined on the probability space $(\tilde{\Omega}, \tilde{P})$, where \tilde{P} denotes the restriction of $q_N^{-1}P$ to the set $\tilde{\Omega}$ and, for every $\omega' \in \tilde{\Omega}$, we have $Y(\omega') = X'(\omega')$. Let Z denote the L^r -valued random element $\{x \rightarrow \psi(x, Y(\omega'))\}$ defined on the probability space $(\tilde{\Omega}, \tilde{P})$.

We can assume that $\mathbb{X} := (X_1, \dots, X_N)$ is a sequence of independent copies of X defined on the probability space (Ω^N, P^N) . Let $\bar{\omega} = (\omega_1, \dots, \omega_N)$ denote an element of Ω^N . Every X_j defines random vector $Z'_j = \psi(\cdot, X_j)$ taking values in L^r . Introduce events $A_j := \{\|Z'_j\|_r \leq N^\alpha\} \subset \Omega^N$ and let $\mathbb{X}' = (X_1, \dots, X_m, Y_{m+1}, \dots, Y_N)$ denote the sequence \mathbb{X} conditioned on the event $\Omega^* = \cap_{j=m+1}^N A_j = \Omega^m \times \tilde{\Omega}^{N-m}$. Clearly, $\mathbb{X}'(\bar{\omega}) = \mathbb{X}(\bar{\omega})$ for every $\bar{\omega} \in \Omega^*$ and \mathbb{X}' is defined on the space $\Omega^m \times \tilde{\Omega}^{N-m}$ equipped with the probability measure $P^m \times \tilde{P}^{N-m}$. In particular, the random variables $X_1, \dots, X_m, Y_{m+1}, \dots, Y_N$ are independent and Y_j , for $m+1 \leq j \leq N$, has the same distribution as Y . Let Z_j denote the L^r -valued random element $\{x \rightarrow \psi(x, Y_j)\}$, for $m+1 \leq j \leq N$.

We are going to replace $\mathbf{E}e^{it\tilde{\mathbb{T}}}$ by $\mathbf{E}e^{itT'}$. For $s > 0$ we have almost surely

$$(38) \quad 1 - \mathbb{I}_{A_j} \leq N^{-\alpha s} \|Z'_j\|_r^s, \quad \|Z'_j\|_r^r = \mathbf{E}(|\psi(X, X_j)|^r | X_j).$$

Therefore, by Chebyshev's inequality, for $s = r$,

$$(39) \quad 0 \leq 1 - q_N \leq N^{-r\alpha} \mathbf{E}|\psi(X, X_j)|^r \leq N^{-r\alpha} M_* \leq c_* N^{-2-3\nu}.$$

We have for $k \leq N$

$$(40) \quad \begin{aligned} q_N^{-k} &\leq (1 - N^{-r\alpha} M_*)^{-k} \leq (1 - N^{-2} M_*)^{-N} \leq c_*, \\ q_N^{-k} - 1 &\leq k q_N^{-k} (1 - q_N) \leq c_* k N^{-2-3\nu} \leq c_* N^{-1-3\nu}. \end{aligned}$$

For a measurable function $f : \mathcal{X}^N \rightarrow \mathbb{R}$, we have

$$(41) \quad \mathbf{E}f(X_1, \dots, X_m, Y_{m+1}, \dots, Y_N) = \mathbf{E}f(X_1, \dots, X_N) \frac{\mathbb{I}_{A_{m+1}} \cdots \mathbb{I}_{A_N}}{q_N^{(N-m)}}.$$

Therefore, for $f \geq 0$, (40) and (41) imply

$$(42) \quad \mathbf{E}f(X_1, \dots, X_m, Y_{m+1}, \dots, Y_N) \leq c_* \mathbf{E}f(X_1, \dots, X_N)$$

Furthermore, for

$$(43) \quad T' := \tilde{\mathbb{T}}(X_1, \dots, X_m, Y_{m+1}, \dots, Y_N)$$

we have, by (40) and (41),

$$(44) \quad \begin{aligned} |\mathbf{E}e^{it(T'-x)} - \mathbf{E}e^{it(\tilde{\mathbb{T}}-x)}| &\leq (q_N^{-(N-m)} - 1) + (1 - \mathbf{P}\{A_{m+1} \cap \cdots \cap A_N\}) \\ &= (q_N^{-(N-m)} - 1) + (1 - q_N^{N-m}) \leq c_* N^{-1-3\nu}. \end{aligned}$$

We replace $\tilde{\mathbb{T}}$ by T' in the exponent in (23). The error of this replacement is $c_*N^{-1-2\nu}$, by (44) and the simple inequality $|K_N(t)| \leq 1$, for every t . In order to prove (23) we shall show that

$$(45) \quad I := \int_{N^{1-\nu} \leq |t| \leq N^{1+\nu}} \mathbf{E} e^{it\hat{T}} v_N(t) dt \leq c_* \frac{1 + \delta_3^{-1}}{N^{1+2\nu}},$$

$$v_N(t) = t^{-1} K_N(t), \quad \hat{T} = T' - x.$$

Step 2. Here we prove (45). Split the integral

$$(46) \quad I = \sum_p I_p, \quad I_p = \mathbf{E} \int_{t \in J_p} e^{it\hat{T}} v_N(t) dt,$$

where $\{J_p, p = 1, 2, \dots\}$ denote a sequence of consecutive intervals of length $\approx \delta_1 N^{1-\nu}$ each. Here δ_1 is a constant defined by (33). In order to prove (45) we show that for every p ,

$$(47) \quad |I_p| \leq c_* N^{-2} + c_* N^{-1-4\nu} (1 + \delta_3^{-1}).$$

Given p let us prove (47). Firstly, we replace I_p by $\mathbf{E} J_*$, where

$$J_* = \int \mathbb{I}_{\{t \in I_*\}} v_N(t) \mathbf{E}_{\mathbb{Y}} e^{it\hat{T}} dt.$$

Here $I_* = I_*(Y_{m+1}, \dots, Y_N) \subset J_p$ is a random subset defined by

$$(48) \quad I_* = \{t \in J_p : |\alpha_t|^2 > 1 - \varepsilon_m^2\}, \quad \varepsilon_m^2 = m^{-1} \ln^2 N.$$

Since, for $t \notin I_*$, we have

$$|\mathbf{E}_{\mathbb{Y}} e^{itT'}| \leq |\alpha_t|^m \leq (1 - \varepsilon_m^2)^{m/2} \leq c_* N^{-3},$$

the error of this replacement is given by

$$(49) \quad |I_p - \mathbf{E} J_*| \leq c_* N^{-2}.$$

Secondly, we shall show that with a high probability the set $I_* \subset J_p$ is a (random) interval. This fact and the fact that $v_N(t)$ is monotone will be used latter to bound the integral J_* . Introduce the L^r -valued random element

$$(50) \quad S = N^{-1/2} (Z_{m+1} + \dots + Z_N) = N^{-1/2} \sum_{j=m+1}^N \psi(\cdot, Y_j).$$

We apply Lemma 12 to the set $N^{-1/2} I_*$ conditionally on the event $SS = \{\|S\|_r < N^{\nu/10}\}$. This lemma shows that $N^{-1/2} I_*$ is an interval of size at most $c_* \varepsilon_m$. That is, we can write $I_* = (a_N, a_N + b_N^{-1})$ and

$$(51) \quad \mathbb{I}_{SS} J_* = \mathbb{I}_{SS} \mathbf{E}_{\mathbb{Y}} \tilde{J}_*, \quad \tilde{J}_* = \int_{a_N}^{a_N + b_N^{-1}} v_N(t) e^{it\hat{T}} dt,$$

where the random variables (functions of Y_{m+1}, \dots, Y_N) satisfy

$$a_N \in J_p \quad \text{and} \quad b_N^{-1} \leq c_* \varepsilon_m \sqrt{N} = c_* \sqrt{N} m^{-1/2} \ln N.$$

By Lemma 13, SS has at least a probability $\mathbf{P}\{SS\} \geq 1 - c_* N^{-3}$. Therefore,

$$(52) \quad |\mathbf{E} J_* - \mathbf{E} \mathbb{I}_{SS} J_*| \leq c_* N^{-2}.$$

Clearly, $I_* \neq \emptyset$ if and only if $\alpha^2 > 1 - \varepsilon_m^2$, where

$$\alpha = \sup\{|\alpha_t| : t \in J_p\}.$$

Therefore, we can write, see also (51),

$$\mathbb{I}_{SS} J_* = \mathbb{I}_{\mathbb{B}} J_* = \mathbb{I}_{\mathbb{B}} \mathbf{E}_{\mathbb{Y}} \tilde{J}_*, \quad \text{where} \quad \mathbb{B} = \{\alpha^2 > 1 - \varepsilon_m^2\} \cap SS.$$

This identity together with (49) and (52) shows

$$(53) \quad |I_p| \leq |\mathbf{E}\mathbb{I}_{\mathbb{B}}\mathbf{E}_{\mathbb{Y}}\tilde{J}_*| + c_*N^{-2}.$$

Using the integration by parts formula we shall show that

$$(54) \quad |\mathbf{E}\mathbb{I}_{\mathbb{B}}\mathbf{E}_{\mathbb{Y}}\tilde{J}_*| \leq \frac{c}{N^{1-\nu}} \left(\mathbf{P}\{\mathbb{B}\} + \int_{b_N}^1 \frac{\mathbf{P}\{\mathbb{B}_\varepsilon\}}{\varepsilon^2} d\varepsilon \right),$$

$$\mathbb{B}_\varepsilon = \mathbb{B} \cap \{|\hat{T}| \leq \varepsilon\}.$$

This inequality in combination with (53) and (55), see below, shows (47),

$$(55) \quad \int_{b_N}^1 \frac{\mathbf{P}\{\mathbb{B}_\varepsilon\}}{\varepsilon^2} d\varepsilon \leq c_* \frac{1 + \delta_3^{-1}}{N^{5\nu}}, \quad \mathbf{P}\{\mathbb{B}\} \leq c_* \frac{1 + \delta_3^{-1}}{N^{5\nu}}.$$

Proof of (55) is rather technical. It is given in subsection **3.3**.

Let us prove (54). Firstly, we show that

$$(56) \quad |\tilde{J}_*| \leq c(|\hat{T}| + b_N)^{-1} a_N^{-1}.$$

The integration by parts formula shows

$$(57) \quad i\hat{T}\tilde{J}_* = v_N(t)e^{it\hat{T}} \Big|_{a_N}^{a_N+b_N^{-1}} - \int_{a_N}^{a_N+b_N^{-1}} v'_N(t)e^{it\hat{T}} dt =: a' + a''.$$

By our choice of the smoothing kernel, $v_N(t)$ is monotone on J_p . Therefore,

$$|a''| \leq \int_{a_N}^{a_N+b_N^{-1}} |v'_N(t)| dt = \left| \int_{a_N}^{a_N+b_N^{-1}} v'_N(t) dt \right| = |v_N(a_N) - v_N(a_N + b_N^{-1})|.$$

Invoking the simple inequality $|a'| \leq |v_N(a_N)| + |v_N(a_N + b_N^{-1})|$ and using $|v_N(t)| \leq |t|^{-1}$ we obtain from (57)

$$|\hat{T}\tilde{J}_*| \leq c(a_N^{-1} + (a_N + b_N^{-1})^{-1}) \leq c a_N^{-1}.$$

For $|\hat{T}| > b_N$, this inequality implies (56). For $|\hat{T}| \leq b_N$ the inequality (56) follows from the inequalities

$$|\tilde{J}_*| \leq \int_{a_N}^{a_N+b_N^{-1}} |v_N(t)| dt \leq \int_{a_N}^{a_N+b_N^{-1}} \frac{c}{|t|} dt \leq c a_N^{-1} b_N^{-1}.$$

Furthermore, it follows from (56) and the inequality $a_N \geq N^{1-\nu}$ that

$$|\tilde{J}_*| \leq c(|\hat{T}| + b_N)^{-1} N^{-1+\nu}.$$

Finally, we apply the inequality (which holds for arbitrary real number v)

$$\frac{1}{|v| + b_N} \leq 2 + 2 \int_{b_N}^1 \frac{d\varepsilon}{\varepsilon^2} \mathbb{I}_{\{|v| \leq \varepsilon\}}$$

to derive

$$|\tilde{J}_*| \leq \frac{c_*}{N^{1-\nu}} \left(1 + \int_{b_N}^1 \frac{d\varepsilon}{\varepsilon^2} \mathbb{I}_{\{|\hat{T}| \leq \varepsilon\}} \right).$$

This inequality gives (54).

3.3. Here we prove (55). The first (respectively second) inequality is proved in *Step A* (respectively *Step B*).

Step A. Here we prove the first inequality of (55). Split

$$\begin{aligned} W &= W_1 + W_2 + W_3, & W_1 &= \frac{1}{N^{1/2}} \sum_{j=m+1}^N g(X_j), \\ W_2 &= \frac{1}{N^{3/2}} \sum_{m < i < j \leq N} \psi(X_i, X_j), & W_3 &= \sum_{|A| \geq 3: A \cap \Omega_m = \emptyset} T_A. \end{aligned}$$

Replacing X_j by Y_j , for $m+1 \leq j \leq N$, we obtain $W' = W'_1 + W'_2 + W'_3$. Therefore, we can write $\hat{T} = L + \Delta + W'_3$, where

$$(58) \quad \begin{aligned} L &= \frac{1}{\sqrt{N}} \sum_{j=1}^m g(X_j) + \frac{1}{\sqrt{N}} \sum_{j=m+1}^N g(Y_j) - x, \\ \Delta &= \frac{1}{N^{3/2}} \sum_{j=1}^m \sum_{l=m+1}^N \psi(X_j, Y_l) + \frac{1}{N^{3/2}} \sum_{m+1 \leq j < l \leq N} \psi(Y_j, Y_l). \end{aligned}$$

The inequalities $|\hat{T}| \leq \varepsilon$ and $|L| \geq 2\varepsilon$ imply $|\Delta + W'_3| > \varepsilon$. Therefore,

$$\mathbf{P}\{\mathbb{B}_\varepsilon\} \leq \mathbf{P}\{\mathbb{B} \cap \{|L| \leq 2\varepsilon\}\} + \mathbf{P}\{|\hat{T}| \leq \varepsilon, |\Delta + W'_3| \geq \varepsilon\} =: I_1(\varepsilon) + I_2(\varepsilon).$$

In order to prove (55) we show that

$$(59) \quad \int_{b_N}^1 \frac{d\varepsilon}{\varepsilon^2} I_1(\varepsilon) \leq c_* N^{-5\nu} (1 + \delta_3^{-1}), \quad \int_{b_N}^1 \frac{d\varepsilon}{\varepsilon^2} I_2(\varepsilon) \leq c_* N^{-5\nu}.$$

Step A.1. Here we prove (59), for the integral over $I_2(\varepsilon)$. We have

$$(60) \quad \begin{aligned} I_2(\varepsilon) &\leq \mathbf{P}\{|W'_3| > \varepsilon/2\} + I_3(\varepsilon), \\ I_3(\varepsilon) &:= \mathbf{P}\{|L + \Delta| < 3\varepsilon/2, |\Delta| > \varepsilon/2\}. \end{aligned}$$

It follows from (42), by Chebyshev's inequality, $\mathbf{P}\{|W'_3| > \varepsilon/2\} \leq c_* \varepsilon^{-2} \mathbf{E}W_3^2$. Furthermore, invoking the inequalities, see (162), (163),

$$\mathbf{E}W_3^2 = \sum_{|A| \geq 3: A \cap \Omega_m = \emptyset} \mathbf{E}T_A^2 \leq \sum_{|A| \geq 3} \mathbf{E}T_A^2 \leq N^{-2} \Delta_3^2 \leq c_* N^{-2}$$

we obtain from (60) $I_2(\varepsilon) \leq I_3(\varepsilon) + c_* \varepsilon^{-2} N^{-2}$. Since

$$\int_{b_N}^1 \frac{d\varepsilon}{\varepsilon^2} \left(\frac{1}{\varepsilon^2 N^2} \right) \leq c_* b_N^{-3} N^{-2} \leq c_* N^{-5\nu},$$

it suffices to show (59) for $I_3(\varepsilon)$.

We have, for $\Lambda_1 = N^{-3/2} \sum_{1 \leq i < j \leq m} \psi(X_i, X_j)$,

$$I_3(\varepsilon) \leq \mathbf{P}\{|\Lambda_1| > \varepsilon/4\} + I_4(\varepsilon), \quad I_4(\varepsilon) := \mathbf{P}\{|L + U| < 2\varepsilon, |U| > \varepsilon/4\},$$

where we denote $U = \Lambda_1 + \Delta$. By Chebyshev's inequality

$$\mathbf{P}\{|\Lambda_1| > \varepsilon/4\} \leq 16\varepsilon^{-2} \mathbf{E}\Lambda_1^2 \leq c_* \varepsilon^{-2} m^2 N^{-3}.$$

Furthermore,

$$\int_{b_N}^1 \frac{d\varepsilon}{\varepsilon^2} \left(\frac{m^2}{\varepsilon^2 N^3} \right) \leq c_* b_N^{-3} m^2 N^{-3} \leq c_* N^{-5\nu}.$$

Therefore, it suffices to show (59) for $I_4(\varepsilon)$.

Let $I'_4(\varepsilon)$ be the same probability as $I_4(\varepsilon)$ but with X_i replaced by Y_i , for $1 \leq i \leq m$,

$$I'_4(\varepsilon) = \mathbf{P}\{|L' + U'| < 2\varepsilon, |U'| > \varepsilon/4\},$$

$$L' = \frac{1}{N^{1/2}} \sum_{1 \leq i \leq N} g(Y_i) - x, \quad U' = \frac{1}{N^{3/2}} \sum_{1 \leq i < j \leq N} \psi(Y_i, Y_j).$$

We have $|I_4(\varepsilon) - I'_4(\varepsilon)| \leq c_* N^{-1-3\nu}$, cf. (44). Since

$$\int_{b_N}^1 \frac{d\varepsilon}{\varepsilon^2} N^{-1-3\nu} \leq c_* b_N^{-1} N^{-1-3\nu} \leq c_* N^{-5\nu},$$

it suffices to show (59) for $I'_4(\varepsilon)$.

In what follows we show the bound (59) for $I'_4(\varepsilon)$. Split the sample

$$\mathbb{Y} := \{Y_1, \dots, Y_N\} = \mathbb{Y}_1 \cup \mathbb{Y}_2 \cup \mathbb{Y}_3,$$

into three groups of nearly equal size. Split $U' = \sum_{i \leq j} U'_{ij}$ so that the sum U'_{ij} depends on observations from \mathbb{Y}_i and \mathbb{Y}_j only. We have

$$(61) \quad I'_4(\varepsilon) \leq \sum_{i \leq j} \mathbf{P}\{|L' + U'| \leq 2\varepsilon, |U'_{ij}| \geq \varepsilon/24\}.$$

In order to prove (59) we shall show this bound for every summand in the right of (61). Let \tilde{U} denote a summand U'_{ij} , say, not depending on \mathbb{Y}_3 . We shall prove (59) for

$$(62) \quad \tilde{I}(\varepsilon) := \mathbf{P}\{|L' + U'| \leq 2\varepsilon, |\tilde{U}| \geq \varepsilon/24\} = \mathbf{E}\mathbb{I}_{\mathcal{U}}\mathbb{I}_{\mathcal{V}},$$

$$\mathcal{U} = \{|\tilde{U}| \geq \varepsilon/24\}, \quad \mathcal{V} = \{|L' + U'| \leq 2\varepsilon\}.$$

By the definition of Y_1, \dots, Y_N , the random function

$$x \rightarrow \bar{S}(x) = N^{-1/2} \sum_{Y_i \in \mathbb{Y} \setminus \mathbb{Y}_3} \psi(x, Y_i), \quad x \in \mathcal{X},$$

defines a random variable with values in L^r such that, for every i , $\|\psi(\cdot, Y_i)\|_r \leq N^\alpha$ for almost all values of Y_i . An application of Lemma 13 gives

$$\mathbf{P}\{\|\bar{S}(\cdot)\|_r > N^\nu\} \leq N^{-3}.$$

Therefore, in (62) we can replace the event \mathcal{V} by $\mathcal{V}_1 = \mathcal{V} \cap \{\|\bar{S}\|_r \leq N^\nu\}$.

Since \tilde{U} does not depend on \mathbb{Y}_3 , the concentration bound for the conditional probability (proof of this bound is given below)

$$(63) \quad p' := \mathbf{E}(\mathbb{I}_{\mathcal{V}_1} | \mathbb{Y}_1, \mathbb{Y}_2) \leq c_*(\varepsilon + N^{-1/2})$$

implies

$$(64) \quad \tilde{I}(\varepsilon) \leq c_*(\varepsilon + N^{-1/2}) \mathbf{P}\{\mathcal{U}\} \leq c_*(\varepsilon + N^{-1/2}) \varepsilon^{-r} N^{-r/2}.$$

In the last step we applied Chebyshev's inequality

$$\mathbf{P}\{\mathcal{U}\} \leq (24/\varepsilon)^r N^{-r/2} \mathbf{E}|N^{1/2} \tilde{U}|^r$$

and the bound $\mathbf{E}|N^{1/2} \tilde{U}|^r \leq c_* \mathbf{E}|N^{1/2} U_{ij}|^r \leq c_*$, which follows from (42) and routine moment inequalities for U -statistics, see Dharmadhikari, S. W., Fabian, V. Jogdeo, K. (1968). Here the random variable U_{ij} is obtained from \tilde{U} after we replace Y_j by X_j for every j .

It follows from (64) and the simple inequality $\varepsilon \geq b_N \geq c_* N^{-1/2}$ that

$$\int_{b_N}^1 \frac{d\varepsilon}{\varepsilon^2} \tilde{I}(\varepsilon) \leq \frac{c_*}{N^{r/2}} \int_{b_N}^1 \frac{d\varepsilon}{\varepsilon^{1+r}} \leq \frac{c_*}{N^{r/2} b_N^r}$$

$$= c_* m^{-r/2} \ln^r N \leq c_* N^{-5\nu},$$

provided that $m^{r/2} \geq N^{6\nu}$. The latter inequality is ensured by (15). Thus we have shown (59) for $\tilde{I}(\varepsilon)$.

It remains to prove (63). Write $L' + U' = L_* + U_* + b - x$, where

$$L_* = \frac{1}{N^{1/2}} \sum_{Y_j \in \mathbb{Y}_3} (g(Y_j) + N^{-1/2} \bar{S}(Y_j)), \quad U_* = \frac{1}{N^{3/2}} \sum_{\{Y_j, Y_k\} \subset \mathbb{Y}_3} \psi(Y_j, Y_k),$$

and where b is a function of $\{Y_i \in \mathbb{Y} \setminus \mathbb{Y}_3\}$. Introduce the random variables \bar{L} and \bar{U} which are obtained from L_* and U_* after we replace every $Y_j \in \mathbb{Y}_3$ by the corresponding observation X_j . We have

$$\begin{aligned} p' &\leq \sup_{v \in R} \mathbf{E}(\mathbb{I}_{\{L_* + U_* \in [v, v+2\varepsilon]\}} | \mathbb{Y}_1, \mathbb{Y}_2) \mathbb{I}_{\{\|\bar{S}\|_r \leq N^\nu\}} \\ &\leq c_* \sup_{v \in R} \mathbf{E}(\mathbb{I}_{\{\bar{L} + \bar{U} \in [v, v+2\varepsilon]\}} | \mathbb{Y}_1, \mathbb{Y}_2) \mathbb{I}_{\{\|\bar{S}\|_r \leq N^\nu\}}. \end{aligned}$$

In the last step we applied (42). An application of the Berry-Esseen bound due to van Zwet (1984) shows (63).

Step A.2. Here we prove (59) for $I_1(\varepsilon)$. Write $I_1(\varepsilon)$ in the form

$$\begin{aligned} I_1(\varepsilon) &= \mathbf{E} \mathbb{I}_{\mathbb{A}} \mathbb{I}_{SS} \mathbb{I}_{\mathbb{W}} \leq \mathbf{E} \mathbb{I}_{\mathbb{A}} \mathbb{I}_{\mathbb{V}} \mathbb{I}_{\mathbb{W}}, \\ \mathbb{A} &= \{\alpha^2 > 1 - \varepsilon_m^2\}, \quad \mathbb{V} = \{\|S\|_r \leq N^\nu\}, \quad \mathbb{W} = \{|L| < 2\varepsilon\}, \end{aligned}$$

where ε_m is defined in (48). Note that, by the Berry-Esseen inequality,

$$(65) \quad \mathbf{P}\{\mathbb{W}\} \leq c_*(\varepsilon + N^{-1/2}).$$

Furthermore, one can show that the probability of the event \mathbb{A} is small, like $\mathbf{P}\{\mathbb{A}\} = O(N^{-6\nu})$. We are going to make use of both of these bounds while constructing an upper bound for $I_1(\varepsilon)$. Since the events \mathbb{A} and \mathbb{W} refer to the same set of random variables Y_{m+1}, \dots, Y_N , we cannot argue directly that $\mathbf{E} \mathbb{I}_{\mathbb{A}} \mathbb{I}_{\mathbb{W}} \approx \mathbf{P}\{\mathbb{A}\} \mathbf{P}\{\mathbb{W}\}$. Nevertheless, invoking a complex conditioning argument we are able to show that

$$(66) \quad I_1(\varepsilon) \leq c_* \mathcal{R}(\varepsilon + N^{-1/2}) + c_* N^{-2}, \quad \mathcal{R} := N^{-6\nu}(1 + \delta_3^{-1}).$$

Since $\varepsilon \geq b_N > N^{-1/2}$, the inequality (66) implies (59).

Let us prove (66). Since the proof is rather involved we start by providing an outline, Let the integers n and M be defined by (24). Split $\{1, \dots, N\} = O_0 \cup O_1 \cup \dots \cup O_n$, where $O_0 = \{1, \dots, m\}$ and where the sets O_i , for $1 \leq i \leq n$, are defined in (25). Split L , see (58),

$$(67) \quad L = \sum_{k=0}^n L_k - x, \quad L_k = N^{-1/2} \sum_{j \in O_k} g(Y_j), \quad \text{for } k = 1, \dots, n,$$

and $L_0 = N^{-1/2} \sum_{j \in O_0} g(X_j)$. Observe, that $\mathbb{I}_{\mathbb{W}}$ is a function of L_0, L_1, \dots, L_n . The random variables $\mathbb{I}_{\mathbb{A}}$ and $\mathbb{I}_{\mathbb{V}}$ are functions of Y_{m+1}, \dots, Y_N and do not depend on X_1, \dots, X_m . Therefore, denoting

$$m(l_1, \dots, l_n) = \mathbf{E}(\mathbb{I}_{\mathbb{A}} \mathbb{I}_{\mathbb{V}} | L_1 = l_1, \dots, L_n = l_n)$$

we obtain from (65)

$$(68) \quad \mathbf{E} \mathbb{I}_{\mathbb{A}} \mathbb{I}_{\mathbb{V}} \mathbb{I}_{\mathbb{W}} = \mathbf{E} \mathbb{I}_{\mathbb{W}} m(L_1, \dots, L_n) \leq c_*(\varepsilon + N^{-1/2}) \mathcal{M},$$

where $\mathcal{M} = \text{ess sup } m(l_1, \dots, l_n)$. Clearly, the bound $\mathcal{M} \leq c_* \mathcal{R}$ would imply (66). Unfortunately, we are not able to establish such a bound directly. In what follows we prove (66) using the argument outlined above. But we shall use a more delicate conditioning which allow to estimate quantities like \mathcal{M} .

Step A.2.1. Firstly we replace L_k , $1 \leq k \leq n$, by smooth random variables

$$(69) \quad g_k = \frac{1}{N} \frac{\xi_k}{n^{1/2}} + L_k,$$

where ξ_1, \dots, ξ_n are symmetric i.i.d. random variables with the density function defined by (18) with $k = 6$ and $a = 1/6$ so that the characteristic function $t \rightarrow \mathbf{E} \exp\{it\xi_1\}$ vanishes outside the unit interval $\{t : |t| < 1\}$. Note that $\mathbf{E}\xi_1^4 < \infty$.

We assume that the sequences ξ_1, ξ_2, \dots and $X_1, \dots, X_m, Y_{m+1}, \dots, Y_N$ are independent. In particular, ξ_k and L_k are independent. Introduce the event

$$\tilde{\mathbb{W}} = \{|L_0 + \sum_{k=1}^n g_k - x| < 3\varepsilon\}.$$

Note that

$$\mathbb{I}_{\mathbb{W}} \leq \mathbb{I}_{\tilde{\mathbb{W}}} + \mathbb{I}_{\{|\xi| \geq \varepsilon N\}}, \quad \text{where} \quad \xi = \frac{1}{n^{1/2}} \sum_{k=1}^n \xi_k.$$

By Chebyshev's inequality and the inequality $\mathbf{E}\xi^4 \leq c$,

$$\mathbf{P}\{|\xi| \geq \varepsilon N\} \leq \frac{\mathbf{E}\xi^4}{\varepsilon^4 N^4} \leq \frac{c}{\varepsilon^4 N^4} \leq \frac{c_*}{N^2}.$$

Here we used the inequality $\varepsilon^2 N \geq b_N^2 N \geq c'_*$. Therefore, we obtain

$$(70) \quad \mathbf{E}\mathbb{I}_{\mathbb{A}}\mathbb{I}_{\mathbb{V}}\mathbb{I}_{\mathbb{W}} \leq \mathbf{E}\mathbb{I}_{\mathbb{A}}\mathbb{I}_{\mathbb{V}}\mathbb{I}_{\tilde{\mathbb{W}}} + c_* N^{-2}.$$

In subsequent steps of the proof we replace the conditioning on L_1, \dots, L_n (in (68)) by the conditioning on the random variables g_1, \dots, g_n . Since the latter random variables have densities (as it is shown in Lemma 7 below) the corresponding conditional distributions are much easier to handle. Moreover, we restrict the conditioning on the event where these densities are positive.

Step A.2.2. Given $w > 0$, consider the events $\{|g_k| \leq n^{-1/2}w\}$ and their indicator functions $\mathbb{I}_k = \mathbb{I}_{\{|g_k| \leq n^{-1/2}w\}}$. Using the simple inequality $n\mathbf{E}g_k^2 \leq c_*$ (where c_* depends on M_* and r) we obtain from Chebyshev's inequality that

$$(71) \quad \mathbf{P}\{\mathbb{I}_k = 1\} = 1 - \mathbf{P}\{|g_k| > n^{-1/2}w\} \geq 1 - w^{-2}n\mathbf{E}|g_k|^2 > 7/8,$$

where the last inequality holds for a sufficiently large constant w (depending on M_* , r). Fix a number w such that (71) holds and introduce the event $\mathbb{B}^* = \{\sum_{k=1}^n \mathbb{I}_k > n/4\}$. Hoeffding's inequality shows $\mathbf{P}\{\mathbb{B}^*\} \geq 1 - \exp\{-n/8\}$. Therefore,

$$(72) \quad \mathbf{E}\mathbb{I}_{\mathbb{A}}\mathbb{I}_{\mathbb{V}}\mathbb{I}_{\tilde{\mathbb{W}}}\mathbb{I}_{\mathbb{B}^*} \leq \mathbf{E}\mathbb{I}_{\mathbb{A}}\mathbb{I}_{\mathbb{V}}\mathbb{I}_{\tilde{\mathbb{W}}}\mathbb{I}_{\mathbb{B}^*} + c_* N^{-2}.$$

Given a binary vector $\theta = (\theta_1, \dots, \theta_n)$ (with $\theta_k \in \{0, 1\}$) write $|\theta| = \sum_k \theta_k$. Introduce the event $\mathbb{B}_\theta = \{\mathbb{I}_k = \theta_k, 1 \leq k \leq n\}$ and the conditional expectation

$$m_\theta(z_1, \dots, z_n) = \mathbf{E}(\mathbb{I}_{\mathbb{A}}\mathbb{I}_{\mathbb{V}}\mathbb{I}_{\mathbb{B}_\theta} | g_1 = z_1, \dots, g_n = z_n).$$

Note that $\mathbb{I}_{\mathbb{B}_\theta}$, the indicator of the event \mathbb{B}_θ , is a function of g_1, \dots, g_n . It follows from the identities

$$\mathbb{B}^* = \cup_{|\theta| > n/4} \mathbb{B}_\theta \quad \text{and} \quad \mathbb{I}_{\mathbb{B}^*} = \sum_{|\theta| > n/4} \mathbb{I}_{\mathbb{B}_\theta}$$

(here $\mathbb{B}_\theta \cap \mathbb{B}_{\theta'} = \emptyset$, for $\theta \neq \theta'$) that

$$\begin{aligned} \mathbf{E}\mathbb{I}_{\mathbb{A}}\mathbb{I}_{\mathbb{V}}\mathbb{I}_{\tilde{\mathbb{W}}}\mathbb{I}_{\mathbb{B}^*} &= \sum_{|\theta| > n/4} \mathbf{E}\mathbb{I}_{\mathbb{A}}\mathbb{I}_{\mathbb{V}}\mathbb{I}_{\tilde{\mathbb{W}}}\mathbb{I}_{\mathbb{B}_\theta} \\ &= \sum_{|\theta| > n/4} \mathbf{E}\mathbb{I}_{\mathbb{B}_\theta}\mathbb{I}_{\tilde{\mathbb{W}}}m_\theta(g_1, \dots, g_n). \end{aligned}$$

Assume that we have already shown that uniformly in θ , satisfying $|\theta| > n/4$, we have

$$(73) \quad M_\theta \leq c_* \mathcal{R}, \quad \text{where} \quad M_\theta := \text{ess sup } m_\theta(z_1, \dots, z_n).$$

This bound in combination with (65), which extends to $\tilde{\mathbb{W}}$ as well, implies

$$\begin{aligned} \mathbf{E} \mathbb{I}_{\mathbb{A}} \mathbb{I}_{\mathbb{V}} \mathbb{I}_{\tilde{\mathbb{W}}} \mathbb{I}_{\mathbb{B}^*} &\leq c_* \mathcal{R} \sum_{|\theta| > n/4} \mathbf{E} \mathbb{I}_{\mathbb{B}_\theta} \mathbb{I}_{\tilde{\mathbb{W}}} = c_* \mathcal{R} \mathbf{E} \mathbb{I}_{\mathbb{B}^*} \mathbb{I}_{\tilde{\mathbb{W}}} \\ &\leq c_* \mathcal{R} \mathbf{P}\{\tilde{\mathbb{W}}\} \leq c_* \mathcal{R} (\varepsilon + N^{-1/2}). \end{aligned}$$

Combining this inequality, (70) and (72) we obtain (66).

Step A.2.3. Here we show (73). Fix $\theta = (\theta_1, \dots, \theta_n)$ satisfying $|\theta| > n/4$. Denote, for brevity, $h = |\theta|$ and assume without loss of generality that $\theta_i = 1$, for $1 \leq i \leq h$, and $\theta_j = 0$, for $h+1 \leq j \leq n$.

Consider the h -dimensional random vector $\bar{g}_{[\theta]} = (g_1, \dots, g_h)$. The random vector $\bar{g}_{[\theta]}$ and the sequences of random variables

$$\mathbb{Y}_\theta = \{Y_j : m + hM < j \leq N\}, \quad \xi_\theta = \{\xi_j : h < j \leq n\}$$

are independent. Furthermore the summands S_θ and S'_θ of the decomposition

$$S = S_\theta + S'_\theta, \quad S_\theta(\cdot) = \frac{1}{\sqrt{N}} \sum_{1 \leq k \leq h} \sum_{j \in O_k} \psi(\cdot, Y_j),$$

are independent, see (50). Moreover, we have $m_\theta(z_1, \dots, z_n) \leq \tilde{m}_\theta(\bar{z}_{[\theta]})$, where

$$\tilde{m}_\theta(\bar{z}_{[\theta]}) = \text{ess sup}_\theta \mathbf{E}(\mathbb{I}_{\mathbb{A}} \mathbb{I}_{\mathbb{V}} \mathbb{I}_{\mathbb{B}_\theta} \mid \bar{g}_{[\theta]} = \bar{z}_{[\theta]}, \mathbb{Y}_\theta, \xi_\theta)$$

denotes the "ess sup" taken with respect to almost all values of \mathbb{Y}_θ and ξ_θ . Here $\bar{z}_{[\theta]} = (z_1, \dots, z_h) \in \mathbb{R}^h$. In order to prove (73) we show that

$$(74) \quad \tilde{m}_\theta(\bar{z}_{[\theta]}) \leq c_* \mathcal{R}.$$

Let us prove (74). Given \mathbb{Y}_θ , denote $f_\theta = S'_\theta$ (note that S'_θ is a function of \mathbb{Y}_θ). Using the notation (36), we have for the interval $J'_p = N^{-1/2} J_p$,

$$(75) \quad \mathbf{E}(\mathbb{I}_{\mathbb{A}} \mathbb{I}_{\mathbb{V}} \mathbb{I}_{\mathbb{B}_\theta} \mid \bar{g}_{[\theta]} = \bar{z}_{[\theta]}, \mathbb{Y}_\theta, \xi_\theta) = \mathbb{I}_{\mathbb{B}_\theta} \mathbf{E}(d_{\varepsilon_m}(f_\theta + S_\theta, J'_p) \mid \bar{g}_{[\theta]} = \bar{z}_{[\theta]}, \mathbb{Y}_\theta, \xi_\theta).$$

Note that the factor $\mathbb{I}_{\mathbb{B}_\theta}$ in the right hand side is non zero only in the case where $\bar{z}_{[\theta]} = (z_1, \dots, z_h)$ satisfies $|z_i| \leq w/\sqrt{n}$, for $i = 1, \dots, h$.

Introduce the L^r valued random variables

$$U_i = N^{-1/2} \sum_{j \in O_i} \psi(\cdot, Y_j), \quad i = 1, \dots, h,$$

and the regular conditional probability

$$P(\bar{z}_{[\theta]}; \mathcal{A}) = \mathbf{E}(\mathbb{I}_{\{(U_1, \dots, U_h) \in \mathcal{A}\}} \mid \bar{g}_{[\theta]} = \bar{z}_{[\theta]}).$$

Here \mathcal{A} denotes a Borel subset of $L^r \times \dots \times L^r$ (h -times). By independence, there exist regular conditional probabilities

$$(76) \quad P_i(z_i; \mathcal{A}_i) = \mathbf{E}(\mathbb{I}_{U_i \in \mathcal{A}_i} \mid g_i = z_i), \quad i = 1, \dots, h,$$

such that for Borel subsets \mathcal{A}_i of L^r we have

$$P(\bar{z}_{[\theta]}; \mathcal{A}_1 \times \dots \times \mathcal{A}_h) = \prod_{1 \leq i \leq h} P_i(z_i; \mathcal{A}_i).$$

In particular, for every $\bar{z}_{[\theta]}$, the regular conditional probability $P(\bar{z}_{[\theta]}; \cdot)$ is the (measure theoretical) extension of the product of the regular conditional probabilities (76). Therefore, denoting by ψ_i a random variable with values in L^r and with the distribution

$$(77) \quad \mathbf{P}\{\psi_i \in \mathcal{B}\} = P_i(z_i; \mathcal{B}), \quad \mathcal{B} \subset L^r - \text{Borel set},$$

we obtain that the distribution of the sum

$$(78) \quad \zeta = \psi_1 + \cdots + \psi_h$$

of independent random variables ψ_1, \dots, ψ_h is the regular conditional distribution of S_θ , given $\bar{g}_{[\theta]} = \bar{z}_{[\theta]}$. In particular, the expectation in the right hand side of (75) equals $\delta_{\varepsilon_m}(f_\theta + \zeta)$, where

$$(79) \quad \delta_s(f_\theta + \zeta) := \mathbf{E}_\zeta d_s(f_\theta + \zeta, J'_p), \quad s > 0,$$

and where \mathbf{E}_ζ denotes the conditional expectation given all the random variables, but ζ . Note that the inequality

$$(80) \quad \varepsilon_m \leq \varepsilon_*$$

implies

$$(81) \quad \delta_{\varepsilon_m}(f_\theta + \zeta) \leq \delta_{\varepsilon_*}(f_\theta + \zeta).$$

We shall apply Lemma 1 to construct an upper bound for $\delta_{\varepsilon_*}(f_\theta + \zeta)$, where $\varepsilon_* = \mu_* |T_0| N^{-1/2} / 20$. The quantity μ_* is defined in (92) and satisfies $c_* \delta_3^2 / n \leq \mu_*^2 \leq c'_* \delta_3^2 / n$, by the inequality (212). Note that for T_0 satisfying (34), for some integers m and n as in (15) and (24), and for the quantity δ_3 (see (37)) which satisfies

$$(82) \quad \delta_3^2 \geq N^{-8\nu},$$

the inequality (80) holds with ε_m defined by (48), provided that N is sufficiently large ($N > C_*$). Moreover, we have

$$(83) \quad \varepsilon_*^2 \leq c_* \delta_3^2 N^{-48\nu}.$$

In order to apply Lemma 1 we invoke the moment inequalities of Lemma 10. Now Lemma 1 shows that

$$(84) \quad \delta_{\varepsilon_*}(f_\theta + \zeta) \leq c_* \kappa_*^{1/2} \varepsilon_*^{(r-2)/(2r)} + c_* N^{-2},$$

where the number κ_* , defined in (92), satisfies $\kappa_* \leq c_* \delta_3^{-r/(r-2)}$, by (213). Denote $\tilde{r} = r^{-1} + (r-2)^{-1}$. It follows from (84), (83) and (81), for $r > 4$, that

$$(85) \quad \begin{aligned} \delta_{\varepsilon_m}(f_\theta + \zeta) &\leq c_* \delta_3^{-\tilde{r}} N^{-6\nu} + c_* N^{-2} \\ &\leq c_*(1 + \delta_3^{-\tilde{r}}) N^{-6\nu} \leq c_* \mathcal{R}. \end{aligned}$$

In the last step we used the simple bound $\delta_3^2 \leq c_*$, see (195), and the inequality $1 + \delta_3^{-\tilde{r}} \leq 2 + \delta_3^{-1}$, which follows from $\tilde{r} < 1$. Note that (85) and (75), (79) imply (74) thus completing the proof of the first inequality (55).

Step B. Here we prove the second bound of (55). It is convenient to write the L^r -valued random variable (50) in the form

$$(86) \quad \begin{aligned} S &= U_1 + \cdots + U_{n-1} + U_n =: S' + U_n, \\ U_i &= N^{-1/2} \sum_{j \in O_i} \psi(\cdot, Y_j). \end{aligned}$$

Observe that U_1, \dots, U_{n-1} are independent and identically distributed L^r -valued random variables.

We are going to apply Lemma 1 conditionally, given U_n , to the probability

$$\mathbf{P}\{\mathbb{B}\} = \mathbf{E}\tilde{p}(U_n), \quad \tilde{p}(f) = \mathbf{E}(d_{\varepsilon_m}(S' + f, N^{-1/2} J_p) | U_n = f).$$

Lemma 9 shows that U_1, \dots, U_{n-1} satisfy the moment conditions of Lemma 1, but now the corresponding quantity μ_* , see (92), satisfies $c_*\delta_3^2/n \leq \mu_*^2 \leq c'_*/n$, by (196). This implies the bound $\varepsilon_* \leq c_*N^{-48\nu}$ instead of (83). As a result we obtain a different power of δ_3 in the upper bound below. Proceeding as in proof of (85), see (81), (83), (84), we obtain

$$\tilde{p}(f) \leq c_*(1 + \delta_3^{-r/2(r-2)})N^{-6\nu} \leq c_*\mathcal{R}.$$

In the last step we used the inequality $1 + \delta_3^{-r/2(r-2)} \leq 2 + \delta_3^{-1}$. This inequality follows from $r/2(r-2) < 1$, for $r > 4$. Therefore, we have $\mathbf{P}\{\mathbb{B}\} \leq \mathbf{E}\tilde{p}(U_n) \leq c_*\mathcal{R}$, where \mathcal{R} is defined in (66). This completes the proof of the second inequality in (55).

3.4. Here we prove the bound $|I_3| \leq c_*N^{-1-\nu}$, see (22). It follows from (44) that

$$(87) \quad |I_3| \leq \int_{t \in \mathbb{J}} \frac{\mathbf{E}|\alpha_t^m|}{|t|} dt + c_*N^{-1-\nu},$$

where $\mathbb{J} = \{t : \sqrt{N}/10^3\beta_3 \leq |t| \leq N^{1-\nu}\}$. For the L_r -valued random element S defined in (50) and the event $\mathbb{S} = \{\|S\|_r < N^{\nu/10}\}$ write

$$(88) \quad \mathbf{E}|\alpha_t^m| \leq \mathbf{E}\mathbb{I}_{\mathbb{S}}|\alpha_t^m| + \mathbf{E}(1 - \mathbb{I}_{\mathbb{S}}).$$

Estimate the second summand as $\mathbf{P}\{\|S\|_r \geq N^{\nu/10}\} \leq c_*N^{-3}$, by Lemma 13. Furthermore, expanding the exponent in α_t we obtain

$$\mathbb{I}_{\mathbb{S}}|\alpha_t| \leq |\mathbf{E} \exp\{itN^{-1/2}g(X_1)\}| + \mathbb{I}_{\mathbb{S}}|t|N^{-1}\|S\|_1.$$

It follows from (5) that the first summand is bounded from above by $1 - v$, for some $v > 0$ depending on A_*, M_*, D_*, δ only, see the proof of (32). Furthermore, the second summand is bounded from above by $N^{-9\nu/10}$ almost surely. Therefore, for sufficiently large $N > C_*$ we have $\mathbb{I}_{\mathbb{S}}|\alpha_t| \leq 1 - v/2$ uniformly in N . Invoking this bound in (88) we obtain $\mathbf{E}|\alpha_t^m| \leq (1 - v/2)^m + c_*N^{-3} \leq c_*N^{-3}$, for m satisfying (15). Finally, we obtain that the integral in (87) is bounded from above by c_*N^{-2} thus completing the proof.

4. COMBINATORIAL CONCENTRATION BOUND

We start the section by introducing some notation and collecting auxiliary inequalities. Then we formulate and prove Lemmas 1 and 2.

Introduce the number

$$(89) \quad \delta_2 = \min\left\{\frac{1}{12c_g}, \frac{(c_r\|g\|_2^2/2^r\|g\|_r^r)^{1/(r-2)}}{1 + 4/\|g\|_2}\right\},$$

where $c_g = 1 + \|g\|_r/\|g\|_2$ and $c_r = (7/24)2^{-(r-1)}$. Denote

$$\rho^* = 1 - \sup\{|\mathbf{E}e^{itg(X_1)}| : 2^{-1}\delta_2 \leq |t| \leq N^{-\nu+1/2}\}.$$

It follows from the identity $\rho^* = \rho(2^{-1}\sigma\delta_2, \sigma N^{-\nu+1/2})$ and the simple inequality $a_1 \leq \delta_2/4$, see (31), that $\rho^* \geq \rho(2\sigma a_1, \sigma N^{-\nu+1/2})$. Furthermore, it follows from (164) and the assumption $\sigma_{\mathbb{T}}^2 = 1$ that $1/2 < \sigma < 2$ for sufficiently large N ($N > C_*$). Therefore, $\rho^* \geq \rho(a_1, 2N^{-\nu+1/2}) \geq \delta'$, where the last inequality follows from (32). We obtain, for $N > C_*$,

$$(90) \quad 1 - \sup\{|\mathbf{E}e^{itg(X_1)}| : 2^{-1}\delta_2 \leq |t| \leq N^{-\nu+1/2}\} \geq \delta',$$

where the number δ' depends on $A_*, D_*, M_*, \nu_1, r, s, \delta$ only. In what follows we use the notation $c_0 = 10$. Let $L_0^r = \{y \in L^r : \int_{\mathcal{X}} y(x)P_X(dx) = 0\}$ denotes a subspace of L^r . Observe, that $\mathbf{E}g(X_1) = 0$ implies $y^*(=p_g(y)) \in L_0^r$, for every $y \in L_0^r$.

4.1. Let ψ_1, \dots, ψ_n denote independent random vectors with values in L_0^r . For $k = 1, \dots, n$, write

$$\zeta_k = \psi_1 + \dots + \psi_k \quad \text{and} \quad \zeta = \zeta_n.$$

Let $\bar{\psi}_i$ denote an independent copy of ψ_i . Write $\psi_i^* = p_g(\psi_i)$ and $\bar{\psi}_i^* = p_g(\bar{\psi}_i)$, see (26). Introduce random vectors

$$\tilde{\psi}_i = 2^{-1}(\psi_i - \bar{\psi}_i), \quad \tilde{\psi}_i^* = 2^{-1}(\psi_i^* - \bar{\psi}_i^*), \quad \hat{\psi}_i = 2^{-1}(\psi_i + \bar{\psi}_i).$$

We shall assume that, for some $c_A \geq c_D \geq c_B > 0$,

$$(91) \quad n^{r/2} \mathbf{E} \|\tilde{\psi}_i\|_r^r \leq c_A^r, \quad c_B^2 \leq n \mathbf{E} \|\tilde{\psi}_i^*\|_2^2 \leq c_D^2,$$

for every $1 \leq i \leq n$. Furthermore, denote $\mu_i^2 = \mathbf{E} \|\tilde{\psi}_i^*\|_2^2$ and $\tilde{\kappa}_i^{r-2} = \frac{8}{3} \frac{\mathbf{E} \|\tilde{\psi}_i\|_r^r}{\mu_i^r}$,

$$(92) \quad \mu_* = \min_{1 \leq i \leq n} \mu_i, \quad \kappa_* = \max_{1 \leq i \leq n} \tilde{\kappa}_i.$$

Observe that, by Hölder's inequality and (27), we have $\kappa_i > 1$, for $i = 1, \dots, n$.

Lemma 1. *Let $4 < r \leq 5$ and $0 < \nu < 10^{-2}(r-4)$. Assume that $n \geq N^{5\nu}$. Suppose that*

$$(93) \quad \kappa_*^4 \leq \frac{9}{256} \frac{n}{\ln N}.$$

Assume that (90), (91) as well as (101), (107) (below) hold. There exist a constant $c_ > 0$ which depends on $r, s, \nu, A_*, D_*, M_*, \delta$ only such that for every T_0 satisfying (34) we have*

$$(94) \quad \delta_{\varepsilon_*}(f + \zeta, I(T_0)) \leq c_*(C_D/C_B)^{1/2} \kappa_*^{1/2} \varepsilon_*^{(r-2)/2r} + c_* N^{-2},$$

for an arbitrary non-random element $f \in L_0^r$. Here $\varepsilon_ = \frac{\mu_*}{2c_0} \frac{|T_0|}{\sqrt{N}}$. The function $\delta_s(\cdot, I(T_0))$, is defined in (36).*

In *Step A.2.3* of Section 3 we apply this lemma to random vector $\zeta = \psi_1 + \dots + \psi_n$, see (78).

In *Step B* of Section 3 we apply this lemma to the random vector S' , see (86).

Proof of Lemma 1. We shall consider the case where $T_0 > 0$. For $T_0 < 0$ the proof is the same. We can assume without loss of generality that $c_0 < N^\nu$. Denote $X = \|\tilde{\psi}_i^*\|_2$ and $Y = \|\tilde{\psi}_i\|_r$ and $\mu = \mu_i$, $\kappa = \tilde{\kappa}_i$. By (27), we have $Y \geq X$.

Step 1. Here we construct the bound (95), see below, for the probability $\mathbf{P}\{B_i\}$, where

$$B_i = \{X \geq \mu/2, Y < \kappa\mu\}.$$

Write

$$\begin{aligned} \mu^2 &= \mathbf{E}X^2 = \mathbf{E}X^2 I_A + \mathbf{E}X^2 I_{B_i} + \mathbf{E}X^2 I_D, \\ A &= \{X < \mu/2\}, \quad D = \{X \geq \mu/2, Y \geq \kappa\mu\}. \end{aligned}$$

Substitution of the bounds

$$\begin{aligned} \mathbf{E}X^2 I_A &\leq \frac{\mu^2}{4}, \\ \mathbf{E}X^2 I_{B_i} &\leq \mathbf{E}Y^2 I_{B_i} \leq (\kappa\mu)^2 \mathbf{P}\{B_i\}, \\ \mathbf{E}X^2 I_D &\leq \mathbf{E}Y^2 I_{\{Y \geq \kappa\mu\}} \leq (\kappa\mu)^{2-r} \mathbf{E}Y^r \end{aligned}$$

gives

$$\mu^2 \leq 4^{-1} \mu^2 + \kappa^2 \mu^2 \mathbf{P}\{B_i\} + (\kappa\mu)^{2-r} \mathbf{E}Y^r.$$

Finally, invoking the identity $\kappa^{r-2} = (8/3) \mathbf{E}Y^r / \mu^r$ we obtain

$$(95) \quad \mathbf{P}\{B_i\} \geq \frac{3}{4\kappa^2} - \frac{\mathbf{E}Y^r}{(\kappa\mu)^r} = \frac{3}{4\kappa^2} \left(1 - \frac{4\mathbf{E}Y^r}{3\mu^r \kappa^{r-2}}\right) = \frac{3}{8\kappa^2} \geq \frac{3}{8\kappa_*^2} =: p$$

Introduce the (random) set $J = \{i : B_i \text{ occurs}\} \subset \{1, \dots, n\}$. Hoeffding's inequality applied to the random variable $|J| = \mathbb{I}_{B_1} + \dots + \mathbb{I}_{B_n}$ shows

$$(96) \quad \mathbf{P}\{|J| \leq \rho n\} \leq \exp\{-np^2/2\} \leq N^{-2}, \quad \rho := p/2 = (3/16)\kappa_*^{-2}.$$

In the last step we invoke (93) and use (95).

Step 2. Here we introduce randomization. Note that for any $\alpha_i \in \{-1, +1\}$, $i = 1, \dots, n$, the distributions of the random vectors

$$(\psi_1, \dots, \psi_n) \quad \text{and} \quad (\alpha_1 \tilde{\psi}_1 + \hat{\psi}_1, \dots, \alpha_n \tilde{\psi}_n + \hat{\psi}_n)$$

coincide. Therefore, denoting

$$\tilde{\zeta}_n = \alpha_1 \tilde{\psi}_1 + \dots + \alpha_n \tilde{\psi}_n, \quad \hat{\zeta}_n = \hat{\psi}_1 + \dots + \hat{\psi}_n,$$

we have for $s > 0$,

$$\delta_s(f + \zeta, I(T_0)) = \delta_s(f + \tilde{\zeta}_n + \hat{\zeta}_n, I(T_0)),$$

for every choice of $\alpha_1, \dots, \alpha_n$. From now on let $\alpha_1, \dots, \alpha_n$ denote a sequence of independent identically distributed Bernoulli random variables independent of $\tilde{\psi}_i, \hat{\psi}_i$, $1 \leq i \leq n$, and with probabilities $\mathbf{P}\{\alpha_1 = 1\} = \mathbf{P}\{\alpha_1 = -1\} = 1/2$. Denoting by \mathbf{E}_α the expectation with respect to the sequence $\alpha_1, \dots, \alpha_n$ we obtain

$$(97) \quad \delta_s(f + \zeta, I(T_0)) = \mathbf{E}_\alpha \delta_s(f + \tilde{\zeta}_n + \hat{\zeta}_n, I(T_0)).$$

We are going to condition on $\tilde{\psi}_i$ and $\hat{\psi}_i$, $1 \leq i \leq n$, while taking expectations with respect to $\alpha_1, \dots, \alpha_n$. It follows from (96), (97) and the fact that the random variable $|J|$ does not depend on $\alpha_1, \dots, \alpha_n$ that

$$(98) \quad \delta_s(f + \zeta, I(T_0)) \leq \mathbf{E} \mathbb{I}_{\{|J| \geq \rho n\}} \gamma_s(\tilde{\psi}_i, \hat{\psi}_i, 1 \leq i \leq n) + N^{-2},$$

where

$$\gamma_s(\tilde{\psi}_i, \hat{\psi}_i, 1 \leq i \leq n) = \mathbf{E}_\alpha \mathbb{I}_{\{|J| \geq \rho n\}} \mathbb{I}_{\{v^2(f + \tilde{\zeta}_n + \hat{\zeta}_n) > 1 - s^2\}} \mathbb{I}_{\{\|f + \tilde{\zeta}_n + \hat{\zeta}_n\|_r \leq N^\nu\}}$$

denotes the conditional expectation given $\tilde{\psi}_i, \hat{\psi}_i$, $1 \leq i \leq n$. Note that (94) is a consequence of (98) and of the bound

$$(99) \quad \gamma_{\varepsilon_*}(\tilde{\psi}_i, \hat{\psi}_i, 1 \leq i \leq n) \leq c_* \kappa_*^{1/2} \varepsilon_*^{(r-2)/(2r)}.$$

Let us prove this bound. Introduce the integers

$$n_0 = l - 1 \quad l = \lfloor \delta_2 \varkappa^{-1} \varepsilon_*^{-(r-2)/r} \rfloor, \quad \varkappa = 2c_0(C_D/C_B)\kappa_*.$$

Let us show that

$$(100) \quad n_0 \leq \rho n.$$

It follows from the inequalities

$$\varepsilon_*^{-1} \leq 2 \frac{c_0}{c_B} N^\nu n^{1/2}, \quad N^{\nu(r-2)/r} \leq N^\nu \leq n^{1/r}, \quad \delta_2 \leq \frac{3}{16} \left(\frac{3}{8}\right)^{1/(r-2)}$$

that

$$l \leq \frac{\delta_2}{C_D} \frac{1}{k_*} \left(\frac{C_B}{2c_0}\right)^{2/r} (N^\nu n^{1/2})^{(r-2)/r} \leq \frac{3}{16} \frac{1}{k_*} \frac{C_B^{2/r}}{C_D} n^{1/2}.$$

Note that (93) implies $k_* \leq n^{1/4}$. Therefore, the inequality

$$(101) \quad C_B^{2/r} C_D^{-1} \leq n^{1/4}$$

implies $l \leq (3/16)k_*^{-2}n = \rho n$. We obtain (100).

Given $\tilde{\psi}_i, \hat{\psi}_i$, $1 \leq i \leq n$, consider the corresponding set J , say $J = \{i_1, \dots, i_k\}$. Assume that $k \geq \rho n$. From the inequality $\rho n \geq n_0$, see (100), it follows that we can choose a subset $J' \subset J$ of size $|J'| = n_0$. Split

$$\tilde{\zeta}_n = \sum_{i \in J'} \alpha_i \tilde{\psi}_i + \sum_{i \in J \setminus J'} \alpha_i \tilde{\psi}_i =: \zeta_* + \zeta'$$

and denote $f + \zeta' + \hat{\zeta}_n = f_*$. Note that $f_* \in L_0^r$ almost surely. The bound (99) would follow if we show that

$$(102) \quad \begin{aligned} \tilde{\delta} &\leq c_* \kappa_*^{1/2} \varepsilon_*^{(r-2)/(2r)}, \\ \tilde{\delta} &:= \mathbf{E}' \mathbb{I}_{\{v^2(f_* + \zeta_*) > 1 - \varepsilon_*^2\}} \mathbb{I}_{\{\|f_* + \zeta_*\|_r \leq N^\nu\}}. \end{aligned}$$

Here \mathbf{E}' denotes the conditional expectation given all the random variables, but $\{\alpha_i, i \in J'\}$.

Step 3. Here we prove (102). Note that for $j \in J'$ the vectors

$$x_j = T_0 N^{-1/2} \tilde{\psi}_j \quad \text{and} \quad x_j^* = p_g(x_j) = T_0 N^{-1/2} \tilde{\psi}_j^*$$

satisfy

$$(103) \quad \|x_j^*\|_2 \geq c_0 \varepsilon_*, \quad \|x_j\|_r \leq \varkappa \varepsilon_*, \quad \varkappa = 2c_0(C_D/C_B)\kappa_*.$$

Given $A \subset J'$ denote

$$x_A = \sum_{i \in A} x_i - \sum_{i \in J' \setminus A} x_i, \quad x_A^* = p_g(x_A).$$

We are going to apply Kleitman's theorem on symmetric partitions (see, e.g. the proof of Theorem 4.2, Bollobas (1986)) to the sequence $\{x_j^*, j \in J'\}$ in L^2 . Since for $j \in J'$ we have $\|x_j^*\|_2 \geq c_0 \varepsilon_*$, it follows from Kleitman's theorem that the collection $\mathcal{P}(J')$ of all subsets of J' splits into non-intersecting non-empty classes $\mathcal{P}(J') = \mathcal{D}_1 \cup \dots \cup \mathcal{D}_s$, such that the corresponding sets of linear combinations $V_t = \{x_A^*, A \in \mathcal{D}_t\}$, $t = 1, 2, \dots, s$, are sparse, i.e., given t , for $A, A' \in \mathcal{D}_t$ and $A \neq A'$ we have

$$(104) \quad \|x_A^* - x_{A'}^*\|_2 \geq c_0 \varepsilon_*.$$

Furthermore, the number of classes s is bounded from above by $\binom{n_0}{\lfloor n_0/2 \rfloor}$.

Next, using Lemma 2 we shall show that given f_* the class \mathcal{D}_t may contain at most one element $A \in \mathcal{D}_t$ such that

$$(105) \quad v^2(f_* + \tilde{x}_A) > 1 - \varepsilon_*^2, \quad \|f_* + \tilde{x}_A\|_r \leq N^\nu, \quad \tilde{x}_A := N^{1/2} T_0^{-1} x_A.$$

This means that there are at most $\binom{n_0}{\lfloor n_0/2 \rfloor}$ different subsets $A \subset J'$ for which (105) holds. This implies (102)

$$\tilde{\delta} \leq 2^{-n_0} \binom{n_0}{\lfloor n_0/2 \rfloor} \leq c n_0^{-1/2} = c \delta_2^{-1/2} \varkappa^{1/2} \varepsilon_*^{\frac{r-2}{2r}}.$$

Finally, (94) follows from (98), (99), (102).

Given $f_* \in L_0^r$ let us show that there is no pair A, A' in \mathcal{D}_t which satisfy (105). Fix $A, A' \in \mathcal{D}_t$. We have, by (103) and the choice of n_0 ,

$$\|x_A - x_{A'}\|_r \leq 2 \sum_{i \in J'} \|x_i\|_r \leq 2n_0 \varkappa \varepsilon_* < 2\delta_2 \varepsilon_*^{2/r}.$$

Denoting $S_A = f_* + \tilde{x}_A$ and $S_{A'} = f_* + \tilde{x}_{A'}$ we obtain

$$(106) \quad \|S_A - S_{A'}\|_r = N^{1/2} T_0^{-1} \|x_A - x_{A'}\|_r \leq 2\delta_2 \varepsilon_*^{2/r} N^{1/2} T_0^{-1}.$$

Assume that S_A and $S_{A'}$ satisfy the second inequality of (105), i.e., $\|S_A\|_r \leq N^\nu$ and $\|S_{A'}\|_r \leq N^\nu$. We are going to apply Lemma 2 to the vectors S_A and $S_{A'}$. In order to check the conditions

of Lemma 2 note that (109) and (110) are verified by (103), (104) and (106). Furthermore, the inequalities $c_0 < N^\nu$ and

$$(107) \quad c_B \geq 2N^{4\nu}(n/N)^{1/2},$$

imply $N^{2\nu-1/2} \leq \varepsilon_*$. Finally, we can assume without loss of generality that $\varepsilon_* \leq c'_* := \min\{(\delta'/4)^{r/2}, (A_*^{1/2}/6)^{r/2}\}$. Otherwise (94) follows from trivial inequalities $\delta_{\varepsilon_*} \leq 1 \leq (\varepsilon_*/c'_*)^{(r-2)/2r} \leq c_*\varepsilon_*^{(r-2)/2r}$ and the inequality $\kappa_* > 1$.

Now Lemma 2 shows that $\min\{v^2(S_A), v^2(S_{A'})\} \leq 1 - \varepsilon_*^2$ thus completing the proof of Lemma 1.

4.2. Here we formulate and prove Lemma 2. Let us introduce first some notation. Given $y \in L^r (= L^r(\mathcal{X}, P_X))$ define the symmetrization $y_s \in L^r(\mathcal{X} \times \mathcal{X}, P_X \times P_X)$ by $y_s(x, x') = y(x) - y(x')$, for $x, x' \in \mathcal{X}$. In what follows X_1, X_2 denote independent random variables with values in \mathcal{X} and with the common distribution P_X . By \mathbf{E} we denote the expectation taken with respect to P_X . For $h \in L^r$ we write

$$\mathbf{E}h = \mathbf{E}h(X_1) = \int_{\mathcal{X}} h(x)P_X(dx), \quad \mathbf{E}e^{ih} = \mathbf{E}e^{ih(X_1)} = \int_{\mathcal{X}} e^{ih(x)}P_X(dx).$$

Furthermore, for $2 \leq p \leq r$, denote

$$\|y_s\|_p^p = \mathbf{E}|y(X_1) - y(X_2)|^p, \quad \|y\|_p^p = \mathbf{E}|y(X_1)|^p.$$

Note that for $y \in L_0^r$ we have $y^*(= p_g(y)) \in L_0^r$ and, therefore,

$$(108) \quad \mathbf{E}|y^*(X_1) - y^*(X_2)|^2 = 2\mathbf{E}|y^*(X_1)|^2.$$

Let y_1, \dots, y_k, f be non-random vectors in L^r . We shall assume that these vectors belong to the linear subspace L_0^r . Given non random vectors $\alpha = \{\alpha_i\}_{i=1}^k$ and $\alpha' = \{\alpha'_i\}_{i=1}^k$, with $\alpha_i, \alpha'_i \in \{-1, +1\}$, denote

$$S_\alpha = f + \sum_{i=1}^k \alpha_i y_i, \quad S_{\alpha'} = f + \sum_{i=1}^k \alpha'_i y_i.$$

Lemma 2. *Let $\varkappa > 0$. Assume that (90) holds and suppose that*

$$N^{\nu-1/2} \leq \varepsilon \leq \min\{(\delta'/4)^{r/2}, (\|g\|_2/6)^{r/2}\}.$$

Given T_0 , satisfying (34), write $T^ = N^{1/2}T_0^{-1}$ and assume that*

$$(109) \quad \|y_j^*\|_2 > c_0 T^* \varepsilon, \quad \|y_j\|_r \leq \varkappa T^* \varepsilon, \quad j = 1, \dots, k.$$

Suppose that $\|S_\alpha\|_r \leq N^\nu$ and $\|S_{\alpha'}\|_r \leq N^\nu$ and

$$(110) \quad \|S_\alpha^* - S_{\alpha'}^*\|_2 \geq c_0 T^* \varepsilon, \quad \|S_\alpha - S_{\alpha'}\|_r \leq 2\delta_2 T^* \varepsilon^{2/r}.$$

Then $\min\{v^2(S_\alpha), v^2(S_{\alpha'})\} \leq 1 - \varepsilon^2$.

Recall that the functionals $\tau(\cdot)$, $|u_t(\cdot)|$ and the interval $I = I(T_0)$ used in proof below are defined in (35).

Proof. Note that $\delta_1 < 1/10$ and $\delta_2 < 1/12$. In particular, we have

$$(111) \quad 9/10 \leq 1 - \delta_1 \leq |s/T_0| \leq 1 + \delta_1 \leq 11/10, \quad \text{for } |s - T_0| < \delta_1 N^{-\nu+1/2}.$$

Step 1. Assume that the inequality $\min\{v^2(S_\alpha), v^2(S_{\alpha'})\} \leq 1 - \varepsilon^2$ fails. Then for some $s, t \in I$ we have

$$(112) \quad 1 - |u_t(S_\alpha)|^2 < \varepsilon^2, \quad 1 - |u_s(S_{\alpha'})|^2 < \varepsilon^2,$$

see also (35). Fix these s, t and denote

$$\tilde{X} = s(g + N^{-1/2}S_{\alpha'}) - t(g + N^{-1/2}S_{\alpha}).$$

We are going to apply the inequality (251),

$$1 - |\mathbf{E}e^{i(Y+Z)}|^2 \geq 2^{-1}(1 - |\mathbf{E}e^{iZ}|^2) - (1 - |\mathbf{E}e^{iY}|^2)$$

to $Z = -\tilde{X}$ and $Y = s(g + N^{-1/2}S_{\alpha'})$. It follows from this inequality and (112) that

$$\varepsilon^2 > 1 - |u_t(S_{\alpha})|^2 = 1 - |\mathbf{E}e^{i(Y+Z)}|^2 \geq 2^{-1}(1 - |\mathbf{E}e^{-i\tilde{X}}|^2) - \varepsilon^2.$$

In view of the identity $|\mathbf{E}e^{-i\tilde{X}}| = |\mathbf{E}e^{i\tilde{X}}|$ we have

$$(113) \quad 1 - |\mathbf{E}e^{i\tilde{X}}|^2 < 4\varepsilon^2.$$

Step 2. Here we shall show that (113) contradicts the second inequality of (110). Firstly, we collect some auxiliary inequalities. Write the decomposition (26) for S_{α} and $S_{\alpha'}$,

$$(114) \quad S_{\alpha} = ag + S_{\alpha}^*, \quad S_{\alpha'} = a'g + S_{\alpha'}^*.$$

Decompose

$$\begin{aligned} \tilde{X} &= vg + h, \\ v &= (s-t)(1 + aN^{-1/2}) + (a' - a)sN^{-1/2}, \\ h &= (s-t)N^{-1/2}S_{\alpha}^* + sN^{-1/2}(S_{\alpha'}^* - S_{\alpha}^*), \end{aligned}$$

where $v \in \mathbb{R}$ and where $h \in L^r$ is L^2 -orthogonal to g . An application of (30) to S_{α}^* and $S_{\alpha'}^* - S_{\alpha}^*$ gives

$$(115) \quad \|h\|_r \leq c_g N^{-1/2} (|s| \|S_{\alpha'} - S_{\alpha}\|_r + |s-t| \|S_{\alpha}\|_r).$$

Furthermore, it follows from the simple inequality

$$\|x+y\|_2^2 \geq 2^{-1}\|x\|_2^2 - \|y\|_2^2$$

that

$$(116) \quad \|h\|_2^2 \geq 2^{-1}s^2N^{-1}\|S_{\alpha'}^* - S_{\alpha}^*\|_2^2 - (s-t)^2N^{-1}\|S_{\alpha}^*\|_2^2.$$

Note that for a and a' defined in (114) we obtain from (28) and (110) that

$$(117) \quad |a| \leq \|S_{\alpha}\|_r \|g\|_2^{-1} \leq N^{\nu} \|g\|_2^{-1},$$

$$(118) \quad |a' - a| \leq \|S_{\alpha'} - S_{\alpha}\|_r \|g\|_2^{-1} \leq 2\delta_2 \varepsilon^{2/r} N^{1/2} T_0^{-1} \|g\|_2^{-1}.$$

Step 4.2.1. Consider the case where, $|s-t| < \delta_2$. Invoking the inequalities $\|S_{\alpha}\|_r \leq N^{\nu}$ and (110) we obtain from (115) that

$$\|h\|_r^r \leq (4c_g)^r \delta_2^r \left(N^{\nu r - r/2} + \varepsilon^2 |s|^r T_0^{-r} \right).$$

Furthermore, using (111), (89), and $N^{\nu-1/2} \leq \varepsilon$, we obtain for $4 \leq r \leq 5$

$$(119) \quad \|h\|_r^r \leq 3^{-r} (\varepsilon^r + \varepsilon^2 (11/10)^r) \leq 3^{1-r} \varepsilon^2.$$

Note that (27) implies $\|S_{\alpha}^*\|_2 \leq \|S_{\alpha}\|_r \leq N^{\nu}$. This inequality in combination with (110) and (116) gives

$$\|h\|_2^2 \geq 2^{-1}(s/T_0)^2 c_0^2 \varepsilon^2 - \delta_2^2 N^{2\nu-1}.$$

Invoking (111) and using $c_0 > 10, \delta_2 < 12^{-1}$, and $N^{\nu-1/2} \leq \varepsilon$ we obtain

$$(120) \quad \|h\|_2^2 \geq (4/10)c_0^2 \varepsilon^2.$$

Now we are going to apply Lemma 12 statement **a)** to $\tilde{X} = vg + h$. For this purpose we verify conditions of the lemma. Firstly, note that (120), (108) imply, $\|h_s\|_2^2 \geq (8/10)c_0^2\varepsilon^2$. Furthermore, it follows from the simple inequality $\mathbf{E}|h(X_1) - h(X_2)|^r \leq 2^r \mathbf{E}|h(X_1)|^r$ and (119) that $\|h_s\|_r^r \leq 3(2/3)^r \varepsilon^2$. Therefore, we obtain, for $4 \leq r \leq 5$,

$$\|h_s\|_r^r \leq \frac{6}{10}\varepsilon^2 \leq c_0^{-2}\|h_s\|_2^2 \leq c_r\|h_s\|_2^2, \quad c_r = (7/24)2^{-(r-1)}.$$

Furthermore, the inequalities (117), (118) and (111) imply

$$|v| \leq \delta_2 + \delta_2\|g\|_2^{-1}(N^{\nu-1/2} + 2\varepsilon^{2/r}(11/10)) \leq \delta_2(1 + 4\|g\|_2^{-1}),$$

for $N^{\nu-1/2} \leq \varepsilon \leq 1$. Invoking (89) and using the inequality $\|g_s\|_r^r \leq 2^r\|g\|_r^r$ and the identity $\|g_s\|_2^2 = 2\|g\|_2^2$ we obtain

$$|v|^{r-2} \leq \frac{c_r}{2^r} \frac{\|g\|_2^2}{\|g\|_r^r} \leq \frac{c_r}{2^r} \frac{2^{-1}\|g_s\|_2^2}{2^{-r}\|g_s\|_r^r} \leq \frac{c_r}{2} \frac{\|g_s\|_2^2}{\|g_s\|_r^r}$$

as required by Lemma 12 **a)**. This lemma implies

$$1 - |\mathbf{E}e^{i\tilde{X}}|^2 \geq 6^{-1}\|h_s\|_2^2 = 3^{-1}\|h\|_2^2.$$

In the last step we used (108). Now (120), for $c_0 \geq 10$, contradicts (113).

Step 4.2.2. Consider the case where $\delta_2 < |s - t| \leq \delta_1 N^{-\nu+1/2}$.

It follows from (115), (110) and (111) that

$$\begin{aligned} \mathbf{E}|h| &\leq \|h\|_r \leq c_g(2\delta_2\varepsilon^{2/r}|s/T_0| + \delta_1) \\ (121) \qquad \qquad \qquad &\leq c_g(\delta_1 + 3\delta_2\varepsilon^{2/r}) \leq c_g\delta_1 + \varepsilon^{2/r}. \end{aligned}$$

In the last step we used $\delta_2 < 1/3$. From (117), (118) and (111), we obtain for $\delta_2 \leq |s - t|$ and $N^{\nu-1/2} \leq \varepsilon$,

$$\begin{aligned} |v| &\geq \delta_2(1 - N^{\nu-1/2}\|g\|_2^{-1}) - 2\delta_2\varepsilon^{2/r}|s/T_0|\|g\|_2^{-1} \\ &= \delta_2(1 - \|g\|_2^{-1}(\varepsilon + \varepsilon^{2/r}(22/10))) \\ &\geq \delta_2(1 - 3\varepsilon^{2/r}\|g\|_2^{-1}) \geq \delta_2/2, \end{aligned}$$

provided that $\varepsilon^{2/r} < \|g\|_2/6$. Similarly, using in addition, $\delta_1, \delta_2 < 1/4$ and $\varepsilon < \|g\|_2$, we obtain, for $|s - t| \leq \delta_1 N^{-\nu+1/2}$,

$$\begin{aligned} |v| &\leq |s - t|(1 + N^{\nu-1/2}\|g\|_2^{-1}) + 2\delta_2\varepsilon^{2/r}|s/T_0|\|g\|_2^{-1} \\ &\leq |s - t|(1 + \varepsilon\|g\|_2^{-1}) + (22/10)\delta_2\varepsilon^{2/r}\|g\|_2^{-1} \\ &\leq 2|s - t| + 1 \leq N^{-\nu+1/2}. \end{aligned}$$

It follows from these inequalities, see (90), that

$$1 - |\mathbf{E}e^{i\tilde{X}}|^2 \geq 1 - |\mathbf{E}e^{i\tilde{X}}| \geq 1 - |\mathbf{E}e^{ivg} - \mathbf{E}|h|| \geq \delta' - \mathbf{E}|h|.$$

Finally, invoking (121) and (33), we get

$$1 - |\mathbf{E}e^{i\tilde{X}}|^2 \geq \delta' - c_g\delta_1 - \varepsilon^{2/r} \geq \delta'/2 > 4\varepsilon^2,$$

Once again we obtain a contradiction to (113), thus completing the proof.

5. EXPANSIONS

Here we prove the bound

$$(122) \quad \int_{|t| \leq t_1} \left| \mathbf{E} e^{it\tilde{\mathbb{T}}} - \hat{G}(t) \right| \frac{dt}{|t|} \leq c_* N^{-1-\nu},$$

where $t_1 = N^{1/2}/10^3\beta_3$. For the definition of $\tilde{\mathbb{T}}$ and \hat{G} see section **2.1**. Here and below c_* denotes a constant depending on $A_*, M_*, D_*, r, s, \nu_1$ only. We prove (122) for sufficiently large N , that is, we shall assume that $N > C_*$, where C_* is a number depending on $A_*, M_*, D_*, r, s, \nu_1$ only. Note that for $N < C_*$, the bound (122) becomes trivial, since in this case the integral is bounded by a constant.

Let us first introduce some notation. Denote $\Omega_m = \{1, \dots, m\}$. For $A \subset \Omega_N$ write $\mathbb{U}_1(A) = \sum_{j \in A} g_1(X_j)$. Given complex valued functions f, h we write $f \prec \mathcal{R}$ if

$$\int_{|t| \leq t_1} |t^{-1} f(t)| dt \leq c_* N^{-1-\nu}$$

and write $f \sim h$ if $f - h \prec \mathcal{R}$. In particular, (122) can be written in short $\mathbf{E} e^{it\tilde{\mathbb{T}}} \sim \hat{G}(t)$. In order to prove (122) we show that

$$(123) \quad \mathbf{E} e^{it\tilde{\mathbb{T}}} \sim \mathbf{E} e^{it\mathbb{T}} \quad \text{and} \quad \mathbf{E} e^{it\mathbb{T}} \sim \hat{G}(t).$$

In what follows we use the notation of Section 2 and assume that (14) holds.

5.1. Let us prove the first part of (123). Write

$$\mathbb{T} = \tilde{\mathbb{T}} + \tilde{\Lambda}_1 + \tilde{\Lambda}_2, \quad \tilde{\Lambda}_1 = \Lambda_1 + \Lambda_4, \quad \tilde{\Lambda}_2 = \Lambda_2 + \Lambda_3 + \Lambda_5,$$

where the random variables Λ_j are introduced in **2.3**. We shall show that

$$(124) \quad \mathbf{E} e^{it\tilde{\mathbb{T}}} \sim \mathbf{E} e^{it(\tilde{\mathbb{T}} + \tilde{\Lambda}_1)} \quad \text{and} \quad \mathbf{E} e^{it(\tilde{\mathbb{T}} + \tilde{\Lambda}_1)} \sim \mathbf{E} e^{it\mathbb{T}}.$$

The second relation follows from the moment bounds of Lemma 5 via Taylor expansion. We have

$$\mathbf{E} e^{it\mathbb{T}} = \mathbf{E} e^{it(\tilde{\mathbb{T}} + \tilde{\Lambda}_1)} + R, \quad |R| \leq |t| \mathbf{E} |\tilde{\Lambda}_2|,$$

By Lyapunov's inequality,

$$\mathbf{E} |\tilde{\Lambda}_2| \leq (\mathbf{E} \Lambda_2^2)^{1/2} + (\mathbf{E} \Lambda_3^2)^{1/2} + (\mathbf{E} \Lambda_5^2)^{1/2}.$$

Invoking the moment bounds of Lemma 5 we obtain $|t| \mathbf{E} |\tilde{\Lambda}_2| \prec \mathcal{R}$, thus, proving the second part of (124).

In order to prove the first part we combine Taylor's expansion with bounds for characteristic functions. Expanding the exponent we obtain

$$\mathbf{E} e^{it(\tilde{\mathbb{T}} + \tilde{\Lambda}_1)} = \mathbf{E} e^{it\tilde{\mathbb{T}}} + it \mathbf{E} e^{it\tilde{\mathbb{T}}} \tilde{\Lambda}_1 + R, \quad |R| \leq t^2 \mathbf{E} |\tilde{\Lambda}_1|^2.$$

Invoking the identities

$$(125) \quad \mathbf{E} \Lambda_1^2 = \binom{m}{2} \frac{\gamma_2}{N^3}, \quad \mathbf{E} \Lambda_4^2 = m \binom{N-m}{2} \frac{\zeta_2}{N^5}$$

we obtain, for $\gamma_2 < c_*$ and $\zeta_2 < c_*$, see (3), and $m \leq N^{1/12}$, that $R \prec \mathcal{R}$. We complete the proof of (124) by showing that

$$(126) \quad t \mathbf{E} e^{it\tilde{\mathbb{T}}} \tilde{\Lambda}_1 \prec \mathcal{R}.$$

Let us prove (126). Split $\mathbb{W} = \mathbb{W}_1 + \mathbb{W}_2 + \mathbb{W}_3 + R_W$, where

$$\mathbb{W}_k = \sum_{A \subset \Omega', |A|=k} T_A, \quad R_W = \sum_{A \subset \Omega', |A| \geq 4} T_A.$$

Here $\Omega' = \{m+1, \dots, N\}$. Denote $\mathbb{R} = \mathbb{U}_2^* + \mathbb{W}_3 + R_W$ and $\mathbb{U}_1 = \sum_{j=1}^N g_1(X_j)$. We have $\tilde{\mathbb{T}} = \mathbb{U}_1 + \mathbb{W}_2 + \mathbb{R}$. Expanding the exponent in powers of $it\mathbb{R}$ we obtain

$$(127) \quad t\mathbf{E}e^{it\tilde{\mathbb{T}}}\tilde{\Lambda}_1 = t\mathbf{E}e^{it(\mathbb{U}_1+\mathbb{W}_2)}\tilde{\Lambda}_1 + t^2R,$$

where

$$|R| \leq \mathbf{E}|\tilde{\Lambda}_1\mathbb{R}| \leq (r_1 + r_2)(r_3 + r_4 + r_5),$$

$$r_1^2 = \mathbf{E}\Lambda_1^2, \quad r_2^2 = \mathbf{E}\Lambda_4^2, \quad r_3^2 = \mathbf{E}(\mathbb{U}_2^*)^2, \quad r_4^2 = \mathbf{E}R_W^2, \quad r_5^2 = \mathbf{E}\mathbb{W}_3^2.$$

In the last step we applied the Cauchy-Schwartz inequality. Combining (125) with the identities

$$\mathbf{E}(\mathbb{U}_2^*)^2 = \frac{m(N-m)}{N^3}\gamma_2, \quad \mathbf{E}\mathbb{W}_3^2 = \frac{\binom{N-m}{3}}{N^5}\zeta_2$$

and invoking the simple bound

$$\mathbf{E}R_W^2 \leq \frac{\Delta_4^2}{N^3} \leq \frac{D_*}{N^{2+2\nu_1}},$$

we obtain $t^2(r_1 + r_2)(r_3 + r_4 + r_5) \prec \mathcal{R}$. Therefore, (127) implies

$$t\mathbf{E}e^{it\tilde{\mathbb{T}}}\tilde{\Lambda}_1 \sim t\mathbf{E}e^{it(\mathbb{U}_1+\mathbb{W}_2)}\tilde{\Lambda}_1.$$

Let us show that $t\mathbf{E}e^{it(\mathbb{U}_1+\mathbb{W}_2)}\tilde{\Lambda}_1 \sim 0$. Expanding the exponent in powers of $it\mathbb{W}_2$ we get

$$t\mathbf{E}e^{it(\mathbb{U}_1+\mathbb{W}_2)}\tilde{\Lambda}_1 = f_1(t) + f_2(t) + f_3(t) + f_4(t),$$

$$f_1(t) = t\mathbf{E}e^{it\mathbb{U}_1}\tilde{\Lambda}_1, \quad f_2(t) = it^2\mathbf{E}e^{it\mathbb{U}_1}\Lambda_1\mathbb{W}_2,$$

$$f_3(t) = t^2\mathbf{E}e^{it\mathbb{U}_1}\Lambda_4\mathbb{W}_2\theta_1, \quad f_4(t) = t^3\mathbf{E}e^{it\mathbb{U}_1}\Lambda_1\mathbb{W}_2^2\theta_2/2,$$

where θ_1, θ_2 are functions of \mathbb{W}_2 satisfying $|\theta_i| \leq 1$.

Let us show that $f_i \prec \mathcal{R}$, for $i = 1, 2, 3, 4$. Split the set $\Omega_m = \{1, \dots, m\}$ in three (non-intersecting) parts $A_1 \cup A_2 \cup A_3 = \Omega_m$ of (almost) equal size $|A_i| \approx m/3$. The set of pairs $\{\{i, j\} \subset \Omega_m\}$ splits into six (non-intersecting) parts B_{kr} , $1 \leq k \leq r \leq 3$ (the pair $\{i, j\}$ belongs to B_{kr} if $i \in A_k$ and $j \in A_r$). Write

$$\Lambda_1 = \sum_{1 \leq k \leq r \leq 3} \Lambda_1(k, r), \quad \Lambda_1(k, r) = \sum_{\{i, j\} \in B_{kr}} g_2(X_k, X_l),$$

$$\Lambda_4 = \sum_{1 \leq k \leq 3} \Lambda_4(k), \quad \Lambda_4(k) = \sum_{i \in A_k} \sum_{m+1 \leq j < l \leq N} g_3(X_i, X_j, X_l).$$

Let us prove $f_4 \prec \mathcal{R}$. We shall show that

$$(128) \quad t^3\mathbf{E}e^{it\mathbb{U}_1}\Lambda_1(k, r)\mathbb{W}_2^2\theta_2 \prec \mathcal{R}.$$

Given a pair (k, r) denote $A_i = \Omega_m \setminus (A_k \cup A_r)$ and write $k_i = |A_i|$. Note that $k_i \approx m/3$. We shall assume that $k_i \geq m/4$. Since the random variable $\mathbb{U}_1(A_i) := \sum_{j \in A_i} g_1(X_j)$ and the random variables $\Lambda_1(k, r)$, \mathbb{W}_2 are independent, we have

$$\mathbf{E}e^{it\mathbb{U}_1}\Lambda_1(k, r)\mathbb{W}_2^2\theta_2 \leq \mathbf{E}e^{it\mathbb{U}_1(A_i)}\mathbf{E}\Lambda_1(k, r)\mathbb{W}_2^2\theta_2.$$

Therefore,

$$(129) \quad |\mathbf{E}e^{it\mathbb{U}_1}\Lambda_1(k, r)\mathbb{W}_2^2\theta_2| \leq |\mathbf{E}e^{it\mathbb{U}_1(A_i)}| \mathbf{E}|\Lambda_1(k, r)\mathbb{W}_2^2|.$$

The first factor on the right is bounded from above by $\exp\{-mt^2/16N\}$, for $k_i \geq m/4$, see (160) below. The second factor is bounded from above by r , where

$$r^2 = \mathbf{E}\Lambda_1^2(k, r)\mathbf{E}\mathbb{W}_2^4 \leq c_*m^2N^{-5}.$$

Here we combined the Cauchy-Schwartz inequality and the bounds

$$\mathbf{E}\Lambda_1^2(k, r) \leq c_*m^2N^{-3}, \quad \mathbf{E}\mathbb{W}_2^4 \leq c_*N^{-2}.$$

Finally, (128) follows from (129)

$$|t^3 \mathbf{E} e^{it\mathbb{U}_1} \Lambda_1(k, r) \mathbb{W}_2^2 \theta_2| \leq c_* |t|^3 e^{-mt^2/16N} m N^{-5/2} \prec \mathcal{R}.$$

The proof of $f_3 \prec \mathcal{R}$ is almost the same as that of $f_4 \prec \mathcal{R}$.

Let us prove $f_2 \prec \mathcal{R}$. Split the set $\Omega' = \{m+1, \dots, N\}$ into three (non-intersecting) parts $B_1 \cup B_2 \cup B_3 = \Omega'$ of (almost) equal sizes $|B_i| \approx (N-m)/3$. Split the set of pairs $\{\{i, j\} : m+1 \leq i < j \leq N\}$ into (non-intersecting) groups $D(k, r)$, for $1 \leq k \leq r \leq 3$. The pair $\{i, j\} \in D(k, r)$ if $i \in B_k$ and $j \in B_r$. Write

$$\begin{aligned} \mathbb{W}_2 &= \sum_{1 \leq k \leq r \leq 3} \mathbb{W}_2(k, r), & \mathbb{W}_2(k, r) &= \sum_{\{i, j\} \in D(k, r)} g_2(X_i, X_j). \\ \Lambda_4 &= \sum_{1 \leq k \leq r \leq 3} \Lambda_4(k, r), & \Lambda_4(k, r) &= \sum_{1 \leq s \leq m} \sum_{\{i, j\} \in D(k, r)} g_3(X_s, X_i, X_j), \end{aligned}$$

In order to prove $f_2 \prec \mathcal{R}$ we shall show that

$$(130) \quad t^2 \mathbf{E} e^{it\mathbb{U}_1} \Lambda_1 \mathbb{W}_2(k, r) \prec \mathcal{R}.$$

Write $B_i = \Omega' \setminus (B_k \cup B_r)$ and denote $m_i = |B_i|$. We shall assume that $m_i \geq N/4$. Since the random variable $\mathbb{U}_1(B_i) = \sum_{j \in B_i} g_1(X_j)$ and the random variables Λ_1 and $\mathbb{W}_2(k, r)$ are independent, we have, cf. (129),

$$(131) \quad |\mathbf{E} e^{it\mathbb{U}_1} \Lambda_1 \mathbb{W}_2(k, r)| \leq |\mathbf{E} e^{it\mathbb{U}_1(B_i)}| \mathbf{E} |\Lambda_1 \mathbb{W}_2(k, r)|.$$

The first factor in the right is the product $|\alpha^{m_i}(t)| \leq e^{-m_i t^2/4N}$, see the argument used in the proof of (128) above. The second factor is bounded from above by r , where

$$r^2 = \mathbf{E} \Lambda_1^2 \mathbf{E} \mathbb{W}_2^2(k, r) \leq c_* m^2 N^{-4}.$$

Finally, we obtain, using the inequality $m_i \geq N/4$,

$$|\mathbf{E} e^{it\mathbb{U}_1} \mathbf{E} |\Lambda_1 \mathbb{W}_2(k, r)| \leq c_* \frac{m}{N^2} \exp\{-t^2 \frac{m_i}{4N}\} \leq c_* \frac{m}{N^2} \exp\{-\frac{t^2}{16}\}.$$

This in combination with (131) shows (130). We obtain $f_2 \prec \mathcal{R}$.

Let us prove $f_1 \prec \mathcal{R}$. We shall show that $f^* \prec \mathcal{R}$ and $f^* \prec \mathcal{R}$, where

$$f^* = t \mathbf{E} e^{it\mathbb{U}_1} \Lambda_1 \quad \text{and} \quad f^* = t \mathbf{E} e^{it\mathbb{U}_1} \Lambda_4$$

satisfy $f^* + f^* = f_1$.

Let us show $f^* \prec \mathcal{R}$. Denote $\mathbb{U}_1^* = \sum_{j=m+1}^N g_1(X_j)$. We obtain, by the independence of \mathbb{U}_1^* and Λ_1 that

$$|\mathbf{E} e^{it\mathbb{U}_1} \Lambda_1| \leq |\mathbf{E} e^{it\mathbb{U}_1^*}| \mathbf{E} |\Lambda_1|.$$

Invoking, for $N-m > N/2$, the bound $|\mathbf{E} e^{it\mathbb{U}_1^*}| \leq e^{-t^2/8}$, see (160) below, and the bound $\mathbf{E} |\Lambda_1| \leq (\mathbf{E} \Lambda_1^2)^{1/2} \leq c_* m N^{-3/2}$ we obtain

$$|f^*(t)| \leq c_* |t| e^{-t^2/8} N^{-3/2} \prec \mathcal{R}.$$

Let us prove $f^* \prec \mathcal{R}$. We shall show that, for $1 \leq k \leq r \leq 3$,

$$(132) \quad t \mathbf{E} e^{it\mathbb{U}_1} \Lambda_4(k, r) \prec \mathcal{R}.$$

Proceeding as in the proof of (130) we obtain the chain of inequalities

$$(133) \quad |\mathbf{E} e^{it\mathbb{U}_1} \Lambda_4(k, r)| \leq e^{-t^2/16} \mathbf{E} |\Lambda_4(k, r)| \leq c_* e^{-t^2/16} m^{1/2} N^{-3/2}.$$

In the last step we applied Cauchy-Schwartz and the simple bound $\mathbf{E} \Lambda_4^2(k, r) \leq c_* m N^{-3}$. Clearly, (133) implies (132).

5.2. Here we prove the second relation of (123). Firstly, we shall show that

$$(134) \quad \mathbf{E}e^{it\mathbb{T}} \sim \mathbf{E} \exp\{it(\mathbb{U}_1 + \mathbb{U}_2 + \mathbb{U}_3)\},$$

$$(135) \quad \mathbf{E} \exp\{it(\mathbb{U}_1 + \mathbb{U}_2 + \mathbb{U}_3)\} \sim \mathbf{E} \exp\{it(\mathbb{U}_1 + \mathbb{U}_2)\} + \binom{N}{3} e^{-t^2/2} (it)^4 w,$$

where $w = \mathbf{E}g_3(X_1, X_2, X_3)g_1(X_1)g_1(X_2)g_1(X_3)$.

Let $m(t)$ be an integer valued function such that

$$(136) \quad m(t) \approx C_1 N t^{-2} \ln(t^2 + 1), \quad C_1 \leq |t| \leq t_1,$$

and put $m(t) \equiv 10$, for $|t| \leq C_1$. Here C_1 denotes a large absolute constant (one can take, e.g., $C_1 = 200$). Assume, in addition, that the numbers $m = m(t)$ are even.

5.2.1. Here we prove (5B.1). Given m write

$$\mathbb{T} = \mathbb{U}_1 + \mathbb{U}_2 + \mathbb{U}_3 + \mathbb{H},$$

where

$$\mathbb{H} = \mathbb{H}_1 + \mathbb{H}_2, \quad \mathbb{H}_1 = \sum_{|A| \geq 4, A \cap \Omega_m = \emptyset} T_A, \quad \mathbb{H}_2 = \sum_{|A| \geq 4, A \cap \Omega_m \neq \emptyset} T_A.$$

In order to show (134) we expand the exponent in powers of $it\mathbb{H}$ and $it\mathbb{U}_3$,

$$\mathbf{E} \exp\{it\mathbb{T}\} = \mathbf{E} \exp\{it(\mathbb{U}_1 + \mathbb{U}_2 + \mathbb{U}_3)\} + \mathbf{E} \exp\{it(\mathbb{U}_1 + \mathbb{U}_2)\} it\mathbb{H} + R,$$

where $|R| \leq t^2(\mathbf{E}\mathbb{H}^2 + \mathbf{E}|\mathbb{U}_3\mathbb{H}|)$. Invoking the bounds, see (161), (162), (3), (4),

$$(137) \quad \mathbf{E}\mathbb{H}^2 \leq N^{-3} \Delta_4^2 \leq c_* N^{-2-2\nu_1}, \quad \mathbf{E}\mathbb{U}_3^2 \leq N^{-2} \zeta_2 \leq c_* N^{-2}$$

we obtain, by Cauchy-Schwartz, $|R| \leq c_* t^2 N^{-2-\nu_1} \prec \mathcal{R}$. We complete the proof of (134) by showing that

$$(138) \quad \mathbf{E} \exp\{it(\mathbb{U}_1 + \mathbb{U}_2)\} it\mathbb{H} \prec \mathcal{R}.$$

Before proving (138) we collect some auxiliary inequalities. For $m = 2k$ write

$$(139) \quad \Omega_m = A_1 \cup A_2, \quad \text{where} \quad A_1 = \{1, \dots, k\}, \quad A_2 = \{k+1, \dots, 2k\}.$$

Furthermore, split the sum

$$(140) \quad \begin{aligned} \mathbb{U}_2 &= \mathbb{Z}_1 + \mathbb{Z}_2 + \mathbb{Z}_3 + \mathbb{Z}_4, \\ \mathbb{Z}_1 &= \sum_{1 \leq i < j \leq m} g_2(X_i, X_j), & \mathbb{Z}_2 &= \sum_{i \in A_1} \sum_{m < j \leq N} g_2(X_i, X_j), \\ \mathbb{Z}_3 &= \sum_{i \in A_2} \sum_{m < j \leq N} g_2(X_i, X_j), & \mathbb{Z}_4 &= \sum_{m < i < j \leq N} g_2(X_i, X_j). \end{aligned}$$

In what follows we shall use the simple bounds, see (3),

$$(141) \quad \begin{aligned} \mathbf{E}\mathbb{Z}_1^2 &\leq \frac{m^2}{N^3} \gamma_2 \leq c_* \frac{m^2}{N^3}, & \mathbf{E}\mathbb{Z}_4^2 &\leq \frac{\gamma_2}{N} \leq \frac{c_*}{N}, \\ \mathbf{E}\mathbb{Z}_i^2 &\leq \frac{m}{N^2} \gamma_2 \leq c_* \frac{m}{N^2}, & \mathbf{E}\mathbb{Z}_i^4 &\leq c \frac{m^2}{N^4} \gamma_4 \leq c_* \frac{m^2}{N^4}, \quad i = 2, 3. \end{aligned}$$

Let us prove (138). Expand the exponent $\exp\{it(\mathbb{U}_1 + \mathbb{Z}_1 + \dots + \mathbb{Z}_4)\}$ in powers of $it\mathbb{Z}_1$ to get

$$\mathbf{E} \exp\{it(\mathbb{U}_1 + \mathbb{U}_2)\} it\mathbb{H} = h_1(t) + R$$

where $h_1(t) = \mathbf{E} \exp\{it(\mathbb{U}_1 + \mathbb{Z}_2 + \dots + \mathbb{Z}_4)\} it\mathbb{H}$ and where

$$|R| \leq t^2 \mathbf{E}|\mathbb{H}\mathbb{Z}_1| \leq t^2 (\mathbf{E}\mathbb{H}^2)^{1/2} (\mathbf{E}\mathbb{Z}_1^2)^{1/2} \leq c_* t^2 m N^{-(5+2\nu_1)/2}.$$

For $m = m(t)$ satisfying (136) we have $R \prec \mathcal{R}$. Therefore, we obtain

$$\mathbf{E} \exp\{it(\mathbb{U}_1 + \mathbb{U}_2)\}it\mathbb{H} \sim h_1.$$

In order to prove $h_1 \prec \mathcal{R}$ we write $h_1 = h_2 + h_3$ and show that $h_2, h_3 \prec \mathcal{R}$, where

$$h_2 = \mathbf{E} \exp\{it(\mathbb{U}_1 + \mathbb{Z}_2 + \cdots + \mathbb{Z}_4)\}it\mathbb{H}_1, \quad h_3 = \mathbf{E} \exp\{it(\mathbb{U}_1 + \mathbb{Z}_2 + \cdots + \mathbb{Z}_4)\}it\mathbb{H}_2.$$

Let us show that $h_2 \prec \mathcal{R}$. Firstly, we prove that

$$(142) \quad h_2 \sim h_{2,1} + h_{2,2} + h_{2,3},$$

where $h_{2,1}(t) = \mathbf{E} \exp\{it(\mathbb{U}_1 + \mathbb{Z}_4)\}it\mathbb{H}_1$ and, for $j = 2, 3$,

$$h_{2,j}(t) = \mathbf{E} \exp\{it(\mathbb{U}_1 + \mathbb{Z}_4)\}(it)^2\mathbb{H}_1\mathbb{Z}_j.$$

Expanding the exponent in powers of $it(\mathbb{Z}_2 + \mathbb{Z}_3)$ we obtain

$$h_2 = h_{2,1} + h_{2,2} + h_{2,3} + R,$$

where $|R| \leq |t|^3 \mathbf{E}|\mathbb{H}_1|(\mathbb{Z}_2 + \mathbb{Z}_3)^2$ is bounded from above by

$$|t|^3 (\mathbf{E}\mathbb{H}_1^2)^{1/2} (\mathbf{E}(\mathbb{Z}_2 + \mathbb{Z}_3)^4)^{1/2} \leq c_* |t|^3 m N^{-3-\nu_1} \prec \mathcal{R}.$$

In the last step we used $\mathbf{E}\mathbb{H}_1^2 \leq \mathbf{E}\mathbb{H}^2$ and applied (137) and (141). Therefore, (142) follows.

Let us show $h_{2,i} \prec \mathcal{R}$, for $i = 1, 2, 3$. The random variable $\mathbb{U}_1(A_1)$ does not depend on the observations X_j , $j \in \Omega \setminus A_1$. Therefore, we can write

$$h_{2,3} = \mathbf{E} \exp\{it\mathbb{U}_1(A_1)\} \mathbf{E} \exp\{it(\mathbb{U}_1(\Omega \setminus A_1) + \mathbb{Z}_4)\}(it)^2\mathbb{H}_1\mathbb{Z}_3.$$

Furthermore, using (160) we obtain, for $|A_1| = m/2$,

$$(143) \quad |h_{2,3}| \leq t^2 |\alpha^{m/2}(t)| \mathbf{E}|\mathbb{H}_1\mathbb{Z}_3| \leq c_* t^2 \exp\{-t^2 \frac{m}{8N}\} \frac{m^{1/2}}{N^{2+\nu_1}}.$$

In the last step we combined the bound $\mathbf{E}\mathbb{H}_1^2 \leq c_* N^{-2-2\nu_1}$ and (141) to get

$$\mathbf{E}|\mathbb{H}_1\mathbb{Z}_3| \leq (\mathbf{E}\mathbb{H}_1^2)^{1/2} (\mathbf{E}\mathbb{Z}_3^2)^{1/2} \leq c_* m^{1/2} N^{-2-\nu_1}.$$

Note that choosing of C_1 in (136) sufficiently large implies, for $|t| \geq C_1$,

$$t^2 m / 12N \approx (C_1/12) \ln(t^2 + 1) \geq 10 \ln(t^2 + 1).$$

An application of this bound to the argument of the exponent in (143) shows $h_{2,3} \prec \mathcal{R}$. The proof of $h_{2,i} \prec \mathcal{R}$, for $i = 1, 2$, is almost the same. Therefore, we obtain $h_2 \prec \mathcal{R}$.

Let us prove $h_3 \prec \mathcal{R}$. Firstly we collect some auxiliary inequalities. Write $m = 2k$ (recall that the number m is even) and split $\Omega_m = B \cup D$, where B denotes the set of odd numbers and D denotes the set of even numbers. Split $\mathbb{H}_2 = \mathbb{H}_B + \mathbb{H}_D + \mathbb{H}_C$. Here, for $A \subset \Omega_N$ and $|A| \geq 4$, we denote by \mathbb{H}_B the sum of T_A such that $A \cap B = \emptyset$ and $A \cap D \neq \emptyset$; \mathbb{H}_D denotes the sum of T_A such that $A \cap B \neq \emptyset$ and $A \cap D = \emptyset$; \mathbb{H}_C denotes the sum of T_A such that $A \cap B \neq \emptyset$ and $A \cap D \neq \emptyset$. It follows from the inequalities (172) and (4) that

$$(144) \quad \mathbf{E}\mathbb{H}_C^2 \leq c_* m^2 N^{-4-2\nu_1}, \quad \mathbf{E}\mathbb{H}_B^2 = \mathbf{E}\mathbb{H}_D^2 \leq c_* m N^{-3-2\nu_1}.$$

Using the notation $z = it \exp\{it(\mathbb{U}_1 + \mathbb{Z}_2 + \mathbb{Z}_3 + \mathbb{Z}_4)\}$ write

$$h_3 = \mathbf{E}z\mathbb{H}_2 = h_{3,1} + h_{3,2} + h_{3,3},$$

$$h_{3,1} = \mathbf{E}z\mathbb{H}_B, \quad h_{3,2} = \mathbf{E}z\mathbb{H}_D, \quad h_{3,3} = \mathbf{E}z\mathbb{H}_C.$$

We shall show that $h_{3,i} \prec \mathcal{R}$, for $i = 1, 2, 3$. The relation $h_{3,3} \prec \mathcal{R}$ follows from (144) and (141), and by Cauchy-Schwartz, $|h_{3,3}| \leq c_* |t| m N^{-2-\nu_1} \prec \mathcal{R}$.

Let us show that $h_{3,2} \prec \mathcal{R}$. Expanding the exponent in powers of $it(\mathbb{Z}_2 + \mathbb{Z}_3)$ we obtain

$$h_{3,2} = h_{3,2}^* + R, \quad h_{3,2}^* := \mathbf{E} \exp\{it(\mathbb{U}_1 + \mathbb{Z}_4)\}it\mathbb{H}_D,$$

where $|R| \leq t^2 \mathbf{E}|\mathbb{H}_D(\mathbb{Z}_2 + \mathbb{Z}_3)|$. Combining the bounds (141) and (144) we obtain, by Cauchy-Schwartz, $|R| \leq c_* t^2 m N^{-(5+2\nu_1)/2} \prec \mathcal{R}$. Next we show that $h_{3,2}^* \prec \mathcal{R}$. The random variable $\mathbb{U}_1(D) = \sum_{j \in D} g_1(X_j)$ and the random variable \mathbb{H}_D are independent. Therefore, we can write

$$|h_{3,2}^*| \leq |t| |\mathbf{E} \exp\{it\mathbb{U}_1(D)\}| |\mathbf{E}|\mathbb{H}_D|.$$

Combining (160) and (144) we obtain using Cauchy-Schwartz,

$$|h_{3,2}^*| \leq c_* |t| e^{-mt^2/8N} m^{1/2} N^{-(3+2\nu_1)/2} \prec \mathcal{R}.$$

The proof of $h_{3,1} \prec \mathcal{R}$ is similar. Therefore, we obtain $h_3 \prec \mathcal{R}$. This together with the relation $h_2 \prec \mathcal{R}$, proved above, implies $h_1 \prec \mathcal{R}$. Thus we arrive at (138) completing the proof of (134).

5.2.2. Here we prove (135). We start with some auxiliary moment inequalities. Split

$$\mathbb{U}_3 = W + Z, \quad W = \sum_{|A|=3, A \cap \Omega_m \neq \emptyset} T_A, \quad Z = \sum_{|A|=3, A \cap \Omega_m = \emptyset} T_A.$$

Using the orthogonality and moment bounds for U -statistics, see, e.g., Dharmadhikari et al (1968), one can show that

$$\mathbf{E}W^2 \leq mN^2 \mathbf{E}g_3^2(X_1, X_2, X_3), \quad \mathbf{E}Z^2 \leq N^3 \mathbf{E}g_3^2(X_1, X_2, X_3),$$

and $\mathbf{E}|Z|^s \leq cN^{3s/2} \mathbf{E}|g_3(X_1, X_2, X_3)|^s$. Invoking (3) we obtain

$$(145) \quad \mathbf{E}W^2 \leq c_* m N^{-3}, \quad \mathbf{E}Z^2 \leq c_* N^{-2}, \quad \mathbf{E}|Z|^s \leq c_* N^{-s}.$$

For the sets $A_1, A_2 \subset \Omega_m$ defined in (139) write

$$\begin{aligned} \mathcal{D} &= \{A \subset \Omega_N : |A| = 3, A \cap \Omega_m \neq \emptyset\}, \\ \mathcal{D}_1 &= \{A \in \mathcal{D} : A \cap A_1 = \emptyset\}, \\ \mathcal{D}_2 &= \{A \in \mathcal{D} : A \cap A_2 = \emptyset\}, \\ \mathcal{D}_3 &= \{A \in \mathcal{D} : A \cap A_1 \neq \emptyset, A \cap A_2 \neq \emptyset\}. \end{aligned}$$

We have $\mathcal{D} = \mathcal{D}_1 \cup \mathcal{D}_2 \cup \mathcal{D}_3$ and $W = \sum_{A \in \mathcal{D}} T_A$. Therefore, we can write $W = W_1 + W_2 + W_3$, where $W_j = \sum_{A \in \mathcal{D}_j} T_A$.

A calculation shows that

$$\mathbf{E}W_1^2 = \mathbf{E}W_2^2 \leq kN^2 \mathbf{E}g_3^2(X_1, X_2, X_3), \quad \mathbf{E}W_3^2 \leq k^2 N \mathbf{E}g_3^2(X_1, X_2, X_3).$$

Therefore, we obtain from (3) that

$$(146) \quad \mathbf{E}W_1^2 = \mathbf{E}W_2^2 \leq c_* m N^{-3}, \quad \mathbf{E}W_3^2 \leq c_* m^2 N^{-4}.$$

Let us prove (135). Write $\mathbb{U}_3 = W + Z$. Expanding the exponent in powers of itW we obtain

$$\begin{aligned} \mathbf{E} \exp\{it(\mathbb{U}_1 + \mathbb{U}_2 + \mathbb{U}_3)\} &= h_4 + h_5 + R, \\ h_4 &= \mathbf{E} \exp\{it(\mathbb{U}_1 + \mathbb{U}_2 + Z)\}, \\ h_5 &= \mathbf{E} \exp\{it(\mathbb{U}_1 + \mathbb{U}_2 + Z)\} itW, \end{aligned}$$

where, by (145), $|R| \leq t^2 \mathbf{E}W^2 \leq c_* t^2 m N^{-3} \prec \mathcal{R}$. This implies

$$\mathbf{E} \exp\{it(\mathbb{U}_1 + \mathbb{U}_2 + \mathbb{U}_3)\} \sim h_4 + h_5.$$

In order to prove (135) we shall show that

$$(147) \quad h_5 \sim \mathbf{E} \exp\{it\mathbb{U}_1\} itW,$$

$$(148) \quad h_4 \sim \mathbf{E} \exp\{it(\mathbb{U}_1 + \mathbb{U}_2)\} + \mathbf{E} \exp\{it\mathbb{U}_1\} itZ,$$

$$(149) \quad \mathbf{E} \exp\{it\mathbb{U}_1\} it\mathbb{U}_3 \sim \binom{N}{3} e^{-t^2/2} (it)^4 w.$$

Let us prove (147). Expanding the exponent (in h_5) in powers of itZ we obtain

$$h_5 = h_6 + R, \quad h_6 = \mathbf{E} \exp\{it(\mathbb{U}_1 + \mathbb{U}_2)\}itW,$$

where, by (145) and Cauchy-Schwartz,

$$|R| \leq t^2 \mathbf{E}|WZ| \leq c_* t^2 m^{1/2} N^{-5/2} \prec \mathcal{R}.$$

We have, $h_5 \sim h_6$.

It remains to show that $h_6 \sim \mathbf{E} \exp\{it\mathbb{U}_1\}itW$. Split

$$(150) \quad \mathbb{U}_2 = \mathbb{U}_2^* + \mathbb{U}_2^*, \quad \mathbb{U}_2^* = \sum_{|A|=2, A \cap \Omega_m \neq \emptyset} T_A, \quad \mathbb{U}_2^* = \sum_{|A|=2, A \cap \Omega_m = \emptyset} T_A.$$

We have, see (141),

$$(151) \quad \mathbf{E}(\mathbb{U}_2^*)^2 \leq c_* m N^{-2}, \quad \mathbf{E}(\mathbb{U}_2^*)^2 \leq c_* N^{-1}.$$

Expanding the exponent (in h_6) in powers of $it\mathbb{U}_2^*$ we obtain

$$h_6 = h_7 + R, \quad \text{where} \quad h_7 = \mathbf{E} \exp\{it(\mathbb{U}_1 + \mathbb{U}_2^*)\}itW,$$

and where, by (145), (151) and Cauchy-Schwartz,

$$|R| \leq t^2 \mathbf{E}|W\mathbb{U}_2^*| \leq c_* t^2 m N^{-5/2} \prec \mathcal{R}.$$

Therefore, we obtain $h_6 \sim h_7$.

We complete the proof of (147) by showing that $h_7 \sim \mathbf{E} \exp\{it\mathbb{U}_1\}itW$. Use the decomposition $W = W_1 + W_2 + W_3$ and write

$$h_7 = h_{7.1} + h_{7.2} + h_{7.3}, \quad h_{7.j} = \mathbf{E} \exp\{it(\mathbb{U}_1 + \mathbb{U}_2^*)\}itW_j.$$

We shall show that

$$(152) \quad h_{7.j} \sim \mathbf{E} \exp\{it\mathbb{U}_1\}itW_j, \quad j = 1, 2, 3.$$

Expanding in powers of $it\mathbb{U}_2^*$ we obtain

$$h_{7.j} = \mathbf{E} \exp\{it\mathbb{U}_1\}itW_j + R_j,$$

where $R_j = (it)^2 \mathbf{E} \exp\{it\mathbb{U}_1\}W_j \mathbb{U}_2^* \theta$ and where θ is a function of \mathbb{U}_2^* satisfying $|\theta| \leq 1$. In order to prove (152) we show that $R_j \prec \mathcal{R}$, for $j = 1, 2, 3$.

Combining (146) and (151) we obtain via Cauchy-Schwartz

$$|R_3| \leq c_* t^2 m N^{-5/2} \prec \mathcal{R}.$$

Furthermore, using the fact that the random variable $\mathbb{U}_1(A_2)$ and the random variables \mathbb{U}_2^* and W_2 are independent, we can write

$$|R_2| \leq t^2 |\mathbf{E} \exp\{it\mathbb{U}_1(A_2)\}| \mathbf{E}|W_2 \mathbb{U}_2^*| \leq c_* t^2 e^{-mt^2/8N} m^{1/2} N^{-2} \prec \mathcal{R}.$$

Here we used (160) and the moment inequalities (146) and (151). The proof of $R_1 \prec \mathcal{R}$ is similar. We arrive at (152) and, thus, complete the proof of (147).

Let us prove (148). We proceed in two steps. Firstly we show

$$(153) \quad h_4 \sim h_8 + h_9, \\ h_8 = \mathbf{E} \exp\{it(\mathbb{U}_1 + \mathbb{U}_2)\}, \quad h_9 = \mathbf{E} \exp\{it(\mathbb{U}_1 + \mathbb{U}_2)\}itZ.$$

Secondly, we show

$$(154) \quad h_9 \sim \mathbf{E} \exp\{it\mathbb{U}_1\}itZ.$$

In order to prove (153) we write

$$h_4 = h_8 + h_9 + R, \quad R = \mathbf{E} \exp\{it(\mathbb{U}_1 + \mathbb{U}_2)\}r, \quad r = \exp\{itZ\} - 1 - itZ,$$

and show that $R \prec \mathcal{R}$. In order to bound the remainder R we write $\mathbb{U}_2 = \mathbb{U}_2^* + \mathbb{U}_2^*$, see (150), and expand the exponent in powers of $it\mathbb{U}_2^*$. We obtain $R = R_1 + R_2$, where

$$R_1 = \mathbf{E} \exp\{it(\mathbb{U}_1 + \mathbb{U}_2^*)\}r \quad \text{and} \quad |R_2| \leq \mathbf{E}|it\mathbb{U}_2^*r|.$$

Note that, for $2 < s \leq 3$, we have $|r| \leq c|tZ|^{s/2}$. Combining (145) and (151) we obtain via Cauchy-Schwartz,

$$|R_2| \leq |t|^{1+s/2} \mathbf{E}|Z|^{s/2} |\mathbb{U}_2^*| \leq c_* |t|^{1+s/2} m^{1/2} N^{-1-s/2} \prec \mathcal{R}.$$

In order to prove $R_1 \prec \mathcal{R}$ we use the fact that the random variable $\mathbb{U}_1(\Omega_m)$ and the random variables \mathbb{U}_2^* and r are independent. Invoking the inequality $|r| \leq t^2 Z^2$ we obtain from (160) and (145)

$$|R_1| \leq t^2 |\alpha^m(t)| \mathbf{E}Z^2 \leq c_* t^2 e^{-mt^2/4N} N^{-2} \prec \mathcal{R}.$$

We thus arrive at (153).

Let us prove (154). Use the decomposition (140) and expand the exponent (in h_9) in powers of $it\mathbb{Z}_1$ to get $h_9 = h_{10} + R$, where

$$h_{10} = \mathbf{E} \exp\{it(\mathbb{U}_1 + \mathbb{Z}_2 + \mathbb{Z}_3 + \mathbb{Z}_4)\}itZ, \quad |R| \leq t^2 \mathbf{E}|ZZ_1|.$$

Combining (141) and (145) we obtain via Cauchy-Schwartz

$$|R| \leq c_* t^2 m N^{-5/2} \prec \mathcal{R}.$$

Therefore, we have

$$h_9 \sim h_{10}.$$

Now we expand the exponent in h_{10} in powers of $it(\mathbb{Z}_2 + \mathbb{Z}_3)$ and obtain $h_{10} = h_{11} + h_{12} + R$, where

$$h_{11} = \mathbf{E} \exp\{it(\mathbb{U}_1 + \mathbb{Z}_4)\}itZ, \quad h_{12} = \mathbf{E} \exp\{it(\mathbb{U}_1 + \mathbb{Z}_4)\}(it)^2 Z(\mathbb{Z}_2 + \mathbb{Z}_3),$$

and where $|R| \leq |t|^3 \mathbf{E}|Z| |\mathbb{Z}_2 + \mathbb{Z}_3|^2$. Combining (141) and (145) we obtain via Cauchy-Schwartz $|R| \leq |t|^3 m N^{-3} \prec \mathcal{R}$. Therefore, we have

$$h_{10} \sim h_{11} + h_{12}.$$

We complete the proof of (154), by showing that

$$(155) \quad h_{11} \sim \mathbf{E} \exp\{it\mathbb{U}_1\}itZ \quad \text{and} \quad h_{12} \prec \mathcal{R}.$$

In order to prove the second bound write

$$h_{12} = R_2 + R_3, \quad \text{where} \quad R_j = \mathbf{E} \exp\{it(\mathbb{U}_1 + \mathbb{Z}_4)\}(it)^2 Z\mathbb{Z}_j.$$

We shall show that $R_3 \prec \mathcal{R}$. Using the fact that the random variable $\mathbb{U}_1(A_1)$ and the random variables Z , \mathbb{Z}_3 and \mathbb{Z}_4 are independent we obtain from (160)

$$|R_3| \leq t^2 |\alpha^{m/2}(t)| \mathbf{E}|Z\mathbb{Z}_3| \leq t^2 e^{-mt^2/8} m^{1/2} N^{-2} \prec \mathcal{R}.$$

In the last step we combined (141), (145) and Cauchy-Schwartz. The proof of $R_1 \prec \mathcal{R}$ is similar. In order to prove the first relation of (155) we expand the exponent in powers of $it\mathbb{Z}_4$ and obtain $h_{11} = \mathbf{E} \exp\{it\mathbb{U}_1\}itZ + R$. Furthermore, combining (160), (141) and (145) we obtain

$$|R| \leq t^2 |\alpha^m(t)| \mathbf{E}|Z\mathbb{Z}_4| \leq c_* t^2 e^{-mt^2/4N} N^{-3/2} \prec \mathcal{R}.$$

Thus the proof of (148) is complete.

Let us prove (149). By symmetry and the independence,

$$(156) \quad \mathbf{E} e^{it\mathbb{U}_1} it\mathbb{U}_3 = \binom{N}{3} h_{13} \mathbf{E} e^{it\mathbb{U}_*}, \quad h_{13} = \mathbf{E} e^{itx_1} e^{itx_2} e^{itx_3} itz.$$

Here we denote $z = g_3(X_1, X_2, X_3)$ and write,

$$\mathbb{U}_1 = x_1 + x_2 + x_3 + \mathbb{U}_*, \quad \mathbb{U}_* = \sum_{4 \leq j \leq N} g_1(X_j), \quad x_j = g_1(X_j).$$

Furthermore, write

$$r_j = e^{itx_j} - 1 - itx_j, \quad v_j = e^{itx_j} - 1.$$

In what follows we expand the exponents in powers of itX_j , $j = 1, 2, 3$ and use the fact that $\mathbf{E}(g_3(X_1, X_2, X_3) | X_1, X_2) = 0$ as well as the obvious symmetry. Thus, we have

$$\begin{aligned} h_{13} &= h_{14} + R_1, & h_{14} &= \mathbf{E}e^{itx_2}e^{itx_3}(it)^2zx_1, & R_1 &= \mathbf{E}e^{itx_2}e^{itx_3}itzr_1, \\ h_{14} &= h_{15} + R_2, & h_{15} &= \mathbf{E}e^{itx_3}(it)^3zx_1x_2, & R_2 &= \mathbf{E}e^{itx_3}(it)^2zx_1r_2 \\ h_{15} &= h_{16} + R_3, & h_{16} &= \mathbf{E}(it)^4zx_1x_2x_3, & R_3 &= \mathbf{E}(it)^3zx_1x_2r_3. \end{aligned}$$

Furthermore, we have

$$R_1 = \mathbf{E}itz_1r_1v_2v_3, \quad R_2 = \mathbf{E}(it)^2zx_1r_2v_3.$$

Invoking the bounds $|r_j| \leq |tx_j|^2$ and $|v_j| \leq |tx_j|$ we obtain

$$(157) \quad h_{13} = h_{16} + R,$$

where $|R| \leq c|t|^5\mathbf{E}|zx_1x_2|x_3^2$. The bound, $|R| \leq c_*|t|^5N^{-9/2}$ (which follows, by Cauchy-Schwartz) in combination with (156) and (157) implies

$$(158) \quad \mathbf{E}e^{it\mathbb{U}_1}it\mathbb{U}_3 \sim \binom{N}{3}\mathbf{E}e^{it\mathbb{U}_*}(it)^4w.$$

Note that $\binom{N}{3}|w| \leq c_*N^{-1}$. In order to show (149) we replace $\mathbf{E}e^{it\mathbb{U}_*}$ by $e^{-t^2/2}$. Therefore, (149) follows from (158) and the inequalities

$$\frac{(it)^4}{N}(\mathbf{E}e^{it\mathbb{U}_*} - e^{-t^2\sigma^2(N-3)/2N}) \prec \mathcal{R}, \quad \frac{(it)^4}{N}(e^{-t^2\sigma^2(N-3)/2N} - e^{-t^2/2}) \prec \mathcal{R}.$$

The second inequality is a direct consequence of (164). The proof of the first inequality is routine and here omitted. Thus the proof of (135) is complete.

5.2.3. Here we show that

$$(159) \quad \mathbf{E} \exp\{it\mathbb{U}_1 + \mathbb{U}_2\} + \binom{N}{3}e^{-t^2/2}(it)^4w \sim \hat{G}(t).$$

This relation in combination with (134) and (135) implies $\mathbf{E}e^{it\mathbb{T}} \sim \hat{G}(t)$.

Let $G_U(t)$ denote the two term Edgeworth expansion of the U -statistic $\mathbb{U}_1 + \mathbb{U}_2$. That is, $G_U(t)$ is defined by (2), but with κ_4 replaced by κ_4^* , where κ_4^* is obtained from κ_4 after removing the summand $4\mathbf{E}g(X_1)g(X_2)g(X_3)\chi(X_1, X_2, X_3)$. Furthermore, let $\hat{G}_U(t)$ denote the Fourier transform of $G_U(t)$. It easy to show that

$$\hat{G}(t) = \hat{G}_U(t) + \binom{N}{3}e^{-t^2/2}(it)^4w.$$

Therefore, in order to prove (159) it suffices to show that $\hat{G}_U(t) \sim \mathbf{E} \exp\{it(\mathbb{U}_1 + \mathbb{U}_2)\}$. The bound

$$\int_{|t| \leq t_1} |\hat{G}_U(t) - \mathbf{E} \exp\{it(\mathbb{U}_1 + \mathbb{U}_2)\}| \frac{dt}{|t|} \leq \varepsilon_N N^{-1}$$

where $\varepsilon_N \downarrow 0$, was shown by Callaert, Janssen and Veraverbeke (1980) [11] and Bickel, Götze and van Zwet (1986) [8]. An inspection of their proofs shows that under the moment conditions (3) one can replace ε_n by $c_*N^{-\nu}$. This completes the proof of (122).

For the reader convenience we formulate in Lemma 3 a known result on upper bounds for characteristic functions.

Lemma 3. *Assume that (14) holds. There exists a constant c_* depending on D_*, M_*, r, s, ν_1 only such that, for $N > c_*$ and $|t| \leq N^{1/2}/10^3\beta_3$ and $B \subset \Omega_N$, we have*

$$(160) \quad |\alpha(t)| \leq 1 - t^2/4N, \quad \mathbf{E} \exp\{it\mathbb{U}_1(B)\} \leq |\alpha(t)|^{|B|} \leq e^{-|B|t^2/4N}.$$

Here $\alpha(t) = \mathbf{E} \exp\{itg_1(X_1)\}$ and $\mathbb{U}_1(B) = \sum_{j \in B} g_1(X_j)$.

Proof. Let us prove the first inequality of (160). Expanding the exponent, see (183), we obtain

$$\begin{aligned} |\alpha(t)| &\leq |1 - 2^{-1}t^2 \mathbf{E}g_1^2(X_1)| + 6^{-1}|t|^3 \mathbf{E}|g_1(X_1)|^3 \\ &= |1 - \sigma^2 t^2/2N| + \beta_3 \sigma^3 |t|^3 / 6N^{3/2} \end{aligned}$$

Invoking the inequality $1 - 10^{-3} \leq \sigma^2 \leq 1$ which follows from (164) for $N > c_*$, where c_* is sufficiently large, we obtain $|\alpha(t)| \leq 1 - t^2/4N$, for $|t| \leq N^{1/2}/10^3\beta_3$.

The second inequality of (160) follows from the first one via the inequality $1 + x \leq e^x$, for $x \in \mathbb{R}$. \square

6. APPENDIX 1

1.1. Here we compare the moments Δ_m^2 and $\mathbf{E}R_m^2$, where R_m denotes the remainder of the expansion (12),

$$\mathbb{T} = \mathbf{E}\mathbb{T} + \mathbb{U}_1 + \cdots + \mathbb{U}_{m-1} + R_m, \quad R_m := \mathbb{U}_m + \cdots + \mathbb{U}_N.$$

For $k = 1, \dots, N$, write $\Omega_k = \{1, 2, \dots, k\}$ and denote $\sigma_k^2 := \mathbf{E}g_k^2(X_1, \dots, X_k) = \mathbf{E}T_{\Omega_k}^2$. It follows from (12), by the orthogonality property (13), that

$$(161) \quad \sigma_{\mathbb{T}}^2 = \sum_{k=1}^N \mathbf{E}U_k^2, \quad \mathbf{E}R_m^2 = \sum_{k=m}^N \mathbf{E}U_k^2, \quad \mathbf{E}U_k^2 = \binom{N}{k} \sigma_k^2.$$

Lemma 4. *Assume that $\mathbf{E}\mathbb{T}^2 < \infty$. Then*

$$(162) \quad \mathbf{E}R_m^2 \leq N^{-(m-1)} \Delta_m^2,$$

$$(163) \quad \Delta_m^2 \leq N^{2m-1} \sigma_m^2 + N^{-1} \Delta_{m+1}^2,$$

Assume that (3) and (4) hold, then there exists a constant $c_ < \infty$ depending on D_*, M_*, r, s, ν_1 such that*

$$(164) \quad 0 \leq 1 - \sigma^2 \sigma_{\mathbb{T}}^{-2} \leq c_* N^{-1}.$$

Remark. For $m = 3$, the inequality (163) yields $\Delta_3^2 \leq \zeta_2 + N^{-1} \Delta_4^2$.

Proof. Let us prove (162). The identity

$$D_1 \cdots D_m \mathbb{T} = \sum_{A: \Omega_m \subset A \subset \Omega_N} T_A = \sum_{m \leq k \leq N} \mathbb{U}_{k|m},$$

where $\mathbb{U}_{k|m} = \sum_{|A|=k, A \supset \Omega_m} T_A$, implies

$$(165) \quad \mathbf{E}(D_1 \cdots D_m \mathbb{T})^2 = \sum_{m \leq k \leq N} \mathbf{E}U_{k|m}^2, \quad \mathbf{E}U_{k|m}^2 = \sigma_k^2 \binom{N-m}{k-m}.$$

Write

$$(166) \quad \mathbf{E}(D_1 D_2 \cdots D_m \mathbb{T})^2 = \sum_{m \leq k \leq N} \sigma_k^2 \binom{N-m}{k-m} = \sum_{m \leq k \leq N} \sigma_k^2 \binom{N}{k} b_k,$$

where $b_k = [k]_m/[N]_m$ satisfies $b_k \geq b_m \geq m!N^{-m}$. Here we denote $[x]_m = x(x-1)\cdots(x-m+1)$. A comparison of (161) and (166) shows (162)

$$\mathbf{E}R_m^2 \leq N^m \mathbf{E}(D_1 \cdots D_m \mathbb{T})^2 = N^{-(m-1)} \Delta_m^2.$$

Let us prove (163). Write

$$\begin{aligned} \mathbf{E}(D_1 \cdots D_m \mathbb{T})^2 &= \sigma_m^2 + \sum_{m < k \leq N} \sigma_k^2 \binom{N-m}{k-m} \\ &= \sigma_m^2 + \sum_{m < k \leq N} \sigma_k^2 \binom{N-m-1}{k-m-1} \tilde{b}_k, \end{aligned}$$

where $\tilde{b}_k = (N-m)/(k-m) \leq N$. We obtain the inequality

$$\mathbf{E}(D_1 \cdots D_m \mathbb{T})^2 \leq \sigma_m^2 + N \mathbf{E}(D_1 \cdots D_{m+1} \mathbb{T})^2$$

which implies (163).

Let us prove (164). From (161), (162) we have, for $\sigma^2 = N\sigma_1^2$,

$$0 \leq 1 - \frac{\sigma^2}{\sigma_{\mathbb{T}}^2} \leq \binom{N}{2} \frac{\sigma_2^2}{\sigma_{\mathbb{T}}^2} + \binom{N}{3} \frac{\sigma_3^2}{\sigma_{\mathbb{T}}^2} + \frac{1}{N^3} \frac{\Delta_4^2}{\sigma_{\mathbb{T}}^2}.$$

Invoking the bounds, which follow from (3),

$$N^3 \sigma_2^2 = \mathbf{E}\psi^2(X_1, X_2) \leq M_*^{2/r} \sigma_{\mathbb{T}}^2, \quad N^5 \sigma_3^2 = \mathbf{E}\chi^2(X_1, X_2, X_3) \leq M_*^{2/s} \sigma_{\mathbb{T}}^2$$

and using (4) we obtain (164).

1.2. Here we prove moments bounds for various parts of Hoeffding decomposition defined in Section 2.

Lemma 5. *Assume that $\sigma_{\mathbb{T}}^2 = 1$. For $3 \leq m \leq N$ and $s > 2$, we have*

$$(167) \quad \mathbf{E}\Lambda_3^2 \leq \frac{m^3}{N^5} \Delta_3^2, \quad \mathbf{E}\Lambda_2^2 \leq \frac{m^2}{N^4} \Delta_3^2, \quad \mathbf{E}|\Lambda_1|^3 \leq c \frac{m^3}{N^{9/2}} \gamma_3,$$

$$(168) \quad \mathbf{E}|\Lambda_4|^s \leq c(s) m^{s/2} N^{-3s/2} \zeta_s, \quad \mathbf{E}\eta_i^2 \leq N^{-4} \Delta_4^2, \quad \mathbf{E}\Lambda_5^2 \leq m N^{-4} \Delta_4^2.$$

Here c denotes an absolute constant and $c(s)$ denotes a constant which depends only on s .

Proof. The inequalities (167) are proved in [3].

Let us prove (168). Split $\Lambda_4 = z_1 + \cdots + z_m$, where

$$z_i = \sum_{|A|=3, A \cap \Omega_m = i} T_A.$$

Let \mathbf{E}' denote the conditional expectation given X_{m+1}, \dots, X_N . It follows from Rosenthal's inequality that almost surely

$$\mathbf{E}'|\Lambda_4|^s \leq c(s) \sum_{i=1}^m \mathbf{E}'|z_i|^s + c(s) \left(\sum_{i=1}^m \mathbf{E}'z_i^2 \right)^{s/2}.$$

Invoking Hölder's inequality we obtain, by symmetry,

$$(169) \quad \mathbf{E}|\Lambda_4|^s = \mathbf{E}\mathbf{E}'|\Lambda_4|^s \leq c(s) m^{s/2} \mathbf{E}|z_1|^s.$$

Using well known martingale moment inequalities (and their applications to U statistics), see Dharmadhikari, Fabian and Jogdeo (1968) [13], one can show the bound $\mathbf{E}|z_1|^s \leq c(s) N^{-3s/2} \zeta_s$. Invoking this bound in (169) we obtain the first bound of (168).

In order to prove the second bound of (168) write

$$\eta_i = \sum_{k=4}^{N-m+1} U_k^*, \quad U_k^* = \sum_{|A|=k, A \cap \Omega_m = \{i\}} T_A.$$

A simple calculation shows $\mathbf{E}(U_k^*)^2 = \binom{N-m}{k-1} \sigma_k^2$. Therefore, by orthogonality,

$$\begin{aligned} \mathbf{E}\eta_i^2 &= \sum_{k=4}^{N-m+1} \binom{N-m}{k-1} \sigma_k^2 = \sum_{k=4}^{N-m+1} \binom{N-4}{k-4} b_k \sigma_k^2 \\ (170) \quad &\leq N^3 \mathbf{E}(D_1 \cdots D_4 \mathbb{T})^2. \end{aligned}$$

In the last step we invoke (165) and use the bound $b_k \leq N^3$, where $b_k = \binom{N-m}{k-1} \binom{N-4}{k-4}^{-1}$. Clearly, (170) implies $\mathbf{E}\eta_i^2 \leq N^{-4} \Delta_4^2$. Finally, using the fact that η_1, \dots, η_m are uncorrelated we obtain

$$\Lambda_5^2 = \mathbf{E}\eta_1^2 + \cdots + \mathbf{E}\eta_m^2 \leq m N^{-4} \Delta_4^2.$$

thus completing the proof. \square

Before formulating next result we introduce some notation. Given m let \mathcal{D} denote the class of subsets $A \subset \Omega_N$ satisfying $|A| \geq 4$ and $\Omega_m \cap A \neq \emptyset$. Introduce the random variable $\mathbb{H}(m) = \sum_{A \in \mathcal{D}} T_A$. Denote $x_i = 2i - 1$ and $y_i = 2i$. For even integer $m = 2k \leq N$ write

$$\Omega_m = A_k \cup B_k, \quad A_k = \{x_1, \dots, x_k\}, \quad B_k = \{y_1, \dots, y_k\}$$

and put $A_0 = B_0 = \emptyset$. Let $\mathcal{A}(k)$ (respectively $\mathcal{B}(k)$) denote the collection of those $A \in \mathcal{D}$ which satisfy $A \cap A_k = \emptyset$ (respectively $A \cap B_k = \emptyset$). Furthermore, let $\mathcal{C}(k)$ denote the collection of $A \in \mathcal{D}$ such that $A \cap A_k \neq \emptyset$ and $A \cap B_k \neq \emptyset$. Write

$$\mathbb{H}_A(k) = \sum_{A \in \mathcal{A}(k)} T_A, \quad \mathbb{H}_B(k) = \sum_{A \in \mathcal{B}(k)} T_A, \quad \mathbb{H}_C(k) = \sum_{A \in \mathcal{C}(k)} T_A.$$

Lemma 6. *There exists an absolute constant c such that,*

$$(171) \quad \mathbf{E}\mathbb{H}^2(m) \leq c \frac{m}{N^4} \Delta_4^2, \quad \text{for } m = 4, 5, \dots, N.$$

For even integer $m = 2k < N$ we have

$$(172) \quad \mathbf{E}\mathbb{H}_A^2(k) = \mathbf{E}\mathbb{H}_B^2(k) \leq c \frac{k}{N^4} \Delta_4^2, \quad \mathbf{E}\mathbb{H}_C^2(k) \leq c \frac{k^2}{N^5} \Delta_4^2.$$

Proof. Let us prove the first bound of (171). For $m = 4$ we have

$$\mathbb{H}(4) = H_1 + H_2 + H_3 + H_4, \quad H_k = \sum_{|A| \geq 4, |A \cap \Omega_4| = k} T_A.$$

A calculation shows that, for $k = 1, 2, 3, 4$,

$$\mathbf{E}H_k^2 = \binom{4}{k} \sum_{j=4}^N \sigma_j^2 \binom{N-4}{j-k} = \binom{4}{k} \sum_{j=4}^N \sigma_j^2 \binom{N-4}{j-4} a_k(j),$$

where the numbers

$$a_k(j) = \frac{\binom{N-4}{j-k}}{\binom{N-4}{j-4}} \leq N^{4-k}.$$

Invoking (166) we obtain

$$(173) \quad \mathbf{E}H_k^2 \leq c N^{4-k} \mathbf{E}(D_1 \cdots D_4 \mathbb{T})^2 = c N^{-3-k} \Delta_4^2.$$

Finally, we obtain (171) for $m = 4$

$$\mathbf{E}\mathbb{H}^2(4) = \mathbf{E}H_1^2 + \cdots + \mathbf{E}H_4^2 \leq c N^{-4} \Delta_4^2.$$

In order to prove (171) for $m = 5, 6, \dots$ we apply a recursive argument. Write

$$(174) \quad \mathbf{E}\mathbb{H}^2(m+1) = \mathbf{E}\mathbb{H}^2(m) + \mathbf{E}d_m^2,$$

where $d_m = \mathbb{H}(m+1) - \mathbb{H}(m)$ is the sum of those T_A with $|A| \geq 4$ satisfying $A \cap \Omega_m = \emptyset$ and $A \cap \Omega_{m+1} \neq \emptyset$. In particular, we have

$$d_m = \sum_{|A| \geq 3, A \cap \Omega_{m+1} = \emptyset} T_{A \cup \{m+1\}}.$$

Therefore,

$$\mathbf{E}d_m^2 = \sum_{j=4}^N \sigma_j^2 \binom{N-m-1}{j-1} = \sum_{j=4}^N \sigma_j^2 \binom{N-4}{j-4} c_j,$$

where the numbers

$$c_j = \frac{\binom{N-m-1}{j-1}}{\binom{N-4}{j-4}} \leq N^3.$$

Invoking (166) we obtain $\mathbf{E}d_m^2 \leq N^{-4} \Delta_4^2$. This bound together with (174) implies (171).

Let us prove (172). Note that for $m = 2k$ we have $\mathbb{H}(m) = \mathbb{H}_A(k) + \mathbb{H}_B(k) + \mathbb{H}_C(k)$ and the summands are uncorrelated. Therefore, the first bound of (172) follows from (171).

Let us show the second inequality of (172). For $k = 2$ we have $\mathcal{C}(2) \subset \mathcal{C}$, where \mathcal{C} denotes the class of subsets $A \subset \Omega_N$ such that $|A| \geq 4$ and $|A \cap \Omega_4| \geq 2$. Write $\mathbb{H}_C = \sum_{A \in \mathcal{C}} T_A$. We have

$$\mathbf{E}\mathbb{H}_C^2(2) \leq \mathbf{E}\mathbb{H}_C^2 = \mathbf{E}H_2^2 + \mathbf{E}H_3^2 + \mathbf{E}H_4^2 \leq cN^{-5} \Delta_4^2.$$

In the last step we applied (173). We obtain (172), for $k = 2$.

In order to prove the bound (172), for $k = 3, 4, \dots$, we apply a recursive argument similar to that used in the proof of (171). Denote

$$d_{[k]} = \mathbb{H}_C(k+1) - \mathbb{H}_C(k) = \sum_{A \in \mathcal{C}(k+1) \setminus \mathcal{C}(k)} T_A.$$

We shall show that

$$(175) \quad \mathbf{E}d_{[k]}^2 \leq ckN^{-5} \Delta_4^2.$$

This bound in combination with the identity $\mathbf{E}\mathbb{H}_C^2(k+1) = \mathbf{E}\mathbb{H}_C^2(k) + \mathbf{E}d_{[k]}^2$ shows (172) for arbitrary k .

In order to show (175) split the set $\mathcal{C}(k+1) \setminus \mathcal{C}(k)$ into $2k+1$ non-intersecting parts

$$\mathcal{C}(k+1) \setminus \mathcal{C}(k) = (\cup_{i=1}^k \mathcal{C}_{x,i}) \cup (\cup_{i=1}^k \mathcal{C}_{y,i}) \cup \mathcal{C}_{x,y},$$

where we denote

$$\mathcal{C}_{x,y} = \{A = \tilde{A} \cup \{x_{k+1}, y_{k+1}\} : \tilde{A} \cap (B_k \cup A_k) = \emptyset, |\tilde{A}| \geq 2\},$$

$$\mathcal{C}_{x,i} = \{A = \tilde{A} \cup \{y_{k+1}, x_i\} : \tilde{A} \cap (B_k \cup A_{i-1}) = \emptyset, |\tilde{A}| \geq 2\},$$

$$\mathcal{C}_{y,i} = \{A = \tilde{A} \cup \{x_{k+1}, y_i\} : \tilde{A} \cap (B_{i-1} \cup A_k) = \emptyset, |\tilde{A}| \geq 2\}.$$

By the orthogonality property ($\mathbf{E}T_A T_V = 0$ for $A \neq V$), the random variables

$$d_{x,i} = \sum_{A \in \mathcal{C}_{x,i}} T_A, \quad d_{y,i} = \sum_{A \in \mathcal{C}_{y,i}} T_A, \quad d_{x,y} = \sum_{A \in \mathcal{C}_{x,y}} T_A$$

are uncorrelated. Therefore, we have

$$(176) \quad \mathbf{E}d_{[k]}^2 = \mathbf{E}d_{x,y}^2 + \sum_{i=1}^k (\mathbf{E}d_{x,i}^2 + \mathbf{E}d_{y,i}^2).$$

A calculation shows that

$$\mathbf{E}d_{x,y}^2 = \sum_{j=4}^N \sigma_j^2 \binom{N-2k-2}{j-2} = \sum_{j=4}^N \sigma_j^2 \binom{N-4}{j-4} v_j,$$

where the coefficients

$$v_j = \frac{\binom{N-2k-2}{j-2}}{\binom{N-4}{j-4}} \leq N^2.$$

Invoking (166) we obtain $\mathbf{E}d_{x,y}^2 \leq N^{-5}\Delta_4^2$. The same argument shows $\mathbf{E}d_{x,i}^2 = \mathbf{E}d_{y,i}^2 \leq N^{-5}\Delta_4^2$. The latter bound in combination with (176) shows (175). The lemma is proved. \square

7. APPENDIX 2

Here we construct bounds for the probability density function (and its derivatives) of random variables $g_k^* = (N/M)^{1/2}g_k$, for $1 \leq k \leq n-1$, where g_k are defined in (69). Since these random variables are identically distributed it suffices to consider

$$g_1^* = \left(\frac{N}{M}\right)^{1/2}g_1 = \frac{1}{\sqrt{M}} \sum_{j=m+1}^{m+M} g(Y_j) + \frac{\xi_1}{R}.$$

Here $R = \sqrt{nMN}$. Introduce the random variables

$$g_2^* = g_1^* - M^{-1/2}g(Y_{m+1}), \quad g_3^* = g_1^* - M^{-1/2}(g(Y_{m+1}) + g(Y_{m+2})).$$

Let $p_i(\cdot)$ denote the probability density function of g_i^* , for $i = 1, 2, 3$. Recall that the integers $n \approx N^{50\nu} \leq N^{\nu_2/10}$ and $M \approx N/n \geq N^{9/10}$ are introduced in (24) and the number $\nu > 0$ is defined by (15).

Lemma 7. *Assume that conditions of Theorem 1 are satisfied. There exist positive constants C_*, c_*, c'_* depending only on M_*, D_*, δ, r and ν_1, ν such that, for $i = 1, 2, 3$, we have uniformly in $u \in \mathbb{R}$ and $N > C_*$*

$$(177) \quad |p_i(u)| \leq c_*, \quad |p'_i(u)| \leq c_*, \quad |p''_i(u)| \leq c_*, \quad |p'''_i(u)| \leq c_*.$$

Furthermore, given $w > 0$ there exists a constant $C_*(w)$ depending on $M_*, D_*, \delta, r, \nu_1, \nu$ and w such that uniformly in $z_* \in [-2w, 2w]$ and $N > C_*(w)$ we have

$$(178) \quad p_i(z_*) \geq c'_*, \quad i = 1, 2, 3.$$

Proof. We shall prove (177) and (178) for $i = 1$. For $i = 2, 3$, the proof is almost the same. Before starting the proof we introduce some notation and collect auxiliary results.

Denote

$$\begin{aligned} \theta &= \mathbf{E}g_1^* = M^{1/2}\theta_1, & \theta_1 &= \mathbf{E}g(Y_{m+1}), \\ s^2 &= \mathbf{E}(g(Y_{m+1}) - \theta_1)^2, & \tilde{\beta}_3 &= s^{-3}\mathbf{E}|g(Y_{m+1}) - \theta_1|^3. \end{aligned}$$

It follows from $\mathbf{E}g(X_{m+1}) = 0$ that

$$\theta_1 = q_N^{-1}\mathbf{E}g(X_{m+1})\mathbb{I}_{A_{m+1}} = -q_N^{-1}\mathbf{E}g(X_{m+1})(1 - \mathbb{I}_{A_{m+1}}).$$

Therefore, by Chebyshev's inequality, for $\alpha = 3/(r+2)$ we have

$$(179) \quad |\theta_1| \leq q_N^{-1}N^{-\alpha(r-1)}\mathbf{E}|g(X_{m+1})| \|Z'_{m+1}\|_r^{r-1} \leq c_*N^{-3/2}.$$

In the last step we invoke the inequalities $\alpha(r-1) \geq 1 + (r-1)/(r+2) \geq 3/2$ and $q_N^{-1} \leq c_*$, see (40), and $\mathbf{E}|g(X_{m+1})| \|Z'_{m+1}\|_r^{r-1} \leq M_*$, where the latter inequality follows from (3) by Hölder inequality.

Similarly, the identities

$$s^2 = q_N^{-1} \mathbf{E}g^2(X_{m+1})\mathbb{I}_{A_{m+1}} - \theta_1^2 = q_N^{-1} \sigma^2 - q_N^{-1} \mathbf{E}g^2(X_{m+1})(1 - \mathbb{I}_{A_{m+1}}) - \theta_1^2$$

in combination with (39) and the inequalities

$$\mathbf{E}g^2(X_{m+1})(1 - \mathbb{I}_{A_{m+1}}) \leq N^{-\alpha(r-2)} \mathbf{E}g^2(X_{m+1}) \|Z'_{m+1}\|_r^{r-2} \leq N^{-\alpha(r-2)} M_*$$

and $\alpha(r-2) = 1 + 2(r-4)/(r-2) \geq 1$ yield

$$(180) \quad |s^2 - \sigma^2| \leq c_* N^{-1}.$$

Introduce the random variables

$$g_* = S + \frac{\xi_1}{sR}, \quad S = w_1 + \dots + w_M, \quad w_j = \frac{g(Y_{m+j}) - \theta_1}{M^{1/2}s}.$$

We have $g_* = s^{-1}(g_1^* - \theta)$. Let $p(\cdot)$ denote the density function of g_* . Note that $p_1(u) = s^{-1}p(s^{-1}(u - \theta))$. Furthermore, we have, by (179), $|\theta| \leq c_* N^{-1}$ and, by (180), (164), $|s^2 - 1| \leq c_* N^{-1}$. Therefore, it suffices to prove (177) and (178) for $p(\cdot)$ (the latter inequality we verify for every $z_* \in [-3w, 3w]$).

In order to prove (177) and (178) we approximate the characteristic function $\hat{p}(t) = \mathbf{E}e^{itg_*}$ by $e^{-t^2/2}$ and then apply a Fourier inversion formula. Write

$$\hat{p}(t) = \mathbf{E}e^{itg_*} = \gamma^M(t) \tau\left(\frac{t}{sR}\right), \quad \gamma(t) := \mathbf{E}e^{itw_1}, \quad \tau(t) := \mathbf{E}e^{it\xi_1}.$$

The fact that $\tau(t) = 0$, for $|t| \geq 1$, implies $\hat{p}(t) = 0$, for $|t| > sR$. Therefore, we obtain from the Fourier inversion formula,

$$p(x) = \frac{1}{2\pi} \int_{-\infty}^{+\infty} e^{-itx} \hat{p}(t) dt = \frac{1}{2\pi} \int_{-sR}^{sR} e^{-itx} \hat{p}(t) dt.$$

Write $\hat{p}(t) - e^{-t^2/2} = r_1(t) + r_2(t)$, where

$$r_1(t) = (\gamma^M(t) - e^{-t^2/2})\tau(t/sR), \quad r_2(t) = e^{-t^2/2}(\tau(t/sR) - 1).$$

We shall show below that

$$(181) \quad \int_{|t| \leq sR} |r_i(t)| dt \leq c_* M^{-1/2}, \quad i = 1, 2.$$

These bounds in combination with the simple inequality

$$\int_{|t| \geq sR} e^{-t^2/2} dt \leq c_* M^{-1/2}$$

show that

$$(182) \quad |p(x) - \varphi(x)| \leq c_* M^{-1/2}, \quad x \in \mathbb{R}.$$

Here φ denotes the standard normal density function

$$\varphi(x) = \frac{1}{\sqrt{2\pi}} e^{-x^2/2} = \frac{1}{2\pi} \int_{-\infty}^{+\infty} e^{-itx} e^{-t^2/2} dt$$

It follows from (182) that

$$|p(x)| \leq c_*, \quad x \in \mathbb{R}.$$

Furthermore, given w we have uniformly in $|z_*| \leq 3w$

$$|p(z_*)| \geq \varphi(3w) - c_* M^{-1/2} \geq c'_* > 0,$$

for sufficiently large M (for $N > C_*(w)$).

In order to prove an upper bounds for the k -th derivative, $|p^{(k)}(x)| \leq c_*$, write

$$p^{(k)}(t) = \frac{1}{2\pi} \int_{-\infty}^{+\infty} (-it)^k \exp\{-itx\} \hat{p}(t) dt, \quad k = 1, 2, 3,$$

and replace $\hat{p}(t)$ by $e^{-t^2/2}$ as in the proof of (182). We obtain

$$p^{(k)}(x) = \frac{1}{2\pi} \int_{-\infty}^{+\infty} (-it)^k \exp\{-itx\} e^{-t^2/2}(t) dt + r, \quad |r| \leq c_* M^{-1/2}.$$

This implies $|p^{(k)}(x) - \varphi^{(k)}(x)| \leq c_* M^{-1/2}$. We arrive at the desired bound $|p^{(k)}(x)| \leq c_*$, for $k = 1, 2, 3$.

In the remaining part of the proof we verify (181). For $i = 2$ this bound follows from $|\tau(t/sR) - 1| \leq ct^2/(sR)^2$. The latter inequality is a consequence of the short expansion

$$|\mathbf{E} \exp\{it\xi_1/sR\} - 1 - \mathbf{E} it\xi_1/sR| \leq \mathbf{E}(t\xi_1)^2/2(sR)^2$$

and $\mathbf{E}\xi_1 = 0$ and $\mathbf{E}\xi_1^2 \leq c$, for some absolute constant c .

Let us prove (181) for $i = 1$. Introduce the sequence of i.i.d. centered Gaussian random variables η_1, η_2, \dots with variances $\mathbf{E}\eta_i^2 = M^{-1}$. Denote

$$f(t) = \mathbf{E} e^{it\eta_1} = e^{-t^2/(2M)} \quad \text{and} \quad \delta(t) = \gamma(t) - f(t).$$

We are going to apply the well known inequality

$$(183) \quad \left| e^{iv} - \left(1 + \frac{iv}{1!} + \frac{(iv)^2}{2!} + \dots + \frac{(iv)^{k-1}}{(k-1)!} \right) \right| \leq \frac{|v|^k}{k!}.$$

It follows from (183) and identities $\mathbf{E}\eta_i^i = \mathbf{E}w_i^i$, $i = 1, 2$, that

$$(184) \quad |\delta(t)| \leq \frac{|t|^3}{3!} (\mathbf{E}|w_1|^3 + \mathbf{E}|\eta_1|^3) \leq c|t|^3 \mathbf{E}|w_1|^3.$$

Here we use the inequality $\mathbf{E}|\eta_1|^3 \leq c\mathbf{E}|w_1|^3$, which follows from $\mathbf{E}\eta_1^2 = \mathbf{E}w_1^2$. Combining (184) and the simple identity

$$\gamma^M(t) - f^M(t) = \delta(t) \sum_{k=1}^M \gamma^{M-k}(t) f^{k-1}(t)$$

we obtain

$$(185) \quad |\gamma^M(t) - f^M(t)| \leq c|t|^3 Z(t) M^{-1/2} \tilde{\beta}_3.$$

Here we denote

$$Z(t) = \max_{r+v=M-1} |f^r(t) \gamma^v(t)|.$$

We shall show below that

$$(186) \quad Z(t) \leq \exp\left\{-\frac{t^2}{3} \frac{M-1}{M}\right\} + \exp\{-\delta''(M-1)/2\}, \quad 0 \leq |t| \leq sR,$$

where $\delta'' > 0$ depends on $\delta, A_*, D_*, M_*, \nu_1$ and it is given in (32). This inequality in combination with (185) proves (181).

Let us prove (186). Clearly, $Z \leq |f^{M-1}(t)| + |\gamma^{M-1}(t)|$. Furthermore, $f^M(t) = e^{-t^2/2}$. In order to prove (186) we shall show

$$(187) \quad |\gamma^M(t)| \leq e^{-t^2/3}, \quad 0 \leq |t| \leq M^{1/2}/\tilde{\beta}_3,$$

$$(188) \quad |\gamma(t)| \leq e^{-\delta''/2}, \quad M^{1/2}/\tilde{\beta}_3 \leq |t| \leq sR.$$

To show (187) we expand e^{itw_1} using (183),

$$\begin{aligned} |\gamma(t)| &= |\mathbf{E}e^{itw_1}| \leq \left| 1 - \frac{t^2}{2}\mathbf{E}w_1^2 \right| + \frac{|t|^3}{3!}\mathbf{E}|w_1|^3 \\ &= \left| 1 - \frac{t^2}{2M} \right| + \frac{|t|^3}{3!} \frac{\tilde{\beta}_3}{M^{3/2}} \\ &= 1 - \frac{t^2}{2M} \left(1 - \frac{|t|}{3} \frac{\tilde{\beta}_3}{\sqrt{M}} \right) \\ &\leq 1 - \frac{t^2}{3M}. \end{aligned}$$

Here we used the identity $|1 - t^2/2M| = 1 - t^2/2M$, which holds for $|t| < M^{1/2}/\tilde{\beta}_3$, since $\tilde{\beta}_3 \geq 1$. Finally, an application of the inequality $1 - x \leq e^{-x}$ to $x = t^2/3M > 0$ completes the proof of (187).

Let us prove (188). For δ'' defined by (32) we shall show $\delta'' \leq 2\tilde{\delta}$, where

$$\begin{aligned} \tilde{\delta} &= 1 - \sup\{|\gamma(t)| : M^{1/2}\tilde{\beta}_3^{-1} \leq |t| \leq sR\} \\ &= 1 - \sup\{|\mathbf{E}\exp\{iu\sigma^{-1}g(Y_{m+1})\}| : \sigma/s\tilde{\beta}_3 \leq |u| \leq \sigma\sqrt{nN}\}. \end{aligned}$$

We are going to replace $g(Y_{m+1})$, $\tilde{\beta}_3$, s^2 by X_{m+1} , β_3 , σ^2 respectively. Write

$$\begin{aligned} \mathbf{E}e^{ivg(Y_{m+1})} &= q_N^{-1}\mathbf{E}e^{ivg(X_{m+1})}\mathbb{I}_{A_{m+1}} = \mathbf{E}e^{ivg(X_{m+1})} + r_1 + r_2, \\ r_1 &= q_N^{-1}\mathbf{E}e^{ivg(X_{m+1})}(\mathbb{I}_{A_{m+1}} - 1), \quad r_2 = (q_N^{-1} - 1)\mathbf{E}e^{ivg(X_{m+1})}. \end{aligned}$$

It follows from (5.A5) that, for every $v \in \mathbb{R}$,

$$|r_1| \leq q_N^{-1}\mathbf{E}|\mathbb{I}_{A_{m+1}} - 1| = q_N^{-1} - 1 \leq c_*N^{-2}, \quad |r_2| \leq q_N^{-1} - 1 \leq c_*N^{-2}.$$

These bounds imply

$$(189) \quad |\mathbf{E}e^{ivg(Y_{m+1})} - \mathbf{E}e^{ivg(X_{m+1})}| \leq c_*N^{-2}, \quad \text{for every } v \in \mathbb{R}.$$

One can show that, for sufficiently large N (i.e., for $N > C_*$), we have

$$(190) \quad |\tilde{\beta}_3/\beta_3 - 1| < 1/5, \quad |s^2/\sigma^2 - 1| < 1/5, \quad |s^2 - 1| \leq 1/5.$$

Using (189), (190) we get, for $N > C_*$,

$$\begin{aligned} \tilde{\delta} &\geq 1 - \sup\{|\mathbf{E}e^{iu\sigma^{-1}g(Y_{m+1})}| : (2\beta_3)^{-1} \leq |u| \leq N^{(1+50\nu)/2}\} \\ &\geq 1 - \sup\{|\mathbf{E}e^{iu\sigma^{-1}g(X_{m+1})}| : (2\beta_3)^{-1} \leq |u| \leq N^{(\nu_2+1)/2}\} - c_*N^{-2} \\ &\geq \delta''/2. \end{aligned}$$

We obtain $|\gamma(t)| \leq 1 - \tilde{\delta} \leq 1 - \delta''/2$ and, therefore, $|\gamma(t)| \leq e^{-\delta''/2}$.

8. APPENDIX 3

The main results of this section are moment inequalities of Lemma 9 and corresponding inequalities for conditional moments of Lemma 10. Lemma 8 provides an auxiliary inequality.

We start with some notation. We call $v = v(\cdot), u = u(\cdot) \in L^r$ orthogonal if $\langle u, v \rangle = 0$, where

$$\langle u, v \rangle = \int_{\mathcal{X}} u(x)v(x)P_X(dx) = \mathbf{E}u(X_1)v(X_1).$$

Given $f \in L^2(P_X)$ we have for the kernel ψ^{**} defined in (37)

$$\mathbf{E}\psi^{**}(X_1, X_2)(f(X_1)g(X_2) + f(X_2)g(X_1)) = 0$$

and almost surely

$$(191) \quad \mathbf{E}(\psi^{**}(X_1, X_2)|X_1) = 0,$$

$$(192) \quad \mathbf{E}(\psi^{**}(X_1, X_2)g(X_1)|X_2) = 0.$$

The latter identity says that almost all values of the L^r -valued random variable $\psi^{**}(\cdot, X_2)$ are orthogonal to the vector $g(\cdot) \in L^r$.

Let $p_g : L^r \rightarrow L^r$ denote the projection on the subspace of elements $u \in L^r$ which are orthogonal to $g = g(\cdot)$. For $v \in L^r$, write $v^* = p_g(v)$. It follows from (192) that

$$(193) \quad \psi^*(\cdot, Y_j) \left(= p_g(\psi(\cdot, Y_j)) \right) = \psi^{**}(\cdot, Y_j) + g(Y_j)b^*(\cdot),$$

where $b^*(\cdot) = p_g(b(\cdot)) = \sigma^{-2}p_g(\mathbf{E}(\psi(\cdot, X_1)g(X_1)))$. Denote

$$U_k^* (= p_g(U_k)) = \frac{1}{\sqrt{N}} \sum_{j \in O_k} \psi^*(\cdot, Y_j), \quad U_k^{**} = N^{-1/2} \sum_{j \in O_k} \psi^{**}(\cdot, Y_j),$$

where the L^r -valued random variables U_k are introduced in (86). For the random variables g_k and L_k introduced in (67) and (69), we have

$$(194) \quad U_k^* = U_k^{**} + L_k b^*(\cdot) = U_k^{**} + \left(g_k - \frac{1}{\sqrt{n}} \frac{\xi_k}{N} \right) b^*(\cdot).$$

Denote $K = \mathbf{E}|\psi(X_1, X_2)|^r$ and $K_s = \mathbf{E}|\psi^{**}(X_1, X_2)|^s$, $s \leq r$.

Lemma 8. *Let $4 < r \leq 5$. For $s \leq r$, we have*

$$(195) \quad K_s^{r/s} \leq K_r \leq cK \left(1 + \frac{\mathbf{E}|g(X_1)|^r}{\sigma^r} \right)^2.$$

Proof. The first inequality of (195) is a consequence of Lyapunov's inequality. Let us prove the second inequality. The inequality $|a + b + c|^r \leq 3^r(|a|^r + |b|^r + |c|^r)$ implies

$$K_r = \mathbf{E}|\psi^{**}(X_1, X_2)|^r \leq 3^r (K + 2\mathbf{E}|b(X_1)|^r \mathbf{E}|g(X_2)|^r).$$

Therefore, (195) is a consequence of the inequalities

$$\begin{aligned} \mathbf{E}|b(X_1)|^r &\leq \frac{2^r}{\sigma^r} K + \frac{|\kappa|^r}{\sigma^{4r}} \mathbf{E}|g(X_1)|^r, \\ \kappa^2 &\leq \sigma^4 \mathbf{E}\psi^2(X_1, X_2) \leq \sigma^4 K^{2/r}. \end{aligned}$$

Here $\kappa = \mathbf{E}\psi(X_1, X_2)g(X_1)g(X_2)$. To prove the first inequality use $|a + b|^r \leq 2^r(|a|^r + |b|^r)$ to get

$$\mathbf{E}|b(X_1)|^r \leq \frac{2^r}{\sigma^{2r}} \mathbf{E}|\mathbf{E}(\psi(X_1, X_2)g(X_2)|X_1)|^r + \frac{\kappa^r}{\sigma^{4r}} \mathbf{E}|g(X_1)|^r.$$

Furthermore, by Cauchy–Schwartz,

$$|\mathbf{E}(\psi(X_1, X_2)g(X_2)|X_1)| \leq (\mathbf{E}(\psi^2(X_1, X_2)|X_1))^{1/2} \sigma.$$

Finally, Lyapunov's inequality implies

$$(\mathbf{E}(\psi^2(X_1, X_2)|X_1))^{r/2} \leq \mathbf{E}(|\psi(X_1, X_2)|^r | X_1).$$

We obtain $\mathbf{E}|\mathbf{E}(\psi(X_1, X_2)g(X_2)|X_1)|^r \leq K\sigma^r$ thus completing the proof. \square

Lemma 9. *Let $1 \leq k \leq n - 1$. For \bar{U}_k^* , an independent copy of U_k^* , we have*

$$(196) \quad 2\delta_3^2 - \frac{c_*}{N^{\alpha(r-4)}} \leq \frac{N}{M} \mathbf{E}\|U_k^* - \bar{U}_k^*\|_2^2 \leq 2\delta_3^2 + c_*,$$

$$(197) \quad \mathbf{E}\|U_k - \bar{U}_k\|_r^r \leq c_* \left(\frac{M}{N} \right)^{r/2}.$$

Recall that $\delta_3^2 = \mathbf{E}|\psi^{**}(X_1, X_2)|^2$.

Proof. Let us prove (196). By symmetry, we have, for $i, j \in O_1$,

$$\begin{aligned} \mathbf{E}\|U_1^* - \bar{U}_1^*\|_2^2 &= 2\frac{M}{N}H_1 - 2\frac{M}{N}H_2, \\ H_1 &:= \mathbf{E}\|\psi^*(\cdot, Y_j)\|_2^2, \quad H_2 := \mathbf{E}\langle \psi^*(\cdot, Y_j), \psi^*(\cdot, Y_i) \rangle, \quad i \neq j. \end{aligned}$$

The inequality (196) follows from the inequalities

$$(198) \quad \delta_3^2 - c_*N^{-\alpha(r-4)} \leq H_1 \leq \delta_3^2 + c_*,$$

$$(199) \quad H_2 \leq c_*N^{-\alpha(r-2)}.$$

Let us prove (198). From (193) we have $H_1 = V_1 + V_2 + 2V_3$, where

$$V_1 = \mathbf{E}\|\psi^{**}(\cdot, Y_j)\|_2^2, \quad V_2 = \|b^*(\cdot)\|_2^2 \mathbf{E}g^2(Y_j), \quad V_3 = \mathbf{E}g(Y_j)\langle \psi^{**}(\cdot, Y_j), b^*(\cdot) \rangle.$$

Let us show that

$$(200) \quad \delta_3^2 - c_*N^{-\alpha(r-2)} \leq V_1 \leq \delta_3^2 + c_*N^{-\alpha r}.$$

This inequality follows from (39), (40) and the identity

$$V_1 = q_N^{-1} \mathbf{E}|\psi^{**}(X_1, X_j)|^2 \mathbb{I}_{A_j} = q_N^{-1} \mathbf{E}|\psi^{**}(X_1, X_j)|^2 - q_N^{-1} V_1',$$

where $V_1' = \mathbf{E}|\psi^{**}(X_1, X_j)|^2(1 - \mathbb{I}_{A_j})$ satisfies, by (38),

$$(201) \quad 0 \leq V_1' \leq N^{-\alpha(r-2)} \mathbf{E}|\psi^{**}(X_1, X_j)|^2 \|Z_j'\|_r^{r-2} \leq c_*N^{-\alpha(r-2)}.$$

In the last step we applied Hölder's inequality and Lemma 8 to get

$$\mathbf{E}|\psi^{**}(X_1, X_j)|^2 \|Z_j'\|_r^{r-2} \leq K_r^{2/r} (\mathbf{E}\|Z_j'\|_r^r)^{(r-2)/r} \leq K_r^{2/r} K^{(r-2)/r} \leq c_*.$$

Let us show that

$$(202) \quad 0 \leq V_2 \leq c_*.$$

For $\tilde{b}(\cdot) := \mathbf{E}\psi(\cdot, X_1)g(X_1)$ we have, by Cauchy-Schwartz,

$$\|\tilde{b}(\cdot)\|_2^2 = \mathbf{E}(\mathbf{E}(\psi(X, X_1)g(X_1) | X))^2 \leq \mathbf{E}\psi^2(X, X_1) \sigma^2 \leq c_*\sigma^2.$$

Now the identity $b^* = \sigma^{-2}p_g(\tilde{b})$ implies

$$(203) \quad \|b^*\|_2^2 \leq \sigma^{-4} \|\tilde{b}\|_2^2 \leq \sigma^{-2} c_*.$$

Invoking the bound $\mathbf{E}g^2(Y_j) \leq c_*\sigma^2$, see (42), we obtain (202).

Finally, write

$$V_3 = q_N^{-1} \mathbf{E}\tilde{V}\mathbb{I}_{A_j}, \quad \tilde{V} = g(X_j)\psi^{**}(X_1, X_j)b^*(X_1).$$

The identity (192) implies $\mathbf{E}\tilde{V} = 0$. Therefore $V_3 = q_N^{-1} \mathbf{E}\tilde{V}(\mathbb{I}_{A_j} - 1)$. Invoking (38) and using $q_N^{-1} \leq c_*$, see (40), we obtain

$$(204) \quad |V_3| \leq c_*N^{-\alpha(r-4)} \mathbf{E}|\tilde{V}| \|Z_j'\|_r^{r-4} \leq c_*N^{-\alpha(r-4)}.$$

In the last step we used the bound $\mathbf{E}|\tilde{V}| \|Z_j'\|_r^{r-4} \leq c_*$. In order to prove this bound we invoke the inequalities

$$|abc| \leq (ab)^2 + c^2 \leq a^4 + b^4 + c^2$$

to show that

$$|\tilde{V}| \leq |g(X_j)|^4 + |\psi^{**}(X_1, X_j)|^4 + |b^*(X_1)|^2.$$

Furthermore, by Hölder's inequality and (195),

$$\mathbf{E}|g(X_j)|^4 \|Z_j'\|_r^{r-4} \leq c_*, \quad \mathbf{E}|\psi^{**}(X_1, X_j)|^4 \|Z_j'\|_r^{r-4} \leq c_*.$$

By the independence and (203),

$$\mathbf{E}|b^*(X_1)|^2 \|Z'_j\|_r^{r-4} = \|b^*\|_2^2 \mathbf{E}\|Z'_j\|_r^{r-2} \leq c_*.$$

Thus we arrive at (204). Combining (200), (202) and (204) we obtain (198).

Let us prove (199). Using (193) write $H_2 = Q_1 + Q_2 + 2Q_3$, where

$$\begin{aligned} Q_1 &= \mathbf{E}\psi^{**}(X_1, Y_j)\psi^{**}(X_1, Y_i), & Q_2 &= \|b^*\|_2^2 \mathbf{E}g(Y_j)g(Y_i), \\ Q_3 &= \mathbf{E}\psi^{**}(X_1, Y_j)g(Y_i)b^*(X_1). \end{aligned}$$

It follows from the identity (191) that

$$Q_1 = q_N^{-2} \mathbf{E}\psi^{**}(X_1, X_j)\psi^{**}(X_1, X_i)(\mathbb{I}_{A_j} - 1)(\mathbb{I}_{A_i} - 1).$$

The simple inequality $|\psi^{**}(X_1, X_j)\psi^{**}(X_1, X_i)| \leq |\psi^{**}(X_1, X_j)|^2 + |\psi^{**}(X_1, X_i)|^2$ yields, by symmetry,

$$(205) \quad |Q_1| \leq 2q_N^{-2} \mathbf{E}|\psi^{**}(X_1, X_j)|^2(1 - \mathbb{I}_{A_j}) \leq c_* N^{-\alpha(r-2)}.$$

In the last step we applied (201) and $q_N^{-1} \leq c_*$, see (39).

Furthermore, using the identity $\mathbf{E}g(X_i) = 0$ we obtain from (38)

$$(206) \quad \begin{aligned} |\mathbf{E}g(Y_i)| &= q_N^{-1} |\mathbf{E}g(X_i)(\mathbb{I}_{A_i} - 1)| \\ &\leq q_N^{-1} N^{-\alpha(r-1)} \mathbf{E}|g(X_i)| \|Z_i\|_r^{r-1} \leq c_* N^{-\alpha(r-1)}. \end{aligned}$$

In the last step we applied Hölder's inequality to show $\mathbf{E}|g(X_i)| \|Z_i\|_r^{r-1} \leq c_*$.

The bounds (206), (39) and (203) together imply

$$(207) \quad |Q_k| \leq c_* N^{-\alpha(r-1)}, \quad k = 2, 3.$$

The bound (199) follows from (205) and (207).

Let us prove (197). For this purpose we shall show that

$$(208) \quad \mathbf{E}\left\| \sum_{j \in O_k} \frac{V_j}{\sqrt{M}} \right\|_r^r \leq c_*, \quad \text{where} \quad V_j = \psi(\cdot, Y_j) - \psi(\cdot, \bar{Y}_j),$$

and where \bar{Y}_j denote independent copies of Y_j , $j \in O_k$. Using

$$\mathbf{E}\|\psi(\cdot, X_j)\|_r^r = \mathbf{E}|\psi(X_1, X_j)|^r \leq c_*$$

we obtain, by symmetry and (42),

$$\mathbf{E}\|V_j\|_r^r \leq 2^r \mathbf{E}\|\psi(\cdot, Y_j)\|_r^r \leq c_* \mathbf{E}\|\psi(\cdot, X_j)\|_r^r \leq c_*.$$

Now (208) follows from the well known inequality

$$(209) \quad \|\xi_1 + \dots + \xi_k\|_r^r \leq c(r) \sum_{i=1}^k \mathbf{E}\|\xi_i\|_r^r + c(r) \left(\sum_{i=1}^k \mathbf{E}\|\xi_i\|_r^2 \right)^{r/2}, \quad k = 1, 2, \dots$$

which is valid for independent centered random elements ξ_i with values in L^r . One can derive this inequality from Hoffmann – Jorgensen's inequality (see e.g., Proposition 6.8 in Ledoux and Talagrand (1991) [21]) using the type 2 property of the Banach space L^r and the symmetrization lemma (see formula (9.8) and Lemma 6.3 ibidem). The proof of the lemma is complete. \square

Before formulating and proving Lemma 10 we introduce some more notation. Let $\mathcal{B}(L^r)$ denote the class of Borel sets of L^r . Consider the regular conditional probability $P_k : \mathbb{R} \times \mathcal{B}(L^r) \rightarrow [0, 1]$, defined, for $z_k \in \mathbb{R}$ and $B \in \mathcal{B}(L^r)$,

$$P_k(z_k; B) := \mathbf{P}(U_k \in B \mid g_k = z_k) = \mathbf{E}(\mathbb{I}_{U_k \in B} \mid g_k = z_k).$$

Recall, see (77), that ψ_k denotes a L^r valued random variable with the distribution $\mathbf{P}\{\psi_k \in B\} = P_k(z_k; B)$. Note that the L^r valued random variable $\psi_k^* = p_g(\psi_k)$ has distribution

$$\begin{aligned} \mathbf{P}\{\psi_k^* \in B\} &= \mathbf{P}\{p_g(\psi_k) \in B\} = \mathbf{P}\{\psi_k \in p_g^{-1}(B)\} \\ (210) \quad &= \mathbf{P}(U_k \in p_g^{-1}(B) \mid g_k = z_k) = \mathbf{P}(U_k^* \in B \mid g_k = z_k). \end{aligned}$$

Furthermore, using (194) we write (210) in the form

$$\mathbf{P}\{\psi_k^* \in B\} = \mathbf{P}\left(U_k^{**} + \left(z_k - \frac{1}{N} \frac{\xi_k}{n^{1/2}}\right) b^* \in B \mid g_k = z_k\right).$$

Let $\bar{\psi}_k$ respectively $\bar{\psi}_k^*$ denote an independent copy of ψ_k respectively ψ_k^* . Denote

$$\tau_N = M^{-(r-4)/2} + N^{-\alpha(r-2)} M.$$

Lemma 10. *Let $k = 1, \dots, n-1$. Let $|z_k| \leq w n^{-1/2}$. There exist positive constants $c_*^{(i)}$, $i = 0, 1, 2, 3$, which depend on $w, r, \nu_1, \nu_2, \delta, A_*, D_*, M_*$ only such that for*

$$(211) \quad \tau_N \leq c_*^{(0)} \delta_3^2,$$

we have

$$(212) \quad c_*^{(1)} \delta_3^2 \leq n \mathbf{E} \|\psi_k^* - \bar{\psi}_k^*\|_2^2 \leq c_*^{(2)} \delta_3^2$$

$$(213) \quad \mathbf{E} \|\psi_k - \bar{\psi}_k\|_r^r \leq c_*^{(3)} n^{-r/2}.$$

The condition (211) requires N to be large enough. A simple calculation shows $\tau_N \leq N^{-75\nu}$, for ν satisfying (15). Therefore, (82) implies $\tau_N \leq N^{-65\nu} \delta_3^2$. In particular, under (82) the inequality (211) is satisfied provided that $N > c_*$, where c_* does not depend on δ_3^2 .

Proof. By $\tilde{c}_*, \tilde{c}'_*$ we denote positive constants which depend only on $w, r, \nu_1, \nu_2, \delta, A_*, D_*, M_*$. These constants can be different in different places of the text. Given $i, j \in O_k$, $i \neq j$, introduce random variables

$$\begin{aligned} g_* &= \eta + \zeta, & \eta &= \frac{\xi_k}{R}, & \zeta &= \frac{1}{\sqrt{M}} \sum_{j \in O_k} g(Y_j), \\ \zeta_i &= \zeta - \frac{g(Y_i)}{\sqrt{M}}, & \zeta_{ij} &= \zeta - \frac{g(Y_i)}{\sqrt{M}} - \frac{g(Y_j)}{\sqrt{M}}. \end{aligned}$$

Here $R = \sqrt{nMN}$ satisfies $N/2 \leq R \leq N$, by the choice of n and M . Let p, p_0, p_1 , and p_2 denote the densities of random variables $\eta, \zeta + \eta, \zeta_i + \eta$, and $\zeta_{ij} + \eta$ respectively.

Note that $g_* = \sqrt{N/M} g_k$. Therefore, the condition $g_k = z_k$ is equivalent to $g_* = z_*$, where $z_* = \sqrt{N/M} z_k$. Furthermore, $|z_k| \leq w n^{-1/2} \Leftrightarrow |z_*| \leq w_*$, where $w_* = w \sqrt{N/Mn} \leq 2w$.

Given a random variable Y , we denote the conditional expectation $\mathbf{E}(Y | g_* = z_*) = \mathbf{E}(Y | g_k = z_k)$ by $\mathbf{E}_* Y$. Similarly, for an event A , we have $P(A | g_k = z_k) = P(A | g_* = z_*)$.

Proof of (212). For the L^r valued random variable $\hat{\psi}^* = \psi_k^* - z_k b^*$ we have

$$(214) \quad \mathbf{P}\{\hat{\psi}^* \in B\} = P\left(U_k^{**} - \frac{1}{N} \frac{\xi_k}{n^{1/2}} b^* \in B \mid g_* = z_*\right).$$

Note that for an independent copy $\bar{\psi}_k^*$ of ψ_k^* the distributions of $\psi_k^* - \bar{\psi}_k^*$ and $\hat{\psi}^* - \hat{\psi}_c^*$ are the same. Here $\hat{\psi}_c^*$ denotes an independent copy of $\hat{\psi}^*$. Therefore,

$$(215) \quad \mathbf{E} \|\psi_k^* - \bar{\psi}_k^*\|_2^2 = \mathbf{E} \|\hat{\psi}^* - \hat{\psi}_c^*\|_2^2 = 2\mathbf{E} \|\hat{\psi}^*\|_2^2 - 2\|\mathbf{E} \hat{\psi}^*\|_2^2.$$

In order to prove (212) we show that

$$(216) \quad \|\mathbf{E} \hat{\psi}^*\|_2^2 \leq \tilde{c}_* N^{-1}$$

and, for $\tau_N \leq c_*^{(0)} \delta_3^2$ (i.e., for sufficiently large N),

$$(217) \quad \tilde{c}_* \delta_3^2 \leq n \mathbf{E} \|\hat{\psi}^*\|_2^2 \leq \tilde{c}'_* \delta_3^2.$$

Since $N^{-1}n < \tau_N$, we can choose $c_*^{(0)}$ small enough such that the inequalities (215), (216) and (217) together imply (212)

Proof of (216). Recall that an element $m = m(\cdot) \in L^2(P_X)$ is called mean of an $L^2(P_X)$ valued random variable $\hat{\psi}^* = \hat{\psi}^*(\cdot)$ if for every $f = f(\cdot) \in L^2(P_X)$

$$\langle f, m \rangle = \mathbf{E} \langle f, \hat{\psi}^* \rangle.$$

We shall show below that $\mathbf{E} \|\hat{\psi}^*\|_2^2 < \infty$. Then, by Fubini,

$$\mathbf{E} \langle f, \hat{\psi}^* \rangle = \int f(x) \mathbf{E} \hat{\psi}^*(x) P_X(dx).$$

Therefore, $m(x) = \mathbf{E} \hat{\psi}^*(x)$, for P_X almost all x .

For $f \in L^2(P_X)$ it follows from (214) that

$$(218) \quad \begin{aligned} \mathbf{E} \langle f, \hat{\psi}^* \rangle &= \mathbf{E}_* \left\langle f, U_k^{**} - \frac{1}{N} \frac{\xi_k}{n^{1/2}} b^* \right\rangle \\ &= \mathbf{E}_* \langle f, U_k^{**} \rangle - \frac{\sqrt{M}}{\sqrt{N}} \langle f, b^* \rangle \mathbf{E}_* \eta. \end{aligned}$$

Fix $i \in O_k$. By symmetry,

$$(219) \quad \mathbf{E}_* \langle f, U_k^{**} \rangle = \frac{M}{\sqrt{N}} \mathbf{E}_* \langle f, \psi^{**}(\cdot, Y_i) \rangle$$

An application of (247) yields

$$(220) \quad \begin{aligned} \mathbf{E}_* \langle f, \psi^{**}(\cdot, Y_i) \rangle &= \frac{1}{p_0(z_*)} \mathbf{E} \langle f, \psi^{**}(\cdot, Y_i) \rangle p_1\left(z_* - \frac{g(Y_i)}{\sqrt{M}}\right) \\ &= \langle f, a_{z_*} \rangle, \end{aligned}$$

where

$$a_{z_*}(\cdot) = \frac{b_{z_*}(\cdot)}{p_0(z_*)}, \quad b_{z_*}(\cdot) = \mathbf{E} \psi^{**}(\cdot, Y_i) p_1\left(z_* - \frac{g(Y_i)}{\sqrt{M}}\right)$$

are non-random elements of L^r .

It follows from (218), (219), (220) that

$$m(\cdot) = \frac{M}{\sqrt{N}} a_{z_*}(\cdot) - \frac{\sqrt{M}}{\sqrt{N}} b^*(\cdot) \mathbf{E}_* \eta.$$

In order to prove (216) we show that, for $|z_*| \leq w_*$,

$$(221) \quad \|b_{z_*}\|_2 \leq c_* M^{-1},$$

$$(222) \quad |\mathbf{E}_* \eta| \leq \tilde{c}_* M^{-1/2} + \tilde{c}'_* R^{-1/2},$$

$$(223) \quad p_i(z_*) \geq \tilde{c}_*, \quad i = 0, 1, 2,$$

and apply (203). Note that, by Lemma 7, there exist positive constants $\tilde{c}_*, \tilde{c}'_*$ such that, for $M, N > \tilde{c}'_*$, the inequality (223) holds.

Let us prove (221). In Lemma 7 we show, for $i = 1, 2$, that p_i and its derivatives are bounded functions. That is,

$$(224) \quad |p_i| \leq c_*, \quad |p'_i| \leq c_*, \quad |p''_i| \leq c_*, \quad |p'''_i| \leq c_*, \quad i = 1, 2.$$

Expanding in powers of $M^{-1/2}g(Y_i)$ we obtain

$$(225) \quad p_1\left(z_* - \frac{g(Y_i)}{\sqrt{M}}\right) = p_1(z_*) - \frac{g(Y_i)}{\sqrt{M}}p_1'(z_*) + \frac{g^2(Y_i)}{M} \frac{p_1''(\theta)}{2}.$$

It follows from the identities (191) and (192) that for P_X almost all x

$$\begin{aligned} \mathbf{E}\psi^{**}(x, Y_i) &= q_N^{-1} \mathbf{E}\psi^{**}(x, X_i) \mathbb{I}_{A_i} \\ &= q_N^{-1} \mathbf{E}\psi^{**}(x, X_i) (\mathbb{I}_{A_i} - 1) \\ &=: q_N^{-1} a_0(x) \\ \mathbf{E}\psi^{**}(x, Y_i) g(Y_i) &= q_N^{-1} \mathbf{E}\psi^{**}(x, X_i) g(X_i) \mathbb{I}_{A_i} \\ &= q_N^{-1} \mathbf{E}\psi^{**}(x, X_i) g(X_i) (\mathbb{I}_{A_i} - 1) \\ &=: q_N^{-1} a_1(x). \end{aligned}$$

Using (224) and the inequality $q_N^{-1} \leq c_*$, see (39), we obtain from (225)

$$\|b_{z_*}(\cdot)\|_2 \leq c_* \|a_0(\cdot)\|_2 + \frac{c_*}{\sqrt{M}} \|a_1(\cdot)\|_2 + \frac{c_*}{M} \|a_2(\cdot)\|_2,$$

where we denote $a_2(\cdot) = \mathbf{E}\psi^{**}(\cdot, Y_i) g^2(Y_i)$. In order to prove (221) we show that

$$(226) \quad \|a_0(\cdot)\|_2 \leq \frac{c_*}{N^{\alpha(r-1)}}, \quad \|a_1(\cdot)\|_2 \leq \frac{c_*}{N^{\alpha(r-2)}}, \quad \|a_2(\cdot)\|_2 \leq c_*.$$

Let us prove (226). Invoking (38) we obtain, by Hölder's inequality,

$$(227) \quad |a_0(x)| \leq \mathbf{E}|\psi^{**}(x, X_i)| \frac{\|Z_i'\|_r^{r-1}}{N^{\alpha(r-1)}} \leq w^{1/r}(x) \frac{K^{(r-1)/r}}{N^{\alpha(r-1)}},$$

where we denote $w(x) = \mathbf{E}|\psi^{**}(x, X_i)|^r$. Furthermore, by Lyapunov's inequality,

$$(228) \quad \|w^{1/r}(\cdot)\|_2^2 = \mathbf{E}w^{2/r}(X) \leq (\mathbf{E}w(X))^{2/r} = K_r^{2/r}.$$

Clearly, the first bound of (226) follows from (227), (228) and (195). A similar argument shows the second bound of (226). We have

$$(229) \quad |a_1(x)| \leq \mathbf{E}|\psi^{**}(x, X_i) g(X_i)| \frac{\|Z_i'\|_r^{r-2}}{N^{\alpha(r-2)}} \leq w^{1/r}(x) \frac{V^{(r-1)/r}}{N^{\alpha(r-2)}},$$

where we denote $V = \mathbf{E}(\|Z_i'\|_r^{r-2} |g(X_i)|)^{r/(r-1)}$. By Hölder's inequality,

$$(230) \quad V \leq (\mathbf{E}|g(X_i)|^r)^{1/(r-1)} (\mathbf{E}\|Z_i'\|_r^r)^{(r-2)/(r-1)} \leq c_*.$$

Clearly, (228), (229) and (230) imply the second bound of (226). The last bound of (226) follows from (42), by Cauchy-Schwartz. Indeed, we have

$$|a_2(x)| \leq c_* \mathbf{E}|\psi^{**}(x, X_i)| g^2(X_i) \leq c_* (\mathbf{E}|\psi^{**}(x, X_i)|^2 \mathbf{E}g^4(X_i))^{1/2}.$$

Therefore, $\|a_2(\cdot)\|_2^2 \leq c_* K_2 \mathbf{E}g^4(X_i) \leq c_*$, by (195).

Let us prove (222). We have, by (246),

$$\mathbf{E}_* \eta = p_0^{-1}(z_*) \mathbf{E}(z_* - \zeta) p(z_* - \zeta).$$

In order to prove (222) it suffices to show in view of (223) that

$$(231) \quad |\mathbf{E}(z_* - \zeta) p(z_* - \zeta)| \leq c_* R^{-1/2} + c_* M^{-1/2}.$$

Let \tilde{p} denotes the density function of ξ_k . Then $p(u) = R \tilde{p}(Ru)$. We have

$$\mathbf{E}(z_* - \zeta) p(z_* - \zeta) = 6c_\xi \mathbf{E} \frac{\sin^6(R(z_* - \zeta)/6)}{(R(z_* - \zeta)/6)^5}.$$

Therefore, denoting $H(z_*) = 1 + |R(z_* - \zeta)|^5$, we obtain

$$(232) \quad \mathbf{E}|(z_* - \zeta)p(z_* - \zeta)| \leq c\mathbf{E}H^{-1}(z_*).$$

On the event $|\zeta - z_*| \geq R^{-1/2}$ we have $H^{-1}(z_*) \leq R^{-5/2}$. Furthermore, a bound for the probability of the complementary event

$$\mathbf{P}\{|\zeta - z_*| \leq R^{-1/2}\} \leq c_*R^{-1/2} + c_*M^{-1/2},$$

follows by the Berry-Esseen bound applied to the sum ζ . Therefore, $\mathbf{E}H^{-1}(z_*)$ is bounded by the right hand side of (231). Now (231) follows from (232).

Proof of (217). Write

$$U_k^{**} - \frac{1}{N} \frac{\xi_k}{\sqrt{n}} b^* = \frac{\sqrt{M}}{\sqrt{N}} (T_1 - T_2),$$

$$T_1 := \frac{1}{\sqrt{M}} \sum_{j \in O_k} \psi^{**}(\cdot, Y_j), \quad T_2 := \eta b^*.$$

It follows from (214), by the inequality $\|u + v\|_2^2 \geq \|u\|_2^2/2 - \|v\|_2^2$, for $u, v \in L^2(P_X)$, that

$$\mathbf{E}\|\hat{\psi}^*\|_2^2 = \frac{M}{N} \mathbf{E}_* \|T_1 - T_2\|_2^2 \geq \frac{M}{2N} \mathbf{E}_* \|T_1\|_2^2 - \frac{M}{N} \mathbf{E}_* \|T_2\|_2^2.$$

We shall show that

$$(233) \quad \mathbf{E}_* \|T_2\|_2^2 \leq p_0^{-1}(z_*) (c_* R^{-1} M^{-1/2} + c_* R^{-3/2}),$$

$$(234) \quad \mathbf{E}_* \|T_1\|_2^2 \geq p_0^{-1}(z_*) (p_1(z_*) \delta_3^2 - c_* \tau_N).$$

$$(235) \quad \mathbf{E}_* \|T_1\|_2^2 \leq p_0^{-1}(z_*) (p_1(z_*) \delta_3^2 + c_* \tau_N).$$

The inequalities (233) and (234) imply the lower bound in (217). Indeed, by (223), we have, for small $c_*^{(0)}$,

$$\frac{c_*}{M^{1/2} R} + \frac{c_*}{R^{3/2}} \leq c_* \tau_N \leq c_* c_*^{(0)} \delta_3^2 \leq p_1(z_*) \delta_3^2 / 4.$$

Similarly, the inequalities (233) and (235) imply the upper bound in (217).

Proof of (233). We have, by (246),

$$\mathbf{E}_* \eta^2 = p_0^{-1}(z_*) W, \quad W := \mathbf{E}(z_* - \zeta)^2 p(z_* - \zeta).$$

Proceeding as in proof of (231), we obtain

$$W = \frac{36}{R} c_\xi \mathbf{E} \frac{\sin^6(R(z_* - \zeta)/6)}{(R(z_* - \zeta)/6)^4} \leq \frac{c}{R} \mathbf{E} \tilde{H}^{-1}(z_*),$$

where $\tilde{H}(z_*) = 1 + |R(z_* - \zeta)|^4$ satisfies

$$\mathbf{E} \tilde{H}^{-1}(z_*) \leq c_* R^{-1/2} + c_* M^{-1/2}.$$

Therefore, $W \leq c_* R^{-3/2} + c_* R^{-1} M^{-1/2}$. This inequality in combination with (203) implies (233).

Proof of (234). Fix $i, j \in O_k$, $i \neq j$. By symmetry,

$$(236) \quad \mathbf{E}_* \|T_1\|_2^2 = \mathbf{E}_* T_{11} + (M - 1) \mathbf{E}_* T_{12},$$

$$T_{11} = \|\psi^{**}(\cdot, Y_i)\|_2^2, \quad T_{12} = \langle \psi^{**}(\cdot, Y_i), \psi^{**}(\cdot, Y_j) \rangle.$$

We have, by (247),

$$\mathbf{E}_* T_{11} = p_0^{-1}(z_*) H_1, \quad \mathbf{E}_* T_{12} = p_0^{-1}(z_*) H_2,$$

$$H_1 = \mathbf{E} T_{11} p_1\left(z_* - \frac{g(Y_i)}{\sqrt{M}}\right), \quad H_2 = \mathbf{E} T_{12} p_2\left(z_* - \frac{g(Y_i) + g(Y_j)}{\sqrt{M}}\right).$$

The inequality (234) follows from (236) and the bounds

$$(237) \quad H_1 \geq p_1(z_*)\delta_3^2 - c_*M^{-1/2},$$

$$(238) \quad |H_2| \leq c_*N^{-\alpha(r-2)} + c_*M^{-(r-2)/2}.$$

Let us prove (237). It follows from (224), by the mean value theorem, that

$$(239) \quad H_1 = p_1(z_*)\mathbf{E}T_{11} + Q, \quad |Q| \leq c_*\mathbf{E}T_{11} \frac{|g(Y_i)|}{\sqrt{M}},$$

where $|Q| \leq c_*M^{-1/2}$. Indeed, by (42) and Cauchy-Schwartz,

$$\mathbf{E}T_{11}|g(Y_i)| \leq \mathbf{E}\|\psi^{**}(\cdot, X_i)\|_2^2|g(X_i)| \leq K_4^{1/2}\sigma \leq c_*.$$

In the last step we applied (195). Furthermore, the identity

$$\begin{aligned} \mathbf{E}T_{11} &= q_N^{-1}\mathbf{E}|\psi^{**}(X, X_i)|^2\mathbb{I}_{A_i} = \mathbf{E}|\psi^{**}(X, X_i)|^2 - b_1 - b_2, \\ b_1 &= (1 - q_N^{-1})\mathbf{E}|\psi^{**}(X, X_i)|^2, \quad b_2 = q_N^{-1}\mathbf{E}|\psi^{**}(X, X_i)|^2(1 - \mathbb{I}_{A_i}) \end{aligned}$$

combined with (38), (39) and (40) yields $\mathbf{E}T_{11} \geq \delta_3^2 - c_*M^{-1/2}$. This bound together with (239) shows (237).

Let us prove (238). Write $y_i = g(Y_i)$ and expand

$$p_2\left(z_* - \frac{y_i + y_j}{\sqrt{M}}\right) = p_2(z_*) - p_2'(z_*)\frac{y_i + y_j}{\sqrt{M}} + \frac{p_2''(z_*)}{2}\frac{(y_i + y_j)^2}{M} + \tilde{Q}.$$

From (224) it follows, for $2 < r - 2 \leq 3$, the bound

$$|\tilde{Q}| \leq c_*|y_i + y_j|^{r-2}/M^{(r-2)/2}.$$

Furthermore, denote

$$\begin{aligned} h_1 &= \mathbf{E}T_{12}, & h_2 &= \mathbf{E}T_{12}g(Y_i), \\ h_3 &= \mathbf{E}T_{12}g^2(Y_i), & h_4 &= \mathbf{E}T_{12}g(Y_i)g(Y_j). \end{aligned}$$

We obtain, by symmetry,

$$(240) \quad \begin{aligned} H_2 &= p_2(z_*)h_1 - 2\frac{p_2'(z_*)}{\sqrt{M}}h_2 + \frac{p_2''(z_*)}{M}(h_3 + h_4) + \mathbf{E}T_{12}\tilde{Q}, \\ \mathbf{E}|T_{12}\tilde{Q}| &\leq c_*M^{-(r-2)/2}\mathbf{E}|g(Y_i)|^{r-2}|T_{12}|. \end{aligned}$$

Denote

$$\tilde{T}_{12} = q_N^{-2}\psi^{**}(X, X_i)\psi^{**}(X, X_j).$$

It follows from (40), by Hölders inequality and (195), that

$$\mathbf{E}|g(Y_i)|^{r-2}|T_{12}| \leq \mathbf{E}|g(X_i)|^{r-2}|\tilde{T}_{12}| \leq c_*.$$

Therefore,

$$(241) \quad \mathbf{E}|T_{12}\tilde{Q}| \leq c_*M^{-(r-2)/2}.$$

Furthermore, (191) and (192) imply

$$\begin{aligned} h_1 &= \mathbf{E}\tilde{T}_{12}(\mathbb{I}_{A_i} - 1)(\mathbb{I}_{A_j} - 1), & h_2 &= \mathbf{E}\tilde{T}_{12}g(X_i)(\mathbb{I}_{A_i} - 1)(\mathbb{I}_{A_j} - 1), \\ h_3 &= \mathbf{E}\tilde{T}_{12}g^2(X_i)\mathbb{I}_{A_i}(\mathbb{I}_{A_j} - 1), & h_4 &= \mathbf{E}\tilde{T}_{12}g(X_i)g(X_j)(\mathbb{I}_{A_i} - 1)(\mathbb{I}_{A_j} - 1). \end{aligned}$$

Invoking the inequalities $q_N^{-2} \leq c_*$, see (39), and $1 - \mathbb{I}_{A_i} \leq V_i^s$, $s > 0$, where $V_i := \|Z'_i\|_r/N^\alpha$, see (38), we obtain, by Hölder's inequality,

$$(242) \quad \begin{aligned} |h_1| &\leq \mathbf{E}|\tilde{T}_{12}|(V_i V_j)^{(r-2)/2} \leq c_* N^{-\alpha(r-2)}, \\ |h_2| &\leq \mathbf{E}|\tilde{T}_{12}g(X_i)|V_i^{(r-4)/2}V_j^{(r-2)/2} \leq c_* N^{-\alpha(r-3)}, \\ |h_3| &\leq \mathbf{E}|\tilde{T}_{12}|g^2(X_i)V_j^{r-4} \leq c_* N^{-\alpha(r-4)}, \\ |h_4| &\leq \mathbf{E}|\tilde{T}_{12}g(X_i)g(X_j)|(V_i V_j)^{(r-4)/2} \leq c_* N^{-\alpha(r-4)}. \end{aligned}$$

Combining (240), (242), (241) and using the simple inequalities

$$\frac{1}{N^{\alpha(r-3)}M^{1/2}} \leq \frac{1}{\tilde{N}}, \quad \frac{1}{N^{\alpha(r-4)}M} \leq \frac{1}{\tilde{N}}, \quad \tilde{N} = \min\{N^{\alpha(r-2)}, M^{(r-2)/2}\}$$

and the inequalities (224), we obtain (238).

Proof of (235). The inequality follows from (236), (238) and the inequality

$$\begin{aligned} H_1 &\leq p_1(z_*)\mathbf{E}T_{11} + c_*M^{-1/2} \leq p_1(z_*)\delta_3^2 + c_*N^{-\alpha r} + c_*M^{-1/2} \\ &\leq p_1(z_*)\delta_3^2 + c_*M^{-1/2}, \end{aligned}$$

which is obtained in the same way as (237) above.

Proof of (213). In order to prove (213) we shall show that

$$(243) \quad \mathbf{E}\|\psi_k\|_r^r \leq \tilde{c}_*n^{-r/2}.$$

Split $O_k = B \cup D$, where $B \cap D = \emptyset$ and $|B| = \lfloor M/2 \rfloor$ and write

$$\begin{aligned} U_k &= \frac{\sqrt{M}}{\sqrt{N}}(U_B + U_D), \quad U_B = \frac{1}{\sqrt{M}} \sum_{j \in B} \psi(\cdot, Y_j), \\ \zeta &= \zeta_B + \zeta_D, \quad \zeta_B = \frac{1}{\sqrt{M}} \sum_{j \in B} g(Y_j). \end{aligned}$$

In particular, we have $g_* = \eta + \zeta_B + \zeta_D$.

The inequality

$$\mathbf{E}\|\psi_k\|_r^r = \mathbf{E}_* \|U_k\|_r^r \leq 2^r \left(\frac{M}{N}\right)^{r/2} (\mathbf{E}_* \|U_B\|_r^r + \mathbf{E}_* \|U_D\|_r^r)$$

combined with the bounds

$$(244) \quad \mathbf{E}_* \|U_B\|_r^r \leq c_*, \quad \mathbf{E}_* \|U_D\|_r^r \leq c_*$$

imply (243). Let us prove the first bound of (244). By (247), we have

$$\mathbf{E}_* \|U_B\|_r^r = p_0^{-1}(z_*)\mathbf{E}\|U_B\|_r^r p_3(z_* - \zeta_B),$$

where p_3 denotes the density of $\eta + \zeta_D$. Furthermore, invoking the bound $\sup_{x \in \mathbb{R}} |p_3(x)| \leq c_*$, (which is obtained using the same argument as in the proof of Lemma 7) and the inequality (223), we obtain $\mathbf{E}_* \|U_B\|_r^r \leq \tilde{c}_* \mathbf{E}\|U_B\|_r^r$. Finally, invoking the bound

$$(245) \quad \mathbf{E}\|U_B\|_r^r \leq c_* \mathbf{E}\left\| \frac{1}{\sqrt{M}} \sum_{j \in B} \psi(\cdot, X_j) \right\|_r^r \leq c_*,$$

see (42) and (209), we obtain the first bound of (244). The second bound is obtained in the same way. This completes the proof of the lemma. \square

We collect some facts about conditional moments in a separate lemma.

Lemma 11. *Let η and ζ be independent random variables. Assume that η is real valued and has a density, say $x \rightarrow p(x)$.*

(i) *Assume that ζ is real valued. Then the function*

$$x \rightarrow \mathbf{E}p(x - \zeta), \quad x \in \mathbb{R},$$

is a density of the distribution $P_{\eta+\zeta}$ of $\eta + \zeta$. Let $w : \mathbb{R} \rightarrow \mathbb{R}$ be a measurable function such that $\mathbf{E}|w(\eta)| < \infty$. For $P_{\eta+\zeta}$ almost all $x \in \mathbb{R}$, we have

$$(246) \quad \mathbf{E}(w(\eta) \mid \eta + \zeta = x) = \frac{\mathbf{E}w(x - \zeta)p(x - \zeta)}{\mathbf{E}p(x - \zeta)}.$$

(ii) *Assume that ζ takes values in a measurable space, say \mathcal{Y} . Assume that $u, v : \mathcal{Y} \rightarrow \mathbb{R}$ are measurable functions and denote $P_{\eta+u(\zeta)}$ the distribution of $\eta + u(\zeta)$. If $\mathbf{E}|v(\zeta)| < \infty$, then for $P_{\eta+u(\zeta)}$ almost all $x \in \mathbb{R}$,*

$$(247) \quad \mathbf{E}(v(\zeta) \mid \eta + u(\zeta) = x) = \frac{\mathbf{E}v(\zeta)p(x - u(\zeta))}{\mathbf{E}p(x - u(\zeta))}.$$

9. APPENDIX 4

In the next lemma we consider independent and identically distributed random vectors (ξ, η) and (ξ', η') with values in \mathbb{R}^2 and the symmetrization (ξ_s, η_s) where $\xi_s = \xi - \xi'$ and $\eta_s = \eta - \eta'$. Note that in the main text we apply this lemma to $\xi = g(X_1)$ and $\eta = N^{-1/2} \sum_{j=m+1}^N \psi(X_1, Y_j)$.

Lemma 12. *Let $0 < \nu < 1/2$ and $r > 2$. Assume that $\mathbf{E}|\xi|^r + \mathbf{E}|\eta|^r < \infty$. The following statements hold.*

a) *For $c_r = (7/12)2^{-r}$ the conditions*

$$|t|^{r-2} \mathbf{E}|\xi_s|^r \leq c_r \mathbf{E}\xi_s^2, \quad \mathbf{E}\xi_s \eta_s = 0, \quad \mathbf{E}|\eta_s|^r \leq c_r \mathbf{E}\eta_s^2$$

imply $1 - |\mathbf{E} \exp\{i(t\xi + \eta)\}|^2 \geq 6^{-1}(t^2 \mathbf{E}\xi_s^2 + \mathbf{E}\eta_s^2)$.

b) *Assume that for some $\tilde{c}_1, \tilde{c}_2 > 0$ we have*

$$(248) \quad \mathbf{E}\xi_s^2/12 - N^{-1} \mathbf{E}\eta_s^2 > \tilde{c}_1^2 \quad \text{and} \quad c_r \mathbf{E}\xi_s^2 / \mathbf{E}|\xi_s|^r \geq \tilde{c}_2^{r-2}.$$

Let $\varepsilon > 0$ be such that

$$(249) \quad \varepsilon < 1/6\tilde{c}_3, \quad \varepsilon^{(r-2)/2} < \sigma_z^2/4A, \quad \varepsilon^{r-2} < \sigma_z^2/4B,$$

where $\tilde{c}_3 = 2 + (5/\tilde{c}_1)^2 \sigma_z^2$ and where the numbers A, B are defined in (260). Here $\sigma_z^2 = \mathbf{E}(\xi_s + N^{-1/2} \eta_s)^2$. Assume that for some $0 < \delta < \tilde{c}_2$ and $\delta' > 10\varepsilon^2$,

$$(250) \quad \sup_{\delta < |t| < N^{-\nu+1/2}} |\mathbf{E}e^{it\xi_s}| \leq 1 - \delta' \quad \text{and} \quad \mathbf{E}|\eta_s| \leq \delta' N^\nu/2.$$

Then for every T^ , satisfying $N^{1/2-\nu} \leq |T^*| \leq N^{\nu+1/2}$, the set*

$$I^* = \{T^* \leq t \leq T^* + N^{1/2-\nu} : |\mathbf{E}e^{it(\xi + N^{-1/2}\eta)}|^2 \geq 1 - \varepsilon^2\}$$

is an interval of size at most $5\tilde{c}_1^{-1}\varepsilon$.

Proof. Proof a). Invoking the inequality $1 - \cos x \geq x^2/2 - x^2/24 - |x|^r$ and using the simple inequality $|a + b|^r \leq 2^{r-1}(|a|^r + |b|^r)$ we obtain

$$\begin{aligned} 1 - |\mathbf{E} \exp\{i(t\xi + \eta)\}|^2 &= 1 - \mathbf{E} \cos(t\xi_s + \eta_s) \\ &\geq \frac{11}{24} \mathbf{E}(t\xi_s + \eta_s)^2 - 2^{r-1}(\mathbf{E}|t\xi_s|^r + \mathbf{E}|\eta_s|^r) \\ &\geq 6^{-1}(t^2 \mathbf{E}\xi_s^2 + \mathbf{E}\eta_s^2). \end{aligned}$$

In the last step we use the conditions a).

Proof **b**). Introduce the function $t \rightarrow \tau_t^* = 1 - |\mathbf{E}e^{it(\xi + N^{-1/2}\eta)}|^2$. Assume that the set I^* is non-empty and choose $s, t \in I^*$, i.e., we have $\tau_t^*, \tau_s^* \leq \varepsilon^2$. Firstly we show that $|s - t| \leq 5\tilde{c}_1^{-1}\varepsilon$, thus proving the bound for the size of the set I^* .

The inequality $1 - \cos(x + y) \geq (1 - \cos x)/2 - (1 - \cos y)$ implies

$$(251) \quad 1 - |\mathbf{E}e^{i(X+Y)}|^2 \geq 2^{-1}(1 - |\mathbf{E}e^{iX}|^2) - (1 - |\mathbf{E}e^{iY}|^2),$$

for arbitrary random variables X, Y . Choosing $\tilde{Y} = t(\xi + N^{-1/2}\eta)$ and $\tilde{X} = (s - t)(\xi + N^{-1/2}\eta)$ shows

$$(252) \quad \tau_s^* \geq (1 - |\mathbf{E}e^{i\tilde{X}}|^2)/2 - \tau_t^*.$$

Now we show that the inequality $|t - s| > 5\tilde{c}_1^{-1}\varepsilon$ implies $1 - |\mathbf{E}e^{i\tilde{X}}|^2 > 5\varepsilon^2$, thus, contradicting to our choice $\tau_s^*, \tau_t^* < \varepsilon^2$ and (252). In what follows the cases of ‘‘large’’ and ‘‘small’’ values of $|t - s|$ are treated separately.

For $5\tilde{c}_1^{-1}\varepsilon < |t - s| \leq \delta$ we shall apply (251) to $\tilde{X} = X + Y$, where $X = (s - t)\xi$ and $Y = (s - t)N^{-1/2}\eta$. Note that the statement **a**) implies

$$(253) \quad 1 - |\mathbf{E}e^{iX}|^2 \geq (t - s)^2 \mathbf{E}\xi_s^2/6.$$

Indeed, in view of the second inequality of (248), the conditions of **a**) are satisfied for $|t - s| \leq \delta \leq \tilde{c}_2$. Furthermore, we have

$$(254) \quad 0 \leq 1 - |\mathbf{E}e^{iY}|^2 = 1 - \cos(N^{-1/2}(s - t)\eta_s) \leq (s - t)^2 N^{-1} \mathbf{E}\eta_s^2.$$

Invoking the bounds (253) and (254) in (251) we obtain

$$1 - |\mathbf{E}e^{i\tilde{X}}|^2 \geq (s - t)^2 \mathbf{E}\xi_s^2/12 - (s - t)^2 N^{-1} \mathbf{E}\eta_s^2 \geq \tilde{c}_1^2 (s - t)^2 \geq 25\varepsilon^2.$$

In the last step we used (248).

For $\delta < |t - s| \leq N^{-\nu+1/2}$ we expand in powers of $a = i(s - t)N^{-1/2}\eta_s$ to get

$$\begin{aligned} 1 - |\mathbf{E}e^{i\tilde{X}}|^2 &= 1 - \mathbf{E} \exp\{i(s - t)\xi_s + a\} \geq 1 - \mathbf{E} \exp\{i(s - t)\xi_s\} - \mathbf{E}|a| \\ &\geq \delta' - \mathbf{E}|(t - s)N^{-1/2}\eta_s| \geq \delta' - N^{-\nu} \mathbf{E}|\eta_s| \\ &\geq \delta'/2 \geq 5\varepsilon^2. \end{aligned}$$

In the last step we applied (250).

Let us prove that I^* is indeed an interval. Assume the contrary, i.e. there exist $s < u < t$ such that $s, t \in I^*$ and $u \notin I^*$. In particular, $\tau_t^* \leq \varepsilon^2 < \tau_u^*$. Clearly, we can choose u to be a local maximum (stationary) point of the function $t \rightarrow \tau_t^*$. Denote

$$z = \xi_s + N^{-1/2}\eta_s, \quad \sigma_z^2 = \mathbf{E}z^2.$$

An application of (251) to $Y' = (t - u)(\xi + N^{-1/2}\eta)$ and $X' = u(\xi + N^{-1/2}\eta)$ gives

$$\tau_t^* \geq \tau_u^*/2 - (1 - \mathbf{E}e^{i(t-u)z}) = \tau_u^*/2 - (1 - \mathbf{E} \cos(t - u)z).$$

Invoking the inequalities $\tau_t^* \leq \varepsilon^2$ and $1 - \cos(t - u)z \leq (t - u)^2 z^2/2$ we obtain

$$(255) \quad \tau_u^* \leq 2\varepsilon^2 + (t - u)^2 \sigma_z^2 \leq \varepsilon^2 \tilde{c}_3, \quad \tilde{c}_3 = 2 + (5/\tilde{c}_1)^2 \sigma_z^2.$$

Here we used the bound $|t - u| \leq |t - s| \leq 5\varepsilon/\tilde{c}_1$ proved above.

Denoting $y = (t - u)z$ we have $\tau_t^* = 1 - \mathbf{E}e^{iuz}e^{iy}$. Invoking the expansion $e^{iy} = 1 + iy + (iy)^2/2 + R'$, where $|R'| \leq y^2/6 + |y|^r$, we obtain

$$(256) \quad \tau_t^* = \tau_u^* - i\mathbf{E}ye^{iuz} + 2^{-1}\mathbf{E}y^2e^{iuz} + R, \quad |R| \leq \mathbf{E}y^2/6 + \mathbf{E}|y|^r =: R_0.$$

For a stationary point u we have $0 = \frac{\partial}{\partial t}\tau_t^*|_{t=u} = -i\mathbf{E}ze^{iuz}$. Therefore, $\mathbf{E}ye^{iuz} = 0$ and (256) implies

$$\tau_t^* \geq \tau_u^* + 2^{-1}(t - u)^2 \mathbf{E}z^2 e^{iuz} - R_0.$$

Write the right hand side in the form $\tau_u^* + 2^{-1}(t-u)^2 R_1$, where

$$R_1 = \mathbf{E}z^2 e^{iuz} - 3^{-1}\sigma_z^2 - 2\mathbf{E}|z|^r |t-u|^{r-2}.$$

Note that the inequality $R_1 > 0$ contradicts to our assumption $\tau_t^* < \tau_u^*$. We complete the proof by showing that $R_1 > 0$.

Since the random variable z is symmetric we have $\mathbf{E}z^2 \sin uz = 0$. Therefore,

$$(257) \quad \mathbf{E}z^2 e^{iuz} = \mathbf{E}z^2 \cos uz = \sigma_z^2 - \mathbf{E}z^2(1 - \cos uz).$$

Given $\lambda > 0$ split

$$(258) \quad \begin{aligned} \mathbf{E}z^2(1 - \cos uz) &= \mathbf{E}z^2(1 - \cos uz) \left(\mathbb{I}_{\{|z|^2 < \lambda^2\}} + \mathbb{I}_{\{|z|^2 \geq \lambda^2\}} \right) \\ &\leq \lambda^2 \mathbf{E}(1 - \cos uz) + 2\mathbf{E}|z|^r \lambda^{2-r}. \end{aligned}$$

In the last step we used Chebyshev's inequality. Furthermore, invoking the inequality $\mathbf{E}(1 - \cos uz) = \tau_u^* \leq \tilde{c}_3 \varepsilon^2$, see (255), we obtain from (257) and (258) for $\lambda^2 = \varepsilon^{-1} \sigma_z^2$

$$(259) \quad \mathbf{E}z^2 e^{iuz} \geq \sigma_z^2 - \varepsilon \tilde{c}_3 \sigma_z^2 - \varepsilon^{(r-2)/2} 2\mathbf{E}|z|^r \sigma_z^{2-r}.$$

Finally, invoking the inequality $|t-u| \leq |t-s| \leq 5\tilde{c}_1^{-1} \varepsilon$ we obtain from (259)

$$R_1 \geq \sigma_z^2(1 - 3^{-1} - \varepsilon \tilde{c}_3) - \varepsilon^{(r-2)/2} A - \varepsilon^{r-2} B,$$

where for random variable $z = \xi_s + N^{-1/2} \eta_s$ we write

$$(260) \quad A = 2\mathbf{E}|z|^r \sigma_z^{2-r} \quad \text{and} \quad B = 2\mathbf{E}|z|^r (5/\tilde{c}_1)^{r-2}.$$

Thus, for ε satisfying (249) we have $R_1 > 0$. □

10. APPENDIX 5

Let Z_1, \dots, Z_N be independent copies of the L^r valued random element $Z = \{x \rightarrow \psi(x, Y)\}$. Recall that almost surely $\|Z\| \leq N^\alpha$. Here $\|\cdot\|$ denotes the norm of the Banach space L^r , where $r > 4$ and $1/2 > \alpha > 0$. Write $M_p = \mathbf{E}|\psi(X_1, X_2)|^p$.

Lemma 13. (i) Assume that $\|\mathbf{E}Z\|^2 \leq \mathbf{E}\|Z\|^2/N$. Then there exists a constant $c(r) > 0$ such that for $k \leq N$ and $x > c(r)$ we have

$$(261) \quad \mathbf{P}\{\|Z_1 + \dots + Z_k\| > k^{1/2} u x\} \leq \exp\{-2^{-5} x^2 (1 + x N^\alpha / k^{1/2} u)^{-1}\}.$$

Here $u^2 = \mathbf{E}\|Z\|^2$.

(ii) The following inequalities hold

$$(262) \quad \|\mathbf{E}Z\| \leq M_r / q_N N^{(r-1)\alpha}$$

$$(263) \quad q_N^{-1} (M_2 - M_r N^{-(r-2)\alpha}) \leq \mathbf{E}\|Z\|^2 \leq q_N^{-1} (M_r^{2/r} + M_r N^{-(r-2)\alpha}).$$

Remark. Assume that

$$M_2 \geq 2M_r N^{-(r-2)\alpha}, \quad M_r^2 \leq (q_N/2) M_2 N^\varkappa, \quad \text{where} \quad \varkappa = 2(r-1)\alpha - 1.$$

Then (262) and (263) imply the inequality $\|\mathbf{E}Z\|^2 \leq \mathbf{E}\|Z\|^2/N$. Note that $r\alpha > 2$ implies $\varkappa > 2$. Furthermore, by (39), the probability $q_N > 1 - M_r N^{-r\alpha}$.

Proof. We derive (i) from Yurinskii's (1976) inequality. Denote $\zeta_k = Z_1 + \dots + Z_k$. Using the type-2 inequality for an L^r valued random variable $\zeta_k - \mathbf{E}\zeta_k$,

$$\mathbf{E}\|\zeta_k - \mathbf{E}\zeta_k\|^2 \leq k\tilde{c}(r)\mathbf{E}\|Z_1 - \mathbf{E}Z_1\|^2,$$

and the inequality $\|Z_1 - \mathbf{E}Z_1\|^2 \leq 2\|Z_1\|^2 + 2\|\mathbf{E}Z_1\|^2$, we obtain

$$\mathbf{E}\|\zeta_k - \mathbf{E}\zeta_k\| \leq (\mathbf{E}\|\zeta_k - \mathbf{E}\zeta_k\|^2)^{1/2} \leq k^{1/2} c'(r)(u + \|\mathbf{E}Z_1\|).$$

We have

$$\begin{aligned} \mathbf{E}\|\zeta_k\| &\leq \mathbf{E}\|\zeta_k - \mathbf{E}\zeta_k\| + k\|\mathbf{E}Z_1\| \\ &\leq c'(r)k^{1/2}u + k(1 + c'(r)k^{-1/2})\|\mathbf{E}Z_1\| =: \beta_k. \end{aligned}$$

It follows from the inequality $\|Z_1\| \leq N^\alpha$ that

$$\mathbf{E}\|Z_1\|^L \leq 2^{-1}L!u^2N^{\alpha(L-2)}, \quad L = 2, 3, \dots$$

Write $B_k^2 = ku^2$. Theorem 2.1 of Yurinskii (1976) shows

$$(264) \quad \mathbf{P}\{\|\zeta_k\| \geq xB_k\} \leq \exp\{-B\}, \quad B = \frac{\bar{x}^2}{8}(1 + (\bar{x}N^\alpha/2B_k))^{-1},$$

provided that $\bar{x} = x - \beta_k/B_k > 0$.

Since $\beta_k/B_k \leq 1 + c'(r)(1 + k^{-1/2})$ we have, for $x > c(r) := 4c'(r) + 2$,

$$x > 2\beta_k/B_k \quad \text{and} \quad x > \bar{x} > x/2.$$

The latter inequality implies

$$B \geq B' := \frac{(x/2)^2}{8}(1 + (xN^\alpha/B_k))^{-1}.$$

Finally, replacing B by B' in (264) we obtain (261).

Let us prove (ii). The mean value $\mathbf{E}Z = \{x \rightarrow \mathbf{E}\psi(x, Y)\}$ is an element of L^r . For P_X almost all $x \in \mathcal{X}$ we have $\mathbf{E}\psi(x, X) = 0$. Therefore,

$$\mathbf{E}Z = q_N^{-1}\mathbf{E}\psi(x, X)\mathbb{I}_A = q_N^{-1}\mathbf{E}\psi(x, X)(\mathbb{I}_A - 1).$$

Invoking (38) and using Chebyshev and Hölder inequalities, we obtain, for P_X almost all x ,

$$|\mathbf{E}Z| \leq \frac{1}{q_N N^{\alpha(r-1)}} \mathbf{E}\|Z'\|_r^{r-1} |\psi(x, X)| \leq \frac{1}{q_N N^{\alpha(r-1)}} (\mathbf{E}\|Z'\|_r^r)^{(r-1)/r} a(x),$$

where $a(x) = (\mathbf{E}|\psi(x, X)|^r)^{1/r}$. Note that $\mathbf{E}\|Z'\|_r^r = M_r$ and $\|a\|^r = M_r$. Finally,

$$\|\mathbf{E}Z\| \leq \|a\| M_r^{(r-1)/r} / q_N N^{\alpha(r-1)} = M_r / q_N N^{\alpha(r-1)}.$$

Let us prove (263). Denote $b_p(x) = (\mathbf{E}_{X_1}|\psi(X_1, x)|^p)^{1/p}$. Here \mathbf{E}_{X_1} denotes the conditional expectation given all the random variables, but X_1 . We have

$$(265) \quad \mathbf{E}\|Z\|^2 = q_N^{-1}\mathbf{E}\mathbb{I}_A b_r^2(X) = q_N^{-1}\mathbf{E}b_r^2(X) + q_N^{-1}R, \quad R = \mathbf{E}(\mathbb{I}_A - 1)b_r^2(X).$$

By Hölder's inequality $b_r(x) \geq b_2(x)$, for P_X almost all x . Therefore,

$$(266) \quad M_2 = \mathbf{E}b_2^2(X) \leq \mathbf{E}b_r^2(X) \leq M_r^{2/r}$$

Combining (266) and (265) and the bound $|R| \leq M_r N^{-(r-2)\alpha}$ we obtain (263). In order to bound $|R|$ we use (38), $|R| \leq N^{-(r-2)\alpha} \mathbf{E}\|Z'\|_r^{r-2} b_r^2(X)$ and apply Hölder inequality,

$$\mathbf{E}\|Z'\|_r^{r-2} b_r^2(X) \leq (\mathbf{E}\|Z'\|_r^r)^{(r-2)/r} (\mathbf{E}b_r^r(X))^{2/r} = M_r.$$

REFERENCES

- [1] Babu, Gutti Jogesh, Bai, Z.D.: Edgeworth expansions of a function of sample means under minimal moment conditions and partial Cramer's condition. *Sankhya, Ser. A* 55, (1993), 244–258.
- [2] Bai, Z.D., Rao, Radhakrishna, C.: Edgeworth expansion of a function of sample means. *Ann. Statist.* 19, (1991), 1295–1315.
- [3] Bentkus, V., Götze, F. and van Zwet, W. R.: An Edgeworth expansion for symmetric statistics *Ann. Statist.* 25, (1997), 851–896.
- [4] Bhattacharya, R.N. and Rao, Ranga R.: *Normal approximation and asymptotic expansions*, Robert E. Krieger Publishing company, Inc. 1986.
- [5] Bhattacharya, Rabi N. and Ghosh, J.K.: On the validity of the formal edgeworth expansion. *Ann. Statist.* 6, (1978), 434–451.

- [6] Bhattacharya, Rabi N. and Ghosh, J.K.: Correction to: On the validity of the formal Edgeworth expansion. *Ann. Statist.* 8, (1980), 1399.
- [7] Bickel, P. J.: Edgeworth expansions in nonparametric statistics. *Ann. Statist.* 2, (1974), 1–20.
- [8] Bickel, P. J., Götze, F. and van Zwet, W. R.: The Edgeworth expansion for U -statistics of degree two. *Ann. Statist.* 14, (1986), 1463–1484.
- [9] Bickel, P. J. and Robinson, J.: Edgeworth expansions and smoothness. *Ann. Probab.* 10, (1982), 500–503.
- [10] Bollobás Béla, *Combinatorics. Set systems, hypergraphs, families of vectors and combinatorial probability*, Cambridge Univ. Press, 1986.
- [11] Callaert, H., Janssen, P. and Veraverbeke N.: An Edgeworth expansion for U -statistics. *Ann. Statist.* 8, (1980), 299–312.
- [12] Chibisov, D.M.: Asymptotic expansion for the distribution of a statistic admitting a stochastic expansion. I. *Teor. Veroyatn. Primen.* 25, (1980), 745–757.
- [13] Dharmadhikari, S. W., Fabian, V. and Jogdeo, K.: Bounds on the moments of martingales *Ann. Math. Statist.* 39, (1968), 1719–1723.
- [14] Efron, B. and Stein, C.: The jackknife estimate of variance *Ann. Statist.* 9, (1981), 586–596.
- [15] Esseen, C.G.: Fourier analysis of distribution functions. A mathematical study of the Laplace-Gaussian law. *Acta Math.* 77, (1945), 1–125.
- [16] Götze, F.: Asymptotic expansions for bivariate von Mises functionals. *Z. Wahrsch. Verw. Gebiete* 50, (1979), 333–355.
- [17] Götze, F. and van Zwet W.R.: Edgeworth expansions for asymptotically linear statistics. Manuscript, 1992, 1–45.
- [18] Hall, P.: Edgeworth expansion for Student’s t statistic under minimal moment conditions. *Ann. Probab.* 15, (1987), 920–931.
- [19] Helmers, R., *Edgeworth expansions for linear combinations of order statistics*. Mathematical Centre Tracts, 105. Amsterdam CWI, 1982.
- [20] Hoeffding, W.: A class of statistics with asymptotically normal distribution. *Ann. Math. Statist.* 19, (1948), 293–325.
- [21] Ledoux, M. and Talagrand M., *Probability in Banach spaces. Isoperimetry and processes*. Springer-Verlag, Berlin, Heidelberg, 1991.
- [22] Petrov, V. V., *Sums of independent random variables*. Springer-Verlag, New York-Heidelberg, 1975.
- [23] Pfanzagl, J., *Asymptotic expansions for general statistical models. With the assistance of W. Wefelmeyer*. Lecture Notes in Statistics, 31. Springer-Verlag, Berlin, 1985.
- [24] Serfling, R. J., *Approximation theorems of mathematical statistics*. Wiley, 1980.
- [25] Yurinskii, V. V.: Exponential inequalities for sums of random vectors. *J. Multivar. Analysis* 6, (1976), 473–499.
- [26] van Zwet, W. R.: A Berry–Esseen bound for symmetric statistics. *Z. Wahrsch. Verw. Gebiete* 66, (1984), 425–440.

FACULTY OF MATHEMATICS, UNIVERSITY OF BIELEFELD, GERMANY
Email address: goetze@math.uni-bielefeld.de

DEPARTMENT OF MATHEMATICS, VILNIUS UNIVERSITY, LITHUANIA
Email address: mindaugas.bloznelis@mif.vu.lt