

# ON GRADED CLASSICAL 2-ABSORBING SECOND SUBMODULES OF GRADED MODULES OVER GRADED COMMUTATIVE RINGS

KHALDOUN AL-ZOUBI\* AND MARIAM AL-AZAIZEH

ABSTRACT. Let  $G$  be a group with identity  $e$ . Let  $R$  be a  $G$ -graded commutative ring and  $M$  a graded  $R$ -module. In this paper, we introduce the concept of graded classical and graded strongly classical 2-absorbing second submodules of graded modules over a graded commutative rings. A number of results concerning of these classes of graded submodules and their homogeneous components are given.

## 1. INTRODUCTION

Badawi in [19] introduced the concept of 2-absorbing ideals of commutative rings. Later on, Anderson and Badawi in [13] generalized the concept of 2-absorbing ideals of commutative rings to the concept of  $n$ -absorbing ideals of commutative rings for every positive integer  $n \geq 2$ . In light of [19] and [13], many authors studied the concept of 2-absorbing submodules and  $n$ -absorbing submodules. In [17], the authors introduced and studied the concepts of 2-absorbing and strongly 2-absorbing second submodules. In [23], the authors introduced and studied the concept of classical 2-absorbing submodules as a generalization of 2-absorbing submodules. Recently, H. Ansari-Toroghy and F. Farshadifar in [14] introduced and studied the concepts of classical and strongly classical 2-absorbing second submodules of modules over commutative rings.

The scope of this paper is devoted to the theory of graded modules over graded commutative rings. One use of rings and modules with gradings is in describing certain topics in algebraic geometry. Here, in particular, we are dealing with graded classical and graded strongly

---

2010 *Mathematics Subject Classification.* 13A02, 16W50.

*Key words and phrases.* graded classical 2-absorbing second submodules, graded strongly classical 2-absorbing second submodules, graded 2-absorbing second submodules.

\* Corresponding author.

classical 2-absorbing second submodules of graded modules over graded commutative rings.

Graded prime ideals of graded commutative rings have been introduced and studied in [10, 30-32]. S.E. Atani in [18] extended graded prime ideals to graded prime submodules. Several authors investigated properties of graded prime submodules, for examples see [1, 9, 11, 29]. The concept of graded 2-absorbing ideals, generalizations of graded prime ideals, were studied by K. Al-Zoubi and R. Abu-Dawwas, and other authors, (see [3, 25].) K. Al-Zoubi and R. Abu-Dawwas in [2] extended graded 2-absorbing ideals to graded 2-absorbing submodules. Later on, M. Hamoda and A. E. Ashour in [24] introduced the concept of graded  $n$ -absorbing submodules that is a generalization of the concept of graded prime ideals. The notion of graded classical prime submodules as a generalization of graded prime submodules was introduced in [21] and studied in [9, 6, 8, 12]. The notion of graded second submodules was introduced in [15] and studied in [7, 16, 20]. In [4], the authors introduced and studied the concept of graded classical 2-absorbing submodules as a generalization of graded 2-absorbing submodules.

Recently, K. Al-Zoubi and M. Al-Azaizeh in [5] introduced and studied the concepts of graded 2-absorbing and graded strongly 2-absorbing second submodules.

Here, we introduce the concept of graded classical (resp. graded strongly classical) 2-absorbing second submodules of graded modules over a commutative graded rings as a generalization of graded 2-absorbing (resp. graded strongly 2-absorbing) second submodules and investigate some properties of these classes of graded submodules.

## 2. PRELIMINARIES

**Convention.** Throughout this paper all rings are commutative with identity and all modules are unitary. First, we recall some basic properties of graded rings and modules which will be used in the sequel. We refer to [22], [26], [27] and [28] for these basic properties and more information on graded rings and modules. Let  $G$  be a multiplicative group and  $e$  denote the identity element of  $G$ . A ring  $R$  is called a graded ring (or  $G$ -graded ring) if there exist additive subgroups  $R_\alpha$  of  $R$  indexed by the elements  $\alpha \in G$  such that  $R = \bigoplus_{\alpha \in G} R_\alpha$  and  $R_\alpha R_\beta \subseteq R_{\alpha\beta}$  for all  $\alpha, \beta \in G$ . The elements of  $R_\alpha$  are called homogeneous of degree  $\alpha$  and all the homogeneous elements are denoted by  $h(R)$ , i.e.  $h(R) = \cup_{\alpha \in G} R_\alpha$ . If  $r \in R$ , then  $r$  can be written uniquely as  $\sum_{\alpha \in G} r_\alpha$ , where  $r_\alpha$  is called a homogeneous component of  $r$  in  $R_\alpha$ . Moreover,  $R_e$  is a subring of

$R$  and  $1 \in R_e$ . Let  $R = \bigoplus_{\alpha \in G} R_\alpha$  be a  $G$ -graded ring. An ideal  $I$  of  $R$  is said to be a graded ideal if  $I = \bigoplus_{\alpha \in G} (I \cap R_\alpha) := \bigoplus_{\alpha \in G} I_\alpha$ . Let  $R = \bigoplus_{\alpha \in G} R_\alpha$  be a  $G$ -graded ring. A Left  $R$ -module  $M$  is said to be a *graded  $R$ -module* (or  *$G$ -graded  $R$ -module*) if there exists a family of additive subgroups  $\{M_\alpha\}_{\alpha \in G}$  of  $M$  such that  $M = \bigoplus_{\alpha \in G} M_\alpha$  and  $R_\alpha M_\beta \subseteq M_{\alpha\beta}$  for all  $\alpha, \beta \in G$ . Here,  $R_\alpha M_\beta$  denotes the additive subgroup of  $M$  consisting of all finite sums of elements  $r_\alpha m_\beta$  with  $r_\alpha \in R_\alpha$  and  $m_\beta \in M_\beta$ . Also if an element of  $M$  belongs to  $\cup_{\alpha \in G} M_\alpha = h(M)$ , then it is called a homogeneous. Note that  $M_\alpha$  is an  $R_e$ -module for every  $\alpha \in G$ . So, if  $I = \bigoplus_{\alpha \in G} I_\alpha$  is a graded ideal of  $R$ , then  $I_\alpha$  is an  $R_e$ -module for every  $\alpha \in G$ . Let  $R = \bigoplus_{\alpha \in G} R_\alpha$  be a  $G$ -graded ring. A submodule  $N$  of  $M$  is said to be a *graded submodule of  $M$*  if  $N = \bigoplus_{\alpha \in G} (N \cap M_\alpha) := \bigoplus_{\alpha \in G} N_\alpha$ . In this case,  $N_\alpha$  is called the  $\alpha$ -component of  $N$ .

Let  $R$  be a  $G$ -graded ring and  $M$  a graded  $R$ -module. A proper graded submodule  $C$  of  $M$  is said to be a *completely graded irreducible* if  $C = \bigcap_{\alpha \in \Delta} C_\alpha$ , where  $\{C_\alpha\}_{\alpha \in \Delta}$  is a family of graded submodules of  $M$ , implies that  $C = C_\beta$  for some  $\beta \in \Delta$  (see [5].) A non-zero graded submodule  $S$  of  $M$  is said to be a *graded 2-absorbing second submodule of  $M$*  if whenever  $r, t \in h(R)$ ,  $C$  is a completely graded irreducible submodule of  $M$ , and  $rtS \subseteq C$ , then  $rS \subseteq C$  or  $tS \subseteq C$  or  $rt \in \text{Ann}_R(S)$  (see [5].) A non-zero graded submodule  $S$  of  $M$  is said to be a *graded strongly 2-absorbing second submodule of  $M$*  if whenever  $r, t \in h(R)$ ,  $C_1, C_2$  are completely graded irreducible submodules of  $M$ , and  $rtS \subseteq C_1 \cap C_2$ , then  $rS \subseteq C_1 \cap C_2$  or  $tS \subseteq C_1 \cap C_2$  or  $rt \in \text{Ann}_R(S)$  (see [5].) A proper graded submodule  $C$  of  $M$  is said to be a *graded classical 2-absorbing submodule of  $M$*  if  $C \neq M$ ; and whenever  $r, s, t \in h(R)$  and  $m \in h(M)$  with  $rstm \in C$ , then either  $rs m \in C$  or  $rt m \in C$  or  $st m \in C$  (see [4].)

### 3. GRADED CLASSICAL 2-ABSORBING SECOND SUBMODULES

**Definition 3.1.** Let  $R$  be a  $G$ -graded ring and  $M$  a graded  $R$ -module. A non-zero graded submodule  $C$  of  $M$  is said to be a *graded classical 2-absorbing second submodule of  $M$* , if whenever  $r_\alpha, s_\beta, t_\gamma \in h(R)$ ,  $U$  is a completely graded irreducible submodule of  $M$  and  $r_\alpha s_\beta t_\gamma C \subseteq U$ , then either  $r_\alpha s_\beta C \subseteq U$  or  $s_\beta t_\gamma C \subseteq U$  or  $r_\alpha t_\gamma C \subseteq U$ . We say that  $M$  is a *graded classical 2-absorbing second module* if  $M$  is a graded classical 2-absorbing second submodule of itself.

**Lemma 3.2.** Let  $R$  be a  $G$ -graded ring,  $M$  a graded  $R$ -module and  $C$  a graded classical 2-absorbing second submodule of  $M$ . Let  $I = \bigoplus_{\gamma \in G} I_\gamma$  be a graded ideal of  $R$ . Then for every  $r_\alpha, s_\beta \in h(R)$ ,  $\gamma \in G$  and

completely graded irreducible submodule  $U$  of  $M$  with  $r_\alpha s_\beta I_\gamma C \subseteq U$ , either  $r_\alpha I_\gamma C \subseteq U$  or  $s_\beta I_\gamma C \subseteq U$  or  $r_\alpha s_\beta C \subseteq U$ .

*Proof.* Let  $r_\alpha, s_\beta \in h(R)$ ,  $\gamma \in G$  and  $U$  be a completely graded irreducible submodule of  $M$  such that  $r_\alpha s_\beta I_\gamma C \subseteq U$ ,  $r_\alpha I_\gamma C \not\subseteq U$  and  $s_\beta I_\gamma C \not\subseteq U$ . We have to show that  $r_\alpha s_\beta C \subseteq U$ . By our assumption there exist  $i_\gamma, i'_\gamma \in I_\gamma$  such that  $r_\alpha i_\gamma C \not\subseteq U$  and  $s_\beta i'_\gamma C \not\subseteq U$ . As  $r_\alpha s_\beta i_\gamma C \subseteq U$  and  $C$  is a graded classical 2-absorbing second submodule, we have either  $r_\alpha s_\beta C \subseteq U$  or  $s_\beta i_\gamma C \subseteq U$ . Similarly, by  $r_\alpha s_\beta i'_\gamma C \subseteq U$ , we get either  $r_\alpha s_\beta C \subseteq U$  or  $r_\alpha i'_\gamma C \subseteq U$ . If  $r_\alpha s_\beta C \subseteq U$ , we are done. Suppose that  $r_\alpha s_\beta C \not\subseteq U$ , which implies that  $s_\beta i_\gamma C \subseteq U$  and  $r_\alpha i'_\gamma C \subseteq U$ . By  $(i_\gamma + i'_\gamma) \in I_\gamma$ , it follows that  $r_\alpha s_\beta (i_\gamma + i'_\gamma) C \subseteq U$ . Then either  $r_\alpha (i_\gamma + i'_\gamma) C \subseteq U$  or  $s_\beta (i_\gamma + i'_\gamma) C \subseteq U$  as  $C$  is a graded classical 2-absorbing second submodule of  $M$ . If  $r_\alpha (i_\gamma + i'_\gamma) C \subseteq U$ , then  $r_\alpha i_\gamma C \subseteq U$  since  $r_\alpha i'_\gamma C \subseteq U$ , which is a contradiction. Similarly, if  $s_\beta (i_\gamma + i'_\gamma) C \subseteq U$ , we get a contradiction. Thus  $r_\alpha s_\beta C \subseteq U$ .  $\square$

**Lemma 3.3.** *Let  $R$  be a  $G$ -graded ring,  $M$  a graded  $R$ -module and  $C$  a graded classical 2-absorbing second submodule of  $M$ . Let  $I = \bigoplus_{\beta \in G} I_\beta$  and  $J = \bigoplus_{\gamma \in G} J_\gamma$  be a graded ideals of  $R$ . Then for every  $r_\alpha \in h(R)$ ,  $\beta, \gamma \in G$  and completely graded irreducible submodule  $U$  of  $M$  with  $r_\alpha I_\beta J_\gamma C \subseteq U$ , either  $r_\alpha I_\beta C \subseteq U$  or  $r_\alpha J_\gamma C \subseteq U$  or  $I_\beta J_\gamma C \subseteq U$ .*

*Proof.* Let  $r_\alpha \in h(R)$ ,  $\beta, \gamma \in G$  and  $U$  be a completely graded irreducible submodule of  $M$  such that  $r_\alpha I_\beta J_\gamma C \subseteq U$ ,  $r_\alpha I_\beta C \not\subseteq U$  and  $r_\alpha J_\gamma C \not\subseteq U$ . We have to show that  $I_\beta J_\gamma C \subseteq U$ . Let  $i_\beta \in I_\beta$  and  $j_\gamma \in J_\gamma$ . By our assumption, there exist  $i'_\beta \in I_\beta$  and  $j'_\gamma \in J_\gamma$  such that  $r_\alpha i'_\beta C \not\subseteq U$  and  $r_\alpha j'_\gamma C \not\subseteq U$ . Since  $r_\alpha i'_\beta J_\gamma C \subseteq U$ ,  $r_\alpha i'_\beta C \not\subseteq U$  and  $r_\alpha J_\gamma C \not\subseteq U$ , by Lemma 3.2, we get  $i'_\beta J_\gamma C \subseteq U$ . Similarly, by  $r_\alpha j'_\gamma I_\beta C \subseteq U$ ,  $r_\alpha j'_\gamma C \not\subseteq U$  and  $r_\alpha I_\beta C \not\subseteq U$ , we get  $j'_\gamma I_\beta C \subseteq U$ . By  $(i_\beta + i'_\beta) \in I_\beta$  and  $(j_\gamma + j'_\gamma) \in J_\gamma$ , it follow that  $r_\alpha (i_\beta + i'_\beta)(j_\gamma + j'_\gamma) C \subseteq U$ . Since  $C$  is a graded classical 2-absorbing second submodule, we have either  $r_\alpha (i_\beta + i'_\beta) C \subseteq U$  or  $r_\alpha (j_\gamma + j'_\gamma) C \subseteq U$  or  $(i_\beta + i'_\beta)(j_\gamma + j'_\gamma) C \subseteq U$ . If  $r_\alpha (i_\beta + i'_\beta) C \subseteq U$ , then  $r_\alpha i_\beta C \not\subseteq U$  since  $r_\alpha i'_\beta C \not\subseteq U$ . Since  $r_\alpha i_\beta J_\gamma C \subseteq U$ ,  $r_\alpha J_\gamma C \not\subseteq U$  and  $r_\alpha i_\beta C \not\subseteq U$ , by Lemma 3.2, we get  $i_\beta J_\gamma C \subseteq U$  and hence  $i_\beta j_\gamma C \subseteq U$ . Similarly, if  $r_\alpha (j_\gamma + j'_\gamma) C \subseteq U$ , we conclude that  $i_\beta j_\gamma C \subseteq U$ . If  $(i_\beta + i'_\beta)(j_\gamma + j'_\gamma) C \subseteq U$ , then  $i_\beta j_\gamma C + i_\beta j'_\gamma C + i'_\beta j_\gamma C + i'_\beta j'_\gamma C \subseteq U$ , this implies that  $i_\beta j_\gamma C \subseteq U$ . Thus  $I_\beta J_\gamma C \subseteq U$ .  $\square$

The following theorem give us a characterization of graded classical 2-absorbing second submodule of a graded module.

**Theorem 3.4.** *Let  $R$  be a  $G$ -graded ring,  $M$  a graded  $R$ -module and  $C$  a non-zero graded submodule of  $M$ . Let  $I = \bigoplus_{\alpha \in G} I_\alpha$ ,  $J = \bigoplus_{\beta \in G} J_\beta$*

and  $K = \bigoplus_{\gamma \in G} K_\gamma$  be a graded ideals of  $R$ . Then the following statements are equivalent:

- (i)  $C$  is a graded classical 2-absorbing second submodule of  $M$ ;
- (ii) For every  $\alpha, \beta, \gamma \in G$  and completely graded irreducible submodule  $U$  of  $M$  with  $I_\alpha J_\beta K_\gamma C \subseteq U$ , either  $J_\beta K_\gamma C \subseteq U$  or  $I_\alpha J_\beta C \subseteq U$  or  $I_\alpha K_\gamma C \subseteq U$ .

*Proof.* (i)  $\Rightarrow$  (ii) Assume that  $C$  is a graded classical 2-absorbing second submodule of  $M$ . Let  $U$  be a completely graded irreducible submodule of  $M$  and  $\alpha, \beta, \gamma \in G$  such that  $I_\alpha J_\beta K_\gamma C \subseteq U$  and  $J_\beta K_\gamma C \not\subseteq U$ . Then by Lemma 3.3, for all  $i_\alpha \in I_\alpha$ , we have either  $i_\alpha J_\beta C \subseteq U$  or  $i_\alpha K_\gamma C \subseteq U$ . If  $i_\alpha J_\beta C \subseteq U$  for all  $i_\alpha \in I_\alpha$ , then  $I_\alpha J_\beta C \subseteq U$ , we are done. Similarly, if  $i_\alpha K_\gamma C \subseteq U$  for all  $i_\alpha \in I_\alpha$ , then  $I_\alpha K_\gamma C \subseteq U$ , we are done. Suppose that there exist  $i_\alpha, i'_\alpha \in I_\alpha$  such that  $i_\alpha J_\beta C \not\subseteq U$  and  $i'_\alpha K_\gamma C \not\subseteq U$ . Since  $i_\alpha J_\beta K_\gamma C \subseteq U$ ,  $i_\alpha J_\beta C \not\subseteq U$  and  $J_\beta K_\gamma C \not\subseteq U$ , by Lemma 3.3, we get  $i_\alpha K_\gamma C \subseteq U$ . Similarly, by  $i'_\alpha J_\beta K_\gamma C \subseteq U$ ,  $i'_\alpha K_\gamma C \not\subseteq U$  and  $J_\beta K_\gamma C \not\subseteq U$ , we get  $i'_\alpha J_\beta C \subseteq U$ . Since  $(i_\alpha + i'_\alpha) J_\beta K_\gamma C \subseteq U$  and  $J_\beta K_\gamma C \not\subseteq U$ , by Lemma 3.3, we get either  $(i_\alpha + i'_\alpha) J_\beta C \subseteq U$  or  $(i_\alpha + i'_\alpha) K_\gamma C \subseteq U$ . If  $(i_\alpha + i'_\alpha) J_\beta C \subseteq U$ , then  $i'_\alpha J_\beta C \subseteq U$  since  $i_\alpha J_\beta C \not\subseteq U$ , which is a contradiction. Similarly, if  $(i_\alpha + i'_\alpha) K_\gamma C \subseteq U$ , we get a contradiction. Therefore either  $I_\alpha J_\beta C \subseteq U$  or  $I_\alpha K_\gamma C \subseteq U$ .

(ii)  $\Rightarrow$  (i) Assume that (ii) holds. Let  $r_\alpha, s_\beta, t_\gamma \in h(R)$  and  $U$  be a completely graded irreducible submodule of  $M$  such that  $r_\alpha s_\beta t_\gamma C \subseteq U$ . Let  $I = r_\alpha R$ ,  $J = s_\beta R$  and  $K = t_\gamma R$  be a graded ideals of  $R$  generated by  $r_\alpha, s_\beta$  and  $t_\gamma$ , respectively. Then  $I_\alpha J_\beta K_\gamma C \subseteq U$ . By our assumption, we have either  $J_\beta K_\gamma C \subseteq U$  or  $I_\alpha J_\beta C \subseteq U$  or  $I_\alpha K_\gamma C \subseteq U$ . This yields that either  $s_\beta t_\gamma C \subseteq U$  or  $r_\alpha s_\beta C \subseteq U$  or  $r_\alpha t_\gamma C \subseteq U$ . Thus  $C$  is a graded classical 2-absorbing second submodule of  $M$ .  $\square$

**Corollary 3.5.** *Let  $R$  be a  $G$ -graded ring,  $M$  a graded  $R$ -module and  $C$  a graded classical 2-absorbing second submodule of  $M$  and  $I = \bigoplus_{\alpha \in G} I_\alpha$  be a graded ideal of  $R$ . Then for each  $\alpha \in G$ ,  $I_\alpha^n C = I_\alpha^{n+1} C$  for all  $n \geq 2$ .*

*Proof.* It is enough to show that  $I_\alpha^2 C = I_\alpha^3 C$ . It is clear that  $I_\alpha^3 C \subseteq I_\alpha^2 C$ . Let  $U$  be a completely graded irreducible submodule of  $M$  such that  $I_\alpha^3 C \subseteq U$ . By Theorem 3.4, we get  $I_\alpha^2 C \subseteq U$ . This yields that  $I_\alpha^2 C \subseteq I_\alpha^3 C$  by [5, Lemma 2.3]. Therefore  $I_\alpha^2 C = I_\alpha^3 C$ .  $\square$

Clearly every graded 2-absorbing second submodule is a graded classical 2-absorbing second submodule. The following example shows that the converse is not true in general.

**Example 3.6.** Let  $G = \mathbb{Z}_2$ , then  $R = \mathbb{Z}$  is a  $G$ -graded ring with  $R_0 = \mathbb{Z}$  and  $R_1 = \{0\}$ . Let  $M = \mathbb{Z}$  be a graded  $R$ -module with  $M_0 = \mathbb{Z}$  and

$M_1 = \{0\}$ . Now, consider the graded submodule  $N = 2\mathbb{Z}$ . Then  $N$  is a graded classical 2-absorbing second submodule of  $R$  which is not a graded 2-absorbing second submodule of  $M$ .

Let  $R$  be a  $G$ -graded ring,  $M$  a graded  $R$ -module and  $N$  a graded submodule of  $M$ . Then  $(N :_R M)$  is defined as  $(N :_R M) = \{r \in R \mid rM \subseteq N\}$ . It is shown in [18, Lemma 2.1] that if  $N$  is a graded submodule of  $M$ , then  $(N :_R M) = \{r \in R : rM \subseteq N\}$  is a graded ideal of  $R$ . The annihilator of  $M$  is defined as  $(0 :_R M)$  and is denoted by  $Ann_R(M)$ . Recall that a proper graded ideal  $I$  of a  $G$ -graded ring  $R$  is said to be a *graded 2-absorbing ideal of  $R$*  if whenever  $r, s, t \in h(R)$  with  $rst \in I$ , then  $rs \in I$  or  $rt \in I$  or  $st \in I$  (see [3].)

**Theorem 3.7.** *Let  $R$  be a  $G$ -graded ring,  $M$  a graded  $R$ -module and  $C$  a graded classical 2-absorbing second submodule of  $M$ . Then for each completely graded irreducible submodule  $U$  of  $M$  with  $C \not\subseteq U$ ,  $(U :_R C)$  is a graded 2-absorbing ideal of  $R$ .*

*Proof.* Let  $r_\alpha, s_\beta, t_\gamma \in h(R)$  such that  $r_\alpha s_\beta t_\gamma \in (U :_R C)$ . Then  $r_\alpha s_\beta t_\gamma C \subseteq U$ . Since  $C$  is a graded classical 2-absorbing second submodule, we have either  $s_\beta t_\gamma C \subseteq U$  or  $r_\alpha s_\beta C \subseteq U$  or  $r_\alpha t_\gamma C \subseteq U$ , it follows that either  $s_\beta t_\gamma \in (U :_R C)$  or  $r_\alpha s_\beta \in (U :_R C)$  or  $r_\alpha t_\gamma \in (U :_R C)$ . Thus  $(U :_R C)$  is a graded 2-absorbing ideal of  $R$ .  $\square$

**Theorem 3.8.** *Let  $R$  be a  $G$ -graded ring,  $M$  a graded  $R$ -module and  $C$  a graded classical 2-absorbing second submodule of  $M$ . Let  $I = \bigoplus_{\lambda \in G} I_\lambda$  be a graded ideal of  $R$  and  $\lambda \in G$  with  $I_\lambda \not\subseteq Ann_R(C)$ . Then  $I_\lambda C$  is a graded classical 2-absorbing second submodule of  $M$ .*

*Proof.* Since  $I_\lambda \not\subseteq Ann_R(C)$ ,  $I_\lambda C \neq 0$ . Now, let  $r_\alpha, s_\beta, t_\gamma \in h(R)$  and  $U$  be a completely graded irreducible submodule of  $M$  such that  $r_\alpha s_\beta t_\gamma I_\lambda C \subseteq U$ . By Lemma 3.2, we have either  $r_\alpha s_\beta I_\lambda C \subseteq U$  or  $t_\gamma I_\lambda C \subseteq U$  or  $r_\alpha s_\beta t_\gamma C \subseteq U$ . If  $r_\alpha s_\beta I_\lambda C \subseteq U$  or  $t_\gamma I_\lambda C \subseteq U$ , then we are done. If  $r_\alpha s_\beta t_\gamma C \subseteq U$ , then either  $s_\beta t_\gamma C \subseteq U$  or  $r_\alpha t_\gamma C \subseteq U$  or  $r_\alpha s_\beta C \subseteq U$  as  $C$  is a graded classical 2-absorbing second submodule, it follows that either  $s_\beta t_\gamma I_\lambda C \subseteq U$  or  $r_\alpha t_\gamma I_\lambda C \subseteq U$  or  $r_\alpha s_\beta I_\lambda C \subseteq U$ . Thus  $I_\lambda C$  is a graded classical 2-absorbing second submodule of  $M$ .  $\square$

Let  $R$  be a  $G$ -graded ring and  $M, M'$  graded  $R$ -modules. Let  $f : M \rightarrow M'$  be an  $R$ -module homomorphism. Then  $f$  is said to be a graded homomorphism if  $f(M_\alpha) \subseteq M'_\alpha$  for all  $\alpha \in G$ , (see [28].) The category of graded  $R$ -modules possesses direct sums, products injective and projective limits. A graded homomorphism that is an injective function will be referred to simply as a monomorphism and a graded homomorphism that is a surjective function will be called an epimorphism.

**Theorem 3.9.** *Let  $R$  be a  $G$ -graded ring and  $M_1, M_2$  be two graded  $R$ -modules. Let  $f : M_1 \rightarrow M_2$  be a graded monomorphism. Then we have the following:*

- (i) *If  $C_1$  is a graded classical 2-absorbing second submodule of  $M_1$ , then  $f(C_1)$  is a graded classical 2-absorbing second submodule of  $f(M_1)$ .*
- (ii) *If  $C_2$  is a graded classical 2-absorbing second submodule of  $f(M_1)$ , then  $f^{-1}(C_2)$  is a graded classical 2-absorbing second submodule of  $M_1$ .*

*Proof.* (i) Assume that  $C_1$  is a graded classical 2-absorbing second submodule of  $M_1$ . It is easy to see  $f(C_1) \neq 0$ . Now, Let  $r_\alpha, s_\beta, t_\gamma \in h(R)$  and  $U_2$  be a completely graded irreducible submodule of  $f(M_1)$  such that  $r_\alpha s_\beta t_\gamma f(C_1) \subseteq U_2$ . Consequently  $r_\alpha s_\beta t_\gamma C_1 \subseteq f^{-1}(U_2)$ . By [5, Lemma 2.11(ii)], we have  $f^{-1}(U_2)$  is a completely graded irreducible submodule of  $M_1$ . Then either  $r_\alpha s_\beta C_1 \subseteq f^{-1}(U_2)$  or  $s_\beta t_\gamma C_1 \subseteq f^{-1}(U_2)$  or  $r_\alpha t_\gamma C_1 \subseteq f^{-1}(U_2)$  as  $C_1$  is a graded classical 2-absorbing second submodule of  $M_1$ . If  $r_\alpha s_\beta C_1 \subseteq f^{-1}(U_2)$ , then  $r_\alpha s_\beta f(C_1) = f(r_\alpha s_\beta C_1) \subseteq f(f^{-1}(U_2)) = U_2 \cap f(M_1) = U_2$ . Similarly, if  $s_\beta t_\gamma C_1 \subseteq f^{-1}(U_2)$ , we get  $s_\beta t_\gamma f(C_1) \subseteq U_2$ , also if  $r_\alpha t_\gamma C_1 \subseteq f^{-1}(U_2)$ , we get  $r_\alpha t_\gamma f(C_1) \subseteq U_2$ . Thus  $f(C_1)$  is a graded classical 2-absorbing second submodule of  $f(M_1)$ .

(ii) Assume that  $C_2$  is a graded classical 2-absorbing second submodule of  $f(M_1)$ . It is easy to see  $f^{-1}(C_2) \neq 0$ . Now, let  $r_\alpha, s_\beta, t_\gamma \in h(R)$  and  $U_1$  be a completely graded irreducible submodule of  $M_1$  such that  $r_\alpha s_\beta t_\gamma f^{-1}(C_2) \subseteq U_1$ . Consequently  $r_\alpha s_\beta t_\gamma C_2 = r_\alpha s_\beta t_\gamma (f(M_1) \cap C_2) = r_\alpha s_\beta t_\gamma f f^{-1}(C_2) = f(r_\alpha s_\beta t_\gamma f^{-1}(C_2)) \subseteq f(U_1)$ . By [5, Lemma 2.11(i)], we have  $f(U_1)$  is a completely graded irreducible submodule of  $f(M_1)$ . Then either  $r_\alpha s_\beta C_2 \subseteq f(U_1)$  or  $s_\beta t_\gamma C_2 \subseteq f(U_1)$  or  $r_\alpha t_\gamma C_2 \subseteq f(U_1)$  as  $C_2$  is a graded classical 2-absorbing second submodule of  $f(M_1)$ . This yields that either  $r_\alpha s_\beta f^{-1}(C_2) \subseteq U_1$  or  $s_\beta t_\gamma f^{-1}(C_2) \subseteq U_1$  or  $r_\alpha t_\gamma f^{-1}(C_2) \subseteq U_1$ . Thus  $f^{-1}(C_2)$  is a graded classical 2-absorbing second submodule of  $M_1$ .  $\square$

#### 4. GRADED STRONGLY CLASSICAL 2-ABSORBING SECOND SUBMODULES

**Definition 4.1.** Let  $R$  be a  $G$ -graded ring and  $M$  a graded  $R$ -module. A non-zero graded submodule  $C$  of  $M$  is said to be a *graded strongly classical 2-absorbing second submodule of  $M$*  if whenever  $r_\alpha, s_\beta, t_\gamma \in h(R)$ ,  $U_1, U_2$  and  $U_3$  are completely graded irreducible submodules of  $M$ , and  $r_\alpha s_\beta t_\gamma C \subseteq U_1 \cap U_2 \cap U_3$ , then  $r_\alpha s_\beta C \subseteq U_1 \cap U_2 \cap U_3$  or  $s_\beta t_\gamma C \subseteq U_1 \cap U_2 \cap U_3$  or  $r_\alpha t_\gamma C \subseteq U_1 \cap U_2 \cap U_3$ . We say that  $M$  is a

graded strongly *classical 2-absorbing second module* if  $M$  is a graded strongly classical 2-absorbing second submodule of itself.

The following results give us a characterization of graded strongly classical 2-absorbing second submodule of a graded module.

**Theorem 4.2.** *Let  $R$  be a  $G$ -graded ring,  $M$  a graded  $R$ -module and  $C$  a non-zero graded submodule of  $M$ . Then the following statements are equivalent:*

- (i)  $C$  is a graded strongly classical 2-absorbing second submodule of  $M$ ;
- (ii) For every  $r_\alpha, s_\beta, t_\gamma \in h(R)$  and every graded submodule  $N$  of  $M$  with  $r_\alpha s_\beta t_\gamma C \subseteq N$ , then either  $r_\alpha s_\beta C \subseteq N$  or  $s_\beta t_\gamma C \subseteq N$  or  $r_\alpha t_\gamma C \subseteq N$ ;
- (iii) For every  $r_\alpha, s_\beta, t_\gamma \in h(R)$ , either  $r_\alpha s_\beta t_\gamma C = r_\alpha s_\beta C$  or  $r_\alpha s_\beta t_\gamma C = s_\beta t_\gamma C$  or  $r_\alpha s_\beta t_\gamma C = r_\alpha t_\gamma C$ .

*Proof.* (i)  $\Rightarrow$  (ii) Assume that  $C$  is a graded strongly classical 2-absorbing second submodule of  $M$ . Let  $r_\alpha, s_\beta, t_\gamma \in h(R)$  and  $N$  be a graded submodule of  $M$  such that  $r_\alpha s_\beta t_\gamma C \subseteq N$ . Assume on the contrary that  $r_\alpha s_\beta C \not\subseteq N$ ,  $s_\beta t_\gamma C \not\subseteq N$  and  $r_\alpha t_\gamma C \not\subseteq N$ . Then there exist completely graded irreducible submodules  $U_1, U_2$  and  $U_3$  of  $M$  such that  $N \subseteq U_1, N \subseteq U_2$  and  $N \subseteq U_3$  but  $r_\alpha s_\beta C \not\subseteq U_1, s_\beta t_\gamma C \not\subseteq U_2$  and  $r_\alpha t_\gamma C \not\subseteq U_3$  by [5, Lemma 2.3]. Hence  $r_\alpha s_\beta t_\gamma C \subseteq U_1 \cap U_2 \cap U_3$ . Since  $C$  is a graded strongly classical 2-absorbing second submodule of  $M$ , we get either  $r_\alpha s_\beta C \subseteq U_1 \cap U_2 \cap U_3$  or  $s_\beta t_\gamma C \subseteq U_1 \cap U_2 \cap U_3$  or  $r_\alpha t_\gamma C \subseteq U_1 \cap U_2 \cap U_3$  which are contradiction.

(ii)  $\Rightarrow$  (iii) Assume (ii) is hold. Let  $r_\alpha, s_\beta, t_\gamma \in h(R)$ . Then  $r_\alpha s_\beta t_\gamma C$  is a graded submodule of  $M$ . Since  $r_\alpha s_\beta t_\gamma C \subseteq r_\alpha s_\beta t_\gamma C$ , by (ii), we have either  $r_\alpha s_\beta C \subseteq r_\alpha s_\beta t_\gamma C$  or  $s_\beta t_\gamma C \subseteq r_\alpha s_\beta t_\gamma C$  or  $r_\alpha t_\gamma C \subseteq r_\alpha s_\beta t_\gamma C$ . This yields that either  $r_\alpha s_\beta t_\gamma C = r_\alpha s_\beta C$  or  $r_\alpha s_\beta t_\gamma C = s_\beta t_\gamma C$  or  $r_\alpha s_\beta t_\gamma C = r_\alpha t_\gamma C$ .

(iii)  $\Rightarrow$  (i) Trivial. □

**Lemma 4.3.** *Let  $R$  be a  $G$ -graded ring,  $M$  a graded  $R$ -module and  $C$  a graded strongly classical 2-absorbing second submodule of  $M$ . Let  $I = \bigoplus_{\gamma \in G} I_\gamma$  be a graded ideal of  $R$ . Then for every  $r_\alpha, s_\beta \in h(R)$ ,  $\gamma \in G$  and graded submodule  $N$  of  $M$  with  $r_\alpha s_\beta I_\gamma C \subseteq N$ , either  $r_\alpha I_\gamma C \subseteq N$  or  $s_\beta I_\gamma C \subseteq N$  or  $r_\alpha s_\beta C \subseteq N$ .*

*Proof.* By using Theorem 4.2 the proof is similar to the poof of Lemma 3.2, so we omit it. □

**Lemma 4.4.** *Let  $R$  be a  $G$ -graded ring,  $M$  a graded  $R$ -module and  $C$  a graded strongly classical 2-absorbing second submodule of  $M$ . Let*



$I = \bigoplus_{\beta \in G} I_\beta$  and  $J = \bigoplus_{\gamma \in G} J_\gamma$  be a graded ideals of  $R$ . Then for every  $r_\alpha \in h(R)$ ,  $\beta, \gamma \in G$  and graded submodule  $N$  of  $M$  with  $r_\alpha I_\beta J_\gamma C \subseteq N$ , either  $r_\alpha I_\beta C \subseteq N$  or  $r_\alpha J_\gamma C \subseteq N$  or  $I_\beta J_\gamma C \subseteq N$ .

*Proof.* By using Theorem 4.2 and Lemma 4.3, the proof is similar to the poof of Lemma 3.3, so we omit it.  $\square$

**Theorem 4.5.** *Let  $R$  be a  $G$ -graded ring,  $M$  a graded  $R$ -module and  $C$  a non-zero graded submodule of  $M$ . Let  $I = \bigoplus_{\alpha \in G} I_\alpha$ ,  $J = \bigoplus_{\beta \in G} J_\beta$  and  $K = \bigoplus_{\gamma \in G} K_\gamma$  be a graded ideals of  $R$ . Then the following statements are equivalent:*

- (i)  $C$  is a graded strongly classical 2-absorbing second submodule of  $M$ .
- (ii) For every  $\alpha, \beta, \gamma \in G$  and graded submodule  $N$  of  $M$  with  $I_\alpha J_\beta K_\gamma C \subseteq N$ , either  $J_\beta K_\gamma C \subseteq N$  or  $I_\alpha J_\beta C \subseteq N$  or  $I_\alpha K_\gamma C \subseteq N$ .

*Proof.* By using Theorem 4.2 and Lemma 4.4, the proof is similar to the proof of Theorem 3.4, so we omit it.  $\square$

**Theorem 4.6.** *Let  $R$  be a  $G$ -graded ring,  $M$  a graded  $R$ -module and  $C$  a non-zero graded submodule of  $M$  and  $N$  a graded submodule of  $M$  with  $C \not\subseteq N$ . Then  $C$  is a graded strongly classical 2-absorbing second submodule of  $M$  if and only if  $(N :_R C)$  is a graded 2-absorbing ideal of  $R$ .*

*Proof.* Assume that  $C$  is a graded strongly classical 2-absorbing second submodule of  $M$ . Since  $C \not\subseteq N$ ,  $(N :_R C) \neq R$ . Now let  $r_\alpha, s_\beta, t_\gamma \in h(R)$  such that  $r_\alpha s_\beta t_\gamma \in (N :_R C)$ . Then  $r_\alpha s_\beta t_\gamma C \subseteq N$ . By Theorem 4.2, we have either  $r_\alpha s_\beta C \subseteq N$  or  $s_\beta t_\gamma C \subseteq N$  or  $r_\alpha t_\gamma C \subseteq N$ . Thus either  $r_\alpha s_\beta \in (N :_R C)$  or  $s_\beta t_\gamma \in (N :_R C)$  or  $r_\alpha t_\gamma \in (N :_R C)$ . Therefore  $(N :_R C)$  is a graded 2-absorbing ideal of  $R$ . Conversely, Let  $r_\alpha, s_\beta, t_\gamma \in h(R)$  and  $N$  be a graded submodule of  $M$  with  $r_\alpha s_\beta t_\gamma C \subseteq N$ . If  $C \subseteq N$ , then we are done. Assume that  $C \not\subseteq N$ . By our assumption, we have  $(N :_R C)$  is a graded 2-absorbing ideal of  $R$ . Since  $r_\alpha s_\beta t_\gamma \in (N :_R C)$ , we conclude that either  $r_\alpha s_\beta C \subseteq N$  or  $s_\beta t_\gamma C \subseteq N$  or  $r_\alpha t_\gamma C \subseteq N$ . Hence  $C$  is a graded strongly classical 2-absorbing second submodule of  $M$  by Theorem 4.2.  $\square$

Clearly every graded strongly 2-absorbing second submodule is a graded strongly classical 2-absorbing second submodule. The following example shows that the converse is not true in general.

**Example 4.7.** Let  $G = \mathbb{Z}_2$ , then  $R = \mathbb{Z}$  is a  $G$ -graded ring with  $R_0 = \mathbb{Z}$  and  $R_1 = \{0\}$ . Let  $M = \mathbb{Z}_6 \times \mathbb{Q}$  be a graded  $R$ -module with

$M_0 = \mathbb{Z}_6 \times \mathbb{Q}$  and  $M_1 = \{(0, 0)\}$ . Then  $M$  is a graded strongly classical 2-absorbing second module which is not a graded strongly 2-absorbing second module.

**Theorem 4.8.** *Let  $R$  be a  $G$ -graded ring and  $M_1, M_2$  be two graded  $R$ -modules. Let  $f : M_1 \rightarrow M_2$  be a graded monomorphism. Then we have the following.*

- (i) *If  $C_1$  is a graded strongly classical 2-absorbing second submodule of  $M_1$ , then  $f(C_1)$  is a graded strongly classical 2-absorbing second submodule of  $M_2$ .*
- (ii) *If  $C_2$  is a graded strongly classical 2-absorbing second submodule of  $f(M_1)$ , then  $f^{-1}(C_2)$  is a graded strongly classical 2-absorbing second submodule of  $M_1$ .*

*Proof.* (i) Assume that  $C_1$  is a graded strongly classical 2-absorbing second submodule of  $M$ . It is easy to see  $f(C_1) \neq 0$ . Let  $r_\alpha, s_\beta, t_\gamma \in h(R)$ . By Theorem 4.2, we have either  $r_\alpha s_\beta t_\gamma C_1 = r_\alpha s_\beta C_1$  or  $r_\alpha s_\beta t_\gamma C_1 = s_\beta t_\gamma C_1$  or  $r_\alpha s_\beta t_\gamma C_1 = r_\alpha t_\gamma C_1$ . We can assume that  $r_\alpha s_\beta t_\gamma C_1 = r_\alpha s_\beta C_1$ . Then  $r_\alpha s_\beta t_\gamma f(C_1) = f(r_\alpha s_\beta t_\gamma C_1) = f(r_\alpha s_\beta C_1) = r_\alpha s_\beta f(C_1)$ . Hence  $f(C_1)$  is a graded strongly classical 2-absorbing second submodule of  $M_2$  by Theorem 4.2.

(ii) Use the technique of Theorem 3.9(ii), and apply Theorem 4.2.  $\square$

**Definition 4.9.** Let  $S$  be a non-zero graded submodule of a graded  $R$ -module  $M$ . We say that  $S$  is a *graded weakly second submodule* of  $M$  if  $r_\alpha s_\beta S \subseteq N$ , where  $r_\alpha, s_\beta \in h(R)$  and  $N$  is a graded submodule of  $M$ , then either  $r_\alpha S \subseteq N$  or  $s_\beta S \subseteq N$ .

**Lemma 4.10.** *Let  $R$  be a  $G$ -graded ring and  $M$  be a graded  $R$ -module. If  $S_1$  and  $S_2$  are graded weakly second submodules of  $M$ , then  $S = S_1 + S_2$  is a graded strongly classical 2-absorbing second submodule of  $M$ .*

*Proof.* Assume that  $S_1$  and  $S_2$  are graded weakly second submodules of  $M$  and  $S = S_1 + S_2$ . Let  $r_\alpha, s_\beta, t_\gamma \in h(R)$ . As  $r_\alpha s_\beta t_\gamma S_1 \subseteq r_\alpha s_\beta t_\gamma S_1$  and  $S_1$  is a graded weakly second submodule, then either  $r_\alpha S_1 \subseteq r_\alpha s_\beta t_\gamma S_1$  or  $s_\beta S_1 \subseteq r_\alpha s_\beta t_\gamma S_1$  or  $t_\gamma S_1 \subseteq r_\alpha s_\beta t_\gamma S_1$ . This yields that either  $r_\alpha S_1 = r_\alpha s_\beta t_\gamma S_1$  or  $s_\beta S_1 = r_\alpha s_\beta t_\gamma S_1$  or  $t_\gamma S_1 = r_\alpha s_\beta t_\gamma S_1$ . Similarly, we have either  $r_\alpha S_2 = r_\alpha s_\beta t_\gamma S_2$  or  $s_\beta S_2 = r_\alpha s_\beta t_\gamma S_2$  or  $t_\gamma S_2 = r_\alpha s_\beta t_\gamma S_2$ . We may assume that  $r_\alpha S_1 = r_\alpha s_\beta t_\gamma S_1$ . Likewise assume that  $s_\beta S_2 = r_\alpha s_\beta t_\gamma S_2$ . Then  $r_\alpha s_\beta t_\gamma S = r_\alpha s_\beta t_\gamma S_1 + r_\alpha s_\beta t_\gamma S_2 = r_\alpha S_1 + s_\beta S_2 = r_\alpha s_\beta S = r_\alpha s_\beta S_1 + r_\alpha s_\beta S_2 = r_\alpha s_\beta S$ . By Theorem 4.2, we have  $S$  is a graded strongly classical 2-absorbing second submodules of  $M$ .  $\square$

Let  $R_i$  be a graded commutative ring with identity and  $M_i$  be a graded  $R_i$ -module, for  $i = 1, 2$ . Let  $R = R_1 \times R_2$ . Then  $M = M_1 \times M_2$  is a graded  $R$ -module and each graded submodule  $C$  of  $M$  is of the form  $C = C_1 \times C_2$  for some graded submodules  $C_1$  of  $M_1$  and  $C_2$  of  $M_2$  (see [28].)

**Theorem 4.11.** *Let  $R = R_1 \times R_2$  be a  $G$ -graded ring and  $M = M_1 \times M_2$  a graded  $R$ -module where  $M_1$  is a graded  $R_1$ -module and  $M_2$  is a graded  $R_2$ -module. Let  $C_1$  and  $C_2$  be a non-zero graded submodules of  $M_1$  and  $M_2$ , respectively.*

- (i)  $C = C_1 \times 0$  is a graded strongly classical 2-absorbing second submodule of  $M$  if and only if  $C_1$  is a graded strongly classical 2-absorbing second submodule of  $M_1$ .
- (ii)  $C = 0 \times C_2$  is a graded strongly classical 2-absorbing second submodule of  $M$  if and only if  $C_2$  is a graded strongly classical 2-absorbing second submodule of  $M_2$ .
- (iii)  $C = C_1 \times C_2$  is a graded strongly classical 2-absorbing submodule of  $M$  if and only if  $C_1$  and  $C_2$  are graded weakly second submodules of  $M_1$  and  $M_2$ , respectively.

*Proof.* (i) Assume that  $C = C_1 \times 0$  is a graded strongly classical 2-absorbing second submodule of  $M$ . From our assumption,  $C$  is a non-zero graded submodule, so  $C_1 \neq 0$ . Set  $M' = M_1 \times 0$ . One can see that  $C = C_1 \times 0$  is a graded strongly classical 2-absorbing second submodule of  $M'$ . Also, observe that  $M' \simeq M_1$  and  $C \simeq C_1$ . Hence  $C_1$  is a graded strongly classical 2-absorbing second submodule of  $M_1$  by Theorem 4.8. The converse is clear.

(ii) It can be easily verified similar to (i).

(iii) Assume that  $C = C_1 \times C_2$  is a graded strongly classical 2-absorbing submodule of  $M$ . We have to show that  $C_1$  is a graded weakly second submodule of  $M_1$ . Since  $C_2 \neq 0$ , by [5, Lemma 2.3], there exist a completely graded irreducible submodule  $U_2$  of  $M_2$  such that  $C_2 \not\subseteq U_2$ . Let  $r_\alpha, s_\beta \in h(R)$ ,  $N$  be a graded submodule of  $M_1$  and  $r_\alpha s_\beta C_1 \subseteq N$ . Thus  $(r_\alpha, 1)(s_\beta, 1)(1, 0)(C_1 \times C_2) = r_\alpha s_\beta C_1 \times 0 \subseteq N \times U_2$ . As  $C = C_1 \times C_2$  is a graded strongly classical 2-absorbing submodule of  $M$ , by Theorem 4.2, we have either  $(r_\alpha, 1)(s_\beta, 1)(C_1 \times C_2) = r_\alpha s_\beta C_1 \times C_2 \subseteq N \times U_2$  or  $(s_\beta, 1)(1, 0)(C_1 \times C_2) = s_\beta C_1 \times 0 \subseteq N \times U_2$  or  $(r_\alpha, 1)(1, 0)(C_1 \times C_2) = r_\alpha C_1 \times 0 \subseteq N \times U_2$ .

If  $r_\alpha s_\beta C_1 \times C_2 \subseteq N \times U_2$ , then  $C_2 \subseteq U_2$  a contradiction. Which implies either  $r_\alpha C_1 \subseteq N$  or  $s_\beta C_1 \subseteq N$ . Thus  $C_1$  is a graded weakly second submodule of  $M_1$ . Similarly, we can show that  $C_2$  is a graded weakly second submodule of  $M_2$ .

Conversely, assume that  $C_1$  and  $C_2$  are graded weakly second submodules of  $M_1$  and  $M_2$ , respectively. Then it is clear that  $C_1 \times 0$ ,  $0 \times C_2$  are graded weakly second submodules of  $M$ . By Lemma 4.10, we have  $C = C_1 \times 0 + 0 \times C_2$  is a graded strongly classical 2-absorbing submodule of  $M$ .  $\square$

### Acknowledgments

The authors wish to thank sincerely the referees for their valuable comments and suggestions.

**Conflict of Interest Statement:** On behalf of all authors, the corresponding author states that there is no conflict of interest.

### REFERENCES

- [1] K. Al-Zoubi and R. Abu-Dawwas, *On graded quasi-prime Submodules*, Kyungpook Math. J., **55** (2) (2015), 259-266.
- [2] K. Al-Zoubi and R. Abu-Dawwas, *On graded 2-absorbing and weakly graded 2-absorbing submodules*, J. Math. Sci. Adv. Appl. **28** (2014), 45-60.
- [3] K. Al-Zoubi, R. Abu-Dawwas and S. Çeken, *On graded 2-absorbing and graded weakly 2-absorbing ideals*, Hacet. J. Math. Stat., **48** (3) (2019), 724-731.
- [4] K. Al-Zoubi and M. Al-Azaizeh, *On graded classical 2-absorbing submodules of graded modules over graded commutative rings*, Rend. Istit. Mat. Univ. Trieste., **50** (2018), 1-10.
- [5] K. Al-Zoubi and M. Al-Azaizeh, *On graded 2-absorbing second submodules of graded modules over graded commutative rings*, submitted.
- [6] K. Al-Zoubi and M. Al-Dolat, *On graded classical primary submodules*, Adv. Pure Appl. Math., **7** (2) (2016), 93-96.
- [7] K. Al-Zoubi and A. Al-Qderat, *Some properties of graded comultiplication modules*, Open Math., **15** (2017), 187-192.
- [8] K. Al-Zoubi and M. Jaradat, *On the graded classical prime spectrum of a graded module*, Proyecciones Journal of Mathematics, **37** (3) (2018), 519-533..
- [9] K. Al-Zoubi, M. Jaradat and R. Abu-Dawwas, *On graded classical prime and graded prime submodules*, Bull. Iranian Math. Soc., **41** (1) (2015), 217-225.
- [10] K. Al-Zoubi and F. Qarqaz, *An intersection condition for graded prime ideals*, Boll Unione Mat. Ital., **11** (4) (2018), 483-488.
- [11] K. Al-Zoubi and F. Qarqaz, *An intersection condition for graded prime submodules in Gr-multiplication modules*, Math. Reports, **20** (3) (2018), 329-336.
- [12] R. Abu-Dawwas and K. Al-Zoubi, *On graded weakly classical prime submodules*, Iran. J. Math. Sci. Inform. **12** (1) (2017), 153-161.
- [13] D. F. Anderson and A. Badawi, *On n-absorbing ideals of commutative rings*, Comm. Algebra **39** (2011), 1646-1672.
- [14] H. Ansari-Toroghy and F. Farshadifar, *Classical and strongly classical 2-absorbing second submodules*, Commun. Fac. Sci. Univ. Ank. Ser. A1 Math. Stat., **69** (1) (2020), 123-136.
- [15] H. Ansari-Toroghy and F. Farshadifar, *Graded comultiplication modules*, Chiang Mai J. Sci., **38** (3) (2011), 311-320.
- [16] H. Ansari-Toroghy and F. Farshadifar, *On graded second modules*, Tamkang J. Math., **43** (4) (2012), 499-505.

- [17] H. Ansari-Toroghy and F. Farshadifar, *Some generalizations of second submodules*, Palestine Journal of Mathematics, **8** (2) (2019), 159–168.
- [18] S.E. Atani, *On graded prime submodules*, Chiang Mai. J. Sci., **33** (1) (2006), 3–7.
- [19] A. Badawi, *On 2-absorbing ideals of commutative rings*, Bull. Austral. Math. Soc. **75** (2007), 417–429.
- [20] S. Çeken and M. Alkan, *On graded second and coprimary modules and graded secondary representations*, Bull. Malays. Math. Sci. Soc., **38** (4) (2015), 1317–1330.
- [21] A. Y. Darani and S. Motmaen, *Zariski topology on the spectrum of graded classical prime submodules*, Appl. Gen. Topol. **14** (2) (2013), 159–169.
- [22] R. Hazrat, *Graded Rings and Graded Grothendieck Groups*, Cambridge University Press, Cambridge, 2016.
- [23] H. Mostafanasab, U. Tekir, and K.H. Oral, *classical 2-absorbing submodules of modules over commutative rings*, Eur. J. Pure Appl. Math., **8** (3) (2015), 417–430.
- [24] H. Mohammad and A. Ashour, *On graded  $n$ -absorbing submodules*, Le Matematiche **70** (2) (2015), 243–254.
- [25] S. R. Naghani and H. F. Moghimi, *On Graded 2-Absorbing and Graded Weakly 2-Absorbing Ideals of a Commutative Ring*, Cankaya University Journal of Science and Engineering, **13** (2) (2016), 011–017.
- [26] C. Nastasescu and F. Van Oystaeyen, *Graded and filtered rings and modules*, Lecture notes in mathematics 758, Berlin-New York: Springer-Verlag, 1982.
- [27] C. Nastasescu, F. Van Oystaeyen, *Graded Ring Theory*, Mathematical Library 28, North Holland, Amsterdam, 1982.
- [28] C. Nastasescu and F. Van Oystaeyen, *Methods of Graded Rings*, LNM 1836. Berlin-Heidelberg: Springer-Verlag, 2004.
- [29] K. H. Oral, U. Tekir and A. G. Agargun, *On graded prime and primary submodules*, Turk. J. Math., **35** (2011), 159–167.
- [30] M. Refai and K. Al-Zoubi, *On graded primary ideals*, Turk. J. Math. **28** (3) (2004), 217–229.
- [31] M. Refai, M. Hailat and S. Obiedat, *Graded radicals on graded prime spectra*, Far East J. of Math. Sci., part I (2000), 59–73.
- [32] R.N. Uregen, U. Tekir and K.H. Oral, *On the union of graded prime ideals*, Open Phys., **14** (2016), 114–118.

*Khaldoun Al-Zoubi, Department of Mathematics and Statistics, Jordan University of Science and Technology, P.O.Box 3030, Irbid 22110, Jordan.*

*Email address: kfzoubi@just.edu.jo*

*Mariam Al-Azaizeh, Department of Mathematics, University of Jordan, Amman, Jordan..*

*Email address: maalazaizeh15@sci.just.edu.jo*