

On Graded ϕ -Prime Submodules

Azzh Saad ALSHEHRY, Malik BATAINEH, and Rashid ABU-DAWWAS

ABSTRACT. Let R be a graded commutative ring with non-zero unity 1 and M be a graded unitary R -module. Let $GS(M)$ be the set of all graded R -submodules of M and $\phi : GS(M) \rightarrow GS(M) \cup \{\emptyset\}$ be a function. A proper graded R -submodule K of M is said to be a graded ϕ -prime R -submodule of M if whenever r is a homogeneous element of R and m is a homogeneous element of M such that $rm \in K - \phi(K)$, then either $m \in K$ or $r \in (K :_R M)$. If $\phi(K) = \emptyset$ for all $K \in GS(M)$, then a graded ϕ -prime submodule is exactly a graded prime submodule. If $\phi(K) = \{0\}$ for all $K \in GS(M)$, then a graded ϕ -prime submodule is exactly a graded weakly prime submodule. Several properties of graded ϕ -prime submodules have been investigated.

1. Introduction

Throughout this article, G will be a group with identity e and R a commutative ring with nonzero unity 1. Then R is said to be G -graded if $R = \bigoplus_{g \in G} R_g$ with

$R_g R_h \subseteq R_{gh}$ for all $g, h \in G$ where R_g is an additive subgroup of R for all $g \in G$. The elements of R_g are called homogeneous of degree g . If $x \in R$, then x can be written uniquely as $\sum_{g \in G} x_g$, where x_g is the component of x in R_g . It is known that

R_e is a subring of R and $1 \in R_e$. The set of all homogeneous elements of R is $h(R) = \bigcup_{g \in G} R_g$. Assume that M is a left unitary R -module. Then M is said to

be G -graded if $M = \bigoplus_{g \in G} M_g$ with $R_g M_h \subseteq M_{gh}$ for all $g, h \in G$ where M_g is an

additive subgroup of M for all $g \in G$. The elements of M_g are called homogeneous of degree g . It is clear that M_g is an R_e -submodule of M for all $g \in G$. If $x \in M$, then x can be written uniquely as $\sum_{g \in G} x_g$, where x_g is the component of x in M_g .

The set of all homogeneous elements of M is $h(M) = \bigcup_{g \in G} M_g$. Let K be an R -

submodule of a graded R -module M . Then K is said to be graded R -submodule if $K = \bigoplus_{g \in G} (K \cap M_g)$, i.e., for $x \in K$, $x = \sum_{g \in G} x_g$ where $x_g \in K$ for all $g \in G$.

An R -submodule of a graded R -module need not be graded. For more details and terminology, see [5, 6].

2010 *Mathematics Subject Classification.* Primary 16W50; Secondary 13A02.

Key words and phrases. Graded prime submodules; graded ϕ -prime submodules.

LEMMA 1.1. ([4], Lemma 2.1) *Let R be a graded ring and M be a graded R -module.*

- (1) *If I and J are graded ideals of R , then $I + J$ and $I \cap J$ are graded ideals of R .*
- (2) *If N and K are graded R -submodules of M , then $N + K$ and $N \cap K$ are graded R -submodules of M .*
- (3) *If N is a graded R -submodule of M , $r \in h(R)$, $x \in h(M)$ and I is a graded ideal of R , then Rx , IN and rN are graded R -submodules of M . Moreover, $(N :_R M) = \{r \in R : rM \subseteq N\}$ is a graded ideal of R .*

Let I be a proper graded ideal of R . Then the graded radical of I is $Grad(I)$, and is defined to be the set of all $r \in R$ such that for each $g \in G$, there exists a positive integer n_g for which $r_g^{n_g} \in I$. One can see that if $r \in h(R)$, then $r \in Grad(I)$ if and only if $r^n \in I$ for some positive integer n . In fact, $Grad(I)$ is a graded ideal of R , see [8]. A graded R -submodule K of M is called a graded radical R -submodule of M if $Grad((K :_R M)) = (K :_R M)$.

Graded prime ideals play a fundamental role in graded ring theory. One of the natural generalizations of graded prime ideals which have attracted the interest of several authors in the last two decades is the concept of graded prime submodules, see for example [1, 3, 4, 7]. A proper graded R -submodule K of M is said to be a graded prime R -submodule of M if whenever $r \in h(R)$ and $m \in h(M)$ such that $rm \in K$, then either $m \in K$ or $r \in (K :_R M)$. It is known that if K is a graded prime R -submodule of M , then $(K :_R M)$ is a graded prime ideal of R . A proper graded R -submodule K of M is said to be a graded weakly prime R -submodule of M if whenever $r \in h(R)$ and $m \in h(M)$ such that $0 \neq rm \in K$, then either $m \in K$ or $r \in (K :_R M)$. A proper graded R -submodule K of M is said to be a graded almost prime R -submodule of M if whenever $r \in h(R)$ and $m \in h(M)$ such that $rm \in K - (K :_R M)K$, then either $m \in K$ or $r \in (K :_R M)$. So, every graded prime submodule is a graded weakly prime submodule, and every graded weakly prime submodule is a graded almost prime submodule. Let $GS(M)$ be the set of all graded R -submodules of M and $\phi : GS(M) \rightarrow GS(M) \cup \{\emptyset\}$ be a function. A proper graded R -submodule K of M is said to be a graded ϕ -prime R -submodule of M if whenever $r \in h(R)$ and $m \in h(M)$ such that $rm \in K - \phi(K)$, then either $m \in K$ or $r \in (K :_R M)$. Since $K - \phi(K) = K - (K \cap \phi(P))$, so without loss of generality, throughout this article, we will consider $\phi(K) \subseteq K$. Let M be a G -graded R -module, $g \in G$ and K be a graded R -submodule of M such that $K_g \neq M_g$. Then K is said to be a $g - \phi$ -prime R -submodule of M if whenever $r \in R_e$ and $m \in M_g$ such that $rm \in K - \phi(K)$, then either $m \in K$ or $r \in (K :_R M)$. Throughout this article, we use the following functions:

$$\begin{aligned} \phi_\emptyset(K) &= \emptyset \text{ for all } K \in GS(M), \\ \phi_0(K) &= \{0\} \text{ for all } K \in GS(M), \\ \phi_1(K) &= (K :_R M)K \text{ for all } K \in GS(M), \\ \phi_2(K) &= (K :_R M)^2 K \text{ for all } K \in GS(M) \text{ and} \\ \phi_w(K) &= \bigcap_{i=1}^{\infty} (K :_R M)^i K \text{ for all } K \in GS(M). \end{aligned}$$

Clearly, graded ϕ_\emptyset -prime, ϕ_0 -prime and ϕ_1 -prime submodules are graded prime, graded weakly prime and graded almost prime submodules respectively. Obviously, for any graded submodule and every positive integer k , we have the following implications:

graded prime submodule \Rightarrow graded ϕ_w -prime submodule \Rightarrow graded ϕ_k -prime submodule \Rightarrow graded ϕ_{k-1} -prime submodule.

For functions $\phi, \varphi : GS(M) \rightarrow GS(M) \cup \{\emptyset\}$, we write $\phi \leq \varphi$ if $\phi(K) \subseteq \varphi(K)$ for all $K \in GS(M)$. So, if $\phi \leq \varphi$, then every graded ϕ -prime submodule is graded φ -prime.

Among several results, we proved that if K is a g - ϕ -prime R -submodule of M such that $(K :_{R_e} M)K_g \not\subseteq \phi(K)$, then K is a g -prime R -submodule of M (Theorem 2.3). We showed that if K is a g - ϕ -prime R -submodule of M which is not g -prime and $\phi(K) \subseteq (K :_{R_e} M)^2 K_g$, then $\phi(K) = (K :_{R_e} M)^i K_g$ for all $i \geq 1$ (Corollary 2.5). We proved that if $0 \neq m \in M_g$ such that $R_e m \neq M_g$, $(0 :_R m) = \{0\}$ and Rm is not a g -prime R -submodule of M , then Rm is not a $g - \phi_1$ -prime R -submodule of M (Theorem 2.6). We showed that if $r \in R_e$ such that $rM_g \neq M_g$ and $(0 :_M r) \subseteq rM$, then rM is a $g - \phi_1$ -prime R -submodule of M if and only if it is a g -prime R -submodule of M (Theorem 2.8). In Theorem 2.9, we introduced a characterization for graded $g - \phi$ -prime R -submodules. In Theorem 2.10, we study graded ϕ -prime R -submodules over graded quotient R -modules. In Theorem 2.11, we examine graded ϕ -prime R -submodules over multiplicative subsets of $h(R)$. Finally, we proposed an interesting question (Question 2.12). Most of the results in this article are inspired from [10].

2. Graded ϕ -Prime Submodules

In this section, we introduce and study the concept of graded ϕ -prime submodules.

DEFINITION 2.1. *Let M be a G -graded R -module and $\phi : GS(M) \rightarrow GS(M) \cup \{\emptyset\}$ be a function.*

- (1) *A proper graded R -submodule K of M is said to be a graded ϕ -prime R -submodule of M if whenever $r \in h(R)$ and $m \in h(M)$ such that $rm \in K - \phi(K)$, then either $m \in K$ or $r \in (K :_R M)$.*
- (2) *Let K be a graded R -submodule of M and $g \in G$ such that $K_g \neq M_g$. Then K is said to be a $g - \phi$ -prime R -submodule of M if whenever $r \in R_e$ and $m \in M_g$ such that $rm \in K - \phi(K)$, then either $m \in K$ or $r \in (K :_R M)$.*

The concept of g -prime ideals have been introduced and investigated in [2]. Let P be a graded ideal of a G -graded ring R and $g \in G$ such that $P_g \neq R_g$. Then P is called a g -prime ideal of R if whenever $x, y \in R_g$ such that $xy \in P$, then either $x \in P$ or $y \in P$. Motivated by g -prime ideals, we introduce the following definitions:

DEFINITION 2.2. *Let M be a G -graded R -module, K be a graded R -submodule of M and $g \in G$ such that $K_g \neq M_g$. Then*

- (1) *K is said to a g -prime R -submodule of M if whenever $r \in R_e$ and $m \in M_g$ such that $rm \in K$, then either $m \in K$ or $r \in (K :_R M)$.*
- (2) *K is said to a g -weakly prime R -submodule of M if whenever $r \in R_e$ and $m \in M_g$ such that $0 \neq rm \in K$, then either $m \in K$ or $r \in (K :_R M)$.*

THEOREM 2.3. *Let M be a G -graded R -module, $g \in G$ and K be a $g - \phi$ -prime R -submodule of M . If $(K :_{R_e} M)K_g \not\subseteq \phi(K)$, then K is a g -prime R -submodule of M .*

PROOF. Let $r \in R_e$ and $m \in M_g$ such that $rm \in K$. If $rm \notin \phi(K)$, then since K is $g - \phi$ -prime, we have $r \in (K :_R M)$ or $m \in K$. Suppose that $rm \in \phi(K)$. If $rK_g \not\subseteq \phi(K)$, then there exists $p \in K_g$ such that $rp \notin \phi(K)$, and then $r(m+p) \in K - \phi(K)$. Therefore, $r \in (K :_R M)$ or $m+p \in K$, and hence $r \in (K :_R M)$ or $m \in K$. So, we may assume that $rK_g \subseteq \phi(K)$. If $(K :_{R_e} M)m \not\subseteq \phi(K)$, then there exists $u \in (K :_{R_e} M)$ such that $um \notin \phi(K)$, and then $(r+u)m \in K - \phi(K)$. Since K is a $g - \phi$ -prime submodule, we have $r+u \in (K :_R M)$ or $m \in K$. So, $r \in (K :_R M)$ or $m \in K$. Therefore, we assume that $(K :_{R_e} M)m \subseteq \phi(K)$. Now, since $(K :_{R_e} M)K_g \not\subseteq \phi(K)$, there exist $s \in (K :_{R_e} M)$ and $t \in K_g$ such that $st \notin \phi(K)$. So, $(r+s)(m+t) \in K - \phi(K)$, and hence $r+s \in (K :_R M)$ or $m+t \in K$. Therefore, $r \in (K :_R M)$ or $m \in K$. \square

COROLLARY 2.4. *Let M be a G -graded R -module, $g \in G$ and K be a g -weakly prime R -submodule of M . If $(K :_{R_e} M)K_g \neq \{0\}$, then K is a g -prime R -submodule of M .*

PROOF. Apply Theorem 2.3 by $\phi = \phi_0$. \square

COROLLARY 2.5. *Let M be a G -graded R -module, $g \in G$ and K be a $g - \phi$ -prime R -submodule of M which is not g -prime. If $\phi(K) \subseteq (K :_{R_e} M)^2 K_g$, then $\phi(K) = (K :_{R_e} M)^i K_g$ for all $i \geq 1$.*

PROOF. Since K is not a g -prime R -submodule of M , we have by Theorem 2.3, $(K :_{R_e} M)K_g \subseteq \phi(K) \subseteq (K :_{R_e} M)^2 K_g \subseteq (K :_{R_e} M)K_g$, which implies that $\phi(K) = (K :_{R_e} M)K_g = (K :_{R_e} M)^2 K_g$. Hence, $\phi(K) = (K :_{R_e} M)^i K_g$ for all $i \geq 1$. \square

THEOREM 2.6. *Let M be a G -graded R -module, $g \in G$ and $0 \neq m \in M_g$ such that $R_e m \neq M_g$ and $(0 :_R m) = \{0\}$. If Rm is not a g -prime R -submodule of M , then Rm is not a $g - \phi_1$ -prime R -submodule of M .*

PROOF. Since Rm is not a g -prime R -submodule of M , there exist $a \in R_e$ and $y \in M_g$ such that $a \notin (Rm :_R M)$, $y \notin Rm$ and $ay \in Rm$. If $ay \notin (Rm :_R M)m$, then Rm is not a $g - \phi_1$ -prime R -submodule of M . Let $ay \in (Rm :_R M)m$. Then $y+m \notin Rm$ and $a(y+m) \in Rm$. If $a(y+m) \notin (Rm :_R M)m$, then Rm is not a $g - \phi_1$ -prime R -submodule of M . Let $a(y+m) \in (Rm :_R M)m$, then $am \in (Rm :_R M)m$, which gives that $am = rm$ for some $r \in (Rm :_R M)$. Since $(0 :_R m) = \{0\}$, it gives that $a = r \in (Rm :_R M)$, which is a contradiction. \square

COROLLARY 2.7. *Let M be a G -graded R -module, $g \in G$ and m be a non-zero element of M_g such that $(0 :_R m) = \{0\}$ and $R_e m \neq M_g$. Then Rm is a g -prime R -submodule of M if and only if Rm is a $g - \phi_1$ -prime R -submodule of M .*

THEOREM 2.8. *Let M be a G -graded R -module, $r \in R_e$ and $g \in G$ such that $rM_g \neq M_g$. If $(0 :_M r) \subseteq rM$, then rM is a $g - \phi_1$ -prime R -submodule of M if and only if it is a g -prime R -submodule of M .*

PROOF. Suppose that rM is a $g - \phi_1$ -prime R -submodule of M . Let $b \in R_e$ and $m \in M_g$ such that $bm \in rM$. If $bm \notin (rM :_R M)rM$, then $b \in (rM :_R M)$ or $m \in rM$, since rM is a $g - \phi_1$ -prime submodule. Suppose that $bm \in (rM :_R M)rM$.

Now, $(b+r)m \in rM$. If $(b+r)m \notin (rM :_R M)rM$, then since rM is a $g - \phi_1$ -prime submodule, $b+r \in (rM :_R M)$ or $m \in rM$, which gives that $b \in (rM :_R M)$ or $m \in rM$. Assume that $(b+r)m \in (rM :_R M)rM$. Then $bm \in (rM :_R M)rM$ gives that $rm \in (rM :_R M)rM$. Hence, there exists $y \in (rM :_R M)M$ such that $rm = ry$ and so $m - y \in (0 :_M r)$. This gives that $m \in (rM :_R M)M + (0 :_M r) \subseteq rM + (0 :_M r) \subseteq rM$. The converse is clear. \square

THEOREM 2.9. *Let M be a G -graded R -module, $g \in G$ and K be a graded R -submodule of M such that $K_g \neq M_g$. Then the following statements are equivalent:*

- (1) K is a $g - \phi$ -prime R -submodule of M .
- (2) For $m \in M_g - K$, $(K :_{R_e} m) = (K :_{R_e} M) \cup (\phi(K) :_{R_e} m)$.
- (3) For $m \in M_g - K$, $(K :_{R_e} m) = (K :_{R_e} M)$ or $(K :_{R_e} m) = (\phi(K) :_{R_e} m)$.
- (4) For any ideal I of R_e and any graded R -submodule N of M , if $IN_g \subseteq K$ and $IN_g \not\subseteq \phi(K)$, then $I \subseteq (K :_R M)$ or $N_g \subseteq K$.

PROOF. (1) \Rightarrow (2) : Let $m \in M_g - K$ and $r \in (K :_{R_e} m) - (\phi(K) :_{R_e} m)$. Then $rm \in K - \phi(K)$. Since K is a $g - \phi$ -prime R -submodule of M , $r \in (K :_{R_e} M)$. As we may assume that $\phi(K) \subseteq K$, the other inclusion always holds.

(2) \Rightarrow (3) : It is known that if a subgroup is the union of two subgroups, then it is equal to one of them.

(3) \Rightarrow (4) : Let I be an ideal of R_e and N be a graded R -submodule of M such that $IN_g \subseteq K$. Suppose that $I \not\subseteq (K :_R M)$ and $N_g \not\subseteq K$. We show that $IN_g \subseteq \phi(K)$. Let $r \in I$ and $m \in N_g$. If $r \notin (K :_{R_e} M)$, then since $rm \in K$, we have $(K :_{R_e} m) \neq (K :_{R_e} M)$. Hence, by our assumption, $(K :_{R_e} m) = (\phi(K) :_{R_e} m)$. So, $rm \in \phi(K)$. Now, assume that $r \in I \cap (K :_{R_e} M)$. Let $u \in I - (K :_{R_e} M)$. Then $r+u \in I - (K :_{R_e} M)$. So, by the first case, for each $m \in N_g$ we have $um \in \phi(K)$ and $(r+u)m \in \phi(K)$. This gives that $rm \in \phi(K)$. Thus in any case $rm \in \phi(K)$. Therefore, $IN_g \subseteq \phi(K)$.

(4) \Rightarrow (1) : Let $r \in R_e$ and $m \in M_g$ such that $rm \in K - \phi(K)$. Suppose that $I = R_e r$ and $N = Rm$. Then I is an ideal of R_e and N is a graded R -submodule of M such that $IN_g = R_e r (Rm)_g = R_e r R_e m_g = R_e r R_e m = R_e R_e r m \subseteq R_e K \subseteq RK \subseteq K$ and $IN_g \not\subseteq \phi(K)$. By our assumption, $I \subseteq (K :_R M)$ or $N_g \subseteq K$, and then $r \in (K :_R M)$ or $m \in N \cap M_g = N_g \subseteq K$. \square

Let M be a G -graded R -module and K be a graded R -submodule of M . Then M/K is a G -graded R -module by $(M/K)_g = (M_g + K)/K$ for all $g \in G$. Let $\phi : GS(M) \rightarrow GS(M) \cup \{\emptyset\}$ be a function. Define $\phi_K : GS(M/K) \rightarrow GS(M/K) \cup \{\emptyset\}$ by $\phi_K(N/K) = (\phi(N) + K)/K$ for $N \supseteq K$ and $\phi_K(N/K) = \emptyset$ for $\phi(N) = \emptyset$.

THEOREM 2.10. *Let M be a graded R -module and K be a graded R -submodule of M . If N is a graded ϕ -prime R -submodule of M and $K \subseteq N$, then N/K is a graded ϕ_K -prime R -submodule of M/K .*

PROOF. By ([9], Lemma 3.2), N/K is a graded R -submodule of M/K . Let $r \in h(R)$ and $m+K \in h(M/K)$ such that $r(m+K) = rm+K \in N/K - \phi_K(N/K)$. Then $m \in h(M)$ such that $rm \in N - (\phi(N) + K)$, and then $rm \in N - \phi(N)$. Since N is graded ϕ -prime, $r \in (N :_R M)$ or $m \in N$, and then $r \in (N/K :_R M/K)$ or $m \in N$. Therefore, N/K is a graded ϕ_K -prime R -submodule of M/K . \square

Let M be a G -graded R -module and $S \subseteq h(R)$ be a multiplicative set. Then $S^{-1}M$ is a G -graded $S^{-1}R$ -module with $(S^{-1}M)_g = \{\frac{m}{s}, m \in M_h, s \in S \cap R_{hg^{-1}}\}$ for all $g \in G$, and $(S^{-1}R)_g = \{\frac{a}{s}, a \in R_h, s \in S \cap R_{hg^{-1}}\}$ for all $g \in G$. If K is a graded R -submodule of M , then $S^{-1}K$ is a graded $S^{-1}R$ -submodule of $S^{-1}M$. It is well known that there is a one-to-one correspondence between the set of all graded prime R -submodules K of M with $(K :_R M) \cap S = \emptyset$ and the set of all graded prime $S^{-1}R$ -submodules $S^{-1}K$ of $S^{-1}M$. For a graded R -submodule K of M , $K(S) = \{m \in M : \text{there exists } s \in S \text{ such that } sm \in K\}$ is a graded R -submodule of M containing K and $S^{-1}(K(S)) = S^{-1}K$. Let $\phi : GS(M) \rightarrow GS(M) \cup \{\emptyset\}$ be a function. Define $(S^{-1}\phi) : GS(S^{-1}M) \rightarrow GS(S^{-1}M) \cup \{\emptyset\}$ by $(S^{-1}\phi)(S^{-1}K) = S^{-1}(\phi(K(S)))$ if $\phi(K(S)) \neq \emptyset$ and $(S^{-1}\phi)(S^{-1}K) = \emptyset$ if $\phi(K(S)) = \emptyset$. Note that, $(S^{-1}\phi_\emptyset) = \phi_\emptyset$ and $(S^{-1}\phi_0) = \phi_0$.

THEOREM 2.11. *Let M be a G -graded R -module and K be a graded ϕ -prime R -submodule of M . Suppose that S is a multiplicative subset of $h(R)$. If $S^{-1}K \neq S^{-1}M$ and $S^{-1}(\phi(K)) \subseteq (S^{-1}\phi)(S^{-1}K)$, then $S^{-1}K$ is a graded $(S^{-1}\phi)$ -prime $S^{-1}R$ -submodule of $S^{-1}M$. Moreover, if $g \in G$ such that $(S^{-1}K) \cap M_g \neq S^{-1}(\phi(K))$, then $K(S) \cap M_g = K$.*

PROOF. Let $r/s \in h(S^{-1}R)$ and $m/t \in h(S^{-1}M)$ such that $(r/s)(m/t) \in S^{-1}K - (S^{-1}\phi)(S^{-1}K)$. Then $rm/st \in S^{-1}K - S^{-1}(\phi(K))$, and then there exists $u \in S$ such that $urm \in K - \phi(K)$ (note that, for each $v \in S$, $vrn \notin \phi(K)$). Since K is graded ϕ -prime and $(K :_R M) \cap S = \emptyset$, it gives that $rm \in K - \phi(K)$ and so $r \in (K :_R M)$ or $m \in K$, and then $r/s \in S^{-1}(K :_R M) \subseteq (S^{-1}K :_{S^{-1}R} S^{-1}M)$ or $m/t \in S^{-1}K$. Hence, $S^{-1}K$ is a graded $(S^{-1}\phi)$ -prime $S^{-1}R$ -submodule of $S^{-1}M$. Moreover, suppose that $(S^{-1}K) \cap M_g \neq S^{-1}(\phi(K))$ for some $g \in G$ and $m \in K(S) \cap M_g$. Then there exists $s \in S$ such that $sm \in K$. If $sm \notin \phi(K)$, then $m \in K$. If $sm \in \phi(K)$, then $m \in \phi(K)(S)$. So, $K(S) \cap M_g = K \cup (\phi(K)(S))$. Hence, either $K(S) \cap M_g = K$ or $K(S) \cap M_g = (\phi(K)(S))$. If $K(S) \cap M_g = (\phi(K)(S))$, then we have $(S^{-1}K) \cap M_g = (S^{-1}K(S)) \cap M_g = S^{-1}(\phi(K)(S)) = S^{-1}(\phi(K))$, which is not the case. So, $K(S) \cap M_g = K$. \square

QUESTION 2.12. *Let M be a graded R -module, S be a multiplicative subset of $h(R)$ and K be a graded R -submodule of M . If $S^{-1}K$ is a graded $S^{-1}\phi$ -prime $S^{-1}R$ -submodule of $S^{-1}M$, then clearly, $(K :_R M) \cap S = \emptyset$. In general, under what conditions K will be a graded ϕ -prime R -submodule of M ? Even in the cases $\phi = \phi_0$, $\phi = \phi_1$ and $\phi = \phi_2$, we could not answer this question.*

Acknowledgement

This research was funded by the Deanship of Scientific Research at Princess Nourah bint Abdulrahman University through the Fast-track Research Funding Program.

References

- [1] R. Abu-Dawwas, K. Al-Zoubi and M. Bataineh, Prime submodules of graded modules, *Proyecciones Journal of Mathematics*, 31 (4) (2012), 355-361.
- [2] R. Abu-Dawwas, E. Yıldız, Ü. Tekir and S. Koç, On graded 1-absorbing prime ideals, accepted in the *Sao Paulo Journal of Mathematical Sciences*, to appear.
- [3] S. E. Atani, On graded prime submodules, *Chiang Mai Journal of Science*, 33 (1) (2006), 3-7.
- [4] F. Farzalipour, P. Ghiasvand, On the union of graded prime submodules, *Thai Journal of Mathematics*, 9 (1) (2011), 49-55.

- [5] R. Hazrat, Graded rings and graded Grothendieck groups, Cambridge University press, 2016.
- [6] C. Nastasescu and F. Oystaeyen, Methods of graded rings, Lecture Notes in Mathematics, 1836, Springer-Verlag, Berlin, 2004.
- [7] K. H. Oral, Ü. Tekir and A. G. Agargün, On graded prime and primary submodules, Turkish Journal of Mathematics, 35 (2) (2011), 159-167.
- [8] M. Refai, M. Hailat and S. Obiedat, Graded radicals and graded prime spectra, Far East Journal of Mathematical Sciences, (2000), 59-73.
- [9] H. Saber, T. Alraqad and R. Abu-Dawwas, On graded s -prime submodules, AIMS Mathematics, 6 (3) (2020), 2510-2524.
- [10] N. Zamani, ϕ -Prime submodules, Glasgow Mathematical Journal, 52 (2010), 253-259.

DEPARTMENT OF MATHEMATICAL SCIENCES, FACULTY OF SCIENCES, PRINCESS NOURAH BINT ABDULRAHMAN UNIVERSITY, P.O. BOX 84428, RIYADH 11671, SAUDI ARABIA.

Email address: asalshihry@pnu.edu.sa

DEPARTMENT OF MATHEMATICS AND STATISTICS, JORDAN UNIVERSITY OF SCIENCE AND TECHNOLOGY, IRBID, JORDAN

Email address: msbataineh@just.edu.jo

DEPARTMENT OF MATHEMATICS, YARMOUK UNIVERSITY, IRBID, JORDAN

Email address: rrashid@yu.edu.jo