

Efficient Data-Driven Optimization with Noisy Data

Bart P.G. Van Parys*¹

¹*Sloan School of Management, MIT*

March 14, 2022

Abstract

Classical Kullback-Leibler or entropic distances have recently been shown [35] to enjoy certain desirable statistical properties in the context of decision-making with noiseless data. However, a major criticism of such distances is that they may result in fragile decisions in the presence of a distributional shift between the training and out-of-sample data. Instead, we study here data-driven prediction problems with data which is corrupted by noise. We derive efficient data-driven formulations in this noisy regime and indicate that they enjoy an entropic optimal transport interpretation. Finally, we show that these efficient robust formulations are tractable in several interesting settings by exploiting a classical representation result by Strassen [34].

1 Introduction

Let \mathcal{P} be a family of probability measures over a space Ξ and let P be an unknown probability measure in this family. Many problems in the machine learning community attempt at heart to learn the unknown data generating probability measure from a finite collection of independent data observations

$$\xi_i \sim P \quad \forall i \in [1, \dots, N]. \quad (1)$$

The most obvious application of this class of learning problems is perhaps density estimation [33]. Depending on whether the family \mathcal{P} is the entire probability simplex $\mathcal{P}(\Xi)$ over Ξ or merely a parametrized subset such methods can be classified as nonparametric or parametric, respectively. That is not to say that all machine learning problems with the data observation model (1) are density estimation problems in disguise. Often, one is not interested in the probability measure P *per se* but rather only a certain statistic. In predictive problems, in which the data consists of both dependent and independent variables, typically only the conditional expectation of the dependent variable given the independent variables is of interest. In prescriptive problems the problem of immediate concern is to find

*vanparys@mit.edu

an $0 < \epsilon$ -suboptimal¹ solution to a stochastic optimization problem, i.e.,

$$z(P) \in \arg_{z \in Z} \inf_{\epsilon} \{ \mathbf{E}_P [\ell(z, \xi)] = \int_{\Xi} \ell(z, \xi) dP(\xi) \}. \quad (2)$$

A wide spectrum of decision problems can be cast as instances of problem (2). Shapiro et al. [32] point out, that (2) can be viewed as the first stage of a two-stage stochastic program, where the loss function $\ell : Z \times \Xi \rightarrow \mathbb{R}$ embodies the optimal value of a subordinate second-stage problem. Alternatively, problem (2) may also be interpreted as a generic learning problem in the spirit of statistical learning theory. Rather than learning the unknown probability measure P , the primary objective in data-driven decision-making is to learn the cost function and perhaps even better an ϵ -suboptimal decision to (2) directly from data.

In this paper we will denote the observational model described in Equation (1) as “noiseless”. That is, the learner has access to uncorrupted independent samples from the probability measure of interest P . Clearly, given only finite amount of data the learner can not expect to learn the data generating probability measure exactly even in this noiseless regime. In case the probability measure P is only known to belong to the probability simplex $\mathcal{P}(\Xi)$, one reasonable substitute for P could be its empirically observed counterpart denoted here as $P_N := \sum_{i=1}^N \delta_{\xi_i} / N$. If on the other hand some prior information is available in the sense that the probability measure P is known to belong to a subset $\mathcal{P} \subset \mathcal{P}(\Xi)$, a maximum likelihood estimate [18] is often used instead. In the machine learning and robust optimization community such point estimates are widely known to be problematic when used naively in subsequent analysis. In particular, it is widely established both empirically as well as in theory that a sample average formulation

$$z(P_N) \in \arg_{z \in Z} \inf_{\epsilon} \mathbf{E}_{P_N} [\ell(z, \xi)] \quad (3)$$

which substitutes P with a mere point estimate P_N tends to disappoint ($\mathbf{E}_P [\ell(z(P_N), \xi)] > \mathbf{E}_{P_N} [\ell(z(P_N), \xi)]$) out of sample. That is, the actual cost ($\mathbf{E}_P [\ell(z(P_N), \xi)]$) observed out of sample breaks the predicted cost ($\mathbf{E}_{P_N} [\ell(z(P_N), \xi)]$) of the data-driven decision $z(P_N)$. This adversarial phenomenon is well known colloquially as the “Optimizer’s Curse” [22] and is akin to the overfitting phenomenon in the context of prediction problems. Such adversarial phenomena related to over-calibration to observed data but poor performance on out-of-sample data can be attributed primarily to the treatment of mere point estimates as exact substitutes for the unknown probability measure.

Ambiguity sets consisting of all probability measures sufficiently compatible with the observed data can offer a better alternative to simple point estimates. As the data observations are here independent and identically distributed, their order is irrelevant, and ambiguity sets $\mathcal{A}_N(P_N) \subseteq \mathcal{P}$ can be made functions of the empirical probability measure P_N rather than the data itself. A large line of work in the robust optimization community, see [27] and references therein, focuses consequently on data-driven formulations of the form

$$z_{\mathcal{A}}(P_N) \in \arg_{z \in Z} \inf_{\epsilon} \sup \{ \mathbf{E}_P [\ell(z, \xi)] : P \in \mathcal{A}_N(P_N) \} \quad (4)$$

¹Here $\arg_{z \in Z} \inf_{\epsilon} \mathbf{E}_P [\ell(z, \xi)]$ with $\epsilon > 0$ corresponds to the set $\{z \in Z : \mathbf{E}_P [\ell(z, \xi)] < \inf_{z \in Z} \mathbf{E}_P [\ell(z, \xi)] + \epsilon\}$.

which can be thought of as robust counterparts to the nominal sample average formulation stated in Equation (3) and where by convention consider the supremum to take the value $-\infty$ in case its feasible set $\mathcal{A}_N(P_N)$ is empty. The recent uptick in popularity of such robust formulations is in no small part due to the fact that they are often just as tractable and typically enjoy superior statistical properties than their nominal counterpart. Much early work [12, 37, 36] focused on ambiguity sets consisting of probability measures sharing certain given moments. More recent approaches [7] however consider ambiguity sets $\mathcal{A}_N(P_N) = \{P \in \mathcal{P} : D(P_N, P) \leq r_N\}$ which are based on some statistical distance $D : \mathcal{P}(\Xi) \times \mathcal{P}(\Xi) \rightarrow \mathbb{R} \cup \{+\infty\}^2$. Such ambiguity sets are interpretable as the set of probability measures sufficiently close to the empirical probability measure P_N . Two qualitatively different distances have recently positioned themselves as the front runners for data-driven decision-making and are now briefly discussed.

The optimal transport distance between a measure μ on Ξ and a measure ν on another set Ξ' can be defined as

$$W_0(\mu, \nu) := \inf_{T \in \mathcal{T}(\mu, \nu)} \int_{\Xi \times \Xi'} d(\xi, \xi') dT(\xi, \xi') \quad (5)$$

for a given the cost function $c : \Xi \times \Xi' \mapsto \mathbb{R} \cup \{+\infty\}$. Here $\mathcal{T}(\mu, \nu) := \{T \in \mathcal{P}(\Xi \times \Xi') : \Pi_{\Xi'} T = \mu, \Pi_{\Xi} T = \nu\}$ is the so called transport polytope and consists of all transport measures with given marginal measures μ and ν . Assume for a moment that $\Xi = \Xi' = \mathbb{R}^m$ and $d(\xi, \xi') = \|\xi - \xi'\|_2$. Then, the optimal transport distance coincides with the classical Wasserstein distance between probability measures [30]. Optimal transport distances [17, 38] have received a lot of attention both in the context of prescriptive analytics as well as in the machine learning community at large [20, 25]. In the context of prescriptive analytics such distances have become very popular after the seminal work [17] pointed out that the resulting robust formulation $z_{W,r}(P_N) \in \arg \inf_{z \in Z} \sup \{\mathbf{E}_P[\ell(z, \xi)] : W_0(P_N, P) \leq r\}$ need not be intractable and enjoys strong out-of-sample guarantees. For $r < 1$ and $2 \leq \dim(\Xi) < \infty$ Kuhn et al. [20] points out that we have for any P with $A = \mathbf{E}_P[\exp(\|\xi\|_2^a)] < \infty$ for some $a > 1$ that

$$\limsup_{N \rightarrow \infty} \frac{1}{N} \log \text{Prob}_P \left[\mathbf{E}_P[\ell(z_{W,r}(P_N), \xi)] > \sup \left\{ \mathbf{E}_Q[\ell(z_{W,r}(P_N), \xi)] : W_0(P_N, Q) \leq (r/C)^{\frac{1}{\dim(\Xi)}} \right\} \right] \leq r \quad (6)$$

where C is a known positive constant depending only on a , A and $\dim(\Xi)$. In other words, the probability of being disappointed (the actual cost $\mathbf{E}_P[\ell(z_{W,r}(P_N), \xi)]$ of the decision $z_{W,r}(P_N)$ exceeds the predicted worst-case cost $\sup \{\mathbf{E}_Q[\ell(z_{W,r}(P_N), \xi)] : W_0(P_N, Q) \leq r\}$) decays exponentially fast in the number of samples.

Optimal transport distances are however computationally quite challenging as even determining the optimal transport distance between two given measures requires the solution of the associated linear optimization problem in Equation (5). In fact, optimal transport distances are often computed numerically, c.f., [10], by considering their embedding in the larger family of entropic optimal transport distances

$$W_\epsilon(\mu, \nu) := \inf_{T \in \mathcal{T}(\mu, \nu)} \int_{\Xi \times \Xi'} d(\xi, \xi') dT(\xi, \xi') + \epsilon \text{KL}(T, \Pi_{\Xi} T \otimes \Pi_{\Xi'} T) \quad (7)$$

as the limit $\epsilon \downarrow 0$ where we denote the entropic divergence between two finite measures μ and ν on the same space

²In this paper we will use *distance* in a rather informal manner as indeed the distances we will encounter are neither symmetric nor satisfy a triangular inequality.

U as

$$\text{KL}(\mu, \nu) = \begin{cases} \int \log \left(\frac{d\mu}{d\nu}(u) \right) d\mu(u) - \int d\mu(u) + \int d\nu(u) & \text{if } \mu \ll \nu, \\ +\infty & \text{otherwise} \end{cases}$$

where the random variable $d\mu/d\nu : U \rightarrow \mathbb{R}$ denotes the Radon-Nikodym derivative between μ and ν .

The previously discussed entropic divergence is a particular member of the class of convex f -divergences which like the Wasserstein distance are well known [19] to yield tractable robust formulations. Moreover, when the loss function $\ell(z, \xi)$ is continuous on the compact set $Z \times \Xi$ the associated robust prescriptive formulation (4) can be specialized to $z_{\text{KL},r}(P_N) \in \min_{z \in Z} \max \{ \mathbf{E}_P [\ell(z, \xi)] : \text{KL}(P_N, P) \leq r \}$ and enjoys strong statistical out-of-sample guarantees. One can prove [35, Theorem 11] for all $P \in \mathcal{P}(\Xi)$ that indeed

$$\limsup_{N \rightarrow \infty} \frac{1}{N} \log \text{Prob}_P [\mathbf{E}_P [\ell(z_{\text{KL},r}(P_N), \xi)] > \sup \{ \mathbf{E}_Q [\ell(z_{\text{KL},r}(P_N), \xi)] + \epsilon : \text{KL}(P_N, Q) \leq r \}] \leq -r. \quad (8)$$

for any $\epsilon > 0$. In other words, the probability of being disappointed (the actual cost $\mathbf{E}_P [\ell(z_{\text{KL},r}(P_N), \xi)]$ of the decision $z_{\text{KL},r}(P_N)$ exceeds the predicted worst-case cost $\sup \{ \mathbf{E}_Q [\ell(z_{\text{KL},r}(P_N), \xi)] : \text{KL}(P_N, Q) \leq r \}$ by any small amount $\epsilon > 0$) decays exponentially fast in the number of samples with rate precisely the size r of the considered ambiguity set.

1.1 Efficient Formulations

Equations (6) and (8) reflect the fact that robust entropic and Wasserstein formulations enjoy essentially the same out-of-sample guarantees. Robust formulations in general enjoy stronger out-of-sample guarantees than the classical sample average formulation by considering slightly inflated or biased cost predictions. The classical sample average formulation can in fact be made to enjoy similar out-of-sample guarantees by naively inflating its objective in Equation (3) by some bias term $b > 0$. Indeed, taking the bias term b sufficiently large we can guarantee that

$$\limsup_{N \rightarrow \infty} \frac{1}{N} \log \text{Prob}_P [\mathbf{E}_P [\ell(z_N(P_N), \xi)] > \mathbf{E}_{P_N} [\ell(z_N(P_N), \xi)] + b] \leq -r \quad \forall P \in \mathcal{P}. \quad (9)$$

No formulations should hence be judged based on its out-of-sample disappointment alone but rather on how it trades off out-of-sample disappointment against biased cost predictions. When several prescription formulations enjoy the same out-of-sample guarantees, we should prefer that formulation which inflates the cost prediction the least. Let $\tilde{c}_r(z, P_N)$ be any continuous and bounded cost prediction function and let $\tilde{z}_r(P_N) \in \arg \min_{z \in Z} \tilde{c}_r(z, P_N)$ be the associated prescriptive formulation. Suppose this formulation satisfies the same out-of-sample guarantee as our entropic robust formulation, i.e.,

$$\limsup_{N \rightarrow \infty} \frac{1}{N} \log \text{Prob}_P [\mathbf{E}_P [\ell(\tilde{z}_r(P_N), \xi)] > \tilde{c}_r(z, P_N)] \leq -r \quad \forall P \in \mathcal{P}. \quad (10)$$




Figure 1: Two distributions μ and ν which are equivalent up to a small distributional shift $\epsilon > 0$ are very dissimilar in terms of their entropic divergence, i.e., $\text{KL}(\mu, \nu) = \infty$.

Then, [35, Theorem 11] shows that for any empirical data distribution P_N we have

$$\tilde{c}_r(\tilde{z}_r(P_N), P_N) \geq \sup \{ \mathbf{E}_Q [\ell(z_{\text{KL},r}(P_N), \xi)] : \text{KL}(P_N, Q) \leq r \} \geq \mathbf{E}_{P_N} [\ell(z(P_N), \xi)]$$

indicating that the entropic formulation is minimally inflated or biased among a very rich class of formulations enjoying the out-of-sample guarantee (10). Hence, the entropic formulation adds a fixed but minimal amount of inflation bias to guarantee that its out-of-sample disappointment decays exponentially fast at the considered rate r . This efficiency of the entropic formulation is technically based on the fact that the sufficient statistic P_N enjoys a large deviation property with rate function precisely the entropic divergence [13, Theorem 6.2.10].

That being said, even a minimal fixed amount of inflation bias still leaves the entropic formulation inconsistent, i.e., $\lim_{N \rightarrow \infty} z_{\text{KL},r}(P_N) \notin \arg \min \mathbf{E}_P [\ell(z, \xi)]$ which may be undesirable in some practical settings. Strong out-of-sample guarantees such as (10) require conservative decisions even as a large amount of data points are observed. Consistent formulations which satisfy slightly weaker out-of-sample guarantees can however readily be derived simply by letting the considered robustness radius r_N shrink at the appropriate rate with N to guarantee $\lim_{N \rightarrow \infty} z_{\text{KL},r_N}(P_N) \in \arg \min \mathbf{E}_P [\ell(z, \xi)]$ whenever the loss function ℓ is lower semi-continuous and P satisfies a mild moment condition [14]. Furthermore, Bennouna and Van Parys [4] have shown via a moderate deviation argument that such consistent formulations with shrinking radius $r_N \downarrow 0$ suffers an inflation bias which decays to zero faster than any other formulation enjoying certain slightly weaker conditions on the out-of-sample performance.

1.2 Noisy Data

Unlike the optimal transport distance the entropic divergence is very sensitive to distributional shifts which may be present when the data is corrupted; see Figure 1. Folklore believe suggests that optimal transport methods derive their superior empirical performance from their ability to guard against such data corruption. We investigate this folklore believe in a framework in which the data ξ_i is not observed directly. Rather,

$$\xi'_i = \xi_i + n_i \sim P' \quad \forall i \in [1, \dots, N], \tag{11}$$

is observed instead where the noise terms n_i are identically distributed following a distribution O_{ξ_i} which may depend on ξ_i . Remark that here the noise distribution $O : \Xi \rightarrow \mathcal{P}(\Xi')$ is assumed known. Our observational process will be required to admit a density function as stated in Assumption 1.1.

Assumption 1.1. The map $O : \Xi \rightarrow \mathcal{P}(\Xi')$ is absolutely continuous with respect to a base measure m' , i.e., $O_\xi \ll m'$ for all $\xi \in \Xi$. Furthermore, there exists a measurable function $d : \Xi \times \Xi' \rightarrow \mathbb{R}$ so that $dO_\xi/dm'(\xi') = \exp(-d(\xi, \xi'))$ for all $\xi' \in \Xi'$.

The relationship between the probability measure P' of the noisy observations and the probability measure P of the unobserved noiseless data can be characterized as the convolution $P' := O \star P$ as $P'(B) = (O \star P)(B) := \int O_\xi(B) dP(\xi)$ for all measurable sets $B \in \mathcal{B}(\Xi')$. We will denote with the set $\mathcal{P}' := \{O \star P \in \mathcal{P}(\Xi') : P \in \mathcal{P}\}$ the family of potential distributions of our noisy data. We point out that this noisy data model is quite flexible and captures a wide variety of settings. Perhaps the most classical of such settings is stated in the following example.

Example 1.2 (Gaussian Noise). We consider the case of zero mean Gaussian additive noise with variance σ^2 as an example. Assumption 1.1 holds here with $\Xi = \Xi' = \mathbb{R}^n$ and $d(\xi, \xi') = \|\xi - \xi'\|_2^2 / (2\sigma^2)$ with noisy power $\sigma \geq 0$ and $m' = \mu' / (\sigma \sqrt{(2\pi)^n})$ with μ' the Lebesgue measure on \mathbb{R}^n . Consequently here for all $\xi \in \Xi$ we have $O_\xi = N(\xi, \sigma^2)$ a normal distribution with mean ξ and variance σ^2 . The noisy data ξ'_i in Equation (11) follows the same distribution as

$$\xi_i + z_i \quad \forall i \in [1, \dots, N],$$

where ξ_1, \dots, ξ_N and z_1, \dots, z_N are independent and identically distributed as P and $N(0, \sigma^2)$, respectively.

We will attempt to infer the unknown probability measure P from the noisy data based on its empirical probability measure $P'_N = \sum_{i=1}^N \delta_{\xi'_i} / N$. Clearly, considering the empirical probability measure rather than the noisy data directly imposes no loss of information as the order of the data points is of no consequence here. Sanov's theorem [13, Theorem 6.2.10] ensures also here that the sufficient statistic P'_N enjoys a large deviation property. That is, the empirical probability measure P'_N satisfies for any open subset $\mathcal{O} \subseteq \mathcal{P}(\Xi')$ the large deviation lower bound

$$-\inf_{\hat{P}' \in \mathcal{O}} \text{KL}(\hat{P}', O \star P) \leq \liminf_{N \rightarrow \infty} \frac{1}{N} \log \text{Prob}_P[P'_N \in \mathcal{O}] \quad (12a)$$

and for any closed subset $\mathcal{C} \subseteq \mathcal{P}(\Xi')$ the large deviation upper bound

$$\limsup_{N \rightarrow \infty} \frac{1}{N} \log \text{Prob}_P[P'_N \in \mathcal{C}] \leq -\inf_{\hat{P}' \in \mathcal{C}} \text{KL}(\hat{P}', O \star P). \quad (12b)$$

for the good³ rate function $I(\hat{P}', P) := \text{KL}(\hat{P}', O \star P)$. We remark that large deviation inequalities generally are quite rough in nature as indeed (12a) and (12b) only pertain to open or closed sets (in the topology of weak convergence), respectively. The rate function is observed to be nonnegative and in fact $I(\hat{P}', P) = 0$ if and only if $\hat{P}' = P'$. For any $\epsilon > 0$, the large deviation inequality (12b) implies

$$\limsup_{N \rightarrow \infty} \frac{1}{N} \log \text{Prob}_P[\text{LP}(P'_N, P) \geq \epsilon] \leq -\min \left\{ I(\hat{P}', P) : \text{LP}(\hat{P}', P) \geq \epsilon \right\} < 0 \quad \forall P \in \mathcal{P}$$

where the minimum is indeed achieved as our good rate function has compact sublevel sets and the set of all

³A rate function I is good if its sublevel sets $\{\hat{P}' \in \mathcal{P}(\Xi') : I(\hat{P}', P) \leq r\}$ for any $r \geq 0$ and $P \in \mathcal{P}$ are compact [13].

$\hat{P}' \in \mathcal{P}(\Xi')$ such that $\text{LP}(\hat{P}', P') \geq \epsilon$ is by definition closed⁴ and does not contain the distribution P' . Hence, the large deviation property immediately implies that the empirical probability measure P'_N converges in probability to P' with an increasing number of observations. In fact, the rate function can be interpreted to quantify the exponential speed with which this convergence in probability takes place.

1.3 Notation

We will assume that Ξ and Ξ' are Polish topological spaces and hence so is the product space $\Xi \times \Xi'$ when equipped with the product topology. Given any set $S \subset \Xi'$ we denote with $S^\delta = \{s' \in \Xi' : \|s - s'\| \leq \delta, s \in S\}$ its δ -inflation. We denote with $\mathcal{M}_+(\Xi)$, $\mathcal{M}_+(\Xi')$ and $\mathcal{M}_+(\Xi \times \Xi')$ the sets of all positive Borel measures on the spaces Ξ , Ξ' and $\Xi \times \Xi'$, respectively. Similarly, we denote with $\mathcal{P}(\Xi)$, $\mathcal{P}(\Xi')$ and $\mathcal{P}(\Xi \times \Xi')$ the sets of all Borel probability measures on the spaces Ξ , Ξ' and $\Xi \times \Xi'$, respectively. Given two measures μ and ν we denote with $\mu \otimes \nu$ as their product measure. Following Dembo and Zeitouni [13, Section 6.2] the probability simplices $\mathcal{P}(\Xi)$, $\mathcal{P}(\Xi')$ and $\mathcal{P}(\Xi \times \Xi')$ when equipped with the topology of weak convergence of probability measures are Polish spaces too.

1.4 Contributions

We present three contributions:

1. We prove in Theorem 2.4 that the family of robust formulations

$$\tilde{z}^\delta(P'_N) \in \arg_\epsilon \inf_{z \in Z} \sup \{ \mathbf{E}_Q [\ell(z, \xi)] : Q \in \mathcal{P}, I^\delta(P'_N, Q) \leq r \} \quad (13)$$

parameterized in $\delta > 0$ essentially dominates all others formulations mirroring the notion of efficiency enjoyed by the entropic distance in the noiseless regime. Perhaps surprisingly we show that the rate function I is in the noisy setting too irregular and its δ -smoothed counterparts

$$I^\delta(\mu, \nu) := \inf \{ I(v, \nu) : v \in \mathcal{P}(\Xi'), \text{LP}(v, \mu) \leq \delta \} \quad (14)$$

must be considered instead.

2. We show in Theorem 2.6 that the rate function admits an interpretation as a marriage between both entropic optimal transport and KL distances, i.e., $I(\mu, \nu) := \inf_{v \in \mathcal{P}(\Xi)} W_1(\mu, v) + \text{KL}(v, \nu) + \text{KL}(\mu, m')$.
3. We state tractable reformulations of the proposed novel family of efficient robust prescriptive formulations (13) under certain technical conditions on the loss function ℓ and space Ξ in Section 3. In particular, we exploit a classical representation result of Strassen [34] which seems to be novel in this context and derive a dual formulation whose size is independent of the event set Ξ' .

⁴We denote here with $\text{LP}(\cdot, \cdot)$ the Lévy-Prokhorov metric [26] which metricizes the weak topology on $\mathcal{P}(\Xi')$.

2 Efficient Prescription with Noisy Data

We consider a prescriptive problem in which we attempt to learn the solution to the stochastic optimization problem stated in Equation (2) from the noisy observational data defined in Equation (11). Let us denote with P_N^{ml} the maximum likelihood estimate for the unobserved probability distribution P which as pointed out by Rigollet and Weed [28] can under certain conditions be computed efficiently as $P_N^{\text{ml}} = \arg \min_{P \in \mathcal{P}} W_1(P'_T, P)$. A straightforward extension of the sample average formulation in Equation (3) to this noisy data would be to consider

$$z(P_N^{\text{ml}}) \in \arg_{\epsilon} \inf_{z \in Z} \mathbf{E}_{P_N^{\text{ml}}} [\ell(z, \xi)]. \quad (15)$$

Many other formulations based on different distributional estimates are evidently possible as well. This naturally leads us to question if between these many alternative data-driven formulations one ought to be preferred over the other from a statistical point of view? To answer this question more broadly we must of course first define precisely what constitutes a data-driven formulation and secondly agree on how its statistical performance should be quantified. We again follow the framework presented in [35] presented in Section 1.1 and define a data-driven formulation as consisting of a predictor and prescriptor.

Definition 2.1 (Data-driven predictors and prescriptors). A measurable function $\tilde{c} : Z \times \mathcal{P}(\Xi') \rightarrow \mathbb{R}$ is called a data-driven predictor. A measurable function $\tilde{z} : \mathcal{P}(\Xi') \rightarrow Z$ is called a data-driven prescriptor if there exists a data-driven predictor \tilde{c} that induces \tilde{z} in the sense that $\tilde{z}(\hat{P}') \in \arg_{\epsilon} \inf_{z \in Z} \tilde{c}(z, \hat{P}')$ for all $\hat{P}' \in \mathcal{P}(\Xi')$. That is, we have $\tilde{c}(\tilde{z}(\hat{P}'), \hat{P}') - \epsilon < \tilde{v}(\hat{P}') := \inf_{z \in Z} \tilde{c}(z, \hat{P}')$ where we denote the function $\tilde{v} : \mathcal{P}(\Xi') \rightarrow \mathbb{R}$ as the optimal value function of the formulation.

The maximum likelihood formulation (15) employs the cost predictor $\mathbf{E}_{P_N^{\text{ml}}} [\ell(z, \xi)]$ to prescribe its decision $z(P_N^{\text{ml}})$. However, the maximum likelihood is a mere point estimate of the unobserved probability distribution P . The maximum likelihood formulation can consequently be expected to suffers similar shortcomings as the sample average formulation in the noiseless regime. That is, the cost budgeted for its prescribed decision is likely to disappoint out of sample. Here we say a formulation based on a predictor prescriptor pair (\tilde{c}, \tilde{z}) disappoints if the event

$$P'_N \in \mathcal{D}(\tilde{c}, \tilde{z}; P) := \left\{ \hat{P}' \in \mathcal{P}(\Xi') : c(\tilde{z}(\hat{P}'), P) > \tilde{c}(\tilde{z}(\hat{P}'), \hat{P}') \right\}$$

occurs with $c(z, P) = \mathbf{E}_P [\ell(z, \xi)]$ the unknown out-of-sample cost. Such disappointment events in which the actual cost of our decision $(c(\tilde{z}(P'_N), P))$ breaks the predicted cost or budget $(\tilde{c}(\tilde{z}(P'_N), P'_N))$ are undesirable and should be avoided by the decision-maker. Consequently, we prefer formulations which keep the disappointment rates

$$\limsup_{N \rightarrow \infty} \frac{1}{N} \log \text{Prob}_P [P_N \in \mathcal{D}(\tilde{c}, \tilde{z}; P)] \quad (16)$$

as small as possible for all $P \in \mathcal{P}$. We will only denote here formulations as feasible if their out-of-sample disappointment probability decays sufficiently fast, i.e., (16) $\leq -r$. Evidently, sufficiently fast disappointment probability decay can be achieved trivially by simply inflating the cost budgeted for each decision by some large nonnegative

amount. We would prefer formulations which promise minimal biased long term cost prediction $\tilde{c}(\tilde{z}(O \star P), O \star P)$ for all $P \in \mathcal{P}$ as indeed we have that the observed random cost $\tilde{c}(\tilde{z}(P'_N), P'_N)$ converges almost surely to $\tilde{c}(\tilde{z}(O \star P), O \star P)$ as the empirical distribution P'_N converges almost surely to $O \star P$ following [15, Theorem 11.4.1] for every $P \in \mathcal{P}$.

We consider here the family of robust formulations utilizing predictor prescriptor pairs

$$\tilde{c}^\delta(z, P'_N) := \sup \{ \mathbf{E}_Q [\ell(z, \xi)] : Q \in \mathcal{P}, I^\delta(P'_N, Q) \leq r \}, \quad \tilde{z}^\delta(P'_N) \in \arg_\epsilon \inf_{z \in Z} \tilde{c}^\delta(z, P'_N) \quad (17)$$

based on our smooth large deviation rate function defined in Equation (14). We will show using a large deviation argument that this family dominates the very rich class of regular formulations.

Definition 2.2 (Regular predictors and prescriptors). A data-driven predictor $\tilde{c} : Z \times \mathcal{P}(\Xi') \rightarrow \mathbb{R}$ is called regular if it is continuous on $Z \times \mathcal{P}(\Xi')$. A data-driven prescriptor $\tilde{z} : \mathcal{P}(\Xi') \rightarrow Z$ is called a regular if it is continuous and there exists a regular data-driven predictor \tilde{c} that induces \tilde{z} in the sense that $\tilde{z}(\hat{P}') \in \arg_\epsilon \inf_{z \in Z} \tilde{c}(z, \hat{P}')$ for all $\hat{P}' \in \mathcal{P}(\Xi')$.

Remark that the class of all regular formulations is very rich as Definition 2.2 imposes only mild structural restrictions. The Berge maximum theorem [5, p. 116] indeed implies that the optimal value function $\tilde{v}(\hat{P}') = \min_{z \in Z} \tilde{c}(z, \hat{P}')$ of any regular formulation is a continuous function on $\mathcal{P}(\Xi)$ already when the constraint set Z is merely compact. The correspondence $\hat{P}' \mapsto \{z \in Z : \tilde{c}(z, \hat{P}') < \tilde{v}(\hat{P}') + \epsilon\}$ of ϵ -suboptimal solutions in a regular formulation is consequently lower semicontinuous [2, Corollary 4.2.4.1] for any $\epsilon > 0$. Hence, for formulations employing a convex predictor \tilde{c} and $\mathcal{P}(\Xi')$ a compact set, an associated regular predictor can always be found [1, Theorem 9.1.]. Should a regular formulation admit unique optimal decisions, such decisions will constitute a regular prescriptor as well following [5, p. 117]. The need to focus on this restricted but nevertheless quite rich class of regular formulations is necessary due to the rough nature of the employed large deviation argument.

Assumption 2.3. The cost function $c : Z \times \mathcal{P} \rightarrow \mathbb{R}$, $(z, P) \mapsto \mathbf{E}_P [\ell(z, \xi)]$ is continuous.

Theorem 2.4. *Let Assumption 2.3 hold. Then, the family of predictor prescriptor pairs $(\tilde{c}^\delta, \tilde{z}^\delta)$ is feasible for any $\delta > 0$, i.e.,*

$$\limsup_{N \rightarrow \infty} \frac{1}{N} \log \text{Prob}_P [P'_N \in \mathcal{D}(\tilde{c}^\delta, \tilde{z}^\delta; P)] \leq -r \quad \forall P \in \mathcal{P}. \quad (18)$$

Furthermore, for any regular predictor prescriptor pair (\tilde{c}, \tilde{z}) that satisfies

$$\limsup_{N \rightarrow \infty} \frac{1}{N} \log \text{Prob}_P [P'_N \in \mathcal{D}(\tilde{c}, \tilde{z}; P)] \leq -r \quad \forall P \in \mathcal{P}, \quad (19)$$

we have that for all $\epsilon > 0$ there exists $0 < \delta'$ so that for all $0 < \delta \leq \delta'$ we have

$$\tilde{c}(\tilde{z}(O \star P), O \star P) + 2\epsilon \geq \tilde{c}^\delta(\tilde{z}^\delta(O \star P), O \star P) \quad \forall P \in \mathcal{P}. \quad (20)$$

Proof. We start by proving that our family of formulations is feasible and satisfies inequality (18). To this end

define the sets

$$\mathcal{D}^\delta(P) = \{\hat{P}' \in \mathcal{P}(\Xi') : \sup_{z \in Z} c(z, P) - \tilde{c}^\delta(z, \hat{P}') > 0\} \quad \text{and} \quad \mathcal{B}^\delta(P) = \{\hat{P}' \in \mathcal{P}(\Xi') : I^\delta(\hat{P}', P) > r\}.$$

We may assume without loss of generality that we have $\mathcal{D}^\delta(P) \neq \emptyset$ for otherwise the out-of-sample disappointment $\text{Prob}_P[P'_N \in \mathcal{D}(\tilde{c}_\delta^\delta, \tilde{z}_\delta^\delta; P) \subseteq \mathcal{D}^\delta(P)]$ clearly vanishes for all $N \geq 1$ and inequality (18) holds trivially. We will now show that $\mathcal{D}^\delta(P) \subseteq \mathcal{B}^\delta(P)$. For the sake of contradiction, choose any $\hat{P}' \in \mathcal{D}^\delta(P)$, and assume that $I^\delta(\hat{P}', P) \leq r$. Thus, we have for some $z \in Z$ that

$$c(z, P) > \tilde{c}^\delta(z, \hat{P}') = \sup \left\{ c(z, Q) : Q \in \mathcal{P}, I^\delta(\hat{P}', Q) \leq r \right\} \geq c(z, P);$$

a contradiction. As $\hat{P}' \in \mathcal{D}^\delta(P)$ was chosen arbitrarily, we have thus shown that $\mathcal{D}^\delta(P) \subseteq \mathcal{B}^\delta(P)$ and hence also $\text{cl } \mathcal{D}^\delta(P) \subseteq \text{cl } \mathcal{B}^\delta(P)$. Next we show that $\hat{P}' \in \text{cl } \mathcal{B}^\delta(P) \implies I(\hat{P}', P) > r$. For the sake of contradiction assume that we have found $\hat{P}' \in \text{cl } \mathcal{B}^\delta(P)$ for which $I(\hat{P}', P) \leq r$. There must exist a sequence $\hat{P}'_k \in \mathcal{B}^\delta(P)$ which converge to \hat{P}' and hence $\text{LP}(\hat{P}'_k, P')$ tends to zero. However, from the definition of $\mathcal{B}^\delta(P)$ we have that in fact for all $Q' \in \mathcal{P}(\Xi')$ such that $\text{LP}(Q', \hat{P}'_k) \leq \delta$ we have that $I(Q', P) > r$. Take now k large enough such that $\text{LP}(\hat{P}'_k, \hat{P}') \leq \delta$ then we must have $I(\hat{P}', P) > r$; a contradiction. The above reasoning implies that

$$\begin{aligned} & \limsup_{N \rightarrow \infty} \frac{1}{N} \log \text{Prob}_P [c(\tilde{z}^\delta(P'_N), P) > \tilde{c}^\delta(\tilde{z}^\delta(P'_N), P'_N)] \\ & \leq \limsup_{N \rightarrow \infty} \frac{1}{N} \log \text{Prob}_P \left[\sup_{z \in Z} c(z, P) - \tilde{c}^\delta(z, P'_N) > 0 \right] \leq - \inf_{\hat{P}' \in \text{cl } \mathcal{B}^\delta(P)} I(\hat{P}', P) \leq -r \end{aligned}$$

establishing inequality (18) as $P \in \mathcal{P}$ was arbitrary.

We will now prove that for any regular predictor prescriptor pair (\tilde{c}, \tilde{z}) which satisfies inequality (19) we have

$$\liminf_{\delta \rightarrow 0} \inf_{P' \in \mathcal{P}'} \tilde{c}(\tilde{z}(P'), P') - \tilde{c}^\delta(\tilde{z}(P'), P') \geq 0. \quad (21)$$

From inequality (21) we can take $\delta' > 0$ sufficiently small so that $\tilde{c}(\tilde{z}(P'), P') \geq \tilde{c}^{\delta'}(\tilde{z}(P'), P') - \epsilon \geq \tilde{c}^{\delta'}(\tilde{z}^{\delta'}(P'), P') - 2\epsilon$ uniformly for all $P' \in \mathcal{P}' = \{O \star P : P \in \mathcal{P}\}$. Remark that we have clearly $\tilde{c}^\delta(z, P') \leq \tilde{c}^{\delta'}(z, P')$ for all $z \in Z$, $P' \in \mathcal{P}(\Xi')$ and $0 < \delta \leq \delta'$ from which inequality (20) follows immediately.

We will now establish inequality (21) by showing that

$$\liminf_{\delta \rightarrow 0} \inf_{P' \in \mathcal{P}'} \tilde{c}(\tilde{z}(P'), P') - \tilde{c}^\delta(\tilde{z}(P'), P') \geq -3\rho \quad (22)$$

for any $\rho > 0$. Assume for the sake of contradiction that $\lim_{\delta \rightarrow 0} \inf_{P' \in \mathcal{P}'} \tilde{c}(\tilde{z}(P'), P') - \tilde{c}^\delta(\tilde{z}(P'), P') < -3\rho$. There must hence exist a $\delta' > 0$ such that for all $\delta \leq \delta'$ we have $\inf_{P' \in \mathcal{P}'} \tilde{c}(\tilde{z}(P'), P') - \tilde{c}^\delta(\tilde{z}(P'), P') < -2\rho$. Consequently,

there exists a distribution $Q''(\delta) \in \mathcal{P}'$ such that

$$\begin{aligned} \tilde{c}(\tilde{z}(Q''(\delta)), Q''(\delta)) + 2\rho &< c^\delta(\tilde{z}(Q''(\delta)), Q''(\delta)) \quad \forall \delta \leq \delta' \\ &< \sup \{c(\tilde{z}(Q''(\delta)), P) : P \in \mathcal{P}, I^\delta(Q''(\delta), P) \leq r\} \quad \forall \delta \leq \delta'. \end{aligned}$$

Hence, there exists for all $\delta \leq \delta'$ a distribution $P^*(\delta) \in \mathcal{P}$ such that $I^\delta(Q''(\delta), P^*(\delta)) \leq r$ and $\tilde{c}(\tilde{z}(Q''(\delta)), Q''(\delta)) + 2\rho < c(\tilde{z}(Q''(\delta)), P^*(\delta))$. From the definition of the smooth rate function I^δ stated in Equation (14) this implies that there exists an auxiliary sequence $Q'''(\delta) \in \mathcal{P}(\Xi')$ such that $I(Q'''(\delta), P^*(\delta)) \leq r$ with $\text{LP}(Q'''(\delta), Q''(\delta)) \leq \delta$ for all $\delta \leq \delta'$. Remark that

$$\begin{aligned} \lim_{\delta \rightarrow 0} \tilde{c}(\tilde{z}(Q'''(\delta)), Q'''(\delta)) &= \lim_{\delta \rightarrow 0} \tilde{c}(\tilde{z}(Q''(\delta)), Q''(\delta)), \\ \lim_{\delta \rightarrow 0} c(\tilde{z}(Q'''(\delta)), P^*(\delta)) &= \lim_{\delta \rightarrow 0} c(\tilde{z}(Q''(\delta)), P^*(\delta)) \end{aligned}$$

as the functions $c : Z \times \mathcal{P} \rightarrow \mathbb{R}$, $\tilde{c} : Z \times \mathcal{P}(\Xi') \rightarrow \mathbb{R}$ and $\tilde{z} : \mathcal{P}(\Xi') \rightarrow Z$ are continuous and $\text{LP}(Q'''(\delta), Q''(\delta)) \leq \delta$ implies $\lim_{\delta \rightarrow 0} Q'''(\delta) = \lim_{\delta \rightarrow 0} Q''(\delta)$. Consequently, there exists a $\delta^* \in (0, \delta']$ so that with $P^* = P^*(\delta^*) \in \mathcal{P}$ and $Q^* = Q'''(\delta^*) \in \mathcal{P}(\Xi')$ we have both

$$|\tilde{c}(\tilde{z}(Q''(\delta^*)), Q''(\delta^*)) - \tilde{c}(\tilde{z}(Q^*), Q^*)| < \rho \quad \text{and} \quad |c(\tilde{z}(Q''(\delta^*)), P^*) - c(\tilde{z}(Q^*), P^*)| < \rho.$$

Hence, we have both $\tilde{c}(\tilde{z}(Q^*), Q^*) < c(\tilde{z}(Q^*), P^*)$ and $I(Q^*, P^*) \leq r$. Define the continuous function $Q' : [0, 1] \rightarrow \mathcal{P}(\Xi')$, $\lambda \mapsto Q^* \star P^* \cdot \lambda + Q^* \cdot (1 - \lambda)$ and recall that $I(O \star P^*, P^*) = 0$. As we have that the functions $c : Z \times \mathcal{P} \rightarrow \mathbb{R}$, $\tilde{c} : Z \times \mathcal{P}(\Xi') \rightarrow \mathbb{R}$ and $\tilde{z} : \mathcal{P}(\Xi') \rightarrow Z$ are continuous there hence exists $Q' = Q'(\lambda')$ for $\lambda' \in (0, 1]$ sufficiently small so that $I(Q', P^*) \leq \lambda I(O \star P^*, P^*) + (1 - \lambda)I(Q^*, P^*) \leq (1 - \lambda') \cdot r = r' < r$ using the convexity of the rate function and $\tilde{c}(\tilde{z}(Q'), Q') < c(\tilde{z}(Q'), P^*)$. Consequently, we have that $Q' \in \mathcal{D}(\tilde{c}, \tilde{z}; P^*) = \text{int } \mathcal{D}(\tilde{c}, \tilde{z}; P^*)$ as again we remark for a final time that the functions $c : Z \times \mathcal{P} \rightarrow \mathbb{R}$, $\tilde{c} : Z \times \mathcal{P}(\Xi') \rightarrow \mathbb{R}$ and $\tilde{z} : \mathcal{P}(\Xi') \rightarrow Z$ are continuous. From the large deviation inequality (12a) it follows now that

$$-r' \leq -I(Q', P^*) < - \inf_{\hat{P}' \in \text{int } \mathcal{D}(\tilde{c}, \tilde{z}; P^*)} I(\hat{P}', P^*) \leq \liminf_{N \rightarrow \infty} \frac{1}{N} \log \text{Prob}_P [P'_N \in \mathcal{D}(\tilde{c}, \tilde{z}; P^*)]$$

which is in direct contradiction with the feasibility inequality (19) as $r' < r$ which establishes our inequality (22) as $\rho > 0$ was arbitrary. \square

Inequality (18) indicates that any formulation in our family is feasible. Recall again that $\tilde{c}(\tilde{z}(O \star P), O \star P)$ and $\tilde{c}^\delta(\tilde{z}^\delta(O \star P), O \star P)$ represent the long term out-of-sample cost of the formulations associated with the cost predictors \tilde{c} and \tilde{c}^δ , respectively. Hence, inequalities (19) and (20) guarantee that any other feasible regular formulation is dominated (modulo the small constant $\epsilon > 0$) by members of our efficient family for all parameters $0 < \delta \leq \delta'$ with $0 < \delta'$ sufficiently small. The previous theorem hence indicates that our family dominates any regular formulation in terms of balancing the desire for small out-of-sample disappointment as well as minimal bias under Assumption

2.3 and are thus *efficient*. Additionally, we remark here that our focus on the asymptotic guarantees (18) and (19) and long term out-of-sample costs allows us to consider a much stronger sense of efficiency than classical minimax efficiency [21]. Minimax efficiency would amount here to guaranteeing that

$$\sup_{P \in \mathcal{P}} \tilde{c}(\tilde{z}(O \star P), O \star P) - \tilde{c}^\delta(\tilde{z}^\delta(O \star P), O \star P) \geq 0.$$

That is, the long term out-of-sample cost of the formulation associated with the cost predictors \tilde{c} is larger than the long term out-of-sample cost of the formulation associated with the cost predictors \tilde{c}^δ for *some distribution* P in \mathcal{P} . Guarantee (20) is much stronger in that we have

$$\tilde{c}(\tilde{z}(O \star P), O \star P) - \tilde{c}^\delta(\tilde{z}^\delta(O \star P), O \star P) \geq -\epsilon \quad \forall P \in \mathcal{P}.$$

That is, the long term out-of-sample cost of the formulation associated with the cost predictors \tilde{c} is larger (again modulo the small constant $\epsilon > 0$) than the long term out-of-sample cost of the formulation associated with the cost predictors \tilde{c}^δ for *any distribution* P in \mathcal{P} . Finally, we remark that Assumption 2.3 is rather mild and is already satisfied when the loss function $\ell : Z \times \Xi \rightarrow \mathbb{R}$ is merely bounded and uniformly continuous.

In view of the previous discussion it is tempting to consider the data-driven formulation with predictor and prescriptor

$$\tilde{c}^0(z, P'_N) := \sup \{ \mathbf{E}_P [\ell(z, \xi)] : Q \in \mathcal{P}, I(P'_N, Q) \leq r \}, \quad \tilde{z}^0(P'_N) \in \arg \min_{z \in Z} \tilde{c}^0(z, P'_N) \quad (23)$$

based directly on our rate function I rather than its δ -smoothed counterpart I^δ . Van Parys et al. [35] prove indeed that when given access to the noiseless data in Equation (1) with empirical distribution P_N the appropriate rate function is precisely the entropic distance $\text{KL}(P_N, P)$ and formulation (23) enjoys the desirable statistical properties as discussed in Section 1.1. However, recall again that for noisy data when the base measure m' defined in Assumption 1.1 fails to be atomic, the ambiguity set $\{Q \in \mathcal{P} : I(P'_N, Q) < \infty\} = \emptyset$ is trivial for any empirical distribution P'_N and consequently the associated data-driven formulation (23) is here obviously infeasible. Hence, considering a somewhat smoothed rate function I^δ instead of the rate function I directly seems unavoidable when faced with noisy observational data.

Remark 2.5 (Confidence Regions). Dembo and Zeitouni [13, Section 7.1] prove that the considered ambiguity sets $\{Q \in \mathcal{P} : I^\delta(P'_T, Q) \leq r\}$ associated with the efficient robust predictor prescriptor pairs $(\tilde{c}^\delta, \tilde{z}^\delta)$ can be interpreted as confidence regions for the unknown distribution P for any $\delta > 0$. They prove furthermore that such confidence regions enjoy a certain *universal optimality* property. Also in this context it can be remarked that in the absence of smoothing, i.e., $\delta = 0$, the ambiguity sets $\{Q \in \mathcal{P} : I(P'_T, Q) \leq r\}$ are empty for any empirical distribution P'_T when m' is nonatomic and hence fail to be a confidence region for the probability distribution P .

We conclude this section by providing an interesting connection between the actual rate function I and the entropic optimal transport which sheds a light on the role entropic optimal transport plays in this noisy observational data regime. We defer the proof of the next result to Appendix A.

Theorem 2.6. *Let Assumption 1.1 hold. The jointly convex rate function can be decomposed as*

$$I(\hat{P}', P) = \inf_{Q \in \mathcal{P}(\Xi)} \text{KL}(Q, P) + \text{KL}(\hat{P}', m') + W_1(\hat{P}', Q) \geq 0. \quad (24)$$

In the proof of Theorem 2.6 we indicate that if $\hat{P}' \ll m'$ the infimum in Problem (24) is achieved at Q^* whose Radon-Nikodym derivative with respect to P is $dQ^*/dP(\xi) = \int_{\Xi'} [\exp(-d(\xi, \xi')) / \int_{\Xi} \exp(-d(\xi'', \xi')) dP(\xi'')] d\hat{P}'(\xi')$ for all $\xi \in \Xi$. The optimization variable Q in Theorem 2.6 can be interpreted to represent the unobserved empirical distribution P_N of the noiseless data points. With this interpretation in mind the first term $\text{KL}(Q, P)$ ensures that $\text{Prob}_P[P_N \approx Q] \asymp \exp(-N \cdot \text{KL}(Q, P))$ following Sanov's theorem and accounts for the fact that the empirical distribution P_N of the noiseless data may differ from unknown the distribution P when the number of training data points is finite. The last two terms accounts for the fact that we only observe the empirical distribution P'_N of the noisy data. One can show that this term quantifies indeed $\text{Prob}_P[P'_N \approx \hat{P}' | P_N \approx Q] \asymp \exp(-N \cdot (\text{KL}(\hat{P}', m') + W_1(\hat{P}', Q)))$. Informally, we have using the law of total probability that

$$\begin{aligned} \text{Prob}_P[P'_N \approx \hat{P}'] &= \int \text{Prob}_P[P'_N \approx \hat{P}' | P_N \approx Q] \text{Prob}_P[P_N \approx Q] \\ &\asymp \int \exp(-N \cdot (\text{KL}(\hat{P}', m') + W_1(\hat{P}', Q))) \exp(-N \cdot \text{KL}(Q, P)) \\ &\asymp \exp(-N \cdot \inf_{Q \in \mathcal{P}} \text{KL}(Q, P) + [\text{KL}(\hat{P}', m') + W_1(\hat{P}', Q)]). \end{aligned}$$

We remark that the entropic optimal transport term $W_1(\hat{P}', Q)$ is defined in Equation (7) where its marginal transport cost d is identified here as the logarithm of the density function of the noise corrupting our data as indicated in Assumption 1.1. The term $\text{KL}(\hat{P}', m')$ can be interpreted as a compensation term for the fact that any density function arbitrarily depends on the base measure m' considered.

Finally, we remark that the optimization problem in Equation (24) is identified as a so called *unbalanced* optimal transport problem initially proposed by [3]. Chizat et al. [8] propose scalable algorithms based on an entropic regularization scheme for such unbalanced optimal transport problems which have recently witnessed a flurry of interest, c.f., Séjourné et al. [31] and references therein.

3 Finite Formulations

Our family of efficient robust formulations in Equation (17) is stated in terms of a saddle-point problem which may be difficult to solve in general. Indeed, even the original stochastic optimization problem (2) may not be easy to solve. For the sake of simplicity we assume here that nothing is known about P and hence $\mathcal{P} = \mathcal{P}(\Xi)$. We remark though that the presented analysis generalizes to the case where \mathcal{P} is a convex subset of $\mathcal{P}(\Xi)$ with only minor modifications. If the loss function $\ell(x, \xi)$ is convex in the decision variable x for any ξ then the robust formulations in Equation (17) only require the solution of a convex optimization problem in the decision variable. Whether or not the convex optimization problem characterizing the optimal decision is tractable depends on whether the

prediction function $\tilde{c}^\delta(z, P'_N)$ can be evaluated efficiently which we will discuss now in more depth.

The maximization problem characterizing the prediction function $\tilde{c}^\delta(z, P'_N)$ in Equation (17) is convex in the distribution P . Indeed, the closed metric balls $\{\hat{P}' \in \mathcal{P}(\Xi') : \text{LP}(\hat{P}', P'_N) \leq \delta\} := \{\hat{P}' \in \mathcal{P}(\Xi') : P'_N(B) \leq \hat{P}'(B^\delta) + \delta \ \forall B \subseteq \Xi'\}$ induced by any metric, and hence in particular the Lévy-Prokhorov distance, are convex. Consequently, we have that the prediction function is characterized here as

$$\begin{aligned} \tilde{c}^\delta(z, P'_N) &= \sup \mathbf{E}_Q [\ell(z, \xi)] \\ \text{s.t. } & Q \in \mathcal{P}(\Xi), \hat{P}' \in \mathcal{P}(\Xi'), \\ & I(\hat{P}', Q) \leq r, \text{LP}(\hat{P}', P'_N) \leq \delta. \end{aligned} \quad (25)$$

However, even in the case where both event sets Ξ and Ξ' have finite cardinality the terminal Lévy-Prokhorov constraint is characterized using $2^{|\Xi'|}$ linear inequalities which becomes prohibitive even for moderately sized event sets Ξ' . This observation may perhaps explain why the Lévy-Prokhorov metric has not been considered extensively in the robust optimization literature. The sole exception is the work of Erdoğan and Iyengar [16] who proposes a stochastic approximation method for chance constrained optimization problems with Lévy-Prokhorov ambiguity.

3.1 Strassen Representation

Surprisingly, by exploiting a classical representation result of Strassen [34] the Lévy-Prokhorov metric need not result in intractable formulations. To the best of our knowledge, the application of the Strassen representation to derive a tractable reformulation of the Lévy-Prokhorov distance is novel.

Theorem 3.1 (Strassen Representation). *Let $\Xi'_N = \text{supp}(P'_N)$ denote the support of P'_N . Then,*

$$\begin{aligned} \tilde{c}^\delta(z, P'_N) &= \sup \mathbf{E}_Q [\ell(z, \xi)] \\ \text{s.t. } & Q \in \mathcal{P}(\Xi), \hat{P}' \in \mathcal{M}_+(\Xi'), T \in \mathcal{M}_+(\Xi' \times \Xi'_N), \\ & I(\hat{P}', Q) \leq r, & [\text{DV} : \beta \geq 0] \\ & \Pi_{\Xi'} T = \hat{P}', & [\text{DV} : u : \Xi' \rightarrow \mathbf{R}] \\ & \Pi_{\Xi'_N} T = P'_N, & [\text{DV} : v : \Xi'_N \rightarrow \mathbf{R}] \\ & \int_{\Xi' \times \Xi'_N} \mathbf{1} \{\|\xi' - \xi'_i\| \leq \delta\} dT(\xi', \xi'_i) \geq 1 - \delta. & [\text{DV} : \gamma \geq 0] \end{aligned} \quad (26)$$

Proof. As we assume that $\int_{\Xi'} P'_N(\xi') = 1$, we have that $T \in \mathcal{M}_+(\Xi' \times \Xi'_N)$, $\Pi_{\Xi'_N} T = P'_N \implies T \in \mathcal{P}(\Xi' \times \Xi'_N)$. Similarly, $\hat{P}' \in \mathcal{M}_+(\Xi')$, $T \in \mathcal{M}_+(\Xi' \times \Xi'_N)$, $\Pi_{\Xi'_N} T = P'_N$, $\Pi_{\Xi'} T = \hat{P}' \implies \hat{P}' \in \mathcal{P}(\Xi')$. The rest of the statement is a simple consequence of the more general equivalence

$$P'_N(B) \leq \hat{P}'(B^{\delta_1}) + \delta_2 \ \forall B \subseteq \Xi' \iff \begin{cases} \exists T \in \mathcal{P}(\Xi' \times \Xi'_N) \text{ s.t. } \Pi_{\Xi'} T = \hat{P}', \Pi_{\Xi'_N} T = P'_N, \\ \int_{\Xi' \times \Xi'_N} \mathbf{1} \{\|\xi' - \xi'_i\| \leq \delta_1\} dT(\xi', \xi'_i) \geq 1 - \delta_2 \end{cases}$$

for any $\delta_1, \delta_2 \geq 0$ [34, Theorem 11]. □

The equivalent formulation stated in Theorem 3.1 can be solved efficiently using an off-the-shelf exponential cone optimization solver [11] when both event sets Ξ and Ξ' have finite cardinality. Perhaps the only complication is that the size of this equivalent formulation counts $\mathcal{O}(|\Xi'| |\Xi'_N|)$ variables for $\mathcal{O}(|\Xi'| + |\Xi'_N|)$ constraints which may limit its practicality when both $|\Xi'|$ and $|\Xi'_N|$ are large.

We now indicate that even if the event set Ξ' is not finite Problem (26) still admits a finite reduction. Consider a finite partition of Ξ' which is generated by the closed balls around the observed data points, i.e.,

$$\Xi'_k = \bigcap_{i=1}^{|\Xi'_N|} \left\{ \xi' \in \Xi' : \begin{array}{l} \|\xi' - \xi'_i\| \leq \delta \quad \text{if } b(k, i) = 1 \\ \|\xi' - \xi'_i\| > \delta \quad \text{if } b(k, i) = 0 \end{array} \right\}$$

for $0 \leq k \leq K - 1$ with $K = 2^{|\Xi'_N|}$ and $b(k, i)$ the i th digit of the natural number k written down in binary notation. Clearly, we have $\Xi'_k \cap \Xi'_{k'} = \emptyset$ if $k \neq k'$ and $\bigcup_{k=0}^{K-1} \Xi'_k = \Xi'$. Note that it may happen that $\Xi'_k = \emptyset$ for some $0 \leq k \leq K - 1$. We defer the proof of the following result to Appendix B.

Lemma 3.2. *Let $\Xi'_N = \text{supp}(P'_N)$ denote the support of P'_N and let Ξ be finite. Then,*

$$\begin{aligned} \tilde{c}^\delta(z, P'_N) = \tilde{c}_f^\delta(z, P'_N) &:= \sup \sum_{\xi \in \Xi} \ell(z, \xi) q_\xi \\ \text{s.t. } & q_\xi \geq 0 \quad \forall \xi \in \Xi, \quad \hat{p}' \in \mathbf{R}^K, \quad t \in \mathbf{R}^{K \times |\Xi'_N|} \\ & \sum_{k=0}^{K-1} \hat{p}'_k \log \left(\hat{p}'_k / \sum_{\xi \in \Xi} O_\xi(\Xi'_k) \cdot q_\xi \right) \leq r, \\ & \sum_{i=1}^{|\Xi'_N|} t_{k,i} = \hat{p}'_k \quad 0 \leq k \leq K - 1, \\ & \sum_{k=0}^{K-1} t_{k,i} = P'_N(\xi'_i) \quad 1 \leq i \leq |\Xi'_N|, \\ & \sum_{i=1}^{|\Xi'_N|} \sum_{k=0}^{K-1} b(k, i) t_{k,i} \geq 1 - \delta. \end{aligned} \tag{27}$$

It should be remarked although Problem (27) is finite it may be very large as indeed we have that $K = \mathcal{O}(2^{|\Xi'_N|})$. This clearly limits the practical usefulness of Problem (27) to situations where the number of observed distinct data points Ξ'_N is very small. Surprisingly, using a dual equivalent formulation instead the computational burden of evaluating $c^\delta(z, P'_N)$ can be further reduced to $\mathcal{O}(|\Xi'_N|)$ variables and $\mathcal{O}(1)$ constraints which is the subject of the following section.

3.2 Dual Representation

Consider the minimization problem

$$\begin{aligned} \inf \quad & \beta r - \int_{\Xi'} v(\xi') dP'_N(\xi') + \gamma(\delta - 1) + \max_{\xi \in \Xi} \left[\ell(z, \xi) + \beta \int_{\Xi'} \exp \left(\frac{u(\xi')}{\beta} - 1 \right) dO_\xi(\xi') \right] \\ \text{s.t. } \quad & \beta \geq 0, \quad u : \Xi' \rightarrow \mathbf{R}, \quad v : \Xi'_N \rightarrow \mathbf{R}, \quad \gamma \geq 0, \\ & \gamma \mathbb{1} \{ \|\xi' - \xi'_i\| \leq \delta \} + v(\xi'_i) \leq u(\xi') \quad \forall \xi' \in \Xi', \quad \xi'_i \in \Xi'_N. \end{aligned} \tag{28}$$

We label the previous problem as the dual problem of the primal problem (26) which is nontrivial if $\tilde{c}^\delta(z, P'_N)$




Figure 2: The associated partition of the event set $\Xi'_N = \cup_{j=1}^{J(v)+1} \Xi'_j(v)$ associated with the vector v given in Example 3.4.

is finite. We remark that the next duality result does not assume Ξ' to have finite cardinality. We defer its proof to Appendix C.

Theorem 3.3 (Dual Representation). *Let Ξ be finite. Suppose that Problem (25) enjoys the Slater constraint qualification condition, i.e., there exist $Q_s \in \mathcal{P}(\Xi)$, $\hat{P}'_s \in \mathcal{P}(\Xi')$ with $I(\hat{P}'_s, Q_s) < r$ and $\text{LP}(\hat{P}'_s, P'_N) < \delta$. Then, we have $\tilde{c}^\delta(z, P'_N) = (28)$.*

First note that the dual characterization of the prediction function in Theorem 3.3 allows for the convex saddle-point formulation (17) to be solved as an ordinary convex minimization problem as the dual formulation (28) is jointly convex in $z \in Z$ and the dual variables. This may be desirable in practice as saddle-point optimization solvers are typically not as mature as solvers addressing standard optimization problems. This dual formulation counts $\mathcal{O}(|\Xi'_N| + |\Xi'|)$ variables for $\mathcal{O}(|\Xi'| |\Xi'_N|)$ constraints.

We now show that this can be further reduced to $\mathcal{O}(|\Xi'_N|)$ variables for $\mathcal{O}(1)$ constraints and in fact allows for a finite optimization characterization independent of the cardinality of the event set Ξ' . Consider $v : \Xi'_N \rightarrow \mathbb{R}$ and define $J(v) := |\{v(\xi') : \xi' \in \Xi'_N\}|$. Let $\xi'_{[j]}(v)$ for $j \in [1, \dots, J(v)]$ denote any partition of the observed data points in Ξ'_N such that the dual variables $v(\xi'_{[j]}(v))$ are non-increasing. That is, we have $\Xi'_N = [\xi'_{[1]}(v), \dots, \xi'_{[J(v)]}(v)]$ and $v(\tilde{\xi}') \geq v(\hat{\xi}')$ for all $\tilde{\xi}' \in \xi'_{[j]}(v)$, $\hat{\xi}' \in \xi'_{[j']}(v)$ with $1 \leq j < j' \leq J(v)$ as well as $v(\tilde{\xi}') = v(\hat{\xi}')$ for all $\tilde{\xi}', \hat{\xi}' \in \xi'_{[j]}(v)$ with $1 \leq j \leq J(v)$. We may now partition the event set Ξ' using the sets $\Xi'_0(v) = \emptyset$, $\Xi'_j(v) = \{\xi' \in \Xi' : \min\{\|\xi' - \tilde{\xi}'\| : \tilde{\xi}' \in \xi'_{[j]}(v)\} \leq \delta\} \setminus \cup_{l \in [1, \dots, j-1]} \Xi'_l(v)$ for $j \in [1, \dots, J(v)]$, and $\Xi'_{J(v)+1}(v) = \Xi' \setminus \cup_{l \in [1, \dots, J(v)]} \Xi'_l(v)$.

Example 3.4. Let $\Xi'_N = [\xi'_1, \dots, \xi'_5]$ and consider $v : \Xi'_N \rightarrow \mathbb{R}$ given as $v(\xi') = 1$ if $\xi' = \xi'_1$, $v(\xi') = 3$ if $\xi' = \xi'_2$, $v(\xi') = 4$ if $\xi' = \xi'_3$, $v(\xi') = 2$ if $\xi' = \xi'_4$ and $v(\xi') = 3$ if $\xi' = \xi'_5$. Then, $J(v) = 4$ and $\xi'_{[1]}(v) = \{\xi'_3\}$, $\xi'_{[2]}(v) = \{\xi'_2, \xi'_5\}$, $\xi'_{[3]}(v) = \{\xi'_4\}$, $\xi'_{[4]}(v) = \{\xi'_1\}$. The associated partition of the event set $\Xi' = \cup_{j=1}^{J(v)+1} \Xi'_j(v)$ is given in Figure 2.

Lemma 3.5 (Finite Convex Dual Representation). *The dual problem admits the following convex formulation:*

$$\begin{aligned}
(28) = \inf \quad & \beta r - \sum_{j=1}^{J(v)} v(\xi'_{[j]}) P'_N(\xi'_{[j]}) + \gamma(\delta - 1) + \max_{\xi \in \Xi} \left[\ell(z, \xi) \right. \\
& \left. + \sum_{j=1}^{J(v)} O_\xi(\Xi'_j(v)) \frac{\beta}{e} \exp\left(\frac{\max(v(\xi'_{[1]}(v)), v(\xi'_{[j]}(v)) + \gamma)}{\beta}\right) + O_\xi(\Xi'_{J(v)+1}(v)) \frac{\beta}{e} \exp\left(\frac{v(\xi'_{[1]}(v))}{\beta}\right) \right] \quad (29) \\
\text{s.t.} \quad & \beta \geq 0, v : \Xi'_N \rightarrow \mathbb{R}, \gamma \geq 0.
\end{aligned}$$

Proof. Note that the optimal dual variable $u^*(\gamma, v)$ in the dual formulation (28) can be determined explicitly as a function of the dual variables v and γ . It is can be easily verified that optimal dual variable is of the form

$$u^*(\xi; \gamma, v) = \begin{cases} \max(v(\xi'_{[1]}(v)), v(\xi'_{[1]}(v)) + \gamma) & \text{if } \xi' \in \Xi'_1(v), \\ \vdots & \vdots \\ \max(v(\xi'_{[1]}(v)), v(\xi'_{[J]}(v)) + \gamma) & \text{if } \xi' \in \Xi'_{J(v)}(v), \\ v(\xi'_{[1]}(v)) & \text{if } \xi' \in \Xi'_{J(v)+1}(v) \end{cases}$$

and is hence a piecewise constant function on the partition $\{\Xi'_j(v)\}_{j=1}^{J(v)+1}$. Indeed, any other feasible solution u in formulation (28) is pointwise larger, i.e., $u^*(\xi; \gamma, v) \leq u(\xi)$ for all $\xi \in \Xi'$. As the objective function in formulation (28) is increasing (in the pointwise order) in u , the choice u^* is without loss of optimality. Consequently, the ultimate term in the objective of the dual formulation (28) can be decomposed as the finite sum

$$\begin{aligned} & \beta \int_{\Xi'} \exp\left(\frac{u^*(\xi', \gamma, v)}{\beta} - 1\right) dO_\xi(\xi') \\ &= \sum_{j=1}^{J(v)} O_\xi(\Xi'_j(v)) \beta \exp\left(\max(v(\xi'_{[1]}(v)), v(\xi'_{[j]}(v)) + \gamma)/\beta - 1\right) + O_\xi(\Xi'_{J(v)+1}(v)) \beta \exp\left(v(\xi'_{[1]}(v))/\beta - 1\right). \end{aligned}$$

The convexity of the reduced formulation is a consequence of the fact [24, Theorem 1.31] that partially minimizing a convex formulation (here formulation (28)) with respect to any variable (here the variable u) results in a reduced formulation (here formulation (29)) which is convex itself. \square

The dual representation (29) can be solved as a finite convex optimization problem using a (stochastic) black-box optimization method [23] as long as we have a (stochastic) oracle which can access the probabilities $O_\xi(\Xi'_j(v)) = \int_{\Xi'} \mathbb{1}\{\Xi'_j(v)\} dO_\xi(\xi') = \int_{\Xi'} \mathbb{1}\{\Xi'_j(v)\} \exp(-d(\xi, \xi')) dm'(\xi')$ for all $\xi \in \Xi$ and $j \in [1, \dots, J(v) + 1]$. The complexity of our efficient formulation (25) is hence reduced to the complexity of integration of the noise distribution over certain intersections and unions of norm balls; see also Figure 2.

4 Conclusions

We introduce a novel class of distributionally robust optimization formulations which are efficient when making decisions based on noisy data in the sense pioneered by [35]. Their associated ambiguity sets are based on both an entropic divergence as well as an entropic optimal transport distance and can be interpreted as a large deviation rate function. Unlike in the noiseless regime, we show that this rate function too irregular and a certain amount of smoothing is necessary. Finally, we show that our efficient formulations are often tractable by exploiting a classical result by Strassen [34] and considering a dual representation.

References

- [1] J.-P. Aubin and H. Frankowska. **Set-valued analysis**. Springer Science & Business Media, 2009.
- [2] B. Bank, J. Guddat, D. Klatte, B. Kummer, and K. Tammer. **Non-linear parametric optimization**. Springer, 1982.
- [3] J.-D. Benamou. Numerical resolution of an “unbalanced” mass transport problem. **ESAIM: Mathematical Modelling and Numerical Analysis**, 37(5):851–868, 2003.
- [4] M. A. Bennouna and B. P. Van Parys. Learning and decision-making with data: Optimal formulations and phase transitions. **ArXiv**, 2021. URL <https://arxiv.org/abs/2109.06911>.
- [5] C. Berge. **Topological Spaces: Including a Treatment of Multi-Valued Functions, Vector Spaces, and Convexity**. Courier Corporation, 1997.
- [6] D. Bertsekas. **Convex Optimization Theory**. Athena Scientific, 2009.
- [7] D. Bertsimas, V. Gupta, and N. Kallus. Data-driven robust optimization. **Mathematical Programming**, 167(2):235–292, 2018.
- [8] L. Chizat, G. Peyré, B. Schmitzer, and F.-X. Vialard. Scaling algorithms for unbalanced optimal transport problems. **Mathematics of Computation**, 87(314):2563–2609, 2018.
- [9] I. Csiszár. A simple proof of sanov’s theorem. **Bulletin of the Brazilian Mathematical Society**, 37(4): 453–459, 2006.
- [10] M. Cuturi. Sinkhorn distances: Lightspeed computation of optimal transport. **Advances in neural information processing systems**, 26:2292–2300, 2013.
- [11] J. Dahl and E. Andersen. A primal-dual interior-point algorithm for nonsymmetric exponential-cone optimization. **Mathematical Programming**, pages 1–30, 2021.
- [12] E. Delage and Y. Ye. Distributionally robust optimization under moment uncertainty with application to data-driven problems. **Operations research**, 58(3):595–612, 2010.
- [13] A. Dembo and O. Zeitouni. **Large Deviations Techniques and Applications**, volume 38. Springer Science & Business Media, 2009.
- [14] J. Duchi, P. Glynn, and H. Namkoong. Statistics of robust optimization: A generalized empirical likelihood approach. **Mathematics of Operations Research**, 2021.
- [15] R. M. Dudley. **Real Analysis and Probability**. CRC Press, 2018.
- [16] E. Erdoğan and G. Iyengar. Ambiguous chance constrained problems and robust optimization. **Mathematical Programming**, 107(1):37–61, 2006.

- [17] P. M. Esfahani and D. Kuhn. Data-driven distributionally robust optimization using the Wasserstein metric: Performance guarantees and tractable reformulations. **Mathematical Programming**, 171(1-2):115–166, 2018.
- [18] P. Groeneboom and J. A. Wellner. **Information bounds and nonparametric maximum likelihood estimation**, volume 19. Springer Science & Business Media, 1992.
- [19] Z. Hu and L. J. Hong. Kullback-Leibler divergence constrained distributionally robust optimization. **Available at Optimization Online**, 2013.
- [20] D. Kuhn, P. M. Esfahani, V. A. Nguyen, and S. Shafieezadeh-Abadeh. Wasserstein distributionally robust optimization: Theory and applications in machine learning. In **Operations Research & Management Science in the Age of Analytics**, pages 130–166. INFORMS, 2019.
- [21] E. Lehmann and G. Casella. **Theory of point estimation**. Springer Science & Business Media, 2006.
- [22] R. O. Michaud. The Markowitz optimization enigma: Is “optimized” optimal? **Financial Analysts Journal**, 45(1):31–42, 1989.
- [23] Y. Nesterov. **Introductory lectures on convex programming volume i: Basic course**. Springer, 1998.
- [24] T. Pennanen. Introduction to convex optimization. <https://nms.kcl.ac.uk/teemu.pennanen/co-new.pdf>, 2019.
- [25] G. Peyré and M. Cuturi. Computational optimal transport: With applications to data science. **Foundations and Trends® in Machine Learning**, 11(5-6):355–607, 2019.
- [26] Y. V. Prokhorov. Convergence of random processes and limit theorems in probability theory. **Theory of Probability & Its Applications**, 1(2):157–214, 1956.
- [27] H. Rahimian and S. Mehrotra. Distributionally robust optimization: A review. **arXiv preprint arXiv:1908.05659**, 2019.
- [28] P. Rigollet and J. Weed. Entropic optimal transport is maximum-likelihood deconvolution. **Comptes Rendus – Mathématique**, 356(11-12):1228–1235, 2018.
- [29] R. Rockafellar and R. J.-B. Wets. **Variational Analysis**, volume 317. Springer Science & Business Media, 2009.
- [30] F. Santambrogio. **Optimal Transport for Applied Mathematicians**. Birkäuser, NY, 2015.
- [31] T. Séjourné, J. Feydy, F.-X. Vialard, A. Trounev, and G. Peyré. Sinkhorn divergences for unbalanced optimal transport. **arXiv preprint arXiv:1910.12958**, 2019.
- [32] A. Shapiro, D. Dentcheva, and A. Ruszczyński. **Lectures on Stochastic Programming: Modeling and Theory**. SIAM, 2014.
- [33] B. W. Silverman. **Density Estimation for Statistics and Data Analysis**, volume 26. CRC press, 1986.

- [34] V. Strassen. The existence of probability measures with given marginals. **The Annals of Mathematical Statistics**, 36(2):423–439, 1965.
- [35] B. P. Van Parys, P. M. Esfahani, and D. Kuhn. From data to decisions: Distributionally robust optimization is optimal. **Management Science**, 2020. doi: <https://doi.org/10.1287/mnsc.2020.3678>.
- [36] B. P. Van Parys, P. M. Esfahani, and D. Kuhn. From data to decisions: Distributionally robust optimization is optimal. **Management Science**, 67(6):3387–3402, 2021.
- [37] W. Wiesemann, D. Kuhn, and M. Sim. Distributionally robust convex optimization. **Operations Research**, 62(6):1358–1376, 2014.
- [38] C. Zhao and Y. Guan. Data-driven risk-averse stochastic optimization with wasserstein metric. **Operations Research Letters**, 46(2):262–267, 2018.

A Proof of Theorem 2.6

Proof. Let $T(P)$ be the joint distribution of the random variables (ξ_i, ξ'_i) where Assumption 1.1 holds. We will show momentarily that $\text{KL}(\hat{P}', O \star P) = \min\{\text{KL}(T', T(P)) : T' \in \mathcal{P}(\Xi \times \Xi'), \Pi_{\Xi'} T' = \hat{P}'\}$. The identity (24) follows then from Lemma A.1. The joint convexity of the rate function $I(\hat{P}', P) = \text{KL}(\hat{P}', O \star P)$ follows from the joint convexity of the relative entropy function and the linearity of $O \star P$ in P . It remains to show that $\text{KL}(\hat{P}', O \star P) = \min\{\text{KL}(T', T(P)) : T' \in \mathcal{P}(\Xi \times \Xi'), \Pi_{\Xi'} T' = \hat{P}'\}$.

Consider first the case where $\hat{P}' \not\ll O \star P$. In this case we have by definition $\text{KL}(\hat{P}', O \star P) = \infty$. Furthermore, $\hat{P}' = \Pi_{\Xi'} T' \not\ll \Pi_{\Xi'} T(P) = O \star P \implies T' \not\ll T(P)$ and consequently $\min\{\text{KL}(T', T(P)) : T' \in \mathcal{P}(\Xi \times \Xi'), \Pi_{\Xi'} T' = \hat{P}'\} = \infty$ as well.

Consider now the case where $\hat{P}' \ll O \star P$. We will show (i) that here $\text{KL}(\hat{P}', O \star P) = \min\{\text{KL}(T', T(P)) : T' \in \mathcal{P}(\Xi \times \Xi'), \Pi_{\Xi'} T' = \hat{P}'\} = \text{KL}(T'^*, T(P))$ and (ii) the minimum is achieved at $T'^* \ll T(P)$ explicitly characterized as

$$\frac{dT'^*}{dT(P)}(\xi, \xi') = \frac{d\hat{P}'}{d(O \star P)}(\xi') \quad \forall \xi \in \Xi, \xi' \in \Xi'. \quad (30)$$

First, observe that we have here the inequalities

$$\begin{aligned} & \text{KL}(\hat{P}', O \star P) \\ &= \int_{\Xi'} \log \left(\frac{d\hat{P}'}{d(O \star P)}(\xi') \right) \frac{d\hat{P}'}{d(O \star P)}(\xi') d(O \star P)(\xi') \\ &= \int_{\Xi'} \left[\max_{v \in \mathbb{R}} v \frac{d\hat{P}'}{d(O \star P)}(\xi') - \exp(v - 1) \right] d(O \star P)(\xi') \\ &= \max_{v: \Xi' \rightarrow \mathbb{R}} \int_{\Xi'} v(\xi') d\hat{P}'(\xi') - \int_{\Xi'} \exp(v(\xi') - 1) d(O \star P)(\xi') \\ &= \max_{v: \Xi' \rightarrow \mathbb{R}} \int_{\Xi'} v(\xi') d\hat{P}'(\xi') - \int_{\Xi \times \Xi'} \exp(v(\xi') - 1) dT(P)(\xi, \xi') \\ &= \max_{v: \Xi' \rightarrow \mathbb{R}} \int_{\Xi'} v(\xi') d\hat{P}'(\xi') - \int_{\Xi \times \Xi'} \max_{R \geq 0} Rv(\xi') - R \log(R) dT(P)(\xi, \xi') \\ &= \max_{v: \Xi' \rightarrow \mathbb{R}} \min_{R: \Xi \times \Xi' \rightarrow \mathbb{R}_+} \int_{\Xi'} v(\xi') d\hat{P}'(\xi') - \int_{\Xi \times \Xi'} R(\xi, \xi')v(\xi') - R(\xi, \xi') \log(R(\xi, \xi')) dT(P)(\xi, \xi') \\ &\leq \max_{v: \Xi' \rightarrow \mathbb{R}} \min_{T' \ll T(P) \in \mathcal{M}_+(\Xi \times \Xi')} \int_{\Xi'} v(\xi') d\hat{P}'(\xi') - \int_{\Xi \times \Xi'} v(\xi') dT'(\xi, \xi') + \log \left(\frac{dT'}{dT(P)}(\xi, \xi') \right) dT'(\xi, \xi') \\ &\leq \min_{T' \ll T(P) \in \mathcal{M}_+(\Xi \times \Xi')} \max_{v: \Xi' \rightarrow \mathbb{R}} \int_{\Xi'} v(\xi') d\hat{P}'(\xi') - \int_{\Xi \times \Xi'} v(\xi') dT'(\xi, \xi') + \log \left(\frac{dT'}{dT(P)}(\xi, \xi') \right) dT'(\xi, \xi') \\ &= \min_{T' \ll T(P) \in \mathcal{M}_+(\Xi \times \Xi'), \Pi_{\Xi'} T' = \hat{P}'} \log \left(\frac{dT'}{dT(P)}(\xi, \xi') \right) dT'(\xi, \xi') \\ &= \min\{\text{KL}(T', T(P)) : T' \in \mathcal{P}(\Xi \times \Xi'), \Pi_{\Xi'} T' = \hat{P}'\}. \end{aligned} \quad (31)$$

The first equality follows from the fact that $\hat{P}' \ll O \star P$ and the definition of the Kullback-Leibler divergence where we remark that here $\int d\hat{P}'(\xi') = \int d(O \star P)(\xi') = 1$. The second equality follows from the convex conjugate identity $\max_v av - \exp(v - 1) = av^*(a) - \exp(v^*(a) - 1) = a \log(a)$ with $v^*(a) = 1 + \log(a)$ for all $a \in \mathbb{R}_+$. The

third equality follows from Rockafellar and Wets [29, Theorem 14.60]. The fourth equality follows from the fact that $\Pi_{\Xi'} T(P) = O \star P$. The fifth equality follows from the convex conjugate identity $\max_{R \geq 0} aR - R \log(R) = aR^*(a) - R^*(a) \log(R^*(a)) = \exp(a - 1)$ with $R^*(a) = \exp(a - 1)$ for all $a \in \mathbb{R}$. The sixth equality follows again from Rockafellar and Wets [29, Theorem 14.60]. The first inequality follows from the fact that with any $T' \ll T(P) \in \mathcal{M}_+(\Xi \times \Xi')$ we can associate the Radon-Nikodym derivative $R(\xi, \xi') = dT'/dT(P)(\xi, \xi')$. The second inequality follows from the standard minimax inequality. The penultimate equality follows from the fact that $\max_{v, \Xi' \rightarrow \mathbb{R}} \int_{\Xi'} v(\xi') d\hat{P}'(\xi') - \int_{\Xi \times \Xi'} v(\xi') dT'(\xi, \xi') = \chi_\infty(\Pi_{\Xi'} T' = \hat{P}')$. The ultimate equality follows from the fact that $T' \ll T(P) \in \mathcal{M}_+(\Xi \times \Xi')$, $\Pi_{\Xi'} T' = \hat{P}'$, $\int d\hat{P}'(\xi') = 1 \implies T' \ll T(P) \in \mathcal{P}(\Xi \times \Xi')$.

Recall the definition of T'^* stated in Equation (30). We have the chain of equalities

$$\begin{aligned} \int_{\Xi \times \Xi'} \mathbb{1}\{\xi' \in B\} dT'^*(\xi, \xi') &= \int_{\Xi \times \Xi'} \mathbb{1}\{\xi' \in B\} d\hat{P}'/d(O \star P)(\xi') dT(P)(\xi, \xi') \\ &= \int_{\Xi'} \mathbb{1}\{\xi' \in B\} d\hat{P}'/d(O \star P)(\xi') d(O \star P)(\xi') = \hat{P}'(B) \end{aligned}$$

for any measurable set $B \subseteq \Xi'$. The first equality follows from Equation (30) and the second equality exploits $\Pi_{\Xi'} T(P) = O \star P$. In particular this implies that $T^* \in \mathcal{P}(\Xi \times \Xi')$ when choosing $B = \Xi'$ and $\Pi_{\Xi'} T^* = \hat{P}'$. As we have here $\hat{P}' \ll O \star P$ we can define the functions $v^*(\xi') = \log(d\hat{P}'/d(O \star P)(\xi')) + 1$ and $R^*(\xi, \xi') := dT'^*/dT(P)(\xi, \xi') = d\hat{P}'/d(O \star P)(\xi') = \exp(v^*(\xi') - 1)$ for all $\xi \in \Xi$ and $\xi' \in \Xi'$. Observe now that

$$\begin{aligned} &\text{KL}(T'^*, T(P)) \\ &= \int_{\Xi \times \Xi'} \log\left(\frac{dT'^*}{dT(P)}(\xi, \xi')\right) dT'^*(\xi, \xi') \\ &= \int_{\Xi'} v^*(\xi') d\hat{P}'(\xi') - \int_{\Xi \times \Xi'} v^*(\xi') + \log\left(\frac{dT'^*}{dT(P)}(\xi, \xi')\right) dT'^*(\xi, \xi') \\ &= \int_{\Xi'} v^*(\xi') d\hat{P}'(\xi') - \int_{\Xi \times \Xi'} v^*(\xi') \frac{dT'^*}{dT(P)}(\xi, \xi') + \log\left(\frac{dT'^*}{dT(P)}(\xi, \xi')\right) \frac{dT'^*}{dT(P)}(\xi, \xi') dT(P)(\xi, \xi') \\ &= \int_{\Xi'} v^*(\xi') d\hat{P}'(\xi') - \int_{\Xi \times \Xi'} R^*(\xi, \xi') v^*(\xi') - R^*(\xi, \xi') \log(R^*(\xi, \xi')) dT(P)(\xi, \xi') \\ &= \int_{\Xi'} v^*(\xi') d\hat{P}'(\xi') - \int_{\Xi \times \Xi'} \exp(v^*(\xi') - 1) dT(P)(\xi, \xi') \\ &= \int_{\Xi'} v(\xi') d\hat{P}'(\xi') - \int_{\Xi'} \exp(v(\xi') - 1) d(O \star P)(\xi') \\ &= \int_{\Xi'} v(\xi') d\hat{P}'(\xi') + \int_{\Xi'} d\hat{P}'(\xi') - \int_{\Xi'} d(O \star P)(\xi') = \text{KL}(\hat{P}', O \star P) \end{aligned}$$

The first equality follows from the definition of the Kullback-Leibler divergence and the fact that both T'^* and $T(P)$ are in $\mathcal{P}(\Xi \times \Xi')$. The second equality follows from the fact that $\int_{\Xi'} v^*(\xi') d\hat{P}'(\xi') - \int_{\Xi \times \Xi'} v^*(\xi') dT'^*(\xi, \xi') = 0$ as $\Pi_{\Xi'} T'^* = \hat{P}'$ and the linearity of integration. The third, fourth and fifth equality follow from the fact that $T^* \ll T(P)$ with Radon-Nikodym derivative $R^*(\xi, \xi') := dT^*/dT(P) = \exp(v^*(\xi') - 1)$. The penultimate equality follows from $\Pi_{\Xi'} T(P) = O \star P$. The final equality is direct consequence of the identity $v^*(\xi') = \log(d\hat{P}'/d(O \star P)(\xi')) + 1$. As we have already shown that $\text{KL}(\hat{P}', O \star P) \leq \min\{\text{KL}(T', T(P)) : T' \in \mathcal{P}(\Xi \times \Xi'), \Pi_{\Xi'} T' = \hat{P}'\}$ in Equality (31) this implies that as $T'^* \in \mathcal{P}(\Xi \times \Xi')$, $\Pi_{\Xi'} T'^* = \hat{P}'$ we have $\text{KL}(\hat{P}', O \star P) = \min\{\text{KL}(T', T(P)) : T' \in$

$\mathcal{P}(\Xi \times \Xi'), \Pi_{\Xi'} T' = \hat{P}'\}$.

We remark that in the case where $\hat{P}' \ll O \star P$ we have

$$\begin{aligned}
\Pi_{\Xi} T'^*(B) &= \int_{\Xi \times \Xi'} \mathbb{1}\{\xi \in B\} dT'^*(\xi, \xi') \\
&= \int_{\Xi \times \Xi'} \mathbb{1}\{\xi \in B\} \frac{d\hat{P}'}{d(O \star P)}(\xi') dT(P)(\xi, \xi') \\
&= \int_{\Xi \times \Xi'} \mathbb{1}\{\xi \in B\} \left(\frac{d\hat{P}'}{d(O \star P)}(\xi') \exp(-d(\xi, \xi')) dm'(\xi') \right) dP(\xi) \\
&= \int_{\Xi} \mathbb{1}\{\xi \in B\} \left(\int_{\Xi'} \frac{\exp(-d(\xi, \xi'))}{\int_{\Xi} \exp(-d(\xi'', \xi')) dP(\xi'')} d\hat{P}'(\xi') \right) dP(\xi)
\end{aligned}$$

for all $B \subseteq \Xi$. Hence, $\Pi_{\Xi} T'^*/dP(\xi) = \int_{\Xi} \exp(-d(\xi, \xi')) / (\int_{\Xi} \exp(-d(\xi'', \xi')) dP(\xi'')) d\hat{P}'(\xi')$ for all $\xi \in \Xi$. \square

Lemma A.1. *We have under Assumption 1.1 that*

$$\inf \left\{ \text{KL}(T', T(P)) : T' \in \mathcal{P}(\Xi \times \Xi'), \Pi_{\Xi'} T' = \hat{P}' \right\} = \inf \left\{ W_1(\hat{P}', Q) + \text{KL}(Q, P) + \text{KL}(\hat{P}', m') : Q \in \mathcal{P}(\Xi) \right\}.$$

Proof. Under Assumption 1.1 we have $T(P) \ll P \otimes m'$ and $P \otimes m' \ll T(P)$ with $dT(P)/d(P \otimes m')(\xi, \xi') = \exp(-d(\xi, \xi'))$ and $d(P \otimes m')/dT(P)(\xi, \xi') = \exp(d(\xi, \xi'))$. Assume that $T' \ll T(P)$ as otherwise $\text{KL}(T', T(P)) = +\infty$. The assumption $T' \ll T(P)$ furthermore implies $\Pi_{\Xi} T' \ll \Pi_{\Xi} T(P) = P$ and $\Pi_{\Xi'} T' \ll \Pi_{\Xi'} T(P) = P' \ll m'$.

We have here hence

$$\begin{aligned}
&\text{KL}(T', T(P)) \\
&= \int_{\Xi \times \Xi'} \log \left(\frac{dT'}{dT(P)}(\xi, \xi') \right) dT'(\xi, \xi') \\
&= \int_{\Xi \times \Xi'} \log \left(\frac{dT'}{dP \otimes m'}(\xi, \xi') \frac{dP \otimes m'}{dT(P)}(\xi, \xi') \right) dT'(\xi, \xi') \\
&= \int_{\Xi \times \Xi'} \log \left(\frac{dT'}{dP \otimes m'}(\xi, \xi') \right) dT'(\xi, \xi') - \int_{\Xi \times \Xi'} \log \left(\frac{dT(P)}{dP \otimes m'}(\xi, \xi') \right) dT'(\xi, \xi') \\
&= \text{KL}(T', P \otimes m') - \int_{\Xi \times \Xi'} \log(\exp(-d(\xi, \xi'))) dT'(\xi, \xi') \\
&= \int_{\Xi \times \Xi'} d(\xi, \xi') dT'(\xi, \xi') + \text{KL}(T', \Pi_{\Xi} T' \otimes \Pi_{\Xi'} T') + \text{KL}(\Pi_{\Xi} T', P) + \text{KL}(\Pi_{\Xi'} T', m') \geq 0. \tag{32}
\end{aligned}$$

The first equality follows from the definition of the Kullback-Leibler divergence and the fact that T' and $T(P)$ are in $\mathcal{P}(\Xi \times \Xi')$. The second equality follows from the fact that $T' \ll P \otimes m'$ and $P \otimes m' \ll T(P)$. The third and fourth equalities follow from $T(P) \ll P \otimes m'$ and $P \otimes m' \ll T(P)$ with $dT(P)/d(P \otimes m')(\xi, \xi') = \exp(-d(\xi, \xi'))$.

The final equality follows from [28, Lemma 6.1]. Hence, we have

$$\inf \left\{ \text{KL}(T', T(P)) : T' \in \mathcal{P}(\Xi \times \Xi'), \Pi_{\Xi'} T' = \hat{P}' \right\}$$

$$\begin{aligned}
&= \inf \left\{ \text{KL}(T', T(P)) : Q \in \mathcal{P}(\Xi), T' \ll T(P) \in \mathcal{P}(\Xi \times \Xi'), \Pi_{\Xi} T' = Q, \Pi_{\Xi'} T' = \hat{P}' \right\} \\
&= \inf \left\{ \int d(\xi, \xi') dT'(\xi, \xi') + \text{KL}(T', \Pi_{\Xi} T' \otimes \Pi_{\Xi'} T') + \text{KL}(Q, P) + \text{KL}(\hat{P}', m') \right. \\
&\quad \left. : Q \in \mathcal{P}(\Xi), T' \ll T(P) \in \mathcal{P}(\Xi \times \Xi'), T' \in \mathcal{T}(\hat{P}', Q) \right\} \\
&= \inf \left\{ \int d(\xi, \xi') dT'(\xi, \xi') + \text{KL}(T', \Pi_{\Xi} T' \otimes \Pi_{\Xi'} T') + \text{KL}(Q, P) + \text{KL}(\hat{P}', m') \right. \\
&\quad \left. : Q \in \mathcal{P}(\Xi), T' \in \mathcal{P}(\Xi \times \Xi'), T' \in \mathcal{T}(\hat{P}', Q) \right\} \\
&= \inf_{Q \in \mathcal{P}(\Xi)} W_1(\hat{P}', Q) + \text{KL}(Q, P) + \text{KL}(\hat{P}', m')
\end{aligned}$$

establishing the claim. The first equality follows here from the fact that $T' \not\ll T(P) \implies \text{KL}(T', T(P)) = \infty$. The second equality follows from Equation (32). Recall that we may assume that $Q \ll P$ and $P' \ll m'$ as otherwise $\text{KL}(Q, P) = \infty$ or $\text{KL}(\hat{P}', m') = \infty$, respectively. Hence, we have $T' \ll \Pi_{\Xi} T' \otimes \Pi_{\Xi'} T' = Q \otimes \hat{P}' \ll P \otimes m' \ll T(P)$ from which the third equality follows immediately. The final equality follows from the definition of entropic optimal transport in Equation (7). \square

B Proof of Lemma 3.2

Proof. We first prove that $\tilde{c}^{\delta}(z, P'_N) \leq \tilde{c}_f^{\delta}(z, P'_N)$. Consider any feasible solution (Q, \hat{P}', T) in Problem (26). Consider $q_{\xi} = Q(\xi)$ for all $\xi \in \Xi$, $\hat{p}'_k = \hat{P}'(\Xi'_k)$ and $t_{k,i} = T(\Xi'_k, \xi'_i)$ for all $1 \leq i \leq |\Xi'_N|$ and $0 \leq k \leq K-1$. We have that

$$\sum_{k=0}^{K-1} \hat{p}'_k \log \left(\hat{p}'_k / \sum_{\xi \in \Xi} O_{\xi}(\Xi'_k) \cdot q_{\xi} \right) = \sum_{k=0}^{K-1} \hat{P}'(\Xi'_k) \log \left(\hat{P}'(\Xi'_k) / \sum_{\xi \in \Xi} O_{\xi}(\Xi'_k) \cdot Q(\xi) \right) \leq I(\hat{P}', Q) \leq r$$

where the penultimate inequality follows from [9]. We have also

$$\sum_{i=1}^{|\Xi'_N|} t_{k,i} = \sum_{i=1}^{|\Xi'_N|} T(\Xi'_k, \xi'_i) = (\Pi_{\Xi'} T)(\Xi'_k) = \hat{P}'(\Xi'_k) = \hat{p}'_k$$

for any $0 \leq k \leq K-1$ as well as

$$\sum_{k=0}^{K-1} t_{k,i} = \sum_{k=0}^{K-1} T(\Xi'_k, \xi'_i) = (\Pi_{\Xi'_N} T)(\xi'_i) = P'_N(\xi'_i)$$

for any $1 \leq i \leq |\Xi'_N|$. Finally, we remark that

$$\sum_{i=1}^{|\Xi'_N|} \sum_{k=0}^{K-1} b(k, i) t_{k,i} = \sum_{i=1}^{|\Xi'_N|} \sum_{k=0}^{K-1} b(k, i) T(\Xi'_k, \xi'_i) = \sum_{i=1}^{|\Xi'_N|} \int_{\Xi'} \mathbb{1}_{\{\|\xi - \xi_i\| \leq \delta\}} T(\xi', \xi'_i) \geq 1 - \delta.$$

Hence, the constructed point (q, \hat{p}', t) is feasible in Problem (27) and attains the same objective value as the point (Q, \hat{P}', T) in Problem (26).

We now prove that $\tilde{c}^{\delta}(z, P'_N) \geq \tilde{c}_f^{\delta}(z, P'_N)$. Consider any feasible point (q, \hat{p}', t) in Problem (27) and let $Q(\xi) = q_{\xi}$

for all $\xi \in \Xi$. Define first the measures $(O \star Q)_{\Xi'_k} \in \mathcal{P}(\Xi'_k)$ as the restriction of the measure $O \star Q$ to the set $\Xi'_k \subseteq \Xi'$ for all $0 \leq k \leq K-1$. That is, these measures are defined through $(O \star Q)_{\Xi'_k}(B) = (O \star Q)(B \cap \Xi'_k) / (O \star Q)(\Xi'_k)$ for all measurable subsets B of Ξ' and $0 \leq k \leq K-1$. Let now $\hat{P}' = \sum_{k=0}^{K-1} \hat{p}'_k \cdot (O \star Q)_{\Xi'_k}$ and $T = \sum_{i=1}^{|\Xi'_N|} \sum_{k=0}^{K-1} t_{k,i} \cdot (\delta_{\xi'_i} \otimes (O \star Q)_{\Xi'_k})$. First, we remark that we have

$$\begin{aligned} (\Pi_{\Xi'} T)(B) &= \sum_{j=1}^{|\Xi'_N|} \sum_{i=1}^{|\Xi'_N|} \sum_{k=0}^{K-1} t_{k,i} \cdot (\delta_{\xi'_i}(\xi'_j) \cdot (O \star Q)_{\Xi'_k}(B)) \\ &= \sum_{i=1}^{|\Xi'_N|} \sum_{k=0}^{K-1} t_{k,i} \cdot (O \star Q)_{\Xi'_k}(B) \\ &= \sum_{k=0}^{K-1} \sum_{i=1}^{|\Xi'_N|} t_{k,i} \cdot (O \star Q)_{\Xi'_k}(B) \\ &= \sum_{k=0}^{K-1} \hat{p}'_k \cdot (O \star Q)_{\Xi'_k}(B) = \hat{P}'(B) \end{aligned}$$

for any measurable subset $B \subseteq \Xi'_k$ for all $0 \leq k \leq K-1$. Hence, $\Pi_{\Xi'} T = \hat{P}'$. Likewise, we have that

$$\begin{aligned} (\Pi_{\Xi'_N} T)(\xi') &= \sum_{l=0}^{K-1} \sum_{i=1}^{|\Xi'_N|} \sum_{k=0}^{K-1} t_{k,i} \cdot (\delta_{\xi'_i}(\xi') \cdot (O \star Q)_{\Xi'_k}(\Xi'_l)) \\ &= \sum_{i=1}^{|\Xi'_N|} \sum_{k=0}^{K-1} t_{k,i} \cdot \delta_{\xi'_i}(\xi') \\ &= \sum_{i=1}^{|\Xi'_N|} P'_N(\xi'_i) \cdot \delta_{\xi'_i}(\xi') = P'_N(\xi') \end{aligned}$$

for all $\xi' \in \Xi'_N$. Hence, $\Pi_{\Xi'_N} T = P'_N$. Furthermore, as by construction the Radon-Nikodym derivative $(d\hat{P}'/d(O \star Q))(\xi) = \hat{p}'_k / (O \star Q)(\Xi'_k)$ for all $\xi \in \Xi_k$, $0 \leq k \leq K-1$ we have

$$I(\hat{P}', Q) = \sum_{k=0}^{K-1} \int_{\Xi'_k} \hat{P}'(\xi') \log \left(\frac{d\hat{P}'}{d(O \star Q)}(\xi') \right) = \sum_{k=0}^{K-1} \hat{p}'_k \log(\hat{p}'_k / (O \star Q)(\Xi'_k)) \leq r.$$

Finally, we have

$$\begin{aligned} \int_{\Xi' \times \Xi'_N} \mathbb{1} \{ \|\xi' - \xi'_i\| \leq \delta \} dT(\xi', \xi'_i) &= \sum_{i=0}^{K-1} \int_{\Xi'_i \times \Xi'_N} \mathbb{1} \{ \|\xi' - \xi'_j\| \leq \delta \} dT(\xi', \xi'_j) \\ &= \sum_{i=0}^{K-1} \int_{\Xi'_i \times \Xi'_N} \mathbb{1} \{ \|\xi' - \xi'_j\| \leq \delta \} d \sum_{i=1}^{|\Xi'_N|} \sum_{k=0}^K t_{k,i} \cdot (\delta_{\xi'_i} \otimes (O \star Q)_{\Xi'_k})(\xi', \xi'_j) \\ &= \sum_{i=0}^{K-1} \sum_{i=1}^{|\Xi'_N|} \sum_{k=0}^K t_{k,i} \int_{\Xi'_i \times \Xi'_N} \mathbb{1} \{ \|\xi' - \xi'_j\| \leq \delta \} d(\delta_{\xi'_i} \otimes (O \star Q)_{\Xi'_k})(\xi', \xi'_j) \\ &= \sum_{i=1}^{|\Xi'_N|} \sum_{k=0}^K t_{k,i} \int_{\Xi'_k \times \Xi'_N} \mathbb{1} \{ \|\xi' - \xi'_j\| \leq \delta \} d(\delta_{\xi'_i} \otimes (O \star Q)_{\Xi'_k})(\xi', \xi'_j) \\ &= \sum_{i=1}^{|\Xi'_N|} \sum_{k=0}^K t_{k,i} \int_{\Xi'_k} \mathbb{1} \{ \|\xi' - \xi'_i\| \leq \delta \} d((O \star Q)_{\Xi'_k})(\xi') \\ &= \sum_{i=1}^{|\Xi'_N|} \sum_{k=0}^K t_{k,i} b(k, i) \geq 1 - \delta. \end{aligned}$$

Hence, the constructed point (Q, \hat{P}', T) is feasible in Problem (26) and attains the same objective value as the point (q, \hat{p}', t) in Problem (27). \square

C Proof of Theorem 3.3

Proof. Introduce the Lagrangian function of the optimization problem (26) as

$$\begin{aligned}
& L(Q, \hat{P}', T ; \beta, u, v, \gamma) \\
&= \int_{\Xi} \ell(z, \xi) dQ(\xi) + \beta(r - I(\hat{P}', Q)) - \int_{\Xi' \times \Xi'_N} u(\xi') dT(\xi', \xi'_i) + \int_{\Xi'} u(\xi') d\hat{P}'(\xi') \\
&\quad + \int_{\Xi' \times \Xi'_N} v(\xi'_i) dT(\xi', \xi'_i) - \int_{\Xi'_N} v(\xi'_i) dP_N(\xi'_i) + \int_{\Xi' \times \Xi'_N} \gamma \mathbb{1} \{ \|\xi' - \xi'_i\| \leq \delta \} dT(\xi', \xi'_i) - \gamma(1 - \delta) \\
&= \beta r - \int_{\Xi'_N} v(\xi'_i) dP_N(\xi'_i) - \gamma(1 - \delta) + \int_{\Xi' \times \Xi'_N} \gamma \mathbb{1} \{ \|\xi' - \xi'_i\| \leq \delta \} + v(\xi'_i) - u(\xi') dT(\xi', \xi'_i) \\
&\quad + \int_{\Xi} \ell(z, \xi) dQ(\xi) + \int_{\Xi'} u(\xi') d\hat{P}'(\xi') - \beta I(\hat{P}', Q).
\end{aligned}$$

We have $\tilde{c}^\delta(z, P'_N) = \max_{Q \in \mathcal{P}(\Xi), \hat{P}' \in \mathcal{M}_+(\Xi'), T \in \mathcal{M}_+(\Xi' \times \Xi'_N)} \min_{\beta \geq 0, u: \Xi' \rightarrow \mathbb{R}, v: \Xi'_N \rightarrow \mathbb{R}, \gamma \geq 0} L(Q, \hat{P}', T ; \beta, u, v, \gamma)$. The associated dual function is

$$\begin{aligned}
& g(\beta, u, v, \gamma) \\
&= \max \left\{ L(Q, \hat{P}', T ; \beta, u, v, \gamma) : Q \in \mathcal{P}(\Xi), \hat{P}' \in \mathcal{M}_+(\Xi'), T \in \mathcal{M}_+(\Xi' \times \Xi'_N) \right\} \\
&= \beta r - \int_{\Xi'_N} v(\xi'_i) dP_N(\xi'_i) - \gamma(1 - \delta) + \chi_\infty \{ \gamma \mathbb{1} \{ \|\xi' - \xi'_i\| \leq \delta \} + v(\xi'_i) \leq u(\xi') \quad \forall \xi' \in \Xi', \xi'_i \in \Xi'_N \} + \\
&\quad \max \left\{ \int_{\Xi} \ell(z, \xi) dQ(\xi) + \int_{\Xi'} u(\xi') d\hat{P}'(\xi') - \beta \int_{\Xi'} \log \left(\frac{d\hat{P}'}{dQ}(\xi') \right) d\hat{P}'(\xi') : Q \in \mathcal{P}(\Xi), \hat{P}' \in \mathcal{M}_+(\Xi') \right\} \\
&= \beta r - \int_{\Xi'_N} v(\xi'_i) dP_N(\xi'_i) - \gamma(1 - \delta) + \chi_\infty \{ \gamma \mathbb{1} \{ \|\xi' - \xi'_i\| \leq \delta \} + v(\xi'_i) \leq u(\xi') \quad \forall \xi' \in \Xi', \xi'_i \in \Xi'_N \} + \\
&\quad \max_{\xi \in \Xi} \left[\ell(x, \xi) + \beta \int_{\Xi'} \exp \left(\frac{u(\xi')}{\beta} + 1 \right) dO_\xi(\xi') \right]
\end{aligned}$$

where we establish the ultimate equality by observing that

$$\begin{aligned}
& \max \left\{ \int_{\Xi} \ell(z, \xi) dQ(\xi) + \int_{\Xi'} u(\xi') d\hat{P}'(\xi') - \beta \int_{\Xi'} \log \left(\frac{d\hat{P}'}{d(O \star Q)}(\xi') \right) d\hat{P}'(\xi') : Q \in \mathcal{P}(\Xi), \hat{P}' \in \mathcal{M}_+(\Xi') \right\} \\
&= \max \left\{ \int_{\Xi} \ell(z, \xi) dQ(\xi) + \int_{\Xi'} u(\xi') R(\xi') - \beta \log(R(\xi')) R(\xi') d(O \star Q)(\xi') : Q \in \mathcal{P}(\Xi), R(\xi') \geq 0 \right\} \\
&= \max \left\{ \int_{\Xi} \ell(z, \xi) dQ(\xi) + \int_{\Xi'} \left[\max_{R \geq 0} u(\xi') R - \beta \log(R) R \right] d(O \star Q)(\xi') : Q \in \mathcal{P}(\Xi) \right\} \\
&= \max \left\{ \int_{\Xi} \ell(z, \xi) dQ(\xi) + \int_{\Xi'} \beta \exp \left(\frac{u(\xi')}{\beta} - 1 \right) d(O \star Q)(\xi') : Q \in \mathcal{P}(\Xi) \right\} \\
&= \max_{\xi \in \Xi} \ell(z, \xi) + \beta \int_{\Xi'} \exp \left(\frac{u(\xi')}{\beta} - 1 \right) dO_\xi(\xi')
\end{aligned}$$

where the first equality follows from the substitution $R = d\hat{P}'/d(O \star Q)$. The fourth equality follows from the identity $\max_{R \geq 0} aR - \beta \log(R) R = aR^*(a, \beta) - \beta \log(R^*(a, \beta)) R^*(a, \beta) = \beta \exp(a/\beta - 1)$ with $R^*(a, \beta) = \exp(a/\beta - 1)$ for all a . Weak duality implies that $\tilde{c}^\delta(z, P'_N) \leq \min \{g(\beta, u, v, \gamma) : \beta \geq 0, u : \Xi' \rightarrow \mathbb{R}, v : \Xi_N \rightarrow \mathbb{R}, \gamma \geq 0\} = (28)$. Strong duality, i.e., $c^\delta(z, P'_N) = (28)$, follows from the assumed existence of the Slater points Q_s and \hat{P}'_s [6, Proposition 5.3.1] in case Ξ' is finite.

In case Ξ' is not finite we consider the slightly weaker dual

$$\begin{aligned} \inf \quad & \beta r - \int_{\Xi'} v(\xi') dP'_N(\xi') + \gamma(\delta - 1) + \max_{\xi \in \Xi} \left[\ell(z, \xi) + \beta \int_{\Xi'} \exp\left(\frac{u(\xi')}{\beta} - 1\right) dO_\xi(\xi') \right] \\ \text{s.t.} \quad & \beta \geq 0, u \in \mathcal{PC}(\Xi'), v : \Xi'_N \rightarrow \mathbb{R}, \gamma \geq 0, \\ & \gamma \mathbb{1}\{\|\xi' - \xi'_i\| \leq \delta\} + v(\xi'_i) \leq u(\xi') \quad \forall \xi' \in \Xi', \xi'_i \in \Xi'_N. \end{aligned} \quad (33)$$

where $u(\xi) \in \mathcal{PC}(\Xi')$ indicates that the dual variable u is now restricted to be piecewise constant on the pieces Ξ'_k for $0 \leq k \leq K - 1$ as defined in Section 3.1. Taking $u(\xi') = u_k$ for all $\xi' \in \Xi'_k$, $0 \leq k \leq K - 1$ and $v(\xi'_i) = v_i$ for $1 \leq i \leq |\Xi'_N|$ the weak dual (33) is clearly equivalent to

$$\begin{aligned} \inf \quad & \beta r - \sum_{i=1}^{|\Xi'_N|} v_i P'_N(\xi'_i) + \gamma(\delta - 1) + \max_{\xi \in \Xi} \left[\ell(z, \xi) + \beta \sum_{k=0}^{K-1} \exp\left(\frac{u_k}{\beta} - 1\right) O_\xi(\Xi'_k) \right] \\ \text{s.t.} \quad & \beta \geq 0, u \in \mathbb{R}^K, v \in \mathbb{R}^{|\Xi'_N|}, \gamma \geq 0, \\ & \gamma b(k, i) + v_i \leq u_k \quad 0 \leq k \leq K - 1, 1 \leq i \leq |\Xi'_N|. \end{aligned} \quad (34)$$

as $\mathbb{1}\{\|\xi' - \xi'_i\| \leq \delta\} = b(k, i)$ for all $\xi' \in \Xi'_k$ and $\xi'_i \in \Xi'_N$. The remainder of the theorem follows from Lemma C.1 and Lemma 3.2. \square

Lemma C.1. *Let $\Xi'_N = \text{supp}(P'_N)$ denote the support of P'_N and let Ξ be finite. Consider the finite convex minimization problem*

$$\begin{aligned} \inf \quad & \beta r - \sum_{i=1}^{|\Xi'_N|} v_i P'_N(\xi'_i) + \gamma(\delta - 1) + \max_{\xi \in \Xi} \left[\ell(z, \xi) + \beta \sum_{k=0}^{K-1} \exp\left(\frac{u_k}{\beta} - 1\right) O_\xi(\Xi'_k) \right] \\ \text{s.t.} \quad & \beta \geq 0, u \in \mathbb{R}^K, v \in \mathbb{R}^{|\Xi'_N|}, \gamma \geq 0, \\ & \gamma b(k, i) + v_i \leq u_k \quad 0 \leq k \leq K - 1, 1 \leq i \leq |\Xi'_N|. \end{aligned} \quad (35)$$

If there exists $Q_s \in \mathcal{P}(\Xi)$, $\hat{P}'_s \in \mathcal{P}(\Xi')$ with $I(\hat{P}'_s, Q_s) < r$ and $\text{LP}(\hat{P}'_s, P'_N) < \delta$ then (27) = (35).

Proof. It can be shown that problem (35) is the Lagrangian dual of Problem (27). As this Lagrangian dual can be derived following the proof of Theorem 3.3 almost verbatim, we omit it here for the sake of brevity. Furthermore, the construction used in the proof of Lemma 3.2 shows that the existence of $Q_s \in \mathcal{P}(\Xi)$, $\hat{P}'_s \in \mathcal{P}(\Xi')$ with $I(\hat{P}'_s, Q_s) < r$ and $\text{LP}(\hat{P}'_s, P'_N) < \delta$ ensures the existence of a Slater point in the finite dimensional optimization Problem (27). Strong duality, i.e., (27) = (35), follows then from [6, Proposition 5.3.1]. \square