

COHOMOLOGY OF SPACES OF HOPF EQUIVARIANT MAPS OF SPHERES

V.A. VASSILIEV

ABSTRACT. For any natural $k \leq n$, the rational cohomology ring of the space of continuous maps $S^{2k-1} \rightarrow S^{2n-1}$ (respectively, $S^{4k-1} \rightarrow S^{4n-1}$) equivariant under the Hopf action of the circle (respectively, of the group S^3 of unit quaternions) is naturally isomorphic to that of the Stiefel manifold $V_k(\mathbb{C}^n)$ (respectively, $V_k(\mathbb{H}^n)$).

1. MAIN THEOREM

The group S^1 of scalars $\lambda \in \mathbb{C}^1$, $|\lambda| = 1$, acts by multiplications on the unit spheres in the complex vector spaces. Denote by $E_{S^1}(k, n)$ the space of continuous maps $f : S^{2k-1} \rightarrow S^{2n-1}$ equivariant under these actions, i.e. of maps such that $f(\lambda x) = \lambda f(x)$ for any $\lambda \in S^1 \subset \mathbb{C}^1$, $x \in S^{2k-1} \subset \mathbb{C}^k$. The Stiefel manifold $V_k(\mathbb{C}^n)$ of orthonormal k -frames in \mathbb{C}^n can be identified with the space of isometric complex linear maps $\mathbb{C}^k \rightarrow \mathbb{C}^n$ of Hermitian spaces, and hence be considered as a subset of $E_{S^1}(k, n)$. In a similar way, the quaternionic Stiefel manifold $V_k(\mathbb{H}^n)$ can be considered as a subset of the space $E_{S^3}(k, n)$ of continuous maps $S^{4k-1} \rightarrow S^{4n-1}$ equivariant under the standard action (by left multiplications) of the group S^3 of quaternions of unit norm.

Theorem 1. *For any natural $k \leq n$, the maps*

$$(1) \quad H^*(E_{S^1}(k, n), \mathbb{Q}) \rightarrow H^*(V_k(\mathbb{C}^n), \mathbb{Q}) ,$$

$$(2) \quad H^*(E_{S^3}(k, n), \mathbb{Q}) \rightarrow H^*(V_k(\mathbb{H}^n), \mathbb{Q})$$

induced by inclusions are isomorphisms.

2. PRELIMINARY REMARKS

Recall (see [1]) that there are ring isomorphisms

$$(3) \quad H^*(V_k(\mathbb{C}^n), \mathbb{Z}) \simeq H^*(S^{2n-1} \times S^{2(n-1)-1} \times \dots \times S^{2(n-k+1)-1}, \mathbb{Z}) ,$$

$$(4) \quad H^*(V_k(\mathbb{H}^n), \mathbb{Z}) \simeq H^*(S^{4n-1} \times S^{4(n-1)-1} \times \dots \times S^{4(n-k+1)-1}, \mathbb{Z}).$$

We replace the space $E_{S^1}(k, n)$ by its subspace $\tilde{E}_{S^1}(k, n)$ of C^∞ -smooth equivariant maps (which is homology equivalent to it by a kind of Weierstrass approximation

2010 *Mathematics Subject Classification.* 55T99, 58D15, 55N25.

Key words and phrases. Equivariant maps, configuration space, Stiefel manifold.

theorem, cf. [4]). Using the standard techniques of finite-dimensional approximations (see e.g. [2], [4]) we will work with these spaces of maps as with manifolds of very large but finite dimensions.

A *singular m -dimensional simplex* in $\tilde{E}_{S^1}(k, n)$ is a continuous map $F : S^{2k-1} \times \Delta^m \rightarrow S^{2n-1}$, where Δ^m is the standard m -dimensional simplex and all maps $F(\cdot, \varkappa) : S^{2k-1} \rightarrow S^{2n-1}$, $\varkappa \in \Delta^m$, are of class $\tilde{E}_{S^1}(k, n)$. Again by approximation theorem, the chain complex of these simplices has the same homology as its subcomplex of C^∞ -smooth maps F .

3. SURJECTIVITY OF THE MAP (1)

Fix an orthonormal frame $\{e_1, \dots, e_k\}$ in \mathbb{C}^k and the corresponding complete flag $\mathbb{C}^1 \subset \dots \subset \mathbb{C}^{k-1} \subset \mathbb{C}^k$ where any \mathbb{C}^j is spanned by e_1, \dots, e_j . Consider \mathbb{C}^k as a subspace in \mathbb{C}^n , so that the source $S^{2k-1} \subset \mathbb{C}^k$ of our equivariant maps is a subset of their target $S^{2n-1} \subset \mathbb{C}^n$. For any $j = 1, 2, \dots, k$, denote by Ξ_j the subset in $\tilde{E}_{S^1}(k, n)$ consisting of all S^1 -equivariant smooth maps $S^{2k-1} \rightarrow S^{2n-1}$ which are identical on some big circle in S^{2k-1} cut by a complex line through the origin in $\mathbb{C}^j \subset \mathbb{C}^k$.

Definition 1. A S^1 -equivariant smooth map $f \in \Xi_j$ is *generic* if it is identical only on finitely many big circles in $S^{2k-1} \cap \mathbb{C}^j$, and for any such circle, any point $*$ of this circle and any $(2k-2)$ -dimensional transversal slice of this circle in S^{2k-1} , the differential of the map $f - \text{Id}$ from this slice to \mathbb{C}^n is injective (i.e. has rank $2k-2$) at this point $*$. (A map f can be generic in Ξ_j and not generic in some $\Xi_l, l > j$.)

Lemma 1. 1. For any $j = 1, \dots, k$, the set of generic maps $f \in \Xi_j$ is a smooth immersed submanifold of codimension $2n - 2j + 1$ in the space $\tilde{E}_{S^1}(k, n)$ in the following exact sense: for any natural m and open domain $U \subset \mathbb{R}^m$ there is a residual subset in the space of m -parametric smooth families $F : S^{2k-1} \times U \rightarrow S^{2n-1}$ of maps of class $\tilde{E}_{S^1}(k, n)$, such that for any family from this subset the points $\varkappa \in U$ such that $F(\cdot, \varkappa) \in \Xi_j$ form a smooth immersed submanifold of codimension $2n - 2j + 1$ in U .

2. For any generic map $f \in \Xi_j$, the local irreducible branches of this immersed manifold at the point f in $\tilde{E}_{S^1}(k, n)$ are in a natural one-to-one correspondence with big circles in $\mathbb{C}^j \cap S^{2k-1}$ on which this map is identical.

3. All these local branches have a natural coorientation (i.e. orientation of the normal bundle) in $\tilde{E}_{S^1}(k, n)$.

4. The set of non-generic maps $f \in \Xi_j$ has codimension at least $(2n - 2j + 1) + 2 + 2(n - k)$ in $\tilde{E}_{S^1}(k, n)$.

Proof. Items 1 and 4 follow from Thom transversality theorem. The local branch of Ξ_j corresponding to a big circle fixed by f , which is assumed in item 2, consists of maps which are identical on certain big circles neighboring to this one.

The coorientation assumed in item 3 is induced from the standard orientations of the target space S^{2n-1} and the space $\mathbb{C}P^{j-1}$ of big circles in $\mathbb{C}^j \cap S^{2k-1}$. Namely, let f be a generic point of Ξ_j and \ast a point of $\mathbb{C}^j \cap S^{2k-1}$ such that $f(\ast) = \ast$. Denote by $\Xi(\ast)$ the local branch of the variety $\Xi_j \subset \tilde{E}_{S^1}(k, n)$ at f which corresponds to the big circle $\{e^{it}\ast\}$ containing \ast . Choose an arbitrary transversal slice L of this big circle in $\mathbb{C}^j \cap S^{2k-1}$. The canonical orientation of L is induced from that of $\mathbb{C}P^{j-1}$ by the Hopf projection $\mathbb{C}^j \cap S^{2k-1} \rightarrow \mathbb{C}P^{j-1}$. By Definition 1 the differential of the map $f - \text{Id} : L \rightarrow \mathbb{C}^n$ is injective at the point \ast ; obviously it acts to the tangent space of S^{2n-1} at this point. The normal space in $\tilde{E}_{S^1}(k, n)$ of the local branch $\Xi(\ast)$ at the point f can be identified with the quotient space of $T_\ast(S^{2n-1})$ by the image of this differential. Indeed, let us realize some tangent vector of the space $\tilde{E}_{S^1}(k, n)$ at f by a one-parameter family of maps f_τ , $\tau \in (-\varepsilon, \varepsilon)$, $f_0 \equiv f$. This vector is tangent to the branch $\Xi(\ast)$ if and only if there is a family of points $\ast(\tau) \in L$, $\ast(0) = \ast$, such that $|f_\tau(\ast(\tau)) - \ast(\tau)| =_{\tau \rightarrow 0} o(|\tau|)$ or, which is the same, the vector $\frac{df_\tau(\ast)}{d\tau}(0) \in T_\ast(S^{2n-1})$ belongs to the image of the tangent space $T_\ast(L)$ under the differential of the map $f - \text{Id}$.

A canonical orientation of the latter image is defined by that of L , and the desired orientation of the quotient space (i.e. of the normal space of $E(\ast)$ at f) is specified as the factor of the standard orientation of S^{2n-1} by this one. \square

Corollary 1. *The intersection index with the set Ξ_j defines an element $[\Xi_j]$ of the group $H^{2n-2j+1}(\tilde{E}_{S^1}(k, n))$ for any $j = 1, \dots, k$. \square*

Proposition 1. *The restrictions of k cohomology classes $[\Xi_j] \in H^{2(n-j)+1}(\tilde{E}_{S^1}(k, n))$, $j = 1, \dots, k$, to the subspace $V_k(\mathbb{C}^n) \subset \tilde{E}_{S^1}(k, n)$ generate multiplicatively the ring $H^*(V_k(\mathbb{C}^n), \mathbb{Z})$.*

Proof. For any $j = 0, 1, \dots, k$ define the subset $A_{k-j} \subset V_k(\mathbb{C}^n)$ as the set of isometric linear maps $\mathbb{C}^k \rightarrow \mathbb{C}^n$ sending the first j vectors e_1, \dots, e_j of the basic frame to ie_1, \dots, ie_j respectively. Obviously, any A_j is diffeomorphic to $V_j(\mathbb{C}^{n-k+j})$, and we have inclusions $\{i \cdot \text{Id}(\mathbb{C}^k)\} \equiv A_0 \subset A_1 \subset A_2 \subset \dots \subset A_{k-1} \subset A_k \equiv V_k(\mathbb{C}^n)$, where any A_{k-j} is embedded into A_{k-j+1} as a fiber of the fiber bundle $A_{k-j+1} \rightarrow S^{2(n-j)+1}$ taking a map $\mathbb{C}^k \rightarrow \mathbb{C}^n$ into the image of the basic vector e_j . The manifolds A_{k-j+1} and Ξ_j meet transversally, and their intersection is equal to a different fiber of the same fiber bundle; hence the restriction of the cohomology class $[\Xi_j]$ to A_{k-j} (respectively, to A_{k-j+1}) is equal to zero (respectively, coincides with the class induced from the basic cohomology class of $S^{2(n-j)+1}$ by the projection map of this fiber bundle). Therefore our proposition follows by induction over the spectral sequences of these fiber bundles (and the multiplicativity of these spectral sequences). \square

Corollary 2. *The map (1) is epimorphic. \square*

4. INJECTIVITY OF (1)

For any topological space X and natural number s , denote by $B(X, s)$ the s -configuration space of X (i.e., the space of s -element subsets of X). Denote by $\pm\mathbb{Z}$ the *sign local system* of groups on $B(X, s)$: it is locally isomorphic to \mathbb{Z} , but loops in $B(X, s)$ act on its fibers by multiplications by 1 or -1 depending on the parity of the permutations of s points defined by these loops. \bar{H}_* denotes the Borel–Moore homology group (i.e. the homology group of the complex of locally finite singular chains).

Proposition 2. *There is a spectral sequence $E_r^{p,q}$ converging to the group $\tilde{H}^*(E_{S^1}(k, n), \mathbb{Z})$, whose term $E_1^{p,q}$ is trivial if $p \geq 0$, and is defined by the formula*

$$(5) \quad E_1^{p,q} \simeq \bar{H}_{-2pn-q}(B(\mathbb{C}P^{k-1}, -p), \pm\mathbb{Z})$$

for $p < 0$.

The construction of this spectral sequence essentially repeats that of the spectral sequence assumed in Theorem 3 of [4], see also [2]. Namely, we consider the vector space of all S^1 -equivariant smooth maps $S^{2k-1} \rightarrow \mathbb{C}^n$, define the discriminant set in it as the set of maps whose images contain the origin, and calculate its ‘‘homology of finite codimension’’ (which is Alexander dual to the usual cohomology group of the space of equivariant maps to $\mathbb{C}^n \setminus 0 \sim S^{2n-1}$) by using a natural simplicial resolution.

□

Proposition 3 (see [3]). *For any $j \geq 0$ and $s \geq 1$ there is a group isomorphism*

$$(6) \quad \bar{H}_j(B(\mathbb{C}P^{k-1}, s), \pm\mathbb{Z} \otimes \mathbb{Q}) \simeq H_{j-s(s-1)}(G_s(\mathbb{C}^k), \mathbb{Q}),$$

where $G_s(\mathbb{C}^k)$ is the Grassmann manifold of s -dimensional complex subspaces in \mathbb{C}^k . In particular, the left-hand group (6) is trivial if $s > k$. □

Therefore the total dimension of the group $\tilde{H}^*(E_{S^1}(k, n), \mathbb{Q})$ does not exceed

$$\sum_{s=1}^k \dim H^*(G_s(\mathbb{C}^k), \mathbb{Q}) = \sum_{s=1}^k \binom{k}{s} = 2^k - 1,$$

which by (3) is equal to the total dimension of the reduced homology group of the space $V_k(\mathbb{C}^n)$. So, (1) is an epimorphic (by Corollary 2) map, the dimension of whose source does not exceed that of the target; hence it is an isomorphism.

The bijectivity of the map (2) can be proved in exactly the same way. In particular, the first term of the spectral sequence calculating the group $\tilde{H}^*(E_{S^3}(k, n), \mathbb{Z})$ and analogous to (5) is given by

$$(7) \quad E_1^{p,q} \simeq \bar{H}_{-4pn-q}(B(\mathbb{H}P^{k-1}, -p), \pm\mathbb{Z})$$

for $p < 0$ and is trivial for $p \geq 0$; the quaternionic version of the equality (6) is

$$(8) \quad \bar{H}_j(B(\mathbb{H}P^{k-1}, s), \pm\mathbb{Z} \otimes \mathbb{Q}) \simeq H_{j-2s(s-1)}(G_s(\mathbb{H}^k), \mathbb{Q}).$$

REFERENCES

- [1] A. Borel, Sur la cohomologie des espaces fibrés principaux et des espaces homogènes de groupes de Lie compacts, *Ann. Math.*, 57 (1953), 115-207.
- [2] V. A. Vassiliev, Complements of discriminants of smooth maps: topology and applications, Second edition, Amer. Math. Soc., Providence, RI, 1994.
- [3] V. A. Vassiliev, How to calculate homology groups of spaces of nonsingular algebraic projective hypersurfaces, *Proc. Steklov Inst. Math.*, 225 (1999), 121–140 arXiv: 1407.7229
- [4] V. A. Vassiliev, Twisted homology of configuration spaces, homology of spaces of equivariant maps, and stable homology of spaces of non-resultant systems of real homogeneous polynomials, to appear, arXiv: 1809.05632

STEKLOV MATHEMATICAL INSTITUTE OF RUSSIAN ACADEMY OF SCIENCES AND NATIONAL RESEARCH UNIVERSITY HIGHER SCHOOL OF ECONOMICS

Email address: vva@mi-ras.ru