

ON 1-ABSORBING δ -PRIMARY IDEALS

ABDELHAQ EL KHALFI, NAJIB MAHDOU, ÜNSAL TEKİR, AND SUAT KOÇ

ABSTRACT. Let R be a commutative ring with nonzero identity. Let $\mathcal{I}(R)$ be the set of all ideals of R and let $\delta : \mathcal{I}(R) \rightarrow \mathcal{I}(R)$ be a function. Then δ is called an expansion function of ideals of R if whenever L, I, J are ideals of R with $J \subseteq I$, we have $L \subseteq \delta(L)$ and $\delta(J) \subseteq \delta(I)$. Let δ be an expansion function of ideals of R . In this paper, we introduce and investigate a new class of ideals that is closely related to the class of δ -primary ideals. A proper ideal I of R is said to be a 1-absorbing δ -primary ideal if whenever nonunit elements $a, b, c \in R$ and $abc \in I$, then $ab \in I$ or $c \in \delta(I)$. Moreover, we give some basic properties of this class of ideals and we study the 1-absorbing δ -primary ideals of the localization of rings, the direct product of rings and the trivial ring extensions.

1. INTRODUCTION

Throughout this paper, all rings are assumed to be commutative with nonzero identity and all modules are nonzero unital. If R is a ring, then \sqrt{I} denotes the radical of an ideal I of R , in the sense of [12, page 17]. Let also $\text{Spec}(R)$ denotes the set of all prime ideals of R .

The prime ideal, which is an important subject of ideal theory, has been widely studied by various authors. Among the many recent generalizations of the notion of prime ideals in the literature, we find the following, due to Badawi [2]. A proper ideal I of R is said to be a 2-absorbing ideal if whenever $a, b, c \in R$ and $abc \in I$, then $ab \in I$ or $ac \in I$ or $bc \in I$. In this case $\sqrt{I} = P$ is a prime ideal with $P^2 \subseteq I$ or $\sqrt{I} = P_1 \cap P_2$ where P_1, P_2 are incomparable prime ideals with $\sqrt{I}^2 \subseteq I$, cf. [2, Theorem 2.4]. Recently, Badawi and Yetkin [4] consider a new class of ideals called the class of 1-absorbing primary ideals. A proper ideal I of a ring R is called a 1-absorbing primary ideal of R if whenever nonunit elements $a, b, c \in R$ and $abc \in I$, then $ab \in I$ or $c \in \sqrt{I}$. In [13], A. Yassine et. al introduced the concept of 1-absorbing prime ideals which is a generalization of prime ideals. A proper ideal I of R is a 1-absorbing prime ideal if whenever we take nonunit elements $a, b, c \in R$ with $abc \in I$, then $ab \in I$ or $c \in I$. In this case $\sqrt{I} = P$ is a prime ideal, cf. [13, Theorem 2.3]. And if R is a ring in

2010 *Mathematics Subject Classification.* 13A99, 13C13.

Key words and phrases. 1-absorbing prime ideal, 1-absorbing δ -primary ideal, δ -primary ideal, trivial ring extension.

which exists a 1-absorbing prime ideal that is not prime, then R is a local ring, that is a ring with one maximal ideal.

Let $\mathcal{I}(R)$ be the set of all ideals of a ring R . Zhao [14] introduced the concept of expansion of ideals of R . We recall from [14] that a function $\delta : \mathcal{I}(R) \rightarrow \mathcal{I}(R)$ is called an expansion function of ideals of R if whenever L, I, J are ideals of R with $J \subseteq I$, we have $L \subseteq \delta(L)$ and $\delta(J) \subseteq \delta(I)$. Note that there are explanatory examples of expansion functions included in [14, Example 1.2] and [3, Example 1]. In addition, recall from [14] that a proper ideal I of R is said to be a δ -primary ideal of R if whenever $a, b \in R$ with $ab \in I$, we have $a \in I$ or $b \in \delta(I)$, where δ is an expansion function of ideals of R . Also, recall from [5] that a proper ideal I of R is called a δ -semiprimary ideal of R if $ab \in I$ implies $a \in \delta(I)$ or $b \in \delta(I)$. In this paper, we introduce and investigate a new concept of ideals that is closely related to the class of δ -primary ideals. A proper ideal I of R is said to be a 1-absorbing δ -primary ideal if whenever nonunit elements $a, b, c \in R$ and $abc \in I$, then $ab \in I$ or $c \in \delta(I)$. For example, let $\delta : \mathcal{I}(R) \rightarrow \mathcal{I}(R)$ such that $\delta(I) = \sqrt{I}$ for each ideal I of R . Then δ is an expansion function of ideals of R , and hence a proper ideal I of R is a 1-absorbing δ -primary ideal of R if and only if I is a 1-absorbing primary ideal of R . Among many results in this paper are given to disclose the relations between this new class and others that already exist. The reader may find it helpful to keep in mind the implications noted in the following figure.

prime ideal \longrightarrow 1-absorbing prime ideal \longrightarrow 1-absorbing δ -primary ideal

Among other things, we give an example of 1-absorbing δ -primary ideal that is not 1-absorbing prime ideal (Example 2.3). Also, we show (Theorem 2.6) that if a ring R admits a 1-absorbing δ -primary ideal of R that is not a δ -primary ideal, then R is a local ring. Moreover, we prove that if R is a chained ring with maximal ideal M , then the only 1-absorbing δ -primary ideals of R are M^2 and the δ -primary ideals of R (Theorem 2.10). Finally, we give an idea about some 1-absorbing δ -primary ideal of the localization of rings, the direct product of rings and the trivial ring extensions.

2. MAIN RESULTS

We start this section by the following definition.

Definition 2.1. *A proper ideal I of a ring R is called a 1-absorbing δ -primary ideal if whenever $abc \in I$ for some nonunit elements $a, b, c \in I$, then $ab \in I$ or $c \in \delta(I)$.*

Remark 2.2. *Let R be a ring, I a proper ideal of R and δ be an expansion function of $\mathcal{I}(R)$.*

- (1) *If $\delta(I) = I$, then I is a 1-absorbing δ -primary ideal of R if and only if it is a 1-absorbing prime ideal.*

- (2) If $\delta(I) = \sqrt{I}$, then I is a 1-absorbing δ -primary ideal of R if and only if it is a 1-absorbing primary ideal.
- (3) Every 1-absorbing prime ideal is a 1-absorbing δ -primary ideal.
- (4) Every δ -primary ideal is a 1-absorbing δ -primary ideal.
- (5) Let γ be an expansion function of the ideal of R such that $\delta(I) \subseteq \gamma(I)$. If I is a 1-absorbing δ -primary ideal of R , then I is a 1-absorbing γ -primary ideal of R .

Next, we give an example of a 1-absorbing δ -primary ideal that is not a 1-absorbing prime ideal.

Example 2.3. Let $R := K[[X_1, X_2, X_3]]$ be a ring of formal power series where K is a field. Consider the expansion function $\delta : \mathcal{I}(R) \rightarrow \mathcal{I}(R)$ defined by $\delta(I) = I + M$ where $M = (X_1, X_2, X_3)$ is the maximal ideal of R . Let $I = (X_1X_2X_3)$ be an ideal of R . Thus, I is not a 1-absorbing prime ideal of R since $X_1X_2X_3 \in I$ but neither $X_1X_2 \in I$ nor $X_3 \in I$. Now, let x, y, z be nonunit elements of R such that $xyz \in I$. Clearly I is a 1-absorbing δ -primary because $z \in \delta(I) = M$.

Proposition 2.4. (i) Every 1-absorbing δ -primary ideals is also a 2-absorbing δ -primary ideal of R .

(ii) Let I be a 1-absorbing δ -primary ideal of R and $\delta(I)$ be a radical ideal, that is, $\sqrt{\delta(I)} = \delta(I)$. Then I is a δ -semiprimary ideal of R .

Proof. (i) Let $abc \in I$ for some $a, b, c \in R$. If at least one of a, b, c is a unit of R , then we are done. So assume that a, b, c are nonunits R . Since I is a 1-absorbing δ -primary ideal of R , we get $ab \in I$ or $c \in \delta(I)$, which implies that $ab \in I$ or $ac \in \delta(I)$ or $bc \in \delta(I)$. Therefore, I is a 2-absorbing δ -primary ideal of R .

(ii) Suppose that $ab \in I$ for some $a, b \in R$. Then we may assume that a, b are nonunits. Thus $a^2b \in I$ implies that $a^2 \in I$ or $b \in \delta(I)$. Then we have $a \in \sqrt{I} \subseteq \sqrt{\delta(I)} = \delta(I)$ or $b \in \delta(I)$. Hence, I is a δ -semiprimary ideal of R . \square

The converse of previous theorem (i) is not true in general. See the following example.

Example 2.5. (2-absorbing δ -primary ideal that is not 1-absorbing δ -primary ideal)

Let $R = \mathbb{Z}$, $I = pq\mathbb{Z}$, where $p \neq q$ are prime numbers, and $\delta(I) = I + p\mathbb{Z}$. Since I is a 2-absorbing ideal, so is 2-absorbing δ -primary. However, it is easy to see that $ppq \in I$, $p^2 \notin I$ and $q \notin \delta(I)$. Thus, I is not a 1-absorbing δ -primary ideal of R .

In the next result, we show that if a ring R admits a 1-absorbing δ -primary ideal that is not a δ -primary ideal, then R is a local ring.

Theorem 2.6. Let δ be an ideal expansion. Suppose that a ring R admits a 1-absorbing δ -primary ideal that is not a δ -primary ideal. Then R is a local ring.

Proof. Assume that I is a 1-absorbing δ -primary ideal that is not a δ -primary ideal of R . Hence there exist nonunit elements $a, b \in R$ such that $ab \in I$, $a \notin I$ and $b \notin \delta(I)$. Let d be a nonunit element of R . As $dab \in I$, I is a 1-absorbing δ -primary ideal of R and $b \notin \delta(I)$, we conclude that $da \in I$. Let c be a unit element of R . Suppose that $d+c$ is a nonunit element of R . Since $(d+c)ab \in I$, I is a 1-absorbing δ -primary ideal of R and $b \notin \delta(I)$, we get that $(d+c)a = da + ca \in I$. Since $da \in I$, we conclude that $a \in I$, which gives a contradiction. Hence, $d+c$ is a unit element of R . Now, the result follows from [4, Lemma 1]. \square

Next, we give a method to construct 1-absorbing δ -primary ideals that are not δ -primary ideals.

Theorem 2.7. *Let R be a local ring with maximal ideal M and δ be an ideal expansion. Let x be a nonzero prime element of R such that $\delta(xM) \subsetneq M$. If $x \in \delta(xM)$, then xM is a 1-absorbing δ -primary ideal of R that is not a δ -primary ideal of R .*

Proof. First, we will show that xM is a 1-absorbing δ -primary ideal of R . Assume that $abc \in xM$ for some nonunit elements $a, b, c \in R$. If $ab \notin xM$, then $a \notin xR$ and $b \notin xR$, so $ab \notin xR$ because x is a prime element of R . Moreover, the fact that $abc \in xR$ and $ab \notin xR$ implies that $c \in xR \subseteq \delta(xM)$. Now, we prove that xM is not a δ -primary ideal of R . By hypothesis, we can pick an element $a \in M \setminus \delta(xM)$, hence $xa \in xM$. However, $x \notin xM$ since x is an irreducible element of R by [4, Lemma 2]. Which implies that xM is not a δ -primary ideal, this completes the proof. \square

Theorem 2.8. *Let I be a 1-absorbing δ -primary ideal of a ring R where δ is an ideal expansion and let $d \in R \setminus I$ be a nonunit element of R . Then $(I : d) = \{x \in R \mid dx \in I\}$ is a δ -primary ideal of R .*

Proof. Suppose that $ab \in (I : d)$ for some elements $a, b \in R$. Without loss of generality, we may assume that a and b are nonunit elements of R . Suppose that $a \notin (I : d)$. Since $dab \in I$ and I is a 1-absorbing δ -primary ideal of R , we conclude that $b \in \delta(I)$. So, $b \in \delta((I : d))$ and this completes the proof. \square

Proposition 2.9. *Let R be a ring, δ an ideal expansion and I be a proper ideal of R . If I is a 1-absorbing δ -primary ideal of R , then either I is a δ -semiprimary ideal of R or R is local, say with maximal ideal M , such that $M^2 \subseteq I$.*

Proof. If R is not local, then Theorem 2.6 implies that I is δ -primary and so I is a δ -semiprimary ideal of R . Now, assume that R is local with maximal ideal M such that I is not a δ -semiprimary ideal of R . Since I is proper, we infer that $I \subseteq M$. Moreover, there are $a, b \in M \setminus \delta(I)$ such that $ab \in I$. To prove that $M^2 \subseteq I$, it suffices to show that $xy \in I$ for all $x, y \in M$. Let $x, y \in M$. Then $xyab \in I$. Since $xy, a, b \in M$, $b \notin \delta(I)$ and I is a 1-absorbing δ -primary ideal, we conclude that $xya \in I$. Again, since

$x, y, a \in M$, $a \notin \delta(I)$ and I is a 1-absorbing δ -primary ideal, we have that $xy \in I$. \square

Recall that a ring R is a chained ring if the set of all ideals of R is linearly ordered by inclusion. Moreover, R is said to be an arithmetical ring if R_M is a chained ring for each maximal ideal M of R . We next determinate the 1-absorbing δ -primary ideals of a chained ring.

Theorem 2.10. *Let R be a chained ring with maximal ideal M , δ an ideal expansion and I be a proper ideal of R such that $I \neq M^2$. Then I is a 1-absorbing δ -primary ideal of R if and only if I is a δ -primary ideal of R .*

Proof. We need only prove the “only if” assertion. Let I be a 1-absorbing δ -primary ideal. Thus, Proposition 2.9 gives that either I is a δ -semiprimary ideal of R or $M^2 \subseteq I$. First, assume that I is a δ -semiprimary ideal of R and $ab \in I$ for some nonunit elements $a, b \in R$ such that $b \notin \delta(I)$. Hence, $a \in \delta(I)$. Now, since R is a chained ring, we conclude that $a \in bR$ and thus $a = br$ for some nonunit element $r \in R$. As $brb \in I$, $b \notin \delta(I)$ and I is a 1-absorbing δ -primary ideal of R , we conclude that $a = br \in I$. Which gives that I is a δ -primary ideal of R . Now, we suppose that $M^2 \subseteq I$. We may assume that $M \neq I$. Thus, we can pick $a \in M \setminus I$ and $b \in I \setminus M^2$. Then $b \in aR$ since R is a chained ring. So, $b = ar$ for some nonunit element $r \in R$ and thus $b \in M^2$, a contradiction. This completes the proof. \square

In view of Theorem 2.10, we have the following result.

Corollary 2.11. *Let R be an arithmetical ring with Jacobson radical M and I be a proper ideal of R such that $I \neq M^2$. Then I is a 1-absorbing δ -primary ideal of R if and only if I is a δ -primary ideal of R .*

Proof. Assume that R is local with maximal ideal M . Since R is an arithmetical ring, we conclude that $R = R_M$ is a chained ring and thus the claim follows from Theorem 2.10. In the remaining case, suppose that R is not a local ring. Then the result follows by Theorem 2.6. \square

Proposition 2.12. *Let R be a local ring with principal maximal ideal M , δ an ideal expansion and I be a proper ideal of R . Then I is a 1-absorbing δ -primary ideal of R if and only if either I is a δ -primary ideal of R or $M^2 \subseteq I$.*

In addition, if $\sqrt{I} \subseteq \delta(I)$ then I is a 1-absorbing δ -primary ideal of R if and only if I is a δ -primary ideal of R .

Proof. By Remark 2.2(4) and Proposition 2.9, we need only prove that if I is a 1-absorbing δ -primary ideal of R which is a δ -semiprimary ideal then I is a δ -primary ideal (along with the hypothesis that R be a local ring with principal maximal ideal M). Also, we may assume that $\delta(I) \neq R$. Set $M = xR$ and let a and b be nonunit elements of R such that $b \notin \delta(I)$ and $ab \in I$. Since I is a δ -semiprimary ideal of R , we get that $a \in \delta(I)$. Moreover, $a = rx$ for some $r \in R$. If r is a unit element of R then $M = \delta(I)$ and thus I

is a δ -primary ideal. If r is a nonunit element of R then $rx = ab \in I$. That implies $a = rx \in I$ since I is a 1-absorbing δ -primary ideal. This completes the proof. The in addition statement is clear. \square

Proposition 2.13. *Let $\{J_i \mid i \in D\}$ be a directed set of 1-absorbing δ -primary ideals of R , where δ is an ideal expansion. Then the ideal $J = \cup_{i \in D} J_i$ is a 1-absorbing δ -primary ideal of R .*

Proof. Let $abc \in J$, then $abc \in J_i$ for some $i \in D$. Since J_i is a 1-absorbing δ -primary ideal of R , $ab \in J_i$ or $c \in \delta(J_i) \subseteq \delta(J)$. Hence, J is a 1-absorbing δ -primary ideal of R . \square

Proposition 2.14. *Let I be a 1-absorbing δ -primary ideal of R such that $\sqrt{\delta(I)} = \delta(\sqrt{I})$, where δ is an ideal expansion. Then, \sqrt{I} is a δ -primary ideal of R .*

Proof. Let $ab \in \sqrt{I}$ such that $a \notin \sqrt{I}$. Hence, there exists a positive integer n such that $(ab)^n \in I$. So, $a^m a^m b^n \in I$ for some positive integer m . Since I is a 1-absorbing δ -primary ideal of R and $a^{2m} \notin I$, we conclude that $b^n \in \delta(I)$. That implies $b \in \sqrt{\delta(I)} = \delta(\sqrt{I})$ and so \sqrt{I} is a δ -primary ideal of R . \square

Proposition 2.15. *Let I be a proper ideal of a ring R and δ be an ideal expansion such that $\delta(\delta(I)) = \delta(I)$. Then $\delta(I)$ is a 1-absorbing δ -primary ideal of R if and only if $\delta(I)$ is a 1-absorbing prime ideal of R .*

Proof. By Remark 2.2(3), we need only prove the “only if” assertion. Let $abc \in \delta(I)$ for some nonunit elements $a, b, c \in R$. Hence $ab \in \delta(I)$ or $c \in \delta(\delta(I)) = \delta(I)$. Thus $\delta(I)$ is a 1-absorbing prime ideal of R . \square

Proposition 2.16. *Let R be a ring, I a proper ideal of R and δ be an ideal expansion. Then I is a 1-absorbing δ -primary ideal if and only if whenever $I_1 I_2 I_3 \subseteq I$ for some proper ideals I_1, I_2 and I_3 of R , then $I_1 I_2 \subseteq I$ or $I_3 \subseteq \delta(I)$.*

Proof. It suffices to prove the “if” assertion. Suppose that I is a 1-absorbing δ -primary ideal and let I_1, I_2 and I_3 be proper ideals of R such that $I_1 I_2 I_3 \subseteq I$ and $I_3 \not\subseteq \delta(I)$. Thus $abc \in I$ for every $a \in I_1, b \in I_2$ and $c \in I_3 \setminus \delta(I)$. Since I is a 1-absorbing δ -primary ideal, we then have $I_1 I_2 \subseteq I$, as desired. \square

Recall from [14] that an ideal expansion δ is said to be intersection preserving if $\delta(I_1 \cap I_2 \cap \dots \cap I_n) = \delta(I_1) \cap \delta(I_2) \cap \dots \cap \delta(I_n)$ for any ideals I_1, \dots, I_n of R .

Proposition 2.17. *Let δ be an intersection preserving ideal expansion. If I_1, I_2, \dots, I_n are 1-absorbing δ -primary ideals of R , and $\delta(I_i) = P$ for all $i \in \{1, 2, \dots, n\}$, then $I_1 \cap I_2 \cap \dots \cap I_n$ is a 1-absorbing δ -primary ideal of R .*

Proof. Let $abc \in J = I_1 \cap I_2 \cap \dots \cap I_n$ such that $ab \notin J$. Let $i \in \{1, 2, \dots, n\}$ such that $ab \notin I_i$. Since $abc \in I_i$ and I_i is a 1-absorbing δ -primary ideal, we conclude that $c \in \delta(I_i) = \delta(J)$. Therefore, J is a 1-absorbing δ -primary ideal of R . \square

Proposition 2.18. *Let R be a ring and δ be an expansion function of $\mathcal{I}(R)$. Then the following statements are equivalent:*

- (1) *Every proper principal ideal is a 1-absorbing δ -primary ideal of R .*
- (2) *Every proper ideal is a 1-absorbing δ -primary ideal of R .*

Proof. Assume that (1) holds and let I be a proper ideal of R . Let a, b, c be nonunit elements of R such that $abc \in I$. Hence $abc \in abcR$ which implies that $ab \in abcR \subseteq I$ or $c \in abcR \subseteq \delta(I)$. Therefore I is a 1-absorbing δ -primary ideal of R . The converse is clear. \square

An expansion function δ of $\mathcal{I}(R)$ is said to satisfy *condition (*)* if $\delta(I) \neq R$ for each proper ideal I of R . Note that the identity function and radical operation are examples of expansion functions satisfying condition (*).

Theorem 2.19. *Let R be a ring and δ an expansion function of $\mathcal{I}(R)$ satisfying condition (*) and $\delta(\text{Jac}(R)) = \text{Jac}(R)$. Suppose that $\delta(xI) = x\delta(I)$ for every proper ideal I of R and every $x \in R$. The following statements are equivalent.*

- (i) *Every proper principal ideal is a 1-absorbing δ -primary ideal of R .*
- (ii) *Every proper ideal is a 1-absorbing δ -primary ideal of R .*
- (iii) *R is local with $\text{Jac}(R)^2 = (0)$, where $\text{Jac}(R) = m$ is the unique maximal ideal.*

Proof. (i) \Leftrightarrow (ii) Follows from Proposition 2.18.

(i) \Rightarrow (iii) Assume that every proper ideal is a 1-absorbing δ -primary ideal R . Choose $x, y \in \text{Jac}(R)$. Now, we will show that $xy = 0$. If x or y is zero, then we are done. Assume that $x, y \neq 0$. Since $x^2y \in (x^2y)$ and (x^2y) is a 1-absorbing δ -primary ideal, we conclude that $x^2 \in (x^2y)$ or $y \in \delta((x^2y)) = y\delta((x^2))$. Suppose that $y \in y\delta((x^2))$. Then there exists $a \in \delta((x^2)) \subseteq \delta(\text{Jac}(R)) = \text{Jac}(R)$ such that $y = ya$. Which implies that $y(1-a) = 0$. Since $1-a$ is unit, we have $y = 0$, which is a contradiction. Thus we have, $x^2 \in (x^2y)$. Then we can write $x^2 = rx^2y$ for some $r \in R$. This implies that $x^2(1-ry) = 0$. Since $1-ry$ is unit, we have $x^2 = 0$. Likewise, we get $y^2 = 0$. Now, choose another $z \in \text{Jac}(R)$. Since $xyz \in (xyz)$ and (xyz) is a 1-absorbing δ -primary, we get $xy \in (xyz)$ or $z \in \delta((xyz)) = z\delta((xy))$. First, assume that $xy \in (xyz)$. Then there exists $r \in R$ such that $xy = rxyz$, which implies that $xy(1-rz) = 0$. Since $1-rz$ is unit, we have $xy = 0$ which completes the proof. Now, assume that $xy \notin (xyz)$, that is, $z \in \delta((xyz)) = z\delta((xy))$. Then there exists $a \in \delta((xy)) \subseteq \text{Jac}(R)$ such that $z = za$. This implies that $z(1-a) = 0$ so that $z = 0$. Now, choose $z = x+y$. Then by above argument, we have either $xy = 0$ or $z = x+y = 0$. If $z = x+y = 0$, then we have $x = -y$ and so $xy = -y^2 = 0$ which completes the proof. Therefore, $\text{Jac}(R)^2 = (0)$.

Now, we will show that R is a local ring. Choose maximal ideals M_1, M_2 of R . Now, put $I = M_1 \cap M_2$. Since $M_1^2 M_2 \subseteq I$ and I is a 1-absorbing δ -primary ideal, we have either $M_1^2 \subseteq I \subseteq M_2$ or $M_2 \subseteq \delta(I) \subseteq \delta(M_1)$. **Case 1:** Suppose that $M_1^2 \subseteq M_2$. Since M_2 is prime, clearly we have $M_1 \subseteq M_2$

which implies that $M_1 = M_2$. **Case 2:** Suppose that $M_2 \subseteq \delta(M_1)$. Since δ satisfies condition (*), $\delta(M_1)$ is proper. As $M_1 \subseteq \delta(M_1)$ and M_1 is a maximal ideal, we have $M_1 = \delta(M_1)$. Then we get $M_2 \subseteq M_1$, which implies that $M_1 = M_2$. Therefore, R is a local ring.

(iii) \Rightarrow (i) Suppose that R is a local ring with $Jac(R)^2 = (0)$. Let I be a proper ideal of R and $abc \in I$ for some nonunits $a, b, c \in R$. Then $a, b, c \in Jac(R)$ since R is local. As $Jac(R)^2 = (0)$, we have $ab = 0 \in I$. Therefore, I is a 1-absorbing δ -primary ideal of R . \square

It can be easily seen that, in Theorem 2.19, (iii) always implies (i) without any assumption on δ . But we give some examples showing that the converse is not true if we drop the aforementioned assumptions on δ .

Example 2.20. Let $R = \mathbb{Z}_{p^3}$, where p is a prime number and $\delta(I) = R$ for every proper ideal I of R . Note that δ does not satisfy condition (*) and note that every ideal I of R is 1-absorbing δ -primary. Thus $Jac(R)^2 \neq (0)$, while R is a local ring.

Example 2.21. Let k be a field and consider the formal power series ring $R = k[[X]]$. Then R is a local ring with unique maximal ideal $m = (X)$. Define expansion function δ as $\delta(I) = \sqrt{I}$ for every ideal I of R . Then it is easy to see that every ideal of R is a 1-absorbing δ -primary ideal. Also, it is clear that δ satisfies condition (*) and $\delta(Jac(R)) = Jac(R)$ but not satisfy the condition $\delta(xI) = x\delta(I)$. Furthermore, $Jac(R)^2 \neq (0)$. Thus Theorem 2.19 fails without assumption $\delta(xI) = x\delta(I)$.

Corollary 2.22. Let R be a ring. The following statements are equivalent.

- (i) Every proper ideal is a 1-absorbing prime ideal of R .
- (ii) Every proper principal ideal is a 1-absorbing prime ideal of R .
- (iii) R is local with $Jac(R)^2 = (0)$.

Proof. (i) \Leftrightarrow (ii) Follows from Proposition 2.18.

(ii) \Rightarrow (iii) Let δ be the identity expansion function, that is, $\delta(I) = I$ for every ideal I of R . Note that δ satisfies all axioms in Theorem 2.19. Then R is a local ring with $Jac(R)^2 = (0)$.

(iii) \Rightarrow (i) It is similar to Theorem 2.19 (iii) \Rightarrow (i). \square

An ideal expansion δ is called a prime expansion if for any 1-absorbing δ -primary ideal I of R , $\delta(I)$ is a prime ideal of R .

Proposition 2.23. Let R be a local ring with maximal ideal M and δ be a prime expansion function of $\mathcal{I}(R)$. Assume that one of the following conditions holds:

- (1) $Spec(R) = \{\delta(0)\}$.
- (2) $Spec(R) = \{\delta(0), M\}$ and $\delta(0)M = 0$.

Then every proper ideal of R is 1-absorbing δ -primary.

Proof. Let I be a proper ideal of R and assume that (1) holds. Since δ is a prime expansion, we conclude that $\delta(I) = \delta(0)$ is the maximal ideal

of R . Clearly I is a 1-absorbing δ -primary ideal of R . Now, assume that $\text{Spec}(R) = \{\delta(0), M\}$ and $\delta(0)M = 0$. If $\delta(I) = M$, we have then I is 1-absorbing δ -primary. In the remaining case, $\delta(I) = \delta(0)$. Let $abc \in I$ for some nonunit elements $a, b, c \in R$ such that $c \notin \delta(0)$. As $I \subseteq \delta(0)$, we get that either $a \in \delta(0)$ or $b \in \delta(0)$. Thus $ab = 0 \in I$ which gives that I is a 1-absorbing δ -primary ideal of R . \square

Let $f : R \rightarrow S$ be a ring homomorphism and δ, γ expansion functions of $\mathcal{I}(R)$ and $\mathcal{I}(S)$ respectively. Recall from [3] that f is called a $\delta\gamma$ -homomorphism if $\delta(f^{-1}(I)) = f^{-1}(\gamma(I))$ for each ideal I of S . Also note that if f is a $\delta\gamma$ -epimorphism and I is an ideal of R containing $\ker(f)$, then $\gamma(f(I)) = f(\delta(I))$.

Theorem 2.24. *Let $f : R \rightarrow S$ be a ring $\delta\gamma$ -homomorphism where δ, γ are expansion functions of $\mathcal{I}(R)$ and $\mathcal{I}(S)$ respectively. Suppose that $f(a)$ is nonunit in S for every nonunit element a in R . Then the following statements hold.*

- (1) *If J is a 1-absorbing γ -primary ideal of S , then $f^{-1}(J)$ is a 1-absorbing δ -primary ideal of R .*
- (2) *If f is an epimorphism and I is a proper ideal of R containing $\ker(f)$, then I is a 1-absorbing δ -primary ideal of R if and only if $f(I)$ is a 1-absorbing γ -primary ideal of S .*

Proof. (1) Assume that $abc \in f^{-1}(J)$, for some nonunit elements $a, b, c \in R$. Then $f(a)f(b)f(c) \in J$. Thus $f(a)f(b) \in J$ or $f(c) \in \gamma(J)$, which implies that $ab \in f^{-1}(J)$ or $c \in f^{-1}(\gamma(J)) = \delta(f^{-1}(J))$. Therefore, $f^{-1}(J)$ is a 1-absorbing δ -primary ideal of R .

(2) Suppose that $f(I)$ is an 1-absorbing γ -primary ideal of S . Since $I = f^{-1}(f(I))$, we conclude that I is a 1-absorbing δ -primary ideal of R by (1). Conversely, let x, y, z be nonunit elements of S with $xyz \in f(I)$. Then there exist $a, b, c \in R$ such that $x = f(a)$, $y = f(b)$ and $z = f(c)$ with $f(abc) = xyz \in f(I)$. Since $\ker(f) \subseteq I$, we then have $abc \in I$. Since I is a 1-absorbing δ -primary ideal of R and $abc \in I$, we conclude that $ab \in I$ or $c \in \delta(I)$ which gives that $xy \in f(I)$ or $z \in f(\delta(I)) = \gamma(f(I))$. Thus $f(I)$ is a 1-absorbing δ -primary ideal of S . \square

Let δ be an expansion function of $\mathcal{I}(\mathcal{R})$ and I an ideal of R . Then the function $\bar{\delta} : \frac{R}{I} \rightarrow \frac{R}{I}$ defined by $\bar{\delta}(\frac{J}{I}) = \frac{\delta(J)}{I}$ for all ideals $I \subseteq J$, becomes an expansion function of $\frac{R}{I}$. Then, we have the following result.

Corollary 2.25. *Let R be a ring, δ an expansion function of $\mathcal{I}(\mathcal{R})$ and $I \subseteq J$ be proper ideals of R . Assume that $a + I$ is a nonunit element of $\frac{R}{I}$ for every nonunit element $a \in R$. Then J is a 1-absorbing δ -primary ideal of R if and only if $\frac{J}{I}$ is a 1-absorbing $\bar{\delta}$ -primary ideal of $\frac{R}{I}$.*

Proposition 2.26. *Let S be a multiplicatively closed subset of a ring R and δ_S an expansion function of $\mathcal{I}(S^{-1}R)$ such that $\delta_S(S^{-1}I) = S^{-1}(\delta(I))$*

for each ideal I of R . If I is a 1-absorbing δ -primary ideal of R such that $I \cap S = \emptyset$, then $S^{-1}I$ is a 1-absorbing δ_S -primary ideal of $S^{-1}R$.

Proof. Let I be a 1-absorbing δ -primary ideal of R such that $I \cap S = \emptyset$ and $\frac{a}{s} \frac{b}{t} \frac{c}{r} \in S^{-1}I$ for some nonunit elements $a, b, c \in R$ and $s, t, r \in S$ such that $\frac{a}{s} \frac{b}{t} \notin S^{-1}I$. Then $abc \in I$ for some $x \in S$. Since I is a 1-absorbing δ -primary and $xab \notin I$, we conclude that $c \in \delta(I)$. Thus $\frac{c}{r} \in S^{-1}(\delta(I)) = \delta_S(S^{-1}I)$ which completes the proof. \square

Let S be a multiplicatively closed subset of a ring R and I an ideal of R . The next example shows that if $S^{-1}I$ is a 1-absorbing δ_S -primary ideal of $S^{-1}R$, then I need not to be a 1-absorbing δ -primary ideal of R .

Example 2.27. Let $p \neq q$ be two prime numbers. Set $I = pq\mathbb{Z}$ and δ be an ideal expansion such that $\delta(I) = I + q\mathbb{Z}$ for each ideal I of \mathbb{Z} . Clearly, I is not a 1-absorbing δ -primary ideal of \mathbb{Z} because $qpp \in I$ but neither $q^2 \in I$ nor $p \in \delta(I)$. Now, let $S = \mathbb{Z} \setminus p\mathbb{Z}$ and note that $S^{-1}I = S^{-1}(p\mathbb{Z})$. Let $\frac{a}{r_1} \frac{b}{r_2} \frac{c}{r_3} \in S^{-1}I$ for some nonunit elements $\frac{a}{r_1}, \frac{b}{r_2}, \frac{c}{r_3} \in S^{-1}\mathbb{Z}$. Note that $\frac{x}{r} \in S^{-1}\mathbb{Z}$ is nonunit if and only if $x \in p\mathbb{Z}$. Thus $a \in p\mathbb{Z}$ and $b \in p\mathbb{Z}$. Which gives that $\frac{a}{r_1} \frac{b}{r_2} \in S^{-1}I$ and hence $S^{-1}I$ is a 1-absorbing δ_S -primary ideal.

Let R_1 and R_2 be two rings, let δ_i be an expansion function of $\mathcal{I}(\mathcal{R}_i)$ for each $i \in \{1, 2\}$ and $R = R_1 \times R_2$. For a proper ideal $I_1 \times I_2$, the function δ_\times defined by $\delta_\times(I_1 \times I_2) = \delta_1(I_1) \times \delta_2(I_2)$ is an expansion function of $\mathcal{I}(R)$. The following result characterizes the 1-absorbing δ -primary ideals of the direct product of rings.

Theorem 2.28. Let R_1 and R_2 be rings, $R = R_1 \times R_2$ and let δ_i be an expansion function of $\mathcal{I}(\mathcal{R}_i)$ for $i = 1, 2$. Then the following statements are equivalent:

- (1) I is a 1-absorbing δ_\times -primary ideal of R .
- (2) I is a δ_\times -primary ideal of R .
- (3) Either $I = I_1 \times R_2$, where I_1 is a δ_1 -primary ideal of R_1 or $I = R_1 \times I_2$, where I_2 is a δ_2 -primary ideal of R_2 or $I = I_1 \times I_2$, where I_1 and I_2 are proper ideals of R_1, R_2 , respectively with $\delta_1(I_1) = R_1$ and $\delta_2(I_2) = R_2$.

Proof. (1) \Leftrightarrow (2). This follows from Theorem 2.6.

(2) \Leftrightarrow (3) Let I be a δ_\times -primary ideal of R . Hence I has the form $I = I_1 \times I_2$ where I_1 and I_2 are ideals of R_1 and R_2 respectively. Without loss of generality, we may assume that $I = I_1 \times R_2$ for some proper ideal I_1 of R_1 . We show that I_1 is a δ -primary ideal of R_1 . Deny. Then there are $a, b \in R_1$ such that $ab \in I_1, a \notin I_1$ and $b \notin \delta_1(I_1)$. Hence $(a, 1)(b, 1) \in I_1 \times R_2$. Which implies that $(a, 1) \in I_1 \times R_2$ or $(b, 1) \in \delta_\times(I_1 \times R_2)$ and so $a \in I_1$ or $b \in \delta_1(I_1)$, which gives a contradiction. Now suppose that both I_1 and I_2 are proper. As $(1, 0)(0, 1) \in I_1 \times I_2$ and $(1, 0), (0, 1) \notin I_1 \times I_2$, we have $(1, 0), (0, 1) \in \delta_\times(I_1 \times I_2) = \delta_1(I_1) \times \delta_2(I_2)$. Therefore $\delta_1(I_1) = R_1$ and $\delta_2(I_2) = R_2$. The converse is clear. \square

The following example proves that the condition “ $\delta(I_i) = P$ for all $i \in \{1, 2, \dots, n\}$ ” is necessary in Proposition 2.17.

Example 2.29. Consider $R = \mathbb{Z} \times \mathbb{Z}$, $I_1 = 4\mathbb{Z} \times \mathbb{Z}$ and $I_2 = \mathbb{Z} \times 9\mathbb{Z}$. Let δ be an expansion function of $\mathcal{I}(\mathbb{Z})$ such that for every ideal I of \mathbb{Z} we have $\delta(I) = \sqrt{I} + J$ where $J = 2\mathbb{Z}$. Thus $\delta_{\times}(I_1) = 2\mathbb{Z} \times \mathbb{Z}$ and $\delta_{\times}(I_2) = \mathbb{Z} \times \mathbb{Z}$. Moreover, I_1 and I_2 are 1-absorbing δ_{\times} -primary ideal. But $I_1 \cap I_2 = 4\mathbb{Z} \times 9\mathbb{Z}$ is not a 1-absorbing δ_{\times} -primary ideal by Theorem 2.28.

Let A be a ring and E an A -module. Then $A \times E$, the *trivial (ring) extension of A by E* , is the ring whose additive structure is that of the external direct sum $A \oplus E$ and whose multiplication is defined by $(a, e)(b, f) := (ab, af + be)$ for all $a, b \in A$ and all $e, f \in E$. (This construction is also known by other terminology and other notation, such as the *idealization* $A(+E)$.) The basic properties of trivial ring extensions are summarized in the books [10], [9]. Trivial ring extensions have been studied or generalized extensively, often because of their usefulness in constructing new classes of examples of rings satisfying various properties (cf. [1, 6, 7, 11]). In addition, for an ideal I of A and a submodule F of E , $I \times F$ is an ideal of $A \times E$ if and only if $IE \subseteq F$. Moreover, for an expansion function δ of A , it is clear that δ_{\times} defined as $\delta_{\times}(I \times F) = \delta(I) \times E$ is an expansion function of $A \times E$. Also as usual, if $c \in A$ then $(F : c) = \{e \in E \mid ce \in F\}$.

Theorem 2.30. Let A be a ring, E an A -module and δ be an expansion function of $\mathcal{I}(A)$. Let I be an ideal of A and F a submodule of E such that $IE \subseteq F$. Then the following statement hold:

- (1) If $I \times F$ is a 1-absorbing δ_{\times} -primary ideal of $A \times E$, then I is a 1-absorbing δ -primary ideal of A .
- (2) Assume that $(F : c) = F$ for every $c \in A \setminus I$. Then $I \times F$ is a 1-absorbing δ_{\times} -primary ideal of $A \times E$ if and only if I is a 1-absorbing δ -primary ideal of A .

Proof. (1) Assume that $I \times F$ is a 1-absorbing δ_{\times} -primary ideal of $A \times E$ and let a, b, c be nonunit elements of A such that $abc \in I$. Thus $(a, 0)(b, 0)(c, 0) = (abc, 0) \in I \times F$ which implies that $(a, 0)(b, 0) \in I \times F$ or $(c, 0) \in \delta_{\times}(I \times F) = \delta(I) \times E$. Therefore $ab \in I$ or $c \in \delta(I)$ and so (1) holds.

(2) By (1), it suffices to prove the "if" assertion. Let $(a, s), (b, t), (c, r)$ be nonunit elements of $A \times E$ such that $(a, s)(b, t)(c, r) = (abc, bcs + act + abr) \in I \times F$. Clearly, $abc \in I$ and so $ab \in I$ or $c \in \delta(I)$ since I is a 1-absorbing δ -primary ideal of A . If $c \in \delta(I)$, then $(c, r) \in \delta(I) \times E = \delta_{\times}(I \times F)$. Hence, we may assume that $c \notin \delta(I)$. Then $ab \in I$. As $bcs + act + abr \in F$ and $abr \in F$, we get that $bcs + act \in F$. This implies $bs + at \in (F : c) = F$ and so $(a, s)(b, t) = (ab, at + bs) \in I \times F$. Therefore $I \times F$ is a 1-absorbing δ_{\times} -primary ideal of $A \times E$. □

Corollary 2.31. Let A be a ring, E an A -module and δ be an expansion function of $\mathcal{I}(A)$. Let I be a proper ideal of A . Then $I \times E$ is a 1-absorbing

δ_{\times} -primary ideal of $A \times E$ if and only if I is a 1-absorbing δ -primary ideal of A .

REFERENCES

- [1] D. D. Anderson and M. Winders, *Idealization of a module*, J. Commut. Algebra 1 (2009), no. 1, 3–56.
- [2] A. Badawi, *On 2-absorbing ideals of commutative rings*, Bull. Austral. Math. Soc. 75(3)(2007), 417–429.
- [3] A. Badawi, B. Fahid, *On weakly 2-absorbing δ -primary ideals of commutative rings*, Georgian. Math. J, 27(4) 2017, 503-516.
- [4] A. Badawi and Y. Celikel, *On 1-absorbing primary ideals of a commutative rings*, J. Algebra Appl. (2020), 2050111.
- [5] A.Badawi, D. Sonmez, G.Yesilot: *On weakly δ -semiprimary ideals of commutative rings*. Algebra Colloq. 25 (2018), 387-398.
- [6] D.E Dobbs, A. Elkhalfi and N. Mahdou, *Trivial extensions satisfying certain valuation-like properties*, Comm. Algebra 47(5) (2019), 2060–2077.
- [7] T. Dumitrescu, N. Mahdou and Y. Zahir, *Radical factorization for trivial extension and amalgamated duplication rings*, J. Algebra Appl., to appear.
- [8] B. Fahid, Z. Dongsheng, *2-absorbing δ -primary ideals in commutative rings*, Kyungpook Mathematical Journal 57(2017), no. 2, 193-198.
- [9] S. Glaz, *Commutative Coherent Rings*, Lecture Notes in Math. 1371, Springer-Verlag, Berlin, 1989.
- [10] J. A. Huckaba, *Commutative Rings with Zero Divisors*, Dekker, New York, 1988.
- [11] S. Kabbaj and N. Mahdou, *Trivial extensions defined by coherent-like conditions*, Comm. Algebra 32 (2004), no. 10, 3937–3953.
- [12] I. Kaplansky, *Commutative Rings*, rev. ed., Univ. Chicago Press, Chicago, 1974.
- [13] A. Yassine, M. J. Nikmehr, R. Nikandish, *On 1-absorbing prime ideals of commutative rings*, J. Algebra Appl., (2020) to appear.
- [14] D. Zhao, *δ -primary ideals of commutative rings*, Kyungpook Math. J. 41 (1) (2001), 17–22.

ABDELHAQ EL KHALFI, LABORATORY OF MODELLING AND MATHEMATICAL STRUCTURES, DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCE AND TECHNOLOGY OF FEZ, BOX 2202, UNIVERSITY S.M. BEN ABDELLAH FEZ, MOROCCO.

E – mail address : abdelhaq.elkhalfi@usmba.ac.ma

NAJIB MAHDOU, LABORATORY OF MODELLING AND MATHEMATICAL STRUCTURES, DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCE AND TECHNOLOGY OF FEZ, BOX 2202, UNIVERSITY S.M. BEN ABDELLAH FEZ, MOROCCO.

E – mail address : mahdou@hotmail.com

ÜNSAL TEKİR, DEPARTMENT OF MATHEMATICS, MARMARA UNIVERSITY, ISTANBUL, TURKEY.

E – mail address : utekir@marmara.edu.tr

SUAT KOÇ, DEPARTMENT OF MATHEMATICS, MARMARA UNIVERSITY, ISTANBUL, TURKEY.

E – mail address : suat.koc@marmara.edu.tr