

MEASURE ESTIMATION ON MANIFOLDS: AN OPTIMAL TRANSPORT APPROACH

VINCENT DIVOL

Courant Institute of Mathematical Science, New York University

Center for Data Science, New York University

ABSTRACT. Assume that we observe i.i.d. points lying close to some unknown d -dimensional C^k submanifold M in a possibly high-dimensional space. We study the problem of reconstructing the probability distribution generating the sample. After remarking that this problem is degenerate for a large class of standard losses (L_p , Hellinger, total variation, etc.), we focus on the Wasserstein loss, for which we build an estimator, based on kernel density estimation, whose rate of convergence depends on d and the regularity $s \leq k - 1$ of the underlying density, but not on the ambient dimension. In particular, we show that the estimator is minimax and matches previous rates in the literature in the case where the manifold M is a d -dimensional cube. The related problem of the estimation of the volume measure of M for the Wasserstein loss is also considered, for which a minimax estimator is exhibited.

1. INTRODUCTION

Density estimation is one of the most fundamental tasks in non-parametric statistics. If efficient methods (from both a theoretical and a practical point of view) exist when the ambient space is of low dimension, minimax rates of estimation become increasingly slow as the dimension increases. To overcome this so-called *curse of dimensionality*, some structural assumptions on the underlying probability are to be made in moderate to high dimensions, which may take different forms, including e.g. the existence of a parametric component [LLW07], the single-index model [LZZL13], sparsity assumptions [Tib96], or constraints on the shape of the support. We focus in this work on the latter, namely on the case where the probability distribution μ generating the observations is assumed to be concentrated around a submanifold M of \mathbb{R}^D , of dimension d smaller than D . This assumption, known as the manifold assumption, has been fruitfully studied, with an emphasis put on reconstructing different geometric quantities related to the manifold, such as M itself [GPPIW12, AL18, AL19, Div21a], its homology groups [NSW08, BRS⁺12], its dimension [HA05, LJM09, KRW19] or its reach [AKC⁺19, BHHS21]. The topic of density estimation in the manifold setting has itself been studied for over thirty years, with the emphasis initially being put on reconstructing the density in the case where the manifold M is given—think for instance of datasets lying on the space of orthogonal matrices—notable works including [Hen90, HJR93, Pel05, CGK⁺20]. Less attention has been dedicated to the more general setting where the manifold M is unknown and acts as a nuisance parameter. Kernel density estimators on manifolds are designed in [BS17, WW20], where rates are exhibited, respectively in the case where the manifold has a boundary and in the case where the density is Hölder continuous. In [BH19], kernel density estimators are shown to be minimax, and an adaptive procedure is designed, based on Lepski’s method, to estimate the unknown density in a point $x \in \mathbb{R}^D$ which is known to belong to the unknown (and possibly nonsmooth) manifold M .

To go beyond the pointwise estimation of μ , even the choice of a relevant loss is nontrivial. Indeed, most standard losses between probability measures (e.g. the L_p distance, the Hellinger distance or the Kullback-Leibler divergence) are degenerate when comparing mutually singular

E-mail address: `firstname.lastname@nyu.edu`.

measures, which will typically be the case for measures on two distinct manifolds, even if they are very close to each other with respect to the Hausdorff distance. This implies that the estimation problem is degenerate from a minimax perspective when choosing such losses (see Theorem 2.11). On the contrary, the Wasserstein distances W_p , $1 \leq p \leq \infty$ are particularly adapted to this problem, as they are by design robust to small metric perturbations of the support of a measure.

Apart from this first motivation, the use of Wasserstein distances, and more generally of the theory of optimal transport, has shown to be an efficient tool in widely different recent problems of machine learning, with fast implementations and sound theoretical results (see e.g. [PC19] for a survey). From a statistical perspective, most of the attention has been dedicated to studying rates of convergence between a probability distribution μ and its empirical counterpart μ_n [Dud69, DSS13, FG15, SP18, WB19a, Lei20]. Unsurprisingly, if more regularity is assumed on μ , then it is possible to build estimators with smaller risks than the empirical measure μ_n . Assume for instance that μ is a probability distribution on the cube $[-1, 1]^D$, with density f of regularity s (measured through the Besov scale $B_{p,q}^s$). In this setting, it has been shown in [WB19b] that, given n i.i.d. points of law μ , the minimax rate (up to logarithmic factors) for the estimation of μ with respect to the Wasserstein distance W_p is of order

$$\begin{cases} n^{-\frac{s+1}{2s+D}} & \text{if } D \geq 3 \\ n^{-\frac{1}{2}} \log n & \text{if } D = 2 \\ n^{-\frac{1}{2}} & \text{if } D = 1, \end{cases} \quad (1.1)$$

and that this rate is attained by a modified linear wavelet density estimator. Our main contribution consists in extending the results of [WB19b] by allowing the support of the probability to be any d -dimensional compact \mathcal{C}^k submanifold $M \subset \mathbb{R}^D$ for $k \geq 2$. More precisely, assume that some probability μ on M has a lower and upper bounded density f which belongs to the Besov space $B_{p,q}^s(M)$ for some $0 < s \leq k-1$, $1 \leq p < \infty$, $1 \leq q \leq \infty$ (see Section 2 for details). We first show (Theorem 3.1) that some weighted kernel density estimator that we integrate against the volume measure vol_M on M attains, for the W_p distance, the rate of estimation

$$\begin{cases} n^{-\frac{s+1}{2s+d}} & \text{if } d \geq 3 \\ n^{-\frac{1}{2}} (\log n)^{\frac{1}{2}} & \text{if } d = 2 \\ n^{-\frac{1}{2}} & \text{if } d = 1. \end{cases} \quad (1.2)$$

In the case where the manifold M is unknown, we do not have access to the volume measure vol_M , so that the latter estimator is not computable. We therefore propose to estimate the volume measure vol_M in a preliminary step. Such an estimator $\widehat{\text{vol}}_M$ is defined by using local polynomial estimation techniques from [AL19]. We show that this estimator is a minimax estimator of the volume measure up to logarithmic factors (Theorem 3.7), with a risk of order $(\log n/n)^{k/d}$. We then show (Theorem 3.8) that a weighted kernel density estimator integrated against $\widehat{\text{vol}}_M$ attains the rate (1.2). Those rates are significantly faster than the rates of (1.1) if $d \ll D$ and are shown to be minimax up to logarithmic factors.

Being able to estimate accurately the volume measure vol_M has also other useful implications, e.g. we provide an algorithm to sample points uniformly on a (possibly unknown) manifold, and we leverage results from [TGHS20] to provide precise estimates of the eigenvalues of the Laplace-Beltrami operator on M (see Section 5).

In Section 2, we define our statistical model and give some preliminary results on Wasserstein distances. In Section 3, we define kernel density estimators on a manifold M , and state our main results. Proofs of the main theorems are then given in Section 4. Section 5 discusses the implementation of our estimators, in particular proving that the local polynomial estimators of Aamari & Levrard [AL19] can be efficiently computed. Additional proofs are found in the Appendix.

2. PRELIMINARIES

2.1. Regularity of manifolds. For any $d > 0$, we write $\langle \cdot, \cdot \rangle$ for the dot product and $|v|$ for the norm of a vector $v \in \mathbb{R}^d$. The open ball centered at $x \in \mathbb{R}^d$ of radius $h > 0$ is denoted by $\mathcal{B}(x, h)$. For $\Omega \subset \mathbb{R}^d$ a set and $x \in \mathbb{R}^d$, we let $d(x, \Omega) := \inf\{|x - y|, y \in \Omega\}$ be the distance from x to Ω and we write $\mathcal{B}_\Omega(x, h)$ for $\mathcal{B}(x, h) \cap \Omega$. Also, we let $\Omega^h := \{x \in \mathbb{R}^d, d(x, \Omega) < h\}$ be the h -tubular neighborhood of Ω . Given a tensor $A : (\mathbb{R}^{d_1})^i \rightarrow \mathbb{R}^{d_2}$ of order $i \geq 0$, the operator norm $\|A\|_{\text{op}}$ is defined as $\|A\|_{\text{op}} := \max\{|A[v_1, \dots, v_i]|, |v_1|, \dots, |v_i| \leq 1\}$. Also, we let $A^\top : \mathbb{R}^{d_2} \rightarrow \mathbb{R}^{d_1}$ denote the adjoint of the operator $A : \mathbb{R}^{d_1} \rightarrow \mathbb{R}^{d_2}$. If $f : \Omega \rightarrow \mathbb{R}^{d_2}$ is a \mathcal{C}^k function defined on an open set Ω of \mathbb{R}^{d_1} , we let $\|f\|_{\mathcal{C}^k(\Omega)} := \max_{0 \leq i \leq k} \sup\{\|d^i f(x)\|_{\text{op}}, x \in \Omega\}$, where $d^i f(x)$ is the i th differential of f at x .

Let $D > 0$ and let \mathcal{M}_d be the set of all smooth d -dimensional connected submanifolds in \mathbb{R}^D without boundary, endowed with the metric induced by the standard metric on \mathbb{R}^D . We denote by d_g the geodesic distance on M . The tangent space at a point $x \in M$ is denoted by $T_x M$. It is identified with a d -dimensional subspace of \mathbb{R}^D , and the orthogonal projection on $T_x M$ is denoted by π_x . We also let $\tilde{\pi}_x : \mathbb{R}^D \rightarrow T_x M$ be defined by $\tilde{\pi}_x(y) = \pi_x(y - x)$. We denote by $T_x M^\perp$ the normal space at $x \in M$. The key quantity used to describe the regularity of a manifold M is its reach $\tau(M)$. It is defined as the distance between M and its medial axis, that is the set of points $x \in \mathbb{R}^D$ for which there are at least two points of M which attain the distance from x to M . In particular, the projection π_M on the manifold M is defined on $M^{\tau(M)}$. Originally introduced in [Fed59], the reach $\tau(M)$ measures both the local regularity of M (namely its curvature) and its global regularity, see e.g. [AKC⁺19, BHHS21] or [DZ01, Section 6.6] for precise results on the relationships between the reach of a manifold and its geometry. We then measure the regularity of M through the regularity of local parametrizations of M (see [AL19]).

Definition 2.1. Let $M \in \mathcal{M}_d$, and $\tau_{\min}, L > 0, k \geq 2$. Let $r_0 = (\tau_{\min} \wedge L)/4$. We say that M is in $\mathcal{M}_{d, \tau_{\min}, L}^k$ if M is closed, of reach larger than τ_{\min} and if, for all $x \in M$, the projection $\tilde{\pi}_x : M \rightarrow T_x M$ is a local diffeomorphism in x , with inverse Ψ_x defined on $\mathcal{B}_{T_x M}(0, r_0)$, satisfying $\|\Psi_x\|_{\mathcal{C}^k(\mathcal{B}_{T_x M}(0, r_0))} \leq L$.

Remark 2.2. (i) Remark that we only consider manifolds M that are smooth in the above definition, with a controlled \mathcal{C}^k norm. As the set of smooth submanifolds is dense in the set of \mathcal{C}^k submanifolds (for an appropriate topology), this is not a strong assumption. Dealing with smooth submanifolds is more convenient for us, as we do not have to deal with tricky existence issues for defining different functional spaces on manifolds.

(ii) For the sake of convenience, we use a definition slightly different from the definition of [AL19], where authors assume the existence of local parametrizations $\tilde{\Psi}_x$ having controlled \mathcal{C}^k norms, with $\tilde{\Psi}_x$ not necessarily equal to the inverse Ψ_x of the orthogonal projection. However, our definition is not restrictive. Indeed, one can write $\Psi_x = \tilde{\Psi}_x \circ (\tilde{\pi}_x \circ \tilde{\Psi}_x)^{-1}$, where the \mathcal{C}^k norm of $(\tilde{\pi}_x \circ \tilde{\Psi}_x)^{-1}$ is controlled by the inverse function theorem. Therefore, the \mathcal{C}^k norm of Ψ_x can always be controlled by the \mathcal{C}^k norms of other parametrizations $\tilde{\Psi}_x$. Both definitions can also be proven to be equivalent to assuming that the function $d^2(\cdot, M)$ has a controlled \mathcal{C}^k norm on $M^{\tau(M)}$, see e.g. [PR84].

(iii) The value of the scale parameter r_0 is used for convenience. Other small scales could be used, or the radius r_0 could also be added as another parameter of the model, without any substantial gain in doing so.

If $M_1 \in \mathcal{M}_{d_1}, M_2 \in \mathcal{M}_{d_2}, x \in M_1$ and $f : M_1 \rightarrow M_2$ is a \mathcal{C}^1 function, then we let $df(x) : T_x M_1 \rightarrow T_{f(x)} M_2$ be the differential of f at x . For $0 \leq l \leq k$, if f is \mathcal{C}^l , we let $\|d^l f(x)\|_{\text{op}} := \max_{0 \leq i \leq l} \|d^i(f \circ \Psi_x)(0)\|_{\text{op}}$ and $\|f\|_{\mathcal{C}^l(M_1)} := \sup_{x \in M_1} \|d^l f(x)\|_{\text{op}}$. If $d_1 \leq d_2$, then we define the Jacobian of f at $x \in M_1$ as $Jf(x) = \sqrt{\det(df(x)^\top df(x))}$. We let $\mathcal{C}^l(M)$ be the space of all \mathcal{C}^l functions $f : M \rightarrow \mathbb{R}$ (with possibly $l = \infty$) and for $f \in \mathcal{C}^l(M)$, we let ∇f denote the gradient of f . We also denote by $\nabla \cdot$ the divergence operator on M .

Let vol_M be the volume measure associated with the Riemannian metric on M . We will denote the integration with respect to $d\text{vol}_M(x)$ by dx when the context is clear. For $1 \leq p \leq \infty$,

we let $L_p(M)$ be the set of measurable functions $f : M \rightarrow \mathbb{R}$ with finite p -norm $\|f\|_{L_p(M)} := (\int f d\text{vol}_M)^{1/p}$ (and usual modification if $p = \infty$). We say that a locally integrable function f is weakly differentiable if there exists a measurable section ∇f of the tangent bundle TM (uniquely defined almost everywhere) such that for all smooth vector fields w on M with compact support, we have

$$\int f(\nabla \cdot w) d\text{vol}_M = - \int (\nabla f) \cdot w d\text{vol}_M.$$

Furthermore, we will denote by $p^* \in [1, \infty]$ the number satisfying $\frac{1}{p} + \frac{1}{p^*} = 1$.

2.2. Besov spaces on manifolds. Let $M \in \mathcal{M}_{d, \tau_{\min}, L}^k$ for some $k \geq 2$, $\tau_{\min}, L > 0$. As stated in the introduction, minimax rates for the estimation of a given probability will depend crucially on the regularity of its density f , which is assumed to belong to some Besov space $B_{p,q}^s(M)$. We first introduce Sobolev spaces $H_p^l(M)$ on M for $l \leq k$ an integer, and Besov spaces on M are then defined by real interpolation.

Definition 2.3 (Sobolev space on a manifold). *Let $0 \leq l \leq k$, $1 \leq p < \infty$ and let $f \in C^\infty(M)$ function. We let*

$$\|f\|_{H_p^l(M)} := \max_{0 \leq i \leq l} \left(\int \|d^i f(x)\|_{\text{op}}^p d\text{vol}_M(x) \right)^{1/p}. \quad (2.1)$$

The space $H_p^l(M)$ is the completion of $C^\infty(M)$ for the norm $\|\cdot\|_{H_p^l(M)}$.

Remark 2.4 (On the case $p = \infty$). The previous definition cannot be extended to the case $p = \infty$. Indeed, the completion of $C^\infty(M)$ for the norm $\|\cdot\|_{H_\infty^l(M)}$ is equal to $\mathcal{C}^l(M)$, whereas for instance $H_\infty^0(M)$ should be equal to $L_\infty(M)$. For $l = 1$, the space $H_p^1(M)$ can equivalently be defined as the space of weakly differentiable functions f with $\|f\|_{H_p^1(M)} < \infty$, while this definition can be easily extended to the case $p = \infty$. In particular, if $f \in H_\infty^1(M)$, then one can verify that $f \circ \Psi_x \in H_\infty^1(\mathcal{B}_{T_x M}(0, r_0))$ for any $x \in M$. It follows from standard results on Sobolev spaces on domains that $f \circ \Psi_x$ is Lipschitz continuous (see e.g. [Bre10, Proposition 9.3]). Hence, f is also locally Lipschitz continuous. By Rademacher theorem, f is therefore almost everywhere differentiable, and its differential coincides with the weak differential. As a consequence, a function $f \in H_\infty^1(M)$ is Lipschitz continuous, with Lipschitz constant for the distance d_g equal to $\|f\|_{H_\infty^1(M)}$.

For $1 \leq p < \infty$, we introduce the negative homogeneous Sobolev norm $\|\cdot\|_{\dot{H}_p^{-1}(M)}$, defined, for $f \in L_p(M)$ with $\int f d\text{vol}_M = 0$, by

$$\|f\|_{\dot{H}_p^{-1}(M)} := \sup \left\{ \int f g d\text{vol}_M, \|\nabla g\|_{L_{p^*}(M)} \leq 1 \right\}, \quad (2.2)$$

where the supremum is taken over all functions $g \in H_{p^*}^1(M)$. For $f \in L_p(M)$, the negative Sobolev norm is defined by

$$\|f\|_{H_p^{-1}(M)} := \sup \left\{ \int f g d\text{vol}_M, \|g\|_{H_{p^*}^1(M)} \leq 1 \right\}, \quad (2.3)$$

and the corresponding Banach space is denoted by $H_p^{-1}(M)$.

Proposition 2.5. *Let $1 \leq p < \infty$ and $f \in H_p^{-1}(M)$ with $\int f d\text{vol}_M = 0$.*

- (i) *We have $C_{d, \tau_{\min}} |\text{vol}_M|^{\frac{d-1}{p} - d} \|f\|_{\dot{H}_p^{-1}(M)} \leq \|f\|_{H_p^{-1}(M)} \leq \|f\|_{\dot{H}_p^{-1}(M)}$ for some positive constant $C_{d, \tau_{\min}}$ depending on d and τ_{\min} .*
- (ii) *We have $\|f\|_{\dot{H}_p^{-1}(M)} = \inf \{ \|w\|_{L_p(M)}, \nabla \cdot w = f \}$, where the infimum is taken over all measurable vector fields w on M with finite p -norm, and where $\nabla \cdot w = f$ means that $\int f g d\text{vol}_M = - \int w \cdot \nabla g d\text{vol}_M$ for all $g \in C^\infty(M)$.*

Following [Tri92], Besov spaces on a manifold M are defined as real interpolation of Sobolev spaces. We refer to [Lun18] for definition of real interpolations of Banach spaces.

Definition 2.6 (Besov space on a manifold). *Let $1 \leq p < \infty$ and $0 < s < k$. The Besov space $B_{p,q}^s(M)$ is defined as $B_{p,q}^s(M) := (L_p(M), H_p^k(M))_{s/k,q}$, the real interpolation space between $L_p(M)$ and $H_p^k(M)$ of parameters s/k and q .*

Basic results from interpolation theory then imply that $\|\cdot\|_{B_{p,q}^s(M)} \leq \|\cdot\|_{B_{p,q}^{s'}(M)}$ if $0 < s \leq s' < k$.

2.3. Wasserstein distances and negative Sobolev distances. Let \mathcal{P} be the set of finite Borel measures μ on \mathbb{R}^D , with $|\mu|$ the total mass of μ . Let \mathcal{P}_1 be the set of measures in \mathcal{P} with $|\mu| = 1$. For $1 \leq p \leq \infty$, let \mathcal{P}^p be the set of measures $\mu \in \mathcal{P}$ satisfying $(\int |x|^p d\mu(x))^{1/p} < \infty$ (with usual modification for $p = \infty$) and let $\mathcal{P}_1^p = \mathcal{P}^p \cap \mathcal{P}_1$. The pushforward of a measure μ by a measurable application $\phi : \mathbb{R}^D \rightarrow \mathbb{R}^D$ is defined by

$$\phi_{\#}\mu(A) := \mu(\phi^{-1}(A)) \quad (2.4)$$

for any Borel set $A \subset \mathbb{R}^D$. For $\rho : \mathbb{R}^D \rightarrow [0, \infty)$ a measurable function, we denote by $\rho \cdot \mu$ the measure having density ρ with respect to μ .

Definition 2.7 (Wasserstein distance). *Let $1 \leq p \leq \infty$ and let $\mu, \nu \in \mathcal{P}^p$ with the same total mass. Let $\Pi(\mu, \nu)$ be the set of transport plans between μ and ν , i.e. measures on $\mathbb{R}^D \times \mathbb{R}^D$ with first marginal π^1 (resp. second marginal π^2) equal to μ (resp. ν). The cost $C_p(\pi)$ of $\pi \in \Pi(\mu, \nu)$ is defined as $\int |x - y|^p d\pi(x, y)$. The p -Wasserstein distance between μ and ν is defined as*

$$W_p(\mu, \nu) := \inf_{\pi \in \Pi(\mu, \nu)} C_p(\pi)^{1/p}, \quad (2.5)$$

with usual modification for $p = \infty$.

A crucial point in the study conducted in the following is the relation between Wasserstein distances and negative Sobolev norms.

Proposition 2.8 (Wasserstein distances and negative Sobolev norms). *Let $1 \leq p < \infty$. Let $M \in \mathcal{M}_d$ be a manifold with reach $\tau(M) \geq \tau_{\min}$, and let $\mu, \nu \in \mathcal{P}_1^p$ be two probability measures supported on M , absolutely continuous with respect to vol_M , with densities f and g . Assume that $f, g \geq f_{\min} \cdot \text{vol}_M$ for some $f_{\min} > 0$. Then, we have*

$$\begin{aligned} W_p(\mu, \nu) &\leq p^{-1/p} f_{\min}^{1/p-1} \|f - g\|_{\dot{H}_p^{-1}(M)} \\ &\leq p^{-1/p} C_{d, \tau_{\min}, f_{\min}} \|f - g\|_{H_p^{-1}(M)}, \end{aligned} \quad (2.6)$$

for some constant $C_{d, \tau_{\min}, f_{\min}}$ depending on d , τ_{\min} and f_{\min} .

In particular, if $p = 1$, then the first inequality in (2.6) is actually an equality by the Kantorovitch duality formula [Vil08, Particular Case 5.16]. This inequality appears in [Pey18] for $p = 2$ and in [San15, Section 5.5.1] for measures having density with respect to the Lebesgue measure. We carefully adapt their proofs in Appendix B.

2.4. Statistical models. We consider the two following models, where points are sampled on a manifold, with possibly tubular noise. We fix in the following some parameters $\tau_{\min}, L_s, L_k > 0$, $1 \leq q \leq \infty$ and $0 < f_{\min} < f_{\max} < \infty$. We also write \mathcal{M}_d^k instead of $\mathcal{M}_{d, \tau_{\min}, L_k}^k$.

Definition 2.9 (Noise-free model). *Let $d \leq D$ be integers, $k \geq 2$, $0 \leq s < k$ and $1 \leq p < \infty$. Let $M \in \mathcal{M}_d^k$. For $s = 0$, the set $\mathcal{Q}^0(M)$ is the set of probability distributions μ on \mathbb{R}^D absolutely continuous with respect to the volume measure vol_M , with a density f satisfying $f_{\min} \leq f \leq f_{\max}$ almost everywhere. For $s > 0$, the set $\mathcal{Q}^s(M)$ is the set of distributions $\mu \in \mathcal{Q}^0(M)$, with density $f \in B_{p,q}^s(M)$ satisfying $\|f\|_{B_{p,q}^s(M)} \leq L_s$. The model $\mathcal{Q}_d^{s,k}$ is equal to the union of the sets $\mathcal{Q}^s(M)$ for $M \in \mathcal{M}_d^k$.*

Remark 2.10. If $\mu \in \mathcal{Q}^s(M)$, then, as $\mu \geq f_{\min} \text{vol}_M$, one has $\text{vol}_M(M) \leq f_{\min}^{-1}$. One can then use standard packing arguments to show that this implies that $\text{diam}(M) \leq C_d / (f_{\min} \tau_{\min}^{d-1})$ for some constant C_d depending only on d . In particular, the manifold M is automatically compact.

Given a set of observations X_1, \dots, X_n sampled according to $\mu \in \mathcal{Q}_d^{s,k}$, the goal of statistical inference is to reconstruct some quantity $\vartheta(\mu)$ related to μ . If \mathcal{L} is a loss function defined on the set of outputs of ϑ , we define the minimax risk for this problem as

$$\mathcal{R}_n(\vartheta, \mathcal{Q}_d^{s,k}, \mathcal{L}) := \inf_{\hat{\vartheta}} \sup_{\mu \in \mathcal{Q}} \mathbb{E} \mathcal{L}(\hat{\vartheta}, \vartheta), \quad (2.7)$$

where $\hat{\vartheta} = \hat{\vartheta}(X_1, \dots, X_n)$ and X_1, \dots, X_n is an i.i.d. sample with law μ . We will focus here on reconstructing (i) the measure $\vartheta(\mu) = \mu$, and (ii) the uniform measure $\vartheta(\mu) = U_M := \text{vol}_M / |\text{vol}_M|$, where M is the support of μ . We first show that task (i) is impossible if the loss function \mathcal{L} is larger than the total variation distance TV, which is defined by $\text{TV}(\mu, \nu) := \sup_A |\mu(A) - \nu(A)|$ for $\mu, \nu \in \mathcal{P}_1$, where the supremum is taken over all measurable sets $A \subset \mathbb{R}^D$.

Theorem 2.11. *Let $d \leq D$ be integers, $k \geq 2$, $0 \leq s < k$, $1 \leq p < \infty$. Let $\mathcal{L} : \mathcal{P} \times \mathcal{P} \rightarrow [0, \infty]$ be a measurable map with respect to the Borel σ -algebra associated with the total variation distance on $\mathcal{P} \times \mathcal{P}$. Assume that $\mathcal{L}(\mu, \nu) \geq g(\text{TV}(\mu, \nu))$ for a convex nondecreasing function $g : \mathbb{R} \rightarrow [0, \infty]$ with $g(0) = 0$. Then, for any $\tau_{\min} > 0$, if f_{\min} is small enough and L_k, L_s, f_{\max} are large enough, we have*

$$\mathcal{R}_n(\mu, \mathcal{Q}_d^{s,k}, \mathcal{L}) \geq g(c_d), \quad (2.8)$$

for some constant $c_d > 0$.

Examples of such losses include the total variation distance, the Hellinger distance (with $g(x) = x$), the Kullback-Leibler divergence (with $g(x) = x^2/2$), and the L_p distance with respect to some dominating measure (with $g(x) = x^p$). We give a proof of Theorem 2.11, based on Assouad's lemma, in Appendix G. A simple example of loss \mathcal{L} which is not degenerate for mutually singular measures is given by the W_p distance. As stated in the introduction, we will therefore choose this loss, and study $\mathcal{R}_n(\mu, \mathcal{Q}_d^{s,k}, W_p)$, the minimax rate of estimation for μ with respect to W_p .

We will also study this problem in the presence of tubular noise.

Definition 2.12 (Tubular noise model). *Let $d \leq D$ be integers, $k \geq 2$, $0 \leq s < k$, $1 \leq p < \infty$ and $\gamma \geq 0$. The set $\mathcal{Q}_d^{s,k}(\gamma)$ is the set of laws ν of random variables $Y + Z$ where $Y \sim \mu \in \mathcal{Q}_d^{s,k}$ and $Z \in \mathcal{B}(0, \gamma)$ is such that $Z \in T_Y M^\perp$.*

Note that the tubular noise model is not identifiable, in the sense that there are several admissible couples (Y, Z) (with possibly different supports) such that $X = Y + Z$ follows the same distribution ν . For each $\nu \in \mathcal{Q}_d^{s,k}(\gamma)$, we will make an arbitrary choice among the admissible couples, while this will not have any impact on the following study (all the admissible couples have their first marginal at most 2γ apart for the Wasserstein distance).

Remark 2.13. For ease of notation, we will write in the following $a \lesssim b$ to indicate that there exists a constant C depending on the parameters $p, k, \tau_{\min}, L_s, L_k, f_{\min}, f_{\max}$, **but not on D** , such that $a \leq Cb$, and write $a \asymp b$ to indicate that $a \lesssim b$ and $b \lesssim a$. Also, we will write c_α to indicate that a constant c depends on some parameter α .

Note that in particular, the risk of the estimators proposed in the next section will not depend on the ambient dimension D , but only on intrinsic parameters such as the regularity of ν and its effective dimension d .

3. KERNEL DENSITY ESTIMATION ON AN UNKNOWN MANIFOLD

Before building an estimator in the model $\mathcal{Q}_d^{s,k}(\gamma)$, let us consider the easier problem of the estimation of μ in the case where $\gamma = 0$ (noise free model) and the support M is known. Let $\mu \in \mathcal{Q}^s(M)$ and Y_1, \dots, Y_n be a n -sample of law μ . Let $\mu_n = \frac{1}{n} \sum_{i=1}^n \delta_{Y_i}$ be the empirical measure of the sample. Identify \mathbb{R}^d with $\mathbb{R}^d \times \{0\}^{D-d}$ and consider a kernel $K : \mathbb{R}^D \rightarrow \mathbb{R}$ satisfying the following conditions:

- **Condition A:** The kernel K is a smooth radial function with support $\mathcal{B}(0, 1)$ such that $\int_{\mathbb{R}^d} K = 1$.

- **Condition $B(m)$:** The kernel K is of order $m \geq 0$ in the following sense. Let $|\alpha| := \sum_{j=1}^d \alpha_j$ be the length of a multiindex $\alpha = (\alpha_1, \dots, \alpha_d)$. Then, for all multiindexes α^0, α^1 with $0 \leq |\alpha^0| < m, 0 \leq |\alpha^1| < m + |\alpha_0|$, and with $|\alpha^1| > 0$ if $\alpha^0 = 0$, we have

$$\int_{\mathbb{R}^d} \partial^{\alpha^0} K(v) v^{\alpha^1} dv = 0, \quad (3.1)$$

where $v^\alpha = \prod_{j=1}^d v_j^{\alpha_j}$ and $\partial^\alpha K$ is the partial derivative of K in the direction α .

- **Condition $C(\beta)$:** The negative part $K_- := -\min(0, K)$ of K satisfies $\int_{\mathbb{R}^d} K_- \leq \beta$.

We show in Appendix H that for every integer $m \geq 0$ and real number $\beta > 0$, there exists a kernel K satisfying conditions $A, B(m)$ and $C(\beta)$. Define the convolution of K with a measure $\rho \in \mathcal{P}$ as

$$K * \rho(x) := \int K(x - y) d\rho(y), \quad x \in \mathbb{R}^D, \quad (3.2)$$

and, for $h > 0$, let $K_h := h^{-d} K(\cdot/h)$. Let $\rho_h := K_h * \text{vol}_M$ and let $\mu_{n,h}$ be the measure with density $f_{n,h} := K_h * (\mu_n/\rho_h)$ with respect to vol_M . Dividing by ρ_h ensures that $\mu_{n,h}$ is a measure of mass 1. Remark that the computation of $\mu_{n,h}$ requires to have access to M , that is $\mu_{n,h}$ is an estimator on $\mathcal{Q}^s(M)$ but not on $\mathcal{Q}_d^{s,k}$. By linearity, the expectation of $\mu_{n,h}$ is given by μ_h , the measure having for density $f_h := K_h * (\mu/\rho_h)$ on M .

Theorem 3.1. *Let $d \leq D$ be integers, $0 < s \leq k - 1$ with $k \geq 2$ and $1 \leq p < \infty$. Let $M \in \mathcal{M}_d^k$ and $\mu \in \mathcal{Q}^s(M)$ with density f . Let Y_1, \dots, Y_n be a n -sample of law μ . There exists a constant β depending on the parameters of the model such that, if K is a kernel satisfying conditions $A, B(k)$ and $C(\beta)$, then the measure $\mu_{n,h}$ satisfies the following:*

- (i) *If $(\log n/n)^{1/d} \lesssim h \lesssim 1$, then, with probability larger than $1 - cn^{-k/d}$, the density $f_{n,h}$ of $\mu_{n,h}$ is larger than $f_{\min}/2$ and smaller than $2f_{\max}$ everywhere on M .*
- (ii) *If $n^{-1/d} \lesssim h \lesssim 1$, then we have*

$$\mathbb{E} \|f - f_{n,h}\|_{H_p^{-1}(M)} \leq \|f - f_h\|_{H_p^{-1}(M)} + \mathbb{E} \|f_{n,h} - f_h\|_{H_p^{-1}(M)} \quad (3.3)$$

$$\lesssim h^{s+1} + \frac{h^{1-d/2} I_d(h)}{\sqrt{n}}, \quad (3.4)$$

where $I_d(h) = 1$ if $d \geq 3$, $(-\log(h))^{1/2}$ if $d = 2$ and $h^{-1/2}$ if $d = 1$.

- (iii) *Let $h \asymp n^{-1/(2s+d)}$ if $d \geq 3$, $h \asymp (\log n/n)^{1/d}$ if $d \leq 2$. Define $\mu_{n,h}^0 = \mu_{n,h}$ if $\mu_{n,h}$ is a nonnegative measure and $\mu_{n,h}^0 = \delta_{X_1}$ otherwise. Then,*

$$\mathbb{E} W_p(\mu_{n,h}^0, \mu) \lesssim \begin{cases} n^{-\frac{s+1}{2s+d}} & \text{if } d \geq 3, \\ n^{-\frac{1}{2}} (\log n)^{\frac{1}{2}} & \text{if } d = 2, \\ n^{-\frac{1}{2}} & \text{if } d = 1. \end{cases} \quad (3.5)$$

- (iv) *Furthermore, for any $0 \leq s < k$ and $\tau_{\min} > 0$, if f_{\min} is small enough and if f_{\max} and L_s are large enough, then there exists a manifold $M \in \mathcal{M}_d^k$ such that*

$$\mathcal{R}_n(\mu, W_p, \mathcal{Q}^s(M)) \gtrsim \begin{cases} n^{-\frac{s+1}{2s+d}} & \text{if } d \geq 3, \\ n^{-\frac{1}{2}} & \text{if } d \leq 2. \end{cases} \quad (3.6)$$

Remark 3.2. The condition $C(\beta)$ on the kernel is only used to ensure that the measure $\mu_{n,h}$ has a lower and upper bounded density on M . An alternative possibility to ensure this property is to assume that the density of μ is Hölder continuous of exponent δ for some $\delta > 0$. Techniques from [BH19] then imply that $\|\mu_{n,h} - \mu\|_{L^\infty(M)} \lesssim h^\delta + n^{-1/2} h^{-d/2} \ll 1$ with high probability, ensuring in particular that the density is lowerbounded. If $sp > d$, then every element of $B_{p,q}^s(M)$ is Hölder continuous [Tri92, Theorem 7.4.2], and condition $C(\beta)$ is no longer required. However, Theorem 3.1 also holds for non-continuous densities.

Remark 3.3. Let K be a nonnegative kernel satisfying conditions A , $B(0)$ and $C(\beta)$. It is straightforward to check that $W_p(\mu_n, \mu_{n,h}) \lesssim h$. Therefore, Theorem 3.1(ii) and Proposition 2.8 imply in particular that $W_p(\mu_n, \mu) \lesssim h + \frac{h^{1-d/2} I_d(h)}{\sqrt{n}}$. By choosing h of the order $n^{-1/d}$, we obtain that

$$W_p(\mu_n, \mu) \lesssim \begin{cases} n^{-\frac{1}{d}} & \text{if } d \geq 3 \\ n^{-\frac{1}{2}} (\log n)^{\frac{1}{2}} & \text{if } d = 2 \\ n^{-\frac{1}{2}} & \text{if } d = 1. \end{cases} \quad (3.7)$$

Such a result was already shown for $p = \infty$ [TGHS20] with additional logarithmic factors, with a proof very different than ours. See also [Div21b] for a short proof of this result when M is the flat torus.

Remark 3.4. There is a logarithmic gap between the minimax lower bound and the upper risk of the proposed estimator for $d = 2$ in Theorem 3.1. When μ is the uniform measure on the square $[0, 1]^2$, it is known that the empirical measure μ_n attains exactly the rate $n^{-\frac{1}{2}} (\log n)^{\frac{1}{2}}$ [Tal14, Section 6.4], suggesting that the factor $(\log n)^{1/2}$ is not a proof artifact. It is however not clear how one can transform Talagrand's tree construction used to lower bound $\mathbb{E}W_1(\mu_n, \mu)$ into a more general minimax lower bound on $\mathcal{R}_n(\mu, W_p, \mathcal{Q}^s(M))$, so that the (non-)existence of estimators attaining a rate of $n^{-1/2}$ for $d = 2$ is still an open problem.

In (3.4), a classical bias-variance trade-off appears. Namely, the bias of the estimator is of order h^{s+1} , whereas its fluctuations are of order $h^{1-d/2}/\sqrt{n}$ (at least for $d \geq 3$). This decomposition can be compared to the classical bias-variance decomposition for a kernel density estimator of bandwidth h , say for the pointwise estimation of a function of class \mathcal{C}^s on the cube $[0, 1]^d$. It is then well-known (see e.g. [Tsy08, Chapter 1]) that the bias of the estimator is of order h^s whereas its variance is of order $h^{-d/2}/\sqrt{n}$. The supplementary factor h appearing both in the bias and fluctuation terms can be explained by the fact that we are using a norm $H_p^{-1}(M)$ instead of a pointwise norm to quantify the risk of the estimator: in some sense, we are estimating the antiderivative of the density rather than the density itself. This is particularly striking when $d = 1$ and $p = 1$, the Wasserstein distance between two measures being then given by the L_1 distance between the cumulative distribution functions of the two measures [San15, Proposition 2.17].

Before giving a proof of Theorem 3.1, let us explain how to extend it to the case where the manifold M is unknown and in the presence of tubular noise. The measure $\mu_{n,h}$ is the measure having density $K_h * (\mu_n/\rho_h)$ with respect to vol_M . Of course, if M is unknown, then so is vol_M , and we therefore propose the following estimation procedure of vol_M , using local polynomial estimation techniques from [AL19]. Let X_1, \dots, X_n be a n -sample in the model with tubular noise $\mathcal{Q}_d^{s,k}(\gamma)$, with $X_i = Y_i + Z_i$, Y_i of law $\mu \in \mathcal{Q}^s(M)$ and $Z_i \in T_{Y_i}M^\perp$ with $|Z_i| \leq \gamma$. Let $\nu_n^{(i)}$ be the empirical measure $\frac{1}{n-1} \sum_{j \neq i} \delta_{X_j - X_i}$. For two positive parameters ℓ, ε , the local polynomial estimator $(\hat{\pi}_i, \hat{V}_{2,i}, \dots, \hat{V}_{m-1,i})$ of order m at X_j is defined as an element of

$$\arg \min_{\pi, \sup_{2 \leq j \leq m-1} \|V_j\|_{\text{op}} \leq \ell} \mu_n^{(i)} \left(\left\| x - \pi(x) - \sum_{j=2}^{m-1} V_j [\pi(x)^{\otimes j}] \right\|^2 \mathbf{1}\{x \in \mathcal{B}(0, \varepsilon)\} \right), \quad (3.8)$$

where the argmin is taken over all orthogonal projectors π of rank d and symmetric tensors $V_j : (\mathbb{R}^D)^j \rightarrow \mathbb{R}^D$ of order j . Let \hat{T}_i be the image of $\hat{\pi}_i$ and $\hat{\Psi}_i : v \in \mathbb{R}^D \mapsto X_i + v + \sum_{j=2}^{m-1} \hat{V}_{j,i}[v^{\otimes j}]$. Let $\angle(T_1, T_2)$ denote the angle between two d -dimensional subspaces, defined by $\|\pi_{T_1} - \pi_{T_2}\|_{\text{op}}$, where π_{T_i} is the orthogonal projection on T_i for $i = 1, 2$. We summarize the results of [AL19] in the following proposition (see Appendix A for details).

Proposition 3.5. *With probability at least $1 - cn^{-k/d}$, if $m \leq k$, $(\log n/n)^{1/d} \lesssim \varepsilon \lesssim 1$, $\gamma \lesssim \varepsilon$ and $1 \lesssim \ell \lesssim \varepsilon^{-1}$, then,*

$$\max_{1 \leq i \leq n} \angle(T_{Y_i}M, \hat{T}_i) \lesssim \varepsilon^{m-1} + \gamma \varepsilon^{-1} \quad (3.9)$$

and, for all $1 \leq i \leq n$, if $v \in \hat{T}_i$ with $|v| \leq 3\varepsilon$, we have

$$|\hat{\Psi}_i(v) - \Psi_{Y_i} \circ \pi_{Y_i}(v)| \lesssim \varepsilon^m + \gamma \quad (3.10)$$

$$\left\| d\hat{\Psi}_i(v) - d(\Psi_{Y_i} \circ \pi_{Y_i})(v) \right\|_{\text{op}} \lesssim \varepsilon^{m-1} + \gamma\varepsilon^{-1}. \quad (3.11)$$

Hence, if γ is of order at most ε^k , then it is possible to approximate the tangent space at Y_i with precision ε^{k-1} and the local parametrization with precision ε^k . In particular, authors in [AL19] show that, with high probability, $\bigcup_{i=1}^n \mathcal{B}_{\hat{\Psi}_i(\hat{T}_i)}(X_i, \varepsilon)$ is at Hausdorff distance of order at most $\varepsilon^k + \gamma$ from M . We now define an estimator $\widehat{\text{vol}}_M$ of vol_M by using an appropriate partition of unity $(\chi_j)_j$, which is built thanks to the next lemma. For $A, B \subset \mathbb{R}^D$, introduce the asymmetric Hausdorff distance $d_H(A|B) := \sup_{x \in A} d(x, B)$ and the Hausdorff distance $d_H(A, B) := d_H(A|B) \vee d_H(B|A)$. We say that a set S is δ -sparse if $|x - y| \geq \delta$ for all distinct points $x, y \in S$.

Lemma 3.6 (Construction of partitions of unity). *Let $\delta \lesssim 1$. Let $S \subset M^\delta$ be a set which is $\frac{7}{3}\delta$ -sparse, with $d_H(M^\delta|S) \leq 4\delta$. Let $\theta : \mathbb{R}^D \rightarrow [0, 1]$ be a smooth radial function supported on $\mathcal{B}(0, 1)$, which is equal to 1 on $\mathcal{B}(0, 1/2)$. Define, for $y \in M^\delta$ and $x \in S$,*

$$\chi_x(y) = \frac{\theta\left(\frac{y-x}{8\delta}\right)}{\sum_{x' \in S} \theta\left(\frac{y-x'}{8\delta}\right)}. \quad (3.12)$$

Then, the sequence of functions $\chi_x : M^\delta \rightarrow [0, 1]$ for $x \in S$, satisfies (i) $\sum_{x \in S} \chi_x \equiv 1$, with at most c_d non zero terms in the sum at any given point of M^δ , (ii) $\|\chi_x\|_{C^l(M^\delta)} \leq C_{l,d}\delta^{-l}$ for any $l \geq 0$ and, (iii) χ_x is supported on $\mathcal{B}_{M^\delta}(x, 8\delta)$.

A proof of Lemma 3.6 is given in Appendix A. Lemma 3.6 requires the set S to be $7\delta/3$ -sparse. Actually, given a set $S_0 \subset M^\delta$ with $d_H(M^\delta|S_0) \leq 5\delta/3$, there always exist a subset $S \subset S_0$ that satisfies the assumptions of Lemma 3.6. Such a subset S can be computed using the farthest point sampling algorithm (see e.g. [AL18, Section 3.3]): initialize S with an arbitrary point of S_0 , and, at each step, add the farthest point $s \in S_0$ from S (that is the point that maximizes $d(s, S)$). We stop the algorithm when every point of $s \in S_0$ satisfies $d(s, S) \leq 7\delta/3$. By construction, the set S is $7\delta/3$ -sparse and satisfies $d_H(S, S_0) \leq 7\delta/3$. We then have $d_H(M^\delta|S) \leq 5\delta/3 + 7\delta/3 = 4\delta$, so that indeed one can construct a partition of unity using S .

The next proposition describes how we may define a minimax estimator $\widehat{\text{vol}}_M$ of the volume measure on M (up to logarithmic factors) using such a partition of unity.

Theorem 3.7 (Minimax estimation of the volume measure on M). *Let $d \leq D$ be integers and $k \geq 2$. Let $\nu \in \mathcal{Q}_d^{0,k}(\gamma)$ and let X_1, \dots, X_n be a n -sample of law ν . Let $(\log n/n)^{1/d} \lesssim \varepsilon \lesssim 1$, $\gamma \lesssim \varepsilon$, $1 \lesssim \ell \lesssim \varepsilon^{-1}$.*

- (i) *Let $\{X_{i_1}, \dots, X_{i_J}\}$ be the output of the farthest point sampling algorithm with parameter $7\varepsilon/24$ and input $\{X_1, \dots, X_n\}$. With probability larger than $1 - cn^{-k/d}$, there exists a sequence of smooth nonnegative functions $\chi_j : M^{\varepsilon/8} \rightarrow [0, 1]$ for $1 \leq j \leq J$, such that χ_j is supported on $\mathcal{B}_{M^{\varepsilon/8}}(X_{i_j}, \varepsilon)$, $\|\chi_j\|_{C^1(M^{\varepsilon/8})} \lesssim \varepsilon^{-1}$ and $\sum_{j=1}^J \chi_j(z) = 1$ for $z \in M^{\varepsilon/8}$, with at most c_d non-zero terms in the sum.*
- (ii) *Let $\hat{\Psi}_i$ be the local polynomial estimator of order $m \leq k$ with parameter ε and ℓ , and \hat{T}_i the associated tangent space. Let $\widehat{\text{vol}}_M$ be the measure defined by, for all continuous bounded functions $\phi : \mathbb{R}^D \rightarrow \mathbb{R}$,*

$$\int \phi(x) d\widehat{\text{vol}}_M(x) = \sum_{j=1}^J \int_{\hat{\Psi}_{i_j}(\hat{T}_{i_j})} \phi(x) \chi_j(x) dx, \quad (3.13)$$

where the integration is taken against the d -dimensional Hausdorff measure on $\hat{\Psi}_{i_j}(\hat{T}_{i_j})$. Then, for $1 \leq r \leq \infty$, with probability larger than $1 - cn^{-k/d}$, we have

$$W_r \left(\frac{\widehat{\text{vol}}_M}{|\widehat{\text{vol}}_M|}, \frac{\text{vol}_M}{|\text{vol}_M|} \right) \lesssim (\gamma + \varepsilon^m)(1 + \gamma\varepsilon^{-2}). \quad (3.14)$$

(iii) In particular, if $m = k$, $\varepsilon \asymp (\log n/n)^{1/d}$ and $\gamma \lesssim \varepsilon^2$, we obtain that

$$\mathbb{E}W_r \left(\frac{\widehat{\text{vol}}_M}{|\widehat{\text{vol}}_M|}, \frac{\text{vol}_M}{|\text{vol}_M|} \right) \lesssim \gamma + \left(\frac{\log n}{n} \right)^{\frac{k}{d}}. \quad (3.15)$$

Also, for any $\tau_{\min} > 0$ and $0 \leq s < k$, if f_{\min} is small enough, and if f_{\max}, L_k, L_s are large enough, then

$$\mathcal{R}_n \left(\frac{\text{vol}_M}{|\text{vol}_M|}, \mathcal{Q}_d^{s,k}, W_r \right) \gtrsim \left(\frac{1}{n} \right)^{\frac{k}{d}}. \quad (3.16)$$

Let $\hat{\rho}_h := K_h * \widehat{\text{vol}}_M$. We define $\hat{\nu}_{n,h}$ as the measure having density $K_h * (\nu_n / \hat{\rho}_h)$ with respect to the measure $\widehat{\text{vol}}_M$, where $\nu_n = \frac{1}{n} \sum_{i=1}^n \delta_{X_i}$ is the empirical measure of the sample (X_1, \dots, X_n) .

Theorem 3.8. *Let $d \leq D$ be integers, $0 < s \leq k-1$ with $k \geq 2$ and $1 \leq p < \infty$. Let $\nu \in \mathcal{Q}_d^{s,k}(\gamma)$ and let X_1, \dots, X_n be a n -sample of law ν . Decompose X_i as $Y_i + Z_i$, where Y_i has law $\mu \in \mathcal{Q}_d^{s,k}$ and Z_i is an orthogonal noise of size γ . There exists a constant β depending on the parameters of the model such that the following holds. Assume that K is a kernel satisfying conditions A, B(k) and C(β), that $(\log n/n)^{1/d} \lesssim \varepsilon \lesssim h \lesssim 1$, $\gamma \lesssim \varepsilon$, $1 \lesssim \ell \lesssim \varepsilon^{-1}$ and consider the estimator $\widehat{\text{vol}}_M$ defined in (3.13) with parameters m , ε and ℓ . We have the following:*

- (i) *The measure $\hat{\nu}_{n,h}$ is a nonnegative measure with probability larger than $1 - cn^{-k/d}$.*
- (ii) *Define $\hat{\nu}_{n,h}^0 = \hat{\nu}_{n,h}$ if $\hat{\nu}_{n,h}$ is a nonnegative measure and $\hat{\nu}_{n,h}^0 = \delta_{X_1}$ otherwise. Then, with probability larger than $1 - cn^{-k/d}$,*

$$W_p(\hat{\nu}_{n,h}^0, \mu_{n,h}^0) \lesssim (\gamma + \varepsilon^m)(1 + \gamma\varepsilon^{-2}). \quad (3.17)$$

- (iii) *In particular, let $m = \lceil s+1 \rceil$, $\varepsilon \asymp (\ln n/n)^{1/d}$, $\gamma \lesssim \varepsilon^2$, $\ell \asymp \varepsilon^{-1}$ and $h \asymp n^{-1/(2s+d)}$ if $d \geq 3$, $h \asymp (\log n/n)^{1/d}$ if $d \leq 2$. Then,*

$$\mathbb{E}W_p(\hat{\nu}_{n,h}^0, \mu) \lesssim \gamma + \begin{cases} n^{-\frac{s+1}{2s+d}} & \text{if } d \geq 3, \\ n^{-\frac{1}{2}} (\log n)^{\frac{1}{2}} & \text{if } d = 2, \\ n^{-\frac{1}{2}} & \text{if } d = 1. \end{cases} \quad (3.18)$$

We thus manage to obtain the same rate of convergence as in Theorem 3.1, although not having access to the manifold M (up to a noise factor γ).

4. PROOFS OF THE MAIN THEOREMS

The proof of Theorem 3.1 relies on the classical bias variance decomposition displayed in (3.3), together with the linearization inequality given in Proposition 2.8. We first bound the bias of the estimator, which can be expressed in term of a convolution operator on M .

4.1. Bias of the kernel density estimator. The bias of the estimator is given by the distance $\|\cdot\|_{H_p^{-1}(M)}$ between f the density of μ and $f_h := K_h * (\mu/\rho_h)$. For $\phi \in L_1(M)$, we write $\tilde{\phi}$ for ϕ/ρ_h . Introduce the operator $A_h : B_{p,q}^s(M) \rightarrow H_p^{-1}(M)$ defined for $\phi \in L_1(M)$ by

$$A_h \phi := K_h * \left(\frac{\phi}{\rho_h} \right) - \phi = \int_M K_h(\cdot - y) (\tilde{\phi}(y) - \tilde{\phi}) \, \text{dvol}_M(y). \quad (4.1)$$

Then,

$$\begin{aligned} \|f_h - f\|_{H_p^{-1}(M)} &= \|A_h f\|_{H_p^{-1}(M)} \leq \|A_h\|_{B_{p,q}^s(M), H_p^{-1}(M)} \|f\|_{B_{p,q}^s(M)} \\ &\leq \|A_h\|_{B_{p,q}^s(M), H_p^{-1}(M)} L_s. \end{aligned} \quad (4.2)$$

Therefore, it suffices to control the operator norm of A_h to bound the bias.

Proposition 4.1. *Let $0 < s \leq k-1$ and $1 \leq p < \infty$. Assume that the kernel K is of order k . Then, if $h \lesssim 1$,*

$$\|A_h\|_{B_{p,q}^s(M), H_p^{-1}(M)} \lesssim h^{s+1}. \quad (4.3)$$

The proof of Proposition 4.1 consists in considering the Taylor expansion of a function $\phi \in B_{p,q}^s(M)$. In this Taylor expansion, all polynomial terms of low order disappear when integrated against K , as the kernel K is of sufficiently large order. Namely, we have the following property, whose proof is given in Appendix C.

Lemma 4.2. *Assume that the kernel K is of order k and let $B : (\mathbb{R}^D)^j \rightarrow \mathbb{R}$ be a tensor of order $1 \leq j < k$. Then, for all $x \in M$,*

$$\left| \int_M K_h(x-y) B[(x-y)^{\otimes j}] dy \right| \lesssim \|B\|_{\text{op}} h^k \quad (4.4)$$

$$|\rho_h(x) - 1| \lesssim h^{k-1} \quad \text{and} \quad \|\rho_h\|_{C^j(M)} \lesssim h^{k-1-j} \quad (4.5)$$

Let us now give a sketch of proof of Proposition 4.1 in the case $0 < s \leq 1$. The $H_p^{-1}(M)$ -norm of $A_h\phi$ is by definition equal to

$$\|A_h\phi\|_{H_p^{-1}(M)} = \sup \left\{ \int (A_h\phi)g d\text{vol}_M, \|g\|_{H_{p^*}^1(M)} \leq 1 \right\}.$$

Let $g \in H_{p^*}^1(M)$ with $\|g\|_{H_{p^*}^1(M)} \leq 1$. We use the following symmetrization trick:

$$\begin{aligned} \int (A_h\phi)(x)g(x)dx &= \iint K_h(x-y)(\tilde{\phi}(y) - \tilde{\phi}(x))g(x)dydx \\ &= \iint K_h(y-x)(\tilde{\phi}(x) - \tilde{\phi}(y))g(y)dydx \quad (\text{by swapping the indexes } x \text{ and } y) \\ &= \frac{1}{2} \iint K_h(x-y)(\tilde{\phi}(y) - \tilde{\phi}(x))(g(x) - g(y))dydx \end{aligned} \quad (4.6)$$

where, at the last line, we averaged the two previous lines and used that K is an even function. Informally, as $K_h(x-y) = 0$ if $|x-y| \geq h$, and as ρ_h is roughly constant, we expect $|\tilde{\phi}(y) - \tilde{\phi}(x)|$ to be of order h^s and $|g(x) - g(y)|$ to be of order h , leading to a bound of $\int (A_h\phi)(x)g(x)dx$ of order h^{s+1} . Such a trick can be generalized to a function ϕ of regularity $s > 1$ by writing a Taylor expansion of ϕ and using that all the polynomial terms $(x-y)^\alpha$ vanish when integrated against $K_h(x-y)$ according to Lemma 4.2. More precisely, we prove in Appendix C the following higher order analogue of the symmetrization trick.

Lemma 4.3 (Symmetrization trick). *There exists $h_0 \lesssim 1$ such that the following holds. Let l be an even number between 0 and $k-1$ and let $K^{(l)}(x) = \int_0^1 K_\lambda(x) \frac{(1-\lambda)^{l-1} \lambda^{-l}}{(l-1)!} d\lambda$ for $x \in \mathbb{R}^D$. Fix $x_0 \in M$ and let $\phi \in C^\infty(M)$ be a function supported in $\mathcal{B}_M(x_0, h_0)$. Define $\tilde{\phi}_l := d^l(\tilde{\phi} \circ \Psi_{x_0}) \circ \tilde{\pi}_{x_0}$. Let $g \in L_{p^*}(M)$ with $\|g\|_{L_{p^*}(M)} \leq 1$.*

Then, for $h \lesssim 1$, $\int (A_h\phi)(x)g(x)dx$ is equal to

$$\frac{1}{2} \iint_{\mathcal{B}_M(x_0, h_0)^2} K_h^{(l)}(x-y)(\tilde{\phi}_l(y) - \tilde{\phi}_l(x)) [\pi_{x_0}(x-y)]^{\otimes l} (g(x) - g(y)) dydx + R, \quad (4.7)$$

where R is a remainder term satisfying $|R| \lesssim \|\tilde{\phi}\|_{H_p^l(M)} h^{l+1}$. Furthermore, if $l \leq k-2$, we have $|R| \lesssim \|\tilde{\phi}\|_{H_p^{l+1}(M)} h^{l+2}$.

There are two restrictions in the above lemma: (i) the function ϕ has to be supported on a small ball and (ii) the regularity s of ϕ is an integer l . We will bypass the first restriction by using a partition of unity, whereas we will use interpolation inequalities to go from integer regularity l to a regularity s being any real number between 0 and $k-1$. Let us first see how one can use Lemma 4.3 to bound the norm of $\int (A_h\phi)(x)g(x)dx$. We use the following technical lemma.

Lemma 4.4. *Let $\eta \in \mathcal{C}^\infty(M)$ and let $0 \leq l \leq k - 2$. Assume that either $l = 0$ or that η is supported on $\mathcal{B}_M(x_0, h_0)$. Let $\eta_l = d^l(\eta \circ \Psi_{x_0}) \circ \tilde{\pi}_{x_0}$. Then, for any $h \lesssim 1$,*

$$\begin{aligned} & \left(h^{-d} \iint_{\mathcal{B}_M(x_0, h_0)^2} \mathbf{1}\{|x - y| \leq h\} \frac{\|\eta_l(x) - \eta_l(y)\|_{\text{op}}^p}{|x - y|^p} dx dy \right)^{1/p} \\ & \lesssim \left(\int_{\mathcal{B}_M(x_0, h_0)} \|\eta_{l+1}(x)\|_{\text{op}}^p dx \right)^{1/p} \lesssim \|\eta\|_{H_p^{l+1}(M)}. \end{aligned} \quad (4.8)$$

Let $\phi \in \mathcal{C}^\infty(M)$ be a function supported in $\mathcal{B}_M(x_0, h_0)$ and $g \in H_{p^*}^1(M)$ with $\|g\|_{H_{p^*}^1(M)} \leq 1$. Case 1: the regularity s is an even number. Let $l = s$. Assume first that $p > 1$ and that g is smooth. We may use that $|K_{h\lambda}(x)| = 0$ if $|x| > h\lambda$ to obtain

$$\begin{aligned} & \iint_{\mathcal{B}_M(x_0, h_0)^2} |K_{\lambda h}(x - y)| \|\tilde{\phi}_l(y) - \tilde{\phi}_l(x)\|_{\text{op}} |g(x) - g(y)| |x - y|^l dx dy \\ & \leq \|K\|_{\mathcal{C}^0(\mathbb{R}^D)} (\lambda h)^{l+1-d} \iint_{\mathcal{B}_M(x_0, h_0)^2} \mathbf{1}\{|x - y| \leq \lambda h\} \|\tilde{\phi}_l(y) - \tilde{\phi}_l(x)\|_{\text{op}} \frac{|g(x) - g(y)|}{|x - y|} dx dy \\ & \lesssim (\lambda h)^{l+1} \left((\lambda h)^{-d} \iint_{\mathcal{B}_M(x_0, h_0)^2} \mathbf{1}\{|x - y| \leq \lambda h\} \|\tilde{\phi}_l(y) - \tilde{\phi}_l(x)\|_{\text{op}}^p dx dy \right)^{1/p} \\ & \quad \times \left((\lambda h)^{-d} \iint_{\mathcal{B}_M(x_0, h_0)^2} \mathbf{1}\{|x - y| \leq \lambda h\} \frac{|g(x) - g(y)|^{p^*}}{|x - y|^{p^*}} dx dy \right)^{1/p^*} \\ & \lesssim (\lambda h)^{l+1} \left((\lambda h)^{-d} \int_{x \in \mathcal{B}_M(x_0, h_0)} \|\tilde{\phi}^l(x)\|_{\text{op}}^p \text{vol}_M(\mathcal{B}_M(x, \lambda h)) dx \right)^{1/p} \|g\|_{H_{p^*}^1(M)} \\ & \lesssim \|\tilde{\phi}\|_{H_p^l(M)} (\lambda h)^{l+1} \lesssim \|\phi\|_{H_p^l(M)} (\lambda h)^{l+1}, \end{aligned} \quad (4.9)$$

where at the last line, we used Lemma A.1(iii) in the Appendix to control the volume of $\mathcal{B}_M(x, \lambda h)$ and, at the second to last line, we used Lemma 4.4. Furthermore, it follows from Leibniz formula for the derivative of a product and Lemma 4.2 that $\|\tilde{\phi}\|_{H_p^l(M)} \lesssim \|\phi\|_{H_p^l(M)}$.

As $\mathcal{C}^\infty(M)$ is dense in $H_{p^*}^1(M)$, inequality (4.9) actually holds for every $g \in H_{p^*}^1(M)$. If $p = 1$, then every function $g \in H_{p^*}^1(M)$ with $\|g\|_{H_{p^*}^1(M)} \leq 1$ is Lipschitz continuous for the distance d_g (see Remark 2.4). Using that $d_g(x, y) \leq 2|x - y|$ if $|x - y| \leq \tau_{\min}/4$ (see [AL18, Proposition 30]), a similar computation than in the case $p < \infty$ shows that inequality (4.9) also holds if $p = \infty$.

By integrating inequality (4.9) against $\lambda \in (0, 1)$ and by using Lemma 4.3, we obtain the inequality $\|A_h \phi\|_{H_p^{-1}(M)} \lesssim h^{s+1} \|\phi\|_{H_p^s(M)}$.

Case 2: the regularity s is an odd number. Similarly, we treat the case where $s \leq k - 1$ is odd. Let $l = s - 1$. Once again, assume first that $p > 1$ and that g is smooth. Then,

$$\begin{aligned} & \iint_{\mathcal{B}_M(x_0, h_0)^2} |K_{\lambda h}(x - y)| \|\tilde{\phi}_l(y) - \tilde{\phi}_l(x)\|_{\text{op}} |g(x) - g(y)| |x - y|^l dx dy \\ & \leq \iint_{\mathcal{B}_M(x_0, h_0)^2} |K_{\lambda h}(x - y)| \frac{\|\tilde{\phi}_l(y) - \tilde{\phi}_l(x)\|_{\text{op}}}{|x - y|} \frac{|g(x) - g(y)|}{|x - y|} |x - y|^{l+2} dx dy \\ & \leq \|K\|_{\mathcal{C}^0(\mathbb{R}^D)} (\lambda h)^{l+2-d} \iint_{\mathcal{B}_M(x_0, h_0)^2} \mathbf{1}\{|x - y| \leq \lambda h\} \frac{\|\tilde{\phi}_l(y) - \tilde{\phi}_l(x)\|_{\text{op}}}{|x - y|} \frac{|g(x) - g(y)|}{|x - y|} dx dy \end{aligned}$$

$$\begin{aligned}
&\lesssim (\lambda h)^{l+2} \left((\lambda h)^{-d} \iint_{\mathcal{B}_M(x_0, h_0)^2} \mathbf{1}\{|x-y| \leq \lambda h\} \|\tilde{\phi}_l(y) - \tilde{\phi}_l(x)\|_{\text{op}}^p dx dy \right)^{1/p} \\
&\quad \times \left((\lambda h)^{-d} \iint_{\mathcal{B}_M(x_0, h_0)^2} \mathbf{1}\{|x-y| \leq \lambda h\} \frac{|g(x) - g(y)|^{p^*}}{|x-y|^{p^*}} dx dy \right)^{1/p^*} \\
&\lesssim (\lambda h)^{l+2} \|\phi\|_{H_p^s(M)}, \tag{4.10}
\end{aligned}$$

where at last line we used Lemma 4.4 and the inequality $\|\tilde{\phi}\|_{H_p^l(M)} \lesssim \|\phi\|_{H_p^l(M)}$. As in the previous case, the same inequality holds for $g \in H_{p^*}^1(M)$ non necessarily smooth and if $p = 1$. By using Lemma 4.3 and by integrating (4.10) against $\lambda \in (0, 1)$, we obtain that $\|A_h \phi\|_{H_p^{-1}(M)} \lesssim h^{s+1} \|\phi\|_{H_p^s(M)}$.

So far, we have proven that

$$\|A_h \phi\|_{H_p^{-1}(M)} \lesssim h^{s+1} \|\phi\|_{H_p^s(M)} \tag{4.11}$$

for all integers $0 \leq s \leq k-1$ and ϕ a smooth function supported on $\mathcal{B}_M(x_0, h_0)$. To obtain the result when ϕ is not supported on some ball $\mathcal{B}_M(x_0, h_0)$, we use an appropriate partition of unity. Indeed, for $\delta = h_0/8$, standard packing arguments show the existence of a set S_0 of cardinality $N \leq c_d \text{vol}_M \delta^{-d}$ with $d_H(M^\delta | S_0) \leq 5\delta/3$. By the remark following Lemma 3.6, the output S of the farthest point sampling algorithm with parameter $7\delta/3$ satisfies the assumption of Lemma 3.6, and is of cardinality smaller than $N \lesssim 1$. We consider such a covering $(\mathcal{B}_M(x, h_0))_{x \in S}$, with associated partition of unity $(\chi_x)_{x \in S}$ given by Lemma 3.6. Then, $\|A_h \phi\|_{H_p^{-1}(M)}$ is bounded by

$$\sum_{x \in S} \|A_h(\chi_x \phi)\|_{H_p^{-1}(M)} \lesssim h^{s+1} \sum_{x \in S} \|\chi_x \phi\|_{H_p^s(M)} \lesssim h^{s+1} \sum_{x \in S} \|\chi_x\|_{C^s(M)} \|\phi\|_{H_p^s(M)} \lesssim h^{s+1} \|\phi\|_{H_p^s(M)},$$

where the second to last inequality follows from Leibniz rule for the derivative of a product. Also, the last inequality follows from the fact that $(\chi_x)|_M = \chi_x \circ i_M$, where $i_M : M \rightarrow M^\delta$ is the inclusion, which is a C^k function with controlled C^k -norm. Hence, $\|\chi_x\|_{C^s(M)} \lesssim \|\chi_x\|_{C^s(M^\delta)} \lesssim 1$ by the chain rule.

As $C^\infty(M)$ is dense in $H_p^s(M)$, this gives the desired bound on the operator norm of $A_h : H_p^s(M) \rightarrow H_p^{-1}(M)$ for $0 \leq s \leq k-1$ an integer. To obtain the conclusion for Besov spaces $B_{p,q}^s(M)$, we use an interpolation inequality [Lun18, Theorem 1.1.6]. By the reiteration theorem [Lun18, Theorem 1.3.5], for $0 < s < k-1$, $B_{p,q}^s(M) = (L_p(M), H_p^{k-1}(M))_{s/(k-1), q}$, with an equivalent norm. Hence, we have, for $0 < s < k-1$, letting $\theta = s/(k-1)$,

$$\begin{aligned}
\|A_h\|_{B_{p,q}^s(M), H_p^{-1}(M)} &\lesssim \|A_h\|_{L_p(M), H_p^{-1}(M)}^{1-\theta} \|A_h\|_{H_p^{k-1}(M), H_p^{-1}(M)}^\theta \\
&\lesssim h^{1-\frac{s}{k-1}} h^{k\frac{s}{k-1}} \lesssim h^{s+1},
\end{aligned}$$

so that Proposition 4.1 is proven for $s < k-1$. It remains to prove the inequality in the case $s = k-1$. By Fatou's lemma and the definition of interpolation spaces [Lun18, Definition 1.1.2], we have, for some constant C not depending on s ,

$$\|A_h f\|_{B_{p,q}^{k-1}(M)} \leq \liminf_{\substack{s \rightarrow k-1 \\ s < k-1}} \|A_h f\|_{B_{p,q}^s(M)} \leq \liminf_{\substack{s \rightarrow k-1 \\ s < k-1}} \left(C h^{s+1} \|f\|_{B_{p,q}^s(M)} \right) \leq C h^k \|f\|_{B_{p,q}^{k-1}(M)},$$

where we used that $\|f\|_{B_{p,q}^s(M)} \leq \|f\|_{B_{p,q}^{k-1}(M)}$. This concludes the proof of Proposition 4.1.

4.2. Fluctuations of the kernel density estimator. The second step in bounding the Sobolev risk $\mathbb{E}\|f_{n,h} - f\|_{H_p^{-1}(M)}$ is to control the fluctuation term $\mathbb{E}\|f_{n,h} - f_h\|_{H_p^{-1}(M)}$. If we were considering a classical L_p -norm instead of a negative Sobolev norm, then we could simply express $f_{n,h} - f_h$ as a sum of the i.i.d. terms to obtain the right order. The key idea to bound the fluctuation term is to show that $\|f_{n,h} - f\|_{H_p^{-1}(M)}$ is smaller than the L_p -norm of a similar sum of i.i.d. functions that can be expressed in term of the Green's function on M . We obtain the following control.

Proposition 4.5. *Let $\mu \in \mathcal{Q}^s(M)$ with Y_1, \dots, Y_n a n -sample of law μ . Assume that $h \lesssim 1$ and that $nh^d \gtrsim 1$. Then,*

$$\mathbb{E} \|f_{n,h} - f_h\|_{H_p^{-1}(M)} \lesssim n^{-1/2} h^{1-d/2} I_d(h), \quad (4.12)$$

where $I_d(h)$ is defined in Theorem 3.1.

Let Δ be the Laplace-Beltrami operator on M and $G : \mathcal{D}_M \rightarrow \mathbb{R}$ be a Green's function, defined on $\mathcal{D}_M := \{(x, y) \in M \times M, x \neq y\}$ (see [Aub82, Chapter 4]). By definition, if $\phi \in \mathcal{C}^\infty(M)$, then the function $G\phi : x \in M \mapsto \int G(x, y)\phi(y)dy$ is a smooth function satisfying $\Delta G\phi = \phi$, with $\nabla G\phi(x) = \int \nabla_x G(x, y)\phi(y)dy$ for $x \in M$. Hence, if $w = \nabla G\phi$, then $\nabla \cdot w = \phi$, so that, Proposition 2.5 yields

$$\|\phi\|_{H_p^{-1}(M)} \leq \|\phi\|_{\dot{H}_p^{-1}(M)} \leq \|\nabla G\phi\|_{L_p(M)}.$$

By linearity, we have

$$\|f_{n,h} - f_h\|_{H_p^{-1}(M)} \leq \left\| \frac{1}{n} \sum_{i=1}^n \nabla G \left(K_h * \left(\frac{\delta_{Y_i}}{\rho_h(Y_i)} \right) \right) - \mathbb{E} \left[\nabla G \left(K_h * \left(\frac{\delta_{Y_i}}{\rho_h(Y_i)} \right) \right) \right] \right\|_{L_p(M)}. \quad (4.13)$$

The expectation of the L_p -norm of the sum of i.i.d. centered functions is controlled using Rosenthal inequality.

Lemma 4.6. *Let U_1, \dots, U_n be i.i.d. functions on $L_p(M)$. Then, $\mathbb{E} \left\| \frac{1}{n} \sum_{i=1}^n (U_i - \mathbb{E}U_i) \right\|_{L_p(M)}^p$ is smaller than*

$$\begin{cases} n^{-p/2} \int (\mathbb{E} [|U_1(z)|^2])^{p/2} dz & \text{if } p \leq 2, \\ C_p n^{-p/2} \int (\mathbb{E} |U_1(z)|^2)^{p/2} dz + C_p n^{1-p} \int_M \mathbb{E} [|U_1(z)|^p] dz & \text{if } p > 2. \end{cases} \quad (4.14)$$

Proof. If $p \leq 2$, one has by Jensen's inequality

$$\mathbb{E} \left| \sum_{i=1}^n (U_i(z) - \mathbb{E}U_i(z)) \right|^p \leq \left(\mathbb{E} \left| \sum_{i=1}^n (U_i(z) - \mathbb{E}U_i(z)) \right|^2 \right)^{p/2} \leq n^{p/2} (\mathbb{E} |U_1(z)|^2)^{p/2}$$

and (4.14) follows by integrating this inequality against $z \in M$. For $p > 2$, we use Rosenthal inequality [Ros70, Theorem 3] for a fixed $z \in M$, and then integrate the inequality against $z \in M$. \square

It remains to bound $\mathbb{E} \left[\left| \nabla G \left(K_h * \left(\frac{\delta_Y}{\rho_h(Y)} \right) \right) (z) \right|^p \right]$ where $Y \sim \mu$, $z \in M$ and $p \geq 2$.

Lemma 4.7. *Let $p \geq 2$. Then, for all $z \in M$ and $h \lesssim 1$,*

$$\mathbb{E} \left[\left| \nabla G \left(K_h * \left(\frac{\delta_Y}{\rho_h(Y)} \right) \right) (z) \right|^p \right] \lesssim \begin{cases} 1 & \text{if } d = 1 \\ -\log h & \text{if } p = d = 2 \\ h^{p+d-dp} & \text{else.} \end{cases} \quad (4.15)$$

A proof of Lemma 4.7 is found in Appendix D. From (4.13), Lemma 4.6 and Lemma 4.7, we obtain, in the case $p \geq 2$ and $d \geq 3$

$$\begin{aligned} \mathbb{E} \|f_{n,h} - f_h\|_{H_p^{-1}(M)} &\leq \left(\mathbb{E} \|f_{n,h} - f_h\|_{H_p^{-1}(M)}^p \right)^{1/p} \\ &\leq C_p n^{-1/2} \left(\int \left(\mathbb{E} \left| \nabla G \left(K_h * \left(\frac{\delta_Y}{\rho_h(Y)} \right) \right) (z) \right|^2 \right)^{p/2} dz \right)^{1/p} \\ &\quad + C_p n^{1/p-1} \left(\int \mathbb{E} \left[\left| \nabla G \left(K_h * \left(\frac{\delta_Y}{\rho_h(Y)} \right) \right) (z) \right|^p \right] dz \right)^{1/p} \\ &\lesssim n^{-1/2} |\text{vol}_M|^{1/p} h^{1-d/2} + n^{1/p-1} |\text{vol}_M|^{1/p} h^{1+d/p-d}. \end{aligned}$$

Recalling that $|\text{vol}_M| \leq f_{\min}^{-1} \lesssim 1$ and that $nh^d \gtrsim 1$, one can check that this quantity is smaller up to a constant than $n^{-1/2}h^{1-d/2}$, proving Proposition 4.5 in the case $p \geq 2$ and $d \geq 3$. A similar computation shows that Proposition 4.5 also holds if $p \leq 2$ or $d \leq 2$.

4.3. Proof of Theorem 3.1. We are now ready to conclude the proof of Theorem 3.1. The proof of (i) follows from a standard control of the ∞ -norm between f and $f_{n,h}$ and is found in Appendix E. Point (ii) is a combination of the results from the two previous sections, and we can obtain (iii) using the linearization inequality (Proposition 2.8). More precisely, let E be the event described in (i). If E is realized, then $\mu_{n,h}^0$ is equal to $\mu_{n,h}$, and it satisfies $\mu_{n,h}^0 \geq \frac{f_{\min}}{2} \text{vol}_M$. Thus, Proposition 2.8 yields $W_p(\mu_{n,h}^0, \mu) \lesssim \|\mu_{n,h} - \mu\|_{H_p^{-1}(M)}$. If E is not realized, we bound $W_p(\mu_{n,h}^0, \mu)$ by $\text{diam}(M)$, which is itself bounded by a constant depending only on the parameters of the model (see [AL18, Lemma 2.2]). Hence,

$$\begin{aligned} \mathbb{E}W_p(\mu_{n,h}^0, \mu) &\leq \mathbb{E} [W_p(\mu_{n,h}^0, \mu)\mathbf{1}\{E\}] + \text{diam}(M)\mathbb{P}(E^c) \\ &\lesssim \mathbb{E}\|\mu_{n,h} - \mu\|_{H_p^{-1}(M)} + n^{-k/d}, \end{aligned}$$

and we conclude thanks to (3.4). Eventually, a proof of the minimax lower bound (iv), based on Assouad's lemma, is given in Appendix G.

4.4. Proofs of Theorem 3.7 and Theorem 3.8. There are three different statement to prove in Theorem 3.7. The first one is a direct application of the construction of partitions of unity proposed in Lemma 3.6.

. *Proof of Theorem 3.7(i).*

Assume that $\gamma \leq \varepsilon/24$. Let $\mathbb{X} = \{X_1, \dots, X_n\}$ and $\mathbb{Y} = \{Y_1, \dots, Y_n\}$. By the remark following Lemma 3.6, the existence of a partition of unity satisfying the requirements of Theorem 3.7(i) is ensured as long as $d_H(M^{\varepsilon/8}|\mathbb{X}) \leq 5\varepsilon/24$. We have $d_H(M^{\varepsilon/8}|\mathbb{X}) \leq d_H(M^{\varepsilon/8}|\mathbb{Y}) + \varepsilon/24 \leq d_H(M|\mathbb{Y}) + 4\varepsilon/24$. Hence, the partition of unity exists if $d_H(M|\mathbb{Y}) \leq \varepsilon/24$. This is satisfied with probability larger than $1 - cn^{-k/d}$ if $\varepsilon \gtrsim (\log n/n)^{1/d}$ by [Aam17, Lemma III.23].

. *Proof of Theorem 3.7(ii).* For ease of notation, we will assume that the output $\{X_{i_1}, \dots, X_{i_J}\}$ of the farthest point sampling algorithm is equal to $\{X_1, \dots, X_J\}$. The measure $\widehat{\text{vol}}_M$ can be written as $\sum_{j=1}^J \xi_j$, where ξ_j is the measure having density χ_j with respect to the d -dimensional Hausdorff measure on $\widehat{\Psi}_j(\widehat{T}_j)$.

Remark that the restriction of each function χ_j on M is supported on a small neighborhood covered by the chart Ψ_{Y_j} . We may therefore write for any continuous bounded function ϕ ,

$$\int \phi(x) d\text{vol}_M(x) = \sum_{j=1}^J \int_{\Psi_{Y_j}(T_{Y_j})} \phi(x) \chi_j(x) dx.$$

As $\widehat{\Psi}_j$ is at distance $\varepsilon^k + \gamma$ from Ψ_{Y_j} and χ_j is Lipschitz, one can hope that the measure $\widehat{\text{vol}}_M$ defined in (3.13) is also at distance $\varepsilon^k + \gamma$ from vol_M . The Wasserstein distance between ξ_j and the measure $\chi_j \cdot \text{vol}_M$ is bounded in two steps. First, we transport ξ_j on M by using the parametrizations $\widehat{\Psi}_j$ and Ψ_{Y_j} . When doing so, we obtain a measure on M , whose density is a modification of χ_j , that is distorted by the Jacobian of the transport map. We then crudely bound the Wasserstein distance between this new measure and $\chi_j \cdot \text{vol}_M$ by their L_1 -distance. We first need a technical result that controls how the density χ_j is impacted by the transport map.

Lemma 4.8. *If $(\log n/n)^{1/d} \lesssim \varepsilon \lesssim 1$ and $\gamma \lesssim \varepsilon$, with probability larger than $1 - cn^{-k/d}$, for all $j = 1, \dots, J$:*

- (a) *The map $\Psi_{Y_j} \circ \pi_{Y_j} : \mathcal{B}_{\widehat{T}_j}(0, 3\varepsilon) \rightarrow M$ is a diffeomorphism on its image, which contains $\mathcal{B}_M(Y_j, 2\varepsilon)$. Let $S_j : \mathcal{B}_M(Y_j, 2\varepsilon) \rightarrow \mathcal{B}_{\widehat{T}_j}(0, 3\varepsilon)$ be the inverse of $\Psi_{Y_j} \circ \pi_{Y_j}$. Then, $\widehat{\Psi}_j \circ S_j : \mathcal{B}_M(Y_j, 2\varepsilon) \rightarrow \widehat{\Psi}_j(\widehat{T}_j)$ is also a diffeomorphism on its image, which contains $\mathcal{B}_{\widehat{\Psi}_j(\widehat{T}_j)}(X_j, \varepsilon)$. Furthermore, for all $z \in \mathcal{B}_M(Y_j, 2\varepsilon)$, we have $|\widehat{\Psi}_j \circ S_j(z) - X_j| \geq \frac{7}{8}|z - Y_j|$.*




FIGURE 1. Illustration of Lemma 4.8(a)

(b) The measure $(\hat{\Psi}_j \circ S_j)_\#^{-1} \xi_j$ has a density $\tilde{\chi}_j$ on M equal to

$$\tilde{\chi}_j(z) = \chi_j(\hat{\Psi}_j \circ S_j(z)) J(\hat{\Psi}_j \circ S_j)(z) \text{ for } z \in M, \quad (4.16)$$

where the function is extended by 0 for $z \in M \setminus \mathcal{B}_M(Y_j, 2\varepsilon)$.

(c) For $z \in \mathcal{B}_M(Y_j, 2\varepsilon)$, we have

$$|\hat{\Psi}_j \circ S_j(z) - z| \lesssim \varepsilon^m + \gamma, \quad (4.17)$$

$$|\tilde{\chi}_j(z) - \chi_j(z)| \lesssim (\varepsilon^m + \gamma)(1 + \gamma\varepsilon^{-2}). \quad (4.18)$$

A proof of Lemma 4.8 is found in Appendix F. Let $\hat{M}_\varepsilon = \bigcup_{j=1}^J \mathcal{B}_{\hat{\Psi}_j(\hat{T}_j)}(X_j, \varepsilon)$ be the support of $\widehat{\text{vol}}_M$. We are now ready to state a stability result between the approximated measure $\widehat{\text{vol}}_M$ and vol_M . Note that we state a lemma that is slightly more general (considering measures having a density ϕ with respect to the volume measures), so that we can also control the distance between the kernel density estimators of Theorem 3.8.

Lemma 4.9. *Let $(\log n/n)^{1/d} \lesssim \varepsilon \lesssim 1$ and $\gamma \lesssim \varepsilon$. Fix $1 \leq r \leq \infty$ and let $\phi : M \rightarrow \mathbb{R}$, $\tilde{\phi} : \hat{M}_\varepsilon \rightarrow \mathbb{R}$ be functions satisfying $\phi_{\min} \leq \phi, \tilde{\phi} \leq \phi_{\max}$ for some positive constants $\phi_{\min}, \phi_{\max} > 0$. Assume further that for all $j = 1, \dots, J$ and for all $z \in M$ we have, $|\tilde{\phi}(\hat{\Psi}_j \circ S_j(z)) - \phi(z)| \leq T \lesssim 1$. Then, with probability larger than $1 - cn^{-k/d}$, we have*

$$W_r \left(\frac{\tilde{\phi} \cdot \widehat{\text{vol}}_M}{|\tilde{\phi} \cdot \widehat{\text{vol}}_M|}, \frac{\phi \cdot \text{vol}_M}{|\phi \cdot \text{vol}_M|} \right) \lesssim C_0(T + (\varepsilon^m + \gamma)(1 + \gamma\varepsilon^{-2})), \quad (4.19)$$

where C_0 depends on ϕ_{\min} and ϕ_{\max} .

In particular, inequality (3.14) is a consequence of Lemma 4.9 with $\phi \equiv \tilde{\phi} \equiv 1$.

Proof. Assume first that $r < \infty$. We have the bound

$$\begin{aligned} W_r \left(\frac{\tilde{\phi} \cdot \widehat{\text{vol}}_M}{|\tilde{\phi} \cdot \widehat{\text{vol}}_M|}, \frac{\phi \cdot \text{vol}_M}{|\phi \cdot \text{vol}_M|} \right) &= \frac{1}{|\tilde{\phi} \cdot \widehat{\text{vol}}_M|^{1/r}} W_r \left(\tilde{\phi} \cdot \widehat{\text{vol}}_M, \phi \cdot \text{vol}_M \frac{|\tilde{\phi} \cdot \widehat{\text{vol}}_M|}{|\phi \cdot \text{vol}_M|} \right) \\ &\leq \frac{1}{|\tilde{\phi} \cdot \widehat{\text{vol}}_M|^{1/r}} \left(W_r \left(\sum_{j=1}^J \tilde{\phi} \cdot \xi_j, \sum_{j=1}^J (\hat{\Psi}_j \circ S_j)_\#^{-1}(\tilde{\phi} \cdot \xi_j) \right) \right. \\ &\quad \left. + W_r \left(\sum_{j=1}^J (\hat{\Psi}_j \circ S_j)_\#^{-1}(\tilde{\phi} \cdot \xi_j), \phi \cdot \text{vol}_M \frac{|\tilde{\phi} \cdot \widehat{\text{vol}}_M|}{|\phi \cdot \text{vol}_M|} \right) \right) \end{aligned} \quad (4.20)$$

We use Proposition 2.8 to bound the second term in (4.20). By a change of variables, the density of $(\hat{\Psi}_j \circ S_j)_\#^{-1}(\tilde{\phi} \cdot \xi_j)$ is given by $\tilde{\phi}_j : z \mapsto \tilde{\phi}(\hat{\Psi}_j \circ S_j(z)) \tilde{\chi}_j(z)$. With probability larger than

$1 - cn^{-k/d}$, we have for $z \in M$, should $(\varepsilon^m + \gamma)(1 + \gamma\varepsilon^{-2})$ be small enough,

$$\sum_{j=1}^J \tilde{\chi}_j(z) \geq \sum_{j=1}^J \chi_j(z) - Cc_d(\varepsilon^m + \gamma)(1 + \gamma\varepsilon^{-2}) \geq 1 - \frac{1}{2} = \frac{1}{2},$$

where c_d is the constant of Lemma 3.6. Therefore, the density of $\sum_{j=1}^J (\hat{\Psi}_j \circ S_j)_{\#}^{-1}(\tilde{\phi} \cdot \xi_j)$ is larger than $\phi_{\min}/2$. Remark also that $\tilde{\chi}_j(z) \leq 2$ for any $z \in M$. Hence, we have according to Lemma 4.8, $|\tilde{\phi}_j(z) - \phi(z)\chi_j(z)| \leq T + 2\phi_{\max}|\chi_j(z) - \tilde{\chi}_j(z)| \lesssim T + \phi_{\max}(\varepsilon^m + \gamma)(1 + \gamma\varepsilon^{-2})$ for some constant C_0 . This gives the bound,

$$\begin{aligned} \left| |\tilde{\phi} \cdot \widehat{\text{vol}}_M| - |\phi \cdot \text{vol}_M| \right| &\leq \left\| \sum_{j=1}^J \tilde{\phi}_j - \phi \right\|_{L_1(M)} \leq \left\| \sum_{j=1}^J \tilde{\phi}_j - \phi \right\|_{L_r(M)} |\text{vol}_M|^{1-1/r} \\ &\leq \left\| \sum_{j=1}^J \tilde{\phi}_j - \phi \right\|_{L_\infty(M)} |\text{vol}_M| \\ &\leq C_0 |\text{vol}_M| (T + \phi_{\max}(\varepsilon^m + \gamma)(1 + \gamma\varepsilon^{-2})). \end{aligned} \quad (4.21)$$

Therefore, $\phi \frac{|\tilde{\phi} \cdot \widehat{\text{vol}}_M|}{|\phi \cdot \text{vol}_M|}$ is larger than

$$\begin{aligned} &\phi_{\min} \left(1 - C_0 |\text{vol}_M| \frac{T + \phi_{\max}(\varepsilon^m + \gamma)(1 + \gamma\varepsilon^{-2})}{\phi_{\min} |\text{vol}_M|} \right) \\ &\geq \phi_{\min} - C_0 (T + \phi_{\max}(\varepsilon^m + \gamma)(1 + \gamma\varepsilon^{-2})) \geq \frac{\phi_{\min}}{2} \end{aligned}$$

if T, ε and γ are small enough. Hence, by Proposition 2.8 and using (4.21),

$$\begin{aligned} &W_r \left(\sum_{j=1}^J (\hat{\Psi}_j \circ S_j)_{\#}^{-1}(\tilde{\phi} \cdot \xi_j), \phi \cdot \text{vol}_M \frac{|\tilde{\phi} \cdot \widehat{\text{vol}}_M|}{|\phi \cdot \text{vol}_M|} \right) \\ &\leq r^{-1/r} \left(\frac{2}{\phi_{\min}} \right)^{1-1/r} \left\| \sum_{j=1}^J \tilde{\phi}_j - \phi \frac{|\tilde{\phi} \cdot \widehat{\text{vol}}_M|}{|\phi \cdot \text{vol}_M|} \right\|_{H_r^{-1}(M)} \\ &\leq \left(\frac{2}{\phi_{\min}} \vee 1 \right) \left\| \sum_{j=1}^n \tilde{\phi}_j - \phi \frac{|\tilde{\phi} \cdot \widehat{\text{vol}}_M|}{|\phi \cdot \text{vol}_M|} \right\|_{L_r(M)} \\ &\leq \left(\frac{2}{\phi_{\min}} \vee 1 \right) \left(\left\| \sum_{j=1}^J \tilde{\phi}_j - \phi \right\|_{L_r(M)} + \frac{\left| |\phi \cdot \text{vol}_M| - |\tilde{\phi} \cdot \widehat{\text{vol}}_M| \right|}{|\phi \cdot \text{vol}_M|} \|\phi\|_{L_r(M)} \right) \\ &\leq \left(\frac{2}{\phi_{\min}} \vee 1 \right) C_0 (T + \phi_{\max}(\varepsilon^m + \gamma)(1 + \gamma\varepsilon^{-2})) \left(|\text{vol}_M|^{1/r} + \frac{|\text{vol}_M|}{\phi_{\min} |\text{vol}_M|} |\text{vol}_M|^{1/r} \phi_{\max} \right) \\ &\leq C_{\phi_{\min}, \phi_{\max}} (T + (\varepsilon^m + \gamma)(1 + \gamma\varepsilon^{-2})), \end{aligned}$$

where we used that $|\text{vol}_M| \leq f_{\min}^{-1} \lesssim 1$, and the constant $C_{\phi_{\min}, \phi_{\max}}$ in the upper bound depending on ϕ_{\min} and ϕ_{\max} , but not on r .

To bound the first term in (4.20), consider the transport plan $\sum_{j=1}^J (\text{id}, (\hat{\Psi}_j \circ S_j)^{-1})_{\#}(\tilde{\phi} \cdot \xi_j)$, which has, according to Lemma 4.8, a cost bounded by

$$\sum_{j=1}^J \int |y - (\hat{\Psi}_j \circ S_j)^{-1}(y)|^r d(\tilde{\phi} \cdot \xi_j)(y) \lesssim \phi_{\max} (\varepsilon^m + \gamma)^r |\widehat{\text{vol}}_M|.$$

As $|\widehat{\text{vol}}_M| \lesssim |\text{vol}_M| + T + \phi_{\max}(\varepsilon^m + \gamma)(1 + \gamma\varepsilon^{-2}) \lesssim 1$, we obtain the desired bound. By letting $r \rightarrow \infty$, and remarking that the different constants involved are independent of r , we observe that the same bound holds for $r = \infty$. \square

Remark 4.10. Inequality (4.21) with $\phi \equiv \phi' \equiv 1$ gives a bound on the distance between the total mass of $\widehat{\text{vol}}_M$ and the volume $|\text{vol}_M|$ of M : choosing $k = m$, it is of order $(\varepsilon^k + \gamma)(1 + \gamma\varepsilon^{-2})$ with probability larger than $1 - cn^{-k/d}$.

. *Proof of Theorem 3.7(iii).*

Inequality (3.15) is a consequence of Theorem 3.7(ii), whereas the lower bound on the minimax risk (3.16) is proven in Appendix G.

. *Proof of Theorem 3.8.*

Note first that $\hat{\nu}_{n,h}$ is indeed a measure of mass 1. We show in Lemma F.2 that

$$T := \max_{j=1 \dots J} \sup_{z \in \mathcal{B}(Y_j, \varepsilon)} \left| K_h * \left(\frac{\nu_n}{\hat{\rho}_h} \right) (\hat{\Psi}_j \circ S_j(z)) - K_h * \left(\frac{\mu_n}{\rho_h} \right) (z) \right|$$

satisfies $T \lesssim (\varepsilon^m + \gamma)(1 + \gamma\varepsilon^{-2})$ with probability larger than $1 - cn^{-k/d}$, as long as $nh^d \gtrsim 1$. As $f_{\min}/2 \leq K_h * \mu_n \leq 2f_{\max}$ on M by Theorem 3.1(i), and as every $y \in \hat{M}_\varepsilon$ is in the image of $\hat{\Psi}_j \circ S_j$ for some $j = 1 \dots J$, we have $f_{\min}/3 \leq K_h * \nu_n \leq 3f_{\max}$ on \hat{M}_ε should $\varepsilon^k + \gamma$ be small enough. This proves Theorem 3.1(i) and, together with Lemma 4.9, this also proves Theorem 3.8(ii). Theorem 3.8(iii) is a consequence of Theorem 3.8(ii).

5. NUMERICAL CONSIDERATIONS

5.1. Computation of the local polynomial estimator. A crucial step in the implementation of the estimators $\widehat{\text{vol}}_M$ and $\hat{\nu}_{n,h}$ is the minimization procedure described in (3.8). Assume for the sake of simplicity that no noise is present in the dataset. Further assume without loss of generality that $0 \in M$, and let X_1, \dots, X_N be N points sampled according to some distribution $\tilde{\mu}$ supported on $\mathcal{B}_M(0, \varepsilon)$, with density lower bounded by f_{\min} (in the setting of Section 3, $\tilde{\mu}$ is the conditional probability of $X \sim \mu$ given that $X \in \mathcal{B}_M(0, \varepsilon)$, where $\mu \in \mathcal{Q}^s(M)$). Our goal is to find a minimizer of the functional

$$\frac{1}{2N} \sum_{i=1}^N \left| X_i - \pi(X_i) - \sum_{j=2}^{m-1} V_j [\pi(X_i)^{\otimes j}] \right|^2, \quad (5.1)$$

where π is an orthogonal projector of rank d and each V_j is a j -tensor with operator norm smaller than ℓ . We propose here a fast procedure with theoretical guarantees to solve this problem, answering a question raised in [AL19]. There are two issues that make this optimization problem not trivial. First, the objective functional is defined on a manifold, and second, it is not globally geodesically convex. We will actually show that the functional is λ -strongly geodesically convex and β -smooth on a small neighborhood of size ε around its minimizer, for both λ and β of order ε^2 . Furthermore, we show that it is possible to find a point in this neighborhood, by letting $V_j = 0$, and π being given by a PCA on the dataset X_1, \dots, X_N . Any standard optimization algorithm on Riemannian manifolds will then converge with such an initialization. As an example, we show that a classical gradient descent converges linearly, although it is expected that more refined algorithms such as a Riemannian SVRG will also converge, with possibly a better behavior in practice [SKM19]. Before going further, we give some background on the Grassmannian manifold that will allow us to rewrite the objective functional in a more practical way. We refer to [EAS98] for a more detailed introduction on the geometry of the Grassmannian. We also provide a short review of convex optimization on Riemannian manifolds in Appendix I.

Let $\mathcal{O}(D)$ be the manifold of orthogonal matrices on \mathbb{R}^D , that we endow with the Riemannian structure induced by the inclusion in $\mathbb{R}^{D \times D}$. For $Q \in \mathcal{O}(D)$, the tangent space of $\mathcal{O}(D)$ at Q is given by $T_Q \mathcal{O}(D) = \{QA : A \text{ skew-symmetric}\}$, whereas the exponential map is given by $\exp_Q(QA) = Q \exp(A)$.

Let $\mathcal{G}(d, D)$ be the Grassmannian manifold. It can be defined as the quotient $\mathcal{O}(D)/(\mathcal{O}(d) \times \mathcal{O}(D-d))$: we identify a d -dimensional subspace U of \mathbb{R}^D with the set of matrices

$$\left\{ Q \begin{pmatrix} Q_1 & 0 \\ 0 & Q_2 \end{pmatrix} : Q_1 \in \mathcal{O}(d), Q_2 \in \mathcal{O}(D-d) \right\}, \quad (5.2)$$

where Q is any orthogonal matrix such that the vector space $\Pi(Q)$ spanned by its first d -columns is equal to U . An orthogonal projector of rank d is then identified with π_U , the projection on the subspace U . The manifold $\mathcal{G}(D, d)$ is endowed with the Riemannian structure given by the quotient map. Fix an element $U_* \in \mathcal{G}(d, D)$, with an arbitrary representant Q_* . Then, the tangent space $T_{U_*}\mathcal{G}(d, D)$ at U_* is given by the set of matrices

$$\Delta = Q_* \begin{pmatrix} 0 & -B^\top \\ B & 0 \end{pmatrix}, \quad (5.3)$$

where B is any $(D-d) \times d$ matrix. The exponential map on the Grassmannian is given by the quotient projection of the exponential on the orthogonal group, that is $\exp_{U_*}(\Delta) = \Pi \circ \exp_{Q_*}(\Delta)$. The exponential map being surjective, any element of the Grassmannian can be written as $\Pi \circ \exp_{Q_*}(\Delta)$ for some $\Delta \in T_{U_*}\mathcal{G}(d, D)$. Furthermore, the quotient map $\Pi : \exp_{Q_*}(T_{U_*}\mathcal{G}(d, D)) \rightarrow \mathcal{G}(d, D)$ is a local isometry. As it will be convenient for us to work with deformations of a fixed orthogonal basis Q_* of \mathbb{R}^D , we will consider the $d(D-d)$ -dimensional submanifold $\mathcal{O}_*(d, D) := \exp_{Q_*}(T_{U_*}\mathcal{G}(d, D))$ of $\mathcal{O}(D)$ instead of $\mathcal{G}(d, D)$. A small neighborhood of U_* in $\mathcal{G}(d, D)$ is isometric to a neighborhood of Q_* in $\mathcal{O}_*(d, D)$, so that both points of view are equivalent. Given $Q \in \mathcal{O}_*(d, D)$, we write $Q = (Q_{[d]} \ Q_{[d,D]})$, where $Q_{[d]} = (e_1, \dots, e_d)$ is a $D \times d$ matrix, and $Q_{[d,D]} = (e_{d+1}, \dots, e_D)$ is a $D \times (D-d)$ matrix.

Given E and F two vector spaces, we let $\mathcal{S}^j(E, F)$ be the set of symmetric j -tensors from E to F . Then, the functional in (5.1) is defined on $\mathcal{G}(d, D) \times \prod_{j=2}^{m-1} \mathcal{S}^j(\mathbb{R}^D, \mathbb{R}^D)$, that is a manifold of possibly very large dimension (of order D^m). However, given a subspace $U \in \mathcal{G}(d, D)$, one can always decrease the loss function by replacing a tensor V_j by the tensor $\pi_U^\top \circ V_j \circ \pi_U$. That is, we may assume that $V_j \in \mathcal{S}^j(U, U^\perp)$. For $Q = (e_1, \dots, e_D) \in \mathcal{O}_*(d, D)$ and $V_j = (V_{j,k})_{d+1 \leq k \leq D} \in \mathcal{S}_j(\mathbb{R}^d, \mathbb{R}^{D-d})$, we let $\iota_j(Q, V)$ be the j -tensor in $\mathcal{S}_j(\Pi(Q), \Pi(Q)^\perp)$ defined by

$$\iota_j(Q, V)[x_1, \dots, x_j] = \sum_{k=d+1}^D e_k V_{j,k} [Q_{[d]}^\top(x_1 \cdots x_j)]$$

for $x_1, \dots, x_j \in \Pi(Q)^j$. Introduce the functional

$$G_{m,x}(Q, V) := \frac{1}{2} \left| x - \sum_{k=1}^d \langle x, e_k \rangle e_k - \sum_{j=2}^{m-1} \iota_j(Q, V)[x^{\otimes j}] \right|^2. \quad (5.4)$$

The function $G_m := \frac{1}{N} \sum_{i=1}^N G_{m, X_i}$ is defined on the manifold

$$\mathcal{M} := \mathcal{O}_*(d, D) \times \prod_{j=2}^{m-1} \mathcal{S}_j(\mathbb{R}^d, \mathbb{R}^{D-d}).$$

Note that \mathcal{M} is a manifold of dimension $(D-d) \sum_{j=1}^{m-1} \binom{d+j-1}{j}$, that is of order $Dd^{m-1} \ll D^m$. We endow \mathcal{M} with the Riemannian metric $g_{\mathcal{M}}$ given by

$$(g_{\mathcal{M}})_{(Q,V)}((\Delta_1, W_1), (\Delta_2, W_2)) = \langle \Delta_1, \Delta_2 \rangle + \sum_{j=2}^{m-1} \varepsilon^{2(j-1)} \langle W_{1,j}, W_{2,j} \rangle, \quad (5.5)$$

for $(Q, V) \in \mathcal{M}$ and $(\Delta_1, W_1), (\Delta_2, W_2) \in T_{(Q,V)}\mathcal{M}$. We also denote by $d_{\mathcal{M}}$ the geodesic distance on \mathcal{M} . Consider the submanifold \mathcal{M}_0 of \mathcal{M} , where we impose that each j -tensor V_j has operator norm smaller than ℓ . Then, solving (5.1) is equivalent to minimizing G_m on \mathcal{M}_0 .

Let us choose (Q_*, V_*) that minimizes G_m on \mathcal{M}_0 . Consider the (geodesically convex) neighborhood Ω of $(Q_*, 0)$ in \mathcal{M}_0 given by matrices $Q \in \mathcal{O}_*(d, D)$ that are r close from Q_* for the geodesic distance.

Proposition 5.1. *Let $\varepsilon \lesssim 1$, $\delta > 0$, and let $r \leq \delta\varepsilon$. If $1 \lesssim \ell \lesssim \varepsilon^{-1}$ is small enough with respect to δ , then, with probability $1 - e^{-cN}$, the functional G_m is geodesically λ -strongly convex and β -smooth on Ω with $\varepsilon^2 \lesssim \lambda \leq \beta \lesssim \varepsilon^2$, where the constant c depends on d, m, τ_{\min} and f_{\min} .*

This implies in particular that a gradient descent with step α of order ε^{-2} will converge at linear rate towards the minimizer of G_m , when initialized in Ω (see Appendix I for details). A number of steps of order $|\log \varepsilon| \simeq \log n$ is needed to attain a point at distance ε^{m-1} from the minimizer. Such a point (Q, V) will then satisfy the same inequalities than the minimizer $(\hat{\pi}, \hat{V})$ from Proposition 3.5. Proposition 3.5 with $m = 2$ implies that a local PCA yields a subspace that is $\delta\varepsilon$ -close from $\Pi(Q^*)$ for some $\delta > 0$. Therefore, choosing ℓ small enough, Proposition 5.1 implies that one can initialize the gradient descent at the local PCA, with linear convergence of the gradient descent.

Let us end this section with a word about the computational complexity of gradient descent. Evaluating the gradient of G_m requires $O(NDd^{m-1})$ operations. Also, one need to compute the exponential map on \mathcal{M} at each step of the gradient descent. This boils down to computing a SVD, which can be made in $O(Dd^2)$ time. In total, each step of the gradient descent takes $O(NDd^{m-1})$ time. The dependence in N is not an issue as N is of order $\log n$ in our setting (that is the expected number of points of a n -sample in a ball of radius $\varepsilon \simeq (\log n/n)^{1/d}$). Still, one could use a stochastic gradient descent algorithm to remove the N factor if needed.

5.2. Sampling from the estimators. The issue of evaluating integrals and sampling from distributions supported defined on manifolds has been addressed in several works using Monte Carlo methods: e.g. authors in [DHS13] propose rejection sampling and Gibbs sampling methods in the case where the manifold can be covered by a single known chart (except on a set of null measure), while authors in [ZHCG18] design a MCMC in the case where the manifolds is defined through equality constraints. We here address the problem of sampling from two estimators: the estimated uniform measure $\hat{U}_M := \widehat{\text{vol}}_M / |\widehat{\text{vol}}_M|$ and the measure $\hat{\mu}_{n,h}$. Our goal is not to propose state-of-the-art procedures but to show that basic sampling algorithms already have good theoretical behaviors.

Let

$$q_j := \hat{U}_M(\hat{\Psi}_{i_j}(\hat{T}_j)) = \int_{\hat{T}_j} \chi_j(\hat{\Psi}_j(u)) J\hat{\Psi}_j(u) du \quad (5.6)$$

and $\hat{U}_M^{(j)} = q_j^{-1} \cdot (\hat{U}_M)_{|\hat{\Psi}_j(\hat{T}_j)}$, the measure \hat{U}_M conditioned on being in $\hat{\Psi}_j(\hat{T}_j)$. Each q_j can be estimated by a Monte Carlo method, for instance by sampling uniform points on $\mathcal{B}_{\hat{T}_j}(0, \varepsilon)$. As the integrand in (5.6) is bounded by 2 (for ε small enough), the variance of the estimator of the integral can be bounded, and δ^{-2} samples are necessary to obtain a precision δ . The volume $|\widehat{\text{vol}}_M|$ can then be approximated by the sum of the q_j s. A N -sample of law \hat{U}_M is built in two steps. First, by sampling (N_1, \dots, N_J) that follows a multinomial distribution of parameters n and (q_1, \dots, q_J) . Second, by sampling N_j points from $\hat{U}_M^{(j)}$ for every j . We propose the following method to produce a sample with approximate distribution $\hat{U}_M^{(j)}$. Let $\tilde{U}_M^{(j)}$ be the pushforward measure of the uniform distribution on $\mathcal{B}_{\hat{T}_j}(0, \varepsilon)$ by $\hat{\Psi}_j$. It is immediate to simulate from $\tilde{U}_M^{(j)}$, while $\hat{U}_M^{(j)} \ll \tilde{U}_M^{(j)}$, with density $\frac{d\hat{U}_M^{(j)}}{d\tilde{U}_M^{(j)}}$ being equal to, for $u \in \hat{T}_j$,

$$\frac{d\hat{U}_M^{(j)}}{d\tilde{U}_M^{(j)}}(\hat{\Psi}_j(u)) = \frac{\omega_d \varepsilon^d}{q_j} \chi_j(\hat{\Psi}_j(u)) J\hat{\Psi}_j(u). \quad (5.7)$$

Rejection sampling with proposal distribution $\tilde{U}_M^{(j)}$ then allows one to create a sample from law $\hat{U}_M^{(j)}$. One can check that the density in (5.7) is upper bounded by some constant depending only on d (for ε small enough), so that the acceptance ratio of the procedure is bounded away from zero.

Another quantity of interest is the number N of samples that are needed to obtain a good approximation of \hat{U}_M based on the empirical distribution $(\hat{U}_M)_N$ of a N -sample from law \hat{U}_M . It

is known [TGHS20] that if a measure is supported on a d -dimensional manifold, then the W_∞ -distance between a measure and a N -sample is of order $(\log N/N)^{1/d}$ (for $d \geq 3$). However, \hat{U}_M is not supported on a manifold, but on a union of overlapping polynomial patches. We however show that the expected rate of convergence still holds in this case. A proof is provided in Appendix F.

Proposition 5.2. *Let \hat{U}_M be the estimator of the uniform measure built on n points sampled from $\mu \in \mathcal{Q}_d^{0,k}$. Let W_1, \dots, W_N be a N -sample of law \hat{U}_M , with associated empirical measure $(\hat{U}_M)_N$. Then,*

$$\mathbb{E}[W_\infty((\hat{U}_M)_N, U_M)] \lesssim \left(\frac{\log n}{n}\right)^{k/d} + \begin{cases} \frac{(\log N)^{3/4}}{N^{1/2}} & \text{if } d = 2, \\ \left(\frac{\log N}{N}\right)^{1/d} & \text{if } d \geq 3. \end{cases} \quad (5.8)$$

Remark 5.3 (Spectral estimation of the Laplace-Beltrami operator on M). García Trillos & al. [TGHS20] study the problem of estimating the spectral properties of the Laplace-Beltrami operator Δ on M . They show that the eigenvalues of a properly tuned graph Laplacian built on top of a uniform sample of points W_1, \dots, W_n on M will converge at rate $(\log n/n)^{1/(2d)}$ towards the eigenvalues of Δ . We may build upon their results using the estimated volume measure $\widehat{\text{vol}}_M$. Let W_1, \dots, W_N be a N -sample of law \hat{U}_M . Then, Theorem 4 in [TGHS20] together with Proposition 5.2 yield that the eigenvalues of an appropriate graph Laplacian built on top of the sample W_1, \dots, W_N will approximate the eigenvalues of Δ at rate (for $d \geq 3$)

$$\sqrt{\left(\frac{\log n}{n}\right)^{k/d} + \left(\frac{\log N}{N}\right)^{1/d}}.$$

In particular, the faster rate of convergence $(\log n/n)^{k/(2d)}$ is attained by such an estimator, at the price of being able to sample $N = n^k$ points according to \hat{U}_M . This tradeoff between statistical accuracy and computational efficiency is a common phenomenon, discussed for instance in [WB19b, Section 6] in the setting of Wasserstein density estimation. Another advantage of our procedure is that, unlike results from [TGHS20], it allows one to recover the eigenvalues of the Laplace-Beltrami operator Δ even in the case where one has access to a sample X_1, \dots, X_n of non-uniform points on M .

We now turn to the problem of sampling from $\hat{\nu}_{n,h}$. Computing the density of $K_h * (\nu_n/\hat{\rho}_h)$ of $\hat{\nu}_{n,h}$ requires first the computation of

$$\hat{\rho}_h(X_i) = \sum_{j=1}^J \int_{\hat{T}_j} K_h(\hat{\Psi}_j(u) - X_i) \chi_j(\hat{\Psi}_j(u)) J \hat{\Psi}_j(u) du \quad (5.9)$$

for every point X_i of the sample. Note that there are only a small number of non-zero terms in this sum. Still, the previous basic Monte Carlo method using uniform samples will not perform well in this case, as the integrand in (5.9) has ∞ -norm of order $h^{-d} \gg 1$. A slight modification of the estimator $\hat{\nu}_{n,h}$ is however possible: consider the measure $\tilde{\nu}_{n,h}$ with density proportional to $K_h * \nu_n$ with respect to $\widehat{\text{vol}}_M$. Then, Lemma 4.9 yields that $\tilde{\nu}_{n,h}$ is at distance ε^k from the measure with density $cK_h * \nu_n$ with respect to vol_M , where $c = \frac{1}{n} \sum_{i=1}^n \int K_h(x - X_i) d\text{vol}_M(x)$ is a normalizing constant. Furthermore, Lemma 4.2 implies that both c and $\hat{\rho}_h(X_i)$ deviates from 1 with an error at most of order h^{k-1} . This implies, from the linearization inequality (2.6) that the distance $W_p(\tilde{\nu}_{n,h}, \hat{\nu}_{n,h})$ is also of order h^{k-1} . In particular, the risk of the estimator $\tilde{\nu}_{n,h}$ is the same as long as $s \leq k-2$ (with a deterioration of the rate for $k-2 \leq s \leq k-1$), while sampling from $\tilde{\nu}_{n,h}$ does not require computing any integrals. Indeed, the measure $\tilde{\nu}_{n,h}$ has a density α with respect to the d -dimensional Hausdorff measure on $\bigcup_{j=1}^J \hat{\Psi}_j(\hat{T}_j)$ equal to, at the point x ,

$$\alpha(x) \propto \sum_{j=1}^J \mathbf{1}\{x \in \hat{\Psi}_j(\hat{T}_j)\} \frac{1}{n} \sum_{i=1}^n K_h(x - X_i) \chi_j(x). \quad (5.10)$$

A Metropolis-Hasting scheme with proposal distribution $\sum_{j=1}^J q_j \tilde{U}_M^{(j)}$ is then implementable. Furthermore, adapting the proof of Theorem 3.1(i), one can check that $K_h * \nu_n(x)$ is upper bounded

by (say) $3f_{\max}$ and lower bounded by $f_{\min}/3$ for x in the support of $\widehat{\text{vol}}_M$. This implies that the acceptance ratio of the Metropolis-Hasting scheme is lower bounded by a positive constant depending only on f_{\min} , f_{\max} and d (for ε small enough). The behavior in practice of those different estimators remains to be investigated and is left for future work.

Acknowledgements. I am grateful to E. Aamari, C. Berenfeld, F. Chazal, C. Levrard and P. Massart for helpful discussions and valuable comments on different mathematical aspects of this work.

APPENDIX A. GEOMETRIC PROPERTIES OF C^k MANIFOLDS WITH POSITIVE REACH AND THEIR ESTIMATORS

Let $M \in \mathcal{M}_{d,\tau_{\min},L}^k$ for some $k \geq 2$ and $\tau_{\min}, L > 0$. Recall that the angle between two d -dimensional subspaces T_1 and T_2 is given by $\angle(T_1, T_2) := \|\pi_{T_1} - \pi_{T_2}\|_{\text{op}} = \|\pi_{T_1}^\perp \circ \pi_{T_2}\|_{\text{op}}$, where π_{T_1} (resp. π_{T_2}) is the orthogonal projection on T_1 (resp. T_2) and $\pi_{T_1}^\perp := \text{id} - \pi_{T_1}$.

Lemma A.1. *Let $x, y \in M$. The following properties hold:*

- (i) *One has $|\pi_y^\perp(x - y)| \leq \frac{|x - y|^2}{2\tau_{\min}}$ and $\angle(T_x M, T_y M) \leq 2\frac{|y - x|}{\tau_{\min}}$.*
- (ii) *If $\pi_M(z) = x$ for some $z \in M^{\tau_{\min}}$, then $z - x \in T_x M^\perp$.*
- (iii) *If $h \leq \tau_{\min}/4$, then $c_d h^d \leq \text{vol}_M(\mathcal{B}_M(x, h)) \leq C_d h^d$.*
- (iv) *If $h \leq r_0$, then $\mathcal{B}_M(x, h) \subset \Psi_x(\mathcal{B}_{T_x M}(0, h)) \subset \mathcal{B}_M(x, 8h/7)$. Also, if $u \in \mathcal{B}_{T_x M}(0, r_0)$, then $|u| \leq |\Psi_x(u) - x| \leq 8|u|/7$.*
- (v) *There exists a map $N_x : \mathcal{B}_{T_x M}(0, r_0) \rightarrow T_x M^\perp$ satisfying $dN_x(0) = 0$, and such that, for $u \in \mathcal{B}_{T_x M}(0, r_0)$, we have $\Psi_x(u) = x + u + N_x(u)$ with $|N_x(u)| \leq L|u|^2$.*
- (vi) *There exist tensors B_x^1, \dots, B_x^{k-1} of operator norm controlled by a constant depending on L, d, k and τ_{\min} , such that, if $u \in T_x M$ satisfies $|u| \leq C_{k,d,L}$, then $J\Psi_x(u) = 1 + \sum_{i=2}^{k-1} B_x^i[u^{\otimes i}] + R_x(u)$, with $|R_x(u)| \leq C'_{k,d,L}|u|^k$.*

Proof. See Theorem 4.18 in [Fed59] and Lemma 6 in [GW03] for (i), Theorem 4.8 in [Fed59] for (ii), and Proposition 8.7 in [AL18] for (iii). See Lemma A.2 in [AL19] for the second inclusion of balls in (iv), which also implies the second inequality in (iv). The first inclusion as well as the first inequality in (iv) follow from the fact that Ψ_x is the local inverse of $\tilde{\pi}_x$, which is 1-Lipschitz.

By a Taylor expansion of Ψ_x at $u = 0$, we have $\Psi_x(u) = x + u + N_x(u)$, with $N_x(u) = \int_0^1 d^2\Psi_x(tu)[u^{\otimes 2}]dt$. Hence, $|N_x(u)| \leq L|u|^2$. Furthermore, as $\tilde{\pi}_x \circ \Psi_x(u) = u$, we have $\pi_x(N_x(u)) = 0$, i.e. N_x takes its values in $T_x M^\perp$. This proves (v).

Eventually, we prove (vi). Let us write $d\Psi_x(u) = \text{id}_{T_x M} + dN_x(u)$ and $d\Psi_x(u)^\top d\Psi_x(u) = \text{id}_{T_x M} + (dN_x(u))^\top dN_x(u)$. We obtain

$$J\Psi_x(u) = \sqrt{\det(d\Psi_x(u)^\top d\Psi_x(u))} = \sqrt{\det(\text{id}_{T_x M} + (dN_x(u))^\top dN_x(u))}.$$

One has $dN_x(u) = dN_x(0) + \sum_{j=2}^{k-1} \frac{d^j N_x(0)}{(j-1)!} [u^{\otimes(j-1)}] + R_x(u)$, with $|R_x(u)| \leq C_{k,L}|u|^{k-1}$ and $dN_x(0) = 0$. Hence, $(dN_x(u))^\top dN_x(u)$ is written as $\sum_{j=2}^{k-1} B_j[u^{\otimes j}] + R'_x(u)$ with $|R'_x(u)| \leq C'_{k,L}|u|^k$, for some j -tensors B_j whose operator norms are bounded in terms of L . The operator norm of this operator is smaller than, say, $1/2$ for $|u|$ sufficiently small, and we conclude the proof by writing a Taylor expansion at 0 of the function $F \mapsto \sqrt{\det(\text{id} + F)}$. \square

We now prove Lemma 3.6, on the construction of smooth partitions of unity based on some set S which is sufficiently sparse and dense over a tubular neighborhood of M .

Proof of Lemma 3.6. Consider the functions θ and $(\chi_x)_{x \in S}$ as in the statement of the lemma, and, for $y \in M^\delta$, let $Z(y) = \sum_{x' \in S} \theta\left(\frac{y - x'}{8\delta}\right)$. As $d_H(M^\delta | S) \leq 4\delta$, we have $Z(y) \geq 1$ and the quantity $\chi_x(y)$ is well defined. The function χ_x is smooth and we have $\sum_{x \in S} \chi_x \equiv 1$ on M^δ . One has $d^l \chi_x(y)$ which is written as a sum of terms of the form $d^{l-j} \theta\left(\frac{y-x}{8\delta}\right) d^j(Z^{-1})(y)$, and $d^j(Z^{-1})(y)$ is equal to a sum of terms of the form $Z^{j'-j-2}(y) d^{j'} Z(y)$ for $1 \leq j' \leq j$. Also,

$\left\|d^j \theta \left(\frac{y-x'}{8\delta}\right)\right\|_{\text{op}} \leq C_j \delta^{-j}$ and $\|d^j Z(y)\|_{\text{op}} \leq C_j \delta^{-j} \sum_{x \in S} \mathbf{1}\{|x-y| \leq 8\delta\}$. Hence, as $Z \geq 1$, we have for any $l \geq 0$

$$\|d^l \chi_x(y)\|_{\text{op}} \leq C'_l \delta^{-l} \sum_{x \in S} \mathbf{1}\{|x-y| \leq 8\delta\}.$$

It remains to bound this sum. If $x \in \mathcal{B}(y, 8\delta)$, then $\pi_M(x) \in \mathcal{B}(\pi_M(y), 10\delta)$. Also, for $x \neq x' \in S$, we have $|\pi_M(x) - \pi_M(x')| \geq |x - x'| - 2\delta \geq 2\delta$. In particular, the balls $\mathcal{B}_M(\pi_M(x), \delta)$ for $x \in S$ are pairwise disjoint, and are all included in $\mathcal{B}_M(\pi_M(y), 11\delta)$. Therefore, if $11\delta \leq \tau(M)/4$, using Lemma A.1(iii) twice, we obtain that $\text{vol}_M(\mathcal{B}_M(\pi_M(x), \delta)) \geq c_d \delta^d$, and that

$$\begin{aligned} \sum_{x \in S} \mathbf{1}\{|x-y| \leq 8\delta\} &\leq \sum_{x \in S} \mathbf{1}\{|x-y| \leq 8\delta\} \frac{\text{vol}_M(\mathcal{B}_M(\pi_M(x), \delta))}{c_d \delta^d} \\ &\leq \frac{\text{vol}_M(\mathcal{B}_M(\pi_M(y), 11\delta))}{c_d \delta^d} \leq c'_d. \end{aligned}$$

This concludes the proof. \square

We end this section by detailing the properties of the local polynomial estimators $\hat{\Psi}_i$ and \hat{T}_i defined in [AL19]. In particular, the next lemma implies Proposition 3.5. Recall that $X_i = Y_i + Z_i$ with $Y_i \in M$ and $|Z_i| \leq \gamma$. Aamari and Levrard introduce tensors $V_{j,i}^*$ which are defined as $d^j \Psi_{X_i}(0)/j!$, where $d^j \Psi_{X_i}(0)$ is the j th differential of Ψ_{X_i} at 0 (see the proof of Lemma 2 in [AL19] for details). In particular, we have $V_{1,i}^* = \pi_{Y_i}$. Furthermore, as $\tilde{\pi}_{Y_j} \circ \Psi_{Y_j} = \text{id}$, we have $\pi_{Y_j} \circ V_{j,i}^* = 0$ for $j \geq 2$.

Lemma A.2. *With probability larger than $1 - cn^{-k/d}$, for any $1 \leq i \leq n$,*

- (i) *We have $\angle(T_{Y_i} M, \hat{T}_i) \lesssim \varepsilon^{m-1} + \gamma \varepsilon^{-1}$.*
- (ii) *For $v \in \hat{T}_i$, we have $\hat{\Psi}_i(v) = X_i + v + \hat{N}_i(v)$, where $\hat{N}_i : \hat{T}_i \rightarrow \hat{T}_i^\perp$ is defined by $\hat{N}_i(v) = \sum_{j=2}^{m-1} \hat{V}_{j,i}[v^{\otimes j}]$.*
- (iii) *For any $2 \leq j < m$, $\|\hat{V}_{j,i} \circ \hat{\pi}_i - V_{j,i}^* \circ \pi_{Y_i}\|_{\text{op}} \lesssim \varepsilon^{m-j} + \gamma \varepsilon^{-j}$.*
- (iv) *For $v \in \mathcal{B}_{\hat{T}_i}(0, 3\varepsilon)$, we have*

$$|\hat{\Psi}_i(v) - \Psi_{Y_i}(\pi_{Y_i}(v))| \lesssim \varepsilon^m + \gamma, \quad (\text{A.1})$$

$$|\hat{N}_i(v) - N_{Y_i}(\pi_{Y_i}(v))| \lesssim \varepsilon^m + \gamma, \quad (\text{A.2})$$

$$\|d\hat{\Psi}_i(v) - d(\Psi_{Y_i} \circ \pi_{Y_i})(v)\|_{\text{op}} \lesssim \varepsilon^{m-1} + \gamma \varepsilon^{-1} \quad (\text{A.3})$$

$$\|d\hat{N}_i(v) - d(N_{Y_i} \circ \pi_{Y_i})(v)\|_{\text{op}} \lesssim \varepsilon^{m-1} + \gamma \varepsilon^{-1}. \quad (\text{A.4})$$

Proof. Lemma A.2(i) is stated in Theorem 2 in [AL19]. Remark that for $x \in \mathcal{B}(X_i, \varepsilon)$, with $\tilde{x} = x - X_i$,

$$\left| \tilde{x} - \pi(\tilde{x}) - \sum_{j=2}^{m-1} V_j[\pi(\tilde{x})^{\otimes j}] \right|^2 = \left| \tilde{x} - \pi(\tilde{x}) - \sum_{j=2}^{m-1} \pi^\perp \circ V_j[\pi(\tilde{x})^{\otimes j}] \right|^2 + \left| \sum_{j=2}^{m-1} \pi \circ V_j[\pi(\tilde{x})^{\otimes j}] \right|^2$$

so that we may always assume that the tensors $\hat{V}_{j,i}$ minimizing the criterion (3.8) satisfy $\hat{\pi}_i \circ \hat{V}_{j,i} = 0$ for $j \geq 2$. This proves Lemma A.2(ii).

We prove Lemma A.2(iii) by induction on $2 \leq j < m$. The result for $j = 2$ is stated in [AL19, Theorem 2]. It is shown in [AL19] (see Equation (3)) that there exist tensors $V'_{j,i}$ for $1 \leq j < m$ satisfying with probability larger than $1 - cn^{-k/d}$,

$$\|V'_{j,i} \circ \pi_{Y_i}\|_{\text{op}} \lesssim \varepsilon^{m-j} + \gamma \varepsilon^{-j}. \quad (\text{A.5})$$

The tensors $V'_{j,i}$ are defined by the relation, for $y \in M$ close enough to Y_i ,

$$y - Y_i - \hat{\pi}_i(y - Y_i) - \sum_{j=2}^{m-1} \hat{V}_{j,i}[\hat{\pi}_i(y - Y_i)^{\otimes j}] = \sum_{j=1}^{m-1} V'_{j,i}[\pi_{Y_i}(y - Y_i)^{\otimes j}] + R(y - Y_i), \quad (\text{A.6})$$

with $|R(y - Y_i)| \lesssim \varepsilon^m$, see the proof of Lemma 3 in [AL19]. We also may write

$$y - Y_i = \pi_{Y_i}(y - Y_i) + \sum_{j=2}^{m-1} V_{j,i}^* [\pi_{Y_i}(y - Y_i)^{\otimes j}] + R'(y - Y_i), \quad (\text{A.7})$$

with $|R'(y - Y_i)| \lesssim \varepsilon^m$. By plugging (A.7) in the left hand side of (A.6) and by noting that $\pi_{Y_i} \circ V_{j,i}^* = 0$ for $j \geq 2$, we see that $V_{j,i}' \circ \pi_{Y_i}$ is written as the sum of $(\pi_{Y_i} - \hat{\pi}_i) \circ V_{j,i}^* + (V_{j,i}^* \circ \pi_{Y_i} - \hat{V}_{j,i} \circ \hat{\pi}_i)$ and of a sum of terms proportional to

$$\hat{V}_{j',i} [\hat{\pi}_i \circ V_{a_1,i}^* \circ \pi_{Y_i}, \dots, \hat{\pi}_i \circ V_{a_{j'},i}^* \circ \pi_{Y_i}], \quad (\text{A.8})$$

where $2 \leq j' < j$ and $a_1 + \dots + a_{j'} = j$, $1 \leq a_1, \dots, a_{j'} < j$. There exists in particular an index in the sum which is larger than 2. Assume without loss of generality that $a_1, \dots, a_l > 1$ and $a_{l+1}, \dots, a_{j'} = 1$, so that $\hat{\pi}_i \circ \hat{V}_{a_u,i} = 0$ for $1 \leq u \leq l$. Then,

$$\begin{aligned} & \left\| \hat{V}_{j',i} [\hat{\pi}_i \circ V_{a_1,i}^* \circ \pi_{Y_i}, \dots, \hat{\pi}_i \circ V_{a_l,i}^* \circ \pi_{Y_i}, \dots, \hat{\pi}_i \circ V_{a_{j'},i}^* \circ \pi_{Y_i}] \right\|_{\text{op}} \\ &= \left\| \hat{V}_{j',i} [\hat{\pi}_i \circ (V_{a_1,i}^* - \hat{V}_{a_1,i}) \circ \pi_{Y_i}, \dots, \hat{\pi}_i \circ (V_{a_l,i}^* - \hat{V}_{a_l,i}) \circ \pi_{Y_i}, \right. \\ & \quad \left. \hat{\pi}_i \circ V_{a_{j'},i}^* \circ \pi_{Y_i}, \dots, \hat{\pi}_i \circ V_{a_{l+1},i}^* \circ \pi_{Y_i}] \right\|_{\text{op}} \\ &\lesssim \ell \prod_{u=1}^l \left\| V_{a_u,i}^* \circ \pi_{Y_i} - \hat{V}_{a_u,i} \circ \pi_{Y_i} \right\|_{\text{op}} \\ &\lesssim \ell \prod_{u=1}^l \left(\left\| V_{a_u,i}^* \circ \pi_{Y_i} - \hat{V}_{a_u,i} \circ \hat{\pi}_i \right\|_{\text{op}} + \ell \|\pi_{Y_i} - \hat{\pi}_i\|_{\text{op}} \right) \\ &\lesssim \varepsilon^{-1} \prod_{u=1}^l (\varepsilon^{m-a_u} + \gamma \varepsilon^{-a_u} + \varepsilon^{m-2} + \gamma \varepsilon^{-2}) \\ &\lesssim \varepsilon^{-1} (\varepsilon^{lm-(j-l)} + \gamma^l \varepsilon^{-(j-l)}) \lesssim \varepsilon^{m-j} + \gamma \varepsilon^{-j}, \end{aligned}$$

where at the last line we use the induction hypothesis as well as Lemma A.2(i), the fact that $\sum_{u=1}^l a_u = j - l$ and that $\ell \lesssim \varepsilon^{-1}$. As $\|(\pi_{Y_i} - \hat{\pi}_i) \circ V_{j,i}^*\|_{\text{op}} \lesssim \varepsilon^{m-1} + \gamma \varepsilon^{-1}$, we obtain that

$$\left\| (V_{j,i}^* \circ \pi_{Y_i} - \hat{V}_{j,i} \circ \hat{\pi}_i) - V_{j,i}' \circ \pi_{Y_i} \right\|_{\text{op}} \lesssim \varepsilon^{m-j} + \gamma \varepsilon^{-j}.$$

Hence, using (A.5),

$$\begin{aligned} \left\| V_{j,i}^* \circ \pi_{Y_i} - \hat{V}_{j,i} \circ \hat{\pi}_i \right\|_{\text{op}} &\leq \left\| (V_{j,i}^* \circ \pi_{Y_i} - \hat{V}_{j,i} \circ \hat{\pi}_i) - V_{j,i}' \circ \pi_{Y_i} \right\|_{\text{op}} + \left\| V_{j,i}' \circ \pi_{Y_i} \right\|_{\text{op}} \\ &\lesssim \varepsilon^{m-j} + \gamma \varepsilon^{-j}. \end{aligned}$$

We now may prove (A.1). Indeed, for $v \in \mathcal{B}_{\hat{T}_i}(0, 3\varepsilon)$, $\hat{\Psi}_i(v) = X_i + v + \sum_{j=2}^{m-1} \hat{V}_{j,i}[v^{\otimes j}]$, whereas by a Taylor expansion, $\Psi_{Y_i} \circ \pi_{Y_i}(v) = Y_i + \pi_{Y_i}(v) + \sum_{j=2}^{m-1} V_{j,i}[\pi_{Y_i}(v)^{\otimes j}] + R(v)$, with $|R(v)| \lesssim \varepsilon^m$. By Lemma A.2(iii), the difference between the two quantities is bounded with high probability by a sum of terms of order $(\varepsilon^{m-j} + \gamma \varepsilon^{-j})|v|^j \lesssim \varepsilon^m + \gamma$. Inequality (A.2) is directly implied by (A.1) and Lemma A.2(i). Inequality (A.3) is proven as (A.1), by noting that, for $h \in \hat{T}_i$,

$$\begin{cases} d(\Psi_{Y_j} \circ \pi_{Y_j})(v)[h] = \pi_{Y_j}(h) + \sum_{j=2}^{m-1} j V_{j,i}^* [\pi_{Y_j}(v), \pi_{Y_j}(h)^{\otimes(j-1)}] + R'(v)h \\ d\hat{\Psi}_j(v)[h] = h + \sum_{j=2}^{m-1} j \hat{V}_{j,i}[v, h^{\otimes(j-1)}], \end{cases}$$

with $\|R'(v)\|_{\text{op}} \lesssim \varepsilon^{m-1}$. Equation (A.4) is shown in a similar way. \square




FIGURE 2. Illustration of the construction in the proof of Lemma B.1

APPENDIX B. PROPERTIES OF NEGATIVE SOBOLEV NORMS

Proof of Proposition 2.5. The second inequality in (i) is trivial. The assertion (ii) is stated in [BCS10, Theorem 2.1] for an open set $\Omega \subset \mathbb{R}^d$, and their proof can be straightforwardly adapted to the manifold setting. It remains to prove the first inequality in (i). Note that for any g with $\|\nabla g\|_{L_{p^*}(M)} \leq 1$, one has $\int fg d\text{vol}_M = \int f(g - \int g d\text{vol}_M) d\text{vol}_M$ as $\int f d\text{vol}_M = 0$. Also, by Poincaré inequality (see [BCH18, Theorem 0.6]),

$$\left\| g - \int_M g \right\|_{L_{p^*}(M)} \leq C^{\frac{1}{p}} R^{\frac{d}{p^*} + \frac{1}{p}} \|\nabla g\|_{L_{p^*}(M)} \leq C^{\frac{1}{p}} R^{\frac{d}{p^*} + \frac{1}{p}},$$

where $R = \max\{d_g(x, y), x, y \in M\}$ and C depends on d and on a lower bound κ on the Ricci curvature of M . Therefore, $\|g - \int_M g\|_{H_{p^*}^1(M)} \leq C^{\frac{1}{p}} R^{\frac{d}{p^*} + \frac{1}{p}}$. The quantity κ can be further lower bounded by a constant depending on τ_{\min} and d . Indeed, a bound on the second fundamental form of M entails a bound on the Ricci curvature according to Gauss equation (see e.g. [dC92, Chapter 6]), and the second fundamental form is controlled by the reach of M , see [NSW08, Proposition 6.1]. As $C^{\frac{1}{p}} \leq C \vee 1$, to conclude, it suffices to bound the geodesic diameter of M . This is done in the following lemma. \square

Lemma B.1. *The geodesic diameter of M satisfies $\sup_{x, y \in M} d_g(x, y) \leq c_d |\text{vol}_M| \tau_{\min}^{1-d}$.*

Proof. Consider a covering of M by N open balls of radius $r_1 = \tau_{\min}/4$ (for the Euclidean distance) and let $x, y \in M$. Such a covering exists with $N \leq c_d |\text{vol}_M| r_1^{-d}$ by standard packing arguments. Let $\gamma : [0, \ell] \rightarrow M$ be a unit speed curve between x and y . Let B_0 be the ball of the covering such that $x \in B_0$. If $y \in B_0$, then $|x - y| \leq 2r_1$, and by [NSW08, Proposition 6.3], we have $d_g(x, y) \leq 4r_1$. Otherwise, let $t_0 = \inf\{t \in [0, \ell], \forall t' \geq t, \gamma(t') \notin B_0\}$. Then $x_1 := \gamma(t_0)$ belong to the boundary of B_0 , and is also in some other ball B_1 . By the previous argument, we have $d_g(x, x_1) \leq 4r_1$. If $y \in B_1$, then $d_g(x_1, y) \leq 4r_1$ and $d_g(x, y) \leq 8r_1$. Otherwise, we define $t_1 = \inf\{t \in [t_0, \ell], \forall t' \geq t, \gamma(t') \notin B_1\}$ and we iterate the same argument. At the end, we obtain a sequence $x = x_0, x_1, \dots, x_I$ of points in M with associated balls B_i which contain x_i , such that $y \in B_I$ and $d_g(x_i, x_{i+1}) \leq 4r_1$. Furthermore, all the balls B_i are pairwise distinct. As $d_g(x_I, y) \leq 4r_1$, we have $\ell \leq (I + 1)4r_1 \leq (N + 1)4r_1 \leq 8Nr_1$. By letting γ be a geodesic, we obtain in particular $\ell = d_g(x, y) \leq 8Nr_1 \leq 8c_d |\text{vol}_M| r_1^{1-d}$. \square

Proof of Proposition 2.8. Given a measurable map $\rho : [0, 1] \rightarrow \mathcal{P}^p$, E_t a vectorial measure absolutely continuous with respect to ρ_t (see [San15, Box 4.2]) and $v(x, t)$ a time-depending vector field, defined as the density of E_t with respect to ρ_t , we define the Benamou-Brenier functional

$$\mathcal{B}_p(\rho, E) := \int |v(x, t)|^p d\rho_t(x) dt. \quad (\text{B.1})$$

The Benamou-Brenier formula [BB00, Bre03] asserts that for $\mu, \nu \in \mathcal{P}_1^p$ supported on some ball of radius R ,

$$W_p^p(\mu, \nu) = \min \{ \mathcal{B}_p(\rho, E), \partial_t \rho_t + \nabla \cdot E_t = 0, \rho_0 = \mu, \rho_1 = \nu \}, \quad (\text{B.2})$$

where ρ_t is supported on the ball of radius R , and the continuity equation $\partial_t \rho + \nabla \cdot E = \mu - \nu$ has to be understood in the distributional sense, i.e.

$$\int_{[0,1] \times \mathbb{R}^D} \partial_t \phi(t, x) d\rho(t, x) + \int_{[0,1] \times \mathbb{R}^D} \nabla \phi(t, x) \cdot dE(t, x) = 0, \quad (\text{B.3})$$

for all $\phi \in \mathcal{C}^1((0, 1) \times \mathcal{B}(0, R))$ with compact support.

Assume that μ has a density f_0 and ν has a density f_1 on M . As $\tau(M) > 0$, the existence of a probability measure of mass 1, supported on M , with density larger than f_{\min} implies that M is compact, see Remark 2.10. It is in particular included in a ball $\mathcal{B}(0, R)$ for some R large enough. Let w be a vector field on M with $\nabla \cdot w = \mu - \nu$ in a distributional sense, i.e. $\int \nabla g \cdot w = -\int g(\mu - \nu)$ for all $g \in \mathcal{C}^1(M)$. Let $\rho_t = (1-t)\mu + t\nu$ and define E the vector measure having density w with respect to $\text{Leb}_1 \times \text{vol}_M$, where Leb_1 is the Lebesgue measure on $[0, 1]$. Then (ρ, E) satisfies the continuity equation and $E = v \cdot \rho$ where $v(t, x) = \frac{w(x)}{(1-t)f_0(x) + tf_1(x)}$ for $t \in [0, 1]$, $x \in M$. Hence,

$$\begin{aligned} W_p^p(\mu, \nu) &\leq \int_0^1 \int \frac{1}{p} |v|^p d\rho \\ &= \frac{1}{p} \int_0^1 \int \frac{|w(x)|^p}{|(1-t)f_0(x) + tf_1(x)|^p} ((1-t)f_0(x) + tf_1(x)) dx dt \\ &\leq \frac{1}{p} \int |w(x)|^p dx \frac{1}{f_{\min}^{p-1}}. \end{aligned}$$

By taking the infimum on vector fields w on M satisfying $\nabla \cdot w = \mu - \nu$ and using Proposition 2.5, we obtain the conclusion. The second inequality in (2.6) follows from Proposition 2.5. \square

APPENDIX C. PROOFS OF SECTION 4.1

Proof of Lemma 4.2. We first prove (4.4). Note that if $|x-y| \geq h$ for $x, y \in M$, then $K_h(x-y) = 0$. Hence, by a change of variable, using that $\mathcal{B}_M(x, h) \subset \Psi_x(\mathcal{B}_{T_x M}(0, h))$ according to Lemma A.1(iv),

$$\begin{aligned} \int_M K_h(x-y) B[(x-y)^{\otimes j}] dy &= \int_{\mathcal{B}_{T_x M}(0, h)} K_h(x - \Psi_x(v)) B[(x - \Psi_x(v))^{\otimes j}] J\Psi_x(v) dv \\ &= \int_{\mathcal{B}_{T_x M}(0, 1)} K\left(\frac{x - \Psi_x(hv)}{h}\right) B[(x - \Psi_x(hv))^{\otimes j}] J\Psi_x(hv) dv. \end{aligned}$$

As the functions Ψ_x and K are \mathcal{C}^k , according to Lemma A.1(v) and Lemma A.1(vi), we can write by a Taylor expansion, for $v, u \in \mathcal{B}_{T_x M}(0, r_0)$,

$$\begin{cases} \Psi_x(v) = x + v + \sum_{i=2}^{k-1} \frac{d^i \Psi_x(0)}{i!} [v^{\otimes i}] + R_1(x, v) \\ J\Psi_x(v) = 1 + \sum_{i=2}^{k-1} B_x^i [v^{\otimes i}] + R_2(x, v) \\ K(v+u) = K(v) + \sum_{i=1}^{k-1} \frac{d^i K(v)}{i!} [u^{\otimes i}] + R_3(v, u) \\ B[(v+u)^{\otimes j}] = B[v^{\otimes j}] + \sum_{\emptyset \neq \sigma \subset \{1, \dots, j\}} B[v^\sigma, u^{\sigma^c}], \end{cases} \quad (\text{C.1})$$

where $|R_j(x, v)| \leq C_j |v|^k$ for $j = 1, 2$, $|R_3(v, u)| \leq C_3 |u|^k$ and (v^σ, u^{σ^c}) is the j -tuple whose l th entry is equal to v if $l \in \sigma$, u otherwise. We obtain that

$$\frac{x - \Psi_x(hv)}{h} = -v - \sum_{i=2}^{k-1} \frac{d^i \Psi_x(0)}{i!} [(hv)^{\otimes i}] h^{-1} - R_1(x, hv) h^{-1},$$

and that the expression $K\left(\frac{x - \Psi_x(hv)}{h}\right) B[(x - \Psi_x(hv))^{\otimes j}] J\Psi_x(hv)$ is written as a sum of terms of the form

$$C_{i_0, i_1, i_2} h^{-i_0} d^{i_0} K(v) [(d^{i_1} \Psi_x(0) [(hv)^{\otimes i_1}])^{\otimes i_0}] F_{i_2} [(hv)^{\otimes i_2}] \quad (\text{C.2})$$

for $0 \leq i_0 \leq k-1$, $2 \leq i_1 \leq k-1$ and $j \leq i_2 \leq k'$, where F_{i_2} is some tensor of order i_2 and k' is some integer depending on k and j , plus a remainder term smaller than $\|B\|_{\text{op}} |hv|^{k-1+j}$ up to a constant depending on k , j , L_k and K . The terms for which $i_0 i_1 + i_2 - i_0 \geq k$ are smaller than $\|B\|_{\text{op}} h^k$ up to a constant, whereas the integrals of the other the terms are null as the kernel is of order k . The first inequality in (4.5) is proven in a similar manner. Let us now bound $\|\rho_h\|_{\mathcal{C}^j(M)}$. Given $x \in M$, we have

$$d^j(\rho_h \circ \Psi_x)(0) = h^{-j} \int_{\mathcal{B}_{T_x M}(0, h)} (d^j K)_h(x - \Psi_x(v)) J\Psi_x(v) dv.$$

Therefore, using the same argument as before, we obtain that $\|d^j(\rho_h \circ \Psi_x)(0)\|_{\text{op}} \lesssim h^{k-1-j}$. \square

Proof of Lemma 4.3. Let $0 \leq l \leq k-1$ be even, $\phi \in \mathcal{C}^\infty(M)$ be supported in $\mathcal{B}_M(x_0, h_0)$ for some h_0 small enough and $g \in L_{p^*}(M)$ with $\|g\|_{L_{p^*}(M)} \leq 1$. Let $x = \Psi_{x_0}(u) \in \mathcal{B}_M(x_0, h_0)$ and let $\tilde{\phi}_{x_0} = \tilde{\phi} \circ \Psi_{x_0}$. Recall that $\tilde{\phi}_l = d^l \tilde{\phi}_{x_0} \circ \tilde{\pi}_{x_0}$. We have $K_h(x - \Psi_{x_0}(v)) \neq 0$ only if $|x - \Psi_{x_0}(v)| \leq h$. Hence, as $|x - \Psi_{x_0}(v)| \geq |u - v|$ (recall that Ψ_{x_0} is the inverse of the projection $\tilde{\pi}_{x_0}$), the function $K_h(x - \Psi_{x_0}(\cdot))$ is supported on $\mathcal{B}_{T_{x_0} M}(u, h) \subset \mathcal{B}_{T_{x_0} M}(0, r_0) =: B_0$ for h, h_0 small enough. Thus,

$$\begin{aligned} A_h \phi(x) &= \int_{\mathcal{B}_M(x, h)} K_h(x - y) (\tilde{\phi}(y) - \tilde{\phi}(x)) dy \\ &= \int_{B_0} K_h(x - \Psi_{x_0}(v)) (\tilde{\phi}_{x_0}(v) - \tilde{\phi}_{x_0}(u)) J\Psi_{x_0}(v) dv. \end{aligned}$$

We may write

$$\tilde{\phi}_{x_0}(v) - \tilde{\phi}_{x_0}(u) = \sum_{i=1}^{l-1} \frac{d^i \tilde{\phi}_{x_0}(u)}{i!} [(v-u)^{\otimes i}] + \int_0^1 d^l \tilde{\phi}_{x_0}(u + \lambda(v-u)) [(v-u)^{\otimes l}] \frac{(1-\lambda)^{l-1}}{(l-1)!} d\lambda.$$

Each term $\int_{B_0} K_h(x - \Psi_{x_0}(v)) \frac{d^i \tilde{\phi}_{x_0}(u)}{i!} [(v-u)^{\otimes i}] J\Psi_{x_0}(v) dv$ is equal to

$$\int_M K_h(x - y) \frac{d^i \tilde{\phi}_{x_0}(\tilde{\pi}_{x_0}(x))}{i!} [(\pi_{x_0}(y-x))^{\otimes i}] dy,$$

and is therefore of order smaller than $h^k \max_{1 \leq i \leq l} \|\tilde{\phi}_i(x)\|_{\text{op}}$ by Lemma 4.2. Hence, $A_h \phi(x)$ is equal to the sum of a remainder term of order $h^k \max_{1 \leq i \leq l} \|\tilde{\phi}_i(x)\|_{\text{op}}$ and of

$$\begin{aligned} &\int_0^1 \int_{B_0} K_h(x - \Psi_{x_0}(v)) d^l \tilde{\phi}_{x_0}(u + \lambda(v-u)) [(v-u)^{\otimes l}] \frac{(1-\lambda)^{l-1}}{(l-1)!} J\Psi_{x_0}(v) dv d\lambda \\ &= \int_0^1 \int_{B_0} K_h(x - \Psi_{x_0}(v)) (d^l \tilde{\phi}_{x_0}(u + \lambda(v-u)) - d^l \tilde{\phi}_{x_0}(u)) [(v-u)^{\otimes l}] \times \frac{(1-\lambda)^{l-1}}{(l-1)!} J\Psi_{x_0}(v) dv d\lambda \\ &\quad + R_1(x), \end{aligned}$$

where $|R_1(x)| \lesssim h^k \max_{1 \leq i \leq l} \|\tilde{\phi}_i(x)\|_{\text{op}}$ by Lemma 4.2. We now fix $\lambda \in (0, 1)$ and write by a change of variables

$$\begin{aligned} U(x) &:= \int_{B_0} K_h(x - \Psi_{x_0}(v)) (d^l \tilde{\phi}_{x_0}(u + \lambda(v-u)) - d^l \tilde{\phi}_{x_0}(u)) [(v-u)^{\otimes l}] J\Psi_{x_0}(v) dv \\ &= \int_{B_0} K_h\left(x - \Psi_{x_0}\left(u + \frac{w-u}{\lambda}\right)\right) (d^l \tilde{\phi}_{x_0}(w) - d^l \tilde{\phi}_{x_0}(u)) \left[\frac{(w-u)^{\otimes l}}{\lambda^l}\right] \\ &\quad \times J\Psi_{x_0}\left(u + \frac{w-u}{\lambda}\right) \frac{dw}{\lambda^d} \end{aligned}$$

Note that $|K_h(u) - K_h(v)| \lesssim h^{-d-1} |u-v| \mathbf{1}\{|u| \leq h \text{ or } |v| \leq h\}$, and that, as Ψ_{x_0} is \mathcal{C}^2 ,

$$\begin{aligned} \left| x - \Psi_{x_0}\left(u + \frac{w-u}{\lambda}\right) - \frac{x - \Psi_{x_0}(w)}{\lambda} \right| &\leq \left| \frac{d\Psi_{x_0}(u)[w-u] - (x - \Psi_{x_0}(w))}{\lambda} \right| + \frac{L_k |w-u|^2}{2\lambda^2} \\ &\leq \frac{L_k |w-u|^2}{\lambda} \lesssim \frac{|w-u|^2}{\lambda}, \end{aligned}$$

whereas, as $J\Psi_{x_0}$ is Lipschitz continuous,

$$\left| J\Psi_{x_0} \left(u + \frac{w-u}{\lambda} \right) - J\Psi_{x_0}(w) \right| \lesssim \left| u + \frac{w-u}{\lambda} - w \right| \lesssim \frac{|w-u|}{\lambda}.$$

Hence, $U(x)$ is equal to the sum of

$$\begin{aligned} & \lambda^{-l} \int_{B_0} K_{h\lambda}(x - \Psi_{x_0}(w)) (d^l \tilde{\phi}_{x_0}(w) - d^l \tilde{\phi}_{x_0}(u)) [(w-u)^{\otimes l}] J\Psi_{x_0}(w) dw \\ &= \lambda^{-l} \int_M K_{h\lambda}(x-y) (\tilde{\phi}_l(y) - \tilde{\phi}_l(x)) [(\pi_{x_0}(y-x))^{\otimes l}] dy, \end{aligned}$$

and of a remainder term smaller than

$$\begin{aligned} & \lambda^{-l} \int_{B_0} \left| \lambda^{-d} K_h \left(x - \Psi_{x_0} \left(u + \frac{w-u}{\lambda} \right) \right) J\Psi_{x_0} \left(u + \frac{w-u}{\lambda} \right) - K_{h\lambda}(x - \Psi_{x_0}(w)) J\Psi_{x_0}(w) \right| \\ & \quad \times \|d^l \tilde{\phi}_{x_0}(w) - d^l \tilde{\phi}_{x_0}(u)\|_{\text{op}} |w-u|^l dw \\ & \lesssim \lambda^{-l} \int_{|w-u| \lesssim \lambda h} \left(\frac{|w-u|^2}{(\lambda h)^{d+1}} J\Psi_{x_0} \left(u + \frac{w-u}{\lambda} \right) + |K_{h\lambda}(x - \Psi_{x_0}(w))| \frac{|w-u|}{\lambda} \right) \\ & \quad \times \|d^l \tilde{\phi}_{x_0}(w) - d^l \tilde{\phi}_{x_0}(u)\|_{\text{op}} |w-u|^l dw \\ & \lesssim h^{l+1} (\lambda h)^{-d} \int_{|w-u| \lesssim \lambda h} \|d^l \tilde{\phi}_{x_0}(w) - d^l \tilde{\phi}_{x_0}(u)\|_{\text{op}} dw. \end{aligned}$$

Putting all the estimates together, we may now write $\int_M A_h \phi(x) g(x) dx$ as $S + R_2$, where, by the symmetrization trick (using that l is even)

$$\begin{aligned} S &= \iint_{M \times M} K_h^{(l)}(x-y) (\tilde{\phi}_l(y) - \tilde{\phi}_l(x)) [(\pi_{x_0}(y-x))^{\otimes l}] g(x) dy dx \\ &= \iint_{M \times M} K_h^{(l)}(x-y) (\tilde{\phi}_l(x) - \tilde{\phi}_l(y)) [(\pi_{x_0}(x-y))^{\otimes l}] g(y) dy dx \\ &= \frac{1}{2} \iint_{M \times M} K_h^{(l)}(x-y) (\tilde{\phi}_l(y) - \tilde{\phi}_l(x)) [(\pi_{x_0}(x-y))^{\otimes l}] (g(x) - g(y)) dy dx, \end{aligned}$$

and, as $A_h \phi$ is supported on $\mathcal{B}_M(x_0, h_0 + h) \subset \mathcal{B}_M(x, 2h_0)$ if h is small enough, R_2 is smaller up to a constant than,

$$h^{l+1} (\lambda h)^{-d} \int_{x \in \mathcal{B}_M(x_0, 2h_0)} \int_{|w - \tilde{\pi}_{x_0}(x)| \lesssim \lambda h} \|d^l \tilde{\phi}_{x_0}(w) - d^l \tilde{\phi}_{x_0}(\tilde{\pi}_{x_0}(x))\|_{\text{op}} |g(x)| dw dx \quad (\text{C.3})$$

$$\begin{aligned} & + \int_M h^k \max_{1 \leq i \leq l} \|\tilde{\phi}_i(x)\|_{\text{op}} |g(x)| dx \\ & \lesssim h^{l+1} (\lambda h)^{-d} \int_{w \in \mathcal{B}_M(x, 3h_0)} \|d^l \tilde{\phi}_{x_0}(w)\|_{\text{op}} \int_{|w - \tilde{\pi}_{x_0}(x)| \lesssim \lambda h} |g(x)| dx dw \quad (\text{C.4}) \\ & + h^{l+1} \int_{x \in \mathcal{B}_M(x, 2h_0)} \|\tilde{\phi}_l(x)\|_{\text{op}} |g(x)| dx + \int_M h^k \max_{1 \leq i \leq l} \|\tilde{\phi}_i(x)\|_{\text{op}} |g(x)| dx, \end{aligned}$$

where we also used Lemma A.1(iii). By the chain rule,

$$\begin{aligned} \max_{1 \leq i \leq l} \|\tilde{\phi}_i(x)\|_{\text{op}} &= \max_{1 \leq i \leq l} \|d^i(\tilde{\phi} \circ \Psi_{x_0}) \circ \tilde{\pi}_{x_0}(x)\|_{\text{op}} \\ &= \max_{1 \leq i \leq l} \|d^i(\tilde{\phi} \circ \Psi_x \circ \tilde{\pi}_x \circ \Psi_{x_0}) \circ \tilde{\pi}_{x_0}(x)\|_{\text{op}} \\ &\lesssim \max_{1 \leq i \leq l} \|d^i(\tilde{\phi} \circ \Psi_x)(\tilde{\pi}_x \circ \Psi_{x_0} \circ \tilde{\pi}_{x_0}(x))\|_{\text{op}} \\ &\lesssim \max_{1 \leq i \leq l} \|d^i(\tilde{\phi} \circ \Psi_x)(0)\|_{\text{op}} = \|d^l \tilde{\phi}(x)\|_{\text{op}}. \end{aligned} \quad (\text{C.5})$$

Hence, applying Hölder's inequality and using that $\|g\|_{L_{p^*}(M)} \leq 1$ show that the two last terms in (C.4) are of order $h^{l+1} \|\tilde{\phi}\|_{H_p^l(M)}$. To bound the first term in (C.4), remark that by Young's

inequality for integral operators [Sog17, Theorem 0.3.1], if $\mathcal{T}_{\lambda h}(g)(y) = (\lambda h)^{-d} \int_{|x-y| \lesssim \lambda h} |g(x)| dx$, then $\|\mathcal{T}_{\lambda h} g\|_{L_{p^*}(M)} \lesssim \|g\|_{L_p(M)}$. This yields, by Hölder's inequality,

$$h^{l+1} \int_{w \in \mathcal{B}_M(x, 3h_0)} \|d^l \tilde{\phi}_{x_0}(w)\|_{\text{op}} \mathcal{T}_{h\lambda}(g)(\Psi_{x_0}(w)) dw \lesssim h^{l+1} \|\tilde{\phi}\|_{H_p^l(M)},$$

which concludes the proof of the first statement of Lemma 4.3. To bound the remainder term in terms of $\|\tilde{\phi}\|_{H_p^{l+1}(M)}$, we bound the second term in (C.3) in the same fashion, while, to bound the first term, we write, by a change of variables,

$$\begin{aligned} & \int_{\mathcal{B}_M(x_0, 2h_0)} \int_{|w - \tilde{\pi}_{x_0}(x)| \lesssim \lambda h} \|d^l \tilde{\phi}_{x_0}(w) - d^l \tilde{\phi}_{x_0}(\tilde{\pi}_{x_0}(x))\|_{\text{op}} |g(x)| dx dw \\ & \leq \int_0^1 \int_{\mathcal{B}_M(x_0, 2h_0)} \int_{|w - \tilde{\pi}_{x_0}(x)| \lesssim \lambda h} \|d^{l+1} \tilde{\phi}_{x_0}(\tilde{\pi}_{x_0}(x) + \lambda'(w - \tilde{\pi}_{x_0}(x)))\|_{\text{op}} \\ & \quad \times |\tilde{\pi}_{x_0}(x) - w| |g(x)| dx dw d\lambda' \\ & \lesssim h \int_0^1 \int_{\mathcal{B}_M(x_0, 2h_0)} \int_{|u - \tilde{\pi}_{x_0}(x)| \lesssim \lambda' \lambda h} \|d^{l+1} \tilde{\phi}_{x_0}(u)\|_{\text{op}} |g(x)| dx \frac{du}{\lambda'^d} d\lambda', \end{aligned}$$

and this term is bounded as the first term in (C.4) by $h(h\lambda)^d \|\tilde{\phi}\|_{H_p^{l+1}(M)}$, concluding the proof of Lemma 4.3. \square

Proof of Lemma 4.4. The second inequality of Lemma 4.4 follows from the definition of the Sobolev norm and of the bound (C.5) applied to $\tilde{\phi} = \eta$. To prove the first inequality, write

$$\begin{aligned} & h^{-d} \iint_{\mathcal{B}_M(x_0, h_0)^2} \mathbf{1}\{|x - y| \leq h\} \frac{\|\eta(x) - \eta(y)\|_{\text{op}}^p}{|x - y|^p} dx dy \\ & \lesssim h^{-d} \iint_{\mathcal{B}_{T_{x_0}M}(0, h_0)^2} \mathbf{1}\{|\Psi_{x_0}(u) - \Psi_{x_0}(v)| \leq h\} \frac{\|d^l(\eta \circ \Psi_{x_0})(u) - d^l(\eta \circ \Psi_{x_0})(v)\|_{\text{op}}^p}{|\Psi_{x_0}(u) - \Psi_{x_0}(v)|^p} du dv \\ & \quad \text{as } J\Psi_{x_0}(u) \lesssim 1 \text{ for } |u| \lesssim 1 \\ & \lesssim h^{-d} \int_0^1 \iint_{\mathcal{B}_{T_{x_0}M}(0, h_0)^2} \mathbf{1}\{|u - v| \leq h\} \|d^{l+1}(\eta \circ \Psi_{x_0})(u + \lambda(v - u))\|_{\text{op}}^p du dv d\lambda \\ & \quad \text{as } |v - u| \leq |\Psi_{x_0}(u) - \Psi_{x_0}(v)| \\ & \lesssim h^{-d} \int_0^1 \iint_{\mathcal{B}_{T_{x_0}M}(0, 2h_0)^2} \mathbf{1}\{|w - u| \leq \lambda h\} \|d^{l+1}(\eta \circ \Psi_{x_0})(w)\|_{\text{op}}^p du dw \lambda^{-d} d\lambda \\ & \lesssim \int_0^1 \int_{\mathcal{B}_{T_{x_0}M}(0, 2h_0)} \|d^{l+1}(\eta \circ \Psi_{x_0})(w)\|_{\text{op}}^p dw \lesssim \int_{\mathcal{B}_M(x_0, h_0)} \|\eta_{l+1}(x)\|_{\text{op}}^p dx, \end{aligned}$$

where at the second to last line, we used that $w = u + \lambda(v - u)$ is of norm smaller than $2h_0$ if $|u| \leq h_0$ and $|v - u| \leq h \leq h_0$, and, at the last line, we used that $J\Psi_{x_0}(w) \geq 1/2$ for $|w|$ small enough. \square

APPENDIX D. PROOF OF LEMMA 4.7

Lemma 4.7 is heavily based on the following classical control on the gradient of the Green function.

Lemma D.1. *Let $x, y \in M$, then*

$$|\nabla_x G(x, y)| \lesssim \frac{1}{d_g(x, y)^{d-1}} \leq \frac{1}{|x - y|^{d-1}}. \quad (\text{D.1})$$

Proof. For $d \geq 2$, a proof of Lemma D.1 is found in [Aub82, Theorem 4.13]. See also [Hir96, Theorem 5.2] for a proof with more explicit constants in the case $d \geq 3$. Constants in their proofs depend on d , bounds on the curvature of M , $|\text{vol}_M|$ and the geodesic diameter of M . As, those three last quantities can be further bounded by constants depending on τ_{\min} , f_{\min} and d , see

Lemma B.1 and [NSW08, Proposition 6.1], this concludes the proof. For $d = 1$, M is isometric to a circle, for which a closed formula for G exists [Bur94], and satisfies $|\nabla_x G(x, y)| \leq 1$. \square

Recall that, by Lemma 4.2, $|\rho_h(x)| \geq 1/2$ for all $x \in M$. Therefore, Lemma D.1 yields

$$\left| \nabla G \left(K_h * \left(\frac{\delta_x}{\rho_h} \right) \right) (z) \right| = \left| \int_M \nabla_z G(z, y) \frac{K_h(x-y)}{\rho_h(x)} dy \right| \lesssim \int_{\mathcal{B}_M(x, h)} \frac{\|K\|_\infty h^{-d}}{|z-y|^{d-1}} dy.$$

If $d = 1$, this quantity is smaller than a constant as $\text{vol}_M(\mathcal{B}_M(x, h)) \lesssim h^d$ by Lemma A.1(iii). We then obtain directly the result in this case by integrating this inequality against $f(x)dx$. If $d \geq 2$, we use the following argument.

- If $|x - z| \geq 2h$ and $y \in \mathcal{B}_M(x, h)$, then $|z - y| \geq |x - z| - h \geq |x - z|/2$. Therefore, by Lemma A.1(iii),

$$\int_{\mathcal{B}_M(x, h)} \frac{\|K\|_\infty h^{-d}}{|z-y|^{d-1}} dy \leq \frac{2^{1-d} \|K\|_\infty h^{-d} \text{vol}_M(\mathcal{B}_M(x, h))}{|x-z|^{d-1}} \lesssim \frac{1}{|x-z|^{d-1}}.$$

- If $|x - z| \leq 2h$, then

$$\begin{aligned} \int_{\mathcal{B}_M(x, h)} \frac{\|K\|_\infty h^{-d}}{|z-y|^{d-1}} dy &\leq \int_{\mathcal{B}_M(z, 3h)} \frac{\|K\|_\infty h^{-d}}{|z-y|^{d-1}} dy \\ &\leq \int_{\mathcal{B}_{T_z M}(0, 3h)} \frac{\|K\|_\infty h^{-d} J\Psi_z(u)}{|z-\Psi_z(u)|^{d-1}} du \\ &\lesssim h^{-d} \int_{\mathcal{B}_{T_z M}(0, 3h)} \frac{du}{|u|^{d-1}} \lesssim h^{1-d}, \end{aligned}$$

where at the last line we used that $|z - \Psi_z(u)| \geq |u|$ and that $J\Psi_z(u) \lesssim 1$ by Lemma A.1.

Hence,

$$\begin{aligned} \mathbb{E} [|\nabla(G(K_h * \delta_X))(z)|^p] &= \int_M f(x) |\nabla(G(K_h * \delta_x))(z)|^p dx \\ &\leq f_{\max} \int_M |\nabla(G(K_h * \delta_x))(z)|^p dx \\ &\lesssim \int_{\mathcal{B}_M(z, 2h)} |\nabla(G(K_h * \delta_x))(z)|^p dx + \int_{M \setminus \mathcal{B}_M(z, 2h)} |\nabla(G(K_h * \delta_x))(z)|^p dx \\ &\lesssim \int_{\mathcal{B}_M(z, 2h)} h^{(1-d)p} dx + \int_{M \setminus \mathcal{B}_M(z, 2h)} |z-x|^{(1-d)p} dx \\ &\lesssim h^{(1-d)p+d} + \int_{M \setminus \mathcal{B}_M(z, 2h)} |z-x|^{(1-d)p} dx. \end{aligned}$$

The latter integral is bounded by

$$\begin{aligned} &\int_{2h \leq |x-z| \leq r_0} |z-x|^{(1-d)p} dx + \int_{|x-z| \geq r_0} |z-x|^{(1-d)p} dx \\ &\leq \int_{2h \leq |\Psi_z(u)-z| \leq r_0} |z-\Psi_z(u)|^{(1-d)p} J\Psi_z(u) du + |\text{vol}_M| r_0^{(1-d)p} \\ &\lesssim \int_{14h/8 \leq |u| \leq r_0} |u|^{(1-d)p} du + 1 \lesssim h^{(1-d)p+d} \text{ if } (1-d)p + d < 0, \end{aligned}$$

where at the last line we use that $|u| \leq |z - \Psi_z(u)| \leq 8|u|/7$ by Lemma A.1. If $d > 2$ or if $d = 2$ and $p > 2$, the condition $(1-d)p + d < 0$ is always satisfied. If $d = 2$ and $p = 2$, then $\int_{14h/8 \leq |u| \leq h_0} |u|^{(1-d)p} du$ is of order $-\log h$, concluding the proof.

APPENDIX E. PROOF OF THEOREM 3.1(I)

Let f be the density of μ and $\tilde{f} = f/\rho_h$. By Lemma 4.2, $f_{\min}(1-c_0h^{k-1}) \leq \tilde{f} \leq f_{\max}(1+c_0h^{k-1})$ for h small enough. We have

$$\begin{aligned} K_h * \tilde{f}(x) &= \int_M K_h(x-y)\tilde{f}(y)dy \\ &= \int_{\mathcal{B}_{T_x M}(0,h)} K_h(x-\Psi_x(v))\tilde{f} \circ \Psi_x(v)J\Psi_x(v)dv \\ &\geq \int_{\mathcal{B}_{T_x M}(0,h)} K_h(v)\tilde{f} \circ \Psi_x(v)J\Psi_x(v)dv - \int_{\mathcal{B}_{T_x M}(0,h)} |K_h(x-\Psi_x(v)) - K_h(v)|\tilde{f} \circ \Psi_x(v)J\Psi_x(v)dv. \end{aligned}$$

By Lemma A.1(v), the quantity $|K_h(x-\Psi_x(v)) - K_h(v)|$ is bounded by

$$\frac{\|K\|_{\mathcal{C}^1(\mathbb{R}^d)}}{h^{d+1}}|x-v-\Psi_x(v)| \lesssim \frac{|v|^2}{h^{d+1}},$$

so that the second term in (E.1) is bounded by $Cf_{\max} \int_{\mathcal{B}_{T_x M}(0,h)} \frac{|v|^2}{h^{d+1}}dv \lesssim h$. Also, using that $|J\Psi_x(v) - 1| \leq c_1|v|$ by Lemma A.1, the first term is larger than

$$\begin{aligned} &f_{\min}(1-c_0h^{k-1})(1-c_1h) \int_{\mathbb{R}^d} K_+ - f_{\max}(1+c_1h)(1+c_0h^{k-1}) \int_{\mathbb{R}^d} K_- \\ &= f_{\min}(1-c_2h) \left(1 + \int_{\mathbb{R}^d} K_-\right) - f_{\max}(1+c_2h) \int_{\mathbb{R}^d} K_- \\ &= f_{\min}(1-c_2h) - (f_{\max}(1+c_2h) - f_{\min}(1-c_2h)) \int_{\mathbb{R}^d} K_-(v)dv \\ &\geq f_{\min}(1-c_2h) - (f_{\max}(1+c_2h) - f_{\min}(1-c_2h))\beta \\ &\geq 3f_{\min}/4, \end{aligned}$$

if $\beta < f_{\min}/(4(f_{\max} - f_{\min}))$ and h is small enough. Likewise, we show that $K_h * \tilde{f}(x) \leq 3f_{\max}/2$. It remains to show that $|K_h * \tilde{f}(x) - K_h * (\mu_n/\rho_h)(x)|$ is small enough for all $x \in M$ with high probability. Note that $K_h * \tilde{f} - K_h * (\mu_n/\rho_h)$ is L -Lipschitz with $L \lesssim h^{-d-1}$. Let $t = f_{\min}/4$ and consider a covering of M by N balls $\mathcal{B}_M(x_j, t/(2L))$. By standard packing arguments, such a covering exists with $N \lesssim (L/t)^d$. If $|K_h * \tilde{f}(x_j) - K_h * \mu_n(x_j)| \leq t/2$ for all $j = 1, \dots, N$, then $\|K_h * \tilde{f} - K_h * \mu_n\|_{L^\infty(M)} \leq t/2 + Lt/(2L) \leq t$. Hence, using Bernstein inequality [GN15, Theorem 3.1.7], as $|K_h(x_j - Y_i)| \leq \|K\|_{\mathcal{C}^0(\mathbb{R}^D)}h^{-d}$ and $\text{Var}(K_h(x_j - Y_i)) \leq \|K^2\|_{\mathcal{C}^0(\mathbb{R}^D)}h^{-d}$, we obtain

$$\begin{aligned} \mathbb{P}(\|K_h * \tilde{f} - K_h * \mu_n\|_{L^\infty(M)} \geq t) &\leq \mathbb{P}(\exists j, |K_h * \tilde{f}(x_j) - K_h * \mu_n(x_j)| \geq t/2) \\ &\lesssim (L/t)^d \mathbb{P}(|K_h * \tilde{f}(x_j) - K_h * \mu_n(x_j)| \geq t/2) \lesssim h^{-d(d+1)} \exp(-Cnh^d). \end{aligned}$$

Choosing $nh^d = C' \log n$ for C' large enough yields the conclusion.

APPENDIX F. PROOFS OF SECTION 4.4

We first prove Lemma 4.8.

. *Proof of (a).* The application $\Psi_{Y_j} \circ \pi_{Y_j} : \mathcal{B}_{\hat{T}_j}(0, 3\varepsilon) \rightarrow M$ is a diffeomorphism on $\mathcal{B}_{\hat{T}_j}(0, 3\varepsilon)$, as the composition of the diffeomorphisms Ψ_{Y_j} and $(\pi_{Y_j})|_{\hat{T}_j}$ (recall that $\angle(\hat{T}_j, T_{Y_j}M) \lesssim \varepsilon^{m-1} + \gamma\varepsilon^{-1} \lesssim 1$ by Proposition 3.5). Furthermore, by Lemma A.1(iv) and the bound on the angle,

$$\mathcal{B}_M(Y_j, 2\varepsilon) \subset \Psi_{Y_j}(\mathcal{B}_{T_{Y_j}M}(0, 2\varepsilon)) \subset (\Psi_{Y_j} \circ \pi_{Y_j})(\mathcal{B}_{\hat{T}_j}(0, 3\varepsilon)).$$

This proves the first part of Lemma 4.8(a). Let $S_j : \mathcal{B}_M(Y_j, 2\varepsilon) \rightarrow \mathcal{B}_{\hat{T}_j}(0, 3\varepsilon)$ be the inverse of $\Psi_{Y_j} \circ \pi_{Y_j}$. By Lemma A.2(ii), $\hat{\Psi}_j$ is injective on \hat{T}_j , while, for $v \in \hat{T}_j$ with $|v| \leq 3\varepsilon$,

$$\left\| \text{id} - d\hat{\Psi}_j(v) \right\|_{\text{op}} \leq \left\| \sum_{a=2}^{m-1} a\hat{V}_{a,j}[\cdot, v^{\otimes(a-1)}] \right\| \lesssim \ell\varepsilon \leq 1/2 \quad (\text{F.1})$$

if $\ell \lesssim \varepsilon^{-1}$ is small enough. Hence, $\hat{\Psi}_j : \mathcal{B}_{\hat{T}_j}(0, 3\varepsilon) \rightarrow \hat{\Psi}_j(\hat{T}_j)$ is a diffeomorphism on its image, and $\hat{\Psi}_j \circ S_j$ is a diffeomorphism as a composition of diffeomorphisms. Note that the inverse of $\hat{\Psi}_j$ is given by $\hat{\pi}_j(\cdot - X_j)$, so that $\mathcal{B}_{\hat{\Psi}_j(\hat{T}_j)}(X_j, \varepsilon) \subset \hat{\Psi}_j(\mathcal{B}_{\hat{T}_j}(0, \varepsilon))$. Furthermore, by Lemma A.1,

$$(\Psi_{Y_j} \circ \pi_{Y_j})(\mathcal{B}_{\hat{T}_j}(0, \varepsilon)) \subset \Psi_{Y_j}(\mathcal{B}_{T_{Y_j}}(0, \varepsilon)) \subset \mathcal{B}_M(Y_j, 8\varepsilon/7),$$

so that $(\hat{\Psi}_j \circ S_j)(\mathcal{B}_M(Y_j, 2\varepsilon))$ contains $\mathcal{B}_{\hat{\Psi}_j(\hat{T}_j)}(X_j, \varepsilon)$. Furthermore, these inclusions of balls also hold for any $\varepsilon' \leq \varepsilon$, proving that $|\hat{\Psi}_j \circ S_j(z) - X_j| \geq (7/8)|z - Y_j|$ for any $z \in \mathcal{B}_M(Y_j, 2\varepsilon)$.

. *Proof of (b).* The formula for the density $\tilde{\chi}_j$ follows from a change of variables.

. *Proof of (c).* The inequality (4.17) follows from Proposition 3.5. We now prove that, for $z \in \mathcal{B}_M(Y_j, 2\varepsilon)$,

$$|\pi_{Y_i}(z - \hat{\Psi}_j \circ S_j(z))| \lesssim (\varepsilon + \gamma\varepsilon^{-1})(\varepsilon^m + \gamma). \quad (\text{F.2})$$

Let $u \in \hat{T}_j$ be such that $z = \Psi_{Y_j} \circ \pi_{Y_j}(u)$ and $y = \hat{\Psi}_j(u)$. Recall that $X_j \in T_{Y_j}M^\perp$ by assumption, so that $\pi_{Y_j}(X_j - Y_j) = 0$. Also, by Lemma A.1(v), we have $\Psi_{Y_j}(\pi_{Y_j}(u)) = Y_j + \pi_{Y_j}(u) + N_{Y_j}(\pi_{Y_j}(u))$ with $N_{Y_j}(\pi_{Y_j}(u)) \in T_{Y_j}M^\perp$, while by Lemma A.2(ii), we have $\hat{\Psi}_j(u) = X_j + u + \hat{N}_j(u)$ with $\hat{N}_j(u) \in \hat{T}_j^\perp$. Hence,

$$\begin{aligned} |\pi_{Y_j}(z - y)| &= |\pi_{Y_j}(Y_j + \pi_{Y_j}(u) + N_{Y_j}(\pi_{Y_j}(u)) - (X_j + u + \hat{N}_j(u)))| \\ &= |\pi_{Y_j}(N_{Y_j}(\pi_{Y_j}(u)) - \hat{N}_j(u))| \\ &\leq \angle(T_{Y_j}M, \hat{T}_j) |N_{Y_j}(\pi_{Y_j}(u)) - \hat{N}_j(u)| + |\hat{\pi}_j(N_{Y_j}(\pi_{Y_j}(u)) - \hat{N}_j(u))| \\ &\lesssim (\varepsilon^{m-1} + \gamma\varepsilon^{-1})(\varepsilon^m + \gamma) + |\hat{\pi}_j(\pi_{Y_j}^\perp(N_{Y_j}(\pi_{Y_j}(u))))| \\ &\lesssim (\varepsilon^{m-1} + \gamma\varepsilon^{-1})(\varepsilon^m + \gamma) + \angle(T_{Y_j}M, \hat{T}_j) |N_{Y_j}(\pi_{Y_j}(u))| \\ &\lesssim (\varepsilon^{m-1} + \gamma\varepsilon^{-1})(\varepsilon^m + \gamma + \varepsilon^2) \lesssim (\varepsilon^{m-1} + \gamma\varepsilon^{-1})(\varepsilon^2 + \gamma), \end{aligned}$$

where we used Proposition 3.5 to bound $\angle(T_{Y_j}M, \hat{T}_j)$, Lemma A.2 to bound $|N_{Y_j}(\pi_{Y_j}(u)) - \hat{N}_j(u)|$ and Lemma A.1 to bound $|N_{Y_j}(\pi_{Y_j}(u))|$. We obtain (F.2).

To prove inequality (4.18), we first bound $|\chi_j(\hat{\Psi}_j \circ S_j(z)) - \chi_j(z)|$ and then bound $|J(\hat{\Psi}_j \circ S_j)(z) - 1|$. The first bound is based on the following elementary lemma.

Lemma F.1. *Let $\theta : \mathbb{R}^D \rightarrow \mathbb{R}$ be a smooth radial function. Then, $|\theta(x) - \theta(y)| \leq \frac{\|\theta\|_{C^2(\mathbb{R}^D)}}{2} \|x\|^2 - \|y\|^2$.*

Proof. As $d\theta(0) = 0$, one can write $\theta(x) = \tilde{\theta}(\|x\|^2)$ for some function $\tilde{\theta}$ which is Lipschitz continuous with Lipschitz constant $\frac{\|d^2\theta\|_{C^0(\mathbb{R}^D)}}{2}$. This implies the conclusion. \square

Recall from the proof of Lemma 3.6 that we have $\chi_j(z) = \zeta_j(z) / \sum_{i=1}^J \zeta_i(z)$ where $\zeta_i = \theta(\frac{z - X_i}{\varepsilon})$ for some smooth radial function θ , and that furthermore, there is at most c_d non-zero terms in the sum in the denominator, which is always larger than 1. Hence, if we control for every $i = 1, \dots, J$ the difference $||z - X_i|^2 - |\hat{\Psi}_j \circ S_j(z) - X_i|^2|$, then we obtain a control on $|\chi_j(z) - \chi_j(\hat{\Psi}_j \circ S_j(z))|$. Let $z \in M$ be such that $|z - X_j| \leq 2\varepsilon$ (for otherwise both $\chi_j(z)$ and $\chi_j(\hat{\Psi}_j \circ S_j(z))$ are equal to zero). We have by (4.17) and (F.2),

$$\begin{aligned} ||\hat{\Psi}_j \circ S_j(z) - X_i|^2 - |z - X_i|^2| &= ||\hat{\Psi}_j \circ S_j(z) - z|^2 + 2\langle \hat{\Psi}_j \circ S_j(z) - z, z - X_i \rangle| \\ &\lesssim (\varepsilon^m + \gamma)^2 + |\langle \hat{\Psi}_j \circ S_j(z) - z, z - Y_i \rangle| + |\langle \hat{\Psi}_j \circ S_j(z) - z, X_i - Y_i \rangle| \\ &\lesssim (\varepsilon^m + \gamma)^2 + |\langle \pi_{Y_j}(\hat{\Psi}_j \circ S_j(z) - z), \pi_{Y_j}(z - Y_i) \rangle| \\ &\quad + |\langle \pi_{Y_j}^\perp(\hat{\Psi}_j \circ S_j(z) - z), \pi_{Y_j}^\perp(z - Y_i) \rangle| + (\varepsilon^m + \gamma)\gamma \\ &\lesssim (\varepsilon^m + \gamma)^2 + (\varepsilon + \gamma\varepsilon^{-1})(\varepsilon^m + \gamma)|z - Y_i| \\ &\quad + (\varepsilon^m + \gamma)|\pi_{Y_j}^\perp(z - Y_i)| + (\varepsilon^m + \gamma)\gamma. \end{aligned} \quad (\text{F.3})$$

By Lemma A.1(i) and the fact that $|z - Y_j| \leq |z - X_j| + \gamma \lesssim \varepsilon$, we have $|\pi_{Y_j}^\perp(z - Y_i)| \leq |\tilde{\pi}_{Y_j}^\perp(z)| + |\tilde{\pi}_{Y_j}^\perp(Y_i)| \lesssim \varepsilon^2 + |Y_i - Y_j|^2$. Hence, we obtain that

$$\|\hat{\Psi}_j \circ S_j(z) - X_i\|^2 - |z - X_i|^2 \lesssim (\varepsilon^m + \gamma)(\varepsilon^2 + \gamma + |Y_i - Y_j|^2). \quad (\text{F.4})$$

Therefore,

$$\begin{aligned} \left| \theta \left(\frac{z - X_i}{\varepsilon} \right) - \theta \left(\frac{\hat{\Psi}_j \circ S_j(z) - X_i}{\varepsilon} \right) \right| &\lesssim \frac{(\varepsilon^m + \gamma)(\varepsilon^2 + \gamma + |Y_i - Y_j|^2)}{\varepsilon^2} \\ &\lesssim (\varepsilon^m + \gamma) \left(1 + \gamma\varepsilon^{-2} + \frac{|Y_i - Y_j|^2}{\varepsilon^2} \right). \end{aligned} \quad (\text{F.5})$$

Note also that if $|Y_i - Y_j| \geq 3\varepsilon$, then $|z - X_i| \geq |X_i - X_j| - |z - X_j| \geq 3\varepsilon - \varepsilon - 3\gamma \geq \varepsilon$, while by the same argument $|\hat{\Psi}_j \circ S_j(z) - X_i| \geq \varepsilon$. Hence, both terms in the left-hand side of (F.5) are null in that case. Thus, we may assume that $|Y_i - Y_j| \leq 3\varepsilon$, so that $\left| \theta \left(\frac{z - X_i}{\varepsilon} \right) - \theta \left(\frac{\hat{\Psi}_j \circ S_j(z) - X_i}{\varepsilon} \right) \right| \lesssim (\varepsilon^m + \gamma)(1 + \gamma\varepsilon^{-2})$. From the definition of $\chi_j(z)$, and as the function $t \mapsto 1/t$ is Lipschitz on $[1, \infty[$, we obtain that

$$|\chi_j(z) - \chi_j(\hat{\Psi}_j \circ S_j(z))| \lesssim (\varepsilon^m + \gamma)(1 + \gamma\varepsilon^{-2}).$$

We now provide a bound on $|J(\hat{\Psi}_j \circ S_j)(z) - 1|$. One has, for $u = S_j(z) \in \hat{T}_j$,

$$|J(\hat{\Psi}_j \circ S_j)(z) - 1| = \frac{|J\hat{\Psi}_j(u) - J(\Psi_{Y_j} \circ \pi_{Y_j})(u)|}{J(\Psi_{Y_j} \circ \pi_{Y_j})(u)}.$$

By Lemma A.1(v) and Lemma A.2(ii), $\|\text{id}_{\hat{T}_j} - d(\Psi_{Y_j} \circ \pi_{Y_j})(u)\|_{\text{op}} \lesssim 1$ and $\|\text{id}_{\hat{T}_j} - d\hat{\Psi}_j(u)\|_{\text{op}} \lesssim 1$ for u small enough. As a consequence, both Jacobians are larger than, say $1/2$ for u small enough, and, as the function $A \in \mathbb{R}^{d \times d} \mapsto \sqrt{\det(A)}$ is Lipschitz continuous on the set of matrices with $\det(A) \geq 1/2$ and $\|A\|_{\text{op}} \leq 2$, we have

$$|J(\hat{\Psi}_j \circ S_j)(z) - 1| \lesssim \left\| d\hat{\Psi}_j(u)^\top d\hat{\Psi}_j(u) - d(\Psi_{Y_j} \circ \pi_{Y_j})(u)^\top d(\Psi_{Y_j} \circ \pi_{Y_j})(u) \right\|_{\text{op}}. \quad (\text{F.6})$$

Recall that $\hat{\Psi}_j(u) = X_j + u + \hat{N}_j(u)$ and $\Psi_{Y_j} \circ \pi_{Y_j}(u) = Y_j + \pi_{Y_j}(u) + N_{Y_j} \circ \pi_{Y_j}(u)$. We may write

$$\begin{aligned} d\hat{\Psi}_j(u)^\top d\hat{\Psi}_j(u) &= \text{id}_{\hat{T}_j} + (d\hat{N}_j(u))^\top d\hat{N}_j(u) \quad \text{and} \\ d(\Psi_{Y_j} \circ \pi_{Y_j})(u)^\top d(\Psi_{Y_j} \circ \pi_{Y_j})(u) &= \hat{\pi}_j \pi_{Y_j} \hat{\pi}_j + (d(N_{Y_j} \circ \pi_{Y_j})(u))^\top d(N_{Y_j} \circ \pi_{Y_j})(u). \end{aligned}$$

One has $\left\| \text{id}_{\hat{T}_j} - \hat{\pi}_j \pi_{Y_j} \hat{\pi}_j \right\|_{\text{op}} = \left\| \hat{\pi}_j \pi_{Y_j}^\perp \pi_{Y_j}^\perp \hat{\pi}_j \right\|_{\text{op}} \leq \angle(T_{Y_j} M, \hat{T}_j)^2 \lesssim (\varepsilon^{m-1} + \gamma\varepsilon^{-1})^2 \leq (\varepsilon^m + \gamma)(1 + \gamma\varepsilon^{-2})$. Furthermore, by Lemma A.2(iv),

$$\begin{aligned} &\left\| (d\hat{N}_j(u))^\top d\hat{N}_j(u) - (d(N_{Y_j} \circ \pi_{Y_j})(u))^\top d(N_{Y_j} \circ \pi_{Y_j})(u) \right\|_{\text{op}} \\ &\leq \left(\left\| d\hat{N}_j(u) \right\|_{\text{op}} + \left\| d(N_{Y_j} \circ \pi_{Y_j})(u) \right\|_{\text{op}} \right) \left\| d\hat{N}_j(u) - d(N_{Y_j} \circ \pi_{Y_j})(u) \right\|_{\text{op}} \\ &\lesssim \varepsilon(\varepsilon^{m-1} + \gamma\varepsilon^{-1}) \lesssim \varepsilon^m + \gamma. \end{aligned}$$

Putting together (F.6) with those two inequalities, we obtain that $|J(\hat{\Psi}_j \circ S_j)(z) - 1| \lesssim (\varepsilon^m + \gamma)(1 + \gamma\varepsilon^{-2})$, concluding the proof of Lemma 4.8.

To conclude the proof of Theorem 3.8, it remains to control the quantity T appearing in Lemma 4.9 for $\phi = K_h * (\nu_n / \hat{\rho}_h)$ and $\phi' = K_h * (\mu_n / \rho_h)$.

Lemma F.2. *The quantity $T = \max_{j=1 \dots J} \sup_{z \in \mathcal{B}(Y_j, \varepsilon)} |\phi(\hat{\Psi}_j \circ S_j(z)) - \phi'(z)|$ satisfies $T \lesssim (\varepsilon^m + \gamma)(1 + \gamma\varepsilon^{-2})$ with probability larger than $1 - cn^{-k/d}$.*

Proof. For $z \in \mathcal{B}(Y_j, \varepsilon)$, we have

$$|\phi(\hat{\Psi}_j \circ S_j(z)) - \phi'(z)| \leq \frac{1}{n} \sum_{i=1}^n \left| \frac{K_h * \delta_{X_i}(\hat{\Psi}_j \circ S_j(z))}{\hat{\rho}_h(X_i)} - \frac{K_h * \delta_{Y_i}(z)}{\rho_h(Y_i)} \right|.$$

The same computation than in (F.3) shows that

$$|\hat{\Psi}_j \circ S_j(z) - Y_i|^2 - |z - Y_i|^2 \lesssim (\varepsilon^m + \gamma)(\varepsilon^2 + \gamma + |Y_i - Y_j|^2).$$

This inequality together with Lemma F.1 yield

$$\begin{aligned} & |K_h(X_i - \hat{\Psi}_j \circ S_j(z)) - K_h(Y_i - z)| \\ & \lesssim h^{-d-2}(\varepsilon^m + \gamma)(\varepsilon^2 + \gamma + |Y_i - Y_j|^2). \end{aligned}$$

We may assume that $|Y_i - Y_j| \leq 3h$ and $|z - Y_i| \leq 2h$, for otherwise both quantities in the left-hand side of the above equation are equal to zero. Hence, as $\varepsilon \lesssim h$ by assumption, we have

$$|K_h(X_i - \hat{\Psi}_j \circ S_j(z)) - K_h(Y_i - z)| \lesssim h^{-d}(\varepsilon^m + \gamma)(1 + \gamma\varepsilon^{-2})\mathbf{1}\{Y_i \in \mathcal{B}_M(z, 2h)\}. \quad (\text{F.7})$$

Let us now bound $|\hat{\rho}_h(X_i) - \rho_h(Y_i)|$. By the triangle inequality, and using (4.18) and (F.7), we obtain that this quantity is smaller than

$$\begin{aligned} & \left| \sum_{j=1}^J \int_{\hat{\Psi}_j(\hat{T}_j)} \chi_j(w) K_h(X_i - w) dw - \sum_{j=1}^J \int_M \chi_j(z) K_h(Y_i - z) dz \right| \\ & \leq \sum_{j=1}^J \int_M |\tilde{\chi}_j(z) K_h(X_i - \hat{\Psi}_j \circ S_j(z)) - \chi_j(z) K_h(Y_i - z)| dz \\ & \lesssim \sum_{j=1}^J \int_M (\mathbf{1}\{z \in \mathcal{B}_M(Y_j, 2\varepsilon)\}) (\varepsilon^m + \gamma)(1 + \gamma\varepsilon^{-2}) |K_h(Y_i - z)| \\ & \quad + \tilde{\chi}_j(z) h^{-d} (\varepsilon^m + \gamma)(1 + \gamma\varepsilon^{-2}) \mathbf{1}\{z \in \mathcal{B}_M(Y_i, 2h)\} dz \\ & \lesssim h^{-d} (\varepsilon^m + \gamma)(1 + \gamma\varepsilon^{-2}) \sum_{j=1}^J \int_M \mathbf{1}\{z \in \mathcal{B}_M(Y_j, 2\varepsilon)\} \mathbf{1}\{z \in \mathcal{B}_M(Y_i, 2h)\} dz \\ & \lesssim \varepsilon^d h^{-d} (\varepsilon^m + \gamma)(1 + \gamma\varepsilon^{-2}) \sum_{j=1}^J \mathbf{1}\{|Y_j - Y_i| \leq 4h\} \\ & \lesssim h^{-d} (\varepsilon^m + \gamma)(1 + \gamma\varepsilon^{-2}) \sum_{j=1}^J \mathbf{1}\{|Y_j - Y_i| \leq 4h\} \text{vol}_M(\mathcal{B}_M(Y_j, \varepsilon/8)) \\ & \lesssim h^{-d} (\varepsilon^m + \gamma)(1 + \gamma\varepsilon^{-2}) \text{vol}_M(\mathcal{B}_M(Y_i, 5h)) \lesssim (\varepsilon^m + \gamma)(1 + \gamma\varepsilon^{-2}), \end{aligned}$$

where we use that $\{X_1, \dots, X_J\}$ is $7\varepsilon/24$ -sparse, so that $\{Y_1, \dots, Y_J\}$ is $\varepsilon/4$ -sparse. Therefore, the balls $\mathcal{B}_M(Y_j, \varepsilon/8)$ for $|Y_j - Y_i| \leq 4h$ are pairwise distincts, and are all included in $\mathcal{B}_M(Y_i, 4h + \varepsilon/8) \subset \mathcal{B}_M(Y_i, 5h)$. We conclude by Lemma A.1(iii). Letting $N(z, 2h)$ be the number of points Y_i belonging to $\mathcal{B}_M(z, 2h)$, we obtain

$$\begin{aligned} & |\phi(\hat{\Psi}_j \circ S_j(z)) - \phi'(z)| \\ & \lesssim \frac{1}{n} \sum_{i=1}^n (|K_h(Y_i - z)| (\varepsilon^m + \gamma)(1 + \gamma\varepsilon^{-2}) + h^{-d} (\varepsilon^m + \gamma)(1 + \gamma\varepsilon^{-2}) \mathbf{1}\{Y_i \in \mathcal{B}_M(z, 2h)\}) \\ & \lesssim \frac{N(z, 2h)}{nh^d} (\varepsilon^m + \gamma)(1 + \gamma\varepsilon^{-2}). \end{aligned}$$

If, for every $z \in M$ and some $\lambda > 0$, $N(z, 2h) \leq \lambda nh^d$, then we have the conclusion. Let us bound

$$P_0 = \mathbb{P}(\exists z \in M, N(z, 2h) > \lambda nh^d).$$

If $N(z, 2h) > \lambda nh^d$, then there exists a point Y_i with $N(Y_i, 4h) \geq N(z, 2h) > \lambda nh^d$. Hence, $P_0 \leq n\mathbb{P}(N(Y_1, 4h) > \lambda nh^d)$. Conditionally on Y_1 , $N(Y_1, 4h) = 1 + U$ with U a binomial random

variable of parameters $n - 1$ and $\mu(\mathcal{B}_M(Y_1, 4h)) \leq f_{\max} \text{vol}_M(\mathcal{B}_M(Y_1, 4h)) \lesssim h^d$ (see Lemma A.1(iii)). In particular, for λ large enough, the probability P_0 is smaller than $n^{-k/d}$ by Bernstein's inequality, as long as $nh^d \gtrsim 1$. \square

We conclude this section by giving a proof of Proposition 5.2.

Proof. Recall that W_1, \dots, W_N is a N -sample of law \hat{U}_M , and that we are in the noiseless regime $\gamma = 0$ with $m = k$. Define j_a the index with $W_a \in \hat{\Psi}_{j_a}(\hat{T}_{j_a})$, and let $H_a = (\hat{\Psi}_{j_a} \circ S_{j_a})^{-1}(W_a)$. Then, Lemma 4.8 implies that $|W_a - H_a| \lesssim \varepsilon^k$. Furthermore, the sample H_1, \dots, H_N has a law μ_H with density $\sum_{j=1}^J \tilde{\chi}_j$ on M . We decompose the distance into

$$W_\infty((\hat{U}_M)_N, U_M) \leq W_\infty((\hat{U}_M)_N, N^{-1} \sum_{a=1}^N \delta_{H_a}) + W_\infty(N^{-1} \sum_{a=1}^N \delta_{H_a}, \mu_H) + W_\infty(\mu_H, U_M).$$

The first term is of order ε^k , while the second term scales as the second term of (5.8) according to [TGHS20]. The third term was already shown to be bounded by ε^k in the proof of Lemma 4.9 (with $\phi = \tilde{\phi} = 1$). As $\varepsilon^k \simeq (\log n/n)^{k/d}$, this concludes the proof. \square

APPENDIX G. LOWER BOUNDS ON MINIMAX RISKS

In this section, we prove the different lower bounds on minimax risks stated in the article. The main tool used will be Assouad's lemma. Fix a statistical model $(\mathcal{Q}, \vartheta, \mathcal{L})$, where we observe a sample of law $\mu \in \mathcal{Q}$, while $\vartheta(\mu)$ is a quantity of interest to be estimated, with risk measured by the loss function \mathcal{L} .

Lemma G.1 (Assouad's lemma [Yu97]). *Let $m \geq 1$ be an integer and let $\mathcal{Q}_m = \{\mu_\sigma, \sigma \in \{-1, 1\}^m\} \subset \mathcal{Q}$ be a set of probability measures. Assume that for all $\sigma, \sigma' \in \{-1, 1\}^m$,*

$$\mathcal{L}(\vartheta(\mu_\sigma), \vartheta(\mu_{\sigma'})) \geq |\sigma - \sigma'| \delta, \quad (\text{G.1})$$

where $|\sigma - \sigma'| = \sum_{i=1}^m \mathbf{1}\{\sigma(i) \neq \sigma'(i)\}$ is the Hamming distance between σ and σ' . Then,

$$\mathcal{R}_n(\vartheta, \mathcal{Q}, \mathcal{L}) \geq m \frac{\delta}{16} (1 - \max\{\text{TV}(\mu_\sigma, \mu_{\sigma'}), |\sigma - \sigma'| = 1\})^{2n}. \quad (\text{G.2})$$

The lower bound on the minimax rates we prove are actually going to hold on the smaller model of uniform distributions on manifolds.

Definition G.2. *Let $k \geq 2$ and $\gamma \geq 0$. The set \mathcal{Q}_d^k is the set of uniform distributions on some manifold $M \in \mathcal{M}_d^k$ with $f_{\max}^{-1} \leq |\text{vol}_M| \leq f_{\min}^{-1}$.*

One can check that $\mathcal{Q}_d^k \subset \mathcal{Q}_d^{k,s}$, with parameter $L_s = f_{\min}^{-1/p} \vee f_{\max}^{1-1/p}$. Therefore, a lowerbound on the minimax risk on the model \mathcal{Q}_d^k yields a lowerbound on the minimax risk on the model $\mathcal{Q}_d^{k,s}$ should the parameter L_s be large enough.

We build a subfamily of manifolds indexed by $\sigma \in \{-1, 1\}^m$ following [AL19]. By [AL19, Section C.2], there exists a d -dimensional manifold $M \subset \mathbb{R}^{d+1}$ of reach $2\tau_{\min}$, of volume $C_d \tau_{\min}^d$ which contains $\mathcal{B}_{\mathbb{R}^d}(0, \tau_{\min}) \times \{0\}$ (that we identify with $\mathcal{B}_{\mathbb{R}^d}(0, \tau_{\min})$). Let $\delta > 0$ and consider a family of m points $x_1, \dots, x_m \in \mathcal{B}_{\mathbb{R}^d}(0, \tau_{\min}/2)$, with $|x_i - x_{i'}| \geq 4\delta$ for $i \neq i'$ and $c_d(\tau_{\min}/\delta)^d \leq m \leq C_d(\tau_{\min}/\delta)^d$. Let $0 < \Lambda < \delta$ and let $\phi: \mathbb{R}^{d+1} \rightarrow [0, 1]$ be a smooth radial function supported on $\mathcal{B}(0, 1)$, with $\phi \equiv 1$ on $\mathcal{B}(0, 1/2)$. Let e be the unit vector in the $(d+1)$ th direction. We then let, for $\sigma \in \{-1, 1\}^m$,

$$\Phi_\sigma^\Lambda(x) = x + \sum_{i=1}^m \frac{\sigma(i) + 1}{2} \Lambda \phi\left(\frac{x - x_i}{\delta}\right) e. \quad (\text{G.3})$$

Let $M_\sigma^\Lambda = \Phi_\sigma^\Lambda(M)$ and μ_σ^Λ be the the uniform measure on M_σ^Λ . Informally, the manifold M_σ^Λ is obtained by adding bumps of height Λ to the base manifold M at locations x_i such that $\sigma(i) = +1$. If $\Lambda \leq c_{k,d,\tau_{\min}} \delta^k$, then $\mu_\sigma^\Lambda \in \mathcal{Q}_d^k$, provided that L_k is large enough [AL19, Lemma

C.13]. If $\sigma(i) = 1$, the volume of $\Phi_\sigma^\Lambda(\mathcal{B}_{\mathbb{R}^d}(x_i, \delta))$ satisfies (with ω_d denoting the volume of the d -dimensional unit ball)

$$\begin{aligned} |\text{vol}_{M_\sigma^\Lambda}(\Phi_\sigma^\Lambda(\mathcal{B}_{\mathbb{R}^d}(x_i, \delta))) - \omega_d \delta^d| &\leq \int_{\mathcal{B}_{\mathbb{R}^d}(x_i, \delta)} |J\Phi_\sigma^\Lambda(x) - 1| dx \\ &\leq \int_{\mathcal{B}_{\mathbb{R}^d}(x_i, \delta)} \left| \sqrt{1 + \Lambda^2 \delta^{-2} \left| \nabla \phi \left(\frac{x - x_i}{\delta} \right) \right|^2} - 1 \right| dx \leq C_d \delta^d \Lambda^2 \delta^{-2}. \end{aligned}$$

Hence, for δ small enough, we have $|\text{vol}_{M_\sigma^\Lambda} - C_d \tau_{\min}^d| \leq m C_d \delta^d \Lambda^2 \delta^{-2} \leq C_d \tau_{\min}^d / 3$, as $m \leq C_d (\tau_{\min} / \delta)^d$ and $\Lambda \leq c_{k,d,\tau_{\min}} \delta^k$. As a consequence, if $|\sigma - \sigma'| = 1$, with for instance $\sigma(i) = 1$ and $\sigma'(i) = -1$, then

$$\text{TV}(\mu_\sigma^\Lambda, \mu_{\sigma'}^\Lambda) \leq \max(\mu_\sigma^\Lambda(\Phi_\sigma^\Lambda(\mathcal{B}_{\mathbb{R}^d}(x_i, \delta))), \mu_{\sigma'}^\Lambda(\mathcal{B}_{\mathbb{R}^d}(x_i, \delta))) \leq C_{d,\tau_{\min}} \delta^d. \quad (\text{G.4})$$

We may now prove the different minimax lower bounds using Assouad's Lemma on the family $\{\mu_\sigma^\Lambda, \sigma \in \{-1, 1\}^m\}$.

Proof of Theorem 2.11. As g is nondecreasing and convex, by Jensen's inequality, we may assume without loss of generality that $\mathcal{L} = \text{TV}$. Let $\Gamma = |(\mu_\sigma^\Lambda - \mu_{\sigma'}^\Lambda)(B_i)|$, where $B_i = \mathcal{B}_{\mathbb{R}^d}(x_i, \delta)$ and $\sigma(i) \neq \sigma'(i)$. Then, $\text{TV}(\mu_\sigma^\Lambda, \mu_{\sigma'}^\Lambda) \geq |\sigma - \sigma'| \Gamma$. Furthermore, if for instance $\sigma'(i) = 1$, $\Gamma \geq \mu_{\sigma'}^\Lambda(B_i) = (\omega_d \delta^d) / |\text{vol}_{M_{\sigma'}^\Lambda}| \geq c_d \delta^d / \tau_{\min}^d$. By Assouad's Lemma,

$$\begin{aligned} \mathcal{R}_n(\mu, \mathcal{Q}_d^{s,k}, \text{TV}) &\geq \mathcal{R}_n(\mu, \mathcal{Q}_d^k, \text{TV}) \geq \frac{m}{16} c_d \frac{\delta^d}{\tau_{\min}^d} (1 - C_{d,\tau_{\min}} \delta^d)^{2n} \\ &\geq C_d (1 - C_{d,\tau_{\min}} \delta^d)^{2n}. \end{aligned}$$

We obtain the conclusion by letting δ go to 0. \square

Proof of Theorem 3.7(iii). As, $W_r \geq W_1$, we may assume that $r = 1$. Let $\sigma, \sigma' \in \{-1, 1\}^m$ with $\sigma(i) \neq \sigma'(i)$. Let $p_{\sigma(i)} = \text{vol}_{M_\sigma^\Lambda}(\mathcal{B}(x_i, \delta))$ and $U_{\sigma,i}^\Lambda = p_{\sigma(i)}^{-1} (\text{vol}_{M_\sigma^\Lambda})|_{\mathcal{B}(x_i, \delta)}$. By the Kantorovitch-Rubinstein duality formula, $W_1(\mu, \nu) = \max \int f d(\mu - \nu)$, where the maximum is taken over all 1-Lipschitz continuous functions $f : \mathbb{R}^D \rightarrow \mathbb{R}$. Recall that e is the unit vector in the $(d+1)$ th direction and let $f : x \mapsto x \cdot e$. Assume for instance that $\sigma(i) = -1$ and $\sigma'(i) = 1$. We have $f(x) = 0$ for $x \in \mathcal{B}_{M_\sigma^\Lambda}(x_i, \delta)$ and $f(x) = \Lambda$ for $x \in \mathcal{B}_{M_{\sigma'}^\Lambda}(x_i, \delta/2)$. Therefore, we have, as $p_{\sigma'(i)} \leq c \delta^{-d}$,

$$W_1(U_{\sigma,i}^\Lambda, U_{\sigma',i}^\Lambda) \geq p_{\sigma'(i)}^{-1} \Lambda \omega_d (\delta/2)^d \geq c_1 \Lambda.$$

Note also that $|p_{\sigma(i)} - p_{\sigma'(i)}| \leq |\text{vol}_{M_\sigma^\Lambda}(\Phi_\sigma^\Lambda(\mathcal{B}_{\mathbb{R}^d}(x_i, \delta))) - \omega_d \delta^d| \leq C_d \delta^d \Lambda^2 \delta^{-2}$. Furthermore, $|\text{vol}_{M_\sigma^\Lambda} - \text{vol}_{M_{\sigma'}^\Lambda}| \leq \sum_{i=1}^m |p_{\sigma(i)} - p_{\sigma'(i)}| \leq |\sigma - \sigma'| C_d \delta^d \Lambda^2 \delta^{-2}$. Let f_i be a 1-Lipschitz continuous function such that $W_1(U_{\sigma,i}^\Lambda, U_{\sigma',i}^\Lambda) = \int f_i d(U_{\sigma,i}^\Lambda - U_{\sigma',i}^\Lambda)$. One can choose f_i such that $f_i(x_i) = 0$, so that the maximum of $|f_i|$ on $\mathcal{B}(x_i, \delta)$ is at most δ . One can then change the value of f_i outside the ball without changing the value of the integral, so that f_i is supported on $\mathcal{B}(x_i, 2\delta)$ and is 1-Lipschitz continuous. Consider the function f obtained by gluing together the different functions f_i . The function f is 1-Lipschitz continuous, so that

$$\begin{aligned} W_1(\mu_\sigma^\Lambda, \mu_{\sigma'}^\Lambda) &\geq \sum_{i=1}^m \left(\frac{p_{\sigma(i)}}{|\text{vol}_{M_\sigma^\Lambda}|} U_{\sigma,i}^\Lambda - \frac{p_{\sigma'(i)}}{|\text{vol}_{M_{\sigma'}^\Lambda}|} U_{\sigma',i}^\Lambda \right) (f) \\ &\geq \sum_{i=1}^m \frac{p_{\sigma(i)}}{|\text{vol}_{M_\sigma^\Lambda}|} (U_{\sigma,i}^\Lambda - U_{\sigma',i}^\Lambda)(f) - |p_{\sigma(i)} - p_{\sigma'(i)}| \frac{|U_{\sigma',i}^\Lambda(f)|}{|\text{vol}_{M_\sigma^\Lambda}|} - p_{\sigma'(i)} |U_{\sigma',i}^\Lambda(f)| \left| \frac{1}{|\text{vol}_{M_\sigma^\Lambda}|} - \frac{1}{|\text{vol}_{M_{\sigma'}^\Lambda}|} \right| \\ &\geq \sum_{i=1}^m \frac{p_{\sigma(i)}}{|\text{vol}_{M_\sigma^\Lambda}|} W_1(U_{\sigma,i}^\Lambda, U_{\sigma',i}^\Lambda) - \sum_{i=1}^m c_4 |p_{\sigma(i)} - p_{\sigma'(i)}| \delta \mathbf{1}\{\sigma(i) \neq \sigma'(i)\} - c_5 \delta |\sigma - \sigma'| \delta^d \Lambda^2 \delta^{-2} \\ &\geq \sum_{i=1}^m \mathbf{1}\{\sigma(i) \neq \sigma'(i)\} (c_6 \delta^d \Lambda - c_4 \delta^d \Lambda^2 \delta^{-1}) - c_5 \delta |\sigma - \sigma'| \delta^d \Lambda^2 \delta^{-2} \geq c_7 \delta^d \Lambda |\sigma - \sigma'|, \end{aligned}$$

where we used at the last line that we choose $\Lambda \leq c\delta^2$ for some constant c small enough. More precisely, we let $\Lambda = c_{k,d,\tau_{\min},L_k} \delta^k$ and $\delta = n^{-1}$, and obtain, by Assouad's Lemma,

$$\mathcal{R}_n \left(\frac{\text{vol}_M}{|\text{vol}_M|}, \mathcal{Q}_d^k(\gamma), W_r \right) \gtrsim n^{-k/d}.$$

□

Proof of Theorem 3.1(iv). Let $a_n = n^{-\frac{s+1}{2s+d}}$ if $d \geq 3$ and $a_n = n^{-1/2}$ if $d \leq 2$. As $W_p \geq W_1$, we may assume without loss of generality that $p = 1$, and up to rescaling, we assume that $\tau_{\min} = \sqrt{d}$. Consider the manifold $M \subset \mathbb{R}^{d+1}$ containing $\mathcal{B}_{\mathbb{R}^d}(0, \sqrt{d})$ of the previous proof. In particular, M contains the cube $[-1, 1]^d$. We adapt the proof of Theorem 3 in [WB19b], where authors consider a family of functions $f_\sigma : [-1, 1]^d \rightarrow M$ indexed by $\sigma \in \{-1, 1\}^m$, with $f_\sigma = 1 + n^{-1/2} \sum_{j=1}^m \sigma_j \psi_j$, where $(\psi_j)_{j=1, \dots, m}$ are elements of a wavelet basis of $[-1, 1]^d$ that satisfy $\int \psi_j = 0$ (see [WB19b, Appendix E] for details on the construction of the wavelet basis). If $m \lesssim n^{d/(2s+d)}$, then $t_0 \leq f_\sigma \leq t_1$ for some positive constants $t_0 < 1 < t_1$, and $\|f_\sigma\|_{B_{p,q}^s([-1, 1]^d)} \lesssim 1$. Note that each ψ_j is supported on a small rectangle inside $[-1, 1]^d$, and can be extended to a smooth function on M (by simply defining $\psi_j = 0$ outside $[-1, 1]^d$). Therefore, we can also consider f_σ as being defined on M . This extension (that we still denote by f_σ) also satisfies $t_0 \leq f_\sigma \leq t_1$ and $\|f_\sigma\|_{B_{p,q}^s(M)} \lesssim 1$ (this last inequality is clear for the $\|\cdot\|_{H_l^s(M)}$ norm for l an integer, while the result follows from interpolation for Besov spaces [Lun18, Corollary 1.1.7]).

As $\int \psi_j = 0$, we have $\int_M f_\sigma = |\text{vol}_M|$. Let $\tilde{f}_\sigma = f_\sigma / |\text{vol}_M|$, that is larger than $f_{\min} = t_0 / |\text{vol}_M|$ and smaller than $f_{\max} = t_1 / |\text{vol}_M|$. Hence, identifying measures with their densities, the set

$$\mathcal{Q}_m = \{\tilde{f}_\sigma, \sigma \in \{-1, 1\}^m\}$$

is a subset of $\mathcal{Q}_d^{s,k}$ for f_{\min} small enough and L_k, L_s, f_{\max} large enough. Furthermore, for $\sigma, \sigma' \in \{-1, 1\}^m$, we have $\text{TV}(\tilde{f}_\sigma, \tilde{f}_{\sigma'}) = \text{TV}(f_\sigma, f_{\sigma'}) / |\text{vol}_M|$. Also, for any function $\phi : \mathbb{R}^{d+1} \rightarrow \mathbb{R}$ that is 1-Lipschitz, we have

$$\int_M \phi(\tilde{f}_\sigma - \tilde{f}_{\sigma'}) = \int_{[-1, 1]^d} \frac{\phi(f_\sigma - f_{\sigma'})}{|\text{vol}_M|},$$

so that $W_1(\tilde{f}_\sigma, \tilde{f}_{\sigma'}) = W_1(f_\sigma, f_{\sigma'}) / |\text{vol}_M|$. Hence, we have reduced our problem to the case of the cube, and applying Assouad's inequality in the same fashion than in [WB19b, Theorem 3] yields that $\mathcal{R}_n(\mu, \mathcal{Q}_d^{s,k}, W_1) \gtrsim a_n$. □

APPENDIX H. EXISTENCE OF KERNELS SATISFYING CONDITIONS A , $B(m)$ AND $C(\beta)$

The goal of the section is to prove the existence of a kernel K satisfying the conditions A , $B(m)$ and $C(\beta)$ stated at the beginning of Section 3.

If K is a radial kernel, we have by integration by parts, as K is smooth with compact support,

$$\begin{aligned} \int_{\mathbb{R}^d} \partial^{\alpha_0} K(v) v^{\alpha_1} dv &= C_{\alpha_0, \alpha_1} \int_{\mathbb{R}^d} K(v) v^{\alpha_1 + \alpha_0} dv \\ &= C'_{\alpha_0, \alpha_1} \int_{\mathbb{R}} K(r) r^{d + |\alpha_0| + |\alpha_1| - 1} dr. \end{aligned}$$

Hence, to show the existence of such a kernel, it suffices to find, for every $m \geq 0$ and every positive constant κ , a smooth even function $K : \mathbb{R} \rightarrow \mathbb{R}$ supported on $[-1, 1]$ satisfying

- **Condition A' :** $\int_{\mathbb{R}} K(r) r^{d-1} dr = \kappa$,
- **Condition $B'(m)$:** $\int_{\mathbb{R}} K(r) r^{d+i-1} dr = 0$ for $i = 1, \dots, m$,
- **Condition $C'(\beta)$:** $\int_{\mathbb{R}} K(r) r^{-d-1} dr \leq \beta$.

We show by recursion on m that for any $\beta > 0$, there exists a such a kernel. For $m = 0$, let K_0 be any smooth even nonnegative function supported on $[-1, 1]$. Then, letting $K = \kappa K_0 / \int_{\mathbb{R}} K_0$, we obtain a kernel K satisfying the desired conditions for any $\beta > 0$. Consider now the case $m > 0$. Let $\beta > 0$.

- If $m + d$ is even, then any K satisfying conditions A' , $B'(m - 1)$ and $C'(\beta)$ will also satisfy $B'(m)$. Indeed, as K is even, we have $\int_{\mathbb{R}} K(r)r^{m+d-1}dr = 0$, so that the induction step is proven.
- If $m + d$ is odd, let K be a kernel satisfying conditions A' , $B'(m - 1)$ and $C'(\beta/2)$. We use the following lemma.

Lemma H.1. *For $i \geq 0$, let $e_i : x \in \mathbb{R} \mapsto x^{i+d-1}$ and fix an integer $m > 0$. For any $a \in \mathbb{R}$, let F_a be the set of smooth functions $f : (1, \infty) \rightarrow \mathbb{R}$ with compact support satisfying $\int f e_i = 0$ for $0 \leq i < m$ and $\int f e_m = a$. Then,*

$$\inf \left\{ \int |f(r)|r^{d-1}dr, f \in F_a \right\} = 0. \quad (\text{H.1})$$

Assume first that the lemma is true. Let $a = -\frac{1}{2} \int_{\mathbb{R}} K(r)r^{m+d-1}$ and $f \in F_a$. Then,

$$\begin{cases} \int (K(r) + f(|r|))r^{d-1}dr = \kappa + \int f(|r|)r^{d-1}dr = \kappa \\ \int (K(r) + f(|r|))r^{i+d-1}dr = \int f(|r|)r^{i+d-1}dr = 0 \text{ for } 0 < i < m \\ \int (K(r) + f(|r|))r^{m+d-1}dr = \int K(r)r^{m+d-1}dr + 2 \int_1^\infty f(r)r^{m+d-1}dr = 0. \end{cases}$$

Hence, the kernel $K + f(|\cdot|)$ satisfies the conditions A and $B'(m)$. Also, we have, as $K(r) = 0$ if $|r| \geq 1$,

$$\begin{aligned} \int_{\mathbb{R}} (K(r) + f(|r|))_+ r^{d-1}dr &= \int_{\mathbb{R}} K(r)_+ dr + 2 \int_1^\infty f(r)_+ r^{d-1}dr \\ &\leq \beta/2 + \int_1^\infty |f(r)|r^{d-1}dr, \end{aligned}$$

where we used at the last line that $\int_1^\infty f(r)_+ r^{d-1}dr = \int_1^\infty f(r)_+ r^{d-1}dr = \frac{1}{2} \int_1^\infty |f(r)|r^{d-1}dr$. Lemma H.1 asserts the existence of $f \in F_a$ with $\int |f(r)|r^{d-1}dr \leq \beta/2$. For such a choice of f , the kernel $\tilde{K} = K + f(|\cdot|)$ satisfies also $C'(\beta)$. Finally, f has a compact support, included in $[0, R]$ for some $R > 0$. The kernel $\tilde{K}_{1/R}$ is supported on $\mathcal{B}(0, 1)$, and satisfies conditions A' , $B'(m)$ and $C'(\beta)$. This concludes the induction step, and the proof of the existence of kernels satisfying conditions A , $B(m)$ and $C(\beta)$.

Proof of Lemma H.1. Consider functions f supported on $[r_0, r_1]$ for some constants $1 < r_0 \leq r_1$ to fix. Let G_{r_0, r_1} be the subspace of $L_2([r_0, r_1])$ spanned by the functions e_i for $0 \leq i \leq m - 1$ and let g_m be the projection of e_m on G_{r_0, r_1}^\perp , the orthogonal space of G_{r_0, r_1} . Let $\ell = \|g_m\|_{L_2([r_0, r_1])}$. The function $f = \frac{a g_m}{\ell^2}$ is a polynomial of degree m restricted to $[r_0, r_1]$ and satisfies $\int f e_i = 0$ for $0 \leq i \leq m - 1$ by construction, with $\int f e_m = \frac{a}{\ell^2} \int e_m g_m = a$. Also, we have for any polynomial $P \in G_{r_0, r_1}$,

$$\begin{aligned} \|e_m - P\|_{L_2([r_0, r_1])}^2 &= \int_{r_0}^{r_1} |r^{m+d-1} - P(r)|^2 dr = \int_1^{\frac{r_1}{r_0}} r_0 |(r_0 r)^{d+m-1} - P(r r_0)|^2 dr \\ &= r_0^{2(d+m)-1} \int_1^{\frac{r_1}{r_0}} |r^{d+m-1} - r_0^{-(d+m-1)} P(r r_0)|^2 dr. \end{aligned}$$

As $r \mapsto r_0^{-(d+m-1)} P(r r_0)$ is an element of $G_{1, r_1/r_0}$, letting $r_1 = 2r_0$, we obtain

$$\begin{aligned} \ell^2 &= \|g_m\|_{L_2([r_0, r_1])}^2 = \min_{P \in G_{r_0, r_1}} \|e_m - P\|_{L_2([r_0, r_1])}^2 \\ &= r_0^{2(d+m)-1} \min_{P \in G_{1, 2}} \|e_m - P\|_{L_2([1, 2])}^2 = C r_0^{2(d+m)-1}, \end{aligned}$$

where $C = C_m > 0$ is the distance between e_m restricted to $[1, 2]$ and $G_{1, 2}$. The function f is not smooth so that it does not belong to F_a . To overcome this issue, we consider a smooth kernel ρ on \mathbb{R} satisfying $\int \rho = 1$ and $\int \rho(r)r^i dr = 0$ for $i = 1, \dots, m + d - 1$, with support included in $\mathcal{B}_{\mathbb{R}}(0, r_0/2)$. See e.g. [BH19, Section 3.2] for the construction of such a kernel ρ . The map $\rho * f$

is supported on $(1, \infty)$ and it is straightforward to check that $\rho * f \in F_a$ for $r_0 > 2$. By Young's inequality, $\|\rho * f\|_{L_2(\mathbb{R})} \leq \|\rho\|_{L_\infty(\mathbb{R})} \|f\|_{L_2(\mathbb{R})}$, so that

$$\begin{aligned} \int |\rho * f(r)| r^{d-1} dr &\leq \left(\int_{r_0/2}^{5r_0/2} r^{2d-2} dr \right)^{1/2} \|\rho * f\|_{L_2(\mathbb{R})} \\ &\leq (c_d r_0^{2d-1})^{1/2} \|\rho\|_{L_\infty(\mathbb{R})} \|f\|_{L_2(\mathbb{R})} \leq C_{d,m} a r_0^{-m}. \end{aligned}$$

By letting r_0 goes to ∞ , we see that $\inf \{ \int |f(r)| r^{d-1} dr, f \in F_a \} = 0$. \square

APPENDIX I. DETAILS ON SECTION 5.1

I.1. Optimization of convex functions on Riemannian manifolds. Let \mathcal{M} be a complete Riemannian manifold endowed with a metric $g_{\mathcal{M}}$. We write Exp_x for the exponential map at $x \in \mathcal{M}$. The geodesic distance is written as $d_{\mathcal{M}}$. We say that a set $\Omega \subset \mathcal{M}$ is geodesically convex if every geodesic joining two points of Ω is included in Ω . We will assume that Ω is small enough so that the logarithmic map Exp_x^{-1} is defined on Ω for every $x \in \Omega$. We say that a \mathcal{C}^2 function $G : \Omega \rightarrow \mathbb{R}$ is λ -strongly geodesically convex and β -smooth if $G \circ \gamma$ is λ -strongly convex and β -smooth for every unit-speed geodesic γ in Ω . In particular, this implies

$$\begin{aligned} G(y) &\geq G(x) + \langle \nabla G(x), \text{Exp}_x^{-1}(y) \rangle + \frac{\lambda}{2} d_{\mathcal{M}}(x, y)^2 \\ G(y) &\leq G(x) + \langle \nabla G(x), \text{Exp}_x^{-1}(y) \rangle + \frac{\beta}{2} d_{\mathcal{M}}(x, y)^2. \end{aligned} \tag{I.1}$$

A fundamental result of convex optimization [Udr13] states that a β -smooth and λ -strongly geodesically convex function can be optimized efficiently through a gradient descent.

Proposition I.1. *Assume that \mathcal{M} has nonnegative curvature. Let $G : \Omega \rightarrow \mathbb{R}$ be β -smooth and λ -strongly geodesically convex, with minimizer x^* . Assume that Ω contains a geodesic ball centered at x^* of radius r_0 . Fix x^0 a point in this geodesic ball and let $0 \leq \alpha \leq \beta^{-1}$.*

- (1) *The sequence of iterates $x^{a+1} = \text{Exp}_{x^a}(-\alpha \nabla G(x^a))$ is well-defined for any $a \in \mathbb{N}$.*
- (2) *The sequence of iterates satisfies*

$$d_{\mathcal{M}}(x^a, x^*)^2 \leq (1 - \lambda\alpha)^t d_{\mathcal{M}}(x^0, x^*)^2. \tag{I.2}$$

Such a result is standard, although we could not find it in this form in the literature. We provide a short proof here.

Proof. The fact that the sequence of iterates is well-defined follows from \mathcal{M} being complete, inequality (I.2), and the fact that Ω contains a geodesic ball centered at x^* . It therefore suffices to show (I.2). As the manifold \mathcal{M} has nonnegative curvature, we have

$$\begin{aligned} d_{\mathcal{M}}(x^{a+1}, x^*)^2 &\leq d_{\mathcal{M}}(x^a, x^*)^2 + d_{\mathcal{M}}(x^a, x^{a+1})^2 - 2 \langle \text{Exp}_{x^a}^{-1}(x^{a+1}), \text{Exp}_{x^a}^{-1}(x^*) \rangle \\ &\leq d_{\mathcal{M}}(x^a, x^*)^2 + \alpha^2 |\nabla G(x^a)|^2 + 2\alpha \langle \nabla G(x^a), \text{Exp}_{x^a}^{-1}(x^*) \rangle \\ &\leq (1 - \lambda\alpha) d_{\mathcal{M}}(x^a, x^*)^2 + \alpha^2 |\nabla G(x^a)|^2 + 2\alpha (G(x^*) - G(x^a)), \end{aligned}$$

where we used (I.1) at the last line. Also, we have by (I.1)

$$\begin{aligned} G(x^*) - G(x^a) &\leq G(\text{Exp}_{x^a}(-\beta^{-1} \nabla G(x^a))) - G(x^a) \\ &\leq \langle \nabla G(x^a), -\beta^{-1} \nabla G(x^a) \rangle + \frac{\beta}{2} |\nabla G(x^a)|^2 \leq -\frac{\beta}{2} |\nabla G(x^a)|^2, \end{aligned}$$

concluding the proof. \square

Proposition 5.1 that is proven just below asserts that G_m is with high probability β -smooth and λ -strongly geodesically convex with both β and λ of order ε^2 on $\Omega = \{(Q, V) \in \mathcal{M} : d_{\mathcal{O}_*(d,D)}(Q, Q^*) \leq \delta\varepsilon, \|V\|_{\text{op}} \leq \ell\}$. Our initialization point is given by $(Q^0, 0)$, where Q^0 is the output of a PCA, that satisfies with high probability $d_{\mathcal{O}_*(d,D)}(Q^0, Q^*) \leq c\varepsilon$ for some constant c . The geodesic distance between $(Q^0, 0)$ and (Q^*, V^*) is smaller than $C\varepsilon$ for some larger constant

C (using the definition of the metric (5.5)). Hence, for δ large enough, Ω contains the geodesic ball centered at (Q^*, V^*) of radius $C\varepsilon$, and we can apply Proposition I.1.

Letting $\alpha = \beta^{-1}$, the iterates of a gradient descent converge at rate

$$d_{\mathcal{M}}((Q^a, V^a), (Q^*, V^*))^2 \leq c^t d_{\mathcal{M}}((Q^0, V^0), (Q^*, V^*))^2, \quad (\text{I.3})$$

where $c \in (0, 1)$ depends on the parameter of the model.

I.2. Convexity of G_m . We prove in this section Proposition 5.1. We assume without loss of generality that $\delta \geq 1/(2\tau_{\min})$ and that $\delta \leq \ell$. Fix $(Q, V) \in \Omega$ and let $(B, W) \in T_{(Q, V)}\mathcal{M}$ be a tangent vector with unit norm. Write U for the vector space spanned by the first d columns of Q . The exponential map $\text{Exp}_{(Q, V)}$ on \mathcal{M} is given by

$$\text{Exp}_{(Q, V)}(B, W) = \left(Q \exp \begin{pmatrix} 0 & -B^\top \\ B & 0 \end{pmatrix}, V + W \right).$$

Introduce the function $F_x : t \mapsto G_{m, x}(\text{Exp}_{(Q, V)}(tB, tW))$. We denote by \mathbb{E}_N the expectation with respect to the empirical distribution associated with X_1, \dots, X_N , so that $\mathbb{E}_N F_X = \frac{1}{N} \sum_{i=1}^N F_{X_i}$. To show that G_m is geodesically λ -strongly convex and β -smooth on Ω , we need to show that

$$\lambda \leq \frac{d^2}{dt^2} \mathbb{E}_N F_X(t)|_{t=0} \leq \beta.$$

To simplify the notation, write $(Q^t, V^t) = \text{Exp}_{(Q, V)}(tB, tW)$, and let $Q^t = (e_1^t \cdots e_D^t)$. We will also write \dot{a} (resp. \ddot{a}) for the first (resp. second) time derivative of a function a evaluated at 0. Let $\mathbb{V}_j^t(x) = \iota_j(Q^t, V^t)[x^{\otimes j}]$ and let $\mathbb{V} = \sum_{j=2}^{m-1} \mathbb{V}_j$. Remark that

$$F_x = \frac{1}{2} \left(|x|^2 - \sum_{k=1}^d \langle e_k, x \rangle^2 + |\mathbb{V}(x)|^2 - 2\langle x, \mathbb{V}(x) \rangle \right). \quad (\text{I.4})$$

One can directly compute

$$\begin{aligned} \ddot{F}_x &= - \sum_{k=1}^d (\langle \dot{e}_k, x \rangle \langle e_k, x \rangle + \langle \dot{e}_k, x \rangle^2) + \langle \ddot{\mathbb{V}}(x), \mathbb{V}(x) - x \rangle + |\dot{\mathbb{V}}(x)|^2 \\ \mathbb{E}_N \ddot{F}_X &= \frac{1}{N} \sum_{i=1}^N \ddot{F}_{X_i}. \end{aligned} \quad (\text{I.5})$$

Also, we have

$$\begin{aligned} \dot{Q} &= (\dot{e}_1 \cdots \dot{e}_D) = Q \begin{pmatrix} 0 & -B^\top \\ B & 0 \end{pmatrix} = (Q_{[d, D]} B \mid -Q_{[d]} B^\top) \\ \ddot{Q} &= (\ddot{e}_1 \cdots \ddot{e}_D) = Q \begin{pmatrix} 0 & -B^\top \\ B & 0 \end{pmatrix}^2 = -(Q_{[d]} B^\top B \mid Q_{[d, D]} B B^\top). \end{aligned} \quad (\text{I.6})$$

Note that (I.6) yields the following identities: for $1 \leq k, l \leq D$,

$$\begin{cases} \langle \dot{e}_k, e_l \rangle = -\langle e_k, \dot{e}_l \rangle \text{ and } \langle \ddot{e}_k, e_l \rangle = \langle e_k, \ddot{e}_l \rangle, \\ \text{for } k \leq d, \dot{e}_k \in U^\perp \text{ and } \ddot{e}_k \in U, \\ \text{for } k > d, \dot{e}_k \in U \text{ and } \ddot{e}_k \in U^\perp. \end{cases} \quad (\text{I.7})$$

We let $\tilde{x} = Q_{[d]}^\top x \in \mathbb{R}^d$. Also, we insist on the distinction between the tensor \mathbb{V}_j (that is a tensor from \mathbb{R}^D to \mathbb{R}^D) and the tensor V_j (that is a tensor from \mathbb{R}^d to \mathbb{R}^{D-d}). The two are related by the identity $\mathbb{V}_j(x) = Q_{[d, D]} V_j[\tilde{x}^{\otimes j}]$. We will also write \mathbb{W}_j for the tensor given by $\mathbb{W}_j = \iota_j(Q, W_j)$ and let $\mathbb{W} = \sum_{j=2}^{m-1} \mathbb{W}_j$. We write $B = u \Sigma v^\top$ for the SVD of B , with u (resp. v) a $(D-d) \times d$ (resp. $d \times d$) matrix with orthogonal columns u_k (resp. v_k) and Σ a $d \times d$ diagonal matrix with nonnegative entries $\sigma_1, \dots, \sigma_d$. In particular, we have $|B|^2 = \sum_{k=1}^d \sigma_k^2$. We will use the following fact.

Lemma I.2. Let $a = (a_{d+1}, \dots, a_D) \in \mathbb{R}^{D-d}$. Then,

$$\left| \sum_{k=d+1}^D \dot{e}_k a_k \right| \leq |a| |B|. \quad (\text{I.8})$$

Proof. We have $\sum_{k=d+1}^D \dot{e}_k a_k = \dot{Q}_{[d,D]} a = Q_{[d]} v \Sigma u^\top a$. As $Q_{[d]}$ and v have orthogonal columns, the squared norm of this vector is equal to the squared norm of $\Sigma u^\top a$, that is equal to

$$\sum_{k=d+1}^D \sigma_k^2 \langle u_k, a \rangle^2 \leq |a|^2 |B|^2,$$

as each u_k is of norm 1. \square

Step 1. We first give bounds on $-\sum_{k=1}^d (\langle \ddot{e}_k, x \rangle \langle e_k, x \rangle + \langle \dot{e}_k, x \rangle^2)$. First, the dot product $|\langle \dot{e}_k, x \rangle|$ is negligible.

Lemma I.3. For $1 \leq k \leq d$, we have $|\langle \dot{e}_k, x \rangle| \leq 2|\dot{e}_k| \varepsilon^2 \delta$.

Proof. Let $x \in M$ with $|x| \leq \varepsilon$. Recall that U is the vector space spanned by $Q_{[d]}$. It holds that

$$\begin{aligned} |\langle \dot{e}_k, x \rangle| &= |\langle \dot{e}_k, \pi_U^\perp(x) \rangle| \leq |\dot{e}_k| (|\pi_U^\perp - \pi_{T_0 M}^\perp|(x)| + |\pi_{T_0 M}^\perp(x)|) \\ &\leq |\dot{e}_k| (r\varepsilon + \varepsilon^2 / (2\tau_{\min})). \end{aligned}$$

The fact that $\delta \geq 1/(2\tau_{\min})$ and that $r = \delta\varepsilon$ gives the conclusion. \square

Lemma I.3 implies that $\sum_{k=1}^d \langle \dot{e}_k, x \rangle^2 \leq 4\delta^2 |B|^2 \varepsilon^4 \leq 4\delta^2 \varepsilon^4$. Also, we have

$$\begin{aligned} -\sum_{k=1}^d \langle \ddot{e}_k, x \rangle \langle e_k, x \rangle &= -\sum_{k=1}^d x^\top \ddot{e}_k e_k^\top x = -x^\top \ddot{Q}_{[d]} Q_{[d]}^\top x \\ &= x^\top Q_{[d]} B^\top B Q_{[d]}^\top x = |B\tilde{x}|^2. \end{aligned}$$

Therefore, we may lower bound the first term in the expression of \ddot{F} :

$$-\mathbb{E}_N \sum_{k=1}^d (\langle \ddot{e}_k, X \rangle \langle e_k, X \rangle + \langle \dot{e}_k, X \rangle^2) \geq \mathbb{E}_N |B\tilde{X}|^2 - 4\delta^2 \varepsilon^4 \quad (\text{I.9})$$

Also, as $\langle e_k, X_i \rangle \leq \varepsilon$ and as (B, W) is of norm 1, we have the upper bound

$$-\mathbb{E}_N \sum_{k=1}^d (\langle \ddot{e}_k, X \rangle \langle e_k, X \rangle + \langle \dot{e}_k, X \rangle^2) \leq c_d \varepsilon^2 |B|^2 \leq c_d \varepsilon^2. \quad (\text{I.10})$$

Step 2. One can compute

$$\begin{aligned} \dot{\mathbb{V}} &= \sum_{j=2}^{m-1} \sum_{k=d+1}^D \left(\dot{e}_k \sum_{1 \leq i_1 \leq \dots \leq i_j \leq d} V_{j,k}^{i_1, \dots, i_j} e_{i_1} \otimes \dots \otimes e_{i_j} \right. \\ &\quad + e_k \sum_{1 \leq i_1 \leq \dots \leq i_j \leq d} W_{j,k}^{i_1, \dots, i_j} e_{i_1} \otimes \dots \otimes e_{i_j} \\ &\quad \left. + e_k \sum_{1 \leq i_1 \leq \dots \leq i_j \leq d} V_{j,k}^{i_1, \dots, i_j} \sum_{a=1}^j e_{i_1} \otimes \dots \otimes \dot{e}_{i_a} \otimes \dots \otimes e_{i_j} \right) \end{aligned}$$

Let us lower bound $|\dot{\mathbb{V}}(x)|^2$. As $\dot{e}_k \in U$ and $e_k \in U^\perp$ for $d+1 \leq k \leq D$,

$$\begin{aligned} |\dot{\mathbb{V}}(x)|^2 &\geq \sum_{k=d+1}^D \left(\sum_{j=2}^{m-1} \sum_{1 \leq i_1 \leq \dots \leq i_j \leq d} W_{j,k}^{i_1, \dots, i_j} \prod_{c=1}^j \langle e_{i_c}, x \rangle \right. \\ &\quad \left. + \sum_{1 \leq i_1 \leq \dots \leq i_j \leq d} V_{j,k}^{i_1, \dots, i_j} \sum_{a=1}^j \langle \dot{e}_{i_a}, x \rangle \prod_{c \neq a} \langle e_{i_c}, x \rangle \right)^2 \end{aligned}$$

$$= \sum_{k=d+1}^D (A_{1,k} + A_{2,k})^2.$$

We lower bound $(A_{1,k} + A_{2,k})^2$ by $A_{1,k}^2 - 2|A_{1,k}||A_{2,k}|$. Notice first that

$$\langle \dot{e}_{i_a}, x \rangle = \sum_{f=d+1}^D \langle \dot{e}_{i_a}, e_f \rangle \langle e_f, x \rangle = - \sum_{f=d+1}^D \langle e_{i_a}, \dot{e}_f \rangle \langle e_f, x \rangle = \langle e_{i_a}, z \rangle, \quad (\text{I.11})$$

where $z = - \sum_{f=d+1}^D \dot{e}_f \langle e_f, x \rangle = -\dot{Q}_{[d,D]} Q_{[d,D]}^\top x = Q_{[d]} B^\top Q_{[d,D]}^\top x$. We have $|z|^2 = \sum_{k=1}^d \sigma_k^2 |\tilde{y}_k|^2$, where \tilde{y}_k is the k th entry of the vector $\tilde{y} = u^\top Q_{[d,D]}^\top x \in \mathbb{R}^d$, that is equal to $\langle x, Q_{[d,D]} u_k \rangle$. As $Q_{[d,D]} u_k \in U^\perp$ and u_k is of unit norm, we have $|\tilde{y}_k| \leq c\delta\varepsilon^2$ by the same argument than in Lemma I.3. Therefore, $|z| \leq c\delta|B|\varepsilon^2$. Write $\tilde{z} = Q_{[d]}^\top z \in \mathbb{R}^d$. This implies

$$\begin{aligned} \left(\sum_{k=d+1}^D |A_{2,k}|^2 \right)^{1/2} &= \left(\sum_{k=d+1}^D \left(\sum_{j=2}^{m-1} \sum_{1 \leq i_1 \leq \dots \leq i_j \leq d} V_{j,k}^{i_1, \dots, i_j} \langle e_{i_a}, z \rangle \prod_{c \neq a} \langle e_{i_c}, x \rangle^2 \right) \right)^{1/2} \\ &= \left(\sum_{k=d+1}^D \sum_{j=2}^{m-1} V_{j,k}[\tilde{z}, \tilde{x}^{\otimes(j-1)}]^2 \right)^{1/2} \leq \sum_{j=2}^{m-1} |V_j[\tilde{z}, \tilde{x}^{\otimes(j-1)}]| \\ &\leq \ell \sum_{j=2}^{m-1} \varepsilon^{j-1} |z| \leq C_{d,m} \delta(\ell\varepsilon) \varepsilon^2 |B|. \end{aligned} \quad (\text{I.12})$$

Also, we have

$$|A_{1,k}| \leq \sum_{j=2}^{m-1} \varepsilon^j \sum_{1 \leq i_1 \leq \dots \leq i_j \leq d} |W_{j,k}^{i_1, \dots, i_j}| \leq C_{d,m} \sum_{j=2}^{m-1} \varepsilon^j |W_{j,k}|,$$

so that

$$\begin{aligned} \sum_{k=d+1}^D |A_{1,k}||A_{2,k}| &\leq C_{d,m} \left(\sum_{k=d+1}^D \left(\sum_{j=2}^{m-1} \varepsilon^j |W_{j,k}| \right)^2 \right)^{1/2} \left(\sum_{k=d+1}^D |A_{2,k}|^2 \right)^{1/2} \\ &\leq C'_{d,m} \sum_{j=2}^{m-1} \varepsilon^j |W_j| \delta(\ell\varepsilon) \varepsilon^2 |B| \leq C''_{d,m} \delta(\ell\varepsilon) (\varepsilon^2 |B|^2 + \sum_{j=2}^{m-1} \varepsilon^{2j+2} |W_j|^2) \\ &\leq C'''_{d,m} \delta(\ell\varepsilon) \varepsilon^2, \end{aligned}$$

where we used at the last line that $|B|^2 + \sum_{j=2}^{m-1} \varepsilon^{2(j-1)} |W_j|^2$ is the norm of the vector (B, W) , that we assume is equal to 1. As $\sum_{k=d+1}^D A_{1,k}^2 = |\mathbb{W}(x)|^2$, we obtain that

$$\mathbb{E}_N |\dot{\mathbb{V}}(X)|^2 \geq \mathbb{E}_N |\mathbb{W}(X)|^2 - C_{d,m} \delta(\ell\varepsilon) \varepsilon^2. \quad (\text{I.13})$$

Let us now upper bound $|\dot{\mathbb{V}}(x)|$. We have $|\sum_{k=d+1}^D \dot{e}_k V_{j,k}[\tilde{x}^{\otimes j}]|^2 = |\dot{Q}_{[d,D]} V_j[\tilde{x}^{\otimes j}]|^2$ (where $V_j[\tilde{x}^{\otimes j}]$ is the vector in \mathbb{R}^{D-d} with entries $V_{j,k}[\tilde{x}^{\otimes j}]$). Therefore,

$$\begin{aligned} \left| \sum_{k=d+1}^D \dot{e}_k V_{j,k}[\tilde{x}^{\otimes j}] \right|^2 &= |Q_{[d]} B^\top V_j[\tilde{x}^{\otimes j}]|^2 = |Q_{[d]} v \Sigma u^\top V_j[\tilde{x}^{\otimes j}]|^2 \\ &= \sum_{l=1}^d \sigma_l^2 \langle u_l, V_j[\tilde{x}^{\otimes j}] \rangle^2 \leq |B|^2 \ell^2 \varepsilon^{2j} \leq |B|^2 \ell^2 \varepsilon^4, \end{aligned}$$

where we used that each u_l is of norm 1. We therefore obtain the upper bound (recalling that $\ell \leq c\varepsilon^{-1}$ for a certain constant c)

$$\begin{aligned} \mathbb{E}_N |\dot{\mathbb{V}}(X)|^2 &\leq C_{d,m} \sum_{j=2}^{m-1} \mathbb{E}_N \left| \sum_{k=d+1}^D \dot{e}_k V_{j,k}[\tilde{X}^{\otimes j}] \right|^2 + 2 \sum_{k=d+1}^D (A_{1,k}^2 + A_{2,k}^2) \\ &\leq C'_{d,m} (|B|^2 \ell^2 \varepsilon^4 + \sum_{j=2}^{m-1} \varepsilon^{2j} |W_j|^2 + \delta^2 (\ell \varepsilon)^2 \varepsilon^4 |B|^2) \leq C''_{d,m} \varepsilon^2, \end{aligned} \quad (\text{I.14})$$

where we used that (B, W) is of norm 1.

Step 3. Eventually, we upper bound $|\ddot{\mathbb{V}}(x)|$. We first compute

$$\begin{aligned} \ddot{\mathbb{V}} &= \sum_{j=2}^{m-1} \sum_{k=d+1}^D \left(\ddot{e}_k \sum_{1 \leq i_1 \leq \dots \leq i_j \leq d} V_{j,k}^{i_1, \dots, i_j} e_{i_1} \otimes \dots \otimes e_{i_j} \right. \\ &\quad + 2\dot{e}_k \sum_{1 \leq i_1 \leq \dots \leq i_j \leq d} W_{j,k}^{i_1, \dots, i_j} e_{i_1} \otimes \dots \otimes e_{i_j} \\ &\quad + 2\dot{e}_k \sum_{1 \leq i_1 \leq \dots \leq i_j \leq d} V_{j,k}^{i_1, \dots, i_j} \sum_{a=1}^j e_{i_1} \otimes \dots \otimes \dot{e}_{i_a} \dots \otimes e_{i_j} \\ &\quad + 2e_k \sum_{1 \leq i_1 \leq \dots \leq i_j \leq d} W_{j,k}^{i_1, \dots, i_j} \sum_{a=1}^j e_{i_1} \otimes \dots \otimes \dot{e}_{i_a} \dots \otimes e_{i_j} \\ &\quad + 2e_k \sum_{1 \leq i_1 \leq \dots \leq i_j \leq d} V_{j,k}^{i_1, \dots, i_j} \sum_{a=1}^j \sum_{b>a} e_{i_1} \otimes \dots \otimes \dot{e}_{i_a} \otimes \dot{e}_{i_b} \dots \otimes e_{i_j} \\ &\quad \left. + e_k \sum_{1 \leq i_1 \leq \dots \leq i_j \leq d} V_{j,k}^{i_1, \dots, i_j} \sum_{a=1}^j e_{i_1} \otimes \dots \otimes \ddot{e}_{i_a} \dots \otimes e_{i_j} \right) \\ &= A_3 + A_4 + A_5 + A_6 + A_7 + A_8. \end{aligned}$$

- Bound on A_3 . We have

$$|A_3(x)| \leq \sum_{j=2}^{m-1} \left(\sum_{k=d+1}^D |\ddot{e}_k|^2 \right)^{1/2} |\mathbb{V}_j(x)| \leq C_{d,m} \ell |B|^2 \varepsilon^2 \leq C_{d,m} \ell \varepsilon^2.$$

- Bound on A_4 . By Lemma I.2 applied to $a = W_j[\tilde{x}^{\otimes j}] \in \mathbb{R}^{D-d}$, we have

$$\begin{aligned} |A_4(x)| &\leq 2|B| \sum_{j=2}^{m-1} \left(\sum_{k=d+1}^D W_{j,k}[\tilde{x}^{\otimes j}]^2 \right)^{1/2} \leq C_{d,m} \sum_{j=2}^{m-1} \varepsilon^j |B| |W_j| \\ &\leq C'_{d,m} \varepsilon (|B|^2 + \sum_{j=2}^{m-1} \varepsilon^{2(j-1)} |W_j|^2) \leq C'_{d,m} \varepsilon. \end{aligned}$$

Note also that

$$A_4(x) = \sum_{j=2}^{m-1} 2\dot{Q}_{[d,D]} W_j[\tilde{x}^{\otimes j}] = -2Q_{[d]} B^\top \sum_{j=2}^{m-1} W_j[\tilde{x}^{\otimes j}].$$

- Bound on A_5 . By Lemma I.2 applied to $a = (A_{2,1}, \dots, A_{2,D-d}) \in \mathbb{R}^{D-d}$ (where the $A_{2,k}$ s were introduced in Step 2) and by (I.12), we have

$$|A_5(x)| \leq 2|B| \left(\sum_{k=d+1}^D A_{2,k}^2 \right)^{1/2} \leq C_{d,m} \delta(\ell \varepsilon) \varepsilon^2 |B|^2 \leq C_{d,m} \delta(\ell \varepsilon) \varepsilon^2.$$

- Bound on A_6 . The quantity $|A_6(x)|$ is smaller than

$$\begin{aligned}
& 2 \left(\sum_{j=2}^{m-1} \sum_{k=d+1}^D \left(\sum_{1 \leq i_1 \leq \dots \leq i_j \leq d} W_{j,k}^{i_1, \dots, i_j} \sum_{a=1}^j \langle \dot{e}_{i_a}, x \rangle \prod_{c \neq a} \langle e_{i_c}, x \rangle \right)^2 \right)^{1/2} \\
& \leq 2 \left(\sum_{j=2}^{m-1} \sum_{k=d+1}^D |W_{j,k}|^2 \left(\sum_{1 \leq i_1 \leq \dots \leq i_j \leq d} \left(\sum_{a=1}^j \langle \dot{e}_{i_a}, x \rangle \prod_{c \neq a} \langle e_{i_c}, x \rangle \right)^2 \right) \right)^{1/2} \\
& \leq C_{d,m} \sum_{j=2}^{m-1} \varepsilon^{j-1} \left(\sum_{k=d+1}^D |W_{j,k}|^2 \sum_{l=1}^d \langle \dot{e}_l, x \rangle^2 \right)^{1/2} \\
& \leq C'_{d,m} \sum_{j=2}^{m-1} \varepsilon^{j-1} \left(\sum_{k=d+1}^D |W_{j,k}|^2 \delta^2 \varepsilon^4 \sum_{l=1}^d |\dot{e}_l|^2 \right)^{1/2} \quad \text{using Lemma I.3} \\
& \leq C'_{d,m} \delta \sum_{j=2}^{m-1} \varepsilon^{j+1} |B| |W_j| \leq C'_{d,m} \delta \varepsilon^2 (|B|^2 + \sum_{j=2}^{m-1} \varepsilon^{2(j-1)} |W_j|^2) \\
& \leq C'_{d,m} \delta \varepsilon^2.
\end{aligned}$$

- Bound on A_7 . Using (I.11), we obtain that $|A_7(x)|$ is smaller than

$$\begin{aligned}
& 2 \left| \sum_{j=2}^{m-1} \sum_{k=d+1}^D e_k \sum_{1 \leq i_1 \leq \dots \leq i_j \leq d} V_{j,k}^{i_1, \dots, i_j} \sum_{a=1}^j \sum_{b>a} \langle e_{i_a}, z \rangle \langle e_{i_b}, z \rangle \prod_{c \neq a, b} \langle e_{i_c}, x \rangle \right| \\
& \leq C_{d,m} \left| \sum_{j=2}^{m-1} V_j[\tilde{z}, \tilde{z}, \tilde{x}^{\otimes(j-2)}] \right| \leq C_{d,m} \ell |z|^2 \leq C'_{d,m} \delta^2 \ell \varepsilon^4 |B|^2 \leq C'_{d,m} \delta^2 \ell \varepsilon^4.
\end{aligned}$$

- Bound on A_8 . We have

$$\langle \ddot{e}_{i_a}, x \rangle = \sum_{f=1}^d \langle \ddot{e}_{i_a}, e_f \rangle \langle e_f, x \rangle = \sum_{f=1}^d \langle e_{i_a}, \ddot{e}_f \rangle \langle e_f, x \rangle = \langle e_{i_a}, y \rangle,$$

where $y = \sum_{f=1}^d \langle e_f, x \rangle \ddot{e}_f$. In particular, $|y| \leq \varepsilon \sum_{f=1}^d |\ddot{e}_f| \leq c_d \varepsilon |B|^2$. Therefore, letting $\tilde{y} = Q_{[d]}^\top y$,

$$\begin{aligned}
|A_8(x)| &= \left| \sum_{j=2}^{m-1} \sum_{k=d+1}^D e_k \sum_{1 \leq i_1 \leq \dots \leq i_j \leq d} V_{j,k}^{i_1, \dots, i_j} \sum_{a=1}^j \langle y, e_{i_a} \rangle \prod_{c \neq a} \langle e_{i_c}, x \rangle \right| \\
&\leq C_{d,m} \left| \sum_{j=2}^{m-1} V_j[\tilde{y}, \tilde{x}^{\otimes(j-1)}] \right| \leq C_{d,m} \ell \varepsilon^2 |B|^2 \leq C_{d,m} \ell \varepsilon^2.
\end{aligned}$$

Putting the different terms together, and recalling that $\ell \geq \delta$, we obtain that

$$\ddot{\Psi}(x) = -2Q_{[d]} B^\top \sum_{j=2}^{m-1} W_j[\tilde{x}^{\otimes j}] + R,$$

where R is a remainder term of norm smaller than $C_{d,m}\ell\varepsilon^2$. Also, we have $|\langle A_4(x), \mathbb{V}(x) \rangle| \leq C_{d,m}(\ell\varepsilon)\varepsilon^2$. We may therefore write

$$\begin{aligned} \langle \check{\mathbb{V}}(x), \mathbb{V}(x) - x \rangle &= 2x^\top Q_{[d]} B^\top \sum_{j=2}^{m-1} W_j[\tilde{x}^{\otimes j}] + R' \\ &= 2(B\tilde{x})^\top \sum_{j=2}^{m-1} W_j[\tilde{x}^{\otimes j}] + R', \end{aligned} \quad (\text{I.15})$$

where R' has norm smaller than $C'_{d,m}(\ell\varepsilon)\varepsilon^2$.

Step 4. Putting the lower bounds (I.9) and (I.13) together with identity (I.15), we obtain the lowerbound

$$\begin{aligned} \mathbb{E}_N \check{F}_X &\geq \mathbb{E}_N |B\tilde{X}|^2 + \mathbb{E}_N \left| \sum_{j=2}^{m-1} W_j[\tilde{X}^{\otimes j}] \right|^2 + 2 \sum_{j=2}^{m-1} \mathbb{E}_N (B\tilde{X})^\top W_j[\tilde{X}^{\otimes j}] - C_{d,m}(\ell\varepsilon)\varepsilon^2 \\ &\geq \mathbb{E}_N |B\tilde{X}|^2 + \sum_{j=2}^{m-1} |W_j[\tilde{X}^{\otimes j}]|^2 - C_{d,m}(\ell\varepsilon)\varepsilon^2. \end{aligned} \quad (\text{I.16})$$

Let us now lower bound the quantity $\mathbb{E} |B\tilde{X} + \sum_{j=2}^{m-1} W_j[\tilde{X}^{\otimes j}]|^2$, where we take the expectation with respect to the density f of the sample X_1, \dots, X_N . Letting $Y = \tilde{X}/\varepsilon$ and $Z_j = \varepsilon^{j-1}W_j$, we have

$$\mathbb{E} |B\tilde{X} + \sum_{j=2}^{m-1} W_j[\tilde{X}^{\otimes j}]|^2 = \varepsilon^2 \mathbb{E} |BY + \sum_{j=2}^{m-1} Z_j[Y^{\otimes j}]|^2,$$

where $|B|^2 + \sum_{j=2}^{m-1} |Z_j|^2 = 1$. We may decompose this expectation as

$$\varepsilon^2 \sum_{k=d+1}^D \mathbb{E} (B_k^\top Y + \sum_{j=2}^{m-1} Z_{j,k}[Y^{\otimes j}])^2.$$

We show in the next lemma that each term in the sum is larger than $c_{d,m}f_{\min}(|B_k|^2 + \sum_{j=2}^{m-1} |Z_{j,k}|^2)$. By summing over k , we obtain that the expectation is larger than $c_{d,m}f_{\min}\varepsilon^2$.

Lemma I.4. *Let S_j be a j -tensor from \mathbb{R}^d to \mathbb{R} for each $j = 1, \dots, m-1$. Then,*

$$\mathbb{E} \left[\left(\sum_{j=1}^{m-1} S_j[Y^{\otimes j}] \right)^2 \right] \geq f_{\min} c_{d,m} \sum_{j=1}^{m-1} |S_j|^2. \quad (\text{I.17})$$

Proof. The random variable \tilde{X} has entries $\langle e_k, X \rangle$ for $1 \leq k \leq d$. As (e_1, \dots, e_d) is an orthonormal basis of U that is $\delta\varepsilon$ -close from T_0M , the random variable \tilde{X} has a density lower bounded by $f_{\min}/2$ on its support, and this support contains $\mathcal{B}(0, \varepsilon/2)$. Therefore, the expectation with respect to Y is larger than

$$f_{\min} \int_{\mathcal{B}(0,1/2)} \left(\sum_{j=1}^{m-1} S_j[y^{\otimes j}] \right)^2 dy.$$

We may also write $\sum_{j=1}^{m-1} S_j[y^{\otimes j}]$ as the dot product $\langle \mathbf{S}, \mathbf{y} \rangle$, where \mathbf{S} and \mathbf{y} are vectors indexed by $\sigma \in \bigcup_{j=1}^{m-1} \{1, \dots, d\}^j$, with the entries corresponding to $\sigma = (i_1, \dots, i_j)$ given by $\mathbf{S}_\sigma = S_j^{i_1, \dots, i_j}$ and $\mathbf{y}_\sigma = \prod_{a=1}^j y_{i_a}$. Therefore, this integral is exactly equal to $\mathbf{S}^\top \mathbf{C} \mathbf{S}$, where \mathbf{C} is the matrix with entries $\mathbf{C}_{\sigma, \sigma'} = \int_{\mathcal{B}(0,1/2)} \mathbf{y}_\sigma \mathbf{y}_{\sigma'} dy$. To conclude, we need to show that this matrix is positive definite. This follows from \mathbf{C} being a Gram matrix for the L_2 dot product on $\mathcal{B}(0, 1/2)$ associated with the L_2 functions $y \mapsto \mathbf{y}_\sigma$ that are linearly independent. Therefore, we have $\mathbf{S}^\top \mathbf{C} \mathbf{S} \geq c_{d,m} |\mathbf{S}|^2 = c_{d,m} \sum_{j=1}^{m-1} |S_j|^2$. \square

Eventually, by Hoeffding's inequality, with probability at least $1 - \exp(-cN)$, the empirical expectation $\mathbb{E}_N |B\tilde{X} + \sum_{j=2}^{m-1} W_j [\tilde{X}^{\otimes j}]|^2$ is larger than $c_{d,m} f_{\min} \varepsilon^2 / 2$. From (I.16), we obtain a lower bound of order ε^2 by choosing $\ell \leq c\varepsilon$ for c small enough.

The upper bound, also of order ε^2 , is obtained by gathering the different upper bounds (I.10), (I.14) obtained in Steps 1 and 2 as well as the identity (I.15).

REFERENCES

- [Aam17] Eddie Aamari. *Vitesses de convergence en inférence géométrique*. PhD thesis, Paris Saclay, 2017.
- [AKC⁺19] Eddie Aamari, Jisu Kim, Frédéric Chazal, Bertrand Michel, Alessandro Rinaldo, and Larry Wasserman. Estimating the reach of a manifold. *Electronic Journal of Statistics*, 13(1):1359–1399, 2019.
- [AL18] Eddie Aamari and Clément Levrard. Stability and minimax optimality of tangential Delaunay complexes for manifold reconstruction. *Discrete & Computational Geometry*, 59(4):923–971, 2018.
- [AL19] Eddie Aamari and Clément Levrard. Nonasymptotic rates for manifold, tangent space and curvature estimation. *The Annals of Statistics*, 47(1):177–204, 2019.
- [Aub82] Thierry Aubin. *Nonlinear Analysis on Manifolds. Monge-Ampère Equations*. Grundlehren der mathematischen Wissenschaften. Springer New York, 1982.
- [BB00] Jean-David Benamou and Yann Brenier. A computational fluid mechanics solution to the monge-kantorovich mass transfer problem. *Numerische Mathematik*, 84:375–393, 01 2000.
- [BCH18] Gérard Besson, Gilles Courtois, and Sa'ar Hersensky. Poincaré inequality on complete Riemannian manifolds with Ricci curvature bounded below. *Mathematical Research Letters*, 25(6):1741–1769, 2018.
- [BCS10] Lorenzo Brasco, Guillaume Carlier, and Filippo Santambrogio. Congested traffic dynamics, weak flows and very degenerate elliptic equations. *Journal de mathématiques pures et appliquées*, 93(6):652–671, 2010.
- [BH19] Clément Berenfeld and Marc Hoffmann. Density estimation on an unknown submanifold. *arXiv preprint arXiv:1910.08477*, 2019.
- [BHHS21] Clément Berenfeld, John Harvey, Marc Hoffmann, and Krishnan Shankar. Estimating the reach of a manifold via its convexity defect function. *Discrete & Computational Geometry*, pages 1–36, 2021.
- [Bre03] Yann Brenier. Extended Monge-Kantorovich theory. In *Optimal transportation and applications*, pages 91–121. Springer, 2003.
- [Bre10] Haim Brezis. *Functional Analysis, Sobolev Spaces and Partial Differential Equations*. Universitext. Springer New York, 2010.
- [BRS⁺12] Sivaraman Balakrishnan, Alesandro Rinaldo, Don Sheehy, Aarti Singh, and Larry Wasserman. Minimax rates for homology inference. In *Artificial Intelligence and Statistics*, pages 64–72, 2012.
- [BS17] Tyrus Berry and Timothy Sauer. Density estimation on manifolds with boundary. *Computational Statistics & Data Analysis*, 107:1–17, 2017.
- [Bur94] Heinrich Burkhardt. Sur les fonctions de Green relatives à un domaine d'une dimension. *Bulletin de la Société Mathématique de France*, 22:71–75, 1894.
- [CGK⁺20] Galatia Cleanthous, Athanasios G. Georgiadis, Gerard Kerkycharian, Pencho Petrushev, and Dominique Picard. Kernel and wavelet density estimators on manifolds and more general metric spaces. *Bernoulli*, 26(3):1832–1862, 2020.
- [dC92] Manfredo P. do Carmo. *Riemannian Geometry*. Mathematics (Boston, Mass.). Birkhäuser, 1992.
- [DHS13] Persi Diaconis, Susan Holmes, and Mehrdad Shahshahani. Sampling from a manifold. In *Advances in modern statistical theory and applications: a Festschrift in honor of Morris L. Eaton*, pages 102–125. Institute of Mathematical Statistics, 2013.
- [Div21a] Vincent Divol. Minimax adaptive estimation in manifold inference. 2021.
- [Div21b] Vincent Divol. A short proof on the rate of convergence of the empirical measure for the wasserstein distance. *arXiv preprint arXiv:2101.08126*, 2021.
- [DSS13] Steffen Dereich, Michael Scheutzow, and Reik Schottstedt. Constructive quantization: Approximation by empirical measures. In *Annales de l'IHP Probabilités et statistiques*, volume 49, pages 1183–1203, 2013.
- [Dud69] Richard Mansfield Dudley. The speed of mean Glivenko-Cantelli convergence. *The Annals of Mathematical Statistics*, 40(1):40–50, 1969.
- [DZ01] Michel C. Delfour and Jean-Paul Zolésio. *Shapes and Geometries: Analysis, Differential Calculus, and Optimization*. Advances in Design and Control. Society for Industrial and Applied Mathematics, 2001.
- [EAS98] Alan Edelman, Tomás A Arias, and Steven T Smith. The geometry of algorithms with orthogonality constraints. *SIAM journal on Matrix Analysis and Applications*, 20(2):303–353, 1998.
- [Fed59] Herbert Federer. Curvature measures. *Transactions of the American Mathematical Society*, 93(3):418–491, 1959.
- [FG15] Nicolas Fournier and Arnaud Guillin. On the rate of convergence in Wasserstein distance of the empirical measure. *Probability Theory and Related Fields*, 162(3-4):707–738, 2015.
- [GN15] Evarist Giné and Richard Nickl. *Mathematical Foundations of Infinite-Dimensional Statistical Models*. Cambridge Series in Statistical and Probabilistic Mathematics. Cambridge University Press, 2015.

- [GPPIW12] Christopher R. Genovese, Marco Perone Pacifico, Verdinelli Isabella, and Larry Wasserman. Minimax manifold estimation. 2012.
- [GW03] Joachim Giesen and Uli Wagner. Shape dimension and intrinsic metric from samples of manifolds with high co-dimension. In *Proceedings of the nineteenth annual symposium on Computational geometry*, pages 329–337, 2003.
- [HA05] Matthias Hein and Jean-Yves Audibert. Intrinsic dimensionality estimation of submanifolds in \mathbb{R}^d . In *Proceedings of the 22nd international conference on Machine learning*, pages 289–296. ACM, 2005.
- [Hen90] Harrie Hendriks. Nonparametric estimation of a probability density on a Riemannian manifold using Fourier expansions. *The Annals of Statistics*, pages 832–849, 1990.
- [Hir96] Tsutomu Hiroshima. Construction of the Green function on Riemannian manifold using harmonic coordinates. *Journal of Mathematics of Kyoto University*, 36(1):1–30, 1996.
- [HJR93] H Hendriks, JHM Janssen, and FH Ruymgaart. Strong uniform convergence of density estimators on compact euclidean manifolds. *Statistics & probability letters*, 16(4):305–311, 1993.
- [KRW19] Jisu Kim, Alessandro Rinaldo, and Larry Wasserman. Minimax rates for estimating the dimension of a manifold. *Journal of Computational Geometry*, 10(1):42–95, 2019.
- [Lei20] Jing Lei. Convergence and concentration of empirical measures under Wasserstein distance in unbounded functional spaces. *Bernoulli*, 26(1):767–798, 2020.
- [LJM09] Anna V. Little, Yoon-Mo Jung, and Mauro Maggioni. Multiscale estimation of intrinsic dimensionality of data sets. In *2009 AAAI Fall Symposium Series*, 2009.
- [LLW07] Han Liu, John Lafferty, and Larry Wasserman. Sparse nonparametric density estimation in high dimensions using the rodeo. In *Artificial Intelligence and Statistics*, pages 283–290, 2007.
- [Lun18] Alessandra Lunardi. *Interpolation Theory*. Publications of the Scuola Normale Superiore. Scuola Normale Superiore, 2018.
- [LZZL13] Jicai Liu, Riquan Zhang, Weihua Zhao, and Yazhao Lv. A robust and efficient estimation method for single index models. *Journal of Multivariate Analysis*, 122:226–238, 2013.
- [NSW08] Partha Niyogi, Stephen Smale, and Shmuel Weinberger. Finding the homology of submanifolds with high confidence from random samples. *Discrete & Computational Geometry*, 39(1-3):419–441, 2008.
- [PC19] Gabriel Peyré and Marco Cuturi. Computational optimal transport: With applications to data science. *Foundations and Trends® in Machine Learning*, 11(5-6):355–607, 2019.
- [Pel05] Bruno Pelletier. Kernel density estimation on Riemannian manifolds. *Statistics & probability letters*, 73(3):297–304, 2005.
- [Pey18] Rémi Peyre. Comparison between W_2 distance and \dot{H}^{-1} norm, and localization of Wasserstein distance. *ESAIM. Control, Optimisation and Calculus of Variations*, 24(4), 2018.
- [PR84] Jean-Baptiste Poly and Gilles Raby. Fonction distance et singularités. *Bulletin de Sciences Mathématiques*, 108:187–195, 1984.
- [Ros70] Haskell P. Rosenthal. On the subspaces of L_p ($p > 2$) spanned by sequences of independent random variables. *Israel Journal of Mathematics*, 8(3):273–303, 1970.
- [San15] Filippo Santambrogio. *Optimal transport for applied mathematicians*. Birkäuser, NY, 2015.
- [SKM19] Hiroyuki Sato, Hiroyuki Kasai, and Bamdev Mishra. Riemannian stochastic variance reduced gradient algorithm with retraction and vector transport. *SIAM Journal on Optimization*, 29(2):1444–1472, 2019.
- [Sog17] Christopher D. Sogge. *Fourier Integrals in Classical Analysis*. Cambridge Tracts in Mathematics. Cambridge University Press, 2 edition, 2017.
- [SP18] Shashank Singh and Barnabás Póczos. Minimax distribution estimation in Wasserstein distance. *arXiv preprint arXiv:1802.08855*, 2018.
- [Tal14] M Talagrand. Upper and lower bounds of stochastic processes. *modern surveys in mathematics* 60, 2014.
- [TGHS20] Nicolás García Trillos, Moritz Gerlach, Matthias Hein, and Dejan Slepčev. Error estimates for spectral convergence of the graph laplacian on random geometric graphs toward the laplace–beltrami operator. *Foundations of Computational Mathematics*, 20(4):827–887, 2020.
- [Tib96] Robert Tibshirani. Regression shrinkage and selection via the lasso. *Journal of the Royal Statistical Society: Series B (Methodological)*, 58(1):267–288, 1996.
- [Tri92] Hans Triebel. *Theory of Function Spaces II*. Monographs in Mathematics. Springer Basel, 1992.
- [Tsy08] Alexandre Tsybakov. *Introduction to Nonparametric Estimation*. Springer Series in Statistics. Springer New York, 2008.
- [Udr13] Constantin Udriste. *Convex functions and optimization methods on Riemannian manifolds*, volume 297. Springer Science & Business Media, 2013.
- [Vil08] Cédric Villani. *Optimal transport: old and new*, volume 338. Springer Science & Business Media, 2008.
- [WB19a] Jonathan Weed and Francis Bach. Sharp asymptotic and finite-sample rates of convergence of empirical measures in Wasserstein distance. *Bernoulli*, 25(4A):2620–2648, 2019.
- [WB19b] Jonathan Weed and Quentin Berthet. Estimation of smooth densities in wasserstein distance. In *Conference on Learning Theory*, pages 3118–3119. PMLR, 2019.
- [WW20] Hau-Tieng Wu and Nan Wu. Strong uniform consistency with rates for kernel density estimators with general kernels on manifolds. *arXiv preprint arXiv:2007.06408*, 2020.

- [Yu97] Bin Yu, Assouad, Fano, and Le Cam. In *Festschrift for Lucien Le Cam*, pages 423–435. Springer, 1997.
- [ZHCG18] Emilio Zappa, Miranda Holmes-Cerfon, and Jonathan Goodman. Monte carlo on manifolds: sampling densities and integrating functions. *Communications on Pure and Applied Mathematics*, 71(12):2609–2647, 2018.