

MARKOV SEMIGROUPS ON UNITARY DUALS GENERATED BY QUANTIZED CHARACTERS

RYOSUKE SATO

ABSTRACT. In this paper, we study Markov dynamics on unitary duals of compact quantum groups. We construct such dynamics from characters of quantum groups. Then we show that the dynamics have generators, and we give an explicit formula of the generators using the representation theory. Moreover, we construct Markov dynamics on the unitary dual of an inductive limit of compact quantum groups.

1. INTRODUCTION

The purpose of this paper can be explained in two ways. The first is to investigate the relationship between stochastic processes on unitary duals of (quantum) groups and representation theory. The second is to study dual objects of stochastic processes on (quantum) groups.

If a continuous function on a topological group G is positive-definite, central, and normalized, it is called a character of G . Then the set of characters is convex, and the set \widehat{G} of extreme points is called the unitary dual of G . If G is a compact group, Fourier analysis gives a one-to-one correspondence between the set of characters of G and the set of probability measures on \widehat{G} . Moreover, we can extend this correspondence to inductive limit groups $G_\infty = \varinjlim_N G_N$ of compact groups G_N . For G_∞ there also exists a one-to-one correspondence between the set of probability measures on $\widehat{G_\infty}$ and the set of so-called *central* probability measures on the *chains* of elements in the $\widehat{G_N}$. In the case of the infinite-dimensional unitary group $U(\infty) = \varinjlim_N U(N)$, this is a one-to-one correspondence between the set of probability measures on $\widehat{U(\infty)}$ and the set of *central* probability measures on the *Gelfand–Tsetlin patterns*. Using these correspondences, we can study the characters of $U(\infty)$ in probabilistic ways. See [10], [9], [29] [31], [36], [40], ... etc. Moreover, we can extend such a probabilistic approach for characters to the inductive limits of compact *quantum* groups. See [34], [35]. If we deal with the quantum unitary groups $U_q(N)$, then there exists a one-to-one correspondence between the set of characters of $U_q(\infty)$ and the set of q^2 -*central* probability measures on the Gelfand–Tsetlin patterns. See also [17].

It is known that Gelfand–Tsetlin patterns involve several statistical mechanical interpretations. See [3], [7] and their references. Thus, a time-evolution of probability measures on the Gelfand–Tsetlin patterns (i.e., on $\widehat{U(\infty)}$) gives a time-evolution of such models in statistical mechanics. The purpose of this paper is giving a general idea to construct Markov dynamics on $\widehat{G_\infty}$ based on the representation theory of the inductive limit $G_\infty = \varinjlim_N G_N$ of compact quantum groups G_N .

For a compact quantum group G , the space $A(G)$ of matrix coefficients of finite-dimensional corepresentations of G has a Hopf-algebra structure. There are several studies about Markov dynamics on $A(G)$. See [11], [16], ... etc. The basic idea in those papers is constructing

Markov dynamics using the comultiplication of $A(G)$ and *conditionally positive* hermitian linear functionals on $A(G)$ (i.e., linear functionals such that positive on the kernel of the counit). In this paper, we will construct dual objects of such Markov dynamics.

Kac algebras are von Neumann algebras with an additional structure like Hopf algebras. Then they give one of the formulations of the duality of locally compact groups (see [12]). For a locally compact quantum group G , *Woronowicz algebras* play a similar role in their duality theory (see [26]). In the duality theory, one of the Woronowicz algebras is given by a von Neumann algebra like $L^\infty(G)$, and another one is given by the group von Neumann algebra $W^*(G)$ that is generated by the left regular representation of G . Therefore, dual objects of Markov dynamics on $A(G) \subset L^\infty(G)$ should be Markov dynamics on $W^*(G)$. In this paper, we construct (quantum) Markov dynamics on $W^*(G)$ using the comultiplication of $W^*(G)$ and conditionally positive hermitian linear functionals on $W^*(G)$. Moreover, some our dynamics induce Markov dynamics on \widehat{G} by an isomorphism between the space $\ell^\infty(\widehat{G})$ and the center $Z(W^*(G))$ of $W^*(G)$.

The first main result in the paper is Theorem 5.1. This theorem gives a construction of *continuous-time* Markov dynamics on the unitary dual \widehat{G} from characters of a compact quantum group G . Then we give an explicit formula of the generator of the dynamics. Moreover, for certain characters of $U(N)$ and $U_q(N)$, we show a determinantal formula of the generators in Theorem 6.1, 7.1. The construction here can be extended to inductive limit quantum groups of compact quantum groups. See 8.

The second main result is Theorem 10.1. In Section 10, we investigate *discrete-time* dynamics on the set of Gelfand–Tsetlin patterns. Our construction is also based on the representation theory. Indeed, it is a generalization of algebraic construction of Markov dynamics due to Kuan [24]. Then our dynamics are given by *Toeplitz-like* transition probabilities due to Borodin and Ferrari [4]. Moreover, it is an algebraic construction of the dynamics in [5].

The organization of this paper is the following: Section 2 serves to prepare necessary notations. We review the basic facts about Woronowicz algebras in Section 3. Through the end of the section and Section 4, we construct Markov dynamics on the center of general Woronowicz algebra with a counit. In Section 5, we apply the construction of Markov dynamics when a Woronowicz algebra is a group von Neumann algebra of compact quantum group. Then we prove Theorem 5.1. In Section 6 and 7, we study the concrete examples from the unitary groups $U(N)$ and the quantum unitary groups $U_q(N)$. In Section 8, we construct Markov dynamics on unitary duals of inductive limit quantum groups of compact quantum groups. In Section 9, we show that the dynamics on $\widehat{U(\infty)}$ is a Feller semigroup. We also discuss discrete-time dynamics on Gelfand–Tsetlin patterns in Section 10. In Appendix A, we review Fourier analysis of compact quantum groups to fix the necessary notations.

2. NOTATIONS

Here we prepare necessary notations in the paper.

Let M be a W^* -algebra. We denote by M_*, M_+ the predual of M and the set of positive elements in M , respectively. For a normal semi-finite weight $\varphi: M_+ \rightarrow [0, \infty]$ we define the left ideal $\mathfrak{n}_\varphi := \{x \in M \mid \varphi(x^*x) < \infty\}$. The triple $(\pi_\varphi, H_\varphi, \eta_\varphi)$ is called the GNS(Gelfand–Naimark–Segal)-triple associated with φ if (π_φ, H_φ) is a non-degenerate $*$ -representation of M and $\eta_\varphi: \mathfrak{n}_\varphi \rightarrow H_\varphi$ is a linear map with a dense range such that $\pi_\varphi(x)\eta_\varphi(y) = \eta_\varphi(xy)$ for any $x \in M$ and $y \in \mathfrak{n}_\varphi$, and $\varphi(y^*x) = \langle \eta_\varphi(x), \eta_\varphi(y) \rangle$ for any $x, y \in \mathfrak{n}_\varphi$.

For a family $\{A_\alpha\}_{\alpha \in I}$ of C^* -algebras we define $\bigoplus_{\alpha \in I} A_\alpha \subset c_0\text{-}\bigoplus_{\alpha \in I} A_\alpha \subset \ell^\infty\text{-}\bigoplus_{\alpha \in I} A_\alpha$ by

$$\begin{aligned} \bigoplus_{\alpha \in I} A_\alpha &:= \left\{ (x_\alpha)_{\alpha \in I} \in \prod_{\alpha \in I} A_\alpha \mid x_\alpha = 0 \text{ without finitely many } \alpha \in I \right\}, \\ c_0\text{-}\bigoplus_{\alpha \in I} A_\alpha &:= \left\{ (x_\alpha)_{\alpha \in I} \in \prod_{\alpha \in I} A_\alpha \mid \lim_{\alpha \in I} \|x_\alpha\|_{A_\alpha} = 0 \right\}, \\ \ell^\infty\text{-}\bigoplus_{\alpha \in I} A_\alpha &:= \left\{ (x_\alpha)_{\alpha \in I} \in \prod_{\alpha \in I} A_\alpha \mid \sup_{\alpha \in I} \|x_\alpha\|_{A_\alpha} < \infty \right\}, \end{aligned}$$

where $\lim_{\alpha \in I} \|x_\alpha\| = 0$ if for any $\epsilon > 0$ there exists a finite subset $I_\epsilon \subset I$ such that $\|x_\alpha\| < \epsilon$ for any $\alpha \in I \setminus I_\epsilon$. Then $c_0\text{-}\bigoplus_{\alpha \in I} A_\alpha$ is a C^* -algebra and $\ell^\infty\text{-}\bigoplus_{\alpha \in I} A_\alpha$ is a W^* -algebra.

For a vector space V we define the flip map $\sigma_V: V \otimes V \rightarrow V \otimes V$ by $\sigma_M(\xi \otimes \eta) = \eta \otimes \xi$ for any $\xi, \eta \in V$. We will use the leg numbering notations. Namely, for a linear map $T: V \otimes V \rightarrow V \otimes V$ we define

$$T_{12} := T \otimes \text{id}_V, \quad T_{23} := \text{id}_V \otimes T, \quad T_{13} := \sigma_{V_{23}} T_{12} \sigma_{V_{23}}$$

as linear maps on $V \otimes V \otimes V$.

For $\lambda = (\lambda_1 \geq \dots \geq \lambda_N) \in \mathbb{Z}^N$ the Schur (Laurent) polynomial $s_\lambda(z_1, \dots, z_N)$ is defined as

$$s_\lambda(z_1, \dots, z_N) = \frac{\det[z_i^{\lambda_j + N - j}]_{i,j=1}^N}{V_N(z_1, \dots, z_N)},$$

where $V_N(z_1, \dots, z_N) := \prod_{1 \leq i < j \leq N} (z_i - z_j)$. Remark that $s_\lambda(z_1, \dots, z_N)$ becomes a Laurent polynomial and it is a polynomial if $\lambda_N \geq 0$.

For a topological space X we denote by $\mathcal{B}(X)$ and $\mathcal{M}_p(X)$ its Borel σ -algebra and the set of Borel probability measures on X , respectively. For two topological spaces X, Y a *Markov kernel* Λ from Y to X is a function $\Lambda: Y \times \mathcal{B}(X) \rightarrow [0, 1]$ such that $\Lambda(y, \cdot)$ is in $\mathcal{M}_p(X)$ for any $y \in Y$ and $\Lambda(\cdot, A)$ is measurable on Y for any $A \in \mathcal{B}(X)$. Remark that Λ induces the mapping $m \in \mathcal{M}_p(Y) \mapsto m\Lambda \in \mathcal{M}_p(X)$ by $m\Lambda(A) = \int_Y m(dy)\Lambda(y, A)$ for any $A \in \mathcal{B}(X)$.

3. WORONOWICZ ALGEBRAS

Here we summarize necessary notions related to Woronowicz algebras due to Masuda and Nakagami [26]. A tuple $\mathcal{M} = (M, \delta, R, \tau, h)$ is a *Woronowicz algebra* if

- M is a W^* -algebra,
- $\delta: M \rightarrow M \otimes M$ is a *comultiplication*, i.e., it is a faithful normal unital $*$ -homomorphism such that $(\delta \otimes \text{id})\delta = (\text{id} \otimes \delta)\delta$,
- $R: M \rightarrow M$ is a *unitary antipode*, i.e., it is an involutive $*$ -anti-automorphism such that $\delta R = \sigma_M(R \otimes R)\delta$,
- $\tau = (\tau_t)_{t \in \mathbb{R}}$ is a *deformation automorphism group*, i.e., it is a one-parameter automorphism group on M such that $(\tau_t \otimes \tau_t)\delta = \delta\tau_t$ and $R\tau_t = \tau_t R$ for any $t \in \mathbb{R}$,
- h is a *left invariant Haar weight*, i.e., it is a τ -invariant faithful normal semi-finite weight on M such that $(\omega \otimes h)\delta = \omega(1)h$ for any $\omega \in M_*$, the modular automorphism groups of h and hR commute, and $(\omega \otimes h)((1 \otimes y^*)\delta(x)) = (\omega\kappa \otimes h)(\delta(y^*)(1 \otimes x))$ for any $x, y \in \mathfrak{n}_h$ and $\omega \in (M_*)_\tau$, where $\kappa := \tau_{-i/2}R$, and $(M_*)_\tau$ is the set of analytic elements with respect to the \mathbb{R} -action on M_* given by $\omega \in M_* \mapsto \omega\tau_t \in M_*$.

Remark 3.1. If we do not assume the existence of Haar weight and assume the existence of σ -weakly dense C^* -algebra A preserved by τ , then (M, A, δ, R, τ) is called a *quantum group W^* -algebra*. We introduced it in the study of inductive limits of compact quantum groups in [35]. See also [42].

Throughout the paper, we assume that h is state, i.e., $h(1) = 1$. Then h is also right-invariant, i.e., $(h \otimes \omega)\delta = \omega(1)h$ for any $\omega \in M_*$. Moreover, h automatically satisfies that $(h \otimes \omega)(\delta(y^*)x \otimes 1) = (h \otimes \omega\kappa)(y^* \otimes 1\delta(x))$ for any $x, y \in M$ and $\omega \in (M_*)_\tau$. See [26].

Let (π_h, H_h, η_h) be the GNS-triple associated with h . Then we define the so-called *multiplicative unitary* $V: H_h \otimes H_h \rightarrow H_h \otimes H_h$ by $V\eta_h(x) \otimes \xi = (\pi_h \otimes \pi_h)(\delta(x))\eta_h(1) \otimes \xi$ for any $x \in M$ and $\xi \in H_h$. It is known that $V \in B(H_h) \bar{\otimes} M$.

We define a multiplication on M_* by $\omega_1 * \omega_2 := (\omega_1 \otimes \omega_2)\delta$ for any $\omega_1, \omega_2 \in M_*$. Then M_* becomes a Banach algebra, and $(M_*)_\tau$ is a Banach subalgebra of M_* with an involution defined by $\omega^\dagger(x) := \overline{\omega(\kappa(x)^*)}$ for $x \in M$ and $\omega \in (M_*)_\tau$. We obtain a representation $(\hat{\pi}_h, H_h)$ of M_* such that $\hat{\pi}_h(\omega) := (\text{id} \otimes \omega)(V)$ for any $\omega \in M_*$. Then the restriction to $(M_*)_\tau$ is a $*$ -representation, i.e., $\hat{\pi}_h(\omega^\dagger) = \hat{\pi}_h(\omega)^*$ for any $\omega \in (M_*)_\tau$. We denote by \hat{M} the von Neumann algebra generated by $\hat{\pi}_h(M_*)$. Then $V \in \hat{M} \bar{\otimes} M$. See [26, Section 2] for more details.

It is known that \hat{M} also has a Woronowicz algebra structure by the following way: We define a comultiplication $\hat{\delta}: \hat{M} \rightarrow \hat{M} \bar{\otimes} \hat{M}$, a unitary antipode $\hat{R}: \hat{M} \rightarrow \hat{M}$, and a deformation automorphism group $\hat{\tau} = (\hat{\tau}_t)_{t \in \mathbb{R}}$ by

$$\hat{\delta}(x) = V^*(1 \otimes x)V, \quad \hat{R}(\hat{\pi}_h(\omega)) = \hat{\pi}_h(\omega R), \quad \hat{\tau}_t(\hat{\pi}_h(\omega)) = \hat{\pi}_h(\omega\tau_{-t})$$

for any $x \in \hat{M}$ and $\omega \in M_*$. Moreover, we obtain a left-invariant Haar weight \hat{h} on \hat{M} using the theory of Hilbert algebras. See [26, Section 3] for more details. Then we call $\hat{\mathcal{M}} := (\hat{M}, \hat{\delta}, \hat{R}, \hat{\tau}, \hat{h})$ the dual Woronowicz algebra of \mathcal{M} .

For a W^* -algebra N a *right action* of \mathcal{M} on N is a faithful normal unital $*$ -homomorphism $\alpha: N \rightarrow N \bar{\otimes} M$ satisfying that $(\alpha \otimes \text{id})\alpha = (\text{id} \otimes \delta)\alpha$. Then we define the fixed-point algebra

$$N^\alpha := \{x \in N \mid \alpha(x) = x \otimes 1\}.$$

A normal linear functional $\varphi \in N_*$ is α -invariant if $(\varphi \otimes \text{id})(\alpha(x)) = \varphi(x)1$ for any $x \in N$.

We define the right action α_0 of \mathcal{M} on \hat{M} by $\alpha_0(x) = V(x \otimes 1)V^*$. Remark that α_0 satisfies

$$(\hat{\delta} \otimes \text{id})\alpha_0 = \text{Ad}V_{13}(\text{id} \otimes \alpha_0)\hat{\delta} \tag{3.1}$$

since V satisfies the pentagonal relation $V_{23}V_{12} = V_{12}V_{13}V_{23}$.

Lemma 3.1. $E := (\text{id} \otimes h)\alpha_0$ is a normal conditional expectation from \hat{M} onto $Z(\hat{M}) := \hat{M} \cap \hat{M}'$.

Proof. By definition, E is normal. Moreover, for any $x \in \hat{M}$ we have

$$\begin{aligned} \alpha_0(E(x)) &= (\text{id} \otimes \text{id} \otimes h)((\alpha \otimes \text{id})(\alpha_0(x))) \\ &= (\text{id} \otimes \text{id} \otimes h)((\text{id} \otimes \delta)(\alpha_0(x))) \\ &= (\text{id} \otimes \text{id} \otimes h)(\alpha_0(x) \otimes 1) \\ &= E(x) \otimes 1. \end{aligned}$$

Thus, $E(x) \in \hat{M}^{\alpha_0} = Z(\hat{M})$ and $E(E(x)) = E(x)$. Namely, E is a conditional expectation from \hat{M} onto $Z(\hat{M})$ by [2, Theorem II.6.10.2]. \square

The following is trivial.

Lemma 3.2. If φ is α_0 -invariant, then $\varphi E = \varphi$.

For $\varphi \in \hat{M}_*$ define $T_\varphi: \hat{M} \rightarrow \hat{M}$ by $T_\varphi := (\text{id} \otimes \varphi)\hat{\delta}$. Then we have the following two lemmas.

Lemma 3.3. If φ is α_0 -invariant, then T_φ preserves the center $Z(\hat{M}) := \hat{M} \cap \hat{M}'$.

Proof. By the definition of α_0 , we have $Z(\hat{M}) = \hat{M}^{\alpha_0}$. Thus it suffices to show that T_φ preserves \hat{M}^{α_0} . By Equation (3.1), for any $x \in \hat{M}^{\alpha_0}$ we have

$$\begin{aligned} \alpha_0(T_\varphi(x)) &= \text{Ad}V((\text{id} \otimes \varphi \otimes \text{id})(\hat{\delta}(x) \otimes 1)) \\ &= \text{Ad}V((\text{id} \otimes \varphi \otimes \text{id})((\text{id} \otimes \alpha_0)(\hat{\delta}(x)))) \\ &= (\text{id} \otimes \varphi \otimes \text{id})(\text{Ad}V((\text{id} \otimes \alpha_0)(\hat{\delta}(x)))) \\ &= (\text{id} \otimes \varphi \otimes \text{id})((\hat{\delta} \otimes \text{id})(\alpha_0(x))) \\ &= T_\varphi(x) \otimes 1. \end{aligned}$$

□

Remark 3.2. There are many investigations of above transformations in probability theory and representation theory of compact (quantum) groups. See [1], [21], [22], [24], ... etc.

Lemma 3.4. If $\varphi, \phi \in \hat{M}_*$ are α_0 -invariant, then $\varphi * \phi$ is also α_0 -invariant.

Proof. By Equation (3.1), for any $x \in \hat{M}$ we have

$$\begin{aligned} (\varphi * \phi \otimes \text{id})(\alpha_0(x)) &= (\varphi \otimes \phi \otimes \text{id})((\hat{\delta} \otimes \text{id})(\alpha_0(x))) \\ &= (\varphi \otimes \phi \otimes \text{id})(\text{Ad}V((\text{id} \otimes \alpha_0)(\hat{\delta}(x)))) \\ &= (\varphi \otimes \text{id})(\text{Ad}V((\text{id} \otimes \phi \otimes \text{id})(\hat{\delta}(x) \otimes 1))) \\ &= (\varphi \otimes \text{id})(\alpha_0((\text{id} \otimes \phi)(\hat{\delta}(x)))) \\ &= (\varphi \otimes \phi)(\hat{\delta}(x))1 \\ &= (\varphi * \phi)(x)1. \end{aligned}$$

□

Since h is a state, we have $1 \in \mathfrak{n}_h$. Thus, we obtain $\hat{\epsilon} \in \hat{M}_*$ by $\hat{\epsilon}(x) := \langle x\eta_h(1), \eta_h(1) \rangle$ for any $x \in \hat{M}$. Then it is known that $\hat{\epsilon}$ becomes a counit of \hat{M} , i.e., $\hat{\epsilon}$ is multiplicative and satisfies that $(\hat{\epsilon} \otimes \text{id})\hat{\delta} = \text{id} = (\text{id} \otimes \hat{\epsilon})\hat{\delta}$. Remark that $\hat{\epsilon}(\hat{\pi}_h(\omega)) = \omega(1)$ for any $\omega \in M_*$.

Lemma 3.5. The counit $\hat{\epsilon}$ is α_0 -invariant.

Proof. By the definition of V , for any $\xi \in H_h$ we have $V(\eta_h(1) \otimes \xi) = \eta_h(1) \otimes \xi$. For any $\xi, \eta \in H_h$ we define $\omega_{\xi, \eta} \in M_*$ by $\omega_{\xi, \eta}(a) := \langle a\xi, \eta \rangle$ for any $a \in M$. Then, for any $x \in \hat{M}$

$$\begin{aligned} \omega_{\xi, \eta}((\hat{\epsilon} \otimes \text{id})(\alpha_0(x))) &= \langle V(x \otimes 1)V^*\eta_h(1) \otimes \xi, \eta_h(1) \otimes \eta \rangle \\ &= \langle x\eta_h(1) \otimes \xi, \eta_h(1) \otimes \eta \rangle \\ &= \hat{\epsilon}(x)\omega_{\xi, \eta}(1). \end{aligned}$$

Namely, we have $\omega((\hat{\epsilon} \otimes \text{id})(\alpha_0(x))) = \hat{\epsilon}(x)\omega(1)$ for any $\omega \in M_*$, i.e., $(\hat{\epsilon} \otimes \text{id})(\alpha_0(x)) = \hat{\epsilon}(x)1$. □

For $\varphi \in \hat{M}_*$ we define $\exp_*(\varphi) := \sum_{n=0}^{\infty} \varphi^{*n}/n!$, where φ^{*n} is the product $\varphi * \cdots * \varphi$ of n copies of φ and $\varphi^{*0} := \hat{\varepsilon}$. We remark that the right-hand side is the limit of $\sum_{n=0}^N \varphi^{*n}/n!$ as $N \rightarrow \infty$ in the Banach space \hat{M}_* .

Corollary 3.1. If $\varphi \in \hat{M}_*$ is α_0 -invariant, then $\exp_*(\varphi)$ is also α_0 -invariant.

Proof. Since the set of α_0 -invariant functionals is closed in the norm topology of \hat{M}_* , it follows from Lemma 3.4 \square

4. CENTER-PRESERVING, RIGHT HAAR WEIGHT INVARIANT CP_0 -SEMIGROUPS

Let M be a W^* -algebra and $(T_t)_{t \geq 0}$ a semigroup of unital completely positive σ -weakly continuous linear maps $T_t: M \rightarrow M$. If $T_0 = \text{id}$ and $\lim_{t \searrow 0} \omega(T_t(x)) = \omega(x)$ for any $x \in M$ and $\omega \in M_*$, then $(T_t)_{t \geq 0}$ is called a CP_0 -semigroup on M . For a weight ψ on M the CP_0 -semigroup $(T_t)_{t \geq 0}$ is said to be ψ -invariant if $\psi T_t = \psi$ for any $t \geq 0$.

Let $\mathcal{M} = (M, \delta, R, \tau, h)$ be a Woronowicz algebra and $\hat{\mathcal{M}} = (\hat{M}, \hat{\delta}, \hat{R}, \hat{\tau}, \hat{h})$ the dual Woronowicz algebra of \mathcal{M} . Assume that h is a state. Thus, $\hat{\mathcal{M}}$ has a counit $\hat{\varepsilon}$. A one-parameter family $(\varphi_t)_{t \geq 0}$ of functionals in \hat{M}_* is called a *convolution semigroup* if $\varphi_0 = \hat{\varepsilon}$ and $\varphi_t * \varphi_s = \varphi_{t+s}$ for any $t, s \geq 0$. If $\lim_{t \searrow 0} \|\varphi_t - \hat{\varepsilon}\| = 0$, then $(\varphi_t)_{t \geq 0}$ is said to be norm-continuous. If $\lim_{t \searrow 0} \varphi_t(x) = \hat{\varepsilon}(x)$ for any $x \in \hat{M}$, then $(\varphi_t)_{t \geq 0}$ is said to be weakly continuous.

Proposition 4.1. Let $(\varphi_t)_{t \geq 0}$ be a convolution semigroup and $T_t := (\text{id} \otimes \varphi_t)\hat{\delta}$. Then the following are equivalent:

- (1) $(\varphi_t)_{t \geq 0}$ is weakly continuous and φ_t is a state for any $t \geq 0$,
- (2) $(T_t)_{t \geq 0}$ is a CP_0 -semigroup on \hat{M} .

Proof. It is clear that (1) implies (2). For any $t \geq 0$ we have $\varphi_t = \hat{\varepsilon} * \varphi_t = \hat{\varepsilon} T_t$ since $(\hat{\varepsilon} \otimes \text{id})\hat{\delta} = \text{id}$. Thus, (2) implies (1). \square

A linear functional $\psi \in \hat{M}_*$ is said to be *conditionally positive* if $\psi(x^*x) \geq 0$ for any $x \in \text{Ker} \hat{\varepsilon}$. It is said to be hermitian if $\psi(x^*) = \overline{\psi(x)}$ for any $x \in \hat{M}_*$.

Theorem 4.1. Let $\psi \in \hat{M}_*$ be α_0 -invariant hermitian conditionally positive with $\psi(1) = 0$, $\varphi_t := \exp_*(t\psi)$, and $T_t := (\text{id} \otimes \varphi_t)\hat{\delta}$ for $t \geq 0$. Then $(T_t)_{t \geq 0}$ is a center-preserving, $\hat{h}\hat{R}$ -invariant CP_0 -semigroup.

Proof. By the Schoenberg correspondence (see [16, Section 1.2]), $(\varphi_t)_{t \geq 0}$ is a convolution semigroup of states. Moreover, φ_t is α_0 -invariant for any $t \geq 0$ by Corollary 3.1. Thus, T_t preserves the center $Z(\hat{M})$ of \hat{M} by Lemma 3.3. Since $(\hat{R} \otimes \hat{R})\hat{\delta} = \sigma_{\hat{M}}\hat{\delta}\hat{R}$, we have the right-invariance with respect to $\hat{h}\hat{R}$. Namely, $\hat{h}\hat{R}T_t = (\hat{h}\hat{R} \otimes \varphi_t)\hat{\delta} = \hat{h}\hat{R}$ for any $t \geq 0$. Moreover, $(T_t)_{t \geq 0}$ is a CP_0 -semigroup since $(\varphi_t)_{t \geq 0}$ is weakly continuous. \square

Example 4.1. Let $\varphi \in M_*$ be a α_0 -invariant state. For $c > 0$ the linear functional $\psi := c(\varphi - \hat{\varepsilon})$ is α_0 -invariant, hermitian, and conditionally positive with $\psi(1) = 0$. Thus, by Theorem 4.1, ψ generates a CP_0 -semigroup. Then its generator $(\text{id} \otimes \psi)\hat{\delta}$ is called a *Poisson generator*.

By the example, for any α_0 -invariant state we can obtain a center-preserving, right Haar weight invariant CP_0 -semigroup. As shown in the next section, the center of the group von Neumann algebra of the compact quantum group G can be identified with the space $\ell^\infty(\hat{G})$ of bounded functions on the unitary dual \hat{G} . Therefore, we obtain a Markov semigroup on \hat{G} .

5. MARKOV SEMIGROUPS ON UNITARY DUALS OF COMPACT QUANTUM GROUPS

In this section, we apply the results in above sections to Woronowicz algebras given by compact quantum groups. Then we construct Markov semigroups on unitary duals of compact quantum groups. See Theorem 5.1. We first recall basic facts about compact quantum groups and give two Woronowicz algebras for a compact quantum group.

Let $G = (C(G), \delta_G)$ be a compact quantum group, i.e., $C(G)$ is a unital C^* -algebra and $\delta_G: C(G) \rightarrow C(G) \otimes C(G)$ is a unital $*$ -homomorphism satisfying that

- $(\delta_G \otimes \text{id})\delta_G = (\text{id} \otimes \delta_G)\delta_G$ as $*$ -homomorphism from $C(G)$ to $C(G) \otimes C(G) \otimes C(G)$,
- $(C(G) \otimes 1)\delta_G(C(G)), (1 \otimes C(G))\delta_G(C(G)) \subset C(G) \otimes C(G)$ are dense,

where \otimes denotes the operation of minimal tensor product of C^* -algebras. Then there exists a unique state h_G on $C(G)$, called the *Haar state* of G , such that

$$(\omega \otimes h_G)\delta_G = (h_G \otimes \omega)\delta_G = \omega(1)h_G$$

for any $\omega \in C(G)^*$. See [27, Theorem 1.2.1]. Let $(\pi_{h_G}, L^2(G), \eta_{h_G})$ be the unique GNS-triple associated with h_G . Then we denote by $L^\infty(G)$ the von Neumann algebra generated by $\pi_{h_G}(C(G))$. In this paper, we assume that h_G is faithful and identify $C(G)$ with a subalgebra of $B(L^2(G))$. There exists a unitary operator V_G on $L^2(G) \otimes L^2(G)$ such that

$$V_G(\eta_{h_G}(a) \otimes \xi) = (\pi_{h_G} \otimes \pi_{h_G})(\delta_G(a))\xi_{h_G} \otimes \xi$$

for any $\xi \in L^2(G)$ and $a \in C(G)$, where $\xi_{h_G} := \eta_{h_G}(1)$. Then we obtain a comultiplication and Haar state of $L^\infty(G)$ by

$$\delta_G(a) := V_G(a \otimes 1)V_G^*, \quad h_G(a) := \langle a\xi_{h_G}, \xi_{h_G} \rangle$$

for any $a \in L^\infty(G)$, respectively. We remark that δ_G and h_G are the extensions of comultiplication and Haar state of $C(G)$, respectively. Thus, we use the same notations.

Let H be a finite-dimensional Hilbert space and U a *unitary corepresentation* of G on H , i.e., U is a unitary element in $B(H) \otimes C(G)$ satisfying that $(\text{id} \otimes \delta_G)(U) = U_{12}U_{13}$. For any linear functional φ on $B(H)$ we call $(\varphi \otimes \text{id})(U)$ a matrix coefficient of U . We denote by $A(G)$ the linear subspace of $C(G)$ generated by all matrix coefficients of finite-dimensional unitary corepresentations of G . Then $A(G)$ becomes $*$ -subalgebra of $C(G)$. In this paper, we assume that $C(G)$ coincides with the universal C^* -algebra generated by $A(G)$. It is known that $A(G)$ has an h_G -invariant one-parameter automorphism group $\tau = (\tau_t)_{t \in \mathbb{R}}$ and a h_G -invariant unitary antipode $R: A(G) \rightarrow A(G)$. See [27, Section 1.6, 1.7]. Thus, there exist unitary operators U_t for any $t \in \mathbb{R}$ and a conjugate linear operator \hat{J} on $L^2(G)$ satisfying

$$U_t \eta_{h_G}(a) = \eta_{h_G}(\tau_t(a)), \quad \hat{J} \eta_{h_G}(a) = \eta_{h_G}(R(a))$$

for any $a \in A(G)$. Therefore, $L^\infty(G)$ has an one-parameter automorphism group $\tau^G = (\tau_t^G)_{t \in \mathbb{R}}$ and a unitary antipode $R_G: L^\infty(G) \rightarrow L^\infty(G)$ defined by

$$\tau_t^G(x) := U_t x U_t^*, \quad R_G(x) := \hat{J} x^* \hat{J}$$

for any $x \in L^\infty(G)$. Then $\mathcal{L}^\infty(G) = (L^\infty(G), \delta_G, R_G, \tau^G, h_G)$ becomes a Woronowicz algebra and V_G is its multiplicative unitary. See [26, Section 5] for more details. We denote by $\mathcal{W}^*(G) = (W^*(G), \hat{\delta}_G, \hat{R}_G, \hat{\tau}_G, \hat{h}_G)$ the dual Woronowicz algebra. Remark that $\mathcal{W}^*(G)$ has a counit, denoted by $\hat{\epsilon}_G$, since h_G is bounded.

Let \widehat{G} be the set of all equivalence classes of irreducible unitary corepresentations of G . For any $\alpha \in \widehat{G}$ we fix a representative (U_α, H_α) of α . We denote by $\{f_z^G\}_{z \in \mathbb{C}}$ the *Woronowicz characters*, which are distinguished multiplicative states on $A(G)$. See [27, Section 1.7]. We define $F_\alpha^z := (\text{id} \otimes f_z^G)(U_\alpha)$ for any $z \in \mathbb{C}$ and $d_q(\alpha) := \text{Tr}_{H_\alpha}(F_\alpha)$. Then there exists a $*$ -isomorphism $\Phi_G: W^*(G) \cong \ell^\infty\text{-}\bigoplus_{\alpha \in \widehat{G}} B(H_\alpha)$. Moreover, the deformation automorphism group $\hat{\tau}^G$ on $W^*(G)$ is given by $\hat{\tau}_t^G = \Phi_G^{-1} \circ (\prod_{\alpha \in \widehat{G}} \text{Ad} F^{\text{it}}) \circ \Phi_G$ for any $t \in \mathbb{R}$.

The following is our formulation of quantization of characters in [34].

Definition 5.1. A normal $\hat{\tau}^G$ -KMS (Kubo–Martin–Schwinger) state χ on $W^*(G)$ with inverse temperature -1 is called a *quantized character* of G . Let $\text{Ch}(G)$ be the set of all quantized characters of G .

Remark 5.1. The $\text{Ch}(G)$ is a convex set and its extreme points $\text{ex}(\text{Ch}(G))$ coincides with \widehat{G} as sets. More precisely, for any $\alpha \in \widehat{G}$ the corresponding extreme quantized character χ_α is given by $\chi_\alpha(x) = \text{Tr}_{H_\alpha}(F_\alpha x_\alpha)/d_q(\alpha)$, where $x = \Phi_G^{-1}((x_\beta)_{\beta \in \widehat{G}}) \in W^*(G)$. For the trivial corepresentation $U_{tr} = 1 \otimes 1_{C(G)}$ we have the associated quantized character χ_{tr} and $\chi_{tr} = \hat{\epsilon}_G$. Moreover, every quantized character is a convex combination of extreme ones (see [34, Lemma 2.2]). Thus, there is an affine bijection between $\text{Ch}(G)$ and $\mathcal{M}_p(\widehat{G})$.

Recall that the right action $\alpha_0: W^*(G) \rightarrow W^*(G) \bar{\otimes} L^\infty(G)$ of $\mathcal{L}^\infty(G)$ on $W^*(G)$ is defined by $\alpha_0(x) := V_G(x \otimes 1)V_G^*$ for any $x \in W^*(G)$. Then we obtain the following lemma by [21, Lemma 2.1.(4)].

Lemma 5.1. Any quantized character $\chi \in \text{Ch}(G)$ is α_0 -invariant.

Corollary 5.1. Let $\chi \in \text{Ch}(G)$ and $\varphi_t := \exp_*(t(\chi - \hat{\epsilon}_G))$ for any $t \geq 0$. Then the CP_0 -semigroup $(T_t)_{t \geq 0}$ defined by $T_t := (\text{id} \otimes \varphi_t)\hat{\delta}_G$ is center-preserving and $\hat{h}_G \hat{R}_G$ -invariant.

Proof. It follows from Lemma 5.1, Lemma 3.5 and Theorem 4.1. \square

This corollary induces Markov dynamics on \widehat{G} . The following gives another algebraic interpretation of those dynamics on \widehat{G} . See also Proposition 5.1.

Lemma 5.2. Let $\chi \in \text{Ch}(G)$ and $\varphi_t := \exp_*(t(\chi - \hat{\epsilon}_G))$ for any $t \geq 0$. Then $\varphi_t \in \text{Ch}(G)$.

Proof. We fix two elements $x, y \in W^*(G)$. Since χ^{*k} is a $\hat{\tau}^G$ -KMS state for any $k \geq 0$ (see [35, Lemma 3.2]), there exists a bonded continuous function F_k on $\bar{D} = \{z \in \mathbb{C} \mid -1 \leq \text{Im}(z) \leq 0\}$, which is analytic in $D = \{z \in \mathbb{C} \mid -1 < \text{Im}(z) < 0\}$ such that for any $s \in \mathbb{R}$

$$F_k(s) = \chi^{*k}(x \hat{\tau}_s^G(y)), \quad F_k(s - i) = \chi^{*k}(\hat{\tau}_s^G(y)x).$$

By [8, Proposition 5.3.7], we have $|F_k(z)| < \|x\| \|y\|$ on \bar{D} . Thus,

$$F := \sum_{n=0}^{\infty} \frac{t^n}{n!} \sum_{k=0}^n (-1)^k \binom{n}{k} F_k.$$

converges uniformly in \bar{D} . Then F satisfies that $|F| \leq e^{2t} \|x\| \|y\|$ on \bar{D} and analytic in D . Moreover, since $(\chi - \hat{\epsilon}_G)^{*n} = \sum_{k=0}^n (-1)^k \binom{n}{k} \chi^{*k}$, we have

$$F(s) = \varphi_t(x \hat{\tau}_s^G(y)), \quad F(s - i) = \varphi_t(\hat{\tau}_s^G(y)x).$$

Namely, φ_t is a normal $\hat{\tau}^G$ -KMS state, i.e., $\varphi_t \in \text{Ch}(G)$. \square

We define $\pi_G: L^\infty(G)_* \rightarrow W^*(G)$ by $\pi_G(\omega) = (\text{id} \otimes \omega)(V_G)$ for any $\omega \in L^\infty(G)_*$. Then for any $\varphi \in W^*(G)_*$ we define $\pi_{G^*}(\varphi) \in L^\infty(G)$ by $\omega(\pi_{G^*}(\varphi)) = \varphi(\pi_G(\omega))$ for any $\omega \in L^\infty(G)$. Namely, $\pi_{G^*}(\varphi) = (\varphi \otimes \text{id})(V_G)$. Since $\pi_{G^*}: W^*(G)_* \rightarrow L^\infty(G)$ is a homomorphism, we have the following lemma.

Lemma 5.3. For any $\psi \in W^*(G)_*$ we have $\pi_{G^*}(\exp_*(\psi)) = e^{\pi_{G^*}(\psi)} = \sum_{n=0}^{\infty} \pi_{G^*}(\psi)^n/n!$.

Let $T_\varphi := (\text{id} \otimes \varphi)\hat{\delta}_G$ for any $\varphi \in W^*(G)_*$. For any $\omega \in L^\infty(G)_*$ and $a \in L^\infty(G)$ we define $a\omega, \omega a \in L^\infty(G)_*$ by $[a\omega](b) := \omega(ba)$ and $[\omega a](b) := \omega(ab)$ for any $b \in L^\infty(G)$.

Lemma 5.4. For any $\varphi \in W^*(G)_*$ and $\omega \in L^\infty(G)_*$ we have $T_\varphi(\pi_G(\omega)) = \pi_G(\omega\pi_{G^*}(\varphi))$.

Proof. By the pentagonal relation of V_G , we have $(\hat{\delta}_G \otimes \text{id})(V_G) = V_{G13}V_{G23}$. Thus,

$$T_\varphi(\pi_G(\omega)) = (\text{id} \otimes \varphi \otimes \omega)(V_{G13}V_{G23}) = (\text{id} \otimes \omega)(1 \otimes \pi_{G^*}(\varphi)V_G) = \pi_G(\omega\pi_{G^*}(\varphi)).$$

□

By Corollary 5.1, for any $\chi \in \text{Ch}(G)$ we obtain the center-preserving CP_0 -semigroup on $W^*(G)$. Now we study the restriction of such CP_0 -semigroups to the center $Z(W^*(G))$. For any $\xi \in L^2(G)$ and $\alpha \in \hat{G}$ we define a linear operator $\hat{\xi}(\alpha)$ on H_α by $\hat{\xi}(\alpha) := (\text{id} \otimes \omega_\xi)(U_\alpha)^*$, where $\omega_\xi \in C(G)^*$ is defined by $\omega_\xi(a) := \langle \eta_G(a), \xi \rangle$ for any $a \in C(G)$. Then the Peter–Weyl theorem (see Theorem A.1) states that for any $\xi, \eta \in L^2(G)$

$$\langle \xi, \eta \rangle = \sum_{\alpha \in \hat{G}} d_q(\alpha) \text{Tr}_{H_\alpha}(F_\alpha \hat{\eta}(\alpha)^* \hat{\xi}(\alpha)).$$

Moreover, there exists a $*$ -isomorphism $\Psi_G: \ell^\infty(\hat{G}) \rightarrow Z(W^*(G))$ such that

$$\langle \Psi_G(\kappa)\xi, \eta \rangle = \sum_{\alpha \in \hat{G}} d_q(\alpha) \kappa(\alpha) \text{Tr}_{H_\alpha}(F_\alpha \hat{\eta}(\alpha)^* \hat{\xi}(\alpha))$$

for any $\xi, \eta \in L^2(G)$ and $\kappa \in \ell^\infty(\hat{G})$.

By Lemma 3.3, the linear map $T_\varphi := (\text{id} \otimes \varphi)\hat{\delta}_G$ preserves $Z(W^*(G))$ if φ is a α_0 -invariant state. Thus, we have a Markov operator $Q_\varphi := \Psi_G^{-1}T_\varphi|_{Z(W^*(G))}\Psi_G$ on $\ell^\infty(\hat{G})$. Then we define the mapping $P \in \mathcal{M}_p(\hat{G}) \rightarrow PQ_\varphi \in \mathcal{M}_p(\hat{G})$ by $(PQ_\varphi, \kappa) = (P, Q_\varphi\kappa)$ for any $\kappa \in \ell^\infty(\hat{G})$, where (\cdot, \cdot) is the natural pairing of $\mathcal{M}_p(\hat{G})$ and $\ell^\infty(\hat{G})$. Recall that there exists an affine homeomorphism from $\mathcal{M}_p(\hat{G})$ to $\text{Ch}(G)$ by irreducible decomposition. More precisely, for any $P \in \mathcal{M}_p(\hat{G})$ the corresponding quantized character $\chi_P \in \text{Ch}(G)$ is given as $\chi_P = \sum_{\alpha \in \hat{G}} P(\alpha)\chi_\alpha$. The following describes the dynamics Q_φ on \hat{G} in terms of quantized characters.

Proposition 5.1. Let $\varphi \in W^*(G)_*$ be a α_0 -invariant state and $P \in \mathcal{M}_p(\hat{G})$. Then $\chi_{PQ_\varphi} = \chi_P * \varphi$.

Proof. Since $\chi_{PQ_\varphi}, \chi_P * \varphi$ are α_0 -invariant, by Lemma 3.2, it suffices to show that $\chi_{PQ_\varphi} = \chi_P * \varphi$ on $Z(W^*(G))$. For any $\alpha \in \hat{G}$ we have $\chi_P * \varphi(\Psi_G(\delta_\alpha)) = \chi_P(\Psi_G(Q_\varphi\delta_\alpha)) = \chi_{PQ_\varphi}(\Psi_G(\delta_\alpha))$. Since $\chi_{PQ_\varphi}, \chi_P * \varphi$ are normal, they are equal each other on $Z(W^*(G))$. Thus, by Lemma 3.2, $\chi_{PQ_\varphi} = \chi_P * \varphi$ on $W^*(G)$. □

Remark 5.2. Let $tr \in \widehat{G}$ be the equivalence class of the trivial corepresentation of G . Then, by the above proposition, we have $\varphi = \chi_{\delta_{tr}Q}$ for any α_0 -invariant state. In particular, any α_0 -invariant states are quantized character of G . This is a natural characterization of quantized characters since the action α_0 is an analog of the adjoint action of group on itself (see [12]).

Remark 5.3. For any $\alpha \in \widehat{G}$ we define $X_\alpha := \pi_{G^*}(\chi_\alpha)$. By Corollary A.1, we have $X_\alpha = (\chi_\alpha^0 \otimes \text{id})(U_\alpha)$, where $\chi_\alpha^0 \in B(H_\alpha)^*$ is given as $\chi_\alpha^0(A) := \text{Tr}_{H_\alpha}(F_\alpha A)/d_q(\alpha)$. Then, by [27, Theorem 1.4.2], for any $\alpha, \beta \in \widehat{G}$ we can show that

$$[h_G X_\alpha^*](X_\beta) = \frac{\delta_{\alpha,\beta}}{d_q(\beta)^2}, \quad \pi_\beta(h_G X_\alpha^*) = \delta_{\alpha,\beta} \frac{1}{d_q(\beta)^2} 1_{H_\beta},$$

where $\pi_\beta(\omega) := (\text{id} \otimes \omega)(U_\beta)$ for any $\omega \in L^\infty(G)_*$. Therefore, by the identification of $W^*(G)$ and $\ell^\infty\text{-}\bigoplus_{\alpha \in \widehat{G}} B(H_\alpha)$ (see Appendix A), if $\omega = \sum_{\alpha \in \widehat{G}} d_q(\alpha)^2 \kappa(\alpha) h_G X_\alpha^*$ in $L^\infty(G)_*$, then $\kappa \in \ell^\infty(\widehat{G})$ and

$$\begin{aligned} \langle \Psi_G(\kappa)\xi, \eta \rangle &= \sum_{\alpha \in \widehat{G}} d_q(\alpha) \kappa(\alpha) \text{Tr}_{H_\alpha}(F_\alpha \hat{\eta}(\alpha)^* \hat{\xi}(\alpha)) \\ &= \sum_{\alpha \in \widehat{G}} d_q(\alpha) \text{Tr}_{H_\alpha}(F_\alpha \hat{\eta}(\alpha)^* \widehat{\pi_G(\omega)} \xi(\alpha)) \\ &= \langle \pi_G(\omega)\xi, \eta \rangle. \end{aligned}$$

Namely, we have $\Psi_G(\kappa) = \pi_G(\omega)$.

Lemma 5.5. Let $\varphi \in W^*(G)_*$ be α -invariant. Then $Q_\varphi(\delta_\beta)(\alpha) = d_q(\beta)^2 h_G(X_\beta^* X_\alpha \pi_{G^*}(\varphi))$ for any $\alpha, \beta \in \widehat{G}$.

Proof. By Proposition 5.1 and Remark 5.3,

$$Q_\varphi(\delta_\beta)(\alpha) = (\chi_\alpha * \varphi)(\Psi_G(\delta_\beta)) = d_q(\beta)^2 (\chi_\alpha * \varphi)(\pi_G(h_G X_\beta^*)) = d_q(\beta)^2 h_G(X_\beta^* X_\alpha \pi_{G^*}(\varphi)).$$

□

By the above lemma, the action Q_φ on $\ell^\infty(\widehat{G})$ coincides with the action of the Markov kernel \mathbb{Q}_φ defined by $\mathbb{Q}_\varphi(\alpha, \beta) = d_q(\beta)^2 h_G(X_\beta^* X_\alpha \pi_{G^*}(\varphi))$.

Lemma 5.6. For any $\omega \in L^\infty(G)_*$ it holds true that $\lim_{\alpha \in \widehat{G}} \|\pi_\alpha(\omega)\| = 0$. In particular, $\kappa \in c_0(\widehat{G})$ if $\Psi_G(\kappa) = \pi_G(\omega)$ for some $\omega \in L^\infty(G)_*$.

Proof. By [2, Theorem III 2.1.4], there exist sequences $(\xi_n)_{n=1}^\infty, (\eta_n)_{n=1}^\infty$ in $L^2(G)$ such that $\sum_{n=1}^\infty \|\xi_n\|_2^2 < \infty$, $\sum_{n=1}^\infty \|\eta_n\|_2^2 < \infty$, and $\omega(x) = \sum_{n=1}^\infty \langle x \xi_n, \eta_n \rangle$ for any $x \in L^\infty(G)$. For any $\epsilon > 0$ we fix sufficiently large N such that

$$\left(\sum_{n \geq N+1} \|\xi_n\|_2^2 \right)^{1/2} \left(\sum_{n \geq N+1} \|\eta_n\|_2^2 \right)^{1/2} < \epsilon/2.$$

Since $\{\eta_G(a) \mid a \in A(G)\}$ is dense in $L^2(G)$, there exist $a_n, b_n \in A(G)$ such that

$$\|\xi_n - \eta_G(a_n)\|_2 < \frac{\epsilon}{4N\|\eta_n\|_2}, \quad \|\eta_n - \eta_G(b_n)\|_2 < \frac{\epsilon}{4N\|\eta_G(a_n)\|_2}.$$

Then $\omega_\epsilon := \sum_{n=1}^N b_n^* h_G a_n$ satisfies that $\|\omega - \omega_\epsilon\|_1 < \epsilon$. Moreover, we have

$$\|\Phi_G(\pi_G(\omega)) - \Phi_G(\pi_G(\omega_\epsilon))\| \leq \|\omega - \omega_\epsilon\|_1 < \epsilon.$$

Since $a_n, b_n \in A(G)$, we have

$$\Phi_G(\pi_G(\omega_\epsilon)) \in \bigoplus_{\alpha \in \widehat{G}} B(H_\alpha) \subset c_0\text{-}\bigoplus_{\alpha \in \widehat{G}} B(H_\alpha).$$

Therefore, we have $\lim_{\alpha \in \widehat{G}} \|\pi_\alpha(\omega)\| = 0$ since $c_0\text{-}\bigoplus_{\alpha \in \widehat{G}} B(H_\alpha)$ is norm-closed.

The rest of the statement immediately follows from Remark 5.3. \square

Here we show the first main theorem.

Theorem 5.1. Let $\chi \in \text{Ch}(G)$, $\chi_t := \exp_*(t(\chi - \hat{\epsilon}_G))$, and $T_t^X := (\text{id} \otimes \chi_t) \hat{\delta}_G$ for any $t \geq 0$. Then the Markov kernels \mathbb{Q}_t^X associated with $Q_t^X := \Psi_G^{-1} T_t^X|_{Z(W^*(G))} \Psi_G$ form a Markov semigroup on \widehat{G} . Moreover, $(\mathbb{Q}_t^X)_{t \geq 0}$ is a Feller semigroup and its generator \mathbb{L}_χ is given by

$$\mathbb{L}_\chi(\alpha, \beta) := d_q(\beta)^2 h_G(X_\beta^* X_\alpha (\pi_{G^*}(\chi) - 1))$$

for any $\alpha, \beta \in \widehat{G}$. In particular, the generator \mathbb{L}_χ is bounded.

Proof. First we show that $(\mathbb{Q}_t^X)_{t \geq 0}$ is a Feller semigroup, i.e., $(\mathbb{Q}_t^X)_{t \geq 0}$ preserves $c_0(\widehat{G})$. Let $\mathcal{A} := \left\{ \kappa \in c_0(\widehat{G}) \mid \sum_{\alpha \in \widehat{G}} d_q(\alpha) |\kappa(\alpha)| < \infty \right\}$. Since $c_0(\widehat{G})$ is norm-closed and \mathcal{A} is dense in $c_0(\widehat{G})$ by the Stone–Weierstrass theorem, it suffices to show that $\mathbb{Q}_t^X \kappa \in c_0(\widehat{G})$ for any $t \geq 0$ and $\kappa \in \mathcal{A}$. Since $\|d_q(\alpha) h_G X_\alpha^*\|_1 \leq \|\eta_G(d_q(\alpha) X_\alpha)\| = 1$, for $\kappa \in \mathcal{A}$ the functional $\omega_\kappa := \sum_{\alpha \in \widehat{G}} d(\alpha)^2 \kappa(\alpha) h_G X_\alpha^*$ converges in $L^1(G)_*$. Then, by Remark 5.3 and Lemma 5.4, 5.6, we have $\mathbb{Q}_t^X \kappa = \Psi_G^{-1} \pi_G(\omega_\kappa \pi_{G^*}(\chi_t)) \in c_0(\widehat{G})$.

By Lemma 5.3, 5.5, for any $\alpha, \beta \in \widehat{G}$ we have

$$\begin{aligned} \frac{1}{t} (\mathbb{Q}_t^X \delta_\beta - \delta_\beta)(\alpha) &= d_q(\beta)^2 h_G(X_\beta^* X_\alpha \frac{e^{t(\pi_G(\chi) - 1)}}{t}) \\ &\rightarrow d_q(\beta)^2 h_G(X_\beta^* X_\alpha (\pi_G(\chi) - 1)) \quad (t \searrow 0). \end{aligned}$$

Therefore, the generator \mathbb{L}_χ of $(\mathbb{Q}_t^X)_{t > 0}$ is given as $\mathbb{L}_\chi(\alpha, \beta) := d(\beta)^2 h_G(X_\beta^* X_\alpha (\pi_{G^*}(\chi) - 1))$ for any $\alpha, \beta \in \widehat{G}$. In particular, $\mathbb{L}_\chi = \mathbb{Q}_\chi - \text{id}$, that is, \mathbb{L}_χ is bounded, \square

By [35, Lemma 3.2], $\chi_\alpha * \chi_\gamma$ is also quantized character of G for any $\alpha, \gamma \in \widehat{G}$. We define $N_{\alpha, \gamma}^\beta$ by $(d_q(\alpha) \chi_\alpha) * (d_q(\gamma) \chi_\gamma) = \sum_{\beta \in \widehat{G}} N_{\alpha, \gamma}^\beta d_q(\beta) \chi_\beta$ be its irreducible decomposition.

Corollary 5.2. Assume that $\chi \in \text{Ch}(G)$ has an irreducible decomposition $\chi = \sum_{\alpha \in \widehat{G}} P_\chi(\alpha) \chi^\alpha$. Then the generator \mathbb{L}_χ of the Feller semigroup in Theorem 5.1 is given by

$$\mathbb{L}_\chi(\alpha, \beta) = \frac{d_q(\beta)}{d_q(\alpha)} \sum_{\gamma \in \widehat{G}} \frac{P_\chi(\gamma) N_{\alpha, \gamma}^\beta}{d_q(\gamma)} - \delta_{\alpha, \beta}.$$

Proof. By Theorem 5.1 and Proposition 5.1, we have

$$\mathbb{L}_\chi(\alpha, \beta) = \mathbb{Q}_\chi(\alpha, \beta) - \delta_{\alpha, \beta} = \chi_\alpha * \chi(\Psi_G(\delta_\beta)) - \delta_{\alpha, \beta} = \frac{d_q(\beta)}{d_q(\alpha)} \sum_{\gamma \in \widehat{G}} \frac{P_\chi(\gamma) N_{\alpha, \gamma}^\beta}{d_q(\gamma)} - \delta_{\alpha, \beta}.$$

\square

If \widehat{G} is separable, we have the following by [14, Theorem 4.2.7].

Corollary 5.3. For any probability measure P on \widehat{G} and $\chi \in \text{Ch}(G)$ there exists a Markov process $(X_t)_{t \geq 0}$ corresponding to $(\mathbb{Q}_t^X)_{t \geq 0}$ with initial distribution P and càdlàg sample paths.

6. EXAMPLES FROM UNITARY GROUPS $U(N)$

In this section, we apply the above results to the case of unitary groups $U(N)$ and certain characters. When we deal with an ordinary topological group G , quantized characters can be regarded as functions on G . A continuous function χ on G is called a *character* of G if χ is positive-definite, $\chi(gh) = \chi(hg)$ for any $g, h \in G$, and $\chi(e) = 1$. We equip the set of characters with the topology of uniform convergence on compact subsets. Then the set of characters is homeomorphic to the set of quantized characters (i.e., the set of normal tracial states on $W^*(U(\infty))$). See [35, Remark 3.2].

Here we recall some results of the representation theory of $U(N)$. It is well known that $\widehat{U(N)}$ can be identified with $\mathbb{S}_N := \{\lambda = (\lambda_1 \geq \dots \geq \lambda_N) \in \mathbb{Z}^N\}$. Then for any $\lambda \in \mathbb{S}_N$ the corresponding irreducible character χ_λ of $U(N)$ is given by the Schur polynomial s_λ , i.e.,

$$\chi_\lambda(U) = \frac{s_\lambda(z_1, \dots, z_N)}{s_\lambda(1, \dots, 1)},$$

where z_1, \dots, z_N are eigenvalues of $U \in U(N)$. See e.g. [43].

We define the *infinite-dimensional unitary group* $U(\infty)$ as the inductive limit $\varinjlim_N U(N)$ by natural embeddings

$$U \in U(N) \hookrightarrow \begin{bmatrix} U & 0 \\ 0 & 1 \end{bmatrix} \in U(N+1).$$

Then the theorem due to Voiculescu [41], Vershik–Kerov [40], and Boyer [10] states that any extreme characters of $U(\infty)$ are parametrized by the set $\Omega \subset \mathbb{R}_{\geq 0}^\infty \times \mathbb{R}_{\geq 0}^\infty \times \mathbb{R}_{\geq 0}^\infty \times \mathbb{R}_{\geq 0}^\infty \times \mathbb{R}_{\geq 0} \times \mathbb{R}_{\geq 0}$ consisting of $\omega = (\alpha^+, \beta^+, \alpha^-, \beta^-, \delta^+, \delta^-)$ such that

$$\begin{aligned} \alpha^\pm &= (\alpha_1^\pm \geq \alpha_2^\pm \geq \dots), & \beta^\pm &= (\beta_1^\pm \geq \beta_2^\pm \geq \dots), \\ \sum_{i=1}^{\infty} (\alpha_i^\pm + \beta_i^\pm) &< \infty, & \beta_1^+ + \beta_1^- &\leq 1, & \gamma^\pm &:= \delta^\pm - \sum_{i=1}^{\infty} (\alpha_i^\pm + \beta_i^\pm) \geq 0. \end{aligned}$$

More precisely, for any $\omega \in \Omega$ the corresponding extreme character χ_ω is given by

$$\chi_\omega(U) = \prod_z \Phi_\omega(z), \tag{6.1}$$

where z runs over all any eigenvalues of $U \in U(\infty)$, which are equal to 1 except for finitely many eigenvalues, and

$$\Phi_\omega(z) := e^{\gamma^+(z-1) + \gamma^-(z^{-1}-1)} \prod_{i=1}^{\infty} \frac{1 + \beta_i^+(z-1)}{1 - \alpha_i^+(z-1)} \frac{1 + \beta_i^-(z^{-1}-1)}{1 - \alpha_i^-(z^{-1}-1)}.$$

Then we define $\varphi_\omega(n)$ by $\Phi_\omega(z) = \sum_{n \in \mathbb{Z}} \varphi_\omega(n) z^n$, where the expansion converges in a neighborhood of $\mathbb{T} := \{z \in \mathbb{C} \mid |z| = 1\}$. It is well known that $(\varphi_\omega(n))_{n \in \mathbb{Z}}$ is totally positive, that is, for any $N \geq 1$ and $m_1 > \dots > m_N$ the determinant $\det[\varphi_\omega(m_i + j)]_{i,j=1}^N$ is nonnegative.

Theorem 6.1. For any $\omega \in \Omega$ let χ_N be the restriction of χ_ω to $U(N)$. Then the generator \mathbb{L}_{χ_N} of Markov semigroup associated with χ_N in Theorem 5.1 is given by

$$\mathbb{L}_{\chi_N}(\lambda, \mu) = \frac{s_\mu(1, \dots, 1)}{s_\lambda(1, \dots, 1)} \det[\varphi_\omega(\mu_j - j - \lambda_i - i)]_{i,j=1}^N - \delta_{\lambda, \mu}$$

for any $\lambda, \mu \in \mathbb{S}_N$, where $s_\lambda(1, \dots, 1)$ is the specialization of the Schur polynomial s_λ with N variables.

Proof. Remark that the dimension $d(\lambda)$ of irreducible representation associated with $\lambda \in \mathbb{S}_N$ coincides with $s_\lambda(1, \dots, 1)$. Thus, by Corollary 5.2,

$$\mathbb{L}_{\chi_N}(\lambda, \mu) = \frac{s_\mu(1, \dots, 1)}{s_\lambda(1, \dots, 1)} \frac{1}{N!} \int_{\mathbb{T}^N} \chi_N(\text{diag}(z_1, \dots, z_N)) s_\lambda(z) \overline{s_\mu(z)} |V_N(z)|^2 \prod_{i=1}^N \frac{dz_i}{2\pi i z_i} - \delta_{\lambda, \mu},$$

where $\text{diag}(z_1, \dots, z_N)$ is the diagonal matrix whose entries are z_1, \dots, z_N , $s_\lambda(z) = s_\lambda(z_1, \dots, z_N)$ and $V_N(z) = V_N(z_1, \dots, z_N)$. By Equation (6.1), we have

$$\begin{aligned} \mathbb{L}_{\chi_N}(\lambda, \mu) &= \frac{s_\mu(1, \dots, 1)}{s_\lambda(1, \dots, 1)} \frac{1}{N!} \int_{\mathbb{T}^N} \left(\prod_{i=1}^N \Phi_\omega(z_i) \right) \det[z_i^{\lambda_j + N - j}]_{i,j=1}^N \det[z_i^{-\mu_j - N + j}]_{i,j=1}^N \prod_{i=1}^N \frac{dz_i}{2\pi i z_i} - \delta_{\lambda, \mu} \\ &= \frac{s_\mu(1, \dots, 1)}{s_\lambda(1, \dots, 1)} \frac{1}{N!} \sum_{\sigma, \tau \in S(N)} \text{sgn}(\sigma\tau) \prod_{i=1}^N \int_{\mathbb{T}} \Phi_\omega(z) z^{-(\mu_{\tau(i)} - \tau(i) - \lambda_{\sigma(i)} + \sigma(i))} \frac{dz}{2\pi i z} - \delta_{\lambda, \mu} \\ &= \frac{s_\mu(1, \dots, 1)}{s_\lambda(1, \dots, 1)} \det[\varphi_\omega(\mu_j - j - \lambda_i - i)]_{i,j=1}^N - \delta_{\lambda, \mu} \end{aligned}$$

□

7. EXAMPLES FROM QUANTUM UNITARY GROUPS $U_q(N)$

In this section, we study the case of quantum unitary groups $U_q(N)$ and certain quantized characters. See [28], [23] for the detailed definition and the representation theory of $U_q(N)$. First we recall some results of the representation theory of $U_q(N)$. It is known that $\widehat{U_q(N)}$ can be identified with \mathbb{S}_N . For $\lambda \in \mathbb{S}_N$ we denote by $\chi_{q, \lambda}$ the corresponding irreducible quantized character. Then the “restriction” of $\chi_{q, \lambda}$ to \mathbb{T}^N is given as

$$\frac{s_\lambda(q^{N-1}z_1, q^{N-3}z_2, \dots, q^{-N+1}z_N)}{s_\lambda(q^{N-1}, q^{N-3}, \dots, q^{-N+1})}.$$

See [28] and [34] for more details.

For any probability measure P_N on \mathbb{S}_N we define its q^2 -Schur generating function $\mathcal{S}(z_1, \dots, z_N; P_N)$ by

$$\mathcal{S}(z_1, \dots, z_N; P_N) := \sum_{\lambda \in \mathbb{S}_N} P_N(\lambda) \frac{s_\lambda(z_1, q^{-2}z_2, \dots, q^{-2(N-1)}z_N)}{s_\lambda(1, q^{-2}, \dots, q^{-2(N-1)})},$$

where the right-hand side absolutely converges on \mathbb{T}^N (see [17, Proposition 4.5]). Remark that q^2 -Schur generating functions uniquely determine probability measures on \mathbb{S}_N . We denote by $U_q(\infty)$ the infinite-dimensional quantum unitary group, which is the inductive limit of $U_q(N)$ in the sense of [35]. Then we denote by $\Theta_N^\infty: W^*(U_q(N)) \rightarrow W^*(U_q(\infty))$ the embeddings of group von Neumann algebras (see Section 8). By [17, Proposition 4.6], [34, Proposition 2.6],

there exists a one-to-one correspondence between $\chi \in \text{Ch}(U_q(\infty))$ and the set of all sequences $(P_N)_{N \geq 1}$ of probability measures P_N on \mathbb{S}_N satisfying that

$$\mathcal{S}(z, \dots, z_N, 1; P_{N+1}) = \mathcal{S}(z_1, \dots, z_N; P_N) \quad (7.1)$$

for any $N \geq 1$. More precisely, the corresponding probability measures P_N are given by the irreducible decompositions $\chi \Theta_N^\infty = \sum_{\lambda \in \mathbb{S}_N} P_N(\lambda) \chi_{q, \lambda}$. The following lemma gives examples of q^2 -Schur generating functions satisfying Equation (7.1).

Lemma 7.1. For $N \geq 1$ let $\omega = (\alpha^+, \beta^+, \alpha^-, \beta^-, \gamma^+, \gamma^-) \in \Omega$ such that $\Phi_\omega(q^{-2(i-1)}) > 0$ for $i = 1, \dots, N$ and the Laurent expansion of $\Phi_\omega(z)$ converges in annulus containing $1, \dots, q^{-2(N-1)}$. Then for $k = 1, \dots, N$

$$\mathcal{F}(z_1, \dots, z_k) := \prod_{i=1}^k \frac{\Phi_\omega(q^{-2(i-1)} z_i)}{\Phi_\omega(q^{-2(i-1)})} \quad (7.2)$$

is a q^2 -Schur generating function of probability measure P_k on \mathbb{S}_k given by

$$P_k(\lambda) := \frac{\det[\varphi_\omega(\lambda_i - i + j)]_{i,j=1}^k}{\prod_{i=1}^k \Phi_\omega(q^{-2(i-1)})} s_\lambda(1, q^{-2}, \dots, q^{-2(k-1)})$$

for any $\lambda \in \mathbb{S}_k$. Moreover, Equation (7.1) holds true for $k = 1, \dots, N$.

Remark 7.1. For instance we have $\Phi_\omega(q^{-2(i-1)}) > 0$ if α_i^\pm and β_i^\pm are zero except for finitely many i and $\beta_1^- < (1 - q^{2(N-1)})^{-1}$, $\alpha_1^+ < (q^{-2(N-1)} - 1)^{-1}$. Moreover, if α_i^- and β_i^+ are zero except for finitely many i , $\alpha^+ = \beta^- = (0, 0, \dots)$, then $\mathcal{F}(z_1, \dots, z_N)$ is a q^2 -Schur generating function satisfying Equation (7.1) for any $N \geq 1$. Thus, there exists a corresponding quantized character $\chi_{q, \omega}$ of $U_q(\infty)$.

Proof. Since Equation (7.1) is clear, it suffices to show that $\mathcal{F}(z_1, \dots, z_k)$ is a q^2 -Schur generating function of some probability measure on \mathbb{S}_k . First we have

$$\begin{aligned} & V(z_1, q^{-2} z_2, \dots, q^{-2(k-1)} z_k) \mathcal{F}(z_1, \dots, z_k) \\ &= \frac{1}{\prod_{i=1}^k \Phi_\omega(q^{-2(i-1)})} \sum_{\sigma \in S(k)} \sum_{k_1, \dots, k_k \in \mathbb{Z}} \text{sgn}(\sigma) \prod_{i=1}^k \varphi_\omega(k_i) (q^{-2(i-1)} z_i)^{k - \sigma(i) + k_i} \\ &= \frac{1}{\prod_{i=1}^k \Phi_\omega(q^{-2(i-1)})} \sum_{\sigma \in S(k)} \sum_{m_1, \dots, m_k \in \mathbb{Z}} \text{sgn}(\sigma) \prod_{i=1}^k \varphi_\omega(m_i + \sigma(i)) (q^{-2(i-1)} z_i)^{k + m_i} \\ &= \frac{1}{\prod_{i=1}^k \Phi_\omega(q^{-2(i-1)})} \sum_{m_1, \dots, m_k \in \mathbb{Z}} \det[\varphi_\omega(m_i + j)]_{i,j=1}^k \prod_{i=1}^k (q^{-2(i-1)} z_i)^{k + m_i} \\ &= \frac{1}{\prod_{i=1}^k \Phi_\omega(q^{-2(i-1)})} \sum_{m_1 > \dots > m_k} \sum_{\tau \in S(k)} \det[\varphi_\omega(m_{\tau(i)} + j)]_{i,j=1}^k \prod_{i=1}^k (q^{-2(i-1)} z_i)^{k + m_{\tau(i)}} \\ &= \frac{1}{\prod_{i=1}^k \Phi_\omega(q^{-2(i-1)})} \sum_{m_1 > \dots > m_k} \det[\varphi_\omega(m_i + j)]_{i,j=1}^k \det[(q^{-2(i-1)} z_i)^{k + m_j}]_{i,j=1}^k. \end{aligned}$$

Thus, we have

$$\mathcal{F}(z_1, \dots, z_k) = \sum_{\lambda \in \mathbb{S}_k} P_k(\lambda) \frac{s_\lambda(z_1, q^{-2}z_2, \dots, q^{-2(k-1)}z_k)}{s_\lambda(1, q^{-2}, \dots, q^{-2(k-1)})},$$

where

$$P_k(\lambda) := \frac{\det[\varphi_\omega(\lambda_i - i + j)]_{i,j=1}^k}{\prod_{i=1}^k \Phi_\omega(q^{-2(i-1)})} s_\lambda(1, q^{-2}, \dots, q^{-2(k-1)}) \geq 0$$

since each factor is nonnegative. Moreover, substituting $(1, \dots, 1)$ for (z_1, \dots, z_k) , we have $\sum_{\lambda \in \mathbb{S}_k} P_k(\lambda) = 1$. \square

By the above lemma, there exists a quantized character $\chi_{q,\omega}^N \in \text{Ch}(U_q(N))$ such that the corresponding q^2 -Schur generating functions are given as Equation (7.2).

Theorem 7.1. Let $\omega = (\alpha^+, \beta^+, \alpha^-, \beta^-, \gamma^+, \gamma^-) \in \Omega$ satisfying the assumption in Lemma 7.1 and let $\chi_{q,\omega}^N$ be the corresponding quantized character of $U_q(N)$. Then the generator $\mathbb{L}_{\chi_{q,\omega}^N}$ of the Markov semigroup associated with $\chi_{q,\omega}^N$ in Theorem 5.1 is given by

$$\mathbb{L}_{\chi_{q,\omega}^N}(\lambda, \mu) = \frac{s_\mu(1, q^{-2}, \dots, q^{-2(N-1)}) \det[\varphi_\omega(\mu_j - j - \lambda_i + i)]_{i,j=1}^N}{s_\lambda(1, q^{-2}, \dots, q^{-2(N-1)}) \prod_{i=1}^N \Phi_\omega(q^{-2(i-1)})} - \delta_{\lambda,\mu}$$

for any $\lambda, \mu \in \mathbb{S}_N$.

Proof. Remark that the quantum dimension $d_q(\lambda)$ of irreducible corepresentation associated with $\lambda \in \mathbb{S}_N$ coincides with $s_\lambda(q^{N-1}, q^{N-3}, \dots, q^{-N+1})$. Since $U_q(N)$ has the same fusion rules of $U(N)$, by Corollary 5.2, we have

$$\begin{aligned} & \mathbb{L}_{\chi_{q,\omega}^N}(\lambda, \mu) \\ &= \frac{s_\mu(q^{N-1}, q^{N-3}, \dots, q^{-N+1})}{s_\lambda(q^{N-1}, q^{N-3}, \dots, q^{-N+1})} \frac{1}{N!} \int_{\mathbb{T}^N} \mathcal{F}(q^{-N+1}z_1, \dots, q^{N-1}z_N) s_\lambda(z) \overline{s_\mu(z)} |V(z)|^2 \prod_{i=1}^N \frac{dz_i}{2\pi iz_i} - \delta_{\lambda,\mu}. \end{aligned}$$

By a similar computation in the proof of Theorem 6.1, we obtain the statement. \square

8. ON INDUCTIVE SYSTEMS OF COMPACT QUANTUM GROUPS

In this section, we extend the previous results in Section 5 to inductive systems of compact quantum groups. We assume that $(G_N)_{N=0}^\infty$ is an inductive system of compact quantum group and G_0 is the trivial quantum group. Namely, $C(G_0) = \mathbb{C}$, and $G_N = (C(G_N), \delta_{G_N})$ are compact quantum groups. Moreover, there are the so-called *restriction maps* $\theta_N: C(G_N) \rightarrow C(G_{N-1})$, i.e., θ_N is a surjective $*$ -homomorphism such that $(\theta_N \otimes \theta_N)\delta_{G_N} = \delta_{G_{N-1}}\theta_N$. We denote by $\mathcal{W}^*(G_N) = (W^*(G_N), \hat{\delta}_{G_N}, \hat{R}_{G_N}, \hat{\tau}^{G_N}, \hat{h}_{G_N})$ the Woronowicz algebras of group von Neumann algebras $W^*(G_N)$. Let h_{G_N} be the Haar state of $\mathcal{L}^\infty(G_N)$ and $(\pi_{h_{G_N}}, L^2(G_N), \eta_{h_{G_N}})$ the associated GNS-triple. We define the *Kac-Takesaki operator* \hat{W}_{G_N} as $\hat{W}_{G_N} := \sigma_{L^2(G_N)} W_{G_N}^* \sigma_{L^2(G_N)}$, where $W_{G_N}: L^2(G_N) \otimes L^2(G_N) \rightarrow L^2(G_N) \otimes L^2(G)$ is given by

$$W_{G_N} \xi \otimes \eta_{h_{G_N}}(a) := (\pi_{h_{G_N}} \otimes \pi_{h_{G_N}})(\delta_{G_N}(a)) \xi \otimes \eta_{h_{G_N}}(1)$$

for any $\xi \in L^2(G_N)$ and $a \in C(G_N)$. By [38, Lemma 2.10], there exist faithful normal unital $*$ -homomorphisms $\Theta_N: W^*(G_N) \rightarrow W^*(G_{N+1})$ such that $\Theta_N(\pi_{G_N}(\omega)) = (\text{id} \otimes \omega\theta_{N+1})(\hat{W}_{G_{N+1}})$ for any $\omega \in L^\infty(G_N)_*$. Moreover, we have

$$(\Theta_N \otimes \Theta_N)\hat{\delta}_{G_N} = \hat{\delta}_{G_{N+1}}\Theta_N, \quad \Theta_N\hat{\tau}_t^{G_N} = \hat{\tau}_t^{G_{N+1}}\Theta_N, \quad \Theta_N\hat{R}_{G_N} = \hat{R}_{G_{N+1}}\Theta_N. \quad (8.1)$$

Therefore, we have the inductive limit quantum group W^* -algebra $(W^*(G_\infty), \mathfrak{A}, \hat{\delta}_{G_\infty}, \hat{R}_{G_\infty}, \hat{\tau}^{G_\infty})$ in the sense of [35], that is,

- $W^*(G_\infty) := \varinjlim_N (W^*(G_N), \Theta_N)$ is the inductive limit W^* -algebra in the sense of Takeda [37],
- \mathfrak{A} is the Stratila–Voiculescu AF-algebra of the inductive system $(G_N)_{N=0}^\infty$,
- $\hat{\delta}_{G_\infty}: W^*(G_\infty) \rightarrow W^*(G_\infty) \otimes W^*(G_\infty)$ is a comultiplication, that is, unital normal $*$ -homomorphism satisfying that $(\text{id} \otimes \hat{\delta}_{G_\infty})\hat{\delta}_{G_\infty} = (\hat{\delta}_{G_\infty} \otimes \text{id})\hat{\delta}_{G_\infty}$,
- \hat{R}_{G_∞} is a unitary antipode, that is, an involutive normal $*$ -anti-automorphism satisfying that $\hat{\delta}_{G_\infty}\hat{R}_{G_\infty} = \sigma_{W^*(G_\infty)}(\hat{R}_{G_\infty} \otimes \hat{R}_{G_\infty})\hat{\delta}_{G_\infty}$,
- $\hat{\tau}^{G_\infty} = \{\hat{\tau}_t^{G_\infty}\}_{t \in \mathbb{R}}$ is a deformation automorphism group, that is, a one-parameter automorphism group on $W^*(G_\infty)$ satisfying that $(\hat{\tau}_t^{G_\infty} \otimes \hat{\tau}_t^{G_\infty})\hat{\delta}_{G_\infty} = \hat{\delta}_{G_\infty}\hat{\tau}_t^{G_\infty}$ and $\hat{\tau}_t^{G_\infty}\hat{R}_{G_\infty} = \hat{R}_{G_\infty}\tau_t^{G_\infty}$ for any $t \in \mathbb{R}$.

Remark that there exist normal unital $*$ -homomorphism $\Theta_N^\infty: W^*(G_N) \rightarrow W^*(G_\infty)$ such that $\Theta_{N+1}^\infty\Theta_N = \Theta_N^\infty$ for any $N \geq 0$. For a von Neumann algebra M , if there are normal unital $*$ -homomorphisms $\Phi_N: W^*(G_N) \rightarrow M$ such that $\Phi_{N+1}\Theta_N = \Phi_N$, then there exists a normal unital $*$ -homomorphism $\Phi_\infty: W^*(G_\infty) \rightarrow M$ such that $\Phi_\infty\Theta_N^\infty = \Phi_N$ for any $N \geq 0$.

We define a *quantized character* of G_∞ as a normal $\hat{\tau}^{G_\infty}$ -KMS state (see [35]). We denote by $\text{Ch}(G_\infty)$ the set of all quantized characters of G_∞ . Then we equip $\text{Ch}(G_\infty)$ with the topology of point-wise convergence on the Stratila–Voiculescu AF-algebra \mathfrak{A} . Let $\mathcal{E} := \text{ex}(\text{Ch}(G_\infty))$ equipped with the relative topology induced by $\text{Ch}(G_\infty)$. We recall that $\text{Ch}(G_\infty)$ is a simplex, that is, there exists a one-to-one correspondence $\chi \in \text{Ch}(G_\infty) \mapsto P_\chi \in \mathcal{M}_p(\mathcal{E})$ such that $\chi = \int_{\mathcal{E}} \chi_\omega dP_\chi(\chi_\omega)$. See [35, Corollary 3.1].

For any $\omega \in L^\infty(G_N)_*$ we have $\hat{\epsilon}_{G_{N+1}}(\Theta_N(\pi_{G_N}(\omega))) = \omega(\theta_{N+1}(1)) = \omega(1) = \hat{\epsilon}_{G_N}(\pi_{G_N}(\omega))$. Thus, there exists a normal $*$ -homomorphism $\hat{\epsilon}_\infty: W^*(G_\infty) \rightarrow \mathbb{C}$ such that $\hat{\epsilon}_\infty\Theta_N^\infty = \hat{\epsilon}_{G_N}$ for any $N \geq 0$. We have $(\text{id} \otimes \hat{\epsilon}_\infty)\hat{\delta}_{G_\infty} = (\hat{\epsilon}_\infty \otimes \text{id})\hat{\delta}_{G_\infty} = \text{id}$ since the equation holds true on $\bigcup_{N \geq 0} \Theta_N^\infty(W^*(G_N))$ and it is σ -weakly dense in $W^*(G_\infty)$. In this section, for any $\chi \in \text{Ch}(G_\infty)$ we study the dynamics $((\text{id} \otimes \varphi_t)\hat{\delta}_{G_\infty})_{t \geq 0}$, where $\varphi_t = \exp_*(t(\chi - \hat{\epsilon}_\infty))$.

Lemma 8.1. Let $\chi \in \text{Ch}(G_\infty)$. Then $\varphi_t = \exp_*(t(\chi - \hat{\epsilon}_\infty)) \in \text{Ch}(G_\infty)$.

Proof. By Lemma 5.2, $\varphi_t\Theta_N^\infty = \exp_*(t(\chi\Theta_N^\infty - \hat{\epsilon}_{G_N})) \in \text{Ch}(G_N)$ since $\chi\Theta_N^\infty \in \text{Ch}(G_N)$. Thus, there exists a unique quantized character $\varphi \in \text{Ch}(G_\infty)$ such that $\varphi\Theta_N^\infty = \varphi_t\Theta_N^\infty$ for any $N \geq 0$. See [34], [35]. Since $\bigcup_{N \geq 0} \Theta_N^\infty(W^*(G_N))$ is σ -weakly dense in $W^*(G_\infty)$, we have $\varphi = \varphi_t$, i.e., $\varphi_t \in \text{Ch}(G_\infty)$. \square

Now we define our main object in this section.

Definition 8.1. By the above lemma and [35, Lemma 3.2], for $\chi \in \text{Ch}(G_\infty)$ we obtain the one-parameter semigroup on $\text{Ch}(G_\infty)$ by $\varphi \mapsto \varphi * \varphi_t$, where $\varphi_t = \exp_*(t(\chi - \hat{\epsilon}_\infty))$ for any $t \geq 0$. Then we have the corresponding semigroup $(Q_t^\chi)_{t \geq 0}$ on $\mathcal{M}_p(\mathcal{E})$ such that $Q_t^\chi(P_\varphi) = P_{\varphi * \varphi_t}$, where $P_\varphi, P_{\varphi * \varphi_t}$ are the Borel probability measures corresponding to $\varphi, \varphi * \varphi_t \in \text{Ch}(G_\infty)$, respectively.

The purpose of this section is studying the above dynamics $(Q_t^x)_{t \geq 0}$, and we particularly show the dynamics $(Q_t^x)_{t \geq 0}$ coincide with the “projective limits” of Markov dynamics on the $\widehat{G_N}$. See Proposition 8.2.

Lemma 8.2. Let $\chi \in \text{Ch}(G_\infty)$, $\varphi_t = \exp_*(t(\chi - \hat{\epsilon}_\infty))$ and $T_t = (\text{id} \otimes \varphi_t)\hat{\delta}_{G_\infty}$. Then T_t preserves $\Theta_N^\infty(W^*(G_N))$ and $\Theta_N^\infty(Z(W^*(G_N)))$ for any $N \geq 0$.

Proof. We have $T_t\Theta_N^\infty = \Theta_N^\infty(\text{id} \otimes \varphi_t\Theta_N^\infty)\hat{\delta}_{G_N}$ and $\varphi_t\Theta_N^\infty = \exp_*(t(\chi\Theta_N^\infty - \hat{\epsilon}_{G_N}))$. Since $\chi\Theta_N^\infty \in \text{Ch}(G_N)$, the linear map $(\text{id} \otimes \varphi_t\Theta_N^\infty)\hat{\delta}_{G_N}$ preserves $W^*(G_N)$ and $Z(W^*(G_N))$. \square

Let V_{G_N} be the multiplicative unitary of $\mathcal{L}^\infty(G_N)$. We define $\alpha_0^{G_N}(x) := V_{G_N}(x \otimes 1)V_{G_N}^*$ and $E_{G_N}(x) := (\text{id} \otimes h_{G_N})(\alpha_0^{G_N}(x))$ for any $x \in W^*(G_N)$. Recall that $\alpha_0^{G_N}$ is a right action of $\mathcal{L}^\infty(G_N)$ on $W^*(G_N)$, and E_{G_N} is a normal conditional expectation from $W^*(G_N)$ onto $Z(W^*(G_N))$ (see Lemma 3.1). By [27, Theorem 1.4.2] and Corollary A.1, for any $x = \Phi_G^{-1}((A_\alpha)_{\alpha \in \widehat{G_N}})$ we have

$$E_{G_N}(x) = \sum_{\alpha \in \widehat{G_N}} \frac{1}{d_q(\alpha)} \sum_{i,j=1}^{\dim H_\alpha} \Phi_{G_N}^{-1}(e_{ij}^\alpha F_\alpha A e_{ji}^\alpha) = \sum_{\alpha \in \widehat{G_N}} \chi_\alpha^0(A) \Phi_{G_N}^{-1}(1_{H_\alpha}),$$

where $(e_{ij}^\alpha)_{i,j=1}^{\dim H_\alpha}$ is a matrix unit system of $B(H_\alpha)$ and 1_{H_α} is the identity map on H_α . See Remark 5.3 for the definition of χ_α^0 .

Here we refer to the paper by Ueda [39]. He reformulated the spherical representation theory (due to Olshanski [30], [31] ... etc) in the setting of C^* -algebras with flows. Then, by replacing $\widehat{G_N}$ with the set of minimal projections of the center of $W^*(G_N)$, we obtain the same formula of E_{G_N} . Moreover, the following investigation of *links* of the $\widehat{G_N}$ also extends to the general setting (see [39, Section 7, 8]): Recall that $Z(W^*(G_N))$ and $\ell^\infty(\widehat{G_N})$ are $*$ -isomorphic. Thus, we obtain Markov kernels Λ_N^{N+1} from $\widehat{G_{N+1}}$ to $\widehat{G_N}$ by $E_{G_{N+1}}\Theta_N|_{W^*(G_N)}: Z(W^*(G_N)) \rightarrow Z(W^*(G_{N+1}))$. Then Λ_N^{N+1} induces the map $\mathcal{M}_p(\widehat{G_{N+1}}) \rightarrow \mathcal{M}_p(\widehat{G_N})$, and $\varprojlim_N \mathcal{M}_p(\widehat{G_N})$ is nothing but the set of so-called *coherent systems*. We equip $\varprojlim_N \mathcal{M}_p(\widehat{G_N})$ with the topology of component-wise weakly convergence. Then we have affine homeomorphism between $\varprojlim_N \mathcal{M}_p(\widehat{G_N})$ and $\mathcal{M}_p(\mathcal{E})$. See [39, Proposition 7.10, Corollary 8.3(3)]. Namely \mathcal{E} is the *boundary* of $(\widehat{G_N})_{N=1}^\infty$ with the links $(\Lambda_N^{N+1})_{N=1}^\infty$ in the sense of Boridin–Olshanski [6]. Then for any $N \geq 1$ we obtain the Markov kernel Λ_N^∞ from \mathcal{E} to $\widehat{G_N}$ such that for any $P \in \mathcal{M}_p(\mathcal{E})$ the corresponding coherent system is $(P\Lambda_N^\infty)_{N \geq 1}$.

Recall that for any $\psi \in W^*(G_N)_*$ we define $Q_\psi := (\text{id} \otimes \psi)\hat{\delta}_{G_N}|_{Z(W^*(G_N))}$.

Lemma 8.3. Let $\psi_{N+1} \in W^*(G_{N+1})_*$ be a $\alpha_0^{G_{N+1}}$ -invariant functional and $\psi_N := \psi_{N+1}\Theta_N$. Then $\Psi_N Q_{\psi_N} = Q_{\psi_{N+1}} \Psi_N$.

Proof. By $\alpha_0^{G_{N+1}}$ -invariance of ψ_{N+1} and Equation (3.1), for any $x \in Z(W^*(G_N))$ we have

$$\begin{aligned} [Q_{\psi_{N+1}} \Psi_N](x) &= (\text{id} \otimes \psi_{N+1} \otimes h_{G_{N+1}})((\hat{\delta}_{G_{N+1}} \otimes \text{id})(\alpha_0^{G_{N+1}}(\Theta_N(x)))) \\ &= (\text{id} \otimes \psi_{N+1} \otimes h_{G_{N+1}})(\text{Ad}V_{G_{N+1}13}((\text{id} \otimes \alpha_0^{G_{N+1}})(\hat{\delta}_{G_{N+1}}(\Theta_N(x)))))) \\ &= (\text{id} \otimes \psi_{N+1} \otimes h_{G_{N+1}})(\text{Ad}V_{G_{N+1}13}(\hat{\delta}_{G_{N+1}}(\Theta_N(x)) \otimes 1)). \end{aligned}$$

Since $V_{G_{N+1}13} = \sigma_{L^2(G_{N+1})23} V_{G_{N+1}12} \sigma_{L^2(G_{N+1})23}$, we have

$$\begin{aligned} [\Psi_N Q_{\psi_N}](x) &= (\text{id} \otimes h_{G_{N+1}} \otimes \psi_{N+1})((\alpha_0^{G_{N+1}} \otimes \text{id})(\hat{\delta}_{G_{N+1}}(\Theta_N(x)))) \\ &= (\text{id} \otimes h_{G_{N+1}} \otimes \psi_{N+1})(\text{Ad}V_{G_{N+1}12} \sigma_{L^2(G_{N+1})23}(\hat{\delta}_{G_{N+1}}(\Theta_N(x)) \otimes 1)) \\ &= (\text{id} \otimes \psi_{N+1} \otimes h_{G_{N+1}})(\text{Ad}V_{G_{N+1}13}(\hat{\delta}_{G_{N+1}}(\Theta_N(x)) \otimes 1)). \end{aligned}$$

Therefore, we have $\Psi_N Q_{\psi_N} = Q_{\psi_{N+1}} \Psi_N$. \square

Proposition 8.1. Let $\chi \in \text{Ch}(G_\infty)$, $\psi_N := \chi \Theta_N^\infty - \hat{\epsilon}_{G_N}$ and $Q_t^N := Q_{\exp_*(t\psi_N)}$ for any $t \geq 0$ and $N \geq 1$. Then $\Psi_N Q_t^N = Q_t^{N+1} \Psi_N$ for any $N \geq 0$.

Proof. Remark that $\chi \Theta_N^\infty \in \text{Ch}(G_N)$ (see [35, Section 3]), and hence ψ_N is α_0^N -invariant by Lemma 5.1, 3.5. Since $\hat{\epsilon}_{G_{N+1}} \Theta_N = \hat{\epsilon}_{G_N}$, we have $\psi_{N+1} \Theta_N = \psi_N$ and $\exp_*(t\psi_{N+1}) \Theta_N = \exp_*(t\psi_N)$ for any $t \geq 0$. Since $\exp_*(t\psi_{N+1})$ is $\alpha_0^{G_{N+1}}$ -invariant, $\Psi_N Q_t^N = Q_t^{N+1} \Psi_N$ by Lemma 8.3. \square

Let Q_t^N be the same as in the above. Then, by [6, Proposition 2.4], we obtain a unique Markov semigroup $(Q_t^\infty)_{t \geq 0}$ on \mathcal{E} such that $Q_t^\infty \Lambda_N^\infty = \Lambda_N^\infty Q_t^N$ for any $t \geq 0$ and $N \geq 1$. Recall that we also obtained the semigroup $(Q_t^X)_{t \geq 0}$ on $\mathcal{M}_p(\mathcal{E})$ such that $P_\varphi Q_t^X = P_{\varphi * \varphi_t}$ for any $\varphi \in \text{Ch}(G_\infty)$, where $\varphi_t = \exp_*(t(\chi - \hat{\epsilon}_\infty))$. The following is the conclusion of this section.

Proposition 8.2. Let $\chi \in \text{Ch}(G_\infty)$ and $(Q_t^\infty)_{t \geq 0}$ the corresponding Markov semigroup on \mathcal{E} . Then $(Q_t^\infty)_{t \geq 0}$ coincides with $(Q_t^X)_{t \geq 0}$, that is, $P Q_t^\infty = P Q_t^X$ for any $P \in \mathcal{M}_p(\mathcal{E})$ and $t \geq 0$.

Proof. We may assume that $P = P_\varphi$ for some $\varphi \in \text{Ch}(G_\infty)$. Thus, it suffices to show that $P_\varphi Q_t^\infty = P_{\varphi * \varphi_t}$ for any $t \geq 0$. Since $P_\varphi Q_t^\infty \Lambda_N^\infty = P_{(\varphi * \varphi_t) \Theta_N^\infty}$ and $(P_{(\varphi * \varphi_t) \Theta_N^\infty})_{N \geq 1}$ is the coherent system corresponding to $P_{\varphi * \varphi_t}$, we have $P_\varphi Q_t^\infty = P_{\varphi * \varphi_t}$ for any $t \geq 0$. \square

9. MARKOV SEMIGROUPS ON THE FELLER BOUNDARY OF THE GELFAND–TSETLIN GRAPH

In this section, we return to the case of unitary groups. Recall that $\text{ex}(\text{Ch}(U(\infty)))$ is completely parametrized by Ω in Section 6. We remark that the set of quantized characters is homeomorphic to the set of ordinary characters of $U(\infty)$ with the topology of uniform convergence on compact subsets. Since any compact subsets in $U(\infty)$ is contained in $U(N)$ for some $N \geq 0$ (see [20, Proposition 6.5(i)]), this topology is equivalent to the topology of uniform convergence on each $U(N)$ for $N \geq 0$. On the other hand, we endow Ω with the relative topology induced by the product topology of $\mathbb{R}_{\geq 0}^\infty \times \mathbb{R}_{\geq 0}^\infty \times \mathbb{R}_{\geq 0}^\infty \times \mathbb{R}_{\geq 0}^\infty \times \mathbb{R}_{\geq 0} \times \mathbb{R}_{\geq 0}$. Then, by [31, Theorem 8.1], $\text{ex}(\text{Ch}(U(\infty)))$ and Ω are homeomorphic. Thus, in what follows, we identify Ω with $\text{ex}(\text{Ch}(U(\infty)))$.

We write $\lambda \prec \mu$ if $\lambda_1 \geq \mu_1 \geq \lambda_2 \geq \mu_2 \geq \dots \geq \lambda_N \geq \mu_N \geq \lambda_{N+1}$ for any $\mu \in \mathbb{S}_N$ and $\lambda \in \mathbb{S}_{N+1}$. Then we have the Markov kernel Λ_N^{N+1} from $U(\widehat{N+1}) \cong \mathbb{S}_{N+1}$ to $U(\widehat{N}) \cong \mathbb{S}_N$ in Section 8 is given by

$$\Lambda_N^{N+1}(\lambda, \mu) = \begin{cases} \frac{s_\mu(1, \dots, 1)}{s_\lambda(1, \dots, 1)} & \mu \prec \lambda, \\ 0 & \text{otherwise,} \end{cases}$$

where $s_\mu(1, \dots, 1)$ and $s_\lambda(1, \dots, 1)$ are specializations of Schur polynomials with $N-1$ and N variables, respectively. Moreover, the Markov kernel $\Lambda_N^\infty: \Omega \times \mathbb{S}_N \rightarrow [0, 1]$ is given by

$$\Lambda_N^\infty(\omega, \lambda) = s_\lambda(1, \dots, 1) \det[\varphi_\omega(\lambda_i - i + j)]_{i,j=1}^N$$

for any $\omega \in \Omega$ and $\lambda \in \mathbb{S}_N$, where the $\varphi_\omega(n)$ is the coefficient of z^n in the Laurent expansion of $\Phi_\omega(z)$. See [6] for more details. Then, by [6, Proposition 3.3, 3.4], the boundary Ω of the sequence $(\mathbb{S}_N)_{N=1}^\infty$ with the Markov kernels $(\Lambda_N^{N+1})_{N=1}^\infty$ is Feller, that is, the Markov kernels Λ_N^{N+1} and Λ_N^∞ gives mappings $c_0(\mathbb{S}_N) \rightarrow c_0(\mathbb{S}_{N+1})$ and $c_0(\mathbb{S}_N) \rightarrow c_0(\Omega)$ by

$$\Lambda_N^{N+1}f(\lambda) = \sum_{\mu \in \mathbb{S}_N} \Lambda_N^{N+1}(\lambda, \mu)f(\mu), \quad \Lambda_N^\infty f(\omega) := \sum_{\mu \in \mathbb{S}_N} \Lambda_N^\infty(\omega, \mu)f(\mu)$$

for any $\lambda \in \mathbb{S}_{N+1}$, $\omega \in \Omega$, and $f \in c_0(\mathbb{S}_N)$.

Proposition 9.1. For any $\chi \in \text{Ch}(U(\infty))$ the Markov semigroup $(Q_t^\chi)_{t \geq 0}$ in Proposition ?? is Feller. Moreover, its generator L^χ is determined by $L^\chi \Lambda_N^\infty f = \Lambda_N^\infty L_N^\chi f$ for any $f \in c_0(\mathbb{S}_N)$ and $N \geq 1$, where L_N^χ is the generator of $(Q_t^N)_{t \geq 0}$.

Proof. By [6, Proposition 2.4], the Markov semigroup $(Q_t^\chi)_{t \geq 0}$ is Feller. Recall that $L_N^\chi = Q^{\chi \Theta_N^\infty} - \text{id}$ (see the proof of Theorem 5.1). Thus, we have $\|L_N^\chi\| \leq 2$ for any $N \geq 1$. By [6, Lemma 2.3], $\bigcup_{N \geq 1} \Lambda_N^\infty(c_0(\mathbb{S}_N)) \subset c_0(\Omega)$ is dense. Thus, L^χ is determined by the relations $L^\chi \Lambda_N^\infty f = \Lambda_N^\infty L_N^\chi f$ for any $f \in c_0(\mathbb{S}_N)$ and $N \geq 1$. \square

10. DISCRETE-TIME DYNAMICS ON THE GELFAND–TSETLIN PATTERNS

In the rest of the paper, we discuss discrete-time Markov dynamics on the set of Gelfand–Tsetlin patterns generated by quantized characters of $U_q(N)$ in Lemma 7.1. Recall that every quantized character $\chi \in \text{Ch}(U_q(N))$ gives the Markov operator $\mathbb{Q}^\chi := (\text{id} \otimes \chi) \hat{\delta}_{U_q(N)}|_{Z(W^*(U_q(N)))}$ on $Z(W^*(U_q(N))) \cong \ell^\infty(\mathbb{S}_N)$. Moreover, they are intertwining by the Markov kernels Λ_N^{N+1} in Section 8. If $G_N = U_q(N)$, we can describe the kernel Λ_N^{N+1} explicitly by

$$\Lambda_{N-1}^N(\lambda, \mu) = \begin{cases} q^{N|\mu| - (N-1)|\lambda|} \frac{s_\mu(q^{N-2}, q^{N-4}, \dots, q^{-N+2})}{s_\lambda(q^{N-1}, q^{N-3}, \dots, q^{-N+1})} & \mu \prec \lambda, \\ 0 & \text{otherwise.} \end{cases}$$

See [34, Section 3].

Remark 10.1. The construction of \mathbb{Q}^χ is the generalization of Markov chains due to Kuan [24]. Let G be a compact quantum group and χ its quantized character. By a similar proof of Corollary 5.2, for any $\alpha, \beta \in \hat{G}$ we have

$$\mathbb{Q}^\chi(\alpha, \beta) = \frac{d_q(\beta)}{d_q(\alpha)} \sum_{\gamma \in \hat{G}} \frac{P_\chi(\gamma) N_{\alpha, \gamma}^\beta}{d_q(\gamma)}$$

if the irreducible decomposition of χ is $\sum_{\gamma \in \hat{G}} P_\chi(\gamma) \chi_\gamma$. In particular, when $G = U_q(N)$ and $\chi_{q, \omega}^N$ is its quantized character in Lemma 7.1, for any $\lambda, \mu \in \mathbb{S}_N$ we have

$$\mathbb{Q}^{\chi_{q, \omega}^N}(\lambda, \mu) = \frac{s_\mu(q^{N-1}, q^{N-3}, \dots, q^{-N+1})}{s_\lambda(q^{N-1}, q^{N-3}, \dots, q^{-N+1})} \frac{1}{N!} \int_{\mathbb{T}^N} \prod_{i=1}^N \frac{\Phi_\omega(q^{-2(i-1)z_i})}{\Phi_\omega(q^{-2(i-1)})} s_\lambda(z) \overline{s_\mu(z)} |V(z)|^2 dz.$$

Therefore, if $\mathcal{S}(z_1, \dots, z_N)$ is the q^2 -Schur generating function of a probability measure P on \mathbb{S}_N , then the q^2 -Schur generating function of $P\mathbb{Q}^\chi$ is given as

$$\mathcal{S}(z_1, \dots, z_N) \prod_{i=1}^N \frac{\Phi_\omega(q^{-2(i-1)z_i})}{\Phi_\omega(q^{-2(i-1)})}.$$

The dynamics above is discussed in [5, Section 2], i.e., we have given a representation-theoretic construction of the dynamics in [5].

We also discuss the relationship to the dynamics in [4]. First we recall the Toeplitz-like transition probabilities in [4]. Let $\alpha_1, \dots, \alpha_N$ be nonzero complex numbers and F an analytic function in an annulus A centered at the origin that contains $\alpha_1^{-1}, \dots, \alpha_N^{-1}$. For $n = 1, \dots, N$ we denote $\mathfrak{X}_n = \{(x_1 < \dots < x_n) \in \mathbb{Z}^n\}$. Assume that $F(\alpha_1^{-1}) \cdots F(\alpha_N^{-1}) \neq 0$ and $\det[\alpha_i^{x_j}]_{i,j=1}^n / \det[\alpha_i^{j-1}]_{i,j=1}^n \neq 0$ for any $X = (x_1, \dots, x_n) \in \mathfrak{X}_n$. We also denote

$$f(m) = \frac{1}{2\pi i} \oint F(z) \frac{dz}{z^{m+1}}$$

for any $m \in \mathbb{Z}$, where the integral is taken over any positively oriented simple loop in A . Then we define kernels on $\mathfrak{X}_n \times \mathfrak{X}_n$ and $\mathfrak{X}_n \times \mathfrak{X}_{n-1}$ by

$$T_n(\alpha_1, \dots, \alpha_n; F)(X, Y) = \frac{\det[\alpha_i^{y_j}]_{i,j=1}^n \det[f(x_i - y_j)]_{i,j=1}^n}{\det[\alpha_i^{x_j}]_{i,j=1}^n \prod_{i=1}^n F(\alpha_i^{-1})} \quad \text{for any } X, Y \in \mathfrak{X}_n,$$

$$T_{n-1}^n(\alpha_1, \dots, \alpha_n; F)(X, Y) = \frac{\det[\alpha_i^{y_j}]_{i,j=1}^{n-1} \det[f(x_i - y_j)]_{i,j=1}^n}{\det[\alpha_i^{x_j}]_{i,j=1}^n \prod_{i=1}^{n-1} F(\alpha_i^{-1})} \quad \text{for any } X \in \mathfrak{X}_n, Y \in \mathfrak{X}_{n-1},$$

where $y_n = \text{virt}$ is *virtual* variable and $f(x_i - \text{virt})$ is defined suitably. See [4, Section 2].

We show that Q^{X_n} and Λ_N^{N+1} are given as the above forms. In what follows, we use new coordinates of \mathbb{S}_n given by $x_k(\lambda) = \lambda_{n-k+1} - n + k - 1$ for $k = 1, \dots, n$ and $\lambda \in \mathbb{S}_n$. Then $X_n(\lambda) := (x_1(\lambda), \dots, x_n(\lambda)) \in \mathfrak{X}_n$. Moreover, $\mu \prec \lambda$ for $\mu \in \mathbb{S}_n$ and $\mu \in \mathbb{S}_{n-1}$ if and only if $x_1(\lambda) < x_1(\mu) \leq x_2(\lambda) < \dots < x_{n-1}(\mu) \leq x_n(\lambda)$. Thus, we write $Y \prec X$ if this condition holds true for $X \in \mathfrak{X}_N$ and $Y \in \mathfrak{X}_{n-1}$.

Theorem 10.1. Let $\omega \in \Omega$ as in Lemma 7.1 and $\chi_n := \chi_{q,\omega} \Theta_n^\infty$. Then

$$\mathbb{Q}^{X_n}(\lambda, \mu) = T_n(1, q^{-2}, \dots, q^{-2(n-1)}; \Psi_\omega)(X_n(\lambda), X_n(\mu))$$

for any $\lambda, \mu \in \mathbb{S}_n$, where $\Psi_\omega(z) := \Phi(z^{-1})$.

Proof. By a similar computation in Theorem 6.1, for any $\lambda, \mu \in \mathbb{S}_n$ we have

$$\mathbb{Q}^{X_n}(\lambda, \mu) = \frac{s_\mu(1, q^{-2}, \dots, q^{-2(n-1)}) \det[\varphi_\omega(\mu_j - j - \lambda_i + i)]_{i,j=1}^n}{s_\lambda(1, q^{-2}, \dots, q^{-2(n-1)}) \prod_{i=1}^n \Phi_\omega(q^{-2(i-1)})}.$$

We remark that the coefficients of z^m in the Laurent expansion of Ψ_ω coincide with $\varphi_\omega(-m)$. Therefore,

$$\begin{aligned} \mathbb{Q}^{X_n}(\lambda, \mu) &= \frac{\det[q^{-2(i-1)(\mu_j+n-j)}]_{i,j=1}^n \det[\varphi_\omega(\mu_j - j - \lambda_i + i)]_{i,j=1}^n}{\det[q^{-2(i-1)(\lambda_j+n-j)}]_{i,j=1}^n \prod_{i=1}^n \Phi_\omega(q^{-2(i-1)})} \\ &= \frac{\det[q^{-2(i-1)x_{n-j+1}(\mu)}]_{i,j=1}^n \det[\varphi_\omega(x_{n-j+1}(\mu) - x_{n-i+1}(\lambda))]_{i,j=1}^n}{\det[q^{-2(i-1)x_{n-j+1}(\lambda)}]_{i,j=1}^n \prod_{i=1}^n \Psi_\omega(q^{2(i-1)})} \\ &= T_n(1, q^{-2}, \dots, q^{-2(n-1)}; \Psi_\omega)(X_n(\lambda), X_n(\mu)). \end{aligned}$$

□

Proposition 10.1. Let $F_n(z) = 1/(1 - q^{-2(n-1)}z)$. Then

$$\Lambda_{n-1}^n(\lambda, \mu) = T_{n-1}^n(1, q^{-2}, \dots, q^{-2(n-1)}; F_n)(X_n(\lambda), X_{n-1}(\mu))$$

for any $\lambda \in \mathbb{S}_n$ and $\mu \in \mathbb{S}_{n-1}$. Remark that

$$f(m) := \frac{1}{2\pi i} \oint F(z) \frac{dz}{z^{m+1}} = \begin{cases} q^{-2(n-1)m} & m \geq 0, \\ 0 & m < 0. \end{cases}$$

Moreover, set $f(m - \text{virt}) = q^{-2(n-1)m}$.

Proof. If $\mu \prec \lambda$, then we have

$$\begin{aligned} \Lambda_{n-1}^n(\lambda, \mu) &= q^{n|\mu| - (n-1)|\lambda|} \frac{s_\mu(q^{n-2}, q^{n-4}, \dots, q^{-n+2})}{s_\lambda(q^{n-1}, q^{n-3}, \dots, q^{-n+1})} \\ &= q^{2(n-1)(|\mu| - |\lambda|)} \frac{\det[q^{-2(i-1)(\mu_j + n-1-j)}]_{i,j=1}^{n-1}}{\det[q^{-2(i-1)(\lambda_i + n-j)}]_{i,j=1}^n} \prod_{i=1}^{n-1} (q^{-2(i-1)} - q^{-2(n-1)}). \end{aligned}$$

By [4, Lemma 2.13(ii)], if $X_{n-1}(\mu) \prec X_n(\lambda)$, we have

$$\begin{aligned} &T_{n-1}^n(1, q^{-2}, \dots, q^{-2(n-1)}; F)(X_n(\lambda), X_{n-1}(\mu)) \\ &= \frac{\det[q^{-2(i-1)y_j}]_{i,j=1}^{n-1}}{\det[q^{-2(i-1)x_j}]_{i,j=1}^n} \frac{(-1)^{n-1} q^{-2(n-1)(\sum_{i=1}^n x_i - \sum_{i=1}^{n-1} y_i)}}{\prod_{i=1}^{n-1} (1 - q^{-2(n-1)} q^{2(i-1)})^{-1}} \\ &= q^{2(n-1)(|\mu| - |\lambda|)} \frac{\det[q^{-2(i-1)(\mu_j + n-1-j)}]_{i,j=1}^{n-1}}{\det[q^{-2(i-1)(\lambda_i + n-j)}]_{i,j=1}^n} \prod_{i=1}^{n-1} (q^{-2(i-1)} - q^{-2(n-1)}). \end{aligned}$$

Otherwise, $T_n(1, q^{-2}, \dots, q^{-2(n-1)}; F_n)(X_n(\lambda), X_{n-1}(\mu)) = 0$. Namely, we have

$$\Lambda_{n-1}^n(\lambda, \mu) = T_n(1, q^{-2}, \dots, q^{-2(n-1)}; F_n)(X_n(\lambda), X_{n-1}(\mu)).$$

□

By Lemma 8.3, we have $\Delta_{n-1}^n := \mathbb{Q}^{X_n} \Lambda_{n-1}^n = \Lambda_{n-1}^n \mathbb{Q}^{X_{n-1}}$. Then, by the above two lemmas and [4, Proposition 2.10], we obtain $\Delta_{n-1}^n = T_{n-1}^n(1, q^{-2}, \dots, q^{-2(n-1)}; \Psi_\omega F_n)$. Let

$$\begin{aligned} \mathcal{S}_N &:= \{ \underline{X} = (X_1 \prec \dots \prec X_N) \in \mathfrak{X}_1 \times \dots \times \mathfrak{X}_N \} \\ &= \left\{ \underline{X} = (X_1, \dots, X_N) \in \mathfrak{X}_1 \times \dots \times \mathfrak{X}_N \left| \prod_{n=2}^N \Lambda_{n-1}^n(X_n, X_{n-1}) \neq 0 \right. \right\}. \end{aligned}$$

We remark that we can identify \mathcal{S}_N with the set of Gelfand–Tsetlin patterns of length N . Then we define a Markov operator P_N on \mathcal{S}_N by

$$P_N(\underline{X}, \underline{Y}) := \begin{cases} \mathbb{Q}^{X_1}(\lambda^{(1)}, \mu^{(1)}) \prod_{n=2}^N \frac{\mathbb{Q}^{X_n}(\lambda^{(n)}, \mu^{(n)}) \Lambda_{n-1}^n(\mu^{(n)}, \mu^{(n-1)})}{\Delta_{n-1}^n(\lambda^{(n)}, \mu^{(n-1)})} & \prod_{n=2}^N \Delta_{n-1}^n(\lambda^{(n)}, \mu^{(n-1)}) > 0, \\ 0 & \text{otherwise,} \end{cases}$$

where $X_n = X_n(\lambda^{(n)})$ and $Y_n = X_n(\mu^{(n)})$. Then the dynamics generated by P_N involve several particle systems as appropriate projections. See [4] for more details. Kuan [24] also discussed the relationship between the above dynamics and the representation theory of $U(n)$. In this paper, we extended this relationship to the quantum unitary groups $U_q(n)$.

APPENDIX A. FOURIER ANALYSIS ON COMPACT QUANTUM GROUPS

The contents in this section may be well known for experts. However, we provide this section to avoid any inconvenience in different notations depending on papers. Let $G = (C(G), \delta_G)$ be a compact quantum group and h_G its Haar state. We denote by $(\pi_{h_G}, L^2(G), \eta_{h_G})$ the GNS-triple associated with h_G . We assume that h_G is faithful. Thus, $\eta_G: C(G) \rightarrow L^2(G)$ is injective. Let \widehat{G} be the set of all equivalence classes of irreducible corepresentation of G . For any $\alpha \in \widehat{G}$ we fix a representatives (U_α, H_α) in α . Let $\{f_z^G\}_{z \in \mathbb{C}}$ the Woronowicz character of $(C(G), \delta_G)$ and define $F_\alpha^z := (\text{id} \otimes f_z^G)(U_\alpha)$, $d_q(\alpha) := \text{Tr}(F_\alpha)$. Since the matrices F_α^z ($z \in \mathbb{C}$) are normal and commute with each other, we may choose a matrix unit system $(e_{ij}^\alpha)_{i,j=1}^{\dim(H_\alpha)}$ of $B(H_\alpha)$ such that F_α^z are diagonal matrices. Then we define $u_{ij}^\alpha \in C(G)$ by $U_\alpha = \sum_{i,j=1}^{\dim(H_\alpha)} e_{ij}^\alpha \otimes u_{ij}^\alpha$. For any $\xi \in L^2(G)$ we define $\omega_\xi \in C(G)^*$ by $\omega_\xi(a) := \langle \eta_G(a), \xi \rangle$ for any $a \in C(G)$. We can show the following Peter–Weyl type theorem:

Theorem A.1. For any $\xi \in L^2(G)$ define $\hat{\xi}(\alpha) := (\text{id} \otimes \omega_\xi)(U_\alpha)^*$. Then the mapping

$$\mathcal{F}_G: \xi \in L^2(G) \mapsto \left(\sqrt{d_q(\alpha)} \hat{\xi}(\alpha) \right)_{\alpha \in \widehat{G}} \in \ell^2\text{-}\bigoplus_{\alpha \in \widehat{G}} B(H_\alpha)$$

is an isometric isomorphism of Hilbert spaces, where $\ell^2\text{-}\bigoplus_{\alpha \in \widehat{G}} B(H_\alpha)$ is the completion of the direct sum $\bigoplus_{\alpha \in \widehat{G}} B(H_\alpha)$ with respect to the norm defined by the following inner product:

$$\langle (A_\alpha)_{\alpha \in \widehat{G}}, (B_\alpha)_{\alpha \in \widehat{G}} \rangle := \sum_{\alpha \in \widehat{G}} \text{Tr}_{H_\alpha}(F_\alpha B_\alpha^* A_\alpha).$$

Proof. Remark that $\eta_G(A(G))$ is dense in $L^2(G)$, where $A(G)$ is the $*$ -subalgebra of $C(G)$ generated by matrix coefficients of finite dimensional unitary corepresentations. Thus, by the Schur orthogonal relation (see [27, Theorem 1.4.2]), $\{\eta_G(u_{ij}^\alpha) \mid i, j = 1, \dots, \dim H_\alpha, \alpha \in \widehat{G}\}$ forms an orthogonal basis for $L^2(G)$ and for any $\xi, \eta \in L^2(G)$ we have

$$\langle \xi, \eta \rangle = \sum_{\alpha \in \widehat{G}} d_q(\alpha) \text{Tr}_{H_\alpha}(F_\alpha \hat{\eta}(\alpha)^* \hat{\xi}(\alpha)).$$

□

Let $L^\infty(G)$ be the von Neumann algebra generated by $\pi_{h_G}(C(G))$. Recall that $L^\infty(G)_*$ is a Banach algebra and there exists a representation $(\pi_G, L^2(G))$ of $L^\infty(G)_*$ such that $\pi_G(\omega)\eta_G(a) = \eta_G((\text{id} \otimes \omega)(\delta_G(a)))$ for any $a \in C(G)$ and $\omega \in L^\infty(G)_*$. Then $W^*(G)$ is the von Neumann algebra generated by $\pi_G(L^\infty(G)_*)$.

We can show that $\mathcal{F}_G \pi_G(\omega) \mathcal{F}_G^{-1} = L_G((\pi_\alpha(\omega))_{\alpha \in \widehat{G}})$ for any $\omega \in L^\infty(G)_*$, where $\pi_\alpha(\omega) := (\text{id} \otimes \omega)(U_\alpha)$ and L_G is the left multiplication of $\ell^\infty\text{-}\bigoplus_{\alpha \in \widehat{G}} B(H_\alpha)$ on $\ell^2\text{-}\bigoplus_{\alpha \in \widehat{G}} B(H_\alpha)$. Therefore, there exists a $*$ -isomorphism from

$$\Phi_G: W^*(G) \rightarrow \ell^\infty\text{-}\bigoplus_{\alpha \in \widehat{G}} B(H_\alpha).$$

Then we can describe the Kac–Takesaki operator \hat{W}_G explicitly.

Corollary A.1. $V_G = \sum_{\alpha \in \widehat{G}} (\Phi_G^{-1} \otimes \pi_{h_G})(U_\alpha)$.

Proof. It immediately follows from Theorem A.1. □

ACKNOWLEDGMENT

The author gratefully acknowledges the useful comments from Sinji Koshida and his supervisor, Professor Yoshimichi Ueda. The author appreciates the comments on the paper [5] by Professor Alexei Borodin and Professor Vadim Gorin. This work was supported by JSPS Research Fellowship for Young Scientists (KAKENHI Grant Number JP 19J21098).

REFERENCES

- [1] P. Biane, Quantum random walk on the dual of $SU(n)$, *Probab. Th. Rel. Fields* **89** (1991) 117–129.
- [2] B. Blackadar, *Operator Algebras : Theory of C^* -Algebras and von Neumann Algebras*, Encyclopedia of Mathematical Sciences **122**, Springer-Verlag Berlin Heidelberg, 2006.
- [3] A. Borodin, I. Corwin, Macdonald processes, *Probab. Theory and Related Fields* **158** (2014), 225–400.
- [4] A. Borodin, P. L. Ferrari, Anisotropic growth of random surfaces in 2+1 dimensions, *Commun. Math. Phys.* **325** (2014), 603–684.
- [5] A. Borodin, V. Gorin, Markov processes of infinitely many nonintersecting random walk, *Probab. Theory Relat. Fields.* **155** (2013), 935–997.
- [6] A. Borodin, G. Olshanski, Markov processes on the path space of the Gelfand–Tsetlin graph and on its boundary, *J. Funct. Anal.* **263** (2012), 248–303.
- [7] A. Borodin, L. Petrov, Integrable probability: From representation theory to Macdonald processes, *Probab. Surv.* **11**, 1–58.
- [8] O. Bratteli, D. W. Robinson, *Operator Algebras and Quantum Statistical Mechanics 2. Equilibrium states. Models in quantum statistical mechanics. Second edition*, Texts and Monographs in Physics, Springer-Verlag, Berlin, Heidelberg, 1997.
- [9] A. Borodin, G. Olshanski, Harmonic analysis on the infinite-dimensional unitary group and determinantal point processes, *Ann. of Math. (2)* **161** (2005), no. 3, 1319–1422
- [10] R. P. Boyer, Infinite traces of AF-algebras and characters of $U(\infty)$, *J. Operator Theory* **9** (1983), 205–236.
- [11] F. Cipriani, U. Franz, A. Kula, Symmetries of Lévy processes on compact quantum groups, their Markov semigroups and potential theory, *Journal of Funct. Anal.*, **266** (2014), no. 5, 2789–2844.
- [12] M. Enock, J.-M. Schwartz, *Kac Algebras and Duality of Locally Compact Groups*, Springer-Verlag Berlin Heidelberg, 1992.
- [13] T. Enomoto, M. Izumi, Indecomposable characters of infinite dimensional groups associated with operator algebras, *J. Math. Soc. Japan* **68** (2016), no. 3, 1231–1270
- [14] S. N. Ethier, T. G. Kurtz, *Markov processes – Characterization and Convergence*, Wiley–Interscience, New York, 1986.
- [15] J. Faraut, *Analysis on Lie groups: An Introduction*, Cambridge stud. adv. math. **110**, Cambridge Univ. Press., 2008.
- [16] U. Franz, Lévy processes on quantum group and dual groups, In: *Quantum independent increment processes II.*, Springer, Berlin, Heidelberg (2006), 161–257.
- [17] V. Gorin, The q -Gelfand–Tsetlin graph, Gibbs measures and q -Toeplitz matrices, *Adv. Math.* **229** (2012), 201–266.
- [18] E. Hewitt, K. A. Ross, *Abstract Harmonic Analysis* vol. 2, Springer-Verlag, New York-Berlin, 1970.
- [19] T. Hirai, E. Hiraï, Positive definite class functions on a topological group and characters of factor representations, *J. Math. Kyoto Univ.* **45** (2005), no. 2, 355–376
- [20] T. Hirai, H. Shimomura, N. Tatsuma, E. Hiraï, Inductive limits of topologies, their direct products, and problems related to algebraic structures, *J. Math. Kyoto Univ.* **41** (2001), 475–505.
- [21] M. Izumi, Non-cocommutative Poisson boundaries and compact quantum group actions, *Adv. in Math.* **169**(2002), no.1, 1–57.
- [22] M. Izumi, S. Neshveyev, L. Tuset, Poisson boundary of the dual of $SU_q(n)$, *Commun. Math. Phys.* **262** (2006), 505–531.
- [23] A. Klimyk, K. Schmüdgen, *Quantum Groups and Their Representations*, Texts and Monographs in Physics, Springer-Verlag, Berlin, 1997.
- [24] J. Kuan, Two construction of Markov chains on the dual of $U(n)$, *Journal of Th. Prob.* **31** (2018), 1411–1428.
- [25] G. W. Mackey, Borel structures on groups and their duals, *Trans. Amer. Math. Soc.* **85**(1957), 134–165.

- [26] T. Masuda, Y. Nakagami, A von Neumann algebra framework for the duality of the quantum groups, *Publ. Res. Inst. Math. Sci.* **30**(1994), no. 5, 799–850
- [27] S. Neshveyev, L. Tuset, *Compact Quantum Groups and Their Representation Categories*, Soc. Math. France., 2013.
- [28] M. Noumi, H. Yamada, K. Mimachi, Finite dimensional representations of the quantum group $GL_q(n; \mathbb{C})$ and the zonal spherical functions on $U_q(n-1) \backslash U_q(n)$, *Japan J. Math.* **19** (1993), no. 1, 31–80
- [29] A. Okounkov, G. Olshanski, Asymptotics of Jack Polynomials as the Number of Variables Goes to Infinity, *Internat. Math. Res. Notices* (1998), no. 13, 641–682
- [30] G. Olshanski, Unitary representations of infinite-dimensional pairs (G, K) and the formalism of R. Howe, In *Representation of Lie groups and related topics* (A. M. Vershik and D. P. Zhelobenko, eds.), Adv. Stud. Comtemp. Math. **7**, Gordon and Breach, New York, 1990, 269–468
- [31] G. Olshanski, The problem of harmonic analysis on the infinite-dimensional unitary group, *Journal of Funct. Anal.*, **205** (2003), 464–524
- [32] R. R. Phelps, *Lectures on Choquet's Theorem*, Lecture Note in Mathematics **1757**, Springer-Verlag, Berlin-New York, 2001.
- [33] P. Podleś, S. L. Woronowicz, Quantum Deformation of Lorentz Group, *Commun. Math. Phys.* **130** (1990), 381–431.
- [34] R. Sato, Quantized Vershik–Kerov theory and quantized central probability measures on branching graphs, *Journal of Funct. Anal.*, **277** (2019), 2522–2557
- [35] R. Sato, Inductive limits of compact quantum groups and their unitary representations, arXiv:1908.03988.
- [36] S. Stratila, D. Voiculescu, *Representations of AF-Algebras and of the Group $U(\infty)$* , Lecture Notes in Mathematics **486**, Springer-Verlag, Berlin-New York, 1975.
- [37] Z. Takeda, Inductive limit and infinite direct product of operator algebras, *Tôhoku Math. J. (2)* **7** (1955), 67–86.
- [38] R. Tomatsu, A characterization of right coideals of quotient type and its application to classification of Poisson boundaries, *Commun. Math. Phys.* **275** (2007), 271–296
- [39] Y. Ueda, Spherical representations of C^* -flows, preprint, arXiv:2010.15324
- [40] A. M. Vershik, S. V. Kerov, Characters and factor representations of the infinite unitary group, *Sov. Math. Dokl.* **26** (1982), 570–574.
- [41] D. Voiculescu, Représentations factorielles de type II de $U(\infty)$, *J. Math. Pures Appl.* **55** (1976), 1–20.
- [42] S. Yamagami, On unitary representation theories of compact quantum groups, *Commun. Math. Phys.*, **167** (1995), 509–529
- [43] D. P. Želobenko, Compact Lie groups and their representations, *Translations of Mathematical Monographs* **40**, Amer. Math. Soc., 1973.

GRADUATE SCHOOL OF MATHEMATICS, NAGOYA UNIVERSITY, CHIKUSAKU, NAGOYA 464-8602, JAPAN
 Email address: d19001r@math.nagoya-u.ac.jp