

On Generalizations of Graded 2-absorbing and Graded 2-absorbing primary submodules

Azzh Saad ALSHEHRY and Rashid ABU-DAWWAS

ABSTRACT. Let R be a graded commutative ring with non-zero unity 1 and M be a graded unitary R -module. In this article, we introduce the concepts of graded ϕ -2-absorbing and graded ϕ -2-absorbing primary submodules as generalizations of the concepts of graded 2-absorbing and graded 2-absorbing primary submodules. Let $GS(M)$ be the set of all graded R -submodules of M and $\phi : GS(M) \rightarrow GS(M) \cup \{\emptyset\}$ be a function. A proper graded R -submodule K of M is said to be a graded ϕ -2-absorbing R -submodule of M if whenever x, y are homogeneous elements of R and m is a homogeneous element of M with $xym \in K - \phi(K)$, then $xm \in K$ or $ym \in K$ or $xy \in (K :_R M)$, and K is said to be a graded ϕ -2-absorbing primary R -submodule of M if whenever x, y are homogeneous elements of R and m is a homogeneous element of M with $xym \in K - \phi(K)$, then xm or ym is in the graded radical of K or $xy \in (K :_R M)$. We investigate several properties of these new types of graded submodules.

1. Introduction

Throughout this article, G will be a group with identity e and R a commutative ring with nonzero unity 1. Then R is said to be G -graded if $R = \bigoplus_{g \in G} R_g$ with

$R_g R_h \subseteq R_{gh}$ for all $g, h \in G$ where R_g is an additive subgroup of R for all $g \in G$. The elements of R_g are called homogeneous of degree g . If $x \in R$, then x can be written uniquely as $\sum_{g \in G} x_g$, where x_g is the component of x in R_g . It is known that

R_e is a subring of R and $1 \in R_e$. The set of all homogeneous elements of R is $h(R) = \bigcup_{g \in G} R_g$. Assume that M is a left unitary R -module. Then M is said to

be G -graded if $M = \bigoplus_{g \in G} M_g$ with $R_g M_h \subseteq M_{gh}$ for all $g, h \in G$ where M_g is an

additive subgroup of M for all $g \in G$. The elements of M_g are called homogeneous of degree g . It is clear that M_g is an R_e -submodule of M for all $g \in G$. If $x \in M$, then x can be written uniquely as $\sum_{g \in G} x_g$, where x_g is the component of x in M_g .

2010 *Mathematics Subject Classification.* Primary 16W50; Secondary 13A02.

Key words and phrases. Graded ϕ -prime submodule, graded ϕ -primary submodule, graded 2-absorbing primary submodule, graded weakly 2-absorbing primary submodule, graded ϕ -2-absorbing submodule, graded ϕ -2-absorbing primary submodule.

The set of all homogeneous elements of M is $h(M) = \bigcup_{g \in G} M_g$. Let K be an R -submodule of a graded R -module M . Then K is said to be graded R -submodule if $K = \bigoplus_{g \in G} (K \cap M_g)$, i.e., for $x \in K$, $x = \sum_{g \in G} x_g$ where $x_g \in K$ for all $g \in G$. An R -submodule of a graded R -module need not be graded. For more details and terminology, see [12, 15].

LEMMA 1.1. ([11], Lemma 2.1) *Let R be a graded ring and M be a graded R -module.*

- (1) *If I and J are graded ideals of R , then $I + J$ and $I \cap J$ are graded ideals of R .*
- (2) *If N and K are graded R -submodules of M , then $N + K$ and $N \cap K$ are graded R -submodules of M .*
- (3) *If N is a graded R -submodule of M , $r \in h(R)$, $x \in h(M)$ and I is a graded ideal of R , then Rx , IN and rN are graded R -submodules of M . Moreover, $(N :_R M) = \{r \in R : rM \subseteq N\}$ is a graded ideal of R .*

In particular, $Ann_R(M) = (0 :_R M)$ is a graded ideal of R . Let I be a proper graded ideal of R . Then the graded radical of I is $Grad(I)$, and is defined to be the set of all $r \in R$ such that for each $g \in G$, there exists a positive integer n_g for which $r_g^{n_g} \in I$. One can see that if $r \in h(R)$, then $r \in Grad(I)$ if and only if $r^n \in I$ for some positive integer n . In fact, $Grad(I)$ is a graded ideal of R , see [17]. Let N be a graded R -submodule of M . Then the graded radical of N is denoted by $Grad_M(N)$ and it is defined to be the intersection of all graded prime submodules of M containing N . If there is no graded prime submodule containing N , then we take $Grad_M(N) = M$.

A graded prime (resp. graded primary) R -submodule is a proper graded R -submodule K of M with the property that for $r \in h(R)$ and $m \in h(M)$ such that $rm \in K$ implies that $m \in K$ or $r \in (K :_R M)$ (resp. $m \in K$ or $r \in Grad((K :_R M))$). As graded prime ideals (submodules) have an important role in graded ring (module) theory, several authors generalized these concepts in different ways, see ([1, 3, 5, 9, 16]). Graded weakly prime submodules have been introduced by Atani in [7]. A proper graded R -submodule K of M is said to be a graded weakly prime R -submodule of M if whenever $r \in h(R)$ and $m \in h(M)$ such that $0 \neq rm \in K$, then $m \in K$ or $r \in (K :_R M)$. The concept of graded ϕ -prime submodules has been introduced in [2]. Let $GS(M)$ be the set of all graded R -submodules of M and $\phi : GS(M) \rightarrow GS(M) \cup \{\emptyset\}$ be a function. A proper graded R -submodule K of M is said to be a graded ϕ -prime R -submodule of M if whenever $r \in h(R)$ and $m \in h(M)$ such that $rm \in K - \phi(K)$, then $m \in K$ or $r \in (K :_R M)$.

The concept of graded 2-absorbing ideals (resp. graded weakly 2-absorbing ideals) is introduced in [4] as a different generalization of graded prime ideals (resp. graded weakly prime ideals). A proper graded ideal I of R is a graded 2-absorbing ideal (resp. graded weakly 2-absorbing ideal) of R if whenever $x, y, z \in h(R)$ and $xyz \in I$ (resp. $0 \neq xyz \in I$), then $xy \in I$ or $xz \in I$ or $yz \in I$. Then introducing graded 2-absorbing submodules (resp. graded weakly 2-absorbing submodules) in [3] generalized the concept of graded 2-absorbing ideals (resp. graded weakly 2-absorbing ideals) to graded submodules as following: A proper graded R -submodule

K of M is said to be a graded 2-absorbing R -submodule (resp. graded weakly 2-absorbing R -submodule) of M if whenever $x, y \in h(R)$ and $m \in h(M)$ with $xym \in K$ (resp. $0 \neq xym \in K$), then $xy \in (K :_R M)$ or $xm \in K$ or $ym \in K$.

Al-Zoubi and Sharafat in [6] introduced the concept of graded 2-absorbing primary ideals, where a proper graded ideal I of R is called graded 2-absorbing primary if whenever $x, y, z \in h(R)$ with $xyz \in I$, then $xy \in I$ or $xz \in Grad(I)$ or $yz \in Grad(I)$. The concept of graded 2-absorbing primary submodules is studied in [9] as a generalization of graded 2-absorbing primary ideals. A proper graded R -submodule K of M is said to be a graded 2-absorbing primary R -submodule (resp. graded weakly 2-absorbing primary R -submodule) of M if whenever $xy \in h(R)$ and $m \in h(M)$ with $xym \in K$ (resp. $0 \neq xym \in K$), then $xy \in (K :_R M)$ or $xm \in Grad_M(K)$ or $ym \in Grad_M(K)$.

A graded R -module M is said to be graded multiplication if for every graded R -submodule N of M , $N = IM$ for some graded ideal I of R . In this case, it is known that $I = (N :_R M)$. Graded multiplication modules were firstly introduced and studied by Escoriza and Torrecillas in [10], and further results were obtained by several authors, see for example [13]. Let N and K be graded R -submodules of a graded multiplication R -module M with $N = IM$ and $K = JM$ for some graded ideals I and J of R . The product of N and K is denoted by NK is defined by $NK = IJM$. Then the product of N and K is independent of presentations of N and K . In fact, as IJ is a graded ideal of R (see [15]), NK is a graded R -submodule of M and $NK \subseteq N \cap K$. Moreover, for $x, y \in h(M)$, by xy , we mean the product of Rx and Ry . Also, it is shown in ([16], Theorem 9) that if N is a proper graded R -submodule of a graded multiplication R -module M , then $Grad_M(N) = Grad((N :_R M))M$.

In this article, our aim is to extend the concept of graded 2-absorbing submodules to graded ϕ -2-absorbing submodules in completely different way from [2], and also to extend graded 2-absorbing primary submodules to graded ϕ -2-absorbing primary submodules. Our study is inspired from [14].

2. Graded ϕ -2-Absorbing and Graded ϕ -2-Absorbing Primary Submodules

In this section, we introduce and study the concepts of graded ϕ -2-absorbing and graded ϕ -2-absorbing primary submodules.

DEFINITION 2.1. *Let M be a G -graded R -module and $\phi : GS(M) \rightarrow GS(M) \cup \{\emptyset\}$ be a function.*

- (1) *A proper graded R -submodule K of M is said to be a graded ϕ -primary R -submodule of M if whenever $r \in h(R)$ and $m \in h(M)$ with $rm \in K - \phi(K)$, then $m \in K$ or $r \in Grad((K :_R M))$.*
- (2) *Let K be a graded R -submodule of M and $g \in G$ such that $K_g \neq M_g$. Then K is said to be a g - ϕ -primary R -submodule of M if whenever $r \in R_e$ and $m \in M_g$ with $rm \in K - \phi(K)$, then $m \in K$ or $r \in Grad((K :_R M))$.*
- (3) *A proper graded R -submodule K of M is said to be a graded ϕ -2-absorbing R -submodule of M if whenever $x, y \in h(R)$ and $m \in h(M)$ with $xym \in K - \phi(K)$, then $xm \in K$ or $ym \in K$ or $xy \in (K :_R M)$.*
- (4) *Let K be a graded R -submodule of M and $g \in G$ such that $K_g \neq M_g$. Then K is said to be a g - ϕ -2-absorbing R -submodule of M if whenever*

$x, y \in R_e$ and $m \in M_g$ with $xym \in K - \phi(K)$, then $xm \in K$ or $ym \in K$ or $xy \in (K :_R M)$.

- (5) A proper graded R -submodule K of M is said to be a graded ϕ -2-absorbing primary R -submodule of M if whenever $x, y \in h(R)$ and $m \in h(M)$ with $xym \in K - \phi(K)$, then $xm \in \text{Grad}_M(K)$ or $ym \in \text{Grad}_M(K)$ or $xy \in (K :_R M)$.
- (6) Let K be a graded R -submodule of M and $g \in G$ such that $K_g \neq M_g$. Then K is said to be a g - ϕ -2-absorbing primary R -submodule of M if whenever $x, y \in R_e$ and $m \in M_g$ with $xym \in K - \phi(K)$, then $xm \in \text{Grad}_M(K)$ or $ym \in \text{Grad}_M(K)$ or $xy \in (K :_R M)$.

REMARK 2.2. (1) Let K be a graded ϕ -primary R -submodule of a graded multiplication R -module M . Then

$$\phi_\emptyset(K) = \emptyset \quad \text{graded primary submodule,}$$

$$\phi_0(K) = \{0\} \quad \text{graded weakly primary submodule,}$$

$$\phi_2(K) = K^2 \quad \text{graded almost primary submodule,}$$

$$\phi_n(K) = K^n \quad \text{graded } n\text{-almost primary submodule, and}$$

$$\phi_\omega(K) = \bigcap_{n=1}^{\infty} K^n \quad \text{graded } \omega\text{-primary submodule.}$$

- (2) Let K be a graded ϕ -2-absorbing (resp. graded ϕ -2-absorbing primary) R -submodule of a graded multiplication R -module M . Then

$$\phi_\emptyset(K) = \emptyset \quad \text{graded 2-absorbing (resp. graded 2-absorbing primary) submodule,}$$

$$\phi_0(K) = \{0\} \quad \text{graded weakly 2-absorbing (resp. graded weakly 2-absorbing primary) submodule,}$$

$$\phi_2(K) = K^2 \quad \text{graded almost 2-absorbing (resp. graded almost 2-absorbing primary) submodule,}$$

$$\phi_n(K) = K^n \quad \text{graded } n\text{-almost 2-absorbing (resp. graded } n\text{-almost 2-absorbing primary) submodule, and}$$

$$\phi_\omega(K) = \bigcap_{n=1}^{\infty} K^n \quad \text{graded } \omega\text{-2-absorbing (resp. graded } \omega\text{-2-absorbing primary) submodule.}$$

- (3) For functions $\phi, \varphi : GS(M) \rightarrow GS(M) \cup \{\emptyset\}$, we write $\phi \leq \varphi$ if $\phi(K) \subseteq \varphi(K)$ for all $K \in GS(M)$. Thus clearly we have the following order:

$$\phi_\emptyset \leq \phi_0 \leq \phi_\omega \leq \dots \leq \phi_{n+1} \leq \phi_n \leq \dots \leq \phi_2 \leq \phi_1.$$

- (4) If $\phi \leq \varphi$, then every graded ϕ -2-absorbing (resp. graded ϕ -2-absorbing primary) R -submodule is graded φ -2-absorbing (resp. graded φ -2-absorbing primary).
- (5) Since $K - \phi(K) = K - (K \cap \phi(K))$ for any graded R -submodule K of M , without loss of generality, throughout this article, we assume that $\phi(K) \subseteq K$.

THEOREM 2.3. Let M be a graded R -module and K be a proper graded R -submodule of M . Then the followings hold:

- (1) K is a graded ϕ -prime R -submodule of $M \Rightarrow K$ is a graded ϕ -2-absorbing R -submodule of $M \Rightarrow K$ is a graded ϕ -2-absorbing primary R -submodule of M .
- (2) If M is a graded multiplication R -module and K is a graded ϕ -primary R -submodule of M , then K is a graded ϕ -2-absorbing primary R -submodule of M .
- (3) For graded multiplication R -module M , K is a graded 2-absorbing R -submodule of $M \Rightarrow K$ is a graded weakly 2-absorbing R -submodule of $M \Rightarrow K$ is a graded ω -2-absorbing R -submodule of $M \Rightarrow K$ is a graded $(n+1)$ -almost 2-absorbing R -submodule of $M \Rightarrow K$ is a graded n -almost 2-absorbing R -submodule of M for all $n \geq 2 \Rightarrow K$ is a graded almost 2-absorbing R -submodule of M .
- (4) For graded multiplication R -module M , K is a graded 2-absorbing primary R -submodule of $M \Rightarrow K$ is a graded weakly 2-absorbing primary R -submodule of $M \Rightarrow K$ is a graded ω -2-absorbing primary R -submodule of $M \Rightarrow K$ is a graded $(n+1)$ -almost 2-absorbing primary R -submodule of $M \Rightarrow K$ is a graded n -almost 2-absorbing primary R -submodule of M for all $n \geq 2 \Rightarrow K$ is a graded almost 2-absorbing primary R -submodule of M .
- (5) Suppose that $\text{Grad}_M(K) = K$. Then K is a graded ϕ -2-absorbing primary R -submodule of M if and only if K is a graded ϕ -2-absorbing R -submodule of M .
- (6) If M is a graded multiplication R -module and K is an idempotent R -submodule of M , then K is a graded ω -2-absorbing R -submodule of M , and K is a graded n -almost 2-absorbing R -submodule of M for every $n \geq 2$.
- (7) Let M be a graded multiplication R -module. Then K is a graded n -almost 2-absorbing (resp. graded n -almost 2-absorbing primary) R -submodule of M for all $n \geq 2$ if and only if K is a graded ω -2-absorbing (resp. graded ω -2-absorbing primary) R -submodule of M .

PROOF. (1) It is straightforward.

- (2) Let $x, y \in h(R)$ and $m \in h(M)$ such that $xym \in K - \phi(K)$. Assume that $ym \notin \text{Grad}_M(K)$. Then $ym \notin K$ and then $x \in \text{Grad}((K :_R M))$ as K is a graded ϕ -primary R -submodule. Therefore, $xm \in \text{Grad}((K :_R M)M) = \text{Grad}_M(K)$. Consequently, K is graded ϕ -2-absorbing primary.
- (3) It is clear by Remark 2.2 (4).
- (4) It is clear by Remark 2.2 (4).
- (5) The claim is obvious.
- (6) Since K is an idempotent R -submodule, $K^n = K$ for all $n > 0$, and then $\phi_\omega(K) = \bigcap_{n=1}^{\infty} K^n = K$. Thus K is a graded ω -2-absorbing R -submodule of M . By (3), we conclude that K is a graded n -almost 2-absorbing R -submodule of M for all $n \geq 2$.
- (7) Suppose that K is a graded n -almost 2-absorbing (resp. graded n -almost 2-absorbing primary) R -submodule of M for all $n \geq 2$. Let $x, y \in h(R)$ and $m \in h(M)$ with $xym \in K$ but $xym \notin \bigcap_{n=1}^{\infty} K^n$. Hence $xym \notin K^n$ for some $n \geq 2$. Since K is graded n -almost 2-absorbing (resp. graded n -almost 2-absorbing primary) for all $n \geq 2$, this implies either $xy \in (K :_R M)$ or $ym \in K$ or $xm \in K$ (resp. $xy \in (K :_R M)$ or $ym \in \text{Grad}_M(K)$ or

$xm \in \text{Grad}_M(K)$). This completes the first implication. The converse is clear from (3) (resp. from (4)). \square

Let M be a G -graded R -module and K be a graded R -submodule of M . Then M/K is G -graded by $(M/K)_g = (M_g + K)/K$ for all $g \in G$ ([15]).

LEMMA 2.4. ([18], Lemma 3.2) *Let M be a graded R -module, K be a graded R -submodule of M , and N be an R -submodules of M such that $K \subseteq N$. Then N is a graded R -submodule of M if and only if N/K is a graded R -submodule of M/K .*

THEOREM 2.5. *Let M be a graded R -module and K be a proper graded R -submodule of M . Then the following hold:*

- (1) *K is a graded ϕ -2-absorbing R -submodule of M if and only if $K/\phi(K)$ is a graded weakly 2-absorbing R -submodule of $M/\phi(K)$.*
- (2) *K is a graded ϕ -2-absorbing primary R -submodule of M if and only if $K/\phi(K)$ is a graded weakly 2-absorbing primary R -submodule of $M/\phi(K)$.*
- (3) *K is a graded ϕ -prime R -submodule of M if and only if $K/\phi(K)$ is a graded weakly prime R -submodule of $M/\phi(K)$.*
- (4) *K is a graded ϕ -primary R -submodule of M if and only if $K/\phi(K)$ is a graded weakly primary R -submodule of $M/\phi(K)$.*

PROOF. (1) If $\phi(K) = \emptyset$, then it is done. Suppose that $\phi(K) \neq \emptyset$. Let $x, y \in h(R)$ and $m + \phi(K) \in h(M/\phi(K))$ such that $\phi(K) \neq xy(m + \phi(K)) = xym + \phi(K) \in K/\phi(K)$. Then $m \in h(M)$ such that $xym \in K$, but $xym \notin \phi(K)$. Hence either $xy \in (K :_R M)$ or $ym \in K$ or $xm \in K$. So, $xy \in (K/\phi(K) :_R M/\phi(K))$ or $y(m + \phi(K)) \in K/\phi(K)$ or $x(m + \phi(K)) \in K/\phi(K)$, as desired. Conversely, let $x, y \in h(R)$ and $m \in h(M)$ such that $xym \in K$ and $xym \notin \phi(K)$. Then $m + \phi(K) \in h(M/\phi(K))$ such that $\phi(K) \neq xy(m + \phi(K)) \in K/\phi(K)$. Hence $xy \in (K/\phi(K) :_R M/\phi(K))$ or $y(m + \phi(K)) \in K/\phi(K)$ or $x(m + \phi(K)) \in K/\phi(K)$. So, $xy \in (K :_R M)$ or $ym \in K$ or $xm \in K$. Thus K is a graded ϕ -2-absorbing R -submodule of M .

- (2) Let $x, y \in h(R)$ and $m + \phi(K) \in h(M/\phi(K))$ such that $\phi(K) \neq xy(m + \phi(K)) = xym + \phi(K) \in K/\phi(K)$. Then $m \in h(M)$ such that $xym \in K$, but $xym \notin \phi(K)$. Hence either $xy \in (K :_R M)$ or $ym \in \text{Grad}_M(K)$ or $xm \in \text{Grad}_M(K)$. So, $xy \in (K :_R M)$ or $y(m + \phi(K)) \in \text{Grad}_M(K)/\phi(K)$ or $x(m + \phi(K)) \in \text{Grad}_M(K)/\phi(K)$. The result holds since $\text{Grad}_M(K)/\phi(K) = \text{Grad}_{M/\phi(K)}(K/\phi(K))$. One can easily prove the converse.

Similarly, one can easily prove (3) and (4). \square

Let M and L be two G -graded R -modules. An R -homomorphism $f : M \rightarrow L$ is said to be a graded R -homomorphism if $f(M_g) \subseteq L_g$ for all $g \in G$ ([15]).

LEMMA 2.6. ([1], Lemma 2.16) *Suppose that $f : M \rightarrow L$ is a graded R -homomorphism. If N is a graded R -submodule of L , then $f^{-1}(N)$ is a graded R -submodule of M .*

LEMMA 2.7. ([8], Lemma 4.8) *Suppose that $f : M \rightarrow L$ is a graded R -homomorphism. If K is a graded R -submodule of M , then $f(K)$ is a graded R -submodule of $f(M)$.*

THEOREM 2.8. *Suppose that $f : M \rightarrow L$ is a graded R -epimorphism. Let $\phi : GS(M) \rightarrow GS(M) \cup \{\emptyset\}$ and $\varphi : GS(L) \rightarrow GS(L) \cup \{\emptyset\}$ be functions. Then the following hold:*

- (1) If N is a graded φ -2-absorbing primary R -submodule of L and $\phi(f^{-1}(N)) = f^{-1}(\varphi(N))$, then $f^{-1}(N)$ is a graded ϕ -2-absorbing primary R -submodule of M .
- (2) If K is a graded ϕ -2-absorbing primary R -submodule of M containing $\text{Ker}(f)$ and $\varphi(f(K)) = f(\phi(K))$, then $f(K)$ is a graded φ -2-absorbing primary R -submodule of L .
- (3) If N is a graded φ -2-absorbing R -submodule of L and $\phi(f^{-1}(N)) = f^{-1}(\varphi(N))$, then $f^{-1}(N)$ is a graded ϕ -2-absorbing R -submodule of M .
- (4) If K is a graded ϕ -2-absorbing R -submodule of M containing $\text{Ker}(f)$ and $\varphi(f(K)) = f(\phi(K))$, then $f(K)$ is a graded φ -2-absorbing R -submodule of L .

PROOF. (1) Since f is epimorphism, $f^{-1}(N)$ is a proper graded R -submodule of M . Let $x, y \in h(R)$ and $m \in h(M)$ such that $xym \in f^{-1}(N)$ and $xym \notin f^{-1}(\varphi(N))$. Since $xym \in f^{-1}(N)$, $xyf(m) \in N$. Also, $\phi(f^{-1}(N)) = f^{-1}(\varphi(N))$ implies that $xyf(m) \notin \varphi(N)$. Thus $xyf(m) \in N - \varphi(N)$. Then $xy \in (N :_R L)$ or $xf(m) \in \text{Grad}_L(N)$ or $yf(m) \in \text{Grad}_L(N)$. Thus $xy \in (f^{-1}(N) :_R M)$ or $xm \in f^{-1}(\text{Grad}_L(N))$ or $ym \in f^{-1}(\text{Grad}_L(N))$. Since $f^{-1}(\text{Grad}_L(N)) \subseteq \text{Grad}_M(f^{-1}(N))$, we conclude that $f^{-1}(N)$ is a graded ϕ -2-absorbing primary R -submodule of M .

- (2) Let $x, y \in h(R)$ and $s \in h(L)$ such that $xys \in f(K) - \varphi(f(K))$. Since f is graded epimorphism, there exists $m \in h(M)$ such that $s = f(m)$. Therefore, $f(xym) \in f(K)$ and so $xym \in K$ as $\text{Ker}(f) \subseteq K$. Since $\varphi(f(K)) = f(\phi(K))$, we have $xym \notin \phi(K)$. Hence $xym \in K - \phi(K)$. It implies that $xy \in (K :_R M)$ or $xm \in \text{Grad}_M(K)$ or $ym \in \text{Grad}_M(K)$. Thus $xy \in (f(K) :_R L)$ or $xs \in f(\text{Grad}_M(K))$ or $ys \in f(\text{Grad}_M(K))$. Since $\text{Ker}(f) \subseteq K$, $f(\text{Grad}_M(K)) = \text{Grad}_L(f(K))$, and then we are done.

Similarly, one can easily prove (3) and (4). □

Let M be a G -graded R -module and $S \subseteq h(R)$ be a multiplicative set. Then $S^{-1}M$ is a G -graded $S^{-1}R$ -module with $(S^{-1}M)_g = \{\frac{m}{s}, m \in M_h, s \in S \cap R_{hg^{-1}}\}$ for all $g \in G$, and $(S^{-1}R)_g = \{\frac{a}{s}, a \in R_h, s \in S \cap R_{hg^{-1}}\}$ for all $g \in G$. If K is a graded R -submodule of M , then $S^{-1}K$ is a graded $S^{-1}R$ -submodule of $S^{-1}M$. Let $\phi : GS(M) \rightarrow GS(M) \cup \{\emptyset\}$ be a function and define $\phi_S : GS(S^{-1}M) \rightarrow GS(S^{-1}M) \cup \{\emptyset\}$ by $\phi_S(S^{-1}K) = S^{-1}\phi(K)$, and $\phi_S(S^{-1}K) = \emptyset$ when $\phi(K) = \emptyset$ for every graded R -submodule K of M .

THEOREM 2.9. *Let M be a graded R -module and $S \subseteq h(R)$ be a multiplicative set.*

- (1) If K is a graded ϕ -2-absorbing primary R -submodule of M and $S^{-1}K \neq S^{-1}M$, then $S^{-1}K$ is a graded ϕ_S -2-absorbing primary $S^{-1}R$ -submodule of $S^{-1}M$.
- (2) If K is a graded ϕ -2-absorbing R -submodule of M and $S^{-1}K \neq S^{-1}M$, then $S^{-1}K$ is a graded ϕ_S -2-absorbing $S^{-1}R$ -submodule of $S^{-1}M$.

PROOF. (1) Let $x/s_1, y/s_2 \in h(S^{-1}R)$ and $m/s \in h(S^{-1}M)$ such that $(x/s_1)(y/s_2)(m/s) \in S^{-1}K - \phi_S(S^{-1}K)$. Then $x, y \in h(R)$ and $m \in h(M)$ such that $uxym \in K - \phi(K)$ for some $u \in S$, and then $uxm \in \text{Grad}_M(K)$ or $uym \in \text{Grad}_M(K)$ or $xy \in (K :_R M)$. So, $(x/s_1)(m/s) = (uxm)/(us_1s) \in S^{-1}(\text{Grad}_M(K)) \subseteq \text{Grad}_{S^{-1}M}(S^{-1}K)$ or $(y/s_2)(m/s) = (uym)/(us_2s) \in$

$\text{Grad}_{S^{-1}M}(S^{-1}K)$ or $(x/s_1)(y/s_2) = (xy)/(s_1s_2) \in S^{-1}(K :_R M) \subseteq (S^{-1}K :_{S^{-1}R} S^{-1}M)$.

Similarly, one can easily prove (2). \square

Let M_1 be a G -graded R_1 -module, M_2 be a G -graded R_2 -module and $R = R_1 \times R_2$. Then $M = M_1 \times M_2$ is G -graded R -module with $M_g = (M_1)_g \times (M_2)_g$ for all $g \in G$, where $R_g = (R_1)_g \times (R_2)_g$ for all $g \in G$, ([15]).

LEMMA 2.10. ([18], Lemma 3.12) *Let M_1 be a G -graded R_1 -module, M_2 be a G -graded R_2 -module, $R = R_1 \times R_2$ and $M = M_1 \times M_2$. Then $L = N \times K$ is a graded R -submodule of M if and only if N is a graded R_1 -submodule of M_1 and K is a graded R_2 -submodule of M_2 .*

LEMMA 2.11. *Let M_1 be a G -graded R_1 -module, M_2 be a G -graded R_2 -module, $R = R_1 \times R_2$ and $M = M_1 \times M_2$. Suppose that $\varphi_1 : GS(M_1) \rightarrow GS(M_1) \cup \{\emptyset\}$, $\varphi_2 : GS(M_2) \rightarrow GS(M_2) \cup \{\emptyset\}$ be functions and $\phi = \varphi_1 \times \varphi_2$. Assume that $K = K_1 \times M_2$ for some proper graded R_1 -submodule K_1 of M_1 . If K is a graded ϕ -2-absorbing R -submodule of M , then K_1 is a graded φ_1 -2-absorbing R_1 -submodule of M_1 .*

PROOF. Let $x_1, y_1 \in h(R_1)$ and $m_1 \in h(M_1)$ such that $x_1y_1m_1 \in K_1 - \varphi_1(K_1)$. Then for any $m \in h(M_2)$, $(x_1, 1), (y_1, 1) \in h(R)$ and $(m_1, m) \in h(M)$ such that $(x_1, 1)(y_1, 1)(m_1, m) \in (K_1 \times M_2) - (\varphi_1(K_1) \times \varphi_2(M_2)) = K - \phi(K)$. Since K is a graded ϕ -2-absorbing R -submodule of M , we get either $(x_1, 1)(y_1, 1) \in ((K_1 \times M_2) :_R M_1 \times M_2)$ or $(x_1, 1)(m_1, m) \in (K_1 \times M_2)$ or $(y_1, 1)(m_1, m) \in (K_1 \times M_2)$. So clearly, we conclude that $x_1y_1 \in (K_1 :_{R_1} M_1)$ or $x_1m_1 \in K_1$ or $y_1m_1 \in K_1$. Therefore, K_1 is a graded φ_1 -2-absorbing R_1 -submodule of M_1 . \square

THEOREM 2.12. *Let M_1 be a G -graded R_1 -module, M_2 be a G -graded R_2 -module, $R = R_1 \times R_2$ and $M = M_1 \times M_2$. Suppose that $\varphi_1 : GS(M_1) \rightarrow GS(M_1) \cup \{\emptyset\}$, $\varphi_2 : GS(M_2) \rightarrow GS(M_2) \cup \{\emptyset\}$ be functions and $\phi = \varphi_1 \times \varphi_2$. Assume that $K = K_1 \times M_2$ for some proper graded R_1 -submodule K_1 of M_1 . Then the following hold:*

- (1) *If $\varphi_2(M_2) = M_2$, then K is a graded ϕ -2-absorbing R -submodule of M if and only if K_1 is a graded φ_1 -2-absorbing R_1 -submodule of M_1 .*
- (2) *If $\varphi_2(M_2) \neq M_2$, then K is a graded ϕ -2-absorbing R -submodule of M if and only if K_1 is a graded 2-absorbing R_1 -submodule of M_1 .*

PROOF. (1) Suppose that K_1 is a graded φ_1 -2-absorbing R_1 -submodule of M_1 . Let $x = (x_1, x_2), y = (y_1, y_2) \in h(R)$ and $m = (m_1, m_2) \in h(M)$ such that $xym \in K - \phi(K)$. Since $\varphi_2(M_2) = M_2$, we get that $x_1, y_1 \in h(R_1)$ and $m_1 \in h(M_1)$ such that $x_1y_1m_1 \in K_1 - \varphi_1(K_1)$, and this implies that either $x_1y_1 \in (K_1 :_{R_1} M_1)$ or $x_1m_1 \in K_1$ or $y_1m_1 \in K_1$. Thus either $xy \in (K :_R M)$ or $xm \in K$ or $ym \in K$. Hence, K is a graded ϕ -2-absorbing R -submodule of M . The converse holds from Lemma 2.11.

- (2) Suppose that K is a graded ϕ -2-absorbing R -submodule of M . Since $\varphi_2(M_2) \neq M_2$, there exists $m_2 \in M_2 - \varphi_2(M_2)$ and then there exists $g \in G$ such that $(m_2)_g \in M_2 - \varphi_2(M_2)$. Assume that K_1 is not a graded 2-absorbing R_1 -submodule of M_1 . By Lemma 2.11, K_1 is a graded φ_1 -2-absorbing R_1 -submodule of M_1 . Hence, there exist $x_1, y_1 \in h(R_1)$ and $m_1 \in h(M_1)$ such that $x_1y_1m_1 \in \varphi_1(K_1)$, $x_1y_1 \notin (K_1 :_{R_1} M_1)$, $x_1m_1 \notin K_1$ and $y_1m_1 \notin K_1$. So, $(x_1, 1)(y_1, 1)(m_1, (m_2)_g) \in (K_1 \times M_2) - (\varphi_1(K_1) \times \varphi_2(M_2)) = (K_1 \times M_2) - \phi(K_1 \times M_2)$ which implies that $x_1y_1 \in (K_1 :_{R_1} M_1)$ or $x_1m_1 \in K_1$

or $y_1 m_1 \in K_1$, which is a contradiction. Thus K_1 is a graded 2-absorbing R_1 -submodule of M_1 . Conversely, if K_1 is a graded 2-absorbing R_1 -submodule of M_1 , then $K = K_1 \times M_2$ is a graded 2-absorbing R -submodule of M by ([5], Theorem 3.3). Hence K is a graded ϕ -2-absorbing R -submodule of M for any ϕ . □

LEMMA 2.13. *Let M_1 be a G -graded R_1 -module, M_2 be a G -graded R_2 -module, $R = R_1 \times R_2$ and $M = M_1 \times M_2$. Suppose that $\varphi_1 : GS(M_1) \rightarrow GS(M_1) \cup \{\emptyset\}$, $\varphi_2 : GS(M_2) \rightarrow GS(M_2) \cup \{\emptyset\}$ be functions and $\phi = \varphi_1 \times \varphi_2$. Assume that $K = K_1 \times M_2$ for some proper graded R_1 -submodule K_1 of M_1 . If K is a graded ϕ -2-absorbing primary R -submodule of M , then K_1 is a graded φ_1 -2-absorbing primary R_1 -submodule of M_1 .*

PROOF. It can be easily proved by using a similar argument in the proof of Lemma 2.11. □

THEOREM 2.14. *Let M_1 be a G -graded R_1 -module, M_2 be a G -graded R_2 -module, $R = R_1 \times R_2$ and $M = M_1 \times M_2$. Suppose that $\varphi_1 : GS(M_1) \rightarrow GS(M_1) \cup \{\emptyset\}$, $\varphi_2 : GS(M_2) \rightarrow GS(M_2) \cup \{\emptyset\}$ be functions and $\phi = \varphi_1 \times \varphi_2$. Assume that $K = K_1 \times M_2$ for some proper graded R_1 -submodule K_1 of M_1 . Then the following hold:*

- (1) *If $\varphi_2(M_2) = M_2$, then K is a graded ϕ -2-absorbing primary R -submodule of M if and only if K_1 is a graded φ_1 -2-absorbing primary R_1 -submodule of M_1 .*
- (2) *If $\varphi_2(M_2) \neq M_2$, then K is a graded ϕ -2-absorbing primary R -submodule of M if and only if K_1 is a graded 2-absorbing primary R_1 -submodule of M_1 .*

PROOF. (1) It can be easily proved by using a similar argument in the proof of Theorem 2.12 (1).

- (2) Suppose that K_1 is a graded 2-absorbing primary R_1 -submodule of M_1 . Then $K = K_1 \times M_2$ is a graded 2-absorbing primary R -submodule of M by ([9], Theorem 18). Hence K is a graded ϕ -2-absorbing R -submodule of M for any ϕ . The remaining of this proof is similar to Theorem 2.12 (2). □

THEOREM 2.15. *Let M be a G -graded R -module $g \in G$ and K be a g - ϕ -primary R -submodule of M . Suppose that $x \in R_e$ and $m \in M_g$ such that $xm \in \phi(K)$, $x \notin \text{Grad}((K :_R M))$ and $m \notin K$. Then*

- (1) $xK_g \subseteq \phi(K)$.
- (2) $(K :_{R_e} M)m \subseteq \phi(K)$.
- (3) $(K :_{R_e} M)K_g \subseteq \phi(K)$.

PROOF. (1) Suppose that $xK_g \not\subseteq \phi(K)$. Then there exists $k \in K_g$ such that $xk \notin \phi(K)$, and then $x(m+k) \notin \phi(K)$. Since $x(m+k) \in K$ and $x \notin \text{Grad}((K :_R M))$, we deduce that $m+k \in K$ as K is a g - ϕ -primary R -submodule of M . So $m \in K$, which is a contradiction. So, $xK_g \subseteq \phi(K)$.

- (2) Suppose that $(K :_{R_e} M)m \not\subseteq \phi(K)$. Then there exists $y \in (K :_{R_e} M)$ such that $ym \notin \phi(K)$, and then $(x+y)m \notin \phi(K)$ as $xm \in \phi(K)$. Since $ym \in K$, we get $(x+y)m \in K$. Since $m \notin K$, we have that $x+y \in \text{Grad}((K :_R M))$. Hence $x \in \text{Grad}((K :_R M))$, which is a contradiction.

- (3) Suppose that $(K :_{R_e} M)K_g \not\subseteq \phi(K)$. Then there exist $y \in (K :_{R_e} M)$ and $k \in K_g$ such that $yk \notin \phi(K)$. By (1) and (2), $(x+y)(m+k) \in K - \phi(K)$.

So, either $x + y \in \text{Grad}((K :_R M))$ or $m + k \in K$. Thus we have either $x \in \text{Grad}((K :_R M))$ or $m \in K$, which is a contradiction. \square

REMARK 2.16. *Note that if K is a g - ϕ -primary R -submodule of M which is not g -primary, then there exist $x \in R_e$ and $m \in M_g$ such that $xm \in \phi(K)$, $x \notin \text{Grad}((K :_R M))$ and $m \notin K$. So, every g - ϕ -primary R -submodule, which is not g -primary, satisfies Theorem 2.15.*

THEOREM 2.17. *Let M be a G -graded R -module, $g \in G$ and K be a g - ϕ -2-absorbing R -submodule of M . Suppose that $x, y \in R_e$ and $m \in M_g$ such that $xym \in \phi(K)$, $xy \notin (K :_R M)$, $xm \notin K$ and $ym \notin K$. Then*

- (1) $xyK_g \subseteq \phi(K)$.
- (2) $x(K :_{R_e} M)m \subseteq \phi(K)$.
- (3) $y(K :_{R_e} M)m \subseteq \phi(K)$.
- (4) $(K :_{R_e} M)^2m \subseteq \phi(K)$.

PROOF. (1) Suppose that $xyK_g \not\subseteq \phi(K)$. Then there exists $k \in K_g$ with $xyk \notin \phi(K)$, and then $xy(m+k) \notin \phi(K)$. Since $xy(m+k) = xym + xyk \in K$ and $xy \notin (K :_R M)$, we conclude that $x(m+k) \in K$ or $y(m+k) \in K$. So, $xm \in K$ or $ym \in K$, which is a contradiction. Thus $xyK \subseteq \phi(K)$.

(2) Suppose that $x(K :_{R_e} M)m \not\subseteq \phi(K)$. Then there exists $a \in (K :_{R_e} M)$ such that $xam \notin \phi(K)$, and then $x(y+a)m \notin \phi(K)$ as $xym \in \phi(K)$. Since $am \in K$, we obtain that $x(y+a)m \in K$. Then $xm \in K$ or $(y+a)m \in K$ or $x(y+a) \in (K :_R M)$. Hence $xm \in K$ or $ym \in K$ or $xy \in (K :_R M)$, which is a contradiction. Hence, $x(K :_{R_e} M)m \subseteq \phi(K)$.

(3) It can be easily proved by using a similar argument in the proof of part (2).

(4) Assume that $(K :_{R_e} M)^2m \not\subseteq \phi(K)$. Then there exist $a, b \in (K :_{R_e} M)$ such that $abm \notin \phi(K)$, and then by parts (2) and (3), $(x+a)(y+b)m \notin \phi(K)$. Clearly, $(x+a)(y+b)m \in K$. Then $(x+a)m \in K$ or $(y+b)m \in K$ or $(x+a)(y+b) \in (K :_R M)$. Therefore, $xm \in K$ or $ym \in K$ or $xy \in (K :_R M)$, which is a contradiction. Consequently, $(K :_{R_e} M)^2m \subseteq \phi(K)$. \square

REMARK 2.18. *Note that if K is a g - ϕ -2-absorbing R -submodule of M which is not g -2-absorbing, then there exist $x, y \in R_e$ and $m \in M_g$ such that $xym \in \phi(K)$, $xy \notin (K :_R M)$, $xm \notin K$ and $ym \notin K$. So, every g - ϕ -2-absorbing R -submodule, which is not g -2-absorbing, satisfies Theorem 2.17.*

THEOREM 2.19. *Let M be a G -graded R -module, $g \in G$ and K be a g - ϕ -2-absorbing primary R -submodule of M . Suppose that $x, y \in R_e$ and $m \in M_g$ such that $xym \in \phi(K)$, $xy \notin (K :_R M)$, $xm \notin \text{Grad}_M(K)$ and $ym \notin \text{Grad}_M(K)$. Then*

- (1) $xyK_g \subseteq \phi(K)$.
- (2) $x(K :_{R_e} M)m \subseteq \phi(K)$.
- (3) $y(K :_{R_e} M)m \subseteq \phi(K)$.
- (4) $(K :_{R_e} M)^2m \subseteq \phi(K)$.

PROOF. It can be easily proved by using a similar argument in the proof of Theorem 2.17. \square

REMARK 2.20. *Note that if K is a g - ϕ -2-absorbing primary R -submodule of M which is not g -2-absorbing primary, then there exist $x, y \in R_e$ and $m \in M_g$ such that $xym \in \phi(K)$, $xy \notin (K :_R M)$, $xm \notin \text{Grad}_M(K)$ and $ym \notin \text{Grad}_M(K)$. So, every*

g - ϕ -2-absorbing primary R -submodule, which is not g -2-absorbing primary, satisfies Theorem 2.19.

THEOREM 2.21. *Let M be a G -graded R -module and $g \in G$. If K is a g - ϕ -2-absorbing primary R -submodule of M that is not g -2-absorbing primary, then $(K :_{R_e} M)^2 K_g \subseteq \phi(K)$.*

PROOF. Since K is a g - ϕ -2-absorbing primary R -submodule of M that is not g -2-absorbing primary, there exist $x, y \in R_e$ and $m \in M_g$ such that $xym \in \phi(K)$, $xy \notin (K :_R M)$, $xm \notin \text{Grad}_M(K)$ and $ym \notin \text{Grad}_M(K)$. Suppose that $(K :_{R_e} M)^2 K_g \not\subseteq \phi(K)$. Then there are $a, b \in (K :_{R_e} M)$ and $k \in K_g$ such that $abk \notin \phi(K)$. By Theorem 2.19, we get $(x+a)(y+b)(m+k) \in K - \phi(K)$. So, $(x+a)(m+k) \in \text{Grad}_M(K)$ or $(y+b)(m+k) \in \text{Grad}_M(K)$ or $(x+a)(y+b) \in (K :_R M)$. Therefore, $xm \in \text{Grad}_M(K)$ or $ym \in \text{Grad}_M(K)$ or $xy \in (K :_R M)$, which is a contradiction. Hence, $(K :_{R_e} M)^2 K_g \subseteq \phi(K)$. \square

PROPOSITION 2.22. *Let M be a graded R -module and K be a graded ϕ -2-absorbing primary R -submodule of M . If $\phi(K)$ is a graded 2-absorbing primary R -submodule of M , then K is a graded-2-absorbing primary R -submodule of M .*

PROOF. Assume that $x, y \in h(R)$ and $m \in h(M)$ such that $xym \in K$. If $xym \in \phi(K)$, then we conclude that $xm \in \text{Grad}_M(\phi(K)) \subseteq \text{Grad}_M(K)$ or $ym \in \text{Grad}_M(\phi(K)) \subseteq \text{Grad}_M(K)$ or $xy \in (\phi(K) :_R M) \subseteq (K :_R M)$ since $\phi(K)$ is graded 2-absorbing primary, and so the result holds. If $xym \notin \phi(K)$, then the result holds easily since K is graded ϕ -2-absorbing primary. \square

THEOREM 2.23. *Let M be a graded R -module and K be a graded R -submodule of M with $\phi(\text{Grad}_M(K)) \subseteq \phi(K)$. If $\text{Grad}_M(K)$ is a graded ϕ -prime R -submodule of M , then K is a graded ϕ -2-absorbing primary R -submodule of M .*

PROOF. Let $x, y \in h(R)$ and $m \in h(M)$ such that $xym \in K - \phi(K)$ and $xm \notin \text{Grad}_M(K)$. Since $\text{Grad}_M(K)$ is a graded ϕ -prime R -submodule and $xym \in \text{Grad}_M(K) - \phi(\text{Grad}_M(K))$, $y \in (\text{Grad}_M(K) :_R M)$. So, $ym \in \text{Grad}_M(K)$. Consequently, K is a graded ϕ -2-absorbing primary R -submodule of M . \square

Acknowledgement

This research was funded by the Deanship of Scientific Research at Princess Nourah bint Abdulrahman University through the Fast-track Research Funding Program.

References

- [1] R. Abu-Dawwas, M. Bataineh and H. Shashan, Graded generalized 2-absorbing submodules, Beiträge zur Algebra und Geometrie / Contributions to Algebra and Geometry, (2020), <https://doi.org/10.1007/s13366-020-00544-1>.
- [2] A. S. Alshehry and R. Abu-Dawwas, On graded ϕ -prime submodules, arXiv:2102.04155v1, submitted.
- [3] K. Al-Zoubi and R. Abu-Dawwas, On graded 2-absorbing and weakly graded 2-absorbing submodules, Journal of Mathematical Sciences: Advances and Applications, 28 (2014), 45-60.
- [4] K. Al-Zoubi, R. Abu-Dawwas and S. Ceken, On graded 2-absorbing and graded weakly 2-absorbing ideals, Hacettepe Journal of Mathematics and Statistics, 48 (3) (2019), 724-731.
- [5] K. Al-Zoubi and M. Al-Azaizeh, Some properties of graded 2-absorbing and graded weakly 2-absorbing submodules, Journal of Nonlinear Sciences and Applications, 12 (2019), 503-508.
- [6] K. Al-Zoubi and N. Sharafat, On graded 2-absorbing primary and graded weakly 2-absorbing primary ideals, Journal of the Korean Mathematical Society, 54 (2) (2017), 675-684.

- [7] S. E. Atani, On graded weakly prime submodules, International Mathematical Forum, 1 (2) (2006), 61-66.
- [8] S. E. Atani and F. E. K. Saraei, Graded modules which satisfy the gr-radical formula, Thai Journal of Mathematics, 8 (1) (2010), 161-170.
- [9] E. Y. Celikel, On graded 2-absorbing primary submodules, International Journal of Pure and Applied Mathematics, 109 (4) (2016), 869-879.
- [10] J. Escoriza and B. Torrecillas, Multiplication objects in commutative Grothendieck category, Communications in Algebra, 26 (1998), 1867-1883.
- [11] F. Farzalipour, P. Ghiasvand, On the union of graded prime submodules, Thai Journal of Mathematics, 9 (1) (2011), 49-55.
- [12] R. Hazrat, Graded rings and graded Grothendieck groups, Cambridge University press, 2016.
- [13] K. Khaksari and F. R. Jahromi, Multiplication graded modules, International Journal of Algebra, 7 (1) (2013), 17-24.
- [14] H. Mostafanasab, Ü. Tekir, E. Y. Celikel, E. A. Ugurlu, G. Ulucak and A. Y. Darani, Generalizations of 2-absorbing and 2-absorbing primary submodules, Hacettepe Journal of Mathematics and Statistics, 48 (4) (2019), 1001-1016.
- [15] C. Nastasescu and F. Oystaeyen, Methods of graded rings, Lecture Notes in Mathematics, 1836, Springer-Verlag, Berlin, 2004.
- [16] K. H. Oral, Ü. Tekir and A. G. Agargün, On graded prime and primary submodules, Turkish Journal of Mathematics, 35 (2011), 159-167.
- [17] M. Refai, M. Hailat and S. Obiedat, Graded radicals and graded prime spectra, Far East Journal of Mathematical Sciences, (2000), 59-73.
- [18] H. Saber, T. Alraqad and R. Abu-Dawwas, On graded s -prime submodules, AIMS Mathematics, 6 (3) (2020), 2510-2524.

DEPARTMENT OF MATHEMATICAL SCIENCES, FACULTY OF SCIENCES, PRINCESS NOURAH BINT ABDULRAHMAN UNIVERSITY, P.O. BOX 84428, RIYADH 11671, SAUDI ARABIA.

Email address: asalshihry@pnu.edu.sa

DEPARTMENT OF MATHEMATICS, YARMOUK UNIVERSITY, IRBID, JORDAN

Email address: rrashid@yu.edu.jo