

Further explorations have shown that convolutions of log-concave real distributions are log-concave (hence unimodal) (see [Mer98]), and that the convolution of a unimodal (but not symmetric) distribution with itself can have any number of modes (see [Sat93]). In the same line of questioning, we prove the following result:

Theorem 1.1. *Let $n \in \mathbb{N}$ be greater or equal to 1. Then there exists a discrete log-concave distribution p_n and a discrete bimodal distribution q_n , both symmetric about 0, such that their convolution $p_n * q_n$ has exactly n modes.*

We also prove a continuous variant, which directly contradicts the main statement¹ in [HLL⁺17]:

Theorem 1.2. *Let $n \in \mathbb{N}$ be greater or equal to 1. Then there exists an absolutely continuous distribution on \mathbb{R} whose density function f_n is smooth, symmetric about 0 and log-concave (hence unimodal), and an absolutely continuous distribution on \mathbb{R} whose density function g_n is smooth, symmetric about 0 and bimodal, such that the convolution $f_n * g_n$ has at least n modes.*

Remark 1.3. *In fact, we can ask that $f_n * g_n$ have exactly n modes; proving it makes the demonstration much more tedious, yet not any deeper.*

In what follows, the reader is reminded of the main definitions in Section 2. Theorem 1.1 is proved in Section 3, and Theorem 1.2 is proved in Section 4.

2. DEFINITIONS

In this note, we will consider absolutely continuous distributions on \mathbb{R} with continuous density functions and discrete distributions on \mathbb{Z} .

The *modes* of an absolutely continuous distribution with continuous density function f are the local maxima of f .

We will say that an absolutely continuous distribution with continuous density function f is *strictly n -modal in $\{a_1, \dots, a_n\}$* if there exists $a_1 < b_1 < a_2 < \dots < b_{n-1} < a_n \in \mathbb{R}$ such that f is strictly increasing on $[-\infty, a_1]$ and $[b_i, a_{i+1}]$ for $i = 1, \dots, n-1$, and strictly decreasing on $[a_n, \infty]$ and $[a_i, b_i]$ for $i = 1, \dots, n-1$.

We will say that a discrete distribution with probability mass function p has a mode in $\{m, m+1, \dots, m+k\} \subset \mathbb{Z}$ if $p(m-1) < p(m) = p(m+1) = \dots = p(m+k) > p(m+k+1)$. We say that it is *n -modal* if it has exactly n modes.

In the discrete case, a mode is often required to be a global maximum of the mass function. This nuance in definition matters not, as all our discrete modes will be global maxima.

The density function f of an absolutely continuous distribution is called *log-concave* if $f(\alpha x + (1-\alpha)y) \geq f(x)^\alpha f(y)^{1-\alpha}$ for all $x, y \in \mathbb{R}$ and all $\alpha \in [0, 1]$. If f is strictly positive, it is equivalent the concavity of $\log \circ f$.

The mass function p of a discrete distribution is called *log-concave* if $p(m)^2 \geq p(m-1)p(m+1)$ for all $m \in \mathbb{Z}$, and if its support is a contiguous

¹The mistake can be found in the proof of Theorem 1 in [HLL⁺17]: Equations (3) and (4) do not imply Equation (5).

interval, *i.e.* if there exists $m_1, m_2 \in \mathbb{Z}$ such that $m_1 < m_2$, $p(m) = 0$ for all $m \leq m_1$ and all $m \geq m_2$, and $p(m) > 0$ for all $m_1 < m < m_2$. The second condition is sometimes omitted.

Remark 2.1. *It is easy to show that log-concavity implies unimodality both in the continuous and the discrete case.*

As usual, the convolution of density functions $f, g : \mathbb{R} \rightarrow \mathbb{R}$ (and by extension the convolution of the two associated absolutely continuous distributions) is defined as

$$f * g(x) = \int_{\mathbb{R}} f(t)g(x-t)dt$$

for all $x \in \mathbb{R}$.

Similarly, the convolution of mass functions $p, q : \mathbb{Z} \rightarrow [0, 1]$ (and by extension the convolution of the two associated discrete distributions) is defined as

$$p * q(m) = \sum_{k \in \mathbb{Z}} p(k)q(m-k)$$

for all $m \in \mathbb{Z}$.

3. DISCRETE CASE

Proof of Theorem 1.1. The case $n = 1$ is almost trivial: consider $p_1 := \frac{1_{\{-1,0,1\}}}{3}$ and $q_1 := \frac{1_{\{-1,1\}}}{2}$, where $1_A : \mathbb{Z} \rightarrow \{0, 1\}$ refers to the indicator function of A for any $A \subset \mathbb{Z}$.

Now for $n \geq 2$, let us define $\tilde{p}_n = 1_{\{-n+2, \dots, 0, \dots, n-2\}}$ and $p_n = \tilde{p}_n \cdot \frac{1}{2n-3}$. The function p_n is clearly a log-concave distribution and symmetric about 0.

If n is even, let \tilde{q}_n be as such: for $k = 1, \dots, \frac{n}{2} - 1$, we let $\tilde{q}_n(2k-1) = \tilde{q}_n(2k) = 2k+1$ and $\tilde{q}_n(2n-2-2k) = \tilde{q}_n(2n-1-2k) = 2k$. We also let $\tilde{q}_n(n-1) = n$, $\tilde{q}_n(0) = 1$ and $\tilde{q}_n(k) = 0$ for any $k > 2n-3$. We define \tilde{q}_n on $\mathbb{Z}_{\leq 0}$ symmetrically.

Let $C_n := \sum_{m \in \mathbb{Z}} \tilde{q}_n(m)$ and $q_n := \frac{\tilde{q}_n}{C_n}$. Then q_n is symmetric about 0, and it is a bimodal distribution whose modes are in $\{-n+1, n-1\}$. It has a minimum in 0.

See Figure 1 to see the case $n = 6$ illustrated.

The convolution $p_n * q_n$ is clearly symmetric about 0.

For any $m \in \mathbb{Z}$, the finite difference satisfies $D(p_n * q_n)(m) = (p_n * q_n)(m) - (p_n * q_n)(m-1) = (D(p_n) * q_n)(m)$. As $D(p_n)(l)$ is equal to $\frac{1}{2n-3}$ if $l = -n+2$, to $\frac{-1}{2n-3}$ if $l = n-1$ and to 0 otherwise, we get that $D(p_n * q_n)(m) = \frac{q_n(m+n-2) - q_n(m-n+1)}{2n-3}$.

We see that $D(p_n * q_n)(m)$ is 0 if $m > (2n-3) + (n-1) = 3n-4$ and is strictly negative if $3n-4 \geq m > n-1$. For $m = 1, \dots, n-1$, the finite difference $D(p_n * q_n)(m)$ is equal to $\frac{1}{(2n-3)C_n}$ if m is odd and to $\frac{-1}{(2n-3)C_n}$ if m is even.




FIGURE 1. Case $n = 6$. The \bullet correspond to \tilde{q}_6 , the \times to \tilde{p}_6 and the $+$ to where they take the same value.

By symmetry of p_n and q_n , if $m \leq 0$, we have that

$$\begin{aligned} D(p_n * q_n)(m) &= \frac{q_n(m+n-2) - q_n(m-n+1)}{2n-3} = \\ &= -\frac{q_n(-m+n-1) - q_n(-m-n+2)}{2n-3} = \\ &= -\frac{q_n(-(m-1)+n-2) - q_n(-(m-1)-n+1)}{2n-3} = -D(p_n * q_n)(-(m-1)). \end{aligned}$$

From this, we get that $p_n * q_n$ has exactly n modes, located in $-n+1, \dots, -3, -1, 1, 3, \dots, n-1$.

The convolution $\tilde{p}_n * \tilde{q}_n$ is illustrated in Figure 2 in the case $n = 6$ (remember that $p_n * q_n = \frac{\tilde{p}_n * \tilde{q}_n}{(2n-3)C_n}$).

The case where n is odd (and strictly greater than 1) is very similar. Let \tilde{q}_n be as such: for $k = 1, \dots, \frac{n-1}{2} - 1$ we let $\tilde{q}_n(2k-1) = \tilde{q}_n(2k) = 2k+1$, and for $k = 1, \dots, \frac{n-1}{2}$ we let $\tilde{q}_n(2n-2-2k) = \tilde{q}_n(2n-1-2k) = 2k$. We also let $\tilde{q}_n(n-2) = n$, $\tilde{q}_n(0) = 1$ and $\tilde{q}_n(k)$ be 0 for any $k > 2n-3$.

As before, we define \tilde{q}_n on $\mathbb{Z}_{\leq 0}$ symmetrically, let $C_n := \sum_{m \in \mathbb{Z}} \tilde{q}_n(m)$ and $q_n := \frac{\tilde{q}_n}{C_n}$. Then q_n is symmetric about 0, and it is a bimodal distribution whose modes are in $\{-n+2, n-2\}$. It has a minimum in 0.

The case $n = 7$ is illustrated in Figure 3.

As above, $D(p_n * q_n)(m) = \frac{q_n(m+n-2) - q_n(m-n+1)}{2n-3}$. Thus we see that $D(p_n * q_n)(m)$ is 0 if $m > (2n-3) + (n-1) = 3n-4$ and is strictly negative if $3n-4 \geq m > n-1$. For $m = 1, \dots, n-1$, the finite difference $D(p_n * q_n)(m)$ is equal to $\frac{1}{(2n-3)C_n}$ if m is even and to $\frac{-1}{(2n-3)C_n}$ if m is odd.




FIGURE 2. The convolution product $\tilde{p}_n * \tilde{q}_n$ in the case $n = 6$.




FIGURE 3. Case $n = 7$. The \bullet correspond to \tilde{q}_7 , the \times to \tilde{p}_7 and the $+$ to where they take the same value.

By symmetry, $D(p_n * q_n)(m) = -D(p_n * q_n)(-(m-1))$ if $m \leq 0$.

From this, we get that $p_n * q_n$ has exactly n modes, located in $-n + 1, \dots, -2, 0, 2, \dots, n - 1$. The proof is complete. \square

Remark 3.1. Note that if we wanted p_n to be such that there is a strict global maximum in 0, we could replace \tilde{p}_n in the proof of Theorem 1.1 by the restriction to \mathbb{Z} of $x \mapsto \exp\left(-\left[\frac{x}{n-2+\frac{1}{2}}\right]^{2i}\right)$ for $i \in \mathbb{N}$ large enough, and then proceed as above.

4. CONTINUOUS CASE

We come up with a smooth variant of the construction used in Section 3. Given a function $p : \mathbb{Z} \rightarrow \mathbb{Z}$, we define

$$\Phi(p) := \sum_{k \in \mathbb{Z}} p(k) 1_{]k-\frac{1}{2}, k+\frac{1}{2}]} : \mathbb{R} \rightarrow \mathbb{R}.$$

Note that $\Phi(p)|_{\mathbb{Z}} = p$.

Lemma 4.1. *Let $p, q : \mathbb{Z} \rightarrow \mathbb{Z}$ be two functions with finite support.*

*Then the convolution $\Phi(p) *_{\mathbb{R}} \Phi(q)$ over \mathbb{R} (where $\Phi(p) *_{\mathbb{R}} \Phi(q)(x) = \int_{\mathbb{R}} \Phi(p)(t) \Phi(q)(x-t) dt$) is a continuous piecewise affine function which is affine on each interval $[l, l+1]$ for $l \in \mathbb{Z}$. Moreover, the convolution $p *_{\mathbb{Z}} q$ over \mathbb{Z} (where $p *_{\mathbb{Z}} q(m) = \sum_{k \in \mathbb{Z}} p(k) q(m-k)$) coincides with the restriction to \mathbb{Z} of $\Phi(p) *_{\mathbb{R}} \Phi(q)$:*

$$(\Phi(p) *_{\mathbb{R}} \Phi(q))|_{\mathbb{Z}} = p *_{\mathbb{Z}} q.$$

*In particular, $p *_{\mathbb{Z}} q$ and $\Phi(p) *_{\mathbb{R}} \Phi(q)$ have the same modes.*

The proof of Lemma 4.1 is straightforward.

Denote by $\| - \|_{\infty} : f \mapsto \|f\|_{\infty} = \sup_{x \in \mathbb{R}} (|f(x)|)$ the uniform norm. We need the following Lemma, which is easy to prove using classical smooth approximation tricks:

Lemma 4.2. *For any $n \in \mathbb{N}$, there exists a family of smooth symmetric about 0 density functions $\{g_n^N\}_{N \in \mathbb{N}}$ such that:*

- (1) *As N goes to ∞ , g_n^N converges to $\Phi(q_n)$ in norm L_1 .*
- (2) *There exists $M > 0$ such $\|g_n^N\|_{\infty} \leq M$ for all N .*
- (3) *If n is even (respectively, odd and strictly greater than 1), g_n^N is strictly increasing from $-\infty$ to $-n+1$ (respectively $-n+2$) and strictly decreasing from $-n+1$ (respectively $-n+2$) to 0 (and symmetrically so on \mathbb{R}_+) for all N . In particular, g_n^N is bimodal.*
- (4) *If $n=1$, g_1^N is strictly increasing from $-\infty$ to -1 and strictly decreasing from -1 to 0 (and symmetrically so on \mathbb{R}_+) for all N . In particular, g_1^N is bimodal.*

Proof of Lemma 4.2. We assume that $n \geq 2$ - the case $n=1$ is similar. Given $N \in \mathbb{N}$, one can for example consider the function

$$h_N := x \mapsto \frac{1}{1 + \exp(\frac{Nx}{x^2-1})} 1_{\{-1 < x < 1\}} + 1_{\{x \geq 1\}},$$

illustrated in Figure 4, which is smooth and approximates the Heavyside step function as $N \rightarrow \infty$.

Let also $b_N : \mathbb{Z} \rightarrow \mathbb{R}$ be such that $b_N(k) = 0$ if $k \notin \{-2n+3, \dots, 2n-3\}$, that $b_N(k) = \frac{1}{N}$ if $k \in \{-2n+3, \dots, 2n-3\}$ is even, and $b_N(k) = -\frac{1}{N}$ if $k \in \{-2n+3, \dots, 2n-3\}$ is odd. Then for any $n \geq 2$ and $N \in \mathbb{N}$, one can define a smooth and symmetric about 0 function \tilde{g}_n^N as follows: let $\tilde{g}_n^N(x) = 0$ for any $x < -2n+2$ or $x > 2n-2$, and let

$$\tilde{g}_n^N(x) = (\tilde{q}_n(k+1) + b_N(k+1) - \tilde{q}_n(k) - b_N(k)) \cdot h_N(2(x-k-0.5)) + \tilde{q}_n(k) + b_N(k)$$




FIGURE 4. The graph of $h_N : x \mapsto \frac{1}{1+\exp(\frac{Nx}{x^2-1})} \mathbf{1}_{\{-1 < x < 1\}} + \mathbf{1}_{\{x \geq 1\}}$ for $N = 10$.




FIGURE 5. The graph of \tilde{g}_n^N , which serves as a smooth approximation of $\Phi(\tilde{q}_n)$, for $n = 5$ and $N = 10$.

for $x \in [k, k+1]$ and $k \in \{-2n+2, 2n-2\}$, where \tilde{q}_n is as in the proof of Theorem 1.1. The case $n = 5$ and $N = 10$ is illustrated in Figure 5. Then \tilde{g}_n^N converges to $\Phi(\tilde{q}_n)$ in norm L^1 as $N \rightarrow \infty$, and the family of functions

$$g_n^N := \frac{\tilde{g}_n^N}{\int_{\mathbb{R}} \tilde{g}_n^N(x) dx}$$

satisfies conditions (1) to (4). □

We can now prove Theorem 1.2.

Proof of Theorem 1.2. Consider the mass functions p_n and q_n defined in the proof of Theorem 1.1.

For any $N \in \mathbb{N}_{>0}$, let f_n^N be defined by

$$f_n^N(x) = \frac{\exp(-(\frac{x}{n-2+\frac{1}{2}})^{2N})}{\int_{\mathbb{R}} \exp(-(\frac{y}{n-2+\frac{1}{2}})^{2N}) dy}$$

if $n \geq 2$ and

$$f_1^N(x) = \frac{\exp(-(\frac{2x}{3})^{2N})}{\int_{\mathbb{R}} \exp(-(\frac{2y}{3})^{2N}) dy}$$

if $n = 1$.

This defines a family of smooth log-concave distributions that are symmetric about 0. Moreover, as N goes to infinity, f_n^N converges in norm L^1 to $\Phi(p_n)$.

Let $\{g_n^N\}_{N \in \mathbb{N}}$ be as in Lemma 4.2. Using properties (1) and (2), we see that

$$\begin{aligned} & \|f_n^N * g_n^N - \Phi(p_n) * \Phi(q_n)\|_\infty \leq \\ & \|f_n^N * g_n^N - \Phi(p_n) * g_n^N\|_\infty + \|\Phi(p_n) * g_n^N - \Phi(p_n) * \Phi(q_n)\|_\infty \\ & \leq M \|f_n^N - \Phi(p_n)\|_{L^1} + 2 \|g_n^N - \Phi(q_n)\|_{L^1} \xrightarrow{N \rightarrow \infty} 0. \end{aligned}$$

In particular, $(f_n^N * g_n^N)|_{\mathbb{Z}}$ converges uniformly to $(\Phi(p_n) * \Phi(q_n))|_{\mathbb{Z}} = p_n * q_n$ (see Lemma 4.1).

We have seen in the proof of Theorem 1.1 that if n is even,

$$\begin{aligned} p_n * q_n(-n) &< p_n * q_n(-n+1) > p_n * q_n(-n+2) < \dots \\ &< p_n * q_n(-1) > p_n * q_n(0) < p_n * q_n(1) > \dots \\ &> p_n * q_n(n-2) < p_n * q_n(n-1) > p_n * q_n(n). \end{aligned}$$

Hence for N large enough, we also have

$$\begin{aligned} f_n^N * g_n^N(-n) &< f_n^N * g_n^N(-n+1) > f_n^N * g_n^N(-n+2) < \dots \\ &< f_n^N * g_n^N(-1) > f_n^N * g_n^N(0) < f_n^N * g_n^N(1) > \dots \\ &> f_n^N * g_n^N(n-2) < f_n^N * g_n^N(n-1) > f_n^N * g_n^N(n), \end{aligned}$$

which means (using Rolle's theorem) that $f_n^N * g_n^N$ has at least n modes. Moreover, the number of modes must be even, since 0 is not a mode (and the convolution is symmetric with respect to 0).

The same reasoning applies if n is odd and strictly greater than 1, in which case

$$\begin{aligned} p_n * q_n(-n) &< p_n * q_n(-n+1) > p_n * q_n(-n+2) < \dots \\ &> p_n * q_n(-1) < p_n * q_n(0) > p_n * q_n(1) < \dots \\ &> p_n * q_n(n-2) < p_n * q_n(n-1) > p_n * q_n(n) \end{aligned}$$

and

$$\begin{aligned} f_n^N * g_n^N(-n) &< f_n^N * g_n^N(-n+1) > f_n^N * g_n^N(-n+2) < \dots \\ &> f_n^N * g_n^N(-1) < f_n^N * g_n^N(0) > f_n^N * g_n^N(1) < \dots \\ &> f_n^N * g_n^N(n-2) < f_n^N * g_n^N(n-1) > f_n^N * g_n^N(n) \end{aligned}$$

for N large enough, which again means that $f_n^N * g_n^N$ has at least n modes. Moreover, as $(g_n^N)'(-n+2-\frac{1}{2}) = -(g_n^N)'(n-2+\frac{1}{2}) > 0$ by property (3) we see that the second derivative of $f_n^N * g_n^N$ in 0 is negative if N is large enough by considering $(f_n^N)'$ (which converges to $\frac{1}{2}(-\delta_{-n+2-\frac{1}{2}} + \delta_{n-2+\frac{1}{2}})$, where δ_x is the Dirac distribution in $x \in \mathbb{R}$). Hence, 0 is a mode of $f_n^N * g_n^N$, which has an odd number of modes.

If $n = 1$, likewise, $p_1 * q_1(-1) < p_1 * q_1(0) > p_1 * q_1(1)$ and $f_1^N * g_1^N(-1) < f_1^N * g_1^N(0) > f_1^N * g_1^N(1)$ for N large enough, which implies that $f_1^N * g_1^N$ has at least one mode. \square

Remark 4.3. *By adding more technical conditions and being more careful than we are in Lemma 4.2 when defining the functions g_n^N , it can be tediously shown that we can make it so that $f_n^N * g_n^N$ has exactly n modes.*

*The idea behind it is to make sure that the convolution alternates between being strictly convex and strictly concave, so that it admits exactly one mode on each interval on which it is convex. To do so, one needs to consider the second derivative $(f_n^N)' * (g_n^N)'$ of $f_n^N * g_n^N$, and use the fact that $(f_n^N)'$ converges nicely to a normalized sum of Dirac distributions and that g_n^N can be chosen so that its derivative also converges to a weighted sum of Dirac distributions. Additional conditions concerning the behavior of g_n^N on certain pairs of segments must also be added.*

REFERENCES

- [AK07] Mohammed I. Ageel and Anwer Khurshid. Simple proofs of two results on convolutions of discrete unimodal distributions. *International Journal of Statistical Sciences*, 6 (Special Issue):39–42, 2007.
- [Bak18] Matthew Baker. Hodge theory in combinatorics. *Bulletin (new series) of the American Mathematical Society*, 55:57–80, 2018.
- [DJD88] S. W. Dharmadhikari and K. Joag-Dev. *Unimodality, Convexity, and Applications*. Academic Press, New York, 1988.
- [HLL⁺17] Linke Hou, Xiaowu Li, Juan Liang, Lin Wang, and Mingsheng Zhang. On convolution of unimodal distribution and multimodal distribution. *Advances and Applications in Statistics*, 50:385–395, 2017.
- [Mer98] Milan Merkle. Convolutions of logarithmically concave functions. *Publikacije Elektrotehnickog fakulteta. Serija Matematika Publikacije Elektrotehnickog fakulteta. Serija Matematika*, 9:113–117, 1998.
- [Sat93] Ken-Iti Sato. Convolution of unimodal distributions can produce any number of mode. *The Annals of Probability*, 21(3):1543–1549, 1993.
- [SW14] Adrien Saumard and Jon A. Wellner. Log-concavity and strong log-concavity: A review. *Statist. Surv.*, 8:45–114, 2014.

CHARLES ARNAL, UNIV. PARIS 6, IMJ-PRG, FRANCE.
Email address: `charles.arnal@imj-prg.fr`