

# Hilbert series of generic ideals in products of projective spaces

Ralf Fröberg

## Abstract

If  $k[x_1, \dots, x_n]/I = R = \sum_{i \geq 0} R_i$ ,  $k$  a field, is a standard graded algebra, the Hilbert series of  $R$  is the formal power series  $\sum_{i \geq 0} \dim_k R_i t^i$ . It is known already since Macaulay which power series are Hilbert series of graded algebras [12]. A much harder question is which series are Hilbert series if we fix the number of generators of  $I$  and their degrees, say for ideals  $I = (f_1, \dots, f_r)$ ,  $\deg f_i = d_i$ ,  $i = 1, \dots, r$ . In some sense "most" ideals with fixed degrees of their generators have the same Hilbert series. There is a conjecture for the Hilbert series of those "generic" ideals, see below. In this paper we make a conjecture, and prove it in some cases, in the case of generic ideals of fixed degrees in the coordinate ring of  $\mathbb{P}^1 \times \mathbb{P}^1$ , which might be easier to prove.

KEYWORDS:  $\mathbb{P}^1 \times \mathbb{P}^1$ , Hilbert series, generic ideals

AMS subject classification: 14N10, 13D40

## 1 Background

The conjecture and the results in section 2 below are inspired by the corresponding conjecture and results in the singly graded case.

**Conjecture 1 ([5])** *Let  $I = (f_1, \dots, f_r) \subset k[x_1, \dots, x_n]$ ,  $k$  an infinite field, be an ideal generated by generic forms  $f_i$  with  $\deg(f_i) = d_i$ , and let  $R = k[x_1, \dots, x_n]/I$ . Then*

$$R(u) = \left[ \prod_{i=1}^r (1 - u^{d_i}) / (1 - u)^n \right]_+$$

here  $[\sum a_i u^i]_+ = \sum b_i u^i$ , where  $b_i = a_i$  if for all  $j \leq i$  we have  $a_j > 0$ , and  $b_i = 0$  otherwise.

We first comment on the use of the word "generic". A polynomial of degree  $d$  in  $k[x_1, \dots, x_n]$  is a linear combination of  $\binom{n+d-1}{d}$  monomials. Thus an ideal  $(f_1, \dots, f_r)$ ,  $\deg(f_i) = d_i$ , can be considered as a point in  $A = \mathbb{A}^N$ ,  $N = \sum_{i=1}^r \binom{n+d_i-1}{d_i}$ . There is a Zariski-open subset of  $A$ , for which the Hilbert series is constant. Ideals corresponding to points in that Zariski-open set are what we call generic, see [8].

The conjecture is proved for  $r \leq n$  (trivial), for  $n \leq 2$  [5], for  $n = 3$  [1], for  $r = n + 1$  [16]. There are partial results in [10], [7], [2], [13], [3], [15], [14].

If  $J = (l_1^{d_1}, \dots, l_r^{d_r})$ , where  $l_i$  are generic linear forms. Sometimes, but not always, the Hilbert series of  $k[x_1, \dots, x_n]/J$  equals the one in the conjecture. There is a conjecture on when it does, [11, 4].

## 2 $\mathbb{P}^1 \times \mathbb{P}^1$

We are considering homogeneous ideals in the coordinate ring of  $\mathbb{P}^1 \times \mathbb{P}^1$ . Thus, let  $k$  be an infinite field,  $S = k[x_0, x_1, y_0, y_1]$  be bigraded,  $\deg(x_i) = (1, 0)$ ,  $\deg(y_j) = (0, 1)$ , and let  $I$  be a bihomogeneous ideal, so generated by bihomogeneous elements. Hence  $R = S/I$  is bigraded,  $R = \bigoplus_{i,j \geq 0} R_{i,j}$ . The Hilbert series of  $R$  is defined as  $R(u, v) = \sum \dim_k R_{i,j} u^i v^j$ . We are interested in the case when the ideal is generated by "generic" elements. Given a sequence of degrees  $\mathbf{d}_1, \dots, \mathbf{d}_r$ , we denote the space of ideals  $I = (f_1, \dots, f_r)$  where  $\deg f_i = \mathbf{d}_i$  by  $\mathbf{I}_{\mathbf{d}_1, \dots, \mathbf{d}_r}$ . An element of degree  $(d, e)$  is a linear combination of  $(d+1)(e+1)$  monomials. Thus an ideal in  $\mathbf{I}_{\mathbf{d}_1, \dots, \mathbf{d}_r}$  can be considered as a point in  $\mathbb{A}_k^N$ ,  $N = \sum_{i=1}^r (d_i+1)(e_i+1)$  where  $\mathbf{d}_i = (d_i, e_i)$ . We partially order Hilbert series termwise, so that  $\sum a_{ij} u^i v^j \geq \sum b_{ij} u^i v^j$  if  $a_{ij} \geq b_{ij}$  for all  $i, j$ .

**Theorem 2** *There are only a finite number of possibilities for Hilbert series of ideals in  $\mathbf{I}_{\mathbf{d}_1, \dots, \mathbf{d}_r}$ . There is a nonempty Zariski open part of  $\mathbf{I}_{\mathbf{d}_1, \dots, \mathbf{d}_r}$  where the Hilbert series is constant. This constant Hilbert series is the smallest possible for ideals in  $\mathbf{I}_{\mathbf{d}_1, \dots, \mathbf{d}_r}$ .*

**Proof** The corresponding theorems in the singly graded case, [8, Theorem 1], [5, Theorem p.120], and [6, Proposition 1] are easily adapted. We call points in this nonempty Zariski open set generic.

We define  $(d, e) \leq (f, g)$  if  $d \leq f$  and  $e \leq g$ , and  $(d, e) \geq (f, g)$  if  $d \geq f$  and  $e \geq g$ . Furthermore  $[\sum_{i,j} a_{ij} u^i v^j]_+ = \sum_{i,j} b_{ij} u^i v^j$ , where  $b_{ij} = a_{ij}$  if  $a_{kl} > 0$  for all  $(k, l) \leq (i, j)$  and  $b_{i,j} = 0$  otherwise.

**Lemma 3** *Let  $R = S/I$  be bigraded and  $f \in R_{i,j}$ . Then  $(R/f)(u, v) \geq [(1 - u^i v^j)(R(u, v))]_+$ ,*

**Proof** Consider the map  $f: R_{d-i, e-j} \rightarrow R_{d,e}$ . The image is largest if the map is of maximal rank, i.e., either injective or surjective, so  $\dim_k (R/f)_{d,e} \geq \max\{0, \dim R_{d,e} - \dim R_{i,j}\}$ . If  $\dim(R/f)_{d,e} = 0$ , then  $\dim(R/f)_{d+f, e+g} = 0$  for all  $(f, g) \geq 0$ .

**Lemma 4**  $[(1 - u^i v^j)[\sum_{i,j} a_{ij} u^i v^j]_+]_+ = [(1 - u^i v^j) \sum_{i,j} a_{ij} u^i v^j]_+$ .

**Proof** Easy calculation.

These two lemmas give the following.

**Theorem 5** *Let  $I = (f_1, \dots, f_r) \subset k[x_0, x_1, y_0, y_1] = S$ ,  $\deg f_i = (d_i, e_i)$ . Then  $S/I(u, v) \geq [\prod_{i=1}^r (1 - u^{d_i} v^{e_i}) / ((1 - u)^2 (1 - v)^2)]_+$ .*

We now give a conjecture in the case when the  $f_i$ 's are generic, c.f. [9]. To prove the conjecture for some fixed  $(\mathbf{d}_1, \dots, \mathbf{d}_r)$  it suffices to give one example with the conjectured series. If the conjecture is true for these parameters, then almost all ideals have the conjectured series, so we must be very unlucky if we miss the series with a random choice of coefficients.

**Conjecture 6** *Let  $I = (f_1, \dots, f_r) \subset k[x_0, x_1, y_0, y_1] = S$ ,  $\deg f_i = (d_i, e_i)$  generic. Then  $(S/I)(u, v) = [\prod_{i=1}^r (1 - u^{d_i} v^{e_i}) / ((1 - u)^2 (1 - v)^2)]_+$ .*

We have checked that the conjecture is true in the following cases. Some of these were checked by Alessandro Oneto. Except for the first class, we have used computer calculations.

1. For small  $r$  the concepts of ideal generated by generic forms and complete intersection agrees. It is well known that the conjecture is true for complete intersections.
2.  $\deg f_i = (1, 1)$  for all  $i$ , any  $r$ .
3. Some  $f_i$  of degree  $(1, 1)$ , some of degree  $(1, 2)$ , any  $r$ .
4.  $\deg f_i = (1, 2)$  for all  $i$ , any  $r$ .
5.  $\deg f_i = (2, 2)$  for all  $i$ , any  $r$ .

We also checked that the corresponding conjecture is true for  $\deg f_i = (1, 1, 1)$  for all  $i$ , any  $r$ , in  $\mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$ .

On the other hand, the corresponding conjecture in  $\mathbb{P}^2 \times \mathbb{P}^2$  cannot be true. For four generic forms of degree  $(1, 1)$  the conjecture would give that  $R_{d,d} = 0$  if  $d \gg 0$ . The correct statement is that  $\dim_k R_{d,d} = 6$  if  $d \gg 0$ .

We think that the conjecture is challenging enough, but we also give some questions.

**Question** What is the Hilbertseries for generic ideals in  $(\mathbb{P}^1)^k = \mathbb{P}^1 \times \dots \times \mathbb{P}^1$ ,  $k > 2$  times?

**Question** What is the Hilbertseries for generic ideals in  $\mathbb{P}^m \times \mathbb{P}^n$ ?

**Question** Let  $f_i$  be generic linear forms in  $k[x_1, x_2]$  and  $g_i$  generic linear forms in  $k[y_1, y_2]$ , and let  $I = (f_1^{d_1} g_1^{e_1}, \dots, f_r^{d_r} g_r^{e_r})$ . What is the Hilbert series of  $k[x_1, x_2, y_1, y_2]/I$ ?

## References

- [1] D. Anick, *Thin algebras of embedding dimension three*, J. Algebra **100**, 235–259 (1986).
- [2] M. Aubry, *Série de Hilbert d'une algèbre de polynômes quotient*, J. Algebra **100**, 235–259 (1995).
- [3] J. Backelin and A. Oneto, *On a class of power ideals*, J. Pure Appl. Algebra **219**, 3158–3180 (2015).

- [4] K. Chandler, *The geometric interpretation of the Fröberg-Iarrobino conjectures on infinitesimal neighbourhoods of points in projective space*, J. Algebra **286**, 421–455 (2005).
- [5] R. Fröberg, *An inequality for Hilbert series*, Math. Scand. **56**, 117–144 (1985).
- [6] R. Fröberg, T. Gulliksen and C. Löfwall, *Flat families of local artinian  $k$ -algebras with infinitely many Poincaré series*, Springer Lect. Notes in Math. **1183**, 170–191 (1983).
- [7] R. Fröberg and J. Hollman, *Hilbert series for ideals generated by generic forms*, J. Symb. Comp. **17** 149–157 (1994).
- [8] R. Fröberg and C. Löfwall, *On Hilbert series for commutative and noncommutative graded algebras*, J. Pure. Appl. Algebra **76**, 33–38 (1990).
- [9] R. Fröberg and S. Lundqvist, *Questions and conjectures on extremal Hilbert series*. Rev. Un. Mat. Argentina **59**(2), 415–429 (2018).
- [10] M. Hochster and D. Laksov, *The linear syzygies of generic forms*, Comm. Algebra **15**, 227–234 (1987).
- [11] A. Iarrobino, *Inverse systems of a symbolic power III. Thin algebras and fat points*, Compositio Math. **108**, 319–336 (1997).
- [12] F. Macaulay, *Some properties of enumeration in the theory of modular systems*, Proc. London Math. Soc. **26**, 531–555 (1927).
- [13] J. Migliore and R. M. Miro-Roig, *Ideals of general forms and the ubiquity of the real Lefschetz property*, J. Pure Appl. Algebra **102**, 79–107 (2003).
- [14] G. Nenashev, *A note on Fröberg’s conjecture for forms of equal degree*, C. R. Acad. Sci. Paris Ser. I **355**, 272–276 (2017).
- [15] L. Nicklasson, *On the Hilbert series of ideals generated by generic forms*, Comm. Algebra **45** no. 8, 3390–3395.
- [16] R. Stanley, *Weyl groups, the hard Lefschetz theorem, and the Sperner property*, SIAM J. Algebraic Discrete Methods **1**(2), 168–184 (1980).