

# Multivariate higher order moments in multi-state life insurance

Jamaal Ahmad<sup>1,\*</sup>

<sup>1</sup>Department of Mathematical Sciences, University of Copenhagen, Universitetsparken 5, DK-2100 Copenhagen Ø, Denmark.

\*Corresponding author. E-mail: jamaal@math.ku.dk.

May 8, 2021

## ABSTRACT

It is well-known that combining life annuities and death benefits introduce opposite effects in payments with respect to the unsystematic mortality risk on the lifetime of the insured. In a general multi-state framework with multiple product types, such joint effects are less trivial. In this paper, we consider a multivariate payment process in multi-state life insurance, where the components are defined in terms of the same Markovian state process. The multivariate present value of future payments is introduced, and we derive differential equations and product integral representations of its conditional moments and moment generating function. Special attention is given to pair-wise covariances between two present values, where results closely connected to Hattendorff type of results for the variance are derived. Numerical examples are shown in a disability model and a spouse model to illustrate applicability of the results on insurance of a single life as well as two dependent lives.

**Keywords:** Multi-state life insurance; Multivariate payment process; Unsystematic biometric risk; Conditional moments; Product integral

**2010 Mathematics Subject Classification:** 60J28; 91B30; 91G99

**JEL Classification:** G22

## 1 Introduction

In this paper, we extend the results of [4] to multivariate payment processes with components defined in terms of the same multi-state Markov process. Our main contributions

## 1 INTRODUCTION

are differential equations and product integral representations of higher order moments of the multivariate present value, which are derived via its moment generating function. The main results appear as natural multivariate generalizations to those of [4, 15], and to some extent also [1], and this is pointed out in a series of remarks throughout the paper. We give special attention to pair-wise covariances between two present values and derive results that reveal close connections to Hattendorff type of results for the variance of a present value.

The paper is motivated by the following. The widespread use of multi-state Markov models in life insurance for the past decades, dating back to at least [9], have provided a unified mathematical framework to model life insurance products with different kinds of biometric risk profiles; see e.g. an overview of these in [3, Chapter V.2]. For valuation and risk management, the present value of future payments is the main quantity of interest, and classic actuarial valuation principles relies on diversification of unsystematic biometric risk, such that the conditional expected present value constitute the prospective reserve; this principle is justified by applying the law of large numbers on a suitably large portfolio of independent and identically distributed insured. The prospective reserve satisfies the well-known Thiele's differential equations, cf. [9, 13].

In order to analyse the unsystematic biometric risk, we are therefore interested in distributional properties of the present value that goes beyond the expected value, and several authors have derived equations for its higher order moments as well as its full probability distribution. The variance in particular have had great attention historically; it is obtained as an integral expression in [9], see also [10, 13], and Hattendorff's theorem, which is formulated in the multi-state framework by [16, 14], provides particularly simple formulae in terms of the associated multivariate counting process and so-called sum at risks. In [15], differential equations are derived for non-central higher order moments, and corresponding central moments are then obtained via binomial formulas. These type of results are extended to the semi-Markovian framework in [1] to allow for duration dependency in payments and transition rates, and more recently, a general matrix-based framework for the calculation of so-called partial reserves and higher order moments using product integrals have been developed in [4] in the multi-state Markovian framework. Here, the moments are also used to approximate the density and distribution function of the present value using (orthogonal) polynomial expansions. Other ways of calculating distribution functions can be found in [8, 2], where integral and differential equations for these are derived.

In practice, an insured typically holds a combination of various product types in order to be covered in case of different kinds of life and health events. In the multi-state framework, this is without further notice handled by considering the aggregated payment process, and reserves, higher order moments and probability distributions can be calculated using the methods outlined above. However, if we wish to understand how


## 1 INTRODUCTION

the unsystematic biometric risk of the individual product types contribute to the overall unsystematic biometric risk of a contract, we need to consider multivariate payment processes and associated properties of the joint distribution. The simplest example is the product combination of life annuities and death benefits, which have opposite effect with respect to the lifetime of the insured. In the general multi-state framework, more complex and non-trivial interactions can be examined. The following two examples illustrates these motivations in relation to insurance of a single life and two dependent lives, respectively. While these two examples have different motivations, they are both mathematically handled within the general multi-state setup of this paper.

**Example 1.1** (Disability model). Consider a single insured that have the following products, with a deterministic retirement time  $T$ :

- (1) Death benefit paying  $S$  upon death before time  $T$ .
- (2) Deferred life annuity paying a benefit rate of  $b$  while alive, starting from time  $T$ .
- (3) Disability annuity paying a benefit rate of  $d$  while disabled until time  $T$ .

The state of the insured is then modelled in the classic disability model with recoveries, as depicted in Figure 1 below.



**Figure 1:** *The classic disability model with recoveries.*

In this example, we have a single insured that have a product combination with three different risk profiles that work in different directions with respect to the unsystematic biometric risk. The dependence structure between the three payment profiles is therefore of particular interest. ◦


**Example 1.2** (Spouse model). Consider a married couple that have the following products to cover each others deaths, with expiry at time  $T$ :

- (1) Widow's pension paying a benefit rate of  $b_1$  after death of her husband until her death or time  $T$ , whichever occurs first.

## 1 INTRODUCTION

- (2) Widower's pension paying a benefit rate of  $b_2$  after death of his wife until his death or time  $T$ , whichever occurs first.

The state of the married couple is then modelled in the spouse model as depicted in Figure 2.



**Figure 2:** *The spouse model.*

In the case where  $\mu \neq \tilde{\mu}$  and/or  $\nu \neq \tilde{\nu}$ , we are able to model a particular form of dependence between two lives, and therefore we are interested in the corresponding dependence between the payments to each of the individuals in the marriage. Hence, this setup provides a different motivation to analyse dependence between payments as that of Example 1.1, even though the mathematical frameworks are the same.  $\circ$

The idea of mixing various product types and analysing their interacting effect has already been discussed extensively in the concept of so-called natural hedging in life insurance, where one considers the effects on the reserves with respect to future changes in mortality rates in a portfolio consisting exactly of life annuities and death benefits; see e.g. [6]. While this concept arose in the survival model at first, an example in the multi-state model have appeared recently in [12] where the authors analyse interactions with life products and long-term care insurance using the disability model. These all relate to changes in reserves with respect to future changes in transition rates, i.e. systematic biometric risk. We think of this paper as a starting point to analyse interacting effects related to unsystematic biometric risk, with the simplest multivariate extension of existing multi-state frameworks, and the motivating Example 1.1 consist of a product combination that is in the same spirit as that of [12].

The paper is structured as follows. Section 2 introduces the setup: the multi-state Markov process governing the state of the insured, the multivariate payment process describing the collection of life insurance contracts held by the insured and the multivariate present value of their payments. In Section 3, we derive product integral representations of the conditional moment generating function of the present value, which is shown to satisfy a system of partial differential equations. In Section 4 we use this to derive

## 2 SETUP

ordinary differential equations and product integral representations for the conditional moments. Finally, in Section 5 we illustrate the methods of the paper in numerical examples of the motivating examples Example 1.1 and 1.2.

## 2 Setup

In the following, we describe the mathematical framework and main purpose of the paper. Subsection 2.1 introduces the process governing the state of the insured and the multivariate payment process describing the collection of products held by the insured. Subsection 2.2 then considers the multivariate present value of future payments, whose components consists of the present value of the individual payments, and we introduce the calculation of their higher order moments as the main purpose of the paper. By considering existing results in [15, 4] for single payment processes, we end the section by explaining our main ideas and give an overview on how we are to obtain the multivariate extension in the remainder of the paper.

**Notation and conventions** For vectors  $\mathbf{x}, \mathbf{y} \in \mathbb{R}^n$ , we say that  $\mathbf{x} \leq \mathbf{y}$  if  $x_\ell \leq y_\ell$  for all  $\ell = 1, \dots, n$ . Furthermore, we denote with  $\bar{\mathbf{x}}$  the sum of the elements in  $\mathbf{x}$ , i.e.  $\bar{\mathbf{x}} = \sum_{\ell=1}^n x_\ell$ . Also,  $\mathbf{e}_\ell$  is the  $\ell$ 'th unit vector in  $\mathbb{R}^n$  and  $E_n = \{\mathbf{e}_1, \dots, \mathbf{e}_n\}$  denotes the set of unit vectors, i.e. the natural basis of  $\mathbb{R}^n$ . Furthermore,  $\langle \mathbf{x}, \mathbf{y} \rangle = \sum_{\ell=1}^n x_\ell y_\ell$  denotes the usual inner product.

For  $n \times n$  matrix valued functions  $\mathbb{R} \ni x \mapsto \mathbf{A}(x)$  we denote its product integral in the (time) interval  $[s, t]$  as

$$\mathbf{F}(s, t) = \prod_s^t (\mathbf{I} + \mathbf{A}(x) dx),$$

whenever it exists; here  $\mathbf{I}$  is the  $n \times n$  identity matrix. For a survey on product integration and its properties, we refer to [7, 11] and the more recent application to life insurance in [4].

For the purpose of computing multivariate higher order moments, we consider the map  $S : \mathbb{N}_0^n \rightarrow \mathcal{P}(\mathbb{N}_0^n)$  defined as

$$S(\mathbf{x}) = \{\boldsymbol{\xi} \in \mathbb{N}_0^n \setminus \{\mathbf{0}\} \mid \boldsymbol{\xi} \leq \mathbf{x}\}, \quad \mathbf{x} \in \mathbb{N}_0^n. \quad (2.1)$$

The set  $S(\mathbf{x})$  then contains all combinations of lower order moments of  $\mathbf{x}$ , excluding the  $\mathbf{0}$ 'th moment, and the cardinality  $|S(\mathbf{x})| = \prod_{\ell=1}^n (x_\ell + 1) - 1$  then represents the amount of these that appear; whenever we wish to include the  $\mathbf{0}$ 'th moment, we write  $\tilde{S}(\mathbf{k}) = S(\mathbf{k}) \cup \{\mathbf{0}\}$ . For notational convenience, we simply define the mapping  $|\cdot| : \mathbb{N}_0^n \rightarrow \mathbb{N}_0$  as

$$|\mathbf{x}| := \prod_{\ell=1}^n (x_\ell + 1) - 1, \quad \mathbf{x} \in \mathbb{N}_0^n.$$

## 2 SETUP

To compute higher order partial derivatives of multivariate scalar functions  $f : \mathbb{R}^n \rightarrow \mathbb{R}$ , we use the following notation, for  $\mathbf{m} \in \mathbb{N}_0^n$ ,

$$\frac{\partial^{\mathbf{m}}}{\partial \mathbf{x}^{\mathbf{m}}} f(\mathbf{x}) = \frac{\partial^{\bar{\mathbf{m}}}}{\partial x_1^{m_1} \partial x_2^{m_2} \dots \partial x_n^{m_n}} f(\mathbf{x})$$

provided they exist.

### 2.1 Life insurance model

The state of the insured is governed by a non-explosive Markov jump process  $Z = \{Z(t)\}_{t \geq 0}$  taking values on a finite state space  $\mathcal{J} = \{0, \dots, J-1\}$ ,  $J \in \mathbb{N}$ , with deterministic initial state  $Z(0) = z(0) \in \mathcal{J}$ . Denote with  $\mathbf{M}(t) = \{\mu_{ij}(t)\}_{i,j \in \mathcal{J}}$  the transition intensity matrix of  $Z$  and let  $\mathbf{P}(s, t) = \{p_{ij}(s, t)\}_{i,j \in \mathcal{J}}$  be the corresponding transition probability matrix. The multivariate counting process associated with  $Z$  is denoted  $\mathbf{N}(t) = \{N_{ij}(t)\}_{i,j \in \mathcal{J}, i \neq j}$ , with components given by

$$N_{ij}(t) = \# \{s \in (0, t] : Z(s-) = i, Z(s) = j\}.$$

We consider a collection of life insurance contracts described by the  $n$ -dimensional payment process  $\mathbf{B}(t) = (B_1(t), \dots, B_n(t))'$  with each component giving accumulated benefits less premiums for the corresponding contract; here  $n \in \mathbb{N}$  is a fixed and finite number of contracts. We suppose that the  $\ell$ 'th payment process,  $\ell = 1, \dots, n$ , consist of suitably regular deterministic sojourn payment rates  $b_i^\ell$  and transition payments  $b_{ij}^\ell$ ,  $i, j \in \mathcal{J}, i \neq j$ , in terms of the state process  $Z$ . Coupling these in vectors,

$$\mathbf{b}_i(t) = (b_i^1(t), \dots, b_i^n(t))', \quad (2.2)$$

$$\mathbf{b}_{ij}(t) = (b_{ij}^1(t), \dots, b_{ij}^n(t))', \quad (2.3)$$

we formally characterize  $\mathbf{B}$  as

$$d\mathbf{B}(t) = \sum_{i \in \mathcal{J}} \mathbb{1}_{(Z(t-)=i)} \left( \mathbf{b}_i(t) dt + \sum_{j:j \neq i} \mathbf{b}_{ij}(t) dN_{ij}(t) \right), \quad \mathbf{B}(0) = \mathbf{0}. \quad (2.4)$$

We think of  $\mathbf{B}$  as describing payments to an insured that have different product types from which the payments naturally are dependent. The simplest example is an insured having a life annuity and death benefit, which have opposite effect with respect to the lifetime of the insured. By considering these mechanisms in a general multi-state framework, we are able to examine product interactions that are less trivial, cf. the motivating examples Example 1.1 and 1.2 considered in the introduction. Similar type of product combination as that of Example 1.1 is considered in [12], however with interacting effects related to systematic biometric risk only. We consider interacting effects related to unsystematic biometric risk in this paper.

## 2 SETUP

**Example 2.1** (Disability model). The setup of Example 1.1 in the disability model corresponds to the having the vector of payment functions (2.2)–(2.3) as

$$\begin{aligned}\mathbf{b}_0(t) &= (0, b \cdot \mathbf{1}_{(t \geq T)}, 0)', \\ \mathbf{b}_1(t) &= (0, b \cdot \mathbf{1}_{(t \geq T)}, d \cdot \mathbf{1}_{(t < T)})', \\ \mathbf{b}_{02}(t) = \mathbf{b}_{12}(t) &= (S \cdot \mathbf{1}_{(t < T)}, 0, 0)',\end{aligned}$$

with the remaining vector of payment functions having purely zero entries.  $\circ$

**Example 2.2** (Spouse model). The setup of Example 1.2 in the spouse model corresponds to the having the vector of payment functions (2.2)–(2.3) as

$$\begin{aligned}\mathbf{b}_1(t) &= (b_1 \cdot \mathbf{1}_{(t \leq T)}, 0)', \\ \mathbf{b}_2(t) &= (0, b_2 \cdot \mathbf{1}_{(t \leq T)})',\end{aligned}$$

with the remaining vector of payment functions having purely zero entries.  $\circ$

### 2.2 Present values and their moments

Let  $r = \{r(t)\}_{t \geq 0}$  be a continuous, suitably regular and deterministic short rate. To account for time value of money, introduce  $v(s, t)$  as the discount factor for a payment at time  $t$  valued at time  $s \leq t$ , i.e.

$$v(s, t) = e^{-\int_s^t r(x) dx}.$$

The present value at time  $s$  of future benefits less premiums up to time  $t$  for the payment process  $\mathbf{B}$  is then given by

$$\mathbf{U}(s, t) = \int_s^t v(s, u) d\mathbf{B}(u) = (U_1(s, t), \dots, U_n(s, t))', \quad (2.5)$$

where the components  $U_\ell(s, t) = \int_s^t v(s, u) dB_\ell(u)$ ,  $\ell = 1, \dots, n$ , are the present values of the individual payment process.

The components of  $\mathbf{U}$  are (by construction) seen to be dependent, which introduces joint effects on the payments with respect to states of the insured. To analyse these dependence structures, we wish to examine its joint distribution by computing its conditional higher order moments given the current state of the insured, that is, by computing

$$V_i^{(k)}(s, t) = \mathbb{E} \left[ \prod_{\ell=1}^n U_\ell(s, t)^{k_\ell} \middle| Z(s) = i \right] \quad (2.6)$$

## 2 SETUP

for any  $i \in \mathcal{J}$  and  $\mathbf{k} = (k_1, \dots, k_n) \in \mathbb{N}_0^n$ . For  $\mathbf{k} = \mathbf{e}_\ell$  we have that  $V_i^{(\mathbf{e}_\ell)}(s, t)$  is the state-wise prospective reserve for the  $\ell$ 'th payment process, calculated using the classic Thiele's differential equation, cf. [9, 13].

As mentioned in the introduction, having multiple product types for the same insured is usually handled by adding all payment functions together as  $\bar{\mathbf{b}}_i$  and  $\bar{\mathbf{b}}_{ij}$ , or, equivalently, by considering the aggregate present value  $\bar{U}(s, t)$ . This effectively gives a one dimensional payment process to examine; in [9, 13] we have Thiele's differential equation for conditional expectations and in [1, 4, 15] we have differential equations for conditional higher order moments, which assess the unsystematic biometric risk in the contract. The present paper is therefore to be thought of as a way of examining how the unsystematic biometric risk of the individual product types affects the total unsystematic biometric risk of the contract.

The following matrices, which we think of as inputs to the computations, are needed in the derivations; for  $\ell \in \{1, \dots, n\}$ ,  $i, j \in \mathcal{J}$ ,  $i \neq j$ , and  $\mathbf{y} \in \mathbb{N}_0^n \setminus (E_n \cup \{\mathbf{0}\})$ , let

$$\mathbf{B}_\ell(s) = \left\{ b_{ij}^\ell(s) \right\}_{i,j \in \mathcal{J}}, \quad (2.7)$$

$$\mathbf{b}^\ell(s) = \left( b_0^\ell(s), \dots, b_{J+1}^\ell(s) \right)', \quad (2.8)$$

$$\mathbf{R}_\ell(s) = \mathbf{M}(s) \bullet \mathbf{B}_\ell(s) + \Delta(\mathbf{b}^\ell(s)), \quad (2.9)$$

$$\mathbf{C}^{(\mathbf{y})}(s) = \mathbf{M}(s) \bullet \mathbf{B}_1^{\bullet y_1}(s) \bullet \dots \bullet \mathbf{B}_n^{\bullet y_n}(s), \quad (2.10)$$

where  $\Delta(\mathbf{b}^\ell(s))$  denotes the diagonal matrix with the vector  $\mathbf{b}^\ell(s)$  as diagonal elements,  $\bullet$  denotes the entry-wise matrix product and e.g.  $\mathbf{B}_1^{\bullet y}(s)$  denotes the entry-wise matrix product of  $\mathbf{B}_1(s)$  with itself  $y$  times; here we use the convention that e.g.  $\mathbf{B}_1^{\bullet 0}(s)$  denotes the matrix with all entries equal to 1. These matrices are defined analogously to those of [4, (3.8)–(3.11)] with the relevant extension to a multivariate payment process.

The remainder of the paper now focuses on computation of  $V_i^{(\mathbf{k})}(s, t)$ , carried out as follows. Following [4], we first compute the conditional higher order moments on events that the insured terminates in specific states, given by, for  $j \in \mathcal{J}$ ,

$$V_{ij}^{(\mathbf{k})}(s, t) = \mathbb{E} \left[ \mathbf{1}_{(Z(t)=j)} \prod_{\ell=1}^n U_\ell(s, t)^{k_\ell} \mid Z(s) = i \right] \quad (2.11)$$

with corresponding matrix

$$\mathbf{V}^{(\mathbf{k})}(s, t) = \left\{ V_{ij}^{(\mathbf{k})}(s, t) \right\}_{i,j \in \mathcal{J}}. \quad (2.12)$$

This is carried out by deriving the relevant moment generating function. As in [4], we shall refer to these as the *conditional partial moments* of  $\mathbf{U}(s, t)$ . From these, the



### 3 MOMENT GENERATING FUNCTIONS

conditional moments  $V_i^{(\mathbf{k})}(s, t)$  are obtained via the relation  $\sum_{j \in \mathcal{J}} V_{ij}^{(\mathbf{k})}(s, t) = V_i^{(\mathbf{k})}(s, t)$ , such that

$$V_i^{(\mathbf{k})}(s, t) = \left\{ \mathbf{V}^{(\mathbf{k})}(s, t) \mathbf{1} \right\}_i, \quad (2.13)$$

where  $\mathbf{1} = (1, 1, \dots, 1)'$ .

While not strictly required, computing the conditional moments via the conditional partial moments as in (2.13) may have its advantages since these quantities have their own interpretations. In [13], similar type of quantities are computed for retrospective reserves.

### 3 Moment generating functions

Consider the multivariate present value  $\mathbf{U}(s, t)$ . The moment generating function of its conditional distribution given  $Z(s) = i$  on the event  $(Z(t) = j)$ ,  $i, j \in \mathcal{J}$ , is given by

$$F_{ij}(\boldsymbol{\theta}; s, t) = \mathbb{E} \left[ e^{\langle \boldsymbol{\theta}, \mathbf{U}(s, t) \rangle} \mathbf{1}_{(Z(t)=j)} \mid Z(s) = i \right], \quad \boldsymbol{\theta} \in \mathbb{R}^n, \quad (3.1)$$

with corresponding matrix

$$\mathbf{F}(\boldsymbol{\theta}; s, t) = \{F_{ij}(\boldsymbol{\theta}; s, t)\}_{i, j \in \mathcal{J}}.$$

We are then able to obtain the moment generating function by a direct application of results in [4].

**Theorem 3.1.** *The joint distribution of the multivariate present value  $\mathbf{U}(s, t)$  has moment generating function given by*

$$\mathbf{F}(\boldsymbol{\theta}; s, t) = \prod_s^t \left( \mathbf{I} + \left[ \mathbf{M}(u) \bullet \left\{ e^{v(s, u) \langle \boldsymbol{\theta}, \mathbf{b}_{ij}(u) \rangle} \right\}_{i, j \in \mathcal{J}} + v(s, u) \sum_{\ell=1}^n \theta_\ell \Delta(\mathbf{b}^\ell(u)) \right] du \right)$$

*Proof.* Let  $\boldsymbol{\theta} \in \mathbb{R}^n$  and  $0 \leq s \leq t$  be given. By linearity of integrals, we note

$$\begin{aligned} \langle \boldsymbol{\theta}, \mathbf{U}(s, t) \rangle &= \int_s^t v(s, u) d\tilde{\mathbf{B}}_{\boldsymbol{\theta}}(u), \\ d\tilde{\mathbf{B}}_{\boldsymbol{\theta}}(u) &= \sum_{i \in \mathcal{J}} \mathbf{1}_{(Z(u^-)=i)} \left( \langle \boldsymbol{\theta}, \mathbf{b}_i(u) \rangle du + \sum_{j: j \neq i} \langle \boldsymbol{\theta}, \mathbf{b}_{ij}(u) \rangle dN_{ij}(u) \right). \end{aligned}$$

Hence,  $\langle \boldsymbol{\theta}, \mathbf{U}(s, t) \rangle$  is the present value of a one-dimensional payment process with sojourn payment rates  $\langle \boldsymbol{\theta}, \mathbf{b}_j(\cdot) \rangle$  and transition payments  $\langle \boldsymbol{\theta}, \mathbf{b}_{jk}(\cdot) \rangle$ . Thus, in the spirit of [4], we may view  $\mathbf{F}(\boldsymbol{\theta}; s, t)$  as the moment generating function of a total (undiscounted) reward evaluated in 1 with the payment functions  $u \mapsto \langle \boldsymbol{\theta}, v(s, u) \mathbf{b}_j(u) \rangle$  and  $u \mapsto \langle \boldsymbol{\theta}, v(s, u) \mathbf{b}_{jk}(u) \rangle$  (for fixed  $s$ ). The result then follows from an application of [4, Theorem 3].  $\square$

### 3 MOMENT GENERATING FUNCTIONS

*Remark 3.2.* Theorem 3.1 is a generalization of [4, Theorem 3] to multivariate payment processes with discounting. Indeed, by letting  $n = 1$  and  $r(s) = 0$  for all  $s \geq 0$ , the result simplifies to, suppressing the superfluous dependency in  $\ell$ ,

$$\mathbf{F}(\boldsymbol{\theta}; s, t) = \prod_s^t \left( \mathbf{I} + \left[ \mathbf{M}(s) \bullet \left\{ e^{\boldsymbol{\theta} b_{ij}(s)} \right\}_{i,j \in \mathcal{J}} + \boldsymbol{\theta} \boldsymbol{\Delta}(\mathbf{b}(s)) \right] du \right),$$

which is the product integral of [4, Theorem 3].  $\triangle$

From the product integral representation of  $\mathbf{F}$ , we are able to obtain a partial differential equation satisfied by  $\mathbf{F}$ . Similar type of equations are also obtained in [1].

**Theorem 3.3.** *The moment generating function  $\mathbf{F}$  satisfies the system of partial differential equations*

$$\begin{aligned} \frac{\partial}{\partial s} \mathbf{F}(\boldsymbol{\theta}; s, t) &= r(s) \sum_{\ell=1}^n \theta_{\ell} \frac{\partial}{\partial \theta_{\ell}} \mathbf{F}(\boldsymbol{\theta}; s, t) \\ &\quad - \left[ \mathbf{M}(s) \bullet \left\{ e^{\langle \boldsymbol{\theta}, \mathbf{b}_{ij}(s) \rangle} \right\}_{i,j \in \mathcal{J}} + \sum_{\ell=1}^n \theta_{\ell} \boldsymbol{\Delta}(\mathbf{b}^{\ell}(s)) \right] \mathbf{F}(\boldsymbol{\theta}; s, t), \end{aligned} \quad (3.2)$$

with the terminal condition  $\mathbf{F}(\boldsymbol{\theta}; t, t) = \mathbf{I}$ .

*Proof.* For  $\boldsymbol{\theta} \in \mathbb{R}^n$  and  $0 \leq s \leq u$ , let

$$\mathbf{A}(s, u; \boldsymbol{\theta}) = \mathbf{M}(u) \bullet \left\{ e^{v(s,u) \langle \boldsymbol{\theta}, \mathbf{b}_{ij}(u) \rangle} \right\}_{i,j \in \mathcal{J}} + v(s, u) \sum_{\ell=1}^n \theta_{\ell} \boldsymbol{\Delta}(\mathbf{b}^{\ell}(u))$$

denote the matrix that is product integrated in Theorem 3.1. It satisfies the relation  $\frac{\partial}{\partial s} \mathbf{A}(s, u; \boldsymbol{\theta}) = r(s) \sum_{\ell=1}^n \theta_{\ell} \frac{\partial}{\partial \theta_{\ell}} \mathbf{A}(s, u; \boldsymbol{\theta})$ , which can be seen from direct calculations. Now, since  $\mathbf{F}$  is a product integral of the matrix function  $\mathbf{A}(s, \cdot; \boldsymbol{\theta})$ , it has the Peano-Baker series representation given by

$$\mathbf{F}(\boldsymbol{\theta}; s, t) = \mathbf{I} + \sum_{n=1}^{\infty} \int_s^t \int_s^{x_n} \cdots \int_s^{x_2} \mathbf{A}(s, x_1; \boldsymbol{\theta}) \cdots \mathbf{A}(s, x_n; \boldsymbol{\theta}) dx_1 \cdots dx_n.$$

Differentiating this w.r.t.  $s$ , we get using Leibniz' integral rule,

$$\begin{aligned} \frac{\partial}{\partial s} \mathbf{F}(\boldsymbol{\theta}; s, t) &= -\mathbf{A}(s, s; \boldsymbol{\theta}) \mathbf{F}(\boldsymbol{\theta}; s, t) \\ &\quad + \sum_{n=1}^{\infty} \int_s^t \int_s^{x_n} \cdots \int_s^{x_2} \frac{\partial}{\partial s} (\mathbf{A}(s, x_1; \boldsymbol{\theta}) \cdots \mathbf{A}(s, x_n; \boldsymbol{\theta})) dx_1 \cdots dx_n \end{aligned}$$

## 4 HIGHER ORDER MOMENTS

In the last differentiation, we can in each combination (when applying the product rule) substitute the derivative w.r.t. to  $s$  with  $r(s) \sum_{\ell=1}^n \theta_\ell \frac{\partial}{\partial \theta_\ell} \mathbf{A}(s, u; \boldsymbol{\theta})$ . This gives

$$\begin{aligned} \frac{\partial}{\partial s} \mathbf{F}(\boldsymbol{\theta}; s, t) &= -\mathbf{A}(s, s; \boldsymbol{\theta}) \mathbf{F}(\boldsymbol{\theta}; s, t) \\ &+ \sum_{n=1}^{\infty} \int_s^t \int_s^{x_n} \cdots \int_s^{x_2} r(s) \sum_{\ell=1}^n \theta_\ell \frac{\partial}{\partial \theta_\ell} (\mathbf{A}(s, x_1; \boldsymbol{\theta}) \cdots \mathbf{A}(s, x_n; \boldsymbol{\theta})) dx_1 \cdots dx_n \\ &= -\mathbf{A}(s, s; \boldsymbol{\theta}) \mathbf{F}(\boldsymbol{\theta}; s, t) + r(s) \sum_{\ell=1}^n \theta_\ell \frac{\partial}{\partial \theta_\ell} \mathbf{F}(\boldsymbol{\theta}; s, t), \end{aligned}$$

as claimed. The boundary condition follows from the definition of product integrals. Throughout the proof, we have used that the Peano-Baker series converges uniformly on compact intervals in order to interchange summation and differentiation.  $\square$

*Remark 3.4.* Consider the univariate case  $n = 1$  and suppress the superfluous dependency in  $\ell$ . By multiplying both sides of (3.2) with  $\mathbf{1} = (1, 1, \dots, 1)'$  and extracting the  $i$ 'th element,  $i \in \mathcal{J}$ , we see that  $F_i(\boldsymbol{\theta}; s, t) := \mathbb{E}[e^{\theta U(s,t)} | Z(s) = i]$  satisfies the partial differential equation

$$\begin{aligned} \frac{\partial}{\partial s} F_i(\boldsymbol{\theta}; s, t) &= r(s) \theta \frac{\partial}{\partial \theta} F_i(\boldsymbol{\theta}; s, t) - \theta b_i(s) F_i(\boldsymbol{\theta}; s, t) \\ &- \sum_{k:k \neq i} \mu_{ik}(s) \left( e^{\theta b_{ik}(s)} F_k(\boldsymbol{\theta}; s, t) - F_i(\boldsymbol{\theta}; s, t) \right), \end{aligned}$$

with the boundary condition  $F_i(\boldsymbol{\theta}; t, t) = 1$ . This is the partial differential equation of [1, Proposition 4.3] in the Markovian special case with no duration dependency.  $\triangle$

Having derived the relevant moment generating function of the multivariate present value  $\mathbf{U}(s, t)$ , we are now ready to compute the corresponding higher order moments.

## 4 Higher order moments

In this section, we derive the conditional (partial) higher order moments (2.6) and (2.12), which constitutes the main contribution of the paper. In Subsection 4.1 we derive differential equations for the conditional partial moments (2.12) using the moment generating function  $\mathbf{F}(\cdot; s, t)$  derived in Section 3. Then we use this result to derive differential equations for the conditional moments (2.6) via the relation (2.13). In Subsection 4.2, we then show how all conditional partial moments up to a given order  $\mathbf{k} \in \mathbb{N}_0^n$  can be obtained from a product integral calculation. Throughout the section, we highlight how the results appear as natural and simple multivariate extensions to results in [4, 15], in particular when notation may appear as cumbersome.

### 4.1 Differential equations of conditional (partial) moments

We now turn our attention to deriving ordinary differential equations for the conditional (partial) moments. For this, recall that  $S(\mathbf{k})$  denotes the set of all lower order moments of the  $\mathbf{k}$ 'th moment, excluding the  $\mathbf{0}$ 'th moment, with elements we may write as

$$S(\mathbf{k}) = \left\{ \mathbf{y}^1, \dots, \mathbf{y}^{|\mathbf{k}|} \right\}.$$

The following main result then take use of the moment generating function derived in Section 3 to obtain differential equations for the conditional partial moments.

**Theorem 4.1.** *The conditional partial moments  $\mathbf{V}^{(\mathbf{k})}(\cdot, t)$  satisfies the backward differential equations given by, for  $\mathbf{k} \in \mathbb{N}_0^n$ ,*

$$\begin{aligned} \frac{\partial}{\partial s} \mathbf{V}^{(\mathbf{k})}(s, t) &= (\bar{\mathbf{k}} r(s) \mathbf{I} - \mathbf{M}(s)) \mathbf{V}^{(\mathbf{k})}(s, t) - \sum_{\ell=1}^n k_\ell \mathbf{R}_\ell(s) \mathbf{V}^{(\mathbf{k}-\mathbf{e}_\ell)}(s, t) \\ &\quad - \sum_{\substack{\mathbf{y} \in S(\mathbf{k}) \\ \mathbf{y} \notin E_n}} \prod_{\ell=1}^n \binom{k_\ell}{y_\ell} \mathbf{C}^{(\mathbf{y})}(s) \mathbf{V}^{(\mathbf{k}-\mathbf{y})}(s, t), \end{aligned} \quad (4.1)$$

with the terminal conditions  $\mathbf{V}^{(\mathbf{k})}(t, t) = \mathbb{1}_{(\mathbf{k}=\mathbf{0})} \mathbf{I}$ .

*Proof.* We can use the moment generating function  $\mathbf{F}(\cdot; s, t)$  to calculate  $\frac{\partial}{\partial s} \mathbf{V}^{(\mathbf{k})}(s, t)$  as

$$\frac{\partial}{\partial s} \mathbf{V}^{(\mathbf{k})}(s, t) = \left. \frac{\partial^{\mathbf{k}}}{\partial \boldsymbol{\theta}^{\mathbf{k}}} \frac{\partial}{\partial s} \mathbf{F}(\boldsymbol{\theta}; s, t) \right|_{\boldsymbol{\theta}=\mathbf{0}}.$$

Differentiating both sides of (3.2), we get by the generalized product rule

$$\begin{aligned} &\frac{\partial^{\mathbf{k}}}{\partial \boldsymbol{\theta}^{\mathbf{k}}} \frac{\partial}{\partial s} \mathbf{F}(\boldsymbol{\theta}; s, t) \\ &= r(s) \sum_{\ell=1}^n \sum_{\mathbf{y} \in \tilde{S}(\mathbf{k})} \prod_{r=1}^n \binom{k_r}{y_r} \left[ \frac{\partial \mathbf{y}}{\partial \boldsymbol{\theta}^{\mathbf{y}}} \theta_\ell \right] \frac{\partial^{\mathbf{k}-\mathbf{y}}}{\partial \boldsymbol{\theta}^{\mathbf{k}-\mathbf{y}}} \frac{\partial}{\partial \theta_\ell} \mathbf{F}(\boldsymbol{\theta}; s, t) \\ &\quad - \sum_{\mathbf{y} \in \tilde{S}(\mathbf{k})} \prod_{r=1}^n \binom{k_r}{y_r} \frac{\partial \mathbf{y}}{\partial \boldsymbol{\theta}^{\mathbf{y}}} \left[ \mathbf{M}(s) \bullet \left\{ e^{\langle \boldsymbol{\theta}, \mathbf{b}_{ij}(s) \rangle} \right\}_{i,j \in \mathcal{J}} + \sum_{\ell=1}^n \theta_\ell \boldsymbol{\Delta}(\mathbf{b}^\ell(s)) \right] \frac{\partial^{\mathbf{k}-\mathbf{y}}}{\partial \boldsymbol{\theta}^{\mathbf{k}-\mathbf{y}}} \mathbf{F}(\boldsymbol{\theta}; s, t). \\ &= r(s) \sum_{\ell=1}^n [\mathbb{1}_{(\mathbf{y}=\mathbf{0})} \theta_\ell + k_\ell \mathbb{1}_{(\mathbf{y}=\mathbf{e}_\ell)}] \frac{\partial^{\mathbf{k}-\mathbf{y}}}{\partial \boldsymbol{\theta}^{\mathbf{k}-\mathbf{y}}} \frac{\partial}{\partial \theta_\ell} \mathbf{F}(\boldsymbol{\theta}; s, t) \\ &\quad - \sum_{\mathbf{y} \in \tilde{S}(\mathbf{k})} \prod_{r=1}^n \binom{k_r}{y_r} \left[ \mathbf{M}(s) \bullet \left\{ \prod_{\ell=1}^n b_{ij}^\ell(s)^{y_\ell} e^{\langle \boldsymbol{\theta}, \mathbf{b}_{ij}(s) \rangle} \right\}_{i,j \in \mathcal{J}} + \sum_{\ell=1}^n \mathbb{1}_{(\mathbf{y}=\mathbf{e}_\ell)} \boldsymbol{\Delta}(\mathbf{b}^\ell(s)) \right] \times \\ &\quad \quad \quad \frac{\partial^{\mathbf{k}-\mathbf{y}}}{\partial \boldsymbol{\theta}^{\mathbf{k}-\mathbf{y}}} \mathbf{F}(\boldsymbol{\theta}; s, t), \end{aligned}$$

#### 4 HIGHER ORDER MOMENTS

In the last equality, we have used that  $\prod_{r=1}^n \binom{k_r}{y_r} = k_\ell$  if  $\mathbf{y} = \mathbf{e}_\ell$  along with  $\frac{\partial \mathbf{y}}{\partial \boldsymbol{\theta}} \boldsymbol{\theta}_\ell = \mathbb{1}_{(\mathbf{y}=\mathbf{0})} \boldsymbol{\theta}_\ell + \mathbb{1}_{(\mathbf{y}=\mathbf{e}_\ell)}$  for  $\ell = 1, \dots, n$ . Now, evaluating in  $\boldsymbol{\theta} = \mathbf{0}$ , we get

$$\begin{aligned} \frac{\partial}{\partial s} \mathbf{V}^{(\mathbf{k})}(s, t) &= r(s) \sum_{\ell=1}^n k_\ell \frac{\partial^{\mathbf{k}-\mathbf{e}_\ell}}{\partial \boldsymbol{\theta}^{\mathbf{k}-\mathbf{e}_\ell}} \frac{\partial}{\partial \theta_\ell} \mathbf{F}(\boldsymbol{\theta}; s, t) \Big|_{\boldsymbol{\theta}=\mathbf{0}} \\ &\quad - \sum_{\mathbf{y} \in \tilde{S}(\mathbf{k})} \prod_{\ell=1}^n \binom{k_\ell}{y_\ell} \left[ \mathbf{M}(s) \bullet \left\{ \prod_{\ell=1}^n b_{ij}^\ell(s)^{y_\ell} \right\}_{i,j \in \mathcal{J}} + \sum_{\ell=1}^n \mathbb{1}_{(\mathbf{y}=\mathbf{e}_\ell)} \boldsymbol{\Delta}(\mathbf{b}^\ell(s)) \right] \times \\ &\quad \mathbf{V}^{(\mathbf{k}-\mathbf{y})}(s, t) \\ &= \bar{\mathbf{k}} r(s) \mathbf{V}^{(\mathbf{k})}(s, t) \\ &\quad - \sum_{\mathbf{y} \in \tilde{S}(\mathbf{k})} \prod_{\ell=1}^n \binom{k_\ell}{y_\ell} \left[ \mathbf{M}(s) \bullet B_1^{\bullet y_1}(s) \bullet \dots \bullet B_n^{\bullet y_n}(s) + \sum_{\ell=1}^n \mathbb{1}_{(\mathbf{y}=\mathbf{e}_\ell)} \boldsymbol{\Delta}(\mathbf{b}^\ell(s)) \right] \times \\ &\quad \mathbf{V}^{(\mathbf{k}-\mathbf{y})}(s, t). \end{aligned}$$

Splitting the last sum into terms corresponding to  $\mathbf{y} = \mathbf{0}$  and  $\mathbf{y} = \mathbf{e}_\ell$ , for  $\ell = 1, \dots, n$ , gives

$$\begin{aligned} \frac{\partial}{\partial s} \mathbf{V}^{(\mathbf{k})}(s, t) &= (\bar{\mathbf{k}} r(s) \mathbf{I} - \mathbf{M}(s)) \mathbf{V}^{(\mathbf{k})}(s, t) - \sum_{\ell=1}^n k_\ell \mathbf{R}_\ell(s) \mathbf{V}^{(\mathbf{k}-\mathbf{e}_\ell)}(s, t) \\ &\quad - \sum_{\substack{\mathbf{y} \in S(\mathbf{k}) \\ \mathbf{y} \notin E_n}} \prod_{\ell=1}^n \binom{k_\ell}{y_\ell} \mathbf{C}^{(\mathbf{y})}(s) \mathbf{V}^{(\mathbf{k}-\mathbf{y})}(s, t), \end{aligned}$$

the desired differential equation. The boundary condition follows from (2.11) with the fact that  $\mathbf{V}^{(\mathbf{0})}(t, t) = \mathbf{P}(t, t) = \mathbf{I}$ .  $\square$

*Remark 4.2.* The differential equation of Theorem 4.1 generalizes [15, Theorem 6] to multivariate present values; for  $n = 1$  we have, supressing the superfluous dependency in  $\ell$ , that  $S(k) = \{1, 2, \dots, k\}$  and  $E_1 = \{1\}$ , and so the differential equations reads

$$\begin{aligned} \frac{\partial}{\partial s} \mathbf{V}^{(k)}(s, t) &= (k r(s) \mathbf{I} - \mathbf{M}(s)) \mathbf{V}^{(k)}(s, t) - k \mathbf{R}(s) \mathbf{V}^{(k-1)}(s, t) \\ &\quad - \sum_{y=2}^k \binom{k}{y} \mathbf{C}^{(y)}(s) \mathbf{V}^{(k-y)}(s, t), \end{aligned}$$

with the terminal conditions  $\mathbf{V}^{(k)}(t, t) = \mathbb{1}_{(k=0)} \mathbf{I}$ .  $\triangle$

From the differential equation of the conditional partial moments, we immediately obtain differential equations for the corresponding conditional moments  $V_i^{(k)}(s, t)$  by multiplying the former with  $\mathbf{1} = (1, 1, \dots, 1)'$ , as explained in Section 2.

#### 4 HIGHER ORDER MOMENTS

**Theorem 4.3.** *The conditional moments  $V_i^{(\mathbf{k})}(\cdot, t)$  satisfies the backward differential equations given by, for  $i \in \mathcal{J}$ ,*

$$\begin{aligned} \frac{\partial}{\partial s} V_i^{(\mathbf{k})}(s, t) &= (\bar{\mathbf{k}}r(s) + \mu_i(s))V_i^{(\mathbf{k})}(s, t) - \sum_{\ell=1}^n k_\ell b_i^\ell(s) V_i^{(\mathbf{k}-\mathbf{e}_\ell)}(s, t) \\ &\quad - \sum_{j:j \neq i} \mu_{ij}(s) \sum_{\mathbf{y} \in \tilde{S}(\mathbf{k})} \prod_{\ell=1}^n \binom{k_\ell}{y_\ell} (b_{ij}^\ell(s))^{y_\ell} V_j^{(\mathbf{k}-\mathbf{y})}(s, t), \end{aligned} \quad (4.2)$$

with the terminal conditions  $V_i^{(\mathbf{k})}(t, t) = \mathbf{1}_{(\mathbf{k}=\mathbf{0})}$ .

*Proof.* Simply multiply the differential equation in Theorem 4.1 with  $\mathbf{1} = (1, 1, \dots, 1)'$  on both sides and extract the  $i$ 'th element of the vectors.  $\square$

*Remark 4.4.* The differential equation of Theorem 4.3 generalizes [15, (3.2)] to multivariate present values; for  $n = 1$  we have, supressing the superfluous dependency in  $\ell$ , that  $S(k) = \{1, 2, \dots, k\}$  and  $E_1 = \{1\}$ , and so the differential equations reads

$$\begin{aligned} \frac{\partial}{\partial s} V_i^{(k)}(s, t) &= (kr(s) + \mu_i(s))V_i^{(k)}(s, t) - kb_i(s)V_i^{(k-1)}(s, t) \\ &\quad - \sum_{j:j \neq i} \mu_{ij}(s) \sum_{y=0}^k \binom{k}{y} (b_{ij}(s))^y V_j^{(k-y)}(s, t), \end{aligned}$$

with the terminal conditions  $V_i^{(k)}(t, t) = \mathbf{1}_{(k=0)}$ .  $\triangle$

*Remark 4.5.* The conditional central moments

$$m_i^{(\mathbf{k})}(s, t) = \mathbb{E} \left[ \prod_{\ell=1}^n \left( U_\ell(s, t) - V_i^{(\mathbf{e}_\ell)}(s, t) \right)^{k_\ell} \middle| Z(s) = i \right]$$

may be obtained from the conditional (non-central) moments  $V_i^{(\mathbf{k})}(s, t)$  via an application of the multidimensional binomial formula, giving

$$m_i^{(\mathbf{k})}(s, t) = \sum_{\mathbf{y} \in \tilde{S}(\mathbf{k})} \prod_{\ell=1}^n (-1)^{k_\ell - y_\ell} \binom{k_\ell}{y_\ell} V_i^{(\mathbf{y})}(s, t) V_i^{(\mathbf{e}_\ell)}(s, t)^{k_\ell - y_\ell}. \quad (4.3)$$

Thus, by first solving the differential equation of Theorem 4.3 we are able to compute  $m_i^{(\mathbf{k})}(s, t)$  via (4.3). Here, we may note that solving the differential equation of Theorem 4.3 to compute the  $\mathbf{k}$ 'th moment immediately gives  $V_i^{(\mathbf{y})}(s, t)$  for all  $\mathbf{y} \in \tilde{S}(\mathbf{k})$ ; details regarding this is explained in Subsection 4.2.  $\triangle$

#### 4 HIGHER ORDER MOMENTS

**Example 4.6** (Conditional covariance). For  $n = 2$  and  $\mathbf{k} = (1, 1)$  we have that  $\tilde{S}(\mathbf{k}) = \{(0, 0), (1, 0), (0, 1), (1, 1)\}$ , and so the differential equation for the conditional product moment  $V_i^{(1,1)}(s, t) = \mathbb{E}[U_1(s, t)U_2(s, t) \mid Z(s) = i]$  is given by

$$\begin{aligned} \frac{\partial}{\partial s} V_i^{(1,1)}(s, t) &= (2r(s) + \mu_i(s))V_i^{(1,1)}(s, t) - b_i^1(s)V_i^{(0,1)}(s, t) - b_i^2(s)V_i^{(1,0)}(s, t) \\ &\quad - \sum_{j:j \neq i} \mu_{ij}(s) \left( V_j^{(1,1)}(s, t) + b_{ij}^1(s)V_j^{(0,1)}(s, t) + b_{ij}^2(s)V_j^{(1,0)}(s, t) \right. \\ &\quad \left. + b_{ij}^1(s)b_{ij}^2(s) \right), \\ V_i^{(1,1)}(t, t) &= 0, \end{aligned}$$

where  $V_i^{(0,1)}$  and  $V_i^{(1,0)}$  are the state-wise prospective reserves of the two payment processes, which can be calculated using Thiele's differential equation. From this, the conditional central moment  $m_i^{(1,1)}(s, t)$  is given as

$$\begin{aligned} m_i^{(1,1)}(s, t) &= \mathbb{E} \left[ \left( U_1(s, t) - V_i^{(1,0)}(s, t) \right) \left( U_2(s, t) - V_i^{(0,1)}(s, t) \right) \mid Z(s) = i \right] \\ &= V_i^{(1,1)}(s, t) - V_i^{(1,0)}(s, t)V_i^{(0,1)}(s, t), \end{aligned} \quad (4.4)$$

which is the conditional covariance of  $U_1(s, t)$  og  $U_2(s, t)$  given  $Z(s) = i$ .  $\circ$

Following up on the example, it turns out that it is possible to derive differential equations for the conditional covariance between two present values in which the sum at risk used in the Thiele differential equations appears. The result is also presented without proof in [3, Proposition 10.3], and we are able to proof the result here through our results on the non-central moments.

**Corollary 4.7.** *The conditional covariance  $m_i^{(e_\ell + e_m)}(s, t)$  between  $U_\ell(s, t)$  and  $U_m(s, t)$  given  $Z(s) = i$  satisfies the system of backward differential equations given by, for  $i \in \mathcal{J}$ ,*

$$\begin{aligned} \frac{\partial}{\partial s} m_i^{(e_\ell + e_m)}(s, t) &= 2r(s)m_i^{(e_\ell + e_m)}(s, t) \\ &\quad - \sum_{j:j \neq i} \mu_{ij}(s) \left( R_{ij}^\ell(s)R_{ij}^m(s) + m_j^{(e_\ell + e_m)}(s, t) - m_i^{(e_\ell + e_m)}(s, t) \right), \end{aligned}$$

with terminal conditions  $m_i^{(e_\ell + e_m)}(t, t) = 0$ . Here,  $R_{ij}^\ell$  denotes the sum at risk in the  $\ell$ 'th payment process for the transition from state  $i$  to  $j$ ,  $i \neq j$ , given by

$$R_{ij}^\ell(s) = b_{ij}^\ell(s) + V_j^{(e_\ell)}(s, t) - V_i^{(e_\ell)}(s, t),$$

and analogously for  $R_{ij}^m$ .

#### 4 HIGHER ORDER MOMENTS

*Proof.* Differentiating (4.4) w.r.t.  $s$  on both sides gives

$$\begin{aligned} \frac{\partial}{\partial s} m_i^{(e_\ell + e_m)}(s, t) &= \frac{\partial}{\partial s} V_i^{(e_\ell + e_m)}(s, t) - V_i^{(e_m)}(s, t) \frac{\partial}{\partial s} V_i^{(e_\ell)}(s, t) \\ &\quad - V_i^{(e_\ell)}(s, t) \frac{\partial}{\partial s} V_i^{(e_m)}(s, t) \end{aligned} \quad (4.5)$$

Then insert the expression for  $\frac{\partial}{\partial s} V_i^{(e_\ell + e_m)}(s, t)$  obtained in Example 4.6 as well as the Thiele differential equations for the state-wise prospective reserves, given by

$$\frac{\partial}{\partial s} V_i^{(e_\ell)}(s, t) = r(s) V_i^{(e_\ell)}(s, t) - b_i^\ell(s) - \sum_{j:j \neq i} \mu_{ij}(s) R_{ij}^\ell(s),$$

and analogously for  $V_i^{(e_m)}$ . By gathering the relevant terms, we obtain the desired differential equation. The boundary condition follows directly from (4.4).  $\square$

*Remark 4.8.* Note that the differential equation of Corollary 4.7 corresponds to a Thiele differential equation with interest rate  $2r$ , no sojourn payments and transition payments  $R_{ij}^\ell R_{ij}^m$  from state  $i$  to state  $j$ . Consequently, the conditional covariance  $m_i^{(e_\ell + e_m)}$  has the representation

$$\begin{aligned} m_i^{(e_\ell + e_m)}(s, t) &= \mathbb{E} \left[ \int_s^t e^{-\int_s^u 2r(v) dv} d\tilde{B}(u) \mid Z(s) = i \right] \\ d\tilde{B}(u) &= \sum_{i \in \mathcal{J}} \mathbb{1}_{(Z(u-) = i)} \sum_{j:j \neq i} R_{ij}^\ell(u) R_{ij}^m(u) dN_{ij}(u). \end{aligned}$$

In the case of the conditional variance, i.e.  $\ell = m$ , these type of representations are known as Hattendorff's theorem, which has been formulated in a multi-state Markovian framework by [16] using martingale techniques and further generalized in [14]. The result here may thus reveal that the probabilistic structures leading to Hattendorff's theorem is not only limited to the variance, but can be carried over to the covariance.  $\triangle$

From the conditional covariances (and variances when  $\ell = m$ ), we are then able to put up the conditional covariance matrix of  $\mathbf{U}(s, t)$  given  $Z(s) = i$ ,

$$\Sigma_i(s, t) = \left\{ m_i^{(e_\ell + e_m)} \right\}_{\ell, m=1, \dots, n} \quad (4.6)$$

as well as the corresponding correlation matrix

$$\rho_i(s, t) = \left\{ \frac{m_i^{(e_\ell + e_m)}(s, t)}{\sqrt{m_i^{(e_\ell + e_\ell)}(s, t) m_i^{(e_m + e_m)}(s, t)}} \right\}_{\ell, m=1, \dots, n} \quad (4.7)$$

which makes us able to analyse pair-wise dependence structures between the present values.

In the following subsection, we derive a product integral representation for the conditional partial moments  $\mathbf{V}(\mathbf{y})$ ,  $\mathbf{y} \in \tilde{S}(\mathbf{k})$ , demonstrating how one may calculate all lower order moments  $\mathbf{0}, \mathbf{y}^1, \dots, \mathbf{y}^{|\mathbf{k}|-1}, \mathbf{k}$  in a single calculation.



## 4.2 Product integral representation of conditional partial moments

From the differential equations for the conditional (partial) moments, we see a structure similar to those of a single payment stream, namely that one must use all lower order moments  $\mathbf{y}^1, \dots, \mathbf{y}^{|\mathbf{k}|-1} \in S(\mathbf{k})$  (each computed using Theorem 4.1 or 4.3) to compute the  $\mathbf{k}$ 'th moment. Since each of these need their corresponding lower order moments  $S(\mathbf{y}^m)$ ,  $m = 1, \dots, |\mathbf{k}| - 1$ , we are able to compute the  $\mathbf{k}$ 'th moment starting from the  $\mathbf{0}$ 'th moment, which is  $\mathbf{V}^{(0)}(s, t) = \mathbf{P}(s, t)$  (or  $V_i^{(0)}(s, t) = 1$ ), and then iteratively for each  $m = 1, \dots, |\mathbf{k}|$  compute the  $\mathbf{y}^m$ 'th moment using the lower order moments  $\mathbf{y}^1, \dots, \mathbf{y}^{m-1}$ . In total, one must then solve  $(|\mathbf{k}| + 1) J \times J$ -dimensional systems of differential equations for  $\mathbf{V}^{(\mathbf{k})}$ , and  $(|\mathbf{k}| + 1) J$ -dimensional systems for  $V_i^{(\mathbf{k})}$ .

However, this approach relies on how we sort the vectors  $\mathbf{y}^1, \dots, \mathbf{y}^{|\mathbf{k}|}$ , or, equivalently, the order at which we solve the differential equations, since we must ensure that the sets  $S(\mathbf{y}^m)$  are increasing, i.e. that  $S(\mathbf{y}^{m-1}) \subseteq S(\mathbf{y}^m)$  for all  $m = 1, \dots, |\mathbf{k}|$ , such that we actually are able to draw upon all lower order moments when calculating a given moment  $\mathbf{y}^m$ . Note that this is a trivial matter in the case of a single payment process, since for  $n = 1$  we have  $S(k) = \{1, \dots, k\}$ .

In this subsection, we derive a product integral representation for the conditional partial moments that demonstrates how one may compute all moments  $\mathbf{V}^{(\mathbf{y})}$ ,  $\mathbf{y} \in \tilde{S}(\mathbf{k})$ , at once. It turns out that the key to obtain the result is to order the vectors  $\mathbf{y}^1, \dots, \mathbf{y}^{|\mathbf{k}|}$  in a way that fortunately is standard in most software packages.

**Assumption 4.9.** *We assume that the lower order moments  $S(\mathbf{k}) = \{\mathbf{y}^1, \dots, \mathbf{y}^{|\mathbf{k}|}\}$  are lexicographical ordered, that is, they satisfy that for all  $m' > m$ ,  $m, m' \in \{1, \dots, |\mathbf{k}|\}$ , there exist  $u \in \{1, \dots, n\}$  (dependent on  $m$  and  $m'$ ), such that*

$$\begin{aligned} y_\ell^m &= y_\ell^{m'} \text{ for all } \ell < u, \text{ and} \\ y_u^m &< y_u^{m'}. \end{aligned}$$

In other words,  $\mathbf{y}^{m'}$  is strictly larger than  $\mathbf{y}^m$  in the first entry where they differ (it may be smaller in the remaining entries). Details regarding this type of ordering and its properties are presented in Appendix A.

Now, define the matrix  $\mathbf{F}_{\mathcal{U}}^{(\mathbf{k})}(x)$ ,  $x \geq 0$ , as

$$\begin{pmatrix} M(x) - \bar{k}r(x)\mathbf{I} & \mathbf{f}_1^{(\mathbf{k})}(x) & \mathbf{f}_2^{(\mathbf{k})}(x) & \dots & \mathbf{f}_{|\mathbf{k}|}^{(\mathbf{k})}(x) \\ \mathbf{0} & M(x) - \bar{\mathbf{y}}^{|\mathbf{k}|-1}r(x)\mathbf{I} & \mathbf{f}_1^{(\mathbf{y}^{|\mathbf{k}|-1})}(x) & \dots & \mathbf{f}_{|\mathbf{k}|-1}^{(\mathbf{y}^{|\mathbf{k}|-1})}(x) \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ \mathbf{0} & \mathbf{0} & \mathbf{0} & M(x) - \bar{\mathbf{y}}^1r(x)\mathbf{I} & \mathbf{f}_1^{(\mathbf{y}^1)}(x) \\ \mathbf{0} & \mathbf{0} & \mathbf{0} & \mathbf{0} & M(x) \end{pmatrix}$$

#### 4 HIGHER ORDER MOMENTS

with  $\mathbf{f}_u^{(\mathbf{y}^m)}(x)$ , for  $m = 1, \dots, |\mathbf{k}|$  and  $u = 1, \dots, m$ , given by, setting  $\mathbf{y}^0 := \mathbf{0}$ ,

$$\mathbf{f}_u^{(\mathbf{y}^m)}(x) = \begin{cases} y_1^m \mathbf{R}_1(x) & \text{if } \mathbf{y}^m - \mathbf{y}^{m-u} = \mathbf{e}_1, \\ y_2^m \mathbf{R}_2(x) & \text{if } \mathbf{y}^m - \mathbf{y}^{m-u} = \mathbf{e}_2, \\ \vdots & \vdots \\ y_n^m \mathbf{R}_n(x) & \text{if } \mathbf{y}^m - \mathbf{y}^{m-u} = \mathbf{e}_n, \\ \prod_{\ell=1}^n \binom{y_\ell^m}{\xi_\ell} C(\boldsymbol{\xi})(x), & \text{if } \mathbf{y}^m - \mathbf{y}^{m-u} = \boldsymbol{\xi} \in S(\mathbf{y}^m) \setminus E_n, \\ \mathbf{0} & \text{Otherwise.} \end{cases} \quad (4.8)$$

Here, one should think of the matrices  $\{\mathbf{f}_u^{(\mathbf{y}^m)}(x)\}_{u=1}^m$  as those needed to compute the  $\mathbf{y}^m$ 'th moment, cf. Theorem 4.1, and so each block row of  $\mathbf{F}_U^{(\mathbf{k})}(x)$  corresponds to a calculation of a lower order moment.

**Example 4.10** (Conditional covariance). For the computation of conditional covariances, we have for  $n = 2$  and  $\mathbf{k} = (1, 1)$  that the matrix  $\mathbf{F}_U^{(1,1)}(x)$  reads as

$$\mathbf{F}_U^{(1,1)}(x) = \begin{pmatrix} \mathbf{M}(x) - 2r(x)\mathbf{I} & \mathbf{R}_2(x) & \mathbf{R}_1(x) & C^{(1,1)}(x) \\ \mathbf{0} & \mathbf{M}(x) - r(x)\mathbf{I} & \mathbf{0} & \mathbf{R}_1(x) \\ \mathbf{0} & \mathbf{0} & \mathbf{M}(x) - r(x)\mathbf{I} & \mathbf{R}_2(x) \\ \mathbf{0} & \mathbf{0} & \mathbf{0} & \mathbf{M}(x) \end{pmatrix},$$

arising from the fact that the lexicographical ordering of  $S((1, 1))$  is given as  $(0, 1)$ ,  $(1, 0)$ , and  $(1, 1)$ .  $\circ$

Now, let

$$\mathbf{G}^{(\mathbf{k})}(s, t) = \prod_s^t \left( \mathbf{I} + \mathbf{F}_U^{(\mathbf{k})}(x) dx \right) \quad (4.9)$$

denote the product integral on the interval  $[s, t]$  of the matrix function  $\mathbf{F}_U^{(\mathbf{k})}$ . Since the matrix  $\mathbf{F}_U^{(\mathbf{k})}(x)$  is a  $(|\mathbf{k}| + 1) \times (|\mathbf{k}| + 1)$  block-partitioned matrix with blocks of sizes  $J \times J$ , we have that  $\mathbf{G}^{(\mathbf{k})}(s, t)$  is of similar size, and we denote with  $\mathbf{G}_{ij}^{(\mathbf{k})}(s, t)$  the  $ij$ 'th block of  $\mathbf{G}^{(\mathbf{k})}(s, t)$  for  $i, j \in \{1, \dots, |\mathbf{k}| + 1\}$ . The following result then demonstrates how all moments of up to order  $\mathbf{k}$  are obtained through this single calculation of a product integral.

**Theorem 4.11.** *For each  $i \in \{0, 1, \dots, |\mathbf{k}|\}$  we have, for  $m = 0, 1, \dots, i$ ,*

$$\mathbf{G}_{|\mathbf{k}|+1-i, |\mathbf{k}|+1-m}^{(\mathbf{k})}(s, t) = \mathbb{1}_{(\mathbf{y}^i \geq \mathbf{y}^m)} \prod_{\ell=1}^n \binom{y_\ell^i}{y_\ell^m} e^{-\bar{\mathbf{y}}^m \int_s^t r(v) dv} \mathbf{V}(\mathbf{y}^i - \mathbf{y}^m)(s, t), \quad (4.10)$$

which in particular gives the conditional  $\mathbf{y}^i$ 'th moment for  $m = 0$ :

$$\mathbf{G}_{|\mathbf{k}|+1-i, |\mathbf{k}|+1}^{(\mathbf{k})}(s, t) = \mathbf{V}(\mathbf{y}^i)(s, t). \quad (4.11)$$

#### 4 HIGHER ORDER MOMENTS

*Proof.* We shall mimic the proof of [4, Theorem 3-4] and modify suitably to the present setup. By multiplying (4.1) with  $\prod_t^s (\mathbf{I} + [\mathbf{M}(u) - \bar{\mathbf{k}}r(u)\mathbf{I}]) du$  on both sides we get

$$\begin{aligned} & \frac{\partial}{\partial s} \left( \prod_t^s (\mathbf{I} + [\mathbf{M}(u) - \bar{\mathbf{k}}r(u)\mathbf{I}]) \mathbf{V}^{(\mathbf{k})}(s, t) \right) \\ &= - \sum_{\ell=1}^n k_\ell \prod_t^s (\mathbf{I} + [\mathbf{M}(u) - \bar{\mathbf{k}}r(u)\mathbf{I}]) \mathbf{R}_\ell(s) \mathbf{V}^{(\mathbf{k}-\mathbf{e}_\ell)}(s, t) \\ & \quad - \sum_{\substack{\mathbf{y} \in S(\mathbf{k}) \\ \mathbf{y} \notin E_n}} \prod_{\ell=1}^n \binom{k_\ell}{y_\ell} \prod_t^s (\mathbf{I} + [\mathbf{M}(u) - \bar{\mathbf{k}}r(u)\mathbf{I}]) \mathbf{C}^{(\mathbf{y})}(s) \mathbf{V}^{(\mathbf{k}-\mathbf{y})}(s, t). \end{aligned}$$

Integrating the equation yields

$$\begin{aligned} \mathbf{V}^{(\mathbf{k})}(s, t) &= \sum_{\ell=1}^n k_\ell \int_s^t \prod_s^x (\mathbf{I} + [\mathbf{M}(u) - \bar{\mathbf{k}}r(u)\mathbf{I}]) \mathbf{R}_\ell(x) \mathbf{V}^{(\mathbf{k}-\mathbf{e}_\ell)}(x, t) dx \\ & \quad + \sum_{\substack{\mathbf{y} \in S(\mathbf{k}) \\ \mathbf{y} \notin E_n}} \prod_{\ell=1}^n \binom{k_\ell}{y_\ell} \int_s^t \prod_s^x (\mathbf{I} + [\mathbf{M}(u) - \bar{\mathbf{k}}r(u)\mathbf{I}]) \mathbf{C}^{(\mathbf{y})}(x) \mathbf{V}^{(\mathbf{k}-\mathbf{y})}(x, t) dx. \end{aligned}$$

We now provide an induction argument to verify the identity (4.10) claimed in the theorem using this integral equation. For  $i = 0$  and  $m = 0$  the identity is trivially true. So assume that the identity is true for some  $i - 1$ ,  $i \in \{1, \dots, |\mathbf{k}| + 1\}$ , and corresponding  $m = 0, 1, \dots, i - 1$ . Then [4, Lemma 1] gives us that

$$\mathbf{G}_{|\mathbf{k}|+1-i, |\mathbf{k}|+1-i}^{(\mathbf{k})}(s, t) = \prod_s^t (\mathbf{I} + [\mathbf{M}(x) - \bar{\mathbf{y}}^i r(x)\mathbf{I}]) dx = e^{-\bar{\mathbf{y}}^i \int_s^t r(v) dv} \mathbf{P}(s, t),$$

which is (4.9) for  $m = i$ . From the Lemma, it also follows that for  $m = 0, 1, \dots, i - 1$ ,

$$\mathbf{G}_{|\mathbf{k}|+1-i, |\mathbf{k}|+1-m}^{(\mathbf{k})}(s, t) = \sum_{j=1}^{i-m} \int_s^t e^{-\bar{\mathbf{y}}^i \int_s^x r(v) dv} \mathbf{P}(s, x) \mathbf{f}_j^{(\mathbf{y}^i)}(x) \mathbf{G}_{|\mathbf{k}|+1-(i-j), |\mathbf{k}|+1-m}^{(\mathbf{k})}(x, t) dx$$

From the induction hypothesis we have that

$$\mathbf{G}_{|\mathbf{k}|+1-(i-j), |\mathbf{k}|+1-m}^{(\mathbf{k})}(x, t) = \mathbb{1}_{(\mathbf{y}^{i-j} \geq \mathbf{y}^m)} \prod_{\ell=1}^n \binom{y_\ell^{i-j}}{y_\ell^m} e^{-\bar{\mathbf{y}}^m \int_x^t r(v) dv} \mathbf{V}^{(\mathbf{y}^{i-j}-\mathbf{y}^m)}(x, t),$$

and furthermore,  $\mathbf{f}_j^{(\mathbf{y}^i)}(x)$  may be written as

$$\mathbf{f}_j^{(\mathbf{y}^i)}(x) = \sum_{\ell=1}^n \mathbb{1}_{(\mathbf{y}^i - \mathbf{y}^{i-j} = \mathbf{e}_\ell)} y_\ell^i \mathbf{R}_\ell(x) + \sum_{\substack{\boldsymbol{\xi} \in S(\mathbf{y}^i) \\ \boldsymbol{\xi} \notin E_n}} \mathbb{1}_{(\mathbf{y}^i - \mathbf{y}^{i-j} = \boldsymbol{\xi})} \prod_{\ell=1}^n \binom{y_\ell^i}{\xi_\ell} \mathbf{C}^{(\boldsymbol{\xi})}(x).$$

#### 4 HIGHER ORDER MOMENTS

Note that when  $\mathbf{y}^i - \mathbf{y}^{i-j} = \boldsymbol{\xi}$  for some  $\boldsymbol{\xi} \in S(\mathbf{y}^i)$  (including the unit vectors) and some  $j = 1, \dots, i - m$ , it holds

$$\begin{pmatrix} y_\ell^{i-j} \\ y_\ell^m \end{pmatrix} \begin{pmatrix} y_\ell^i \\ \xi_\ell \end{pmatrix} = \begin{pmatrix} y_\ell^i \\ y_\ell^m \end{pmatrix} \begin{pmatrix} y_\ell^i - y_\ell^m \\ \xi_\ell \end{pmatrix}$$

for all  $\ell = 1, \dots, n$ . Thus, we now have

$$\begin{aligned} & \mathbf{G}_{|\mathbf{k}|+1-i, |\mathbf{k}|+1-m}^{(\mathbf{k})}(s, t) \\ &= \prod_{\ell=1}^n \begin{pmatrix} y_\ell^i \\ y_\ell^m \end{pmatrix} \int_s^t e^{-\bar{\mathbf{y}}^i \int_s^x r(v) dv} \mathbf{P}(s, x) e^{-\bar{\mathbf{y}}^m \int_x^t r(v) dv} \sum_{j=1}^{i-m} \mathbb{1}_{(\mathbf{y}^{i-j} \geq \mathbf{y}^m)} \times \\ & \quad \left\{ \sum_{\ell=1}^n \mathbb{1}_{(\mathbf{y}^i - \mathbf{y}^{i-j} = \mathbf{e}_\ell)} (y_\ell^i - y_\ell^m) \mathbf{R}_\ell(x) \mathbf{V}^{(\mathbf{y}^i - \mathbf{y}^m - \mathbf{e}_\ell)}(x, t) \right. \\ & \quad \left. + \sum_{\substack{\boldsymbol{\xi} \in S(\mathbf{y}^i) \\ \boldsymbol{\xi} \notin E_n}} \mathbb{1}_{(\mathbf{y}^i - \mathbf{y}^{i-j} = \boldsymbol{\xi})} \prod_{\ell=1}^n \begin{pmatrix} y_\ell^i - y_\ell^m \\ \xi_\ell \end{pmatrix} \mathbf{C}(\boldsymbol{\xi})(x) \mathbf{V}^{(\mathbf{y}^i - \mathbf{y}^m - \boldsymbol{\xi})}(x, t) \right\} dx \end{aligned}$$

Then observe that the discount factors can be factored as, for  $x \in [s, t]$ ,

$$e^{-\bar{\mathbf{y}}^i \int_s^x r(v) dv} e^{-\bar{\mathbf{y}}^m \int_x^t r(v) dv} = e^{-\bar{\mathbf{y}}^m \int_s^t r(v) dv} e^{-(\bar{\mathbf{y}}^i - \bar{\mathbf{y}}^m) \int_s^x r(v) dv}.$$

Furthermore, when  $\mathbf{y}^{i-j} \geq \mathbf{y}^m$  and  $\mathbf{y}^i - \mathbf{y}^{i-j} = \boldsymbol{\xi}$  for some  $\boldsymbol{\xi} \in S(\mathbf{y}^i)$  and  $j = 1, \dots, i - m$ , we have that  $\boldsymbol{\xi} \leq \mathbf{y}^i - \mathbf{y}^m$ , and so the last sum can be carried out over  $S(\mathbf{y}^i - \mathbf{y}^m) \setminus E_n$ . Thus, we now have

$$\begin{aligned} & \mathbf{G}_{|\mathbf{k}|+1-i, |\mathbf{k}|+1-m}^{(\mathbf{k})}(s, t) \\ &= \mathbb{1}_{(\mathbf{y}^i \geq \mathbf{y}^m)} \prod_{\ell=1}^n \begin{pmatrix} y_\ell^i \\ y_\ell^m \end{pmatrix} e^{-\bar{\mathbf{y}}^m \int_s^t r(v) dv} \times \\ & \quad \int_s^t e^{-(\bar{\mathbf{y}}^i - \bar{\mathbf{y}}^m) \int_s^x r(v) dv} \mathbf{P}(s, x) \left\{ \sum_{\ell=1}^n (y_\ell^i - y_\ell^m) \mathbf{R}_\ell(x) \mathbf{V}^{(\mathbf{y}^i - \mathbf{y}^m - \mathbf{e}_\ell)}(x, t) \right. \\ & \quad \left. + \sum_{\substack{\boldsymbol{\xi} \in S(\mathbf{y}^i - \mathbf{y}^m) \\ \boldsymbol{\xi} \notin E_n}} \prod_{\ell=1}^n \begin{pmatrix} y_\ell^i - y_\ell^m \\ \xi_\ell \end{pmatrix} \mathbf{C}(\boldsymbol{\xi})(x) \mathbf{V}^{(\mathbf{y}^i - \mathbf{y}^m - \boldsymbol{\xi})}(x, t) \right\} dx \\ &= \mathbb{1}_{(\mathbf{y}^i \geq \mathbf{y}^m)} \prod_{\ell=1}^n \begin{pmatrix} y_\ell^i \\ y_\ell^m \end{pmatrix} e^{-\bar{\mathbf{y}}^m \int_s^t r(v) dv} \mathbf{V}^{(\mathbf{y}^i - \mathbf{y}^m)}(s, t), \end{aligned}$$

as claimed. In the first equality, we have used that for all  $\boldsymbol{\xi} \in S(\mathbf{y}^i - \mathbf{y}^m)$ ,

$$\sum_{j=1}^{i-m} \mathbb{1}_{(\mathbf{y}^i - \mathbf{y}^{i-j} = \boldsymbol{\xi})} = \mathbb{1}_{(\mathbf{y}^i \geq \mathbf{y}^m)},$$

which follows from Lemma A.1. □

## 5 NUMERICAL EXAMPLES

The theorem gives us that the right-block column of  $\mathbf{G}^{(\mathbf{k})}(s, t)$  contains all moments of order up to  $\mathbf{k}$ , including the  $\mathbf{0}$ 'th moment, as follows:

$$\prod_s^t \left( \mathbf{I} + \mathbf{F}_U^{(\mathbf{k})}(x) dx \right) = \begin{pmatrix} * & * & * & * & \dots & * & \mathbf{V}^{(\mathbf{k})}(s, t) \\ * & * & * & * & \dots & * & \mathbf{V}^{(\mathbf{y}^{|\mathbf{k}|-1})}(s, t) \\ * & * & * & * & \dots & * & \mathbf{V}^{(\mathbf{y}^{|\mathbf{k}|-2})}(s, t) \\ \vdots & \vdots & \vdots & \vdots & & \vdots & \vdots \\ * & * & * & * & \dots & * & \mathbf{V}^{(\mathbf{y}^1)}(s, t) \\ * & * & * & * & \dots & * & \mathbf{P}(s, t) \end{pmatrix},$$

and so all lower order up to order  $\mathbf{k}$  are obtained through this product integral calculation. From this, the conditional moments  $V_i^{(\mathbf{k})}$  and corresponding central moments  $m_i^{(\mathbf{k})}$  are obtained via (2.6) and (4.3), respectively.

## 5 Numerical examples

In this section, we illustrate the methods presented in the previous sections in a numerical example of the motivating examples Example 1.1 and 1.2 considered in the introduction.

### 5.1 Disability model


We consider a 40-year old male today at time 0 with retirement age 65, such that  $T = 25$ . We suppose that the valuation basis is taken to be the technical basis in the numerical example of [5], which is given by the following:

$$\begin{aligned} r(s) &= 0.01, \\ \mu_{01}(s) &= \left( 0.0004 + 10^{4.54+0.06(s+40)-10} \right) \mathbf{1}_{(s \leq 25)}, \\ \mu_{10}(s) &= \left( 2.0058 \cdot e^{-0.117(s+40)} \right) \mathbf{1}_{(s \leq 25)}, \\ \mu_{02}(s) &= 0.0005 + 10^{5.88+0.038(s+40)-10}, \\ \mu_{12}(s) &= \mu_{02}(s) \left( 1 + \mathbf{1}_{(s \leq 25)} \right). \end{aligned}$$


Some of the transition intensities are inspired by the Danish G82M technical basis. We examine the pair-wise covariance and correlation structures of the product combinations in the active state, i.e. the entries of the conditional covariance and correlation matrix given in (4.6)–(4.7) with  $i = 0$ ; this is illustrated in Figure 3–4.

We see that the biggest dependence is between the life annuity and death benefit, which as one would have expected, but we also see a similar (but opposite) dependence structure between the disability annuity and death benefit respectively life annuity.

## 5 NUMERICAL EXAMPLES



**Figure 3:** Pair-wise conditional covariances  $[0, 25] \in s \mapsto \Sigma_0(s, 70)_{\ell m}$ ,  $\ell, m \in \{1, 2, 3\}$ ,  $\ell \leq m$ , between the three products until retirement conditional on the insured being active.



**Figure 4:** Pair-wise conditional correlations  $[0, 25] \in s \mapsto \rho_0(s, 70)_{\ell m}$ ,  $\ell, m \in \{1, 2, 3\}$ ,  $\ell < m$ , between the three products until retirement conditional on the insured being active.

## 5 NUMERICAL EXAMPLES

The death benefit seem to be slightly more correlated with the disability annuity than the life annuity is, which is the most non-trivial observation encountered here.

### 5.2 Spouse model


We consider a married couple where both individuals are 25 years old at time 0 with expiry on their life insurances at retirement age 65; thus  $T = 40$ . Again, we take as inspiration the Danish G82 technical basis as valuation elements for the mortality risk of the husband and wife (before each of them are dead). Thus, we have  $r(s) = 0.01$ , and

$$\begin{aligned}\mu(s) &= 0.0005 + 10^{5.88+0.038(s+25)-10}, \\ \nu(s) &= 0.0005 + 10^{5.728+0.038(s+25)-10}.\end{aligned}$$

To illustrate an example of dependence between their two lives, we suppose that the mortality of the husband increases upon death of his wife, while the mortality of the wife remains unchanged upon death of her husband; this phenomena has been observed in many statistical studies of mortality. For simplicity, we assume that the increase in the husbands mortality happens doubles constantly in all ages. Formally, this reads as

$$\begin{aligned}\tilde{\nu}(s) &= \nu(s), \\ \tilde{\mu}(s) &= 2\mu(s).\end{aligned}$$

Note that we have assumed the husbands mortality as widower is the same as his mortality as disabled in Subsection 5.1.



**Figure 5:** *Pair-wise covariances*  $[0, 40] \in s \mapsto \Sigma_0(s, 40)_{\ell m}$ ,  $\ell, m \in \{1, 2\}$ ,  $\ell \leq m$ , and *correlations*  $[0, 40] \in s \mapsto \rho_0(s, 40)_{\ell m}$ ,  $\ell < m$ , between the coverages to the husband and wife given they both are alive.

## REFERENCES

In Figure 5 we illustrate the conditional variance of the coverages to each of the individuals along with their covariance and correlation given they both are alive.

We see that the coverage between the husband and wife are negatively correlated throughout the contract period, which as we would expect since payments work in the opposite directions w.r.t. their individual lifetimes. However, we also see that the correlation becomes more negative when they are young, and then increases towards zero when they come closer to retirement.

## Acknowledgments

I should like to thank my PhD supervisors Mogens Bladt and Mogens Steffensen for valuable suggestions and fruitful discussions during this work.

## References

- [1] F. Adékambi and M.C. Christiansen. Integral and differential equations for the moments of multistate models in health insurance. *Scandinavian Actuarial Journal*, 2017(1):29–50, 2017.
- [2] F. Adékambi and M.C. Christiansen. Probability distributions of multi-states insurance models under semi-markov assumptions. *Markov Processes and Related Fields*, 26(3):517–534, 2020.
- [3] S. Asmussen and M. Steffensen. *Risk and Insurance*. Series on Probability Theory and Stochastic Modelling. Springer, 2020.
- [4] M. Bladt, S. Asmussen, and M. Steffensen. Matrix representations of life insurance payments. *European Actuarial Journal*, pages 1–39, 01 2020.
- [5] K. Buchardt and T. Møller. Life Insurance Cash Flows with Policyholder Behavior. *Risks*, 3(3):290–317, 2015.
- [6] S.H. Cox and Y. Lin. Natural hedging of life and annuity mortality risks. *North American actuarial journal*, 11(3):1–15, 2007.
- [7] R.D. Gill and S. Johansen. A survey of product-integration with a view toward application in survival analysis. *The Annals of statistics*, 18(4):1501–1555, 1990.
- [8] O. Hesselager and R. Norberg. On probability distributions of present values in life insurance. *Insurance, mathematics and economics*, 18(1):35–42, 1996.



## A LEXICOGRAPHICAL ORDERING

- [9] J.M. Hoem. Markov chain models in life insurance. *Blätter der DGVFM*, 9:91–107, 1969.
- [10] J.M. Hoem and O.O. Aalen. Actuarial values of payment streams. *Scandinavian Actuarial Journal*, 1978(1):38–47, 1978.
- [11] S. Johansen. Product integrals and markov processes. *CWI Newsletter*, (12):3–13, 1986.
- [12] S. Levantesi and M. Menzietti. Natural hedging in long-term care insurance. *ASTIN Bulletin*, 48(1):233–274, 2018.
- [13] R. Norberg. Reserves in Life and Pension Insurance. *Scandinavian Actuarial Journal*, 1991:3–24, 1991.
- [14] R. Norberg. Hattendorff’s theorem and Thiele’s differential equation generalized. *Scandinavian Actuarial Journal*, 1992(1):2–14, 1992.
- [15] R. Norberg. Differential equations for moments of present values in life insurance. *Insurance: Mathematics and Economics*, 17(2):171–180, 1995.
- [16] H. Ramlau-Hansen. Hattendorff’s theorem: A markov chain and counting process approach. *Scandinavian Actuarial Journal*, 1988(1-3):143–156, 1988.

## A Lexicographical ordering

Let  $\mathbf{k} = (k_1, \dots, k_n) \in \mathbb{N}_0^n$  be a  $n$ -dimensional vector of natural numbers including zero. We think of this vector as representing some multivariate higher order moment we wish to calculate for the multivariate present value  $\mathbf{U}$ . Specific to this, we need to consider all combinations of lower order moments represented by the elements of  $S(\mathbf{k})$ ; recall (2.1) for its definition. Write these elements as

$$S(\mathbf{k}) = \left\{ \mathbf{y}^1, \dots, \mathbf{y}^{|\mathbf{k}|} \right\}.$$

The vectors  $\mathbf{y}^1, \dots, \mathbf{y}^{|\mathbf{k}|}$  are then said to be *lexicographical ordered* if for all  $m' > m$ , there exist  $u$  (dependent on  $m$  and  $m'$ ) such that

$$\begin{aligned} y_\ell^m &= y_\ell^{m'} \text{ for all } \ell < u, \text{ and} \\ y_u^m &< y_u^{m'}. \end{aligned}$$

In other words  $y^{m'}$  must be strictly larger than  $y^m$  in the first entry where they differ (it may be smaller in the remaining entries), and one commonly writes  $\mathbf{y}^m <_{\text{lex}} \mathbf{y}^{m'}$ . An illustration of the ordering is presented in Table 1.

## A LEXICOGRAPHICAL ORDERING

$$\begin{array}{lll}
\mathbf{y}^1 = (0, \dots, 0, 1)' & \mathbf{y}^{k_n+1} = (0, \dots, 1, 0) & \mathbf{y}^{|\mathbf{k}|-k_n} = (k_1, \dots, k_{n-1}, 0)' \\
\mathbf{y}^2 = (0, \dots, 0, 2)' & \mathbf{y}^{k_n+2} = (0, \dots, 1, 1) & \mathbf{y}^{|\mathbf{k}|-k_n+1} = (k_1, \dots, k_{n-1}, 1)' \\
\vdots & \vdots & \vdots \\
\mathbf{y}^{k_n} = (0, \dots, 0, k_n)' & \mathbf{y}^{2(k_n+1)-1} = (0, \dots, 1, k_n)' & \mathbf{y}^{|\mathbf{k}|} = (k_1, \dots, k_n)'
\end{array}$$

**Table 1:** Illustration of lexicographical ordering of vectors in  $S(\mathbf{k})$ .

The following result on this type of ordering provides the foundation to successively compute multivariate higher order moments based on already computed lower order moments.

**Theorem A.1.** Assume that  $\mathbf{y}^1, \dots, \mathbf{y}^{|\mathbf{k}|}$  are lexicographical ordered. Then for all  $i = 1, \dots, |\mathbf{k}|$  and  $m = 0, 1, \dots, i-1$  such that  $\mathbf{y}^i \geq \mathbf{y}^m$ , we have

$$S(\mathbf{y}^i - \mathbf{y}^m) \subseteq \bigcup_{j=m}^{i-1} \{\mathbf{y}^i - \mathbf{y}^j\}.$$

*Sketch of proof.* We show the result for the two-dimensional case  $n = 2$ , and the generalization to higher dimension then follows using same (but notationally cumbersome) techniques. Let  $i = 1, \dots, |\mathbf{k}|$  be given. From the illustration of the lexicographic ordering of vectors in  $S(\mathbf{k})$  shown in Table 1, we may realize that the vectors take the form

$$\mathbf{y}^i = (a, i - a(k_2 + 1))', \quad \text{if } i \in [a(k_2 + 1), (a + 1)(k_2 + 1) - 1]$$

where  $a = 1, \dots, k_1$ . This can be verified by exploiting the structure of which the ordering is carried out. Now let  $m = 0, 1, \dots, i-1$  be given such that  $\mathbf{y}^i \geq \mathbf{y}^m$ . Then similarly there exist  $b = 0, 1, \dots, k_1$  such that  $m \in [b(k_2 + 1), (b + 1)(k_2 + 1) - 1]$ , which then gives

$$\mathbf{y}^i - \mathbf{y}^m = (a - b, i - m - (a - b)(k_2 + 1))', \quad (\text{A.1})$$

By similar reasonings, we have for  $j = m, \dots, i-1$  there exist  $c = 1, \dots, k_1$  such that  $j \in [c(k_2 + 1), (c + 1)(k_2 + 1) - 1]$ , giving

$$\mathbf{y}^i - \mathbf{y}^j = (a - c, i - j - (a - c)(k_2 + 1))'.$$

Due to the indices having the order  $m \leq j \leq i$ , we have that  $b \leq c \leq a$ . Consequently, the set  $\bigcup_{j=m}^{i-1} \{\mathbf{y}^i - \mathbf{y}^j\}$  consists of vectors on the form  $(a - c, i - j - (a - c)(k_2 + 1))'$  where  $j$  varies on  $m, \dots, i-1$  and  $c$  varies on  $b, b + 1, \dots, a$ .

Now take  $\boldsymbol{\xi} \in S(\mathbf{y}^i - \mathbf{y}^m)$ . Since  $\boldsymbol{\xi} \leq \mathbf{y}^i - \mathbf{y}^m$ , it follows from (A.1) that there exist  $d \leq a - b$  and  $r \leq i - m - (a - b)(k_2 + 1)$ ,  $d, r \in \mathbb{N}_0$ , such that

$$\begin{aligned}
\boldsymbol{\xi} &= (a - (b + d), i - (m + r) - (a - b)(k_2 + 1))' \\
&= (a - (b + d), i - (m + r + d(k_2 + 1)) - (a - (b + d))(k_2 + 1))',
\end{aligned}$$

## A LEXICOGRAPHICAL ORDERING

where we have added and subtracted the term  $d(k_2+1)$  in the second coordinate to obtain the last equality. We now see that  $\xi$  is on the form  $(a - c, i - j - (a - c)(k_2 + 1))'$  with  $c = b + d$  and  $j = m + r + d(k_2 + 1)$ , and so we have  $\xi \in \bigcup_{j=m}^{i-1} \{\mathbf{y}^i - \mathbf{y}^j\}$ , as claimed. This concludes the sketch of the proof.  $\square$