

Convergence rates for gradient descent in the training of overparameterized artificial neural networks with biases

Arnulf Jentzen^{1,2} and Timo Kröger³

¹ Faculty of Mathematics and Computer Science, University of Münster, Münster, Germany; e-mail: ajentzen@uni-muenster.de

² School of Data Science and Shenzhen Research Institute of Big Data, The Chinese University of Hong Kong, Shenzhen, China; e-mail: ajentzen@cuhk.edu.cn

³ Faculty of Mathematics and Computer Science, University of Münster, Münster, Germany; e-mail: timo.kroeger@uni-muenster.de

November 25, 2021

Abstract

In recent years, artificial neural networks have developed into a powerful tool for dealing with a multitude of problems for which classical solution approaches reach their limits. However, it is still unclear why randomly initialized gradient descent optimization algorithms, such as the well-known batch gradient descent, are able to achieve zero training loss in many situations even though the objective function is non-convex and non-smooth. One of the most promising approaches to solving this problem in the field of supervised learning is the analysis of gradient descent optimization in the so-called overparameterized regime. In this article we provide a further contribution to this area of research by considering overparameterized fully-connected rectified artificial neural networks with biases. Specifically, we show that for a fixed number of training data the mean squared error using batch gradient descent optimization applied to such a randomly initialized artificial neural network converges to zero at a linear convergence rate as long as the width of the artificial neural network is large enough, the learning rate is small enough, and the training input data are pairwise linearly independent.

Keywords: artificial neural network, ANN, gradient descent, GD, overparameterization, empirical risk minimization, optimization, random initialization, Gram matrix

Contents

1	Introduction	2
2	Mathematical framework for gradient descent (GD) optimization algorithms	4
2.1	Mathematical description of artificial neural networks (ANNs)	4
2.2	Gradients of the considered risk functions	4
2.3	Mathematical description of GD processes	5
2.4	Connection between deterministic and stochastic Gramian matrices	6
2.5	Positive definiteness of deterministic Gramian matrices for nondegenerated data	6
3	Analysis of eigenvalues of stochastic Gramian matrices	9
3.1	Probabilistic error analysis at initialization	9
3.2	Analysis of weights and biases of ANNs during training	11
3.3	Properties of subexponential random variables	13
3.4	Concentration type inequalities for stochastic Gramian matrices at initialization	16
3.5	Analysis of stochastic Gramian matrices during training	17
3.6	Analysis of eigenvalues of stochastic Gramian matrices during training	19

4	Error analysis for GD optimization algorithms in the training of ANNs	21
4.1	Quantitative pathwise error analysis for GD optimization algorithms	21
4.2	Quantitative probabilistic error analysis for GD optimization algorithms	28
4.3	Qualitative probabilistic error analysis for GD optimization algorithms	33

1 Introduction

Many problems, such as recognizing faces, handwritten text, or natural language, as well as performing various activities, such as driving a car, seem very simple to humans, but pose a major challenge to computers. The reason probably lies in the variety of ways in which the data to be processed can exist, combined with the fact that computers, unlike humans, are not able to intuitively recognize complex patterns in them due to their structurally precise data processing. As a result, it is very difficult to hard-code a program or provide a function which attaches to the input – be it a picture, an audio file, a game situation, or a traffic situation of an autonomous driving car – a desired meaning or guidance. Nevertheless, artificial neural networks (ANNs) have become a powerful tool in dealing with such problems in recent times. Although many ANN training algorithms have demonstrated great success in practice, the reasons for this are generally not known, as no mathematically rigorous analysis exists for most algorithms. An overview of different gradient descent (GD) variations commonly used for ANN optimization is given in Ruder [25].

There are several promising attempts in the scientific literature which intend to mathematically analyze GD optimization algorithms in the training of ANNs. In particular, there are various convergence results for GD optimization algorithms in the training of ANNs that assume convexity of the considered objective functions (cf., e.g., [4, 5, 24] and the references mentioned therein), there are general abstract convergence results for GD optimization algorithms that do not assume convexity of the considered objective functions (cf., e.g., [1, 6, 10, 14, 19, 20, 22] and the references mentioned therein), there are divergence results and lower bounds for GD optimization algorithms in the training of ANNs (cf., e.g., [8, 18, 23] and the references mentioned therein), there are mathematical analyzes regarding the initialization in the training of ANNs with GD optimization algorithms (cf., e.g., [15, 16, 23, 27] and the references mentioned therein), and there are convergence results for GD optimization algorithms in the training of ANNs in the case of constant target functions (cf. [7]).

One of the most promising approaches in the field supervised learning is the analysis of GD optimization in the so-called overparameterized regime. In this regime the number of parameters determining the ANN highly exceeds the number of available training data. Under this condition the convergence of the GD optimization method and some of its variants has been proved in several settings even though the objective function is non-convex and non-smooth (cf., e.g., [2, 3, 9, 11, 12, 13, 29] and the references mentioned therein). In this overparameterized regime progress was also made in the analysis of the popular stochastic GD optimization algorithm regarding the influence of the architecture of an ANN on the so-called gradient confusion, which directly affects the convergence speed (cf. Sankararaman et al. [26]), regarding the characterization of stability properties of global minima (cf. Wu et al. [28]), and regarding the multi-class classification with additional considerations for the generalization error (cf. Li & Liang [21]). Another promising approach in the context of overparameterization is the use of kernel methods. In particular, it was proved that the realization of an ANN during GD optimization follows the kernel gradient of the functional costs with respect to the neural tangent kernel (cf., e.g., Jacot et al. [17]).

In this work we provide a further contribution by extending Du et al. [12] in the following way. We extend the considered ANNs with biases on the hidden layer and output layer and initialize all weights normally distributed – as usual in practice with mean 0 and the same variance on each layer. In addition, we relax the requirements on the training data. This approach is in contrast to other articles in this area of research that extend Du et al. [12], for example, by considering ANNs with multiple layers without biases and smooth activation functions instead of the rectifier function (cf. Du et al. [11]) or consider ANNs with multiple layers in the context of stochastic GD (cf. Zou et al. [29]). To illustrate the findings of this article in a special case we now present Theorem 1.1 and we refer to Section 4.2 below for the more general convergence results which we develop in this article. Below Theorem 1.1 we also provide some explanations regarding the mathematical objects that are introduced within Theorem 1.1.

Theorem 1.1. Let $d, m \in \mathbb{N}$, $x_1, x_2, \dots, x_m \in \mathbb{R}^d \setminus \{0\}$, $y_1, y_2, \dots, y_m \in \mathbb{R}$ satisfy for all $\lambda \in \mathbb{R}$, $i, j \in \{1, 2, \dots, m\}$ with $i \neq j$ that $x_i \neq \lambda x_j$, let $\varphi_t: \mathbb{R} \rightarrow \mathbb{R}$, $t \in (0, \infty]$, and $\mathcal{P}: (-\infty, \infty] \rightarrow \mathbb{N}$ satisfy for all $t \in (0, \infty)$, $z, \mathfrak{d} \in \mathbb{R}$ that $\varphi_t(z) = \frac{1}{t} \ln(1 + t \exp(tz))$, $\varphi_\infty(z) = \max\{z, 0\}$, and $\mathcal{P}(\mathfrak{d}) \geq \mathfrak{d}d + 2\mathfrak{d} + 1$, let $\mathcal{N}_\theta^{\mathfrak{d},t}: \mathbb{R}^d \rightarrow \mathbb{R}$, $\mathfrak{d}, t \in [0, \infty]$, $\theta \in \mathbb{R}^{\mathcal{P}(\mathfrak{d})}$, and $\mathcal{E}^{\mathfrak{d},t}: \mathbb{R}^{\mathcal{P}(\mathfrak{d})} \rightarrow [0, \infty)$, $\mathfrak{d}, t \in [0, \infty]$, satisfy for all $\mathfrak{d} \in \mathbb{N}$, $t \in (0, \infty)$, $\theta = (\theta_1, \theta_2, \dots, \theta_{\mathcal{P}(\mathfrak{d})}) \in \mathbb{R}^{\mathcal{P}(\mathfrak{d})}$, $z = (z_1, z_2, \dots, z_d) \in \mathbb{R}^d$ that $\mathcal{N}_\theta^{\mathfrak{d},t}(z) = \sum_{k=1}^{\mathfrak{d}} \theta_{\mathfrak{d}d+\mathfrak{d}+k} \varphi_t(\sum_{\ell=1}^d \theta_{(k-1)d+\ell} z_\ell + \theta_{\mathfrak{d}d+k}) + \theta_{\mathfrak{d}d+2\mathfrak{d}+1}$ and $\mathcal{E}^{\mathfrak{d},t}(\theta) = \frac{1}{m} \sum_{i=1}^m |\mathcal{N}_\theta^{\mathfrak{d},t}(x_i) - y_i|^2$, let $(\Omega, \mathcal{F}, \mathbb{P})$ be a probability space, let $\Theta^{\mathfrak{d},\eta} = (\Theta_1^{\mathfrak{d},\eta}, \Theta_2^{\mathfrak{d},\eta}, \dots, \Theta_{\mathcal{P}(\mathfrak{d})}^{\mathfrak{d},\eta}): \mathbb{N}_0 \times \Omega \rightarrow \mathbb{R}^{\mathcal{P}(\mathfrak{d})}$, $\mathfrak{d}, \eta \in \mathbb{R}$, be measurable, assume for all $\mathfrak{d}, \eta \in \mathbb{R}$ that $\Theta_1^{\mathfrak{d},\eta}(0), \Theta_2^{\mathfrak{d},\eta}(0), \dots, \Theta_{\mathfrak{d}d}^{\mathfrak{d},\eta}(0), \sqrt{\mathfrak{d}}\Theta_{\mathfrak{d}d+\mathfrak{d}+1}^{\mathfrak{d},\eta}(0), \sqrt{\mathfrak{d}}\Theta_{\mathfrak{d}d+\mathfrak{d}+2}^{\mathfrak{d},\eta}(0), \dots, \sqrt{\mathfrak{d}}\Theta_{\mathfrak{d}d+2\mathfrak{d}}^{\mathfrak{d},\eta}(0)$ are independent and standard normal, and assume for all $\mathfrak{d}, n \in \mathbb{N}_0$, $\eta \in \mathbb{R}$, $i \in \{1, 2, \dots, \mathcal{P}(\mathfrak{d})\}$ that $\Theta_{\mathfrak{d}d+1}^{\mathfrak{d},\eta}(0) = \Theta_{\mathfrak{d}d+2}^{\mathfrak{d},\eta}(0) = \dots = \Theta_{\mathfrak{d}d+\mathfrak{d}}^{\mathfrak{d},\eta}(0) = \Theta_{\mathfrak{d}d+2\mathfrak{d}+1}^{\mathfrak{d},\eta}(0) = 0$ and $\Theta_i^{\mathfrak{d},\eta}(n+1) = \Theta_i^{\mathfrak{d},\eta}(n) - \eta [\lim_{t \rightarrow \infty} (\frac{\partial}{\partial \theta_i} \mathcal{E}^{\mathfrak{d},t})(\Theta^{\mathfrak{d},\eta}(n))] \mathbb{1}_{\mathbb{R} \setminus \{\mathfrak{d}d+\mathfrak{d}, \mathfrak{d}d+2\mathfrak{d}\}}(i)$. Then there exists $\lambda \in (0, 1)$ such that for all $\varepsilon \in (0, 1)$, $\eta \in (0, \lambda \varepsilon^2)$, $\mathfrak{d} \in \mathbb{N} \cap [\lambda^{-1} \varepsilon^{-4}, \infty)$ it holds that

$$\mathbb{P}(\forall n \in \mathbb{N}_0: \mathcal{E}^{\mathfrak{d},\infty}(\Theta^{\mathfrak{d},\eta}(n)) \leq (1 - \eta \lambda)^n \mathcal{E}^{\mathfrak{d},\infty}(\Theta^{\mathfrak{d},\eta}(0))) \geq 1 - \varepsilon. \quad (1)$$

Theorem 1.1 is an immediate consequence of Corollary 4.10 in Section 4 below. Corollary 4.10 follows from Corollary 4.8 which, in turn, builds on a series of intermediate results which are all based on Theorem 4.3. In the following we add some comments and explanations regarding the mathematical objects which appear in Theorem 1.1 above. The natural number $d \in \mathbb{N} = \{1, 2, 3, \dots\}$ in Theorem 1.1 above specifies the dimension of the input data of the considered training data. The natural number $m \in \mathbb{N}$ specifies the number of input-output data pairs which are used in the training of the considered ANN. The vectors $x_1, x_2, \dots, x_m \in \mathbb{R}^d \setminus \{0\}$ constitute the input data used for the training, whereas the real numbers $y_1, y_2, \dots, y_m \in \mathbb{R}$ constitute the output data used for the training. In Theorem 1.1 above we employ fully connected ANNs with 1 hidden layer and the rectifier function $\mathbb{R} \ni z \mapsto \max\{z, 0\} \in \mathbb{R}$ as the activation function in front of the hidden layer. Note that the function $\mathbb{R} \ni z \mapsto \max\{z, 0\} = \varphi_\infty(z) \in \mathbb{R}$ is not differentiable at $0 \in \mathbb{R}$ and, as a consequence, the associated risk function is not always differentiable. To overcome this obstacle we modify the rectifier function by defining the GD method on smooth approximations of the risk function and then consider the limit. In particular, observe that the real valued functions $\varphi_t: \mathbb{R} \rightarrow \mathbb{R}$, $t \in (0, \infty]$, which are introduced for this purpose, satisfy that for all $z \in \mathbb{R}$ we have that

$$\lim_{t \rightarrow \infty} \varphi_t(z) = \varphi_\infty(z) = \max\{z, 0\}. \quad (2)$$

Moreover, note that for all $z \in \mathbb{R}$, $t \in (0, \infty)$ it holds that φ_t is differentiable at z and $\lim_{t \rightarrow \infty} (\varphi_t)'(z) = \mathbb{1}_{[0, \infty)}(z)$. Observe that for all $z \in \mathbb{R} \setminus \{0\}$ we have that this limit coincides with the derivative of the rectifier function φ_∞ .

The function $\mathcal{P}: (-\infty, \infty] \rightarrow \mathbb{N}$ specifies the number of parameters that form an ANN with $\mathfrak{d} \in \mathbb{N}$ neurons on the hidden layer and must therefore satisfy for all $\mathfrak{d} \in \mathbb{N}$ that $\mathcal{P}(\mathfrak{d}) \geq \mathfrak{d}d + 2\mathfrak{d} + 1$. Next observe that for every $\mathfrak{d} \in \mathbb{N}$, $t \in (0, \infty]$, $\theta = (\theta_1, \theta_2, \dots, \theta_{\mathcal{P}(\mathfrak{d})}) \in \mathbb{R}^{\mathcal{P}(\mathfrak{d})}$ we have that the function $\mathcal{N}_\theta^{\mathfrak{d},t}: \mathbb{R}^d \rightarrow \mathbb{R}$ in Theorem 1.1 above describes the realization of a fully connected ANN with \mathfrak{d} neurons on the hidden layer and φ_t as the activation function in front of the hidden layer. The vector $\theta = (\theta_1, \theta_2, \dots, \theta_{\mathcal{P}(\mathfrak{d})}) \in \mathbb{R}^{\mathcal{P}(\mathfrak{d})}$ stores the real parameters (weights and biases) for the concrete considered ANN. Moreover, note that for every $\mathfrak{d} \in \mathbb{N}$, $t \in (0, \infty]$ we have that the function $\mathcal{E}^{\mathfrak{d},t}: \mathbb{R}^{\mathcal{P}(\mathfrak{d})} \rightarrow [0, \infty)$ measures the average squared difference (mean squared error) between the estimated values $\mathcal{N}_\theta^{\mathfrak{d},t}(x_1), \mathcal{N}_\theta^{\mathfrak{d},t}(x_2), \dots, \mathcal{N}_\theta^{\mathfrak{d},t}(x_m)$ and the actual values y_1, y_2, \dots, y_m . Observe that (2) implies for all $\mathfrak{d} \in \mathbb{N}$, $\theta \in \mathbb{R}^{\mathcal{P}(\mathfrak{d})}$ that

$$\lim_{t \rightarrow \infty} \mathcal{E}^{\mathfrak{d},t}(\theta) = \mathcal{E}^{\mathfrak{d},\infty}(\theta). \quad (3)$$

Next note that for every $\mathfrak{d} \in \mathbb{N}_0$, $\eta \in \mathbb{R}$ we have that the stochastic process $\Theta^{\mathfrak{d},\eta}: \mathbb{N}_0 \times \Omega \rightarrow \mathbb{R}^{\mathcal{P}(\mathfrak{d})}$ is associated to the GD process of an ANN with \mathfrak{d} neurons on the hidden layer, which is randomly initialized and optimized with GD using the constant learning rate η . To be more precise, the parameters of the considered ANN in Theorem 1.1 above are initialized as follows. All parameters are initialized independently, the weights from the input layer to the hidden layer are standard normal distributed, the weights from the hidden layer to the output layer are normal distributed with mean 0 and variance $1/\mathfrak{d}$, and all biases are 0. After that, all parameters except for the weights from the hidden layer to the output layer are optimized with GD using the learning rate η .

Under these conditions, Theorem 1.1 above states that there exists a convergence rate $\lambda \in (0, 1)$ such that for every arbitrary small error probability $\varepsilon \in (0, 1)$ and every ANN with $\mathfrak{d} \in \mathbb{N} \cap [\lambda^{-1}\varepsilon^{-4}, \infty)$ neurons on the hidden layer, initialized and trained as above with learning rate $\eta \in (0, \lambda\varepsilon^2)$, it holds with probability at least $1 - \varepsilon$ that in every iteration step $n \in \mathbb{N}_0$ the associated empirical risk is bounded by the initial empirical risk and the factor $(1 - \lambda\eta)^n$. In particular, this implies that the associated empirical risk converges to zero at a linear convergence rate. For simplicity, in Theorem 1.1 above, we only assert the existence of such a convergence rate λ and give very rough bounds on the learning rate η and network width \mathfrak{d} . However, in the more detailed results in Section 4 below, we show how λ can be computed using only the input training data x_1, x_2, \dots, x_m (cf. (161)) and we also give much tighter bounds on η and \mathfrak{d} . The arguments in this work are partially inspired by the existing techniques in the overparameterized regime, in particular, by the arguments in Du et al. [12].

The remainder of this articles is organized as follows. In Section 2 we recall a common approach to mathematically describe ANNs and we introduce the setting of GD processes that we often use in our article. In Section 3 we analyze some of the mathematical objects that appear in this setting. In particular, in Section 3 we deal with two Gramian matrices and their eigenvalues, which are essential for the proof of the main result. In Section 4 we combine the results from Section 3 to establish error analyzes for the considered GD processes.

2 Mathematical framework for gradient descent (GD) optimization algorithms

This section is devoted to the mathematical description of ANNs (cf. Section 2.1 below) and to the introduction of the mathematical framework for GD optimization algorithms (cf. Section 2.3 below). For the understanding of our main setting, which we often impose in this article, Setting 2.7 below, we show in Lemma 2.5 that the specific algorithm used in Setting 2.7 results directly from the gradients of the considered risk functions. In Lemma 2.8 we explain the connection between the deterministic and stochastic Gramian matrices and in Lemma 2.11 we prove that the deterministic Gramian matrices only have positive eigenvalues if the considered input training data is nondegenerate, by which we mean that the input training data are pairwise linearly independent. In our proof of Lemma 2.11 we use a well-known property of an appropriate Gramian matrix which we establish in Lemma 2.9. Only for completeness we include in this section the proof of Lemma 2.9.

2.1 Mathematical description of artificial neural networks (ANNs)

Definition 2.1 (Standard scalar product and norm). We denote by $\langle \cdot, \cdot \rangle: (\bigcup_{n \in \mathbb{N}} (\mathbb{R}^n \times \mathbb{R}^n)) \rightarrow \mathbb{R}$ and $\|\cdot\|: (\bigcup_{n \in \mathbb{N}} \mathbb{R}^n) \rightarrow [0, \infty)$ the functions which satisfy for all $n \in \mathbb{N}$, $x = (x_1, x_2, \dots, x_n)$, $y = (y_1, y_2, \dots, y_n) \in \mathbb{R}^n$ that $\langle x, y \rangle = \sum_{i=1}^n x_i y_i$ and $\|x\| = (\sum_{i=1}^n |x_i|^2)^{1/2}$.

Definition 2.2 (ANNs). Let $d, \mathfrak{d} \in \mathbb{N}$. Then we denote by $\mathbf{N}_{d, \mathfrak{d}}$ the set given by $\mathbf{N}_{d, \mathfrak{d}} = ((\mathbb{R}^{\mathfrak{d} \times d} \times \mathbb{R}^{\mathfrak{d}}) \times (\mathbb{R}^{\mathfrak{d}} \times \mathbb{R}))$.

Definition 2.3 (Rectifier function). We denote by $\mathfrak{r}: (\bigcup_{d \in \mathbb{N}} \mathbb{R}^d) \rightarrow (\bigcup_{d \in \mathbb{N}} \mathbb{R}^d)$ the function which satisfies for all $d \in \mathbb{N}$, $x = (x_1, x_2, \dots, x_d) \in \mathbb{R}^d$ that $\mathfrak{r}(x) = (\max\{x_1, 0\}, \max\{x_2, 0\}, \dots, \max\{x_d, 0\})$.

Definition 2.4 (Realization associated to a rectified ANN). We denote by $\mathcal{R}: (\bigcup_{d, \mathfrak{d} \in \mathbb{N}} \mathbf{N}_{d, \mathfrak{d}}) \rightarrow (\bigcup_{d \in \mathbb{N}} C(\mathbb{R}^d, \mathbb{R}))$ the function which satisfies for all $d, \mathfrak{d} \in \mathbb{N}$, $\Phi = ((W, B), (\mathcal{W}, \mathfrak{B})) \in \mathbf{N}_{d, \mathfrak{d}}$, $x \in \mathbb{R}^d$ that $\mathcal{R}(\Phi) \in C(\mathbb{R}^d, \mathbb{R})$ and $(\mathcal{R}(\Phi))(x) = \langle \mathcal{W}, \mathfrak{r}(Wx + B) \rangle + \mathfrak{B}$ (cf. Definitions 2.1–2.3).

2.2 Gradients of the considered risk functions

Lemma 2.5. Let $d, \mathfrak{d}, m \in \mathbb{N}$, $x_1, x_2, \dots, x_m \in \mathbb{R}^d$, $y_1, y_2, \dots, y_m \in \mathbb{R}$, $e_1 = (1, 0, \dots, 0)$, $e_2 = (0, 1, 0, \dots, 0)$, \dots , $e_d = (0, \dots, 0, 1) \in \mathbb{R}^d$, let $\mathcal{E}: \mathbf{N}_{d, \mathfrak{d}} \rightarrow \mathbb{R}$ satisfy for all $\Phi \in \mathbf{N}_{d, \mathfrak{d}}$ that $\mathcal{E}(\Phi) = \frac{1}{m} \sum_{i=1}^m |(\mathcal{R}(\Phi))(x_i) - y_i|^2$, let $D_i \subseteq \mathbf{N}_{d, \mathfrak{d}}$, $i \in \{1, 2, \dots, m\}$, satisfy for all $i \in \{1, 2, \dots, m\}$ that $D_i = \{((W, B), (\mathcal{W}, \mathfrak{B})) \in \mathbf{N}_{d, \mathfrak{d}}: (Wx_i + B) \in (\mathbb{R} \setminus \{0\})^{\mathfrak{d}}\}$, let $W = (W_1, W_2, \dots, W_{\mathfrak{d}}) = (W_{k, \ell})_{(k, \ell) \in \{1, 2, \dots, \mathfrak{d}\} \times \{1, 2, \dots, d\}} \in \mathbb{R}^{\mathfrak{d} \times d}$, $B = (B_1, B_2, \dots, B_{\mathfrak{d}})$, $\mathcal{W} = (\mathcal{W}_1, \mathcal{W}_2, \dots, \mathcal{W}_{\mathfrak{d}}) \in \mathbb{R}^{\mathfrak{d}}$, $\mathfrak{B} \in \mathbb{R}$, $\Phi \in \mathbf{N}_{d, \mathfrak{d}}$ satisfy $\Phi = ((W, B), (\mathcal{W}, \mathfrak{B}))$ and $\Phi \in (\bigcap_{i=1}^m D_i)$, and let $k \in \{1, 2, \dots, \mathfrak{d}\}$, $\ell \in \{1, 2, \dots, d\}$ (cf. Definitions 2.2 and 2.4). Then

(i) it holds that $\mathbb{R} \ni x \mapsto \mathcal{E}(((W_{1,1}, \dots, W_{1,d}), \dots, (W_{k,1}, \dots, W_{k,\ell-1}, x, W_{k,\ell+1}, \dots, W_{k,d}), \dots, (W_{\mathfrak{d},1}, \dots, W_{\mathfrak{d},d}), B), (\mathcal{W}, \mathfrak{B})) \in \mathbb{R}$ is differentiable at $W_{k,\ell}$,

(ii) it holds that $\mathbb{R} \ni x \mapsto \mathcal{E}((W, (B_1, \dots, B_{k-1}, x, B_{k+1}, \dots, B_{\mathfrak{d}})), (\mathcal{W}, \mathfrak{B})) \in \mathbb{R}$ is differentiable at B_k ,

(iii) it holds that $\mathbb{R} \ni x \mapsto \mathcal{E}((W, B), (\mathcal{W}, x)) \in \mathbb{R}$ is differentiable at \mathfrak{B} ,

(iv) it holds that $\frac{\partial \mathcal{E}(\Phi)}{\partial W_{k,\ell}} = \frac{2}{m} \sum_{j=1}^m ((\mathcal{R}(\Phi))(x_j) - y_j) \mathcal{W}_k \mathbb{1}_{[0,\infty)}(\langle W_k, x_j \rangle + B_k) \langle x_j, e_s \rangle$,

(v) it holds that $\frac{\partial \mathcal{E}(\Phi)}{\partial B_k} = \frac{2}{m} \sum_{j=1}^m ((\mathcal{R}(\Phi))(x_j) - y_j) \mathcal{W}_k \mathbb{1}_{[0,\infty)}(\langle W_k, x_j \rangle + B_k)$, and

(vi) it holds that $\frac{\partial \mathcal{E}(\Phi)}{\partial \mathfrak{B}} = \frac{2}{m} \sum_{j=1}^m ((\mathcal{R}(\Phi))(x_j) - y_j)$

(cf. Definition 2.1).

Proof of Lemma 2.5. First, observe that

$$\begin{aligned} \mathcal{E}(\Phi) &= \frac{1}{m} \sum_{j=1}^m |(\mathcal{R}(\Phi))(x_j) - y_j|^2 = \frac{1}{m} \sum_{j=1}^m |\langle \mathcal{W}, \mathfrak{r}(W x_j + B) \rangle + \mathfrak{B} - y_j|^2 \\ &= \frac{1}{m} \sum_{j=1}^m \left| \sum_{r=1}^{\mathfrak{d}} \mathcal{W}_r \mathfrak{r}(\langle W_r, x_j \rangle + B_r) + \mathfrak{B} - y_j \right|^2 = \frac{1}{m} \sum_{j=1}^m \left| \sum_{r=1}^{\mathfrak{d}} \mathcal{W}_r \mathfrak{r} \left(\sum_{s=1}^d W_{r,s} \langle x_j, e_s \rangle + B_r \right) + \mathfrak{B} - y_j \right|^2 \end{aligned} \quad (4)$$

(cf. Definitions 2.1 and 2.3). This, the fact that for all $x \in \mathbb{R} \setminus \{0\}$ it holds that $\mathbb{R} \ni y \mapsto \mathfrak{r}(y) \in \mathbb{R}$ is differentiable at x , the fact that for all $r \in \{1, 2, \dots, \mathfrak{d}\}$, $j \in \{1, 2, \dots, m\}$ it holds that $\langle W_r, x_j \rangle + B_r \in \mathbb{R} \setminus \{0\}$, and the fact that sums and compositions of differentiable functions are differentiable establish items (i), (ii), and (iii). Moreover, observe that (4) and the fact that for all $x \in \mathbb{R} \setminus \{0\}$ it holds that $\frac{\partial \mathfrak{r}(x)}{\partial x} = \mathbb{1}_{[0,\infty)}(x)$ imply that it holds that

$$\begin{aligned} \frac{\partial \mathcal{E}(\Phi)}{\partial W_{k,\ell}} &= \frac{2}{m} \sum_{j=1}^m ((\mathcal{R}(\Phi))(x_j) - y_j) \frac{\partial}{\partial W_{k,\ell}} \left(\sum_{r=1}^{\mathfrak{d}} \mathcal{W}_r \mathfrak{r} \left(\sum_{s=1}^d W_{r,s} \langle x_j, e_s \rangle + B_r \right) + \mathfrak{B} - y_j \right) \\ &= \frac{2}{m} \sum_{j=1}^m ((\mathcal{R}(\Phi))(x_j) - y_j) \mathcal{W}_k \mathbb{1}_{[0,\infty)}(\langle W_k, x_j \rangle + B_k) \langle x_j, e_\ell \rangle, \end{aligned} \quad (5)$$

$$\begin{aligned} \frac{\partial \mathcal{E}(\Phi)}{\partial B_k} &= \frac{2}{m} \sum_{j=1}^m ((\mathcal{R}(\Phi))(x_j) - y_j) \frac{\partial}{\partial B_k} \left(\sum_{r=1}^{\mathfrak{d}} \mathcal{W}_r \mathfrak{r}(\langle W_r, x_j \rangle + B_r) + \mathfrak{B} - y_j \right) \\ &= \frac{2}{m} \sum_{j=1}^m ((\mathcal{R}(\Phi))(x_j) - y_j) \mathcal{W}_k \mathbb{1}_{[0,\infty)}(\langle W_k, x_j \rangle + B_k), \end{aligned} \quad (6)$$

and

$$\begin{aligned} \frac{\partial \mathcal{E}(\Phi)}{\partial \mathfrak{B}} &= \frac{2}{m} \sum_{j=1}^m ((\mathcal{R}(\Phi))(x_j) - y_j) \frac{\partial}{\partial \mathfrak{B}} \left(\sum_{r=1}^{\mathfrak{d}} \mathcal{W}_r \mathfrak{r}(\langle W_r, x_j \rangle + B_r) + \mathfrak{B} - y_j \right) \\ &= \frac{2}{m} \sum_{j=1}^m ((\mathcal{R}(\Phi))(x_j) - y_j). \end{aligned} \quad (7)$$

This establishes items (iv), (v), and (vi). The proof of Lemma 2.5 is thus completed. \square

2.3 Mathematical description of GD processes

Definition 2.6 (Smallest eigenvalue). We denote by $\lambda_{\min}: (\bigcup_{n \in \mathbb{N}} \mathbb{R}^{n \times n}) \rightarrow [-\infty, \infty]$ the function which satisfies for all $n \in \mathbb{N}$, $A \in \mathbb{R}^{n \times n}$ that $\lambda_{\min}(A) = \min(\{\lambda \in \mathbb{R}: [\exists v \in \mathbb{R}^n \setminus \{0\}: Av = \lambda v]\} \cup \{\infty\})$.

Setting 2.7. Let $d, \mathfrak{d}, m \in \mathbb{N}$, $x_1, x_2, \dots, x_m \in \mathbb{R}^d \setminus \{0\}$, $y = (y_1, y_2, \dots, y_m) \in \mathbb{R}^m$, $\eta \in (0, \infty)$, $c, C \in \mathbb{R}$ satisfy $c = \min_{i \in \{1, 2, \dots, m\}} \|x_i\|$ and $C = \max_{i \in \{1, 2, \dots, m\}} \|x_i\|$ (cf. Definition 2.1), let $(\Omega, \mathcal{F}, \mathbb{P})$ be a probability space, let $W = (W_1, W_2, \dots, W_{\mathfrak{d}}): \mathbb{N}_0 \times \Omega \rightarrow \mathbb{R}^{\mathfrak{d} \times d}$, $B = (B_1, B_2, \dots, B_{\mathfrak{d}}): \mathbb{N}_0 \times \Omega \rightarrow \mathbb{R}^{\mathfrak{d}}$, $\mathcal{W} = (\mathcal{W}_1, \mathcal{W}_2, \dots, \mathcal{W}_{\mathfrak{d}}): \Omega \rightarrow \mathbb{R}^{\mathfrak{d}}$, and $\mathfrak{B}: \mathbb{N}_0 \times \Omega \rightarrow \mathbb{R}$ be measurable, let $\Phi: \mathbb{N}_0 \times \Omega \rightarrow \mathbf{N}_{d, \mathfrak{d}}$ and $f = (f_1, f_2, \dots, f_m): \mathbb{N}_0 \times \Omega \rightarrow \mathbb{R}^m$ satisfy for all $i \in \{1, 2, \dots, m\}$, $n \in \mathbb{N}_0$, $\omega \in \Omega$ that $\Phi(n, \omega) = ((W(n, \omega), B(n, \omega)), (\mathcal{W}(\omega), \mathfrak{B}(n, \omega)))$ and $f_i(n, \omega) =$

$(\mathcal{R}(\Phi(n, \omega)))(x_i)$ (cf. Definitions 2.2 and 2.4), assume that $W_1(0), W_2(0), \dots, W_{\mathfrak{d}}(0), \sqrt{\mathfrak{d}}\mathcal{W}_1, \sqrt{\mathfrak{d}}\mathcal{W}_2, \dots, \sqrt{\mathfrak{d}}\mathcal{W}_{\mathfrak{d}}$ are independent and standard normal, assume for all $k \in \{1, 2, \dots, \mathfrak{d}\}$, $n \in \mathbb{N}_0$, $\omega \in \Omega$ that

$$W_k(n+1, \omega) = W_k(n, \omega) - \frac{2\eta}{m} \left(\sum_{j=1}^m (f_j(n, \omega) - y_j) \mathcal{W}_k(\omega) \mathbb{1}_{\{(W_k(n), x_j) + B_k(n) \geq 0\}}(\omega) x_j \right), \quad (8)$$

$$B_k(n+1, \omega) = B_k(n, \omega) - \frac{2\eta}{m} \left(\sum_{j=1}^m (f_j(n, \omega) - y_j) \mathcal{W}_k(\omega) \mathbb{1}_{\{(W_k(n), x_j) + B_k(n) \geq 0\}}(\omega) \right), \quad (9)$$

$$\mathfrak{B}(n+1, \omega) = \mathfrak{B}(n, \omega) - \frac{2\eta}{m} \left(\sum_{j=1}^m (f_j(n, \omega) - y_j) \right), \quad (10)$$

and $\|B(0, \omega)\| = |\mathfrak{B}(0, \omega)| = 0$, let $\mathcal{G} = (\mathcal{G}_{i,j})_{(i,j) \in \{1,2,\dots,m\}^2} : \mathbb{N}_0 \times \Omega \rightarrow \mathbb{R}^{m \times m}$ and $\mathcal{H} = (\mathcal{H}_{i,j})_{(i,j) \in \{1,2,\dots,m\}^2} : \mathbb{N}_0 \times \Omega \rightarrow \mathbb{R}^{m \times m}$ satisfy for all $i, j \in \{1, 2, \dots, m\}$, $n \in \mathbb{N}_0$, $\omega \in \Omega$ that

$$\mathcal{G}_{i,j}(n, \omega) = \sum_{k=1}^{\mathfrak{d}} |\mathcal{W}_k(\omega)|^2 \mathbb{1}_{\{(W_k(n), x_i) + B_k(n) \geq 0, \langle W_k(n), x_j \rangle + B_k(n) \geq 0\}}(\omega) \quad (11)$$

and $\mathcal{H}_{i,j}(n, \omega) = \langle x_i, x_j \rangle \mathcal{G}_{i,j}(n, \omega)$, and let $\mathbf{G} = (\mathbf{G}_{i,j})_{(i,j) \in \{1,2,\dots,m\}^2}$, $\mathbf{H} = (\mathbf{H}_{i,j})_{(i,j) \in \{1,2,\dots,m\}^2} \in \mathbb{R}^{m \times m}$, $\lambda, \mu \in [0, \infty)$ satisfy for all $i, j \in \{1, 2, \dots, m\}$ that $\mathbf{G}_{i,j} = \mathbb{P}(\langle W_1(0), x_i \rangle \geq 0, \langle W_1(0), x_j \rangle \geq 0)$, $\mathbf{H}_{i,j} = \langle x_i, x_j \rangle \mathbf{G}_{i,j}$, $\lambda = \lambda_{\min}(\mathbf{G})$, and $\mu = \lambda_{\min}(\mathbf{H})$ (cf. Definition 2.6).

2.4 Connection between deterministic and stochastic Gramian matrices

Lemma 2.8. Assume Setting 2.7. Then it holds for all $i, j \in \{1, 2, \dots, m\}$ that $\mathbf{G}_{i,j} = \mathbb{E}[\mathcal{G}_{i,j}(0)]$ and $\mathbf{H}_{i,j} = \mathbb{E}[\mathcal{H}_{i,j}(0)]$.

Proof of Lemma 2.8. Observe that the assumption that $W_1(0), W_2(0), \dots, W_{\mathfrak{d}}(0), \sqrt{\mathfrak{d}}\mathcal{W}_1, \sqrt{\mathfrak{d}}\mathcal{W}_2, \dots, \sqrt{\mathfrak{d}}\mathcal{W}_{\mathfrak{d}}$ are independent implies that for all $i, j \in \{1, 2, \dots, m\}$, $k \in \{1, 2, \dots, \mathfrak{d}\}$ it holds that $\mathbb{1}_{\{(W_k(0), x_i) \geq 0, \langle W_k(0), x_j \rangle \geq 0\}}$ and \mathcal{W}_k are independent. Combining this with (11), the assumption that $\|B(0)\| = 0$, the assumption that $W_1(0), W_2(0), \dots, W_{\mathfrak{d}}(0)$ are identically distributed, and the fact that for all $k \in \{1, 2, \dots, \mathfrak{d}\}$ it holds that \mathcal{W}_k is a centered normal random variable with $\text{Var}[\mathcal{W}_k] = 1/\mathfrak{d}$ ensures that for all $i, j \in \{1, 2, \dots, m\}$ it holds that

$$\begin{aligned} \mathbb{E}[\mathcal{G}_{i,j}(0)] &= \mathbb{E} \left[\sum_{k=1}^{\mathfrak{d}} |\mathcal{W}_k|^2 \mathbb{1}_{\{(W_k(0), x_i) + B_k(0) \geq 0, \langle W_k(0), x_j \rangle + B_k(0) \geq 0\}} \right] \\ &= \sum_{k=1}^{\mathfrak{d}} \mathbb{E} [|\mathcal{W}_k|^2 \mathbb{1}_{\{(W_k(0), x_i) \geq 0, \langle W_k(0), x_j \rangle \geq 0\}}] = \sum_{k=1}^{\mathfrak{d}} \mathbb{E} [|\mathcal{W}_k|^2] \mathbb{E} [\mathbb{1}_{\{(W_k(0), x_i) \geq 0, \langle W_k(0), x_j \rangle \geq 0\}}] \\ &= \sum_{k=1}^{\mathfrak{d}} \frac{1}{\mathfrak{d}} \mathbb{P}(\langle W_1(0), x_i \rangle \geq 0, \langle W_1(0), x_j \rangle \geq 0) = \mathbb{P}(\langle W_1(0), x_i \rangle \geq 0, \langle W_1(0), x_j \rangle \geq 0) = \mathbf{G}_{i,j} \end{aligned} \quad (12)$$

(cf. Definition 2.1). Hence, we obtain for all $i, j \in \{1, 2, \dots, m\}$ that $\mathbb{E}[\mathcal{H}_{i,j}(0)] = \mathbb{E}[\langle x_i, x_j \rangle \mathcal{G}_{i,j}(0)] = \langle x_i, x_j \rangle \mathbb{E}[\mathcal{G}_{i,j}(0)] = \langle x_i, x_j \rangle \mathbf{G}_{i,j} = \mathbf{H}_{i,j}$. This completes the proof of Lemma 2.8. \square

2.5 Positive definiteness of deterministic Gramian matrices for nondegenerated data

Lemma 2.9. Let V be a vector space over \mathbb{R} , let $\varphi: V \times V \rightarrow \mathbb{R}$ be an inner product on V , let $n \in \mathbb{N}$, $v_1, v_2, \dots, v_n \in V$, and let $G = (G_{i,j})_{(i,j) \in \{1,2,\dots,n\}^2} \in \mathbb{R}^{n \times n}$ satisfy for all $i, j \in \{1, 2, \dots, n\}$ that $G_{i,j} = \varphi(v_i, v_j)$. Then

(i) it holds that G is symmetric and positive semidefinite and

(ii) it holds that G is positive definite if and only if $\{v_1, v_2, \dots, v_n\}$ is linearly independent.

Proof of Lemma 2.9. First, observe that the assumption that φ is symmetric implies that G is symmetric. In addition, note that the assumption that for all $i, j \in \{1, 2, \dots, n\}$ it holds that $G_{i,j} = \varphi(v_i, v_j)$ and the assumption that φ is an inner product on V ensure that for all $x = (x_1, x_2, \dots, x_n) \in \mathbb{R}^n$ it holds that

$$\langle x, Gx \rangle = \sum_{i=1}^n x_i \sum_{j=1}^n G_{i,j} x_j = \sum_{i=1}^n \sum_{j=1}^n x_i x_j \varphi(v_i, v_j) = \sum_{i=1}^n \sum_{j=1}^n \varphi(x_i v_i, x_j v_j) = \varphi \left(\sum_{i=1}^n x_i v_i, \sum_{j=1}^n x_j v_j \right) \geq 0 \quad (13)$$

(cf. Definition 2.1). This establishes item (i). Moreover, observe that (13) and the assumption that φ is positive definite establish that it holds that

$$\begin{aligned}
G \text{ is positive definite} &\iff \forall x = (x_1, x_2, \dots, x_n) \in \mathbb{R}^n \setminus \{0\}: \langle x, Gx \rangle > 0 \\
&\iff \forall x = (x_1, x_2, \dots, x_n) \in \mathbb{R}^n \setminus \{0\}: \varphi\left(\sum_{i=1}^n x_i v_i, \sum_{i=1}^n x_i v_i\right) > 0 \\
&\iff \forall x = (x_1, x_2, \dots, x_n) \in \mathbb{R}^n \setminus \{0\}: \sum_{i=1}^n x_i v_i \neq 0 \\
&\iff \{v_1, v_2, \dots, v_n\} \text{ is linearly independent.}
\end{aligned} \tag{14}$$

This establishes item (ii). The proof of Lemma 2.9 is thus completed. \square

Lemma 2.10. *Let $d, m \in \mathbb{N}$, $x_1, x_2, \dots, x_m \in \mathbb{R}^d \setminus \{0\}$ satisfy for all $\lambda \in \mathbb{R}$, $i, j \in \{1, 2, \dots, m\}$ with $i \neq j$ that $x_i \neq \lambda x_j$, and let $D_i \subseteq \mathbb{R}^d$, $i \in \{1, 2, \dots, m\}$, satisfy for all $i \in \{1, 2, \dots, m\}$ that $D_i = \{w \in \mathbb{R}^d: \langle w, x_i \rangle = 0\}$ (cf. Definition 2.1). Then it holds for all $i \in \{1, 2, \dots, m\}$ that $D_i \not\subseteq \bigcup_{j \in \{1, 2, \dots, m\}, j \neq i} D_j$.*

Proof of Lemma 2.10. Throughout this proof let μ be the Lebesgue measure on $(\mathbb{R}^{d-1}, \mathcal{B}(\mathbb{R}^{d-1}))$ and let $A^{i,j} = (A_{k,l}^{i,j})_{(k,l) \in \{1,2\} \times \{1,2,\dots,d\}} \in \mathbb{R}^{2 \times d}$, $(i,j) \in \{1, 2, \dots, m\}^2$, satisfy for all $i, j \in \{1, 2, \dots, m\}$, $k \in \{1, 2\}$, $l \in \{1, 2, \dots, d\}$ that

$$(A_{1,1}^{i,j}, A_{1,2}^{i,j}, \dots, A_{1,d}^{i,j}) = x_i \quad \text{and} \quad (A_{2,1}^{i,j}, A_{2,2}^{i,j}, \dots, A_{2,d}^{i,j}) = x_j. \tag{15}$$

First, note that the fact that for all $i \in \{1, 2, \dots, m\}$ it holds that $x_i \neq 0$ implies for all $i \in \{1, 2, \dots, m\}$ that $\dim(D_i) = d - 1$. This ensures that for all $i \in \{1, 2, \dots, m\}$ there exists a linear isomorphism $T_i: D_i \rightarrow \mathbb{R}^{d-1}$. Furthermore, observe that for all $i, j \in \{1, 2, \dots, m\}$ it holds that $D_i \cap D_j = \{w \in \mathbb{R}^d: \langle w, x_i \rangle = \langle w, x_j \rangle = 0\} = \ker(A^{i,j})$. This, the assumption that for all $\lambda \in \mathbb{R}$, $i, j \in \{1, 2, \dots, m\}$ with $i \neq j$ it holds that $x_i \neq \lambda x_j$, and the rank-nullity theorem ensure that for all $i, j \in \{1, 2, \dots, m\}$ with $i \neq j$ it holds that

$$\dim(T_i(D_i \cap D_j)) = \dim(D_i \cap D_j) = \dim(\ker(A^{i,j})) = d - 2. \tag{16}$$

Hence, we obtain that for all $i, j \in \{1, 2, \dots, m\}$ with $i \neq j$ it holds that $\mu(T_i(D_i \cap D_j)) = 0$. This establishes for all $i \in \{1, 2, \dots, m\}$ that

$$0 \leq \mu\left(T_i\left(D_i \cap \bigcup_{j \neq i} D_j\right)\right) = \mu\left(\bigcup_{j \neq i} T_i(D_i \cap D_j)\right) \leq \sum_{j \neq i} \mu(T_i(D_i \cap D_j)) = 0. \tag{17}$$

Thus, we obtain for all $i \in \{1, 2, \dots, m\}$ that $\mu(T_i(D_i \cap (\bigcup_{j \neq i} D_j))) = 0$. This and the fact that for all $i \in \{1, 2, \dots, m\}$ it holds that $\mu(T_i(D_i)) = \mu(\mathbb{R}^{d-1}) = \infty \neq 0$ demonstrate that for all $i \in \{1, 2, \dots, m\}$ it holds that $D_i \not\subseteq \bigcup_{j \in \{1, 2, \dots, m\}, j \neq i} D_j$. The proof of Lemma 2.10 is thus completed. \square

Lemma 2.11. *Let $d, m \in \mathbb{N}$, $x_1, x_2, \dots, x_m \in \mathbb{R}^d \setminus \{0\}$ satisfy for all $\lambda \in \mathbb{R}$, $i, j \in \{1, 2, \dots, m\}$ with $i \neq j$ that $x_i \neq \lambda x_j$, let $(\Omega, \mathcal{F}, \mathbb{P})$ be a probability space, let $W: \Omega \rightarrow \mathbb{R}^d$ be standard normal, and let $\mathbf{G} = (\mathbf{G}_{i,j})_{(i,j) \in \{1,2,\dots,m\}^2}$, $\mathbf{H} = (\mathbf{H}_{i,j})_{(i,j) \in \{1,2,\dots,m\}^2} \in \mathbb{R}^{m \times m}$, $\lambda, \mu \in [0, \infty)$ satisfy for all $i, j \in \{1, 2, \dots, m\}$ that $\mathbf{G}_{i,j} = \mathbb{P}(\langle W, x_i \rangle \geq 0, \langle W, x_j \rangle \geq 0)$, $\mathbf{H}_{i,j} = \langle x_i, x_j \rangle \mathbf{G}_{i,j}$, $\lambda = \lambda_{\min}(\mathbf{G})$, and $\mu = \lambda_{\min}(\mathbf{H})$ (cf. Definitions 2.1 and 2.6). Then $\lambda, \mu \in (0, \infty)$.*

Proof of Lemma 2.11. Throughout this proof let $D_i \subseteq \mathbb{R}^d$, $i \in \{1, 2, \dots, m\}$, satisfy for all $i \in \{1, 2, \dots, m\}$ that $D_i = \{w \in \mathbb{R}^d: \langle w, x_i \rangle = 0\}$, let $A_r^i(z), B_r^i(z) \subseteq \mathbb{R}^d$, $k \in (0, \infty)$, $i \in \{1, 2, \dots, m\}$, $z \in \mathbb{R}^d$, satisfy for all $r \in (0, \infty)$, $i \in \{1, 2, \dots, m\}$, $z \in \mathbb{R}^d$ that

$$A_r^i(z) = \{w \in \mathbb{R}^d: \|w - z\| \leq r, \langle w, x_i \rangle > 0\} \quad \text{and} \quad B_r^i(z) = \{w \in \mathbb{R}^d: \|w - z\| \leq r, \langle w, x_i \rangle < 0\}, \tag{18}$$

let \mathcal{G}, \mathcal{H} satisfy $\mathcal{G} = L^2(\mathbb{R}^d, \mathcal{B}(\mathbb{R}^d), \mathbb{P}_W; \mathbb{R})$ and $\mathcal{H} = L^2(\mathbb{R}^d, \mathcal{B}(\mathbb{R}^d), \mathbb{P}_W; \mathbb{R}^d)$, let $\varphi: \mathcal{G} \times \mathcal{G} \rightarrow \mathbb{R}$ and $\psi: \mathcal{H} \times \mathcal{H} \rightarrow \mathbb{R}$ satisfy for all $g, h \in \mathcal{G}$ and $\mathcal{g}, \mathcal{h} \in \mathcal{H}$ that

$$\varphi(g, h) = \int_{\mathbb{R}^d} gh \, d\mathbb{P}_W \quad \text{and} \quad \psi(\mathcal{g}, \mathcal{h}) = \int_{\mathbb{R}^d} \langle \mathcal{g}, \mathcal{h} \rangle \, d\mathbb{P}_W, \tag{19}$$

let $g_x: \mathbb{R}^d \rightarrow \mathbb{R}$, $x \in \mathbb{R}^d$, and $h_x: \mathbb{R}^d \rightarrow \mathbb{R}^d$, $x \in \mathbb{R}^d$, satisfy for all $x, w \in \mathbb{R}^d$ that

$$g_x(w) = \mathbb{1}_{[0, \infty)}(\langle w, x \rangle) \quad \text{and} \quad h_x(w) = x \mathbb{1}_{[0, \infty)}(\langle w, x \rangle), \quad (20)$$

let $\alpha_1, \alpha_2, \dots, \alpha_m \in \mathbb{R}$, $\beta_1, \beta_2, \dots, \beta_m \in \mathbb{R}$, and assume $\sum_{j=1}^m \alpha_j g_{x_j} = 0 \in \mathcal{G}$ and $\sum_{j=1}^m \beta_j h_{x_j} = 0 \in \mathcal{H}$. First, observe that φ is an inner product on \mathcal{G} and ψ is an inner product on \mathcal{H} . Moreover, observe that for all $x \in \mathbb{R}^d$ it holds that $g_x \in \mathcal{G}$ and $h_x \in \mathcal{H}$. Next note that (19) and (20) ensure that for all $i, j \in \{1, 2, \dots, m\}$ it holds that

$$\begin{aligned} \mathbf{G}_{i,j} &= \mathbb{P}(\langle W, x_i \rangle \geq 0, \langle W, x_j \rangle \geq 0) = \mathbb{E}[\mathbb{1}_{\{\langle W, x_i \rangle \geq 0, \langle W, x_j \rangle \geq 0\}}] = \mathbb{E}[\mathbb{1}_{\{\langle W, x_i \rangle \geq 0\}} \mathbb{1}_{\{\langle W, x_j \rangle \geq 0\}}] \\ &= \int_{\mathbb{R}^d} \mathbb{1}_{[0, \infty)}(\langle w, x_i \rangle) \mathbb{1}_{[0, \infty)}(\langle w, x_j \rangle) d\mathbb{P}_W(w) = \int_{\mathbb{R}^d} g_{x_i}(w) g_{x_j}(w) d\mathbb{P}_W(w) = \varphi(g_{x_i}, g_{x_j}) \end{aligned} \quad (21)$$

and

$$\begin{aligned} \mathbf{H}_{i,j} &= \langle x_i, x_j \rangle \mathbf{G}_{i,j} = \langle x_i, x_j \rangle \int_{\mathbb{R}^d} \mathbb{1}_{[0, \infty)}(\langle w, x_i \rangle) \mathbb{1}_{[0, \infty)}(\langle w, x_j \rangle) d\mathbb{P}_W(w) \\ &= \int_{\mathbb{R}^d} \langle x_i, x_j \rangle \mathbb{1}_{[0, \infty)}(\langle w, x_i \rangle) \mathbb{1}_{[0, \infty)}(\langle w, x_j \rangle) d\mathbb{P}_W(w) \\ &= \int_{\mathbb{R}^d} \langle x_i \mathbb{1}_{[0, \infty)}(\langle w, x_i \rangle), x_j \mathbb{1}_{[0, \infty)}(\langle w, x_j \rangle) \rangle d\mathbb{P}_W(w) = \int_{\mathbb{R}^d} h_{x_i}(w) h_{x_j}(w) d\mathbb{P}_W(w) = \psi(h_{x_i}, h_{x_j}). \end{aligned} \quad (22)$$

In the next step, we show that $g_{x_1}, g_{x_2}, \dots, g_{x_m}$ respectively $h_{x_1}, h_{x_2}, \dots, h_{x_m}$ are linearly independent. Note that Lemma 2.10 ensures that there exist $z_1, z_2, \dots, z_m \in \mathbb{R}^d$ which satisfy for all $i \in \{1, 2, \dots, m\}$ that $z_i \in D_i \setminus \cup_{j \neq i} D_j$. This and the fact that for all $i \in \{1, 2, \dots, m\}$ it holds that $x_i \neq 0$ ensure that there exist $R_1, R_2, \dots, R_m \in (0, \infty)$ which satisfy for all $i \in \{1, 2, \dots, m\}$ that

$$R_i = \min \left\{ \frac{|\langle z_i, x_j \rangle|}{2 \|x_j\|} : j \in \{1, 2, \dots, m\}, j \neq i \right\}. \quad (23)$$

Combining this and the Cauchy Schwarz inequality ensures that for all $i, j \in \{1, 2, \dots, m\}$, $w \in \mathbb{R}^d$ with $i \neq j$ and $\|z_i - w\| \leq R_i$ it holds that

$$|\langle z_i, x_j \rangle - \langle w, x_j \rangle| = |\langle z_i - w, x_j \rangle| \leq \|z_i - w\| \|x_j\| \leq R_i \|x_j\| \leq \frac{|\langle z_i, x_j \rangle|}{2 \|x_j\|} \|x_j\| = \frac{1}{2} |\langle z_i, x_j \rangle|. \quad (24)$$

This and (19) therefore establish that for all $i, j \in \{1, 2, \dots, m\}$, $w \in \mathbb{R}^d$ with $i \neq j$ and $\|z_i - w\| \leq R_i$ it holds that

$$g_{x_j}(w) = \mathbb{1}_{[0, \infty)}(\langle w, x_j \rangle) = \mathbb{1}_{[0, \infty)}(\langle z_i, x_j \rangle) = g_{x_j}(z_i) \quad (25)$$

and

$$h_{x_j}(w) = x_j \mathbb{1}_{[0, \infty)}(\langle w, x_j \rangle) = x_j \mathbb{1}_{[0, \infty)}(\langle z_i, x_j \rangle) = h_{x_j}(z_i). \quad (26)$$

Combining this and (18) ensures that for all $i, j \in \{1, 2, \dots, m\}$ with $i \neq j$ it holds that

$$\begin{aligned} & \frac{1}{\mathbb{P}_W(A_{R_i}^i(z_i))} \int_{A_{R_i}^i(z_i)} g_{x_j}(w) d\mathbb{P}_W(w) - \frac{1}{\mathbb{P}_W(B_{R_i}^i(z_i))} \int_{B_{R_i}^i(z_i)} g_{x_j}(w) d\mathbb{P}_W(w) \\ &= \frac{1}{\mathbb{P}_W(A_{R_i}^i(z_i))} \int_{A_{R_i}^i(z_i)} g_{x_j}(z_i) d\mathbb{P}_W(w) - \frac{1}{\mathbb{P}_W(B_{R_i}^i(z_i))} \int_{B_{R_i}^i(z_i)} g_{x_j}(z_i) d\mathbb{P}_W(w) \\ &= g_{x_j}(z_i) - g_{x_j}(z_i) = 0 \end{aligned} \quad (27)$$

and

$$\begin{aligned} & \frac{1}{\mathbb{P}_W(A_{R_i}^i(z_i))} \int_{A_{R_i}^i(z_i)} h_{x_j}(w) d\mathbb{P}_W(w) - \frac{1}{\mathbb{P}_W(B_{R_i}^i(z_i))} \int_{B_{R_i}^i(z_i)} h_{x_j}(w) d\mathbb{P}_W(w) \\ &= \frac{1}{\mathbb{P}_W(A_{R_i}^i(z_i))} \int_{A_{R_i}^i(z_i)} h_{x_j}(z_i) d\mathbb{P}_W(w) - \frac{1}{\mathbb{P}_W(B_{R_i}^i(z_i))} \int_{B_{R_i}^i(z_i)} h_{x_j}(z_i) d\mathbb{P}_W(w) \\ &= h_{x_j}(z_i) - h_{x_j}(z_i) = 0. \end{aligned} \quad (28)$$

Moreover, note that (18) implies that for all $i \in \{1, 2, \dots, m\}$ it holds that

$$\begin{aligned} & \frac{1}{\mathbb{P}_W(A_{R_i}^i(z_i))} \int_{A_{R_i}^i(z_i)} g_{x_i}(w) d\mathbb{P}_W(w) - \frac{1}{\mathbb{P}_W(B_{R_i}^i(z_i))} \int_{B_{R_i}^i(z_i)} g_{x_i}(w) d\mathbb{P}_W(w) \\ &= \frac{1}{\mathbb{P}_W(A_{R_i}^i(z_i))} \int_{A_{R_i}^i(z_i)} 1 d\mathbb{P}_W(w) - \frac{1}{\mathbb{P}_W(B_{R_i}^i(z_i))} \int_{B_{R_i}^i(z_i)} 0 d\mathbb{P}_W(w) = 1 \end{aligned} \quad (29)$$

and

$$\begin{aligned} & \frac{1}{\mathbb{P}_W(A_{R_i}^i(z_i))} \int_{A_{R_i}^i(z_i)} h_{x_i}(w) d\mathbb{P}_W(w) - \frac{1}{\mathbb{P}_W(B_{R_i}^i(z_i))} \int_{B_{R_i}^i(z_i)} h_{x_i}(w) d\mathbb{P}_W(w) \\ &= \frac{1}{\mathbb{P}_W(A_{R_i}^i(z_i))} \int_{A_{R_i}^i(z_i)} x_i d\mathbb{P}_W(w) - \frac{1}{\mathbb{P}_W(B_{R_i}^i(z_i))} \int_{B_{R_i}^i(z_i)} 0 d\mathbb{P}_W(w) = x_i. \end{aligned} \quad (30)$$

Combining this, (27), and (28) with the assumption that $\sum_{j=1}^m \alpha_j g_{x_j} = 0 \in \mathcal{G}$ and $\sum_{j=1}^m \beta_j h_{x_j} = 0 \in \mathcal{H}$ establishes that for all $i \in \{1, 2, \dots, m\}$ it holds that

$$\begin{aligned} 0 &= \frac{1}{\mathbb{P}_W(A_{R_i}^i(z_i))} \int_{A_{R_i}^i(z_i)} \sum_{j=1}^m \alpha_j g_{x_j}(w) d\mathbb{P}_W(w) - \frac{1}{\mathbb{P}_W(B_{R_i}^i(z_i))} \int_{B_{R_i}^i(z_i)} \sum_{j=1}^m \alpha_j g_{x_j}(w) d\mathbb{P}_W(w) \\ &= \sum_{j=1}^m \alpha_j \left(\frac{1}{\mathbb{P}_W(A_{R_i}^i(z_i))} \int_{A_{R_i}^i(z_i)} g_{x_j}(w) d\mathbb{P}_W(w) - \frac{1}{\mathbb{P}_W(B_{R_i}^i(z_i))} \int_{B_{R_i}^i(z_i)} g_{x_j}(w) d\mathbb{P}_W(w) \right) = \alpha_i \end{aligned} \quad (31)$$

and

$$\begin{aligned} 0 &= \frac{1}{\mathbb{P}_W(A_{R_i}^i(z_i))} \int_{A_{R_i}^i(z_i)} \sum_{j=1}^m \beta_j h_{x_j}(w) d\mathbb{P}_W(w) - \frac{1}{\mathbb{P}_W(B_{R_i}^i(z_i))} \int_{B_{R_i}^i(z_i)} \sum_{j=1}^m \beta_j h_{x_j}(w) d\mathbb{P}_W(w) \\ &= \sum_{j=1}^m \beta_j \left(\frac{1}{\mathbb{P}_W(A_{R_i}^i(z_i))} \int_{A_{R_i}^i(z_i)} h_{x_j}(w) d\mathbb{P}_W(w) - \frac{1}{\mathbb{P}_W(B_{R_i}^i(z_i))} \int_{B_{R_i}^i(z_i)} h_{x_j}(w) d\mathbb{P}_W(w) \right) = \beta_i x_i. \end{aligned} \quad (32)$$

This and the assumption that for all $i \in \{1, 2, \dots, m\}$ it holds that $x_i \neq 0$ ensure that for all $i \in \{1, 2, \dots, m\}$ it holds that $\alpha_i = \beta_i = 0$. Hence, we obtain that $g_{x_1}, g_{x_2}, \dots, g_{x_m}$ are linearly independent and $h_{x_1}, h_{x_2}, \dots, h_{x_m}$ are linearly independent. Item (ii) in Lemma 2.9 (applied with $V \curvearrowright \mathcal{G}$, $\varphi \curvearrowright \varphi$, $n \curvearrowright m$, $(v_i)_{i \in \{1, 2, \dots, n\}} \curvearrowright (g_{x_i})_{i \in \{1, 2, \dots, m\}}$, $G \curvearrowright \mathbf{G}$ in the notation of Lemma 2.9) and item (ii) in Lemma 2.9 (applied with $V \curvearrowright \mathcal{H}$, $\varphi \curvearrowright \psi$, $n \curvearrowright m$, $(v_i)_{i \in \{1, 2, \dots, n\}} \curvearrowright (h_{x_i})_{i \in \{1, 2, \dots, m\}}$, $G \curvearrowright \mathbf{H}$ in the notation of Lemma 2.9) therefore establish that \mathbf{G} and \mathbf{H} are positive definite. This and the assumption that it holds that $\boldsymbol{\lambda} = \lambda_{\min}(\mathbf{G})$ and $\boldsymbol{\mu} = \lambda_{\min}(\mathbf{H})$ ensure that it holds that $\boldsymbol{\lambda} > 0$ and $\boldsymbol{\mu} > 0$. The proof of Lemma 2.11 is thus completed. \square

3 Analysis of eigenvalues of stochastic Gramian matrices

In this section we study the error at initialization (cf. Section 3.1 below), the evolution of the weights and biases of the considered ANNs during training (cf. Section 3.2 below), and the eigenvalues of the considered stochastic Gramian matrices (cf. Section 3.6 below). In particular, the main result of Section 3.6, Lemma 3.22 below, establishes a lower bound for the eigenvalues of the matrices $\mathcal{G}(n)$, $n \in \mathbb{N}_0$, and $\mathcal{H}(n)$, $n \in \mathbb{N}_0$, from Setting 2.7 in Section 2.3 above. In order to achieve these estimates, we analyze the distance from \mathcal{G} (resp. \mathcal{H}) at initialization to its deterministic counterpart (cf. Section 3.4 below), as well as the distance from \mathcal{G} (resp. \mathcal{H}) during training to its initialization (cf. Section 3.5 below) with respect to the spectral norm. These estimates will be used in our error analysis for GD optimization algorithms in Section 4 below. We also note that the results in Lemma 3.3, Lemma 3.4, Lemma 3.14, Lemma 3.17, Lemma 3.20, and Lemma 3.21 are all well-known and we observe that the results in Lemma 3.8, Lemma 3.9, and Lemma 3.12 are all elementary. Only for completeness we include in this section the detailed proofs for these lemmas.

3.1 Probabilistic error analysis at initialization

Lemma 3.1. *Let $(\Omega, \mathcal{F}, \mathbb{P})$ be a probability space, let $\sigma \in \mathbb{R}$, and let $X : \Omega \rightarrow \mathbb{R}$ be a normal random variable with $\text{Var}[X] = \sigma^2$. Then $\mathbb{E}[|\mathfrak{r}(X - \mathbb{E}[X])|^2] = \sigma^2/2$ (cf. Definition 2.3).*

Proof of Lemma 3.1. Observe that the assumption that $\text{Var}[X] = \sigma^2$ and the fact that $X - \mathbb{E}[X]$ and $\mathbb{E}[X] - X$ are identically distributed establish that

$$\begin{aligned}\sigma^2 &= \mathbb{E}[|X - \mathbb{E}[X]|^2] = \mathbb{E}[|X - \mathbb{E}[X]|^2(\mathbb{1}_{\{X - \mathbb{E}[X] \geq 0\}} + \mathbb{1}_{\{X - \mathbb{E}[X] \leq 0\}})] \\ &= \mathbb{E}[|X - \mathbb{E}[X]|^2 \mathbb{1}_{\{X - \mathbb{E}[X] \geq 0\}}] + \mathbb{E}[|X - \mathbb{E}[X]|^2 \mathbb{1}_{\{X - \mathbb{E}[X] \leq 0\}}] \\ &= \mathbb{E}[|X - \mathbb{E}[X]|^2 \mathbb{1}_{\{X - \mathbb{E}[X] \geq 0\}}] + \mathbb{E}[|\mathbb{E}[X] - X|^2 \mathbb{1}_{\{\mathbb{E}[X] - X \geq 0\}}] = 2 \mathbb{E}[|X - \mathbb{E}[X]|^2 \mathbb{1}_{\{X - \mathbb{E}[X] \geq 0\}}].\end{aligned}\quad (33)$$

Hence, we obtain that

$$\mathbb{E}[|\mathfrak{r}(X - \mathbb{E}[X])|^2] = \mathbb{E}[|(X - \mathbb{E}[X]) \mathbb{1}_{\{X - \mathbb{E}[X] \geq 0\}}|^2] = \mathbb{E}[|X - \mathbb{E}[X]|^2 \mathbb{1}_{\{X - \mathbb{E}[X] \geq 0\}}] = \frac{\sigma^2}{2} \quad (34)$$

(cf. Definition 2.3). This completes the proof of Lemma 3.1. \square

Lemma 3.2. *Assume Setting 2.7 and let $\varepsilon \in (0, 1)$. Then $\mathbb{P}(\|f(0) - y\|^2 \leq (\frac{1}{2} \sum_{i=1}^m \|x_i\|^2 + \|y\|^2) \varepsilon^{-1}) \geq 1 - \varepsilon$ (cf. Definition 2.1).*

Proof of Lemma 3.2. Note that the assumption that $\|B(0)\| = |\mathfrak{B}(0)| = 0$ ensures that for all $i \in \{1, 2, \dots, m\}$ it holds that

$$f_i(0) = (\mathcal{R}(\Phi(0)))(x_i) = \langle \mathcal{W}, \mathfrak{r}(W(0)x_i + B(0)) \rangle + \mathfrak{B}(0) = \langle \mathcal{W}, \mathfrak{r}(W(0)x_i) \rangle = \sum_{k=1}^{\mathfrak{d}} \mathcal{W}_k \mathfrak{r}(\langle W_k(0), x_i \rangle) \quad (35)$$

(cf. Definitions 2.1, 2.3, and 2.4). Combining this with the assumption that $W_1(0), W_2(0), \dots, W_{\mathfrak{d}}(0), \sqrt{\mathfrak{d}}\mathcal{W}_1, \sqrt{\mathfrak{d}}\mathcal{W}_2, \dots, \sqrt{\mathfrak{d}}\mathcal{W}_{\mathfrak{d}}$ are independent and standard normal demonstrates that

$$\mathbb{E}[f_i(0)] = \sum_{k=1}^{\mathfrak{d}} \mathbb{E}[\mathcal{W}_k \mathfrak{r}(\langle W_k(0), x_i \rangle)] = \sum_{k=1}^{\mathfrak{d}} \mathbb{E}[\mathcal{W}_k] \mathbb{E}[\mathfrak{r}(\langle W_k(0), x_i \rangle)] = 0. \quad (36)$$

Moreover, note that (35) ensures that for all $i \in \{1, 2, \dots, m\}$ it holds that

$$\begin{aligned}|f_i(0)|^2 &= \left| \sum_{k=1}^{\mathfrak{d}} \mathcal{W}_k \mathfrak{r}(\langle W_k(0), x_i \rangle) \right|^2 = \sum_{k=1}^{\mathfrak{d}} \sum_{\ell=1}^{\mathfrak{d}} \mathcal{W}_k \mathfrak{r}(\langle W_k(0), x_i \rangle) \mathcal{W}_{\ell} \mathfrak{r}(\langle W_{\ell}(0), x_i \rangle) \\ &= \sum_{k=1}^{\mathfrak{d}} |\mathcal{W}_k|^2 |\mathfrak{r}(\langle W_k(0), x_i \rangle)|^2 + \sum_{\substack{k, \ell \in \{1, 2, \dots, \mathfrak{d}\}, \\ k \neq \ell}} \mathcal{W}_k \mathcal{W}_{\ell} \mathfrak{r}(\langle W_k(0), x_i \rangle) \mathfrak{r}(\langle W_{\ell}(0), x_i \rangle).\end{aligned}\quad (37)$$

In addition, observe that the assumption that $W_1(0), W_2(0), \dots, W_{\mathfrak{d}}(0)$ are standard normal ensures that for all $k \in \{1, 2, \dots, \mathfrak{d}\}$, $i \in \{1, 2, \dots, m\}$ it holds that $\langle W_k(0), x_i \rangle$ is a centered normal random variable with $\text{Var}[\langle W_k(0), x_i \rangle] = \|x_i\|^2$. Combining this with Lemma 3.1 (applied for every $k \in \{1, 2, \dots, \mathfrak{d}\}$, $i \in \{1, 2, \dots, m\}$ with $(\Omega, \mathcal{F}, \mathbb{P}) \curvearrowright (\Omega, \mathcal{F}, \mathbb{P})$, $X \curvearrowright \langle W_k(0), x_i \rangle$, $\sigma \curvearrowright \|x_i\|$ in the notation of Lemma 3.1), the assumption that $W_1(0), W_2(0), \dots, W_{\mathfrak{d}}(0), \sqrt{\mathfrak{d}}\mathcal{W}_1, \sqrt{\mathfrak{d}}\mathcal{W}_2, \dots, \sqrt{\mathfrak{d}}\mathcal{W}_{\mathfrak{d}}$ are independent and standard normal, and (37) establishes that for all $i \in \{1, 2, \dots, m\}$ it holds that

$$\begin{aligned}\mathbb{E}[|f_i(0)|^2] &= \sum_{k=1}^{\mathfrak{d}} \mathbb{E}[|\mathcal{W}_k|^2 |\mathfrak{r}(\langle W_k(0), x_i \rangle)|^2] + \sum_{\substack{k, \ell \in \{1, 2, \dots, \mathfrak{d}\}, \\ k \neq \ell}} \mathbb{E}[\mathcal{W}_k \mathcal{W}_{\ell} \mathfrak{r}(\langle W_k(0), x_i \rangle) \mathfrak{r}(\langle W_{\ell}(0), x_i \rangle)] \\ &= \sum_{k=1}^{\mathfrak{d}} \mathbb{E}[|\mathcal{W}_k|^2] \mathbb{E}[|\mathfrak{r}(\langle W_k(0), x_i \rangle)|^2] + \sum_{\substack{k, \ell \in \{1, 2, \dots, \mathfrak{d}\}, \\ k \neq \ell}} \mathbb{E}[\mathcal{W}_k] \mathbb{E}[\mathcal{W}_{\ell} \mathfrak{r}(\langle W_k(0), x_i \rangle) \mathfrak{r}(\langle W_{\ell}(0), x_i \rangle)] \\ &= \sum_{k=1}^{\mathfrak{d}} \frac{1}{2\mathfrak{d}} \|x_i\|^2 = \frac{1}{2} \|x_i\|^2.\end{aligned}\quad (38)$$

This and (36) therefore demonstrate that

$$\begin{aligned}\mathbb{E}[\|f(0) - y\|^2] &= \mathbb{E}\left[\sum_{i=1}^m |f_i(0) - y_i|^2\right] = \sum_{i=1}^m \mathbb{E}[|f_i(0) - y_i|^2] = \sum_{i=1}^m \mathbb{E}[|f_i(0)|^2 - 2f_i(0)y_i + |y_i|^2] \\ &= \sum_{i=1}^m (\mathbb{E}[|f_i(0)|^2] - 2y_i \mathbb{E}[f_i(0)] + |y_i|^2) = \sum_{i=1}^m (\frac{1}{2} \|x_i\|^2 + |y_i|^2) = \frac{1}{2} \sum_{i=1}^m \|x_i\|^2 + \|y\|^2.\end{aligned}\quad (39)$$

The Markov inequality thus establishes that

$$\begin{aligned} \mathbb{P}\left(\|f(0) - y\|^2 \leq \left(\frac{1}{2} \sum_{i=1}^m \|x_i\|^2 + \|y\|^2\right)\varepsilon^{-1}\right) &\geq 1 - \mathbb{P}\left(\|f(0) - y\|^2 \geq \left(\frac{1}{2} \sum_{i=1}^m \|x_i\|^2 + \|y\|^2\right)\varepsilon^{-1}\right) \\ &\geq 1 - \frac{\mathbb{E}[\|f(0) - y\|^2]}{\left(\frac{1}{2} \sum_{i=1}^m \|x_i\|^2 + \|y\|^2\right)\varepsilon^{-1}} = 1 - \varepsilon. \end{aligned} \quad (40)$$

This completes the proof of Lemma 3.2. \square

3.2 Analysis of weights and biases of ANNs during training

Lemma 3.3. *Let $n \in \mathbb{N}$, $x = (x_1, x_2, \dots, x_n) \in \mathbb{R}^n$. Then $\|x\| \leq \sum_{i=1}^n |x_i| \leq \sqrt{n}\|x\|$ (cf. Definition 2.1).*

Proof of Lemma 3.3. Throughout this proof let $x = (x_1, x_2, \dots, x_n)$, $y = (y_1, y_2, \dots, y_n) \in \mathbb{R}^n$ satisfy for all $i \in \{1, 2, \dots, n\}$ that $x_i = |x_i|$ and $y_i = 1$. Note that the fact that for all $a, b \in [0, \infty)$ it holds that $(a + b)^{1/2} \leq a^{1/2} + b^{1/2}$ inductively ensures that $\|x\| = (\sum_{i=1}^n |x_i|^2)^{1/2} \leq \sum_{i=1}^n (|x_i|^2)^{1/2} = \sum_{i=1}^n |x_i|$ (cf. Definition 2.1). Moreover, observe that the Cauchy Schwarz inequality establishes that $\sum_{i=1}^n |x_i| = \langle x, y \rangle \leq \|x\| \|y\| = (\sum_{i=1}^n |x_i|^2)^{1/2} (\sum_{i=1}^n 1^2)^{1/2} = \sqrt{n}\|x\|$. This completes the proof of Lemma 3.3. \square

Lemma 3.4. *Let $(\Omega, \mathcal{F}, \mathbb{P})$ be a probability space, let $X: \Omega \rightarrow \mathbb{R}$ be a standard normal random variable, and let $\varepsilon \in (0, \infty)$. Then $\mathbb{P}(|X| \geq \varepsilon) \leq 2 \exp(-\varepsilon^2/2)$.*

Proof of Lemma 3.4. Throughout this proof let $f: (0, \infty) \rightarrow \mathbb{R}$ satisfy for all $\lambda \in (0, \infty)$ that $f(\lambda) = \frac{1}{2}\lambda^2 - \lambda\varepsilon$. Observe that the assumption that X is a standard normal random variable ensures that for all $\lambda \in \mathbb{R}$ it holds that

$$\begin{aligned} \mathbb{E}[\exp(\lambda X)] &= \int_{-\infty}^{\infty} \frac{1}{\sqrt{2\pi}} \exp\left(-\frac{x^2}{2}\right) \exp(\lambda x) dx = \int_{-\infty}^{\infty} \frac{1}{\sqrt{2\pi}} \exp\left(-\frac{x^2 - 2\lambda x}{2}\right) dx \\ &= \exp\left(\frac{\lambda^2}{2}\right) \int_{-\infty}^{\infty} \frac{1}{\sqrt{2\pi}} \exp\left(-\frac{(x - \lambda)^2}{2}\right) dx = \exp\left(\frac{\lambda^2}{2}\right) < \infty. \end{aligned} \quad (41)$$

This and the Markov inequality ensure that for all $\lambda \in (0, \infty)$ it holds that

$$\begin{aligned} \mathbb{P}(|X| \geq \varepsilon) &= \mathbb{P}(X \geq \varepsilon) + \mathbb{P}(-X \geq \varepsilon) = \mathbb{P}(\exp(\lambda X) \geq \exp(\lambda\varepsilon)) + \mathbb{P}(\exp(-\lambda X) \geq \exp(\lambda\varepsilon)) \\ &\leq \frac{\mathbb{E}[\exp(\lambda X)]}{\exp(\lambda\varepsilon)} + \frac{\mathbb{E}[\exp(-\lambda X)]}{\exp(\lambda\varepsilon)} = \exp\left(\frac{\lambda^2}{2} - \lambda\varepsilon\right) + \exp\left(\frac{(-\lambda)^2}{2} - \lambda\varepsilon\right) \\ &= 2 \exp\left(\frac{\lambda^2}{2} - \lambda\varepsilon\right) = 2 \exp(f(\lambda)). \end{aligned} \quad (42)$$

Combining this with the fact that $\mathbb{R} \ni x \mapsto \exp(x) \in \mathbb{R}$ is strictly increasing establishes that

$$\mathbb{P}(|X| \geq \varepsilon) \leq \inf_{\lambda \in (0, \infty)} 2 \exp(f(\lambda)) = 2 \exp(\inf_{\lambda \in (0, \infty)} f(\lambda)). \quad (43)$$

Moreover, note that for all $\lambda \in (0, \infty)$ it holds that

$$f(\lambda) = \frac{\lambda^2}{2} - \lambda\varepsilon = \frac{1}{2}[\lambda^2 - 2\lambda\varepsilon] = \frac{1}{2}[(\lambda - \varepsilon)^2 - \varepsilon^2] = \frac{1}{2}(\lambda - \varepsilon)^2 - \frac{1}{2}\varepsilon^2 \geq -\frac{1}{2}\varepsilon^2. \quad (44)$$

This and the fact that $f(\varepsilon) = -\varepsilon^2/2$ ensure that it holds that $\inf_{\lambda \in (0, \infty)} f(\lambda) = -\varepsilon^2/2$. Combining this with (43) demonstrates that $\mathbb{P}(|X| \geq \varepsilon) \leq 2 \exp(\inf_{\lambda \in (0, \infty)} f(\lambda)) = 2 \exp(-\varepsilon^2/2)$. The proof of Lemma 3.4 is thus completed. \square

Lemma 3.5. *Assume Setting 2.7 and let $\varepsilon \in (0, 1)$. Then $\mathbb{P}(\bigcap_{k=1}^{\mathfrak{d}} \{|\mathcal{W}_k| \leq (\frac{2\mathfrak{d}}{\varepsilon})^{1/2}\}) \geq 1 - \varepsilon$.*

Proof of Lemma 3.5. Note that the assumption that $\sqrt{\mathfrak{d}}\mathcal{W}_1, \sqrt{\mathfrak{d}}\mathcal{W}_2, \dots, \sqrt{\mathfrak{d}}\mathcal{W}_{\mathfrak{d}}$ are standard normal random variables and Lemma 3.4 (applied for every $k \in \{1, 2, \dots, \mathfrak{d}\}$ with $(\Omega, \mathcal{F}, \mathbb{P}) \curvearrowright (\Omega, \mathcal{F}, \mathbb{P})$, $X \curvearrowright \sqrt{\mathfrak{d}}\mathcal{W}_k$,

$\varepsilon \curvearrowright (2 \ln(\frac{2\mathfrak{d}}{\varepsilon}))^{1/2}$ in the notation of Lemma 3.4) prove that

$$\begin{aligned} & \mathbb{P}\left(\bigcap_{k=1}^{\mathfrak{d}} \left\{ |\mathcal{W}_k| \leq \left(\frac{2}{\mathfrak{d}} \ln\left(\frac{2\mathfrak{d}}{\varepsilon}\right)\right)^{1/2} \right\}\right) \geq 1 - \mathbb{P}\left(\bigcup_{k=1}^{\mathfrak{d}} \left\{ |\mathcal{W}_k| \geq \left(\frac{2}{\mathfrak{d}} \ln\left(\frac{2\mathfrak{d}}{\varepsilon}\right)\right)^{1/2} \right\}\right) \\ & \geq 1 - \sum_{k=1}^{\mathfrak{d}} \mathbb{P}\left(|\mathcal{W}_k| \geq \left(\frac{2}{\mathfrak{d}} \ln\left(\frac{2\mathfrak{d}}{\varepsilon}\right)\right)^{1/2}\right) = 1 - \sum_{k=1}^{\mathfrak{d}} \mathbb{P}\left(|\sqrt{\mathfrak{d}}\mathcal{W}_k| \geq (2 \ln(\frac{2\mathfrak{d}}{\varepsilon}))^{1/2}\right) \\ & \geq 1 - \sum_{k=1}^{\mathfrak{d}} 2 \exp\left(-\frac{1}{2}(2 \ln(\frac{2\mathfrak{d}}{\varepsilon}))\right) = 1 - \sum_{k=1}^{\mathfrak{d}} 2 \exp(\ln(\frac{\varepsilon}{2\mathfrak{d}})) = 1 - \sum_{k=1}^{\mathfrak{d}} \frac{\varepsilon}{\mathfrak{d}} = 1 - \varepsilon. \end{aligned} \quad (45)$$

This completes the proof of Lemma 3.5. \square

Lemma 3.6. *Assume Setting 2.7, assume $\lambda + \mu \in (0, \infty)$, assume $\eta < \frac{m}{\lambda + \mu}$, let $\varepsilon \in (0, 1)$, $A \in \mathcal{F}$ satisfy $A = \bigcap_{k=1}^{\mathfrak{d}} \{|\mathcal{W}_k| \leq (\frac{2}{\mathfrak{d}} \ln(\frac{2\mathfrak{d}}{\varepsilon}))^{1/2}\}$, let $\omega \in A$, $N \in \mathbb{N}_0$, and assume for all $n \in \{0, 1, 2, \dots, N\}$, $k \in \{1, 2, \dots, \mathfrak{d}\}$ that $\|f(n, \omega) - y\|^2 \leq (1 - \frac{\eta(\lambda + \mu)}{m})^n \|f(0, \omega) - y\|^2$ (cf. Definition 2.1). Then*

- (i) *it holds for all $k \in \{1, 2, \dots, \mathfrak{d}\}$ that $\|W_k(N+1, \omega) - W_k(0, \omega)\| \leq \frac{4C\|f(0, \omega) - y\|}{\lambda + \mu} (\frac{2m}{\mathfrak{d}} \ln(\frac{2\mathfrak{d}}{\varepsilon}))^{1/2}$, and*
- (ii) *it holds for all $k \in \{1, 2, \dots, \mathfrak{d}\}$ that $|B_k(N+1, \omega) - B_k(0, \omega)| \leq \frac{4\|f(0, \omega) - y\|}{\lambda + \mu} (\frac{2m}{\mathfrak{d}} \ln(\frac{2\mathfrak{d}}{\varepsilon}))^{1/2}$.*

Proof of Lemma 3.6. Observe that for all $q \in (0, 1)$ it holds that $(1 - q)^{1/2} \leq 1 - \frac{q}{2}$ and

$$\sum_{n=0}^N (1 - q)^{n/2} \leq \sum_{n=0}^{\infty} (1 - q)^{n/2} = \frac{1}{1 - (1 - q)^{1/2}} \leq \frac{1}{1 - (1 - \frac{q}{2})} = \frac{2}{q}. \quad (46)$$

This, the fact that it holds that $\frac{1}{m}\eta(\lambda + \mu) \in (0, 1)$, Lemma 3.3 (applied for every $n \in \{0, 1, 2, \dots, N\}$ with $n \curvearrowright m$, $x \curvearrowright f(n, \omega) - y$ in the notation of Lemma 3.3), the assumption that for all $n \in \{0, 1, 2, \dots, N\}$ it holds that $\|f(n, \omega) - y\|^2 \leq (1 - \frac{\eta(\lambda + \mu)}{m})^n \|f(0, \omega) - y\|^2$, and the assumption that $\omega \in A$ ensure that it holds for all $k \in \{1, 2, \dots, \mathfrak{d}\}$ that

$$\begin{aligned} & |\mathcal{W}_k(\omega)| \sum_{n=0}^N \sum_{j=1}^m |f_j(n, \omega) - y_j| \leq |\mathcal{W}_k(\omega)| \sum_{n=0}^N \sqrt{m} \|f(n, \omega) - y\| = |\mathcal{W}_k(\omega)| \sqrt{m} \sum_{n=0}^N (\|f(n, \omega) - y\|^2)^{1/2} \\ & \leq |\mathcal{W}_k(\omega)| \sqrt{m} \sum_{n=0}^N \left(1 - \frac{\eta(\lambda + \mu)}{m}\right)^{n/2} \|f(0, \omega) - y\| = |\mathcal{W}_k(\omega)| \sqrt{m} \|f(0, \omega) - y\| \sum_{n=0}^N \left(1 - \frac{\eta(\lambda + \mu)}{m}\right)^{n/2} \\ & \leq |\mathcal{W}_k(\omega)| \sqrt{m} \|f(0, \omega) - y\| \frac{2m}{\eta(\lambda + \mu)} = |\mathcal{W}_k(\omega)| \frac{2m^{3/2} \|f(0, \omega) - y\|}{\eta(\lambda + \mu)} \leq \left(\frac{2}{\mathfrak{d}} \ln\left(\frac{2\mathfrak{d}}{\varepsilon}\right)\right)^{1/2} \frac{2m^{3/2} \|f(0, \omega) - y\|}{\eta(\lambda + \mu)}. \end{aligned} \quad (47)$$

Combining this with (8) establishes for all $k \in \{1, 2, \dots, \mathfrak{d}\}$ that

$$\begin{aligned} & \|W_k(N+1, \omega) - W_k(0, \omega)\| = \left\| \sum_{n=0}^N [W_k(n+1, \omega) - W_k(n, \omega)] \right\| \\ & = \left\| \sum_{n=0}^N -\frac{2\eta}{m} \sum_{j=1}^m (f_j(n, \omega) - y_j) \mathcal{W}_k(\omega) \mathbb{1}_{\{(W_k(n), x_j) + B_k(n) \geq 0\}}(\omega) x_j \right\| \\ & \leq \frac{2\eta}{m} \sum_{n=0}^N \sum_{j=1}^m |(f_j(n, \omega) - y_j) \mathcal{W}_k(\omega) \mathbb{1}_{\{(W_k(n), x_j) + B_k(n) \geq 0\}}(\omega)| \|x_j\| \\ & \leq \frac{2\eta C}{m} |\mathcal{W}_k(\omega)| \sum_{n=0}^N \sum_{j=1}^m |f_j(n, \omega) - y_j| \leq \frac{4C\|f(0, \omega) - y\|}{\lambda + \mu} \left(\frac{2m}{\mathfrak{d}} \ln\left(\frac{2\mathfrak{d}}{\varepsilon}\right)\right)^{1/2}. \end{aligned} \quad (48)$$

This establishes item (i). Next note that (9) and (47) demonstrate that

$$\begin{aligned} & |B_k(N+1, \omega) - B_k(0, \omega)| = \left| \sum_{n=0}^N [B_k(n+1, \omega) - B_k(n, \omega)] \right| \\ & = \left| \sum_{n=0}^N -\frac{2\eta}{m} \sum_{j=1}^m (f_j(n, \omega) - y_j) \mathcal{W}_k(\omega) \mathbb{1}_{\{(W_k(n), x_j) + B_k(n) \geq 0\}}(\omega) \right| \\ & \leq \frac{2\eta}{m} \sum_{n=0}^N \sum_{j=1}^m |(f_j(n, \omega) - y_j) \mathcal{W}_k(\omega) \mathbb{1}_{\{(W_k(n), x_j) + B_k(n) \geq 0\}}(\omega)| \\ & \leq \frac{2\eta}{m} |\mathcal{W}_k(\omega)| \sum_{n=0}^N \sum_{j=1}^m |f_j(n, \omega) - y_j| \leq \frac{4\|f(0, \omega) - y\|}{\lambda + \mu} \left(\frac{2m}{\mathfrak{d}} \ln\left(\frac{2\mathfrak{d}}{\varepsilon}\right)\right)^{1/2}. \end{aligned} \quad (49)$$

This establishes item (ii). The proof of Lemma 3.6 is thus completed. \square

3.3 Properties of subexponential random variables

Definition 3.7 (Subexponential random variable). Let $(\Omega, \mathcal{F}, \mathbb{P})$ be a probability space and let $\nu, b \in (0, \infty)$. Then we say that X is (ν, b) -subexponential with respect to \mathbb{P} (we say that X is (ν, b) -subexponential) if and only if

- (i) it holds that $X: \Omega \rightarrow \mathbb{R}$ is a function from Ω to \mathbb{R} ,
- (ii) it holds that X is measurable,
- (iii) it holds that $\mathbb{E}[|X|] < \infty$, and
- (iv) it holds for all $\lambda \in (-1/b, 1/b)$ that $\mathbb{E}[\exp(\lambda(X - \mathbb{E}[X]))] \leq \exp(\frac{1}{2}\lambda^2\nu^2)$.

Lemma 3.8. Let $(\Omega, \mathcal{F}, \mathbb{P})$ be a probability space, let $\nu, b, a \in (0, \infty)$, and let X be (ν, b) -subexponential (cf. Definition 3.7). Then aX is $(a\nu, ab)$ -subexponential.

Proof of Lemma 3.8. Note that for all $\lambda \in (-\frac{1}{ab}, \frac{1}{ab})$ it holds that $a\lambda \in (-1/b, 1/b)$. Hence, we obtain for all $\lambda \in (-\frac{1}{ab}, \frac{1}{ab})$ that $\mathbb{E}[\exp(\lambda(aX - \mathbb{E}[aX]))] = \mathbb{E}[\exp(a\lambda(X - \mathbb{E}[X]))] \leq \exp(\frac{1}{2}(a\lambda)^2\nu^2) = \exp(\frac{1}{2}\lambda^2(a\nu)^2)$. This ensures that aX is $(a\nu, ab)$ -subexponential. The proof of Lemma 3.8 is thus completed. \square

Lemma 3.9. Let $(\Omega, \mathcal{F}, \mathbb{P})$ be a probability space, let $n \in \mathbb{N}$, $\nu = (\nu_1, \nu_2, \dots, \nu_n)$, $b = (b_1, b_2, \dots, b_n) \in (0, \infty)^n$, let $X_i: \Omega \rightarrow \mathbb{R}$, $i \in \{1, 2, \dots, n\}$, be independent random variables, and assume for all $i \in \{1, 2, \dots, n\}$ that X_i is (ν_i, b_i) -subexponential (cf. Definition 3.7). Then $\sum_{i=1}^n X_i$ is $(\|\nu\|, \max\{b_1, b_2, \dots, b_n\})$ -subexponential (cf. Definition 2.1).

Proof of Lemma 3.9. Throughout this proof let $B \in (0, \infty)$ satisfy $B = \max\{b_1, b_2, \dots, b_n\}$. The fact that X_1, X_2, \dots, X_n are independent and the fact that for all $i \in \{1, 2, \dots, n\}$ it holds that $(-1/B, 1/B) \subseteq (-1/b_i, 1/b_i)$ ensure that for all $\lambda \in (-1/B, 1/B)$ it holds that

$$\begin{aligned} \mathbb{E}\left[\exp\left(\lambda\left(\sum_{i=1}^n X_i - \mathbb{E}\left[\sum_{i=1}^n X_i\right]\right)\right)\right] &= \mathbb{E}\left[\exp\left(\lambda\sum_{i=1}^n (X_i - \mathbb{E}[X_i])\right)\right] = \mathbb{E}\left[\prod_{i=1}^n \exp(\lambda(X_i - \mathbb{E}[X_i]))\right] \\ &= \prod_{i=1}^n \mathbb{E}\left[\exp(\lambda(X_i - \mathbb{E}[X_i]))\right] \leq \prod_{i=1}^n \exp\left(\frac{\lambda^2|\nu_i|^2}{2}\right) \\ &= \exp\left(\frac{\lambda^2\sum_{i=1}^n |\nu_i|^2}{2}\right) = \exp\left(\frac{\lambda^2\|\nu\|^2}{2}\right) \end{aligned} \quad (50)$$

(cf. Definition 2.1). This establishes that $\sum_{i=1}^n X_i$ is $(\|\nu\|, B)$ -subexponential. The proof of Lemma 3.9 is thus completed. \square

Lemma 3.10. Let $(\Omega, \mathcal{F}, \mathbb{P})$ be a probability space, let $X: \Omega \rightarrow \mathbb{R}$ be a standard normal random variable, let $A \in \mathcal{F}$, and assume that X and $\mathbb{1}_A$ are independent. Then $X^2\mathbb{1}_A$ is $(2, 4)$ -subexponential (cf. Definition 3.7).

Proof of Lemma 3.10. Throughout this proof let $Y: \Omega \rightarrow \mathbb{R}$ satisfy $Y = X^2\mathbb{1}_A$, let $p, q \in [0, 1]$ satisfy $p = \mathbb{P}(A)$ and $q = 1 - \mathbb{P}(A)$, and let $f: (-\infty, 1/2) \rightarrow \mathbb{R}$ and $g: (-\infty, 1/2) \rightarrow \mathbb{R}$ satisfy for all $\lambda \in (-\infty, 1/2)$ that

$$f(\lambda) = 2\lambda^2 + \lambda + \frac{1}{2}\ln(1 - 2\lambda) \quad \text{and} \quad g(\lambda) = q\lambda + \ln(p + q\sqrt{1 - 2\lambda}). \quad (51)$$

Observe that for all $\lambda \in (-1/4, 1/4)$ it holds that $1 - 2\lambda > 0$, $1/4 - \lambda > 0$, and

$$\begin{aligned} f'(\lambda) &= 4\lambda + 1 - \frac{1}{(1 - 2\lambda)} = \frac{(4\lambda + 1)(1 - 2\lambda) - 1}{(1 - 2\lambda)} \\ &= \frac{4\lambda + 1 - 8\lambda^2 - 2\lambda - 1}{(1 - 2\lambda)} = \frac{2\lambda - 8\lambda^2}{(1 - 2\lambda)} = \frac{8\lambda(\frac{1}{4} - \lambda)}{(1 - 2\lambda)}. \end{aligned} \quad (52)$$

This ensures for all $\lambda \in (-1/4, 0)$ that $f'(\lambda) < 0$. Hence, we obtain that $f|_{(-1/4, 0)}$ is strictly decreasing. Moreover, note that (52) ensures that for all $\lambda \in (0, 1/4)$ it holds that $f'(\lambda) > 0$. Hence, we obtain that

$f|_{(0,1/4)}$ is strictly increasing. Combining this, the fact that $f|_{(-1/4,0)}$ is strictly decreasing, and the fact that $f(0) = 0$ ensures that for all $\lambda \in (-1/4, 1/4)$ it holds that

$$f(\lambda) \geq f(0) = 0. \quad (53)$$

Next observe that (51) establishes that for all $\lambda \in (-\infty, 1/2)$ it holds that

$$g'(\lambda) = q + \frac{\frac{q}{2}[(1-2\lambda)^{-1/2}(-2)]}{p + q\sqrt{1-2\lambda}} = q - q[p(1-2\lambda)^{1/2} + q(1-2\lambda)]^{-1} \quad (54)$$

and

$$g''(\lambda) = \frac{q[\frac{p}{2}(1-2\lambda)^{-1/2}(-2) - 2q]}{(p(1-2\lambda)^{1/2} + q(1-2\lambda))^2} = -\frac{q[p(1-2\lambda)^{-1/2} + 2q]}{(p(1-2\lambda)^{1/2} + q(1-2\lambda))^2} \leq 0. \quad (55)$$

Moreover, note that the fundamental theorem of calculus and the fact that $g(0) = g'(0) = 0$ establish that for all $\lambda \in (-\infty, 1/2)$ it holds that

$$g(\lambda) = g(0) + \int_0^\lambda g'(s) ds = g(0) + \int_0^\lambda \left(g'(0) + \int_0^s g''(r) dr \right) ds = \int_0^\lambda \int_0^s g''(r) dr ds. \quad (56)$$

Combining this with (55) shows that for all $\lambda \in [0, 1/2)$ it holds that

$$g(\lambda) = \int_0^\lambda \int_0^s g''(r) dr ds \leq 0. \quad (57)$$

Furthermore, note that (55) and (56) establish for all $\lambda \in (-\infty, 0)$ that

$$g(\lambda) = \int_0^\lambda \int_0^s g''(r) dr ds = -\int_\lambda^0 \int_0^s g''(r) dr ds = \int_\lambda^0 \int_s^0 g''(r) dr ds \leq 0. \quad (58)$$

Next note that the fact that X is a standard normal random variable ensures that for all $\lambda \in (-\infty, 1/2)$ it holds that

$$\begin{aligned} \mathbb{E}[\exp(\lambda X^2)] &= \int_{-\infty}^{\infty} \frac{1}{\sqrt{2\pi}} \exp\left(-\frac{x^2}{2}\right) \exp(\lambda x^2) dx = \int_{-\infty}^{\infty} \frac{1}{\sqrt{2\pi}} \exp\left(-\frac{x^2(1-2\lambda)}{2}\right) dx \\ &= \frac{1}{\sqrt{1-2\lambda}} \int_{-\infty}^{\infty} \frac{1}{\sqrt{2\pi}} \exp\left(-\frac{y^2}{2}\right) dy = \frac{1}{\sqrt{1-2\lambda}} < \infty. \end{aligned} \quad (59)$$

In addition, observe that the assumption that X and $\mathbb{1}_A$ are independent and the fact that X is a standard normal random variable ensure that it holds that $\mathbb{E}[Y] = \mathbb{E}[X^2 \mathbb{1}_A] = \mathbb{E}[X^2] \mathbb{E}[\mathbb{1}_A] = \mathbb{E}[\mathbb{1}_A] = \mathbb{P}(A) = p$. This, the assumption that X and $\mathbb{1}_A$ are independent, (53), (57), (58), and (59) demonstrate that for all $\lambda \in (-1/4, 1/4)$ it holds that

$$\begin{aligned} \ln(\mathbb{E}[\exp(\lambda(Y - \mathbb{E}[Y]))]) &= \ln(\mathbb{E}[\exp(\lambda X^2 \mathbb{1}_A - p\lambda)]) = \ln(\mathbb{E}[\exp(\lambda X^2 \mathbb{1}_A)]) - p\lambda \\ &= \ln(\mathbb{E}[\exp(\lambda X^2 \mathbb{1}_A)(\mathbb{1}_A + \mathbb{1}_{\Omega \setminus A})]) - p\lambda \\ &= \ln(\mathbb{E}[\exp(\lambda X^2) \mathbb{1}_A + \mathbb{1}_{\Omega \setminus A}]) - p\lambda \\ &= \ln(p\mathbb{E}[\exp(\lambda X^2)] + q) - p\lambda = \ln\left(\frac{p}{\sqrt{1-2\lambda}} + q\right) - p\lambda \\ &= \ln\left(\frac{p + q\sqrt{1-2\lambda}}{\sqrt{1-2\lambda}}\right) - p\lambda = \ln(p + q\sqrt{1-2\lambda}) - \frac{1}{2} \ln(1-2\lambda) - p\lambda \\ &= \left(q\lambda + \ln(p + q\sqrt{1-2\lambda})\right) - \left(\frac{1}{2} \ln(1-2\lambda) + \lambda + 2\lambda^2\right) + 2\lambda^2 \\ &= g(\lambda) - f(\lambda) + 2\lambda^2 \leq g(\lambda) + 2\lambda^2 \leq 2\lambda^2. \end{aligned} \quad (60)$$

This implies for all $\lambda \in (-1/4, 1/4)$ that $\mathbb{E}[\exp(\lambda(Y - \mathbb{E}[Y]))] \leq \exp(2\lambda^2) = \exp(\frac{1}{2}\lambda^2 2^2)$. Hence, we obtain that Y is $(2, 4)$ -subexponential (cf. Definition 3.7). The proof of Lemma 3.10 is thus completed. \square

Lemma 3.11. *Let $(\Omega, \mathcal{F}, \mathbb{P})$ be a probability space, let $n \in \mathbb{N}$, $\sigma = (\sigma_1, \sigma_2, \dots, \sigma_n) \in \mathbb{R}^n$, $A_1, A_2, \dots, A_n \in \mathcal{F}$, let $X_i: \Omega \rightarrow \mathbb{R}$, $i \in \{1, 2, \dots, n\}$, be normal random variables, assume for all $i \in \{1, 2, \dots, n\}$ that $\text{Var}[X_i] = |\sigma_i|^2$, and assume that $X_1, X_2, \dots, X_n, \mathbb{1}_{A_1}, \mathbb{1}_{A_2}, \dots, \mathbb{1}_{A_n}$ are independent. Then $\sum_{i=1}^n |X_i - \mathbb{E}[X_i]|^2 \mathbb{1}_{A_i}$ is $(2(\sum_{i=1}^n |\sigma_i|^4)^{1/2}, 4 \max\{|\sigma_1|^2, |\sigma_2|^2, \dots, |\sigma_n|^2\})$ -subexponential (cf. Definition 3.7).*

Proof of Lemma 3.11. Throughout this proof let $Y_i: \Omega \rightarrow \mathbb{R}$, $i \in \{1, 2, \dots, n\}$, satisfy for all $i \in \{1, 2, \dots, n\}$ that $Y_i = 1/\sigma_i(X_i - \mathbb{E}[X_i])$. Observe that the assumption that for all $i \in \{1, 2, \dots, n\}$ it holds that X_i is a normal random variable with $\text{Var}[X_i] = |\sigma_i|^2$ ensures that for all $i \in \{1, 2, \dots, n\}$ it holds that Y_i is a standard normal random variable. This, the fact that for all $i \in \{1, 2, \dots, n\}$ it holds that Y_i and $\mathbb{1}_{A_i}$ are independent, and Lemma 3.10 (applied for every $i \in \{1, 2, \dots, n\}$ with $(\Omega, \mathcal{F}, \mathbb{P}) \curvearrowright (\Omega, \mathcal{F}, \mathbb{P})$, $X \curvearrowright Y_i$, $A \curvearrowright A_i$ in the notation of Lemma 3.10) ensure that for all $i \in \{1, 2, \dots, n\}$ it holds that $|Y_i|^2 \mathbb{1}_{A_i}$ is $(2, 4)$ -subexponential (cf. Definition 3.7). Lemma 3.8 (applied for every $i \in \{1, 2, \dots, n\}$ with $(\Omega, \mathcal{F}, \mathbb{P}) \curvearrowright (\Omega, \mathcal{F}, \mathbb{P})$, $\nu \curvearrowright 2$, $b \curvearrowright 4$, $a \curvearrowright |\sigma_i|^2$, $X \curvearrowright |Y_i|^2 \mathbb{1}_{A_i}$ in the notation of Lemma 3.8) thus demonstrates that for all $i \in \{1, 2, \dots, n\}$ it holds that $|\sigma_i Y_i|^2 \mathbb{1}_{A_i}$ is $(2|\sigma_i|^2, 4|\sigma_i|^2)$ -subexponential. Combining this, the fact that $|\sigma_1 Y_1|^2 \mathbb{1}_{A_1}, |\sigma_2 Y_2|^2 \mathbb{1}_{A_2}, \dots, |\sigma_n Y_n|^2 \mathbb{1}_{A_n}$ are independent, and Lemma 3.9 (applied with $(\Omega, \mathcal{F}, \mathbb{P}) \curvearrowright (\Omega, \mathcal{F}, \mathbb{P})$, $n \curvearrowright n$, $\nu = (2|\sigma_i|^2)_{i \in \{1, 2, \dots, n\}}$, $b = (4|\sigma_i|^2)_{i \in \{1, 2, \dots, n\}}$, $(X_i)_{i \in \{1, 2, \dots, n\}} \curvearrowright (|\sigma_i Y_i|^2 \mathbb{1}_{A_i})_{i \in \{1, 2, \dots, n\}}$ in the notation of Lemma 3.9) establishes that $\sum_{i=1}^n |\sigma_i Y_i|^2 \mathbb{1}_{A_i}$ is $(2(\sum_{i=1}^n |\sigma_i|^4)^{1/2}, 4 \max\{|\sigma_1|^2, |\sigma_2|^2, \dots, |\sigma_n|^2\})$ -subexponential. This and the fact that $\sum_{i=1}^n |X_i - \mathbb{E}[X_i]|^2 \mathbb{1}_{A_i} = \sum_{i=1}^n |\sigma_i Y_i|^2 \mathbb{1}_{A_i}$ therefore show that $\sum_{i=1}^n |X_i - \mathbb{E}[X_i]|^2 \mathbb{1}_{A_i}$ is $(2(\sum_{i=1}^n |\sigma_i|^4)^{1/2}, 4 \max\{|\sigma_1|^2, |\sigma_2|^2, \dots, |\sigma_n|^2\})$ -subexponential. The proof of Lemma 3.11 is thus completed. \square

Lemma 3.12. *Let $(\Omega, \mathcal{F}, \mathbb{P})$ be a probability space, let $\nu, b \in (0, \infty)$, and let X be (ν, b) -subexponential (cf. Definition 3.7). Then it holds for all $\varepsilon \in (0, \infty)$ that $\mathbb{P}(|X - \mathbb{E}[X]| \geq \varepsilon) \leq 2 \exp(-\frac{1}{2} \min\{\varepsilon^2/\nu^2, \varepsilon/b\})$.*

Proof of Lemma 3.12. Throughout this proof let $f_\varepsilon: (0, \infty) \rightarrow \mathbb{R}$, $\varepsilon \in (0, \infty)$, satisfy for all $\varepsilon, \lambda \in (0, \infty)$ that $f_\varepsilon(\lambda) = \frac{1}{2} \lambda^2 \nu^2 - \lambda \varepsilon$. Observe that the Markov inequality ensures for all $\varepsilon \in (0, \infty)$, $\lambda \in (0, \frac{1}{b})$ that

$$\begin{aligned} \mathbb{P}(|X - \mathbb{E}[X]| \geq \varepsilon) &= \mathbb{P}(X - \mathbb{E}[X] \geq \varepsilon) + \mathbb{P}(-(X - \mathbb{E}[X]) \geq \varepsilon) \\ &= \mathbb{P}(\exp(\lambda(X - \mathbb{E}[X])) \geq \exp(\lambda \varepsilon)) + \mathbb{P}(\exp(-\lambda(X - \mathbb{E}[X])) \geq \exp(\lambda \varepsilon)) \\ &\leq \frac{\mathbb{E}[\exp(\lambda(X - \mathbb{E}[X]))]}{\exp(\lambda \varepsilon)} + \frac{\mathbb{E}[\exp(-\lambda(X - \mathbb{E}[X]))]}{\exp(\lambda \varepsilon)} \\ &\leq \exp\left(\frac{\lambda^2 \nu^2}{2} - \lambda \varepsilon\right) + \exp\left(\frac{(-\lambda)^2 \nu^2}{2} - \lambda \varepsilon\right) \\ &= 2 \exp\left(\frac{\lambda^2 \nu^2}{2} - \lambda \varepsilon\right) = 2 \exp(f_\varepsilon(\lambda)). \end{aligned} \tag{61}$$

This and the fact that $\mathbb{R} \ni x \mapsto \exp(x) \in \mathbb{R}$ is strictly increasing ensure that for all $\varepsilon \in (0, \infty)$ it holds that

$$\mathbb{P}(|X - \mathbb{E}[X]| \geq \varepsilon) \leq \inf_{\lambda \in (0, \frac{1}{b})} 2 \exp(f_\varepsilon(\lambda)) = 2 \exp(\inf_{\lambda \in (0, \frac{1}{b})} f_\varepsilon(\lambda)). \tag{62}$$

In addition, note that the fact that for all $\varepsilon, \lambda \in (0, \infty)$ it holds that $f'_\varepsilon(\lambda) = \lambda \nu^2 - \varepsilon$ proves that for all $\varepsilon \in (0, \infty)$, $\lambda \in (0, \varepsilon/\nu^2)$ it holds that $f'_\varepsilon(\lambda) < 0$. Therefore, we obtain for all $\varepsilon \in (0, \infty)$ that $f_\varepsilon|_{(0, \varepsilon/\nu^2)}$ is strictly decreasing. This and the fact that for all $\varepsilon \in (\nu^2/b, \infty)$ it holds that $1/b < \varepsilon/\nu^2$ ensure that for all $\varepsilon \in (\nu^2/b, \infty)$ it holds that

$$\inf_{\lambda \in (0, \frac{1}{b})} f_\varepsilon(\lambda) = f_\varepsilon\left(\frac{1}{b}\right) = \frac{\nu^2}{2b^2} - \frac{\varepsilon}{b} \leq \frac{\varepsilon}{2b} - \frac{\varepsilon}{b} = -\frac{\varepsilon}{2b}. \tag{63}$$

Moreover, note that the fact that for all $\varepsilon, \lambda \in (0, \infty)$ it holds that $f'_\varepsilon(\lambda) = \lambda \nu^2 - \varepsilon$ implies that for all $\varepsilon \in (0, \infty)$, $\lambda \in (\varepsilon/\nu^2, \infty)$ it holds that $f'_\varepsilon(\lambda) > 0$. Therefore, we obtain for all $\varepsilon \in (0, \infty)$ that $f_\varepsilon|_{(\varepsilon/\nu^2, \infty)}$ is strictly increasing. This, the fact that for all $\varepsilon \in (0, \infty)$ it holds that $f_\varepsilon|_{(0, \varepsilon/\nu^2)}$ is strictly decreasing, and the fact that for all $\varepsilon \in (0, \nu^2/b]$ it holds that $1/b \geq \varepsilon/\nu^2$ establish that for all $\varepsilon \in (0, \nu^2/b]$ it holds that

$$\inf_{\lambda \in (0, \frac{1}{b})} f_\varepsilon(\lambda) = f_\varepsilon\left(\frac{\varepsilon}{\nu^2}\right) = \frac{\varepsilon^2}{2\nu^2} - \frac{\varepsilon^2}{\nu^2} = -\frac{\varepsilon^2}{2\nu^2}. \tag{64}$$

Combining this with (62) and (63) establishes that for all $\varepsilon \in (0, \infty)$ it holds that

$$\mathbb{P}(|X - \mathbb{E}[X]| \geq \varepsilon) \leq \max\left\{2 \exp\left(-\frac{\varepsilon^2}{2\nu^2}\right), 2 \exp\left(-\frac{\varepsilon}{2b}\right)\right\} = 2 \exp\left(-\frac{1}{2} \min\left\{\frac{\varepsilon^2}{\nu^2}, \frac{\varepsilon}{b}\right\}\right). \tag{65}$$

The proof of Lemma 3.12 is thus completed. \square

3.4 Concentration type inequalities for stochastic Gramian matrices at initialization

Definition 3.13 (Spectral norm). We denote by $\|\cdot\|$: $(\bigcup_{m,n \in \mathbb{N}} \mathbb{R}^{m \times n}) \rightarrow [0, \infty)$ the function which satisfies for all $m, n \in \mathbb{N}$, $A \in \mathbb{R}^{m \times n}$ that $\|A\| = \sup_{x \in \mathbb{R}^n \setminus \{0\}} \|Ax\|/\|x\|$ (cf. Definition 2.1).

Lemma 3.14. Let $m, n \in \mathbb{N}$, $A = (A_{i,j})_{(i,j) \in \{1,2,\dots,m\} \times \{1,2,\dots,n\}} \in \mathbb{R}^{m \times n}$. Then $\|A\| \leq (\sum_{i=1}^m \sum_{j=1}^n |A_{i,j}|^2)^{1/2} \leq \sum_{i=1}^m \sum_{j=1}^n |A_{i,j}|$ (cf. Definition 3.13).

Proof of Lemma 3.14. Throughout this proof let $e_1 = (1, 0, \dots, 0)$, $e_2 = (0, 1, 0, \dots, 0)$, \dots , $e_n = (0, \dots, 0, 1) \in \mathbb{R}^n$. Observe that the Cauchy-Schwarz inequality ensures that for all $x = (x_1, x_2, \dots, x_n) \in \mathbb{R}^n$ it holds that

$$\begin{aligned} \|Ax\| &= \left\| A \left(\sum_{j=1}^n x_j e_j \right) \right\| = \left\| \sum_{j=1}^n x_j A e_j \right\| \leq \sum_{j=1}^n |x_j| \|A e_j\| \\ &\leq \left(\sum_{j=1}^n |x_j|^2 \right)^{1/2} \left(\sum_{j=1}^n \|A e_j\|^2 \right)^{1/2} = \|x\| \left(\sum_{j=1}^n \sum_{i=1}^m |A_{i,j}|^2 \right)^{1/2} \end{aligned} \quad (66)$$

(cf. Definition 2.1). Hence, we obtain for all $x \in \mathbb{R}^n \setminus \{0\}$ that $\|Ax\|/\|x\| \leq (\sum_{i=1}^m \sum_{j=1}^n |A_{i,j}|^2)^{1/2}$. This implies that $\|A\| = \sup_{x \in \mathbb{R}^n \setminus \{0\}} \|Ax\|/\|x\| \leq (\sum_{i=1}^m \sum_{j=1}^n |A_{i,j}|^2)^{1/2}$ (cf. Definition 3.13). Moreover, the fact that for all $a, b \in [0, \infty)$ it holds that $(a+b)^{1/2} \leq a^{1/2} + b^{1/2}$ inductively ensures that $(\sum_{i=1}^m \sum_{j=1}^n |A_{i,j}|^2)^{1/2} \leq \sum_{i=1}^m \sum_{j=1}^n |A_{i,j}|$. The proof of Lemma 3.14 is thus completed. \square

Lemma 3.15. Assume Setting 2.7, assume $\lambda, \mu \in (0, \infty)$, let $\varepsilon \in (0, 1)$, and assume $\mathfrak{d} \geq 32 \ln(\frac{2m^2}{\varepsilon}) \max\{\frac{4m^2}{\lambda^2}, \frac{4m^2 C^4}{\mu^2}, \frac{m}{\lambda}, \frac{mC^2}{\mu}\}$. Then $\mathbb{P}(\bigcap_{i=1}^m \bigcap_{j=1}^m \{|\mathcal{G}_{i,j}(0) - \mathbf{G}_{i,j}|\} \leq \frac{\min\{\lambda, \mu C^{-2}\}}{4m}) \geq 1 - \varepsilon$.

Proof of Lemma 3.15. Observe that (11) and the assumption that $\|B(0)\| = 0$ ensure that for all $i, j \in \{1, 2, \dots, m\}$ it holds that

$$\mathcal{G}_{i,j}(0) = \sum_{k=1}^{\mathfrak{d}} |\mathcal{W}_k|^2 \mathbb{1}_{\{\langle W_k(0), x_i \rangle + B_k(0) \geq 0, \langle W_k(0), x_j \rangle + B_k(0) \geq 0\}} = \sum_{k=1}^{\mathfrak{d}} |\mathcal{W}_k|^2 \mathbb{1}_{\{\langle W_k(0), x_i \rangle \geq 0, \langle W_k(0), x_j \rangle \geq 0\}} \quad (67)$$

(cf. Definition 2.1). Moreover, note that the assumption that $W_1(0), W_2(0), \dots, W_{\mathfrak{d}}(0), \sqrt{\mathfrak{d}}\mathcal{W}_1, \sqrt{\mathfrak{d}}\mathcal{W}_2, \dots, \sqrt{\mathfrak{d}}\mathcal{W}_{\mathfrak{d}}$ are independent implies that for all $i, j \in \{1, 2, \dots, m\}$, $k \in \{1, 2, \dots, \mathfrak{d}\}$ it holds that \mathcal{W}_k and $\mathbb{1}_{\{\langle W_k(0), x_i \rangle \geq 0, \langle W_k(0), x_j \rangle \geq 0\}}$ are independent. Combining this, Lemma 3.11 (applied for every $i, j \in \{1, 2, \dots, m\}$ with $(\Omega, \mathcal{F}, \mathbb{P}) \curvearrowright (\Omega, \mathcal{F}, \mathbb{P})$, $n \curvearrowright \mathfrak{d}$, $\sigma \curvearrowright (1/\sqrt{\mathfrak{d}}, 1/\sqrt{\mathfrak{d}}, \dots, 1/\sqrt{\mathfrak{d}})$, $(X_i)_{i \in \{1, 2, \dots, n\}} \curvearrowright (\mathcal{W}_k)_{k \in \{1, 2, \dots, \mathfrak{d}\}}$, $(A_i)_{i \in \{1, 2, \dots, n\}} \curvearrowright (\{\langle W_k(0), x_i \rangle \geq 0, \langle W_k(0), x_j \rangle \geq 0\})_{k \in \{1, 2, \dots, \mathfrak{d}\}}$ in the notation of Lemma 3.11), and the fact that for all $k \in \{1, 2, \dots, \mathfrak{d}\}$ it holds that \mathcal{W}_k is a centered normal random variable with $\text{Var}[\mathcal{W}_k] = 1/\mathfrak{d}$ ensures that for all $i, j \in \{1, 2, \dots, m\}$ it holds that $\mathcal{G}_{i,j}(0)$ is $(2/\sqrt{\mathfrak{d}}, 4/\mathfrak{d})$ -subexponential (cf. Definition 3.7). Moreover, observe that Lemma 2.8 demonstrates for all $i, j \in \{1, 2, \dots, m\}$ that $\mathbb{E}[\mathcal{G}_{i,j}(0)] = \mathbf{G}_{i,j}$. Lemma 3.12 (applied for every $i, j \in \{1, 2, \dots, m\}$ with $(\Omega, \mathcal{F}, \mathbb{P}) \curvearrowright (\Omega, \mathcal{F}, \mathbb{P})$, $\nu \curvearrowright 2/\sqrt{\mathfrak{d}}$, $b \curvearrowright 4/\mathfrak{d}$, $X \curvearrowright \mathcal{G}_{i,j}(0)$, $\varepsilon \curvearrowright \min\{\lambda, \mu C^{-2}\}(4m)^{-1}$ in the notation of Lemma 3.12), the fact that for all $i, j \in \{1, 2, \dots, m\}$ it holds that $\mathcal{G}_{i,j}(0)$ is $(2/\sqrt{\mathfrak{d}}, 4/\mathfrak{d})$ -subexponential, and the assumption that $\mathfrak{d} \geq 32 \ln(\frac{2m^2}{\varepsilon}) \max\{\frac{4m^2}{\lambda^2}, \frac{4m^2 C^4}{\mu^2}, \frac{m}{\lambda}, \frac{mC^2}{\mu}\}$ hence establish that for all $i, j \in \{1, 2, \dots, m\}$ it holds that

$$\begin{aligned} &\mathbb{P}\left(|\mathcal{G}_{i,j}(0) - \mathbf{G}_{i,j}| \geq \frac{\min\{\lambda, \mu C^{-2}\}}{4m}\right) \\ &\leq 2 \exp\left(-\frac{1}{2} \min\left\{\frac{\mathfrak{d}}{4} \left(\frac{\min\{\lambda, \mu C^{-2}\}}{4m}\right)^2, \frac{\mathfrak{d}}{4} \frac{\min\{\lambda, \mu C^{-2}\}}{4m}\right\}\right) \\ &= 2 \exp\left(-\frac{\mathfrak{d}}{8} \min\left\{\left(\frac{\min\{\lambda, \mu C^{-2}\}}{4m}\right)^2, \frac{\min\{\lambda, \mu C^{-2}\}}{4m}\right\}\right) \\ &= 2 \exp\left(-\frac{\mathfrak{d}}{8} \min\left\{\min\left\{\frac{\lambda^2}{16m^2}, \frac{\mu^2}{16m^2 C^4}\right\}, \min\left\{\frac{\lambda}{4m}, \frac{\mu}{4m C^2}\right\}\right\}\right) \\ &= 2 \exp\left(-\frac{\mathfrak{d}}{32} \min\left\{\frac{\lambda^2}{4m^2}, \frac{\mu^2}{4m^2 C^4}, \frac{\lambda}{m}, \frac{\mu}{m C^2}\right\}\right) \\ &\leq 2 \exp\left(-\ln\left(\frac{2m^2}{\varepsilon}\right) \max\left\{\frac{4m^2}{\lambda^2}, \frac{4m^2 C^4}{\mu^2}, \frac{m}{\lambda}, \frac{m C^2}{\mu}\right\} \min\left\{\frac{\lambda^2}{4m^2}, \frac{\mu^2}{4m^2 C^4}, \frac{\lambda}{m}, \frac{\mu}{m C^2}\right\}\right) \\ &= 2 \exp\left(-\ln\left(\frac{2m^2}{\varepsilon}\right)\right) = 2 \exp\left(\ln\left(\frac{\varepsilon}{2m^2}\right)\right) = \frac{\varepsilon}{m^2}. \end{aligned} \quad (68)$$

Therefore, we obtain that

$$\begin{aligned}
\mathbb{P}\left(\bigcap_{i=1}^m \bigcap_{j=1}^m \left\{ |\mathcal{G}_{i,j}(0) - \mathbf{G}_{i,j}| \leq \frac{\min\{\lambda, \mu C^{-2}\}}{4m} \right\}\right) &\geq 1 - \mathbb{P}\left(\bigcup_{i=1}^m \bigcup_{j=1}^m \left\{ |\mathcal{G}_{i,j}(0) - \mathbf{G}_{i,j}| \geq \frac{\min\{\lambda, \mu C^{-2}\}}{4m} \right\}\right) \\
&\geq 1 - \sum_{i=1}^m \sum_{j=1}^m \mathbb{P}\left(|\mathcal{G}_{i,j}(0) - \mathbf{G}_{i,j}| \geq \frac{\min\{\lambda, \mu C^{-2}\}}{4m}\right) \\
&\geq 1 - \sum_{i=1}^m \sum_{j=1}^m \frac{\varepsilon}{m^2} = 1 - \varepsilon.
\end{aligned} \tag{69}$$

This completes the proof of Lemma 3.15. \square

Lemma 3.16. *Assume Setting 2.7, let $\varepsilon \in (0, 1)$, $A \in \mathcal{F}$ satisfy $A = (\bigcap_{i,j=1}^m \{|\mathcal{G}_{i,j}(0) - \mathbf{G}_{i,j}| \leq \frac{\min\{\lambda, \mu C^{-2}\}}{4m}\})$, and let $\omega \in A$. Then*

(i) *it holds that $\|\mathcal{G}(0, \omega) - \mathbf{G}\| \leq \lambda/4$, and*

(ii) *it holds that $\|\mathcal{H}(0, \omega) - \mathbf{H}\| \leq \mu/4$*

(cf. Definition 3.13).

Proof of Lemma 3.16. Note that Lemma 3.14 (applied with $m \curvearrowright m$, $n \curvearrowright m$, $A \curvearrowright \mathcal{G}(0, \omega) - \mathbf{G}$ in the notation of Lemma 3.14) and the assumption that $\omega \in A$ ensure that

$$\|\mathcal{G}(0, \omega) - \mathbf{G}\|^2 \leq \sum_{i=1}^m \sum_{j=1}^m |\mathcal{G}_{i,j}(0, \omega) - \mathbf{G}_{i,j}|^2 \leq \sum_{i=1}^m \sum_{j=1}^m \left(\frac{\min\{\lambda, \mu C^{-2}\}}{4m}\right)^2 = \frac{\min\{\lambda^2, \mu^2 C^{-4}\}}{16} \leq \frac{\lambda^2}{16}. \tag{70}$$

(cf. Definition 3.13). Thus, we obtain that $\|\mathcal{G}(0, \omega) - \mathbf{G}\| \leq \lambda/4$. This establishes item (i). In addition, observe that Lemma 3.14 (applied with $m \curvearrowright m$, $n \curvearrowright m$, $A \curvearrowright \mathcal{H}(0, \omega) - \mathbf{H}$ in the notation of Lemma 3.14), the fact that for all $i, j \in \{1, 2, \dots, m\}$ it holds that $|\langle x_i, x_j \rangle|^2 \leq \|x_i\|^2 \|x_j\|^2 \leq C^4$, and the assumption that $\omega \in A$ demonstrate that

$$\begin{aligned}
\|\mathcal{H}(0, \omega) - \mathbf{H}\|^2 &\leq \sum_{i=1}^m \sum_{j=1}^m |\mathcal{H}_{i,j}(0, \omega) - \mathbf{H}_{i,j}|^2 = \sum_{i=1}^m \sum_{j=1}^m |\langle x_i, x_j \rangle|^2 |\mathcal{G}_{i,j}(0, \omega) - \mathbf{G}_{i,j}|^2 \\
&\leq C^4 \sum_{i=1}^m \sum_{j=1}^m |\mathcal{G}_{i,j}(0, \omega) - \mathbf{G}_{i,j}|^2 \leq C^4 \sum_{i=1}^m \sum_{j=1}^m \left(\frac{\min\{\lambda, \mu C^{-2}\}}{4m}\right)^2 = \frac{C^4 \min\{\lambda^2, \mu^2 C^{-4}\}}{16} \leq \frac{\mu^2}{16}.
\end{aligned} \tag{71}$$

Thus, we obtain that $\|\mathcal{H}(0, \omega) - \mathbf{H}\| \leq \mu/4$. This establishes item (ii). The proof of Lemma 3.16 is thus completed. \square

3.5 Analysis of stochastic Gramian matrices during training

Lemma 3.17. *Let $(\Omega, \mathcal{F}, \mathbb{P})$ be a probability space, let $X : \Omega \rightarrow \mathbb{R}$ be a standard normal random variable, and let $\varepsilon \in (0, \infty)$. Then $\mathbb{P}(|X| \leq \varepsilon) \leq 2\varepsilon/\sqrt{2\pi}$.*

Proof of Lemma 3.17. Observe that the fact that X is a standard normal random variable and the fact that for all $x \in \mathbb{R}$ it holds that $\exp(-x^2/2) \leq 1$ establish that it holds that

$$\mathbb{P}(|X| \leq \varepsilon) = \mathbb{P}(-\varepsilon \leq X \leq \varepsilon) = \int_{-\varepsilon}^{\varepsilon} \frac{1}{\sqrt{2\pi}} \exp\left(-\frac{x^2}{2}\right) dx \leq \int_{-\varepsilon}^{\varepsilon} \frac{1}{\sqrt{2\pi}} dx = \frac{2\varepsilon}{\sqrt{2\pi}}. \tag{72}$$

The proof of Lemma 3.17 is thus completed. \square

Lemma 3.18. *Assume Setting 2.7, assume $\lambda, \mu \in (0, \infty)$, and let $\varepsilon \in (0, 1)$, $R \in (0, \infty)$ satisfy $R \leq \sqrt{2\pi}\varepsilon \min\{\lambda, \mu C^{-2}\} 2^{-4} m^{-2}$. Then it holds that $\mathbb{P}(\sum_{i,j=1}^m \sum_{k=1}^{\mathfrak{d}} |\mathcal{W}_k|^2 (\mathbb{1}_{\{|\langle W_k(0), x_i \rangle| \leq R\}} + \mathbb{1}_{\{|\langle W_k(0), x_j \rangle| \leq R\}})) \leq \min\{\lambda, \mu C^{-2}\}/4) \geq 1 - \varepsilon$.*

Proof of Lemma 3.18. Throughout this proof let $X_{k,i}: \Omega \rightarrow \mathbb{R}$, $k \in \{1, 2, \dots, \mathfrak{d}\}$, $i \in \{1, 2, \dots, m\}$, satisfy for all $k \in \{1, 2, \dots, \mathfrak{d}\}$, $i \in \{1, 2, \dots, m\}$ that $X_{k,i} = \langle W_k(0), x_i \rangle / \|x_i\|$. Observe that the assumption that $W_1(0), W_2(0), \dots, W_{\mathfrak{d}}(0)$ are standard normal ensures that for all $k \in \{1, 2, \dots, \mathfrak{d}\}$, $i \in \{1, 2, \dots, m\}$ it holds that $\langle W_k(0), x_i \rangle$ is a centered normal random variable with $\text{Var}[\langle W_k(0), x_i \rangle] = \|x_i\|^2$. Hence, we obtain for all $k \in \{1, 2, \dots, \mathfrak{d}\}$, $i \in \{1, 2, \dots, m\}$ that $X_{k,i}$ is a standard normal random variable. This and Lemma 3.17 (applied for every $k \in \{1, 2, \dots, \mathfrak{d}\}$, $i \in \{1, 2, \dots, m\}$ with $(\Omega, \mathcal{F}, \mathbb{P}) \curvearrowright (\Omega, \mathcal{F}, \mathbb{P})$, $X \curvearrowright X_{k,i}$, $\varepsilon \curvearrowright R/c$ in the notation of Lemma 3.17) ensure that for all $k \in \{1, 2, \dots, \mathfrak{d}\}$, $i \in \{1, 2, \dots, m\}$ it holds that

$$\begin{aligned} \mathbb{E}[\mathbb{1}_{\{|\langle W_k(0), x_i \rangle| \leq R\}}] &= \mathbb{P}(|\langle W_k(0), x_i \rangle| \leq R) = \mathbb{P}(|X_{k,i}| \leq R/\|x_i\|) \\ &\leq \mathbb{P}(|X_{k,i}| \leq R/c) \leq \frac{2R}{\sqrt{2\pi}c} \leq \frac{\varepsilon \min\{\lambda, \mu C^{-2}\}}{8m^2}. \end{aligned} \quad (73)$$

Moreover, note that the assumption that $W_1(0), W_2(0), \dots, W_{\mathfrak{d}}(0), \sqrt{\mathfrak{d}}W_1, \sqrt{\mathfrak{d}}W_2, \dots, \sqrt{\mathfrak{d}}W_{\mathfrak{d}}$ are independent ensures that for all $k \in \{1, 2, \dots, \mathfrak{d}\}$, $i \in \{1, 2, \dots, m\}$ it holds that \mathcal{W}_k and $\mathbb{1}_{\{|\langle W_k(0), x_i \rangle| \leq R\}}$ are independent. Combining this, (73), and the fact that for all $k \in \{1, 2, \dots, \mathfrak{d}\}$ it holds that \mathcal{W}_k is a centered normal random variable with $\text{Var}[\mathcal{W}_k] = 1/\mathfrak{d}$ establishes that it holds that

$$\begin{aligned} &\mathbb{E}\left[\sum_{i=1}^m \sum_{j=1}^m \sum_{k=1}^{\mathfrak{d}} |\mathcal{W}_k|^2 (\mathbb{1}_{\{|\langle W_k(0), x_i \rangle| \leq R\}} + \mathbb{1}_{\{|\langle W_k(0), x_j \rangle| \leq R\}})\right] \\ &= \sum_{i=1}^m \sum_{j=1}^m \sum_{k=1}^{\mathfrak{d}} \mathbb{E}[|\mathcal{W}_k|^2] \left(\mathbb{E}[\mathbb{1}_{\{|\langle W_k(0), x_i \rangle| \leq R\}}] + \mathbb{E}[\mathbb{1}_{\{|\langle W_k(0), x_j \rangle| \leq R\}}] \right) \\ &\leq \sum_{i=1}^m \sum_{j=1}^m \sum_{k=1}^{\mathfrak{d}} \frac{1}{\mathfrak{d}} \left(\frac{\varepsilon \min\{\lambda, \mu C^{-2}\}}{8m^2} + \frac{\varepsilon \min\{\lambda, \mu C^{-2}\}}{8m^2} \right) = \sum_{i=1}^m \sum_{j=1}^m \frac{\varepsilon \min\{\lambda, \mu C^{-2}\}}{4m^2} = \frac{\varepsilon \min\{\lambda, \mu C^{-2}\}}{4}. \end{aligned} \quad (74)$$

The Markov inequality thus demonstrates that

$$\begin{aligned} &\mathbb{P}\left(\sum_{i=1}^m \sum_{j=1}^m \sum_{k=1}^{\mathfrak{d}} |\mathcal{W}_k|^2 (\mathbb{1}_{\{|\langle W_k(0), x_i \rangle| \leq R\}} + \mathbb{1}_{\{|\langle W_k(0), x_j \rangle| \leq R\}}) \leq \frac{\min\{\lambda, \mu C^{-2}\}}{4}\right) \\ &\geq 1 - \mathbb{P}\left(\sum_{i=1}^m \sum_{j=1}^m \sum_{k=1}^{\mathfrak{d}} |\mathcal{W}_k|^2 (\mathbb{1}_{\{|\langle W_k(0), x_i \rangle| \leq R\}} + \mathbb{1}_{\{|\langle W_k(0), x_j \rangle| \leq R\}}) \geq \frac{\min\{\lambda, \mu C^{-2}\}}{4}\right) \\ &\geq 1 - \frac{\mathbb{E}[\sum_{i=1}^m \sum_{j=1}^m \sum_{k=1}^{\mathfrak{d}} |\mathcal{W}_k|^2 (\mathbb{1}_{\{|\langle W_k(0), x_i \rangle| \leq R\}} + \mathbb{1}_{\{|\langle W_k(0), x_j \rangle| \leq R\}})]}{\min\{\lambda, \mu C^{-2}\} 4^{-1}} \geq 1 - \varepsilon. \end{aligned} \quad (75)$$

This completes the proof of Lemma 3.18. \square

Lemma 3.19. *Assume Setting 2.7, assume $\lambda, \mu \in (0, \infty)$, let $\varepsilon \in (0, 1)$, $R \in (0, \infty)$, $A \in \mathcal{F}$ satisfy $A = \{\sum_{i,j=1}^m \sum_{k=1}^{\mathfrak{d}} |\mathcal{W}_k|^2 (\mathbb{1}_{\{|\langle W_k(0), x_i \rangle| \leq R\}} + \mathbb{1}_{\{|\langle W_k(0), x_j \rangle| \leq R\}}) \leq \min\{\lambda, \mu C^{-2}\}/4\}$, let $\omega \in A$, $n \in \mathbb{N}_0$, and assume $\max_{k \in \{1, 2, \dots, \mathfrak{d}\}} [C \|W_k(n, \omega) - W_k(0, \omega)\| + |B_k(n, \omega) - B_k(0, \omega)|] \leq R$ (cf. Definition 2.1). Then*

(i) *it holds that $\|\mathcal{G}(n, \omega) - \mathcal{G}(0, \omega)\| \leq \lambda/4$, and*

(ii) *it holds that $\|\mathcal{H}(n, \omega) - \mathcal{H}(0, \omega)\| \leq \mu/4$*

(cf. Definition 3.13).

Proof of Lemma 3.19. Observe that (11) and the assumption that $\|B(0)\| = 0$ ensure that for all $i, j \in \{1, 2, \dots, m\}$ it holds that

$$\begin{aligned} &|\mathcal{G}_{i,j}(n, \omega) - \mathcal{G}_{i,j}(0, \omega)| \\ &= \left| \sum_{k=1}^{\mathfrak{d}} |\mathcal{W}_k(\omega)|^2 \mathbb{1}_{\{\langle W_k(n), x_i \rangle + B_k(n) \geq 0, \langle W_k(n), x_j \rangle + B_k(n) \geq 0\}}(\omega) \right. \\ &\quad \left. - \sum_{k=1}^{\mathfrak{d}} |\mathcal{W}_k(\omega)|^2 \mathbb{1}_{\{\langle W_k(0), x_i \rangle + B_k(0) \geq 0, \langle W_k(0), x_j \rangle + B_k(0) \geq 0\}}(\omega) \right| \\ &= \left| \sum_{k=1}^{\mathfrak{d}} |\mathcal{W}_k(\omega)|^2 (\mathbb{1}_{\{\langle W_k(n), x_i \rangle + B_k(n) \geq 0, \langle W_k(n), x_j \rangle + B_k(n) \geq 0\}}(\omega) - \mathbb{1}_{\{\langle W_k(0), x_i \rangle \geq 0, \langle W_k(0), x_j \rangle \geq 0\}}(\omega)) \right| \\ &\leq \sum_{k=1}^{\mathfrak{d}} |\mathcal{W}_k(\omega)|^2 |\mathbb{1}_{\{\langle W_k(n), x_i \rangle + B_k(n) \geq 0, \langle W_k(n), x_j \rangle + B_k(n) \geq 0\}}(\omega) - \mathbb{1}_{\{\langle W_k(0), x_i \rangle \geq 0, \langle W_k(0), x_j \rangle \geq 0\}}(\omega)|. \end{aligned} \quad (76)$$

Moreover, observe that for all $a, b, \mathfrak{a}, \mathfrak{b} \in \mathbb{R}$ it holds that

$$\begin{aligned}
|\mathbb{1}_{[0,\infty)^2}(a, b) - \mathbb{1}_{[0,\infty)^2}(\mathfrak{a}, \mathfrak{b})| &= |\mathbb{1}_{[0,\infty)^2}(a, b) - \mathbb{1}_{[0,\infty)^2}(\mathfrak{a}, b) + \mathbb{1}_{[0,\infty)^2}(\mathfrak{a}, b) - \mathbb{1}_{[0,\infty)^2}(\mathfrak{a}, \mathfrak{b})| \\
&= |\mathbb{1}_{[0,\infty)}(b)(\mathbb{1}_{[0,\infty)}(a) - \mathbb{1}_{[0,\infty)}(\mathfrak{a})) + \mathbb{1}_{[0,\infty)}(\mathfrak{a})(\mathbb{1}_{[0,\infty)}(b) - \mathbb{1}_{[0,\infty)}(\mathfrak{b}))| \\
&\leq |\mathbb{1}_{[0,\infty)}(b)(\mathbb{1}_{[0,\infty)}(a) - \mathbb{1}_{[0,\infty)}(\mathfrak{a}))| + |\mathbb{1}_{[0,\infty)}(\mathfrak{a})(\mathbb{1}_{[0,\infty)}(b) - \mathbb{1}_{[0,\infty)}(\mathfrak{b}))| \quad (77) \\
&\leq |\mathbb{1}_{[0,\infty)}(a) - \mathbb{1}_{[0,\infty)}(\mathfrak{a})| + |\mathbb{1}_{[0,\infty)}(b) - \mathbb{1}_{[0,\infty)}(\mathfrak{b})| \\
&\leq \mathbb{1}_{[0,\infty)}(|a - \mathfrak{a}| - |\mathfrak{a}|) + \mathbb{1}_{[0,\infty)}(|b - \mathfrak{b}| - |\mathfrak{b}|).
\end{aligned}$$

This ensures that for all $k \in \{1, 2, \dots, \mathfrak{d}\}$, $i, j \in \{1, 2, \dots, m\}$ it holds that

$$\begin{aligned}
&|\mathbb{1}_{\{\langle W_k(n), x_i \rangle + B_k(n) \geq 0, \langle W_k(n), x_j \rangle + B_k(n) \geq 0\}}(\omega) - \mathbb{1}_{\{\langle W_k(0), x_i \rangle \geq 0, \langle W_k(0), x_j \rangle \geq 0\}}(\omega)| \\
&= |\mathbb{1}_{[0,\infty)^2}(\langle W_k(n), x_i \rangle + B_k(n), \langle W_k(n), x_j \rangle + B_k(n)) - \mathbb{1}_{[0,\infty)^2}(\langle W_k(0), x_i \rangle, \langle W_k(0), x_j \rangle)| \\
&\leq \mathbb{1}_{[0,\infty)}(|\langle W_k(n), x_i \rangle + B_k(n) - \langle W_k(0), x_i \rangle| - |\langle W_k(0), x_i \rangle|) \\
&\quad + \mathbb{1}_{[0,\infty)}(|\langle W_k(n), x_j \rangle + B_k(n) - \langle W_k(0), x_j \rangle| - |\langle W_k(0), x_j \rangle|). \quad (78)
\end{aligned}$$

In addition, note that the assumption that $\max_{k \in \{1, 2, \dots, \mathfrak{d}\}} [C\|W_k(n, \omega) - W_k(0, \omega)\| + |B_k(n, \omega) - B_k(0, \omega)|] \leq R$ and the Cauchy Schwarz inequality ensure that for all $k \in \{1, 2, \dots, \mathfrak{d}\}$, $i \in \{1, 2, \dots, m\}$ it holds that

$$\begin{aligned}
&\mathbb{1}_{[0,\infty)}(|\langle W_k(n), x_i \rangle + B_k(n) - \langle W_k(0), x_i \rangle| - |\langle W_k(0), x_i \rangle|) \\
&= \mathbb{1}_{[0,\infty)}(|\langle W_k(n) - W_k(0), x_i \rangle + B_k(n) - B_k(0) - \langle W_k(0), x_i \rangle|) \\
&\leq \mathbb{1}_{[0,\infty)}(|\langle W_k(n) - W_k(0), x_i \rangle| + |B_k(n) - B_k(0)| - |\langle W_k(0), x_i \rangle|) \\
&\leq \mathbb{1}_{[0,\infty)}(\|W_k(n) - W_k(0)\| \|x_i\| + |B_k(n) - B_k(0)| - |\langle W_k(0), x_i \rangle|) \quad (79) \\
&\leq \mathbb{1}_{[0,\infty)}(C\|W_k(n) - W_k(0)\| + |B_k(n) - B_k(0)| - |\langle W_k(0), x_i \rangle|) \\
&\leq \mathbb{1}_{[0,\infty)}(R - |\langle W_k(0), x_i \rangle|) = \mathbb{1}_{\{|\langle W_k(0), x_i \rangle| \leq R\}}(\omega).
\end{aligned}$$

Combining this, (76), (78), and the assumption that $\omega \in A$ therefore demonstrates that

$$\begin{aligned}
&\sum_{i=1}^m \sum_{j=1}^m |\mathcal{G}_{i,j}(n, \omega) - \mathcal{G}_{i,j}(0, \omega)| \\
&\leq \sum_{i=1}^m \sum_{j=1}^m \sum_{k=1}^{\mathfrak{d}} |\mathcal{W}_k(\omega)|^2 |\mathbb{1}_{\{\langle W_k(n), x_i \rangle + B_k(n) \geq 0, \langle W_k(n), x_j \rangle + B_k(n) \geq 0\}}(\omega) - \mathbb{1}_{\{\langle W_k(0), x_i \rangle \geq 0, \langle W_k(0), x_j \rangle \geq 0\}}(\omega)| \quad (80) \\
&\leq \sum_{i=1}^m \sum_{j=1}^m \sum_{k=1}^{\mathfrak{d}} |\mathcal{W}_k(\omega)|^2 (\mathbb{1}_{\{|\langle W_k(0), x_i \rangle| \leq R\}}(\omega) + \mathbb{1}_{\{|\langle W_k(0), x_j \rangle| \leq R\}}(\omega)) \leq \frac{\min\{\lambda, \mu C^{-2}\}}{4}.
\end{aligned}$$

Lemma 3.14 (applied with $m \curvearrowright m$, $n \curvearrowright m$, $A \curvearrowright \mathcal{G}(n, \omega) - \mathcal{G}(0, \omega)$ in the notation of Lemma 3.14) hence ensures that $\|\mathcal{G}(n, \omega) - \mathcal{G}(0, \omega)\| \leq \sum_{i=1}^m \sum_{j=1}^m |\mathcal{G}_{i,j}(n, \omega) - \mathcal{G}_{i,j}(0, \omega)| \leq \min\{\lambda, \mu C^{-2}\}/4 \leq \lambda/4$ (cf. Definition 3.13). This establishes item (i). Moreover, note that Lemma 3.14 (applied with $m \curvearrowright m$, $n \curvearrowright m$, $A \curvearrowright \mathcal{H}(n, \omega) - \mathcal{H}(0, \omega)$ in the notation of Lemma 3.14), the fact that for all $i, j \in \{1, 2, \dots, m\}$ it holds that $|\langle x_i, x_j \rangle| \leq \|x_i\| \|x_j\| \leq C^2$, and (80) ensure that

$$\begin{aligned}
\|\mathcal{H}(n, \omega) - \mathcal{H}(0, \omega)\| &\leq \sum_{i=1}^m \sum_{j=1}^m |\mathcal{H}_{i,j}(n, \omega) - \mathcal{H}_{i,j}(0, \omega)| = \sum_{i=1}^m \sum_{j=1}^m |\langle x_i, x_j \rangle| |\mathcal{G}_{i,j}(n, \omega) - \mathcal{G}_{i,j}(0, \omega)| \\
&\leq C^2 \sum_{i=1}^m \sum_{j=1}^m |\mathcal{G}_{i,j}(n, \omega) - \mathcal{G}_{i,j}(0, \omega)| \leq C^2 \frac{\min\{\lambda, \mu C^{-2}\}}{4} \leq \frac{\mu}{4}. \quad (81)
\end{aligned}$$

This establishes item (ii). The proof of Lemma 3.19 is thus completed. \square

3.6 Analysis of eigenvalues of stochastic Gramian matrices during training

Lemma 3.20. *Let $n \in \mathbb{N}$ and let $A \in \mathbb{R}^{n \times n}$ be a symmetric matrix. Then*

(i) *it holds for all $x \in \mathbb{R}^n$ that $\langle x, Ax \rangle \geq \lambda_{\min}(A) \|x\|^2$ and*

(ii) it holds that $\lambda_{\min}(A) = \min\{\langle x, Ax \rangle : [x \in \mathbb{R}^n, \|x\| = 1]\}$

(cf. Definitions 2.1 and 2.6).

Proof of Lemma 3.20. Note that the spectral theorem ensures that there exist $\lambda_1, \lambda_2, \dots, \lambda_n \in \mathbb{R}$ and a basis $\{v_1, v_2, \dots, v_n\}$ of the Euclidean space $(\mathbb{R}^n, \langle \cdot, \cdot \rangle)$ which satisfy for all $i, j \in \{1, 2, \dots, n\}$ that

$$\{\lambda \in \mathbb{R} : [\exists v \in \mathbb{R}^n \setminus \{0\} : Av = \lambda v]\} = \{\lambda_1, \lambda_2, \dots, \lambda_n\}, \quad Av_i = \lambda_i v_i, \quad \text{and} \quad \langle v_i, v_j \rangle = \begin{cases} 1 & : i = j, \\ 0 & : i \neq j \end{cases} \quad (82)$$

(cf. Definition 2.1). This demonstrates that for all $x \in \mathbb{R}^n$ there exist $\alpha_1^x, \alpha_2^x, \dots, \alpha_n^x \in \mathbb{R}$ which satisfy $x = \sum_{i=1}^n \alpha_i^x v_i$. Combining this with (82) and the fact that for all $i \in \{1, 2, \dots, n\}$ it holds that $\lambda_i \geq \min\{\lambda_1, \lambda_2, \dots, \lambda_n\} = \min\{\lambda \in \mathbb{R} : [\exists v \in \mathbb{R}^n : Av = \lambda v]\} = \lambda_{\min}(A)$ ensures that for all $x \in \mathbb{R}^n$ it holds that

$$\begin{aligned} \langle x, Ax \rangle &= \left\langle \sum_{i=1}^n \alpha_i^x v_i, A \left(\sum_{j=1}^n \alpha_j^x v_j \right) \right\rangle = \left\langle \sum_{i=1}^n \alpha_i^x v_i, \sum_{j=1}^n \alpha_j^x \lambda_j v_j \right\rangle = \sum_{i=1}^n \sum_{j=1}^n \alpha_i^x \alpha_j^x \lambda_j \langle v_i, v_j \rangle = \sum_{i=1}^n \lambda_i |\alpha_i^x|^2 \\ &\geq \lambda_{\min}(A) \sum_{i=1}^n |\alpha_i^x|^2 = \lambda_{\min}(A) \sum_{i=1}^n \sum_{j=1}^n \alpha_i^x \alpha_j^x \langle v_i, v_j \rangle = \lambda_{\min}(A) \left\langle \sum_{i=1}^n \alpha_i^x v_i, \sum_{j=1}^n \alpha_j^x v_j \right\rangle \\ &= \lambda_{\min}(A) \langle x, x \rangle = \lambda_{\min}(A) \|x\|^2 \end{aligned} \quad (83)$$

(cf. Definition 2.6). This establishes item (i). Next observe that item (i) implies for all $x \in \mathbb{R}^n$ with $\|x\| = 1$ that $\langle x, Ax \rangle \geq \lambda_{\min}(A)$. Hence, we obtain that

$$\min\{\langle x, Ax \rangle : [x \in \mathbb{R}^n, \|x\| = 1]\} \geq \lambda_{\min}(A). \quad (84)$$

In addition, note that (82) ensures that there exists $i \in \{1, 2, \dots, n\}$ which satisfies $Av_i = \lambda_{\min}(A)v_i$. Thus, we obtain that there exists $v \in \mathbb{R}^n$ which satisfies $v = v_i/\|v_i\|$, $\|v\| = 1$, and $Av = \lambda_{\min}(A)v$. This demonstrates that

$$\lambda_{\min}(A) = \lambda_{\min}(A)\|v\|^2 = \lambda_{\min}(A)\langle v, v \rangle = \langle v, \lambda_{\min}(A)v \rangle = \langle v, Av \rangle \geq \min\{\langle x, Ax \rangle : [x \in \mathbb{R}^n, \|x\| = 1]\}. \quad (85)$$

Combining this with (84) ensures that $\lambda_{\min}(A) = \min\{\langle x, Ax \rangle : [x \in \mathbb{R}^n, \|x\| = 1]\}$. This establishes item (ii). The proof of Lemma 3.20 is thus completed. \square

Lemma 3.21. *Let $n \in \mathbb{N}$ and let $A, B \in \mathbb{R}^{n \times n}$ be symmetric matrices. Then $\lambda_{\min}(A) \geq \lambda_{\min}(B) - \|A - B\|$ (cf. Definitions 2.6 and 3.13).*

Proof of Lemma 3.21. Note that the Cauchy-Schwarz inequality ensures for all $x \in \mathbb{R}^n \setminus \{0\}$ that

$$-\langle x, (A - B)x \rangle \leq |\langle x, (A - B)x \rangle| \leq \|x\| \|(A - B)x\| = \|x\|^2 \frac{\|(A - B)x\|}{\|x\|} \leq \|x\|^2 \|A - B\| \quad (86)$$

(cf. Definitions 2.1 and 3.13). Hence, we obtain for all $x \in \mathbb{R}^n$ with $\|x\| = 1$ that $\langle x, (A - B)x \rangle \geq -\|A - B\|$. This and item (i) in Lemma 3.20 (applied with $n \curvearrowright n$, $A \curvearrowright B$ in the notation of Lemma 3.20) demonstrate for all $x \in \mathbb{R}^n$ with $\|x\| = 1$ that

$$\langle x, Ax \rangle = \langle x, (A - B)x \rangle + \langle x, Bx \rangle \geq \lambda_{\min}(B) - \|A - B\|. \quad (87)$$

Combining this with item (ii) in Lemma 3.20 (applied with $n \curvearrowright n$, $A \curvearrowright A$ in the notation of Lemma 3.20) establishes that

$$\lambda_{\min}(A) = \min\{\langle Ax, x \rangle : [x \in \mathbb{R}^n, \|x\| = 1]\} \geq \lambda_{\min}(B) - \|A - B\|. \quad (88)$$

The proof of Lemma 3.21 is thus completed. \square

Lemma 3.22. *Assume Setting 2.7, assume $\lambda, \mu \in (0, \infty)$, let $\varepsilon \in (0, 1)$, $R \in (0, \infty)$, $A_1, A_2 \in \mathcal{F}$ satisfy $A_1 = \bigcap_{i,j=1}^m \{|\mathcal{G}_{i,j}(0) - \mathbf{G}_{i,j}| \leq \frac{\min\{\lambda, \mu C^{-2}\}}{4m}\}$ and $A_2 = \{\sum_{i,j=1}^m \sum_{k=1}^{\mathfrak{d}} |\mathcal{W}_k|^2 (\mathbb{1}_{\{|\langle W_k(0), x_i \rangle| \leq R\}} + \mathbb{1}_{\{|\langle W_k(0), x_j \rangle| \leq R\}})\} \leq \min\{\lambda, \mu C^{-2}\}/4\}$, let $\omega \in A_1 \cap A_2$, $n \in \mathbb{N}_0$, and assume $\max_{k \in \{1, 2, \dots, \mathfrak{d}\}} [C\|W_k(n, \omega) - W_k(0, \omega)\| + |B_k(n, \omega) - B_k(0, \omega)|] \leq R$ (cf. Definition 2.1). Then*

(i) it holds that $\lambda_{\min}(\mathcal{G}(n, \omega)) \geq \lambda/2$ and

(ii) it holds that $\lambda_{\min}(\mathcal{H}(n, \omega)) \geq \mu/2$

(cf. Definition 2.6).

Proof of Lemma 3.22. Observe that Lemma 3.21 (applied with $n \curvearrowright m$, $A \curvearrowright \mathcal{G}(n, \omega)$, $B \curvearrowright \mathbf{G}$ in the notation of Lemma 3.21), the assumption that $\lambda_{\min}(\mathbf{G}) = \lambda$, item (i) in Lemma 3.16 (applied with $\varepsilon \curvearrowright \varepsilon$, $A \curvearrowright A_1$, $\omega \curvearrowright \omega$ in the notation of Lemma 3.16), and item (i) in Lemma 3.19 (applied with $\varepsilon \curvearrowright \varepsilon$, $R \curvearrowright R$, $A \curvearrowright A_2$, $\omega \curvearrowright \omega$, $n \curvearrowright n$ in the notation of Lemma 3.16) ensure that it holds that

$$\begin{aligned} \lambda_{\min}(\mathcal{G}(n, \omega)) &\geq \lambda_{\min}(\mathbf{G}) - \|\mathcal{G}(n, \omega) - \mathbf{G}\| = \lambda - \|\mathcal{G}(n, \omega) - \mathcal{G}(0, \omega) + \mathcal{G}(0, \omega) - \mathbf{G}\| \\ &\geq \lambda - \|\mathcal{G}(n, \omega) - \mathcal{G}(0, \omega)\| - \|\mathcal{G}(0, \omega) - \mathbf{G}\| \geq \lambda - \frac{\lambda}{4} - \frac{\lambda}{4} = \frac{\lambda}{2}. \end{aligned} \quad (89)$$

This establishes item (i). Moreover, note that Lemma 3.21 (applied with $n \curvearrowright m$, $A \curvearrowright \mathcal{H}(n, \omega)$, $B \curvearrowright \mathbf{H}$ in the notation of Lemma 3.21), the assumption that $\lambda_{\min}(\mathbf{H}) = \mu$, item (ii) in Lemma 3.16 (applied with $\varepsilon \curvearrowright \varepsilon$, $A \curvearrowright A_1$, $\omega \curvearrowright \omega$ in the notation of Lemma 3.16), and item (ii) in Lemma 3.19 (applied with $\varepsilon \curvearrowright \varepsilon$, $R \curvearrowright R$, $A \curvearrowright A_2$, $\omega \curvearrowright \omega$, $n \curvearrowright n$ in the notation of Lemma 3.16) ensure that it holds that

$$\begin{aligned} \lambda_{\min}(\mathcal{H}(n, \omega)) &\geq \lambda_{\min}(\mathbf{H}) - \|\mathcal{H}(n, \omega) - \mathbf{H}\| = \mu - \|\mathcal{H}(n, \omega) - \mathcal{H}(0, \omega) + \mathcal{H}(0, \omega) - \mathbf{H}\| \\ &\geq \mu - \|\mathcal{H}(n, \omega) - \mathcal{H}(0, \omega)\| - \|\mathcal{H}(0, \omega) - \mathbf{H}\| \geq \mu - \frac{\mu}{4} - \frac{\mu}{4} = \frac{\mu}{2}. \end{aligned} \quad (90)$$

This establishes item (ii). The proof of Lemma 3.22 is thus completed. \square

4 Error analysis for GD optimization algorithms in the training of ANNs

In this section we combine the estimates of the errors at initialization, the analysis of the evolution of the weights and biases of the considered ANNs during training, and the lower bounds for the eigenvalues of the considered stochastic Gramian matrices to obtain an error analysis for the considered GD optimization algorithms. Specifically, in Proposition 4.2 below we prove a pathwise upper bound on the squared error at each iteration step on a particular measurable set. In the main result of this article, Theorem 4.3 in Section 4.2 below, we combine this result with the analysis of the probability of this measurable set in Lemma 4.1 to obtain a quantitative probabilistic error analysis for the considered GD optimization algorithms. In the remainder of Section 4.2 we progressively weaken the constraints on the width $\mathfrak{d} \in \mathbb{N}$ and the learning rate $\eta \in (0, \infty)$ and additionally consider nondegenerated input training data in order to obtain more easily accessible results. In the proof of these convergence results we use among other things the well-known upper bound on the natural logarithm in Lemma 4.5 below, whose proof we only add for reasons of completeness. We conclude Section 4 with the qualitative error analysis for the considered GD optimization algorithms which we provide in Corollary 4.10 below.

4.1 Quantitative pathwise error analysis for GD optimization algorithms

Lemma 4.1. *Assume Setting 2.7, assume $\lambda, \mu \in (0, \infty)$, let $\varepsilon \in (0, 1)$, $R \in (0, \sqrt{2\pi\varepsilon c} \min\{\lambda, \mu C^{-2}\} 2^{-4} m^{-2}]$, assume $\mathfrak{d} \geq 32 \ln(\frac{2m^2}{\varepsilon}) \max\{\frac{4m^2}{\lambda^2}, \frac{4m^2 C^4}{\mu^2}, \frac{m}{\lambda}, \frac{m C^2}{\mu}\}$, and let $A_1, A_2, A_3, A_4, A_5, A_6 \in \mathcal{F}$ satisfy*

$$\begin{aligned} A_1 &= \left\{ \|f(0) - y\|^2 \leq \left(\frac{1}{2} \sum_{i=1}^m \|x_i\|^2 + \|y\|^2\right) \varepsilon^{-1} \right\}, & A_2 &= \bigcap_{k=1}^{\mathfrak{d}} \left\{ |\mathcal{W}_k| \leq \left(\frac{2}{\mathfrak{d}} \ln\left(\frac{2\mathfrak{d}}{\varepsilon}\right)\right)^{1/2} \right\}, \\ A_3 &= \bigcap_{i,j=1}^m \left\{ |\mathcal{G}_{i,j}(0) - \mathbf{G}_{i,j}| \leq \frac{\min\{\lambda, \mu C^{-2}\}}{4m} \right\}, \\ A_4 &= \left\{ \sum_{i,j=1}^m \sum_{k=1}^{\mathfrak{d}} |\mathcal{W}_k|^2 (\mathbb{1}_{\{|\langle W_k(0), x_i \rangle| \leq R\}} + \mathbb{1}_{\{|\langle W_k(0), x_j \rangle| \leq R\}}) \leq \frac{\min\{\lambda, \mu C^{-2}\}}{4} \right\}, \\ A_5 &= \left\{ \sum_{i=1}^m \sum_{k=1}^{\mathfrak{d}} |\mathcal{W}_k|^2 \mathbb{1}_{\{|\langle W_k(0), x_i \rangle| \leq R\}} \leq \frac{2mR}{\sqrt{2\pi\varepsilon c}} \right\}, & \text{and} & & A_6 &= \left\{ \sum_{k=1}^{\mathfrak{d}} |\mathcal{W}_k|^2 \leq \frac{1}{\varepsilon} \right\} \end{aligned} \quad (91)$$

(cf. Definition 2.1). Then $\mathbb{P}(\bigcap_{i=1}^6 A_i) \geq 1 - 6\varepsilon$.

Proof of Lemma 4.1. Throughout this proof let $X_{k,i}: \Omega \rightarrow \mathbb{R}$, $k \in \{1, 2, \dots, \mathfrak{d}\}$, $i \in \{1, 2, \dots, m\}$, satisfy for all $k \in \{1, 2, \dots, \mathfrak{d}\}$, $i \in \{1, 2, \dots, m\}$ that $X_{k,i} = \langle W_k(0), x_i \rangle / \|x_i\|$. Observe that the assumption that $W_1(0), W_2(0), \dots, W_{\mathfrak{d}}(0)$ are standard normal ensures that for all $k \in \{1, 2, \dots, \mathfrak{d}\}$, $i \in \{1, 2, \dots, m\}$ it holds that $\langle W_k(0), x_i \rangle$ is a centered normal random variable with $\text{Var}[\langle W_k(0), x_i \rangle] = \|x_i\|$. Hence, we obtain for all $k \in \{1, 2, \dots, \mathfrak{d}\}$, $i \in \{1, 2, \dots, m\}$ that $X_{k,i}$ is a standard normal random variable. This and Lemma 3.17 (applied for every $k \in \{1, 2, \dots, \mathfrak{d}\}$, $i \in \{1, 2, \dots, m\}$ with $(\Omega, \mathcal{F}, \mathbb{P}) \curvearrowright (\Omega, \mathcal{F}, \mathbb{P})$, $X \curvearrowright X_{k,i}$, $\varepsilon \curvearrowright R/c$ in the notation of Lemma 3.17) ensure that for all $k \in \{1, 2, \dots, \mathfrak{d}\}$, $i \in \{1, 2, \dots, m\}$ it holds that

$$\mathbb{E}[\mathbb{1}_{\{|\langle W_k(0), x_i \rangle| \leq R\}}] = \mathbb{P}(|\langle W_k(0), x_i \rangle| \leq R) = \mathbb{P}(|X_{k,i}| \leq R/\|x_i\|) \leq \mathbb{P}(|X_{k,i}| \leq R/c) \leq \frac{2R}{\sqrt{2\pi}c}. \quad (92)$$

Moreover, note that the assumption that $W_1(0), W_2(0), \dots, W_{\mathfrak{d}}(0), \sqrt{\mathfrak{d}}W_1, \sqrt{\mathfrak{d}}W_2, \dots, \sqrt{\mathfrak{d}}W_{\mathfrak{d}}$ are independent ensures that for all $k \in \{1, 2, \dots, \mathfrak{d}\}$, $i \in \{1, 2, \dots, m\}$ it holds that \mathcal{W}_k and $\mathbb{1}_{\{|\langle W_k(0), x_i \rangle| \leq R\}}$ are independent. Combining this, (92), and the fact that for all $k \in \{1, 2, \dots, \mathfrak{d}\}$ it holds that \mathcal{W}_k is a centered normal random variable with $\text{Var}[\mathcal{W}_k] = 1/\mathfrak{d}$ establishes that

$$\mathbb{E}\left[\sum_{i=1}^m \sum_{k=1}^{\mathfrak{d}} |\mathcal{W}_k|^2 \mathbb{1}_{\{|\langle W_k(0), x_i \rangle| \leq R\}}\right] = \sum_{i=1}^m \sum_{k=1}^{\mathfrak{d}} \mathbb{E}[|\mathcal{W}_k|^2] \mathbb{E}[\mathbb{1}_{\{|\langle W_k(0), x_i \rangle| \leq R\}}] \leq \sum_{i=1}^m \sum_{k=1}^{\mathfrak{d}} \frac{1}{\mathfrak{d}} \frac{2R}{\sqrt{2\pi}c} = \frac{2mR}{\sqrt{2\pi}c}. \quad (93)$$

The Markov inequality thus demonstrates that

$$\begin{aligned} \mathbb{P}(A_5) &= \mathbb{P}\left(\sum_{i=1}^m \sum_{k=1}^{\mathfrak{d}} |\mathcal{W}_k|^2 \mathbb{1}_{\{|\langle W_k(0), x_i \rangle| \leq R\}} \leq \frac{2mR}{\sqrt{2\pi}c}\right) \geq 1 - \mathbb{P}\left(\sum_{i=1}^m \sum_{k=1}^{\mathfrak{d}} |\mathcal{W}_k|^2 \mathbb{1}_{\{|\langle W_k(0), x_i \rangle| \leq R\}} \geq \frac{2mR}{\sqrt{2\pi}c}\right) \\ &\geq 1 - \frac{\mathbb{E}[\sum_{i=1}^m \sum_{k=1}^{\mathfrak{d}} |\mathcal{W}_k|^2 \mathbb{1}_{\{|\langle W_k(0), x_i \rangle| \leq R\}}]}{2mR(\sqrt{2\pi}c)^{-1}} \geq 1 - \varepsilon. \end{aligned} \quad (94)$$

Next observe that the Markov inequality and the fact that for all $k \in \{1, 2, \dots, \mathfrak{d}\}$ it holds that \mathcal{W}_k is a centered normal random variable with $\text{Var}[\mathcal{W}_k] = 1/\mathfrak{d}$ establish that

$$\begin{aligned} \mathbb{P}(A_6) &= \mathbb{P}\left(\sum_{k=1}^{\mathfrak{d}} |\mathcal{W}_k|^2 \leq \frac{1}{\varepsilon}\right) \geq 1 - \mathbb{P}\left(\sum_{k=1}^{\mathfrak{d}} |\mathcal{W}_k|^2 \geq \frac{1}{\varepsilon}\right) \geq 1 - \frac{\mathbb{E}[\sum_{k=1}^{\mathfrak{d}} |\mathcal{W}_k|^2]}{\varepsilon^{-1}} \\ &= 1 - \varepsilon \sum_{k=1}^{\mathfrak{d}} \mathbb{E}[|\mathcal{W}_k|^2] = 1 - \varepsilon \sum_{k=1}^{\mathfrak{d}} \frac{1}{\mathfrak{d}} = 1 - \varepsilon. \end{aligned} \quad (95)$$

In addition, note that Lemma 3.2 (applied with $\varepsilon \curvearrowright \varepsilon$ in the notation of Lemma 3.2) and Lemma 3.5 (applied with $\varepsilon \curvearrowright \varepsilon$ in the notation of Lemma 3.5) ensure that $\mathbb{P}(A_1) \geq 1 - \varepsilon$ and $\mathbb{P}(A_2) \geq 1 - \varepsilon$. Moreover, observe that the assumption that $\mathfrak{d} \geq 32 \ln\left(\frac{2m^2}{\varepsilon}\right) \max\left\{\frac{4m^2}{\lambda^2}, \frac{4m^2 C^4}{\mu^2}, \frac{m}{\lambda}, \frac{mC^2}{\mu}\right\}$ and Lemma 3.15 (applied with $\varepsilon \curvearrowright \varepsilon$ in the notation of Lemma 3.15) demonstrate that $\mathbb{P}(A_3) \geq 1 - \varepsilon$. Furthermore, note that the assumption that $R \in (0, \sqrt{2\pi}c \min\{\lambda, \mu C^{-2}\} 2^{-4} m^{-2}]$ and Lemma 3.18 (applied with $\varepsilon \curvearrowright \varepsilon$, $R \curvearrowright R$ in the notation of Lemma 3.18) ensure that $\mathbb{P}(A_4) \geq 1 - \varepsilon$. Combining this, the fact that for all $i \in \{1, 2, 3\}$ it holds that $\mathbb{P}(A_i) \geq 1 - \varepsilon$, (94), and (95) establishes that for all $i \in \{1, 2, \dots, 6\}$ it holds that $\mathbb{P}(A_i) \geq 1 - \varepsilon$. Hence, we obtain that

$$\mathbb{P}\left(\bigcap_{i=1}^6 A_i\right) = 1 - \mathbb{P}\left(\bigcup_{i=1}^6 (\Omega \setminus A_i)\right) \geq 1 - \sum_{i=1}^6 \mathbb{P}(\Omega \setminus A_i) = 1 - \sum_{i=1}^6 (1 - \mathbb{P}(A_i)) \geq 1 - \sum_{i=1}^6 \varepsilon = 1 - 6\varepsilon. \quad (96)$$

This completes the proof of Lemma 4.1. \square

Proposition 4.2. *Assume Setting 2.7, assume $\lambda, \mu \in (0, \infty)$, let $\varepsilon \in (0, 1)$, $R \in (0, \infty)$ satisfy $R \leq \sqrt{2\pi}c(\lambda + \mu)(16(1 + C^2)(\sqrt{m} + m)m)^{-1}$, let $A_1, A_2, A_3, A_4, A_5, A_6 \in \mathcal{F}$ satisfy*

$$\begin{aligned} A_1 &= \left\{ \|f(0) - y\|^2 \leq \left(\frac{1}{2} \sum_{i=1}^m \|x_i\|^2 + \|y\|^2\right) \varepsilon^{-1} \right\}, & A_2 &= \bigcap_{k=1}^{\mathfrak{d}} \left\{ |\mathcal{W}_k| \leq \left(\frac{2}{\mathfrak{d}} \ln\left(\frac{2\mathfrak{d}}{\varepsilon}\right)\right)^{1/2} \right\}, \\ A_3 &= \bigcap_{i,j=1}^m \left\{ |\mathcal{G}_{i,j}(0) - \mathbf{G}_{i,j}| \leq \frac{\min\{\lambda, \mu C^{-2}\}}{4m} \right\}, \\ A_4 &= \left\{ \sum_{i,j=1}^m \sum_{k=1}^{\mathfrak{d}} |\mathcal{W}_k|^2 (\mathbb{1}_{\{|\langle W_k(0), x_i \rangle| \leq R\}} + \mathbb{1}_{\{|\langle W_k(0), x_j \rangle| \leq R\}}) \leq \frac{\min\{\lambda, \mu C^{-2}\}}{4} \right\}, \\ A_5 &= \left\{ \sum_{i=1}^m \sum_{k=1}^{\mathfrak{d}} |\mathcal{W}_k|^2 \mathbb{1}_{\{|\langle W_k(0), x_i \rangle| \leq R\}} \leq \frac{2mR}{\sqrt{2\pi}c} \right\}, & \text{and} & \quad A_6 = \left\{ \sum_{k=1}^{\mathfrak{d}} |\mathcal{W}_k|^2 \leq \frac{1}{\varepsilon} \right\}, \end{aligned} \quad (97)$$

assume $\eta < \min\left\{\frac{\lambda+\mu}{8((1+C^2)\varepsilon^{-1}+1)^2m}, \frac{m}{\lambda+\mu}\right\}$, assume

$$\frac{1}{\mathfrak{d}} \ln\left(\frac{2\mathfrak{d}}{\varepsilon}\right) \leq \frac{\varepsilon R^2(\lambda + \mu)^2}{128 \max\{1, C^4\} m \left(\frac{1}{2} \sum_{i=1}^m \|x_i\|^2 + \|y\|^2\right)}, \quad (98)$$

and let $\omega \in (\bigcap_{i=1}^{\mathfrak{d}} A_i)$ (cf. Definition 2.1). Then

(i) it holds for all $n \in \mathbb{N}_0$, $k \in \{1, 2, \dots, \mathfrak{d}\}$ that $\|W_k(n, \omega) - W_k(0, \omega)\| \leq \frac{R}{2C}$,

(ii) it holds for all $n \in \mathbb{N}_0$, $k \in \{1, 2, \dots, \mathfrak{d}\}$ that $|B_k(n, \omega) - B_k(0, \omega)| \leq \frac{R}{2}$, and

(iii) it holds for all $n \in \mathbb{N}_0$ that $\|f(n, \omega) - y\|^2 \leq (1 - \frac{\eta(\lambda+\mu)}{m})^n \|f(0, \omega) - y\|^2$.

Proof of Proposition 4.2. Throughout this proof let $I = (I_i)_{i \in \{1, 2, \dots, m\}}: \mathbb{N}_0 \rightarrow \mathbb{R}^m$, $J = (J_i)_{i \in \{1, 2, \dots, m\}}: \mathbb{N}_0 \rightarrow \mathbb{R}^m$, $P = (P_{i,j})_{(i,j) \in \{1, 2, \dots, m\}^2}: \mathbb{N}_0 \rightarrow \mathbb{R}^{m \times m}$, and $Q = (Q_{i,j})_{(i,j) \in \{1, 2, \dots, m\}^2} \in \mathbb{R}^{m \times m}$ satisfy for all $i, j \in \{1, 2, \dots, m\}$, $n \in \mathbb{N}_0$ that

$$I_i(n) = \sum_{k=1}^{\mathfrak{d}} \mathcal{W}_k(\omega) \left[\mathfrak{r}(\langle W_k(n+1, \omega), x_i \rangle + B_k(n+1, \omega)) - \mathfrak{r}(\langle W_k(n, \omega), x_i \rangle + B_k(n, \omega)) \right] \mathbb{1}_{\{|\langle W_k(0), x_i \rangle| > R\}}(\omega), \quad (99)$$

$$J_i(n) = \sum_{k=1}^{\mathfrak{d}} \mathcal{W}_k(\omega) \left[\mathfrak{r}(\langle W_k(n+1, \omega), x_i \rangle + B_k(n+1, \omega)) - \mathfrak{r}(\langle W_k(n, \omega), x_i \rangle + B_k(n, \omega)) \right] \mathbb{1}_{\{|\langle W_k(0), x_i \rangle| \leq R\}}(\omega), \quad (100)$$

$$P_{i,j}(n) = (1 + \langle x_i, x_j \rangle) \sum_{k=1}^{\mathfrak{d}} |\mathcal{W}_k(\omega)|^2 \mathbb{1}_{\{\langle W_k(n), x_i \rangle + B_k(n) \geq 0, \langle W_k(n), x_j \rangle + B_k(n) \geq 0, |\langle W_k(0), x_i \rangle| \leq R\}}(\omega), \quad (101)$$

and $Q_{i,j} = 1$ (cf. Definition 2.3). Observe that Lemma 3.14 (applied for every $n \in \mathbb{N}_0$ with $m \curvearrowright m$, $n \curvearrowright m$, $A \curvearrowright M(n)$ in the notation of Lemma 3.14), (101), the fact that for all $i, j \in \{1, 2, \dots, m\}$ it holds that $|1 + \langle x_i, x_j \rangle| \leq 1 + |\langle x_i, x_j \rangle| \leq 1 + \|x_i\| \|x_j\| \leq 1 + C^2$, and the assumption that $\omega \in A_5$ ensure that for all $n \in \mathbb{N}_0$ it holds that

$$\begin{aligned} \|P(n)\| &\leq \sum_{i=1}^m \sum_{j=1}^m |P_{i,j}(n)| \\ &\leq \sum_{i=1}^m \sum_{j=1}^m |1 + \langle x_i, x_j \rangle| \sum_{k=1}^{\mathfrak{d}} |\mathcal{W}_k(\omega)|^2 \mathbb{1}_{\{\langle W_k(n), x_i \rangle + B_k(n) \geq 0, \langle W_k(n), x_j \rangle + B_k(n) \geq 0, |\langle W_k(0), x_i \rangle| \leq R\}}(\omega) \\ &\leq (1 + C^2) \sum_{i=1}^m \sum_{j=1}^m \sum_{k=1}^{\mathfrak{d}} |\mathcal{W}_k(\omega)|^2 \mathbb{1}_{\{|\langle W_k(0), x_i \rangle| \leq R\}}(\omega) \\ &= (1 + C^2) m \sum_{i=1}^m \sum_{k=1}^{\mathfrak{d}} |\mathcal{W}_k(\omega)|^2 \mathbb{1}_{\{|\langle W_k(0), x_i \rangle| \leq R\}}(\omega) \leq \frac{2m^2(1 + C^2)R}{\sqrt{2\pi\varepsilon c}} \end{aligned} \quad (102)$$

(cf. Definition 3.13). Next observe that (8) and (9) demonstrate that for all $k \in \{1, 2, \dots, \mathfrak{d}\}$, $i \in \{1, 2, \dots, m\}$, $n \in \mathbb{N}_0$ it holds that

$$\begin{aligned} &\left[\langle W_k(n+1, \omega), x_i \rangle + B_k(n+1, \omega) \right] - \left[\langle W_k(n, \omega), x_i \rangle + B_k(n, \omega) \right] \\ &= \langle W_k(n+1, \omega) - W_k(n, \omega), x_i \rangle + B_k(n+1, \omega) - B_k(n, \omega) \\ &= \left\langle -\frac{2\eta}{m} \sum_{j=1}^m (f_j(n, \omega) - y_j) \mathcal{W}_k(\omega) \mathbb{1}_{\{\langle W_k(n), x_j \rangle + B_k(n) \geq 0\}}(\omega) x_j, x_i \right\rangle \\ &\quad - \frac{2\eta}{m} \sum_{j=1}^m (f_j(n, \omega) - y_j) \mathcal{W}_k(\omega) \mathbb{1}_{\{\langle W_k(n), x_j \rangle + B_k(n) \geq 0\}}(\omega) \\ &= -\frac{2\eta}{m} \sum_{j=1}^m (f_j(n, \omega) - y_j) \mathcal{W}_k(\omega) \mathbb{1}_{\{\langle W_k(n), x_j \rangle + B_k(n) \geq 0\}}(\omega) (1 + \langle x_i, x_j \rangle). \end{aligned} \quad (103)$$

This, the fact that $\mathbb{R} \ni x \mapsto \mathfrak{r}(x) \in \mathbb{R}$ is 1-Lipschitz continuous, the fact that for all $i, j \in \{1, 2, \dots, m\}$ it holds that $|1 + \langle x_i, x_j \rangle| \leq 1 + |\langle x_i, x_j \rangle| \leq 1 + \|x_i\| \|x_j\| \leq 1 + C^2$, and Lemma 3.3 (applied for every

$n \in \mathbb{N}_0$ with $n \curvearrowright m$, $x \curvearrowright f(n, \omega) - y$ in the notation of Lemma 3.3) establish that for all $k \in \{1, 2, \dots, \mathfrak{d}\}$, $i \in \{1, 2, \dots, m\}$, $n \in \mathbb{N}_0$ it holds that

$$\begin{aligned}
& \left| \mathfrak{r}(\langle W_k(n+1, \omega), x_i \rangle + B_k(n+1, \omega)) - \mathfrak{r}(\langle W_k(n, \omega), x_i \rangle + B_k(n, \omega)) \right| \\
& \leq \left| \left[\langle W_k(n+1, \omega), x_i \rangle + B_k(n+1, \omega) \right] - \left[\langle W_k(n, \omega), x_i \rangle + B_k(n, \omega) \right] \right| \\
& \leq \frac{2\eta}{m} \sum_{j=1}^m |(f_j(n, \omega) - y_j) \mathcal{W}_k(\omega) \mathbb{1}_{\{\langle W_k(n), x_j \rangle + B_k(n) \geq 0\}}(\omega) (1 + \langle x_i, x_j \rangle)| \\
& \leq \frac{2\eta}{m} (1 + C^2) |\mathcal{W}_k(\omega)| \sum_{j=1}^m |f_j(n, \omega) - y_j| \leq \frac{2\eta}{m} (1 + C^2) |\mathcal{W}_k(\omega)| \sqrt{m} \|f(n, \omega) - y\| \\
& = \frac{2\eta(1 + C^2)}{\sqrt{m}} |\mathcal{W}_k(\omega)| \|f(n, \omega) - y\|.
\end{aligned} \tag{104}$$

Combining this, Lemma 3.3 (applied for every $n \in \mathbb{N}_0$ with $n \curvearrowright m$, $x \curvearrowright J(n)$ in the notation of Lemma 3.3), (100), and the assumption that $\omega \in A_5$ demonstrates that for all $n \in \mathbb{N}_0$ it holds that

$$\begin{aligned}
\|J(n)\| & \leq \sum_{i=1}^m |J_i(n)| \\
& \leq \sum_{i=1}^m \sum_{k=1}^{\mathfrak{d}} |\mathcal{W}_k(\omega)| \left| \mathfrak{r}(\langle W_k(n+1, \omega), x_i \rangle + B_k(n+1, \omega)) \right. \\
& \quad \left. - \mathfrak{r}(\langle W_k(n, \omega), x_i \rangle + B_k(n, \omega)) \right| \mathbb{1}_{\{|\langle W_k(0), x_i \rangle| \leq R\}}(\omega) \\
& \leq \frac{2\eta(1 + C^2)}{\sqrt{m}} \|f(n, \omega) - y\| \sum_{i=1}^m \sum_{k=1}^{\mathfrak{d}} |\mathcal{W}_k(\omega)|^2 \mathbb{1}_{\{|\langle W_k(0), x_i \rangle| \leq R\}}(\omega) \\
& \leq \frac{2\eta(1 + C^2)}{\sqrt{m}} \|f(n, \omega) - y\| \frac{2mR}{\sqrt{2\pi\epsilon c}} = \frac{4\eta(1 + C^2)\sqrt{m}R}{\sqrt{2\pi\epsilon c}} \|f(n, \omega) - y\|.
\end{aligned} \tag{105}$$

Next we claim that for all $N \in \mathbb{N}_0$, $n \in \{0, 1, 2, \dots, N\}$, $k \in \{1, 2, \dots, \mathfrak{d}\}$ it holds that

$$\begin{aligned}
\|W_k(n, \omega) - W_k(0, \omega)\| & \leq \frac{R}{2C}, \quad |B_k(n, \omega) - B_k(0, \omega)| \leq \frac{R}{2}, \quad \text{and} \\
\|f(n, \omega) - y\|^2 & \leq \left(1 - \frac{\eta(\lambda + \mu)}{m}\right)^n \|f(0, \omega) - y\|^2.
\end{aligned} \tag{106}$$

We now prove (106) by induction on $N \in \mathbb{N}_0$. Note that for all $k \in \{1, 2, \dots, \mathfrak{d}\}$ it holds that

$$\|W_k(0, \omega) - W_k(0, \omega)\| = 0 \leq \frac{R}{2C} \quad \text{and} \quad |B_k(0, \omega) - B_k(0, \omega)| = 0 \leq \frac{R}{2}. \tag{107}$$

This and the fact that $\|f(0, \omega) - y\|^2 = \left(1 - \frac{\eta(\lambda + \mu)}{m}\right)^0 \|f(0, \omega) - y\|^2$ establish (106) in the base case $N = 0$. For the induction step $\mathbb{N}_0 \ni N \mapsto N + 1 \in \mathbb{N}$ assume for all $n \in \{0, 1, 2, \dots, N\}$, $k \in \{1, 2, \dots, \mathfrak{d}\}$ that

$$\begin{aligned}
\|W_k(n, \omega) - W_k(0, \omega)\| & \leq \frac{R}{2C}, \quad |B_k(n, \omega) - B_k(0, \omega)| \leq \frac{R}{2}, \quad \text{and} \\
\|f(n, \omega) - y\|^2 & \leq \left(1 - \frac{\eta(\lambda + \mu)}{m}\right)^n \|f(0, \omega) - y\|^2.
\end{aligned} \tag{108}$$

Observe that this, the assumption that $\eta < \frac{m}{\lambda + \mu}$, and item (i) and item (ii) in Lemma 3.6 (applied with $\varepsilon \curvearrowright \varepsilon$, $A \curvearrowright A_2$, $\omega \curvearrowright \omega$, $N \curvearrowright N$ in the notation of Lemma 3.6) ensure that for all $k \in \{1, 2, \dots, \mathfrak{d}\}$ it holds that

$$\|W_k(N+1, \omega) - W_k(0, \omega)\| \leq \frac{4C\|f(0, \omega) - y\|}{\lambda + \mu} \left(\frac{2m}{\mathfrak{d}} \ln\left(\frac{2\mathfrak{d}}{\varepsilon}\right)\right)^{1/2} \tag{109}$$

and

$$|B_k(N+1, \omega) - B_k(0, \omega)| \leq \frac{4\|f(0, \omega) - y\|}{\lambda + \mu} \left(\frac{2m}{\mathfrak{d}} \ln\left(\frac{2\mathfrak{d}}{\varepsilon}\right)\right)^{1/2}. \tag{110}$$

Moreover, the assumption that $\omega \in A_1$ and (98) ensure that it holds that

$$\begin{aligned}
& \frac{4\|f(0, \omega) - y\|}{\lambda + \mu} \left(\frac{2m}{\mathfrak{d}} \ln\left(\frac{2\mathfrak{d}}{\varepsilon}\right) \right)^{1/2} \\
& \leq \frac{4}{\lambda + \mu} \left(\left(\frac{1}{2} \sum_{i=1}^m \|x_i\|^2 + \|y\|^2 \right) \frac{2m}{\varepsilon \mathfrak{d}} \ln\left(\frac{2\mathfrak{d}}{\varepsilon}\right) \right)^{1/2} \\
& \leq \frac{4}{\lambda + \mu} \left(\left(\frac{1}{2} \sum_{i=1}^m \|x_i\|^2 + \|y\|^2 \right) \frac{2m}{\varepsilon} \frac{\varepsilon R^2 (\lambda + \mu)^2}{128 \max\{1, C^4\} m \left(\frac{1}{2} \sum_{i=1}^m \|x_i\|^2 + \|y\|^2 \right)} \right)^{1/2} \\
& = \frac{4}{\lambda + \mu} \left(\frac{R^2 (\lambda + \mu)^2}{64 \max\{1, C^4\}} \right)^{1/2} = \frac{4}{\lambda + \mu} \frac{R (\lambda + \mu)}{8 \max\{1, C^2\}} = \frac{R}{2 \max\{1, C^2\}}.
\end{aligned} \tag{111}$$

Combining this with (109) and (110) ensures that for all $k \in \{1, 2, \dots, \mathfrak{d}\}$ it holds that

$$\|W_k(N+1, \omega) - W_k(0, \omega)\| \leq \frac{4C\|f(0, \omega) - y\|}{\lambda + \mu} \left(\frac{2m}{\mathfrak{d}} \ln\left(\frac{2\mathfrak{d}}{\varepsilon}\right) \right)^{1/2} \leq C \frac{R}{2 \max\{1, C^2\}} \leq \frac{R}{2C} \tag{112}$$

and

$$|B_k(N+1, \omega) - B_k(0, \omega)| \leq \frac{4\|f(0, \omega) - y\|}{\lambda + \mu} \left(\frac{2m}{\mathfrak{d}} \ln\left(\frac{2\mathfrak{d}}{\varepsilon}\right) \right)^{1/2} \leq \frac{R}{2 \max\{1, C^2\}} \leq \frac{R}{2}. \tag{113}$$

To complete the induction step it remains to prove that $\|f(N+1, \omega) - y\| \leq \left(1 - \frac{\eta(\lambda + \mu)}{m}\right)^{N+1} \|f(0, \omega) - y\|^2$. Observe that it holds that

$$\begin{aligned}
& \|f(N+1, \omega) - y\|^2 \\
& = \sum_{i=1}^m \left(f_i(N+1, \omega) - y_i \right)^2 = \sum_{i=1}^m \left(f_i(N+1, \omega) - f_i(N, \omega) + f_i(N, \omega) - y_i \right)^2 \\
& = \sum_{i=1}^m \left[\left(f_i(N+1, \omega) - f_i(N, \omega) \right)^2 + 2 \left(f_i(N+1, \omega) - f_i(N, \omega) \right) \left(f_i(N, \omega) - y_i \right) + \left(f_i(N, \omega) - y_i \right)^2 \right] \\
& = \|f(N+1, \omega) - f(N, \omega)\|^2 + 2 \sum_{i=1}^m \left(f_i(N+1, \omega) - f_i(N, \omega) \right) \left(f_i(N, \omega) - y_i \right) + \|f(N, \omega) - y\|^2.
\end{aligned} \tag{114}$$

Moreover, note that (10) ensures that for all $i \in \{1, 2, \dots, m\}$, $n \in \mathbb{N}_0$ it holds that

$$\begin{aligned}
& f_i(n+1) - f_i(n) \\
& = (\mathcal{R}(\Phi(n+1)))(x_i) - (\mathcal{R}(\Phi(n)))(x_i) \\
& = \left[\langle \mathcal{W}, \mathfrak{r}(W(n+1)x_i + B(n+1)) \rangle + \mathfrak{B}(n+1) \right] - \left[\langle \mathcal{W}, \mathfrak{r}(W(n)x_i + B(n)) \rangle + \mathfrak{B}(n) \right] \\
& = \left[\sum_{k=1}^{\mathfrak{d}} \mathcal{W}_k \mathfrak{r}(\langle W_k(n+1), x_i \rangle + B_k(n+1)) + \mathfrak{B}(n+1) \right] - \left[\sum_{k=1}^{\mathfrak{d}} \mathcal{W}_k \mathfrak{r}(\langle W_k(n), x_i \rangle + B_k(n)) + \mathfrak{B}(n) \right] \\
& = \sum_{k=1}^{\mathfrak{d}} \mathcal{W}_k \left[\mathfrak{r}(\langle W_k(n+1), x_i \rangle + B_k(n+1)) - \mathfrak{r}(\langle W_k(n), x_i \rangle + B_k(n)) \right] + \mathfrak{B}(n+1) - \mathfrak{B}(n) \\
& = \sum_{k=1}^{\mathfrak{d}} \mathcal{W}_k \left[\mathfrak{r}(\langle W_k(n+1), x_i \rangle + B_k(n+1)) - \mathfrak{r}(\langle W_k(n), x_i \rangle + B_k(n)) \right] - \frac{2\eta}{m} \left(\sum_{j=1}^m (f_j(n) - y_j) \right)
\end{aligned} \tag{115}$$

(cf. Definition 2.4). This, the fact that $\mathbb{R} \ni x \mapsto \mathfrak{r}(x) \in \mathbb{R}$ is 1-Lipschitz continuous, (104), Lemma 3.3 (applied for every $n \in \mathbb{N}_0$ with $n \frown m$, $x \frown f(n, \omega) - y$ in the notation of Lemma 3.3), and the assumption that $\omega \in A_5$ ensure that for all $i \in \{1, 2, \dots, m\}$, $n \in \mathbb{N}_0$ it holds that

$$\begin{aligned}
& |f_i(n+1, \omega) - f_i(n, \omega)| \\
& \leq \sum_{k=1}^{\mathfrak{d}} |\mathcal{W}_k(\omega)| \left| \mathfrak{r}(\langle W_k(n+1, \omega), x_i \rangle + B_k(n+1, \omega)) - \mathfrak{r}(\langle W_k(n, \omega), x_i \rangle + B_k(n, \omega)) \right| + \frac{2\eta}{m} \sum_{j=1}^m |f_j(n, \omega) - y_j| \\
& \leq \frac{2\eta(1+C^2)}{\sqrt{m}} \|f(n, \omega) - y\| \sum_{k=1}^{\mathfrak{d}} |\mathcal{W}_k(\omega)|^2 + \frac{2\eta}{m} \sqrt{m} \|f(n, \omega) - y\| \\
& = \frac{2\eta}{\sqrt{m}} \left((1+C^2) \sum_{k=1}^{\mathfrak{d}} |\mathcal{W}_k(\omega)|^2 + 1 \right) \|f(n, \omega) - y\| \leq \frac{2\eta}{\sqrt{m}} \left(\frac{1+C^2}{\varepsilon} + 1 \right) \|f(n, \omega) - y\|.
\end{aligned} \tag{116}$$

Thus, we obtain that

$$\begin{aligned} \|f(N+1, \omega) - f(N, \omega)\|^2 &= \sum_{i=1}^m |f_i(N+1, \omega) - f_i(N, \omega)|^2 \leq \sum_{i=1}^m \frac{4\eta^2}{m} \left(\frac{1+C^2}{\varepsilon} + 1\right)^2 \|f(N, \omega) - y\|^2 \\ &= 4\eta^2 \left(\frac{1+C^2}{\varepsilon} + 1\right)^2 \|f(N, \omega) - y\|^2. \end{aligned} \quad (117)$$

Next note that (115), the fact that it holds that $1 = \mathbb{1}_{\{|\langle W_k(0), x_i \rangle| > R\}}(\omega) + \mathbb{1}_{\{|\langle W_k(0), x_i \rangle| \leq R\}}(\omega)$, (99), and (100) demonstrate that for all $i \in \{1, 2, \dots, m\}$, $n \in \mathbb{N}_0$ it holds that

$$\begin{aligned} &f_i(n+1, \omega) - f_i(n, \omega) \\ &= \sum_{k=1}^{\mathfrak{d}} \mathcal{W}_k(\omega) \left[\mathfrak{r}(\langle W_k(n+1, \omega), x_i \rangle + B_k(n+1, \omega)) - \mathfrak{r}(\langle W_k(n, \omega), x_i \rangle + B_k(n, \omega)) \right] \mathbb{1}_{\{|\langle W_k(0), x_i \rangle| > R\}}(\omega) \\ &\quad + \sum_{k=1}^{\mathfrak{d}} \mathcal{W}_k(\omega) \left[\mathfrak{r}(\langle W_k(n+1, \omega), x_i \rangle + B_k(n+1, \omega)) - \mathfrak{r}(\langle W_k(n, \omega), x_i \rangle + B_k(n, \omega)) \right] \mathbb{1}_{\{|\langle W_k(0), x_i \rangle| \leq R\}}(\omega) \\ &\quad - \frac{2\eta}{m} \sum_{j=1}^m (f_j(n, \omega) - y_j) \\ &= I_i(n) + J_i(n) - \frac{2\eta}{m} \sum_{j=1}^m (f_j(n, \omega) - y_j). \end{aligned} \quad (118)$$

Moreover, note that the assumption that $\|B(0)\| = 0$, the Cauchy Schwarz inequality, (108), (112), and (113) demonstrate that for all $n \in \{0, 1, 2, \dots, N, N+1\}$, $i \in \{1, 2, \dots, m\}$ it holds that

$$\begin{aligned} |\langle W_k(n, \omega), x_i \rangle + B_k(n, \omega) - \langle W_k(0, \omega), x_i \rangle| &= |\langle W_k(n, \omega) - W_k(0, \omega), x_i \rangle + B_k(n, \omega) - B_k(0, \omega)| \\ &\leq |\langle W_k(n, \omega) - W_k(0, \omega), x_i \rangle| + |B_k(n, \omega) - B_k(0, \omega)| \\ &\leq \|W_k(n, \omega) - W_k(0, \omega)\| \|x_i\| + |B_k(n, \omega) - B_k(0, \omega)| \\ &\leq C \|W_k(n, \omega) - W_k(0, \omega)\| + |B_k(n, \omega) - B_k(0, \omega)| \\ &\leq C \frac{R}{2C} + \frac{R}{2} = \frac{R}{2} + \frac{R}{2} = R. \end{aligned} \quad (119)$$

In addition, note that for all $a, b, z \in \mathbb{R}$ with $|a - z| < |z|$ and $|b - z| < |z|$ it holds that $\mathbb{1}_{[0, \infty)}(a) = \mathbb{1}_{[0, \infty)}(b)$. This demonstrates that for all $a, b, z \in \mathbb{R}$ with $|a - z| < |z|$ and $|b - z| < |z|$ it holds that $\mathfrak{r}(a) - \mathfrak{r}(b) = a \mathbb{1}_{[0, \infty)}(a) - b \mathbb{1}_{[0, \infty)}(b) = a \mathbb{1}_{[0, \infty)}(b) - b \mathbb{1}_{[0, \infty)}(b) = (a - b) \mathbb{1}_{[0, \infty)}(b)$. Combining this, (119), and (103) therefore demonstrates that for all $i \in \{1, 2, \dots, m\}$, $n \in \{0, 1, 2, \dots, N\}$ it holds that

$$\begin{aligned} I_i(n) &= \sum_{k=1}^{\mathfrak{d}} \mathcal{W}_k(\omega) \left[\mathfrak{r}(\langle W_k(n+1, \omega), x_i \rangle + B_k(n+1, \omega)) - \mathfrak{r}(\langle W_k(n, \omega), x_i \rangle + B_k(n, \omega)) \right] \mathbb{1}_{\{|\langle W_k(0), x_i \rangle| > R\}}(\omega) \\ &= \sum_{k=1}^{\mathfrak{d}} \mathcal{W}_k(\omega) \left[\langle W_k(n+1, \omega), x_i \rangle + B_k(n+1, \omega) \right. \\ &\quad \left. - \langle W_k(n, \omega), x_i \rangle - B_k(n, \omega) \right] \mathbb{1}_{[0, \infty)}(\langle W_k(n, \omega), x_i \rangle + B_k(n, \omega)) \mathbb{1}_{\{|\langle W_k(0), x_i \rangle| > R\}}(\omega) \\ &= \sum_{k=1}^{\mathfrak{d}} \mathcal{W}_k(\omega) \left[-\frac{2\eta}{m} \sum_{j=1}^m (f_j(n, \omega) - y_j) \mathcal{W}_k(\omega) \mathbb{1}_{\{\langle W_k(n), x_j \rangle + B_k(n) \geq 0\}}(\omega) (1 + \langle x_i, x_j \rangle) \right] \\ &\quad \times \mathbb{1}_{\{\langle W_k(n), x_i \rangle + B_k(n) \geq 0, |\langle W_k(0), x_i \rangle| > R\}}(\omega) \\ &= -\frac{2\eta}{m} \sum_{j=1}^m (f_j(n, \omega) - y_j) (1 + \langle x_i, x_j \rangle) \sum_{k=1}^{\mathfrak{d}} |\mathcal{W}_k(\omega)|^2 \mathbb{1}_{\{\langle W_k(n), x_i \rangle + B_k(n) \geq 0, \langle W_k(n), x_j \rangle + B_k(n) \geq 0, |\langle W_k(0), x_i \rangle| > R\}}(\omega) \\ &= -\frac{2\eta}{m} \sum_{j=1}^m (f_j(n, \omega) - y_j) (1 + \langle x_i, x_j \rangle) \sum_{k=1}^{\mathfrak{d}} |\mathcal{W}_k(\omega)|^2 \mathbb{1}_{\{\langle W_k(n), x_i \rangle + B_k(n) \geq 0, \langle W_k(n), x_j \rangle + B_k(n) \geq 0\}}(\omega) \\ &\quad \times \left[1 - \mathbb{1}_{\{|\langle W_k(0), x_i \rangle| \leq R\}}(\omega) \right] \\ &= -\frac{2\eta}{m} \sum_{j=1}^m (f_j(n, \omega) - y_j) (\mathcal{G}_{i,j}(n, \omega) + \mathcal{H}_{i,j}(n, \omega) - P_{i,j}(n)). \end{aligned} \quad (120)$$

Thus, we obtain that

$$2\langle I(N), f(N, \omega) - y \rangle = -\frac{4\eta}{m}\langle f(N, \omega) - y, (\mathcal{G}(N, \omega) + \mathcal{H}(N, \omega) - P(N))(f(N, \omega) - y) \rangle. \quad (121)$$

Combining this with (118) and the assumption that for all $i, j \in \{1, 2, \dots, m\}$ it holds that $Q_{i,j} = 1$ therefore establishes that

$$\begin{aligned} & 2 \sum_{i=1}^m (f_i(N+1, \omega) - f_i(N, \omega))(f_i(N, \omega) - y_i) \\ &= 2 \sum_{i=1}^m \left[I_i(N) + J_i(N) - \frac{2\eta}{m} \sum_{j=1}^m (f_j(N, \omega) - y_j) \right] (f_i(N, \omega) - y_i) \\ &= 2 \sum_{i=1}^m I_i(N)(f_i(N, \omega) - y_i) + 2 \sum_{i=1}^m J_i(N)(f_i(N, \omega) - y_i) - \frac{4\eta}{m} \sum_{i=1}^m \sum_{j=1}^m (f_j(N, \omega) - y_j)(f_i(N, \omega) - y_i) \quad (122) \\ &= 2\langle I(N), f(N, \omega) - y \rangle + 2\langle J(N), f(N, \omega) - y \rangle - \frac{4\eta}{m}\langle f(N, \omega) - y, Q(f(N, \omega) - y) \rangle \\ &= 2\langle J(N), f(N, \omega) - y \rangle - \frac{4\eta}{m}\langle f(N, \omega) - y, (\mathcal{G}(N, \omega) + \mathcal{H}(N, \omega) - P(N) + Q)(f(N, \omega) - y) \rangle. \end{aligned}$$

In addition, observe that (108) ensures that $\max_{k \in \{1, 2, \dots, \mathfrak{d}\}} [C\|W_k(N, \omega) - W_k(0, \omega)\| + |B_k(N, \omega) - B_k(0, \omega)|] \leq \max_{k \in \{1, 2, \dots, \mathfrak{d}\}} [R/2 + R/2] = R$. Item (i) and item (ii) in Lemma 3.22 (applied with $\varepsilon \curvearrowright \varepsilon$, $R \curvearrowright R$, $A_1 \curvearrowright A_3$, $A_2 \curvearrowright A_4$, $n \curvearrowright N$, $\omega \curvearrowright \omega$ in the notation of Lemma 3.22) and the assumption that $\omega \in A_3 \cap A_4$ hence demonstrate that

$$\lambda_{\min}(\mathcal{G}(N, \omega)) \geq \lambda/2 \quad \text{and} \quad \lambda_{\min}(\mathcal{H}(N, \omega)) \geq \mu/2. \quad (123)$$

Combining this, item (i) in Lemma 3.20 (applied with $n \curvearrowright m$, $A \curvearrowright \mathcal{G}(N, \omega)$, $x \curvearrowright f(N, \omega) - y$ in the notation of Lemma 3.20), item (i) in Lemma 3.20 (applied with $n \curvearrowright m$, $A \curvearrowright \mathcal{H}(N, \omega)$, $x \curvearrowright f(N, \omega) - y$ in the notation of Lemma 3.20), the Cauchy Schwarz inequality, (102), and the fact that Q is positive semidefinite ensures that

$$\begin{aligned} & \langle f(N, \omega) - y, (\mathcal{G}(N, \omega) + \mathcal{H}(N, \omega) - P(N) + Q)(f(N, \omega) - y) \rangle \\ &= \langle f(N, \omega) - y, \mathcal{G}(N, \omega)(f(N, \omega) - y) \rangle + \langle f(N, \omega) - y, \mathcal{H}(N, \omega)(f(N, \omega) - y) \rangle \\ & \quad - \langle f(N, \omega) - y, P(N)(f(N, \omega) - y) \rangle + \langle f(N, \omega) - y, Q(f(N, \omega) - y) \rangle \\ &\geq \lambda_{\min}(\mathcal{G}(N, \omega))\|f(N, \omega) - y\|^2 + \lambda_{\min}(\mathcal{H}(N, \omega))\|f(N, \omega) - y\|^2 - \|f(N, \omega) - y\|\|P(N)(f(N, \omega) - y)\| \\ &\geq \frac{\lambda}{2}\|f(N, \omega) - y\|^2 + \frac{\mu}{2}\|f(N, \omega) - y\|^2 - \|f(N, \omega) - y\|\|P(N)\| \\ &\geq \left(\frac{\lambda}{2} + \frac{\mu}{2} - \frac{2m^2(1+C^2)R}{\sqrt{2\pi\varepsilon c}} \right) \|f(N, \omega) - y\|^2. \end{aligned} \quad (124)$$

Furthermore, note that the Cauchy Schwarz inequality and (105) establish that

$$2\langle J(N), f(N, \omega) - y \rangle \leq 2\|J(N)\|\|f(N, \omega) - y\| \leq \frac{8\eta(1+C^2)\sqrt{m}R}{\sqrt{2\pi\varepsilon c}}\|f(N, \omega) - y\|^2. \quad (125)$$

Combining this, (114), (117), (122), and (124) demonstrates that

$$\begin{aligned}
& \|f(N+1, \omega) - y\|^2 \\
&= \|f(N+1, \omega) - f(N, \omega)\|^2 + 2 \sum_{i=1}^m (f_i(N+1, \omega) - f_i(N, \omega))(f_i(N, \omega) - y_i) + \|f(N, \omega) - y\|^2 \\
&= \|f(N+1, \omega) - f(N, \omega)\|^2 + 2\langle J(N), f(N, \omega) - y \rangle \\
&\quad - \frac{4\eta}{m} \langle f(N, \omega) - y, (\mathcal{G}(N, \omega) + \mathcal{H}(N, \omega) - P(N) + Q)(f(N, \omega) - y) \rangle + \|f(N, \omega) - y\|^2 \\
&\leq 4\eta^2 \left(\frac{1+C^2}{\varepsilon} + 1\right)^2 \|f(N, \omega) - y\|^2 + \frac{8\eta(1+C^2)\sqrt{m}R}{\sqrt{2\pi\varepsilon c}} \|f(N, \omega) - y\|^2 \\
&\quad - \frac{4\eta}{m} \left(\frac{\lambda}{2} + \frac{\mu}{2} - \frac{2m^2(1+C^2)R}{\sqrt{2\pi\varepsilon c}}\right) \|f(N, \omega) - y\|^2 + \|f(N, \omega) - y\|^2 \\
&= \left[4\eta^2 \left(\frac{1+C^2}{\varepsilon} + 1\right)^2 + \frac{8\eta(1+C^2)\sqrt{m}R}{\sqrt{2\pi\varepsilon c}} - \frac{2\eta(\lambda + \mu)}{m} + \frac{8\eta m(1+C^2)R}{\sqrt{2\pi\varepsilon c}} + 1\right] \|f(N, \omega) - y\|^2 \\
&= \left[1 - \frac{2\eta(\lambda + \mu)}{m} + 4\eta^2 \left(\frac{1+C^2}{\varepsilon} + 1\right)^2 + \frac{8\eta(1+C^2)(\sqrt{m} + m)R}{\sqrt{2\pi\varepsilon c}}\right] \|f(N, \omega) - y\|^2.
\end{aligned} \tag{126}$$

Moreover, observe that the assumption that $\eta \leq \frac{\lambda + \mu}{8((1+C^2)\varepsilon^{-1} + 1)^2 m}$ and $R \leq \frac{\sqrt{2\pi\varepsilon c}(\lambda + \mu)}{16(1+C^2)(\sqrt{m} + m)m}$ ensures that

$$4\eta \left(\frac{1+C^2}{\varepsilon} + 1\right)^2 + \frac{8(1+C^2)(\sqrt{m} + m)R}{\sqrt{2\pi\varepsilon c}} \leq \frac{\lambda + \mu}{2m} + \frac{\lambda + \mu}{2m} = \frac{\lambda + \mu}{m}. \tag{127}$$

Combining this with (126) therefore demonstrates that

$$\|f(N+1, \omega) - y\|^2 \leq \left[1 - \frac{2\eta(\lambda + \mu)}{m} + \frac{\eta(\lambda + \mu)}{m}\right] \|f(N, \omega) - y\|^2 = \left[1 - \frac{\eta(\lambda + \mu)}{m}\right] \|f(N, \omega) - y\|^2. \tag{128}$$

This and (108) therefore establish that $\|f(N+1, \omega) - y\|^2 \leq (1 - \frac{\eta(\lambda + \mu)}{m})^{N+1} \|f(0, \omega) - y\|^2$. Induction thus proves (106). Note that (106) implies for all $n \in \mathbb{N}_0$ that $\|f(n, \omega) - y\|^2 \leq (1 - \frac{\eta(\lambda + \mu)}{m})^n \|f(0, \omega) - y\|^2$. This establishes item (iii) and therefore completes the proof of Proposition 4.2. \square

4.2 Quantitative probabilistic error analysis for GD optimization algorithms

Theorem 4.3. *Assume Setting 2.7, assume $\lambda, \mu \in (0, \infty)$, let $\varepsilon \in (0, 1)$, assume $\eta < \min\{\frac{\lambda + \mu}{8(6(1+C^2)\varepsilon^{-1} + 1)^2 m}, \frac{m}{\lambda + \mu}\}$, $\mathfrak{d} \geq 32 \ln(\frac{12\mathfrak{d}}{\varepsilon}) \max\{\frac{4m^2}{\lambda^2}, \frac{4m^2 C^4}{\mu^2}, \frac{m}{\lambda}, \frac{mC^2}{\mu}\}$, and*

$$\frac{1}{\mathfrak{d}} \ln\left(\frac{12\mathfrak{d}}{\varepsilon}\right) \leq \frac{\pi \varepsilon^3 c^2 (\lambda + \mu)^2}{2^{17} 3^3 \max\{1, C^4\} m^3 (\frac{1}{2} \sum_{i=1}^m \|x_i\|^2 + \|y\|^2)} \min\left\{\frac{\lambda^2}{m^2}, \frac{\mu^2}{m^2 C^4}, \frac{(\lambda + \mu)^2}{(1 + C^2)^2 (\sqrt{m} + m)^2}\right\}, \tag{129}$$

and let $\mathcal{E}: \mathbf{N}_{d, \mathfrak{d}} \rightarrow [0, \infty)$ satisfy for all $\Psi \in \mathbf{N}_{d, \mathfrak{d}}$ that $\mathcal{E}(\Psi) = \frac{1}{m} \sum_{i=1}^m |(\mathcal{R}(\Psi))(x_i) - y_i|^2$ (cf. Definitions 2.1, 2.2, and 2.4). Then

$$\mathbb{P}\left(\forall n \in \mathbb{N}_0: \mathcal{E}(\Phi(n)) \leq (1 - \frac{\eta(\lambda + \mu)}{m})^n \mathcal{E}(\Phi(0))\right) \geq 1 - \varepsilon. \tag{130}$$

Proof of Theorem 4.3. Throughout this proof let $\delta \in (0, 1)$ satisfy $\delta = \varepsilon/6$, let $R \in (0, \infty)$ satisfy

$$R = \min\left\{\frac{\sqrt{2\pi\delta c} \min\{\lambda, \mu C^{-2}\}}{16m^2}, \frac{\sqrt{2\pi\delta c}(\lambda + \mu)}{16(1+C^2)(\sqrt{m} + m)m}\right\}, \tag{131}$$

and let $A_1, A_2, A_3, A_4, A_5, A_6 \in \mathcal{F}$ satisfy

$$\begin{aligned}
A_1 &= \left\{\|f(0) - y\|^2 \leq \left(\frac{1}{2} \sum_{i=1}^m \|x_i\|^2 + \|y\|^2\right) \delta^{-1}\right\}, & A_2 &= \bigcap_{k=1}^{\mathfrak{d}} \left\{|\mathcal{W}_k| \leq \left(\frac{2}{\mathfrak{d}} \ln\left(\frac{2\mathfrak{d}}{\delta}\right)\right)^{1/2}\right\}, \\
A_3 &= \bigcap_{i=1}^m \bigcap_{j=1}^m \left\{|\mathcal{G}_{i,j}(0) - \mathbf{G}_{i,j}| \leq \frac{\min\{\lambda, \mu C^{-2}\}}{4m}\right\}, \\
A_4 &= \left\{\sum_{i,j=1}^m \sum_{k=1}^{\mathfrak{d}} |\mathcal{W}_k|^2 (\mathbb{1}_{\{|\langle \mathcal{W}_k(0), x_i \rangle| \leq R\}} + \mathbb{1}_{\{|\langle \mathcal{W}_k(0), x_j \rangle| \leq R\}}) \leq \frac{\min\{\lambda, \mu C^{-2}\}}{4}\right\}, \\
A_5 &= \left\{\sum_{i=1}^m \sum_{k=1}^{\mathfrak{d}} |\mathcal{W}_k|^2 \mathbb{1}_{\{|\langle \mathcal{W}_k(0), x_i \rangle| \leq R\}} \leq \frac{2mR}{\sqrt{2\pi\delta c}}\right\}, & \text{and} & & A_6 &= \left\{\sum_{k=1}^{\mathfrak{d}} |\mathcal{W}_k|^2 \leq \frac{1}{\delta}\right\}.
\end{aligned} \tag{132}$$

First, observe that

$$\mathfrak{d} \geq 32 \ln\left(\frac{12m^2}{\varepsilon}\right) \max\left\{\frac{4m^2}{\lambda^2}, \frac{4m^2C^2}{\mu^2}, \frac{m}{\lambda}, \frac{mC^2}{\mu}\right\} = 32 \ln\left(\frac{2m^2}{\delta}\right) \max\left\{\frac{4m^2}{\lambda^2}, \frac{4m^2C^2}{\mu^2}, \frac{m}{\lambda}, \frac{mC^2}{\mu}\right\}. \quad (133)$$

Lemma 4.1 (applied with $\varepsilon \curvearrowright \delta$, $R \curvearrowright R$, $(A_i)_{i \in \{1,2,\dots,6\}} \curvearrowright (A_i)_{i \in \{1,2,\dots,6\}}$ in the notation of Lemma 4.1) and the fact that $R \leq \sqrt{2\pi}\delta c \min\{\lambda, \mu C^{-2}\} 2^{-4}m^{-2}$ therefore establish that

$$\mathbb{P}\left(\bigcap_{i=1}^6 A_i\right) \geq 1 - 6\delta = 1 - \varepsilon. \quad (134)$$

Next note that

$$\begin{aligned} R^2 &= \left(\min\left\{\frac{\sqrt{2\pi}\delta c \min\{\lambda, \mu C^{-2}\}}{16m^2}, \frac{\sqrt{2\pi}\delta c(\lambda + \mu)}{16(1+C^2)(\sqrt{m}+m)m}\right\}\right)^2 \\ &= \left(\frac{\sqrt{2\pi}\delta c}{16m} \min\left\{\frac{\lambda}{m}, \frac{\mu}{mC^2}, \frac{\lambda + \mu}{(1+C^2)(\sqrt{m}+m)}\right\}\right)^2 \\ &= \frac{\pi\delta^2 c^2}{2^7 m^2} \min\left\{\frac{\lambda^2}{m^2}, \frac{\mu^2}{m^2 C^4}, \frac{(\lambda + \mu)^2}{(1+C^2)^2(\sqrt{m}+m)^2}\right\}. \end{aligned} \quad (135)$$

Hence, we obtain that

$$\begin{aligned} \frac{1}{\mathfrak{d}} \ln\left(\frac{2\mathfrak{d}}{\delta}\right) &= \frac{1}{\mathfrak{d}} \ln\left(\frac{12\mathfrak{d}}{\varepsilon}\right) \\ &\leq \frac{\pi\varepsilon^3 c^2 (\lambda + \mu)^2}{2^{17} 3^3 \max\{1, C^4\} m^3 \left(\frac{1}{2} \sum_{i=1}^m \|x_i\|^2 + \|y\|^2\right)} \min\left\{\frac{\lambda^2}{m^2}, \frac{\mu^2}{m^2 C^4}, \frac{(\lambda + \mu)^2}{(1+C^2)^2(\sqrt{m}+m)^2}\right\} \\ &= \frac{\pi\delta^3 c^2 (\lambda + \mu)^2}{2^{14} \max\{1, C^4\} m^3 \left(\frac{1}{2} \sum_{i=1}^m \|x_i\|^2 + \|y\|^2\right)} \min\left\{\frac{\lambda^2}{m^2}, \frac{\mu^2}{m^2 C^4}, \frac{(\lambda + \mu)^2}{(1+C^2)^2(\sqrt{m}+m)^2}\right\} \\ &= \frac{\delta R^2 (\lambda + \mu)^2}{2^7 \max\{1, C^4\} m \left(\frac{1}{2} \sum_{i=1}^m \|x_i\|^2 + \|y\|^2\right)} = \frac{\delta R^2 (\lambda + \mu)^2}{128 \max\{1, C^4\} m \left(\frac{1}{2} \sum_{i=1}^m \|x_i\|^2 + \|y\|^2\right)}. \end{aligned} \quad (136)$$

Moreover, observe that

$$\eta < \min\left\{\frac{\lambda + \mu}{8(6(1+C^2)\varepsilon^{-1} + 1)^2 m}, \frac{m}{\lambda + \mu}\right\} = \min\left\{\frac{\lambda + \mu}{8((1+C^2)\delta^{-1} + 1)^2 m}, \frac{m}{\lambda + \mu}\right\}. \quad (137)$$

Combining this, (136), and the fact that $R \leq \sqrt{2\pi}\delta c(\lambda + \mu)(16(1+C^2)(\sqrt{m}+m)m)^{-1}$ with item (iii) in Proposition 4.2 (applied with $\varepsilon \curvearrowright \delta$, $R \curvearrowright R$, $(A_i)_{i \in \{1,2,\dots,6\}} \curvearrowright (A_i)_{i \in \{1,2,\dots,6\}}$ in the notation of Proposition 4.2) ensures that for all $\omega \in \bigcap_{i=1}^6 A_i$, $n \in \mathbb{N}_0$ it holds that

$$\|f(n, \omega) - y\|^2 \leq \left(1 - \frac{\eta(\lambda + \mu)}{m}\right)^n \|f(0, \omega) - y\|^2. \quad (138)$$

Thus, we obtain for all $\omega \in \bigcap_{i=1}^6 A_i$, $n \in \mathbb{N}_0$ that

$$\begin{aligned} \mathcal{E}(\Phi(n, \omega)) &= \frac{1}{m} \sum_{i=1}^m |(\mathcal{R}(\Phi(n, \omega)))(x_i) - y_i|^2 = \frac{1}{m} \sum_{i=1}^m |f_i(n, \omega) - y_i|^2 = \frac{1}{m} \|f(n, \omega) - y\|^2 \\ &\leq \frac{1}{m} \left(1 - \frac{\eta(\lambda + \mu)}{m}\right)^n \|f(0, \omega) - y\|^2 = \frac{1}{m} \left(1 - \frac{\eta(\lambda + \mu)}{m}\right)^n \sum_{i=1}^m |f_i(0, \omega) - y_i|^2 \\ &= \frac{1}{m} \left(1 - \frac{\eta(\lambda + \mu)}{m}\right)^n \sum_{i=1}^m |(\mathcal{R}(\Phi(0, \omega)))(x_i) - y_i|^2 = \left(1 - \frac{\eta(\lambda + \mu)}{m}\right)^n \mathcal{E}(\Phi(0, \omega)). \end{aligned} \quad (139)$$

Combining this and (134) establishes that

$$\mathbb{P}\left(\forall n \in \mathbb{N}_0: \mathcal{E}(\Phi(n)) \leq \left(1 - \frac{\eta(\lambda + \mu)}{m}\right)^n \mathcal{E}(\Phi(0))\right) \geq 1 - \varepsilon. \quad (140)$$

The proof of Theorem 4.3 is thus completed. \square

Corollary 4.4. Assume Setting 2.7, assume $\lambda, \mu \in (0, \infty)$, let $\varepsilon \in (0, 1)$, assume

$$\eta < \frac{\varepsilon^2 \min\{\lambda, \mu, \lambda^{-1}, \mu^{-1}\}}{1568 \max\{1, C^4\}m} \quad \text{and} \quad \frac{1}{\delta} \ln\left(\frac{12\delta}{\varepsilon}\right) \leq \frac{\pi \varepsilon^3 c^2 \min\{1, \lambda^4, \mu^4\}}{2^{22} 3^3 \max\{1, C^{16}, \|y\|^4\}m^6}, \quad (141)$$

and let $\mathcal{E}: \mathbf{N}_{d,\delta} \rightarrow [0, \infty)$ satisfy for all $\Psi \in \mathbf{N}_{d,\delta}$ that $\mathcal{E}(\Psi) = \frac{1}{m} \sum_{i=1}^m |(\mathcal{R}(\Psi))(x_i) - y_i|^2$ (cf. Definitions 2.1, 2.2, and 2.4). Then

$$\mathbb{P}\left(\forall n \in \mathbb{N}_0: \mathcal{E}(\Phi(n)) \leq \left(1 - \frac{\eta(\lambda + \mu)}{m}\right)^n \mathcal{E}(\Phi(0))\right) \geq 1 - \varepsilon. \quad (142)$$

Proof of Corollary 4.4. First, observe that

$$(1 + C^2)^2 = 1 + 2C^2 + C^4 \leq 1 + 2 \max\{1, C^2\} + C^4 \leq 1 + 2 \max\{1, C^4\} + C^4 \leq 4 \max\{1, C^4\}. \quad (143)$$

Combining this with the fact that $(1 + C^2)\varepsilon^{-1} \geq 1$ demonstrates that

$$\begin{aligned} 8(6(1 + C^2)\varepsilon^{-1} + 1)^2 &\leq 8(6(1 + C^2)\varepsilon^{-1} + (1 + C^2)\varepsilon^{-1})^2 = 8(7(1 + C^2)\varepsilon^{-1})^2 \\ &= 392(1 + C^2)^2 \varepsilon^{-2} \leq 1568 \max\{1, C^4\} \varepsilon^{-2}. \end{aligned} \quad (144)$$

Hence, we obtain that

$$\begin{aligned} \min\left\{\frac{\lambda + \mu}{8(6(1 + C^2)\varepsilon^{-1} + 1)^2 m}, \frac{m}{\lambda + \mu}\right\} &\geq \min\left\{\frac{\varepsilon^2(\lambda + \mu)}{1568 \max\{1, C^4\}m}, \frac{m}{\lambda + \mu}\right\} \\ &\geq \min\left\{\frac{\varepsilon^2 \min\{\lambda, \mu\}}{1568 \max\{1, C^4\}m}, \frac{m}{2 \max\{\lambda + \mu\}}\right\} \\ &\geq \min\left\{\frac{\varepsilon^2 \min\{\lambda, \mu\}}{1568 \max\{1, C^4\}m}, \frac{\varepsilon^2}{1568 \max\{1, C^4\}m \max\{\lambda, \mu\}}\right\} \\ &= \frac{\varepsilon^2}{1568 \max\{1, C^4\}m} \min\{\min\{\lambda, \mu\}, 1/\max\{\lambda, \mu\}\} \\ &= \frac{\varepsilon^2 \min\{\lambda, \mu, \lambda^{-1}, \mu^{-1}\}}{1568 \max\{1, C^4\}m} > \eta. \end{aligned} \quad (145)$$

Next note that (143) and the fact that $(\sqrt{m} + m)^2 \leq (2m)^2 = 4m^2$ demonstrate that

$$\begin{aligned} \min\left\{\frac{\lambda^2}{m^2}, \frac{\mu^2}{m^2 C^4}, \frac{(\lambda + \mu)^2}{(1 + C^2)^2 (\sqrt{m} + m)^2}\right\} &\geq \min\left\{\frac{\lambda^2}{m^2}, \frac{\mu^2}{m^2 C^4}, \frac{(\lambda + \mu)^2}{16 \max\{1, C^4\}m^2}\right\} \\ &\geq \min\left\{\frac{\lambda^2}{16 \max\{1, C^4\}m^2}, \frac{\mu^2}{16 \max\{1, C^4\}m^2}, \frac{(\lambda + \mu)^2}{16 \max\{1, C^4\}m^2}\right\} \\ &= \frac{\min\{\lambda^2, \mu^2, (\lambda + \mu)^2\}}{16 \max\{1, C^4\}m^2} = \frac{\min\{\lambda^2, \mu^2\}}{16 \max\{1, C^4\}m^2}. \end{aligned} \quad (146)$$

In addition, observe that

$$\frac{1}{2} \sum_{i=1}^m \|x_i\|^2 + \|y\|^2 \leq \frac{1}{2} \sum_{i=1}^m C^2 + \|y\|^2 = \frac{mC^2}{2} + \|y\|^2 = m\left(\frac{C^2}{2} + \frac{\|y\|^2}{m}\right) \leq m(C^2 + \|y\|^2) \leq 2m \max\{C^2, \|y\|^2\}. \quad (147)$$

Thus, we obtain that

$$\begin{aligned} \max\{1, C^8\} \left(\frac{1}{2} \sum_{i=1}^m \|x_i\|^2 + \|y\|^2\right) &\leq 2m \max\{1, C^8\} \max\{C^2, \|y\|^2\} \\ &\leq 2m \max\{1, C^8, \|y\|^2\} \max\{1, C^8, \|y\|^2\} \\ &= 2m \max\{1, C^{16}, \|y\|^4\}. \end{aligned} \quad (148)$$

Combining this, (146), and (141) demonstrates that

$$\begin{aligned}
& \frac{\pi \varepsilon^3 c^2 (\lambda + \mu)^2}{2^{17} 3^3 \max\{1, C^4\} m^3 (\frac{1}{2} \sum_{i=1}^m \|x_i\|^2 + \|y\|^2)} \min \left\{ \frac{\lambda^2}{m^2}, \frac{\mu^2}{m^2 C^4}, \frac{(\lambda + \mu)^2}{(1 + C^2)^2 (\sqrt{m} + m)^2} \right\} \\
& \geq \frac{\pi \varepsilon^3 c^2 (\lambda + \mu)^2}{2^{17} 3^3 \max\{1, C^4\} m^3 (\frac{1}{2} \sum_{i=1}^m \|x_i\|^2 + \|y\|^2)} \frac{\min\{\lambda^2, \mu^2\}}{16 \max\{1, C^4\} m^2} \\
& = \frac{\pi \varepsilon^3 c^2 (\lambda + \mu)^2 \min\{\lambda^2, \mu^2\}}{2^{21} 3^3 \max\{1, C^8\} m^5 (\frac{1}{2} \sum_{i=1}^m \|x_i\|^2 + \|y\|^2)} \geq \frac{\pi \varepsilon^3 c^2 \min\{\lambda^2, \mu^2\} \min\{\lambda^2, \mu^2\}}{2^{21} 3^3 \max\{1, C^8\} m^5 (\frac{1}{2} \sum_{i=1}^m \|x_i\|^2 + \|y\|^2)} \quad (149) \\
& = \frac{\pi \varepsilon^3 c^2 \min\{\lambda^4, \mu^4\}}{2^{21} 3^3 \max\{1, C^8\} m^5 (\frac{1}{2} \sum_{i=1}^m \|x_i\|^2 + \|y\|^2)} \geq \frac{\pi \varepsilon^3 c^2 \min\{\lambda^4, \mu^4\}}{2^{22} 3^3 \max\{1, C^{16}, \|y\|^4\} m^6} \\
& \geq \frac{\pi \varepsilon^3 c^2 \min\{1, \lambda^4, \mu^4\}}{2^{22} 3^3 \max\{1, C^{16}, \|y\|^4\} m^6} \geq \frac{1}{8} \ln\left(\frac{12\mathfrak{d}}{\varepsilon}\right).
\end{aligned}$$

Moreover note that (136) and the fact that for all $x \in [1, \infty)$ it holds that $\ln(x) \geq 1$ establish that

$$\mathfrak{d} \geq \ln\left(\frac{12\mathfrak{d}}{\varepsilon}\right) \frac{2^{22} 3^3 \max\{1, C^{16}, \|y\|^4\} m^6}{\pi \varepsilon^3 c^2 \min\{1, \lambda^4, \mu^4\}} \geq \frac{2^{22} 3^3 \max\{1, C^{16}, \|y\|^4\} m^6}{\pi \varepsilon^3 c^2 \min\{1, \lambda^4, \mu^4\}}. \quad (150)$$

The fact that $C^2/c^2 \geq 1$ and the fact that for all $x \in (0, \infty)$ it holds that $\ln(x) \leq x$ therefore ensure that

$$\begin{aligned}
& 32 \ln\left(\frac{12m^2}{\varepsilon}\right) \max \left\{ \frac{4m^2}{\lambda^2}, \frac{4m^2 C^2}{\mu^2}, \frac{m}{\lambda}, \frac{m C^2}{\mu} \right\} \leq 32 \left(\frac{12m^2}{\varepsilon}\right) \max \left\{ \frac{4m^2}{\lambda^2}, \frac{4m^2 C^2}{\mu^2}, \frac{m}{\lambda}, \frac{m C^2}{\mu} \right\} \\
& \leq \frac{384m^2}{\varepsilon} \max \left\{ \frac{4m^2 \max\{1, C^2\}}{\lambda^2}, \frac{4m^2 \max\{1, C^2\}}{\mu^2}, \frac{4m^2 \max\{1, C^2\}}{\lambda}, \frac{4m^2 \max\{1, C^2\}}{\mu} \right\} \\
& = \frac{1536 \max\{1, C^2\} m^4}{\varepsilon} \max \left\{ \frac{1}{\lambda^2}, \frac{1}{\mu^2}, \frac{1}{\lambda}, \frac{1}{\mu} \right\} = \frac{1536 \max\{1, C^2\} m^4}{\varepsilon \min\{\lambda^2, \mu^2, \lambda, \mu\}} \leq \frac{1536 \max\{1, C^2\} m^4}{\varepsilon \min\{1, \lambda^2, \mu^2, \lambda, \mu\}} \quad (151) \\
& = \frac{1536 \max\{1, C^2\} m^4}{\varepsilon \min\{1, \lambda^2, \mu^2\}} \leq \frac{1536 C^2 \max\{1, C^2\} m^4}{\varepsilon c^2 \min\{1, \lambda^2, \mu^2\}} \leq \frac{1536 \max\{1, C^4\} m^4}{\varepsilon c^2 \min\{1, \lambda^2, \mu^2\}} \\
& \leq \frac{2^{22} 3^3 \max\{1, C^{16}, \|y\|^4\} m^6}{\pi \varepsilon^3 c^2 \min\{1, \lambda^4, \mu^4\}} \leq m.
\end{aligned}$$

Combining this, (145), (149) and Theorem 4.3 (applied with $\varepsilon \curvearrowright \varepsilon$, $\mathcal{E} \curvearrowright \mathcal{E}$ in the notation of Theorem 4.3) establishes that

$$\mathbb{P}\left(\forall n \in \mathbb{N}_0: \mathcal{E}(\Phi(n)) \leq \left(1 - \frac{\eta(\lambda + \mu)}{m}\right)^n \mathcal{E}(\Phi(0))\right) \geq 1 - \varepsilon. \quad (152)$$

The proof of Corollary 4.4 is thus completed. \square

Lemma 4.5. *Let $\varepsilon \in (0, \infty)$. Then it holds for all $x \in (0, \infty)$ that $\ln(x) \leq x^\varepsilon \varepsilon^{-1}$.*

Proof of Lemma 4.5. Observe that for all $x \in (0, 1)$ it holds that $\ln(x) \leq 0 \leq x^\varepsilon \varepsilon^{-1}$. Moreover, note that the fundamental theorem of calculus and the fact that for all $s \in [1, \infty)$ it holds that $1 \leq s^\varepsilon$ ensure that for all $x \in [1, \infty)$ it holds that

$$\ln(x) = \ln(x) - \ln(1) = \int_1^x s^{-1} ds \leq \int_1^x s^{\varepsilon-1} ds = \left[\frac{1}{\varepsilon} s^\varepsilon\right]_1^x = \frac{1}{\varepsilon} x^\varepsilon - \frac{1}{\varepsilon} \leq \frac{1}{\varepsilon} x^\varepsilon. \quad (153)$$

The proof of Lemma 4.5 is thus completed. \square

Corollary 4.6. *Assume Setting 2.7, assume $\lambda, \mu \in (0, \infty)$, let $\varepsilon \in (0, 1)$, assume $\eta \leq 2^{-11} \min\{1, C^{-4}\} \min\{\lambda, \mu, \lambda^{-1}, \mu^{-1}\} \varepsilon^2 m^{-1}$ and $\mathfrak{d} \geq 2^{36} \max\{1, C^{20}, \|y\|^5\} c^{-5/2} \max\{1, \lambda^{-5}, \mu^{-5}\} \varepsilon^{-4} m^8$, and let $\mathcal{E}: \mathbf{N}_{d, \mathfrak{d}} \rightarrow [0, \infty)$ satisfy for all $\Psi \in \mathbf{N}_{d, \mathfrak{d}}$ that $\mathcal{E}(\Psi) = \frac{1}{m} \sum_{i=1}^m |(\mathcal{R}(\Psi))(x_i) - y_i|^2$ (cf. Definitions 2.1, 2.2, and 2.4). Then*

$$\mathbb{P}\left(\forall n \in \mathbb{N}_0: \mathcal{E}(\Phi(n)) \leq \left(1 - \frac{\eta(\lambda + \mu)}{m}\right)^n \mathcal{E}(\Phi(0))\right) \geq 1 - \varepsilon. \quad (154)$$

Proof of Corollary 4.6. First, observe that

$$\eta \leq 2^{-11} \min\{1, C^{-4}\} \min\{\boldsymbol{\lambda}, \boldsymbol{\mu}, \boldsymbol{\lambda}^{-1}, \boldsymbol{\mu}^{-1}\} \varepsilon^2 m^{-1} < \frac{\varepsilon^2 \min\{\boldsymbol{\lambda}, \boldsymbol{\mu}, \boldsymbol{\lambda}^{-1}, \boldsymbol{\mu}^{-1}\}}{1568 \max\{1, C^4\} m}. \quad (155)$$

Next note that the fact that $3^{1/5} 5 \leq 2\pi$ and the fact that $2^5 \geq 3^3$ ensure that

$$\frac{12^{1/5} 5}{2^{144/5}} = \frac{3^{1/5} 5}{2^{142/5}} \leq \frac{2\pi}{2^{142/5}} = \frac{\pi}{2^{137/5}} \leq \frac{\pi}{2^{27}} = \frac{\pi}{2^{22} 2^5} \leq \frac{\pi}{2^{22} 3^3}. \quad (156)$$

Lemma 4.5 (applied with $\varepsilon \curvearrowright 1/5$, $x \curvearrowright 12\mathfrak{d}/\varepsilon$ in the notation of Lemma 4.5) and the assumption that $\mathfrak{d} \geq 2^{36} \max\{1, C^{20}, \|y\|^5\} c^{-5/2} \max\{1, \boldsymbol{\lambda}^{-5}, \boldsymbol{\mu}^{-5}\} \varepsilon^{-4} m^8$ therefore establish that

$$\begin{aligned} \frac{1}{\mathfrak{d}} \ln\left(\frac{12\mathfrak{d}}{\varepsilon}\right) &\leq \frac{5}{\mathfrak{d}} \left(\frac{12\mathfrak{d}}{\varepsilon}\right)^{1/5} = \frac{12^{1/5} 5}{\varepsilon^{1/5} \mathfrak{d}^{4/5}} \leq \frac{12^{1/5} 5}{\varepsilon^{1/5}} \left[\frac{\varepsilon^4 c^{5/2} \min\{1, \boldsymbol{\lambda}^5, \boldsymbol{\mu}^5\}}{2^{36} \max\{1, C^{20}, \|y\|^5\} m^8} \right]^{4/5} \\ &= \frac{12^{1/5} 5}{\varepsilon^{1/5}} \left[\frac{\varepsilon^{16/5} c^2 \min\{1, \boldsymbol{\lambda}^4, \boldsymbol{\mu}^4\}}{2^{144/5} \max\{1, C^{16}, \|y\|^4\} m^{32/5}} \right] = \frac{12^{1/5} 5 \varepsilon^3 c^2 \min\{1, \boldsymbol{\lambda}^4, \boldsymbol{\mu}^4\}}{2^{144/5} \max\{1, C^{16}, \|y\|^4\} m^{32/5}} \\ &\leq \frac{\pi \varepsilon^3 c^2 \min\{1, \boldsymbol{\lambda}^4, \boldsymbol{\mu}^4\}}{2^{22} 3^3 \max\{1, C^{16}, \|y\|^4\} m^{32/5}} \leq \frac{\pi \varepsilon^3 c^2 \min\{1, \boldsymbol{\lambda}^4, \boldsymbol{\mu}^4\}}{2^{22} 3^3 \max\{1, C^{16}, \|y\|^4\} m^6}. \end{aligned} \quad (157)$$

Combining this with Corollary 4.4 (applied with $\varepsilon \curvearrowright \varepsilon$, $\mathcal{E} \curvearrowright \mathcal{E}$ in the notation of Corollary 4.4) demonstrates that

$$\mathbb{P}\left(\forall n \in \mathbb{N}_0: \mathcal{E}(\Phi(n)) \leq \left(1 - \frac{\eta(\boldsymbol{\lambda} + \boldsymbol{\mu})}{m}\right)^n \mathcal{E}(\Phi(0))\right) \geq 1 - \varepsilon. \quad (158)$$

This completes the proof of Corollary 4.6. \square

Corollary 4.7. *Assume Setting 2.7, assume for all $\lambda \in \mathbb{R}$, $i, j \in \{1, 2, \dots, m\}$ with $i \neq j$ that $x_i \neq \lambda x_j$, let $\varepsilon \in (0, 1)$, assume $\eta \leq 2^{-11} \min\{1, C^{-4}\} \min\{\boldsymbol{\lambda}, \boldsymbol{\mu}, \boldsymbol{\lambda}^{-1}, \boldsymbol{\mu}^{-1}\} \varepsilon^2 m^{-1}$ and $\mathfrak{d} \geq 2^{36} \max\{1, C^{20}, \|y\|^5\} c^{-5/2} \max\{1, \boldsymbol{\lambda}^{-5}, \boldsymbol{\mu}^{-5}\} \varepsilon^{-4} m^8$, and let $\mathcal{E}: \mathbf{N}_{d, \mathfrak{d}} \rightarrow [0, \infty)$ satisfy for all $\Psi \in \mathbf{N}_{d, \mathfrak{d}}$ that $\mathcal{E}(\Psi) = \frac{1}{m} \sum_{i=1}^m |(\mathcal{R}(\Psi))(x_i) - y_i|^2$ (cf. Definitions 2.1, 2.2, and 2.4). Then*

$$\mathbb{P}\left(\forall n \in \mathbb{N}_0: \mathcal{E}(\Phi(n)) \leq \left(1 - \frac{\eta(\boldsymbol{\lambda} + \boldsymbol{\mu})}{m}\right)^n \mathcal{E}(\Phi(0))\right) \geq 1 - \varepsilon. \quad (159)$$

Proof of Corollary 4.7. Note that Lemma 2.11 (applied with $d \curvearrowright d$, $m \curvearrowright m$, $(x_i)_{i \in \{1, 2, \dots, m\}} \curvearrowright (x_i)_{i \in \{1, 2, \dots, m\}}$, $(\Omega, \mathcal{F}, \mathbb{P}) \curvearrowright (\Omega, \mathcal{F}, \mathbb{P})$, $W \curvearrowright W_1(0)$, $\mathbf{G} \curvearrowright \mathbf{G}$, $\mathbf{H} \curvearrowright \mathbf{H}$, $\boldsymbol{\lambda} \curvearrowright \boldsymbol{\lambda}$, $\boldsymbol{\mu} \curvearrowright \boldsymbol{\mu}$ in the notation of Lemma 2.11) and the assumption that for all $\lambda \in \mathbb{R}$, $i, j \in \{1, 2, \dots, m\}$ with $i \neq j$ it holds that $x_i \neq \lambda x_j$ ensure that it holds that $\boldsymbol{\lambda}, \boldsymbol{\mu} \in (0, \infty)$. Combining this with Corollary 4.6 (applied with $\varepsilon \curvearrowright \varepsilon$, $\mathcal{E} \curvearrowright \mathcal{E}$ in the notation of Corollary 4.6) establishes that

$$\mathbb{P}\left(\forall n \in \mathbb{N}_0: \mathcal{E}(\Phi(n)) \leq \left(1 - \frac{\eta(\boldsymbol{\lambda} + \boldsymbol{\mu})}{m}\right)^n \mathcal{E}(\Phi(0))\right) \geq 1 - \varepsilon. \quad (160)$$

This completes the proof of Corollary 4.7. \square

Corollary 4.8. *Assume Setting 2.7, assume for all $\lambda \in \mathbb{R}$, $i, j \in \{1, 2, \dots, m\}$ with $i \neq j$ that $x_i \neq \lambda x_j$, let $\Lambda \in \mathbb{R}$, $\varepsilon \in (0, 1)$ satisfy*

$$\Lambda = \min\left\{ \frac{\boldsymbol{\lambda} + \boldsymbol{\mu}}{m}, \frac{\min\{\boldsymbol{\lambda}, \boldsymbol{\mu}, \boldsymbol{\lambda}^{-1}, \boldsymbol{\mu}^{-1}\}}{2^{11} \max\{1, C^4\} m}, \frac{c^{5/2} \min\{1, \boldsymbol{\lambda}^5, \boldsymbol{\mu}^5\}}{2^{36} \max\{1, C^{20}, \|y\|^5\} m^8} \right\}, \quad (161)$$

assume $\eta \leq \Lambda \varepsilon^2$ and $\mathfrak{d} \geq \Lambda^{-1} \varepsilon^{-4}$, and let $\mathcal{E}: \mathbf{N}_{d, \mathfrak{d}} \rightarrow [0, \infty)$ satisfy for all $\Psi \in \mathbf{N}_{d, \mathfrak{d}}$ that $\mathcal{E}(\Psi) = \frac{1}{m} \sum_{i=1}^m |(\mathcal{R}(\Psi))(x_i) - y_i|^2$ (cf. Definitions 2.1, 2.2, and 2.4). Then

$$\mathbb{P}\left(\forall n \in \mathbb{N}_0: \mathcal{E}(\Phi(n)) \leq (1 - \eta \Lambda)^n \mathcal{E}(\Phi(0))\right) \geq 1 - \varepsilon. \quad (162)$$

Proof of Corollary 4.8. Note that Lemma 2.11 (applied with $d \curvearrowright d$, $m \curvearrowright m$, $(x_i)_{i \in \{1,2,\dots,m\}} \curvearrowright (x_i)_{i \in \{1,2,\dots,m\}}$, $(\Omega, \mathcal{F}, \mathbb{P}) \curvearrowright (\Omega, \mathcal{F}, \mathbb{P})$, $W \curvearrowright W_1(0)$, $\mathbf{G} \curvearrowright \mathbf{G}$, $\mathbf{H} \curvearrowright \mathbf{H}$, $\boldsymbol{\lambda} \curvearrowright \boldsymbol{\lambda}$, $\boldsymbol{\mu} \curvearrowright \boldsymbol{\mu}$ in the notation of Lemma 2.11) and the assumption that for all $\lambda \in \mathbb{R}$, $i, j \in \{1, 2, \dots, m\}$ with $i \neq j$ it holds that $x_i \neq \lambda x_j$ ensure that it holds that $\boldsymbol{\lambda}, \boldsymbol{\mu} \in (0, \infty)$. Thus, we obtain that

$$\frac{\boldsymbol{\lambda} + \boldsymbol{\mu}}{m} \in (0, \infty), \quad \frac{\min\{\boldsymbol{\lambda}, \boldsymbol{\mu}, \boldsymbol{\lambda}^{-1}, \boldsymbol{\mu}^{-1}\}}{2^{11} \max\{1, C^4\}m} \in (0, 1), \quad \text{and} \quad \frac{c^{5/2} \min\{1, \boldsymbol{\lambda}^5, \boldsymbol{\mu}^5\}}{2^{36} \max\{1, C^{20}, \|y\|^5\}m^8} \in (0, \infty). \quad (163)$$

Combining this with (161) establishes that $\Lambda \in (0, 1)$. Next note that (161), the assumption that $\eta \leq \Lambda \varepsilon^2$, and the assumption that $\mathfrak{d} \geq \Lambda^{-1} \varepsilon^{-4}$ ensure that

$$\eta \leq \Lambda \varepsilon^2 \leq \frac{\min\{\boldsymbol{\lambda}, \boldsymbol{\mu}, \boldsymbol{\lambda}^{-1}, \boldsymbol{\mu}^{-1}\}}{2^{11} \max\{1, C^4\}m} \varepsilon^2 = 2^{-11} \min\{1, C^{-4}\} \min\{\boldsymbol{\lambda}, \boldsymbol{\mu}, \boldsymbol{\lambda}^{-1}, \boldsymbol{\mu}^{-1}\} \varepsilon^2 m^{-1}, \quad (164)$$

and

$$\mathfrak{d} \geq \Lambda^{-1} \varepsilon^{-4} \geq \left[\frac{c^{5/2} \min\{1, \boldsymbol{\lambda}^5, \boldsymbol{\mu}^5\}}{2^{36} \max\{1, C^{20}, \|y\|^5\}m^8} \right]^{-1} \varepsilon^{-4} = 2^{36} \max\{1, C^{20}, \|y\|^5\} c^{-5/2} \max\{1, \boldsymbol{\lambda}^{-5}, \boldsymbol{\mu}^{-5}\} \varepsilon^{-4} m^8. \quad (165)$$

Corollary 4.7 (applied with $\varepsilon \curvearrowright \varepsilon$, $\mathcal{E} \curvearrowright \mathcal{E}$ in the notation of Corollary 4.7) hence establishes that

$$\mathbb{P}(\forall n \in \mathbb{N}_0: \mathcal{E}(\Phi(n)) \leq (1 - \frac{\eta(\boldsymbol{\lambda} + \boldsymbol{\mu})}{m})^n \mathcal{E}(\Phi(0))) \geq 1 - \varepsilon. \quad (166)$$

Combining this with the fact that $\Lambda \leq \frac{\boldsymbol{\lambda} + \boldsymbol{\mu}}{m}$ demonstrates that

$$\mathbb{P}(\forall n \in \mathbb{N}_0: \mathcal{E}(\Phi(n)) \leq (1 - \eta \Lambda)^n \mathcal{E}(\Phi(0))) \geq \mathbb{P}(\forall n \in \mathbb{N}_0: \mathcal{E}(\Phi(n)) \leq (1 - \frac{\eta(\boldsymbol{\lambda} + \boldsymbol{\mu})}{m})^n \mathcal{E}(\Phi(0))) \geq 1 - \varepsilon. \quad (167)$$

The proof of Corollary 4.8 is thus completed. \square

4.3 Qualitative probabilistic error analysis for GD optimization algorithms

Lemma 4.9. *Let $\varphi_t: \mathbb{R} \rightarrow \mathbb{R}$, $t \in (0, \infty)$, satisfy for all $t \in (0, \infty)$, $z \in \mathbb{R}$ that $\varphi_t(z) = \frac{1}{t} \ln(1 + t \exp(tz))$. Then*

- (i) *it holds for all $z \in \mathbb{R}$ that $\lim_{t \rightarrow \infty} \varphi_t(z) = \max\{z, 0\}$,*
- (ii) *it holds for all $t \in (0, \infty)$ that φ_t is differentiable, and*
- (iii) *it holds for all $z \in \mathbb{R}$ that $\lim_{t \rightarrow \infty} (\varphi_t)'(z) = \mathbb{1}_{[0, \infty)}(z)$.*

Proof of Lemma 4.9. Note that for all $t \in (0, \infty)$, $z \in \mathbb{R}$ it holds that

$$\begin{aligned} \varphi_t(z) &= \frac{1}{t} \ln(1 + t \exp(tz)) = \frac{1}{t} \ln(t \exp(tz) (\frac{1}{t} \exp(-tz) + 1)) \\ &= \frac{1}{t} \left[\ln(t) + tz + \ln(\frac{1}{t} \exp(-tz) + 1) \right] = z + \frac{\ln(t)}{t} + \frac{1}{t} \ln(\frac{1}{t} \exp(-tz) + 1). \end{aligned} \quad (168)$$

Combining this, the fact that for all $z \in [0, \infty)$ it holds that $\lim_{t \rightarrow \infty} \frac{1}{t} \exp(-tz) = 0$, and the fact that $\lim_{t \rightarrow \infty} \frac{\ln(t)}{t} = 0$ ensures that for all $z \in [0, \infty)$ it holds that

$$\lim_{t \rightarrow \infty} \varphi_t(z) = \lim_{t \rightarrow \infty} \left[z + \frac{\ln(t)}{t} + \frac{1}{t} \ln(\frac{1}{t} \exp(-tz) + 1) \right] = z. \quad (169)$$

Moreover, observe that the fact that for all $z \in (-\infty, 0)$ it holds that $\lim_{t \rightarrow \infty} t \exp(tz) = 0$ establishes that for all $z \in (-\infty, 0)$ it holds that

$$\lim_{t \rightarrow \infty} \varphi_t(z) = \lim_{t \rightarrow \infty} \left[\frac{1}{t} \ln(1 + t \exp(tz)) \right] = 0. \quad (170)$$

Combining this and (169) demonstrates that for all $z \in \mathbb{R}$ it holds that $\lim_{t \rightarrow \infty} \varphi_t(z) = \max\{z, 0\}$. This establishes item (i). Next note that the fact that for all $t \in (0, \infty)$ it holds that φ_t is a composition of

differentiable functions implies that for all $t \in (0, \infty)$ it holds that φ_t is differentiable at all $z \in \mathbb{R}$. This establishes item (ii). Observe that for all $t \in (0, \infty)$, $z \in \mathbb{R}$ it holds that

$$(\varphi_t)'(z) = \frac{1}{t} \frac{t^2 \exp(tz)}{1 + t \exp(tz)} = \frac{t \exp(tz)}{1 + \exp(tz)} = \frac{1}{1 + t^{-1} \exp(-tz)}. \quad (171)$$

Combining this with the fact that for all $z \in (-\infty, 0)$ it holds that $\lim_{t \rightarrow \infty} t^{-1} \exp(-tz) = \infty$ and the fact that for all $z \in [0, \infty)$ it holds that $\lim_{t \rightarrow \infty} t^{-1} \exp(-tz) = 0$ ensures that for all $z \in \mathbb{R}$ it holds that

$$\lim_{t \rightarrow \infty} (\varphi_t)'(z) = \mathbb{1}_{[0, \infty)}(z). \quad (172)$$

This establishes item (iii). The proof of Lemma 4.9 is thus completed. \square

Corollary 4.10. *Let $d, m \in \mathbb{N}$, $x_1, x_2, \dots, x_m \in \mathbb{R}^d \setminus \{0\}$, $y_1, y_2, \dots, y_m \in \mathbb{R}$ satisfy for all $\lambda \in \mathbb{R}$, $i, j \in \{1, 2, \dots, m\}$ with $i \neq j$ that $x_i \neq \lambda x_j$, let $\varphi_t: \mathbb{R} \rightarrow \mathbb{R}$, $t \in \mathbb{N}$, and $\mathcal{P}: \mathbb{R} \rightarrow \mathbb{N}$ satisfy for all $t, \mathfrak{d} \in \mathbb{N}$, $z \in \mathbb{R}$ that $\varphi_t(z) = \frac{1}{t} \ln(1 + t \exp(tz))$ and $\mathcal{P}(\mathfrak{d}) \geq \mathfrak{d}d + 2\mathfrak{d} + 1$, let $\mathcal{N}_\theta^{\mathfrak{d}, t}: \mathbb{R}^d \rightarrow \mathbb{R}$, $\mathfrak{d}, t \in \mathbb{N}$, $\theta \in \mathbb{R}^{\mathcal{P}(\mathfrak{d})}$, and $\mathcal{E}^{\mathfrak{d}, t}: \mathbb{R}^{\mathcal{P}(\mathfrak{d})} \rightarrow [0, \infty)$, $\mathfrak{d}, t \in \mathbb{N}_0$, satisfy for all $\mathfrak{d}, t \in \mathbb{N}$, $\theta = (\theta_1, \theta_2, \dots, \theta_{\mathcal{P}(\mathfrak{d})}) \in \mathbb{R}^{\mathcal{P}(\mathfrak{d})}$, $z = (z_1, z_2, \dots, z_d) \in \mathbb{R}^d$ that $\mathcal{N}_\theta^{\mathfrak{d}, t}(z) = \sum_{k=1}^{\mathfrak{d}} \theta_{\mathfrak{d}d+\mathfrak{d}+k} \varphi_t(\sum_{\ell=1}^d \theta_{(k-1)d+\ell} z_\ell + \theta_{\mathfrak{d}d+k}) + \theta_{\mathfrak{d}d+2\mathfrak{d}+1}$ and $\mathcal{E}^{\mathfrak{d}, t}(\theta) = \frac{1}{m} \sum_{i=1}^m |\mathcal{N}_\theta^{\mathfrak{d}, t}(x_i) - y_i|^2$, let $(\Omega, \mathcal{F}, \mathbb{P})$ be a probability space, let $\Theta^{\mathfrak{d}, \eta} = (\Theta_1^{\mathfrak{d}, \eta}, \Theta_2^{\mathfrak{d}, \eta}, \dots, \Theta_{\mathcal{P}(\mathfrak{d})}^{\mathfrak{d}, \eta}): \mathbb{N}_0 \times \Omega \rightarrow \mathbb{R}^{\mathcal{P}(\mathfrak{d})}$, $\mathfrak{d}, \eta \in \mathbb{R}$, be measurable, assume for all $\mathfrak{d}, \eta \in \mathbb{R}$ that $\Theta_1^{\mathfrak{d}, \eta}(0), \Theta_2^{\mathfrak{d}, \eta}(0), \dots, \Theta_{\mathfrak{d}}^{\mathfrak{d}, \eta}(0), \sqrt{\mathfrak{d}} \Theta_{\mathfrak{d}d+\mathfrak{d}+1}^{\mathfrak{d}, \eta}(0), \sqrt{\mathfrak{d}} \Theta_{\mathfrak{d}d+\mathfrak{d}+2}^{\mathfrak{d}, \eta}(0), \dots, \sqrt{\mathfrak{d}} \Theta_{\mathfrak{d}d+2\mathfrak{d}}^{\mathfrak{d}, \eta}(0)$ are independent and standard normal, and assume for all $\mathfrak{d}, n \in \mathbb{N}_0$, $\eta \in \mathbb{R}$, $i \in \{1, 2, \dots, \mathcal{P}(\mathfrak{d})\}$ that $\Theta_{\mathfrak{d}d+1}^{\mathfrak{d}, \eta}(0) = \Theta_{\mathfrak{d}d+2}^{\mathfrak{d}, \eta}(0) = \dots = \Theta_{\mathfrak{d}d+\mathfrak{d}}^{\mathfrak{d}, \eta}(0) = \Theta_{\mathfrak{d}d+2\mathfrak{d}+1}^{\mathfrak{d}, \eta}(0) = 0$ and $\Theta_i^{\mathfrak{d}, \eta}(n+1) = \Theta_i^{\mathfrak{d}, \eta}(n) - \eta [\lim_{t \rightarrow \infty} (\frac{\partial}{\partial \theta_i} \mathcal{E}^{\mathfrak{d}, t})(\Theta^{\mathfrak{d}, \eta}(n))] \mathbb{1}_{\mathbb{R} \setminus (\mathfrak{d}d+\mathfrak{d}, \mathfrak{d}d+2\mathfrak{d})}(i)$. Then there exists $\lambda \in (0, 1)$ such that for all $\varepsilon \in (0, 1)$, $\eta \in (0, \lambda \varepsilon^2)$, $\mathfrak{d} \in \mathbb{N} \cap [\lambda^{-1} \varepsilon^{-4}, \infty)$ it holds that*

$$\mathbb{P}(\forall n \in \mathbb{N}_0: [\lim_{t \rightarrow \infty} \mathcal{E}^{\mathfrak{d}, t}(\Theta^{\mathfrak{d}, \eta}(n))] \leq (1 - \eta \lambda)^n [\lim_{t \rightarrow \infty} \mathcal{E}^{\mathfrak{d}, t}(\Theta^{\mathfrak{d}, \eta}(0))]) \geq 1 - \varepsilon. \quad (173)$$

Proof of Corollary 4.10. Throughout this proof let $Z: \Omega \rightarrow \mathbb{R}^d$ satisfy $Z = (\Theta_1^{0,0}(0), \Theta_2^{0,0}(0), \dots, \Theta_d^{0,0}(0))$, let $\mathbf{G} = (\mathbf{G}_{i,j})_{(i,j) \in \{1,2,\dots,m\}^2}$, $\mathbf{H} = (\mathbf{H}_{i,j})_{(i,j) \in \{1,2,\dots,m\}^2} \in \mathbb{R}^{m \times m}$ satisfy for all $i, j \in \{1, 2, \dots, m\}$ that $\mathbf{G}_{i,j} = \mathbb{P}(\langle Z, x_i \rangle \geq 0, \langle Z, x_j \rangle \geq 0)$ and $\mathbf{H}_{i,j} = \langle x_i, x_j \rangle \mathbf{G}_{i,j}$ (cf. Definition 2.1), let $c, C \in \mathbb{R}$, $e_1 = (1, 0, \dots, 0)$, $e_2 = (0, 1, 0, \dots, 0)$, \dots , $e_d = (0, \dots, 0, 1) \in \mathbb{R}^d$ satisfy $c = \min_{i \in \{1,2,\dots,m\}} \|x_i\|$ and $C = \max_{i \in \{1,2,\dots,m\}} \|x_i\|$, let $\lambda, \mu \in [0, \infty)$ satisfy $\lambda = \lambda_{\min}(\mathbf{G})$ and $\mu = \lambda_{\min}(\mathbf{H})$ (cf. Definition 2.6), let $\Lambda \in \mathbb{R}$ satisfy

$$\Lambda = \min \left\{ \frac{\lambda + \mu}{m}, \frac{\min\{\lambda, \mu, \lambda^{-1}, \mu^{-1}\}}{2^{11} \max\{1, C^4\} m}, \frac{c^{5/2} \min\{1, \lambda^5, \mu^5\}}{2^{36} \max\{1, C^{20}, \|y\|^5\} m^8} \right\}, \quad (174)$$

let $\varepsilon \in (0, 1)$, $\eta \in (0, \Lambda \varepsilon^2)$, $\mathfrak{d} \in \mathbb{N} \cap [\Lambda^{-1} \varepsilon^{-4}, \infty)$, let $W = (W_1, W_2, \dots, W_{\mathfrak{d}}): \mathbb{N}_0 \times \Omega \rightarrow \mathbb{R}^{\mathfrak{d} \times d}$, $B = (B_1, B_2, \dots, B_{\mathfrak{d}}): \mathbb{N}_0 \times \Omega \rightarrow \mathbb{R}^{\mathfrak{d}}$, $\mathcal{W} = (\mathcal{W}_1, \mathcal{W}_2, \dots, \mathcal{W}_{\mathfrak{d}}): \Omega \rightarrow \mathbb{R}^{\mathfrak{d}}$, and $\mathfrak{B}: \mathbb{N}_0 \times \Omega \rightarrow \mathbb{R}$ satisfy for all $n \in \mathbb{N}_0$, $\omega \in \Omega$ that

$$W(n, \omega) = \begin{pmatrix} \Theta_1^{\mathfrak{d}, \eta}(n, \omega) & \Theta_2^{\mathfrak{d}, \eta}(n, \omega) & \dots & \Theta_d^{\mathfrak{d}, \eta}(n, \omega) \\ \Theta_{\mathfrak{d}d+1}^{\mathfrak{d}, \eta}(n, \omega) & \Theta_{\mathfrak{d}d+2}^{\mathfrak{d}, \eta}(n, \omega) & \dots & \Theta_{2\mathfrak{d}}^{\mathfrak{d}, \eta}(n, \omega) \\ \vdots & \vdots & \ddots & \vdots \\ \Theta_{(\mathfrak{d}-1)d+1}^{\mathfrak{d}, \eta}(n, \omega) & \Theta_{(\mathfrak{d}-1)d+2}^{\mathfrak{d}, \eta}(n, \omega) & \dots & \Theta_{\mathfrak{d}d}^{\mathfrak{d}, \eta}(n, \omega) \end{pmatrix}, \quad B(n, \omega) = \begin{pmatrix} \Theta_{\mathfrak{d}d+1}^{\mathfrak{d}, \eta}(n, \omega) \\ \Theta_{\mathfrak{d}d+2}^{\mathfrak{d}, \eta}(n, \omega) \\ \vdots \\ \Theta_{\mathfrak{d}d+\mathfrak{d}}^{\mathfrak{d}, \eta}(n, \omega) \end{pmatrix}, \quad (175)$$

$\mathcal{W}(\omega) = (\Theta_{\mathfrak{d}d+\mathfrak{d}+1}^{\mathfrak{d}, \eta}(0, \omega), \Theta_{\mathfrak{d}d+\mathfrak{d}+2}^{\mathfrak{d}, \eta}(0, \omega), \dots, \Theta_{\mathfrak{d}d+2\mathfrak{d}}^{\mathfrak{d}, \eta}(0, \omega))$, and $\mathfrak{B}(n, \omega) = \Theta_{\mathfrak{d}d+2\mathfrak{d}+1}^{\mathfrak{d}, \eta}(n, \omega)$, let $\Phi: \mathbb{N}_0 \times \Omega \rightarrow \mathbf{N}_{d, \mathfrak{d}}$ and $f = (f_1, f_2, \dots, f_m): \mathbb{N}_0 \times \Omega \rightarrow \mathbb{R}^m$ satisfy for all $i \in \{1, 2, \dots, m\}$, $n \in \mathbb{N}_0$, $\omega \in \Omega$ that $\Phi(n, \omega) = ((W(n, \omega), B(n, \omega)), (\mathcal{W}(\omega), \mathfrak{B}(n, \omega)))$ and $f_i(n, \omega) = (\mathcal{R}(\Phi(n, \omega)))(x_i)$ (cf. Definitions 2.2 and 2.4), and let $\mathfrak{E}: \mathbf{N}_{d, \mathfrak{d}} \rightarrow [0, \infty)$ satisfy for all $\Psi \in \mathbf{N}_{d, \mathfrak{d}}$ that $\mathfrak{E}(\Psi) = \frac{1}{m} \sum_{i=1}^m |(\mathcal{R}(\Psi))(x_i) - y_i|^2$. First, observe that the assumption that $\Theta_i^{0,0}(0): \Omega \rightarrow \mathbb{R}$, $i \in \{1, 2, \dots, d\}$, are independent and standard normal random variables implies that Z is standard normal. Next note that Lemma 2.11 and the assumption that for all $\lambda \in \mathbb{R}$, $i, j \in \{1, 2, \dots, n\}$ with $i \neq j$ it holds that $x_i \neq \lambda x_j$ ensure that it holds that $\lambda, \mu \in (0, \infty)$. Thus, we obtain that $\Lambda \in (0, 1)$. In addition observe that the assumption that $\Theta_1^{\mathfrak{d}, \eta}(0), \Theta_2^{\mathfrak{d}, \eta}(0), \dots, \Theta_{\mathfrak{d}}^{\mathfrak{d}, \eta}(0), \sqrt{\mathfrak{d}} \Theta_{\mathfrak{d}d+\mathfrak{d}+1}^{\mathfrak{d}, \eta}(0), \sqrt{\mathfrak{d}} \Theta_{\mathfrak{d}d+\mathfrak{d}+2}^{\mathfrak{d}, \eta}(0), \dots, \sqrt{\mathfrak{d}} \Theta_{\mathfrak{d}d+2\mathfrak{d}}^{\mathfrak{d}, \eta}(0)$ are independent and standard normal establishes that $W_1(0), W_2(0), \dots, W_{\mathfrak{d}}(0), \sqrt{\mathfrak{d}} \mathcal{W}_1, \sqrt{\mathfrak{d}} \mathcal{W}_2, \dots, \sqrt{\mathfrak{d}} \mathcal{W}_{\mathfrak{d}}$ are independent and standard normal.

Moreover, note that the assumption that $\Theta_{\mathfrak{d}d+1}^{\mathfrak{d},\eta}(0) = \Theta_{\mathfrak{d}d+2}^{\mathfrak{d},\eta}(0) = \dots = \Theta_{\mathfrak{d}d+\mathfrak{d}}^{\mathfrak{d},\eta}(0) = \Theta_{\mathfrak{d}d+2\mathfrak{d}+1}^{\mathfrak{d},\eta}(0) = 0$ implies that $\|B(0)\| = |\mathfrak{B}(0)| = 0$. Next note that for all $i \in \{1, 2, \dots, \mathfrak{P}(\mathfrak{d})\}$, $n \in \mathbb{N}_0$, $t \in \mathbb{N}$ it holds that

$$\frac{\partial}{\partial \theta_i} \mathcal{E}^{\mathfrak{d},t}(\Theta^{\mathfrak{d},\eta}(n)) = \frac{\partial}{\partial \theta_i} \left(\frac{1}{m} \sum_{p=1}^m |\mathcal{N}_{\Theta^{\mathfrak{d},\eta}(n)}^{\mathfrak{d},t}(x_p) - y_p|^2 \right) = \frac{2}{m} \sum_{p=1}^m (\mathcal{N}_{\Theta^{\mathfrak{d},\eta}(n)}^{\mathfrak{d},t}(x_p) - y_p) \left(\frac{\partial}{\partial \theta_i} \mathcal{N}_{\Theta^{\mathfrak{d},\eta}(n)}^{\mathfrak{d},t}(x_p) \right). \quad (176)$$

Moreover, note that item (i) in Lemma 4.9 and the fact that for all $k \in \{1, 2, \dots, \mathfrak{d}\}$, $n \in \mathbb{N}_0$ it holds that $\Theta_{\mathfrak{d}d+\mathfrak{d}+k}^{\mathfrak{d},\eta}(n) = \Theta_{\mathfrak{d}d+\mathfrak{d}+k}^{\mathfrak{d},\eta}(0) = \mathcal{W}_k$ demonstrates that for all $n \in \mathbb{N}_0$, $z = (z_1, z_2, \dots, z_d) \in \mathbb{R}^d$ it holds that

$$\begin{aligned} \lim_{t \rightarrow \infty} \mathcal{N}_{\Theta^{\mathfrak{d},\eta}(n)}^{\mathfrak{d},t}(z) &= \lim_{t \rightarrow \infty} \left[\sum_{k=1}^{\mathfrak{d}} \Theta_{\mathfrak{d}d+\mathfrak{d}+k}^{\mathfrak{d},\eta}(n) \varphi_t \left(\sum_{\ell=1}^d \Theta_{(k-1)d+\ell}^{\mathfrak{d},\eta}(n) z_\ell + \Theta_{\mathfrak{d}d+k}^{\mathfrak{d},\eta}(n) \right) + \Theta_{\mathfrak{d}d+2\mathfrak{d}+1}^{\mathfrak{d},\eta}(n) \right] \\ &= \sum_{k=1}^{\mathfrak{d}} \Theta_{\mathfrak{d}d+\mathfrak{d}+k}^{\mathfrak{d},\eta}(n) \mathfrak{r} \left(\sum_{\ell=1}^d \Theta_{(k-1)d+\ell}^{\mathfrak{d},\eta}(n) z_\ell + \Theta_{\mathfrak{d}d+k}^{\mathfrak{d},\eta}(n) \right) + \Theta_{\mathfrak{d}d+2\mathfrak{d}+1}^{\mathfrak{d},\eta}(n) \\ &= \sum_{k=1}^{\mathfrak{d}} \mathcal{W}_k \mathfrak{r}(\langle W_k(n), z \rangle + B_k(n)) + \mathfrak{B}(n) \\ &= \langle \mathcal{W}, \mathfrak{r}(W(n)z + B(n)) \rangle + \mathfrak{B}(n) = (\mathcal{R}(\Phi(n)))(z) \end{aligned} \quad (177)$$

(cf. Definition 2.3). In addition, observe that item (ii) in Lemma 4.9 shows that for all $z \in \mathbb{R}^d$, $k \in \{1, 2, \dots, \mathfrak{d}\}$, $\ell \in \{1, 2, \dots, d\}$, $i \in \{1, 2, \dots, \mathfrak{d}d\}$, $j \in \{\mathfrak{d}d+1, \mathfrak{d}d+2, \dots, \mathfrak{d}d+\mathfrak{d}\}$, $b \in \mathbb{N}$ with $i = (k-1)d + \ell$, $j = \mathfrak{d}d+k$, and $b = \mathfrak{d}d+2\mathfrak{d}+1$ it holds that

$$\begin{aligned} \frac{\partial}{\partial \theta_i} \mathcal{N}_{\Theta^{\mathfrak{d},\eta}(n)}^{\mathfrak{d},\eta}(z) &= \frac{\partial}{\partial \theta_i} \left(\sum_{r=1}^{\mathfrak{d}} \Theta_{\mathfrak{d}d+\mathfrak{d}+r}^{\mathfrak{d},\eta}(n) \varphi_t \left(\sum_{s=1}^d \Theta_{(r-1)d+s}^{\mathfrak{d},\eta}(n) \langle z, e_s \rangle + \Theta_{\mathfrak{d}d+r}^{\mathfrak{d},\eta}(n) \right) + \Theta_{\mathfrak{d}d+2\mathfrak{d}+1}^{\mathfrak{d},\eta}(n) \right) \\ &= \Theta_{\mathfrak{d}d+\mathfrak{d}+k}^{\mathfrak{d},\eta}(n) (\varphi_t)' \left(\sum_{s=1}^d \Theta_{(k-1)d+s}^{\mathfrak{d},\eta}(n) \langle z, e_s \rangle + \Theta_{\mathfrak{d}d+k}^{\mathfrak{d},\eta}(n) \right) \langle z, e_\ell \rangle \\ &= \mathcal{W}_k (\varphi_t)'(\langle W_k(n), z \rangle + B_k(n)) \langle z, e_\ell \rangle, \end{aligned} \quad (178)$$

$$\begin{aligned} \frac{\partial}{\partial \theta_j} \mathcal{N}_{\Theta^{\mathfrak{d},\eta}(n)}^{\mathfrak{d},\eta}(z) &= \frac{\partial}{\partial \theta_j} \left(\sum_{r=1}^{\mathfrak{d}} \Theta_{\mathfrak{d}d+\mathfrak{d}+r}^{\mathfrak{d},\eta}(n) \varphi_t \left(\sum_{s=1}^d \Theta_{(r-1)d+s}^{\mathfrak{d},\eta}(n) \langle z, e_s \rangle + \Theta_{\mathfrak{d}d+r}^{\mathfrak{d},\eta}(n) \right) + \Theta_{\mathfrak{d}d+2\mathfrak{d}+1}^{\mathfrak{d},\eta}(n) \right) \\ &= \Theta_{\mathfrak{d}d+\mathfrak{d}+k}^{\mathfrak{d},\eta}(n) (\varphi_t)' \left(\sum_{s=1}^d \Theta_{(k-1)d+s}^{\mathfrak{d},\eta}(n) \langle z, e_s \rangle + \Theta_{\mathfrak{d}d+k}^{\mathfrak{d},\eta}(n) \right) \\ &= \mathcal{W}_k (\varphi_t)'(\langle W_k(n), z \rangle + B_k(n)), \end{aligned} \quad (179)$$

and

$$\frac{\partial}{\partial \theta_b} \mathcal{N}_{\Theta^{\mathfrak{d},\eta}(n)}^{\mathfrak{d},\eta}(z) = \frac{\partial}{\partial \theta_b} \left(\sum_{r=1}^{\mathfrak{d}} \Theta_{\mathfrak{d}d+\mathfrak{d}+r}^{\mathfrak{d},\eta}(n) \varphi_t \left(\sum_{s=1}^d \Theta_{(r-1)d+s}^{\mathfrak{d},\eta}(n) \langle z, e_s \rangle + \Theta_{\mathfrak{d}d+r}^{\mathfrak{d},\eta}(n) \right) + \Theta_{\mathfrak{d}d+2\mathfrak{d}+1}^{\mathfrak{d},\eta}(n) \right) = 1. \quad (180)$$

Combining this with (176), (177), and item (iii) in Lemma 4.9 establishes that for all $k \in \{1, 2, \dots, \mathfrak{d}\}$, $\ell \in \{1, 2, \dots, d\}$, $i \in \{1, 2, \dots, \mathfrak{d}d\}$, $j \in \{\mathfrak{d}d+1, \mathfrak{d}d+2, \dots, \mathfrak{d}d+\mathfrak{d}\}$, $b \in \mathbb{N}$ with $i = (k-1)d + \ell$, $j = \mathfrak{d}d+k$, and $b = \mathfrak{d}d+2\mathfrak{d}+1$ it holds that

$$\begin{aligned} \lim_{t \rightarrow \infty} \frac{\partial}{\partial \theta_i} \mathcal{E}^{\mathfrak{d},t}(\Theta^{\mathfrak{d},\eta}(n)) &= \lim_{t \rightarrow \infty} \left[\frac{2}{m} \sum_{p=1}^m (\mathcal{N}_{\Theta^{\mathfrak{d},\eta}(n)}^{\mathfrak{d},t}(x_p) - y_p) \left(\frac{\partial}{\partial \theta_i} \mathcal{N}_{\Theta^{\mathfrak{d},\eta}(n)}^{\mathfrak{d},t}(x_p) \right) \right] \\ &= \frac{2}{m} \sum_{p=1}^m ((\mathcal{R}(\Phi(n)))(x_p) - y_p) \mathcal{W}_k \mathbb{1}_{[0,\infty)}(\langle W_k(n), x_p \rangle + B_k(n)) \langle x_p, e_\ell \rangle \\ &= \frac{2}{m} \sum_{p=1}^m (f_p(n) - y_p) \mathcal{W}_k \langle x_p, e_\ell \rangle \mathbb{1}_{\{\langle W_k(n), x_p \rangle + B_k(n) \geq 0\}}, \end{aligned} \quad (181)$$

$$\begin{aligned} \lim_{t \rightarrow \infty} \frac{\partial}{\partial \theta_j} \mathcal{E}^{\mathfrak{d},t}(\Theta^{\mathfrak{d},\eta}(n)) &= \lim_{t \rightarrow \infty} \left[\frac{2}{m} \sum_{p=1}^m (\mathcal{N}_{\Theta^{\mathfrak{d},\eta}(n)}^{\mathfrak{d},t}(x_p) - y_p) \left(\frac{\partial}{\partial \theta_j} \mathcal{N}_{\Theta^{\mathfrak{d},\eta}(n)}^{\mathfrak{d},t}(x_p) \right) \right] \\ &= \frac{2}{m} \sum_{p=1}^m ((\mathcal{R}(\Phi(n)))(x_p) - y_p) \mathcal{W}_k \mathbb{1}_{[0,\infty)}(\langle W_k(n), x_p \rangle + B_k(n)) \\ &= \frac{2}{m} \sum_{p=1}^m (f_p(n) - y_p) \mathcal{W}_k \mathbb{1}_{\{\langle W_k(n), x_p \rangle + B_k(n) \geq 0\}}, \end{aligned} \quad (182)$$

and

$$\begin{aligned} \lim_{t \rightarrow \infty} \frac{\partial}{\partial \theta_b} \mathcal{E}^{\mathfrak{d},t}(\Theta^{\mathfrak{d},\eta}(n)) &= \lim_{t \rightarrow \infty} \left[\frac{2}{m} \sum_{p=1}^m (\mathcal{N}_{\Theta^{\mathfrak{d},\eta}(n)}^{\mathfrak{d},t}(x_p) - y_p) \left(\frac{\partial}{\partial \theta_b} \mathcal{N}_{\Theta^{\mathfrak{d},\eta}(n)}^{\mathfrak{d},t}(x_p) \right) \right] \\ &= \frac{2}{m} \sum_{p=1}^m ((\mathcal{R}(\Phi(n)))(x_p) - y_p) = \frac{2}{m} \sum_{p=1}^m (f_p(n) - y_p). \end{aligned} \quad (183)$$

This and the assumption that for all $i \in \{1, 2, \dots, \mathcal{P}(\mathfrak{d})\}$, $n \in \mathbb{N}_0$ it holds that $\Theta_i^{\mathfrak{d},\eta}(n+1) = \Theta_i^{\mathfrak{d},\eta}(n) - \eta [\lim_{t \rightarrow \infty} (\frac{\partial}{\partial \theta_i} \mathcal{E}^{\mathfrak{d},t})(\Theta^{\mathfrak{d},\eta}(n))] \mathbb{1}_{\mathbb{R} \setminus (\mathfrak{d}d+\mathfrak{d}, \mathfrak{d}d+2\mathfrak{d})}(i)$ therefore establish that for all $k \in \{1, 2, \dots, \mathfrak{d}\}$, $n \in \mathbb{N}_0$, $\omega \in \Omega$ it holds that

$$W_k(n+1, \omega) = W_k(n, \omega) - \frac{2\eta}{m} \left(\sum_{j=1}^m (f_j(n, \omega) - y_j) \mathcal{W}_k(\omega) \mathbb{1}_{\{(W_k(n), x_j) + B_k(n) \geq 0\}}(\omega) x_j \right), \quad (184)$$

$$B_k(n+1, \omega) = B_k(n, \omega) - \frac{2\eta}{m} \left(\sum_{j=1}^m (f_j(n, \omega) - y_j) \mathcal{W}_k(\omega) \mathbb{1}_{\{(W_k(n), x_j) + B_k(n) \geq 0\}}(\omega) \right), \quad (185)$$

and

$$\mathfrak{B}(n+1, \omega) = \mathfrak{B}(n, \omega) - \frac{2\eta}{m} \left(\sum_{j=1}^m (f_j(n, \omega) - y_j) \right). \quad (186)$$

Corollary 4.8 (applied with $\Lambda \curvearrowright \Lambda$, $\mathcal{E} \curvearrowright \mathfrak{E}$ in the notation of Corollary 4.8) hence demonstrates that

$$\mathbb{P}(\forall n \in \mathbb{N}_0: \mathfrak{E}(\Phi(n)) \leq (1 - \eta\Lambda)^n \mathfrak{E}(\Phi(0))) \geq 1 - \varepsilon. \quad (187)$$

In addition, observe that (177) implies for all $n \in \mathbb{N}_0$ that

$$\lim_{t \rightarrow \infty} \mathcal{E}^{\mathfrak{d},t}(\Theta^{\mathfrak{d},\eta}(n)) = \lim_{t \rightarrow \infty} \frac{1}{m} \sum_{j=1}^m |\mathcal{N}_{\Theta^{\mathfrak{d},\eta}(n)}^{\mathfrak{d},t}(x_j) - y_j|^2 = \frac{1}{m} \sum_{j=1}^m |(\mathcal{R}(\Phi(n)))(x_j) - y_j|^2 = \mathfrak{E}(\Phi(n)). \quad (188)$$

Combining this with (187) therefore ensures that

$$\begin{aligned} \mathbb{P}(\forall n \in \mathbb{N}_0: [\lim_{t \rightarrow \infty} \mathcal{E}^{\mathfrak{d},t}(\Theta^{\mathfrak{d},\eta}(n))] \leq (1 - \eta\lambda)^n [\lim_{t \rightarrow \infty} \mathcal{E}^{\mathfrak{d},t}(\Theta^{\mathfrak{d},\eta}(0))]) \\ = \mathbb{P}(\forall n \in \mathbb{N}_0: \mathfrak{E}(\Phi(n)) \leq (1 - \eta\Lambda)^n \mathfrak{E}(\Phi(0))) \geq 1 - \varepsilon. \end{aligned} \quad (189)$$

The proof of Corollary 4.10 is thus completed. \square

Acknowledgements

This work has been funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) under Germany's Excellence Strategy EXC 2044–390685587, Mathematics Münster: Dynamics–Geometry–Structure.

References

- [1] AKYILDIZ, Ö. D., AND SABANIS, S. Nonasymptotic analysis of Stochastic Gradient Hamiltonian Monte Carlo under local conditions for nonconvex optimization. *arXiv:2002.05465* (2021), 26 pages.
- [2] ALLEN-ZHU, Z., LI, Y., AND LIANG, Y. Learning and Generalization in Overparameterized Neural Networks, Going Beyond Two Layers. *arXiv:1811.04918* (2020), 84 pages.
- [3] ALLEN-ZHU, Z., LI, Y., AND SONG, Z. A Convergence Theory for Deep Learning via Overparameterization. *arXiv:1811.03962* (2019), 53 pages.
- [4] BACH, F. Breaking the curse of dimensionality with convex neural networks. *Journal of Machine Learning Research* 18, 19 (2017), 1–53.
- [5] BACH, F., AND MOULINES, E. Non-strongly-convex smooth stochastic approximation with convergence rate $O(1/n)$. In *Advances in Neural Information Processing Systems* (2013), C. J. C. Burges, L. Bottou, M. Welling, Z. Ghahramani, and K. Q. Weinberger, Eds., vol. 26, Curran Associates, Inc., pp. 773–781.

- [6] BECK, C., JENTZEN, A., AND KUCKUCK, B. Full error analysis for the training of deep neural networks. *arXiv:1910.00121* (2020), 53 pages.
- [7] CHERIDITO, P., JENTZEN, A., RIEKERT, A., AND ROSSMANNEK, F. A proof of convergence for gradient descent in the training of artificial neural networks for constant target functions. *arXiv:2102.09924* (2021), 23 pages.
- [8] CHERIDITO, P., JENTZEN, A., AND ROSSMANNEK, F. Non-convergence of stochastic gradient descent in the training of deep neural networks. *Journal of Complexity* (2020), 101540.
- [9] CHIZAT, L., AND BACH, F. On the Global Convergence of Gradient Descent for Over-parameterized Models using Optimal Transport. *arXiv:1805.09545* (2018), 32 pages.
- [10] DEREICH, S., AND KASSING, S. Convergence of stochastic gradient descent schemes for Lojasiewicz-landscapes. *arXiv:2102.09385* (2021), 21 pages.
- [11] DU, S. S., LEE, J. D., LI, H., WANG, L., AND ZHAI, X. Gradient Descent Finds Global Minima of Deep Neural Networks. *arXiv:1811.03804* (2019), 45 pages.
- [12] DU, S. S., ZHAI, X., POCZOS, B., AND SINGH, A. Gradient Descent Provably Optimizes Over-parameterized Neural Networks. *arXiv:1810.02054* (2019), 19 pages.
- [13] E, W., MA, C., AND WU, L. A comparative analysis of optimization and generalization properties of two-layer neural network and random feature models under gradient descent dynamics. *Science China Mathematics* 63, 7 (2020), 1235–1258.
- [14] FEHRMAN, B., GESS, B., AND JENTZEN, A. Convergence rates for the stochastic gradient descent method for non-convex objective functions. *J. Mach. Learn. Res.* 21 (2020), Paper No. 136, 48.
- [15] HANIN, B. Which Neural Net Architectures Give Rise to Exploding and Vanishing Gradients? In *Advances in Neural Information Processing Systems* (2018), S. Bengio, H. Wallach, H. Larochelle, K. Grauman, N. Cesa-Bianchi, and R. Garnett, Eds., vol. 31, Curran Associates, Inc., pp. 582–591.
- [16] HANIN, B., AND ROLNICK, D. How to Start Training: The Effect of Initialization and Architecture. In *Advances in Neural Information Processing Systems* (2018), S. Bengio, H. Wallach, H. Larochelle, K. Grauman, N. Cesa-Bianchi, and R. Garnett, Eds., vol. 31, Curran Associates, Inc., pp. 571–581.
- [17] JACOT, A., GABRIEL, F., AND HONGLER, C. Neural Tangent Kernel: Convergence and Generalization in Neural Networks. *arXiv:1806.07572* (2020), 19 pages.
- [18] JENTZEN, A., AND VON WURSTEMBERGER, P. Lower error bounds for the stochastic gradient descent optimization algorithm: Sharp convergence rates for slowly and fast decaying learning rates. *Journal of Complexity* 57 (2020), 101438.
- [19] JENTZEN, A., AND WELTI, T. Overall error analysis for the training of deep neural networks via stochastic gradient descent with random initialisation. *arXiv:1910.00121* (2020), 51 pages.
- [20] LEI, Y., HU, T., LI, G., AND TANG, K. Stochastic Gradient Descent for Nonconvex Learning Without Bounded Gradient Assumptions. *IEEE Transactions on Neural Networks and Learning Systems* 31, 10 (2020), 4394–4400.
- [21] LI, Y., AND LIANG, Y. Learning Overparameterized Neural Networks via Stochastic Gradient Descent on Structured Data. *arXiv:1808.01204* (2019), 28 pages.
- [22] LOVAS, A., LYTRAS, I., RÁSONYI, M., AND SABANIS, S. Taming neural networks with TUSLA: Non-convex learning via adaptive stochastic gradient Langevin algorithms. *arXiv:2006.14514* (2020), 29 pages.
- [23] LU, L., SHIN, Y., SU, Y., AND KARNIAKAKIS, G. E. Dying ReLU and Initialization: Theory and Numerical Examples. *Communications in Computational Physics* 28, 5 (2020), 1671–1706.

- [24] MOULINES, E., AND BACH, F. Non-Asymptotic Analysis of Stochastic Approximation Algorithms for Machine Learning. In *Advances in Neural Information Processing Systems* (2011), J. Shawe-Taylor, R. Zemel, P. Bartlett, F. Pereira, and K. Q. Weinberger, Eds., vol. 24, Curran Associates, Inc., pp. 451–459.
- [25] RUDER, S. An overview of gradient descent optimization algorithms. *arXiv:1609.04747* (2017), 14 pages.
- [26] SANKARARAMAN, K. A., DE, S., XU, Z., HUANG, W. R., AND GOLDSTEIN, T. The Impact of Neural Network Overparameterization on Gradient Confusion and Stochastic Gradient Descent. *arXiv:1904.06963* (2020), 28 pages.
- [27] SHIN, Y., AND KARNIADAKIS, G. E. Trainability of ReLU networks and Data-dependent Initialization. *Journal of Machine Learning for Modeling and Computing* 1, 1 (2020), 39–74.
- [28] WU, L., MA, C., AND E, W. How SGD Selects the Global Minima in Over-parameterized Learning: A Dynamical Stability Perspective. In *Advances in Neural Information Processing Systems* (2018), S. Bengio, H. Wallach, H. Larochelle, K. Grauman, N. Cesa-Bianchi, and R. Garnett, Eds., vol. 31, Curran Associates, Inc., pp. 8279–8288.
- [29] ZOU, D., CAO, Y., ZHOU, D., AND GU, Q. Stochastic Gradient Descent Optimizes Over-parameterized Deep ReLU Networks. *arXiv:1811.08888* (2018), 54 pages.