

# Simplified quasi-likelihood analysis for a locally asymptotically quadratic random field <sup>\*</sup>

Nakahiro Yoshida

Graduate School of Mathematical Sciences, University of Tokyo <sup>†</sup>  
 Japan Science and Technology Agency CREST  
 The Institute of Statistical Mathematics

December 27, 2021

*Summary* The asymptotic decision theory by Le Cam and Hájek has been given a lucid perspective by the Ibragimov-Hasminskii theory on convergence of the likelihood random field. Their scheme has been applied to stochastic processes by Kutoyants, and today this plot is called the IHK program. This scheme ensures that asymptotic properties of an estimator follow directly from the convergence of the random field if a large deviation estimate exists. The quasi-likelihood analysis (QLA) proved a polynomial type large deviation (PLD) inequality to go through a bottleneck of the program. A conclusion of the QLA is that if the quasi-likelihood random field is asymptotically quadratic and if a key index reflecting identifiability the random field has is non-degenerate, then the PLD inequality is always valid, and as a result, the IHK program can run. Many studies already took advantage of the QLA theory. However, not a few of them are using it in an inefficient way yet. The aim of this paper is to provide a reformed and simplified version of the QLA and to improve accessibility to the theory. As an example of the effects of the program and the PLD, the user can obtain asymptotic properties of the quasi-Bayesian estimator by only verifying non-degeneracy of the key index.

*Keywords and phrases* Ibragimov-Has'minskii theory, quasi-likelihood analysis, polynomial type large deviation, random field, asymptotic decision theory, non-ergodic statistics.

## 1 Introduction

The asymptotic decision theory by Le Cam and Hájek has been given a lucid perspective by the Ibragimov-Has'minskii theory ([3, 4, 5]) on convergence of the likelihood random field. Their

---

<sup>\*</sup>This work was in part supported by Japan Science and Technology Agency CREST JPMJCR14D7, JPMJCR2115; Japan Society for the Promotion of Science Grants-in-Aid for Scientific Research No. 17H01702 (Scientific Research); and by a Cooperative Research Program of the Institute of Statistical Mathematics. The author thanks Junichiro Yoshida for valuable comments.

<sup>†</sup>Graduate School of Mathematical Sciences, University of Tokyo: 3-8-1 Komaba, Meguro-ku, Tokyo 153-8914, Japan. e-mail: nakahiro@ms.u-tokyo.ac.jp

scheme has been applied to stochastic processes by Kutoyants ([9, 10, 11, 12]), and today this plot is called the IHK program. This scheme ensures that asymptotic properties of an estimator follow directly from the convergence of the random field if a large deviation estimate exists.

The theory of statistical inference for stochastic processes is heading toward heavily dependent stochastic systems: nonlinear, non-Markovian, non-stationary, non-ergodic, discrete and/or dependent sampling. A formal extension of the likelihood analysis to a quasi-likelihood analysis is inevitable. However, the large deviation inequality is an issue even within the likelihood analysis for stochastic processes. The quasi-likelihood analysis (QLA) proved a polynomial type large deviation (PLD) inequality to go through a bottleneck of the program (Yoshida [30]).<sup>1</sup> As a conclusion of the QLA theory, if the quasi-likelihood random field is locally asymptotically quadratic (LAQ) and if a key index reflecting identifiability the random field has is non-degenerate, then the polynomial type large deviation inequality is always valid, and as a result, the IHK program can run.

Since an ad hoc model-dependent method is not necessary, the QLA is universal and can apply to various dependent models. Many studies are based on and taking advantage of the QLA. These applications include sampled ergodic diffusion processes (Yoshida [30]), adaptive estimation for diffusion processes (Uchida and Yoshida [25]), adaptive Bayes type estimators for ergodic diffusion processes (Uchida and Yoshida [28]), approximate self-weighted LAD estimation of discretely observed ergodic Ornstein-Uhlenbeck processes (Masuda [13]), parametric estimation of Lévy processes (Masuda [15]), Gaussian quasi-likelihood random fields for ergodic Lévy driven SDE (Masuda [14]), and ergodic point processes for limit order book (Clinet and Yoshida [1]). Thanks to its flexibility, the QLA is also applicable to non-ergodic statistics: volatility parameter estimation in regular sampling of finite time horizon (Uchida and Yoshida [27]) and in non-synchronous sampling (Ogihara and Yoshida [19]), a non-ergodic point process regression model (Ogihara and Yoshida [20]). Analysis of complex algorithms is possible by relying on the universal design of the QLA: hybrid multi-step estimators (Kamatani and Uchida [7]), adaptive Bayes estimators and hybrid estimators for small diffusion processes based on sampled data (Nomura and Uchida [17]). Information criteria, sparse estimation and regularization methods are recently understood in the framework of the QLA: contrast-based information criterion for diffusion processes (Uchida [24]), AIC for non-concave penalized likelihood method (Umezumi et al. [29]), Schwarz type model comparison for LAQ models (Eguchi and Masuda [2]), moment convergence of regularized least-squares estimator for linear regression model (Shimizu [21]), moment convergence in regularized estimation under multiple and mixed-rates asymptotics (Masuda and Shimizu [16]), penalized method and polynomial type large deviation inequality (Kinoshita and Yoshida [8]) and related Suzuki and Yoshida ([23]). Jump filtering problems: jump diffusion processes Ogihara and Yoshida([18]), threshold estimation for stochastic processes with small noise (Shimizu [22]), global jump filters (Inatsugu and Yoshida [6]). Partial quasi-likelihood analysis: Yoshida [31]. Such variety of applications are demonstrating the universality of the framework of the QLA. Since the IHK program runs there, we can obtain limit theorems and the  $L^p$ -boundedness of the QL estimators (quasi-maximum likelihood estimator and the quasi-Bayesian estimator), which is indispensable

---

<sup>1</sup>The term “quasi-likelihood” is not in the sense of GLM. We use “quasi-likelihood analysis” because statistical inference for sampled stochastic processes cannot avoid a quasi-likelihood function for estimation. The method is relatively new, but not because of “quasi”. The difficulty in large deviation estimates already existed in the likelihood analysis for stochastic processes.

to develop statistical theories.

The essence of the QLA is the polynomial type large deviation inequality that was proved in a general setting (Yoshida [30]). Since the LAQ property quite often appears when the model is differentiable, Yoshida [30] was based on this structure. Because of it, the limit distribution of the associated estimators has an explicit expression. The paper [30] gave it, but due to a general way of writing, not a few users are apt to avoid following that passage after the PLD's theorem and try to reconstruct it in each situation. However, such a task is unnecessary in fact. Besides, four time differentiability is often assumed in many applications of the QLA. It may be only because a handy condition in [30] assumed an estimate of the supremum of the third-order derivative of the quasi-log likelihood random field  $\mathbb{H}_T$ , though the paper gave a condition ([A1']) to treat  $\mathbb{H}_T$  of class  $C^2$ .

The aim of this paper is to provide a simplified version of the QLA theory directly connecting the assumptions with the limit theorems in order to improve accessibility to the theory. Essentially, the user is only requested to verify non-degeneracy of a key index, and this task is trivial in particular in ergodic statistics. We will give handy conditions for the quasi-likelihood random field of class  $C^2$ , based on [30], in order to reach the asymptotic properties of the estimators at a single leap. Some assumptions in [30] are arranged and replaced by simple-looking ones in this paper. This simplification will serve for future progress e.g. in analysis of regularization methods. The LAQ property we adopted here is just one principle of separation, and it is possible to develop a similar theory for a non-LAQ type random field; see Kinoshita and Yoshida [8] for a case of regularization.

A smart way of presenting the theory is to use the convergence of the quasi-likelihood random field  $\mathbb{Z}_T$  to a random field  $\mathbb{Z}$  in the function space  $\widehat{C}(\mathbb{R}^p)$ , the separable Banach space of continuous functions  $f$  on  $\mathbb{R}^p$  satisfying  $\lim_{|u| \rightarrow \infty} f(u) = 0$ , equipped with the supremum norm. This plot is possible but to carry out it, one needs a suitable measurable extension of  $\mathbb{Z}_T$  to the outside of the originally given local parameter space and an argument about tightness of random fields on the non-compact  $\mathbb{R}^p$ . In this article, we dared avoid this approach to give priority to simplicity. As a result, the presentation of the theory is now much more elementary than Yoshida [30]. Additionally, though they are classical, some basic properties like the first order efficiency and asymptotic equivalence between the maximum likelihood estimator and the Bayesian estimator will be given within the QLA framework. For convenience of use, we will give and detail several versions of theorems and conditions in different situations.

## 2 A simplified QLA in non-ergodic statistics

Given a probability space  $(\Omega, \mathcal{F}, P)$  and a bounded open set  $\Theta$  in  $\mathbb{R}^p$ , we consider a random field  $\mathbb{H}_T : \Omega \times \overline{\Theta} \rightarrow \mathbb{R}$ , a function measurable with respect to the product  $\sigma$ -field  $\mathcal{F} \times \mathbb{B}(\Theta)$ ,  $\mathbb{B}(\Theta)$  being the Borel  $\sigma$ -field of  $\Theta$ .<sup>2</sup> Here  $t \in \mathbb{T}$ , a subset of  $\mathbb{R}_+ = [0, \infty)$  satisfying  $\sup \mathbb{T} = \infty$ . We suppose that  $\mathbb{H}_T$  is continuous and of class  $C^2$ , that is, for every  $\omega \in \Omega$ , the mapping  $\Theta \ni \theta \mapsto \mathbb{H}_T(\omega, \theta) \in \mathbb{R}$  is of class  $C^2$  and that  $\mathbb{H}_T$  is continuously extended to  $\partial\Theta$ . We shall present a simplified version of the polynomial type large deviation inequality of Yoshida [30] under a handy set of sufficient conditions.

---

<sup>2</sup>Because of the assumptions below about the continuity of  $\mathbb{H}_T$  and the separability of  $\Theta$ , this is equivalent to that the function  $\mathbb{H}_T(\cdot, \theta)$  is measurable for each  $\theta \in \Theta$ .

Let  $\theta^* \in \Theta$ . Define  $\Delta_T$  and  $\Gamma_T(\theta)$  by

$$\Delta_T = \partial_\theta \mathbb{H}_T(\theta^*) a_T \quad \text{and} \quad \Gamma_T(\theta) = -a_T^* \partial_\theta^2 \mathbb{H}_T(\theta) a_T \quad (2.1)$$

respectively, where  $\star$  denotes the matrix transpose. Let  $a_T \in GL(\mathbb{R}^p)$  be a scaling matrix such that  $|a_T| \rightarrow 0$  as  $n \rightarrow \infty$ . We suppose that  $\Gamma$  is a  $p \times p$  symmetric random matrix. Let  $U(\theta, r) = \{\theta' \in \mathbb{R}^p; |\theta' - \theta| < r\}$  for  $\theta \in \Theta$  and  $r > 0$ . There exists a positive constant  $r_0$  such that  $U(\theta^*, r_0) \subset \Theta$ .

The minimum and maximum eigenvalues of the symmetric matrix  $M$  are denoted by  $\lambda_{\min}(M)$  and  $\lambda_{\max}(M)$ , respectively. Let  $b_T = \{\lambda_{\min}(a_T^* a_T)\}^{-1}$ . In particular,  $b_T \rightarrow \infty$  as  $n \rightarrow \infty$ . Moreover, we assume that

$$b_T^{-1} \leq \lambda_{\max}(a_T^* a_T) \leq C_0 b_T^{-1} \quad (T \in \mathbb{T}) \quad (2.2)$$

for some constant  $C_0 \in [1, \infty)$ . A typical case is  $n$  for  $b_T$ , and  $n^{-1/2} I_p$  for  $a_T$ , where  $I_p$  is the identity matrix.

**Remark 2.1.** In an ergodic diffusion model, the parameter  $\theta_1$  of the diffusion coefficient and the parameter  $\theta_2$  of the drift coefficient have different convergence rates in estimation with high frequency data. Then Condition (2.2) may seem restrictive, but it is incorrect. The random field  $\mathbb{H}_n$  is not necessarily the same as a quasi-log likelihood function  $\Psi_n$  used for estimation in reality, where  $n = T \in \mathbb{T} = \mathbb{N}$ . The random field  $\mathbb{H}_n$  is rather "living in the proof" in various manners. Consider a joint quasi-maximum likelihood estimator  $(\hat{\theta}_{1,n}, \hat{\theta}_{2,n})$  for  $(\theta_1, \theta_2)$ . To analyze the asymptotic behavior of  $\hat{\theta}_{1,n}$ , the random field  $\mathbb{H}_n(\theta_1) = \Psi_n(\theta_1, \hat{\theta}_{2,n})$  can be used.  $\mathbb{H}_n(\theta_1)$  is estimated by taking supremum about the second argument of  $\Psi_n$  at some stage. For  $\hat{\theta}_{2,n}$ , one can switch  $\mathbb{H}_n$  to a different random field  $\mathbb{H}_n(\theta_2) = \Psi_n(\hat{\theta}_{1,n}, \theta_2)$ . Such a stepwise application of the QLA in the present article's form can be observed in many studies; see Yoshida [30], Uchida and Yoshida [26, 28] and the papers listed in Introduction.

Define  $\Upsilon_T : \Omega \times \Theta \rightarrow \mathbb{R}$  by

$$\Upsilon_T(\theta) = \frac{1}{b_T} \{\mathbb{H}_T(\theta) - \mathbb{H}_T(\theta^*)\} \quad (\theta \in \Theta)$$

for  $T \in \mathbb{T}$ . Let  $\Upsilon : \Omega \times \Theta \rightarrow \mathbb{R}$  be a continuous random field. Let  $L > 0$  be a positive number.

[S1] Parameters  $\alpha, \beta_1, \beta_2, \rho_1$  and  $\rho_2$  satisfy the following inequalities:

$$\begin{aligned} 0 < \beta_1 < 1/2, \quad 0 < \rho_1 < \min\{1, \alpha/(1-\alpha), 2\beta_1/(1-\alpha)\}, \\ 0 < 2\alpha < \rho_2, \quad \beta_2 \geq 0, \quad 1 - 2\beta_2 - \rho_2 > 0. \end{aligned}$$

[S2] (i) There exists a positive random variable  $\chi_0$  and the following conditions are fulfilled.

(i-1)  $\Upsilon(\theta) = \Upsilon(\theta) - \Upsilon(\theta^*) \leq -\chi_0 |\theta - \theta^*|^2$  for all  $\theta \in \Theta$ .

(i-2) For some constant  $C_L$ , it holds that

$$P[\chi_0 \leq r^{-(\rho_2 - 2\alpha)}] \leq \frac{C_L}{r^L} \quad (r > 0).$$

(ii) For some constant  $C_L$ , it holds that

$$P[\lambda_{\min}(\Gamma) < r^{-\rho_1}] \leq \frac{C_L}{r^L} \quad (r > 0)$$

Let  $\beta = \alpha/(1 - \alpha)$ . Let  $\|V\|_p = (E[|V|^p])^{1/p}$  for  $p > 0$  and a matrix valued random variable  $V$ .

[S3] (i) For  $M_1 = L(1 - \rho_1)^{-1}$ ,  $\sup_{T \in \mathbb{T}} \|\Delta_T\|_{M_1} < \infty$ .

(ii) For  $M_2 = L(1 - 2\beta_2 - \rho_2)^{-1}$ ,

$$\sup_{T \in \mathbb{T}} \left\| \sup_{\theta \in \Theta \setminus U(\theta^*, b_T^{-\alpha/2})} b_T^{\frac{1}{2} - \beta_2} |\Upsilon_T(\theta) - \Upsilon(\theta)| \right\|_{M_2} < \infty.$$

(iii) For  $M_3 = L(\beta - \rho_1)^{-1}$

$$\sup_{T \in \mathbb{T}} \left\| \sup_{u \in \delta^{-1}(\Theta - \theta^*), |u| \leq 1} |\Gamma_T(\theta^* + \delta u) - \Gamma_T(\theta^*)| \right\|_{M_3} = O(\delta) \quad (\delta \downarrow 0).$$

(iv) For  $M_4 = L(2\beta_1(1 - \alpha)^{-1} - \rho_1)^{-1}$ ,

$$\sup_{T \in \mathbb{T}} \|b_T^{\beta_1} |\Gamma_T(\theta^*) - \Gamma|\|_{M_4} < \infty.$$

**Remark 2.2.** (i) In the above conditions, each constant  $C_L$  is independent of  $r$  and  $T$ , but may depend on the parameters appearing in [S1] as well as  $L$ . (ii) In applications, we often need to estimate the supremum of a sequence of martingales depending on  $\theta$  to verify the above moment conditions. Use of Sobolev's embedding inequality is a simple solution. The GRR inequality is an alternative if one wants to reduce differentiability assumptions. (iii) The random matrix  $\Gamma$  is positive-definite a.s. if [S2] (ii) is satisfied.

Let  $\mathcal{U}_T = \{u \in \mathbb{R}^p; \theta^* + a_T u \in \Theta\}$  and  $\mathcal{V}_T(r) = \{u \in \mathcal{U}_T; |u| \geq r\}$  for  $r > 0$ . Define the random field  $\mathbb{Z}_T$  on  $\mathcal{U}_T$  by

$$\mathbb{Z}_T(u) = \exp(\mathbb{H}_T(\theta^* + a_T u) - \mathbb{H}_T(\theta^*))$$

for  $u \in \mathcal{U}_T$ . Following Yoshida [30], we give a polynomial type large deviation inequality for the random field  $\mathbb{Z}_T$ .

**Theorem 2.3.** *Given a positive constant  $L$ , suppose that [S1], [S2] and [S3] are fulfilled. Then there exists a constant  $C_L$  such that*

$$P \left[ \sup_{u \in \mathcal{V}_T(r)} \mathbb{Z}_T(u) \geq \exp(-2^{-1} r^{2 - (\rho_1 \vee \rho_2)}) \right] \leq \frac{C_L}{r^L} \quad (2.3)$$

for all  $r > 0$  and  $T \in \mathbb{T}$ . The supremum of the empty set should read  $-\infty$ .

*Proof.* Suppose that the constants  $\alpha, \beta_1, \beta_2, \rho_1, \rho_2$  satisfy Condition [S1]. We will apply Theorem 1 of [30] with  $\rho = 2$  for  $\mathbb{H}_T$  of class  $C^2$ . According to Section 3.1 of [30], it suffices to verify Conditions [A1'] and [A2]-[A6] therein. Condition [A4] of [30] with  $\rho = 2$  and the condition that  $\alpha \in (0, 1)$  are satisfied under [S1] since  $\rho_2 < 1$ .

Condition [A1'] of [30] requires the estimate

$$1_{\{r \leq b_T^{(1-\alpha)/2}\}} P[S'_T(r)^c] \leq \frac{C_L}{r^L} \quad (r > 0, T \in \mathbb{T}) \quad (2.4)$$

for some constant  $C_L$ , where the event  $S'_T(r)$  is defined by

$$S'_T(r) = \left\{ \sup_{\substack{h: \theta^* + h \in \Theta, \\ |h| \leq C_0^{1/2} b_T^{-\alpha/2}}} |\Gamma_T(\theta^* + h) - \Gamma| < r^{-\rho_1} \right\}; \quad (2.5)$$

see Remark 2.4. To verify (2.4), we may consider the case where  $r \leq b_T^{(1-\alpha)/2}$ , equivalently,

$$b_T^{-1} \leq r^{-2/(1-\alpha)}. \quad (2.6)$$

We have

$$1_{\{r \leq b_T^{(1-\alpha)/2}\}} P[S'_T(r)^c] \leq \mathbb{P}_1(T, r) + \mathbb{P}_2(T, r), \quad (2.7)$$

where

$$\mathbb{P}_1(T, r) = P \left[ \sup_{h \in \Theta - \theta^*: |h| \leq C_0^{1/2} r^{-\beta}} |\Gamma_T(\theta^* + h) - \Gamma_T(\theta^*)| \geq \frac{1}{2} r^{-\rho_1} \right]$$

and

$$\mathbb{P}_2(T, r) = P \left[ |\Gamma_T(\theta^*) - \Gamma| \geq \frac{1}{2} r^{-\rho_1} \right] 1_{\{b_T^{-1} \leq r^{-2/(1-\alpha)}\}}$$

in view of (2.6). For sufficiently large  $r$ , by Condition [S3] (iii),

$$\begin{aligned} \sup_{T \in \mathbb{T}} \mathbb{P}_1(T, r) &\lesssim r^{M_3 \rho_1} \sup_{T \in \mathbb{T}} E \left[ \sup_{h \in \Theta - \theta^*: |h| \leq C_0^{1/2} r^{-\beta}} |\Gamma_T(\theta^* + h) - \Gamma_T(\theta^*)|^{M_3} \right] \\ &\lesssim r^{-M_3(\beta - \rho_1)} \\ &= r^{-L}. \end{aligned} \quad (2.8)$$

Next, by Condition [S3] (iv), we have

$$\begin{aligned} \sup_{T \in \mathbb{T}} \mathbb{P}_2(T, r) &\lesssim \sup_{T \in \mathbb{T}} \left( b_T^{-M_4 \beta_1} r^{M_4 \rho_1} 1_{\{b_T^{-1} \leq r^{-2/(1-\alpha)}\}} \right) \\ &\lesssim r^{-M_4(2\beta_1/(1-\alpha) - \rho_1)} \\ &= r^{-L}. \end{aligned} \quad (2.9)$$

From (2.7), (2.8) and (2.9), we obtain (2.4), therefore [A1'] of [30] has been verified.

Condition [A6] of [30] follows from Condition [S3] (i) and Condition [S3] (ii). Condition [S2] (i) ensures Conditions [A3] for  $\rho = 2$  and [A5] of [30]. Moreover, [S2] (ii) verifies [A2] of [30]. Now, as already mentioned, we apply Theorem 1 of [30] to  $\mathbb{Z}_T$  for  $\rho = 2$  in order to obtain (2.3).  $\square$

**Remark 2.4.** In the proof of Theorem 2.3, we changed  $S'_T(r; \xi_0)$  of Yoshida [30] to (2.5), and Condition [A1'] therein to (2.4). The latter apparently weakens Condition [A1'], however it does not affect the proof in Yoshida [30] because  $S_T(r; \xi_0) = \Omega$  when  $r > b_T(\xi_0)^{(1-\alpha)/2}$  in the notation of the paper.

Define  $r_T(u)$  by

$$r_T(u) = \begin{cases} \mathbb{H}_T(\theta^* + a_T u) - \mathbb{H}_T(\theta^*) - \left( \Delta_T[u] - \frac{1}{2}\Gamma[u^{\otimes 2}] \right) & (u \in \mathbb{U}_T) \\ 1 & (u \notin \mathbb{U}_T) \end{cases} \quad (2.10)$$

**Proposition 2.5.** *Suppose that*

$$\sup_{u \in \mathbb{U}_T \cap U(0, K)} |\Gamma_T(\theta^* + a_T u) - \Gamma| \xrightarrow{p} 0 \quad (T \rightarrow \infty) \quad (2.11)$$

for every  $K > 0$ . Then the random field  $\mathbb{Z}_T$  is locally asymptotically quadratic at  $\theta^*$ , that is,

$$\mathbb{Z}_T(u) = \exp \left( \Delta_T[u] - \frac{1}{2}\Gamma[u^{\otimes 2}] + r_T(u) \right) \quad (u \in \mathbb{U}_T) \quad (2.12)$$

and  $r_T(u) \xrightarrow{p} 0$  as  $T \rightarrow \infty$  for every  $u \in \mathbb{R}^p$ .

*Proof.* By definition of  $r_T(u)$ , Equation (2.12) holds for  $u \in \mathbb{U}_T$ . For each  $u \in \mathbb{R}^p$ , there is a number  $T_u$  such that  $a_T u \in U(0, \mathbf{r}_0)$  for all  $T \geq T_u$ . Then  $r_T(u)$  admits the expression

$$r_T(u) = - \int_0^1 (1-s) \{ \Gamma_T(\theta^* + s a_T u) - \Gamma \} ds [u^{\otimes 2}]. \quad (2.13)$$

Therefore  $r_T(u) \xrightarrow{p} 0$  as  $T \rightarrow \infty$  by (2.11).  $\square$

**Remark 2.6.** We have

$$\sup_{u \in \mathbb{U}_T \cap U(0, K)} |\Gamma_T(\theta^* + a_T u) - \Gamma_T(\theta^*)| \xrightarrow{p} 0 \quad (2.14)$$

as  $T \rightarrow \infty$  under [S3] (iii) since

$$\begin{aligned} & \limsup_{T \rightarrow \infty} \left\| \sup_{u \in \mathbb{U}_T \cap U(0, K)} |\Gamma_T(\theta^* + a_T u) - \Gamma_T(\theta^*)| \right\|_{M_3} \\ & \leq \limsup_{T \rightarrow \infty} \left\| \sup_{v \in \mathbb{R}^p: |v| \leq 1} |\Gamma_T(\theta^* + K|a_T|v) - \Gamma_T(\theta^*)| \right\|_{M_3} \\ & \leq \limsup_{T \rightarrow \infty} O(K|a_T|) = 0. \end{aligned}$$

On the other hand,

$$\Gamma_T(\theta^*) - \Gamma \xrightarrow{p} 0 \quad (2.15)$$

as  $T \rightarrow \infty$  under [S3] (iv). Therefore,  $\mathbb{Z}_T$  is locally asymptotically quadratic at  $\theta^*$  if [S3] (iii) and (iv) are satisfied since the convergence (2.11) holds under [S3] (iii) and (iv), though these conditions are too sufficient for (2.11).

Let  $\Delta$  be a  $p$ -dimensional random vector on some extension of  $(\Omega, \mathcal{F}, P)$ . Define a random field  $\mathbb{Z}$  on  $\mathbb{R}^p$  by

$$\mathbb{Z}(u) = \exp\left(\Delta[u] - \frac{1}{2}\Gamma[u^{\otimes 2}]\right) \quad (2.16)$$

for  $u \in \mathbb{R}^p$ . Let  $\hat{u} = \Gamma^{-1}\Delta$ .

Any measurable mapping  $\hat{\theta}_T^M : \Omega \rightarrow \bar{\Theta}$  is called a quasi-maximum likelihood estimator (QMLE) for  $\mathbb{H}_T$  if

$$\mathbb{H}_T(\hat{\theta}_T^M) = \max_{\theta \in \bar{\Theta}} \mathbb{H}_T(\theta). \quad (2.17)$$

Since  $\mathbb{H}_T$  is continuous on the compact  $\bar{\Theta}$ , such a measurable function always exists, which is ensured by the measurable selection theorem. Uniqueness of  $\hat{\theta}_T^M$  is not assumed. Let  $\hat{u}_T^M = a_T^{-1}(\hat{\theta}_T^M - \theta^*)$  for the QMLE  $\hat{\theta}_T^M$ .

Let  $\mathcal{G}$  be a  $\sigma$ -field such that  $\sigma[\Gamma] \subset \mathcal{G} \subset \mathcal{F}$ . It is said that a sequence  $(V_T)_{T \in \mathbb{T}}$  of random variables taking values in a metric space  $S$  equipped with the Borel  $\sigma$ -field converges  $\mathcal{G}$ -stably to an  $S$ -valued random variable  $V_\infty$  defined on an extension of  $(\Omega, \mathcal{F}, P)$  if  $(V_T, \Psi) \xrightarrow{d} (V_\infty, \Psi)$  as  $T \rightarrow \infty$  for any  $\mathcal{G}$ -measurable random variable  $\Psi$ . The  $\mathcal{G}$ -stable convergence is denoted by  $\xrightarrow{d_s(\mathcal{G})}$ .

**Theorem 2.7.** *Let  $L > p > 0$ . Suppose that Conditions [S1], [S2] and [S3] are satisfied and that*

$$\Delta_T \xrightarrow{d_s(\mathcal{G})} \Delta \quad (2.18)$$

as  $T \rightarrow \infty$ . Then

(a) As  $T \rightarrow \infty$ ,

$$\hat{u}_T^M - \Gamma^{-1}\Delta_T \xrightarrow{p} 0. \quad (2.19)$$

(b) As  $T \rightarrow \infty$ ,

$$E[f(\hat{u}_T^M)\Phi] \rightarrow \mathbb{E}[f(\hat{u})\Phi] \quad (2.20)$$

for any bounded  $\mathcal{G}$ -measurable random variable  $\Phi$  and any  $f \in C(\mathbb{R}^p)$  satisfying  $\limsup_{|u| \rightarrow \infty} |u|^{-p}|f(u)| < \infty$ .



*Proof.* As mentioned in Remark 2.6, the convergence (2.11) holds for every  $K > 0$  under [S3] (iii) and (iv). Then the representation (2.13) ensures

$$\sup_{u: |u| \leq R} |r_T(u)| \xrightarrow{p} 0 \quad (T \rightarrow \infty) \quad (2.21)$$

for every  $R > 0$ . The space  $C(\overline{U(0, R)})$  of continuous function on  $\overline{U(0, R)}$  is equipped with the supremum norm. Combining the representation (2.12) of  $\mathbb{Z}_T$  with the convergences (2.18) and (2.21), by estimating the modulus of continuity of  $\log \mathbb{Z}_T$  on  $\overline{U(0, R)}$ , we obtain tightness of the family  $\{\mathbb{Z}_T|_{\overline{U(0, R)}}\}_{T \geq T_0}$  for some  $T_0 \in \mathbb{T}$ , which yields the convergence

$$\mathbb{Z}_T|_{\overline{U(0, R)}} \xrightarrow{d} \mathbb{Z}|_{\overline{U(0, R)}} \quad (2.22)$$

in  $C(\overline{U(0, R)})$  as  $T \rightarrow \infty$  for every  $R > 0$  because the finite dimensional convergence  $\mathbb{Z}_T \xrightarrow{df} \mathbb{Z}$  is given by (2.18), (2.21) and (2.12).

Let  $F$  be any closed set in  $\mathbb{R}^p$ . Then

$$\begin{aligned} \limsup_{T \rightarrow \infty} P[\hat{u}_T^M \in F] &\leq \limsup_{T \rightarrow \infty} P[\hat{u}_T^M \in F \cap \overline{U(0, R)}] + \limsup_{T \rightarrow \infty} P[\hat{u}_T^M \in \overline{\mathbb{V}_T(R)}] \\ &\leq \limsup_{T \rightarrow \infty} P \left[ \sup_{u \in F \cap \overline{U(0, R)}} \mathbb{Z}_T(u) - \sup_{u \in F^c \cap \overline{U(0, R)}} \mathbb{Z}_T(u) \geq 0 \right] \\ &\quad + \limsup_{T \rightarrow \infty} P \left[ \sup_{u \in \mathbb{V}_T(R)} \mathbb{Z}_T(u) \geq 1 \right] \\ &\leq P \left[ \sup_{u \in F \cap \overline{U(0, R)}} \mathbb{Z}(u) - \sup_{u \in F^c \cap \overline{U(0, R)}} \mathbb{Z}(u) \geq 0 \right] + \frac{C_L}{R^L} \end{aligned} \quad (2.23)$$

by the convergence (2.22) and the polynomial type large deviation inequality (2.3) given by Theorem 2.3. Let  $R \rightarrow \infty$  in (2.23) to obtain

$$\limsup_{T \rightarrow \infty} P[\hat{u}_T^M \in F] \leq P \left[ \sup_{u \in F} \mathbb{Z}(u) - \sup_{u \in F^c} \mathbb{Z}(u) \geq 0 \right] \leq P[\hat{u} \in F]. \quad (2.24)$$

Here the positivity of  $\Gamma$  given by [S2] (ii) (Remark 2.2) was used for the first inequality, and the last inequality is by the uniqueness of the maximum point of the random field  $\mathbb{Z}$  defined by (2.16). Inequality (2.24) shows the convergence  $\hat{u}_T^M \xrightarrow{d} \hat{u}$  as  $T \rightarrow \infty$ .

From the convergence of  $\hat{u}_T^M$ , in particular  $\hat{\theta}_T^M \xrightarrow{p} \theta^*$ , and when  $\hat{\theta}_T^M \in U(\theta^*, \mathbf{r}_0)$ , one has

$$\Delta_T = \int_0^1 \Gamma_T(\theta_T^* + s(\hat{\theta}_T^M - \theta^*)) ds \hat{u}_T^M$$

since  $\partial_\theta \mathbb{H}_T(\hat{\theta}_T^M) = 0$ . Then we obtain (2.19) from [S3] (iii) and (iv). The  $\mathcal{G}$ -stable convergence

$$\hat{u}_T^M \xrightarrow{d_s(\mathcal{G})} \hat{u} \quad (2.25)$$

follows from (2.18).

As already used in the above argument,

$$P[|\hat{u}_T^M| \geq r] \leq P\left[\sup_{u \in \mathbb{V}_T(r)} \mathbb{Z}_T(u) \geq 1\right] \leq \frac{C_L}{r^L} \quad (2.26)$$

for all  $T \in \mathbb{T}$  and  $r > 0$ . Therefore,

$$\sup_{T \in \mathbb{T}} E[|\hat{u}_T^M|^q] < \infty$$

for any constant  $q$  such that  $L > q > p$ . This means the family  $\{f(\hat{u}_T^M)\}_{T>0}$  is uniformly integrable. Consequently, we obtain (2.20) from (2.25).  $\square$

**Remark 2.8.** (i) The convergence (2.20) holds for non-bounded  $\Phi$  if  $\Phi$  has the dual integrability for  $f(\hat{u}_T^M)$ . For example, the convergence holds for  $\Phi \in L^r(\mathcal{G})$  for some  $r > 1$  if  $\limsup_{|u| \rightarrow \infty} |u|^{-p(r-1)/r} |f(u)| < \infty$ . (ii) The asymptotic equivalence (2.19) between  $\hat{u}_T^M$  and  $\Gamma^{-1}\Delta_T$  is called the first-order efficiency in particular for the maximum likelihood estimator. This relation is useful when one considers a joint convergence of  $\hat{u}_T^M$  with other variables. Such an asymptotic representation of the error is useful in analysis of a model having multi-scaled parameters.

The quasi-likelihood analysis enables us to derive asymptotic properties of the Bayesian estimator, as well as the quasi-maximum likelihood estimator. The mapping

$$\hat{\theta}_T^B = \left[ \int_{\Theta} \exp(\mathbb{H}_T(\theta)) \varpi(\theta) d\theta \right]^{-1} \int_{\Theta} \theta \exp(\mathbb{H}_T(\theta)) \varpi(\theta) d\theta \quad (2.27)$$

is called a quasi-Bayesian estimator (QBE) with respect to the prior density  $\varpi$ . The QBE  $\hat{\theta}_T^B$  takes values in the convex-hull of  $\Theta$ . When the  $\mathbb{H}_T$  is the log likelihood function, the QBE coincides with the Bayesian estimator with respect to the quadratic loss function. We will assume  $\varpi$  is continuous and  $0 < \inf_{\theta \in \Theta} \varpi(\theta) \leq \sup_{\theta \in \Theta} \varpi(\theta) < \infty$ . Let  $\hat{u}_T^B = a_T^{-1}(\hat{\theta}_T^B - \theta^*)$ .

**Theorem 2.9. (I)** *Let  $L > 1$ . Suppose that Conditions [S1], [S2] and [S3] are satisfied and that the convergence (2.18) holds as  $T \rightarrow \infty$ . Then*

(a) *As  $T \rightarrow \infty$ ,*

$$\hat{u}_T^B - \Gamma^{-1}\Delta_T \xrightarrow{p} 0. \quad (2.28)$$

(b) *As  $T \rightarrow \infty$ ,*

$$\hat{u}_T^B \xrightarrow{d_s(\mathcal{G})} \hat{u}. \quad (2.29)$$

**(II)** *Let  $p \geq 0$  and  $L > (p+1) \vee 2$ . Suppose that Conditions [S1], [S2] and [S3] are satisfied and that the convergence (2.18) holds as  $T \rightarrow \infty$ . Moreover, suppose that there exist positive constants  $q$ ,  $c_0$ ,  $\delta$  and  $T_0 \in \mathbb{T}$  such that  $q > p$  and*

$$\sup_{T \geq T_0} E\left[|\mathbb{H}_T(\theta^* + a_T u) - \mathbb{H}_T(\theta^*)|^q\right] \leq c_0 |u|^q \quad (2.30)$$

for all  $u \in U(0, \delta)$ . Then (2.28) holds, and moreover,

$$E[f(\hat{u}_T^B)\Phi] \rightarrow \mathbb{E}[f(\hat{u})\Phi] \quad (2.31)$$

as  $T \rightarrow \infty$  for any  $\mathcal{G}$ -measurable bounded random variable  $\Phi$  and any  $f \in C(\mathbb{R}^p)$  satisfying  $\limsup_{|u| \rightarrow \infty} |u|^{-p}|f(u)| < \infty$ .

**Remark 2.10.** In Theorem 2.9, we implicitly assume that  $T_0$  is sufficiently large so that  $U(0, \delta) \subset \mathcal{U}_T$  for all  $T \geq T_0$ , and the left-hand side of (2.30) makes sense.

**Remark 2.11.** Condition (2.30) holds under any one of the following conditions:

(i) There exist constants  $q > p$ ,  $\delta > 0$  and  $T_0 \in \mathbb{T}$  such that

$$\sup_{T \geq T_0} \sup_{u \in \mathcal{U}_T \cap U(0, \delta)} \|\Gamma_T(\theta^* + a_T u)\|_q < \infty$$

and  $\sup_{T \geq T_0} \|\Delta_T\|_q < \infty$ .

(ii)  $|\Gamma| \in L^q$ , and  $M_1$ ,  $M_3$ , and  $M_4$  appearing in [S3] satisfy  $M_1 \wedge M_3 \wedge M_4 \geq q$ .

This follows from the representation (2.12) of  $\mathbb{Z}_T(u)$  and the representation (2.13) of  $r_T(u)$ .

*Proof of Theorem 2.9.* (I) We obtain a polynomial type large deviation inequality from Theorem 2.3: for any  $D > 0$ , there exist positive constants  $C_1$  and  $C_2$  such that

$$P \left[ \sup_{u \in \mathbb{V}_T(r)} \mathbb{Z}_T(u) \geq C_1 r^{-D} \right] \leq C_2 r^{-L} \quad (2.32)$$

for all  $T$  and  $r > 0$ . Choose a number  $D$  such that  $D > p + (p \vee 1)$ . If we take a sufficiently large constant  $C'_1$ , then

$$\begin{aligned} & P \left[ \int_{\mathbb{V}_T(r)} (1 + |u|) \mathbb{Z}_T(u) du > C'_1 \sum_{\ell=0}^{\infty} (r + \ell)^{p-D} \right] \\ & \leq \sum_{\ell=0}^{\infty} P \left[ \int_{\{r+\ell \leq |u| < (r+\ell+1)\} \cap \mathcal{U}_T} (1 + |u|) \mathbb{Z}_T(u) du > C'_1 (r + \ell)^{p-D} \right] \\ & \leq \sum_{\ell=0}^{\infty} P \left[ \sup_{\{r+\ell \leq |u| < (r+\ell+1)\} \cap \mathcal{U}_T} \mathbb{Z}_T(u) > C_1 (r + \ell)^{-D} \right] \\ & \leq C_2 \sum_{\ell=0}^{\infty} (r + \ell)^{-L} \end{aligned}$$

for  $T \in \mathbb{T}$  and  $r > 1$ , by (2.32). Since  $D - p > 1$  and  $L > 1$  by assumption, there exist positive constants  $C_3$  and  $\epsilon$  (independent of  $(r, T)$ ) such that

$$P \left[ \int_{\mathbb{V}_T(r)} (1 + |u|) \mathbb{Z}_T(u) du > C_3 r^{-\epsilon} \right] \leq C_3 r^{-\epsilon} \quad (r > 0, T \in \mathbb{T}) \quad (2.33)$$

The variable  $\hat{u}_T^B$  has the expression

$$\hat{u}_T^B = \left( \int_{\mathbb{U}_T} \mathbb{Z}_T(u) \varpi(\theta^* + a_T u) du \right)^{-1} \int_{\mathbb{U}_T} u \mathbb{Z}_T(u) \varpi(\theta^* + a_T u) du. \quad (2.34)$$

For  $g(u) = (1, u)$ , let

$$\begin{aligned} X_T &= \int_{\mathbb{U}_T} g(u) \mathbb{Z}_T(u) \varpi(\theta_T^\dagger(u)) du, & X_{T,r} &= \int_{\mathbb{U}_T \cap U(0,r)} g(u) \mathbb{Z}_T(u) \varpi(\theta_T^\dagger(u)) du, \\ W_{T,r} &= \int_{\mathbb{V}_T(r)} g(u) \mathbb{Z}_T(u) \varpi(\theta_T^\dagger(u)) du, \\ X_\infty &= \int_{\mathbb{R}^p} g(u) \mathbb{Z}(u) \varpi(\theta^*) du, & X_{\infty,r} &= \int_{U(0,r)} g(u) \mathbb{Z}(u) \varpi(\theta^*) du, \end{aligned}$$

where  $\theta_T^\dagger(u) = \theta^* + a_T u$  and  $\mathbb{Z}$  is given by (2.16). Then  $X_T = X_{T,r} + W_{T,r}$  and the following properties hold.

- (i) For any  $\eta > 0$ , there exists  $r_0 > 0$  such that  $\sup_{T \in \mathbb{T}} P[|W_{T,r}| > \eta] < \eta$  for all  $r \geq r_0$ .
- (ii) For every  $r > 1$ ,  $X_{T,r} \rightarrow^d X_{\infty,r}$  as  $T \rightarrow \infty$ .
- (iii)  $X_{\infty,r} \rightarrow^d X_\infty$  as  $r \rightarrow \infty$ .

Indeed, (i) follows from (2.33), (ii) from the convergence (2.22), and (iii) is obvious. Therefore

$$X_T \rightarrow^d X_\infty \quad (2.35)$$

as  $T \rightarrow \infty$ .

Denote  $X_T = (X_T^{(0)}, X_T^{(1)})$  and  $X_{T,r} = (X_{T,r}^{(0)}, X_{T,r}^{(1)})$ . We will consider sufficiently large  $T$  such that  $\mathbb{U}_T \supset U(0, 1)$ . Let

$$A_T = (|X_T^{(1)}| + X_T^{(0)}) \left( X_T^{(0)} \int_{U(0,1)} \mathbb{Z}_T(u) du \inf_{\theta \in \Theta} \varpi(\theta) \right)^{-1}.$$

Then

$$A_T \geq \frac{|X_T^{(1)}| + X_T^{(0)}}{X_T^{(0)} X_{T,r}^{(0)}}$$

for all  $r \geq 1$ . We have

$$\left| (X_T^{(0)})^{-1} X_T^{(1)} - (X_{T,r}^{(0)})^{-1} X_{T,r}^{(1)} \right| \leq |W_{T,r}| A_T$$

for all  $r \geq 1$ . Let  $\epsilon > 0$ . Then there exists a positive number  $\eta > 0$  such that

$$\limsup_{T \rightarrow \infty} P \left[ A_T > \frac{1}{\eta} \right] < \frac{\epsilon}{4}$$

since the family  $\{A_T\}_{T \geq T_1}$  is tight for some  $T_1 \in \mathbb{T}$  by (2.35) and (2.22). For the pair  $(\epsilon, \eta)$ , there exists  $r_0 = r_0(\epsilon, \eta) \geq 1$  such that

$$\limsup_{T \rightarrow \infty} P \left[ |W_{T,r}| > \frac{\epsilon \eta}{4} \right] < \frac{\epsilon}{4} \quad (r \geq r_0)$$

by the property (i) mentioned just before (2.35). In what follows, we fix an  $r \geq r_0$ . Then

$$\begin{aligned} & \limsup_{T \rightarrow \infty} P \left[ \left| (X_T^{(0)})^{-1} X_T^{(1)} - (X_{T,r}^{(0)})^{-1} X_{T,r}^{(1)} \right| > \frac{\epsilon}{4} \right] \\ & \leq \limsup_{T \rightarrow \infty} P \left[ |W_{T,r}| > \frac{\epsilon \eta}{4} \right] + \limsup_{T \rightarrow \infty} P \left[ A_T > \frac{1}{\eta} \right] < \frac{\epsilon}{2}. \end{aligned} \quad (2.36)$$

We have

$$\begin{aligned} & \left| \int_{U(0,r)} u^i \mathbb{Z}_T(u) \varpi(\theta_T^\dagger(u)) du - \int_{U(0,r)} u^i \mathbb{Z}_T(u) \varpi(\theta^*) du \right| \\ & \leq \int_{U(0,r)} (1 + |u|) \mathbb{Z}_T(u) du \\ & \quad \times \sup \left\{ |\varpi(\theta) - \varpi(\theta^*)|; \theta \in \Theta, |\theta - \theta^*| \leq |a_T| r \right\} \end{aligned}$$

where  $u^0 = 1$  and  $u^1 = u$ . Therefore

$$\begin{aligned} & \left( \int_{U(0,r)} \mathbb{Z}_T(u) \varpi(\theta_T^\dagger(u)) du \right)^{-1} \int_{U(0,r)} u \mathbb{Z}_T(u) \varpi(\theta_T^\dagger(u)) du \\ & \quad - \left( \int_{U(0,r)} \mathbb{Z}_T(u) du \right)^{-1} \int_{U(0,r)} u \mathbb{Z}_T(u) du \rightarrow^p 0 \end{aligned} \quad (2.37)$$

as  $T \rightarrow \infty$ .

Moreover, we have

$$\begin{aligned} & \left| \int_{U(0,r)} u^i \mathbb{Z}_T(u) du - \int_{U(0,r)} u^i \hat{\mathbb{Z}}_T(u) du \right| \\ & = \left| \int_{U(0,r)} u^i \exp(\Delta_T[u] - 2^{-1} \Gamma[u^{\otimes 2}] + r_T(u)) du - \int_{U(0,r)} u^i \exp(\Delta_T[u] - 2^{-1} \Gamma[u^{\otimes 2}]) du \right| \\ & \leq \int_{U(0,r)} (1 + |u|) \hat{\mathbb{Z}}_T(u) du \sup_{u \in U(0,r)} |e^{r_T(u)} - 1| \end{aligned}$$

where

$$\hat{\mathbb{Z}}_T(u) = \exp(\Delta_T[u] - 2^{-1} \Gamma[u^{\otimes 2}]).$$

Therefore

$$\left( \int_{U(0,r)} \mathbb{Z}_T(u) du \right)^{-1} \int_{U(0,r)} u \mathbb{Z}_T(u) du - \left( \int_{U(0,r)} \hat{\mathbb{Z}}_T(u) du \right)^{-1} \int_{U(0,r)} u \hat{\mathbb{Z}}_T(u) du \rightarrow^p 0 \quad (2.38)$$

as  $T \rightarrow \infty$  thanks to the convergence (2.21).

Let

$$\begin{aligned}\hat{X}_T &= (\hat{X}_T^{(0)}, \hat{X}_T^{(1)}) = \int_{\mathbb{R}^p} g(u) \hat{Z}_T(u) du, & \hat{X}_{T,r} &= (\hat{X}_{T,r}^{(0)}, \hat{X}_{T,r}^{(1)}) = \int_{U(0,r)} g(u) \hat{Z}_T(u) du, \\ \hat{W}_{T,r} &= \int_{\mathbb{R}^p \setminus U(0,r)} g(u) \hat{Z}_T(u) du.\end{aligned}$$

Then

$$\left| (\hat{X}_T^{(0)})^{-1} \hat{X}_T^{(1)} - (\hat{X}_{T,r}^{(0)})^{-1} \hat{X}_{T,r}^{(1)} \right| \leq |\hat{W}_{T,r}| \hat{A}_T$$

for all  $r \geq 1$ , where

$$\begin{aligned}\hat{A}_T &= \left( |\hat{X}_T^{(1)}| + \hat{X}_T^{(0)} \right) \left( \hat{X}_T^{(0)} \int_{U(0,1)} \hat{Z}_T(u) du \inf_{\theta \in \Theta} \varpi(\theta) \right)^{-1} \\ &\geq \frac{|\hat{X}_T^{(1)}| + \hat{X}_T^{(0)}}{\hat{X}_T^{(0)} \hat{X}_{T,r}^{(0)}}\end{aligned}$$

Positive-definiteness of  $\Gamma$  (Remark 2.2 (iii)) and the tightness due to the convergence of  $\Delta_T$  show that for any  $\eta > 0$ , there exist  $r_1 > 0$  and  $T_2 \in \mathbb{T}$  such that

$$\sup_{T \geq T_2} P[\hat{W}_{T,r} > \eta] < \eta \quad (r \geq r_1). \quad (2.39)$$

In the same way as we showed (2.36),

$$\limsup_{T \rightarrow \infty} P \left[ \left| (\hat{X}_T^{(0)})^{-1} \hat{X}_T^{(1)} - (\hat{X}_{T,r}^{(0)})^{-1} \hat{X}_{T,r}^{(1)} \right| > \frac{\epsilon}{4} \right] < \frac{\epsilon}{2} \quad (2.40)$$

for some  $r$ . To obtain (2.40) by using (2.39) and the tightness of  $\{\hat{A}_T\}_{T \geq T_3}$  for some  $T_3 \in \mathbb{T}$ , we replace  $r$  by a larger number, if necessary.

Combining (2.36), (2.37), (2.38) and (2.40), we obtain

$$\limsup_{T \rightarrow \infty} P \left[ \left| (X_T^{(0)})^{-1} X_T^{(1)} - (\hat{X}_T^{(0)})^{-1} \hat{X}_T^{(1)} \right| > \epsilon \right] < \epsilon. \quad (2.41)$$

This completes the proof of (2.28) since  $\hat{u}_T^B = (X_T^{(0)})^{-1} X_T^{(1)}$  and  $(\hat{X}_T^{(0)})^{-1} \hat{X}_T^{(1)} = \Gamma^{-1} \Delta_T$ . From (2.28), we obtain (2.29).

(II) There exists a number  $p_*$  such that

$$p_* \geq 1, \quad p < p_* < (D - \mathbf{p}) \wedge (L - 1)$$

since  $p \geq 0$  and  $L > (p + 1) \vee 2$ . Then the following estimates are standard:

$$\begin{aligned}E[|\hat{u}_T^B|^{p_*}] &\leq E \left[ \left( \int_{\mathcal{U}_T} \mathbb{Z}_T(u) \varpi(\theta_T^\dagger(u)) du \right)^{-1} \int_{\mathcal{U}_T} |u|^{p_*} \mathbb{Z}_T(u) \varpi(\theta_T^\dagger(u)) du \right] \\ &\leq C(\varpi) \sum_{r=0}^{\infty} (r+1)^{p_*} E \left[ \left( \int_{\mathcal{U}_T} \mathbb{Z}_T(u) du \right)^{-1} \int_{\{u; r < |u| \leq r+1\} \cap \mathcal{U}_T} \mathbb{Z}_T(u) du \right] \\ &\leq C(\varpi) (1 + \Phi_{1,T} + \Phi_{2,T})\end{aligned}$$

for some constant  $C(\varpi)$ , where

$$\begin{aligned} \Phi_{1,T} = & \sum_{r=1}^{\infty} (r+1)^{p^*} E \left[ \left( \int_{\cup_T} \mathbb{Z}_T(u) du \right)^{-1} \int_{\{u; r < |u| \leq r+1\} \cap \cup_T} \mathbb{Z}_T(u) du \right. \\ & \left. \times 1 \left\{ \int_{\{u; r < |u| \leq r+1\} \cap \cup_T} \mathbb{Z}_T(u) du > \frac{C'_1}{r^{D-p+1}} \right\} \right] \end{aligned}$$

and

$$\begin{aligned} \Phi_{2,T} = & \sum_{r=1}^{\infty} (r+1)^{p^*} E \left[ \left( \int_{\cup_T} \mathbb{Z}_T(u) du \right)^{-1} \int_{\{u; r < |u| \leq r+1\} \cap \cup_T} \mathbb{Z}_T(u) du \right. \\ & \left. \times 1 \left\{ \int_{\{u; r < |u| \leq r+1\} \cap \cup_T} \mathbb{Z}_T(u) du \leq \frac{C'_1}{r^{D-p+1}} \right\} \right]. \end{aligned}$$

Take a sufficiently large number  $C'_1$ . Since the integrand of the expectation of  $\Phi_{1,T}$  is not greater than one, we obtain

$$\begin{aligned} \Phi_{1,T} & \leq \sum_{r=1}^{\infty} (r+1)^{p^*} P \left[ \int_{\{u; r < |u| \leq r+1\} \cap \cup_T} \mathbb{Z}_T(u) du > \frac{C'_1}{r^{D-p+1}} \right] \\ & \lesssim \sum_{r=1}^{\infty} r^{-(L-p^*)} \end{aligned}$$

thanks to the polynomial type large deviation inequality (2.32). For  $\Phi_{2,T}$ ,

$$\Phi_{2,T} \lesssim \sum_{r=1}^{\infty} r^{-(D-p^*-p+1)} E \left[ \left( \int_{\cup_T} \mathbb{Z}_T(u) du \right)^{-1} \right].$$

Therefore, the family  $\{|\hat{u}_T^B|^p\}_{T \geq T_0}$  is uniformly integrable if

$$\sup_{T \geq T_0} E \left[ \left( \int_{U(0,\delta)} \mathbb{Z}_T(u) du \right)^{-1} \right] < \infty \quad (2.42)$$

since  $U(0,\delta) \subset \cap_{T \geq T_0} \cup_T$  (see Remark 2.10) and then

$$\sup_{T \geq T_0} E [|\hat{u}_T^B|^{p^*}] < \infty.$$

[We note that the family  $\{\hat{u}_T^B\}_{T \in \mathbb{T}, T < T_0}$  is not necessarily uniformly bounded when  $\#\{T < T_0\} = \infty$  because  $\inf_{T < T_0} |a_T|$  may be zero for example, though  $\hat{\theta}_T^B$  takes values in the convex hull of  $\Theta$ , that is bounded.] We obtain the inequality (2.42) from (2.30) by applying e.g. Lemma 2 of [30]. The convergence (2.31) is now obtained from (2.29) and the uniform integrability of the family  $\{|\hat{u}_T^B|^p\}_{T \geq T_0}$ .  $\square$

**Remark 2.12.** From (2.19) and (2.28), the QMLE and the QBE are asymptotically equivalent in that  $\hat{u}_T^M - \hat{u}_T^B \xrightarrow{p} 0$  as  $T \rightarrow \infty$ .

The following conditions [T1] and [T2] strengthen [S2] and [S3], respectively.

[T1] (i) There exists a positive random variable  $\chi_0$  and the following conditions are fulfilled.

(i-1)  $\Upsilon(\theta) = \Upsilon(\theta) - \Upsilon(\theta^*) \leq -\chi_0 |\theta - \theta^*|^2$  for all  $\theta \in \Theta$ .

(i-2) For every  $L > 0$ , there exists a constant  $C$  such that

$$P[\chi_0 \leq r^{-1}] \leq \frac{C}{r^L} \quad (r > 0).$$

(ii) For every  $L > 0$ , there exists a constant  $C$  such that

$$P[\lambda_{\min}(\Gamma) < r^{-1}] \leq \frac{C}{r^L} \quad (r > 0)$$

**Remark 2.13.**  $\chi_0^{-1} \in L^{\infty-} = \cap_{p>1} L^p$  under [T1] (i-2).  $|\Gamma^{-1}| \in L^{\infty-}$  under [T1] (ii) since  $(\lambda_{\min}(\Gamma))^{-1} \in L^{\infty-}$  and  $|\Gamma^{-1}| \leq C_p (\lambda_{\min}(\Gamma))^{-1}$  for a constant only depending on  $p$ . The  $L^q$ -integrability of  $\Gamma$  will be assumed when we verify (2.30).

[T2] There exist positive numbers  $\epsilon_1$  and  $\epsilon_2$  such that the following conditions are satisfied for all  $p > 1$ :

(i)  $\sup_{T \in \mathbb{T}} \|\Delta_T\|_p < \infty$ .

(ii)  $\sup_{T \in \mathbb{T}} \left\| \sup_{\theta \in \Theta} b_T^{\epsilon_1} |\Upsilon_T(\theta) - \Upsilon(\theta)| \right\|_p < \infty$ .

(iii)  $\sup_{T \in \mathbb{T}} \left\| \sup_{u \in \delta^{-1}(\Theta - \theta^*), |u| \leq 1} |\Gamma_T(\theta^* + \delta u) - \Gamma_T(\theta^*)| \right\|_p = O(\delta) \quad (\delta \downarrow 0)$ .

(iv)  $\sup_{T \in \mathbb{T}} \left\| b_T^{\epsilon_2} |\Gamma_T(\theta^*) - \Gamma| \right\|_p < \infty$ .

Condition [T2] requires any order of moments of variables. In applications of the QLA to inference for stochastic differential equations, usually one needs not be nervous about the existence of moments of arbitrary order. Denote by  $f \in C_p(\mathbb{R}^p)$  the set of continuous functions of at most polynomial growth. We can further simplify Theorems 2.7 and 2.9 under Conditions [T1] and [T2].

**Theorem 2.14.** *Suppose that Conditions [T1] and [T2] are satisfied and that the convergence (2.18) holds as  $T \rightarrow \infty$ . Then*

(a) As  $T \rightarrow \infty$ ,

$$\hat{u}_T^M - \Gamma^{-1} \Delta_T \xrightarrow{p} 0. \quad (2.43)$$



(b) As  $T \rightarrow \infty$ ,

$$E[f(\hat{u}_T^M)\Phi] \rightarrow \mathbb{E}[f(\hat{u})\Phi] \quad (2.44)$$

for any  $f \in C_p(\mathbb{R}^p)$  and any  $\mathcal{G}$ -measurable random variable  $\Phi \in \cup_{p>1} L^p$ .

*Proof.* There exist values of the parameters  $\alpha, \beta_1, \beta_2, \rho_1$  and  $\rho_2$  satisfying  $\beta_1 \in (0, \min\{\epsilon_2, 1/2\})$ ,  $1/2 - \beta_2 \leq \epsilon_1$  and Condition [S1]. Then [S2] is verified for any given  $L > 0$  by [T1]. Condition [T2] is sufficient for [S3] for any  $L > 0$ . Therefore we can apply Theorems 2.7. This concludes the proof.  $\square$

**Theorem 2.15.** *Suppose that Conditions [T1] and [T2] are satisfied and that the convergence (2.18) holds as  $T \rightarrow \infty$ . Moreover, suppose that  $|\Gamma| \in L^q$  for some  $q > p$ . Then*

(a) As  $T \rightarrow \infty$ ,

$$\hat{u}_T^B - \Gamma^{-1}\Delta_T \xrightarrow{p} 0. \quad (2.45)$$

(b) As  $T \rightarrow \infty$ ,

$$E[f(\hat{u}_T^B)\Phi] \rightarrow \mathbb{E}[f(\hat{u})\Phi] \quad (2.46)$$

for any  $f \in C_p(\mathbb{R}^p)$  and any  $\mathcal{G}$ -measurable random variable  $\Phi \in \cup_{p>1} L^p$ .

*Proof.* We apply Theorem 2.9. In particular, (2.30) holds now according to Remark 2.11.  $\square$

### 3 Further simplification in ergodic statistics

When the limit  $\mathbb{Y}$  of  $\mathbb{Y}_T$  is deterministic, more simplification of the theory is possible. In this section, we suppose that the random field  $\mathbb{Y}$  and the  $p \times p$  positive-definite symmetric matrix  $\Gamma$  are deterministic. Let  $L$  be a positive number. We will consider the following conditions. Some simplification has been made in Condition [U2], that is slightly different from [S1] plus [S3].

[U1]  $\Gamma$  is positive-definite, in addition, there is a positive number  $\chi_0$  such that

$$\mathbb{Y}(\theta) = \mathbb{Y}(\theta) - \mathbb{Y}(\theta^*) \leq -\chi_0|\theta - \theta^*|^2$$

for all  $\theta \in \Theta$ .

[U2] The numbers  $\alpha, \beta_1, \beta_2$  and  $\rho_2$  satisfy the inequalities

$$0 < 2\alpha < \rho_2, \quad \beta_2 \geq 0, \quad 1 - 2\beta_2 - \rho_2 > 0, \quad 0 < \beta_1 < 1/2, \quad (3.1)$$

and the following conditions are fulfilled.

(i) For some  $M_1 > L$ ,  $\sup_{T \in \mathbb{T}} \|\Delta_T\|_{M_1} < \infty$ .

(ii) For  $M_2 = L(1 - 2\beta_2 - \rho_2)^{-1}$ ,

$$\sup_{T \in \mathbb{T}} \left\| \sup_{\theta \in \Theta \setminus U(\theta^*, b_T^{-\alpha/2})} b_T^{\frac{1}{2} - \beta_2} |\Upsilon_T(\theta) - \Upsilon(\theta)| \right\|_{M_2} < \infty.$$

(iii) For some  $M_3 > L\beta^{-1}$ ,

$$\sup_{T \in \mathbb{T}} \left\| \sup_{u \in \delta^{-1}(\Theta - \theta^*), |u| \leq 1} |\Gamma_T(\theta^* + \delta u) - \Gamma_T(\theta^*)| \right\|_{M_3} = O(\delta) \quad (\delta \downarrow 0).$$

(iv) For some  $M_4 > L(2\beta_1)^{-1}(1 - \alpha)$ ,

$$\sup_{T \in \mathbb{T}} \|b_T^{\beta_1} |\Gamma_T(\theta^*) - \Gamma|\|_{M_4} < \infty.$$

**Remark 3.1.** Condition [U1] is almost trivial because the function  $\Upsilon$  should be of  $C^2$  on  $\Theta$  and continuous on the compact set  $\bar{\Theta}$  and then local non-degeneracy of the information implies the global identifiability.

**Theorem 3.2.** *Suppose that Conditions [U1] and [U2] are fulfilled for a positive constant  $L$ . Then there exists a constant  $C_L$  such that*

$$P \left[ \sup_{u \in \bar{\mathbb{V}}_T(r)} \mathbb{Z}_T(u) \geq \exp(-2^{-1}r^{2-\rho_2}) \right] \leq \frac{C_L}{r^L}$$

for all  $T \in \mathbb{T}$  and  $r > 0$ . Here the supremum on the empty set should read  $-\infty$  by convention.

*Proof.* Choose a positive constant  $\rho_1$  such that

$$\begin{aligned} 0 < \rho_1 &< \min \{1, \alpha/(1 - \alpha), 2\beta_1/(1 - \alpha)\}, \quad \rho_1 \leq \rho_2, \\ M_1 &\geq M'_1 := L(1 - \rho_1)^{-1}, \quad M_3 \geq M'_3 := L(\beta - \rho_1)^{-1}, \\ M_4 &\geq M'_4 := L(2\beta_1(1 - \alpha)^{-1} - \rho_1)^{-1} \end{aligned} \quad (3.2)$$

for  $M_1, M_2, M_3$  given in Condition [U2]. Such a positive number  $\rho_1$  exists. It is sufficient to verify the conditions of Theorem 2.3. Condition [S1] is fulfilled by [U2] and a choice of  $\rho_1$  in (3.2). Condition [S2] (i-1) is satisfied with [U1], and Conditions (i-2) and (ii) of [S2] are trivial because  $\chi_0$  is a deterministic positive number and  $\Gamma$  is positive-definite, deterministic in the present situation, respectively. Conditions (i)-(iv) of [S3] are verified by [U2] with  $(M'_1, M_2, M'_3, M'_4)$  for  $(M_1, M_2, M_3, M_4)$  in [S3].  $\square$

As before, the quasi-maximum likelihood estimator (QMLE) for  $\mathbb{H}_T$  is characterized by (2.17). Theorem 2.7 is rephrased as follows with the trivial  $\sigma$ -field for  $\mathcal{G}$ .

**Theorem 3.3.** *Let  $L > p > 0$ . Suppose that Conditions [U1] and [U2] are satisfied and that*

$$\Delta_T \rightarrow^d \Delta \quad (3.3)$$

as  $T \rightarrow \infty$ . Then,

- (a)  $\hat{u}_T^M - \Gamma^{-1}\Delta_T \xrightarrow{p} 0$  as  $T \rightarrow \infty$ .
- (b)  $E[f(\hat{u}_T^M)] \rightarrow E[f(\hat{u})]$  as  $T \rightarrow \infty$  for any  $f \in C(\mathbb{R}^p)$  satisfying  $\limsup_{|u| \rightarrow \infty} |u|^{-p}|f(u)| < \infty$ .

*Proof.* Take  $\rho_1$  as (3.2) and apply Theorem 2.7.  $\square$

Consider the quasi-Bayesian estimator  $\hat{\theta}_T^B$  defined by (2.27). We can rephrase Theorem 2.9 as follows.

**Theorem 3.4.** *Let  $p \geq 0$  and  $L > (p + 1) \vee 2$ . Suppose that Conditions [U1] and [U2] are satisfied and that the convergence (3.3) holds as  $T \rightarrow \infty$ . Moreover, suppose that there exist positive constants  $q, \delta, T_0 \in \mathbb{T}$  and  $c_0$  such that  $q > p$  and the inequality (2.30) holds for all  $u \in U(0, \delta)$ . Then*

- (a)  $\hat{u}_T^B - \Gamma^{-1}\Delta_T \xrightarrow{p} 0$  as  $T \rightarrow \infty$ .
- (b)  $E[f(\hat{u}_T^B)] \rightarrow E[f(\hat{u})]$  as  $T \rightarrow \infty$  for any  $f \in C(\mathbb{R}^p)$  satisfying  $\limsup_{|u| \rightarrow \infty} |u|^{-p}|f(u)| < \infty$ .

As a corollary of Theorem 2.14 (or Theorem 3.3) and Theorem 2.15 (or Theorem 3.4), we obtain the following result.

**Theorem 3.5.** *Suppose that Conditions [U1] and [T2] are satisfied and that the convergence (3.3) holds as  $T \rightarrow \infty$ . Then*

- (a)  $\hat{u}_T^A - \Gamma^{-1}\Delta_T \xrightarrow{p} 0$  as  $T \rightarrow \infty$  for  $A \in \{M, B\}$ .
- (b)  $E[f(\hat{u}_T^A)] \rightarrow E[f(\hat{u})]$  as  $T \rightarrow \infty$  for  $A \in \{M, B\}$  and any  $f \in C_p(\mathbb{R}^p)$ .

## References

- [1] Clinet, S., Yoshida, N.: Statistical inference for ergodic point processes and application to limit order book. arXiv preprint arXiv:1512.01899 (2015)
- [2] Eguchi, S., Masuda, H.: Schwarz type model comparison for laq models. arXiv preprint arXiv:1606.01627 (2016)
- [3] Ibragimov, I., Khas'minskii, R.: Asymptotic behavior of statistical estimators in the smooth case. I. Study of the likelihood ratio. *Theory of Probability & Its Applications* **17**(3), 445–462 (1973)
- [4] Ibragimov, I., Khas'minskii, R.: Asymptotic behavior of some statistical estimators II. Limit theorems for the a posteriori density and Bayes' estimators. *Theory of Probability & Its Applications* **18**(1), 76–91 (1973)
- [5] Ibragimov, I.A., Has'minskiĭ, R.Z.: Statistical estimation, Applications of Mathematics, vol. 16. Springer-Verlag, New York (1981). Asymptotic theory, Translated from the Russian by Samuel Kotz

- [6] Inatsugu, H., Yoshida, N.: Global jump filters and quasi-likelihood analysis for volatility. *Annals of the Institute of Statistical Mathematics*: updated arXiv:1806.10706v3 pp. 1–44 (2021)
- [7] Kamatani, K., Uchida, M.: Hybrid multi-step estimators for stochastic differential equations based on sampled data. *Statistical Inference for Stochastic Processes* **18**(2), 177–204 (2014)
- [8] Kinoshita, Y., Yoshida, N.: Penalized quasi likelihood estimation for variable selection. arXiv preprint arXiv:1910.12871 (2019)
- [9] Kutoyants, Y.A.: Parameter estimation for stochastic processes, Research and Exposition in Mathematics, vol. 6. Heldermann Verlag, Berlin (1984). Translated from the Russian and edited by B. L. S. Prakasa Rao
- [10] Kutoyants, Y.A.: Identification of dynamical systems with small noise, Mathematics and its Applications, vol. 300. Kluwer Academic Publishers Group, Dordrecht (1994)
- [11] Kutoyants, Y.A.: Statistical inference for spatial Poisson processes, Lecture Notes in Statistics, vol. 134. Springer-Verlag, New York (1998)
- [12] Kutoyants, Y.A.: Statistical inference for ergodic diffusion processes. Springer Series in Statistics. Springer-Verlag London Ltd., London (2004)
- [13] Masuda, H.: Approximate self-weighted LAD estimation of discretely observed ergodic ornstein-uhlenbeck processes. *Electronic Journal of Statistics* **4**, 525–565 (2010)
- [14] Masuda, H.: Convergence of Gaussian quasi-likelihood random fields for ergodic lévy driven SDE observed at high frequency. *The Annals of Statistics* **41**(3), 1593–1641 (2013)
- [15] Masuda, H.: Parametric estimation of Lévy processes. In: *Lévy Matters IV*, pp. 179–286. Springer (2015)
- [16] Masuda, H., Shimizu, Y.: Moment convergence in regularized estimation under multiple and mixed-rates asymptotics. *Mathematical Methods of Statistics* **26**(2), 81–110 (2017)
- [17] Nomura, R., Uchida, M.: Adaptive bayes estimators and hybrid estimators for small diffusion processes based on sampled data. *Journal of the Japan Statistical Society* **46**(2), 129–154 (2016)
- [18] Ogihara, T., Yoshida, N.: Quasi-likelihood analysis for the stochastic differential equation with jumps. *Stat. Inference Stoch. Process.* **14**(3), 189–229 (2011). DOI 10.1007/s11203-011-9057-z. URL <http://dx.doi.org/10.1007/s11203-011-9057-z>
- [19] Ogihara, T., Yoshida, N.: Quasi-likelihood analysis for nonsynchronously observed diffusion processes. *Stochastic Processes and their Applications* **124**(9), 2954–3008 (2014)
- [20] Ogihara, T., Yoshida, N.: Quasi likelihood analysis of point processes for ultra high frequency data. arXiv preprint arXiv:1512.01619 (2015)

- [21] Shimizu, Y.: Moment convergence of regularized least-squares estimator for linear regression model. *Annals of the Institute of Statistical Mathematics* **69**(5), 1141–1154 (2017)
- [22] Shimizu, Y.: Threshold estimation for stochastic processes with small noise. *Scandinavian Journal of Statistics* **44**(4), 951–988 (2017)
- [23] Suzuki, T., Yoshida, N.: Penalized least squares approximation methods and their applications to stochastic processes. *Japanese Journal of Statistics and Data Science* pp. 1–29 (2020)
- [24] Uchida, M.: Contrast-based information criterion for ergodic diffusion processes from discrete observations. *Ann. Inst. Statist. Math.* **62**(1), 161–187 (2010). DOI 10.1007/s10463-009-0245-1. URL <http://dx.doi.org/10.1007/s10463-009-0245-1>
- [25] Uchida, M., Yoshida, N.: Adaptive estimation of an ergodic diffusion process based on sampled data. *Stochastic Process. Appl.* **122**(8), 2885–2924 (2012). DOI 10.1016/j.spa.2012.04.001. URL <http://dx.doi.org/10.1016/j.spa.2012.04.001>
- [26] Uchida, M., Yoshida, N.: Adaptive estimation of an ergodic diffusion process based on sampled data. *Stochastic Processes and their Applications* **122**(8), 2885–2924 (2012)
- [27] Uchida, M., Yoshida, N.: Quasi likelihood analysis of volatility and nondegeneracy of statistical random field. *Stochastic Process. Appl.* **123**(7), 2851–2876 (2013). DOI 10.1016/j.spa.2013.04.008. URL <http://dx.doi.org/10.1016/j.spa.2013.04.008>
- [28] Uchida, M., Yoshida, N.: Adaptive Bayes type estimators of ergodic diffusion processes from discrete observations. *Statistical Inference for Stochastic Processes* **17**(2), 181–219 (2014)
- [29] Umezū, Y., Shimizu, Y., Masuda, H., Ninomiya, Y.: AIC for the non-concave penalized likelihood method. *Annals of the Institute of Statistical Mathematics* **71**(2), 247–274 (2019)
- [30] Yoshida, N.: Polynomial type large deviation inequalities and quasi-likelihood analysis for stochastic differential equations. *Ann. Inst. Statist. Math.* **63**(3), 431–479 (2011). DOI 10.1007/s10463-009-0263-z. URL <http://dx.doi.org/10.1007/s10463-009-0263-z>
- [31] Yoshida, N.: Partial quasi-likelihood analysis. *Japanese Journal of Statistics and Data Science* **1**(1), 157–189 (2018)