

Thoughts on the Proteins Native State

Jorge A. Vila

IMASL-CONICET, Universidad Nacional de San Luis, Ejército de Los Andes 950, 5700 San Luis, Argentina.

Abstract

The presence of metamorphism in the protein's native state is not yet fully understood. In an attempt to throw light on this issue here we present an assessment, in terms of the amide hydrogen exchange protection factor, that aims to determine the likely existence of structural fluctuations in the native-state consistent with both the upper bound marginal stability of proteins and the metamorphism presence. The preliminary results enable us to conclude that the native-state metamorphism is, indeed, more probable than thought.

Introduction

Let us start recalling the research that lead Anfinsen (1973) to obtain the Nobel price “... *The work that my colleagues and I have carried out on the nature of the process that controls the folding of polypeptide chains into the unique three-dimensional structures of proteins was, indeed, strongly influenced by observations on the ribonuclease molecule...*” Now, what a “unique three-dimensional structure” for Anfinsen is? In his own words “...*is the one in which the Gibbs free energy of the whole system is lowest...*” (Anfinsen, 1973). There is no contradiction between Anfinsen's unique structure concept and the existence of an ensemble of folded conformations coexisting with the native state (McCammon, 1977; Matthews, 1987; Albert, 1989; Lange *et al.*, 2008; Martin & Vila, 2020), essentially, because all those conformations must possess higher Gibbs free energy, although comparable, to that of the native state (Vila 2019). From a statistical-mechanics point of view, this means that the native form will be the most probable form of the ensemble and any other conformation will have a representation consistent with their Boltzmann factor.

From the above analysis two basic questions arise. Firstly, which is the origin of the ensemble of folded conformations in equilibrium with the native state? Secondly, how large the structural difference among the conformations of the ensemble could be?

Regarding the first question, it is well known that the native state is marginally stable (Matthews, 1987; Albert, 1989; Dill, 1990) and, hence, stabilized by weak interactions, such as those due to the interplay of pairwise and many-body interactions on both the proteins and the solvent (Frauenfelder *et al.*, 1991; Fernández *et al.* 2002; Ejtehadi *et al.*, 2004; Martin & Vila, 2020); therefore, one may expect transient fluctuations in the native structure to occur at a sizeable level. This is a crucial issue since it contains the origin of the protein native-state metamorphism (Tuinstra *et al.*, 2008; Murzin, 2008; Luo & Yu, 2008; Lella *et al.*, 2017), which is a highly important phenomenon in structural biology (Dishman *et al.*, 2018) and protein evolvability (Murzin, 2008; Tokuriki & Tawfik, 2009; Yadid *et al.*, 2010; Goldstein, 2011; Dishman, *et al.*, 2021). Illustrations of this phenomenon are abundant and some of them have been recently reviewed in light of Anfinsen's dogma (Vila, 2020); indeed, we have been able to show that the coexistence among highly-similar folded states, *e.g.*, for ubiquitin (Lange *et al.*, 2008), α -chymotrypsin, ribonuclease A, cytochrome *c*, etc. (Privalov & Tsalkova, 1979), or between highly-dissimilar folded states, *e.g.*, Mad2 (Luo & Yu, 2008) and lymphotactin (Ltn) (Tuinstra *et al.*, 2008), did take place and satisfied the thermodynamic hypothesis (Anfinsen, 1973).

An explanation to the second question, stated above, is not so straightforward and, hence, it will be investigated below in terms of the amide hydrogen exchange protection factor (Bahar *et al.*, 1998; Persson & Halle, 2015). Before we embark on elaborate a possible solution to this problem, let us remember that proteins that fulfill the Anfinsen dogma possess an upper bound marginal stability, namely $\Delta G \sim 7.4$ kcal/mol (Vila, 2019; Martin & Vila, 2020). This implies that the largest free-energy difference among any conformation of the ensemble of folded states and the native-state (see Figure 1) must be lower than such upper bound, otherwise, the conformation will unfold, *i.e.*, will become nonfunctional (Martin & Vila, 2020). As to whether such upper bound protein marginal stability limit ($\Delta G \sim 7.4$ Kcal/mol) is large enough to assure the existence of significant topological differences in the ensemble of conformations, *i.e.*, to account for the existence of proteins native state metamorphisms, will be here examined in terms of the amide Hydrogen eXchange (HX) rate which is a sensitive probe to the stability of proteins (Privalov &

Tsalkova, 1979; Mayo & Baldwin, 1993; Bai *et al.*, 1994; Bahar *et al.*, 1998; Persson & Halle, 2015).

Materials and Methods

The use of hydrogen exchange for the purpose of determine protein stability (Privalov & Tsalkova, 1979; Huyghues-Despointes *et al.*, 1999; Craig *et al.*, 2011; Balasubramaniam & Komives, 2013; Persson & Halle, 2015) is well known since pioneer experiments of Linderstrøm-Lang and coworkers (Hvidt & Linderstrøm-Lang, 1954; Berger & Linderstrøm-Lang, 1957; Englander *et al.*, 1997; Schellman & Schellman, 1997) and hence will not be here reviewed.

When a protein is transferred from water to deuterium there is a fast exchange of protons exposed to the solvent, namely an isotopic exchange. However, the backbone amide protons exchange will not take place if they are participating in intramolecular hydrogen bonding unless a transient opening in the native state of the protein occurs. Since both the intramolecular hydrogen bonding and the solvent shielding are straightforwardly related to the protein native state structure, it is reasonable to assume that the amide hydrogen exchange could be used as a sensitive probe to appraise changes in the protein topology (Privalov & Tsalkova, 1979). Hence, adopting the kinetic scheme of Bahar *et al.* (1998) in the EX2 limit, which describe the amide HX rate in the native state, the following relation should hold:

$$\Delta G_{\text{HX}} = -RT \ln K_{\text{op}} \quad (1)$$

where ΔG_{HX} represent the Gibbs free-energy change for the opening/closing equilibrium (Bai *et al.*, 1994; Bahar *et al.*, 1998; Craig *et al.*, 2011); the conversion of HX rates to Gibbs free-energies is possible because in the EX2 limit “...*the observed exchange rates are a measure of the equilibrium fraction of molecules that are available for exchange...*” (Mayo & Baldwin, 1993), R the gas constant, T the absolute temperature, $K_{\text{op}} = 1/P_{\text{f}}$ is the equilibrium constant of the opening process (Bahar *et al.*, 1998; Craig *et al.*, 2011) and P_{f} the protection factor, which provides a measure of the resistance of the amide hydrogen to exchange in the native state relative to that of a random polypeptide chain (Bahar *et al.*, 1998). However, our interest is focused on a particular region of the protein conformational space, namely, the ensemble of folded states in equilibrium with the native state (the yellow region in Figure 1). Therefore, the foresee value for the largest

Gibbs free-energy change should be $\Delta G_{\text{HX}} \approx \Delta G$ (~ 7.4 kcal/mol). Under this conjecture, the protection factor (P_f) computed with equation (1) should represent the resistance of amide hydrogen to exchange in the native state relative to that of the highest free-energy conformation in the ensemble of folded states. Consequently, at room temperature, eq. (1) will give a $P_f \sim 10^5$.

Results and Conclusions

On average, the amide HX rate in proteins would be reduced by a factor $\geq 10^8$ from their exchange rates in unfolded peptides (Zhang & Smith, 1993). Therefore, it is reasonable that the amide HX rate for the native state should have a smaller protection factor, such as the computed $\sim 10^5$. In this regard, it is worth noting that the computed upper bound for the protection factor ($\sim 10^5$) does not reflect what can be observed at ‘residue-level’. Indeed, the β_2 region of staphylococcal nuclease can reach P_f ‘peak-values’ of $\sim 10^5$ (Bahar *et al.*, 1998); while, a much larger ‘local P_f ’ can be found in other proteins, such as for the β_1 region of Bovine Pancreatic Trypsin Inhibitor (BPTI) or the α_5 region of cytochrome *c*, where P_f ’s can reach peak-values of $\sim 10^9$ and $\sim 10^{10}$, respectively (Bahar *et al.*, 1998). This contrast, in terms of the higher local P_f ’s, seen on staphylococcal nuclease, BPTI and cytochrome *c* (*i*) illustrate that the ‘local’ protection factors can certainly vary significantly along the amino-acid sequence, *i.e.*, this phenomenon merely indicates the high or low resistance of the amide hydrogens to exchange; though, the structural changes, when occur, do not need to be necessarily large because hydrogens are small atoms. However, as noted by Bahar *et al.* (1998), it is conceivable that some ‘largest’ structural shift could arise as a consequence of several small individual changes acting cooperatively; and (*ii*) suggest that the ‘local structural changes’ in the native state depends on many factors, among others, the local packing density, the topology of tertiary contacts, changes in the milieu, *etc.* (Bahar *et al.*, 1998), making the forecast of the extent of local structural changes, for a given sequence, very hard to determine accurately; a problem that will be exacerbated in the absence of a three-dimensional structure. Certainly, the discussion of this problem is mattering, among other reasons, because it could be straightforwardly related to (*i*) the so-called ‘fold-switching regions in proteins’, a phenomenon that seems to be more common than currently believed (Bryan & Orban, 2010; Porter & Looger, 2018); and (*ii*) pieces of evidence showing that the native state could be a metastable state with respect to the formation of other molecular complexes (Honeycutt & Thirumalai, 1990; Dinner & Karplus, 1998; Thirumalai &

Reddy, 2011; Baldwin, *et al.*, 2011); although, discussion of these issues is beyond the scope of the current work.

The main drawback of our analysis, so far, is that we fail to provide conclusive proof on the ‘magnitude’ of the local structural changes that could take place in the ensemble of folded states in equilibrium with the native state. The reason for this failure obeys, as explained above, the fact the largest local structural changes in native folds are not a universal feature of protein’s but one depending, mainly, on the interactions of the residues with their nearest neighbors; in other words, it is determined by the fold-class and amino-acid sequence. This means that the only known universal feature of the native state of proteins is the existence of marginal stability upper bound ($\Delta G \sim 7.4$ kcal/mol). Therefore, the computed amide HX protection factor upper bound ($\sim 10^5$) should also be, by definition, a universal feature of proteins and, consequently, enable us to suggest the existence of an ensemble of folded states that may exhibit significant structural differences, relative to their native state, independent of the fold-class, sequence or whether the protein contains, or could form, (quasi) independent domains (Vila, 2019); with the warning that the folded states ratio is determined by their Boltzmann factors. Then, how is that more than one native fold could be energetically favored? (Lella *et al.*, 2017). A simple and feasible solution to this enigma is the following. Let us assume that small spontaneous changes in the milieu may indeed happens (Murzin, 2008; Bryan & Orban, 2010), *i.e.*, on the solvent, pH, ionic strength, metal ions, temperature, etc. This could allow redistribution of the Boltzmann factors and, hence, a new thermodynamic equilibrium between highly-dissimilar (metamorphic) folded states would in fact be possible, *e.g.*, the relative abundance of coexistent folded states on the aforementioned metamorphic proteins Mad2 and Ltn is, indeed, a thermodynamic-driven process (Luo & Yu, 2008). It is worth noting that a switch between native-folds can also be trigger by mutations or cellular stimuli (Porter & Looger, 2018).

Overall, the take-home message of this work is twofold. Firstly, there is not a unique conformation representing protein's native states, *i.e.*, proteins are not monomorphic as is recurrently assume, but an ensemble of native-folds that coexist in a narrow range of free-energy variations determined by the physical-origin of the marginal stability, which is a universal property of proteins and molecular complexes (Vila, 2019; Martin & Vila 2020). Secondly, the aforementioned protein marginal stability is large enough to enable us suggest that native state metamorphism in proteins could be a highly probable phenomena; as inferred from the computed

amide hydrogen exchange protection factor upper bound for the ensemble of folded states ($\sim 10^5$). This conclusion, in addition, enables us rationalize the origin of the widespread observation, in the structural biology field, that “...*Proteins that can adopt more than one native folded conformation may be more common than previously thought...*” Murzin (2008).

Acknowledgments

I am honored to dedicate this manuscript to the memory of Harold A. Scheraga, Professor of Chemistry, Emeritus at Cornell University. He achieved leadership in the world of science, and high respect among colleagues, as a result of his colossal experience in Experimental and Theoretical Chemistry, Physics and Mathematics, research in Protein Chemistry and, in particular, due to his tireless efforts in the search of possible solutions to the Protein Folding problem. Harold Scheraga passed away at the age of 98 in Ithaca, NY, on August 1st, 2020.

The author acknowledges financial support from the IMASL-CONICET (PIP-0087) and ANPCyT (Grants PICT-0767; PICT-02212), Argentina.

Corresponding author

Jorge A. Vila jv84@cornell.edu

<https://orcid.org/0000-0001-7557-9350>




Figure 1. The shown schematic representation of the proteins conformational space aims to highlight the relative free-energy difference between the ensemble of folded states (yellow zone) and that of the unfolded states (green zone). The relative size of both ensembles is far from real and have only an illustrative purpose. The ‘unique three-dimensional structure’ representing the native state, according to Anfinsen's dogma, is highlighted in blue.

References

- Albert, T. Mutational effects on protein stability. *Annu. Rev. Biochem.* 58, 765-798 (1989)
- Anfinsen, C.B. Principles that govern the folding of protein chains. *Science* 181, 223-230 (1973).
- Bahar, I., Wallqvist, A., Covell, D.G. and Jernigan, R.L. Correlation between native-state hydrogen exchange and cooperative residue fluctuation from a simple model. *Biochemistry* 37, 4, 1067–1075 (1998)
- Bai, Y., Milne, J.S., Mayne, L., Englander, S.W. Protein stability parameters measured by hydrogen exchange. *Proteins*, 20, 4-14 (1994).
- Balasubramaniam D, Komives E.A. Hydrogen-exchange mass spectrometry for the study of intrinsic disorder in proteins. *Biochim. Biophys. Acta* 1834(6), 1202-1209 (2013)
- Baldwin, A.J. *et al.* Metastability of native proteins and the phenomenon of amyloid formation. *J. Am. Chem. Soc.* 133, 14160-14163 (2011)
- Berger A., Linderstrøm-Lang K. Deuterium exchange of poly-DL-alanine in aqueous solution. *Arch Biochem Biophys* 69, 106-118 (1957)
- Bryan, P.N. and Orban, J. Proteins that switch folds. *Current Opinion in Structural Biology* 20, 482-488 (2010)
- Craig, P.O., Lätzer, J., Weinkam, P., Hoffman, R.M.B., Ferreira, D.U., Komives, E.A. and Wolynes, P.G. Prediction of Native-State Hydrogen Exchange from Perfectly Funneled Energy Landscapes. *JACS*, 133(43), 17463-17472 (2011)
- Dill, K.A. Dominant forces in protein folding. *Biochemistry*, 29(31), 7133-7154 (1990)
- Dinner, A., Karplus, M.A. Metastable state in folding simulations of a protein model. *Nat. Struct. Mol. Biol.* 5, 236-241 (1998)
- Dishman, A.F. and Volkman, B.F. Unfolding the Mysteries of Protein Metamorphosis. *ACS Chemical Biology* 13(6) 1438-1446 (2018)
- Dishman, A.F., Tyler, R.C., Fox, J., Kleist, A.B., Prehoda, K.E., Babu, M.M., Peterson, F.C. and Volkman, B.F. Evolution of fold switching in a metamorphic protein. *Science* 371, Issue 6524, pp. 86-90 (2021)

- Ejtehadi M.R., Avall S.P., and Plotkin S.S. Three-body interactions improve the prediction of rate and mechanism in protein folding models. *Proc Natl Acad Sci USA*, 101 (42) 15088-15093 (2004)
- Englander S.W., Mayne L., Bai Y. and Sosnick T.R. Hydrogen exchange: the modern legacy of Linderstrøm-Lang. *Protein Sci* 6, 1101-1109 (1997)
- Fernández A., Colubri A., Stephen Berry R. Three-body correlations in protein folding: the origin of cooperativity, *Physica A: Statistical Mechanics and its Applications*, 307, 235-259 (2002)
- Frauenfelder H., Sligar S.G., Wolynes P.G. The energy landscapes and motions of proteins. *Science* 254(5038), 1598-1603 (1991)
- Goldstein, R.A. The evolution and evolutionary consequences of marginal thermostability in proteins. *Proteins* 79, 1396-1407 (2011)
- Honeycutt, J.D. and Thirumalai, D. Metastability of the folded states of globular proteins. *Proc. Natl Acad. Sci.* 87, 3526-3529 (1990)
- Huyghues-Despointes, B., Scholtz, J. and Pace, C. Protein conformational stabilities can be determined from hydrogen exchange rates. *Nat. Struct. Mol. Biol.* 6, 910–912 (1999)
- Hvidt A, Linderstrøm-Lang K. Exchange of hydrogen atoms in insulin with deuterium atoms in aqueous solutions. *Biochim. Biophys Acta* 14, 574-575 (1954)
- Lange, O.F., Lakomek, N.-A., Farès, C., Schröder, G.F., Walter, K.F.A., Becker, S., Meiler, J., Grubmüller, H., Griesinger, C. and de Groot, B.L. Recognition Dynamics Up to Microseconds Revealed from an RDC-Derived Ubiquitin Ensemble in Solution. *Science*, 13, 1471-1475 (2008)
- Lella, M. and Mahalakshmi, R. Metamorphic Proteins: Emergence of Dual Protein Folds from One Primary Sequence. *Biochemistry* 56(24), 2971-2984 (2017)
- Luo, X. and Yu, H. Protein Metamorphosis: The two-state behavior of Mad2. *Structure*, 16 (11), 1616-1625 (2008).
- Martin, A.O. and Vila, J.A. The Marginal Stability of Proteins: How the Jiggling and Wiggling of Atoms is Connected to Neutral Evolution. *Journal Molecular Evolution* 88, 424-426 (2020)
- Matthews, B.W. Genetic and structural analysis of the protein stability problem. *Biochemistry*, 26(22), 6885-6888 (1987)

- Mayo, S., and Baldwin, R. Guanidinium Chloride Induction of Partial Unfolding in Amide Proton Exchange in RNase A. *Science*, 262(5135), 873-876 (1993)
- McCammon J.A., Gelin B.R. and Karplus M. Dynamics of folded proteins. *Nature*. 267, 585-590 (1977)
- Murzin, A.G. Biochemistry. Metamorphic proteins. *Science* 320, 1725-1726 (2008)
- Persson F. and Halle B. How amide hydrogens exchange in native proteins *Proc. Natl. Acad. Sci. USA*, 112 (33) 10383-10388 (2015)
- Porter L.L, Looger L.L. Extant fold-switching proteins are widespread. *Proc. Natl. Acad. Sci. USA* 115(23), 5968-5973 (2018)
- Privalov, P.L. & Tsalkova, T.N. Micro- and macro-stabilities of globular proteins. *Nature* 280, 693-696 (1979)
- Schellman J.A. and Schellman C.G. Kaj Ulrik Linderstrøm-Lang (1896- 1959) *Protein Science*, 6, 1092-1100, (1997)
- Sheu, S.-Y., Yang, D.-Y., Selzle, H.L. and Schlag, E.W. Energetic of hydrogen bonds in peptides. *Proc. Natl. Acad. Sci USA* 100, 12683-12687 (2003)
- The Feynman Lectures on Physics, Volume 1, Chapter 3, The Relation of Physics to Other Sciences (<http://www.feynmanlectures.caltech.edu/info/>).
- Thirumalai, D., Reddy, G. Are native proteins metastable? *Nature Chem* 3, 910-911 (2011)
- Tokuriki, N. and Tawfik, D.S. Stability effects of mutations and protein evolvability *Current Opinion in Structural Biology* 19, 596-604 (2009)
- Tuinstra, R.L., Peterson, F.C., Kutlesa, S., Elgin, E.S., Kron, M.A. and Volkman, B.F. Interconversion between two unrelated protein folds in the lymphotactin native state. *Proc. Natl. Acad. Sci. USA* 105, 5057-5062 (2008)
- Vila, J.A. Forecasting the upper bound free energy difference between protein native-like structures. *Physics A* 533, 122053 (2019)
- Vila, J.A. Metamorphic proteins in light of Anfinsen's dogma. *J. Phys. Chem. Lett.* 11, 4998-4999, (2020)
- Yadid, I., Kirshenbaum, N., Sharon, M., Dym, O. and Tawfik, D.S. Metamorphic proteins mediate evolutionary transitions of structure. *Proc Natl Acad Sci USA* 107(16), 7287-7292 (2010)
- Zhang Z., Smith, D.L. Determination of amide hydrogen exchange by mass spectrometry: A new tool for protein structure elucidation. *Protein Science*, 2, 522-531 (1993)