

NECESSARY AND SUFFICIENT CONDITIONS FOR DOMINATION RESULTS FOR PROPER SCORING RULES

ALEXANDER R. PRUSS

ABSTRACT. Scoring rules measure the deviation between a probabilistic forecast and reality. Strictly proper scoring rules have the property that for any forecast, the mathematical expectation of the score of a forecast p by the lights of p is strictly better than the mathematical expectation of any other forecast q by the lights of p . Probabilistic forecasts need not satisfy the axioms of the probability calculus, but Predd, et al. (2009) have shown that given a finite sample space and any strictly proper *additive and continuous* scoring rule, the score for any forecast that does not satisfy the axioms of probability is strictly dominated by the score for some probabilistically consistent forecast. Recently, this result has been extended to non-additive continuous scoring rules. In this paper, a condition weaker than continuity is given that suffices for the result, and the condition is proved to be optimal.

1. THE MAIN RESULTS

Scoring rules measure the deviation between a probabilistic forecast and reality. Strictly proper scoring rules have the property that for any forecast, the mathematical expectation of the score of a forecast p by the lights of p is strictly better than the mathematical expectation of any other forecast q by the lights of p . Forecasts need not satisfy the axioms of probability, but under some conditions whose discussion is the main purpose of this paper, the score of a forecast that does not satisfy the axioms of probability is strictly dominated by the score of a forecast by that does satisfy these axioms. This result has been interpreted by epistemologists as supporting the idea that reasonable forecasts will always be probabilistically consistent (e.g., Joyce 1998; Greaves and Wallace 2006; Pettigrew 2016).

To be precise, let Ω be a finite sample space, encoding the possible situations that the forecasts concern. Let \mathcal{C} be the set of all functions from the power set of Ω to the reals: these we call credence functions. Let \mathcal{P} be the subset of \mathcal{C} which consists of the functions satisfying the axioms of probability. Members of \mathcal{C} can be thought of as forecasts regarding Ω . An *inaccuracy* (respectively, *accuracy*) *scoring rule* is a function s from a set $\mathcal{F} \supseteq \mathcal{P}$ of credence function to $[M, \infty]^\Omega$ (respectively, $[-\infty, M]^\Omega$) for some finite M , where A^B is the set of functions from B to A . Then $s(c)(\omega)$ for $c \in \mathcal{F}$ measures the accuracy of the forecast c when we are in fact at $\omega \in \Omega$, with lower (respectively, higher) values being more accurate. One can go

back and forth between inaccuracy and accuracy scoring rules, since if s is one, then $-s$ is the other. By a *scoring rule*, we shall mean an accuracy or inaccuracy scoring rule.

Given a probability $p \in \mathcal{P}$ and an extended real function f on Ω , let $E_p f$ be the expected value with respect to p defined in the following way to avoid multiplying infinity by zero:

$$E_p f = \sum_{\omega \in \Omega, p(\{\omega\}) \neq 0} p(\{\omega\}) f(\omega).$$

We then say that an inaccuracy (respectively, accuracy) scoring rule s is *proper* on $\mathcal{F} \supseteq \mathcal{P}$ provided that for every $p \in \mathcal{P}$ and every $c \in \mathcal{F}$, we have $E_p s(p) \leq E_p s(c)$ (respectively, $E_p s(p) \geq E_p s(c)$), that it is *strictly proper* provided the inequality is always strict, and that it is *quasi-strictly proper* provided that it is proper and the inequality is strict when $p \in \mathcal{P}$ and $c \in \mathcal{C} \setminus \mathcal{P}$.

Propriety captures the idea that if an agent adopts a probability function p as their forecast, then by the agent's lights there can be no improvement in the expected score from switching to a different forecast. Strict propriety captures the idea that an agent who has adopted a probability function p as their forecast will think other forecasts to be inferior.

A scoring rule is said to be *additive* provided that $\mathcal{F} = \mathcal{C}$ and there is a collection of extended-real functions $(s_A)_{A \subseteq \Omega}$ each defined on $\mathbb{R} \times \{0, 1\}$ such that for all $c \in \mathcal{F}$ and $\omega \in \Omega$:

$$s(c)(\omega) = \sum_{A \subseteq \Omega} s_A(c(A), 1_A(\omega)).$$

Here, $s_A(c(A), 1_A(\omega))$ can be thought of as measuring the accuracy of the forecast for the probability of A at ω .

The set of probabilities \mathcal{P} can be equipped with the topology resulting from its natural embedding ψ in the product space $[0, 1]^\Omega$, where $\psi(p)(\omega) = p(\{\omega\})$. Thus, a sequence of probabilities (p_n) converges to a probability p just in case $p_n(\{\omega\}) \rightarrow p(\{\omega\})$ for all $\omega \in \Omega$.

A scoring rule is *probability-continuous* provided that the restriction of s to the probabilities is a continuous function from \mathcal{P} to $[-\infty, \infty]^\Omega$ equipped with the product topology.

Say that an inaccuracy score $s(c_1)$ is *weakly dominated* by an inaccuracy score $s(c_2)$ provided that $s(c_2)(\omega) \leq s(c_1)(\omega)$ for all $\omega \in \Omega$, with the inequality reversed for accuracy scores. If the inequality is strict, we shall talk of *strict domination*.

Predd, et al. (2009) showed that if s is a probability-continuous additive inaccuracy strictly proper scoring rule, then for any non-probability c , there is a probability p such that $s(c)$ is strictly dominated by $s(p)$. Obviously, the same is true for accuracy scoring rules. Recently, Pettigrew (forthcoming) announced that this result holds without the assumption of additivity, merely assuming probability-continuity. Pettigrew's proof had flaws (2021),

but correct proofs were given by Nielsen (2021) and Pruss (2021). Nielsen's proof also extended the result to the quasi-proper case.

The Pettigrew-Nielsen-Pruss theorem still assumes the continuity of the scoring rule. The purpose of the present paper is to identify the weakest assumption on a strictly proper scoring rule as restricted to the probability functions that guarantees the strict domination property.

The following is our main result. It holds for both accuracy and inaccuracy scores, with domination defined in the respectively appropriate way.

Say that an $s \in [-\infty, \infty]^\Omega$ is finite provided that $|s(\omega)| < \infty$ for all ω .

Fix $Q \subseteq \mathbb{R}^\Omega$. Let $\langle \cdot, \cdot \rangle$ be the usual scalar product on \mathbb{R}^Ω : $\langle f, g \rangle = \sum_\omega f(\omega)g(\omega)$ for any $f, g \in \mathbb{R}^\Omega$. We say that a boundary point z of a set $G \subseteq \mathbb{R}^\Omega$ is Q -facing provided that G has a non-zero normal at z lying in Q , i.e., there is a $v \in Q \setminus \{0\}$ such that $\langle v, w \rangle \leq \langle v, z \rangle$ for all $w \in G$. Write $\partial_Q G$ for the set of all Q -facing boundary points of G . Write $\text{Conv } F$ for the convex hull of F . The positive and negative orthants of \mathbb{R}^Ω are the set of functions on Ω that are everywhere strictly positive or negative, respectively.

Theorem 1. *Consider a proper scoring rule s on \mathcal{P} . Then the following conditions are equivalent:*

- (a) *for every extension of s to a quasi-strictly proper scoring rule $s' : \mathcal{C} \rightarrow [0, \infty]^\Omega$, if $c \in \mathcal{C} \setminus \mathcal{P}$, then there is a $p \in \mathcal{P}$ such that $s'(p) = s(p)$ strictly dominates $s'(c)$*
- (b) *either $E_p s(p)$ is infinite for some $p \in \mathcal{P}$ or both:*
 - (i) *for any sequence (p_n) in \mathcal{P} that converges to some probability function p such that $s(p_n)$ is finite for all n while $s(p)$ is not finite, we have $\lim_n E_{p_n} s(p_n) = E_p s(p)$, and*
 - (ii) *if $F = s[\mathcal{P}] \cap \mathbb{R}^\Omega$ is the set of finite scores, then F is dense in $\partial_Q \text{Conv } F$, where Q is the positive orthant in the accuracy score case and the negative orthant in the inaccuracy score case.*

Combining the above with following yields a new proof of the Pettigrew-Nielsen-Pruss theorem.

Proposition 1. *Suppose that s is quasi-strictly proper and continuous on \mathcal{C} . Then condition (b) of Theorem 1 is fulfilled.*

It follows from Lemma 3, below, that for any proper scoring rule s , the function $p \mapsto E_p s(p)$ is continuous on the probabilities with finite score. Thus, in condition (i) we can drop the restriction that $s(p)$ is not finite.

Condition (ii) intuitively says that the set of finite scores doesn't have any open holes in the Q -direction.

We now give an example of an everywhere-finite proper scoring rule that satisfies (i) and (ii) but is not continuous.

Let \mathcal{R} be the set of *regular* probabilities, i.e., probability functions p such that $p\{\omega\} > 0$ for all ω .

Let b be the Brier score, which is a continuous, additive, bounded, strictly proper inaccuracy score with values in $[0, 1]^\Omega$. Define a new scoring rule s as

follows. If $c \notin \mathcal{P} \setminus \mathcal{R}$, let $s(c) = b(c)$. If $c \in \mathcal{P} \setminus \mathcal{R}$, then let $s(c)(\omega) = b(c)(\omega)$ if $c\{\omega\} > 0$ and $s(c)(\omega) = \infty$ if $c\{\omega\} = 0$. It is easy to see that s is strictly proper but it is clearly not continuous. Note that $E_p s(p) = E_p b(p)$ for all $p \in \mathcal{P}$, so (i) is satisfied for s just as by Proposition 1 it is for b . Moreover, the set of finite s -scores is a subset of the set of finite Brier scores, but the closure of the two sets is the same (since \mathcal{R} is dense in \mathcal{P} and s is continuous), and so (ii) holds for s -scores given its holding for Brier scores.

2. PROOFS

From now on we will assume we are dealing only with accuracy scores with values in $[-\infty, M]$ for some finite M .

Given a probability function p , let \hat{p} be the function from Ω to $[0, 1]$ defined by $\hat{p}(\omega) = p(\{\omega\})$. Define $\hat{\mathcal{P}} = \{\hat{p} : p \in \mathcal{P}\}$ and $\hat{\mathcal{R}} = \{\hat{p} : p \in \mathcal{R}\}$. Thus, $\hat{\mathcal{P}}$ is the set of non-negative functions on Ω the sum of whose values is 1, and $\hat{\mathcal{R}}$ is the subset of the strictly positive ones.

Given $p \in \hat{\mathcal{P}}$, let \check{p} be the probability function such that $\check{p}(\{\omega\}) = p(\omega)$ for all $\omega \in \Omega$.

Given a scoring rule s defined on \mathcal{P} , by abuse of notation let $s(p) = s(\check{p})$ for $p \in \hat{\mathcal{P}}$.

Given two functions f and g from Ω to the extended reals, say that g strictly (weakly) dominates f provided that $f < g$ ($f \leq g$) everywhere. The following is an easy fact.

Let \cdot be a multiplication operation on the extended reals with the stipulation that $a \cdot 0 = 0 \cdot a$ for any a , finite or not. With this stipulation, multiplication is upper semicontinuous on $[-\infty, \infty) \times [0, 1]$, and continuous at all (x, y) where x is finite or y is positive. Now define the *extended scalar product* on $X = [0, 1]^\Omega \times [-\infty, \infty)^\Omega$ by

$$\langle f, g \rangle = \sum_{\omega \in \Omega} f(\omega) \cdot g(\omega),$$

using the above stipulation.

Lemma 1. *The extended scalar product is an upper semicontinuous function from X to $[-\infty, \infty)$. Moreover, it is continuous at any (f, g) such that either f is strictly positive everywhere or g is finite everywhere. Finally, for a fixed $f \in [0, 1]^\Omega$, the function $g \mapsto \langle f, g \rangle$ is continuous on $[-\infty, \infty)^\Omega$.*

Observe that $E_p f = \langle \hat{p}, f \rangle$. This fact will allow us to go back and forth between probabilistic concepts and geometric concepts.

For a subset F of \mathbb{R}^Ω and a vector v , let

$$\sigma_F(v) = \sup_{z \in F} \langle z, v \rangle$$

be the support function of F at v .

Lemma 2. *Assume s is a proper scoring rule on \mathcal{P} with $E_p s(p)$ finite for all p . Let F be the set of finite scores. Suppose that for every convergent sequence (p_n) of members of $\hat{\mathcal{P}}$ with $s(p_n) \in F$ for all n , we have $\lim_n \langle p_n, s(p_n) \rangle = \langle p, s(\lim_n p_n) \rangle$. Then $\sigma_F(p) = \langle p, s(p) \rangle$ for all $p \in \hat{\mathcal{P}}$.*

Proof of Lemma 2. First, suppose $s(p) \in F$. Then for every $z \in F$ we have $\langle p, z \rangle \leq \langle p, s(p) \rangle$ by propriety. Since $s(p) \in F$, it follows that $\sigma_F(p) = \langle p, s(p) \rangle$.

Now suppose $s(p)$ is not finite. Let (p_n) be a sequence in $\hat{\mathcal{R}}$ converging to p . For $q \in \hat{\mathcal{R}}$, the fact that $E_q s(q)$ is finite implies that $s(q)$ is finite, so $s(p_n) \in F$ for all n . Using compactness and passing to a subsequence if necessary, assume that $s(p_n)$ converges to some value $z \in [-\infty, M]^\Omega$. Then

$$\sigma_F(p) \geq \lim_n \langle p, s(p_n) \rangle = \langle p, z \rangle \geq \limsup_n \langle p_n, s(p_n) \rangle = \langle p, s(p) \rangle,$$

where the relations follow respectively by definition of σ_F , the continuity of the extended scalar product for a fixed first argument (Lemma 1), the upper semicontinuity of the extended scalar product on X (same Lemma), and the assumptions of our lemma.

On the other hand:

$$\langle p, s(p) \rangle \geq \langle p, s(q) \rangle$$

by propriety for all $q \in \hat{\mathcal{P}}$. Hence $\langle p, s(p) \rangle \geq \sigma_F(p)$. Thus, $\sigma_F(p) = \langle p, s(p) \rangle$. \square

Lemma 3. *Let s be a proper scoring rule on \mathcal{P} . Let $H = \{p \in \hat{\mathcal{P}} : s(p) \text{ is finite}\}$. For any $p \in \bar{H}$, the limit of $\langle q, s(q) \rangle$ as q tends to p within H exists. If $s(p)$ is finite, the limit equals $\langle p, s(p) \rangle$. Finally, if p_n is a sequence in H converging to $p \in \bar{H}$ such that $\lim_n s(p_n) = r$, then $\langle p, r \rangle = \lim_n \langle p_n, s(p_n) \rangle$.*

Proof. Fix $p \in \bar{H}$. To show that a sequence converges to some member of a compact set, it suffices to show that every convergent subsequence of it converges to the same point. The set $[-\infty, M]$ is compact and $\langle q, s(q) \rangle \in [-\infty, M]$ for all q . Thus to show the existence of our limit, all we need to show is that if (p_n) and (p'_n) are two sequences in H converging to p with $\langle p_n, s(p_n) \rangle \rightarrow L$ and $\langle p'_n, s(p'_n) \rangle \rightarrow L'$, then $L = L'$. Moreover, if we can show this, then letting $p'_n = p$ for all n , it will follow that $L = s(p)$.

Thus, suppose (p_n) and (p'_n) are two sequences in H converging to p with $\langle p_n, s(p_n) \rangle$ and $\langle p'_n, s(p'_n) \rangle$ convergent. Passing to subsequences if necessary, assume that $s(p_n)$ and $s(p'_n)$ converge respectively to r and r' in $[-\infty, M]^\Omega$.

I now claim that $\langle p, r \rangle = \lim_n \langle p_n, s(p_n) \rangle$. First, note that

$$\limsup_n \langle p_n, s(p_n) \rangle \leq \langle p, r \rangle$$

by Lemma 1. Next observe that

$$\langle p_n, s(p_n) \rangle \geq \langle p_n, s(p_m) \rangle$$

by propriety. Since $s(p_m)$ is finite as $p_m \in H$, by Lemma 1 the right-hand-side converges to $\langle p, s(p_m) \rangle$ as $n \rightarrow \infty$. Thus:

$$\liminf_n \langle p_n, s(p_n) \rangle \geq \langle p, s(p_m) \rangle.$$

Taking the limit as $m \rightarrow \infty$ and using Lemma 1 we get

$$\liminf_n \langle p_n, s(p_n) \rangle \geq \langle p, r \rangle.$$

Thus we have $\langle p, r \rangle = \lim_n \langle p_n, s(p_n) \rangle$, which is the final remark in the statement of our Lemma.

Thus:

$$\begin{aligned} \langle p, r' \rangle &= \lim_m \langle p, s(p'_m) \rangle \\ &= \lim_m \lim_n \langle p_n, s(p'_m) \rangle \\ &\leq \lim_m \lim_n \langle p_n, s(p_n) \rangle \\ &= \lim_m \langle p, r \rangle = \langle p, r \rangle, \end{aligned}$$

where the first equality was due to Lemma 1, the second due to Lemma 1 and the finiteness of $s(p'_m)$, and the inequality was due to propriety. In exactly the same way, $\langle p, r \rangle \leq \langle p, r' \rangle$. Thus, $\langle p, r \rangle = \langle p, r' \rangle$. But $\langle p, r' \rangle = \lim_n \langle p'_n, s(p'_n) \rangle$, just as we proved in the unprimed case. Thus, $\lim_n \langle p_n, s(p_n) \rangle = \lim_n \langle p'_n, s(p'_n) \rangle$. \square

Lemma 4. *Let F be a nonempty subset of $[-\infty, \infty)^\Omega$. Suppose $z_0 \in [-\infty, \infty)^\Omega$ is such that $\langle p, z_0 \rangle < \sigma_F(p)$ for all $p \in \hat{\mathcal{P}}$. Then z_0 is strictly dominated by some member of $\text{Conv}(F)$.*

Proof of Lemma 4. Let z_0 be as in the statement of the Lemma. Let $Q = \{z \in \mathbb{R}^\Omega : \forall \omega (z(\omega) > z_0(\omega))\}$ be the set of points of \mathbb{R}^Ω strictly dominating z_0 . We need to show that $Q \cap \text{Conv}(F) \neq \emptyset$.

Suppose that Q does not intersect $\text{Conv}(F)$. Both Q and $\text{Conv}(D)$ are convex sets. Thus by hyperplane separation, there is a non-zero $p \in \mathbb{R}^\Omega$ and an $\alpha \in \mathbb{R}$ such that $\langle p, z \rangle \geq \alpha$ for all $z \in Q$ and $\langle p, z \rangle \leq \alpha$ for all $z \in \text{Conv}(F)$.

I claim that $p(\omega) \geq 0$ for all ω . To see this, suppose $p(\omega_0) < 0$ for some $\omega_0 \in \Omega$. Let z be any member of Q . Let $\beta = \langle p, z \rangle$. Let $z'(\omega) = z(\omega)$ for $\omega \neq \omega_0$ and $z'(\omega_0) = z(\omega_0) + (\alpha - \beta - 1)/p(\omega_0)$. Then $z' \in Q$ since $z \in Q$ while $\beta \geq \alpha$ and $p(\omega_0) < 0$. Observe that

$$\langle p, z' \rangle = \langle p, z \rangle + (\alpha - \beta - 1) = \alpha - 1,$$

which is impossible as $\langle p, z' \rangle \geq \alpha$ since $z' \in Q$.

Rescaling if necessary, we may assume that $\sum_\omega p(\omega) = 1$ and hence $p \in \hat{\mathcal{P}}$.

Since z_0 is on the boundary of $Q \subseteq [-\infty, \infty)^\Omega$, by the upper semicontinuity of the extended scalar product (Lemma 1) we have $\langle p, z_0 \rangle \geq \alpha$. Moreover, since $\langle p, z \rangle \leq \alpha$ for all $z \in \text{Conv}(F)$, we must have $\sigma_F(p) \leq \alpha$. Thus, $\langle p, z_0 \rangle \geq \sigma_F(p)$, which contradicts the assumptions of the Lemma. \square

Say that a vector v is normal to a convex set G at a point $z_1 \in G$ provided that $\langle v, z \rangle \leq \langle v, z_1 \rangle$ for all $z \in G$. The following Lemma is due to a *MathOverflow* user (fedja 2021).

Lemma 5. *Let G be a closed convex subset of \mathbb{R}^n whose intersection with the positive orthant $\mathbb{R}_+^n = (0, \infty)^n$ is non-empty and bounded. Then there is a vector $v \in \mathbb{R}_+^n$ that is normal to G at some point $z \in G \cap \mathbb{R}_+^n$.*

Proof. Let $f(x_1, \dots, x_n) = x_1 \cdots x_n$ be the product of coordinates function. Let $U = G \cap \mathbb{R}_+^n$. Then f is a continuous function on the closure \bar{U} of U in \mathbb{R}^n . Moreover, \bar{U} is compact by the boundedness requirement, so f attains a maximum on \bar{U} . This maximum is not attained on the boundary of \mathbb{R}_+^n , i.e., on any point with a zero coordinate, since f is zero at any such point, while there is at least one point in \bar{U} where f is strictly positive (since f is non-zero everywhere on $U \neq \emptyset$). Hence, the maximum is attained at a point z of U . That point cannot be an interior point (since then $z + (\varepsilon, \dots, \varepsilon)$ would be in U and f would be bigger then), so it must be a boundary point of U that isn't a boundary point of \mathbb{R}_+^n . Thus, z must be a boundary point of G . By the fundamental normal cone condition for optimality, the gradient of f must be normal to U at z . Since \mathbb{R}_+^n is a neighborhood of z , and the intersection of U with that neighborhood is the same as that of G with it, it follows that the gradient of f must be normal to G at z . But the gradient of f on \mathbb{R}_+^n always lies in \mathbb{R}_+^n , which completes the proof. \square

Let $\|z\|_\infty = \max_{\omega \in \Omega} |z(\omega)|$ for $z \in [-\infty, \infty]^\Omega$ and let

$$B(z, r) = \{z' : \|z' - z\|_\infty < r\}$$

be the open ball of radius r around a finite z' in this norm.

Lemma 6. *Fix $p \in \hat{\mathcal{R}}$ and $\alpha \in \mathbb{R}$, and let $K = \{y \in [-\infty, \infty]^\Omega : \langle p, y \rangle \leq \alpha\}$. Fix $x \in \mathbb{R}^\Omega$ such that $\langle p, x \rangle = \alpha$ and $\varepsilon > 0$. Then there is a $\delta \in (0, \varepsilon)$ such that no point of $K \cap B(x, \delta)$ is weakly dominated by any point of $K \setminus B(x, \varepsilon)$.*

Proof of Lemma 6. Translating if necessary, without loss of generality assume $x = 0$ and $\alpha = 0$. Fix $y \in K \cap B(0, \delta)$. Suppose that y is weakly dominated by $z \in K$.

Then $z(\omega) \leq y(\omega)$ for all ω and $\sum_\omega z(\omega)p(\omega) \leq 0$ as well as $\sum_\omega y(\omega)p(\omega) \leq 0$. Let $c = \max(1, (|\Omega| - 1)/\min_\omega p(\omega))$. Then for any ω :

$$-\delta \leq z(\omega) \leq -\frac{\sum_{\omega' \neq \omega} z(\omega')p(\omega')}{p(\omega)} \leq -\frac{\sum_{\omega' \neq \omega} y(\omega')p(\omega')}{p(\omega)} \leq c\delta.$$

Hence, $\|z\|_\infty \leq c\delta$. Thus, we have shown that if $y \in K \cap B(0, \delta)$ is dominated by $z \in K \setminus B(0, \varepsilon)$, then $\varepsilon \geq c\delta$. Hence, if $\varepsilon > 0$ is fixed, any choice of $\delta \in (0, c^{-1}\varepsilon)$ will complete the proof. \square

Proof of Theorem 1. If $E_p s(p)$ is infinite (i.e., equal to $-\infty$ since we are working with accuracy scores) for some probability function p , then s has

no extension to a quasi-strictly proper scoring rule on \mathcal{C} , as no point $z \in [-\infty, \infty]^\Omega$ will be such that $E_p z < E_p s(p)$. Thus, we may assume for all our proofs that $E_p s(p)$ is finite for all probabilities p , and hence that so is $\langle p, s(p) \rangle$.

Without loss of generality, assume we have accuracy scores with ranges in $[-\infty, -1]$.

Recall that $E_p s(q) = \langle \hat{p}, s(\hat{q}) \rangle$. If (i) fails, then there is a sequence $p_n \rightarrow p$ in $\hat{\mathcal{P}}$ such that $\langle p_n, s(p_n) \rangle$ does not converge to $\langle p, s(p) \rangle$. By Lemma 3, $\lim_n \langle p_n, s(p_n) \rangle = L$ exists and cannot equal $\langle p, s(p) \rangle$. Passing to a subsequence if necessary, we may assume that $s(p_n)$ has a limit $r \in [-\infty, 0]^\Omega$, and then

$$(1) \quad \lim_n \langle p_n, s(p_n) \rangle = \langle p, r \rangle$$

by the same lemma.

Note that

$$\langle q, r \rangle = \lim_n \langle q, s(p_n) \rangle \leq \langle q, s(q) \rangle$$

for all $q \in \hat{\mathcal{P}}$ by Lemma 1 and propriety. In particular, $L = \langle p, r \rangle \leq \langle p, s(p) \rangle$. Thus, $L < \langle p, s(p) \rangle$ since $\langle p_n, s(p_n) \rangle$ does not converge to $\langle p, s(p) \rangle$. Further, $\langle p, s(p) \rangle$ is negative since our accuracy scores are negative. Choose $\alpha \in (1, L/\langle p, s(p) \rangle)$ so that

$$(2) \quad \alpha \langle p, s(p) \rangle > L = \langle p, r \rangle$$

Let $x = \alpha s(p)$.

Infinite scores cannot strictly dominate any score. Now I claim that x is not weakly dominated by any finite score. For suppose that $s(q)$ is finite. Then by Lemma 1, propriety, (1) and (2) we have:

$$\langle p, s(q) \rangle = \lim_n \langle p_n, s(q) \rangle \leq \lim_n \langle p_n, s(p_n) \rangle = \langle p, r \rangle < \alpha \langle p, s(p) \rangle = \langle p, x \rangle.$$

And this implies that x is not dominated by $s(q)$.

On the other hand,

$$(3) \quad \langle q, x \rangle = \alpha \langle q, s(p) \rangle \leq \alpha \langle q, s(q) \rangle < \langle q, s(q) \rangle$$

for any $q \in \hat{\mathcal{P}}$ since $\alpha > 0$.

Now define $s'(c) = s(c)$ if $c \in \mathcal{P}$ and $s'(c) = x$ if $c \in \mathcal{C} \setminus \mathcal{P}$. Then s' is quasi-strictly proper by (3), but $s'(c)$ is not dominated by any score of a probability. Hence condition (a) fails.

Now suppose (ii) fails. Thus there is a $z_0 \in \partial_Q \text{Conv } F$ and $\varepsilon > 0$ such that $F \cap B(z_0, \varepsilon) = \emptyset$. Then $\text{Conv } F$ has a normal in $Q \setminus \{0\}$ at z_0 . Rescaling if necessary, we can assume that normal is some $p \in \hat{\mathcal{R}}$. Let $K = \{z \in [-\infty, \infty]^\Omega : \langle p, z \rangle \leq \langle p, z_0 \rangle\}$. By Lemma 6, there is a $\delta \in (0, \varepsilon)$ such that no point in $B(z_0, \delta)$ is strictly dominated by any point in $K \setminus B(z_0, \varepsilon)$. Choose a point z_1 in $B(z_0, \delta)$ that is strictly dominated by z_0 . Note that z_0 is a limit of convex combinations of points of F . If q is any member of $\hat{\mathcal{P}}$ and u is any point of F , then $\langle q, u \rangle \leq \langle q, s(q) \rangle$ by propriety. Thus, the same is true if u

is a convex combination of points of F , and by Lemma 1 also for limits of convex combinations of points of F .

Hence, $\langle q, z_0 \rangle \leq \langle q, s(q) \rangle$ for all $q \in \hat{\mathcal{P}}$. Since z_1 is strictly dominated by z_0 , it follows that $\langle q, z_1 \rangle < \langle q, s(q) \rangle$ for all $q \in \hat{\mathcal{P}}$.

On the other hand, since z_1 is not strictly dominated by anything in $K \setminus (z_0, \varepsilon)$, it's not strictly dominated by anything in $\supseteq F$. Much as before, let $s'(c) = s(c)$ if c is a probability function and $s'(c) = z_1$ if c is not a probability function. As before, we have strict quasi-proprity and yet no credence that isn't a probability is strictly s' -dominated by any probability function.

Now suppose that (i) and (ii) hold. Let s' be an extension of s to a quasi-strictly proper scoring rule defined for all credences. Fix a non-probability credence c and let $z_0 = s'(c)$. By (i) and Lemma 2, $\langle p, s(p) \rangle = \sigma_F(p)$ for all p . By Lemma 4 and quasi-strict propriety, z_0 is strictly dominated by some member of $\text{Conv}(F)$.

Let G be the closure of $\text{Conv}(F)$ in \mathbb{R}^Ω . Then there is a point $z_1 \in [-\infty, \infty)^\Omega$ such that z_0 is strictly dominated by z_1 and z_1 is strictly dominated by some member of G (e.g., if z_2 is a point of G that dominates z_0 , then let $z_1(\omega) = z_2(\omega) - 1$ if $z_0(\omega)$ is infinite, and $z_1(\omega) = (z_0(\omega) + z_2(\omega))/2$ otherwise). Let Q be the set of points of $[-\infty, \infty)^\Omega$ that strictly dominate z_1 . Then $Q \cap G$ is non-empty. Note that every point z of F satisfies

$$(4) \quad \langle u, z \rangle \leq \langle u, s(u) \rangle$$

by propriety, where $u(\omega) = 1/|\Omega|$ for all ω , and hence every point z of G satisfies (4) as well (a convex combination of points z satisfying (4) will satisfy it as well, and by Lemma 1, so will every point in the closure of $\text{Conv}(F)$). Moreover, $\langle u, s(u) \rangle$ is finite. The set of points $z \in Q$ such that $\langle u, z \rangle \leq \langle u, s(u) \rangle$ is bounded, and hence $Q \cap G$ is bounded.

By Lemma 5 (letting $n = |\Omega|$), there is a $z_3 \in Q \cap G$ such that $z \in G$ has a normal in the positive orthant. Thus, $z_3 \in \partial_Q G = \partial_Q \text{Conv} F$. By condition (ii), there are points of F arbitrarily close to z_2 . Since z_2 strictly dominates z_0 , so do points that are sufficiently close to z_2 , and hence some point of F strictly dominates z_0 . \square

Proof of Proposition 1. We must have $E_p s(p)$ finite for all probability functions p or else quasi-strict propriety is impossible. It follows that the score of any regular probability is finite.

By Lemma 1, $p \mapsto \langle p, s(p) \rangle$ is upper semicontinuous on $\hat{\mathcal{P}}$ if the scoring rule is continuous.

Moreover, $\langle p, s(p) \rangle = \sup_{q \in \mathcal{P}} \langle p, s(q) \rangle$. Since $\hat{\mathcal{R}}$ is dense in $\hat{\mathcal{P}}$ and $q \mapsto \langle p, s(q) \rangle$ is continuous for a fixed p , it follows that $\sup_{q \in \mathcal{P}} \langle p, s(q) \rangle = \sup_{q \in \mathcal{R}} \langle p, s(q) \rangle$. Moreover, since every score of a regular probability is finite, $p \mapsto \langle p, s(q) \rangle$ is continuous by Lemma 1. Thus, $p \mapsto \sup_{q \in \mathcal{R}} \langle p, s(q) \rangle$ is the supremum of continuous functions and hence it is lower semicontinuous at these points. (This observation is due to Nielsen [2021].)

Hence, $p \mapsto \langle p, s(p) \rangle$ is continuous at every point of $\hat{\mathcal{P}}$, and so we have (i).

It remains to prove (ii). Assume first that s is strictly proper. Let F be the set of finite scores of probabilities. Then F considered as a subset of \mathbb{R}^Ω is closed, since it is the intersection with \mathbb{R}^Ω of the set of scores of probability functions, and the set of probability functions is compact while s is continuous. Moreover, for any $p \in \hat{\mathcal{P}}$, we have $\langle p, s(p) \rangle > \langle p, z \rangle$ for all $z \in F \setminus \{s(p)\}$. It follows that for any p , the only point z in the closed convex hull of F such that $\langle p, s(p) \rangle = \langle p, z \rangle$ is $s(p)$ itself, from which (ii) follows.

Now, suppose s is merely quasi-strictly proper. Let $b : \mathcal{C} \rightarrow [-1, 0]$ be any strictly proper continuous accuracy score, for instance the negative of the Brier score. Let $s_\varepsilon = s + \varepsilon b$ for any $\varepsilon > 0$. Then s_ε is a strictly proper continuous score, and (ii) must hold for it. Let $F_\varepsilon = s_\varepsilon[\mathcal{P}] \cap \mathbb{R}^\Omega$.

Now, consider a point $z_0 \in \partial_Q F$. Fix $\varepsilon > 0$. We will show that there is a point of F within distance ε of z_0 . Let $\delta = \varepsilon/4$. Thus there is a $p \in \hat{\mathcal{R}}$ (since any vector in Q can be rescaled to get a vector in $\hat{\mathcal{R}}$) such that $\langle p, z \rangle \leq \langle p, z_0 \rangle$ for all $z \in F$. The point z_0 is a convex combination $z_0 = c_1 w_1 + \dots + c_n w_n$ of points of F , where $c_i > 0$ and $\sum_i c_i = 1$. We then have to have $\langle p, w_i \rangle = \langle p, z_0 \rangle$ for each i . Choose $p_i \in \hat{\mathcal{P}}$ such that $w_i = s(p_i)$. Since s_δ is continuous and $\hat{\mathcal{R}}$ is dense in $\hat{\mathcal{P}}$, for each i choose p'_i of $\hat{\mathcal{R}}$ such that $\|s_\delta(p'_i) - s_\delta(p_i)\|_\infty \leq \delta$. Then $\|w_i - s_\delta(p_i)\|_\infty \leq 2\delta$ since s_δ is never more than δ away from s .

Let $z'_0 = \sum_i c_i s_\delta(p'_i)$. Then, $\|z'_0 - z_0\|_\infty \leq 2\delta$. Note that $s_\delta(p'_i) \in \partial_Q F_\varepsilon$, since $p'_i \in Q$ and s_δ is proper. Applying the strictly proper case to s_δ , we conclude that there is a $p \in \hat{\mathcal{P}}$ such that $\|s_\delta(p) - z'_0\|_\infty \leq \delta$ and $s_\delta(p) \in F_\varepsilon$. Then, $\|s(p) - z'_0\|_\infty \leq 2\delta$, and so $\|s(p) - z_0\|_\infty \leq 4\delta = \varepsilon$, which completes the proof.¹ \square

REFERENCES

- [1] “fedja”. 2021. “Answer To: If a Convex Compact Set Meets the Positive Orthant Does It Meet It At a Point With a Normal in the Positive Orthant”. *Mathoverflow*. <https://mathoverflow.net/questions/384140/if-a-compact-convex-set-meets-the-positive-orthant-does-it-meet-it-at-a-point-with-a-normal-in-the-positive-orthant>
- [2] H. Greaves, and D. Wallace. 2006. “Justifying Conditionalization: Conditionalization Maximizes Expected Epistemic Utility”, *Mind* 115:607–632.
- [3] J. M. Joyce. 1998. “A Nonpragmatic Vindication of Probabilism”, *Philosophy of Science* 65:575–603.
- [4] Michael Nielsen. 2021. “On the Best Accuracy Arguments for Probabilism”. Manuscript.
- [5] R. Pettigrew. 2016. *Accuracy and the Laws of Credence*. Oxford: Oxford University Press.
- [6] R. Pettigrew. Forthcoming. “Accuracy-First Epistemology Without the Additivity Axiom”, *Philosophy of Science*.
- [7] Joel B. Predd, Robert Seiringer, Elliott H. Lieb, Daniel N. Osherson, H. Vincent Poor, and Sanjeev R. Kulkarni. 2009. “Probabilistic Coherence and Proper Scoring Rules”, *IEEE Transactions on Information Theory* 55:4786–4792.

¹I am grateful to Michael Nielsen for many discussions and much encouragement.

- [8] Alexander R. Pruss. 2021. “Proper Scoring Rules and Domination”.
<https://arxiv.org/abs/2102.02260>