

Inversion formula for infinitely divisible distributions

E.V. Burnaev

The aim of this note is to prove the inversion formula, which can be used to compute the Levi measure of an infinitely divisible distribution from its characteristic function. Obtained formula is similar to the well-known inversion formula [2], which is used to compute the distribution function of a random variable from the corresponding characteristic function.

THEOREM. *Let $\phi(z) = e^{\psi(z)}$ be the characteristic function of an infinitely divisible distribution with the cumulant function $\psi(z) = i\gamma z - 1/2 \cdot \sigma^2 z^2 + \int_{\mathbb{R}} (e^{izx} - 1 - izxh(x)) U(dx)$, where $h(x)$ is a truncation function and $U(dx)$ is the Levy measure. Let $\hat{\rho}(z) = \int_{-1}^1 \log \frac{\phi(z)}{\phi(z+\lambda)} d\lambda - 1/3 \cdot \sigma^2$ and $B = [a, b]$, where $-\infty < a < b < \infty$ and $U(\{a\} \cup \{b\}) = 0$. We then have*

$$\int_B \left(\frac{x - \sin x}{x} \right) U(dx) = \frac{1}{4\pi} \cdot \text{v.p.} \int_{\mathbb{R}} \hat{\rho}(z) \left(\int_B e^{-izx} dx \right) dz. \quad (1)$$

Moreover, if $\int_{|x|>1} x^2 U(dx) < \infty$, then

$$\int_B x^2 U(dx) = -\frac{1}{2\pi} \cdot \text{v.p.} \int_{\mathbb{R}} (\psi''(z) + \sigma^2) \left(\int_B e^{-izx} dx \right) dz. \quad (2)$$

If $\int_{\mathbb{R}} |\hat{\rho}(z)| dz < \infty$, then the Levy measure is absolutely continuous with respect to the Lebesgue measure, has a bounded continuous density $u(x)$, and

$$u(x) = \frac{x}{x - \sin x} \cdot \frac{1}{4\pi} \int_{\mathbb{R}} e^{-izx} \hat{\rho}(z) dz. \quad (3)$$

If $\int_{|x|>1} x^2 U(dx) < \infty$ and $\int_{\mathbb{R}} |\psi''(z) + \sigma^2| dz < \infty$, then the Levy measure is absolutely continuous with respect to the Lebesgue measure, has a bounded continuous density $u(x)$, and

$$u(x) = -\frac{1}{2\pi x^2} \int_{\mathbb{R}} e^{-izx} (\psi''(z) + \sigma^2) dz \quad (4)$$

PROOF. First let us prove formulas (1) and (3). Let $f(z) = \log \phi(z) + 1/2 \cdot \sigma^2 z^2$. Therefore, we have $f(z) = i\gamma z + \int_{\mathbb{R}} (e^{izx} - 1 - izxh(x)) U(dx)$ and $f(z) - f(z + \lambda) = \log \frac{\phi(z)}{\phi(z+\lambda)} + 1/2 \cdot \sigma^2 (-\lambda^2 - 2\lambda z) = -i\gamma \lambda + \int_{\mathbb{R}} (e^{izx} - e^{i(z+\lambda)x} + i\lambda x h(x)) U(dx)$. Let us integrate the both sides of the last equality over $[-1, 1]$ with respect to λ . On the one hand, $\hat{\rho}(z) = \int_{-1}^1 (f(z) - f(z + \lambda)) d\lambda = \int_{-1}^1 \log \frac{\phi(z)}{\phi(z+\lambda)} d\lambda - 1/3 \cdot \sigma^2$. On the other hand, $\hat{\rho}(z) = \int_{-1}^1 \left\{ \int_{\mathbb{R}} (e^{izx} (1 - e^{i\lambda x}) + i\lambda x h(x)) U(dx) \right\} d\lambda$. Since $h(x) \sim 1 + o(x)$ for $x \rightarrow 0$ and $h(x) \sim O(1/x)$ for $x \rightarrow \infty$, it follows that $e^{izx} (1 - e^{i\lambda x}) + i\lambda x h(x) \sim o(x^2)$ for $x \rightarrow 0$ and $e^{izx} (1 - e^{i\lambda x}) + i\lambda x h(x) \sim O(1)$ for $x \rightarrow \infty$. Hence we can use Fubini's theorem and get $\hat{\rho}(z) = \int_{\mathbb{R}} \left(e^{izx} \int_{-1}^1 (1 - e^{i\lambda x}) d\lambda + ixh(x) \int_{-1}^1 \lambda d\lambda \right) U(dx) = \int_{\mathbb{R}} e^{izx} 2(1 - \sin x/x) U(dx)$. Therefore $\hat{\rho}(z) = \int_{\mathbb{R}} e^{izx} \rho(dx)$, where $\rho(dx) = 2(1 - \sin x/x) U(dx)$. This means that the function $\hat{\rho}(z)$ is the Fourier transform of the measure $\rho(dx)$. Using the inversion formula for the function $\hat{\rho}(z)$ [2], we obtain formula (1). If $\int_{\mathbb{R}} |\hat{\rho}(z)| dz < \infty$, then the Levy measure is absolutely continuous with respect to the Lebesgue measure, has a bounded continuous density $u(x)$, and $2(1 - \sin x/x) u(x) = 1/2\pi \cdot \int_{\mathbb{R}} e^{-izx} \hat{\rho}(z) dz$. Thus we have $u(x) = \frac{x}{4\pi(x - \sin x)} \int_{\mathbb{R}} e^{-izx} \hat{\rho}(z) dz$.

Now let us prove formulas (2) and (4). By assumption $\int_{|x|>1} x^2 U(dx) < \infty$, so that, using the monotone convergence theorem, we get $\psi''(z) = -\sigma^2 + \int_{\mathbb{R}} (-x^2 e^{izx}) U(dx)$. It now follows that $\int_{\mathbb{R}} e^{izx} x^2 U(dx) = -(\psi''(z) + \sigma^2)$. Arguing as above, we obtain formulas (2) and (4).

The following examples illustrate the application of the obtained results for the case of typical infinitely divisible distributions.

(a) The characteristic function of the *normal distribution* has the form $\phi(z) = e^{imz - 1/2 \cdot \sigma^2 z^2}$. It follows that $\hat{\rho}(z) = \int_{-1}^1 (-\sigma^2/2 (z^2 - (z + \lambda)^2) - im\lambda) d\lambda - 1/3 \cdot \sigma^2 = 0$. Using (3), we obtain $u(x) = 0$.

(b) The characteristic function of the *Cauchy distribution* has the form $\phi(z) = e^{-c|z| + i\gamma z}$, $c > 0$. It follows that $\hat{\rho}(z) = \int_{-1}^1 (-c|z| + c|z + \lambda| - \gamma\lambda) d\lambda = -2c|z| + c \int_{-1}^1 |z + \lambda| d\lambda = \begin{cases} c(|z| - 1)^2, & \text{for } z \in [-1, 1]; \\ 0, & \text{for } z \notin [-1, 1]. \end{cases}$ Using (3), we get $u(x) = \frac{cx}{4\pi(x - \sin x)} \int_{-1}^1 e^{-izx} (|z| - 1)^2 dz = \frac{cx}{4\pi(x - \sin x)} \frac{4(x - \sin x)}{x^3} = \frac{c}{\pi x^2}$.

(c) The characteristic function of the *compound Poisson* random variable has the form $\phi(z) = \exp\left\{c(\hat{\theta}(z) - 1)\right\}$, where c is an intensity of jumps, $\hat{\theta}(z)$ is the characteristic function of the distribution $\theta(dx)$ of the jump size. Since the distribution $\theta(dx)$ does not need to have a density, formula (1) is used. We have $\hat{\rho}(z) = \int_{-1}^1 c(\hat{\theta}(z) - \hat{\theta}(z + \lambda)) d\lambda = \int_{-1}^1 c(\int_{\mathbb{R}} e^{izx} \theta(dx) - \int_{\mathbb{R}} e^{i(z+\lambda)x} \theta(dx)) d\lambda = 2c \int_{\mathbb{R}} e^{izx} \theta(dx) - c \int_{\mathbb{R}} \left(\int_{-1}^1 e^{i\lambda x} d\lambda\right) e^{izx} \theta(dx) = 2c \int_{\mathbb{R}} e^{izx} \left(\frac{x - \sin x}{x}\right) \theta(dx)$. Using (1), we get $\int_B \left(\frac{x - \sin x}{x}\right) U(dx) = \frac{2c}{4\pi} \text{v.p.} \int_{\mathbb{R}} \left(\int_{\mathbb{R}} e^{izx} \left(\frac{x - \sin x}{x}\right) \theta(dx) \int_B e^{-izy} dy\right) dz = c \int_B \left(\int_{\mathbb{R}} \left(\frac{x - \sin x}{x}\right) \theta(dx) \frac{1}{2\pi} \text{v.p.} \int_{\mathbb{R}} e^{iz(x-y)} dz\right) dy = c \int_B \left(\int_{\mathbb{R}} \left(\frac{x - \sin x}{x}\right) \theta(dx) \delta(x - y)\right) dy = c \int_B \left(\frac{x - \sin x}{x}\right) \theta(dx)$. Hence we obtain $\int_B \left(\frac{x - \sin x}{x}\right) U(dx) = \int_B \left(\frac{x - \sin x}{x}\right) (c \cdot \theta(dx))$. Since $0 < \frac{x - \sin x}{x} < \infty$ for all $x \neq 0$, it now follows that $U(dx) = c \cdot \theta(dx)$.

(d) The characteristic function of the *negative binomial distribution* has the form $\phi(z) = \frac{p^c}{(1 - qe^{iz})^c}$, where $c > 0$, $p \in (0, 1)$, $q = 1 - p$. We have $\hat{\rho}(z) = c \int_{-1}^1 \log(1 - qe^{i(z+\lambda)}) d\lambda - 2c \log(1 - qe^{iz})$. It is well-known that $\log(1 - x) = -\sum_{k=1}^{\infty} \frac{x^k}{k}$. Let $x = qe^{i(z+\lambda)}$, then $\log(1 - qe^{i(z+\lambda)}) = -\sum_{k=1}^{\infty} \frac{q^k e^{ik(z+\lambda)}}{k}$. Integrating with respect to λ we obtain $\int_{-1}^1 \log(1 - qe^{i(z+\lambda)}) d\lambda = -\sum_{k=1}^{\infty} \frac{q^k e^{ikz}}{k} \cdot \int_{-1}^1 e^{ik\lambda} d\lambda = -2 \sum_{k=1}^{\infty} \frac{q^k e^{ikz} \sin k}{k}$. Therefore $\hat{\rho}(z) = 2c \left(-\sum_{k=1}^{\infty} \frac{q^k e^{ikz} \sin k}{k} + \sum_{k=1}^{\infty} \frac{q^k e^{ikz}}{k}\right) = 2c \cdot \sum_{k=1}^{\infty} \frac{q^k (k - \sin k)}{k^2} e^{ikz}$. Using (1), we get $\int_B \left(\frac{x - \sin x}{x}\right) U(dx) = \frac{1}{4\pi} \text{v.p.} \int_{\mathbb{R}} \left(2c \cdot \sum_{k=1}^{\infty} \frac{q^k (k - \sin k)}{k^2} e^{ikz}\right) \left(\int_B e^{-izx} dx\right) dz = c \int_B \left(\sum_{k=1}^{\infty} \frac{q^k (k - \sin k)}{k^2} \frac{1}{2\pi} \cdot \text{v.p.} \int_{\mathbb{R}} e^{iz(k-x)} dz\right) dx = c \sum_{k=1}^{\infty} \frac{q^k (k - \sin k)}{k^2} \int_B \delta(k - x) dx$. Therefore $\int_B \left(\frac{x - \sin x}{x}\right) U(dx) = \int_B \sum_{k=1}^{\infty} \left(\frac{k - \sin k}{k}\right) \frac{cq^k}{k} \delta(k - x) dx$. It now follows that $U(\{k\}) = \frac{cq^k}{k}$.

(e) The characteristic function of the *hyperbolic cosine distribution* has the form $\phi(z) = 1/\cosh(\pi z/2)$ [1]. Thus we have $\psi''(z) = (\log \phi(z))'' = -(\pi^2/4)\phi^2(z)$. The density of the hyperbolic cosine distribution equals to $f(x) = 1/(\pi \cosh x)$. It is obvious that the convolution product has the form $f * f(x) = \int_{\mathbb{R}} f(x - y)f(y)dy = (4/\pi^2)e^{-x} \int_{\mathbb{R}} (1 + e^{2y-2x})^{-1}(1 + e^{-2y})^{-1}dy = \frac{4}{\pi^2} \frac{x}{e^x - e^{-x}}$. Therefore $\int_{\mathbb{R}} e^{izx} \frac{x}{e^x - e^{-x}} dx = (\pi^2/4)\phi^2(z)$. Using (4), we get $u(x) = \frac{1}{x^2} \cdot \frac{1}{2\pi} \int_{\mathbb{R}} e^{-izx} (\pi^2/4)\phi^2(z) dz = \frac{1}{x^2} \cdot \frac{x}{e^x - e^{-x}} = \frac{1}{x(e^x - e^{-x})}$.

(f) The characteristic function of the Γ -*distribution* has the form $\phi(z) = (1 - i\alpha^{-1}z)^{-c}$, $c > 0$, $\alpha > 0$. It follows that $\psi(z) = \log \phi(z) = -c \log(1 - i\alpha^{-1}z)$, $\psi''(z) = \frac{c}{(i\alpha + z)^2}$. Using (4), we obtain $u(x) = -\frac{1}{x^2} \cdot \frac{1}{2\pi} \int_{\mathbb{R}} e^{-izx} \frac{c}{(i\alpha + z)^2} dz = (c/x)e^{-\alpha x} \mathbf{I}_{\{x > 0\}}$, since $\int_{\mathbb{R}} e^{izx} cxe^{-\alpha x} \mathbf{I}_{\{x > 0\}} dx = -\frac{c}{(i\alpha + z)^2}$.

The author is grateful to Professor A.N. Shiryaev for constant attention to this work.

References

- [1] W. Feller, *An introduction to probability theory and its applications*, Vol.2, 2nd ed., Wiley, New York, 1971. [2] E. Lukacs, *Characteristic functions*, 2nd ed., Griffin, London, 1970.

E.V. Burnaev
Skoltech

Presented by A.V. Bulinskiy
Accepted by editorial board

E-mail: e.burnaev@skoltech.ru