

Unramified Cohomology of Quadrics in Characteristic Two

Yong HU and Peng SUN

Abstract

Let F be a field of characteristic 2 and let X be a smooth projective quadric of dimension ≥ 1 over F . We study the unramified cohomology groups with 2-primary torsion coefficients of X in degrees 2 and 3. We determine completely the kernel and the cokernel of the natural map from the cohomology of F to the unramified cohomology of X . This extends the results in characteristic different from 2 obtained by Kahn, Rost and Sujatha in the nineteen-nineties.

Key words: Quadratic forms, quadrics, unramified cohomology, cycle class map

MSC classification 2020: 11E04, 14F20, 19E15

1 Introduction

Let F be a field. Let m be a positive integer not divisible by the characteristic of F . For each $j \geq 1$, the tensor product $\mathbb{Z}/m(j-1) := \mu_m^{\otimes(j-1)}$ of m -th roots of unity can be viewed as étale sheaves on F -schemes. Let X be a proper smooth connected variety over F . The unramified cohomology group $H_{nr}^j(X, \mathbb{Z}/m(j-1))$ is the group $H_{Zar}^0(X, \mathcal{H}_m^j(j-1))$, where $\mathcal{H}_m^j(j-1)$ denotes the Zariski sheaf associated to the presheaf $U \mapsto H_{\acute{e}t}^j(U, \mathbb{Z}/m(j-1))$. By taking the direct limit, we can also define

$$H_{nr}^j(X, (\mathbb{Q}/\mathbb{Z})'(j-1)) := \varinjlim_{\text{char}(F) \nmid m} H_{nr}^j(X, \mathbb{Z}/m(j-1)).$$

These groups can also be described in terms of residue maps in Galois cohomology, thanks to the Bloch–Ogus theorem on Gersten’s conjecture ([BO74]). As important birational invariants, they have found many important applications, for instance to the rationality problem (see e.g. [Sal84], [CTO89], [CTP16]), and have been extensively studied in the literature. With the development of the motivic cohomology theory (by Beilinson, Lichtenbaum, Suslin, Voevodsky, et al.), even more machinery can be applied to compute unramified cohomology nowadays.

When F has characteristic different from 2 and X is a smooth projective quadric, the above unramified cohomology groups are computed up to degree $j \leq 4$ by Kahn, Rost and Sujatha in a series of papers ([Kah95], [KRS98], [KS00], [KS01]). Some of their results are further developed and used by Izhboldin [Izh01] to solve a number of problems

on quadratic forms, including a construction of fields of u -invariant 9 in characteristic $\neq 2$.

It has been noticed for decades that unramified cohomology theory can be formulated in a more general setting (see [CT95], [CTHK97], [Kah04]). In particular, when F has positive characteristic p , the aforementioned groups have p -primary torsion variants. Indeed, the unramified cohomology functors $H_{nr}^j(\cdot, \mathbb{Z}/p^r(j-1))$ for all $r \geq 1$ and their limit $H_{nr}^j(\cdot, \mathbb{Q}_p/\mathbb{Z}_p(j-1))$ can be defined by using the Hodge–Witt cohomology (see (3.1) and (3.10)). In contrast to the prime-to- p case, there has been much less work on these p -primary unramified cohomology groups.

In this paper, we are interested in the case of a smooth projective quadric X over a field F of characteristic 2. We investigate the unramified cohomology groups via the natural maps

$$\eta_r^j : H^j(F, \mathbb{Z}/2^r(j-1)) \longrightarrow H_{nr}^j(X, \mathbb{Z}/2^r(j-1)), \quad r \geq 1$$

and

$$\eta_\infty^j : H^j(F, \mathbb{Q}_2/\mathbb{Z}_2(j-1)) \longrightarrow H_{nr}^j(X, \mathbb{Q}_2/\mathbb{Z}_2(j-1)).$$

For each $j \geq 1$, it is not difficult to see that the maps η_r^j for different $r \geq 1$ have essentially the same behavior (Lemma 4.3). So we may focus on the two maps $\eta^j := \eta_1^j$ and η_∞^j . They are both isomorphisms if $j = 1$ (Prop. 3.4) or X is isotropic (Prop. 4.1 (1)). In our main results, we determine completely the kernel and the cokernel of η^j and η_∞^j for $j = 2, 3$.

The following theorem extends Kahn’s results in [Kah95] to characteristic 2.

Theorem 1.1. *Let F be a field of characteristic 2 and let X be the smooth projective quadric defined by a nondegenerate quadratic form φ with $\dim \varphi \geq 3$. Assume that φ is anisotropic.*

1. (See 5.1, 5.3, 5.6 and 5.7) *Suppose $\dim \varphi = 3$, so that X is the conic associated to a quaternion division algebra D , or $\dim \varphi = 4$ and $e_1(\varphi) = 0 \in H^1(F, \mathbb{Z}/2)$, that is, φ is similar to the reduced norm of a quaternion division algebra D .*

Then

$$\begin{aligned} \text{Coker}(\eta^2) &\cong \text{Ker}(\eta^2) = \{0, (D)\} \cong \mathbb{Z}/2, \\ \text{Coker}(\eta^3) &\cong \text{Ker}(\eta^3) = \{(a) \cup (D) \mid a \in F^*\}, \end{aligned}$$

where (D) denotes the Brauer class of D .

2. (See 5.8 and 6.7) *If $\dim \varphi = 4$ and $e_1(\varphi) \neq 0 \in H^1(F, \mathbb{Z}/2)$, then*

$$\begin{aligned} \text{Coker}(\eta^2) &= \text{Ker}(\eta^2) = 0, \quad \text{Coker}(\eta^3) \cong \text{Ker}(\eta^3), \\ \text{Ker}(\eta^3) &= \{0\} \cup \{(a) \cup (b) \cup (c) \mid \varphi \text{ is similar to a subform of } \langle\langle a, b; c \rangle\rangle\}. \end{aligned}$$

3. (See 6.7, 6.11, 6.12 and 8.4) *Suppose $\dim \varphi > 4$. Then $\text{Coker}(\eta^2) = \text{Ker}(\eta^2) = 0$ and*

- (a) if φ is an Albert form (i.e. a 6-dimensional form with trivial Arf invariant), then $\text{Coker}(\eta^3) \cong \mathbb{Z}/2$ and $\text{Ker}(\eta^3) = 0$;
- (b) if φ is a neighbor of a 3-Pfister form $\langle\langle a, b; c \rangle\rangle$, then

$$\text{Coker}(\eta^3) \cong \text{Ker}(\eta^3) = \{0, (a) \cup (b) \cup (c)\} \cong \mathbb{Z}/2.$$

- (c) in all the other cases (e.g. $\dim \varphi > 8$), $\text{Coker}(\eta^3) = \text{Ker}(\eta^3) = 0$.

The counterpart in characteristic $\neq 2$ of the next theorem appeared in [KRS98, Thms. 4 and 5].

Theorem 1.2 (See 4.3, 6.8 and 8.5). *Let F be a field of characteristic 2 and let X be the smooth projective quadric defined by a nondegenerate quadratic form φ with $\dim \varphi \geq 3$.*

Then $\text{Ker}(\eta_r^j) = \text{Ker}(\eta_\infty^j)$ for all $r \geq 1, j \geq 1$, and $\text{Coker}(\eta_\infty^2) = 0$.

If φ is an anisotropic Albert form, then $\text{Coker}(\eta^2) \cong \mathbb{Z}/2$. Otherwise $\text{Coker}(\eta^3) = 0$.

Main tools in our proofs include the Bloch–Ogus and the Hochschild–Serre spectral sequences. A key difference between the p -primary torsion cohomology and the prime-to- p case is the lack of homotopy invariance. This results in the phenomenon that our spectral sequences look different from their analogues in characteristic different from 2. Fortunately, due to vanishing theorems for the local cohomology and the fact that the field F has 2-cohomological dimension at most 1, these spectral sequences still have many vanishing terms.

The cycle class maps with finite or divisible coefficients are also studied and used in the paper. In this respect we need information about the structure of Chow groups in low codimension. This information can be found in Karpenko’s work [Kar90] in characteristic $\neq 2$, and recently the paper [HLS21] has provided the corresponding results in characteristic 2.

In the above two theorems the case of Albert quadrics is more subtle than the others. In that case we have to utilize more techniques from the algebraic theory of quadratic forms, especially residue maps on Witt groups of discrete valuation fields of characteristic 2 ([Ara18]).

Notation and conventions. For any field k , denote by \bar{k} a separable closure of k .

For an algebraic variety Y over k , we write $Y_L = Y \times_k L$ for any field extension L/k , and $\bar{Y} = Y \times_k \bar{k}$. We say Y is k -rational if it is integral and birational to the projective space $\mathbb{P}_k^{\dim Y}$ over k . We say Y is *geometrically rational* if Y_L is L -rational for the algebraic closure L of k .

Milnor K -groups of a field k are denoted by $K_i^M(k)$, $i \in \mathbb{N}$.

For an abelian group M , we denote by M_{tors} the subgroup of torsion elements in M . For any positive integer n , we define $M[n]$ and M/n via the exact sequence

$$0 \longrightarrow M[n] \longrightarrow M \xrightarrow{\times n} M \longrightarrow M/n \longrightarrow 0.$$

For any scheme X , let $\text{Br}(X) = H_{\text{ét}}^2(X, \mathbb{G}_m)$ denote its cohomological Brauer group. In the rest of the paper, F denotes a field of characteristic 2.

2 Quadrics and their Chow groups

(2.1) We recall some basic definitions and facts about quadratic forms in characteristic 2. For general reference we refer to [EKM08].

We work over a field F , which has characteristic 2 according to our convention.

Let $a \in F^*$. We denote by $\langle a \rangle$ the 1-dimensional quadratic form $x \mapsto ax^2$, and let $[1, a]$ or $\langle\langle a \rangle\rangle$ denote the binary quadratic form $(x, y) \mapsto x^2 + xy + ay^2$. A (quadratic) 1-Pfister form is a binary quadratic form isomorphic to $\langle\langle a \rangle\rangle = [1, a]$ for some $a \in F^*$. Let $\langle 1, a \rangle_{\text{bil}}$ denote the binary bilinear form $((x_1, x_2), (y_1, y_2)) \mapsto x_1y_1 + ax_2y_2$. For $n \geq 2$, a quadratic form is called an n -Pfister form if it is isomorphic to

$$\langle\langle a_1, \dots, a_{n-1}; a_n \rangle\rangle := \langle 1, a_1 \rangle_{\text{bil}} \otimes \dots \otimes \langle 1, a_{n-1} \rangle_{\text{bil}} \otimes \langle\langle a_n \rangle\rangle$$

for some $a_1, \dots, a_{n-1} \in F^*$ and $a_n \in F$. If $\lambda \in F^*$ and φ is a Pfister form, the scalar multiple $\lambda\varphi$ is called a *general Pfister form*.

For two quadratic forms φ and ψ over F , we say ψ is a *subform* of φ if $\psi \cong \varphi|_W$ for some subspace W in the vector space V of φ . For $n \geq 2$, an n -Pfister neighbor is a subform of dimension $> 2^{n-1}$ of a general n -Pfister form.

We write $I_q(F)$ or $I_q^1(F)$ for the Witt group of even-dimensional nondegenerate quadratic forms over F . For $n \geq 2$, let $I_q^n(F)$ denote the subgroup of $I_q(F)$ generated by the n -Pfister forms. For a quadratic form φ over F , we will write $\varphi \in I_q^n(F)$ if φ is nondegenerate, of even dimension, and its Witt class lies in $I_q^n(F)$.

We also have the Witt ring $W(F)$ of nondegenerate symmetric bilinear forms over F , in which the classes of even-dimensional forms form an ideal $I(F)$, called the *fundamental ideal*. For each $n \geq 1$, let $I^n(F)$ be the n -th power of the ideal $I(F)$ and put $I^0(F) = W(F)$. The group $I_q(F)$ has a $W(F)$ -module structure, and we have $I_q^n(F) = I^{n-1}(F) \cdot I_q(F)$ for all $n \geq 1$.

The Galois cohomology group $H^1(F, \mathbb{Z}/2)$ can be identified with $F/\wp(F)$ by Artin–Schreier theory, where \wp denotes the map $x \mapsto x^2 - x$. For any $b \in F$, we write (b) for its canonical image in $F/\wp(F) = H^1(F, \mathbb{Z}/2)$. The map

$$e_1 : I_q^1(F) \longrightarrow H^1(F, \mathbb{Z}/2); \quad \langle\langle a \rangle\rangle \longmapsto (a)$$

is a well defined homomorphism, often called the *discriminant* or *Arf invariant*. It is well known that e_1 is surjective with $\text{Ker}(e_1) = I_q^2(F)$. A 6-dimensional form in $I_q^2(F)$ (i.e. a 6-dimensional nondegenerate form with trivial Arf invariant) is called an *Albert form*.

For $n \geq 2$, by using the Kato–Milne group $H^n(F, \mathbb{Z}/2(n-1))$ (cf. [Kat80], [Mil76]), a generalization of which will be discussed in (3.1), one can also define a functorial homomorphism (see [Sah72] for $n = 2$ and [Kat82b] for general n)

$$e_n : I_q^n(F) \longrightarrow H^n(F, \mathbb{Z}/2(n-1))$$

which is surjective with $\text{Ker}(e_n) = I_q^{n+1}(F)$ such that

$$e_n(\langle\langle a_1, \dots, a_{n-1}; a_n \rangle\rangle) = (a_1) \cup \dots \cup (a_{n-1}) \cup (a_n),$$

where for any $a \in F^*$, (a) denotes its canonical image in F^*/F^{*2} . (The maps e_2 and e_3 are more classical, called the *Clifford invariant* and the *Arason invariant* respectively.)

Note that for all $n \geq 1$,

$$(2.1.1) \quad e_n(\varphi) = e_n(c.\varphi) \quad \text{for all } \varphi \in I_q^n(F) \quad \text{and all } c \in F^*$$

because $\varphi - c\varphi = \langle\langle c \rangle\rangle\varphi \in I_q^{n+1}(F) = \text{Ker}(e_n)$.

The group $H^2(F, \mathbb{Z}/2(1))$ may be identified with the 2-torsion subgroup of the Brauer group $\text{Br}(F)$ of F . For any $a \in F^*$ and $b \in F$, let $(a, b]$ be the quaternion algebra generated by two elements i, j subject to the relations

$$i^2 = a, \quad j^2 + j = b, \quad ij = ji + i.$$

Its reduced norm is the 2-Pfister form $\langle\langle a; b \rangle\rangle$, and its Brauer class is $(a) \cup (b)$, the Clifford invariant of $\langle\langle a; b \rangle\rangle$. The plane conic defined by the ternary form $\langle a \rangle \perp \langle\langle b \rangle\rangle$ is called the conic associated to the quaternion algebra $(a, b]$.

For any $a, c \in F^*$ and $b, d \in F$, the form $[1, b+d] \perp a.[1, b] \perp c.[1, d]$ is an Albert form. Its Clifford invariant is the Brauer class of the biquaternion algebra $(a, b] \otimes_F (c, d]$.

(2.2) Now we recall some known facts about Chow groups of projective quadrics (which are valid in arbitrary characteristic). More details can be found in [Kar90, §2], [EKM08, §68] and [HLS21, §5].

Let φ be a quadratic form defined on a (finite dimensional) F -vector space V of dimension ≥ 3 . Let $X = X_\varphi$ be the projective quadric defined by φ . It is a closed subvariety in the projective space $\mathbb{P}(V)$. Unless otherwise stated, we always assume φ is nondegenerate, which means the projective quadric $X = X_\varphi$ is smooth as an algebraic variety over F . In fact, X is a smooth geometrically integral F -variety. It is well known that if K/F is a field extension such that φ_K is isotropic, then X_K is K -rational. In particular, \overline{X} is rational over \overline{F} .

For each $i \in \mathbb{N}$, let $\text{CH}^i(X)$ denote the Chow group of codimension i cycles of X . Let $h \in \text{CH}^1(X)$ be the class of a hyperplane section (i.e., the pullback of the hyperplane class in $\text{CH}^1(\mathbb{P}(V))$). Using the intersection pairing as multiplication in the Chow ring ([EKM08, §57]), we get elements $h^i \in \text{CH}^i(X)$ for each i .

Set $d = \dim X$. For every integer $j \in [0, d/2]$, let $\ell_j \in \text{CH}^{d-j}(\overline{X})$ be the class of a j -dimensional linear subspace contained in \overline{X} . Then, for each $0 \leq i \leq d$, we have

$$\text{CH}^i(X) = \text{CH}_{d-i}(X) = \begin{cases} \mathbb{Z}.h^i & \text{if } 0 \leq i < \frac{d}{2}, \\ \mathbb{Z}.\ell_{d-i} & \text{if } \frac{d}{2} < i \leq d, \\ \mathbb{Z}.h^i \oplus \mathbb{Z}.\ell_i & \text{if } i = \frac{d}{2}. \end{cases}$$

If $\dim X = 2m$ is even, there are exactly two different classes of m -dimensional linear subspaces ℓ_m, ℓ'_m in $\text{CH}^m(X)$ and the sum of these two classes is equal to h^m .

The next two propositions can be proved in arbitrary characteristic.

Proposition 2.3 ([Kah99, Lemma 8.2]). *Let $X = X_\varphi$ be a smooth projective quadric of even dimension $\dim X = 2m \geq 2$.*

Then $\mathrm{CH}^m(\overline{X})$ is a trivial Galois module if and only if $e_1(\varphi) = 0$. When $e_1(\varphi) \neq 0$, the Galois action permutes the two classes ℓ_m and ℓ'_m .

Proposition 2.4 ([Kar90, (2.4)]). *Let X be the smooth projective quadric defined by an anisotropic form φ with $\dim \varphi \geq 3$.*

1. *If $\dim X \geq 3$, then*

$$\mathrm{CH}^1(X) = \mathrm{CH}^1(\overline{X})^{\mathrm{Gal}(\overline{k}/k)} = \mathrm{CH}^1(\overline{X}) = \mathbb{Z}.h.$$

2. *If $\dim X = 2$ and $e_1(\varphi) \neq 0$, then*

$$\mathrm{CH}^1(X) = \mathbb{Z}.h = \mathrm{CH}^1(\overline{X})^{\mathrm{Gal}(\overline{k}/k)} \subseteq \mathrm{CH}^1(\overline{X}) = \mathbb{Z}.h \oplus \mathbb{Z}.\ell_1.$$

3. *If $\dim X = 2$ and $e_1(\varphi) = 0$ is trivial, then*

$$\mathrm{CH}^1(X) = \mathbb{Z}.2\ell_1 \oplus \mathbb{Z}.h \subseteq \mathrm{CH}^1(\overline{X})^{\mathrm{Gal}(\overline{k}/k)} = \mathrm{CH}^1(\overline{X}) = \mathbb{Z}.\ell_1 \oplus \mathbb{Z}.h.$$

4. *If $\dim X = 1$, then*

$$\mathrm{CH}^1(X) = \mathbb{Z}.h = \mathbb{Z}.2\ell_0 \subseteq \mathrm{CH}^1(\overline{X}) = \mathrm{CH}^1(\overline{X})^{\mathrm{Gal}(\overline{k}/k)} = \mathbb{Z}.\ell_0.$$

3 Unramified cohomology in positive characteristic

Throughout this section, we fix a positive integer r and a field k of characteristic $p > 0$.

(3.1) For each $i \in \mathbb{N}$, let $\nu_r(i) = W_r\Omega_{\log}^i$ be the i -th logarithmic Hodge–Witt sheaf on the big étale site of k ([Ill79], [Shi07]). Define $\mathbb{Z}/p^r(i) := \nu_r(i)[-i]$, as an object in the derived category of étale sheaves. This object can also be viewed as an étale motivic complex ([GL00]). For every integer b , we have the cohomology functor on k -schemes

$$H^b(\cdot, \mathbb{Z}/p^r(i)) := H_{\text{ét}}^b(\cdot, \mathbb{Z}/p^r(i)) = H_{\text{ét}}^{b-i}(\cdot, \nu_r(i)).$$

For short hand, we sometimes write $H_{p^r}^b(\cdot, i)$ instead of the precise notation $H^b(\cdot, \mathbb{Z}/p^r(i))$. The Zariski sheaf associated to the presheaf $U \mapsto H^b(U, \mathbb{Z}/p^r(i))$ is denoted by $\mathcal{H}_{p^r}^b(i)$.

For a smooth connected k -variety X , we define the *unramified cohomology group*

$$H_{nr}^b(X, \mathbb{Z}/p^r(i)) := H_{\text{Zar}}^0(X, \mathcal{H}_{p^r}^b(i)).$$

This group can also be described by using the Cousin complex of X . It is naturally a subgroup of $H^b(k(X), \mathbb{Z}/p^r(i))$.

In the theorem below, we collect some well known results that are needed in this paper.

Theorem 3.2. *Let X be a smooth connected k -variety.*

1. *We have the Bloch–Ogus spectral sequence*

$$(3.2.1) \quad E_2^{a,b} = H_{Zar}^a(X, \mathcal{H}_{p^r}^b(i)) \implies E^{a+b} = H^{a+b}(X, \mathbb{Z}/p^r(i))$$

with

$$(3.2.2) \quad E_2^{a,b} = 0 \quad \text{if } b \notin \{i, i+1\}, \text{ or if } a > b = i$$

and

$$(3.2.3) \quad E_2^{i,i} = H_{Zar}^i(X, \mathcal{H}_{p^r}^i(i)) \cong \text{CH}^i(X)/p^r.$$

2. *There are natural isomorphisms*

$$(3.2.4) \quad \begin{aligned} H^i(X, \mathbb{Z}/p^r(i)) &\xrightarrow{\sim} H_{Zar}^0(X, \mathcal{H}_{p^r}^i(i)) = H_{nr}^i(X, \mathbb{Z}/p^r(i)), \\ H^{2i+j}(X, \mathbb{Z}/p^r(i)) &\xrightarrow{\sim} H_{Zar}^{j+i-1}(X, \mathcal{H}_{p^r}^{i+1}(i)) \quad \text{for } j \geq 1, \end{aligned}$$

and

$$(3.2.5) \quad H_{nr}^2(X, \mathbb{Z}/p^r(1)) = H_{Zar}^0(X, \mathcal{H}_{p^r}^2(1)) \cong \text{Br}(X)[p^r].$$

3. *For smooth proper connected k -varieties, the group $H_{nr}^b(X, \mathbb{Z}/p^r(i))$ is a k -birational invariant.*

4. *Let $\pi : X \rightarrow Y$ be a proper morphism between smooth connected k -varieties whose generic fiber is $k(Y)$ -rational.*

Then the natural map $\pi^ : H_{nr}^b(Y, \mathbb{Z}/p^r(i)) \rightarrow H_{nr}^b(X, \mathbb{Z}/p^r(i))$ is an isomorphism.*

Proof. (1) These are standard consequences of Gersten’s conjecture for smooth varieties ([CTHK97, Cor. 5.1.11 and § 7.4 (3)], [Shi07, Thm. 4.1]) and the Bloch–Kato–Gabber theorem ([GS17, Thm. 9.5.2]). The nontrivial part of (3.2.2) is proved in [Gro85, (3.5.3)] and [Shi07, Cor. 3.4].

(3) The isomorphisms in (3.2.4) follow easily from the vanishing results in (3.2.2). The isomorphism (3.2.5) is a consequence of purity for Brauer groups ([Č19, Thm. 1.2]).

(4) See [CTHK97, Thm. 8.5.1].

(5) See [CTHK97, Thm. 8.6.1]. □

(3.3) Let X be a smooth connected k -variety. For every $j \geq 1$, we have a natural restriction map

$$(3.3.1) \quad \eta_r^j : H^j(k, \mathbb{Z}/p^r(j-1)) \longrightarrow H_{nr}^j(X, \mathbb{Z}/p^r(j-1)).$$

Let $i \in \mathbb{N}$ be such that $i \leq j$. We have the cup product map

$$(3.3.2) \quad H_{nr}^{j-i}(X, \mathbb{Z}/p^r(j-i)) \otimes H_{Zar}^i(X, \mathcal{H}_{p^r}^i(i)) \xrightarrow{\cup} H_{Zar}^j(X, \mathcal{H}_{p^r}^j(j))$$

and a natural map

$$(3.3.3) \quad H^{j-i}(k, \mathbb{Z}/p^r(j-i)) \otimes \mathrm{CH}^i(X)/p^r \longrightarrow H_{nr}^{j-i}(X, \mathbb{Z}/p^r(j-i)) \otimes H_{Zar}^i(X, \mathcal{H}_{p^r}^i(i))$$

induced from the identification $\mathrm{CH}^i(X)/p^r = H_{Zar}^i(X, \mathcal{H}_{p^r}^i(i))$ in (3.2.3) and the natural map

$$H^{j-i}(k, \mathbb{Z}/p^r(j-i)) \longrightarrow H_{Zar}^0(X, \mathcal{H}_{p^r}^{j-i}(j-i)) = H_{nr}^{j-i}(X, \mathbb{Z}/p^r(j-i)).$$

Composing (3.3.2) with (3.3.3) yields a natural map

$$(3.3.4) \quad \mu_r^{i,j} : H^{j-i}(k, \mathbb{Z}/p^r(j-i)) \otimes \mathrm{CH}^i(X)/p^r \longrightarrow H_{Zar}^i(X, \mathcal{H}_{p^r}^j(j)).$$

For $i = 0$ this coincides with the natural map

$$\tilde{\eta}_r^j : H^j(k, \mathbb{Z}/p^r(j)) \rightarrow H_{nr}^j(X, \mathbb{Z}/p^r(j)) = H^j(X, \mathbb{Z}/p^r(j)).$$

Proposition 3.4. *With notation as in (3.3), suppose that X is proper and geometrically rational.*

Then the natural map $\eta_r^1 : H^1(k, \mathbb{Z}/p^r) \rightarrow H_{nr}^1(X, \mathbb{Z}/p^r)$ is an isomorphism.

Proof. By (3.2.4), we have $H_{nr}^1(X, \mathbb{Z}/p^r) = H^1(X, \mathbb{Z}/p^r)$. The geometric rationality of X implies that $\overline{X} = X \times_k \overline{k}$ (where \overline{k} denotes a separable closure of k) has trivial étale fundamental group. By the homotopy exact sequence for étale fundamental groups ([Gro03, (IX.6.1)]) we get $\pi_1(X) \cong \mathrm{Gal}(\overline{k}/k)$. Hence the natural map

$$H^1(k, \mathbb{Z}/p^r) = \mathrm{Hom}(\mathrm{Gal}(\overline{k}/k), \mathbb{Z}/p^r) \longrightarrow H^1(X, \mathbb{Z}/p^r) = \mathrm{Hom}(\pi_1(X), \mathbb{Z}/p^r)$$

is an isomorphism. □

Proposition 3.5. *Let X be a smooth, proper, k -rational variety.*

Then for every $j \geq 1$, the map

$$\eta_r^j : H^j(k, \mathbb{Z}/p^r(j-1)) \longrightarrow H_{nr}^j(X, \mathbb{Z}/p^r(j-1))$$

is an isomorphism.

Proof. This is a special case of Theorem 3.2 (4). □

(3.6) Let X be a smooth connected k -variety. The Bloch–Ogus spectral sequence (3.2.1) is concentrated in two horizontal lines (cf. (3.2.2)). So we can obtain natural homomorphisms

$$e^a(i) : E_2^{a,i} = H_{Zar}^a(X, \mathcal{H}_{p^r}^i(i)) \longrightarrow E^{i+a} = H^{i+a}(X, \mathbb{Z}/p^r(i)), \quad 1 \leq a \leq i$$

which fit into a long exact sequence

$$(3.6.1) \quad \begin{array}{ccccccc} 0 & \longrightarrow & E_2^{1,i} & \xrightarrow{e^1(i)} & E^{i+1} & \longrightarrow & E_2^{0,i+1} = H_{Zar}^0(X, \mathcal{H}_{p^r}^{i+1}(i)) \\ & & \xrightarrow{d} & E_2^{2,i} & \xrightarrow{e^2(i)} & E^{i+2} & \longrightarrow E_2^{1,i+1} = H_{Zar}^1(X, \mathcal{H}_{p^r}^{i+1}(i)) \rightarrow \dots \\ & & & \dots & & & \\ & & \xrightarrow{d} & E_2^{a,i} & \xrightarrow{e^a(i)} & E^{i+a} & \longrightarrow E_2^{a-1,i+1} = H_{Zar}^{a-1}(X, \mathcal{H}_{p^r}^{i+1}(i)) \rightarrow \dots \\ & & & \dots & & & \\ & & \xrightarrow{d} & E_2^{i,i} & \xrightarrow{e^i(i)} & E^{2i} & \longrightarrow E_2^{i-1,i+1} = H_{Zar}^{i-1}(X, \mathcal{H}_{p^r}^{i+1}(i)) \longrightarrow 0. \end{array}$$

In particular, we have a natural map

$$(3.6.2) \quad \mathrm{cl}_X^i = \mathrm{cl}^i := e^i(i) : E_2^{i,i} = \mathrm{CH}^i(X)/p^r \longrightarrow E^{2i} = H^{2i}(X, \mathbb{Z}/p^r(i))$$

called the *cycle class map*.

Let $j \in \mathbb{N}$ be another integer with $j \geq i$. By composing $e^i(j)$ with the map $\mu_r^{i,j}$ in (3.3.4) we get a natural map

$$(3.6.3) \quad \nu_r^{i,j} : H^{j-i}(k, \mathbb{Z}/p^r(j-i)) \otimes \mathrm{CH}^i(X)/p^r \longrightarrow H^{i+j}(X, \mathbb{Z}/p^r(j)).$$

Compatibility of the Bloch–Ogus spectral sequence with cup products (cf. [KRS98, p.868]) gives the commutative diagram

$$(3.6.4) \quad \begin{array}{ccc} & H^{j-i}(k, \mathbb{Z}/p^r(j-i)) \otimes \mathrm{CH}^i(X)/p^r & \\ \tilde{\eta}_r^{j-i} \otimes \mathrm{Id} \swarrow & & \searrow \tilde{\eta}_r^{j-i} \otimes \mathrm{cl}_X^i \\ S & \xrightarrow{\mathrm{Id} \otimes e^i(i)} & T \\ \mu_r^{i,j} \swarrow & & \searrow \nu_r^{i,j} \\ H_{\mathrm{Zar}}^i(X, \mathcal{H}_{p^r}^j(j)) & \xrightarrow{e^i(j)} & H^{i+j}(X, \mathbb{Z}/p^r(j)) \end{array}$$

(Note: The diagram above is a simplified representation of the commutative diagram in the image. The top node is $H^{j-i}(k, \mathbb{Z}/p^r(j-i)) \otimes \mathrm{CH}^i(X)/p^r$. The left node is S and the right node is T . The bottom-left node is $H_{\mathrm{Zar}}^i(X, \mathcal{H}_{p^r}^j(j))$ and the bottom-right node is $H^{i+j}(X, \mathbb{Z}/p^r(j))$. Arrows: $S \xrightarrow{\tilde{\eta}_r^{j-i} \otimes \mathrm{Id}} \text{Top}$, $T \xrightarrow{\tilde{\eta}_r^{j-i} \otimes \mathrm{cl}_X^i} \text{Top}$, $S \xrightarrow{\mathrm{Id} \otimes e^i(i)} T$, $S \xrightarrow{\mu_r^{i,j}} \text{Bottom-Left}$, $T \xrightarrow{\nu_r^{i,j}} \text{Bottom-Right}$, $\text{Bottom-Left} \xrightarrow{e^i(j)} \text{Bottom-Right}$, $S \xrightarrow{\cup} \text{Bottom-Left}$, $T \xrightarrow{\cup} \text{Bottom-Right}$.

where

$$\begin{aligned} S &= H_{nr}^{j-i}(X, \mathbb{Z}/p^r(j-i)) \otimes H_{\mathrm{Zar}}^i(X, \mathcal{H}_{p^r}^i(i)), \\ T &= H_{nr}^{j-i}(X, \mathbb{Z}/p^r(j-i)) \otimes H^{2i}(X, \mathbb{Z}/p^r(i)). \end{aligned}$$

Proposition 3.7. *Let X be a smooth connected k -variety. Then there is an exact sequence*

$$0 \longrightarrow \mathrm{CH}^1(X)/p^r \xrightarrow{\mathrm{cl}_X^1} H_{\mathrm{et}}^2(X, 1) \longrightarrow \mathrm{Br}(X)[p^r] \longrightarrow 0.$$

In particular, the cycle class map cl_X^1 is injective, and if X is proper and k -rational, we have $\mathrm{Coker}(\mathrm{cl}_X^1) \cong \mathrm{Br}(k)[p^r]$.

Proof. Note that $H_{\mathrm{Zar}}^2(X, \mathcal{H}^1(1)) = 0$ by (3.2.2). Thus, taking $i = 1$ in (3.6.1) and noticing the isomorphism (3.2.5) we get the desired exact sequence.

Alternatively, we can use the long cohomology sequence for the distinguished triangle

$$\mathbb{Z}(1) = \mathbb{G}_m[-1] \xrightarrow{\times p^r} \mathbb{Z}(1) \longrightarrow \mathbb{Z}/p^r(1) \xrightarrow{+1}$$

of motivic complexes, which is an analogue of the Kummer sequence in characteristic $\neq p$. \square

Proposition 3.8. *Let X be a smooth proper k -rational variety. Then there is an exact sequence*

$$(3.8.1) \quad \begin{aligned} 0 &\longrightarrow H_{Zar}^1(X, \mathcal{H}_{p^r}^2(2)) \longrightarrow H^3(X, \mathbb{Z}/p^r(2)) \longrightarrow H^3(k, \mathbb{Z}/p^r(2)) \\ &\longrightarrow \mathrm{CH}^2(X)/p^r \xrightarrow{\mathrm{cl}_X^2} H^4(X, \mathbb{Z}/p^r(2)) \longrightarrow H_{Zar}^1(X, \mathcal{H}_{p^r}^3(2)) \longrightarrow 0. \end{aligned}$$

In particular, if $H^3(k, \mathbb{Z}/p^r(2)) = 0$ (e.g. if k is separably closed), then the cycle class map

$$\mathrm{cl}_X^2 : \mathrm{CH}^2(X)/p^r \longrightarrow H^4(X, \mathbb{Z}/p^r(2))$$

is injective.

Proof. Taking $i = 2$ in (3.6.1) we obtain an exact sequence

$$(3.8.2) \quad \begin{aligned} 0 &\longrightarrow H_{Zar}^1(X, \mathcal{H}_{p^r}^2(2)) \longrightarrow H^3(X, \mathbb{Z}/p^r(2)) \longrightarrow H_{nr}^3(X, \mathbb{Z}/p^r(2)) \\ &\longrightarrow \mathrm{CH}^2(X)/p^r \xrightarrow{\mathrm{cl}^2} H^4(X, \mathbb{Z}/p^r(2)) \longrightarrow H_{Zar}^1(X, \mathcal{H}_{p^r}^3(2)) \longrightarrow 0. \end{aligned}$$

The desired exact sequence follows from (3.8.2), because for the (smooth proper) k -rational variety X we have $H_{nr}^3(X, \mathbb{Z}/p^r(2)) \cong H^3(k, \mathbb{Z}/p^r(2))$ (Prop. 3.5). \square

(3.9) Let X be a smooth connected k -variety. For each $i \in \mathbb{N}$ we have the Hochschild–Serre spectral sequence

$$(3.9.1) \quad H^a(k, H^b(\overline{X}, \mathbb{Z}/p^r(i))) \implies H^{a+b}(X, \mathbb{Z}/p^r(i)).$$

Since $\mathrm{cd}_p(k) \leq 1$, we have $H^a(k, H^b(\overline{X}, \mathbb{Z}/p^r(i))) = 0$ for all $a > 1$. Thus, the spectral sequence (3.9.1) yields an isomorphism

$$(3.9.2) \quad H^0(k, H^i(\overline{X}, \mathbb{Z}/p^r(i))) \cong H^i(X, \mathbb{Z}/p^r(i))$$

and an exact sequence

$$(3.9.3) \quad \begin{aligned} 0 &\longrightarrow H^1(k, H^j(\overline{X}, \mathbb{Z}/p^r(i))) \longrightarrow H^{j+1}(X, \mathbb{Z}/p^r(i)) \\ &\longrightarrow H^0(k, H^{j+1}(\overline{X}, \mathbb{Z}/p^r(i))) \longrightarrow 0 \end{aligned}$$

for each $j \geq i$.

Now assume X is proper and geometrically rational (e.g. X is a smooth projective quadric of dimension ≥ 1). Then by (3.2.4) and Thm. 3.2 (4), we have

$$(3.9.4) \quad H^i(\overline{X}, \mathbb{Z}/p^r(i)) \cong H_{nr}^i(\overline{X}, \mathbb{Z}/p^r(i)) = H^i(\overline{k}, \mathbb{Z}/p^r(i)) = K_i^M(\overline{k})/p^r.$$

It is proved in [Lzh91, Cor. 6.5] that

$$(3.9.5) \quad \begin{aligned} H^i(k, \mathbb{Z}/p^r(i)) &= K_i^M(k)/p^r = H^0(k, K_i^M(\overline{k})/p^r), \\ H^{i+1}(k, \mathbb{Z}/p^r(i)) &= H^1(k, K_i^M(\overline{k})/p^r). \end{aligned}$$

Hence, combining (3.9.4), (3.9.5) and the case $j = i$ of (3.9.3), we obtain an exact sequence

$$(3.9.6) \quad 0 \longrightarrow H_{p^r}^{i+1}(k, i) \longrightarrow H^{i+1}(X, \mathbb{Z}/p^r(i)) \longrightarrow H^0(k, H^{i+1}(\overline{X}, \mathbb{Z}/p^r(i))) \longrightarrow 0.$$

On the other hand, since

$$H_{Zar}^0(\overline{X}, \mathcal{H}_{p^r}^{i+1}(i)) = H_{nr}^{i+1}(\overline{X}, \mathbb{Z}/p^r(i)) = H^{i+1}(\overline{k}, \mathbb{Z}/p^r(i)) = 0$$

by Thm. 3.2 (4), it follows from (3.6.1) that

$$(3.9.7) \quad H^{i+1}(\overline{X}, \mathbb{Z}/p^r(i)) = H_{Zar}^1(\overline{X}, \mathcal{H}_{p^r}^i(i)).$$

For $i = 0$, we can deduce from (3.9.7) and (3.2.2) that $H^1(\overline{X}, \mathbb{Z}/p^r) = 0$. Hence (3.9.6) yields $H^1(k, \mathbb{Z}/p^r) \cong H^1(X, \mathbb{Z}/p^r)$ in this case, recovering the result of Prop. 3.4.

For $i = 1$, from (3.9.7) and (3.2.3) we get

$$(3.9.8) \quad H^2(\overline{X}, \mathbb{Z}/p^r(1)) \cong \text{CH}^1(\overline{X})/p^r.$$

(This can also be deduced from Prop. 3.7.) So the case $i = 1$ of (3.9.6) gives an exact sequence

$$(3.9.9) \quad 0 \longrightarrow H^2(k, \mathbb{Z}/p^r(1)) \longrightarrow H^2(X, \mathbb{Z}/p^r(1)) \longrightarrow H^0(k, \text{CH}^1(\overline{X})/p^r) \longrightarrow 0.$$

Let \mathcal{H}_i denote the Zariski sheaf defined by Quillen's K -theory. Using the Bloch–Kato–Gabber theorem, we can deduce an exact sequence ([GS88, Thm. 4.13])

$$(3.9.10) \quad 0 \longrightarrow H_{Zar}^{i+1}(X, \mathcal{H}_i)/p^r \longrightarrow H_{Zar}^{i-1}(X, \mathcal{H}_{p^r}^i(i)) \longrightarrow \text{CH}^i(X)[p^r] \longrightarrow 0$$

for X and similarly for \overline{X} .

Now assume further that X is a projective homogeneous variety. Then $\text{CH}^i(\overline{X})$ is torsion-free and

$$H_{Zar}^{i-1}(\overline{X}, \mathcal{H}_i) = K_1(\overline{k}) \otimes \text{CH}^{i-1}(\overline{X})$$

by [Mer95, § 1, Prop. 1]. So, in this case from (3.9.10) we get

$$(3.9.11) \quad H_{Zar}^{i-1}(\overline{X}, \mathcal{H}_{p^r}^i(i)) \cong H_{Zar}^{i-1}(\overline{X}, \mathcal{H}_i)/p^r \cong K_1(\overline{k})/p^r \otimes \text{CH}^{i-1}(\overline{X})/p^r.$$

For $i = 2$ we can combine (3.9.11) and (3.9.7) to get an isomorphism

$$(3.9.12) \quad H^3(\overline{X}, \mathbb{Z}/p^r(2)) \cong K_1(\overline{k})/p^r \otimes \text{CH}^1(\overline{X})/p^r.$$

In particular, this holds when X is a smooth projective quadric of dimension ≥ 1 .

We end this section with a few remarks on cohomology with divisible coefficients.

(3.10) Given integers $b, i \in \mathbb{N}$, by taking direct limits we can define the functors

$$\begin{aligned} H^b(\cdot, \mathbb{Q}_p/\mathbb{Z}_p(i)) &:= \varinjlim_r H^b(\cdot, \mathbb{Z}/p^r(i)), \\ H_{nr}^b(\cdot, \mathbb{Q}_p/\mathbb{Z}_p(i)) &:= \varinjlim_r H_{nr}^b(\cdot, \mathbb{Z}/p^r(i)). \end{aligned}$$

It is easy to extend all the previous discussions in this section to these cohomology groups with divisible coefficients. In particular, for a smooth connected k -variety X , by taking the limits of (3.3.1) and (3.6.2) we obtain a natural map

$$\eta_\infty^j : H^j(k, \mathbb{Q}_p/\mathbb{Z}_p(j-1)) \longrightarrow H_{nr}^j(X, \mathbb{Q}_p/\mathbb{Z}_p(j-1)).$$

for each $j \geq 1$, and a cycle class map with divisible coefficients

$$\text{cl}_\infty^i : \text{CH}^i(X) \otimes (\mathbb{Q}_p/\mathbb{Z}_p) \longrightarrow H^{2i}(X, \mathbb{Q}_p/\mathbb{Z}_p(i))$$

for each $i \geq 1$.

A useful standard fact (which follows from [Lzh91, Lemma 6.6]) is that the sequence

$$(3.10.1) \quad 0 \rightarrow H_{p^r}^j(K, j-1) \rightarrow H^j(K, \mathbb{Q}_p/\mathbb{Z}_p(j-1)) \xrightarrow{\times p^r} H^j(K, \mathbb{Q}_p/\mathbb{Z}_p(j-1)) \rightarrow 0$$

is exact for any field extension K/k . As a consequence, we have an identification

$$(3.10.2) \quad \text{Ker}(\eta_r^j) = \text{Ker}(\eta_\infty^j)[p^r].$$

4 Some general observations for quadrics

From now on, we work over a field F of characteristic 2, and let X denote a smooth projective quadric of dimension ≥ 1 over F . Notation and results in §3 will be applied with $p = 2$.

For each $j \geq 1$, we have the natural maps (cf. (3.3.1) and (3.10))

$$\eta_r^j = \eta_{r,X}^j : H^j(F, \mathbb{Z}/2^r(j-1)) \longrightarrow H_{nr}^j(X, \mathbb{Z}/2^r(j-1)), \quad r \geq 1$$

and

$$\eta_\infty^j = \eta_{\infty,X}^j : H^j(F, \mathbb{Q}_2/\mathbb{Z}_2(j-1)) \longrightarrow H_{nr}^j(X, \mathbb{Q}_2/\mathbb{Z}_2(j-1)).$$

Observe that the proof of the following proposition is characteristic free.

Proposition 4.1 ([KRS98, Prop. 2.5]). *With notation as above, the following statements hold:*

1. *If X is isotropic over F , then the maps $\eta_{r,X}^j$ and $\eta_{\infty,X}^j$ are all isomorphisms.*
2. *In general, the maps $\eta_{r,X}^j$ and $\eta_{\infty,X}^j$ all have 2-torsion kernel and cokernel.*
3. *Let Y be another smooth projective quadric of dimension ≥ 1 over F . If X is isotropic over the function field $F(Y)$, then there is a natural commutative diagram*

$$\begin{array}{ccc} & H^j(F, \mathbb{Z}/2^r(j-1)) & \\ \eta_{r,X}^j \swarrow & & \searrow \eta_{r,Y}^j \\ H_{nr}^j(X, \mathbb{Z}/2^r(j-1)) & \xrightarrow{\rho} & H_{nr}^j(Y, \mathbb{Z}/2^r(j-1)) \end{array}$$

If moreover Y is isotropic over $F(X)$, the map ρ in the above diagram is an isomorphism.

Similar results hold in the case of divisible coefficients.

Corollary 4.2 ([CTS95, Lemma 1.3]). *Suppose φ is a Pfister form of dimension ≥ 4 and let ψ be a neighbor of φ . Let X and Y be the projective quadrics defined by φ and ψ respectively.*

Then for every $j \geq 1$ we have isomorphisms

$$\mathrm{Ker}(\eta_{r,X}^j) = \mathrm{Ker}(\eta_{r,Y}^j) \quad \text{and} \quad \mathrm{Coker}(\eta_{r,X}^j) \cong \mathrm{Coker}(\eta_{r,Y}^j), \quad r \geq 1,$$

and similarly in the case of divisible coefficients.

Proof. Apply Prop. 4.1 (3). □

Lemma 4.3. *For every $r \geq 1$ and $j \geq 1$, $\mathrm{Ker}(\eta_r^j) = \mathrm{Ker}(\eta_\infty^j)$ and there is an exact sequence*

$$0 \longrightarrow \mathrm{Ker}(\eta_r^j) \longrightarrow \mathrm{Coker}(\eta_r^j) \longrightarrow \mathrm{Coker}(\eta_\infty^j) \longrightarrow 0.$$

Proof. The first assertion is immediate from (3.10.2) and Prop. 4.1 (2).

By functoriality and (3.10.1) we have the commutative diagram with exact rows

$$\begin{array}{ccccccc} 0 & \longrightarrow & H^j(F, \mathbb{Z}/2^r(j-1)) & \longrightarrow & H^j(F, \mathbb{Q}_2/\mathbb{Z}_2(j-1)) & \xrightarrow{\times 2^r} & H^j(F, \mathbb{Q}_2/\mathbb{Z}_2(j-1)) & \longrightarrow & 0 \\ & & \eta_r^j \downarrow & & \eta_\infty^j \downarrow & & \eta_\infty^j \downarrow & & \\ 0 & \longrightarrow & H_{nr}^j(X, \mathbb{Z}/2^r(j-1)) & \longrightarrow & H_{nr}^j(X, \mathbb{Q}_2/\mathbb{Z}_2(j-1)) & \xrightarrow{\times 2^r} & H_{nr}^j(X, \mathbb{Q}_2/\mathbb{Z}_2(j-1)) & & \end{array}$$

Applying the snake lemma to this diagram yields the desired exact sequence, noticing that $\mathrm{Coker}(\eta_\infty^j)$ is 2-torsion. □

Thanks to the above lemma, when studying the kernel and the cokernel of η_r^j we may restrict to the case $r = 1$.

The following result is a special case of [Kah99, Prop. 5.2].

Proposition 4.4. *There is a natural isomorphism $\mathrm{Coker}(\eta_\infty^3) \cong \mathrm{Ker}(\mathrm{cl}_\infty^2)$ for the smooth projective quadric X .*

5 Results for conic curves

To simplify the notation, we henceforth write

$$\begin{aligned} H^b(\cdot, i) &= H_{\text{ét}}^b(\cdot, \mathbb{Z}/2(i)), \quad \mathcal{H}^b(i) = \mathcal{H}_2^b(i), \\ H^{i+1}(\cdot) &= H^{i+1}(\cdot, i), \quad H_{nr}^{i+1}(\cdot) = H_{nr}^{i+1}(\cdot, i). \end{aligned}$$

In this section and the next, we study the maps

$$\begin{aligned} \eta^j &:= \eta_1^j : H^j(F) \longrightarrow H_{nr}^j(X), \\ \eta_\infty^j &: H^j(F, \mathbb{Q}_2/\mathbb{Z}_2(j-1)) \longrightarrow H_{nr}^j(X, \mathbb{Q}_2/\mathbb{Z}_2(j-1)) \end{aligned}$$

for the quadric X . They are both isomorphisms if X is isotropic (Prop. 4.1 (1)) or $j = 1$ (Prop. 3.4). So we assume X is anisotropic and $j \geq 2$ in the sequel.

For a conic curve an explicit description is known for the kernel of the map η^j .

Proposition 5.1. *Let $X \subseteq \mathbb{P}_F^2$ be the conic associated to a quaternion F -algebra D . Then for all $j \geq 2$,*

$$\text{Ker}(\eta^j : H^j(F) \longrightarrow H_{nr}^j(X)) = \{(D) \cup \xi \mid \xi \in K_{j-2}^M(F)\}$$

where $(D) \in H^2(F) = \text{Br}(F)[2]$ denotes the Brauer class of D .

Proof. Since $H_{nr}^j(X) = H_{Zar}^0(X, \mathcal{H}^j(j-1))$ is a subgroup of $H^j(F(X))$, $\text{Ker}(\eta^j)$ coincides with the kernel of the natural map $H^j(F) \rightarrow H^j(F(X))$. The result thus follows from [AJ09, Thm. 3.6]. \square

Remark 5.2. In Prop. 5.1, the case $j = 2$ amounts to an exact sequence

$$0 \longrightarrow \mathbb{Z}/2 \xrightarrow{\delta} H^2(F) \longrightarrow H^2(F(X))$$

where the map δ sends 1 to the Brauer class (D) . This is in fact a special case of Amitsur's theorem (cf. [GS17, Thm. 5.4.1]).

For $j = 3$, the proposition gives a characteristic 2 analogue of a theorem of Arason [Ara75, Satz 5.4], i.e., we have an exact sequence

$$F^* \xrightarrow{\cup(D)} H^3(F) \longrightarrow H^3(F(X)).$$

The kernel of the first map is the group $\text{Nrd}(D^*)$ of reduced norms of D by [Gil00, p.94, Thm. 6]. So in this case we have an isomorphism $F^*/\text{Nrd}(D^*) \cong \text{Ker}(\eta^3)$.

The following result is slightly different from its counterpart in characteristic $\neq 2$ (cf. [Kah95, p.246, Remarks (4)]).

Proposition 5.3. *Let $X \subseteq \mathbb{P}_F^2$ be a smooth anisotropic conic. Then $\text{Coker}(\eta^2) \cong \mathbb{Z}/2$.*

Proof. By (3.2.5), we can identify η^2 with the natural map $\text{Br}(F)[2] \rightarrow \text{Br}(X)[2]$. Consider the exact sequence

$$0 \rightarrow \text{Pic}(X) \rightarrow \text{Pic}(\overline{X})^{\text{Gal}(\overline{F}/F)} \rightarrow \text{Br}(F) \longrightarrow \text{Ker}(\text{Br}(X) \rightarrow \text{Br}(\overline{X})) \longrightarrow H^1(F, \text{Pic}(\overline{X}))$$

obtained from the Hochschild–Serre spectral sequence

$$H^p(F, H^q(\overline{X}, \mathbb{G}_m)) \implies H^{p+q}(X, \mathbb{G}_m).$$

Note that $\text{Br}(\overline{X}) = \text{Br}(\mathbb{P}_{\overline{F}}^1) = 0$, and $H^1(F, \text{Pic}(\overline{X})) = 0$ since $\text{Pic}(\overline{X}) = \mathbb{Z}$ as a Galois module. The conic X being anisotropic, the natural map $\text{Pic}(X) \rightarrow \text{Pic}(\overline{X})^{\text{Gal}(\overline{F}/F)}$ can be identified with the inclusion $2\mathbb{Z} \hookrightarrow \mathbb{Z}$. So the above exact sequence gives an exact sequence

$$0 \longrightarrow \mathbb{Z}/2 \longrightarrow \text{Br}(F) \longrightarrow \text{Br}(X) \longrightarrow 0.$$

The Brauer group $\text{Br}(F)$ is 2-divisible since $\text{char}(F) = 2$ (cf. (3.10.1)). Thus, applying the snake lemma to the diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & \mathbb{Z}/2 & \longrightarrow & \text{Br}(F) & \longrightarrow & \text{Br}(X) \longrightarrow 0 \\ & & \downarrow 0 & & \downarrow \times 2 & & \downarrow \times 2 \\ 0 & \longrightarrow & \mathbb{Z}/2 & \longrightarrow & \text{Br}(F) & \longrightarrow & \text{Br}(X) \longrightarrow 0 \end{array}$$

we obtain an exact sequence

$$0 \longrightarrow \mathbb{Z}/2 \longrightarrow \mathrm{Br}(F)[2] \xrightarrow{\eta^2} \mathrm{Br}(X)[2] \longrightarrow \mathbb{Z}/2 \longrightarrow 0$$

proving $\mathrm{Coker}(\eta^2) \cong \mathbb{Z}/2$ as claimed. \square

(5.4) Let $X \subseteq \mathbb{P}_F^2$ be a smooth conic. Fix $i \in \mathbb{N}$. In the Hochschild–Serre spectral sequence $H^p(F, H^q(\overline{X}, i)) \Rightarrow H^{p+q}(X, i)$ we have

$$H^q(\overline{X}, i) = H^q(\mathbb{P}_F^1, i) = \begin{cases} H^{i-1}(\overline{F}, i-1) = K_{i-1}^M(\overline{F})/2 & \text{if } q = i+1, \\ H^i(\overline{F}, i) = K_i^M(\overline{F})/2 & \text{if } q = i, \\ 0 & \text{otherwise} \end{cases}$$

by [Gro85, (2.1.15)] (and the Bloch–Kato–Gabber theorem). This combined with (3.9.2), (3.9.3), (3.9.5) and (3.9.6) yields isomorphisms

$$\begin{aligned} H^i(F, i) &= H^0(F, H^i(\overline{X}, i)) \cong H^i(X, i), \\ H^i(F) &= H^1(F, H^{i+1}(\overline{X})) \cong H^{i+2}(X, i) \end{aligned}$$

and an exact sequence

$$(5.4.1) \quad 0 \longrightarrow H^{i+1}(F) \longrightarrow H^{i+1}(X) \xrightarrow{\rho} H^{i-1}(F, i-1) \longrightarrow 0.$$

Since $\dim X = 1$, (3.6.1) gives a short exact sequence

$$(5.4.2) \quad 0 \longrightarrow H_{Zar}^1(X, \mathcal{H}^i(i)) \xrightarrow{\iota} H^{i+1}(X) \longrightarrow H_{nr}^{i+1}(X) \longrightarrow 0.$$

Also, by the Gersten resolution of $\mathcal{H}^i(i)$ we have

$$H_{Zar}^1(X, \mathcal{H}^i(i)) = \mathrm{Coker} \left(H^i(F(X), i) \longrightarrow \bigoplus_{x \in X^{(1)}} H^{i-1}(\kappa(x), i-1) \right).$$

In particular, there is a natural surjection

$$\bigoplus_{x \in X^{(1)}} H^{i-1}(\kappa(x), i-1) \longrightarrow H_{Zar}^1(X, \mathcal{H}^i(i)).$$

The same arguments as in the proof of [Pey95, p.377, Lemma 2.1] show that the diagram

$$(5.4.3) \quad \begin{array}{ccc} \bigoplus_{x \in X^{(1)}} H^{i-1}(\kappa(x), i-1) & \longrightarrow & H_{Zar}^1(X, \mathcal{H}^i(i)) \\ \oplus \mathrm{Cor}_{\kappa(x)/F} \downarrow & & \downarrow \iota \\ H^{i-1}(F, i-1) & \xleftarrow{\rho} & H^{i+1}(X) \end{array}$$

is commutative. Let D be the quaternion F -algebra corresponding to the conic X . For each $x \in X^{(1)}$ we have $(D)_{\kappa(x)} = 0$ in $H^2(\kappa(x)) = \mathrm{Br}(\kappa(x))[2]$. Therefore,

$$\mathrm{Cor}_{\kappa(x)/F}(\alpha) \cup (D) = \mathrm{Cor}_{\kappa(x)/F}(\alpha \cup (D)_{\kappa(x)}) = 0 \quad \text{for all } \alpha \in H^{i-1}(\kappa(x), i-1).$$

This together with the commutative diagram (5.4.3) shows that the composite map

$$H_{Zar}^1(X, \mathcal{H}^i(i)) \xrightarrow{\iota} H^{i+1}(X) \xrightarrow{\rho} H^{i-1}(F, i-1) \xrightarrow{\cup(D)} H^{i+1}(F)$$

is 0. Now, using (5.4.2) we can define

$$(5.4.4) \quad N : H_{nr}^{i+1}(X) \longrightarrow H^{i+1}(F)$$

to be the unique homomorphism making the following diagram commute:

$$\begin{array}{ccc} H^{i+1}(X) & \longrightarrow & H_{nr}^{i+1}(X) \\ \rho \downarrow & & \downarrow N \\ H^{i-1}(F, i-1) & \xrightarrow{\cup(D)} & H^{i+1}(F). \end{array}$$

Note that (5.4.2) also yields a homomorphism

$$\tau : \text{Ker}(\eta^{i+1}) = \text{Ker}(H^{i+1}(F) \rightarrow H^{i+1}(F(X))) \longrightarrow H_{Zar}^1(X, \mathcal{H}^i(i))$$

such that the diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & \text{Ker}(\eta^{i+1}) & \longrightarrow & H^{i+1}(F) & \xrightarrow{\eta^{i+1}} & H_{nr}^{i+1}(X) \\ & & \tau \downarrow & & \downarrow & & \parallel \\ 0 & \longrightarrow & H_{Zar}^1(X, \mathcal{H}^i(i)) & \xrightarrow{\iota} & H^{i+1}(X) & \longrightarrow & H_{nr}^{i+1}(X) \longrightarrow 0 \end{array}$$

is commutative with exact rows.

Now we have the following complex

$$(5.4.5) \quad \begin{array}{ccccccc} 0 & \longrightarrow & \text{Ker}(\eta^{i+1}) & \xrightarrow{\tau} & H_{Zar}^1(X, \mathcal{H}^i(i)) & \xrightarrow{\rho \circ \iota} & H^{i-1}(F, i-1) \\ & & & & \xrightarrow{\cup(D)} & H^{i+1}(F) & \xrightarrow{\eta^{i+1}} & H_{nr}^{i+1}(X) & \xrightarrow{N'} & H^{i-1}(F, i-1) \cup (D) \longrightarrow 0 \end{array}$$

where

$$H^{i-1}(F, i-1) \cup (D) := \text{Im} \left(H^{i-1}(F, i-1) \xrightarrow{\cup(D)} H^{i+1}(F) \right)$$

and N' is induced by the map N in (5.4.4).

We can now prove a characteristic 2 counterpart of [Pey95, p.379, Prop. 2.2].

Proposition 5.5. *The complex (5.4.5) is exact except possibly at the third term $H^{i-1}(F, i-1)$ and the fifth term $H_{nr}^{i+1}(X)$.*

Moreover,

$$\frac{\text{Ker}(N')}{\text{Im}(\eta^{i+1})} \cong \frac{\text{Ker}(\cup(D))}{\text{Im}(\rho \circ \iota)} = \frac{\text{Ker}(\cup(D))}{\text{Im} \left(\oplus \text{Cor}_{\kappa(x)/F} : \bigoplus_{x \in X^{(1)}} H^{i-1}(\kappa(x), i-1) \rightarrow H^{i-1}(F, i-1) \right)}$$

If $i \leq 2$, then (5.4.5) is exact everywhere.

Proof. The injectivity of τ is a consequence of the injectivity of $H^{i+1}(F) \rightarrow H^{i+1}(X)$, and the surjectivity of N' follows from that of the map ρ (cf. (5.4.1)). The equality $\text{Im}(\tau) = \text{Ker}(\rho \circ \iota)$ can be easily shown by a diagram chase, and the equality $\text{Im}(\cup(D)) = \text{Ker}(\eta^{i+1})$ was Prop. 5.1.

To get the isomorphism $\frac{\text{Ker}(N')}{\text{Im}(\eta^{i+1})} \cong \frac{\text{Ker}(\cup(D))}{\text{Im}(\rho \circ \iota)}$, it suffices to apply the snake lemma to the following commutative diagram

$$\begin{array}{ccccccccc} 0 & \longrightarrow & H_{Zar}^1(X, \mathcal{H}^i(i)) & \xrightarrow{\iota} & H^{i+1}(X) & \longrightarrow & H_{nr}^{i+1}(X) & \longrightarrow & 0 \\ & & \downarrow (\rho \circ \iota)' & & \downarrow \rho & & \downarrow N' & & \\ 0 & \longrightarrow & \text{Ker}(\cup(D)) & \longrightarrow & H^{i-1}(F, i-1) & \xrightarrow{\cup(D)} & H^{i-1}(F, i-1) \cup (D) & \longrightarrow & 0. \end{array}$$

From the diagram (5.4.3) we find

$$\text{Im}(\rho \circ \iota) = \text{Im} \left(\bigoplus \text{Cor}_{\kappa(x)/F} : \bigoplus_{x \in X^{(1)}} H^{i-1}(\kappa(x), i-1) \rightarrow H^{i-1}(F, i-1) \right).$$

If $i = 0$, then $H^{i-1}(F, i-1) = 0$. If $i = 1$, then

$$\text{Im}(\rho \circ \iota) = \text{Im}(\text{deg} : \text{CH}^1(X)/2 \rightarrow \mathbb{Z}/2)$$

and it is equal to $\text{Ker}(\mathbb{Z}/2 \xrightarrow{\cup(D)} H^2(F))$. If $i = 2$, the equality $\text{Im}(\rho \circ \iota) = \text{Ker}(\cup(D))$ follows from [Gil00, p.94, Thm. 6]. \square

Corollary 5.6. *Let $X \subseteq \mathbb{P}_F^2$ be a smooth conic with associated quaternion algebra D . Then we have isomorphisms*

$$\text{Coker}(\eta^3) \cong H^1(F, 1) \cup (D) \cong F^*/\text{Nrd}(D^*) \cong \text{Ker}(\eta^3).$$

Proof. The first isomorphism follows from the $i = 2$ case of Prop. 5.5. The other isomorphisms have been discussed in Remark 5.2. \square

Corollary 5.7. *Suppose φ is the reduced norm form of a quaternion division algebra D . Then for the quadric $X = X_\varphi$ we have isomorphisms*

$$\begin{aligned} \text{Ker}(\eta^j) &= H^{j-2}(F, j-2) \cup (D), \quad \text{for all } j \geq 2, \\ \text{Ker}(\eta^2) &\cong \text{Coker}(\eta^2) \cong \mathbb{Z}/2, \\ \text{Ker}(\eta^3) &\cong \text{Coker}(\eta^3) \cong H^1(F, 1) \cup (D) \cong F^*/\text{Nrd}(D^*). \end{aligned}$$

Proof. This follows from Cor. 4.2 and the corresponding results for conics (Props. 5.2, 5.3 and Cor. 5.6). \square

Remark 5.8. Suppose $3 \leq \dim \varphi \leq 4$. Then the cycle class map with divisible coefficients

$$\text{cl}_\infty^2 : \text{CH}^2(X) \otimes (\mathbb{Q}_2/\mathbb{Z}_2) \rightarrow H^4(X, \mathbb{Q}_2/\mathbb{Z}_2(2))$$

is injective and the map $\eta_\infty^3 : H^3(F, \mathbb{Q}_2/\mathbb{Z}_2(2)) \rightarrow H_{nr}^3(X, \mathbb{Q}_2/\mathbb{Z}_2(2))$ is surjective.

By Prop. 4.4, if one of the two groups $\text{Ker}(\text{cl}_\infty^2)$ and $\text{Coker}(\eta_\infty^3)$ is trivial, then so is the other. If $\dim \varphi = 3$, then $\dim X = 1$ and $\text{CH}^2(X) = 0$, so trivially $\text{Ker}(\text{cl}_\infty^2) = 0$. If $\dim \varphi = 4$, as in the proof of Cor. 5.7 we may reduce to the case where φ is a 2-Pfister form. Then we can apply Cor. 4.2 to obtain $\text{Coker}(\eta_\infty^3) = 0$ by passing to the case of conics.

As a corollary, we have an isomorphism $\text{Ker}(\eta^3) \cong \text{Coker}(\eta^3)$ in view of Lemma 4.3. In fact, it is shown in [HLS21, Thm. 5.6] that

$$\text{Ker}(\eta^3) = \{0\} \cup \{(a) \cup (b) \cup (c) \mid a, b, c \in F^*, \varphi \text{ is similar to a subform of } \langle\langle a, b; c \rangle\rangle\}.$$

6 Quadrics of dimension ≥ 2

In this section, we prove our main results about low degree unramified cohomology for quadrics of dimension at least 2. These extend the main theorems of [Kah95] to characteristic 2.

Throughout this section, let φ be a nondegenerate quadratic form of dimension ≥ 3 over the field F and let $X = X_\varphi$ be the projective quadric defined by φ .

Lemma 6.1. *For each $i \in \mathbb{N}$, let $\xi^i : \text{CH}^i(X)/2 \rightarrow H^0(F, \text{CH}^i(\overline{X})/2)$ be the canonical map. Assume φ is anisotropic.*

1. *If $\dim \varphi = 4$ and $e_1(\varphi) = 0$, then $\text{Ker}(\xi^1) \cong \text{Coker}(\xi^1) \cong \mathbb{Z}/2$.*

Otherwise ξ^1 is an isomorphism.

2. *Suppose $4 \leq \dim \varphi \leq 5$. Then $\xi^2 = 0$, $\text{Coker}(\xi^2) \cong \mathbb{Z}/2$ and*

$$\text{Ker}(\xi^2) \cong \begin{cases} (\mathbb{Z}/2)^2 & \text{if } \dim \varphi = 5 \text{ and } \varphi \text{ is a Pfister neighbor,} \\ \mathbb{Z}/2 & \text{otherwise.} \end{cases}$$

3. *Suppose $\dim \varphi \geq 6$.*

(a) *If φ is an Albert form, then $\text{Ker}(\xi^2) \cong \text{Coker}(\xi^2) \cong \mathbb{Z}/2$.*

(b) *If φ is a neighbor of a 3-Pfister form, then $\text{Ker}(\xi^2) \cong \mathbb{Z}/2$ and $\text{Coker}(\xi^2) = 0$.*

(c) *In all the other cases, ξ^2 is an isomorphism.*

Proof. (1) We use Prop. 2.4. If $\dim \varphi = 4$ and $e_1(\varphi) = 0$, then ξ^1 can be identified with the map

$$(2\mathbb{Z}/4\mathbb{Z}).\ell_1 \oplus (\mathbb{Z}/2\mathbb{Z}).h \xrightarrow{(0, \text{Id})} (\mathbb{Z}/2\mathbb{Z}).\ell_1 \oplus (\mathbb{Z}/2\mathbb{Z}).h.$$

If $\dim \varphi = 4$ and $e_1(\varphi) \neq 0$, then the Galois action on $\text{CH}^1(\overline{X}) = \mathbb{Z}.h \oplus \mathbb{Z}.\ell = \mathbb{Z}.\ell'_1 \oplus \mathbb{Z}.\ell_1$ permutes the two line classes ℓ_1 and ℓ'_1 (Prop. 2.3). Thus ξ^1 can be identified with the natural isomorphism

$$(\mathbb{Z}/2\mathbb{Z}).h \xrightarrow{\sim} (\mathbb{Z}/2\mathbb{Z}).h = (\mathbb{Z}/2\mathbb{Z}).(\ell'_1 + \ell_1).$$

If $\dim \varphi > 4$, then $\mathrm{CH}^1(X) = \mathrm{CH}^1(\overline{X}) = \mathbb{Z}.h$ and ξ^1 is clearly the identity map on $(\mathbb{Z}/2\mathbb{Z}).h$.

(2) If $\dim \varphi = 4$, then $\mathrm{CH}^2(X) = \mathbb{Z}.2\ell_0 \subseteq \mathrm{CH}^2(\overline{X}) = \mathrm{CH}^2(\overline{X})^{\mathrm{Gal}(\overline{F}/F)} = \mathbb{Z}.\ell_0$, and ξ^2 is the zero map from $2\mathbb{Z}/4\mathbb{Z}$ to $\mathbb{Z}/2\mathbb{Z}$.

Now suppose $\dim \varphi = 5$. We have

$$\mathrm{CH}^2(X) = \mathrm{CH}^2(X)_{\mathrm{tors}} \oplus \mathbb{Z}.2\ell_1 \quad \text{and} \quad \mathrm{CH}^2(\overline{X}) = \mathrm{CH}^2(\overline{X})^{\mathrm{Gal}(\overline{F}/F)} = \mathbb{Z}.\ell_1 .$$

The map ξ^2 is thus therefore the zero map. By [Kar90, (5.3)] (see [HLS21, Thm. 5.1] in characteristic 2), $\mathrm{CH}^2(X)_{\mathrm{tors}} \cong \mathbb{Z}/2$ if φ is a Pfister neighbor, otherwise $\mathrm{CH}^2(X)_{\mathrm{tors}} = 0$. In the former case, $\mathrm{CH}^2(X)/2 \cong (\mathbb{Z}/2)^2$ and $H^0(F, \mathrm{CH}^2(\overline{X})/2) = \mathbb{Z}/2$. In the other case, $\mathrm{CH}^2(X)/2 \cong \mathbb{Z}/2 \cong H^0(F, \mathrm{CH}^2(\overline{X})/2)$.

(3) We will use the results in [Kar90, (5.5)] (see also [HLS21, Thm. 5.2]) if $\dim \varphi = 6$ and [HLS21, Thm. 5.3] if $\dim \varphi \geq 7$.

If φ is an Albert form, we have

$$\mathrm{CH}^2(X) = \mathbb{Z}.h^2 \oplus \mathbb{Z}.4\ell_2 \subseteq \mathrm{CH}^2(\overline{X}) = \mathrm{CH}^2(\overline{X})^{\mathrm{Gal}(\overline{F}/F)} = \mathbb{Z}.h^2 \oplus \mathbb{Z}.\ell_2 .$$

In this case, ξ^2 can be identified with the natural map

$$(\mathbb{Z}/2\mathbb{Z}).h^2 \oplus (4\mathbb{Z}/8\mathbb{Z}).\ell_2 \xrightarrow{(\mathrm{Id}, 0)} (\mathbb{Z}/2\mathbb{Z}).h^2 \oplus (\mathbb{Z}/2\mathbb{Z}).\ell_2 .$$

If φ is a 6-dimensional Pfister neighbor, we have

$$\mathrm{CH}^2(X) = \mathrm{CH}^2(X)_{\mathrm{tors}} \oplus \mathbb{Z}.\ell'_2 \oplus \mathbb{Z}.\ell_2 \quad \text{and} \quad \mathrm{CH}^2(\overline{X}) = \mathbb{Z}.\ell'^2 \oplus \mathbb{Z}.\ell_2 .$$

The Galois action on $\mathrm{CH}^2(\overline{X})$ permutes ℓ_2 and ℓ'_2 . In this case, $H^0(F, \mathrm{CH}^2(\overline{X})/2) = (\mathbb{Z}/2\mathbb{Z}).(\ell'_2 + \ell_2)$ and ξ^2 can be identified with the map

$$\mathbb{Z}/2\mathbb{Z} \oplus (\mathbb{Z}/2\mathbb{Z}).(\ell'_2 + \ell_2) \xrightarrow{0 \oplus \mathrm{Id}} (\mathbb{Z}/2\mathbb{Z}).(\ell'_2 + \ell_2) .$$

If φ is a 7 or 8 dimensional Pfister neighbor, then

$$\mathrm{CH}^2(X) = \mathrm{CH}^2(X)_{\mathrm{tors}} \oplus \mathbb{Z}.h^2 \cong \mathbb{Z}/2\mathbb{Z} \oplus \mathbb{Z} \quad \text{and} \quad \mathrm{CH}^2(\overline{X}) = \mathrm{CH}^2(\overline{X})^{\mathrm{Gal}(\overline{F}/F)} = \mathbb{Z}.h^2 .$$

The map ξ^2 is the identity on $(\mathbb{Z}/2\mathbb{Z}).h^2$ and 0 on $\mathrm{CH}^2(X)_{\mathrm{tors}}$.

If $\dim \varphi = 6$ and φ is neither an Albert form nor a Pfister neighbor, we have

$$\mathrm{CH}^2(X) = \mathbb{Z}.h^2 = \mathbb{Z}.\ell'_2 \oplus \mathbb{Z}.\ell_2 = \mathrm{CH}^2(\overline{X})^{\mathrm{Gal}(\overline{F}/F)} \subseteq \mathrm{CH}^2(\overline{X}) = \mathbb{Z}.\ell'_2 \oplus \mathbb{Z}.\ell_2 .$$

Now the map ξ^2 is the identity map on $(\mathbb{Z}/2\mathbb{Z}).(\ell'_2 + \ell_2) = H^0(F, \mathrm{CH}^2(\overline{X})/2)$.

In all the remaining case, we have $\mathrm{CH}^2(X) = \mathbb{Z}.h^2 = \mathrm{CH}^2(\overline{X})$ and ξ^2 is the identity on $(\mathbb{Z}/2).h$. \square

Recall that we have defined the cycle class maps in (3.6.2). Here we only need the mod 2 case.

Corollary 6.2. *The cycle class map $\mathrm{cl}_X^2 : \mathrm{CH}^2(X)/2 \rightarrow H^4(X, 2)$ is injective in the following cases:*

1. $\dim \varphi = 6$ and φ is neither an Albert form nor a Pfister neighbor.
2. $7 \leq \dim \varphi \leq 8$ and φ is not a Pfister neighbor.
3. $\dim \varphi > 8$.

Proof. We have the following commutative diagram

$$\begin{array}{ccc} \mathrm{CH}^2(X)/2 & \xrightarrow{\mathrm{cl}_X^2} & H^4(X, 2) \\ \downarrow & & \downarrow \\ \mathrm{CH}^2(\overline{X})/2 & \xrightarrow{\mathrm{cl}_{\overline{X}}^2} & H^4(\overline{X}, 2) \end{array}$$

The left vertical map is injective, by Lemma 6.1 (3). We have seen in Prop. 3.8 that the cycle class map $\mathrm{cl}_{\overline{X}}^2$ for \overline{X} is injective. So the above diagram yields the desired result. \square

The next result is a characteristic two version of [Kah95, Lemma 2].

Lemma 6.3. *Suppose $X = X_\varphi$ is defined by the reduced norm form φ of a quaternion division algebra D over F . Let P be a closed point in X and let $C \subseteq X$ be a smooth hyperplane section.*

Then the cycle class map $\mathrm{cl}_X^2 : \mathrm{CH}^2(X)/2 \rightarrow H^4(X, 2)$ sends the class $[P] \in \mathrm{CH}^2(X)/2$ to the cup product $(D) \cup \mathrm{cl}_X^1(C) \in H^4(X, 2)$.

Proof. We may assume $P \in C$ and consider its image under the map $\mathrm{cl}_C^1 : \mathrm{CH}^1(C)/2 \rightarrow H^2(C, 1)$. For every field extension K/F such that C has a K -rational point, the natural map $\mathrm{CH}^1(C)/2 \rightarrow \mathrm{CH}^1(C_K)/2$ is the zero map. So we have $\mathrm{cl}_C^1(P) \in \mathrm{Ker}(H^2(C) \rightarrow H^2(C_K))$. Taking $K = F(C)$ and $K = \overline{F}$ respectively we find

$$(6.3.1) \quad \mathrm{cl}_C^1(P) \in \mathrm{Ker}(H^2(C) \rightarrow H^2(F(C))) \cap \mathrm{Ker}(H^2(C) \rightarrow H^2(\overline{C})) .$$

On the other hand, from the Hochschild–Serre spectral sequence we see that $H^2(F) = \mathrm{Ker}(H^2(C) \rightarrow H^2(\overline{C}))$ (cf. (3.9.6)). This together with (6.3.1) shows

$$\mathrm{cl}_C^1(P) \in \mathrm{Ker}(H^2(F) \rightarrow H^2(F(C))) = \{0, (D)\} .$$

Since the map cl_C^1 is injective (Prop. 3.7), we have $\mathrm{cl}_C^1(P) = (D) \in H^2(F) \subseteq H^2(C)$. In particular, $\mathrm{cl}^1(P)$ is the pullback of the class $(D) \in H^2(X)$.

Now let $i_* : H^2(C, 1) \rightarrow H^4(X, 2)$ be the composition of the Gysin map $H^2(C, 1) \rightarrow H_C^4(X, 2)$ and the canonical map $H_C^4(X, 2) \rightarrow H^4(X, 2)$. By [Gro85, Chap. II, Prop. 4.2.3], we have the commutative diagram

$$\begin{array}{ccc} \mathrm{CH}^1(C)/2 & \longrightarrow & \mathrm{CH}^2(X)/2 \\ \mathrm{cl}_C^1 \downarrow & & \downarrow \mathrm{cl}_X^2 \\ H^2(C, 1) & \xrightarrow{i_*} & H^4(X, 2) \end{array}$$

Therefore,

$$\mathrm{cl}_X^2(P) = i_* \mathrm{cl}_C^1(P) = i_*(D) = i_*(1 \cdot i^*(D)),$$

where $1 \in \mathbb{Z}/2 = H^0(C, 0)$ and $i^* : H^2(X, 1) \rightarrow H^2(C, 1)$ is the pullback map. By the projection formula for the Gysin maps (cf. [Gro85, Chap. II, Cor. 2.2.5]) we get

$$(6.3.2) \quad \mathrm{cl}_X^2(P) = i_*(1 \cdot i^*(D)) = i'_*(1) \cup (D) \in H^4(X, 2).$$

Here $i'_* : H^0(C, 0) \rightarrow H^2(X, 1)$ is defined in the same way as i_* . That is, i'_* is the composite map

$$H^0(C, 0) \xrightarrow{\mathrm{Gysin}} H_C^2(X, 1) \longrightarrow H^2(X, 1).$$

From the commutative diagram

$$\begin{array}{ccc} \mathbb{Z}/2 = \mathrm{CH}^0(C)/2 & \xrightarrow{1 \rightarrow [C]} & \mathrm{CH}^1(X)/2 \\ \cong \downarrow & & \downarrow \mathrm{cl}_X^1 \\ \mathbb{Z}/2 = H^0(C, 0) & \xrightarrow{i'_*} & H^2(X, 1) \end{array}$$

it follows that $i'_*(1) = \mathrm{cl}_X^1(C) \in H^2(X, 1)$. In view of (6.3.2) we have thus proved the lemma. \square

Lemma 6.4. *Let $X = X_\varphi$ be as in Lemma 6.3. Then we have natural isomorphisms*

$$H^2(F, H^2(\overline{X}, 1) \otimes \mathbb{Z}/2(1)) \cong H^2(F) \oplus H^2(F) \cong H^1(F, H^3(\overline{X}, 2)).$$

Proof. First, we have seen in (3.9.8) that $H^2(\overline{X}, 1) \cong \mathrm{CH}^1(\overline{X})/2$. The Galois action on $\mathrm{CH}^1(\overline{X})/2$ being trivial (by Prop. 2.3), the first isomorphism follows.

To obtain the second isomorphism, note that

$$H^3(\overline{X}, 2) \cong K_1(\overline{F})/2 \otimes \mathrm{CH}^1(\overline{X})/2 \cong (K_1(\overline{F})/2)^2$$

by (3.9.12). Applying the second formula in (3.9.5) with $i = 1$, we get

$$H^1(F, H^3(\overline{X}, 2)) \cong H^1(F, K_1(\overline{F})/2)^2 \cong H^2(F) \oplus H^2(F).$$

This completes the proof. \square

Proposition 6.5. *The cycle class map $\mathrm{cl}_X^2 : \mathrm{CH}^2(X)/2 \rightarrow H^4(X, 2)$ is injective in the following two cases:*

1. $\dim \varphi = 4$.
2. $\dim \varphi = 5$ and φ is not a Pfister neighbor.

Proof. First we show that the second case follows from the first one. Indeed, $\mathrm{CH}^2(X) \cong \mathbb{Z} \cdot h^2$ by [Kar90, (5.3)] (or [HLS21, Thm. 5.1]), and if $Y \subseteq X$ is a smooth hyperplane section, the generator h^2 of $\mathrm{CH}^2(X)$ restricts to the generators h_Y^2 of $\mathrm{CH}^2(Y)$. So the

natural pullback map $i^* : \text{CH}^2(X)/2 \rightarrow \text{CH}^2(Y)/2$ is an isomorphism. The commutative diagram

$$\begin{array}{ccc} \text{CH}^2(X)/2 & \xrightarrow{i^*} & \text{CH}^2(Y)/2 \\ \text{cl}_X^2 \downarrow & & \downarrow \text{cl}_Y^2 \\ H^4(X, 2) & \xrightarrow{i^*} & H^4(Y, 2) \end{array}$$

shows that the injectivity of cl_X^2 follows from that of cl_Y^2 .

Now we can assume $\dim \varphi = 4$. If φ is isotropic, then the natural map $\text{CH}^2(X) \rightarrow \text{CH}^2(\overline{X})$ is an isomorphism. Since $\text{cl}_{\overline{X}}^2$ is injective by Prop. 3.8, it follows that cl_X^2 is also injective. We may thus assume φ is anisotropic.

If $e_1(\varphi)$ is nonzero in $H^1(F)$, it correspond to a separable quadratic extension K/F . We claim that φ_K remains anisotropic. Indeed, since $e_1(\varphi_K) = 0$, if φ_K is isotropic, then it is hyperbolic. By [EKM08, (34.8)], we would have $\varphi = \langle a, b \rangle \otimes N_{K/F}$ for some $a, b \in F^*$. But this contradicts the assumption $e_1(\varphi) \neq 0$. Our claim is thus proved. It follows that $\text{CH}^2(X) \rightarrow \text{CH}^2(X_K)$ is an isomorphism. Passing to the extension K/F if necessary, we may assume $e_1(\varphi) = 0$. Then we may further assume that φ is the reduced norm form of a quaternion division algebra D over F .

By Lemma 6.3, the generator of $\text{CH}^2(X)/2 \cong \mathbb{Z}/2$ is mapped to the cup product $(D) \cup \text{cl}_X^1(h) \in H^4(X, 2)$ by the cycle class map cl_X^2 . So it suffices to show $(D) \cup \text{cl}_X^1(h) \neq 0$ in $H^4(X, 2)$.

Taking $j = 3$ and $i = 2$ in (3.9.3) we obtain an exact sequence

$$0 \longrightarrow H^1(F, H^3(\overline{X}, 2)) \longrightarrow H^4(X, 2) \longrightarrow H^0(F, H^4(\overline{X}, 2)) \longrightarrow 0.$$

Thus,

$$(D) \cup \text{cl}_X^1(h) \in \text{Ker}(H^4(X, 2) \longrightarrow H^4(\overline{X}, 2)) = H^1(F, H^3(\overline{X}, 2)).$$

By Lemma 6.4, we have $H^1(F, H^3(\overline{X}, 2)) = H^2(F, H^2(\overline{X}) \otimes \mathbb{Z}/2(1))$. Therefore, we need only to show $(D) \cup \text{cl}_X^1(h)$ is nonzero in $H^2(F, H^2(\overline{X}) \otimes \mathbb{Z}/2(1))$.

Consider the commutative diagram

$$\begin{array}{ccc} \text{CH}^1(X)/2 & \longrightarrow & \text{CH}^1(\overline{X})/2 \\ \text{cl}_X^1 \downarrow & & \downarrow \text{cl}_{\overline{X}}^1 \\ H^2(X, 1) & \longrightarrow & H^2(\overline{X}, 1) \end{array}$$

The top horizontal map can be identified with the map

$$(0, \text{Id}) : (2\mathbb{Z}/4\mathbb{Z}).\ell_1 \oplus (\mathbb{Z}/2).h \longrightarrow (\mathbb{Z}/2).\ell_1 \oplus (\mathbb{Z}/2).h.$$

The map $\text{cl}_{\overline{X}}^1$ being an isomorphism, it follows that $\text{cl}_X^1(h) \neq 0$ even in $H^2(\overline{X})$. Now the map

$$\mathbb{Z}/2 \longrightarrow H^2(\overline{X}) ; \quad 1 \longmapsto \text{cl}_X^1(h)$$

can be identified with the map

$$\iota : \mathbb{Z}/2 \longrightarrow \mathbb{Z}/2 \oplus \mathbb{Z}/2 = \text{CH}^1(\overline{X})/2 ; \quad 1 \longmapsto (0, 1) = h.$$

We have the commutative diagram

$$(6.5.1) \quad \begin{array}{ccc} \mathbb{Z}/2 = H^0(F, 0) & \xrightarrow{1 \mapsto h} & H^0(F, \text{CH}^1(\overline{X})/2) \\ (D) \cup \cdot \downarrow & & \downarrow (D) \cup \text{cl}_X^1(\cdot) \\ H^2(F, 1) & \xrightarrow{\iota_*} & H^2(F, H^2(\overline{X}) \otimes \mathbb{Z}/2(1)) \end{array}$$

The left vertical map in (6.5.1) is injective since $(D) \neq 0$ in $H^2(F)$. The bottom horizontal map ι_* in (6.5.1) can be identified with the map

$$H^2(F) \longrightarrow H^2(F) \oplus H^2(F); \quad \alpha \longmapsto (0, \alpha)$$

via the first isomorphism given in Lemma 6.3. Hence ι_* is injective. Thus, the diagram (6.5.1) implies that $(D) \cup \text{cl}_X^1(h) \neq 0$ in $H^2(F, H^2(\overline{X}) \otimes \mathbb{Z}/2(1))$. As we have said before, this completes the proof. \square

Lemma 6.6 (Compare [Kah95, Lemma 3]).

1. If either $\dim \varphi > 4$, or $\dim \varphi = 4$ and $e_1(\varphi) \neq 0$, then the map

$$\text{cl}^1 = \nu^{1,1} : \text{CH}^1(X)/2 = H^0(F, 0) \otimes \text{CH}^1(X)/2 \longrightarrow H^2(X)$$

induces an isomorphism

$$H^2(F) \oplus (\text{CH}^1(X)/2) \xrightarrow{\sim} H^2(X).$$

2. If $\dim \varphi > 4$, then the map

$$\nu^{1,2} : H^1(F, 1) \otimes (\text{CH}^1(X)/2) \longrightarrow H^3(X)$$

(defined in (3.6.3)) induces an isomorphism

$$H^3(F) \oplus (H^1(F, 1) \otimes \text{CH}^1(X)/2) \xrightarrow{\sim} H^3(X).$$

Proof. (1) We have the following commutative diagram

$$(6.6.1) \quad \begin{array}{ccccccc} & & \text{CH}^1(X) & \xlongequal{\quad} & H^0(F, 0) \otimes \text{CH}^1(X)/2 & & \\ & & \text{cl}_X^1 \downarrow & & \downarrow & & \\ 0 & \longrightarrow & H^2(F) & \longrightarrow & H^2(X) & \longrightarrow & H^0(F, \text{CH}^1(\overline{X})/2) \longrightarrow 0 \end{array}$$

where the square commutes and the bottom row is the exact sequence given by (3.9.9). The right vertical map in (6.6.1) is an isomorphism by Lemma 6.1 (1). The result follows easily from these observations.

(2) The method of proof is similar. This time we use the following commutative diagram

$$(6.6.2) \quad \begin{array}{ccccccc} & & H^1(F, 1) \otimes \text{CH}^1(X)/2 & \xlongequal{\quad} & H^0(F, K_1(\overline{F})/2) \otimes \text{CH}^1(X)/2 & & \\ & & \nu^{1,2} \downarrow & & \delta \downarrow & & \\ 0 & \longrightarrow & H^3(F) & \longrightarrow & H^3(X) & \longrightarrow & H^0(F, K_1(\overline{F})/2 \otimes \text{CH}^1(\overline{X})/2) \longrightarrow 0 \end{array}$$

Here the bottom row is obtained by combining (3.9.6) with (3.9.12). The right vertical map δ is an isomorphism since $\text{CH}^1(X)/2 \cong H^0(F, \text{CH}^1(\bar{X})/2) = \text{CH}^1(\bar{X})/2 \cong \mathbb{Z}/2$. \square

The following theorem is the characteristic 2 version of [Kah95, Thm. 1]. Notice however that unlike the case of characteristic different from 2, the maps $\mu^{0,j}$ and η^j are different in our situation.

Theorem 6.7. *If $\dim \varphi > 4$, or $\dim \varphi = 4$ and $e_1(\varphi) \neq 0$, then the map $\eta^2 : H^2(F) \rightarrow H_{nr}^2(X)$ is an isomorphism.*

Proof. By Prop. 3.7, we have a natural exact sequence

$$0 \longrightarrow \text{CH}^1(X)/2 \xrightarrow{\text{cl}^1} H^2(X) \longrightarrow H_{nr}^2(X) \longrightarrow 0.$$

This sequence together with Lemma 6.6 (1) proves the theorem immediately. \square

Corollary 6.8. *We have $\text{Coker}(\eta_\infty^2) = 0$ for any smooth projective quadric of dimension ≥ 1 .*

Proof. Combine Lemma 4.3, Prop. 5.3, Cor. 5.7 and Thm. 6.7. \square

Theorem 6.9. *If $\dim \varphi > 4$, then the map*

$$\mu^{1,2} : H^1(F, 1) \otimes (\text{CH}^1(X)/2) \rightarrow H_{Zar}^1(X, \mathcal{H}^2(2))$$

is injective and there are isomorphisms

$$\text{Coker}(\mu^{1,2}) \cong \text{Ker}(\eta^3) \quad \text{and} \quad \text{Coker}(\eta^3) \cong \text{Ker}(\text{cl}_X^2).$$

Proof. Note that the map $\nu^{1,2}$ factors through $\mu^{1,2}$ by construction (cf. (3.6.4)). So we have the following commutative diagram

$$(6.9.1) \quad \begin{array}{ccccccc} 0 & \longrightarrow & H^1(F, 1) \otimes (\text{CH}^1(X)/2) & \xlongequal{\quad} & H^1(F, 1) \otimes (\text{CH}^1(X)/2) & \longrightarrow & 0 \longrightarrow 0 \\ & & \mu^{1,2} \downarrow & & \nu^{1,2} \downarrow & & \downarrow \\ 0 & \longrightarrow & H_{Zar}^1(X, \mathcal{H}^2(2)) & \longrightarrow & H^3(X, 2) & \longrightarrow & M \longrightarrow 0 \end{array}$$

where the second row is obtained from the exact sequence (3.8.2) and $M := \text{Ker}(H_{nr}^3(X, 2) \rightarrow \text{CH}^2(X)/2)$. Applying the snake lemma to the above diagram and using Lemma 6.6 (2) (and (3.8.2)), we get an exact sequence

$$(6.9.2) \quad 0 \rightarrow H^1(F, 1) \otimes (\text{CH}^1(X)/2) \xrightarrow{\mu^{1,2}} H_{Zar}^1(X, \mathcal{H}^2(2)) \rightarrow H^3(F) \xrightarrow{\eta^3} \text{Ker}(\text{cl}_X^2) \rightarrow 0.$$

The theorem follows immediately from the above sequence. \square

Remark 6.10. About the map $\mu^{1,2}$, the cases not covered by Theorem 6.9 are:

- (a) $\dim \varphi = 3$; (b) $\dim \varphi = 4$ and $e_1(\varphi) = 0$; (c) $\dim \varphi = 4$ and $e_1(\varphi) \neq 0$.

In all these cases, we can still use the diagram (6.9.1) to get an exact sequence

$$0 \longrightarrow \text{Coker}(\mu^{1,2}) \longrightarrow \text{Coker}(\nu^{1,2}) \longrightarrow \text{Ker}(H_{nr}^3(X) \rightarrow \text{CH}^2(X)/2) \longrightarrow 0$$

and an equality $\text{Ker}(\mu^{1,2}) = \text{Ker}(\nu^{1,2})$.

To get further information about $\text{Ker}(\mu^{1,2})$, we need to analyze the right vertical map δ in (6.6.2). If φ is isotropic, then $\text{CH}^1(X)/2 \cong \text{CH}^1(\overline{X})/2$ and δ is an isomorphism. So in this case the exact sequence (6.9.2) remains valid, and we get the same conclusions as in Theorem 6.9.

Now we assume φ is anisotropic. Then we claim

$$\begin{cases} \text{Ker}(\mu^{1,2}) = \text{Ker}(\nu^{1,2}) \cong \text{Nrd}(D^*)/F^{*2}, & \text{in Cases (a) and (b),} \\ \text{Ker}(\mu^{1,2}) = \text{Ker}(\nu^{1,2}) = 0, & \text{in Case (c).} \end{cases}$$

In Case (a), the map δ in (6.6.2) can be identified with the zero map from $H^1(F, 1)$ to itself, since the map $\text{CH}^1(X)/2 \rightarrow \text{CH}^1(\overline{X})/2$ is the zero map from $2\mathbb{Z}/4\mathbb{Z}$ to $\mathbb{Z}/2\mathbb{Z}$. Thus, (6.6.2) yields an exact sequence

$$0 \longrightarrow \text{Ker}(\nu^{1,2}) \longrightarrow H^1(F, 1) \xrightarrow{\cup(D)} H^3(F) \longrightarrow \text{Coker}(\nu^{1,2}) \longrightarrow H^1(F, 1) \longrightarrow 0.$$

In Case (b), we can get an exact sequence of the same form, because the map δ can be viewed as the map

$$(0, \text{Id}) : H^1(F, 1) \oplus H^1(F, 1) \longrightarrow H^1(F, 1) \oplus H^1(F, 1).$$

This proves our claim in Cases (a) and (b).

In Case (c), the Galois action on $K_1(\overline{F})/2 \otimes \text{CH}^1(\overline{X})/2 \cong (K_1(\overline{F})/2)^2$ is given by

$$\sigma.(a, b) = (\sigma(b), \sigma(a)), \quad a, b \in K_1(\overline{F})/2.$$

The map δ in this case can be identified with the map

$$H^0(F, K_1(\overline{F})/2) \longrightarrow H^0(F, K_1(\overline{F})/2) \oplus K_1(\overline{F})/2$$

induced from the diagonal map

$$H^0(F, K_1(\overline{F})/2) \longrightarrow (K_1(\overline{F})/2)^2; \quad x \longmapsto (x, x).$$

Therefore, δ is injective in this case. We conclude that $\text{Ker}(\mu^{1,2}) = \text{Ker}(\nu^{1,2}) = 0$ in Case (c).

Theorem 6.11. *Assume $\dim \varphi > 4$. The maps $\mu^{1,2}$ and η^3 are both isomorphisms in each of the following cases:*

1. φ is isotropic.
2. φ is neither an Albert form nor a Pfister neighbor.
3. $\dim \varphi > 8$.

Proof. We shall use Theorem 6.9. In Case (1) the result follows from Prop. 3.5. In the other cases we have $\text{Coker}(\eta^3) \cong \text{Ker}(\text{cl}_X^2) = 0$ by Cor. 6.2 and Prop. 6.5. The injectivity of η^3 is immediate from [HLS21, Thm. 5.6]. \square

Theorem 6.12. *Suppose that φ is a neighbor of an anisotropic 3-fold Pfister form $\langle\langle a, b; c \rangle\rangle$.*

Then $\text{Ker}(\eta^3)$ is generated by $(a) \cup (b) \cup (c) \in H^3(F)$ (so $\text{Ker}(\eta^3) \cong \mathbb{Z}/2$), and $\text{Coker}(\eta^3) \cong \mathbb{Z}/2$.

Proof. The assertion about $\text{Ker}(\eta^3)$ is immediate from [HLS21, Thm. 5.6]. To prove the other assertion, we may reduce to the 5-dimensional case thanks to Cor. 4.2. We may thus assume $\varphi = \langle a \rangle \perp \langle\langle b; c \rangle\rangle$. Since there is an injection from $\text{Ker}(\eta^3)$ to $\text{Coker}(\eta^3)$ by Lemma 4.3, it suffices to show $|\text{Coker}(\eta^3)| \leq 2$.

Let Y be the projective quadric defined by $\langle\langle b; c \rangle\rangle$. It is a smooth hyperplane section in X . We have the commutative diagram

$$\begin{array}{ccc} \text{CH}^2(X)/2 & \xrightarrow{i^*} & \text{CH}^2(Y)/2 \\ \text{cl}_X^2 \downarrow & & \downarrow \text{cl}_Y^2 \\ H^4(X, 2) & \longrightarrow & H^4(Y, 2) \end{array}$$

The map cl_Y^2 is injective by Prop. 6.5. So the above diagram shows that $\text{Ker}(\text{cl}_X^2) \subseteq \text{Ker}(i^*)$. Here

$$\begin{aligned} \text{CH}^2(X)/2 &= \text{CH}^2(X)_{\text{tors}} \oplus (\mathbb{Z}/2).h^2 = (\mathbb{Z}/2).(h^2 - 2\ell_1) \oplus (\mathbb{Z}/2).2\ell_1, \\ \text{CH}^2(Y)/2 &= (\mathbb{Z}/2).h_Y^2 = (\mathbb{Z}/2).2\ell_0, \end{aligned}$$

and the map i^* sends $h^2 - 2\ell_1$ to 0 and $2\ell_1$ to $2\ell_0 = h_Y^2$. Hence $\text{Ker}(i^*) = (\mathbb{Z}/2).(h^2 - 2\ell_1) = \text{CH}^2(X)_{\text{tors}}$, and by Thm. 6.9 we have

$$|\text{Coker}(\eta^3)| = |\text{Ker}(\text{cl}_X^2)| \leq |\text{Ker}(i^*)| = 2$$

as desired. \square

Now we have computed $\text{Ker}(\eta^3)$ and $\text{Coker}(\eta^3)$ for the quadric $X = X_\varphi$ except in the case where φ is an Albert form. This last case will be treated in §8.

7 Unramified Witt groups in characteristic 2

We need to use residue maps on the Witt group of a discrete valuation field of characteristic 2. We recall some key definitions and facts that will be used in the next section.

Throughout this section, let K be a field extension of F (so $\text{char}(K) = 2$) and let R be the valuation ring of a nontrivial discrete valuation v on K . Let $\pi \in R$ be a uniformizer and let k be the residue field of R .

(7.1) Let $I_q(R) = W_q(R)$ be the Witt group of nonsingular quadratic spaces over R as defined in [Bae78, p.18, (I.4.8)]. It is naturally a subgroup of $I_q(K)$. For $n \geq 2$, let $I_q^n(R)$ be the subgroup of $I_q(R)$ generated by Pfister forms of the following type:

$$\langle\langle a_1, \dots, a_{n-1}; b \rangle\rangle \quad \text{where } a_i \in R^*, b \in R.$$

Put $I_q^1(R) = I_q(R)$. There is a natural homomorphism $I_q^n(R) \rightarrow I_q^n(k)$, $\varphi \mapsto \bar{\varphi}$ for each $n \geq 1$.

Following [Ara18], we define the *tame* (or *tamely ramified*) subgroup of $I_q(K)$ to be the subgroup $I_q(K)_{tr} := W(K) \cdot I_q(R)$. For general $n \geq 1$, we put

$$I_q^n(K)_{tr} := I_q^{n-1}(K) \cdot I_q(R) = I_q^n(R) + \langle 1, \pi \rangle_{\text{bil}} \cdot I_q^{n-1}(R).$$

By [Ara18, Props. 1.1 and 1.2], there is a well defined *residue map*

$$\partial : I_q(K)_{tr} \longrightarrow I_q(k); \quad \varphi_0 + \pi \cdot \varphi_1 \longmapsto \bar{\varphi}_1 \quad \text{for } \varphi_0, \varphi_1 \in I_q(R)$$

and the sequence

$$0 \longrightarrow I_q(R) \longrightarrow I_q(K)_{tr} \xrightarrow{\partial} I_q(k) \longrightarrow 0$$

is exact. For each $n \geq 1$ we have an induced residue map $\partial : I_q^n(K)_{tr} \rightarrow I_q^{n-1}(k)$ (with $I_q^0(k) = I_q^1(k)$ by convention), and putting

$$I_q^n(K)_{nr} := \text{Ker}(\partial : I_q^n(K)_{tr} \rightarrow I_q^{n-1}(k)) = I_q^n(K)_{tr} \cap I_q(R),$$

we get an exact sequence

$$0 \longrightarrow I_q^n(K)_{nr} \longrightarrow I_q^n(K)_{tr} \xrightarrow{\partial} I_q^{n-1}(k) \longrightarrow 0.$$

Note that the term “tame” and the residue map depend on the discrete valuation v (or the valuation ring R).

(7.2) We use shorthand notation for cohomology functors introduced at the beginning of §5.

Given $i \in \mathbb{N}$, localization theory in étale cohomology theory gives rise to a long exact sequence

$$\dots \longrightarrow H^{q+1}(R, i+1) \longrightarrow H^{q+1}(F, i+1) \xrightarrow{\delta} H_{\mathfrak{m}}^{q+2}(R, i+1) \longrightarrow \dots$$

where \mathfrak{m} denotes the unique closed point of $\text{Spec}(R)$. By [Shi07, Cor. 3.4] we have

$$H_{\mathfrak{m}}^{q+2}(R, i+1) = 0 \quad \text{if } q \notin \{i, i+1\}.$$

Moreover, [Shi07, Thm. 3.2] tells us that $H^i(k, i) \xrightarrow{\sim} H_{\mathfrak{m}}^{i+2}(R, i+1)$. So the localization sequence above yields an exact sequence

$$\begin{aligned} 0 &\longrightarrow H^{i+1}(R, i+1) \longrightarrow H^{i+1}(K, i+1) \xrightarrow{\delta_0} H^i(k, i) \longrightarrow \\ &\longrightarrow H^{i+2}(R) \longrightarrow H^{i+2}(K) \xrightarrow{\delta_1} H_{\mathfrak{m}}^{i+3}(R, i+1) \longrightarrow H^{i+3}(R, i+1) \longrightarrow 0. \end{aligned}$$

By the Bloch–Kato–Gabber theorem, the map δ_0 in the above sequence can be identified with the residue map $K_{i+1}^M(K)/2 \rightarrow K_i^M(k)/2$ in Milnor K -theory, which is surjective. Hence, we have an exact sequence

$$0 \longrightarrow H^{i+2}(R) \longrightarrow H^{i+2}(K) \xrightarrow{\delta_1} H_{\mathfrak{m}}^{i+3}(R, i+1) \longrightarrow H^{i+3}(R, i+1) \longrightarrow 0.$$

An element of $H^{i+2}(K)$ is called *unramified* at v if it lies in the subgroup $H^{i+2}(R) = \text{Ker}(\delta_1)$.

For each $n \geq 2$, we define the *tame* (or *tamely ramified*) part of $H^n(K)$ to be the subgroup

$$H_{tr}^n(K) := \text{Ker} \left(H^n(K) \longrightarrow H^n(\hat{K}) \longrightarrow H^n(\hat{K}^{nr}) \right),$$

where \hat{K} denotes the v -adic completion of K and \hat{K}^{nr} is the maximal unramified extension of \hat{K} . Kato constructed (cf. [Kat82a], [Kat86, §1]) a *residue map* $\partial^H : H_{tr}^n(K^h) \rightarrow H^{n-1}(k)$ satisfying

$$(7.2.1) \quad \partial^H((u\pi) \cup (a_1) \cdots (a_{n-2}) \cup (b)) = (\bar{a}_1) \cup \cdots (\bar{a}_{n-2}) \cup (\bar{b}), \quad u, a_i \in R^*, b \in R,$$

such that the sequence

$$0 \longrightarrow H^n(R) \longrightarrow H^n(K)_{tr} \xrightarrow{\partial^H} H^{n-1}(k) \longrightarrow 0$$

is exact. Therefore, an element $\alpha \in H^n(K)$ is unramified if and only if $\alpha \in H^n(K)_{tr}$ and $\partial^H(\alpha) = 0$.

For each $n \geq 2$, using the formula (7.2.1) we can check through calculation that

$$e_n(I_q^n(R)) \subseteq H^n(R), \quad e_n(I_q^n(K)_{tr}) \subseteq H_{tr}^n(K)$$

and that the following diagram with exact rows is commutative:

$$(7.2.2) \quad \begin{array}{ccccccc} 0 & \longrightarrow & I_q^n(K)_{nr} & \longrightarrow & I_q^n(K)_{tr} & \xrightarrow{\partial} & I_q^{n-1}(k) \longrightarrow 0 \\ & & e_n \downarrow & & e_n \downarrow & & \downarrow e_{n-1} \\ 0 & \longrightarrow & H^n(R) & \longrightarrow & H_{tr}^n(K) & \xrightarrow{\partial^H} & H^{n-1}(k) \longrightarrow 0 \end{array}$$

8 Nontrivial unramified class for Albert quadrics

In this section we investigate the case of Albert quadrics and complete our study of the map η^3 .

Recall some standard notation. The hyperbolic plane \mathbb{H} is the binary quadratic form $(x, y) \mapsto xy$. For any nondegenerate form q of dimension ≥ 3 over F , let $F(q)$ denote the function field of the projective quadric defined by q .

Lemma 8.1. *Let q be an Albert form over F which represents 1, and let q_1 be a form over $F(q)$ such that $q_{F(q)} = q_1 \perp \mathbb{H}$.*

If q_1 represents 1, then q must be isotropic over F .

Proof. We may write $q = a[1, b] \perp c[1, d] \perp [1, b + d]$, where $a, c \in F^*$ and $b, d \in F$. Set $q' = q \perp \langle 1 \rangle$ and $q'' = a[1, b] \perp c[1, d] \perp \langle 1 \rangle$. Since

$$[1, b + d] \perp \langle 1 \rangle \cong [0, b + d] \perp \langle 1 \rangle \cong \mathbb{H} \perp \langle 1 \rangle$$

we have $q' \cong \mathbb{H} \perp q''$ over F . Now

$$\mathbb{H} \perp q_1 \perp \langle 1 \rangle = q_{F(q)} \perp \langle 1 \rangle = q'_{F(q)} \cong \mathbb{H} \perp q''_{F(q)}.$$

By Witt cancellation, $q_1 \perp \langle 1 \rangle \cong q''_{F(q)}$. The assumption that q_1 represents 1 implies that $q''_{F(q)}$ is isotropic.

Assume that q is anisotropic. Then q'' is also anisotropic because it is a subform of q . By [Lag02, Thm. 1.2 (2)], q'' must be a neighbor of a 3-Pfister form π . The isotropy of $q''_{F(q)}$ shows that $\pi_{F(q)}$ is hyperbolic. Thus, by [Lag02, Prop. 3.4], q must be a neighbor of π . But this is absurd, because an anisotropic Albert form cannot be a Pfister neighbor (see e.g. [HLS21, Lemma 3.4]). \square

(8.2) Now assume X is defined by an anisotropic Albert form φ over F . Let $K = F(X)$ be its function field. We shall normalize φ so that it represents 1. Thus

$$\varphi = [1, b + d] \perp a.[1, b] \perp c.[1, d] \quad \text{for some } a, b, c, d \in F^*.$$

We have $\varphi = \langle\langle a; b \rangle\rangle - \langle\langle c; d \rangle\rangle \in I_q^2(F)$.

Let φ_1/K be the anisotropic part of φ_K . Since φ is anisotropic of dimension < 8 , the Hauptsatz [EKM08, (23.7)] implies $\varphi \notin I_q^3(F)$. This means that φ has degree 2 (by [EKM08, (40.10)]). By [EKM08, (25.7)], φ_1 is a general 2-Pfister form (and the height of φ is 2). Let τ/K be the quadratic 2-Pfister form similar to φ_1 , say $\varphi_1 = f \cdot \tau$, where $f \in K^*$. We have $\tau - \varphi_1 = \langle 1, f \rangle_{\text{bil}} \cdot \tau \in I_q^3(K)$, so that the cohomology class $e_3(\tau - \varphi_1) \in H^3(K)$ can be defined.

We claim that $e_3(\tau - \varphi_1)$ lies in the unramified cohomology group $H_{nr}^3(X)$. In fact, we can prove the following:

For every discrete valuation v of K that is trivial on F , $e_3(\tau - \varphi_1)$ is unramified at v .

Proof. Let R be the valuation ring of v in $K = F(X)$. By (7.2.2), it is sufficient to show

$$\tau - \varphi_1 \in I_q^3(K)_{nr} = I_q(R) \cap I_q^3(K)_{tr}.$$

First note that the Witt class of φ_1 is equal to that of φ_K . Hence

$$\varphi_1 = \langle\langle a; b \rangle\rangle - \langle\langle c; d \rangle\rangle \in \text{Im}(I_q^2(F) \rightarrow I_q^2(K)) \subseteq I_q^2(R).$$

If f is a unit for v , then $\tau - \varphi_1 = -\langle 1, f^{-1} \rangle_{\text{bil}} \cdot \varphi_1 \in I_q^3(R) \subseteq I_q^3(K)_{nr}$. We may thus assume $\pi := f^{-1}$ is a uniformizer for v . Then

$$\tau - \varphi_1 = -\langle 1, f^{-1} \rangle_{\text{bil}} \cdot \varphi_1 = -\langle 1, \pi \rangle_{\text{bil}} \cdot \varphi_1 \in \langle 1, \pi \rangle_{\text{bil}} \cdot I_q^2(R) \subseteq I_q^3(K)_{tr}.$$

It remains to prove $\tau - \varphi_1 \in I_q(R)$. We already know $\varphi_1 \in I_q^2(R) \subseteq I_q(R)$. So it suffices to show $\tau \in I_q(R)$. This is equivalent to $\tau_{\hat{K}} \in I_q(\hat{R})$, where \hat{R} denotes the completion of R and \hat{K} is the fraction field of \hat{R} (see e.g. [Ara18, p.106]).

Indeed,

$$e_2(\varphi_1) = e_2(\langle\langle a; b \rangle\rangle - \langle\langle c; d \rangle\rangle) \in \text{Im}(H^2(F) \longrightarrow H^2(K)) \subseteq H^2(R).$$

Therefore, the cohomology class $e_2(\varphi_1)$ is unramified at v . Since $e_2(\tau) = e_2(\varphi_1)$ by (2.1.1), $e_2(\tau)$ is also unramified at v . As a 2-Pfister form, τ is the reduced norm form of a quaternion division K -algebra D . The cohomology class $e_2(\tau)$ is the Brauer class of D . Thus, the fact that $e_2(\tau)$ is unramified means that the quaternion algebra D is unramified. Thus, $D_{\hat{K}} \cong (\alpha, \beta]$ for some $\alpha \in \hat{R}^*$, $\beta \in \hat{R}$. It follows that $\tau_{\hat{K}} \cong \langle\langle \alpha; \beta \rangle\rangle$ ([EKM08, (12.5)]), proving that the Witt class of $\tau_{\hat{K}}$ lies in $I_q^2(\hat{R})$. In particular, $\tau_{\hat{R}} \in I_q(\hat{R})$ as desired. \square

Proposition 8.3. *With notation as in (8.2), $e_3(\tau - \varphi_1) \notin \text{Im}(\eta^3 : H^3(F) \rightarrow H_{nr}^3(X))$.*

Proof. Since $e_3 : I_q^3(F) \rightarrow H^3(F)$ is a surjection and $I_q^3(F)$ is additively generated by 3-Pfister forms, every element of $H^3(F)$ is a sum of finitely many symbols (by a *symbol* in $H^3(F)$ we mean the class $e_3(\pi)$ of a 3-fold Pfister form π). We may define the *symbol length* of an element $\beta \in H^3(F)$ to be the smallest positive integer n such that β is a sum of n symbols.

We use induction to show:

For every $n \geq 1$, there is no element $\beta \in H^3(F)$ of symbol length n such that $\beta_{F(X)} = e_3(\tau - \varphi_1)$.

Assume the contrary. Let $\beta \in H^3(F)$ be an element of symbol length n such that $\beta_{F(X)} = e_3(\tau - \varphi_1)$.

First assume $n = 1$, i.e., $\beta = e_3(\pi)$ for some 3-Pfister form π over F . Since $\tau \perp -\varphi_1$ and $\pi_{F(X)}$ are both 3-Pfister forms, from $e_3(\tau - \varphi_1) = e_3(\pi_{F(X)})$ we can conclude that $\tau \perp -\varphi_1 \cong \pi_{F(X)}$, by [Kat82a, p.237, Prop. 3]. Letting $E = F(\pi)$, we get that $\tau - \varphi_1 = 0$ in $I_q^3(E(X))$. Hence $(\varphi_1)_{E(X)} = \tau_{E(X)}$ is a 2-Pfister form. In particular, $(\varphi_1)_{E(X)}$ represents 1. By Lemma 8.1, $\varphi_E = \varphi_{F(\pi)}$ must be isotropic. But this contradicts an index reduction theorem of Merkurjev (cf. [EKM08, (30.9)]).

Now consider the general case and write $\beta = \pi + \delta$, where $\pi \in H^3(F)$ is a symbol and $\delta \in H^3(F)$ has symbol length $n - 1$. Consider again the function field $E = F(\pi)$. As before $\varphi_E = \varphi_{F(\pi)}$ is anisotropic by Merkurjev's index reduction theorem. But

$$\delta_{E(X)} = \beta_{E(X)} = e_3(\tau_{E(X)} - (\varphi_1)_{E(X)}) \in H^3(E(X)).$$

This contradicts the induction hypothesis. \square

Theorem 8.4. *Let X be the projective quadric defined by an anisotropic Albert form over F .*

Then the map $\eta^3 : H^3(F) \rightarrow H_{nr}^3(X)$ is injective and $\text{Coker}(\eta^3) \cong \mathbb{Z}/2$.

Proof. The injectivity of η^3 follows from [HLS21, Thm. 5.6]. We have seen in Prop. 8.3 that $\text{Coker}(\eta^3) \neq 0$. It remains to show $|\text{Coker}(\eta^3)| \leq 2$.

To this end, we use the isomorphism $\text{Coker}(\eta^2) \cong \text{Ker}(\text{cl}_X^2)$ obtained in Thm. 6.9. By functoriality and the injectivity of cl_X^2 , it follows that $\text{Ker}(\text{cl}_X^2)$ is contained in the kernel of the restriction map $\xi^2 : \text{CH}^2(X)/2 \rightarrow \text{CH}^2(\bar{X})/2$. By Lemma 6.1, we have $\text{Ker}(\xi^2) \cong \mathbb{Z}/2$. This completes the proof. \square

Together with Prop. 4.4, the following corollary also describes the kernel of the cycle class map

$$\mathrm{cl}_\infty^2 : \mathrm{CH}^2(X) \otimes (\mathbb{Q}_2/\mathbb{Z}_2) \longrightarrow H^4(X, \mathbb{Q}_2/\mathbb{Z}_2(2)) .$$

Corollary 8.5. *Let φ be a nondegenerate quadratic form of dimension ≥ 3 over F and let $X = X_\varphi$.*

Then the map $\eta_\infty^3 : H^3(F, \mathbb{Q}_2/\mathbb{Z}_2(2)) \rightarrow H_{nr}^3(X, \mathbb{Q}_2/\mathbb{Z}_2(2))$ is surjective unless φ is an anisotropic Albert form. In the latter case $\mathrm{Coker}(\eta_\infty^3) \cong \mathbb{Z}/2$.

Proof. If $\dim \varphi \leq 4$, this has been discussed in Remark 5.8. When $\dim \varphi > 4$, we have already computed $\mathrm{Ker}(\eta^3)$ and $\mathrm{Coker}(\eta^3)$ in Theorems 6.11, 6.12 and 8.4. So it suffices to apply Lemma 4.3. \square

Acknowledgements. We thank Yang Cao, Zhengyao Wu and Yigeng Zhao for help discussions. Yong Hu is supported by a grant from the National Natural Science Foundation of China (Project No. 11801260). Peng Sun is supported by the Fundamental Research Funds for the Central Universities.

References

- [AJ09] Roberto Aravire and Bill Jacob. $H^1(X, \nu)$ of conics and Witt kernels in characteristic 2. In *Quadratic forms—algebra, arithmetic, and geometry*, volume 493 of *Contemp. Math.*, pages 1–19. Amer. Math. Soc., Providence, RI, 2009.
- [Ara75] Jón Kr. Arason. Cohomologische invarianten quadratischer Formen. *J. Algebra*, 36(3):448–491, 1975.
- [Ara18] Jón Kr. Arason. The Witt group of a discretely valued field. *J. Algebra*, 511:102–113, 2018.
- [Bae78] Ricardo Baeza. *Quadratic forms over semilocal rings*. Lecture Notes in Mathematics, Vol. 655. Springer-Verlag, Berlin-New York, 1978.
- [BO74] Spencer Bloch and Arthur Ogus. Gersten’s conjecture and the homology of schemes. *Ann. Sci. École Norm. Sup. (4)*, 7:181–201 (1975), 1974.
- [CT95] J.-L. Colliot-Thélène. Birational invariants, purity and the Gersten conjecture. In *K-theory and algebraic geometry: connections with quadratic forms and division algebras (Santa Barbara, CA, 1992)*, volume 58 of *Proc. Sympos. Pure Math.*, pages 1–64. Amer. Math. Soc., Providence, RI, 1995.
- [CTHK97] Jean-Louis Colliot-Thélène, Raymond T. Hoobler, and Bruno Kahn. The Bloch-Ogus-Gabber theorem. In *Algebraic K-theory (Toronto, ON, 1996)*, volume 16 of *Fields Inst. Commun.*, pages 31–94. Amer. Math. Soc., Providence, RI, 1997.

- [CTO89] Jean-Louis Colliot-Thélène and Manuel Ojanguren. Variétés unirationnelles non rationnelles: au-delà de l'exemple d'Artin et Mumford. *Invent. Math.*, 97(1):141–158, 1989.
- [CTP16] Jean-Louis Colliot-Thélène and Alena Pirutka. Hypersurfaces quartiques de dimension 3: non-rationalité stable. *Ann. Sci. Éc. Norm. Supér. (4)*, 49(2):371–397, 2016.
- [CTS95] J.-L. Colliot-Thélène and R. Sujatha. Unramified Witt groups of real anisotropic quadrics. In *K-theory and algebraic geometry: connections with quadratic forms and division algebras (Santa Barbara, CA, 1992)*, volume 58 of *Proc. Sympos. Pure Math.*, pages 127–147. Amer. Math. Soc., Providence, RI, 1995.
- [EKM08] Richard Elman, Nikita Karpenko, and Alexander Merkurjev. *The algebraic and geometric theory of quadratic forms*, volume 56 of *American Mathematical Society Colloquium Publications*. American Mathematical Society, Providence, RI, 2008.
- [Gil00] Philippe Gille. Invariants cohomologiques de Rost en caractéristique positive. *K-Theory*, 21(1):57–100, 2000.
- [GL00] Thomas Geisser and Marc Levine. The K -theory of fields in characteristic p . *Invent. Math.*, 139(3):459–493, 2000.
- [Gro85] Michel Gros. Classes de Chern et classes de cycles en cohomologie de Hodge-Witt logarithmique. *Mém. Soc. Math. France (N.S.)*, (21):87, 1985.
- [Gro03] A. Grothendieck. *Revêtements étales et groupe fondamental (SGA 1)*. Société Mathématique de France, 2003.
- [GS88] Michel Gros and Noriyuki Suwa. La conjecture de Gersten pour les faisceaux de Hodge-Witt logarithmique. *Duke Math. J.*, 57(2):615–628, 1988.
- [GS17] Philippe Gille and Tamás Szamuely. *Central simple algebras and Galois cohomology*, volume 165 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 2017. Second edition of [MR2266528].
- [HLS21] Yong Hu, Ahmed Laghribi, and Peng Sun. Chow groups of quadrics in characteristic two. arXiv: 2101.0300, 2021.
- [Ill79] Luc Illusie. Complexe de de Rham-Witt et cohomologie cristalline. *Ann. Sci. École Norm. Sup. (4)*, 12(4):501–661, 1979.
- [Izh91] O. Izhboldin. On p -torsion in K_*^M for fields of characteristic p . In *Algebraic K-theory*, volume 4 of *Adv. Soviet Math.*, pages 129–144. Amer. Math. Soc., Providence, RI, 1991.

- [Izh01] Oleg T. Izhboldin. Fields of u -invariant 9. *Ann. of Math. (2)*, 154(3):529–587, 2001.
- [Kah95] Bruno Kahn. Lower \mathcal{H} -cohomology of higher-dimensional quadrics. *Arch. Math. (Basel)*, 65(3):244–250, 1995.
- [Kah99] Bruno Kahn. Motivic cohomology of smooth geometrically cellular varieties. In *Algebraic K-theory (Seattle, WA, 1997)*, volume 67 of *Proc. Sympos. Pure Math.*, pages 149–174. Amer. Math. Soc., Providence, RI, 1999.
- [Kah04] Bruno Kahn. Cohomologie non ramifiée des quadriques. In *Geometric methods in the algebraic theory of quadratic forms*, volume 1835 of *Lecture Notes in Math.*, pages 1–23. Springer, Berlin, 2004.
- [Kar90] N. A. Karpenko. Algebro-geometric invariants of quadratic forms. *Algebra i Analiz*, 2(1):141–162, 1990.
- [Kat80] Kazuya Kato. A generalization of local class field theory by using K -groups. II. *J. Fac. Sci. Univ. Tokyo Sect. IA Math.*, 27(3):603–683, 1980.
- [Kat82a] Kazuya Kato. Galois cohomology of complete discrete valuation fields. In *Algebraic K-theory, Part II (Oberwolfach, 1980)*, volume 967 of *Lecture Notes in Math.*, pages 215–238. Springer, Berlin, 1982.
- [Kat82b] Kazuya Kato. Symmetric bilinear forms, quadratic forms and Milnor K -theory in characteristic two. *Invent. Math.*, 66(3):493–510, 1982.
- [Kat86] Kazuya Kato. A Hasse principle for two-dimensional global fields. *J. Reine Angew. Math.*, 366:142–183, 1986. With an appendix by Jean-Louis Colliot-Thélène.
- [KRS98] Bruno Kahn, Markus Rost, and R. Sujatha. Unramified cohomology of quadrics. I. *Amer. J. Math.*, 120(4):841–891, 1998.
- [KS00] Bruno Kahn and R. Sujatha. Motivic cohomology and unramified cohomology of quadrics. *J. Eur. Math. Soc. (JEMS)*, 2(2):145–177, 2000.
- [KS01] Bruno Kahn and R. Sujatha. Unramified cohomology of quadrics. II. *Duke Math. J.*, 106(3):449–484, 2001.
- [Lag02] Ahmed Laghribi. Certaines formes quadratiques de dimension au plus 6 et corps des fonctions en caractéristique 2. *Israel J. Math.*, 129:317–361, 2002.
- [Mer95] A. S. Merkurjev. The group $H^1(X, \mathcal{K}_2)$ for projective homogeneous varieties. *Algebra i Analiz*, 7(3):136–164, 1995.
- [Mil76] J. S. Milne. Duality in the flat cohomology of a surface. *Ann. Sci. École Norm. Sup. (4)*, 9(2):171–201, 1976.

- [Pey95] Emmanuel Peyre. Products of Severi-Brauer varieties and Galois cohomology. In *K-theory and algebraic geometry: connections with quadratic forms and division algebras (Santa Barbara, CA, 1992)*, volume 58 of *Proc. Sympos. Pure Math.*, pages 369–401. Amer. Math. Soc., Providence, RI, 1995.
- [Sah72] Chih Han Sah. Symmetric bilinear forms and quadratic forms. *J. Algebra*, 20:144–160, 1972.
- [Sal84] David J. Saltman. Noether’s problem over an algebraically closed field. *Invent. Math.*, 77(1):71–84, 1984.
- [Shi07] Atsushi Shiho. On logarithmic Hodge-Witt cohomology of regular schemes. *J. Math. Sci. Univ. Tokyo*, 14(4):567–635, 2007.
- [Č19] Kestutis Česnavičius. Purity for the Brauer group. *Duke Math. J.*, 168(8):1461–1486, 2019.

Contact information of the authors:

Yong HU

Department of Mathematics
Southern University of Science and Technology
Shenzhen 518055, China
Email: huy@sustech.edu.cn

Peng SUN

School of Mathematics
Hunan University
Changsha 410082, China
Email: sunpeng@hnu.edu.cn