

Submitted to *Bernoulli*

Minimum spanning trees of random geometric graphs with location dependent weights

GHURUMURUHAN GANESAN^{1,*}

¹*Institute of Mathematical Sciences, Chennai. E-mail: *gganesan82@gmail.com*

Consider n nodes $\{X_i\}_{1 \leq i \leq n}$ independently distributed in the unit square S , each according to a distribution f . Nodes X_i and X_j are joined by an edge if the Euclidean distance $d(X_i, X_j)$ is less than r_n , the adjacency distance and the resulting random graph G_n is called a random geometric graph (RGG). We now assign a location dependent weight to each edge of G_n and define MST_n to be the sum of the weights of the minimum spanning trees of all components of G_n . For values of r_n above the connectivity regime, we obtain upper and lower bound deviation estimates for MST_n and L^2 -convergence of MST_n appropriately scaled and centred.

AMS 2000 Subject Classification: Primary: 60J10, 60K35; Secondary: 60C05, 62E10, 90B15, 91D30.

Keywords:

Minimum spanning tree; random geometric graphs; location dependent edge weights

1. Introduction

The study of minimum spanning trees (MSTs) of a graph arise in many applications and many analytical results have been derived regarding the weight of the MST for various types of weighted graphs. For MSTs of complete Euclidean graphs with edge weights being Euclidean length, [2] study convergence of the weight of the travelling salesman path and describe modifications that allow for the study of MSTs appropriately scaled and centred. Also, [9] [10] [12] study MSTs of complete Euclidean graphs where the weights of the edges grow as a power of the Euclidean length of the edges. Asymptotic convergence of the MSTs (in probability and in the sense of CLT) together with auxiliary results are studied as the number of nodes $n \rightarrow \infty$ by estimating the expected weight using the bounded degree property of MSTs (see also [11] [13] [6] and references therein for more details). For material on the algorithmic and application oriented aspects of MSTs, we refer to [3] [5] and references therein.

In this paper, we consider random geometric graphs (RGGs) [7] [8] where nodes are distributed randomly across the unit square and nodes close enough to each other are connected to each other by edges. Also, the weight of an edge in the RGG might depend on the *individual* locations of the endvertices.

The scenario described above arises frequently in wireless networks and for example, suppose it is required to establish a fully connected wireless communication network among the set of nodes distributed randomly in a certain geographical area. The cost of setting up a communication link between any two nodes (e.g. length of the cables etc) is usually assumed to be directly proportional to the distance between the nodes [4]. Additionally, the cost could also depend on the location of the nodes since it might happen that some parts of area are “remote” and so it might cost more to install links among nodes located in these parts.

In the rest of this section, we describe the model under consideration and state our main results regarding the MSTs of RGGs with location dependent weights. In our main result (Theorem 1), we obtain deviation and variance estimates for the weight of the MST, appropriately scaled and centred.

Model Description

Let n nodes $\{X_i\}_{1 \leq i \leq n}$ be independently distributed in the unit square S , each according to a density f satisfying

$$\epsilon_1 \leq \inf_{x \in S} f(x) \leq \sup_{x \in S} f(x) \leq \epsilon_2 \quad (1.1)$$

for some positive constants $\epsilon_1 \leq 1 \leq \epsilon_2$, since $\int_S f(x) dx = 1$. Throughout constants do not depend on n . The Euclidean distance between nodes X_i and X_j is denoted by $d(X_i, X_j)$ and nodes X_i and X_j are connected by an edge if $d(X_i, X_j) < r_n$. The term r_n is called the adjacency distance and the resulting random graph G_n is called a random geometric graph (RGG) (Penrose (2003)).

Let $Y_1, \dots, Y_t \subset \{X_k\}_{1 \leq k \leq n}$ be $t \geq 2$ distinct nodes. A path $\mathcal{Q} = (Y_1, \dots, Y_t)$ is a subgraph of G_n with vertex set $\{Y_j\}_{1 \leq j \leq t}$ and edge set $\{(Y_j, Y_{j+1})\}_{1 \leq j \leq t-1}$. The nodes Y_1 and Y_t are said to be *connected* by edges of the path \mathcal{Q} . For $t \geq 3$, the subgraph $\mathcal{C} = (Y_1, Y_2, \dots, Y_t, Y_1)$ with vertex set $\{Y_j\}_{1 \leq j \leq t}$ and edge set $\{(Y_j, Y_{j+1})\}_{1 \leq j \leq t-1} \cup \{(Y_t, Y_1)\}$ is said to be a *cycle*.

A subgraph \mathcal{T} of G_n with vertex set $\{Y_i\}_{1 \leq i \leq t}$ and edge set $E_{\mathcal{T}}$ is said to be a *tree* if the following two conditions hold:

- (1) The graph \mathcal{T} is connected; i.e., any two nodes in \mathcal{T} are connected by a path containing only edges in $E_{\mathcal{T}}$.
- (2) The graph \mathcal{T} is acyclic; i.e., no subgraph of \mathcal{T} is a cycle.

The tree \mathcal{T} is said to be a *spanning tree* of a component \mathcal{C} of the graph G_n if \mathcal{T} contains all the nodes of \mathcal{C} . If \mathcal{T} contains all the n nodes $\{X_k\}_{1 \leq k \leq n}$ then \mathcal{T} is said to be a spanning tree of the RGG G_n .

We are interested in studying minimum spanning trees of G_n each of whose edges is assigned a random weight as described below.

Edge weights

Let $\xi : S \times S \rightarrow (0, \infty)$ be any measurable function satisfying $\xi(x, y) = \xi(y, x)$ for all $x, y \in S$. For $1 \leq i < j \leq n$ we define

$$w(X_i, X_j) := d^\alpha(X_i, X_j) \cdot \xi(X_i, X_j)$$

to be the weight of the edge (X_i, X_j) , associated with the (deterministic) edge weight factor ξ and exponent $\alpha > 0$. Throughout we assume that $\xi(x, y) \in [\xi_{min}, \xi_{max}]$ for some positive constants ξ_{min} and ξ_{max} . For a component \mathcal{C} of the RGG G_n with vertex set $\{Y_1, \dots, Y_t\}$ and for a spanning tree \mathcal{T} of \mathcal{C} , the *weight* of \mathcal{T} is the sum of the weights of the edges in \mathcal{T} ; i.e.,

$$W(\mathcal{T}) = W(\mathcal{T}, \xi) := \sum_{e \in \mathcal{T}} w(e) = \frac{1}{2} \sum_{i=1}^t w(Y_i, \mathcal{T}), \quad (1.2)$$

where $w(Y_i, \mathcal{T})$ is the sum of the weights of edges in \mathcal{T} containing Y_i as an endvertex. Let

$$MST_n = MST_n(G_n, \xi) := \sum_{\mathcal{C} \in G_n} \min_{\mathcal{T}} W(\mathcal{T}) \quad (1.3)$$

where the sum is taken over all components of G_n and for each component \mathcal{C} , the minimum is taken over all spanning trees \mathcal{T} of \mathcal{C} . If G_n is connected, then we denote the spanning tree \mathcal{T}_n with weight MST_n to be the *minimal spanning tree* (MST). If there is more than one choice for \mathcal{T}_n , we choose one according to a deterministic rule.

We emphasize here that the edge weight factor ξ is deterministic and the only randomness in MST_n comes from the location of the nodes $\{X_i\}_{1 \leq i \leq n}$. Let ϵ_1, ϵ_2 be as in (1.1) and set $\delta = \delta(\alpha) = \epsilon_1$ if the edge weight exponent $\alpha \leq 1$ and $\delta = \epsilon_2$ if $\alpha > 1$. For $A > 0$ we define $C_1(A) = C_1(A, \epsilon_1, \epsilon_2, \alpha)$ and $C_2(A) = C_2(A, \epsilon_1, \epsilon_2, \alpha)$ as

$$\begin{aligned} C_1(A) &:= \frac{1}{2} \xi_{min} A^{\alpha-2} (1 - e^{-\epsilon_1 A^2}) e^{-8\epsilon_2 A^2} \text{ and} \\ C_2(A) &:= \xi_{max} (2A)^\alpha \left(1 + \frac{\mathbb{E} \tilde{T}^\alpha}{A^2} \right), \end{aligned} \quad (1.4)$$

where \tilde{T} is a geometric random variable with success parameter $p = 1 - e^{-\delta A^2}$. We have the following result.

Theorem 1. Suppose the following conditions are satisfied:

(i) The adjacency distance $r_n \geq \sqrt{\frac{M \log n}{n}}$ and $r_n \rightarrow 0$ as $n \rightarrow \infty$, where $M > \left(\frac{1600}{\epsilon_1}\right)$ is a constant.

(ii) The edge weight exponent $\alpha > 0$ and the edge weight factor $\xi(x, y) \in [\xi_{min}, \xi_{max}]$ for all x, y and some positive constants ξ_{min}, ξ_{max} .

There is a constant $C > 0$ such that

$$\text{var} \left(\frac{MST_n}{n^{1-\frac{\alpha}{2}}} \right) \leq C r_n^2 (n r_n^2)^\alpha. \quad (1.5)$$

Moreover, for every $A > 0$ there exists $A_n \in \left[A + \frac{1}{(\log n)^{1/4}}, A + \frac{2}{(\log n)^{1/4}} \right)$ and a constant $D > 0$ such that

$$\mathbb{P} \left(MST_n \geq C_1(A_n) n^{1-\frac{\alpha}{2}} \left(1 - \frac{36\sqrt{A_n}}{n^{1/4}} \right) \right) \geq 1 - \frac{1}{n^2} \quad (1.6)$$

and

$$\mathbb{P} \left(MST_n \leq C_2(A_n) n^{1-\frac{\alpha}{2}} \left(1 + \frac{1}{n^{1/17}} \right) \right) \geq 1 - \frac{D}{n^{5/4}} \quad (1.7)$$

for all n large. Consequently,

$$C_1(A_n) \left(1 - \frac{37\sqrt{A_n}}{n^{1/4}} \right) \leq \mathbb{E} \left(\frac{MST_n}{n^{1-\frac{\alpha}{2}}} \right) \leq C_2(A_n) \left(1 + \frac{2}{n^{1/17}} \right) \quad (1.8)$$

for all n large.

We remark here that we have chosen r_n to be at least of the order of $\sqrt{\frac{\log n}{n}}$ so that the resulting RGG is connected with high probability (see discussion following (2.4) in Section 2). From the variance estimate (1.5) in Theorem 1, we see that if

$$n^{\frac{1}{2} \cdot \frac{\alpha}{1+\alpha}} \cdot r_n \rightarrow 0 \quad (1.9)$$

as $n \rightarrow \infty$, then

$$\frac{MST_n - \mathbb{E} MST_n}{n^{1-\frac{\alpha}{2}}} \rightarrow 0$$

in L^2 as $n \rightarrow \infty$. For example if $\alpha = 1$ and $r_n = \frac{1}{n^{1/3}}$, then both condition (i) in Theorem 1 and (1.9) are satisfied.

We use (1.7) to evaluate the asymptotic numeric bounds for $\mathbb{E}MST_n$ as follows. Recalling the definition of A_n in the statement of Theorem 1, we prove in the Appendix that for $A > 0$, the limits

$$\lim_n C_i(A_n) = C_i(A) \text{ for } i = 1, 2. \quad (1.10)$$

Therefore defining

$$\beta_{up} := \inf_{A>0} C_2(A) \text{ and } \beta_{low} := \sup_{A>0} C_1(A),$$

we get from (1.8) that

$$\beta_{low} \leq \liminf_n \mathbb{E} \left(\frac{MST_n}{n^{1-\frac{\alpha}{2}}} \right) \leq \limsup_n \mathbb{E} \left(\frac{MST_n}{n^{1-\frac{\alpha}{2}}} \right) \leq \beta_{up}.$$

For example, for the homogenous case of $\epsilon_1 = \epsilon_2 = \xi_{min} = \xi_{max} = 1$ and $\alpha = 1$, we have that

$$\beta_{low} \approx 0.0735633 \text{ and } \beta_{up} \approx 4.46256.$$

We briefly outline the methods in the proof of Theorem 1. The variance bound (1.5) is obtained via the martingale difference method together with one node difference estimates that measures the change in MST lengths after adding or removing a single node, retaining the same adjacency distance.

To prove the deviation estimates, we use Poissonization and prove results for the Poissonized RGG and then dePoissonize to obtain the corresponding results for the Binomial RGG. In the rest of the two paragraphs we simply denote the Poissonized RGG as RGG. First we show that the event E_{conn} that the (Poissonized) RGG is connected occurs with high probability, i.e., with probability converging to one as $n \rightarrow \infty$ and consider only subevents of E_{conn} . For the lower deviation estimate, we tile the unit square into roughly n small squares $\{R_k\}$ of side length of order $\frac{1}{\sqrt{n}}$ each and look for special type of occupied squares, whose neighbouring squares are all vacant. This results in a vacant annulus around such squares and since the RGG is connected, nodes within these special squares must have edges crossing over the vacant annulus to reach other nodes. This allows us to construct edges of length of order $\frac{1}{\sqrt{n}}$ in the MST and determining that there are order of n such special squares with high probability, we obtain the lower deviation bound.

For the upper deviation bound, we join nodes within each square in $\{R_k\}$ to get a collection of subtrees. We then join all these subtrees together by adding extra edges to get a spanning tree whose weight is no more than the MST. Further the weight of this constructed tree has two parts: (1) the sum length of the edges of the subtrees, each of which is of order $\frac{1}{\sqrt{n}}$ by construction and (2) the sum length of the extra edges. To estimate this second sum, we use coupling with homogenous Poisson process and deviation estimates for Geometric random variables to obtain the desired upper bound. For more details, we refer to Section 3 and finally, we remark that the methods used in this paper can also be used to analyze RGGs in d -dimensions and in regular shapes other than the unit square.

The paper is organized as follows. In Section 2, we prove the variance estimate (1.5) in Theorem 1 and in Section 3, we prove the deviation estimates (1.6) and (1.7) and the expectation bounds (1.8) in Theorem 1.

2. Proof of the variance estimate in Theorem 1

Throughout we use the following standard deviation estimates [1] for sums of independent Poisson and Bernoulli random variables. Suppose $W_i, 1 \leq i \leq m$ are independent Bernoulli random variables satisfying $\mu_1 \leq \mathbb{P}(W_1 = 1) = 1 - \mathbb{P}(W_1 = 0) \leq \mu_2$. For any $0 < \epsilon < \frac{1}{2}$,

$$\mathbb{P}\left(\sum_{i=1}^m W_i > \mu_2(1 + \epsilon)\right) \leq \exp\left(-\frac{\epsilon^2}{4}\mu_2\right) \quad (2.1)$$

and

$$\mathbb{P}\left(\sum_{i=1}^m W_i < \mu_1(1 - \epsilon)\right) \leq \exp\left(-\frac{\epsilon^2}{4}\mu_1\right) \quad (2.2)$$

Estimates (2.1) and (2.2) also hold if $\{W_i\}$ are independent Poisson random variables with $\mu_1 \leq \mathbb{E}W_1 \leq \mu_2$.

One node difference estimates

In this subsection we find estimates for changes in the length of the MST upon adding or removing a single node. By definition, the adjacency distance r_n depends on the number of nodes n in consideration and so strictly speaking, adding or removing a node changes the adjacency distance and hence the underlying RGG. However, for our purposes we obtain below difference estimates *retaining* the same adjacency distance.

As in (1.3), let MST_{n+1} be the sum of the weights of the MSTs of all the components of the RGG G_{n+1} , formed by the nodes $\{X_i\}_{1 \leq i \leq n+1}$ with adjacency distance r_{n+1} . For $1 \leq i \leq n+1$, let $G(i, r_{n+1})$ be the RGG formed by the n nodes $\{X_j\}_{1 \leq j \neq i \leq n+1}$ with the *same* adjacency distance r_{n+1} and let $MST_n(i, r_{n+1})$ be the sum of the weights of the MSTs of all the components of the RGG $G(i, r_{n+1})$, as defined in (1.3). We are interested in estimating $|MST_{n+1} - MST(i, r_{n+1})|$ for any $1 \leq i \leq n+1$.

We have a couple of preliminary definitions. Tile the unit square S into W^2 disjoint $t_n \times t_n$ squares $\{S_l\}_{1 \leq l \leq W^2}$ as in Figure 1 where

$$\frac{1}{20}\sqrt{\frac{M \log n}{n}} \leq \frac{1}{10}\sqrt{\frac{M \log(n+1)}{n+1}} \leq \frac{r_{n+1}}{10} \leq t_n := \frac{r_{n+1}}{2\sqrt{2} + \delta_n} \leq r_{n+1} \quad (2.3)$$

for all n large and $\delta_n = \lfloor \sqrt{r_{n+1}}, 2\sqrt{r_{n+1}} \rfloor$ is such that $\frac{1}{t_n}$ is an integer for all n large. This is possible since $r_n \rightarrow 0$ and so the difference

$$\frac{2\sqrt{2} + 2\sqrt{r_{n+1}}}{r_{n+1}} - \frac{2\sqrt{2} + \sqrt{r_{n+1}}}{r_{n+1}} = \frac{1}{\sqrt{r_{n+1}}} \rightarrow \infty$$

as $n \rightarrow \infty$. The estimates in (2.3) follow from the fact that $nr_n^2 \geq M \log n$ for all n large (see bounds for r_n in the statement of the Theorem).

S_1					
S_2					
S_3	•				
•	•				•
•	•				•
•	S_{W+2}				•
•	S_{W+1}				•
S_W	S_{W+1}				S_{W^2}

Figure 1: Tiling the unit square into $W^2 = \frac{1}{t_n^2}$ smaller $t_n \times t_n$ squares $\{S_l\}_{1 \leq l \leq W^2}$.

For $1 \leq j \leq n+1$ and $1 \leq i \leq W^2$, let $E_j(i)$ be the event that the square S_i contains between $\frac{\epsilon_1 n t_n^2}{2}$ and $2\epsilon_2 n t_n^2$ nodes of $\{X_k\}_{1 \leq k \neq j \leq n+1}$ where ϵ_1 and ϵ_2 are as in (1.1) and let

$$E_{dense}(n+1) := \bigcap_{j=1}^{n+1} \bigcap_{i=1}^{W^2} E_j(i). \quad (2.4)$$

If $E_{dense}(n+1)$ occurs, then each $t_n \times t_n$ square in $\{S_l\}$ contains at least one node of $\{X_j\}_{1 \leq j \leq n+1}$. Since $2t_n\sqrt{2} < r_{n+1}$ (see (2.3)), nodes present in squares of $\{S_l\}$ sharing a corner are attached to each other by edges in the RGG G_{n+1} . Thus G_{n+1} is connected and we let \mathcal{T}_{n+1} be the (unique) MST of G_{n+1} .

For $1 \leq i \leq n+1$, we denote d_i to be the degree of the node X_i in the MST of the component containing the node X_i in the graph G_{n+1} so that if the event $E_{dense}(n+1)$ occurs, then d_i is simply the degree of the node X_i in the MST \mathcal{T}_{n+1} . Recalling that ϵ_1 and ϵ_2 are the bounds for the distribution f as described in (1.1), we have the following result.

Lemma 2. Suppose conditions (i) – (ii) in Theorem 1 hold. For all n large,

$$\mathbb{P}(E_{dense}(n+1)) \geq 1 - \frac{1}{n^3} \quad (2.5)$$

and if $E_{dense}(n+1)$ occurs, then for any $1 \leq i \leq n+1$ the degree $1 \leq d_i \leq 200\epsilon_2 n r_{n+1}^2$. Moreover, for any $1 \leq i \leq n+1$, the difference

$$\begin{aligned} |MST_{n+1} - MST(i, r_{n+1})| &\leq \xi_{max} d_i \cdot r_{n+1}^\alpha \mathbb{1}(E_{dense}(n+1)) \\ &\quad + \xi_{max} n \cdot r_{n+1}^\alpha \mathbb{1}(E_{dense}^c(n+1)), \end{aligned}$$

(2.6)

where ξ_{max} is the maximum weight factor of an edge as defined prior to (1.2).

Proof of Lemma 2: We first prove the estimate (2.5). For $1 \leq i \leq W^2 = \frac{1}{t_n^2}$, let N_i be the random number of nodes of $\{X_k\}_{1 \leq k \leq n}$ in the $t_n \times t_n$ square S_i . Since each node occurs independently with probability $\int_{S_i} f(x) dx$ within the square S_i , the random variable N_i is Binomially distributed and so from the bounds on the distribution $f(\cdot)$ in (1.1), the average number of nodes $\mathbb{E}N_i = n \int_{S_i} f(x) dx$ satisfies

$$8 \leq \frac{\epsilon_1 M \log n}{20} \leq \frac{\epsilon_1 n r_{n+1}^2}{10} \leq \epsilon_1 n t_n^2 \leq \mathbb{E}N_i \leq \epsilon_2 n t_n^2 \leq \epsilon_2 n r_{n+1}^2 \quad (2.7)$$

for all n large, where $\epsilon_1, \epsilon_2 > 0$ are as in (1.1). The estimates in (2.7) follow from the definition of t_n in (2.3) and the fact that $n r_n^2 \geq M \log n$ (see statement of Theorem 1).

Plugging the bounds obtained in (2.7) into the standard deviation estimates (2.1) and (2.2) with $\mu_1 = \epsilon_1 t_n^2$ and $\mu_2 = \epsilon_2 t_n^2$ and letting $\epsilon = \frac{1}{2}$ there, we therefore get

$$\mathbb{P}\left(\frac{\epsilon_1 n t_n^2}{2} \leq N_i \leq \frac{3\epsilon_2 n t_n^2}{2}\right) \geq 1 - 2 \exp\left(-\frac{\epsilon_1 n t_n^2}{16}\right)$$

for all n large, not depending on the choice of i . Thus $\mathbb{P}(E_j(i)) \geq 1 - 2e^{-\frac{\epsilon_1 n t_n^2}{16}}$ and since there are $W^2 = \frac{1}{t_n^2}$ squares in $\{S_l\}$ we get from the definition of the event $E_{dense}(n+1)$ in (2.4) that

$$\begin{aligned} \mathbb{P}(E_{dense}(n+1)) &\geq 1 - (n+1) \cdot \frac{1}{t_n^2} \cdot \exp\left(-\frac{\epsilon_1 n t_n^2}{16}\right) \\ &\geq 1 - (n+1) \cdot \frac{10}{r_{n+1}^2} \cdot \exp\left(-\frac{\epsilon_1 n r_{n+1}^2}{160}\right) \end{aligned}$$

since $t_n^2 \geq \frac{r_{n+1}^2}{10}$ (see definition of t_n in (2.3)). Using the bounds in (2.3), we then get that

$$\mathbb{P}(E_{dense}(n+1)) \geq 1 - (n+1) \cdot \frac{20n}{M \log n} \cdot \exp\left(-\frac{\epsilon_1 M \log n}{320}\right) \quad (2.8)$$

which is at least $1 - \frac{1}{n^3}$ for all n large, provided $M > \frac{1600}{\epsilon_1}$. We henceforth fix such an M .

We now prove the difference estimate (2.6). Let \mathcal{T}_{n+1} be the weighted MST containing all the $n+1$ nodes $\{X_j\}_{1 \leq j \leq n+1}$ in the RGG G_{n+1} with adjacency distance r_{n+1} . Similarly, for $1 \leq i \leq n+1$ let $\mathcal{T}(i, r_{n+1})$ be the weighted MST of the RGG $G(i, r_{n+1})$ formed by the nodes $\{X_k\}_{1 \leq k \neq i \leq n+1}$ with adjacency distance r_{n+1} .

Suppose first that the event $E_{dense}(n+1)$ does not occur. There are at most n edges in either \mathcal{T}_{n+1} or $\mathcal{T}(i, r_{n+1})$ and each such edge has length at most r_{n+1} . Since the weight factor of any edge is at most ξ_{max} , we therefore get

$$\max(MST_{n+1}, MST(i, r_{n+1})) \mathbb{1}(E_{dense}^c(n+1)) \leq \xi_{max} n r_{n+1}^\alpha \mathbb{1}(E_{dense}^c(n+1)).$$

Henceforth we assume that $E_{dense}(n+1)$ occurs so that from the discussion prior to the statement of Lemma 2, we get that the RGG G_{n+1} is connected. Arguing similarly, we get that $G(i, r_{n+1}) \subseteq$




Figure 2: The spanning tree \mathcal{T}_{n+1} where the node X_i has three neighbours $u_i, 1 \leq i \leq 3$. The corresponding subtrees $\mathcal{T}(u_i), 1 \leq i \leq 3$ are denoted by the dotted lines.

G_{n+1} is also connected and wherever the location of the node X_i , there is always a node of $X_{l(i)} \in \{X_k\}_{1 \leq k \neq i \leq n+1}$ present within a distance of $t_n \sqrt{2} < r_{n+1}$ from X_i . This means that $\mathcal{T}(i, r_{n+1}) \cup \{(X_i, X_{l(i)})\}$ is a (not necessarily minimal) spanning tree containing all the nodes $\{X_j\}_{1 \leq j \leq n+1}$. The edge $(X_i, X_{l(i)})$ has weight factor at most ξ_{max} and so the weight of the spanning tree \mathcal{T}_{n+1} does not exceed the weight of $\mathcal{T}(i, r_{n+1})$ by more than $\xi_{max} r_{n+1}^\alpha$. Consequently

$$\begin{aligned} MST_{n+1} \mathbb{1}(E_{dense}(n+1)) &\leq MST(i, r_{n+1}) \mathbb{1}(E_{dense}(n+1)) \\ &\quad + \xi_{max} r_{n+1}^\alpha \mathbb{1}(E_{dense}(n+1)). \end{aligned} \quad (2.9)$$

To get an inequality in the reverse direction, let $u_1, \dots, u_{d_i}, d_i \geq 1$ be the neighbours of X_i in the MST \mathcal{T}_{n+1} containing all the nodes $\{X_k\}_{1 \leq k \leq n+1}$. Removing the node X_i from \mathcal{T}_{n+1} , we therefore obtain d_i vertex disjoint trees $\mathcal{T}(u_1), \dots, \mathcal{T}(u_{d_i})$ with edges in $G(i, r_{n+1})$. This is illustrated in Figure 2 where $d_i = 3$.

Since $G(i, r_{n+1})$ is connected, the node u_1 is connected to u_2 by a path $P_{12} = (e_1, \dots, e_g) \subseteq G(i, r_{n+1})$. Without loss of generality, we assume that the path P_{12} contains only edges of two types: Each edge e_k is either an edge of a tree in $\{\mathcal{T}(u_j)\}_{1 \leq j \leq d_i}$ or e_k has one endvertex in $\mathcal{T}(u_a)$ and another vertex in a different tree $\mathcal{T}(u_b)$. This is illustrated in Figure 3 where e_{j_1} is the first edge that “leaves” the tree \mathcal{T}_{u_1} and therefore contains an endvertex v not in $\mathcal{T}(u_1)$. Since the trees $\{\mathcal{T}(u_w)\}$ contain all the nodes $\{X_k\}_{1 \leq k \neq i \leq n+1}$, there exists $1 \leq w_1 \neq u_1 \leq d_i$ such that v belongs to $\mathcal{T}(u_{w_1})$. We denote the edge e_{j_1} to be a bridge and adding the edge e_{j_1} to the collection of trees $\{\mathcal{T}(u_w)\}$, we get that the graph $e_{j_1} \cup \cup \{\mathcal{T}(u_w)\}_{1 \leq w \leq d_i}$ contains $d_i - 1$ trees.

We continue the above procedure of identifying bridges in the path P_{12} and merging trees together until all edges of P_{12} are exhausted. The resulting final graph $G_{merge}^{(1)} \subseteq G(i, r_{n+1})$ is a forest containing at most $d_i - 1$ trees and the nodes u_1 and u_2 belong to the same tree of $G_{merge}^{(1)}$. If there still exist nodes u_k and u_l that belong to different trees of $G_{merge}^{(1)}$, we repeat the above procedure and




Figure 3: The path $P_{12} \subset G(i, r_{n+1})$ connecting u_1 and u_2 is formed by the union of e_{j_1}, e_{j_2} and the wavy curves. We merge the trees $\mathcal{T}(u_i), 1 \leq i \leq 3$ together by iteratively adding the bridges e_{j_1} and e_{j_2} .

connect u_k and u_l to get a new forest $G_{merge}^{(2)}$ containing at most $d_i - 2$ trees. Continuing this way iteratively for a finite number of steps s , we finally obtain a spanning tree $G_{merge}^{(s)}$ of $G(i, r_{n+1})$.

In Figure 3, we have illustrated the tree merging procedure described in the previous paragraph for the case when the number of neighbours $d_i = 3$. For $1 \leq j \leq 3$, the dotted triangle containing the vertex u_j represents the tree $\mathcal{T}(u_j)$. The wavy curves together with the two edges e_{j_1} and e_{j_2} (which are bridges) form the path P_{12} . Adding the edge e_{j_1} merges the trees $\mathcal{T}(u_1)$ and $\mathcal{T}(u_3)$. Adding the edge e_{j_2} to the resulting graph merges all the trees. In this example, all the three trees have been merged at the end of the first iteration.

The number of bridges added in the above tree merging procedure is $d_i - 1$ (one less than the number of trees in $\{\mathcal{T}(u_w)\}_{1 \leq w \leq d_i}$) and since any bridge has length at most r_{n+1} and weight factor at most ξ_{max} , we get that the weight $W(G_{merge}^{(s)})$ of the final spanning tree of $G(i, r_{n+1})$ obtained satisfies

$$\begin{aligned} MST(i, r_{n+1})\mathbb{1}(E_{dense}(n+1)) &\leq W(G_{merge}^{(s)})\mathbb{1}(E_{dense}(n+1)) \\ &\leq MST_{n+1}\mathbb{1}(E_{dense}(n+1)) \\ &\quad + \xi_{max}d_i \cdot r_{n+1}^\alpha \mathbb{1}(E_{dense}(n+1)). \end{aligned} \tag{2.10}$$

Combining (2.9) and (2.10) and using the fact that $d_i \geq 1$ if $E_{dense}(n+1)$ occurs (see discussion prior to statement of Lemma 2), we then get (2.6).

Finally, to upper bound d_i , we proceed as follows. Suppose $X_i \in S_l$ and let $\mathcal{N}_1(S_l)$ be the set of all squares in $\{S_k\}$ sharing a corner with S_l . Similarly, for $k \geq 2$ we let $\mathcal{N}_k(S_l)$ be the set of squares sharing a corner with some square in $\mathcal{N}_{k-1}(S_l)$. Since every edge in G_{n+1} has length at

most $r_{n+1} < \frac{5t_n}{\sqrt{2}}$ (see definition of t_n in (2.3)), the neighbours of X_i , $\{u_w\}_{1 \leq w \leq t}$, are all present in $\mathcal{N}_4(S_l)$.

There are $81 < 100$ squares of $\{S_k\}$ in the set $\mathcal{N}_4(S_l)$ and because $E_{dense}(n+1)$ occurs, every square in $\mathcal{N}_4(S_l)$ has at most $2\epsilon_2 n t_n^2$ nodes of $\{X_j\}_{1 \leq j \neq i \leq n+1}$ (see definition of $E_{dense}(n+1)$ in (2.4)). Therefore $d_i \leq 200\epsilon_2 n t_n^2 \leq 200\epsilon_2 n r_{n+1}^2$ (see definition of t_n in (2.3)). ■

Proof of the variance estimate (1.5)

We use the martingale difference method and for $1 \leq j \leq n+1$, let $\mathcal{F}_j = \sigma(\{X_k\}_{1 \leq k \leq j})$ denote the σ -field generated by the node positions $\{X_k\}_{1 \leq k \leq j}$. Defining the martingale difference

$$H_j = \mathbb{E}(MST_{n+1} | \mathcal{F}_j) - \mathbb{E}(MST_{n+1} | \mathcal{F}_{j-1}),$$

we then have that $MST_{n+1} - \mathbb{E}MST_{n+1} = \sum_{j=1}^{n+1} H_j$ and so by the martingale property

$$\text{var}(MST_{n+1}) = \mathbb{E} \left(\sum_{j=1}^{n+1} H_j \right)^2 = \sum_{j=1}^{n+1} \mathbb{E} H_j^2.$$

Using the one node difference estimate from Lemma 2, we prove below that

$$\sum_{j=1}^{n+1} \mathbb{E} H_j^2 \leq C n^2 r_{n+1}^{2+2\alpha} \quad (2.11)$$

for some constant $C > 0$ and this obtains (1.5).

Proof of (2.11): We first rewrite the martingale difference H_j in a more convenient form. Let $\eta = (x_1, \dots, x_{n+1})$ and $\eta' = (y_1, \dots, y_{n+1})$ be two vectors in $(\mathbb{R}^2)^{n+1}$ so that $\{x_k\}_{1 \leq k \leq n+1}$ denote the node positions in the configuration η . For $1 \leq j \leq n$, we let $\eta_j = (x_1, \dots, x_j, y_{j+1}, \dots, y_{n+1})$ be the configuration obtained by considering the first j entries from η and the remaining from η' . Let $G(\eta, r_{n+1})$ be the RGG formed by the nodes of η with adjacency distance r_{n+1} and let $M(\eta)$ be the total MST weight of $G(\eta, r_{n+1})$ as in (1.3). Using Fubini's theorem with the notation $\eta_0 = \eta'$ and $\eta_{n+1} = \eta$, we then get that the martingale difference

$$H_j = \int (M(\eta_j) - M(\eta_{j-1})) \prod_{k=j}^{n+1} f(y_k) dy_k$$

and so we have that $|H_j| \leq L_j$, where

$$L_j := \int |M(\eta_j) - M(\eta_{j-1})| \prod_{k=j}^{n+1} f(y_k) dy_k. \quad (2.12)$$

Recalling the event $E_{dense}(n+1)$ defined in (2.4) we write $L_j = I_1 + I_2$ where I_1 equals

$$\int |M(\eta_j) - M(\eta_{j-1})| \mathbb{1}(\eta_j \in E_{dense}(n+1)) \mathbb{1}(\eta_{j-1} \in E_{dense}(n+1)) \prod_{k=j}^{n+1} f(y_k) dy_k \quad (2.13)$$

and $I_2 = L_j - I_1$ and estimate $\mathbb{E}I_1^2$ and $\mathbb{E}I_2^2$ below, separately.

Estimate for I_1 : Suppose the configuration $\eta_j \in E_{dense}(n+1)$ and let θ_j be the configuration obtained by removing the node x_j from η_j . By definition θ_j is also the configuration obtained by removing the node y_j from η_{j-1} . We then recall from the discussion prior to and in Lemma 2 that the RGGs formed by the nodes in η_j and in θ_j , with adjacency distance r_{n+1} , are both connected. Denoting the corresponding weighted MST lengths as defined in (1.3) to be $M(\eta_j)$ and $M(\theta_j)$, respectively, we get from the one node difference estimate (2.6) in Lemma 2 that

$$|M(\eta_j) - M(\theta_j)| \leq \xi_{max} d_j(\eta_j) r_{n+1}^\alpha, \quad (2.14)$$

where $1 \leq d_j(\eta_j) \leq 200\epsilon_2 n r_{n+1}^2$ is the degree of the node x_j in the weighted MST formed by the nodes of η_j . Similarly we have

$$|M(\eta_{j-1}) - M(\theta_j)| \leq \xi_{max} d_j(\eta_{j-1}) r_{n+1}^\alpha, \quad (2.15)$$

where $d_j(\eta_{j-1})$ is the degree of the node y_j in the weighted MST formed by the nodes of η_{j-1} .

From (2.14), (2.15) and the triangle inequality, we get that

$$|M(\eta_j) - MST(\eta_{j-1})| \leq \xi_{max} (d_j(\eta_j) + d_j(\eta_{j-1})) r_{n+1}^\alpha.$$

Using

$$(d_j(\eta_j) + d_j(\eta_{j-1}))^2 \leq 2(d_j^2(\eta_j) + d_j^2(\eta_{j-1})) \leq 400\epsilon_2 n r_{n+1}^2 (d_j(\eta_j) + d_j(\eta_{j-1}))$$

and the definition of I_1 in (2.13), we then get that

$$\begin{aligned} I_1^2 &\leq 400\epsilon_2 \xi_{max}^2 n r_{n+1}^{2+2\alpha} \int (d_j(\eta_j) + d_j(\eta_{j-1})) \prod_{k=j}^{n+1} f(y_k) dy_k \\ &= 400\epsilon_2 \xi_{max}^2 n r_{n+1}^{2+2\alpha} (\mathbb{E}(d_j | \mathcal{F}_j) + \mathbb{E}(d_j | \mathcal{F}_{j-1})) \end{aligned} \quad (2.16)$$

where we recall from the paragraph prior to Lemma 2 that d_j is the degree of the node X_j in the MST of the component containing the node X_j in the graph G_{n+1} . Taking expectations in (2.16), we get

$$\mathbb{E}I_1^2 \leq 800\epsilon_2 \xi_{max}^2 n r_{n+1}^{2+2\alpha} \mathbb{E}(d_j). \quad (2.17)$$

Estimate for I_2 : The total number of edges in the spanning trees of all the components formed by the RGG containing the $n+1$ nodes of η_j is at most n and each such edge has length at most r_{n+1} . Since all edge weight factors are bounded above by ξ_{max} , we use the estimate $\mathbb{1}(A^c \cup B^c) \leq \mathbb{1}(A^c) + \mathbb{1}(B^c)$ with

$$A = \{\eta_j \in E_{dense}(n+1)\} \text{ and } B = \{\eta_{j-1} \in E_{dense}(n+1)\}$$

to get that $I_2 \leq J_1 + J_2$ where

$$\begin{aligned} J_1 &= n r_{n+1}^\alpha \xi_{max} \int \mathbb{1}(\eta_j \notin E_{dense}(n+1)) \prod_{k=j}^{n+1} f(y_k) dy_k \\ &= n r_{n+1}^\alpha \xi_{max} \mathbb{P}(E_{dense}^c(n+1) | \mathcal{F}_j) \end{aligned}$$

and $J_2 = n r_{n+1}^\alpha \xi_{max} \mathbb{P}(E_{dense}^c(n+1) | \mathcal{F}_{j-1})$.

Thus

$$J_1^2 \leq n^2 r_{n+1}^{2\alpha} \xi_{max}^2 \left(\mathbb{P}(E_{dense}^c(n+1) | \mathcal{F}_j) \right)^2 \leq n^2 r_{n+1}^{2\alpha} \xi_{max}^2 \mathbb{P}(E_{dense}^c(n+1) | \mathcal{F}_j)$$

and analogously, we also have

$$J_2^2 \leq n^2 r_{n+1}^{2\alpha} \xi_{max}^2 \mathbb{P}(E_{dense}^c(n+1) | \mathcal{F}_{j-1}).$$

Using $I_2^2 \leq (J_1 + J_2)^2 \leq 2(J_1^2 + J_2^2)$ and taking expectations, we then get

$$\mathbb{E}(I_2^2) \leq 4n^2 r_{n+1}^{2\alpha} \xi_{max}^2 \mathbb{P}(E_{dense}^c(n+1)) \leq 4n^2 r_{n+1}^{2\alpha} \xi_{max}^2 \frac{1}{n^3} \quad (2.18)$$

using the estimate for $E_{dense}(n+1)$ from Lemma 2.

Adding the estimates for $\mathbb{E}I_1^2$ and $\mathbb{E}I_2^2$ in (2.17) and (2.18), respectively, we get

$$\mathbb{E}I_1^2 + \mathbb{E}I_2^2 \leq 800\epsilon_2 \xi_{max}^2 n r_{n+1}^{2+2\alpha} \mathbb{E}(d_j) + 4n^2 r_{n+1}^{2\alpha} \xi_{max}^2 \frac{1}{n^3}$$

and since $\mathbb{E}H_j^2 \leq \mathbb{E}I_j^2 = \mathbb{E}(I_1 + I_2)^2 \leq 2(\mathbb{E}I_1^2 + \mathbb{E}I_2^2)$ we get that

$$\sum_{j=1}^{n+1} \mathbb{E}H_j^2 \leq 800\epsilon_2 \xi_{max}^2 n r_{n+1}^{2+2\alpha} \mathbb{E} \left(\sum_{j=1}^{n+1} d_j \right) + 4n^2 r_{n+1}^{2\alpha} \xi_{max}^2 \frac{(n+1)}{n^3}.$$

The sum $\sum_{j=1}^{n+1} d_j \leq 2n$ always since there are at most n edges in total among all the MSTs of the components of the RGG G_{n+1} and so

$$\sum_{j=1}^{n+1} \mathbb{E}H_j^2 \leq \left(1600\epsilon_2 \xi_{max}^2 n^2 r_{n+1}^{2+2\alpha} + 4n^2 r_{n+1}^{2\alpha} \xi_{max}^2 \frac{(n+1)}{n^3} \right).$$

Since $r_{n+1}^2 \geq \frac{M \log(n+1)}{n+1} \geq \frac{n+1}{\epsilon_2 n^3}$ for all n large (see bounds for r_n in the statement of the Theorem), we get that $\sum_{j=1}^{n+1} \mathbb{E}H_j^2 \leq 1604\epsilon_2 \xi_{max}^2 n^2 r_{n+1}^{2+2\alpha}$ for all n large, proving (2.11). \blacksquare

3. Proof of the deviation estimates in Theorem 1

Throughout we use Poissonization and so we construct a Poisson process \mathcal{P} in the unit square S with intensity $nf(\cdot)$ as follows. We tile the unit square S into $W^2 = \frac{1}{t_n^2}$ disjoint $t_n \times t_n$ squares $\{S_i\}_{1 \leq i \leq W^2}$ (see Figure 1), where we redefine $t_n = \frac{r_n}{2\sqrt{2+\delta_n}}$ and choose $\delta_n \in [\sqrt{r_n}, 2\sqrt{r_n})$ such that $W = \frac{1}{t_n}$ is an odd integer for all n large. This is possible by an analogous argument following (2.3).

Each square $S_i, 1 \leq i \leq W^2$ has Poisson number of nodes present in it and for a formal definition we let $\{V_{i,k}\}_{1 \leq i \leq W^2, k \geq 1}$ be i.i.d. random vectors in \mathbb{R}^2 with density $\frac{f(x)}{\int_{S_i} f(x) dx} \mathbb{1}(x \in S_i)$ and let $\{N_i\}_{1 \leq i \leq W^2}$ be independent Poisson random variables such that N_i has mean $n \int_{S_i} f(x) dx$ for $1 \leq i \leq W^2$. The random variables $\{N_i\}$ are independent of $\{V_{i,k}\}$ and we define $(\{V_{i,k}\}, \{N_i\})$ on the probability space $(\Omega_0, \mathcal{F}_0, \mathbb{P}_0)$.

For $1 \leq i \leq W^2$, we set $\{V_{i,k}\}_{1 \leq k \leq N_i}$ to be the nodes of \mathcal{P} in the square S_i . If u, v are any two nodes of \mathcal{P} that are within a Euclidean distance of r_n of each other, we join them by an edge and denote the resulting random graph as $G_n^{(P)}$. As in (1.3) we let $MST_n^{(P)}$ denote the sum of the weights of the minimum spanning trees of all components of $G_n^{(P)}$. We first obtain deviation estimates for $MST_n^{(P)}$ and then convert via dePoissonization to obtain the corresponding estimates for MST_n , the MST length for the Binomial process.

Analogous to (2.7), we have for every $1 \leq i \leq W^2$ that

$$8 \leq \frac{\epsilon_1 M \log n}{20} \leq \frac{\epsilon_1 n r_n^2}{10} \leq \epsilon_1 n t_n^2 \leq \mathbb{E}_0 N_i \leq \epsilon_2 n t_n^2 \leq \epsilon_2 n r_n^2, \quad (3.1)$$

where $\epsilon_1, \epsilon_2 > 0$ are as in (1.1). If $E_i := \{\frac{\epsilon_1 n t_n^2}{2} \leq N_i \leq 2\epsilon_2 n t_n^2\}$ then using the standard deviation estimates (2.1) and (2.2) with $m = 1, \epsilon = \frac{1}{2}, \mu_1 = \epsilon_1 n t_n^2$ and $\epsilon_2 = n t_n^2$, we get that $\mathbb{P}_0(E_i) \geq 1 - \exp(-\frac{1}{16}\epsilon_1 n t_n^2) \geq 1 - \exp(-\frac{\epsilon_1 M \log n}{320})$ and so if $E_{poi} := \bigcap_{1 \leq i \leq W^2} E_i$ then using $W^2 = \frac{1}{t_n^2} \leq \frac{10}{r_n^2} \leq \frac{10n}{M \log n}$ (see bounds on r_n in the statement of the theorem) we get

$$\mathbb{P}_0(E_{poi}) \geq 1 - \frac{10Mn}{\log n} \cdot \exp\left(-\frac{\epsilon_1 M \log n}{320}\right) \geq 1 - \frac{1}{n^3} \quad (3.2)$$

for all n large, provided $M > \frac{1600}{\epsilon_1}$. Henceforth, we fix such an M .

If the event E_{poi} occurs, then each $t_n \times t_n$ square in $\{S_l\}$ contains a node and since $2t_n\sqrt{2} < r_n$, nodes in squares of $\{S_l\}$ sharing a corner are connected to each other by edges in the graph $G_n^{(P)}$. Thus $G_n^{(P)}$ is connected.


Lower bounds

We recall that we have divided the unit square S into $t_n \times t_n$ squares where $t_n = \frac{r_n}{2\sqrt{2} + \delta_n}$ and we had chosen $\delta_n \in [\sqrt{r_n}, 2\sqrt{r_n}]$ so that $W = \frac{1}{t_n}$ is an odd integer. For a real number $A > 0$, we now divide each $t_n \times t_n$ square $S_i, 1 \leq i \leq W^2 = \frac{1}{t_n^2}$ into smaller disjoint $\frac{A(n)}{\sqrt{n}} \times \frac{A(n)}{\sqrt{n}}$ squares $\{R_j\}$ where $A(n) \in \left[A + \frac{1}{(\log n)^{1/4}}, A + \frac{2}{(\log n)^{1/4}}\right)$ is chosen so that $L = \frac{t_n}{A(n)/\sqrt{n}}$ is an *odd* integer for all n large. This is possible since $n t_n^2 \geq \frac{\epsilon_1 M \log n}{20}$ (see (3.1)) and so

$$\begin{aligned} \frac{t_n \sqrt{n}}{A + \frac{1}{(\log n)^{1/4}}} - \frac{t_n \sqrt{n}}{A + \frac{2}{(\log n)^{1/4}}} &= \frac{t_n \sqrt{n}}{(\log n)^{1/4}} \left(\frac{1}{A + \frac{1}{(\log n)^{1/4}}} \right) \left(\frac{1}{A + \frac{2}{(\log n)^{1/4}}} \right) \\ &\geq \frac{t_n \sqrt{n}}{8A^2 (\log n)^{1/4}} \end{aligned}$$

which $\rightarrow \infty$ as $n \rightarrow \infty$. For notational simplicity we henceforth refer to $A(n)$ simply as A .

The total number of $\frac{A}{\sqrt{n}} \times \frac{A}{\sqrt{n}}$ squares $\{R_j\}$ obtained as above is $(WL)^2 = \frac{n}{A^2}$ and we label these squares in such a way that the squares R_j and $R_{j+1}, 1 \leq j \leq \frac{n}{A^2} - 1$, always share a common edge. This is possible since both W and L are odd integers and is illustrated in Figure 4 where the square labelled 1 is R_1 square labelled 2 is R_2 and so on.

Figure 4: Dividing each $t_n \times t_n$ square into smaller subsquares.

The number of nodes $N(R_i)$ in the square R_i , $1 \leq i \leq \frac{n}{A^2}$ is Poisson distributed with mean $\epsilon_1 A^2 \leq n \int_{R_i} f(x) dx \leq \epsilon_2 A^2$, using (1.1). Let $E(R_i)$ denote the event that R_i is occupied i.e., contains at least one node of \mathcal{P} , and all squares sharing a corner with R_i are empty. We remark that the event $E(R_i)$ is not restricted to subsquares within the same $t_n \times t_n$ square containing R_i . We use the particular structure of the tiling in Figure 4 in the next subsection for the upper bound. We now recall that if the event E_{poi} occurs, then the graph $G_n^{(P)}$ is connected (see discussion following (3.2)) and so if $E_{poi} \cap E(R_i)$ occurs, then there is an edge e in the MST of $G_n^{(P)}$ with one endvertex in R_i and the other endvertex in a square not sharing a corner with R_i . The edge length of e at least $\frac{A}{\sqrt{n}}$ and edge weight factor of e is at least ξ_{min} , implying that

$$MST_n^{(P)} \mathbb{1}(E_{poi}) \geq \frac{1}{2} \xi_{min} \sum_{i=1}^{\frac{n}{A^2}} \left(\frac{A}{\sqrt{n}} \right)^\alpha \mathbb{1}(E(R_i) \cap E_{poi}) \quad (3.3)$$

where the factor $\frac{1}{2}$ occurs, since each edge is counted at most twice in the summation in (3.3).

To estimate $H_\alpha := \sum_{i=1}^{\frac{n}{A^2}} \left(\frac{A}{\sqrt{n}} \right)^\alpha \mathbb{1}(E(R_i))$, we would like to split it into sums of independent r.v.s using the following construction. For a square R_i , let $\mathcal{N}(R_i)$ be the set of all squares sharing a corner with R_i , including R_i . If R_i does not intersect the sides of the unit square S , then there are 9 squares in $\mathcal{N}(R_i)$ and if R_j is another square such that $\mathcal{N}(R_i) \cap \mathcal{N}(R_j) = \emptyset$, then the corresponding events $E(R_i)$ and $E(R_j)$ are independent, by Poisson property. We now extract nine disjoint subsets $\{\mathcal{U}_l\}_{1 \leq l \leq 9}$ of $\{R_i\}$ with the following properties:

(A) If $R_i, R_j \in \mathcal{U}_l$, then $\#\mathcal{N}(R_i) = \#\mathcal{N}(R_j) = 9$ and $\mathcal{N}(R_i) \cap \mathcal{N}(R_j) = \emptyset$.

(B) The number of squares $\#\mathcal{U}_l \geq \frac{n}{9A^2} - \frac{4\sqrt{n}}{A}$ for each $1 \leq l \leq 9$.

This is possible since there are at most $\frac{4\sqrt{n}}{A} - 4 < \frac{4\sqrt{n}}{A}$ squares in $\{R_k\}$ intersecting the sides of the unit square S and the total number of squares in $\{R_k\}$ is $\frac{n}{A^2}$.

We now write $H_\alpha = \sum_{i=1}^{\frac{n}{A^2}} \mathbb{1}(E(R_i)) \geq \sum_{l=1}^9 \sum_{R_i \in \mathcal{U}_l} \mathbb{1}(E(R_i))$, where each inner summation on the right side is a sum of independent Bernoulli random variables, which we bound via standard deviation estimates. Indeed for $1 \leq l \leq 9$ and $R_i \in \mathcal{U}_l$, the number of nodes $N(R_i)$ is Poisson distributed with mean $n \int_{R_i} f(x) dx \in [\epsilon_1 A^2, \epsilon_2 A^2]$ (see (1.1)) and so R_i is occupied with probability at least $1 - e^{-\epsilon_1 A^2}$. Also each of the eight squares sharing a corner with R_i is empty with probability at least $e^{-\epsilon_2 A^2}$, implying that $\mathbb{P}(E(R_i)) \geq (1 - e^{-\epsilon_1 A^2})e^{-8\epsilon_2 A^2}$. Using the deviation estimate (2.2) with $\mu_1 = (1 - e^{-\epsilon_1 A^2})e^{-8\epsilon_2 A^2}$, $m = \frac{n}{9A^2} - \frac{4\sqrt{n}}{A}$ and $\epsilon = \frac{1}{m^{1/4}}$ we then get that

$$\mathbb{P}_0 \left(\sum_{R_i \in \mathcal{U}_l} \mathbb{1}(E(R_i)) \geq (1 - \epsilon) \left(\frac{n}{9A^2} - \frac{4\sqrt{n}}{A} \right) (1 - e^{-\epsilon_1 A^2})e^{-8\epsilon_2 A^2} \right) \geq 1 - e^{-D_1 \epsilon^2 n} \quad (3.4)$$

for some constant $D_1 > 0$ not depending on l . Since $m^{1/4} < \left(\frac{n}{9A^2} \right)^{1/4}$, we get that $D_1 \epsilon^2 n \geq 2D_2 \sqrt{n}$

for some constant $D_2 > 0$ and since $m^{1/4} > \left(\frac{n}{10A^2} \right)^{1/4}$ for all n large, we have

$$(1 - \epsilon) \left(\frac{n}{9A^2} - \frac{4\sqrt{n}}{A} \right) \geq \frac{n}{9A^2} - \frac{4\sqrt{n}}{A} - \frac{n}{A^2 m^{1/4}} \geq \frac{n}{9A^2} \left(1 - \frac{36\sqrt{A}}{n^{1/4}} \right)$$

for all n large.

Letting

$$E_{low} := \left\{ H_\alpha \geq (1 - e^{-\epsilon_1 A^2})e^{-8\epsilon_2 A^2} \frac{n}{A^2} \left(1 - \frac{36\sqrt{A}}{n^{1/4}} \right) \right\},$$

we get from (3.4) and a union bound that $\mathbb{P}_0(E_{low}) \geq 1 - 9e^{-2D_2 \sqrt{n}}$ and moreover, from (3.3) we also get that if $E_{low} \cap E_{poi}$ occurs, then

$$MST_n^{(P)} \geq \Delta_n := C_1(A) n^{1 - \frac{\alpha}{2}} \left(1 - \frac{36\sqrt{A}}{n^{1/4}} \right),$$

where $C_1(A)$ is as defined in (1.4). From the estimate for the probability of the event E_{low} above and the estimate for E_{poi} from (3.2), we therefore get

$$\mathbb{P}_0 \left(MST_n^{(P)} \geq \Delta_n \right) \geq 1 - \frac{1}{n^3} - 9e^{-2D_2 \sqrt{n}} \geq 1 - \frac{2}{n^3}$$

for all n large.

To convert the above estimate from Poisson to the Binomial process, we let $E_P := \{MST_n^{(P)} \geq \Delta_n\}$, $E := \{MST_n \geq \Delta_n\}$ and use the dePoissonization formula

$$\mathbb{P}(E) \geq 1 - D\sqrt{n}\mathbb{P}(E_P^c) \quad (3.5)$$

for some constant $D > 0$ to get that $\mathbb{P}(E) \geq 1 - 2D\frac{\sqrt{n}}{n^3} \geq 1 - \frac{1}{n^2}$ for all n large. This proves the bound in (1.6). Also

$$\mathbb{E}MST_n \geq \mathbb{E}MST_n \mathbb{1}(MST_n \geq \Delta_n) \geq \Delta_n \left(1 - \frac{1}{n^2}\right)$$

and using $\left(1 - \frac{36\sqrt{A}}{n^{1/4}}\right) \left(1 - \frac{1}{n^2}\right) \geq 1 - \frac{37\sqrt{A}}{n^{1/4}}$ for all n large, we get the lower bound on the expectation in (1.8).

To prove (3.5), we let N_P denote the random number of nodes of \mathcal{P} in all the squares $\{S_j\}$ so that $\mathbb{E}_0 N_P = n$ and $\mathbb{P}_0(N_P = n) = e^{-n\frac{n^n}{n!}} \geq \frac{D_1}{\sqrt{n}}$ for some constant $D_1 > 0$, using the Stirling formula. Given $N_P = n$, the nodes of \mathcal{P} are i.i.d. with distribution $f(\cdot)$ as defined in (1.1); i.e., $\mathbb{P}_0(E_P^c | N_P = n) = \mathbb{P}(E^c)$ and so

$$\mathbb{P}_0(E_P^c) \geq \mathbb{P}_0(E_P^c | N_P = n) \mathbb{P}_0(N_P = n) = \mathbb{P}(E^c) \mathbb{P}_0(N_P = n) \geq \mathbb{P}(E^c) \frac{D_1}{\sqrt{n}},$$

proving (3.5). ■

Upper bounds

As before, we let \mathcal{P} be the Poisson process with intensity $nf(\cdot)$ on the unit square, denote $G_n^{(P)}$ to be the RGG obtained from the nodes of \mathcal{P} with adjacency distance r_n and set $MST_n^{(P)}$ to be the weighted MST length of $G_n^{(P)}$ as defined in (1.3). The main idea for obtaining the upper bound for $MST_n^{(P)}$ is to connect all the nodes within a square R_j to form a subtree and then connect all these subtrees together to get an overall spanning tree. Suppose the event E_{poi} as defined prior to (3.2) occurs so that each $t_n \times t_n$ square in $\{S_l\}$ contains at least one node of the Poisson process \mathcal{P} and let $R_{i_1}, R_{i_2}, \dots, R_{i_Q}$ with $1 \leq i_1 < i_2 < \dots < i_Q \leq \frac{n}{A^2}$, $Q \leq \frac{n}{A^2}$ be the $\frac{A}{\sqrt{n}} \times \frac{A}{\sqrt{n}}$ subsquares containing all the nodes of \mathcal{P} .

For $1 \leq j \leq Q$, any two nodes in R_{i_j} are connected by an edge (of length at most $\frac{A\sqrt{2}}{\sqrt{n}}$) in the RGG $G_n^{(P)}$ since they belong to the same $t_n \times t_n$ square and $t_n\sqrt{2} < r_n$ (see definition of t_n in (2.3)). Therefore any spanning tree \mathcal{T}_{i_j} containing all the nodes of R_{i_j} has total weight of at most $N(R_{i_j}) \cdot \xi_{max} \cdot \left(\frac{A\sqrt{2}}{\sqrt{n}}\right)^\alpha$, where $N(R_i)$ is the number of points of the Poisson process \mathcal{P} in the square R_i . Moreover, since the event E_{poi} occurs, the squares R_{i_j} and $R_{i_{j+1}}$, $1 \leq j \leq Q - 1$ either belong to the same $t_n \times t_n$ square or belong to two different $t_n \times t_n$ squares sharing a common side. In any case, since $2t_n\sqrt{2} < r_n$, every node in R_{i_j} is connected to every node of $R_{i_{j+1}}$ by an edge of $G_n^{(P)}$ of length at most $\frac{2T_{j+1}A}{\sqrt{n}}$ where $T_{j+1} := i_{j+1} - i_j$. We pick one such edge and call it e_{j+1} . By definition the edge e_{j+1} has weight at most $\xi_{max} \cdot \left(\frac{2T_{j+1}A}{\sqrt{n}}\right)^\alpha$ and for future convenience we also set $T_1 := i_1 - 1$ and $T_{Q+1} := \frac{n}{A^2} - i_Q$.

The union $\mathcal{T}_{uni} := \cup_{1 \leq j \leq Q} \mathcal{T}_{i_j} \cup \cup_{2 \leq l \leq Q} \{e_l\}$ is a spanning tree of the graph G_n and has weight

$$W(\mathcal{T}_{uni}) \leq \sum_{j=1}^Q N(R_{i_j}) \cdot \xi_{max} \cdot \left(\frac{A\sqrt{2}}{\sqrt{n}}\right)^\alpha + \sum_{j=2}^Q \xi_{max} \cdot \left(\frac{2T_j A}{\sqrt{n}}\right)^\alpha$$

$$= \sum_{i=1}^{\frac{n}{A^2}} N(R_i) \cdot \xi_{max} \cdot \left(\frac{A\sqrt{2}}{\sqrt{n}} \right)^\alpha + \sum_{j=2}^Q \xi_{max} \cdot \left(\frac{2T_j A}{\sqrt{n}} \right)^\alpha$$

since $\{R_{i_j}\}_{1 \leq j \leq Q}$ are all the squares containing at least one node of \mathcal{P} . Letting $Y_\alpha := \sum_{j=1}^{Q+1} T_j^\alpha$ we then get

$$MST_n^{(P)} \mathbb{1}(E_{poi}) \leq \xi_{max} \left(\frac{2A}{\sqrt{n}} \right)^\alpha \left(\sum_{i=1}^{\frac{n}{A^2}} N(R_i) + Y_\alpha \right). \quad (3.6)$$

The first term $\sum_{i=1}^{\frac{n}{A^2}} N(R_i)$ is a Poisson random variable with mean n since this denotes the total number of nodes of the Poisson process in the unit square. From the deviation estimate (2.1) with $m = 1$, $\mu_2 = n$ and $\epsilon = \frac{\log n}{\sqrt{n}}$, we have that

$$\mathbb{P}_0 \left(\sum_{i=1}^{\frac{n}{A^2}} N(R_i) \leq n \left(1 + \frac{\log n}{\sqrt{n}} \right) \right) \geq 1 - e^{-C(\log n)^2} \quad (3.7)$$

for some constant $C > 0$, using (1.10). Setting $E_{node} := \left\{ \sum_{i=1}^{\frac{n}{A^2}} N(R_i) \leq n \left(1 + \frac{\log n}{\sqrt{n}} \right) \right\}$, we get from (3.6) that

$$MST_n \mathbb{1}(E_{poi} \cap E_{node}) \leq \xi_{max} \left(\frac{2A}{\sqrt{n}} \right)^\alpha \left(n \left(1 + \frac{\log n}{\sqrt{n}} \right) + Y_\alpha \right). \quad (3.8)$$

To evaluate the second term Y_α in (3.8) (which is not an i.i.d. sum), we consider two different cases depending on whether $\alpha \leq 1$ or $\alpha > 1$. The term Y_α is well-defined for any configuration ω of the Poisson process provided we set $Y_\alpha(\omega_0) = \left(\frac{n}{A^2} - 1 \right)^\alpha$ for the configuration ω_0 that contains no node of the Poisson process. We now show that $Y_\alpha(\omega)$ is monotonic in ω in the sense that adding more points increases Y_α if $\alpha \leq 1$ and decreases Y_α if $\alpha > 1$. This then allows us to use coupling and upper bound Y_α by simply considering homogenous Poisson processes.

Monotonicity of Y_α : For a configuration $\omega \neq \omega_0$ let $1 \leq i_1(\omega) < \dots < i_Q(\omega) \leq \frac{n}{A^2}$ be the indices of the squares in $\{R_j\}$ containing at least one node of the Poisson process \mathcal{P} . Letting $i_0(\omega) = 1$ and $i_{Q+1}(\omega) = \frac{n}{A^2}$ we have $Y_\alpha(\omega) = \sum_{j=0}^Q (i_{j+1}(\omega) - i_j(\omega))^\alpha$. Suppose $\omega' = \omega \cup \{x\}$ is obtained by adding a single extra node at $x \in R_{j_0}$ for some $1 \leq j_0 \leq \frac{n}{A^2}$. If $j_0 \in \{i_k(\omega)\}_{0 \leq k \leq Q+1}$, then $Y_\alpha(\omega') = Y_\alpha(\omega)$. Else there exists $0 \leq a \leq Q$ such that $i_a(\omega) < j_0 < i_{a+1}(\omega)$ and so

$$Y_\alpha(\omega') = Y_\alpha(\omega) + (i_{a+1}(\omega) - j_0)^\alpha + (j_0 - i_a(\omega))^\alpha - (i_{a+1}(\omega) - i_a(\omega))^\alpha.$$

If $\alpha \leq 1$ then using $a^\alpha + b^\alpha \geq (a+b)^\alpha$ for positive numbers a, b we get that $Y_\alpha(\omega') \geq Y_\alpha(\omega)$. If $\alpha > 1$ then $a^\alpha + b^\alpha \leq (a+b)^\alpha$ and so $Y_\alpha(\omega') \leq Y_\alpha(\omega)$. This monotonicity property together with coupling allows us to upper bound Y_α as follows. Letting $\delta = \epsilon_2$ if $\alpha \leq 1$ and $\delta = \epsilon_1$ if $\alpha > 1$, we let \mathcal{P}_δ be a homogenous Poisson process of intensity δn on the unit square S , defined on the probability space $(\Omega_\delta, \mathcal{F}_\delta, \mathbb{P}_\delta)$.

Let F_δ denote the event that there is at least one node of \mathcal{P}_δ in the unit square S and set $Y_\alpha^{(\delta)} := \left(\frac{n}{A^2} - 1\right)^\alpha$ if there is no node of \mathcal{P}_δ in S . Suppose now that F_δ occurs and as before let $\{i_j^{(\delta)}\}_{1 \leq j \leq Q_\delta}$ be the indices of the squares in $\{R_j\}$ containing at least one node of \mathcal{P}_δ . Moreover, let $T_{j+1}^{(\delta)} := i_{j+1}^{(\delta)} - i_j^{(\delta)}$ for $1 \leq j \leq Q_\delta$ and set $T_1^{(\delta)} := i_1^{(\delta)} - 1$ and $T_{Q_\delta+1}^{(\delta)} := \frac{n}{A^2} - i_{Q_\delta}^{(\delta)}$. Defining $Y_\alpha^{(\delta)} := \sum_{j=1}^{Q_\delta+1} \left(T_j^{(\delta)}\right)^\alpha$, we now show for any $x > 0$ that

$$\mathbb{P}_\delta \left(Y_\alpha^{(\delta)} < x \right) \leq \mathbb{P}_0 (Y_\alpha < x). \quad (3.9)$$

Proof of (3.9): For $\alpha \leq 1$ we couple the original Poisson process \mathcal{P} and the homogenous process \mathcal{P}_δ in the following way. Let $V_i, i \geq 1$ be i.i.d. random variables each with density $f(\cdot)$ and let N_V be a Poisson random variable with mean n , independent of $\{V_i\}$. The nodes $\{V_i\}_{1 \leq i \leq N_V}$ form a Poisson process with intensity $nf(\cdot)$ which we denote as \mathcal{P} and colour green.

Let $U_i, i \geq 1$ be i.i.d. random variables each with density $\epsilon_2 - f(\cdot)$ where $\epsilon_2 \geq 1$ is as in (1.1) and let N_U be a Poisson random variable with mean $n(\epsilon_2 - 1)$. The random variables $(\{U_i\}, N_U)$ are independent of $(\{V_i\}, N_V)$ and the nodes $\{U_i\}_{1 \leq i \leq N_U}$ form a Poisson process with intensity $n(\epsilon_2 - f(\cdot))$ which we denote as \mathcal{P}_{ext} and colour red. The nodes of \mathcal{P} and \mathcal{P}_{ext} together form a homogenous Poisson process with intensity $n\epsilon_2$, which we denote as \mathcal{P}_δ and define it on the probability space $(\Omega_\delta, \mathcal{F}_\delta, \mathbb{P}_\delta)$.

Let $\omega_\delta \in \Omega_\delta$ be any configuration and as above let $\{i_j^{(\delta)}\}_{1 \leq j \leq Q_\delta}$ be the indices of the squares in $\{R_j\}$ containing at least one node of \mathcal{P}_δ and let $\{i_j\}_{1 \leq j \leq Q}$ be the indices of the squares in $\{R_j\}$ containing at least one node of \mathcal{P} . The indices in $\{i_j^{(\delta)}\}$ and $\{i_j\}$ depend on ω_δ . Defining $Y_\alpha = Y_\alpha(\omega_\delta)$ and $Y_\alpha^{(\delta)} = Y_\alpha^{(\delta)}(\omega_\delta)$ as before, we have that Y_α is determined only by the green nodes of ω_δ while $Y_\alpha^{(\delta)}$ is determined by both green and red nodes of ω_δ .

From the monotonicity property, we therefore have that $Y_\alpha(\omega_\delta) \leq Y_\alpha^{(\delta)}(\omega_\delta)$ and so for any $x > 0$ we have

$$\mathbb{P}_\delta(Y_\alpha^{(\delta)} < x) \leq \mathbb{P}_\delta(Y_\alpha < x) = \mathbb{P}_0(Y_\alpha < x), \quad (3.10)$$

proving (3.9).

If $\alpha > 1$, we perform a slightly different analysis. Letting $\epsilon_1 \leq 1$ be as in (1.1), we construct a Poisson process \mathcal{P}_{ext} with intensity $n(f(\cdot) - \epsilon_1)$ and colour nodes of \mathcal{P}_{ext} red. Letting \mathcal{P}_δ be another independent Poisson process with intensity $n\epsilon_1$, we colour nodes of \mathcal{P}_δ green. The superposition of \mathcal{P}_{ext} and \mathcal{P}_δ is a Poisson process with intensity $nf(\cdot)$, which we define on the probability space $(\Omega_\delta, \mathcal{F}_\delta, \mathbb{P}_\delta)$. In this case, the sum Y_α is determined by both green and red nodes while $Y_\alpha^{(\delta)}$ is determined only by the green nodes. Again using the monotonicity property of Y_α , we get (3.10). ■

To estimate $Y_\alpha^{(\delta)}$ we let $N^{(\delta)}(R_i), 1 \leq i \leq \frac{n}{A^2}$, be the random number of nodes of \mathcal{P}_δ in the square R_i . The random variables $\{N^{(\delta)}(R_i)\}$ are i.i.d. Poisson distributed each with mean $A^2\delta$. For $i \geq \frac{n}{A^2} + 1$, we define $N^{(\delta)}(R_i)$ to be i.i.d. Poisson random variables with mean $A^2\delta$, that are also independent of $\{N^{(\delta)}(R_i)\}_{1 \leq i \leq \frac{n}{A^2}}$. Without loss of generality, we associate the probability measure \mathbb{P}_δ for the random variables $\{N^{(\delta)}(R_i)\}_{i \geq \frac{n}{A^2} + 1}$ as well.

Let $\tilde{T}_1 := \min\{j \geq 1 : N^{(\delta)}(R_j) \geq 1\}$ and for $j \geq 2$, let

$$\tilde{T}_j := \min\{k \geq \tilde{T}_{j-1} + 1 : N^{(\delta)}(R_k) \geq 1\} - \tilde{T}_{j-1}.$$

The random variables $\{\tilde{T}_i\}$ are nearly the same as $\{T_i^{(\delta)}\}$ in the following sense: Suppose the event F_δ occurs so that there is at least one node of \mathcal{P}_δ in the unit square. This means that $1 \leq Q_\delta \leq \frac{n}{A^2}$ and so $T_1^{(\delta)} = i_1 - 1 = \tilde{T}_1 - 1, T_j^{(\delta)} = \tilde{T}_j$ for $2 \leq j \leq Q_\delta$ and $T_{Q_\delta+1}^{(\delta)} \leq \tilde{T}_{Q_\delta+1}$. Consequently

$$Y_\alpha^{(\delta)} \mathbb{1}(F_\delta) \leq \sum_{i=1}^{Q_\delta+1} \tilde{T}_i^\alpha \mathbb{1}(F_\delta) \leq \sum_{i=1}^{\frac{n}{A^2}+1} \tilde{T}_i^\alpha \mathbb{1}(F_\delta) \leq \sum_{i=1}^{\frac{n}{A^2}+1} \tilde{T}_i^\alpha, \quad (3.11)$$

since $Q_\delta \leq \frac{n}{A^2}$.

The advantage of the above construction is that $\{\tilde{T}_i\}$ are i.i.d. geometric random variables with success parameter $p = 1 - e^{-A^2\delta}$ and so all moments of \tilde{T}_i^α exist. Letting $\beta_i = (\tilde{T}_i^\alpha - \mathbb{E}_\delta \tilde{T}_i^\alpha)$ and $\beta_{tot} = \sum_{i=1}^{\frac{n}{A^2}+1} \beta_i$ we then have that

$$\mathbb{E}_\delta \beta_{tot}^4 = \sum_i \mathbb{E}_\delta \beta_i^4 + \sum_{i < j} \mathbb{E}_\delta \beta_i^2 \beta_j^2 \leq Cn^2$$

for some constant $C > 0$, by the relation (3.1) in Appendix. For $\epsilon > 0$ we therefore get from Chebyshev's inequality that

$$\mathbb{P}_\delta \left(|\beta_{tot}| > \epsilon \left(\frac{n}{A^2} + 1 \right) \mathbb{E}_\delta \tilde{T}_1^\alpha \right) \leq D_1 \frac{\mathbb{E}_\delta (|\beta_{tot}|^4)}{n^4 \epsilon^4} \leq \frac{D_2}{n^2 \epsilon^4}$$

for some constants $D_1, D_2 > 0$. Setting $\epsilon = \frac{1}{n^{1/16}}$ we get

$$\mathbb{P}_\delta \left(\sum_{i=1}^{\frac{n}{A^2}+1} \tilde{T}_i^\alpha \leq \left(1 + \frac{1}{n^{1/16}} \right) \frac{n}{A^2} \mathbb{E}_\delta \tilde{T}_1^\alpha \right) \geq 1 - \frac{D_2}{n^{7/4}} \quad (3.12)$$

From the upper bound for $Y_\alpha^{(\delta)}$ in (3.11) and the fact that there is at least one node of \mathcal{P}_δ in the unit square S with probability $1 - e^{-\delta n}$ we further get

$$\mathbb{P}_\delta \left(Y_\alpha^{(\delta)} \leq \left(1 + \frac{1}{n^{1/16}} \right) \frac{n}{A^2} \mathbb{E}_\delta \tilde{T}_1^\alpha \right) \geq 1 - \frac{D_2}{n^{7/4}} - e^{-\delta n} \geq 1 - \frac{D_3}{n^{7/4}}$$

for all n large and some constant $D_3 > 0$. Denoting $\mathbb{E}_\delta \tilde{T}_1^\alpha$ as simply $\mathbb{E} \tilde{T}_1^\alpha$ and using the coupling relation (3.9) we finally get that

$$\mathbb{P}_0 \left(Y_\alpha \leq \left(1 + \frac{1}{n^{1/16}} \right) \frac{n}{A^2} \mathbb{E} \tilde{T}_1^\alpha \right) \geq 1 - \frac{D_3}{n^{7/4}}. \quad (3.13)$$

Letting E_{til} be the event on the left side of (3.13) we have from (3.8) that the term $MST_n^{(P)} \mathbb{1}(E_{poi} \cap E_{node} \cap E_{til})$ is bounded above by

$$\xi_{max} \left(\frac{2A}{\sqrt{n}} \right)^\alpha \left(n \left(1 + \frac{\log n}{\sqrt{n}} \right) + \left(1 + \frac{1}{n^{1/16}} \right) \frac{n}{A^2} \mathbb{E} \tilde{T}_1^\alpha \right) = C_2(A) n^{1-\frac{\alpha}{2}} + b n^{-\frac{\alpha}{2}}, \quad (3.14)$$

where $C_2(A)$ is as in (1.4) and

$$b = \xi_{max}(2A)^\alpha \left(\log n \cdot \sqrt{n} + \left(\frac{1}{A^2} \mathbb{E} \tilde{T}_1^\alpha \right) n^{15/16} \right) \leq n^{16/17}$$

for all n large, again using (3.1) from Appendix. From (3.14) and the estimates for the events E_{poi} , E_{node} and E_{til} from (3.2), (3.7) and (3.13), respectively, we then have

$$\mathbb{P}_0 \left(MST_n^{(P)} \leq C_2(A) n^{1-\frac{\alpha}{2}} + n^{\frac{16}{17}-\frac{\alpha}{2}} \right) \geq 1 - \frac{1}{n^3} - e^{-D_4(\log n)^2} - \frac{D_3}{n^{7/4}} \geq 1 - \frac{D_5}{n^{7/4}} \quad (3.15)$$

for all n large and some constants $D_4, D_5 > 0$.

Proof of (1.7): From (3.15) and the dePoissonization formula (3.5), we obtain

$$\mathbb{P} \left(MST_n \leq C_2(A) n^{1-\frac{\alpha}{2}} + n^{\frac{16}{17}-\frac{\alpha}{2}} \right) \geq 1 - \frac{D_6 \sqrt{n}}{n^{7/4}} = 1 - \frac{D_6}{n^{5/4}} \quad (3.16)$$

proving the estimate in (1.7). For bounding the expectation, we let

$$\Delta_n := C_2(A) n^{1-\frac{\alpha}{2}} + n^{\frac{16}{17}-\frac{\alpha}{2}}$$

and write

$$\begin{aligned} \mathbb{E} MST_n &= \mathbb{E} MST_n \mathbb{1}(MST_n \leq \Delta_n) + \mathbb{E} MST_n \mathbb{1}(MST_n > \Delta_n) \\ &\leq \Delta_n + \mathbb{E} MST_n \mathbb{1}(MST_n > \Delta_n) \end{aligned} \quad (3.17)$$

For the second term in (3.17), we use the estimate $MST_n \leq n \cdot \xi_{max} \cdot r_n^\alpha \leq \xi_{max} n$ since each edge has length at most $r_n \leq 1$ (see bounds on r_n in the statement of the Theorem) and weight at most ξ_{max} . Using the probability estimate (3.16), we then get

$$\mathbb{E} MST_n \leq \Delta_n + \frac{D_6 \xi_{max} n}{n^{5/4}} \leq C_2(A) n^{1-\frac{\alpha}{2}} + 2n^{\frac{16}{17}-\frac{\alpha}{2}}$$

for all n large. This proves the expectation upper bound in (1.8). \blacksquare

Appendix

By the continuity of the function $C_1(\cdot)$ defined in (1.4), we have that $C_1(A_n) \rightarrow C_1(A)$ as $n \rightarrow \infty$. To prove (1.10) for $i = 2$ we let T_n be a geometric random variable with success probability $p_n = 1 - q_n = 1 - e^{-\delta A_n^2}$ and show that

$$\lim_n \mathbb{E} T_n^\alpha = \mathbb{E} T^\alpha \quad (3.1)$$

where T is a geometric random variable with success probability $p = 1 - q = 1 - e^{-\delta A^2}$. Indeed letting $L \geq 1$ be fixed, we write

$$\mathbb{E} T_n^\alpha = \sum_{k \geq 1} k^\alpha e^{-\delta A_n^2(k-1)} (1 - e^{-\delta A_n^2}) = Y_n(L) + Z_n(L)$$

and $\mathbb{E}T^\alpha = Y(L) + Z(L)$ where

$$Y_n(L) := \sum_{1 \leq k \leq L} k^\alpha e^{-\delta A_n^2(k-1)} (1 - e^{-\delta A_n^2}) \longrightarrow \sum_{1 \leq k \leq L} k^\alpha e^{-\delta A^2(k-1)} (1 - e^{-\delta A^2}) =: Y(L) \quad (3.2)$$

as $n \rightarrow \infty$ and

$$Z_n(L) := \sum_{k \geq L+1} k^\alpha e^{-\delta A_n^2(k-1)} (1 - e^{-\delta A_n^2}) \leq \sum_{k \geq L+1} k^\alpha e^{-\delta A_n^2(k-1)}.$$

Using $A_n \geq A$ we further have for $\epsilon > 0$ that

$$Z_n(L) \leq \sum_{k \geq L+1} k^\alpha e^{-\delta A^2(k-1)} \leq \epsilon \quad (3.3)$$

if $L = L(\epsilon, \delta, \alpha, A)$ is large. For the same choice of L , we argue as above to get that

$$Z(L) := \sum_{k \geq L+1} k^\alpha e^{-\delta A^2(k-1)} (1 - e^{-\delta A^2}) \leq \epsilon. \quad (3.4)$$

Finally, writing

$$|\mathbb{E}T_n^\alpha - \mathbb{E}T^\alpha| \leq |Y_n(L) - Y(L)| + |Z_n(L) - Z(L)| \leq |Y_n(L) - Y(L)| + Z_n(L) + Z(L)$$

and using (3.2), (3.3) and (3.4), we get for all n large that $|\mathbb{E}T_n^\alpha - \mathbb{E}T^\alpha| \leq 3\epsilon$. Since $\epsilon > 0$ is arbitrary, this proves (1.10) for $i = 2$. \blacksquare

Acknowledgement

I thank Professors Rahul Roy, Thomas Mountford, Federico Camia, C. R. Subramanian and the referee for crucial comments that led to an improvement of the paper. I also thank IMSc for my fellowships.

References

- [1] Alon, N. and Spencer, J. (2008) *The probabilistic method* Wiley interscience, 1st edition.
- [2] Beardwood, J., Halton, J. H. and Hammersley, J. M. (1959), The shortest path through many points, *Proceedings Cambridge Philosophical Society*, **55**, 299–327.
- [3] Djahhari, M. and Gan, S. (2015), Optimality problem of network topology in stocks market analysis, *Physica A: Statistical Mechanics and Its Applications*, **419**, 108–114.
- [4] Gupta, P. and Kumar, P. R. (1998), Critical Power for Asymptotic Connectivity in Wireless Networks, *Stochastic Analysis, Control, Optimization and Applications*, 2203–2214.
- [5] Jothi, R. and Raghavachari, B. (2005), Approximation algorithms for the capacitated minimum spanning tree problem and its variants in network design, *ACM Transactions on Algorithms*, **1(2)**, 265–282.
- [6] Kesten, H. and Lee, S. (1996), The central limit theorem for weighted minimal spanning trees on random points, *Annals of Applied Probability*, **6**, pp. 495–527.
- [7] Penrose, M. (2003), *Random Geometric Graphs*, Oxford University Press.

- [8] Penrose, M. (2016), Connectivity of soft random geometric graphs, *Annals of Applied Probability*, **26**, 986–1028.
- [9] Penrose, M. and Yukich, J. (2003), Weak laws of large numbers in geometric probability, *Annals of Applied Probability*, **13**, 277–303.
- [10] Steele, J. M. (1988), Growth rates of Euclidean minimal spanning trees with power weighted edges, *Annals of Probability*, **16**, 1767–1787.
- [11] Steele, J. M. (1993), Probability and Problems in Euclidean Combinatorial Optimization, *Statistical Science*, **8**, pp. 48–56.
- [12] Yukich, J. (1998), *Probability Theory of Classical Euclidean Optimization Problems*, Lecture Notes in Mathematics, **1675**, Springer.
- [13] Yukich, J. (2000), Asymptotics for weighted minimal spanning trees on random points, *Stochastic Processes and Applications*, **85**, pp. 123-138.