

# General Bayesian $L^2$ calibration of mathematical models

Antony M. Overstall

School of Mathematical Sciences, University of Southampton,  
Southampton SO17 1BJ, U.K., A.M.Overstall@soton.ac.uk

and

James M. McGree

School of Mathematical Sciences, Queensland University of Technology,  
Brisbane 4001, Australia, james.mcgree@qut.edu.au

June 30, 2021

## Abstract

A general Bayesian method for  $L^2$  calibration of a mathematical model is presented. General Bayesian inference starts with the specification of a loss function. Then, the log-likelihood in Bayes' theorem is replaced by the negative loss. While the minimiser of the loss function is unchanged by, for example, multiplying the loss by a constant, the same is not true of the resulting general posterior distribution. To address this problem in the context of  $L^2$  calibration of mathematical models, different automatic scalings of the general Bayesian posterior are proposed. These are based on equating asymptotic properties of the general Bayesian posterior and the minimiser of the  $L^2$  loss.

*Keywords:* loss functions, composite likelihood, curvature scaling, magnitude scaling

# 1 Introduction

A mathematical model is a representation of a physical system often underpinned by scientific theory. Mathematical models are used to understand, predict and control physical systems. When such models are evaluated by complex, computationally expensive code, they are known as computer models.

A mathematical model is considered to be a function,  $\eta(\mathbf{x}, \boldsymbol{\theta})$ , taking arguments and returning a theoretical prediction of a feature (or features) of the physical system. Following Plumlee (2017), the arguments  $\mathbf{x}$  and  $\boldsymbol{\theta}$  can be split into two groups. They are (a) *general inputs*: controllable or measurable variables of the system denoted  $\mathbf{x} = (x_1, \dots, x_k)^\top \in \mathcal{X} \subset \mathbb{R}^k$ ; and (b) *calibration parameters*: unknown attributes of the physical system that cannot be controlled or directly measured denoted  $\boldsymbol{\theta} = (\theta_1, \dots, \theta_p)^\top \in \Theta \subset \mathbb{R}^p$ .

This paper addresses *calibration*: the task of inferring values of the calibration parameters from observations of the physical system. That is, for  $i = 1, \dots, n$ , a response  $y_i$  is observed of the physical system under inputs  $\mathbf{x}_i$ . Calibration aims to estimate values for the calibration parameters using  $\{\mathbf{x}_i, y_i\}_{i=1}^n$ . It is assumed that the true data-generating process for  $\mathbf{y} = (y_1, \dots, y_n)^\top$  is given by

$$y_i = \mu(\mathbf{x}_i) + e_i, \tag{1}$$

where  $\mu(\mathbf{x})$  is a function giving the true value of the physical system at input  $\mathbf{x}$ , and  $e_1, \dots, e_n$  are independent and identically distributed random variables, with  $E(e_i) = 0$  and  $\text{var}(e_i) = \sigma^2 > 0$ , representing observational error. The principle challenge of calibration is that the mathematical model is an incomplete (or inexact) representation of the physical system, i.e. there do not exist values of the calibration parameters  $\boldsymbol{\theta}$  such that  $\eta(\mathbf{x}, \boldsymbol{\theta}) = \mu(\mathbf{x})$  for all values of  $\mathbf{x} \in \mathcal{X}$ . Thus calibration refers to the search for the values of  $\boldsymbol{\theta}$  that result in the mathematical model  $\eta(\mathbf{x}, \boldsymbol{\theta})$  being as “close” to the physical system  $\mu(\mathbf{x})$  as possible.

In their seminal work, Kennedy and O’Hagan (2001) proposed a calibration framework under which a Gaussian process prior is assumed for the bias function,  $\delta_{\boldsymbol{\theta}}(\mathbf{x}) = \mu(\mathbf{x}) - \eta(\mathbf{x}, \boldsymbol{\theta})$ : the difference between the physical system and the mathematical model. However, the Kennedy and O’Hagan framework does not imply a unique definition of the values of the calibration parameters, i.e. in the frequentist sense, the resulting model is non-identifiable (e.g. Tuo and Wu, 2016; Wong et al., 2017). Under Kennedy and O’Hagan’s Bayesian

approach, this problem is alleviated by informative prior distributions for the calibration parameters and the bias function. However, the influence of the prior distribution does not diminish as the number of observations of the physical system increases.

To address this problem, Tuo and Wu (2015) proposed an alternative frequentist  $L^2$  calibration approach. In this case, let  $\boldsymbol{\theta}_{L^2}$  denote the values of the calibration parameters that minimise the squared norm (in the associated  $L^2$  space) of the bias function. Correspondingly, Tuo and Wu (2015) defined an estimator given by the values of the calibration parameters that minimise a loss function given by the  $L^2$  norm of the difference between a non-parametric estimator of  $\mu(\mathbf{x})$  and the mathematical model  $\eta(\mathbf{x}, \boldsymbol{\theta})$ . Tuo and Wu (2015) showed that this estimator is consistent for  $\boldsymbol{\theta}_{L^2}$ . Alternatively, Wong et al. (2017) showed that the ordinary least squares estimator of the calibration parameters is also consistent for  $\boldsymbol{\theta}_{L^2}$ . Furthermore, for uncertainty quantification, Wong et al. (2017) proposed a bootstrapping procedure to provide confidence intervals for the calibration parameters.

Due to the natural way that Bayesian inference deals with uncertainty quantification, several authors have recently proposed Bayesian analogues of  $L^2$  calibration. Plumlee (2017) noted that minimising the  $L^2$  norm imposes orthogonality of the bias function and the gradient of the mathematical model. Under the Kennedy and O’Hagan framework, this orthogonality places constraints on the Gaussian process prior for the bias function. Gu and Wang (2018) pointed out that orthogonality of the bias and mathematical model gradient is a necessary, but not sufficient condition, for minimising the  $L^2$  norm. Instead, they assumed a prior distribution on the  $L^2$  norm penalising “large” bias functions. Xie and Xu (2020) proposed a different approach, not under the Kennedy and O’Hagan framework. They assumed a Gaussian process prior for the physical system,  $\mu(\mathbf{x})$ . The resulting posterior distribution for the  $L^2$  norm induces a posterior distribution for  $\boldsymbol{\theta}_{L^2}$ .

This paper considers a novel general Bayesian framework for  $L^2$  calibration of mathematical models. Briefly, general Bayesian inference begins with specification of a loss function, denoted  $\ell(\boldsymbol{\theta}; \mathbf{y})$ , identifying desirable values for the calibration parameters  $\boldsymbol{\theta}$ , in light of observed responses  $\mathbf{y}$ . General Bayesian inference proceeds via the *generalised posterior distribution* given by

$$\pi_G(\boldsymbol{\theta}|\mathbf{y}) \propto \exp(-n\ell(\boldsymbol{\theta}; \mathbf{y})) \pi(\boldsymbol{\theta}). \quad (2)$$

In (2),  $\pi(\boldsymbol{\theta})$  is the probability density function of the prior distribution for the calibration parameters, and  $\exp(-n\ell(\boldsymbol{\theta}; \mathbf{y}))$  is known as the *generalised likelihood*. Bissiri et al. (2016) show that the generalised posterior distribution provides a coherent representation

of subjective uncertainty about the values of the calibration parameters that minimise the expectation of the loss with respect to the true data-generating process of the responses.

In this paper, we focus on generalised Bayesian calibration of mathematical models using the Tuo & Wu loss, i.e. general Bayesian  $L^2$  calibration. Specifically, we consider loss functions given by functionals of the Tuo & Wu loss, the simplest being

$$\ell_{TW}^\gamma(\boldsymbol{\theta}, \mathbf{y}) = \gamma \ell_{TW}(\boldsymbol{\theta}, \mathbf{y}) \tag{3}$$

for a specified constant  $\gamma > 0$ .

The advantages of the resulting general Bayesian framework, when compared with existing Bayesian approaches to calibration described above, are that it is conceptually and computationally simpler, and allows more transparent incorporation of prior information.

However the major impediment to applying general Bayesian  $L^2$  calibration is that the scale of the general posterior of  $\boldsymbol{\theta}$  is essentially arbitrary. For example, consider the loss given by (3). For any  $\gamma > 0$ , the minimiser of the function  $\ell_{TW}^\gamma(\boldsymbol{\theta}, \mathbf{y}) = \gamma \ell_{TW}(\boldsymbol{\theta}, \mathbf{y})$  remains unchanged yet the same is not true of the generalised posterior.

In this paper, to address this problem, automatic adjustments of the loss are developed based on the asymptotic behaviour of the resulting generalised posterior distribution. Woody et al. (2019) considered this loss and specified  $\gamma$  to maintain frequentist coverage properties of credible sets of  $\boldsymbol{\theta}$ . They devise a computational approach to achieve this using a bootstrapping procedure. We consider more general forms of the functional of the Tuo & Wu loss but choose the exact form to maintain certain asymptotical properties, i.e. to equate i) the expectation of a proxy of the likelihood ratio statistic to  $p$  (the number of parameters), or ii) the generalised posterior variance to the frequentist variance of  $\hat{\boldsymbol{\theta}}_{TW}$ .

The remainder of this paper is organised as follows. Section 2 provides a more in depth description of existing approaches to calibration. In Section 3, the new framework is presented including results on the asymptotic behaviour of the generalised posterior and loss function adjustment. In Section 4, the methodology is demonstrated on several synthetic examples.

## 2 Existing approaches to calibration

### 2.1 Kennedy and O’Hagan calibration

Under the Kennedy and O’Hagan (2001) framework it is assumed that the observational errors  $(e_1, \dots, e_n)$  have a normal distribution and

$$\mu(\mathbf{x}) = \eta(\boldsymbol{\theta}, \mathbf{x}) + \delta_{\boldsymbol{\theta}}(\mathbf{x}).$$

With the subscript on the bias function, we follow Plumlee (2017) by making explicit that this function depends on  $\boldsymbol{\theta}$ . Kennedy and O’Hagan imposed a zero mean Gaussian process prior on the unknown bias function, i.e.  $\delta_{\boldsymbol{\theta}}(\cdot) \sim \text{GP}(0, \tau^2 \kappa(\cdot, \cdot; \boldsymbol{\rho}))$  where  $\kappa(\cdot, \cdot; \boldsymbol{\rho})$  is a specified correlation function depending on an  $r \times 1$  vector of parameters  $\boldsymbol{\rho}$ . A fully Bayesian approach was taken by evaluating the marginal posterior distribution of  $\boldsymbol{\theta}$  after specifying prior distributions for  $\boldsymbol{\theta}$ ,  $\sigma^2$ ,  $\boldsymbol{\rho}$  and  $\tau^2$ .

Tuo and Wu (2016) showed that, as  $n \rightarrow \infty$ , the maximum likelihood estimators,  $\hat{\boldsymbol{\theta}}_{KO}$ , of  $\boldsymbol{\theta}$  under the Kennedy and O’Hagan model converge to  $\boldsymbol{\theta}_{KO}^*$ ; the values of  $\boldsymbol{\theta}$  that minimise the norm of  $\delta_{\boldsymbol{\theta}}(\mathbf{x})$  in the associated reproducing kernel Hilbert space norm with kernel  $\kappa$ . Under a Bayesian approach the influence of the prior distributions will diminish as  $n$  grows meaning the posterior distribution will converge to a point mass at  $\boldsymbol{\theta}_{KO}^*$ . The convergence to quantities which depend on the choice of correlation function has been criticised by several authors (e.g. Tuo and Wu, 2016, 2015; Plumlee, 2017; Wong et al., 2017).

### 2.2 $L^2$ calibration

To address the asymptotic correlation function dependency of the Kennedy and O’Hagan framework, Tuo and Wu (2015) defined the “best” values of the calibration parameters to be those that minimise the squared norm of the bias function in the associated  $L^2$  space. That is

$$\boldsymbol{\theta}_{L^2} = \arg \min_{\boldsymbol{\theta} \in \Theta} L_{L^2}(\boldsymbol{\theta}),$$

where

$$\begin{aligned} L_{L^2}(\boldsymbol{\theta}) &= \int_{\mathcal{X}} \delta_{\boldsymbol{\theta}}(\mathbf{x})^2 d\mathbf{x}, \\ &= \int_{\mathcal{X}} (\mu(\mathbf{x}) - \eta(\boldsymbol{\theta}, \mathbf{x}))^2 d\mathbf{x}. \end{aligned} \quad (4)$$

They then define the estimates  $\hat{\boldsymbol{\theta}}_{TW}$  to be the values of  $\boldsymbol{\theta}$  that minimise a loss given by squared  $L^2$  norm of a non-parametric estimate of the bias function, i.e.

$$\hat{\boldsymbol{\theta}}_{TW} = \arg \min_{\boldsymbol{\theta} \in \Theta} \ell_{TW}(\boldsymbol{\theta}; \mathbf{y}),$$

where

$$\ell_{TW}(\boldsymbol{\theta}; \mathbf{y}) = \int_{\mathcal{X}} (\hat{\mu}(\mathbf{x}) - \eta(\boldsymbol{\theta}, \mathbf{x}))^2 d\mathbf{x}. \quad (5)$$

In (5),  $\hat{\mu}(\mathbf{x})$  is a non-parametric regression estimate of  $\mu(\mathbf{x})$  constructed using the observed data  $\{\mathbf{x}_i, y_i\}_{i=1}^n$ . Specifically,

$$\hat{\mu} = \arg \min_{\mu \in \mathcal{F}} R(\mu), \quad (6)$$

where

$$R(\mu) = \sum_{i=1}^n (y_i - \mu(\mathbf{x}_i))^2 + J(\mu),$$

$\mathcal{F}$  is a space of functions, and  $J(\mu)$  is a penalty function. Suppose  $\mathcal{F} = \mathcal{H}_{\kappa}$ ; the reproducing kernel Hilbert space (RKHS; e.g. Lange 2010, Section 17.5) with kernel  $\kappa(\cdot, \cdot; \boldsymbol{\rho})$ ; and  $J(\mu) = \lambda \|\mu\|_{\mathcal{H}_{\kappa}}$ , where  $\|\cdot\|_{\mathcal{H}_{\kappa}}$  is the RKHS norm. It can be shown (e.g. Wahba, 1990) that  $\hat{\mu}(\mathbf{x})$  is given by

$$\hat{\mu}(\mathbf{x}) = \sum_{i=1}^n u_i \kappa(\mathbf{x}, \mathbf{x}_i; \boldsymbol{\rho}),$$

where  $\mathbf{u} = (u_1, \dots, u_n)^T$  is

$$\mathbf{u} = \Phi^{-1} \mathbf{y},$$

and  $\Phi = K + \lambda I_n$ , with  $K$  the  $n \times n$  matrix with  $ij$ th element  $\kappa(\mathbf{x}_i, \mathbf{x}_j; \boldsymbol{\rho})$ . The values of  $\lambda$  and  $\boldsymbol{\rho}$  can be determined using using generalised cross validation (e.g. Tuo and Wu, 2015). Under certain regularity conditions, the most stringent being that  $\mathbf{x}_1, \dots, \mathbf{x}_n$  are an independent and identically distributed sample from  $U(\mathcal{X})$ , Tuo and Wu (2015) show that

$\hat{\boldsymbol{\theta}}_{TW}$  is a consistent estimator of  $\boldsymbol{\theta}_{L^2}$ . Moreover,  $\hat{\boldsymbol{\theta}}_{TW}$  has the following asymptotic normal distribution

$$\hat{\boldsymbol{\theta}}_{TW} \sim \text{N}(\boldsymbol{\theta}_{L^2}, V^{-1}WV^{-1}), \quad (7)$$

where

$$V = \frac{\partial^2 L_{L^2}(\boldsymbol{\theta}_{L^2})}{\partial \boldsymbol{\theta} \partial \boldsymbol{\theta}^T} \quad W = \frac{4\sigma^2}{n \text{Vol}(\mathcal{X})} \int_{\mathcal{X}} \frac{\partial \eta(\boldsymbol{\theta}_{L^2}, \mathbf{x})}{\partial \boldsymbol{\theta}} \frac{\partial \eta(\boldsymbol{\theta}_{L^2}, \mathbf{x})}{\partial \boldsymbol{\theta}^T} d\mathbf{x},$$

with  $\text{Vol}(\mathcal{X})$  denoting the volume of  $\mathcal{X}$ . Note that the distribution (7) is derived by marginalising over  $\mathbf{y}$  and  $\mathbf{x}_1, \dots, \mathbf{x}_n$ .

Alternatively, Tuo and Wu (2015) and Wong et al. (2017) considered the ordinary least squares estimator given by

$$\hat{\boldsymbol{\theta}}_{OLS} = \arg \min_{\boldsymbol{\theta} \in \Theta} \ell_{OLS}(\boldsymbol{\theta}; \mathbf{y}),$$

where

$$\ell_{OLS}(\boldsymbol{\theta}; \mathbf{y}) = \frac{1}{n} \sum_{i=1}^n (y_i - \eta(\boldsymbol{\theta}, \mathbf{x}))^2. \quad (8)$$

Tuo and Wu (2015) and Wong et al. (2017) showed that  $\hat{\boldsymbol{\theta}}_{OLS}$  is also a consistent estimator of  $\boldsymbol{\theta}_{L^2}$ . Furthermore,  $\hat{\boldsymbol{\theta}}_{OLS}$  has the following asymptotic normal distribution

$$\hat{\boldsymbol{\theta}}_{OLS} \sim \text{N}(\boldsymbol{\theta}_{L^2}, V^{-1}W_{OLS}V^{-1}), \quad (9)$$

where  $W_{OLS} = W + W_E$  with

$$W_E = \frac{4}{n \text{vol}(\mathcal{X})} \int (\mu(\mathbf{x}) - \eta(\boldsymbol{\theta}_{L^2}, \mathbf{x}))^2 \frac{\partial \eta(\boldsymbol{\theta}_{L^2}, \mathbf{x})}{\partial \boldsymbol{\theta}} \frac{\partial \eta(\boldsymbol{\theta}_{L^2}, \mathbf{x})}{\partial \boldsymbol{\theta}^T} d\mathbf{x}.$$

Wong et al. (2017) also proposed a bootstrapping scheme for uncertainty quantification for small  $n$ .

The comparative advantage of the ordinary least squares estimator is that it is computationally less demanding. The integration required to evaluate  $\ell_{L^2}(\boldsymbol{\theta}; \mathbf{y})$  is rarely available in closed form and requires numerical evaluation. Additionally, for the non-parametric regression, the values of  $\boldsymbol{\rho}$  and  $\lambda$  need to be determined. This allows the bootstrapping procedure proposed in Wong et al. (2017) to be implemented efficiently. However, Tuo and Wu (2015) show that the asymptotic variance of  $\hat{\boldsymbol{\theta}}_{OLS}$  is larger than for  $\hat{\boldsymbol{\theta}}_{L^2}$ , i.e.  $W_E \succcurlyeq 0$ , where  $A \succcurlyeq B$ , means that  $A - B$  is positive-definite.

Bayesian approaches allow a coherent approach to uncertainty quantification. Due to this, several authors have recently proposed Bayesian  $L^2$  calibration approaches.

Consider the definition of  $\boldsymbol{\theta}_{L^2}$  as the values of  $\boldsymbol{\theta}$  minimising  $L_{L^2}(\boldsymbol{\theta})$  given by (4). This implies that  $\partial L_{L^2}(\boldsymbol{\theta}_{L^2})/\partial \boldsymbol{\theta} = \mathbf{0}$ , where

$$\frac{\partial L_{L^2}(\boldsymbol{\theta})}{\partial \boldsymbol{\theta}} = \int_{\mathcal{X}} \frac{\partial \eta(\boldsymbol{\theta}, \mathbf{x})}{\partial \boldsymbol{\theta}} \delta_{\boldsymbol{\theta}}(\mathbf{x}) d\mathbf{x}.$$

Therefore, the bias function is orthogonal to the gradient (with respect to  $\boldsymbol{\theta}$ ) of the mathematical model, at  $\boldsymbol{\theta} = \boldsymbol{\theta}_{L^2}$ . Under the Kennedy and O’Hagan framework, this orthogonality condition introduces constraints on the Gaussian process prior for the bias function, i.e. it modifies the form of the chosen correlation function  $\kappa$ . Imposition of these constraints leads to the Plumlee (2017) Bayesian  $L^2$  calibration approach.

However, note that, due to local minima, the orthogonality condition can be true for values of  $\boldsymbol{\theta}$  not equal to  $\boldsymbol{\theta}_{L^2}$ . For this reason, Gu and Wang (2018) assumed a prior distribution for  $L_{L^2}(\boldsymbol{\theta})$ , in addition to the Kennedy and O’Hagan Gaussian process prior on the bias function, which penalises large values of the bias function.

An alternative approach begins by assuming a zero mean Gaussian process for  $\mu(\mathbf{x})$ , i.e.  $\mu(\cdot) \sim \text{GP}(0, \tau^2(\lambda + \kappa(\cdot, \cdot; \boldsymbol{\rho})))$ . The resulting posterior distribution for  $\mu(\mathbf{x})$  is a normal distribution with mean  $\hat{\mu}(\mathbf{x})$  given by (6) and variance  $\nu(\mathbf{x}) = \tau^2(1 + \lambda - \mathbf{u}^T \mathbf{k}(\mathbf{x}))$ , where  $\mathbf{k}(\mathbf{x})$  is an  $n \times 1$  vector with  $i$ th element  $\kappa(\mathbf{x}, \mathbf{x}_i; \boldsymbol{\rho})$ . By writing  $L(\boldsymbol{\theta})$  as a functional of  $\mu(\mathbf{x})$ , i.e.

$$L(\boldsymbol{\theta}, \mu(\mathbf{x})) = \int_{\mathcal{X}} (\mu(\mathbf{x}) - \eta(\boldsymbol{\theta}, \mathbf{x}))^2 d\mathbf{x},$$

Xie and Xu (2020) noted that this implies a posterior distribution for  $L(\boldsymbol{\theta}, \mu(\mathbf{x}))$  and hence for the values of  $\boldsymbol{\theta}$  minimising  $L(\boldsymbol{\theta}, \mu(\mathbf{x}))$ . Xie and Xu (2020) proved 1) the resulting posterior distribution converges to a point mass at  $\boldsymbol{\theta}_{L^2}^*$  and 2) a limiting normal approximation applies.

With most relevance, Woody et al. (2019) consider general Bayesian inference (2) with  $\ell(\boldsymbol{\theta}; \mathbf{y}) = \gamma \ell_{TW}(\boldsymbol{\theta}; \mathbf{y})$  where  $\gamma > 0$  is chosen to maintain frequentist coverage properties of credible sets for  $\boldsymbol{\theta}$ . Such a  $\gamma$  is analytically intractable but Woody et al. (2019) devise a bootstrap procedure to evaluate it numerically.



### 3 General Bayesian calibration of mathematical models

#### 3.1 The idea

Generalised posterior distributions in the form of (2) have been proposed based on a variety of different loss functions. Standard Bayesian inference occurs under the self-information loss given by  $\ell(\boldsymbol{\theta}; \mathbf{y}) = -n^{-1} \log \pi(\mathbf{y}|\boldsymbol{\theta})$  where  $\pi(\mathbf{y}|\boldsymbol{\theta})$  is the (marginal) likelihood function. For example, under the Kennedy and O’Hagan model,

$$\pi(\mathbf{y}|\boldsymbol{\theta}) \propto \int \int \int |\sigma^2 I_n + \tau^2 K|^{-\frac{1}{2}} \exp \left[ -\frac{1}{2} (\mathbf{y} - \boldsymbol{\eta}(\boldsymbol{\theta}))^\top (\sigma^2 I_n + \tau^2 K)^{-1} (\mathbf{y} - \boldsymbol{\eta}(\boldsymbol{\theta})) \right] \times \pi(\sigma^2, \tau^2, \boldsymbol{\rho}|\boldsymbol{\theta}) d\boldsymbol{\rho} d\sigma^2 d\tau^2,$$

where  $\boldsymbol{\eta}(\boldsymbol{\theta}) = (\eta(\boldsymbol{\theta}, \mathbf{x}_1), \dots, \eta(\boldsymbol{\theta}, \mathbf{x}_n))^\top$  and  $\pi(\sigma^2, \tau^2, \boldsymbol{\rho}|\boldsymbol{\theta})$  is the probability density function of the joint prior distribution of  $\boldsymbol{\rho}$ ,  $\sigma^2$  and  $\tau^2$  (conditional on  $\boldsymbol{\theta}$ ).

However, more generally and flexibly, generalised posterior distributions have been proposed for loss functions given by, for example, composite likelihood (e.g. Smith and Stephenson, 2009; Pauli et al., 2011; Ribatet et al., 2012), restricted likelihood (e.g. Pettitt, 1983), partial likelihood (e.g. Raftery et al., 1996) and quasi-likelihood (e.g. Ventura et al., 2010). In particular, we consider a general Bayesian  $L^2$  calibration approach using (2) where the loss function  $\ell(\boldsymbol{\theta}; \mathbf{y})$  is a functional of  $\ell_{TW}(\boldsymbol{\theta}; \mathbf{y})$ .

The idea is demonstrated with a simple example adapted from Plumlee (2017). Suppose the true physical system is  $\mu(x) = 4x + x \sin 5x$  with  $k = 1$  input  $x \in \mathcal{X} = [0, 1]$ . The mathematical model is  $\eta(\theta, x) = \theta x$  with  $p = 1$  calibration parameter  $\theta \in \Theta = \mathbb{R}$ . In this case,  $\theta_{L^2} = 3.565$ . The Tuo and Wu estimator is

$$\hat{\theta}_{TW} = 3 \int_0^1 x \hat{\mu}(x) dx.$$

Under (1), the true mean and variance (conditional on  $\mathbf{d} = \{x_1, \dots, x_n\}$ ,  $\sigma^2$ ,  $\lambda$ , and  $\rho$ ) of

$\hat{\theta}_{TW}$  are

$$\begin{aligned} \mathbb{E}(\hat{\theta}_{TW}) &= 3 \int_0^1 x \mathbf{k}(x)^\top dx \Phi^{-1} \boldsymbol{\mu} \\ \text{var}(\hat{\theta}_{TW}) &= 9\sigma^2 \int_0^1 x \mathbf{k}(x)^\top dx \Phi^{-2} \int_0^1 x \mathbf{k}(x) dx \end{aligned} \quad (10)$$

respectively, where  $\mathbf{k}(x) = (\kappa(x_1, x; \rho), \dots, \kappa(x_n, x; \rho))^\top$ .

Under a normal prior distribution for  $\theta$  (with mean 0 and variance  $\tau^2$ ) and  $\ell(\boldsymbol{\theta}, x) = \gamma \ell_{TW}(\boldsymbol{\theta}, \mathbf{y})$ , the generalised posterior distribution (conditional on  $\mathbf{d} = \{x_1, \dots, x_n\}$ ,  $\sigma^2$ ,  $\lambda$ , and  $\rho$ ) is a normal distribution with mean and variance

$$\begin{aligned} \mathbb{E}_G(\theta|\mathbf{y}) &= \frac{\hat{\theta}_{TW}}{1 + \frac{3}{2n\tau^2\gamma}}, \\ \text{var}_G(\theta|\mathbf{y}) &= \frac{\frac{3}{2n\gamma}}{1 + \frac{3}{2n\tau^2\gamma}} \end{aligned}$$

respectively.

We consider  $n = 4$  and  $n = 8$ . Suppose  $x_1, \dots, x_n$  are evenly-spaced in  $\mathcal{X}$  and we generate responses under (1), with  $e_i$  being normally distributed, for  $i = 1, \dots, n$ , with  $\sigma^2 = 0.25^2$  and suppose  $\tau^2 = 1$ . For  $n = 4$  and  $n = 8$ , Figures 1 (a) and (b), respectively, show plots of  $y_i$  against  $x_i$ , for  $i = 1, \dots, n$ , as well as  $\mu(x)$  (solid) and  $\hat{\mu}(x)$  (dashed) against  $x$ . For  $n = 4$  and  $n = 8$ , Figures 1 (c) and (d), respectively, show plots of the generalised posterior density  $\pi_G(\theta|\mathbf{y})$  against  $\theta$  for  $\gamma = 1$  (solid) and  $\gamma = 15$  (dotted). Lastly,  $\hat{\theta}_{TW}$  (circle) and the true  $\theta_{L^2} = 3.565$  (triangle) are identified on the  $x$ -axes of Figures 1 (c) and (d).

As  $n$  and  $\gamma$  increase the generalised posterior precision increases and the distribution converges to a point mass at  $\hat{\theta}_{TW}$ . Note that under a uniform prior distribution ( $\tau^2 \rightarrow \infty$ ), the generalised posterior mode would be equal to the maximum generalised likelihood estimate. However, in this case the posterior mode is shrunk toward the prior mode of zero. Although multiplying the loss by a constant  $\gamma$  will not change the minimiser,  $\hat{\theta}_{TW}$ , it clearly has a significant effect on the generalised posterior distribution. The choice of  $\gamma$  controls the relative weight of information from the responses and from the prior distribution. Bissiri et al. (2016) discuss various strategies for automatic adjustment of the generalised posterior distribution including 1) specifying  $\gamma$  so that the responses provide a factor of  $n$  times more




Figure 1: A caption

Table 1: Coverage and average length of 95% credible intervals for  $\theta_{L^2}$  for different values of  $\gamma$ .

$\gamma$	Coverage (%)		Average length	
	$n = 4$	$n = 8$	$n = 4$	$n = 8$
1	93	98	2.3	1.7
$\hat{\gamma}$	93	93	0.9	0.6
15	81	84	0.6	0.4

information than the prior; 2) allowing  $\gamma$  to be unknown with its own loss and prior so the resulting generalised posterior is more robust to its misspecification and 3) choosing  $\gamma$  so that the generalised posterior has the desired frequentist properties.

Consider the third approach. As long as the prior variance  $\tau^2$  is sufficiently large, we specify  $\gamma$  to be the value that equates the generalised posterior variance and the frequentist variance of  $\hat{\theta}_{TW}$ , yielding

$$\tilde{\gamma} = \frac{3}{2n} \left( \frac{1}{\text{var}(\hat{\theta}_{TW})} - \frac{1}{\tau^2} \right),$$

where  $\text{var}(\hat{\theta}_{TW})$  is given by (10).

For the responses shown in Figures 1 (a) and (b),  $\tilde{\gamma} = 6.37$  and  $\tilde{\gamma} = 8.04$  for  $n = 4$  and  $n = 8$ , respectively. For  $n = 4$  and  $n = 8$ , Figures 1 (c) and (d), respectively, show plots of the generalised posterior density (dashed) for  $\gamma = \tilde{\gamma}$ , respectively.

We extend this example by performing a simulation study. For each value of  $n = 4$  and  $n = 8$ , we repeat the above experiment 10,000 times and calculate the coverage and average length of 95% highest density credible intervals for  $\theta_{L^2}$ . The results are shown in Table 1. For  $\gamma = 1$ , the coverage is close to the nominal 95% but the intervals are the widest (on average). For  $\gamma = 15$ , the intervals are the narrowest, but the coverage is significantly less than 95%. For  $\gamma = \hat{\gamma}$ , the coverage is close to 95% and the intervals are only 50% wider (on average) than when  $\gamma = 15$ .

The above approach to select  $\tilde{\gamma}$  is only appropriate if  $p = 1$  since we cannot equate  $p > 1$

variances using scalar  $\gamma$ . Additionally, it will not be possible to, in general, calculate the exact generalised posterior variance and frequentist variance of  $\hat{\boldsymbol{\theta}}_{TW}$ . To solve these problems, we consider alternative ways of equating generalised posterior and frequentist properties, more complex functionals of  $\ell_{TW}(\boldsymbol{\theta}, \mathbf{y})$  and equating asymptotic generalised posterior and frequentist properties as opposed to exact.

## 3.2 Theoretical details

In this section we establish the asymptotic behaviour of the generalised posterior under the Tuo and Wu loss  $\ell(\boldsymbol{\theta}; \mathbf{y}) = \ell_{TW}(\boldsymbol{\theta}; \mathbf{y})$  given by (5). Initially, in Section 3.2.1, we consider the case, as assumed by Tuo and Wu (2015), that the design points  $\mathbf{x}_1, \dots, \mathbf{x}_n$  are an independent and identically distributed sample from a uniform distribution on  $\mathcal{X}$ . Then, in Section 3.2.2, we consider the behaviour conditional on  $\mathbf{d} = \{\mathbf{x}_1, \dots, \mathbf{x}_n\}$ .

### 3.2.1 Asymptotic behaviour of generalised posterior marginal of design

We assume the regularity conditions A1 to A5 and B1 to B4 in Tuo and Wu (2015).

In addition, we assume the following conditions.

- C1  $E \subset \Theta$  is open, convex and bounded with  $\boldsymbol{\theta}_{L^2} \in E$  and  $\hat{\boldsymbol{\theta}}_{TW} \in E$ .
- C2  $\eta(\boldsymbol{\theta}, \mathbf{x})$ ,  $\partial\eta(\boldsymbol{\theta}, \mathbf{x})/\partial\theta_u$ ,  $\partial^2\eta(\boldsymbol{\theta}, \mathbf{x})/\partial\theta_u\partial\theta_v$ , and  $\partial^3\eta(\boldsymbol{\theta}, \mathbf{x})/\partial\theta_u\partial\theta_v\partial\theta_w$  are continuous and uniformly bounded on  $E \times \mathcal{X}$ .
- C3  $\|\partial^2\eta(\boldsymbol{\theta}, \mathbf{x})/\partial\boldsymbol{\theta}\partial\boldsymbol{\theta}^T\|_{\mathcal{H}_\kappa(\mathcal{X})} < \infty$  for  $\boldsymbol{\theta} \in B_\epsilon(\boldsymbol{\theta}_{L^2})$  for some  $\epsilon > 0$ .
- C4  $\pi(\boldsymbol{\theta})$  is continuous and positive at  $\boldsymbol{\theta}_{L^2}$ .

Let  $\|\cdot\|_F$  denote the Frobenius norm, i.e. if  $U$  is an  $p \times q \times r$  array, then  $\|U\|_F = (\sum_{u=1}^p \sum_{v=1}^q \sum_{w=1}^r U_{uvw}^2)^{1/2}$ . Additionally, let  $B_\epsilon(\mathbf{u}) = \{\mathbf{u} : \|\mathbf{u}\|_F < \epsilon\}$  denote the open ball of radius  $\epsilon > 0$  at  $\mathbf{u}$ .

*THEOREM 1*

Under conditions A1 to A5, B1 to B4 and C1 to C4, the following results hold.

(a) (2nd order Taylor series expansion) The loss  $\ell_{TW}(\boldsymbol{\theta}; \mathbf{y})$  can be written as

$$\ell_{TW}(\boldsymbol{\theta}; \mathbf{y}) = \ell_{TW}(\hat{\boldsymbol{\theta}}_{TW}; \mathbf{y}) + \frac{1}{2} \left( \boldsymbol{\theta} - \hat{\boldsymbol{\theta}}_{TW} \right)^{\top} \hat{V}_{TW} \left( \boldsymbol{\theta} - \hat{\boldsymbol{\theta}}_{TW} \right) + r_n(\boldsymbol{\theta} - \hat{\boldsymbol{\theta}}_{TW}) \quad (11)$$

where

$$\hat{V}_{TW} = \frac{\partial^2 \ell(\hat{\boldsymbol{\theta}}_{TW}; \mathbf{y})}{\partial \boldsymbol{\theta} \partial \boldsymbol{\theta}^{\top}},$$

and  $r_n : \mathbb{R}^p \rightarrow \mathbb{R}$  where there exist  $\epsilon_0, c_0 > 0$  such that  $|r_n(\mathbf{u})| \leq c_0 \|\mathbf{u}\|_F^3$  for all  $\mathbf{u} \in B_{\epsilon_0}(\mathbf{0})$ .

(b) (Generalised posterior concentration) For all  $\epsilon > 0$ ,

$$\int_{B_{\epsilon}(\boldsymbol{\theta}_{L^2})} \pi_G(\boldsymbol{\theta} | \mathbf{y}) d\boldsymbol{\theta} \rightarrow 1,$$

as  $n \rightarrow \infty$ .

(c) (Asymptotic normal distribution) The generalised posterior distribution of (2) converges to

$$\mathbf{N} \left( \hat{\boldsymbol{\theta}}_{TW}, \frac{1}{n} V^{-1} \right)$$

in total variation.

*Proof*

Result (a) follows by Theorem 3.3 of Miller (2019) if the following conditions (i) - (iii) are true.

(i) There exist  $\hat{\boldsymbol{\theta}}_{TW} \in E$  with  $\partial \ell(\hat{\boldsymbol{\theta}}_{TW}; \mathbf{y}) / \partial \boldsymbol{\theta} = \mathbf{0}$  such that  $\hat{\boldsymbol{\theta}}_{TW} \xrightarrow{p} \boldsymbol{\theta}_{L^2}$ , as  $n \rightarrow \infty$ .

This follows from the definition of  $\hat{\boldsymbol{\theta}}_{TW}$  and, under conditions A1 to A5 and B1 to B4, Theorem 1 of Tuo and Wu (2015) (which shows  $\hat{\boldsymbol{\theta}}_{TW} \xrightarrow{p} \boldsymbol{\theta}_{L^2}$ ).

$$(ii) \frac{\partial^2 \ell(\hat{\boldsymbol{\theta}}_{TW}; \mathbf{y})}{\partial \boldsymbol{\theta} \partial \boldsymbol{\theta}^T} \xrightarrow{p} V.$$

Let  $S_n(\mu) = \frac{1}{n} \sum_{i=1}^n (y_i - \mu(\mathbf{x}_i))^2 + \lambda_n \|\mu\|_{\mathcal{H}_\kappa(\mathcal{X})}^2$ . Now  $\hat{\mu}(\cdot)$  minimises  $S_n$  over  $\mathcal{H}_\kappa(\mathcal{X})$ . Under C3, and since  $\hat{\boldsymbol{\theta}}_{TW}$  is consistent for  $\boldsymbol{\theta}_{L^2}$ , it means  $\partial^2 \eta(\hat{\boldsymbol{\theta}}_{TW}, \cdot) / \partial \theta_u \partial \theta_v \in \mathcal{H}_\kappa(\mathcal{X})$  for sufficiently large  $n$ . Let  $\langle \mu_1, \mu_2 \rangle_{\mathcal{H}_\kappa(\mathcal{X})}$  denote the inner product of  $\mathcal{H}_\kappa(\mathcal{X})$ , with  $\|\mu_1\|_{\mathcal{H}_\kappa(\mathcal{X})}^2 = \langle \mu_1, \mu_1 \rangle_{\mathcal{H}_\kappa(\mathcal{X})}$ . For  $u, v = 1, \dots, p$ , we have

$$\begin{aligned} 0 &= \left. \frac{\partial}{\partial t} S_n \left( \hat{\mu}(\cdot) + t \frac{\partial^2 \eta(\hat{\boldsymbol{\theta}}_{TW}, \cdot)}{\partial \theta_u \partial \theta_v} \right) \right|_{t=0}, \\ &= \frac{2}{n} \sum_{i=1}^n (\hat{\mu}(\mathbf{x}_i) - y_i) \frac{\partial^2 \eta(\hat{\boldsymbol{\theta}}_{TW}, \mathbf{x}_i)}{\partial \theta_u \partial \theta_v} + 2\lambda_n \left\langle \hat{\mu}(\cdot), \frac{\partial^2 \eta(\hat{\boldsymbol{\theta}}_{TW}, \cdot)}{\partial \theta_u \partial \theta_v} \right\rangle_{\mathcal{H}_\kappa(\mathcal{X})}, \\ &= \frac{2}{n} \sum_{i=1}^n (\hat{\mu}(\mathbf{x}_i) - \mu(\mathbf{x}_i)) \frac{\partial^2 \eta(\hat{\boldsymbol{\theta}}_{TW}, \mathbf{x}_i)}{\partial \theta_u \partial \theta_v} - \frac{2}{n} \sum_{i=1}^n e_i \frac{\partial^2 \eta(\hat{\boldsymbol{\theta}}_{TW}, \mathbf{x}_i)}{\partial \theta_u \partial \theta_v} \\ &\quad + 2\lambda_n \left\langle \hat{\mu}(\cdot), \frac{\partial^2 \eta(\hat{\boldsymbol{\theta}}_{TW}, \cdot)}{\partial \theta_u \partial \theta_v} \right\rangle_{\mathcal{H}_\kappa(\mathcal{X})}, \\ &= 2(C_n + D_n + E_n). \end{aligned}$$

Following the same arguments as in the proof of Theorem 1 in Tuo and Wu (2015), with  $\partial^2 \eta(\hat{\boldsymbol{\theta}}_{TW}, \cdot) / \partial \theta_u \partial \theta_v$  in place of  $\partial \eta(\hat{\boldsymbol{\theta}}_{TW}, \cdot) / \partial \theta_u$ , we obtain

$$\begin{aligned} C_n &= \int_{\mathcal{X}} (\hat{\mu}(\mathbf{x}) - \mu(\mathbf{x})) \frac{\partial^2 \eta(\hat{\boldsymbol{\theta}}_{TW}, \mathbf{x})}{\partial \theta_u \partial \theta_v} d\mathbf{x} + o_p(n^{-1/2}), \\ D_n &= \frac{1}{n} \sum_{i=1}^n e_i \frac{\partial^2 \eta(\boldsymbol{\theta}_{L^2}, \mathbf{x}_i)}{\partial \theta_u \partial \theta_v} + o_p(n^{-1/2}), \\ &= o_p(1), \quad \text{by the weak law of large numbers,} \\ E_n &\leq o_p(n^{-1/2}). \end{aligned}$$

Therefore

$$\begin{aligned} \int_{\mathcal{X}} \hat{\mu}(\mathbf{x}) \frac{\partial^2 \eta(\hat{\boldsymbol{\theta}}_{TW}, \mathbf{x})}{\partial \theta_u \partial \theta_v} d\mathbf{x} &= \int_{\mathcal{X}} \mu(\mathbf{x}) \frac{\partial^2 \eta(\hat{\boldsymbol{\theta}}_{TW}, \mathbf{x})}{\partial \theta_u \partial \theta_v} d\mathbf{x} + o_p(n^{-1/2}), \\ &\xrightarrow{p} \int_{\mathcal{X}} \mu(\mathbf{x}) \frac{\partial^2 \eta(\boldsymbol{\theta}_{L^2}, \mathbf{x})}{\partial \theta_u \partial \theta_v} d\mathbf{x}, \end{aligned}$$

by the consistency of  $\hat{\boldsymbol{\theta}}_{TW}$ . Hence

$$\frac{\partial^2 \ell(\hat{\boldsymbol{\theta}}_{TW}; \mathbf{y})}{\partial \boldsymbol{\theta} \partial \boldsymbol{\theta}^T} \xrightarrow{p} \frac{\partial^2 L_{L^2}(\boldsymbol{\theta}_{L^2})}{\partial \boldsymbol{\theta} \partial \boldsymbol{\theta}^T} = V,$$

as required.

- (iii) The sequence of functions  $\partial^3 \ell_{TW}(\boldsymbol{\theta}; \mathbf{y}) / \partial \theta_i \partial \theta_j \partial \theta_h$  is uniformly bounded for  $i, j, h = 1, \dots, p$  and  $\boldsymbol{\theta} \in E$ .

This follows from condition C2.

Results (b) and (c) follow by Theorem 3.1 of Miller (2019), if, for any  $\epsilon > 0$ ,

$$\liminf_{n \rightarrow \infty} \inf_{\boldsymbol{\theta} \in B_\epsilon(\hat{\boldsymbol{\theta}}_{TW})^c} \left\{ \ell_{TW}(\boldsymbol{\theta}; \mathbf{y}) - \ell_{TW}(\hat{\boldsymbol{\theta}}_{TW}; \mathbf{y}) \right\} > 0, \quad (12)$$

and the assumption of condition C4. Since  $\hat{\boldsymbol{\theta}}_{TW}$  is a unique minimiser of  $\ell_{TW}(\boldsymbol{\theta}; \mathbf{y})$  and is an interior point of  $\Theta$ , then (12) follows. ■

### 3.2.2 Asymptotic behaviour of generalised posterior conditional on design

Following Wong et al. (2017), we establish results for fixed values of  $\lambda$  and  $\boldsymbol{\rho}$ .

Define

$$\bar{\boldsymbol{\mu}}(\mathbf{x}) = \mathbf{k}(\mathbf{x})^T \Phi^{-1} \boldsymbol{\mu},$$

where  $\boldsymbol{\mu} = (\mu(\mathbf{x}_1), \dots, \mu(\mathbf{x}_n))^T$ . Furthermore, define

$$\bar{L}_{L^2}(\boldsymbol{\theta}) = \int_{\mathcal{X}} (\bar{\boldsymbol{\mu}}(\mathbf{x}) - \eta(\boldsymbol{\theta}, \mathbf{x}))^2 d\mathbf{x}$$

and let  $\bar{\boldsymbol{\theta}}_{L^2}$  be the unique minimiser of  $\bar{L}_{L^2}(\boldsymbol{\theta})$  over  $\Theta$ .

In addition to conditions A1 to A5 and B1 to B4, we assume the following conditions.



D1  $E \subset \Theta$  is open, convex and bounded with  $\bar{\boldsymbol{\theta}}_{L^2} \in E$  and  $\hat{\boldsymbol{\theta}}_{TW} \in E$ .

D2  $\eta(\boldsymbol{\theta}, \mathbf{x})$ ,  $\partial\eta(\boldsymbol{\theta}, \mathbf{x})/\partial\theta_u$ ,  $\partial^2\eta(\boldsymbol{\theta}, \mathbf{x})/\partial\theta_u\partial\theta_v$ , and  $\partial^3\eta(\boldsymbol{\theta}, \mathbf{x})/\partial\theta_u\partial\theta_v\partial\theta_w$  are continuous and uniformly bounded on  $E \times \mathcal{X}$ .

D3  $\|\partial^2\eta(\boldsymbol{\theta}, \mathbf{x})/\partial\boldsymbol{\theta}\partial\boldsymbol{\theta}^\top\|_{\mathcal{H}_\kappa(\mathcal{X})} < \infty$  for  $\boldsymbol{\theta} \in B_\epsilon(\bar{\boldsymbol{\theta}}_{L^2})$  for some  $\epsilon > 0$ .

D4  $\pi(\boldsymbol{\theta})$  is continuous and positive at  $\bar{\boldsymbol{\theta}}_{L^2}$ .

D5 The design  $\mathbf{d} = \{\mathbf{x}_1, \dots, \mathbf{x}_n\}$  is such that

$$\lim_{n \rightarrow \infty} \sum_{i=1}^n \frac{g_i^2}{i^2} < \infty$$

where  $g_i$  is the  $i$ th element of  $\mathbf{g} = \Phi^{-1}\mathbf{k}(\mathbf{x})$ .

D6 For  $i = 1, \dots, n$ ,  $e_i$  is normally distributed.

Condition D6 was also assumed by Xie and Xu (2020).

### *THEOREM 2*

Under conditions A1 to A5, B1 to B4 and D1 to D6, the following results hold.

(a)  $\hat{\boldsymbol{\theta}}_{L^2}$  is a consistent estimator of  $\bar{\boldsymbol{\theta}}_{L^2}$ .

(b)  $\hat{\boldsymbol{\theta}}_{L^2}$  has the following asymptotic distribution

$$\hat{\boldsymbol{\theta}}_{L^2} \sim \text{N}(\bar{\boldsymbol{\theta}}_{L^2}, \bar{V}^{-1}\bar{W}\bar{V}^{-1}),$$

where

$$\begin{aligned} \bar{V} &= \frac{\partial L_{L^2}(\bar{\boldsymbol{\theta}}_{L^2}, \mathbf{x})}{\partial\boldsymbol{\theta}\partial\boldsymbol{\theta}^\top} \\ \bar{W} &= 4\sigma^2 \int_{\mathcal{X}} \frac{\partial\eta(\boldsymbol{\theta}, \mathbf{x})}{\partial\boldsymbol{\theta}} \frac{\partial\eta(\boldsymbol{\theta}, \mathbf{x})}{\partial\boldsymbol{\theta}^\top} \sum_{i=1}^n g_i^2. \end{aligned}$$

(c) (Generalised posterior concentration) For all  $\epsilon > 0$ ,

$$\int_{B_\epsilon(\bar{\boldsymbol{\theta}}_{L^2})} \pi_G(\boldsymbol{\theta}|\mathbf{y})d\boldsymbol{\theta} \rightarrow 1,$$

as  $n \rightarrow \infty$ .

(d) (Asymptotic normal distribution) The generalised posterior distribution of (2) converges to

$$\mathbf{N}\left(\hat{\boldsymbol{\theta}}_{TW}, \frac{1}{n}\bar{V}^{-1}\right)$$

in total variation.

*Proof*

By the strong law of large numbers and D5,  $\hat{\mu}(\mathbf{x}) \xrightarrow{p} \bar{\mu}(\mathbf{x})$ , as  $n \rightarrow \infty$ . Due to the definitions of  $\hat{\boldsymbol{\theta}}_{L^2}$  and  $\bar{\boldsymbol{\theta}}_{L^2}$ , to prove (a) it suffices to show that  $\ell(\boldsymbol{\theta}, \mathbf{y})$  converges to  $\bar{L}_{L^2}(\boldsymbol{\theta})$  uniformly in probability for all  $\boldsymbol{\theta} \in \Theta$ . This follows from

$$\begin{aligned} \ell(\boldsymbol{\theta}, \mathbf{y}) - \bar{L}_{L^2}(\boldsymbol{\theta}) &= \int_{\mathcal{X}} (\hat{\mu}(\mathbf{x}) - \bar{\mu}(\mathbf{x})) (\hat{\mu}(\mathbf{x}) + \bar{\mu}(\mathbf{x}) - 2\eta(\boldsymbol{\theta}, \mathbf{x})) d\mathbf{x}, \\ &\leq \|\hat{\mu}(\mathbf{x}) - \bar{\mu}(\mathbf{x})\|_2 \|\hat{\mu}(\mathbf{x}) + \bar{\mu}(\mathbf{x}) - 2\eta(\boldsymbol{\theta}, \mathbf{x})\|_2, \\ &\leq \|\hat{\mu}(\mathbf{x}) - \bar{\mu}(\mathbf{x})\|_2 (\|\hat{\mu}(\mathbf{x})\|_2 + \|\bar{\mu}(\mathbf{x})\|_2 + 2\|\eta(\boldsymbol{\theta}, \mathbf{x})\|_2), \end{aligned}$$

where the first and second inequalities follow from the Cauchy-Schwartz and triangle inequalities, respectively. Since  $\hat{\mu}(\mathbf{x}) \xrightarrow{p} \bar{\mu}(\mathbf{x})$ , the right hand side converges in probability to 0.

As  $\hat{\boldsymbol{\theta}}_{L^2}$  is the unique minimiser of  $\ell(\boldsymbol{\theta}, \mathbf{y})$ , then by the mean-value theorem,

$$\begin{aligned} \mathbf{0} &= \frac{\partial \ell(\hat{\boldsymbol{\theta}}_{L^2}, \mathbf{y})}{\partial \boldsymbol{\theta}} \\ &= \frac{\partial \ell(\bar{\boldsymbol{\theta}}_{L^2}, \mathbf{y})}{\partial \boldsymbol{\theta}} + \frac{\partial^2 \ell(\boldsymbol{\zeta}, \mathbf{y})}{\partial \boldsymbol{\theta} \partial \boldsymbol{\theta}^T} (\hat{\boldsymbol{\theta}}_{L^2} - \bar{\boldsymbol{\theta}}_{L^2}), \end{aligned} \tag{13}$$

where  $\boldsymbol{\zeta}$  lies between  $\hat{\boldsymbol{\theta}}_{L^2}$  and  $\bar{\boldsymbol{\theta}}_{L^2}$ . Rearranging (14)

$$\hat{\boldsymbol{\theta}}_{L^2} = \bar{\boldsymbol{\theta}}_{L^2} + \left[ -\frac{\partial^2 \ell(\boldsymbol{\zeta}, \mathbf{y})}{\partial \boldsymbol{\theta} \partial \boldsymbol{\theta}^T} \right]^{-1} \frac{\partial \ell(\bar{\boldsymbol{\theta}}_{L^2}, \mathbf{y})}{\partial \boldsymbol{\theta}}.$$

From condition D6, and that  $\boldsymbol{\theta}_0 \xrightarrow{p} \bar{\boldsymbol{\theta}}_{L^2}$ , (b) follows.

Proof of (c) and (d) follow the same arguments as in the proof of (b) and (c) from Theorem 1, respectively. ■

### 3.3 Scaling the generalised posterior distribution

The scaling procedure used in Section 3.1 is to choose  $\gamma$  in  $\ell(\boldsymbol{\theta}; \mathbf{y}) = \gamma \ell_{TW}(\boldsymbol{\theta}; \mathbf{y})$  so that the generalised posterior variance of  $\boldsymbol{\theta}$  is equal to the asymptotic variance of  $\hat{\boldsymbol{\theta}}_{TW}$ . However this is only suitable for a  $p = 1$  parameter as it would be impossible to choose  $\gamma$  to equate  $p$  general posterior and asymptotic variances. Instead, we consider two approaches previously used in the composite likelihood literature (Ribatet et al., 2012). The first chooses  $\gamma$  to ensure the expectation of a quantity analogous to the likelihood ratio statistic converges to  $p$  (the expectation of the likelihood ratio statistic). The second scales the curvature of the generalised likelihood so that the asymptotic generalised posterior variance is equal to the asymptotic variance of  $\hat{\boldsymbol{\theta}}_{TW}$ .

#### 3.3.1 Magnitude scaling

Suppose  $\ell(\boldsymbol{\theta}; \mathbf{y})$  is the self-information loss, i.e.  $\ell(\boldsymbol{\theta}; \mathbf{y}) = -\frac{1}{n} \pi(\mathbf{y}|\boldsymbol{\theta})$ . The likelihood ratio statistic is

$$\Lambda(\boldsymbol{\theta}; \mathbf{y}) = 2n \left( \ell(\boldsymbol{\theta}; \mathbf{y}) - \ell(\hat{\boldsymbol{\theta}}; \mathbf{y}) \right),$$

where  $\hat{\boldsymbol{\theta}}$  are the maximum likelihood estimates of  $\boldsymbol{\theta}$ . It can be shown that  $\Lambda(\boldsymbol{\theta}; \mathbf{y})$ , asymptotically, has a  $\chi_p^2$  distribution when  $\boldsymbol{\theta} = \boldsymbol{\theta}_0$  with  $\boldsymbol{\theta}_0$  the true value of  $\boldsymbol{\theta}$ . Therefore,  $E(\Lambda(\boldsymbol{\theta}_0; \mathbf{y})) \xrightarrow{p} p$ , as  $n \rightarrow \infty$ .

Now define

$$\Lambda_{TW}^\gamma(\boldsymbol{\theta}; \mathbf{y}) = 2n \left( \ell_{TW}^\gamma(\boldsymbol{\theta}; \mathbf{y}) - \ell_{TW}^\gamma(\hat{\boldsymbol{\theta}}; \mathbf{y}) \right).$$

The idea is to choose  $\gamma$  so that the expectation of  $\Lambda_{TW}^\gamma(\boldsymbol{\theta}_{L^2}; \mathbf{y})$  converges to  $p$  as  $n \rightarrow \infty$ .

By Theorem 1(a),

$$\Lambda_{TW}^\gamma(\boldsymbol{\theta}_{L^2}; \mathbf{y}) = \gamma n \left( \boldsymbol{\theta}_{L^2} - \hat{\boldsymbol{\theta}}_{TW} \right)^\top \hat{V}_{TW} \left( \boldsymbol{\theta}_{L^2} - \hat{\boldsymbol{\theta}}_{TW} \right).$$

Initially consider marginalising over the design (see Section 3.2.1). By Theorem 1(c),

$$\mathbb{E}(\Lambda_{TW}^\gamma(\boldsymbol{\theta}_{L^2}; \mathbf{y})) \rightarrow^p \gamma n \text{tr} [V^{-1}W],$$

as  $n \rightarrow \infty$ . Thus setting

$$\tilde{\gamma} = \frac{p}{n \text{tr} [V^{-1}W]},$$

maintains the desired asymptotic behaviour of  $\Lambda_{TW}^\gamma(\boldsymbol{\theta}_{L^2}; \mathbf{y})$ .

If we condition on the design (see Section 3.2.1), then

$$\tilde{\gamma} = \frac{p}{n \text{tr} [\tilde{V}^{-1}\tilde{W}]},$$

### 3.3.2 Curvature scaling

Now consider the following loss

$$\ell_{TW}^\Gamma(\boldsymbol{\theta}, \mathbf{y}) = \ell_{TW} \left( \hat{\boldsymbol{\theta}}_{TW} + \Gamma \left( \boldsymbol{\theta} - \hat{\boldsymbol{\theta}}_{TW} \right), \mathbf{y} \right),$$

where  $\Gamma$  is a  $p \times p$  positive semi-definite matrix requiring specification. It is straightforward to see that  $\ell_{TW}^\Gamma(\boldsymbol{\theta}, \mathbf{y})$  is minimised by  $\boldsymbol{\theta} = \hat{\boldsymbol{\theta}}_{TW}$ . Minor modification of Theorem 1 leads to the generalised posterior distribution converging to a normal distribution with mean  $\hat{\boldsymbol{\theta}}_{TW}$  and variance  $(\Gamma^\top V \Gamma)^{-1}$ . Then  $\Gamma$  is chosen to equate the asymptotic general posterior variance with the asymptotic variance of  $\hat{\boldsymbol{\theta}}_{TW}$ , i.e. set  $\Gamma = \tilde{\Gamma}$  where

$$\tilde{\Gamma}^\top V \tilde{\Gamma} = V W^{-1} V.$$

Initially, consider marginalising over the design. Following Chandler and Bate (2007), we set

$$\tilde{\Gamma} = \tilde{\Gamma}_1^{-1} \tilde{\Gamma}_2$$

where  $\tilde{\Gamma}_2^\top \tilde{\Gamma}_2 = V W^{-1} V$  and  $\tilde{\Gamma}_1^\top \tilde{\Gamma}_1 = V$ . Such choices of  $\tilde{\Gamma}_1$  and  $\tilde{\Gamma}_2$  are not unique. It may be important that the mapping defined by  $\tilde{\Gamma}$  preserve direction of asymmetries. For this

reason, following Ribatet et al. (2012), we use singular value decompositions to obtain  $\tilde{\Gamma}_1$  and  $\tilde{\Gamma}_2$ .

If we condition on the design then  $\tilde{\Gamma}_2^T \tilde{\Gamma}_2 = \bar{V} \bar{W}^{-1} \bar{V}$  and  $\tilde{\Gamma}_1^T \tilde{\Gamma}_1 = \bar{V}$ .

### 3.3.3 Implementation details

The results above rely on knowing the values of  $\boldsymbol{\theta}_{L^2}$ ,  $\bar{\boldsymbol{\theta}}_{L^2}$ ,  $\lambda$ ,  $\boldsymbol{\rho}$  and  $\sigma^2$ . To implement the above we replace  $\boldsymbol{\theta}_{L^2}$  and  $\bar{\boldsymbol{\theta}}_{L^2}$  by estimates given by  $\hat{\boldsymbol{\theta}}_{TW}$ . Additionally, we replace  $\lambda$ ,  $\boldsymbol{\rho}$  and  $\sigma^2$  by estimates  $\hat{\lambda}$ ,  $\hat{\boldsymbol{\rho}}$  and  $\hat{\sigma}^2$  found via generalised cross-validation.

## 4 Examples

We consider three illustrative examples from Xie and Xu (2020). In each case there is  $k = 1$  controllable variable and  $\mathcal{X} = [0, 1]$ .

### Scenario 1

The model is

$$\eta(\boldsymbol{\theta}, x) = 7 [\sin(2\pi\theta_1 - \pi)]^2 + 2 [(2\pi\theta_2 - \pi)^2 \sin(2\pi x - \pi)],$$

where  $\boldsymbol{\theta} = (\theta_1, \theta_2)^T \in [0, 0.25] \times [0, 0.5]$ . The true physical system is

$$\mu(x) = \eta(\boldsymbol{\theta}_0, x),$$

where  $\boldsymbol{\theta}_0 = (0.2, 0.3)^T$ . Thus  $\boldsymbol{\theta}_{L^2} = \boldsymbol{\theta}_0$ .

The number of responses is  $n = 50$  and  $\mathbf{d} = \{x_1, \dots, x_n\}$  are generated from a  $U(\mathcal{X})$ . The errors are generated from a normal distribution with variance  $\sigma^2 = 0.2^2$ .

### Scenario 2

The model is

$$\eta(\theta, x) = \sin(5\theta x) + 5x,$$

where  $\theta \in [0, 3]$ . The true physical system is

$$\mu(x) = 5x \cos(15x/2) + 5x.$$

It can be found that  $\theta_{L^2} = 1.8771$ .

The number of responses is  $n = 30$  and  $\mathbf{d} = \{x_1, \dots, x_n\}$  are equidistant on  $\mathcal{X}$ . The errors are generated from a normal distribution with variance  $\sigma^2 = 0.2^2$ .

### Scenario 3

The model is

$$\eta(\theta, x) = \theta x,$$

where  $\theta \in [2, 4]$ . The true physical system is

$$\mu(x) = 4x + x \sin(5x),$$

with  $\theta_{L^2} = 3.5609$ .

The number of responses is  $n = 17$  and  $\mathbf{d} = \{x_1, \dots, x_n\}$  are equidistant on  $[0, 0.8]$ . The errors are generated from a normal distribution with variance  $\sigma^2 = 0.02^2$ .

In each case, we perform 200 simulations. In each simulation, we generate  $n$  responses from the true physical system and perform four analyses: 1) marginal on design, magnitude scaling; 2) marginal on design, curvature scaling; 3) conditional on design, magnitude scaling; and 4) conditional on design, curvature scaling.

Table 2 shows the results from the simulation study. Shown are the mean of the generalised posterior mean and standard deviation. Also shown are the coverage of the 95% (0.025, 0.975) quantile credible intervals.

For Scenarios 1 and 2 the results under the four approaches are broadly similar. The conditional approaches result in slightly higher generalised posterior standard deviation but have slightly improved coverage. However, in scenario 3, where the controllable variables are not generated from  $U(\mathcal{X})$ , the coverage under the conditional approach are far improved.

Table 2: Results from the simulation study. Shown are the mean of the generalised posterior mean and standard deviation (SD). Also shown are the coverage of the 95% (0.025, 0.975) quantile credible intervals

Scenario		Marginal				Conditional			
		Magnitude		Curvature		Magnitude		Curvature	
1	Mean	0.2000	0.3000	0.2000	0.3001	0.2000	0.3001	0.2001	0.3001
	SD	0.0011	0.0013	0.0011	0.0013	0.0012	0.0013	0.0012	0.0013
	Coverage	0.9250	0.9000	0.9150	0.9050	0.9350	0.9100	0.9600	0.9100
2	Mean	1.8771		1.8771		1.8771		1.8771	
	SD	0.0046		0.0046		0.0045		0.0045	
	Coverage	0.9400		0.9450		0.9350		0.9450	
3	Mean	3.5622		3.5622		3.5619		3.5619	
	SD	0.0078		0.0078		0.0571		0.0571	
	Coverage	0.1400		0.1400		0.7700		0.7700	

## 5 Discussion

We have presented a general Bayesian method to calibrate a mathematical model under the  $L^2$  loss. This is based on automatic adjustment of the  $L^2$  loss to maintain certain asymptotic properties of the general Bayesian posterior of the calibration parameters.

## References

- Bissiri, P., Holmes, C. and Walker, S. (2016) A general framework for updating belief distributions. *Journal of the Royal Statistical Society Series B* **78**, 1103–1130.
- Chandler, R. and Bate, S. (2007) Inference for clustered data using the independence log-likelihood. *Biometrika* **94**, 167–183.
- Gu, M. and Wang, L. (2018) Scaled Gaussian stochastic process for computer model calibration and prediction. *SIAM/ASA Journal of Uncertainty Quantification* **6**, 1555–1583.

- Kennedy, M. and O’Hagan, A. (2001) Bayesian calibration of computer models (with discussion). *Journal of the Royal Statistical Society Series B* **63**, 425–464.
- Lange, K. (2010) *Numerical Analysis for Statisticians*. Springer-Verlag, New York 2nd edn.
- Miller, J. (2019) Asymptotic normality, concentration, and coverage of generalized posteriors. *arXiv:1907.09611v1*.
- Pauli, F., Racugno, W. and Ventura, L. (2011) Bayesian composite marginal likelihoods. *Statistica Sinica* **21**, 149–164.
- Pettitt, A. N. (1983) Likelihood based inference using signed ranks for matched pairs. *Journal of the Royal Statistical Society Series B* **45**, 287–296.
- Plumlee, M. (2017) Bayesian calibration of inexact computer models. *Journal of the American Statistical Association* **112**, 1274–1285.
- Raftery, A., Madigan, D. and Volinsky, C. (1996) Accounting for model uncertainty in survival analysis improves predictive performance. In *Bayesian Statistics 5* (eds. J. Bernardo, J. Berger, A. Dawid and A. Smith) 323–349. Oxford University Press.
- Ribatet, M., Cooley, D. and Davison, A. (2012) Bayesian inference from composite likelihoods with an application to spatial extremes. *Statistica Sinica* **22**, 813–845.
- Smith, E. and Stephenson, A. (2009) An extended Gaussian max-stable process for spatial extremes. *Journal of Statistical Planning and Inference* **139**, 1266–1275.
- Tuo, R. and Wu, C. (2015) Efficient calibration for imperfect computer models. *Annals of Statistics* **43**, 2331–2352.
- (2016) A theoretical framework for calibration in computer models: Parametrization, estimation and convergence properties. *SIAM/ASA Journal of Uncertainty Quantification* **4**, 767–795.
- Ventura, L., Cabras, S. and Racugno, W. (2010) Default prior distributions from quasi- and quasi-profile likelihoods. *Journal of Statistical Planning and Inference* **140**, 2937–2942.
- Wahba, G. (1990) *Spline models for observational data* vol. 59. Society for Industrial Mathematics.



Wong, R., Storlie, C. and Lee, T. (2017) A frequentist approach to computer model calibration. *Journal of the Royal Statistical Society Series B* **79**, 635–648.

Woody, S., Ghaffari, N. and Hund, L. (2019) Bayesian model calibration for extrapolative prediction via Gibbs posteriors. *arXiv:1909.05428v1*.

Xie, F. and Xu, Y. (2020) Bayesian projected calibration of computer models. *Journal of the American Statistical Association* **To appear**.