

# Renormalization in combinatorially non-local field theories: the BPHZ momentum scheme

Johannes Thürigen

Mathematisches Institut der WWU Münster, Einsteinstr. 62, 48149 Münster, Germany, EU  
Institut für Physik/Mathematik, HU Berlin, Unter den Linden 6, 10099 Berlin, Germany, EU

E-mail: johannes.thuerigen@uni-muenster.de

URL: <http://www.uni-muenster.de/mathphys>

Dedicated to Dirk Kreimer on the occasion of his 60th birthday

**Abstract.** Various combinatorially non-local field theories are known to be renormalizable. Still, explicit calculations of amplitudes are very rare and restricted to matrix field theory. In this contribution I want to demonstrate how the BPHZ momentum scheme in terms of the Connes-Kreimer Hopf algebra applies to any combinatorially non-local field theory which is renormalizable. This algebraic method improves the understanding of known results in noncommutative field theory in its matrix formulation. Furthermore, I use it to provide the first explicit perturbative calculations of amplitudes in tensorial field theories of rank  $r > 2$ .

*Key words:* non-local field theory; renormalization; Hopf algebras; multiple polylogarithms

## 1 Introduction

Since Dirk Kreimer's seminal work [29] it is well known that a Hopf algebra of divergent Feynman diagrams is the universal structure underlying renormalization in perturbative quantum field theory [17, 18, 30]. A classical example is BPHZ momentum subtraction as used for the general proof of renormalization [8, 27, 49]. In [47] I have shown that the Connes-Kreimer Hopf algebra describes renormalization not only in field theories with point-like interactions but also with full generality in combinatorially non-local field theories (cNLFT) such as non-commutative and matrix field theory [23, 48, 28], tensorial field theories [4, 2] and group field theory [35, 20, 14].

So far, explicit calculations of renormalized amplitudes in cNLFT are only known for non-commutative field theory and its matrix-field representation using the BPHZ momentum scheme [6, 7, 22, 28]. In fact, exact solutions for matrix field theory have been found recently [38, 21, 12] and the perturbative results are mainly used as a check of consistency [22, 28]. For other examples of cNLFT, renormalizability has been proven for various field theories with tensorial interactions [4, 3, 2, 15, 16, 14] but no explicit results of amplitudes, let alone of full renormalized correlation functions, are known<sup>1</sup>. Using the Hopf-algebraic structure of renormalization I will give explicit amplitudes of perturbative tensorial field theory here for the first time.

Explicit perturbative calculations in matrix field theory use a version of Zimmermann's forest formula [49] adapted to ribbon graphs [22]. While agreement with exact non-perturbative solutions shows that this implementation of the BPHZ momentum scheme is perfectly valid, it uses a very peculiar definition of subdiagrams which are not themselves ribbon graphs, thus strictly speaking in a different class of diagrams. Here I will show how BPHZ momentum subtraction can be defined on any cNLFT in Kreimer's way using the Hopf algebra of divergent diagrams and a subtraction operator as a Taylor expansion which is a Rota-Baxter operator on the algebra of amplitudes. This gives two results: First, it reproduces in a very natural way the known perturbative calculations in matrix field theory [22, 28]. Second, it provides the algorithm to calculate

---

<sup>1</sup>Exact solutions similar to matrix field theory [38] have been found in [39] for a rank-3 tensor field; due to the specific structure of its tensorial interaction, a closer look reveals that it is still a matrix field theory in disguise.

renormalized amplitudes in any cNLFT in a straightforward way. I will demonstrate this for the example of tadpole and sunrise diagrams in field theory with tensor-invariant interactions.

The physical meaning of amplitudes in cNLFT is not as obvious as for scattering events of high energy particles described by standard-model quantum field theories. Matrix field theory can be understood as a representation of field theory of elementary particles on non-commutative (Moyal-deformed) spacetime which still allows for such scattering interpretation. On the other hand, one main application of theories with tensor fields of higher rank  $r > 2$  are models of quantum gravity as their perturbative series is a sum over  $r$ -dimensional combinatorial (pseudo) manifolds [24] and one can build models which have gravitational amplitudes on such discrete manifolds as lattice gauge theories [35, 20, 14], known as spin foam models [40]. Then, the main challenge is to recover continuous  $D = 4$  dimensional spacetime in some critical regime of the theory [43, 36] in the sense of continuous random geometries [1, 34, 33]; to this end, non-perturbative results like [41] are necessary. However, also perturbation theory might be the right starting point towards this goal, either by identification of appropriate linear combinations of amplitudes in the perturbative series related to non-perturbative structures as found for example via topological recursion [12, 11] or exploiting the Hopf-algebraic structure of perturbative renormalization leading to Kreimer's combinatorial Dyson-Schwinger equations [13, 31, 32, 5]. Thus, the explicit perturbative results presented here might also be seen as a starting point to investigate the non-perturbative regime of cNLFT.

## 2 Feynman rules in cNLFT

In the following I introduce the general structure of renormalization as captured by a Hopf algebra [29, 30]. In parallel, I explain its application to combinatorially non-local field theory. This is largely based on [47] where one can find the precise details of combinatorial and algebraic definitions and statements. Here, the emphasis is on the practical use for perturbative calculations in field theory.

A perturbative expansion in coupling constants is a formal power series labelled by Feynman diagrams. Feynman rules associate to each diagram  $\Gamma$  an amplitude  $A_\Gamma$ , i.e. the coefficient in this power series. This can be understood as a map from diagrams to amplitudes

$$A : \mathcal{G} \rightarrow \mathcal{A}, \quad \Gamma \mapsto A(\Gamma) = A_\Gamma. \quad (1)$$

In the perturbative series these amplitudes are summed and for renormalization it will also be necessary to multiply them which demands an algebra  $\mathcal{A}$  of amplitudes. This structure is mirrored in a  $\mathbb{Q}$ -algebra  $\mathcal{G}$  generated by the set of diagrams in which multiplication is simply the disjoint union of diagrams,  $m_{\mathcal{G}}(\Gamma_1 \otimes \Gamma_2) := \Gamma_1 \sqcup \Gamma_2$ . Both algebras are unital with unit  $u_{\mathcal{G}} : \mathbb{Q} \rightarrow \mathcal{G}, q \mapsto q\mathbb{1}$  where  $\mathbb{1} \in \mathcal{G}$  is the empty diagram and  $u_{\mathcal{A}}$  maps to the amplitude  $A(\mathbb{1}) = 1$ . The map  $A$  must then be an algebra homomorphism.

A combinatorially non-local field theory (cNLFT) is characterized by a pairing of arguments at interaction vertices which leads to stranded Feynman diagrams, also called *2-graphs* [47]. That is, order- $n$  interactions are of the form

$$S_{\text{IA}}[\phi] = \lambda_\gamma \int \prod_{i=1}^n d\mathbf{p}_i \prod_{(ia,jb)} \delta(p_i^a - p_j^b) \prod_{i=1}^n \phi(\mathbf{p}_i) \quad (2)$$

where the field  $\phi$  depends on  $r$  arguments  $\mathbf{p} = (p^1, \dots, p^r)$  of which each  $p^a$  is in a  $d$ -dimensional manifold. For simplicity I will consider  $p^a \in \mathbb{R}^d$  throughout this paper. The product over pairs  $(ia, jb)$  means that for each argument  $p_i^a$  there is a convolution in terms of the Dirac distribution  $\delta(p_i^a - p_j^b)$  with exactly one other argument  $p_j^b$ . Diagrammatically, these convolutions cannot be

fully captured by a vertex in a graph but it is necessary to add a second layer called strands. For example,




is the diagrammatic representation of an interaction vertex of  $n = 4$  fields (red) with  $r = 3$  arguments each (green) which are convoluted pairwise (blue dots); this is equivalent to, on the right-hand side of the equation, a vertex (black) with half-edges (red) and additional edges (green) adjacent to these half-edges which characterize the interaction vertex by a graph. Thus, interactions are labelled in general by *vertex graphs*  $\gamma$  and so are their coupling constants  $\lambda_\gamma$ . A set of graphs  $\mathbf{G}_1^v$  determines the diagrammatics of a given theory.

Feynman diagrams follow as usual by Wick contraction of the fields according to a quadratic Gaussian part of the action. The arguments  $p^a$  of the field  $\phi$  are not necessarily ordered, i.e. the quadratic part might pair the  $r$  arguments according to an arbitrary permutation  $\sigma_2$ ,

$$S_0(\phi) = \frac{1}{2} \int d\mathbf{p}_i d\mathbf{p}_j \phi(\mathbf{p}_i) \frac{\prod_{a=1}^r \delta(p_i^a - p_j^{\sigma_2(a)})}{\sum_{a=1}^r |p_i^a|^{2\zeta} + \mu} \phi(\mathbf{p}_j), \quad (4)$$

where  $\mu$  is a mass parameter and the scaling  $2\zeta$  of the propagator is in general parametrized by a  $\zeta > 0$ , usually  $\zeta \in ]0, 1]$ . This leads to the general class of 2-graphs defined in [47] based on [37, 46]. Still, in many theories this permutation is fixed. For example, in Hermitian matrix field theory [48, 28], the 2-graphs are combinatorial maps (ribbon graphs) and  $\sigma_2$  is fixed by the orientation of the vertices, e.g. in



the pairing of strands adjacent to half edges  $i \in \{1, 2, 3, 4, 5, 6, 7\}$  along the edges  $\{1, 2\}$ ,  $\{3, 5\}$  and  $\{4, 6\}$  is determined by the orientation of the vertices (1), (234) and (576) (see Fig. 2 in [47] for further details). Following a strand through the 2-graph defines a face which is an internal face if the strand is closed and external otherwise. In the example  $\Gamma$  there is one internal face (3564) and one external face (764212357). The external structure of a 2-graph  $\Gamma$  is the vertex graph  $\gamma$  obtained by deleting all internal edges and internal faces which is the boundary  $\gamma = \partial\Gamma$ .

The above Gaussian and interaction parts of the action, Eq.(4) and Eq.(2), lead to an identification of variables  $p^a$  along strands and thus to the following Feynman rules: For a 2-graph  $\Gamma$ , the amputated amplitude  $A_\Gamma$  is an integral consisting of

1. a Lebesgue integral  $\int_{\mathbb{R}^d} dq_f$  for each internal face  $f \in \mathcal{F}_\Gamma^{\text{int}}$ , integrating over the variable identifying  $q_f = p_i^a$  for all strands  $a$  at half edges  $i$  along  $f$ ,
2. a propagator factor  $\frac{1}{\sum_{a=1}^r |p_i^a|^{2\zeta} + \mu}$  for each internal edge  $e = \{i, j\} \in \mathcal{E}_\Gamma$ ,
3. a coupling constant  $\lambda_\gamma$  for each vertex  $v \in \mathcal{V}_\Gamma$  with vertex graph  $\gamma$  and
4. the symmetry factor  $\frac{1}{|\text{Aut } \Gamma|}$  given by the group of automorphisms  $\text{Aut } \Gamma$  under relabelling of internal strands, half edges and vertices in  $\Gamma$ .

These rules define the algebra homomorphism  $A : \mathcal{G} \rightarrow \mathcal{A}$  to the algebra  $\mathcal{A}$  of integral functions with rational functions as integrands,

$$A_\Gamma : \{p_f\}_{f \in \tilde{\mathcal{F}}_\Gamma^{\text{ext}}} \mapsto A_\Gamma(\{p_f\}) := \frac{1}{|\text{Aut } \Gamma|} \prod_{v \in \mathcal{V}_\Gamma} \lambda_\gamma \prod_{f \in \mathcal{F}_\Gamma^{\text{int}}} \int_{\mathbb{R}_+^d} dq_f \prod_{\{i,j\} \in \mathcal{E}} \frac{1}{\sum_{a=1}^r |p_i^a|^{2\zeta} + \mu}. \quad (6)$$

They depend on an external variables  $p_f$  for each external face  $f \in \tilde{\mathcal{F}}_\Gamma^{\text{ext}}$  adjacent to *at least one internal edge*, given again by identifying  $p_f = p_i^a$  for all half edges  $i$  and adjacent strands  $a$  along  $f$ . Importantly, only faces in this subset  $\tilde{\mathcal{F}}_\Gamma^{\text{ext}} \subset \mathcal{F}_\Gamma^{\text{ext}}$  of external faces play a role: If a face is not adjacent to any internal edge, there is no propagator in which the corresponding variable occurs and the amputated amplitude  $A_\Gamma$  does not depend on it. This is in contrast to local field theory where loop amplitudes always depend on all external variables.

Multiplication in the algebra  $\mathcal{A}$  of amplitudes is given by multiplication of integrands. That is, the product of two amplitudes  $A, B \in \mathcal{A}$  with  $A(p_1, \dots, p_m) = c_A \int \prod_i dq_i I_A(p_1, \dots, p_m, q_1, \dots, q_k)$  and  $B(r_1, \dots, r_n) = c_B \int \prod_j ds_j I_B(r_1, \dots, r_n, s_1, \dots, s_l)$  where the integrands  $I_A$  and  $I_B$  are rational functions is

$$(A \cdot B)(p_a, r_b) := c_{ACB} \int \prod_{i=1}^k dq_i \prod_{j=1}^l ds_j I_A(p_1, \dots, p_m, q_1, \dots, q_k) I_B(r_1, \dots, r_n, s_1, \dots, s_l). \quad (7)$$

This is relevant since some external variables of  $p_a$  of the amplitude  $A$  might coincide with some integration variables  $s_j$  of  $B$ , i.e.  $p_a = s_j$  for some  $a \in \{1, \dots, n\}$  and  $j \in \{1, \dots, l\}$ , and vice versa. Accordingly, these are not external variables of the product  $A \cdot B$ .

To give an example of an amplitude, for  $\Gamma$  in Eq. (5) there is one external variable  $p$  for the external face and one integration variable  $q$  for the internal face leading to the amputated amplitude

$$A_\Gamma(p) = A \left( \text{Diagram} \right) (p) = \frac{1}{2} \lambda_\gamma \lambda_\gamma^2 \int_{\mathbb{R}^d} dq \frac{1}{2|p|^{2\zeta} + \mu} \frac{1}{(|p|^{2\zeta} + |q|^{2\zeta} + \mu)^2}. \quad (8)$$

For  $d \geq 2\zeta$  this integral does not converge. To render the algebra of formal amplitudes  $\mathcal{A}$  an algebra of well defined functions it is necessary to introduce a regularization cutoff  $\Lambda$ , that is consider the algebra  $\mathcal{A}^\Lambda$  of the same integrals but integrating all internal variables over the domain  $|q|^{2\zeta} \leq \Lambda^{2\zeta}$ ,

$$A_\Gamma^\Lambda(\{p_f\}) := \frac{1}{|\text{Aut } \Gamma|} \prod_{v \in \mathcal{V}_\Gamma} \lambda_\gamma \prod_{f \in \mathcal{F}_\Gamma^{\text{int}}} \int_\Lambda dq_f \prod_{\{i,j\} \in \mathcal{E}} \frac{1}{\sum_{a=1}^r |p_i^a|^{2\zeta} + \mu}. \quad (9)$$

For a well-defined limit  $\Lambda \rightarrow \infty$  renormalization is needed.

### 3 BPHZ renormalization using the Hopf algebra of 2-graphs

To renormalize a divergent amplitude one subtracts a counter term which exactly eliminates the divergent part and leaves a finite result. To this end one needs a measure of how much a divergent amplitude is divergent. A useful starting point is the power  $\omega_\Gamma$  of the asymptotic scaling in the cutoff  $\Lambda$  of the regularized amplitude Eq. (9), i.e.  $A_\Gamma^\Lambda \sim \Lambda^{\omega_\Gamma}$ . A simple way to determine this quantity is to count the number of integrals and propagators in the integration variables in the amplitude. For Eq. (9) this gives the *superficial degree of divergence*

$$\omega^{\text{sd}}(\Gamma) = d F_\Gamma - 2\zeta E_\Gamma + \sum_{v \in \mathcal{V}_\Gamma} \omega(\gamma_v) \quad (10)$$

where  $F_\Gamma = |\mathcal{F}_\Gamma^{\text{int}}|$  is the number of internal faces,  $E_\Gamma = |\mathcal{E}_\Gamma|$  is the number of (internal) edges and a scaling  $\lambda_\gamma \sim \Lambda^{\omega(\gamma)}$  of the couplings with vertex graph  $\gamma = \gamma_v$  for each vertex  $v \in \mathcal{V}_\Gamma$  in the 2-graph  $\Gamma$  is assumed in addition. A 2-graph  $\Gamma$  is *superficially divergent* iff  $\omega^{\text{sd}}(\Gamma) \geq 0$ . These properties are specific to a given combinatorially non-local field theory  $T$  which is defined by a set of interactions given by its vertex graphs  $\mathbf{G}_1^v$  together with their scaling weights  $\omega : \mathbf{G}_1^v \rightarrow \mathbb{Z}$ , the propagators with scaling  $2\zeta$  and the dimension  $d$ .

The theorem of Bogoliubov and Parasiuk [8] with proof completed by Hepp [27] and further improvements by Zimmermann leading to the forest formula [49] states that in a renormalizable local quantum field theory all divergences can be renormalized by counter terms for each superficially divergent one-particle irreducible (1PI) diagram. Kreimer [29] has further significantly improved the understanding of the combinatorics of the forest formula (in particular, and renormalization in much more general) showing that it has an algebraic structure given by a Hopf algebra of divergent Feynman diagrams. In [47] I have shown that this Hopf algebra exists not only in field theories with point-like interactions but also for theories with any combinatorially non-local interactions, in this way generalizing and improving similar statements for some specific examples [45, 44, 42]. Here I demonstrate how this explicitly works using the BPHZ momentum scheme.

The challenge to define the right counter term to renormalize a divergent amplitude lies in the nested structure of divergent subdiagrams. In cNLFT, just like in local quantum field theory, a 2-graph  $\Theta$  is a *subgraph*  $\Theta \subset \Gamma$  of a 2-graph  $\Gamma$  iff it has a subset of edges  $\mathcal{E}_\Theta \subset \mathcal{E}_\Gamma$ , including the adjacent strands, and the same structure otherwise [47]. A *primitively divergent* 2-graph is then one which has no proper subgraphs which are superficially divergent. On a primitively divergent 2-graphs  $\Gamma$  one defines the renormalized amplitude  $A_R(\Gamma)$  directly by a subtraction operation which is a linear operator  $R : \mathcal{A} \rightarrow \mathcal{A}$  as

$$A_R : \mathcal{G} \rightarrow A_R(\Gamma) := (A - R \circ A)(\Gamma). \quad (11)$$

The explicit form of  $R$  is exactly what specifies a particular renormalization scheme. For concreteness, let us here define  $R$  for the BPHZ momentum scheme of cNLFT:

**Definition 1** (BPHZ momentum scheme). The operator  $R : \mathcal{A} \rightarrow \mathcal{A}$  on the unital commutative algebra  $\mathcal{A}$  of functions of the form  $A_\Gamma^\Lambda$ , Eq. (9), with degree of divergence  $\omega(A_\Gamma^\Lambda) := \omega^{\text{sd}}(\Gamma)$ , Eq. (10), is given by the multivariate Taylor expansion  $T^\omega$  of order  $\omega = \omega(A_\Gamma^\Lambda)$ ,

$$R(A_\Gamma^\Lambda)(\{p_f\}) := \left( T_{\{p_f\}}^\omega A_\Gamma^\Lambda \right) (\{p_f\}) = \sum_{|\vec{k}| \leq \omega(A_\Gamma^\Lambda)} \frac{1}{\vec{k}!} \frac{\partial^{|\vec{k}|} A_\Gamma^\Lambda}{\prod_f \partial p_f^{k_f}} (\{p_f = 0\}) \prod_{f \in \tilde{\mathcal{F}}_\Gamma^{\text{ext}}} p_f^{k_f}, \quad (12)$$

where the sum is over multi-indices  $\vec{k}$  of length  $|\vec{k}| = \sum_{f \in \tilde{\mathcal{F}}_\Gamma^{\text{ext}}} k_f$ . Note that the underlying 2-graph structure is only used to encode the form of the integral function  $A(\{p_f\}) = A_\Gamma^\Lambda(\{p_f\})$  but the definition of  $R$  is otherwise independent of it.

For divergent diagrams  $\Gamma$  with subdivergences the whole counter term is a nested sum over counter terms for the various superficially divergent subgraphs according to Zimmermann's forest formula [49]. These terms are products of counter terms of subdiagrams  $\Theta \subset \Gamma$  and amplitudes of remaining diagrams in which the subdiagram is shrunken to a vertex, that is contractions  $\Gamma/\Theta$ . For 2-graphs  $\Theta \subset \Gamma$  the definition of such contraction  $\Gamma/\Theta$  is that there is a vertex for each connected component of  $\Theta$  and all internal edges and internal strands (and thus internal faces) of  $\Theta$  are deleted (see Def. 3.2 in [47] for further detail). This defines a coproduct  $\Delta$  on the algebra of superficially divergent 2-graphs  $\mathcal{H}_T^{\text{f2g}}$  of a given renormalizable theory  $T$  as

$$\Delta : \mathcal{H}_T^{\text{f2g}} \rightarrow \mathcal{H}_T^{\text{f2g}} \otimes \mathcal{H}_T^{\text{f2g}}, \quad \Delta \Gamma := \sum_{\Theta \subset \Gamma, \Theta \in \mathcal{H}_T^{\text{f2g}}} \Theta \otimes \Gamma/\Theta. \quad (13)$$

For this it is crucial that  $\mathcal{H}_T^{\text{f2g}}$  is closed under contractions which is satisfied due to the property of renormalizability  $\omega(\partial\Gamma) = \omega^{\text{sd}}(\Gamma)$  in  $T$  [47]. Together with the counit  $\epsilon : \mathcal{H}_T^{\text{f2g}} \rightarrow \mathbb{Q}$  defined as the projector to 2-graphs without any edges,  $\mathcal{H}^{\text{f2g}}$  is a coalgebra and due to compatibility with multiplication also a bialgebra ([47], Prop. 4.2).

To implement the forest formula, the crucial object is an inverse to the algebra homomorphism of amplitudes  $A : \mathcal{H}^{\text{f2g}} \rightarrow \mathcal{A}$  with respect to the convolution product, i.e.

$$\phi * \psi := m_{\mathcal{A}} \circ (\phi \otimes \psi) \circ \Delta \quad (14)$$

for arbitrary such homomorphisms  $\phi, \psi : \mathcal{H}^{\text{f2g}} \rightarrow \mathcal{A}$ . Such inverse is a natural consequence if  $\mathcal{H}^{\text{f2g}}$  is a Hopf algebra, i.e. has a *coinverse*, also called *antipode*,  $S : \mathcal{H}^{\text{f2g}} \rightarrow \mathcal{H}^{\text{f2g}}$  defined by  $S * \text{id} = \text{id} * S = u \circ \epsilon$ . In fact, one can show that the bialgebra  $\mathcal{H}_T^{\text{f2g}}$  of superficially divergent 2-graphs in  $T$  is a Hopf algebra ([47] Thm. 5.1) using the possibility to express  $S$  recursively as

$$S = -(S * P) \quad (15)$$

where  $P = \text{id} - u \circ \epsilon$  is the projector to the augmentation ideal, i.e. all 2-graphs in  $\mathcal{H}_T^{\text{f2g}}$  with at least on edge [10]. This works because  $\mathcal{H}^{\text{f2g}}$  is graded with respect to the number of edges. As a direct consequence, the inverse of an algebra homomorphism  $\phi$  is  $S^\phi = S \circ \phi$ . Both facts can now be used to recursively define the counter term

$$\begin{aligned} S_{\text{R}}^{\text{A}} : \mathcal{H}^{\text{f2g}} \rightarrow \mathcal{A}, \quad S_{\text{R}}^{\text{A}}(r) = 1, \quad S_{\text{R}}^{\text{A}}(\Gamma) &= -R[(S_{\text{R}}^{\text{A}} * A \circ P)(\Gamma)] \\ &= - \sum_{\Theta \subsetneq \Gamma, \Theta \in \mathcal{H}_T^{\text{f2g}}} R[S_{\text{R}}^{\text{A}}(\Theta)A(\Gamma/\Theta)] \end{aligned} \quad (16)$$

for any  $\Gamma$  in the augmentation ideal and  $r$  edgeless. Then, the renormalized amplitude is simply

$$\boxed{A_{\text{R}} := S_{\text{R}}^{\text{A}} * A.} \quad (17)$$

The counter term map  $S_{\text{R}}^{\text{A}}$  is an algebra homomorphism, in particular  $S_{\text{R}}^{\text{A}}(\Gamma_1\Gamma_2) = S_{\text{R}}^{\text{A}}(\Gamma_1)S_{\text{R}}^{\text{A}}(\Gamma_2)$ , if the subtraction map  $R$  is a Rota-Baxter operator, that is satisfies

$$R[AB] + R[A]R[B] = R[R[A]B + AR[B]]. \quad (18)$$

**Proposition 1.** *The BPHZ momentum subtraction operator  $R = T^\omega$  (Def. 1) is a Rota-Baxter operator.*

The property Eq. (18) can be checked by explicit calculations using the product rule for multivariate derivatives.

## 4 BPHZ renormalization in tensorial $\phi_{2,2}^4$ and $\phi_{1,3}^4$ theory

To demonstrate how BPHZ renormalization works, I give the example of tadpole and sunrise amplitudes in the just renormalizable tensorial field theories of rank  $r = 2$  and  $r = 3$ . A tensorial  $\phi_{d,r}^n$  theory is a cNLFT of a scalar field  $\phi$  with  $r$  arguments in a  $d$ -dimensional manifold with tensor-invariant interactions of maximal order  $n$  (degree of the vertex) [2]. An interaction is tensor-invariant if its vertex graph  $\gamma$  is  $r$ -regular and edge-coloured, i.e. each vertex in  $\gamma$  is adjacent to exactly  $r$  edges with distinct labels  $c = 1, 2, \dots, r$  [9]. This corresponds to fixing the position of each argument in the field and pairing only arguments in the same position. As a consequence, also the 2-graphs  $\Gamma$  labelling the perturbative expansion can be represented as

coloured graphs<sup>2</sup>, now with  $r + 1$  colours. In the following I will consider a complex field  $\phi$ , just for the reason that the vertex graphs then have to be furthermore bipartite which simplifies the theory space further.

With weight  $\omega(\gamma_v) = d_r - d^v(d_r - 2\zeta)/2$  for each  $d^v$ -valent vertex with vertex graph  $\gamma_v$ , the superficial degree of divergence in tensorial  $\phi_{d,r}^n$  theory is [2, 47]

$$\omega^{\text{sd}}(\Gamma) = d_r - \frac{d_r - 2\zeta}{2} V_{\partial\Gamma} - d \left( \frac{2\omega_\Gamma^{\text{G}} - 2\omega_{\partial\Gamma}^{\text{G}}}{(r-1)!} + K_{\partial\Gamma} - 1 \right) \quad (19)$$

where  $d_r = d(r-1)$  is the equivalent of dimension compared to local field theory,  $K_{\partial\Gamma}$  is the number of connected components of the boundary graph and  $\omega^{\text{G}} \in \mathbb{N}$  is the Gurau degree of a coloured graph [25, 26]. It is the topological quantity generalizing the genus  $g$  in the  $r = 2$  case, though not a topological invariant. Crucially, the overall Gurau degree is conserved under the coproduct [42] such that renormalizability is given even though  $\omega^{\text{sd}}(\Gamma) = \omega(\partial\Gamma)$  only up to  $\omega_\Gamma^{\text{G}}$ . One can further show that  $\omega_\Gamma^{\text{G}} \geq \omega_{\partial\Gamma}^{\text{G}}$  [4] such that the whole term in brackets is always greater or equal to zero and renormalizable theories are determined by the first two terms. Thus, a tensorial  $\phi_{d,r}^n$  theory is super renormalizable if  $d_r = 2\zeta$  and just renormalizable if  $n = \frac{2d_r}{d_r - 2\zeta}$ .







Just-renormalizable quartic theories are those with  $d_r = d(r-1) = 4\zeta$ . The tensorial equivalent to standard local  $\phi_4^4$  theory with quadratic propagator  $\zeta = 1$  are therefore theories with  $d_r = 4$ , that is theories with  $(d, r) = (4, 2), (2, 3)$  or  $(1, 5)$ . Let us here simplify explicit renormalization even further and chose  $\zeta = 1/2$ . The two just renormalizable theories are then tensorial  $\phi_{2,2}^4$  and  $\phi_{1,3}^4$  theory. The divergence degree in these two theories is

$$\omega^{\text{sd}}(\Gamma) = dF - E = 2 - \frac{1}{2} V_{\partial\Gamma} - \begin{cases} 2(2g_\Gamma + K_{\partial\Gamma} - 1) & \text{for } \phi_{2,2}^4 \text{ theory} \\ (\omega_\Gamma^{\text{G}} - g_{\partial\Gamma} + K_{\partial\Gamma} - 1) & \text{for } \phi_{1,3}^4 \text{ theory} \end{cases} \quad (20)$$

In these two cases I will present the explicit BPHZ-renormalized amplitudes  $A_\Gamma$  for the tadpole and sunrise diagrams. Complex  $\phi_{2,2}^4$  theory is very similar to the quartic Hermitian matrix field theory known as the Grosse-Wulkenhaar model or quartic Kontsevich model [23, 48, 28], i.e. has the same power counting and shares most diagrams for which the amplitudes agree; for this theory I simply reproduce known results [28] with the Hopf algebra method. In the rank  $r = 3$  tensorial case, to the best of my knowledge, the calculations presented are the first explicit results of amplitudes in tensorial field theory.

### The tadpole in $\phi_{2,2}^4$ theory

The only primitive 2-point amplitude in  $\phi_{2,2}^4$  theory is the one for the tadpole 2-graph. There

are two versions,   $\cong c$   with colour  $c = 1$  and   $\cong c$   with  $c = 2$  when using the convention that in the representation via combinatorial maps  and  one has always the external variable  $p_1$  associated to the face of colour  $c = 1$  on the upper face and  $p_2$  respectively on the lower face. Its divergence degree is  $\omega^{\text{sd}}(\text{tadpole}) = \omega^{\text{sd}}(\text{tadpole}) = 2 - 1 = 1$ . There are no

<sup>2</sup>Strictly speaking, this is only true for theories without multi-trace vertices, that is vertices whose vertex graphs have more than one connected component. Otherwise, the full coloured 2-graph structure is necessary [47].

subdivergences and thus by definition Eq. (9)

$$\begin{aligned}
A_{\text{R}}\left(\text{diagram}_1\right) &= (S_{\text{R}}^{\text{A}} * A)\left(\text{diagram}_1\right) \\
&= m_{\mathcal{A}}(S_{\text{R}}^{\text{A}} \otimes A)\Delta\left(\text{diagram}_1\right) = m_{\mathcal{A}}(S_{\text{R}}^{\text{A}} \otimes A)\left(\text{diagram}_2 \otimes \text{diagram}_3 + \text{diagram}_4 \otimes \text{diagram}_5\right) \\
&= A\left(\text{diagram}_1\right) + S_{\text{R}}^{\text{A}}\left(\text{diagram}_1\right)A\left(\text{diagram}_6\right) = A\left(\text{diagram}_1\right) - R\left[A\left(\text{diagram}_1\right)\right]
\end{aligned} \tag{21}$$

since  $A(\text{diagram}_6) = 1$  as there are no edges in a single vertex. This calculation works for any primitive diagram and thus validates Eq. (11). Explicit calculation in the BPHZ momentum scheme according to Def. 1 gives then

$$\begin{aligned}
A_{\text{R}}\left(\text{diagram}_1\right)(p_1) &\equiv A_{\text{R}}\left(p_1 \text{diagram}_1^{\tilde{q}_2}\right) = \lambda_{\text{diagram}_7} (1 - T_{p_1}^1) \int_{\mathbb{R}^2} dq_2 \frac{1}{|p_1| + |q_2| + \mu} \\
&= \lambda_{\text{diagram}_7} \int_0^\infty \pi |q_2| d|q_2| \left( \frac{1}{|p_1| + |q_2| + \mu} - \frac{1}{|q_2| + \mu} + \frac{|p_1|}{(|q_2| + \mu)^2} \right) \\
&= \pi \mu \lambda_{\text{diagram}_7} \left( (\tilde{p}_1 + 1) \log(\tilde{p}_1 + 1) - \tilde{p}_1 \right)
\end{aligned} \tag{22}$$

with  $\tilde{p}_1 = |p_1|/\mu$ . The tadpole amplitude for colour  $c = 2$  gives the same with  $p_2$  instead of  $p_1$ . This result is exactly the same as in Hermitian matrix field theory [28].

### Tadpole diagrams in $\phi_{1,3}^4$ theory

In  $\phi_{1,3}^4$  theory, the tadpole has not only versions for each colour but also two possibilities where to have the edge. In particular, the two differ in degree of divergence (using  $\omega^{\text{sd}}(\Gamma) = F - E$ ),

$$\omega^{\text{sd}}\left(\text{diagram}_8\right) = 2 - 1 = 1 \quad \text{and} \quad \omega^{\text{sd}}\left(\text{diagram}_9\right) = 1 - 1 = 0. \tag{23}$$

The renormalized amplitude of the first one for  $c = 1$  is

$$\begin{aligned}
A_{\text{R}}\left(p_1 \text{diagram}_8\right) &= \lambda_{\text{diagram}_{10}} (1 - T_{p_1}^1) \int_{\mathbb{R}} dq_2 \int_{\mathbb{R}} dq_3 \frac{1}{|p_1| + |q_2| + |q_3| + \mu} \\
&= \lambda_{\text{diagram}_{10}} \int_0^\infty 2dq_2 \int_0^\infty 2dq_3 \left( \frac{1}{|p_1| + q_2 + q_3 + \mu} - \frac{1}{q_2 + q_3 + \mu} + \frac{|p_1|}{(q_2 + q_3 + \mu)^2} \right) \\
&= 4\mu \lambda_{\text{diagram}_{10}} \left( (\tilde{p}_1 + 1) \log(\tilde{p}_1 + 1) - \tilde{p}_1 \right)
\end{aligned} \tag{24}$$

with  $\tilde{p}_1 = |p_1|/\mu$ . Up to a constant it is the same as in the  $r = 2$  matrix case, Eq. (22). This is not surprising as the external variable is the same and the two internal faces with one-dimensional integration variables  $q_2$  and  $q_3$  behaves effectively like one face with a two-dimensional variable.

The first amplitude specific to a higher rank  $r = 3$  is

$$\begin{aligned}
A_{\text{R}}\left(p_2, p_3 \text{diagram}_9^{\tilde{q}_1}\right) &= \lambda_{\text{diagram}_{11}} (1 - T_{p_2, p_3}^0) \int_{\mathbb{R}} dq_1 \frac{1}{|q_1| + |p_2| + |p_3| + \mu} \\
&= \lambda_{\text{diagram}_{11}} \int_0^\infty 2dq_1 \left( \frac{1}{q_1 + |p_2| + |p_3| + \mu} - \frac{1}{q_1 + \mu} \right) = -2\lambda_{\text{diagram}_{11}} \log(\tilde{p}_{23} + 1)
\end{aligned} \tag{25}$$

with  $\tilde{p}_{23} = (|p_2| + |p_3|)/\mu$ . In the full 1PI 2-point Green's function one has a sum over the different colours  $c = 1, 2, 3$  for which in general the couplings  $\lambda_{\text{diagram}_{11}^c}$  are distinct.



### The sunrise diagram in $\phi_{2,2}^4$ theory

The 2-graph version of the sunrise diagram is the first example which shows the Hopf-algebraic formulation of Zimmermann's forest formula in its full beauty. In  $\phi_{2,2}^4$  theory the sunrise combinatorial map corresponds to the colouring

$$\text{Sunrise} \cong \text{Sunrise}^c \quad \text{and has } \omega^{\text{sd}}(\text{Sunrise}) = 2 \cdot 2 - 3 = 1. \quad (26)$$

(The other choice of colouring of this 2-graph corresponds to a non-planar map of cylinder topology which is not divergent,  $\omega^{\text{sd}} = 0 - 3 = -3$ .) The sunrise 2-graph has three 1PI subgraphs of which only two are divergent,

$$\omega^{\text{sd}}(\text{Sunrise}^c) = \omega^{\text{sd}}(\text{Sunrise}^c) = 2 \cdot 1 - 2 = 0, \quad \omega^{\text{sd}}(\text{Sunrise}^c) = 0 - 2 = -2. \quad (27)$$

As they do not have any divergent 1PI subgraphs, they are primitives. Accordingly, the coproduct in  $\mathcal{H}^{\text{f2g}}$  is

$$\begin{aligned} \Delta(\text{Sunrise}^c) &= c \cdot \text{Sunrise}^c \otimes c \cdot \text{Sunrise}^c + c \cdot \text{Sunrise}^c \otimes c \cdot \text{Sunrise}^c \\ &+ c \cdot \text{Sunrise}^c \otimes c \cdot \text{Sunrise}^c + c \cdot \text{Sunrise}^c \otimes \text{Sunrise}^c \end{aligned} \quad (28)$$

which yields the renormalized amplitude

$$\begin{aligned} A_{\text{R}}(\text{Sunrise}^c) &= A(\text{Sunrise}^c) + S_{\text{R}}^{\text{A}}(\text{Sunrise}^c) A(\text{Sunrise}^c) \\ &+ S_{\text{R}}^{\text{A}}(\text{Sunrise}^c) A(\text{Sunrise}^c) + S_{\text{R}}^{\text{A}}(\text{Sunrise}^c) A(\text{Sunrise}^c). \end{aligned} \quad (29)$$

In the last term,  $S_{\text{R}}^{\text{A}}$  acts on a divergent diagram which is not primitive such that according to Eq. (16) there is a nontrivial sum

$$\begin{aligned} S_{\text{R}}^{\text{A}}(\text{Sunrise}^c) &= -R \left[ A(\text{Sunrise}^c) - R \left[ A(\text{Sunrise}^c) \right] A(\text{Sunrise}^c) \right. \\ &\quad \left. - R \left[ A(\text{Sunrise}^c) \right] A(\text{Sunrise}^c) \right]. \end{aligned} \quad (30)$$

Putting everything together and applying  $R = T^\omega$  one arrives at the finite integral

$$\begin{aligned}
A_R(p_1, p_2) &= \lambda_{\text{sunrise}}^2 (1 - T_{p_1, p_2}^1) \int_{\mathbb{R}^2} dq_1 \int_{\mathbb{R}^2} dq_2 \left( \frac{1}{|p_1| + |q_2| + \mu} \frac{1}{|q_1| + |q_2| + \mu} \frac{1}{|q_1| + |p_2| + \mu} \right. \\
&\quad + \frac{1}{|q_1| + |p_2| + \mu} (-T_{p_1, q_1}^0) \frac{1}{|p_1| + |q_2| + \mu} \frac{1}{|q_1| + |q_2| + \mu} \\
&\quad \left. + \frac{1}{|p_1| + |q_1| + \mu} (-T_{q_2, p_2}^0) \frac{1}{|q_1| + |q_2| + \mu} \frac{1}{|q_2| + |p_2| + \mu} \right) \\
&= \lambda_{\text{sunrise}}^2 \frac{\pi^2 \mu}{\tilde{p}_1 + \tilde{p}_2 + 1} \left[ \tilde{p}_1 \tilde{p}_2 \zeta_2 + (\tilde{p}_1 + \tilde{p}_2 + 1) \sum_{i=1,2} \left( (\tilde{p}_i + 1) \log(\tilde{p}_i + 1) - \tilde{p}_i \right) \right. \\
&\quad \left. - \prod_{i=1,2} (\tilde{p}_i + 1) \log(\tilde{p}_i + 1) + \sum_{i=1,2} \tilde{p}_i (\tilde{p}_i + 1) \text{Li}_2(-\tilde{p}_i) \right] \quad (31)
\end{aligned}$$

involving the polylogarithm  $\text{Li}_2$  and the zeta value  $\zeta_2 = \pi^2/6$  and again  $\tilde{p}_i = |p_i|/\mu$ . In general, any kind of multiple polylogarithms are to be expected due to the form of the integral. The result reproduces Eq. D.9 in [28]. As the diagram is symmetric under colour change, the amplitude is symmetric under exchange of  $p_1$  and  $p_2$ , i.e.  $A_R(p_2, p_1) = A_R(p_1, p_2)$ .

### The sunrise diagram in $\phi_{1,3}^4$ theory

Also in  $\phi_{1,3}^4$  there is only one divergent sunrise 2-graph out of four possibilities to join two quartic vertices by three edges (without tadpoles). This diagram has only a logarithmic divergence,

$$\omega^{\text{sd}} \left( \text{diagram} \right) = 3 - 3 = 0. \quad (32)$$

Out of the three 1PI subgraphs, in analogy to Eq. (27), only one is divergent which is

$$\omega^{\text{sd}} \left( \text{diagram} \right) = 2 - 2 = 0. \quad (33)$$

Thus, in rank  $r = 3$  there is one counter term less than in the  $r = 2$  case,

$$\begin{aligned}
A_R(p_1, p_2, p_3) &= A_R \left( \text{diagram} \right) = A \left( \text{diagram} \right) \\
&\quad + S_R^A \left( \text{diagram} \right) A \left( \text{diagram} \right) + S_R^A \left( \text{diagram} \right). \quad (34)
\end{aligned}$$

Expanding these counter terms in  $R$  expressions analogously to Eq. (30) and using the explicit subtraction operation  $R = T^\omega$ , this amplitude is

$$\begin{aligned}
A_R(p_1, p_2, p_3) &= \lambda_{\text{sunrise}}^2 \int_{\mathbb{R}} dq_1 \frac{1}{|q_1| + |p_2| + |p_3| + \mu} \\
&\quad \times (1 - T_{p_1, q_1}^0) \int_{\mathbb{R}} dq_2 \int_{\mathbb{R}} dq_3 \frac{1}{|q_1| + |q_2| + |q_3| + \mu} \frac{1}{|p_1| + |q_2| + |q_3| + \mu} \\
&= \lambda_{\text{sunrise}}^2 \frac{6}{1 + \tilde{p}_1 + \tilde{p}_{23}} \left[ \log(\tilde{p}_{23} + 1) \left( (\tilde{p}_1 + 1) \log(\tilde{p}_1 + 1) - \tilde{p}_1 - \tilde{p}_{23} - 1 \right) \right. \\
&\quad - \log(\tilde{p}_1) (\tilde{p}_1 + 1) \log(\tilde{p}_1 + 1) + \tilde{p}_{23} \left( \text{Li}_2 \left( -\frac{1}{\tilde{p}_{23}} \right) + \frac{1}{2} \log(\tilde{p}_{23}) + 2\zeta_2 \right) \\
&\quad \left. + (\tilde{p}_1 + 1) \left( \text{Li}_2 \left( \frac{1}{\tilde{p}_1 + 1} \right) + \frac{1}{2} \log(\tilde{p}_1 + 1) - \zeta_2 \right) \right] \quad (35)
\end{aligned}$$

where again  $\tilde{p}_1 = |p_1|/\mu$  and  $\tilde{p}_{23} = (|p_2| + |p_3|)/\mu$ . Note that the Taylor operator  $T_{p_1, p_2, p_3}^0$  in the first line acts on the whole expression which is following. One might use shuffle relations of multiple polylogarithms to further simplify the terms of weight two in  $\tilde{p}_1 + 1$  and  $\tilde{p}_{23}$ .

Interestingly, the calculation of the sunrise diagram in the  $r = 3$  theory is slightly simpler than in the  $r = 2$  case since the integral factorizes. This might be seen as a hint that individual amplitudes in higher rank tensorial theories are not more involved than in the matrix case. The general expectation is that they can always be expanded in multiple polylogarithms. However, for higher rank  $r > 2$  the number of divergent diagrams grows much faster. Thus one expects that the whole perturbative series of Green's functions is indeed more intricate than in matrix field theory.

## 5 Outlook

The main purpose of this contribution has been to demonstrate how the Connes-Kreimer Hopf algebra [29, 17, 18, 30] can be used for explicit calculations in cNLFT. Feynman diagrams in cNLFT are best described by the notion of 2-graph which generalizes usual Feynman graphs of point-like interactions by adding a second layer of strands along which the degrees of freedom in cNLFT propagate. The crucial difference to standard quantum field theory is that the external structure of such diagrams is not just given by the number of legs of a given field type but is captured by graphs. Nevertheless, with the right definitions, the Hopf-algebraic structure relying on the usual coproduct of subgraph contraction is exactly the same as for local quantum field theory [47]. Thus, it directly gives a convenient algorithm to renormalize Green's functions in cNLFT, such as non-commutative field theory and matrix field theory [23, 48, 28] as well as higher rank tensorial and group field theories [4, 2, 35, 20, 14].

While the Hopf algebra unveils the structure of renormalization independent of a specific scheme, here I have demonstrated how it works for BPHZ momentum subtraction [8, 27, 49] in two simple examples of tensorial field theory on  $(\mathbb{R}^d)^{\otimes r}$ . As a first example, in  $\phi_{2,2}^4$  theory, i.e. complex matrix field theory on a 2-dimensional configuration space, the tadpole and sunrise diagram have the same amplitude as in the Grosse-Wulkenhaar model [23, 48] and the Hopf-algebraic calculations reproduce the known results [28]. Second, as a first example of a field theory of higher rank  $r > 2$  I have calculated the amplitudes for the same diagrams in  $\phi_{1,3}^4$  theory. As to be expected, these calculations give first hints that also in tensorial field theories the algebra of amplitudes is spanned by multiple polylogarithms. A deeper understanding of this space of amplitudes for tensorial theories would be an interesting research topic in its own.

The renormalized amplitudes of the tensorial versions of the tadpole and sunrise diagrams give examples showing that the UV regime of tensorial field theories [2] is much richer than the large- $N$  limit of the corresponding tensor models [25, 26]. The latter is completely described by so-called melonic diagrams, i.e. diagrams which have vanishing Gurau degree  $\omega^G = 0$ . There have been considerable efforts to find random geometries beyond the melonic regime, for example by enhancement of subleading classes of diagrams [19, 33]. In tensorial field theories, melonic diagrams are the most divergent but not the only divergent ones. In fact, the tadpole and sunrise amplitudes Eq.(25) and Eq.(35) are examples of divergent, non-melonic amplitudes. Thus, in renormalized quantities they contribute as much as the melonic contributions which makes tensorial field theory a perfect candidate to find new universality classes beyond the melonic regime.

## Acknowledgments

I would like to thank A. Hock and R. Wulkenhaar for discussions on the forest formula in matrix field theory. I am deeply grateful to D. Kreimer for advice and support, in particular in

developing the research project this work is part of which is funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) under the project number 418838388. Further funds come from the DFG under Germany's Excellence Strategy EXC 2044–390685587, Mathematics Münster: Dynamics–Geometry–Structure.

## References

- [1] Aldous D., The Continuum Random Tree. I, *Ann. Probab.* **19** (1991), 1–28.
- [2] Ben Geloun J., Renormalizable Models in Rank  $d \geq 2$  Tensorial Group Field Theory, *Comm. Math. Phys.* **332** (2014), 117–188, arXiv:1306.1201.
- [3] Ben Geloun J., Ousmane Samary D., 3D Tensor Field Theory: Renormalization and One-Loop  $\beta$ -Functions, *Ann. Henri Poincaré* **14** (2013), 1599–1642, arXiv:1201.0176.
- [4] Ben Geloun J., Rivasseau V., A Renormalizable 4-Dimensional Tensor Field Theory, *Comm. Math. Phys.* **318** (2013), 69–109, arXiv:1111.4997.
- [5] Bergbauer C., Kreimer D., Hopf algebras in renormalization theory: Locality and Dyson-Schwinger equations from Hochschild cohomology, in *Physics and Number Theory*, Editor L. Nyssen, European Mathematical Society Publishing House, 2006, 133–164, arXiv:hep-th/0506190.
- [6] Blaschke D.N., Garschall T., Gieres F., Heindl F., Schweda M., Wohlgenannt M., On the Renormalization of Non-Commutative Field Theories, *Eur. Phys. J. C* **73** (2013), 2262, arXiv:1207.5494.
- [7] Blaschke D.N., Gieres F., Heindl F., Schweda M., Wohlgenannt M., BPHZ renormalization and its application to non-commutative field theory, *Eur. Phys. J. C* **73** (2013), 2566, arXiv:1307.4650.
- [8] Bogoliubow N.N., Parasiuk O.S., Über die Multiplikation der Kausalfunktionen in der Quantentheorie der Felder, *Acta Math.* **97** (1957), 227–266.
- [9] Bonzom V., Gurau R., Rivasseau V., Random tensor models in the large N limit: Uncoloring the colored tensor models, *Phys. Rev. D* **85** (2012), 084037, arXiv:1202.3637.
- [10] Borinsky M., *Graphs in Perturbation Theory: Algebraic Structure and Asymptotics*, Springer Theses Recognizing Outstanding Ph.D. Research, Springer International Publishing, 2018, arXiv:1807.02046.
- [11] Branahl J., Hock A., Wulkenhaar R., Blobbed topological recursion of the quartic Kontsevich model I: Loop equations and conjectures, *preprint* (2020),, arXiv:2008.12201.
- [12] Branahl J., Hock A., Wulkenhaar R., Perturbative and Geometric Analysis of the Quartic Kontsevich Model, *preprint* (2020),, arXiv:2012.02622.
- [13] Broadhurst D.J., Kreimer D., Exact solutions of Dyson-Schwinger equations for iterated one-loop integrals and propagator-coupling duality, *Nuclear Physics B* **600** (2001), 403–422, arXiv:hep-th/0012146.
- [14] Carrozza S., *Tensorial methods and renormalization in Group Field Theories*, Ph.D. thesis, Université Paris-Sud 11, Paris Orsay, 2013, arXiv:1310.3736.
- [15] Carrozza S., Oriti D., Rivasseau V., Renormalization of a SU(2) Tensorial Group Field Theory in Three Dimensions, *Comm. Math. Phys.* **330** (2014), 581–637, arXiv:1303.6772.
- [16] Carrozza S., Oriti D., Rivasseau V., Renormalization of Tensorial Group Field Theories: Abelian U(1) Models in Four Dimensions, *Comm. Math. Phys.* **327** (2014), 603–641, arXiv:1207.6734.
- [17] Connes A., Kreimer D., Renormalization in Quantum Field Theory and the Riemann-Hilbert Problem I: The Hopf Algebra Structure of Graphs and the Main Theorem, *Comm. Math. Phys.* **210** (2000), 249–273, arXiv:hep-th/9912092.
- [18] Connes A., Kreimer D., Renormalization in Quantum Field Theory and the Riemann-Hilbert Problem II: The  $\beta$ -Function, Diffeomorphisms and the Renormalization Group, *Comm. Math. Phys.* **216** (2001), 215–241, arXiv:hep-th/0003188.
- [19] Delepouve T., Gurau R., Phase transition in tensor models, *JHEP* **06** (2015), 178, arXiv:1504.05745.
- [20] Freidel L., Group Field Theory: An Overview, *Int. J. Theor. Phys.* **44** (2005), 1769, arXiv:hep-th/0505016.
- [21] Grosse H., Hock A., Wulkenhaar R., Solution of the self-dual  $\Phi^4$  QFT-model on four-dimensional Moyal space, *JHEP* **01** (2020), 081, arXiv:1908.04543.
- [22] Grosse H., Sako A., Wulkenhaar R., Exact solution of matricial  $\Phi_2^3$  quantum field theory, *Nuclear Physics* **925** (2017), 319–347, arXiv:1610.00526.

- [23] Grosse H., Wulkenhaar R., Renormalisation of  $\phi^4$ -Theory on Non-Commutative  $\mathbb{R}^4$  to All Orders, *Letters in Mathematical Physics* **71** (2005), 13–26, arXiv:hep-th/0403232.
- [24] Gurau R., Colored Group Field Theory, *Comm. Math. Phys.* **304** (2011), 69–93, arXiv:0907.2582.
- [25] Gurau R., The Complete  $1/N$  Expansion of Colored Tensor Models in Arbitrary Dimension, *Ann. Henri Poincaré* **13** (2012), 399–423, arXiv:1102.5759.
- [26] Gurau R., Random Tensors, Oxford University Press, 2016.
- [27] Hepp K., Proof of the Bogoliubov-Parasiuk theorem on renormalization, *Comm. Math. Phys.* **2** (1966), 301–326.
- [28] Hock A., Matrix Field Theory, Ph.D. thesis, WWU Münster, 2020, arXiv:2005.07525.
- [29] Kreimer D., On the Hopf algebra structure of perturbative quantum field theories, *Adv. Theor. Math. Phys.* **2** (1998), 303–334, arXiv:q-alg/9707029.
- [30] Kreimer D., Combinatorics of (perturbative) quantum field theory, *Phys. Rept.* **363** (2002), 387–424, arXiv:hep-th/0010059.
- [31] Kreimer D., Anatomy of a gauge theory, *Annals Phys.* **321** (2006), 2757–2781, arXiv:hep-th/0509135.
- [32] Kreimer D., Yeats K., An Étude in non-linear Dyson–Schwinger Equations, *Nucl Phys B-Proc Sup* **160** (2006), 116–121, arXiv:hep-th/0605096.
- [33] Lionni L., Marckert J.F., Iterated foldings of discrete spaces and their limits: candidates for the role of Brownian map in higher dimensions, *preprint* (2019),, arXiv:1908.02259.
- [34] Marckert J.F., Mokkadem A., Limit of normalized quadrangulations: The Brownian map, *Ann. Probab.* **34** (2006), 2144–2202, arXiv:math/0403398.
- [35] Oriti D., Spin Foam Models of Quantum Spacetime, Ph.D. thesis, Cambridge University, 2003, arXiv:gr-qc/0311066.
- [36] Oriti D., Group field theory as the microscopic description of the quantum spacetime fluid: A New perspective on the continuum in quantum gravity, in PoS QG-PH, 2007, 030, arXiv:0710.3276.
- [37] Oriti D., Ryan J.P., Thürigen J., Group field theories for all loop quantum gravity, *New J. Phys.* **17** (2015), 023042, arXiv:1409.3150.
- [38] Panzer E., Wulkenhaar R., Lambert-W solves the noncommutative  $\Phi^4$ -model, *Comm. Math. Phys.* **374** (2020), 1935–1961, arXiv:1807.02945.
- [39] Pascalie R., A Solvable Tensor Field Theory, *Letters in Mathematical Physics* **110** (2020), 925–943, arXiv:1903.02907.
- [40] Pérez A., The Spin-Foam Approach to Quantum Gravity, *Living Rev. Relativity* **16** (2013), 3, arXiv:1205.2019.
- [41] Pithis A.G.A., Thürigen J., Phase transitions in TGFT: Functional renormalization group in the cycli-melonic potential approximation and equivalence to  $O(N)$  models, *JHEP* **12** (2020), 159, arXiv:2009.13588.
- [42] Raasakka M., Tanasa A., Combinatorial Hopf algebra for the Ben Geloun-Rivasseau tensor field theory, *Sém. Lothar. Combin.* **70** (2014), B70d, arXiv:1306.1022.
- [43] Rivasseau V., Quantum gravity and renormalization: The tensor track, in 8th International Conference on Progress in theoretical physics (ICPTP 2011). AIP Conference Proceedings, Vol. 1444, 2012, 18–29, arXiv:1112.5104.
- [44] Tanasa A., Kreimer D., Combinatorial Dyson-Schwinger equations in noncommutative field theory, *Journal of Noncommutative Geometry* **7** (2013), 255–289, arXiv:0907.2182.
- [45] Tanasa A., Vignes-Tourneret F., Hopf algebra of non-commutative field theory, *Journal of Noncommutative Geometry* **2** (2008), 125–139, arXiv:0707.4143.
- [46] Thürigen J., Discrete quantum geometries and their effective dimension, Ph.D. thesis, Humboldt-Universität zu Berlin, 2015, arXiv:1510.08706.
- [47] Thürigen J., Renormalization in combinatorially non-local field theories: the Hopf algebra of 2-graphs, *preprint* (2021),, arXiv:2102.12453.
- [48] Wulkenhaar R., Quantum field theory on noncommutative spaces, in Advances in Noncommutative Geometry, Editors A.H. Chamseddine, C. Consani, N. Higson, M. Khalkhali, H. Moscovici, G. Yu, Springer International Publishing, 2019, 607–690.
- [49] Zimmermann W., Convergence of Bogoliubov’s method of renormalization in momentum space, *Comm. Math. Phys.* **15** (1969), 208–234.