

BUSEMANN PROCESS AND SEMI-INFINITE GEODESICS IN BROWNIAN LAST-PASSAGE PERCOLATION

TIMO SEPPÄLÄINEN AND EVAN SORENSEN

ABSTRACT. We prove the existence of semi-infinite geodesics for Brownian last-passage percolation (BLPP). Specifically, on a single event of probability one, there exist semi-infinite geodesics started from every space-time point and traveling in every asymptotic direction. Properties of these geodesics include uniqueness for a fixed initial point and direction, non-uniqueness for fixed direction but random initial points, and coalescence of all geodesics traveling in a common, fixed direction. Along the way, we prove that for fixed northeast and southwest directions, there almost surely exist no bi-infinite geodesics in the given directions. The semi-infinite geodesics are constructed from Busemann functions. Our starting point is a result of Alberts, Rassoul-Agha and Simper that established Busemann functions for fixed points and directions. Out of this, we construct the global process of Busemann functions simultaneously for all initial points and directions, and then the family of semi-infinite Busemann geodesics. The uncountable space of the semi-discrete setting requires extra consideration and leads to new phenomena, compared to discrete models.

CONTENTS

1. Introduction	2
1.1. Brownian last-passage percolation	2
1.2. BLPP in the Kardar-Parisi-Zhang universality class	2
1.3. Semi-infinite geodesics in discrete models	2
1.4. Contributions of the paper	3
1.5. Future work	3
1.6. Organization of the paper	4
1.7. Acknowledgements	4
2. Definitions and previous results	4
2.1. Preliminaries	4
2.2. Geodesics in Brownian last-passage percolation	4
2.3. Busemann functions	6
3. Main results	7
3.1. Existence and properties of the Busemann process	7
3.2. Heuristic for construction of semi-infinite geodesics	10
3.3. Semi-infinite geodesics	10
3.4. Connection to infinite-length polymer measures for the Brownian polymer	13
3.5. Connection to semi-infinite geodesics in discrete last-passage percolation	14
3.6. Connection to queuing theory	15
4. Construction of the Busemann process	16
4.1. Construction of Busemann functions for a fixed direction	16
4.2. Global construction of the Busemann process	18
5. Proofs of the results for semi-infinite geodesics	20
5.1. Key lemmas	21
5.2. Proofs of Theorems 3.8-3.12	23
5.3. The dual environment	27
5.4. Dual geodesics	28
5.5. Proof of Lemma 3.15 and Theorem 3.17	32

Date: Wednesday 1st September, 2021.

2020 Mathematics Subject Classification. 60K30,60K35,60K37.

Key words and phrases. Brownian motion, Busemann function, coalescence, geodesic, last-passage percolation, midpoint problem.

T. Seppäläinen was partially supported by National Science Foundation grant DMS-1854619 and by the Wisconsin Alumni Research Foundation.

E. Sorensen was partially supported by T. Seppäläinen, through National Science Foundation grants DMS-1602846 and DMS-1854619.

Appendix A. Technical lemmas and theorems	34
A.1. Deterministic facts about continuous functions	34
A.2. Probabilistic results	35
A.3. The Brownian queue and stationary last-passage process	37
A.4. Time reversal	40
References	41

1. INTRODUCTION

1.1. Brownian last-passage percolation. Brownian last-passage percolation (BLPP) dates back to the 1991 work of Glynn and Whitt [GW91], where the model appeared as a large-scale limit of multiple queues in series under heavy traffic conditions. Harrison and Williams [Har85, HW90, HW92] also studied what is known as the Brownian queue and developed a stability result for the model. More specifically, if the arrivals process is given by increments of Brownian motion and the service process is given by increments of an independent Brownian motion with drift, the departures process is also given by increments of Brownian motion. The connection between BLPP and the Brownian queue is expounded on by O’Connell and Yor [OY01], who also introduced the positive temperature version of this model, known as the Brownian polymer. We discuss the connection to queuing theory in Sections 3.6 and A.3.

1.2. BLPP in the Kardar-Parisi-Zhang universality class. In the early 2000s, Gravner, Tracy, and Widom [GTW01] and Baryshnikov [Bar01] made the remarkable discovery that the Brownian last-passage value $L_{(0,0),(n,1)}(\mathbf{B})$ (to be defined in (2.1) below) has the same distribution as the largest eigenvalue of an $n \times n$ GUE random matrix. Soon after, O’Connell and Yor [OY02] provided an alternate proof relying on the queuing interpretation of the model. Since then, Brownian last-passage percolation and the Brownian polymer have been widely studied as gateways to properties of the KPZ universality class. As a semi-discrete model with one continuous and one discrete parameter, BLPP serves as an intermediary between discrete models such as LPP on the planar integer lattice and continuum models such as the stochastic heat equation and the KPZ equation. In [CH14], Corwin and Hammond constructed the Airy line ensemble and proved that certain statistics of Brownian last-passage percolation converge in distribution to the Airy line ensemble. Dauvergne, Nica, and Virág ([DNV19], Corollary 6.4) provided an alternate method to prove this fact, which couples Brownian last-passage percolation with geometric random walks.

Recently, Dauvergne, Ortmann, and Virág [DOV18] constructed the directed landscape, a central object in the KPZ universality class, as the scaling limit of Brownian last-passage percolation. Another central object is the KPZ fixed point, constructed by Matetski, Quastel, and Remenik in [MQR16], as the limit of the totally asymmetric simple exclusion process. In [DOV18], a variational duality is described between the directed landscape and KPZ fixed point. Even more recently, convergence to the KPZ fixed point was shown for a larger class of models, including the height function of the KPZ equation. This was done independently by Quastel and Sarkar [QS20] and Virág [Vir20].

1.3. Semi-infinite geodesics in discrete models. The study of infinite geodesics in planar random growth models has gone through a number of stages over the last 40 years, beginning with the work of Licea and Newman [LN96, New95] on first-passage percolation with i.i.d. edge weights. Under a global curvature assumption on the limit shape, for continuously distributed edge weights, they proved existence of a deterministic full-Lebesgue measure set of directions in which there is a unique semi-infinite geodesic out of every lattice point. They also showed that, for each direction in this set, the semi-infinite geodesics in that direction all coalesce.

A separate strand of work consists of long-term efforts to prove the nonexistence of bi-infinite geodesics. In first-passage percolation, Licea and Newman [LN96] showed that there are no bi-infinite geodesics in fixed northeast and southwest directions. Howard and Newman [HN01] later proved similar results for Euclidean last-passage percolation. Around this time, Wehr and Woo [WW98] proved that, under a first moment assumption on the edge weights, there are no bi-infinite geodesics for first-passage percolation that lie entirely on the upper-half plane.

In 2016, Damron and Hanson [DH17] strengthened the result of Licea and Newman by proving that, if the weights have continuous distribution and the boundary of the limit shape is differentiable, for each fixed direction, there are no bi-infinite geodesics with one end having that direction. The conjectured nonexistence was finally resolved in exponential last-passage percolation, known also as the exponential corner growth model (CGM). The proofs came in two independent works: first by Basu, Hoffman, and Sly [BHS18] and shortly thereafter by Balázs, Busani, and the first author [BBS20]. The latter proof is in spirit aligned with

the development in the present paper, as it rested on understanding the joint distribution of the Busemann functions from [FS20].

Another focus of research has been the coalescence structure of geodesics in discrete last-passage percolation. Ferrari and Pimentel [FP05] imported the Licea–Newman approach [LN96, New95] to exponential last-passage percolation. Later Pimentel [Pim16] developed a probabilistic duality between the coalescence time of two semi-infinite geodesics and their last exit times from the initial boundary, in exponential LPP. A key idea was the equality in distribution of the tree of directed semi-infinite geodesics and the dual tree of southwest-directed geodesics. From this, he obtained a lower bound on tail probabilities of the coalescence time for two geodesics starting from $(-\lfloor k^{2/3} \rfloor, \lfloor k^{2/3} \rfloor)$ and $(\lfloor k^{2/3} \rfloor, -\lfloor k^{2/3} \rfloor)$. Coalescence bounds have seen significant recent improvement in [BSS19] and [SS20].

Hoffman [Hof08] was the first to use ideas of Busemann functions to study geodesics in first-passage percolation. In this paper, he showed the existence of disjoint semi-infinite geodesics. Later, Damron and Hanson [DH14] constructed generalized Busemann functions from weak subsequential limits of first-passage times. This allowed them to develop results for semi-infinite geodesics under weaker assumptions than the global curvature. Under the assumption that the limit shape is strictly convex and differentiable, they proved that every semi-infinite geodesic has an asymptotic direction and that, in every direction, there exists a semi-infinite geodesic out of every lattice point.

Georgiou, Rassoul-Agha, and the first author [GRAS17b, GRAS17a] used an approach based on the stationary LPP process to prove existence of Busemann functions in discrete last-passage percolation with general i.i.d. weights, under mild moment conditions. The Busemann functions were then used to construct semi-infinite geodesics. They also showed that if the shape function is strictly concave, every semi-infinite geodesic has an asymptotic direction. Further in this direction, in [Sep20] the first author introduced a new proof of the coalescence of semi-infinite geodesics that utilizes the stationary LPP process.

Our work is related to this last approach. We use Busemann functions to construct and study semi-infinite geodesics in BLPP. Existence of Busemann functions in BLPP, for fixed initial points and directions, was recently established by Alberts, Rassoul-Agha and Simper [ARAS20], along with Busemann functions and infinite polymer measures for the semi-discrete Brownian polymer. Their BLPP result is the starting point of our study.

1.4. Contributions of the paper. This paper develops the global setting of Busemann functions and semi-infinite geodesics in the semi-discrete BLPP, with a view to future study of their finer properties. The novelty lies in going beyond the familiar discrete set-up. Specific items proved below include the following:

- (1) The Busemann functions of BLPP from [ARAS20] are extended to a global process on a single event of probability one, for all the uncountably many initial points and directions.
- (2) Once the Busemann process is in place, we use it to construct semi-infinite geodesics for BLPP. With probability one, there exists a family of semi-infinite geodesics, starting from each initial point and in each direction.
- (3) With probability one, all semi-infinite geodesics, whether constructed by the Busemann functions or not, have an asymptotic direction. For a fixed initial point and direction, there is almost surely a unique semi-infinite geodesic starting from the given point and traveling asymptotically in the given direction. We also show that for a fixed direction, there is a countably infinite set of initial points whose geodesic in that direction is not unique.
- (4) For each fixed direction, we prove that all semi-infinite geodesics, traveling in that common direction coalesce.
- (5) For fixed northeast and southwest directions, we prove the almost sure nonexistence of bi-infinite geodesics in those directions.

Many of our results have direct analogues to those in discrete models, but resolving the issue of uncountably many initial points for geodesics requires additional care. For example, to construct an infinite up-right path in a discrete model, one iteratively chooses at each step whether to move upward or to the right. In the BLPP setting one chooses a real-valued increment to the right and then takes an upward step.

Section 3.2 informally demonstrates the idea behind this construction. The proofs of the results for semi-infinite geodesics then rely on understanding the structure of the locations of the upward steps. Furthermore, as seen in point (3) above, a uniqueness statement for semi-infinite geodesics starting from a fixed initial point does not translate to a uniqueness statement for all initial points simultaneously, unlike in discrete models. This leads to interesting questions related to the exceptional sets of initial points.

1.5. Future work. The present paper lays the foundation for future work on Busemann functions and semi-infinite geodesics. In the short term, we will use the tools developed in this paper in the following directions.

- (1) To study the full family of semi-infinite geodesics in BLPP, we derive the joint distribution of Busemann functions across all directions. The analogous result was achieved in [FS20] for the exponential CGM, which led to important advances in the global structure of geodesic trees [JRS19], coalescence [SS20], convergence [BBS21, BF20], and bi-infinite geodesics [BBS20].
- (2) We will define the competition interface for each initial point of BLPP. This has a straightforward definition in discrete models, but requires more care in the semi-discrete setting. We will use this to demonstrate non-uniqueness of semi-infinite geodesics in the BLPP setting, somewhat analogously to what was done in the discrete setting in [JRS19]. Specifically, we will show that, on each level m , outside of a random set of Hausdorff dimension $\frac{1}{2}$, all points (m, t) have a degenerate competition interface. For each of the points in this set of Hausdorff dimension $\frac{1}{2}$, there exists a random direction θ^* such that there are two semi-infinite geodesics starting at (m, t) in direction θ^* . These two geodesics only share the common initial point.

As can be seen from (2), there are many rich questions to explore related to non-uniqueness of semi-infinite geodesics for Brownian last-passage percolation. Other questions related to Hausdorff dimension have recently been explored for the KPZ fixed point and the directed landscape in [BGH21, BGH19, CHHM21]. Our longer-term goal is to extend the approach of this paper to the continuum models of the KPZ class.

1.6. Organization of the paper. Section 2 defines the Brownian last-passage percolation model and related objects. We also state results from other papers, including the existence of Busemann functions from [ARAS20]. Section 3 contains our main results on the existence and properties of the global Busemann process and semi-infinite geodesics. At the end of Section 3 we discuss connections to infinite polymer measures in the Brownian polymer, semi-infinite geodesics in discrete last-passage percolation, and to queuing theory. Section 4 constructs the global Busemann process and derives its properties. The results for semi-infinite geodesics are proved in Section 5. We prove coalescence through a family of dual southwest geodesics, developed in Section 5.4. The Appendix collects some standard material, background from the literature, and some technical theorems.

1.7. Acknowledgements. The authors thank Tom Alberts and Firas Rassoul-Agha for helpful discussions on [ARAS20] and Neil O’Connell for valuable discussions related to the paper [OY01].

2. DEFINITIONS AND PREVIOUS RESULTS

2.1. Preliminaries. The following notation and conventions are used throughout the paper.

- (i) For a function $f : \mathbb{R} \rightarrow \mathbb{R}$, we write $f(s, t) = f(t) - f(s)$ and $\tilde{f}(t) = -f(-t)$.
- (ii) \mathbb{Z} , \mathbb{Q} and \mathbb{R} are restricted by subscripts, as in for example $\mathbb{Z}_{>0} = \{1, 2, 3, \dots\}$.
- (iii) Whenever $m \leq n \in \mathbb{Z}$ and $s \leq t \in \mathbb{R}$, we say that $(m, s) \leq (n, t)$.
- (iv) Let $X \sim \mathcal{N}(\mu, \sigma^2)$ indicate that the random variable X has normal distribution with mean μ and variance σ^2 . For $\alpha > 0$, let $X \sim \text{Exp}(\alpha)$ indicate that X has exponential distribution with rate α , or equivalently, mean α^{-1} .
- (v) Equality in distribution between random variables and processes is denoted by $\stackrel{d}{=}$.
- (vi) A two-sided Brownian motion is a continuous random process $\{B(t) : t \in \mathbb{R}\}$ such that $B(0) = 0$ almost surely and such that $\{B(t) : t \geq 0\}$ and $\{B(-t) : t \geq 0\}$ are two independent standard Brownian motions on $[0, \infty)$.
- (vii) For $\lambda \in \mathbb{R}$, $\{Y(t) : t \in \mathbb{R}\}$ is a two-sided Brownian motion with drift λ if the process $\{Y(t) - \lambda t : t \in \mathbb{R}\}$ is a two-sided Brownian motion.
- (viii) The square \square as a superscript represents a sign $+$ or $-$.

2.2. Geodesics in Brownian last-passage percolation. The Brownian last-passage process is defined as follows. On a probability space $(\Omega, \mathcal{F}, \mathbb{P})$, let $\mathbf{B} = \{B_r\}_{r \in \mathbb{Z}}$ be a field of independent, two-sided Brownian motions. For $(m, s) \leq (n, t)$, define the set

$$\Pi_{(m,s),(n,t)} := \{\mathbf{s}_{m,n} = (s_{m-1}, s_m, \dots, s_n) \in \mathbb{R}^{n-m+2} : s = s_{m-1} \leq s_m \leq \dots \leq s_n = t\}.$$

Denote the energy of a sequence $\mathbf{s}_{m,n} \in \Pi_{(m,s),(n,t)}$ by

$$\mathcal{E}(\mathbf{s}_{m,n}) = \sum_{r=m}^n B_r(s_{r-1}, s_r).$$

Now, for $\mathbf{x} = (m, s) \leq (n, t) = \mathbf{y}$, define the Brownian last-passage time as

$$(2.1) \quad L_{\mathbf{x}, \mathbf{y}}(\mathbf{B}) = \sup\{\mathcal{E}(\mathbf{s}_{m,n}) : \mathbf{s}_{m,n} \in \Pi_{\mathbf{x}, \mathbf{y}}\}.$$

Whenever the specific field of Brownian motions used is either clear from or not important in the context, we write $L_{(m,s),(n,t)}$.




FIGURE 2.1. Maximizing Brownian increments from $(0, s)$ to $(4, t)$. Each discrete level has an associated Brownian motion.

The elements of $\Pi_{\mathbf{x}, \mathbf{y}}$ are in bijection with paths in \mathbb{R}^2 between \mathbf{x} and \mathbf{y} , that move to the right in real-valued increments, and move upwards in integer increments. For a given $\mathbf{s}_{m,n} \in \Pi_{(m,s),(n,t)}$, the path consists of the following points:

$$(2.2) \quad \bigcup_{r=m}^n \{(u, r) : u \in [s_{r-1}, s_r]\} \cup \bigcup_{r=m}^{n-1} \{(s_r, v) : v \in [r, r + 1]\}.$$

This set consists of horizontal and vertical line segments such that the vertical segments occur at the points s_r . Because of this bijection, we sometimes say $\Gamma \in \Pi_{\mathbf{x}, \mathbf{y}}$ for such an up-right path. For $(m, t) \in \mathbb{Z} \times \mathbb{R}$, we graphically represent the t -coordinate as the horizontal coordinate and the m -coordinate as the vertical coordinate in the plane. Since $\Pi_{\mathbf{x}, \mathbf{y}}$ is a compact set and Brownian motion is continuous, on a single event of probability one, for all $(m, t) = \mathbf{x} \leq \mathbf{y} = (n, t) \in \mathbb{Z} \times \mathbb{R}$, there exists a sequence $\tau_{m,n} \in \Pi_{\mathbf{x}, \mathbf{y}}$ such that $\mathcal{E}(\tau_{m,n}) = L_{\mathbf{x}, \mathbf{y}}$. The associated path is called a *geodesic* between the points.




FIGURE 2.2. Example of a BLPP geodesic from $(0, s)$ to $(4, t)$

To an infinite sequence, $s = s_{m-1} \leq s_m \leq s_{m+1} \leq \dots$ we similarly associate a semi-infinite path. It is possible that $s_r = \infty$ for some $r \geq m$, in which case the last segment of the path is the ray $[s_{r-1}, \infty) \times \{r\}$, where r is the first index with $s_r = \infty$. The infinite path has direction $\theta \in [0, \infty]$ if

$$\lim_{n \rightarrow \infty} \frac{s_n}{n} \quad \text{exists and equals } \theta.$$

We call an up-right semi-infinite path a *semi-infinite geodesic* if, for any two points $\mathbf{x} \leq \mathbf{y} \in \mathbb{Z} \times \mathbb{R}$ that lie along the path, the portion of the path between the two points is a geodesic between the two points. Similarly, a *bi-infinite geodesic* is a bi-infinite path that forms a geodesic between any two of its points. Two up-right, semi-infinite paths *coalesce* if there exists $\mathbf{z} \in \mathbb{Z} \times \mathbb{R}$ such that the two paths agree above and to the right of \mathbf{z} , as shown in Figure 2.3. Alternatively, if the paths are defined by sequences of jump times $s^1 = s^1_{m_1-1} \leq s^1_{m_1} \dots$ and $s^2 = s^2_{m_2-1} \leq s^2_{m_2} \leq \dots$, then the two paths coalesce if and only if there exists $N \in \mathbb{Z}$ such that $s^1_r = s^2_r$ for all $r \geq N$.

The following Theorem, due to Hammond [Ham19], establishes uniqueness of geodesics for a fixed initial and terminal point.

Lemma 2.1 ([Ham19], Theorem B.1). *Fix endpoints $\mathbf{x} \leq \mathbf{y} \in \mathbb{Z} \times \mathbb{R}$. Then, there is almost surely a unique path whose energy achieves $L_{\mathbf{x}, \mathbf{y}}(\mathbf{B})$.*

However, it is also true that for each fixed initial point $\mathbf{x} \in \mathbb{Z} \times \mathbb{R}$, with probability one, there exist points $\mathbf{y} \geq \mathbf{x}$, such that the geodesic between \mathbf{x} and \mathbf{y} is not unique. Hence, the following lemma, found in [DOV18] is important for our understanding. It is a deterministic statement which holds for last-passage percolation across any field of continuous functions.





FIGURE 2.3. Coalescence of semi-infinite paths

FIGURE 2.4. Two geodesics with a common terminal point $(n, n\theta)$.

Lemma 2.2 ([DOV18], Lemma 3.5). *Between any two points $(m, s) \leq (n, t) \in \mathbb{Z} \times \mathbb{R}$ there is a right-most and a left-most Brownian last-passage geodesic between the two points. That is, there exist $\mathbf{s}_{m,n}^L, \mathbf{s}_{m,n}^R \in \Pi_{(m,s),(n,t)}$, that are maximal for $\mathcal{E}(\mathbf{s}_{m,n})$, such that, for any other maximal sequence $\mathbf{s}_{m,n}$, and for $m \leq r \leq n$,*

$$s_r^L \leq s_r \leq s_r^R.$$

For two fields of Brownian motions, \mathbf{B} and $\overline{\mathbf{B}}$, and an initial point $(m, s) \in \mathbb{Z} \times \mathbb{R}$, define the point-to-line last passage time:

$$(2.3) \quad \overline{L}_{(m,s),n}^\lambda(\mathbf{B}, \overline{\mathbf{B}}) = \sup \left\{ \sum_{r=m}^n B_r(s_{k-1}, s_k) - \overline{B}_{n+1}(s_n) - \lambda s_n : \mathbf{s}_{m,n} \in \Pi_{(m,s),n} \right\},$$

where $\Pi_{(m,s),n}$ denotes the set of sequences $s = s_{m-1} \leq s_m \leq \dots \leq s_n$ (no constraint on s_n). This object will be essential for our understanding of semi-infinite geodesics. The following is also due to Hammond [Ham19]

Lemma 2.3 ([Ham19], Lemma B.2). *Let \mathbf{B} be a field of independent, two-sided Brownian motions and $\overline{\mathbf{B}}$ be any other field of Brownian motions such that, for each n , \overline{B}_{n+1} is independent of $\{B_m\}_{m \leq n}$. Fix an initial point (m, s) and let $n \geq m$. Then, with probability one, the quantity in (2.3) is finite, and there is a unique sequence $\mathbf{s}_{m,n} \in \Pi_{(m,s),n}$ that is maximal for (2.3).*

Remark 2.4. Lemma 2.4 is stated slightly differently in [Ham19]. In that paper, $-\overline{B}_{n+1}(\cdot) - \lambda \cdot$ is replaced by an arbitrary deterministic and measurable function $h : \mathbb{R} \rightarrow \mathbb{R} \cup \{-\infty\}$ satisfying $h(t) > -\infty$ for some $t > s$ and $\limsup_{t \rightarrow \infty} h(t)/t < 0$. The assumption that $\{\overline{B}_{n+1}\}_{n+1}$ is independent of $\{B_m\}_{m \leq n}$ allows us to condition on \overline{B}_{n+1} and obtain the desired result.

2.3. Busemann functions. In the present paper, semi-infinite geodesics are constructed from Busemann functions. Busemann functions are defined to be the asymptotic difference of last-passage times from two different starting points to a common terminal point that is traveling to ∞ in a given direction. The direction is indexed by a parameter $\theta > 0$. Existence of these Busemann functions was proven in [ARAS20], both for the positive temperature and zero temperature cases. We state their zero-temperature result:

Theorem 2.5 ([ARAS20], Theorem 4.2). *Fix $\theta > 0$ and $\mathbf{x}, \mathbf{y} \in \mathbb{Z} \times \mathbb{R}$. Then, there exists a random variable $\mathcal{B}^\theta(\mathbf{x}, \mathbf{y})$ and an event $\Omega_{\mathbf{x}, \mathbf{y}}^{(\theta)}$ of probability one, on which*

$$(2.4) \quad \mathcal{B}^\theta(\mathbf{x}, \mathbf{y}) = \lim_{n \rightarrow \infty} [L_{\mathbf{x}, (n, t_n)} - L_{\mathbf{y}, (n, t_n)}]$$

holds for any sequence $\{t_n\} \subseteq \mathbb{R}$ satisfying $t_n/n \rightarrow \theta$. Further, if

$$(2.5) \quad v_m^\theta(t) := \mathcal{B}^\theta((m-1, t), (m, t)), \text{ and}$$

$$(2.6) \quad h_m^\theta(t) := \mathcal{B}^\theta((m, 0), (m, t)),$$

then $v_m^\theta(t) \sim \text{Exp}(\frac{1}{\sqrt{\theta}})$ and $h_m^\theta(s, t) \sim \mathcal{N}(\frac{t-s}{\sqrt{\theta}}, |t-s|)$ for all $s, t \in \mathbb{R}$ and $m \in \mathbb{Z}$.

3. MAIN RESULTS

3.1. Existence and properties of the Busemann process. To construct semi-infinite geodesics starting from every point and in every asymptotic direction, we extend the individual Busemann functions of Theorem 2.5 to a global *Busemann process*. The following transformations are used to understand the structure of this process. For functions $Z, B : \mathbb{R} \rightarrow \mathbb{R}$ satisfying $Z(0) = B(0) = 0$ and $\limsup_{t \rightarrow \infty} (B(t) - Z(t)) = -\infty$, define

$$(3.1) \quad Q(Z, B)(t) = \sup_{t \leq s < \infty} \{B(t, s) - Z(t, s)\},$$

$$(3.2) \quad D(Z, B)(t) = Z(t) + Q(Z, B)(0) - Q(Z, B)(t),$$

$$(3.3) \quad R(Z, B)(t) = B(t) + Q(Z, B)(t) - Q(Z, B)(0).$$

Reverse-time analogues of these transformations are defined for continuous functions $Y, C : \mathbb{R} \rightarrow \mathbb{R}$ satisfying $Y(0) = C(0) = 0$ and $\limsup_{t \rightarrow -\infty} (Y(t) - C(t)) = -\infty$:

$$(3.4) \quad \overleftarrow{Q}(Y, C)(t) = \sup_{-\infty < s \leq t} \{C(s, t) - Y(s, t)\},$$

$$(3.5) \quad \overleftarrow{D}(Y, C)(t) = Y(t) + \overleftarrow{Q}(Y, C)(t) - \overleftarrow{Q}(Y, C)(0),$$

$$(3.6) \quad \overleftarrow{R}(Y, C)(t) = C(t) + \overleftarrow{Q}(Y, C)(0) - \overleftarrow{Q}(Y, C)(t).$$

These transformations originate from the Brownian queue, first studied by Glynn and Whitt [GW91], and further expounded on by Harrison and Williams [Har85, HW90, HW92] and O'Connell and Yor [OY01]. See Sections 3.6 and A.3 for more about the queuing interpretation. The following lemma is a straightforward exercise. We state it for completeness as we will refer to it later in the paper.

Lemma 3.1. *Let $B, Z : \mathbb{R} \rightarrow \mathbb{R}$ be continuous functions satisfying $\limsup_{t \rightarrow \infty} (B(t) - Z(t)) = -\infty$. Then, $Q(Z, B), D(Z, B)$, and $R(Z, B)$ are continuous. Similarly, if $Y, C : \mathbb{R} \rightarrow \mathbb{R}$ are continuous functions satisfying $\limsup_{t \rightarrow \infty} (Y(t) - C(t)) = -\infty$, then $\overleftarrow{Q}(Y, C), \overleftarrow{D}(Y, C)$, and $\overleftarrow{R}(Y, C)$ are continuous.*

The next theorem summarizes the existence and properties of the Busemann process. This process has discontinuities in the direction parameter θ . Instead of a single cadlag process, it is useful to retain both a left- and a right-continuous version indicated by $\theta-$ and $\theta+$ because this distinction captures spatial limits.

Theorem 3.2. *There exists a process,*

$$\{\mathcal{B}^{\theta \square}(\mathbf{x}, \mathbf{y}) : \theta > 0, \square \in \{+, -\}, \mathbf{x}, \mathbf{y} \in \mathbb{Z} \times \mathbb{R}\},$$

and for $\theta > 0$, there exist events $\Omega^{(\theta)} \subseteq \Omega_1 \subseteq \Omega$, each of probability one, such that the following hold. Here, $v_{m+1}^{\theta \square}, h_m^{\theta \square}$ are defined as in (2.5) and (2.6), placing \square in the appropriate superscripts.

(i) (*Additivity*) On Ω_1 , whenever $\mathbf{x}, \mathbf{y}, \mathbf{z} \in (\mathbb{Z} \times \mathbb{R}), \theta > 0$, and $\square \in \{+, -\}$,

$$\mathcal{B}^{\theta \square}(\mathbf{x}, \mathbf{y}) + \mathcal{B}^{\theta \square}(\mathbf{y}, \mathbf{z}) = \mathcal{B}^{\theta \square}(\mathbf{x}, \mathbf{z}).$$

(ii) (*Monotonicity*) On Ω_1 , whenever $0 < \gamma < \theta < \infty, m \in \mathbb{Z}$, and $s < t \in \mathbb{R}$,

$$0 \leq v_m^{\gamma-}(s) \leq v_m^{\gamma+}(s) \leq v_m^{\theta-}(s) \leq v_{m+1}^{\theta+}(s), \text{ and}$$

$$B_m(s, t) \leq h_m^{\theta+}(s, t) \leq h_m^{\theta-}(s, t) \leq h_m^{\gamma+}(s, t) \leq h_m^{\gamma-}(s, t).$$

(iii) (*Convergence*) On Ω_1 , for every $m \in \mathbb{Z}, \theta > 0$ and $\square \in \{+, -\}$,

- (a) As $\gamma \nearrow \theta$, $h_m^{\gamma\Box}$ and $v_m^{\gamma\Box}$ converge uniformly, on compact subsets of \mathbb{R} , to $h_m^{\theta-}$ and $v_m^{\theta-}$, respectively.
- (b) As $\delta \searrow \theta$, $h_m^{\delta\Box}$ and $v_m^{\delta\Box}$ converge uniformly, on compact subsets of \mathbb{R} , to $h_m^{\theta+}$ and $v_m^{\theta+}$, respectively.
- (c) As $\gamma \rightarrow \infty$, $h_m^{\gamma\Box}$ converges uniformly, on compact subsets of \mathbb{R} , to B_m .
- (d) As $\delta \searrow 0$, $v_m^{\delta\Box}$ converges uniformly, on compact subsets of \mathbb{R} , to 0.
- (iv) (Continuity) On Ω_1 , for any $r, m \in \mathbb{Z}, \theta > 0$, and $\Box \in \{+, -\}$, $(s, t) \mapsto \mathcal{B}^{\theta\Box}((m, s), (r, t))$ is a continuous function $\mathbb{R}^2 \rightarrow \mathbb{R}$.
- (v) (Limits) On Ω_1 , for each $\theta > 0$ and $\Box \in \{+, -\}$,

$$\lim_{s \rightarrow \pm\infty} B_m(s) - h_{m+1}^{\theta\Box}(s) = \mp\infty.$$

- (vi) (Queuing relationships between Busemann functions) On Ω_1 , for all $m \in \mathbb{Z}, \theta > 0$, and signs $\Box \in \{+, -\}$,

$$v_{m+1}^{\theta\Box} = Q(h_{m+1}^{\theta\Box}, B_m) \quad \text{and} \quad h_m^{\theta\Box} = D(h_{m+1}^{\theta\Box}, B_m).$$

- (vii) (Independence) For any $m \in \mathbb{Z}$,

$$\{h_r^{\theta\Box} : \theta > 0, \Box \in \{+, -\}, r > m\} \text{ is independent of } \{B_r : r \leq m\}.$$

- (viii) (Equality for fixed directions) For each $\theta > 0$, on the event $\Omega^{(\theta)}$, for all $\mathbf{x}, \mathbf{y} \in \mathbb{Z} \times \mathbb{R}$ and all sequences $\{t_n\}$ with $t_n/n \rightarrow \theta$,

$$\mathcal{B}^{\theta-}(\mathbf{x}, \mathbf{y}) = \lim_{n \rightarrow \infty} [L_{\mathbf{x}, (n, t_n)} - L_{\mathbf{y}, (n, t_n)}] = \mathcal{B}^{\theta+}(\mathbf{x}, \mathbf{y}).$$

Remark 3.3. On account of part (viii), when working on the event $\Omega^{(\theta)}$ we write $\mathcal{B}^\theta = \mathcal{B}^{\theta-} = \mathcal{B}^{\theta+}$. The fact that the limits exist for all initial points $\mathbf{x}, \mathbf{y} \in \mathbb{Z} \times \mathbb{R}$ and fixed $\theta > 0$ on a single event of probability one gives a generalization of Theorem 2.5. As will be seen in Section 4, the key is Lemma A.16, which generalizes a proof in [ARAS20].

The following theorem describes the distribution of the Busemann function for a fixed θ . We introduce a dual field of Brownian motions used later in the proof of coalescence of semi-infinite geodesics. Recall the definition (3.3) of the mapping R , and set

$$(3.7) \quad X_m^\theta = R(h_m^\theta, B_{m-1}).$$

Denote the field of these random functions by $\mathbf{X}^\theta := \{X_m^\theta\}_{m \in \mathbb{Z}}$.

Theorem 3.4. *Fix $\theta > 0$.*

- (i) *The process $t \mapsto h_m^\theta(t)$ is a two-sided Brownian motion with drift $\frac{1}{\sqrt{\theta}}$.*
- (ii) *\mathbf{X}^θ is a field of independent two-sided Brownian motions. For each $n \in \mathbb{Z}$, $\{X_m^\theta\}_{m > n}$ is independent of $\{h_m^\theta\}_{m \leq n}$.*
- (iii) *The process $t \mapsto v_m^\theta(t)$ is a stationary and reversible strong Markov process. For each $t \in \mathbb{R}$, $v_m^\theta(t) \sim \text{Exp}(\frac{1}{\sqrt{\theta}})$.*
- (iv) *(Burke property) Fix $n < m \in \mathbb{Z}$ and $-\infty < t_n \leq t_{n-1} \leq \dots \leq t_{m+1} < \infty$. Then, the following random variables and processes are mutually independent:*

$$\{B_m(u, t_{m+1}) : u \leq t_{m+1}\}, \{h_m^\theta(t_{m+1}, u) : u \geq t_{m+1}\},$$

$v_r^\theta(t_r), \{X_r^\theta(t_r, u) : u \geq t_r\}, \{B_{r-1}(u, t_r) : u \leq t_r\}$, and $\{h_r^\theta(t_{r+1}, u) : t_{r+1} \leq u \leq t_r\}$ for $m+1 \leq r \leq n-1$,

$$\text{and } v_n^\theta(t_n), \{X_n^\theta(t_n, u) : u \geq t_n\} \{h_n^\theta(u, t_n) : u \leq t_n\}.$$

As a special case, $\{v_m^\theta(t)\}_{m \in \mathbb{Z}}$ is an i.i.d sequence for each $t \in \mathbb{R}$. Refer to Figure 3.1 for clarity.

Remark 3.5. Using the representation of Equation (3.10) below, many formulas for the process $t \mapsto v_m^\theta(t)$ are well known. See [BS02], page 321 and [SN01], specifically equations (4) and (5), for more on this process, including the transition density and infinitesimal generator. However, we caution that when comparing formulas, in the setting of [BS02, SN01], the process is

$$t \mapsto \sup_{-\infty < s \leq t} \{B(s, t) - \mu(t - s)\},$$

where B is a two-sided Brownian motion and $\mu > 0$. In our setting, there is a factor of $\sqrt{2}$ multiplied to B .




FIGURE 3.1. Independence structure for Busemann functions

The construction of the Busemann process and the proof of Theorem 3.2 can be found in Section 4. We prove Theorem 3.4 here, assuming Theorem 3.2 and with the help of the results of the appendix.

Proof of Theorem 3.4. Part (i): As will be seen from the construction in Section 4, $t \mapsto h_m^\theta(t)$ has the proper finite-dimensional distributions and is continuous by Theorem 3.2(iv).

Part (ii): Fix integers $p < n < m$. By Theorem 3.2(vii),

$$(B_p, \dots, B_n, \dots, B_{m-2}, B_{m-1}, h_m^\theta) \text{ are independent.}$$

Theorem 3.2(vi) and definition (3.7) give $h_{m-1}^\theta = D(h_m^\theta, B_{m-1})$ and $X_m^\theta = R(h_m^\theta, B_{m-1})$, so by Theorem A.13, h_{m-1}^θ and X_m^θ are independent and X_m^θ is a two-sided Brownian motion. In particular, now

$$(B_p, \dots, B_n, \dots, B_{m-2}, h_{m-1}^\theta, X_m^\theta) \text{ are independent.}$$

Continue inductively by applying the transformation (D, R) to successive pairs (B_{j-1}, h_j^θ) for $j = m-1, m-2, \dots, n+1$ after which

$$(B_p, \dots, B_{n-1}, h_n^\theta, X_{n+1}^\theta, \dots, X_m^\theta) \text{ are independent.}$$

To conclude, note that $(h_p^\theta, \dots, h_n^\theta)$ is a function of $(B_p, \dots, B_{n-1}, h_n^\theta)$ through iteration of $h_k^\theta = D(h_{k+1}^\theta, B_k)$.

Part (iii): By Theorem 3.2(vi) and (3.7), on the event $\Omega^{(\theta)}$ we have these relations $\forall m \in \mathbb{Z}$:

$$(3.8) \quad h_{m-1}^\theta = D(h_m^\theta, B_{m-1}), \quad X_m^\theta = R(h_m^\theta, B_{m-1}), \quad \text{and} \quad v_m^\theta = Q(h_m^\theta, B_{m-1}).$$

The fact that $v_m^\theta(t)$ is exponential with rate $\frac{1}{\sqrt{\theta}}$ then follows from Lemma A.6. Theorem A.17 allows us to reverse these mappings, so $\forall m \in \mathbb{Z}$:


$$(3.9) \quad h_m^\theta = \overleftarrow{D}(h_{m-1}^\theta, X_m^\theta), \quad B_{m-1} = \overleftarrow{R}(h_{m-1}^\theta, X_m^\theta), \quad \text{and} \quad v_m^\theta = \overleftarrow{Q}(h_{m-1}^\theta, X_m^\theta).$$

Then, for $t \in \mathbb{R}$,

$$(3.10) \quad v_m^\theta(t) = \sup_{-\infty < u \leq t} \{X_m^\theta(u, t) - h_{m-1}^\theta(u, t)\}.$$

By Parts (i) and (ii), $t \mapsto X_m^\theta(t) - h_{m-1}^\theta(t)$ is equal in distribution to a two-sided Brownian motion with negative drift, multiplied by a factor of $\sqrt{2}$. Represented this way, $t \mapsto v_m^\theta(t)$ is known as a reflected Brownian motion with drift. Stationarity follows from the stationarity of increments. The fact that X is a reversible strong Markov process is proven in [Har85], pg. 81 (see also pg. 49-50 in [Har85] and Equations (4) and (5) in [SN01]).

Part (iv): By Part (ii) and (3.9), for any initial level m , the process $\{h_r^\theta, v_{r+1}^\theta, X_{r+1}^\theta, B_r\}_{r \geq m}$ has the same distribution as $\{Y_r^{\frac{1}{\sqrt{\theta}}}, q_{r+1}^{\frac{1}{\sqrt{\theta}}}, B_{r+1}, W_r^\lambda\}_{r \geq 0}$ as defined in (A.2). Note that B_r now plays the role of the W_r^λ , as stated in the definition. Therefore, the independence structure of Theorem A.14 holds. \square

FIGURE 3.2. Example of an element of $\mathbf{T}_{(m,t)}^\theta$

3.2. Heuristic for construction of semi-infinite geodesics. The next task is the construction of semi-infinite geodesics from each initial point and in each asymptotic direction. For each given point $(m, t) \in \mathbb{Z} \times \mathbb{R}$ and direction parameter $\theta > 0$, we want to find a semi-infinite geodesic, defined by jump times $t = \tau_{m-1} \leq \tau_m \leq \dots$ that satisfy

$$\lim_{n \rightarrow \infty} \frac{\tau_n}{n} = \theta.$$

We argue heuristically to motivate the useful construction. Start by finding a maximal path for $L_{(m,t),(n,n\theta)}$ for a large value of n . Note that

$$L_{(m,t),(n,n\theta)} = \max_{s \in [t, n\theta]} (B_m(t, s) + L_{(m+1,s),(n,n\theta)}),$$

and the maximizer $s = \tau_m$ gives the location of the vertical segment of the geodesic from m to $m+1$. For all $t \leq s \leq n\theta$,

$$B_m(t, s) + L_{(m+1,s),(n,n\theta)} \leq B_m(t, \tau_m) + L_{(m+1,\tau_m),(n,n\theta)}.$$

Rearranging yields

$$B_m(\tau_m, s) \leq L_{(m+1,\tau_m),(n,n\theta)} - L_{(m+1,s),(n,n\theta)}.$$

As n changes, so could τ_m , but for the sake of heuristic we hold τ_m constant. Take limits as $n \rightarrow \infty$ and rearrange again to get, for some sign $\square \in \{+, -\}$,

$$B_m(s) - h_{m+1}^{\theta\square}(s) \leq B_m(\tau_m) - h_{m+1}^{\theta\square}(\tau_m).$$

3.3. Semi-infinite geodesics. The discussion of the previous section motivates this rigorous definition.

Definition 3.6. On the event Ω_1 , for all $(m, t) \in \mathbb{Z}$, $\theta > 0$ and $\square \in \{+, -\}$, let $\mathbf{T}_{(m,t)}^{\theta\square}$ denote the set of sequences

$$t = \tau_{m-1} \leq \tau_m \leq \tau_{m+1} \leq \dots$$

that satisfy

$$(3.11) \quad B_r(\tau_r) - h_{r+1}^{\theta\square}(\tau_r) = \sup_{s \in [\tau_{r-1}, \infty)} \{B_r(s) - h_{r+1}^{\theta\square}(s)\} \quad \text{for each } r \geq m.$$

Theorem 3.2, Parts (iv) and (v) imply that such sequences exist. At each level r there exist left-most and right-most maximizers. Let

$$t = \tau_{(m,t),m-1}^{\theta\square,L} \leq \tau_{(m,t),m}^{\theta\square,L} \leq \tau_{(m,t),m+1}^{\theta\square,L} \leq \dots \quad \text{and} \quad t = \tau_{(m,t),m-1}^{\theta\square,R} \leq \tau_{(m,t),m}^{\theta\square,R} \leq \tau_{(m,t),m+1}^{\theta\square,R} \leq \dots$$

denote the left-most and right-most sequences in $\mathbf{T}_{(m,t)}^{\theta\square}$. Furthermore, define

$$\mathbf{T}_{(m,t)}^\theta := \mathbf{T}_{(m,t)}^{\theta+} \cup \mathbf{T}_{(m,t)}^{\theta-}.$$

Remark 3.7. Since every non-decreasing sequence in $\mathbf{T}_{(m,t)}^\theta$ defines a semi-infinite up-right path, $\mathbf{T}_{(m,t)}^\theta$ will be used to denote the set of up-right paths constructed in this way. By Theorem 3.2(viii), on the event $\Omega^{(\theta)} \subseteq \Omega_1$, $\mathbf{T}_\mathbf{x}^{\theta-} = \mathbf{T}_\mathbf{x}^{\theta+} = \mathbf{T}_\mathbf{x}^\theta$ for all $\mathbf{x} \in \mathbb{Z} \times \mathbb{R}$. However, this does not mean that $\mathbf{T}_\mathbf{x}^\theta$ contains only one element, so the left-most and right-most distinction is still necessary in general for a fixed θ . See Theorem 3.12 below. However, as will be seen in Theorem 3.10, for fixed $(m, t) \in \mathbb{Z} \times \mathbb{R}$, $\mathbf{T}_{(m,t)}^\theta$ almost surely contains one element. In this case, theorem A.8 gives the distribution of the first jump time τ_m .

Two elements $\{\tau_r\}_{r \geq m-1}$ and $\{\tau'_r\}_{r \geq m-1}$ of $\mathbf{T}_\mathbf{x}^{\theta\square}$ are distinct if $\tau_r \neq \tau'_r$ for at least one index r . Uniqueness of a semi-infinite geodesic in $\mathbf{T}_\mathbf{x}^{\theta\square}$ means that this set contains exactly one sequence.

The following theorems collect the properties of the semi-infinite geodesics, to be proved in Section 5.

Theorem 3.8. *Let Ω_1 be the event given in Theorem 3.2.*

- (i) (*Existence*) On Ω_1 , for all $\mathbf{x} \in \mathbb{Z} \times \mathbb{R}$, $\theta > 0$, and $\square \in \{+, -\}$, every element of $\mathbf{T}_{\mathbf{x}}^{\theta, \square}$ defines a semi-infinite geodesic starting from \mathbf{x} . More specifically, for any two points $\mathbf{y} \leq \mathbf{z}$ along a path in $\mathbf{T}_{\mathbf{x}}^{\theta}$, the energy of this path between \mathbf{y} and \mathbf{z} is $\mathcal{B}^{\theta, \square}(\mathbf{y}, \mathbf{z})$, and this energy is maximal over all paths between \mathbf{y} and \mathbf{z} .
- (ii) (*Left-most and right-most finite geodesics along paths*) If, for some $\theta > 0$, the points $\mathbf{y} \leq \mathbf{z} \in \mathbb{Z} \times \mathbb{R}$ both lie on the left-most semi-infinite geodesic in $\mathbf{T}_{\mathbf{x}}^{\theta, \square}$, then the portion of this geodesic between \mathbf{y} and \mathbf{z} coincides with the left-most finite geodesic between these two points. Similarly, the right-most semi-infinite geodesic is the right-most geodesic between any two of its points.
- (iii) (*Monotonicity*) On Ω_1 , for all $0 < \gamma < \theta$, all $(m, t) \in \mathbb{Z} \times \mathbb{R}$, and $r \geq m$,

$$\tau_{(m,t),r}^{\gamma-,L} \leq \tau_{(m,t),r}^{\gamma+,L} \leq \tau_{(m,t),r}^{\theta-,L} \leq \tau_{(m,t),r}^{\theta+,L} \quad \text{and} \quad \tau_{(m,t),r}^{\gamma-,R} \leq \tau_{(m,t),r}^{\gamma+,R} \leq \tau_{(m,t),r}^{\theta-,R} \leq \tau_{(m,t),r}^{\theta+,R}.$$

For all $\theta > 0$, $m \leq r \in \mathbb{Z}$, $s < t \in \mathbb{R}$, and $\square \in \{+, -\}$,

$$\tau_{(m,s),r}^{\theta, L} \leq \tau_{(m,t),r}^{\theta, L} \quad \text{and} \quad \tau_{(m,s),r}^{\theta, R} \leq \tau_{(m,t),r}^{\theta, R}.$$

- (iv) (*Convergence*) On Ω_1 , for all $(m, t) \in \mathbb{Z} \times \mathbb{R}$, $r \geq m$, $\theta > 0$, and $\square \in \{+, -\}$,

$$\lim_{\gamma \nearrow \theta} \tau_{(m,t),r}^{\gamma, L} = \tau_{(m,t),r}^{\theta, L} \quad \text{and} \quad \lim_{\delta \searrow \theta} \tau_{(m,t),r}^{\delta, R} = \tau_{(m,t),r}^{\theta, R}.$$

Additionally,

$$\lim_{s \nearrow t} \tau_{(m,s),r}^{\theta, L} = \tau_{(m,t),r}^{\theta, L} \quad \text{and} \quad \lim_{u \searrow t} \tau_{(m,u),r}^{\theta, R} = \tau_{(m,t),r}^{\theta, R}.$$

- (v) (*Directedness*) There exists a full-probability event $\Omega_2 \subset \Omega_1$ on which, for all $\mathbf{x} \in \mathbb{Z} \times \mathbb{R}$, $\theta > 0$, $\square \in \{+, -\}$, and all $\{\tau_r\}_{r \geq m} \in \mathbf{T}_{\mathbf{x}}^{\theta, \square}$,

$$\lim_{n \rightarrow \infty} \frac{\tau_n}{n} = \theta.$$

Remark 3.9 (A look ahead). In a subsequent paper, we will derive the joint distribution of the Busemann functions for varying θ and use this to strengthen the first line of Part (iv). Specifically, we will show that, there exists an event of full probability on which, for all $\theta > 0$, $m \leq r \in \mathbb{Z}$, $s < t \in \mathbb{R}$, $S \in \{L, R\}$ and $\square \in \{+, -\}$, there exists $\varepsilon > 0$ such that

$$\tau_{(m,s),r}^{\gamma, S} = \tau_{(m,s),r}^{\theta-, S} \quad \text{for all } \theta - \varepsilon < \gamma < \theta \quad \text{and} \quad \tau_{(m,s),r}^{\delta, S} = \tau_{(m,s),r}^{\theta+, S} \quad \text{for all } \theta < \delta < \theta + \varepsilon.$$

This will be used to strengthen the second line of Part (iii) in that, on an event of probability one, for each $m \in \mathbb{Z}$, $m \leq r \in \mathbb{Z}$, $s < t \in \mathbb{R}$, $\theta > 0$, and $\square \in \{+, -\}$,

$$\tau_{(m,s),r}^{\theta, R} \leq \tau_{(m,t),r}^{\theta, L}.$$

See Corollary 3.14 below for a statement of this form for fixed θ . We will also use the joint distribution of the Busemann functions to show that the first line of Part (iii) cannot be strengthened to compare $\tau_{(m,t),r}^{\gamma, R}$ and $\tau_{(m,t),r}^{\theta, L}$ for general $\gamma < \theta$. Specifically, we will derive examples such that $\gamma < \theta$ and

$$\tau_{(m,t),r}^{\gamma-, R} = \tau_{(m,t),r}^{\gamma+, R} > \tau_{(m,t),r}^{\theta-, L} = \tau_{(m,t),r}^{\theta+, L}.$$

The following theorems give control over the collection of all semi-infinite geodesics, whether constructed from the Busemann functions or not.

Theorem 3.10. *Let Ω_2 be the event of Theorem 3.8(v).*

- (i) (*Control of finite geodesics*) On Ω_2 , let $\theta > 0$, $(m, t) \in \mathbb{Z} \times \mathbb{R}$, and let $\{t_n\}$ be any sequence that has direction θ . For all n sufficiently large so that $n \geq m$ and $t_n \geq t$, let $t = t_{n, m-1} \leq t_{n, m} \leq \dots \leq t_{n, n} = t_n$ be any sequence that defines a (finite) geodesic between (m, t) and (n, t_n) . Then, for each $r \geq m$,

$$\tau_{(m,s),r}^{\theta-, L} \leq \liminf_{n \rightarrow \infty} t_{n,r} \leq \limsup_{n \rightarrow \infty} t_{n,r} \leq \tau_{(m,s),r}^{\theta+, R}.$$

- (ii) (*Control of semi-infinite geodesics*) On Ω_2 , if, for some $\theta > 0$ and $(m, t) \in \mathbb{Z} \times \mathbb{R}$, any other geodesic (constructed from the Busemann functions or not) is defined by the sequence $t = t_{m-1} \leq t_m \leq \dots$, starts at (m, t) , and has direction θ , then for all $r \geq m$,

$$\tau_{(m,t),r}^{\theta-, L} \leq t_r \leq \tau_{(m,t),r}^{\theta+, R}.$$

- (iii) (*Uniqueness for fixed points and directions*) For each fixed $\mathbf{x} \in \mathbb{Z} \times \mathbb{R}$ and $\theta > 0$, there exists an event, $\Omega_{\mathbf{x}}^{(\theta)} \subseteq \Omega^{(\theta)} \cap \Omega_2$, of probability one, on which there is exactly one semi-infinite geodesic starting from \mathbf{x} , in direction θ .

- (iv) (*Convergence of finite geodesics*) On $\Omega_{\mathbf{x}}^{(\theta)}$, let $\{\tau_r\}_{r \geq m-1}$ be the unique sequence in $\mathbf{T}_{\mathbf{x}}^\theta$. Then, if $\{t_n\}$ is a θ -directed sequence and, for each n , the sequence $t = t_{n,m-1} \leq t_{n,m} \leq \dots \leq t_{n,n}$ defines a finite geodesic between (m, t) and (n, t_n) ,

$$\lim_{n \rightarrow \infty} t_{n,r} = \tau_r \quad \text{for all } r \geq m.$$

Theorem 3.11. *The following hold on the event Ω_2 , of full probability, defined in Theorem 3.8(v).*

- (i) *For any infinite sequence $t = t_{m-1} \leq t_m \leq t_{m+1} \leq \dots$ defining a semi-infinite geodesic starting from some point $(m, t) \in \mathbb{Z} \times \mathbb{R}$, the limit*

$$\lim_{n \rightarrow \infty} \frac{t_n}{n} \text{ exists in } [0, \infty].$$

- (ii) *If for any $(m, t) \in \mathbb{Z} \times \mathbb{R}$ and any such sequence, the above limit equals 0, then $t_r = t$ for all $r \geq m$. Similarly, if this limit is ∞ , then $t_m = \infty$. That is, the only semi-infinite geodesics that are asymptotically vertical or horizontal are trivial (i.e. straight lines).*
- (iii) *For each $(m, t) \in \mathbb{Z} \times \mathbb{R}$, $r \geq m$, $S \in \{L, R\}$, and $\square \in \{+, -\}$, we have the monotone limits*

$$\lim_{\theta \searrow 0} \tau_{(m,t),r}^{\theta, S} = t, \quad \text{and} \quad \lim_{\theta \rightarrow \infty} \tau_{(m,t),r}^{\theta, S} = \infty.$$

The following theorem tells us about the set of initial points with a non-unique θ -directed semi-infinite geodesic, for a fixed direction $\theta > 0$.

Theorem 3.12. *For each $\theta > 0$, there exists an event $\tilde{\Omega}^{(\theta)} \subseteq \Omega^{(\theta)} \cap \Omega_2$, of probability one, on which the following hold*

- (i) *The set*

$$\text{NU}^\theta := \{\mathbf{x} \in \mathbb{Z} \times \mathbb{R} : \text{the } \theta\text{-directed semi-infinite geodesic starting from } \mathbf{x} \text{ is not unique}\}$$

is countably infinite. For every $(m, t) \in \text{NU}^\theta$, at most one of the θ -directed semi-infinite geodesics passes through the point $(m, t + \varepsilon)$ for some $\varepsilon > 0$. All the others pass through $(m + 1, t)$.

- (ii) *The set NU^θ can be decomposed as the union*

$$\bigcup_{m \in \mathbb{Z}, r \geq m} \text{NU}_{m,r}^\theta,$$

where

$$\text{NU}_{m,r}^\theta = \{(m, t) : t \in \mathbb{R} \text{ and } t = \tau_{(m,t),k}^{\theta, L} = \tau_{(m,t),k}^{\theta, R} \text{ for } m \leq k \leq r - 1 \text{ and } t = \tau_{(m,t),r}^{\theta, L} < \tau_{(m,t),r}^{\theta, R}\}.$$

That is, for each level $m \in \mathbb{Z}$, $\text{NU}_{m,r}^\theta$ is the set of initial points (m, t) such that, for all $m \leq k \leq r - 1$, all geodesics starting from (m, t) travel vertically to (r, t) , then the right-most geodesic moves to the right while the left-most geodesic moves vertically to $(r + 1, t)$.

- (iii) *The set*

$$\text{NU}_1^\theta := \bigcup_{m \in \mathbb{Z}} \text{NU}_{m,1}^\theta$$

is the set of points $(m, t) \in \mathbb{Z} \times \mathbb{R}$ such that the left-most θ -directed semi-infinite geodesic passes through $(m + 1, t)$ and the right-most such geodesic passes through $(m, t + \varepsilon)$ for some $\varepsilon > 0$. This set is neither discrete nor dense in $\mathbb{Z} \times \mathbb{R}$. More specifically, for each point $(m, t) \in \text{NU}_1^\theta$ and every $\varepsilon > 0$, there exists $s \in (t - \varepsilon, t)$ such that $(m, s) \in \text{NU}_1^\theta$. For each $(m, t) \in \text{NU}_1^\theta$, there exists $\delta > 0$ such that, for all $s \in (t, t + \delta)$, $(m, s) \notin \text{NU}^\theta$.

Remark 3.13. In general, there are two types of non-uniqueness of semi-infinite geodesics. The first is stated in Theorem 3.12(i) for a fixed direction θ . This motivates the necessity of the L/R distinction. In this case, while for some points \mathbf{x} , there are θ -directed geodesics that disagree near the initial point, Theorem 3.17 below implies that the geodesics will eventually come back together. To see an example of this non-uniqueness discussed in Theorem 3.12(i), fix $\theta > 0$ and $m \in \mathbb{Z}$, and let $s^* \geq 0$ be such that

$$X(s^*) = \sup_{0 \leq s < \infty} \{B_m(s) - h_{m+1}^\theta(s)\}.$$

By Theorem 3.2(vii) and Lemma 2.3, with probability one, the maximizer s^* is unique. By Theorem A.8, $s^* > 0$ with probability one. By Theorem 3.2(v),

$$t^* := \sup\{t < 0 : B_m(t) - h_{m+1}^\theta(t) = X(s^*)\} \quad \text{exists in } \mathbb{R}.$$

Then, both t^* and s^* are maximizers for $B_m(s) - h_{m+1}^\theta(s)$ on $[t^*, \infty)$, giving at least two sequences in the set $\mathbf{T}_{(m,t^*)}^\theta$.

The second type of non-uniqueness of semi-infinite geodesics requires investigation of the collection of all semi-infinite geodesics in all directions. This will motivate the $\theta \pm$ distinction. In future work, we will show that there exists a random, Hausdorff dimension $\frac{1}{2}$ set of points $\mathbf{x} \in \mathbb{Z} \times \mathbb{R}$, and to each of these points an associated direction θ^* , such that the right-most (resp. left-most) geodesic in $\mathbf{T}_{\mathbf{x}}^{\theta^*-}$ and the right-most (resp. left-most) geodesic in $\mathbf{T}_{\mathbf{x}}^{\theta^*+}$ only share the initial point \mathbf{x} .

Corollary 3.14. *On the θ -dependent full-probability event $\tilde{\Omega}^{(\theta)}$ of Theorem 3.12, the second line of Theorem 3.8(iii) can be strengthened in the following way: for all pairs of initial points (m, s) and (m, t) in $\mathbb{Z} \times \mathbb{R}$ that satisfy $s < t$, we have*

$$\tau_{(m,s),r}^{\theta,R} \leq \tau_{(m,t),r}^{\theta,L} \quad \text{for all } r \geq m.$$

A key result for proving the coalescence of semi-infinite geodesics is the nonexistence of bi-infinite geodesics for fixed directions. This is proven by first solving the ‘‘midpoint problem.’’ This problem first appeared in the context of first-passage percolation in a paper of Benjamini, Kalai, Schramm [BKS03]. In that context the problem asks whether

$$\lim_{n \rightarrow \infty} \mathbb{P}(\lfloor \frac{n}{2} \rfloor e_1 \text{ lies on some geodesic between } 0 \text{ and } ne_1) = 0.$$

In 2016, Damron and Hanson [DH17] proved that this is true, under the assumption that the boundary of the limit shape is differentiable. Later Ahlberg and Hoffman [AH16] proved this result without the assumption of differentiability. The following formulation of the midpoint problem more closely matches that for exponential last-passage percolation in the arXiv version of [Sep20], Theorem 4.12.

Lemma 3.15 (Midpoint problem). *Let $\theta, \eta > 0$ and $(m, t) \in \mathbb{Z} \times \mathbb{R}$. Then, the following subset of Ω is contained in an event of probability zero:*

$$\left\{ \begin{array}{l} \text{There exists a sequence } \{t_n\}_{n \in \mathbb{Z}} \text{ satisfying } \lim_{n \rightarrow \infty} \frac{t_n}{n} = \theta \text{ and } \lim_{n \rightarrow \infty} \frac{t_{-n}}{-n} = \eta \\ \\ \text{such that, for each } n \in \mathbb{Z}_{>0}, \text{ some geodesic between } (-n, t_{-n}) \text{ and } (n, t_n) \text{ passes through } (m, t) \end{array} \right\}$$

The following is a direct corollary of Lemma 3.15.

Theorem 3.16. *Fix $\theta, \eta > 0$. Then, there exists an event, $\Omega^{(\theta, \eta)}$, of probability one, on which there are no bi-infinite geodesics defined by jump times $\dots \leq \tau_{-1} \leq \tau_0 \leq \tau_1 \leq \dots$ such that*

$$(3.12) \quad \lim_{n \rightarrow \infty} \frac{\tau_n}{n} = \theta \quad \text{and} \quad \lim_{n \rightarrow \infty} \frac{\tau_{-n}}{n} = -\eta.$$

Proof. Let $\Omega^{(\theta)}$ be a full probability event on which, for every $(m, q) \in \mathbb{Z} \times \mathbb{Q}$, there exists no sequences satisfying the conditions (3.12) and such that, for every $n \in \mathbb{Z}$, there exists a geodesic between $(-n, \tau_{-n})$ and (n, τ_n) that passes through (m, q) . Then, on this event, if a bi-infinite geodesic that satisfies (3.12) exists, it cannot pass through (m, q) for any $q \in \mathbb{Q}$. But then this bi-infinite geodesic cannot ever move horizontally and so there must exist $t \in \mathbb{R}$ such that the bi-infinite geodesic consists only of points (r, t) . Now (3.12) fails. \square

Our last main result gives the coalescence in a fixed direction.

Theorem 3.17. *Let $\tilde{\Omega}^{(\theta)}$ be the full-probability event of Theorem 3.12. For each $\theta > 0$, there exists an event $\hat{\Omega}^{(\theta)} \subseteq \tilde{\Omega}^{(\theta)}$, of probability one, on which all θ -directed semi-infinite geodesics coalesce.*

3.4. Connection to infinite-length polymer measures for the Brownian polymer. Recalling the definitions at the beginning of Section 2.2, for $(m, s) \leq (n, t) \in \mathbb{Z} \times \mathbb{R}$, the point-to-point partition function of the Brownian polymer with unit temperature is

$$Z_{(m,s),(n,t)}(\mathbf{B}) = \int e^{\mathcal{E}(\mathbf{s}_{m,n})} \mathbf{1}\{\mathbf{s}_{m,n-1} \in \Pi_{(m,s),(n,t)}\} d\mathbf{s}_{m,n-1}.$$

The associated quenched polymer measure on $\Pi_{(m,s),(n,t)}$ is

$$\mathbb{Q}^{\mathbf{B}}(\tau_m \in ds_m, \dots, \tau_{n-1} \in ds_{n-1}) = \frac{1}{Z_{(m,s),(n,t)}(\mathbf{B})} \exp\left(\sum_{k=m}^n B_k(s_{k-1}, s_k)\right) \mathbf{1}\{\mathbf{s}_{m,n-1} \in \Pi_{(m,s),(n,t)}\} d\mathbf{s}_{m,n-1}.$$

Brownian last-passage percolation is the zero-temperature analogue of the Brownian or O’Connell-Yor polymer. This is made precise by the limit

$$\lim_{\beta \rightarrow \infty} \frac{1}{\beta} \log Z_{(m,s),(n,t)}(\beta \mathbf{B}) = L_{(m,s),(n,t)}(\mathbf{B}),$$

which, by convergence of L^p norms, holds in a deterministic sense as long as \mathbf{B} is a field of continuous functions $\mathbb{R} \rightarrow \mathbb{R}$. The parameter β is used to denote inverse temperature.

Alberts, Rassoul-Agha, and Simper [ARAS20] showed the existence of Busemann functions and infinite-length limits of the quenched measures when the right endpoint (n, t_n) satisfies $t_n/n \rightarrow \theta$ for some fixed $\theta > 0$. The Busemann functions are defined as

$$\widehat{\mathcal{B}}^\theta(\mathbf{x}, \mathbf{y}) := \lim_{n \rightarrow \infty} \log \frac{Z_{\mathbf{x},(n,t_n)}(\mathbf{B})}{Z_{\mathbf{y},(n,t_n)}(\mathbf{B})}.$$

Similarly as in (2.6), define $\widehat{h}_m^\theta(t) = \widehat{\mathcal{B}}^\theta((m, 0), (m, t))$. Using the “Proof of Theorem 2.5, assuming Theorem 3.1” on page 1937 of [ARAS20] and an analogous construction of the Busemann functions for all initial points as in Section 4 of the present paper, it can be shown that, the process $t \mapsto \widehat{h}_m^\theta(t)$ is a two-sided Brownian motion with drift, independent of $\{B_r\}_{r \leq m}$. By Theorems 3.2(vii) and 3.4(i), the same is true for the zero temperature Busemann process $t \mapsto h_m^\theta(t)$. An infinite, up-right path under the quenched infinite-length polymer measure is a continuous-time Markov chain, starting from a point (m, t) and defined by jump times $t = \tau_{m-1} \leq \tau_m \leq \dots$. By Equation (2.3) in [ARAS20], the quenched conditional distribution of τ_r given τ_{r-1} is

$$Q_{(m,s),(n,t_n)}^\mathbf{B}(\tau_r \in ds_r | \tau_{r-1} = s_{r-1}) = e^{B_r(s_{r-1}, s_r)} \frac{Z_{(r+1, s_r), (n, t_n)}(\mathbf{B})}{Z_{(r, s_{r-1}), (n, t_n)}(\mathbf{B})} \mathbf{1}\{s_{r-1} \leq s_r\} ds_r.$$

The proof of existence of infinite length measures requires a rigorous tightness argument, but to motivate the connection to BLPP, we formally take limits as $n \rightarrow \infty$ to yield the conditional measure

$$\begin{aligned} & \exp(B_r(s_{r-1}, s_r) + \widehat{\mathcal{B}}^\theta((r+1, s_r), (r, s_{r-1}))) \mathbf{1}\{s_{r-1} \leq s_r\} ds_r \\ &= \exp(B_r(s_{r-1}, s_r) - \widehat{h}_{r+1}(s_r) + \widehat{\mathcal{B}}^\theta((r+1, 0), (r, s_{r-1}))) \mathbf{1}\{s_{r-1} \leq s_r\} ds_r. \end{aligned}$$

Now, note that the term $\widehat{\mathcal{B}}^\theta((r+1, 0), (r, s_{r-1}))$ does not depend on s_r . We can then think of the connection between BLPP and the Brownian polymer in the following sense: In the positive temperature case, given the environment \mathbf{B} , the transition density from τ_{r-1} to τ_r is given by

$$C \exp(B_r(s) - \widehat{h}_{r+1}^\theta(s)) \mathbf{1}\{\tau_{r-1} \leq s\} ds,$$

where C is a normalizing constant depending on r and τ_{r-1} . On the other hand, by Definition 3.6, in the zero-temperature case, given the environment \mathbf{B} and the previous jump τ_{r-1} , the jump τ_r is chosen in a deterministic fashion by maximizing $B_r(s) - h_{r+1}^\theta(s)$ over $s \in [\tau_{r-1}, \infty)$.

3.5. Connection to semi-infinite geodesics in discrete last-passage percolation. The corner growth model, or discrete last-passage percolation, is defined as follows. Let $\{Y_{\mathbf{x}}\}_{\mathbf{x} \in \mathbb{Z}^2}$ be a collection of nonnegative i.i.d random variables, each associated to a vertex on the integer lattice. For $\mathbf{x} \leq \mathbf{y} \in \mathbb{Z} \times \mathbb{Z}$, define the last-passage time as

$$G_{\mathbf{x}, \mathbf{y}} = \sup_{\mathbf{x} \in \Pi_{\mathbf{x}, \mathbf{y}}} \sum_{k=0}^{|\mathbf{y}-\mathbf{x}|_1} Y_{\mathbf{x}_k},$$

where $\Pi_{\mathbf{x}, \mathbf{y}}$ is the set of up-right paths $\{\mathbf{x}_k\}_{k=0}^n$ that satisfy $\mathbf{x}_0 = \mathbf{x}$, $\mathbf{x}_n = \mathbf{y}$, and $\mathbf{x}_k - \mathbf{x}_{k-1} \in \{\mathbf{e}_1, \mathbf{e}_2\}$. Under the assumption that Y_0 has finite second moment, Theorem 3.1 and Corollary 3.1 of [GW91] introduced BLPP as a universal scaling limit of the corner growth model, where one variable is scaled and the other is held constant. That is, if Y_0 is normalized to have unit mean and variance,

$$\{L_{(m,s),(n,t)} : (m, s) \leq (n, t) \in \mathbb{Z} \times \mathbb{R}\}$$

is the functional limit, as $k \rightarrow \infty$, of the properly interpolated version of the process

$$\left\{ \frac{1}{\sqrt{k}} G_{(m, \lfloor sk \rfloor), (n, \lfloor tk \rfloor)} - (t-s)k : (m, s) \leq (n, t) \in \mathbb{Z} \times \mathbb{R} \right\}.$$

The most tractable case of discrete last-passage percolation is the case where Y_0 has the exponential distribution with rate 1. In this case, Busemann functions exist and are indexed by a direction vector \mathbf{u} . They are defined by

$$U^\mathbf{u}(\mathbf{x}, \mathbf{y}) := \lim_{n \rightarrow \infty} G_{(\mathbf{x}, z_n)} - G_{(\mathbf{y}, z_n)},$$

where \mathbf{z}_n satisfies $\mathbf{z}_n/n \rightarrow \mathbf{u}$ for a fixed \mathbf{u} . For a given $\omega \in \Omega$, $\mathbf{x} \in \mathbb{Z}^2$, and direction \mathbf{u} , a semi-infinite geodesic γ is defined by the sequence $\{\gamma_k^{\mathbf{u}, \mathbf{x}}\}_{k \in \mathbb{Z}_{\geq 0}}$. At each step, a choice is made to move upward or to the right. First, set $\gamma_0^{\mathbf{u}, \mathbf{x}} = \mathbf{x}$, and for $k \geq 0$,

$$(3.13) \quad \gamma_{k+1}^{\mathbf{u}, \mathbf{x}} = \begin{cases} \gamma_k^{\mathbf{u}, \mathbf{x}} + \mathbf{e}_1, & \text{if } U^{\mathbf{u}}(\gamma_k^{\mathbf{u}, \mathbf{x}}, \gamma_k^{\mathbf{u}, \mathbf{x}} + \mathbf{e}_1) \leq U^{\mathbf{u}}(\gamma_k^{\mathbf{u}, \mathbf{x}}, \gamma_k^{\mathbf{u}, \mathbf{x}} + \mathbf{e}_2), \\ \gamma_k^{\mathbf{u}, \mathbf{x}} + \mathbf{e}_2, & \text{if } U^{\mathbf{u}}(\gamma_k^{\mathbf{u}, \mathbf{x}}, \gamma_k^{\mathbf{u}, \mathbf{x}} + \mathbf{e}_2) < U^{\mathbf{u}}(\gamma_k^{\mathbf{u}, \mathbf{x}}, \gamma_k^{\mathbf{u}, \mathbf{x}} + \mathbf{e}_1). \end{cases}$$

In the case of exponential weights, this sequence is a semi-infinite geodesic with direction \mathbf{u} . This construction is inherently discrete, so it does not extend directly to the case of BLPP. However, this construction is equivalent to taking a sequence of maximizers of the appropriate function, analogous to Definition 3.6. The following is a discrete analogue of Theorem 3.2(vi) that holds in the case of exponential weights.

$$(3.14) \quad U^{\mathbf{u}}((m, r), (m, r + 1)) = \max_{\substack{m \leq k < \infty \\ k \in \mathbb{Z}}} \left\{ \sum_{i=m}^k Y_{(i, r)} + U^{\mathbf{u}}((k, r + 1), (m, r + 1)) \right\}.$$

Now, suppose that, starting at the point (m, r) , the semi-infinite geodesic constructed in (3.13) makes an \mathbf{e}_2 step from (k, r) to $(k, r + 1)$ for some $k \geq m$. Then, we show that k is maximal for (3.14). Indeed, by (3.13), if the path γ makes an \mathbf{e}_2 step from (k, r) to $(k, r + 1)$, then

$$U^{\mathbf{u}}((i, r), (i + 1, r)) \leq U^{\mathbf{u}}((i, r), (i, r + 1)) \quad \text{for } m \leq i \leq k - 1, \quad \text{and}$$

$$U^{\mathbf{u}}((k, r), (k + 1, r)) > U^{\mathbf{u}}((k, r), (k, r + 1)).$$

Using the identity $Y_{\mathbf{x}} = U^{\mathbf{u}}(\mathbf{x}, \mathbf{x} + \mathbf{e}_1) \wedge U^{\mathbf{u}}(\mathbf{x}, \mathbf{x} + \mathbf{e}_2)$ and additivity of the Busemann functions,

$$\begin{aligned} & \sum_{i=m}^k Y_{(i, r)} + U^{\mathbf{u}}((k, r + 1), (m, r + 1)) \\ &= \sum_{i=m}^{k-1} U^{\mathbf{u}}((i, r), (i + 1, r)) + U^{\mathbf{u}}((k, r), (k, r + 1)) + U^{\mathbf{u}}((k, r + 1), (m, r + 1)) \\ &= U^{\mathbf{u}}((m, r), (m, r + 1)), \end{aligned}$$

so k is indeed maximal in for the right hand-side of (3.14). The inductive step follows in the same manner. Hence, we see that the construction of semi-infinite geodesics in Definition 3.6 is a continuous analogue of the procedure for the discrete case when viewed from the perspective of maximizers.

3.6. Connection to queuing theory. Fix $\theta > 0$, and consider the almost surely unique θ -directed semi-infinite geodesic (Theorem 3.10(iii)) starting from $(0, 0)$ and defined by the sequence of jump times $\{\tau_r\}_{r \geq -1}$ with $\tau_{-1} = 0$. For each $r \geq 0$, associate a queuing station as follows. For $s < t$, let $\frac{1}{\sqrt{\theta}}(t - s) - X_{r+1}^\theta(s, t)$ denote the service available in the interval $(s, t]$ and let $\frac{1}{\sqrt{\theta}}(t - s) - h_r^\theta(s, t)$ denote the arrivals to the queue in the interval $(s, t]$. Here, the parameter $\frac{1}{\sqrt{\theta}}$ dictates the rate of service. See Section A.3 for more details about the queuing setup. By (3.9), $h_{r+1}^\theta = \overleftarrow{D}(h_r^\theta, X_{r+1}^\theta)$, so as in the proof of Theorem A.13, $\frac{1}{\sqrt{\theta}}(t - s) - h_{r+1}^\theta(s, t)$ gives the departures from the r th station in the interval $(s, t]$, and the departures process from the r th station becomes the arrivals process for the $r + 1$ st station. In other words, once a customer is served at the r th station, they move into the queue at the $r + 1$ st station. By Equations (3.8) and (3.9), for all $r \in \mathbb{Z}$ and $t \in \mathbb{R}$,

$$v_{r+1}^\theta(t) = \sup_{t \leq u < \infty} \{B_r(t, u) - h_{r+1}^\theta(t, u)\} = \sup_{-\infty < u \leq s} \{X_{r+1}^\theta(t, u) - h_r^\theta(t, u)\} = \overleftarrow{Q}(h_r^\theta, X_{r+1}^\theta)(t).$$

In queuing terms, $v_{r+1}^\theta(t)$ gives the length of the r th queue at time t . As τ_r is the maximizer of $B_r(u) - h_{r+1}^\theta(u)$ over $s \in [\tau_{r-1}, \infty)$ it is the first time greater than τ_{r-1} that $v_{r+1}^\theta(s) = 0$ (see Lemma 5.2). In queuing terms, τ_r is the first time greater than τ_{r-1} that the queue is empty. Thus, the semi-infinite geodesic represents the movement of a customer through the infinite series of queuing stations: the customer starts at station 0 at time 0 and is served at the r th station at a time no later than τ_r .

4. CONSTRUCTION OF THE BUSEMANN PROCESS

4.1. Construction of Busemann functions for a fixed direction.

Definition 4.1 (Definition of Busemann functions for fixed θ and a countable dense set of points). For a fixed $\theta > 0$, and $\mathbf{x}, \mathbf{y} \in \mathbb{Z} \times \mathbb{Q}$, let

$$\Omega_0^{(\theta)} = \bigcap_{\mathbf{x}, \mathbf{y} \in \mathbb{Z} \times \mathbb{Q}} \Omega_{\mathbf{x}, \mathbf{y}}^{(\theta)},$$

where the $\Omega_{\mathbf{x}, \mathbf{y}}^{(\theta)}$ are the events of Theorem 2.5. Then, $\mathbb{P}(\Omega_0^{(\theta)}) = 1$. On this event, define $\mathcal{B}^\theta(\mathbf{x}, \mathbf{y})$ by (2.4). By definition in terms of limits, it is clear that the Busemann functions are additive, that is, for $\omega \in \Omega_0^{(\theta)}$ and $\mathbf{x}, \mathbf{y}, \mathbf{z} \in \mathbb{Z} \times \mathbb{Q}$,

$$(4.1) \quad \mathcal{B}^\theta(\mathbf{x}, \mathbf{y}) + \mathcal{B}^\theta(\mathbf{y}, \mathbf{z}) = \mathcal{B}^\theta(\mathbf{x}, \mathbf{z}).$$

Hence, the entire collection of Busemann functions is constructed from the collection of horizontal and vertical Busemann functions $\{h_m^\theta, v_m^\theta\}_{m \in \mathbb{Z}}$ (Equations (2.6) and (2.5)).

Set $Y(t) = -B_0(t) + \frac{1}{\sqrt{\theta}}t$. By Lemma A.16, as elements of the space of functions $\mathbb{Q} \rightarrow \mathbb{R}$, equipped with the standard product σ -algebra,

$$\{h_0^\theta(t) : t \in \mathbb{Q}\} \stackrel{d}{=} \{Y(t) : t \in \mathbb{Q}\} \quad \text{and} \quad \{v_1^\theta(t) : t \in \mathbb{Q}\} \stackrel{d}{=} \{\overleftarrow{Q}(Y, B_1)(t) : t \in \mathbb{Q}\}.$$

For $\omega \in \Omega_0^{(\theta)}$ and each $m \in \mathbb{Z}$,

$$h_m^\theta(t) = \lim_{n \rightarrow \infty} L_{(m,0), (n, n\theta)}(\mathbf{B}) - L_{(m,t), (n, n\theta)}(\mathbf{B}) = \lim_{n \rightarrow \infty} L_{(m,0), (n+m, (n+m)\theta)}(\mathbf{B}) - L_{(m,t), (n+m, (n+m)\theta)}(\mathbf{B}).$$

Since the environment $\mathbf{B} = \{B_m\}_{m \in \mathbb{Z}}$ is a field of i.i.d. two-sided Brownian motions, the right-hand side implies that $h_m^\theta(t)$ has the same distribution for all m . By this same reasoning, for each $m \in \mathbb{Z}$,

$$\{h_m^\theta(t), v_{m+1}^\theta(t) : t \in \mathbb{Q}\} \stackrel{d}{=} \{h_0^\theta(t), v_1^\theta(t) : t \in \mathbb{Q}\}.$$

Since Brownian motion satisfies $\lim_{s \rightarrow \infty} \frac{B(s)}{s} = 0$ almost surely, for all $m \in \mathbb{Z}$, the following limit also holds almost surely:

$$\lim_{\mathbb{Q} \ni s \rightarrow \infty} B_m(s) - h_{m+1}^\theta(s) = -\infty.$$

Set

$$\Omega^{(\theta)} = \bigcap_{m, r \in \mathbb{Z}} \{(s, t) \mapsto \mathcal{B}^\theta((m, s), (r, t)) \text{ is uniformly continuous on all bounded subsets of } \mathbb{Q} \times \mathbb{Q}\}$$

$$(4.2) \quad \cap \bigcap_{m \in \mathbb{Z}} \{\limsup_{\mathbb{Q} \ni s \rightarrow \infty} [B_m(s) - h_{m+1}^\theta(s)] = -\infty\}.$$

By almost sure continuity of the functions Y and $\overleftarrow{Q}(Y, B_1)$ (Lemma 3.1), $\mathbb{P}(\Omega^{(\theta)}) = 1$.

Definition 4.2 (Definition of Busemann functions for fixed θ , arbitrary points). On the event $\Omega^{(\theta)}$, for arbitrary $\mathbf{x}, \mathbf{y} \in \mathbb{Z} \times \mathbb{R}$, define $\mathcal{B}^\theta(\mathbf{x}, \mathbf{y})$ such that, for each $m, r \in \mathbb{Z}$,

$$(s, t) \mapsto \mathcal{B}^\theta((m, s), (r, t))$$

is the unique continuous extension of this function from $\mathbb{Q} \times \mathbb{Q}$ to \mathbb{R}^2 .

The following lemma states properties for a fixed θ , as a precursor to the more general Theorems 3.2 and Theorem 3.4.

Lemma 4.3. *The following hold.*

(i) *On the event $\Omega^{(\theta)}$, for all $\mathbf{x}, \mathbf{y} \in \mathbb{Z} \times \mathbb{R}$ and all sequences $\{t_n\}$ satisfying $t_n/n \rightarrow \theta$,*

$$\mathcal{B}^\theta(\mathbf{x}, \mathbf{y}) = \lim_{n \rightarrow \infty} [L_{\mathbf{x}, (n, t_n)} - L_{\mathbf{y}, (n, t_n)}].$$

(ii) *On the event $\Omega^{(\theta)}$ (which has full probability), whenever $\mathbf{x}, \mathbf{y}, \mathbf{z} \in \mathbb{Z} \times \mathbb{R}$,*

$$\mathcal{B}^\theta(\mathbf{x}, \mathbf{y}) + \mathcal{B}^\theta(\mathbf{y}, \mathbf{z}) = \mathcal{B}^\theta(\mathbf{x}, \mathbf{z}).$$

(iii) *For $0 < \gamma < \theta < \infty$, on the event $\Omega^{(\theta)} \cap \Omega^{(\gamma)}$, for all $m \in \mathbb{Z}$ and $s < t \in \mathbb{R}$,*

$$0 \leq v_m^\gamma(s) \leq v_m^\theta(s), \text{ and } B_m(s, t) \leq h_m^\theta(s, t) \leq h_m^\gamma(s, t).$$

(iv) *On the event $\Omega^{(\theta)}$, for each $m \in \mathbb{Z}$,*

$$\lim_{s \rightarrow \pm\infty} B_m(s) - h_{m+1}^\theta(s) = \mp\infty.$$

- (v) For every $m \in \mathbb{Z}$, the process $t \mapsto h_m^\theta(t)$ is a two-sided Brownian motion with drift $\frac{1}{\sqrt{\theta}}$. For each $m \in \mathbb{Z}$ and $t \in \mathbb{R}$, $v_m^\theta(t) \sim \text{Exp}(\frac{1}{\sqrt{\theta}})$.
- (vi) For any $m \in \mathbb{Z}$, $\{h_r^\theta\}_{r>m}$ is independent of $\{B_r\}_{r \leq m}$.
- (vii) There exists an event $\Omega_1^{(\theta)} \subseteq \Omega^{(\theta)}$, of full probability, on which the following hold for all $t \in \mathbb{R}$:

$$v_{m+1}^\theta(t) = Q(h_{m+1}^\theta, B_m)(t) \quad \text{and} \quad h_m^\theta(t) = D(h_{m+1}^\theta, B_m)(t).$$

Proof. **Part (i):** On $\Omega^{(\theta)}$, let $(m, s), (r, t) \in \mathbb{Z} \times \mathbb{R}$, and let $\{t_n\}$ be any sequence with $t_n/n \rightarrow \theta$. Note that, whenever $t_1 < t_2, m \in \mathbb{Z}$ and $\mathbf{y} \geq (m, t_2)$,

$$L_{(m, t_1), \mathbf{y}} \geq B_m(t_1, t_2) + L_{(m, t_2), \mathbf{y}}.$$

Then, let $q_1, q_2 \in \mathbb{Q}$ be such that $q_1 < s$ and $q_2 > t$. Then,

$$L_{(m, s), (n, t_n)} - L_{(r, t), (n, t_n)} \leq L_{(m, q_1), (n, t_n)} - B_m(q_1, s) - L_{(r, q_2), (n, t_n)} - B_r(t, q_2),$$

and therefore,

$$\limsup_{n \rightarrow \infty} [L_{(m, s), (n, t_n)} - L_{(r, t), (n, t_n)}] \leq \mathcal{B}^\theta((m, q_1), (r, q_2)) - B_m(q_1, s) - B_r(t, q_2).$$

Taking $q_1 \nearrow s$ and $q_2 \searrow t$ and using the continuity of Brownian motion and the Busemann functions,

$$\limsup_{n \rightarrow \infty} [L_{(m, s), (n, t_n)} - L_{(r, t), (n, t_n)}] \leq \mathcal{B}^\theta((m, s), (r, t)).$$

A similar procedure gives the appropriate lower bound.

Part (ii): This follows from the additivity of the process on the countable dense set (Equation (4.1)) and the construction of Definition 4.2 as the unique continuous extension.

Part (iii): By Lemma A.4, for $s, t \in \mathbb{R}$ and all n such that $n \geq m$ and $n\gamma \geq s \wedge t$,

$$0 \leq L_{(m, s), (n, n\gamma)} - L_{(m+1, s), (n, n\gamma)} \leq L_{(m, s), (n, n\theta)} - L_{(m+1, s), (n, n\theta)}$$

and $B_m(s, t) \leq L_{(m, s), (n, n\theta)} - L_{(m, t), (n, n\theta)} \leq L_{(m, s), (n, n\gamma)} - L_{(m, t), (n, n\gamma)}.$

The proof is complete by Part (i), taking limits as $n \rightarrow \infty$.

Part (iv): This follows from the definition of $\Omega^{(\theta)}$ and continuity.

Part (v): Observe that $t \mapsto h_m^\theta(t)$ is a continuous process with the correct finite-dimensional distributions by Lemma A.16, taking limits when necessary. Part (i) and Theorem 2.5 guarantee that $v_m^\theta(t) \sim \text{Exp}(\frac{1}{\sqrt{\theta}})$ for all $t \in \mathbb{R}$.

Part (vi): By Part (i),

$$\{h_r^\theta(t) : r > m, t \in \mathbb{R}\} = \left\{ \lim_{n \rightarrow \infty} [L_{(r, 0), (n, n\theta)} - L_{(r, t), (n, n\theta)}] : r > m, t \in \mathbb{R} \right\}.$$

The right-hand side is a function of $\{B_r\}_{r>m}$, which is independent of $\{B_r\}_{r \leq m}$.

Part (vii): We start with the first statement. We need to show that

$$(4.3) \quad v_{m+1}^\theta(t) = \sup_{t \leq s < \infty} \{B_m(t, s) - h_{m+1}^\theta(t, s)\}.$$

By Part (i),

$$(4.4) \quad \sup_{t \leq s < \infty} \{B_m(t, s) - h_{m+1}^\theta(t, s)\} = \sup_{t \leq s < \infty} \{B_m(t, s) + \lim_{n \rightarrow \infty} [L_{(m+1, s), (n, n\theta)} - L_{(m+1, t), (n, n\theta)}]\}.$$

On the other hand,

$$(4.5) \quad v_{m+1}^\theta(t) = \lim_{n \rightarrow \infty} [L_{(m, t), (n, n\theta)} - L_{(m+1, t), (n, n\theta)}].$$

For each $s \geq t$ and all n sufficiently large so that $n\theta \geq s$ and $n \geq m+1$,

$$B_m(t, s) + L_{(m+1, s), (n, n\theta)} - L_{(m+1, t), (n, n\theta)} \leq L_{(m, t), (n, n\theta)} - L_{(m+1, t), (n, n\theta)}.$$

Taking limits as $n \rightarrow \infty$ and comparing (4.4) and (4.5) establishes

$$v_{m+1}^\theta(t) \geq \sup_{t \leq s < \infty} \{B_m(t, s) - h_{m+1}^\theta(t, s)\}.$$

Next, by Part (v), $v_{m+1}^\theta(t) \sim \text{Exp}(\frac{1}{\sqrt{\theta}})$. By Parts (v) and (vi) and Lemma A.6,

$$\sup_{t \leq s < \infty} \{B_m(t, s) - h_{m+1}^\theta(t, s)\} \stackrel{d}{=} \sup_{0 \leq s < \infty} \left\{ \sqrt{2}B(s) - \frac{1}{\sqrt{\theta}}s \right\} \stackrel{d}{=} v_{m+1}^\theta(t),$$

where B is a standard Brownian motion. Thus, since $v_{m+1}^\theta(t) \geq \sup_{t \leq s < \infty} \{B_m(t, s) - h_{m+1}^\theta(t, s)\}$, equality holds with probability one for each fixed $t \in \mathbb{R}$. Let $\Omega_1^{(\theta)} \subseteq \Omega^{(\theta)}$, be the event of full probability on which (4.3) holds for all $t \in \mathbb{Q}$. Continuity of both sides of (4.3) (Definition 4.2 and Lemma 3.1) extend the result to all $t \in \mathbb{R}$ on $\Omega_1^{(\theta)}$. The equality for h_m^θ then follows by the definitions and additivity of the Busemann functions, as shown below.

$$\begin{aligned} D(h_{m+1}^\theta, B_m)(t) &= h_{m+1}^\theta(t) + Q(h_{m+1}^\theta, B_m)(0) - Q(h_{m+1}^\theta, B_m)(t) \\ &= h_{m+1}^\theta(t) + v_{m+1}^\theta(0) - v_{m+1}^\theta(t) \\ &= \mathcal{B}^\theta((m+1, 0), (m+1, t)) + \mathcal{B}^\theta((m, 0), (m+1, 0)) - \mathcal{B}^\theta((m, t), (m+1, t)) \\ &= \mathcal{B}^\theta((m, 0), (m, t)) = h_m^\theta(t). \end{aligned} \quad \square$$

4.2. Global construction of the Busemann process. We now have the proper framework to define global Busemann the process of Theorem 3.2. Fix a countable subset D of $(0, \infty)$. Let

$$\Omega_0 = \bigcap_{\theta \in D} \Omega_1^{(\theta)},$$

and on the event Ω_0 , define \mathcal{B}^θ for all $\theta \in D$, as in Definition 4.2. Then, the conclusions of Lemma 4.3 hold for all $\theta \in D$.

Lemma 4.4. *On Ω_0 , for each $\theta \in D$, $m \in \mathbb{Z}$, and $t \in \mathbb{R}$, let $\tau_{(m,t)}^\theta$ denote the right-most maximizer of $B_m(s) - h_{m+1}^\theta(s)$ over $s \in [t, \infty)$. Such a maximizer exists by continuity and Lemma 4.3(iv). Let Ω_1 be the subset of Ω_0 on which, for each $\theta \in D$, $m, N \in \mathbb{Z}$, and $t \in \mathbb{R}$, the following limits hold:*

- (i) $\lim_{D \ni \gamma \rightarrow \theta} h_m^\gamma(N) = h_m^\theta(N)$
- (ii) $\lim_{D \ni \gamma \rightarrow \infty} h_m^\gamma(N) = B_m(N)$.
- (iii) $\lim_{D \ni \delta \searrow 0} \tau_{(m,t)}^\delta = t$.
- (iv) $\lim_{D \ni \gamma \rightarrow \infty} \tau_{(m,t)}^\gamma = \infty$.

Then, $\mathbb{P}(\Omega_1) = 0$.

Proof. The almost sure uniqueness of maximizers follows from Lemma 4.3(vi) and the $n = m$ case of Lemma 2.3. By Lemma 4.3(iii), $h_m^\gamma(N) = h_m^\gamma(0, N)$ is monotone as $D \ni \gamma \nearrow \theta$, and for $N > 0$,

$$(4.6) \quad \lim_{D \ni \gamma \nearrow \theta} h_m^\gamma(N) \geq h_m^\theta(N),$$

For negative N , the inequality flips. By Theorem 2.5, $h_m^\gamma(N) \sim \mathcal{N}(\frac{N}{\sqrt{\gamma}}, |N|)$. As $\gamma \nearrow \theta$, this converges in distribution to $\mathcal{N}(\frac{N}{\sqrt{\theta}}, |N|)$. Then, by (4.6), $\lim_{D \ni \gamma \nearrow \theta} h_m^\gamma(N) = h_m^\theta(N)$ with probability one. Analogous argument proves the almost sure convergence for limits from the right and the convergence to $B_m(N)$.

Next, we show that, on Ω_0 , Parts (iii) and (iv) hold if and only if they hold for all $t \in \mathbb{Q}$. Assume the statement holds for all $t \in \mathbb{Q}$. By Lemmas 4.3(iii) and A.1, $\tau_{(m,t)}^\delta$ is monotone as $\delta \searrow 0$, and so the limit exists. By definition of $\tau_{(m,t)}^\theta$, we have the inequality $\tau_{(m,s)}^\theta \leq \tau_{(m,t)}^\theta$ whenever $s < t$. Then, for any $t \in \mathbb{R}$ and any $q_1, q_2 \in \mathbb{Q}$ with $q_1 < t < q_2$,

$$q_1 = \lim_{D \ni \delta \searrow 0} \tau_{(m,q_1)}^\delta \leq \lim_{D \ni \delta \searrow 0} \tau_{(m,t)}^\delta \leq \lim_{D \ni \delta \searrow 0} \tau_{(m,q_2)}^\delta = q_2.$$

Taking limits as $q_1 \nearrow t$ and $q_2 \searrow t$ shows the statement for all $t \in \mathbb{R}$. The same argument can be applied to the limits as $\gamma \rightarrow \infty$.

Lastly, we show that the limit in (iii) holds with probability one for a fixed $t \in \mathbb{R}$. By Lemma 4.3, Parts (v) and (vi),

$$\{B_m(s) - h_{m+1}^\delta(s) : s \in \mathbb{R}\} \stackrel{d}{=} \{\sqrt{2}B(s) - \frac{s}{\sqrt{\delta}} : s \in \mathbb{R}\},$$

where B is a standard, two-sided Brownian motion. Then, by Theorem A.8, for $(m, t) \in \mathbb{Z} \times \mathbb{R}$ and $s \geq 0$,

$$\mathbb{P}(\tau_{(m,t)}^\delta > s + t) = \left(2 + \frac{t}{\delta}\right) \Phi\left(-\sqrt{\frac{t}{2\delta}}\right) - \sqrt{\frac{t}{\pi\delta}} e^{-\frac{t}{4\delta}},$$

Taking limits as $\delta \searrow 0$, it follows that $\tau_{(m,t)}^\theta$ converges weakly to the constant t . Since the limit exists almost surely by monotonicity, the desired conclusion follows. A similar argument applies to show $\lim_{\gamma \rightarrow \infty} \tau_{(m,t)}^\gamma = \infty$ with probability one. \square

On Ω_1 , we extend the definition of \mathcal{B}^θ to all $\theta > 0$. We proceed in a similar manner as shown for the exponential corner growth model in [Sep18]. For BLPP, there is an additional step that must be taken to guarantee the convergence and continuity of Theorem 3.2, Parts (iii) and (iv).

Lemma 4.5. *On Ω_1 , for each $\theta \in D$, $m \in \mathbb{Z}$, and $t \in \mathbb{R}$,*

$$\lim_{D \ni \gamma \rightarrow \theta} h_m^\gamma(t) = h_m^\theta(t) \quad \text{and} \quad \lim_{D \ni \gamma \rightarrow \theta} v_m^\gamma(t) = v_m^\theta(t).$$

For each $m \in \mathbb{Z}$, the convergence is uniform in t on compact subsets of \mathbb{R} . Additionally, for each $m \in \mathbb{Z}$,

$$\lim_{D \ni \gamma \rightarrow \infty} h_m^\gamma(t) = B_m(t), \quad \text{and} \quad \lim_{D \ni \delta \searrow 0} v_m^\delta(t) = 0,$$

uniformly in t on compact subsets of \mathbb{R} .

Proof. We first prove the statements for the h_m . We show that

$$\lim_{D \ni \gamma \nearrow \theta} h_m^\gamma(t)$$

exists and equals $h_m^\theta(t)$, uniformly in t on compact subsets of \mathbb{R} . The limits from the right (as well as the case of B_m in place of h_m^θ) follow by analogous arguments.

By rearranging the inequality of Lemma 4.3(iii), for $0 < \gamma < \theta < \infty$ and $a < b$ and any $t \in [a, b]$,

$$(4.7) \quad h_m^\gamma(a) - h_m^\theta(a) \leq h_m^\gamma(t) - h_m^\theta(t) \leq h_m^\gamma(b) - h_m^\theta(b),$$

and the inequality still holds if we replace h_m^θ with B_m . Thus, on the event Ω_1 , h_m^γ converges to h_m^θ , uniformly in t on compact subsets of \mathbb{R} . Now, we prove the convergence statements for v_m . By Lemma 4.3(vii), on Ω_0 , for all $t \in \mathbb{R}$, and all $\theta \in D$.

$$(4.8) \quad v_m^\theta(t) = Q(h_m^\theta, B_{m-1})(t) = \sup_{t \leq s < \infty} \{B_{m-1}(t, s) - h_m^\theta(t, s)\}.$$

Let $\omega \in \Omega_1$, and $t \in \mathbb{R}$. By the monotonicity of Lemma 4.3(iii) and Lemma A.1, since $\tau_{(m,t)}^{\theta+1}$ is a maximizer of $B_{m-1}(s) - h_m^{\theta+1}(s)$ over $s \in [t, \infty)$, for all $\gamma \in D$ with $\gamma < \theta + 1$,

$$v_m^\gamma(t) = \sup_{t \leq s < \infty} \{B_{m-1}(t, s) - h_m^\gamma(t, s)\} = \sup_{t \leq s \leq \tau_{(m,t)}^{\theta+1}} \{B_{m-1}(t, s) - h_m^\gamma(t, s)\}.$$

Since h_m^γ converges uniformly on compact sets to h_m^θ , $v_m^\gamma(t)$ converges pointwise to $v_m^\theta(t)$. By Lemma 4.3(iii), the convergence from both right and left is monotone, so by the continuity of Definition 4.2 and Dini's Theorem, the convergence is uniform.

Lastly, for limits as $\delta \searrow 0$, we again apply Lemmas 4.3(iii) and A.1 so that, for $\delta \in D$ with $\delta \leq 1$,

$$\begin{aligned} v_m^\delta(t) &= \sup_{t \leq s < \infty} \{B_{m-1}(t, s) - h_m^\delta(t, s)\} = \sup_{t \leq s \leq \tau_{(m,t)}^\delta} \{B_{m-1}(t, s) - h_m^\delta(t, s)\} \\ &\leq \sup_{t \leq s \leq \tau_{(m,t)}^\delta} \{B_{m-1}(t, s) - h_m^1(t, s)\}, \end{aligned}$$

and the right-hand side converges to 0 as $\delta \searrow 0$ by the continuity of $B_{m-1} - h_m^1$ and the convergence of $\tau_{(m,t)}^\delta$ to t given in the definition of Ω_1 . Dini's Theorem again strengthens the pointwise convergence to uniform convergence. \square

By the additivity of Lemma 4.3(ii), for arbitrary $(m, s), (r, t) \in \mathbb{Z} \times \mathbb{R}$ with $m \leq r$,

$$\mathcal{B}^\theta((m, s), (r, t)) = h_m^\theta(s, t) + \sum_{k=m}^{r-1} v_{k+1}(t).$$

For $m > r$, $\mathcal{B}^\theta((m, s), (r, t)) = -\mathcal{B}^\theta((r, t), (m, s))$, so $\mathcal{B}^\theta((m, s), (r, t))$ is still a sum of horizontal and vertical increments. Then, by Lemma 4.5, on the event Ω_1 , for all $\theta \in D$ and $\mathbf{x}, \mathbf{y} \in \mathbb{Z} \times \mathbb{R}$,

$$(4.9) \quad \lim_{D \ni \gamma \rightarrow \theta} \mathcal{B}^\gamma(\mathbf{x}, \mathbf{y}) = \mathcal{B}^\theta(\mathbf{x}, \mathbf{y}).$$

Definition 4.6. On the event Ω_1 , for an arbitrary $\theta > 0$, $\mathbf{x}, \mathbf{y} \in \mathbb{Z} \times \mathbb{R}$, define the following.

$$\mathcal{B}^{\theta-}(\mathbf{x}, \mathbf{y}) = \lim_{D \ni \gamma \nearrow \theta} \mathcal{B}^\gamma(\mathbf{x}, \mathbf{y}), \quad \text{and} \quad \mathcal{B}^{\theta+}(\mathbf{x}, \mathbf{y}) = \lim_{D \ni \delta \searrow \theta} \mathcal{B}^\delta(\mathbf{x}, \mathbf{y}).$$

Remark 4.7. By additivity of Lemma 4.3(ii) and the monotonicity of Lemma 4.3(iii), these limits exist for all $\mathbf{x}, \mathbf{y} \in \mathbb{Z} \times \mathbb{R}$ and $\theta > 0$. By (4.9), on Ω_1 , for all $\theta \in D$ and $\mathbf{x}, \mathbf{y} \in \mathbb{Z} \times \mathbb{R}$, $\mathcal{B}^{\theta+}(\mathbf{x}, \mathbf{y}) = \mathcal{B}^{\theta-}(\mathbf{x}, \mathbf{y}) = \mathcal{B}^{\theta}(\mathbf{x}, \mathbf{y})$.

Proof of Theorem 3.2. Parts (i)-(ii): These follow by taking limits in Parts (ii) and (iii) of Lemma 4.3.

Parts (iii)-(iv): By Part (ii), it suffices to take limits along the countable dense set D . Then, Lemma 4.5 establishes Parts (c) and (d). The monotonicity of Part (ii) can be rearranged, just as in Equation (4.7), which strengthens the pointwise convergence of the h_m^γ to uniform convergence on compact sets. Thus, since h_m^γ is continuous for $\gamma \in D$, $h_m^{\theta\Box}$ is also continuous for $\Box \in \{+, -\}$. Now, for the convergence of the $v_m^{\theta\Box}$, recall that on the event Ω_1 , for all $t \in \mathbb{R}$ and $\gamma \in D$,

$$v_m^\gamma(t) = Q(h_m^\gamma, B_{m-1})(t) = \sup_{t \leq s < \infty} \{B_{m-1}(t, s) - h_m^\gamma(t, s)\}.$$

By Part (ii) and Lemma A.1, this supremum may be restricted to a common compact set for all $\gamma < \theta$. Then, since h_m^γ converges uniformly to $h_m^{\theta-}$ as $D \ni \gamma \nearrow \theta$, $v_m^\gamma(t)$ converges pointwise to $Q(h_m^{\theta-}, B_{m-1})(t)$ as $\gamma \nearrow \theta$. But, by definition, $v_m^{\theta-}(t) = \lim_{D \ni \gamma \nearrow \theta} v_m^\gamma(t)$, so $v_m^{\theta-}(t) = Q(h_m^{\theta-}, B_{m-1})(t)$, which is continuous by continuity of $h_m^{\theta-}$ and B_{m-1} . Since the convergence is monotone, Dini's Theorem implies the convergence is uniform on compact sets. The proof for $\delta \searrow \theta$ is analogous. Part (iv) follows by continuity of the Busemann functions from Definition 4.2, the uniform convergence of Part (iii), and the additivity in Part (i).

Part (v): This follows from Lemma 4.3(iv) and the monotonicity of Part (ii).

Part (vi): The equality for $v_{m+1}^{\theta\Box}$ was shown in the proof of Part (iii). The equality for h_m^θ follows by the additivity of Part (i) and the same argument as in the proof of Lemma 4.3(vii).

Part (vii): This follows because $\{h_r^{\theta\Box} : \theta > 0, \Box \in \{+, -\}, r > m\}$ is a function of $\{B_r\}_{r > m}$.

Part (viii): Fix $\theta > 0$. We first show that, for each $\theta > 0$ and $\mathbf{x}, \mathbf{y} \in \mathbb{Z} \times \mathbb{R}$, there exists an event $\Omega_{\mathbf{x}, \mathbf{y}}^{(\theta)}$ on which

$$(4.10) \quad \mathcal{B}^{\theta-}(\mathbf{x}, \mathbf{y}) = \lim_{n \rightarrow \infty} [L_{\mathbf{x}, (n, t_n)} - L_{\mathbf{y}, (n, t_n)}] = \mathcal{B}^{\theta+}(\mathbf{x}, \mathbf{y}),$$

for all sequences $\{t_n\}$ with $\lim_{n \rightarrow \infty} \frac{t_n}{n} = \theta$. By Theorem 2.5, for each \mathbf{x}, \mathbf{y} , and $\theta > 0$, there exists an event of full probability on which the limit in (4.10) exists and is independent of the choice of sequence. Then, by additivity, it is sufficient to show that, for each fixed $\theta > 0$, $m \in \mathbb{Z}$, and $t \in \mathbb{R}$, with probability one,

$$h_m^{\theta-}(t) = \lim_{n \rightarrow \infty} [L_{(m, 0), (n, n\theta)} - L_{(m, t), (n, n\theta)}] = h_m^{\theta+}(t), \text{ and}$$

$$v_m^{\theta-}(t) = \lim_{n \rightarrow \infty} [L_{(m-1, t), (n, n\theta)} - L_{(m, t), (n, n\theta)}] = v_m^{\theta+}(t).$$

We show that

$$(4.11) \quad h_m^{\theta-}(t) = \lim_{n \rightarrow \infty} [L_{(m, 0), (n, n\theta)} - L_{(m, t), (n, n\theta)}] \quad \text{a.s.},$$

and the other statements follow by analogous arguments. By Theorem 2.5, the right-hand side of (4.11) has distribution $\mathcal{N}(\frac{t}{\sqrt{\theta}}, |t|)$. By definition, $h_m^{\theta-}(t)$ is the limit, as $\gamma \nearrow \theta$, of $h_m^\gamma(t)$, and $h_m^\gamma(t) \sim \mathcal{N}(\frac{t}{\sqrt{\gamma}}, |t|)$, which converges weakly to $\mathcal{N}(\frac{t}{\sqrt{\theta}}, |t|)$. Using Theorem 4.3(i) and Lemma A.4, for each fixed $t > 0$ and all $D \ni \gamma < \theta$,

$$h_m^\gamma(t) = \lim_{n \rightarrow \infty} [L_{(m, 0), (n, n\gamma)} - L_{(m, t), (n, n\gamma)}] \geq \lim_{n \rightarrow \infty} [L_{(m, 0), (n, n\theta)} - L_{(m, t), (n, n\theta)}] \quad \text{a.s.}$$

The inequality flips for $t < 0$. Hence, both sides in (4.11) have the same distribution, while one dominates the other, so they are equal with probability one.

Next, set

$$\Omega^{(\theta)} = \bigcap_{\mathbf{x}, \mathbf{y} \in \mathbb{Z} \times \mathbb{Q}} \Omega_{\mathbf{x}, \mathbf{y}}^{(\theta)}.$$

Then, using the continuity of Part (iv), the desired conclusion follows by the same reasoning as in the proof of Lemma 4.3(i). \square

5. PROOFS OF THE RESULTS FOR SEMI-INFINITE GEODESICS

Throughout this section, Ω_1 is the event given in Theorem 3.2.

5.1. Key lemmas.

Lemma 5.1. *Let $\omega \in \Omega_1$, $(m, t) \in \mathbb{Z} \times \mathbb{R}$, $\theta > 0$, $\square \in \{+, -\}$ and $\{\tau_r\}_{r=m-1}^\infty$ be any sequence in $\mathbf{T}_{(m,t)}^{\theta\square}$. Then, for each $n \geq m$, the jump times $t = \tau_{m-1} \leq \tau_m \leq \dots \leq \tau_n$ are a maximizing sequence for*

$$(5.1) \quad \bar{L}_{(m,t),n}^\lambda(\mathbf{B}, \bar{\mathbf{B}}) := \sup \left\{ \sum_{r=m}^n B_r(s_{r-1}, s_r) - h_{n+1}^{\theta\square}(s_n) : \mathbf{s}_{m,n} \in \Pi_{(m,t),n} \right\}.$$

Here, $\bar{L}_{(m,t),n}^\lambda(\mathbf{B}, \bar{\mathbf{B}})$ is as in definition (2.3), where $\lambda = \frac{1}{\sqrt{\theta}}$, $\mathbf{B} = \{B_r\}_{r \in \mathbb{Z}}$, and $\bar{\mathbf{B}} = \{h_r^{\theta\square}(t) - \frac{t}{\sqrt{\theta}} : t \in \mathbb{R}\}_{r \in \mathbb{Z}}$.

Proof. Recall by (3.11), that $\tau_m \geq t$ is a maximizer of

$$B_m(t, s) - h_{m+1}^{\theta\square}(s) \quad \text{over} \quad s \in [t, \infty).$$

Hence, the statement holds for $n = m$. Now, assume that the statement holds for some $n \geq m$. Then, $t = \tau_{m-1} \leq \dots \leq \tau_n$ satisfies

$$(5.2) \quad \sum_{r=m}^n B_r(\tau_{r-1}, \tau_r) - h_{n+1}^{\theta\square}(\tau_n) = \sup \left\{ \sum_{r=m}^n B_r(s_{r-1}, s_r) - h_{n+1}^{\theta\square}(s_n) : \mathbf{s}_{m,n} \in \Pi_{(m,t),n} \right\}.$$

Using Theorem 3.2(vi) and rearranging the terms in the definition (3.2) of the operator D ,

$$(5.3) \quad h_{n+1}^{\theta\square}(s_n) = B_{n+1}(s_n) + \sup_{0 \leq s < \infty} \{B_{n+1}(s) - h_{n+2}^{\theta\square}(s)\} - \sup_{s_n \leq s_{n+1} < \infty} \{B_{n+1}(s_{n+1}) - h_{n+2}^{\theta\square}(s_{n+1})\},$$

for $s_n \in \mathbb{R}$. Specifically, since τ_{n+1} is a maximizer of $B_{n+1}(s) - h_{n+2}^{\theta\square}(s)$ over $s \in [\tau_n, \infty)$,

$$(5.4) \quad h_{n+1}^{\theta\square}(\tau_n) = -B_{n+1}(\tau_n, \tau_{n+1}) + h_{n+2}^{\theta\square}(\tau_{n+1}) + \sup_{0 \leq s < \infty} \{B_{n+1}(s) - h_{n+2}^{\theta\square}(s)\}$$

Substituting (5.3) and (5.4) into (5.2) and discarding the term $\sup_{0 \leq s < \infty} \{B_{n+1}(s) - h_{n+2}^{\theta\square}(s)\}$ on both sides

$$\begin{aligned} & \sum_{r=m}^{n+1} B_r(\tau_{r-1}, \tau_r) - h_{n+2}^{\theta\square}(\tau_{n+1}) \\ &= \sup \left\{ \sum_{r=m}^n B_r(s_{r-1}, s_r) - B_{n+1}(s_n) + \sup_{s_n \leq s_{n+1} < \infty} \{B_{n+1}(s_{n+1}) - h_{n+2}^{\theta\square}(s_{n+1})\} : \mathbf{s}_{m,n} \in \Pi_{(m,t),n} \right\} \\ &= \sup \left\{ \sum_{r=m}^{n+1} B_r(s_{r-1}, s_r) - h_{n+2}^{\theta\square}(s_{n+1}) : \mathbf{s}_{m,n+1} \in \Pi_{(m,t),n+1} \right\}. \quad \square \end{aligned}$$

Lemma 5.2. *Let $\omega \in \Omega_1$, $(m, t) \in \mathbb{Z} \times \mathbb{R}$, $\theta > 0$, and $\square \in \{+, -\}$, and $\{\tau_r\}_{r \geq m-1} \in \mathbf{T}_{(m,t)}^{\theta\square}$. Then, for all $r \geq m$,*

$$v_{r+1}^{\theta\square}(\tau_r) = 0, \quad \text{and} \quad h_r^{\theta\square}(u, v) = B_r(u, v) \text{ for all } u, v \in [\tau_{r-1}, \tau_r].$$

Furthermore, the following identities hold for $r \geq m$.

$$(5.5) \quad \tau_{(m,t),r}^{\theta\square,L} = \inf \{u \geq \tau_{(m,t),r-1}^{\theta\square,L} : v_{r+1}^{\theta\square}(u) = 0\} \text{ and}$$

$$(5.6) \quad \tau_{(m,t),r}^{\theta\square,R} = \sup \{u \geq \tau_{(m,t),r-1}^{\theta\square,R} : h_r^{\theta\square}(\tau_{(m,t),r-1}^{\theta\square,R}, u) = B_r(\tau_{(m,t),r-1}^{\theta\square,R}, u)\}$$

More specifically, if $u \geq \tau_{(m,t),r-1}^{\theta\square,R}$, then $h_r^{\theta\square}(\tau_{(m,t),r-1}^{\theta\square,R}, u) = B_r(\tau_{(m,t),r-1}^{\theta\square,R}, u)$ if and only if $u \leq \tau_{(m,t),r}^{\theta\square,R}$.

Proof. By Theorem 3.2(vi),

$$\begin{aligned} 0 &\leq v_{r+1}^{\theta\square}(\tau_r) = \sup_{\tau_r \leq s < \infty} \{B_r(\tau_r, s) - h_{r+1}^{\theta\square}(\tau_r, s)\} \\ &= \sup_{\tau_r \leq s < \infty} \{B_r(s) - h_{r+1}^{\theta\square}(s)\} - [B_r(\tau_r) - h_{r+1}^{\theta\square}(\tau_r)] \\ &\leq \sup_{\tau_{r-1} \leq s < \infty} \{B_r(s) - h_{r+1}^{\theta\square}(s)\} - [B_r(\tau_r) - h_{r+1}^{\theta\square}(\tau_r)] = 0. \end{aligned}$$

To establish (5.5), assume, by way of contradiction, that for some $\tau_{(m,t),r-1}^{\theta\square,L} \leq u < \tau_{(m,t),r}^{\theta\square,L}$,

$$0 = v_{r+1}^{\theta\square}(u) = \sup_{u \leq s < \infty} \{B_r(u, s) - h_{r+1}^{\theta\square}(u, s)\}.$$

Then,

$$B_m(u) - h_{m+1}^{\theta\Box}(u) \geq B_m(s) - h_{m+1}^{\theta\Box}(s)$$

for all $s \geq u$, and specifically for $s = \tau_{(m,t),r}^{\theta\Box,L}$. This contradicts the definition of $\tau_{(m,t),r}^{\theta\Box,L}$ as the left-most maximizer of $B_r(s) - h_{r+1}^{\theta\Box}(s)$ over $s \in [\tau_{(m,t),r-1}^{\theta\Box,L}, \infty)$.

By Theorem 3.2(vi),

$$(5.7) \quad h_r^{\theta\Box}(u, v) = B_r(u, v) + \sup_{u \leq s < \infty} \{B_r(s) - h_{r+1}^{\theta\Box}(s)\} - \sup_{v \leq s < \infty} \{B_r(s) - h_{r+1}^{\theta\Box}(s)\},$$

so, since τ_r maximizes $B_r(s) - h_{r+1}^{\theta\Box}(s)$ on $[\tau_{r-1}, \infty)$, the two supremum terms above are both equal to $B_r(\tau_r) - h_{r+1}^{\theta\Box}(\tau_r)$ whenever $u, v \in [\tau_{r-1}, \tau_r]$. Therefore, $h_r^{\theta\Box}(u, v) = B_r(u, v)$. To establish (5.6), if, for some $r \geq m$, (5.6) fails, then by (5.7), for some $u > \tau_{(m,t),r}^{\theta\Box,R}$,

$$\sup_{\tau_{(m,t),r-1}^{\theta\Box,R} \leq s < \infty} \{B_r(s) - h_{r+1}^{\theta\Box}(s)\} = \sup_{u \leq s < \infty} \{B_r(s) - h_{r+1}^{\theta\Box}(s)\}.$$

Thus, all maximizers of $B_r(s) - h_{r+1}^{\theta\Box}(s)$ over $s \in [u, \infty)$ are also maximizers over the larger set $[\tau_{(m,t),r-1}^{\theta\Box,R}, \infty)$. But this is a contradiction because $\tau_{(m,t),r}^{\theta\Box,R}$ is the right-most maximizer over $s \in [\tau_{(m,t),r-1}^{\theta\Box,R}, \infty)$. \square

Recall, by Remark 3.7, that on the event $\Omega^{(\theta)} \subseteq \Omega_1$, $\mathbf{T}_\mathbf{x}^{\theta+} = \mathbf{T}_\mathbf{x}^{\theta-} = \mathbf{T}_\mathbf{x}^\theta$ for all $\mathbf{x} \in \mathbb{Z} \times \mathbb{R}$.

Lemma 5.3. *Fix $\theta > 0$ and $\mathbf{x} = (m, t) \in \mathbb{Z} \times \mathbb{R}$. Then there exists an event $\tilde{\Omega}_\mathbf{x}^{(\theta)} \subseteq \Omega^{(\theta)}$, of probability one, on which the set $\mathbf{T}_\mathbf{x}^\theta$ contains exactly one sequence $\{\tau_r\}_{r \geq m-1}$. This sequence satisfies*

$$\lim_{n \rightarrow \infty} \frac{\tau_n}{n} = \theta.$$

To prove this lemma, we need some machinery from [ARAS20]. For two fields of Brownian motions \mathbf{B} and $\bar{\mathbf{B}}$ and a subset $A \subset \mathbb{R}$, define

$$(5.8) \quad \bar{L}_{(m,s),n}^\lambda(\mathbf{B}, \bar{\mathbf{B}}; s_n \in A) = \sup \left\{ \sum_{r=m}^n B_r(s_{r-1}, s_r) - \bar{B}_{n+1}(s_n) - \lambda s_n : \mathbf{s}_{m,n} \in \Pi_{(m,s),n}, s_n \in A \right\}.$$

The only difference between this definition and (2.3) is the restriction on the s_n .

Lemma 5.4 ([ARAS20], page 1949). *Let \mathbf{B} be a field of independent, two-sided Brownian motions and $\bar{\mathbf{B}}$ a field of Brownian motions (not necessarily independent). Fix $s \in \mathbb{R}$, $0 \leq S \leq T \leq \infty$, and $m \in \mathbb{Z}$. Then, with probability one,*

$$\lim_{n \rightarrow \infty} n^{-1} \bar{L}_{(m,s),n}^\lambda(\mathbf{B}, \bar{\mathbf{B}}; s + nS \leq s_n \leq s + nT) = \sup_{S \leq t \leq T} \{2\sqrt{t} - \lambda t\}.$$

Lemma 5.5 ([ARAS20], Lemma 4.7). *Let \mathbf{B} and $\bar{\mathbf{B}}$ satisfy the same conditions of Lemma 5.4. Fix $\lambda > 0$, $s \in \mathbb{R}$, and $m \in \mathbb{Z}$. If $0 < \theta < \lambda^{-2}$, then with probability one, for all sufficiently large n ,*

$$\bar{L}_{(m,s),n}^\lambda(\mathbf{B}, \bar{\mathbf{B}}; s_n \leq s + n\theta) < \bar{L}_{(m,s),n}^\lambda(\mathbf{B}, \bar{\mathbf{B}}),$$

If $\theta > \lambda^{-2}$, then with probability one, for all sufficiently large n ,

$$\bar{L}_{(m,s),n}^\lambda(\mathbf{B}, \bar{\mathbf{B}}; s_n \geq s + n\theta) < \bar{L}_{(m,s),n}^\lambda(\mathbf{B}, \bar{\mathbf{B}}).$$

Lemma 5.5 is slightly stronger than the result stated in [ARAS20], so we include a proof.

Proof. We prove the first statement, and the second follows by the same reasoning. The unique maximum of $2\sqrt{t} - \lambda t$ for $t \in [0, \infty)$ is achieved at $t = \lambda^{-2}$. Since $\theta < \lambda^{-2}$, Lemma 5.4 implies that

$$\lim_{n \rightarrow \infty} n^{-1} \bar{L}_{(m,s),n}^\lambda(\mathbf{B}, \bar{\mathbf{B}}; s_n \leq s + n\theta) = \sup_{0 \leq t \leq \theta} \{2\sqrt{t} - \lambda t\} < \sup_{0 \leq t < \infty} \{2\sqrt{t} - \lambda t\} = \lim_{n \rightarrow \infty} n^{-1} \bar{L}_{(m,s),n}^\lambda(\mathbf{B}, \bar{\mathbf{B}}).$$

Hence, for sufficiently large n ,

$$\bar{L}_{(m,s),n}^\lambda(\mathbf{B}, \bar{\mathbf{B}}; s_n \leq s + n\theta) < \bar{L}_{(m,s),n}^\lambda(\mathbf{B}, \bar{\mathbf{B}}). \quad \square$$

Proof of Lemma 5.3. By Lemma 5.1, for a sequence $\{\tau_r\}_{r \geq m-1} \in \mathbf{T}_{(m,t)}^\theta$, the jump times $t = \tau_{m-1} \leq \dots \leq \tau_n$ are a maximizing sequence for the point-to-line last passage time (5.1). By Theorem 3.2(vii), h_{n+1}^θ and $\{B_r\}_{r \leq n}$ are independent for each n . This allows us to apply the almost sure uniqueness of point-to-line last-passage maximizers in Lemma 2.3. Let $\tilde{\Omega}_\mathbf{x}^{(\theta)}$ be the event on which these maximizers are unique and on which

$$\lim_{n \rightarrow \infty} \frac{\tau_n}{n} = \theta.$$

for the almost surely unique sequence $\{\tau_r\}_{r \geq m-1} \in \mathbf{T}_{(m,t)}^\theta$. It remains to show that

$$\mathbb{P}\left(\lim_{n \rightarrow \infty} \frac{\tau_n}{n} = \theta\right) = 1.$$

For $\gamma < \theta = \lambda^{-2}$, Lemma 5.5 guarantees that, with probability one, for all sufficiently large n ,

$$\bar{L}_{(m,t),n}^\lambda(\mathbf{B}, \bar{\mathbf{B}}; s_n \leq s + n\gamma) < \bar{L}_{(m,t),n}^\lambda(\mathbf{B}, \bar{\mathbf{B}}).$$

Therefore, by Lemma 5.1, $\tau_n > s + n\gamma$ for all sufficiently large n . Thus, for $\gamma < \theta$,

$$\mathbb{P}\left(\liminf_{n \rightarrow \infty} \frac{\tau_n}{n} \geq \gamma\right) = 1, \quad \text{so by taking } \gamma \nearrow \theta, \quad \mathbb{P}\left(\liminf_{n \rightarrow \infty} \frac{\tau_n}{n} \geq \theta\right) = 1.$$

Using the second statement of Lemma 5.5, a symmetric argument shows that

$$\mathbb{P}\left(\limsup_{n \rightarrow \infty} \frac{\tau_n}{n} \leq \theta\right) = 1. \quad \square$$

5.2. Proofs of Theorems 3.8-3.12. We now begin to prove the theorems of Section 3.

Proof of Theorem 3.8. Part (i): On the event Ω_1 , let $(m, t) \in \mathbb{Z} \times \mathbb{R}$ $\theta > 0$ and $\square \in \{+, -\}$, and let $t = \tau_{m-1} \leq \tau_m \leq \dots$ be a sequence in $\mathbf{T}_{(m,t)}^{\theta\square}$. Let Γ be the associated path. By Lemma 5.2,

$$(5.9) \quad v_{r+1}^{\theta\square}(\tau_r) = 0, \quad \text{and} \quad h_r^{\theta\square}(s, t) = B_r(s, t) \text{ for } \tau_{r-1} \leq s \leq t \leq \tau_r.$$

We take $\mathbf{x} = \mathbf{y} = (m, t)$ and $\mathbf{z} = (n, \tau_n)$. The case for general $\mathbf{y} \leq \mathbf{z}$ along the path Γ follows by the same argument. By (5.9) and additivity of Busemann functions (Theorem 3.2(i)), the energy of path Γ between \mathbf{x} and \mathbf{y} is given by

$$\sum_{r=m}^n (B_r(\tau_{r-1}, \tau_r)) = \sum_{r=m}^{n-1} (h_r^{\theta\square}(\tau_{r-1}, \tau_r) + v_{r+1}^{\theta\square}(\tau_r)) + h_n^{\theta\square}(\tau_{n-1}, \tau_n) = \mathcal{B}^{\theta\square}((m, t), (n, \tau_n)).$$

Let jump times $t = s_{m-1} \leq s_m \leq s_{m+1} \leq \dots \leq s_n = \tau_n$ define any other path between (m, t) and (n, τ_n) . Then, by Theorem 3.2, Parts (i) and (ii), the energy of this path is

$$(5.10) \quad \sum_{r=m}^n B_r(s_{r-1}, s_r) \leq \sum_{r=m}^{n-1} (h_r^{\theta\square}(s_{r-1}, s_r) + v_{r+1}^{\theta\square}(s_r)) + h_n^{\theta\square}(s_{n-1}, \tau_n) = \mathcal{B}^{\theta\square}((m, t), (n, \tau_n)).$$

Part (ii): By Theorem 3.2(ii), equality holds in Equation (5.10) only if $B_r(s_{r-1}, s_r) = h_r^{\theta\square}(s_{r-1}, s_r)$ for $m \leq r \leq n$ and $v_{r+1}^{\theta\square}(s_r) = 0$ for $m \leq r \leq n-1$. Then, the statement follows by the Equations (5.5) and (5.6) of Lemma 5.2.

Part (iii): The key is Theorem 3.2(ii). We show that, for $r \geq m$,

$$\tau_r^- := \tau_{(m,t),r}^{\theta-,L} \leq \tau_{(m,t),r}^{\theta+,L} =: \tau_r^+,$$

and all other statements on the first line of Part (iii) follow by the same reasoning. By definition, $\tau_{m-1}^- = \tau_{m-1}^+ = t$. Inductively, assume that $\tau_r^- \leq \tau_r^+$ for some $r \geq m$. We use the notation $L \arg \sup$ to denote leftmost maximizer. Then,

$$\begin{aligned} \tau_{r+1}^- &= L \arg \sup_{\tau_r^- \leq s < \infty} \{B_r(s) - h_{r+1}^{\theta-,L}(s)\} \leq L \arg \sup_{\tau_r^+ \leq s < \infty} \{B_r(s) - h_{r+1}^{\theta-,L}(s)\} \\ &\leq L \arg \sup_{\tau_r^+ \leq s < \infty} \{B_r(s) - h_{r+1}^{\theta+,L}(s)\} = \tau_{r+1}^+. \end{aligned}$$

The first inequality above holds because $\tau_r^- \leq \tau_r^+$, and the second inequality is an application of Lemma A.1, using the fact that $h_m^{\theta-}(s, t) \leq h_m^{\theta+}(s, t)$ for all $s \leq t$ by Theorem 3.2(ii). For the second line of Part (iii), let $s < t$. We show that

$$\tau_{s,r} := \tau_{(m,s),r}^{\theta\square,L} \leq \tau_{(m,t),r}^{\theta\square,L} =: \tau_{t,r},$$

and the statement with “ L ” replaced by “ R ” has an analogous proof. Again, the base case of $r = m-1$ follows by definition. Assume the inequality holds for some $r \geq m-1$. Then,

$$\tau_{s,r+1} = L \arg \sup_{\tau_{s,r} \leq u < \infty} \{B_m(u) - h_{m+1}^{\theta\square}(u)\} \leq L \arg \sup_{\tau_{t,r} \leq u < \infty} \{B_m(u) - h_{m+1}^{\theta\square}(u)\} = \tau_{t,r+1}.$$

Part (iv): The monotonicity of Part (iii) ensures that the limits exist. Additionally, the first limit satisfies

$$(5.11) \quad \tau_r := \lim_{\gamma \nearrow \theta} \tau_{(m,t),r}^{\gamma\square,L} \leq \tau_{(m,t),r}^{\theta-,L}.$$

We prove equality in the above expression, and the other statements follow analogously. By Lemma 5.1, for any $n \geq m$, $t = \tau_{(m,t),m-1}^{\gamma \square, L} \leq \dots \leq \tau_{(m,t),n}^{\gamma \square, L}$ is a maximizing sequence for

$$\overline{L}_{(m,t),n}^{\frac{1}{\sqrt{\gamma}}}(\mathbf{B}, \overline{\mathbf{B}}) := \sup \left\{ \sum_{r=m}^n B_r(s_{r-1}, s_r) - h_{n+1}^{\square}(s_n) : \mathbf{s}_{m,n} \in \Pi_{(m,t),n} \right\}.$$

Using the monotonicity of Part (iii) again, for all $\gamma \leq \theta$, the supremum may be restricted to the compact subset of $\Pi_{(m,t),n}$ such that $s_n \leq \tau_{(m,t),n}^{(\theta+1)^-, L}$. Then, by Theorem 3.2(iii) and Lemma A.2, $t = \tau_{m-1} \leq \dots \leq \tau_n$ is a maximizing sequence for

$$\overline{L}_{(m,t),n}^{\frac{1}{\sqrt{\theta}}}(\mathbf{B}, \overline{\mathbf{B}}) := \sup \left\{ \sum_{r=m}^n B_r(s_{r-1}, s_r) - h_{n+1}^{\theta-}(s_n) : \mathbf{s}_{m,n} \in \Pi_{(m,t),n} \right\}.$$

Thus the inequality (5.11) must be an equality since the $\tau_{(m,t),r}^{\theta-, L}$ are the left-most maximizers. The statement for limits as $\delta \searrow \theta$ follow by the same reasoning. To prove the statement for limits as $s \nearrow t$ and $u \searrow t$, simply replace the use of Lemma A.2 with Lemma A.3.

Part (v): Recall the events of probability one, $\tilde{\Omega}_{\mathbf{y}}^{(\theta)} \subseteq \Omega_1$ from Lemma 5.3, and for the countable dense set $D \subseteq (0, \infty)$ of Section 4, set

$$\Omega_2 := \bigcap_{\gamma \in D, \mathbf{y} \in \mathbb{Z} \times \mathbb{Q}} \tilde{\Omega}_{\mathbf{y}}^{(\gamma)}$$

Then, $\mathbb{P}(\Omega_2) = 1$ and $\Omega_2 \subseteq \Omega_1$. Let $\omega \in \Omega_2$, $\theta > 0$, $(m, t) \in \mathbb{Z} \times \mathbb{Q}$, $(\tau_r)_{r \geq m-1} \in \mathbf{T}_{(m,t)}^{\theta \square}$, $\varepsilon > 0$, and let $\gamma, \delta \in D$ be such that $\theta - \varepsilon < \gamma < \theta < \delta < \theta + \varepsilon$. Part (iii) and Lemma 5.3 imply that

$$\limsup_{n \rightarrow \infty} \frac{\tau_n}{n} \leq \limsup_{n \rightarrow \infty} \frac{\tau_{(m,t),n}^{\theta+, R}}{n} \leq \limsup_{n \rightarrow \infty} \frac{\tau_{(m,t),n}^{\delta+, R}}{n} = \delta < \theta + \varepsilon.$$

By a similar argument, on Ω_2 ,

$$\liminf_{n \rightarrow \infty} \frac{\tau_n}{n} \geq \theta - \varepsilon.$$

An analogous application of the second line of Part (iii) extends Part (v) to all $\mathbf{x} \in \mathbb{Z} \times \mathbb{R}$, on the event Ω_2 . \square

Proof of Theorem 3.10. Part (i): Let $\omega \in \Omega_2$, $\theta > 0$, $(m, t) \in \mathbb{Z} \times \mathbb{R}$, and $\{t_{n,r}\}_{n \geq m, m-1 \leq r \leq n}$ satisfy the given assumptions. Let $0 < \gamma < \theta < \delta$ be arbitrary. Then, by assumption that $t_n/n \rightarrow \theta$ and Part (v), there exists $N \in \mathbb{Z}$ such that for all $n \geq N$,

$$(5.12) \quad \tau_{(m,t),n}^{\gamma-, L} < t_n < \tau_{(m,t),n}^{\delta+, R}.$$

By Part (ii), the sequence $t = \tau_{(m,t),m-1}^{\gamma-, L} \leq \tau_{(m,t),m}^{\gamma-, L} \leq \dots \leq \tau_{(m,t),n}^{\gamma-, L}$ defines the left-most geodesic between (m, t) and $(n, \tau_{(m,t),n}^{\gamma-, L})$, and $t = \tau_{(m,t),m-1}^{\delta+, R} \leq \tau_{(m,t),m}^{\delta+, R} \leq \dots \leq \tau_{(m,t),n}^{\delta+, R}$ defines the right-most geodesic between (m, t) and $(n, \tau_{(m,t),n}^{\delta+, R})$. Let $t = t_{m-1}^L \leq t_m^L \leq \dots \leq t_n^L = t_n$ and $t = t_{m-1}^R \leq t_m^R \leq \dots \leq t_n^R = t_n$ define the left-most (resp. right-most) geodesic between the points (m, t) and (n, t_n) . Then, by (5.12) and Lemma 2.2, for all $n \geq N$ and $m \leq r \leq n$,

$$\tau_{(m,t),r}^{\gamma-, L} \leq t_r^L \leq t_{n,r} \leq t_r^R \leq \tau_{(m,t),r}^{\delta+, R}.$$

Taking limits as $n \rightarrow \infty$ produces

$$\tau_{(m,t),n}^{\gamma-, L} \leq \liminf_{n \rightarrow \infty} t_{n,r} \leq \limsup_{n \rightarrow \infty} t_{n,r} \leq \tau_{(m,t),r}^{\delta+, R}.$$

Taking limits as $\gamma \nearrow \theta$ and $\delta \searrow \theta$ and using Part (iv) completes the proof of Part (i).

Part (ii): This is an immediate consequence of Part (i), setting $t_{n,r} = t_r$ for any θ -directed semi-infinite geodesic defined by the sequence $\{t_r\}_{r \geq m-1}$.

Parts (iii)-(iv): Using the uniqueness of Lemma 5.3, define $\Omega_{\mathbf{x}}^{(\theta)} = \tilde{\Omega}_{\mathbf{x}}^{(\theta)} \cap \Omega_2$, and the desired conclusions follow from Parts (i) and (ii). \square

Proof of Theorem 3.11. Part (i): Assume to the contrary, that, for some $\omega \in \Omega_1$, there exists a point $(m, t) \in \mathbb{Z} \times \mathbb{R}$ and a sequence $t = t_{m-1} \leq t_m \leq t_{m+1} \leq \dots$ defining a semi-infinite geodesic Γ , starting from (m, t) and satisfying

$$0 \leq \underline{\theta} := \liminf_{n \rightarrow \infty} \frac{t_n}{n} < \limsup_{n \rightarrow \infty} \frac{t_n}{n} =: \overline{\theta} \leq \infty.$$




FIGURE 5.1. Crossing of geodesic paths

Choose some $\theta \in (\underline{\theta}, \bar{\theta})$. By Theorem 3.8(v),

$$\lim_{n \rightarrow \infty} \frac{\tau_{(m,t),n}^{\theta-,L}}{n} = \theta$$

Therefore, there are infinitely many values of n with $t_n < \tau_{(m,t),n}^{\theta-,L}$ and another infinitely many values of n with $t_n > \tau_{(m,t),n}^{\theta-,L}$. Let n_1 be such that $t_{n_1} < \tau_{(m,t),n_1}^{\theta-,L}$ and let $n_2 > n_1$ be such that $t_{n_2} > \tau_{(m,t),n_2}^{\theta-,L}$. At the point $(n_1 + 1, t_{n_1})$, the path Γ lies strictly to the left of the leftmost semi-infinite geodesic in $\mathbf{T}_{(m,t)}^{\theta-}$. However, at the point (n_2, t_{n_2}) , the path Γ lies strictly to the right of this left-most semi-infinite geodesic. Thus, there must be points $\mathbf{x} \leq \mathbf{y} \in \mathbb{Z} \times \mathbb{R}$, such that the two geodesics intersect at both points and such that the portion of Γ between \mathbf{x} and \mathbf{y} is a geodesic between the two points, lying strictly to the left of the leftmost geodesic in $\mathbf{T}_{(m,t)}^{\theta-}$. See Figure 5.1 for clarity. This contradicts Theorem 3.8(ii).

Part (ii): Next, assume that for some $\omega \in \Omega_2$, there exists $(m, t) \in \mathbb{Z} \times \mathbb{R}$, and a sequence $t = t_{m-1} \leq t_m \leq \dots$ that defines a semi-infinite geodesic Γ , starting from (m, t) and satisfying

$$\lim_{n \rightarrow \infty} \frac{t_n}{n} = 0.$$

To show that $t_r = t$ for all $r \geq m$, it is sufficient to show that $t_m = t$. For then, this semi-infinite geodesic travels vertically to $(m + 1, t)$, and the remaining part of the geodesic is a geodesic starting at $(m + 1, t)$, so the result follows by induction. By Theorem 3.8(v), for every $\theta > 0$ and any $(m, t) \in \mathbb{Z} \times \mathbb{R}$,

$$\lim_{n \rightarrow \infty} \frac{\tau_{(m,t),n}^{\theta+,R}}{n} = \theta.$$

Hence, for all $\theta > 0$ and all sufficiently large n , $t_n < \tau_{(m,t),n}^{\theta+,R}$. However, by an analogous argument as in Part (i), again using Theorem 2.5(ii), $t_r \leq \tau_{(m,t),r}^{\theta+,R}$ for all $r \geq m$. Specifically, the inequality holds for $r = m$. By monotonicity of Theorem 3.8(iii), $\lim_{\theta \searrow 0} \tau_{(m,t),m}^{\theta+,R}$ exists. By definition of the event Ω_1 from Lemma 4.4, this limit equals t on the event $\Omega_1 \supseteq \Omega_2$. The case where $\lim_{n \rightarrow \infty} \frac{t_n}{n} = \infty$ is handled similarly.

Part (iii): By the monotonicity of Theorem 3.8(iii), it suffices to show that

$$\lim_{\theta \searrow 0} \tau_{(m,t),r}^{\theta+,R} = t \quad \text{and} \quad \lim_{\theta \rightarrow \infty} \tau_{(m,t),r}^{\theta-,L} = \infty \quad \text{for } r \geq m.$$

The proof of Part (ii), established the statement for $r = m$. For the limits as $\theta \rightarrow \infty$, this implies the statement holds for all $r \geq m$ since $\tau_{(m,t),m}^{\theta-,L} \leq \tau_{(m,t),r}^{\theta-,L}$ for all $r \geq m$. By the monotonicity of Theorem 3.8(iii), the limit

$$\tau_{(m,t),r}^0 := \lim_{\theta \searrow 0} \tau_{(m,t),r}^{\theta+,R}$$

exists, and satisfies $\tau_{(m,t),r}^0 \leq \tau_{(m,t),r}^{\theta+,R}$ for all $\theta > 0$. Furthermore, since $\{\tau_{(m,t),r}^{\theta+,R}\}_{r \geq m-1}$ is a nondecreasing sequence for each $\theta > 0$, the sequence $\{\tau_{(m,t),r}^0\}_{r \geq m-1}$ is also nondecreasing. By Theorem 3.8(v),

$$0 \leq \limsup_{n \rightarrow \infty} \frac{\tau_{(m,t),n}^0}{n} \leq \lim_{n \rightarrow \infty} \frac{\tau_{(m,t),n}^{\theta+,R}}{n} = \theta \text{ for all } \theta > 0 \implies \lim_{n \rightarrow \infty} \frac{\tau_{(m,t),n}^0}{n} = 0.$$

Therefore, by Part (ii), if the sequence of jump times $t = \tau_{(m,t),m-1}^{0,R} \leq \tau_{(m,t),m}^{0,R} \leq \dots$ defines a semi-infinite geodesic starting from (m, t) , the desired conclusion follows. It is sufficient to show that, for any $n \geq m$, the sequence $t = \tau_{(m,t),m-1}^0 \leq \tau_{(m,t),m}^0 \leq \dots \leq \tau_{(m,t),n}^0$ defines jump times for a finite geodesic between

(m, t) and $(n, \tau_{(m,t),n}^0)$. By Theorem 3.8(i), For each $\theta > 0$, the sequence $t = \tau_{(m,t),m-1}^{\theta+,R} \leq \dots \leq \tau_{(m,t),n}^{\theta+,R}$ is a maximizing sequence for

$$L_{(m,t),(n,\tau_{(m,t),n}^{\theta+,R})} = \sup \left\{ \sum_{r=m}^n B_r(s_r, s_{r-1}) : t = s_{m-1} \leq s_m \leq \dots \leq s_n = \tau_{(m,t),n}^{\theta+,R} \right\},$$

so since $\lim_{\theta \searrow 0} \tau_{(m,t),n}^{\theta+,R} = \tau_{(m,t),n}^0$, Lemma A.3 completes the proof. \square

Proof of Theorem 3.12. By Theorem 3.2, Parts (vii) and Theorem 3.4(i), for each $r \in \mathbb{Z}$, $s \mapsto B_r(s) - h_{r+1}^\theta(s)$ is a scaled, two-sided Brownian motion with strictly negative drift. Then, by Theorem A.10, there exists an event, $\text{CM}_r^{(\theta)}$, of probability one, on which, the set of $t \in \mathbb{R}$ such that $B_r(s) - h_{r+1}^\theta(s)$ for $s \in [t, \infty)$ has a non-unique maximum at $s = t$ is countably infinite. Then, for $\theta > 0$, set

$$(5.13) \quad \tilde{\Omega}^{(\theta)} := \bigcap_{\mathbf{x} \in \mathbb{Z} \times \mathbb{Q}} \Omega_{\mathbf{x}}^{(\theta)} \cap \bigcap_{r \in \mathbb{Z}} \text{CM}_r^{(\theta)},$$

where $\Omega_{\mathbf{x}}^{(\theta)}$ are the events of Theorem 3.10(iii). By construction, $\tilde{\Omega}^{(\theta)} \subseteq \Omega^{(\theta)} \cap \Omega_2$ and $\mathbb{P}(\tilde{\Omega}^{(\theta)}) = 1$. Since θ is fixed, we drop the \pm distinction for θ in the exponent for the remainder of the proof.

Parts (i)-(ii): By Theorem 3.10 (ii), on Ω_2 , there exist multiple θ -directed geodesics starting from (m, t) if and only if the set $\mathbf{T}_{(m,t)}^\theta$ contains more than one element (Recall that $\mathbf{T}_{(m,t)}^\theta$ was defined as $\mathbf{T}_{(m,t)}^{\theta-} \cup \mathbf{T}_{(m,t)}^{\theta+}$). If $\omega \in \tilde{\Omega}^{(\theta)} \subseteq \Omega_2$ and (m, t) is a point whose θ -directed geodesic is not unique, then at most one such semi-infinite geodesic can pass through $(m, t + \varepsilon)$ for some $\varepsilon > 0$. Otherwise, two different geodesics would pass through (m, q) for some $q \in \mathbb{Q}$, giving two θ -directed geodesics starting at (m, q) . Inductively, to get two different θ -directed geodesics starting from (m, t) , there must be some level $r \geq m$ such that all θ -directed geodesics pass through (r, t) and then one geodesic passes through $(r + 1, t)$, and the other passes through $(r, t + \varepsilon)$ for some $\varepsilon > 0$. Therefore, by Definition 3.6, on the event $\tilde{\Omega}^{(\theta)}$, there exists a point $(m, t) \in \mathbb{Z} \times \mathbb{R}$ whose semi-infinite geodesic in direction θ is not-unique if and only if there exists $r \geq m$ such that, for $m \leq k \leq r - 1$, $B_r(s) - h_{r+1}^\theta(s)$ over $s \in [t, \infty)$ has a unique maximum at $s = t$, and $B_r(s) - h_{r+1}^\theta(s)$ over $s \in [t, \infty)$ has a non-unique maximum at $s = t$. This proves the decomposition of Part (ii), and Part (i) follows from Theorem A.10(ii).

Part (iii): This follows from Theorem A.10(iv). \square

Proof of Corollary 3.14. Let $\omega \in \tilde{\Omega}^{(\theta)}$, $m \in \mathbb{Z}$, and $s < t \in \mathbb{R}$. We use a modified induction, in the following manner.

- (a) First, note by definition that $s = \tau_{(m,s),m-1}^{\theta,R} < \tau_{(m,t),m-1}^{\theta,L} = t$.
- (b) For each $r \geq m$, we assume that $\tau_{(m,s),r-1}^{\theta,R} < \tau_{(m,t),r-1}^{\theta,L}$.
- (c) Under assumption (b), we show that if $\tau_{(m,s),r}^{\theta,R} \geq \tau_{(m,t),r}^{\theta,L}$, then $\tau_{(m,s),k}^{\theta,R} = \tau_{(m,t),k}^{\theta,L}$ for all $k \geq r$.

With this procedure mapped out, assume that, for some $r \geq m$,

$$\tau_{(m,s),r-1}^{\theta,R} < \tau_{(m,t),r-1}^{\theta,L} \quad \text{and} \quad \tau_{(m,s),r}^{\theta,R} \geq \tau_{(m,t),r}^{\theta,L}.$$

The second inequality above follows from definition. By definition, $\tau_{(m,s),r}^{\theta,R}$ is a maximizer of $B_r(u) - h_{r+1}^\theta(u)$ over $u \in [\tau_{(m,s),r-1}^{\theta,R}, \infty)$. But since

$$\tau_{(m,s),r}^{\theta,R} \geq \tau_{(m,t),r}^{\theta,L} \geq \tau_{(m,t),r-1}^{\theta,L},$$

$\tau_{(m,s),r}^{\theta,R}$ is also a maximizer over $u \in [\tau_{(m,t),r-1}^{\theta,L}, \infty)$. By definition, $\tau_{(m,t),r}^{\theta,L}$ is another maximizer over this set, so

$$B_r(\tau_{(m,s),r}^{\theta,R}) - h_{r+1}^\theta(\tau_{(m,s),r}^{\theta,R}) = B_r(\tau_{(m,t),r}^{\theta,L}) - h_{r+1}^\theta(\tau_{(m,t),r}^{\theta,L}),$$

and both $\tau_{(m,s),r}^{\theta,R}$ and $\tau_{(m,t),r}^{\theta,L}$ are maximizers of $B_r(u) - h_{r+1}^\theta(u)$ over $u \in [q, \infty)$ for any rational $q \in [\tau_{(m,s),r-1}^{\theta,R}, \tau_{(m,t),r-1}^{\theta,L}]$. By assumption (b), such a rational q exists. Then, the sequence $\{\tau_{(m,s),k}^{\theta,R}\}_{k \geq r}$ and the sequence $\{\tau_{(m,t),k}^{\theta,L}\}_{k \geq r}$ both define jump times for a semi-infinite geodesic starting from (r, q) . By construction of $\Omega^{(\theta)}$, since $\omega \in \tilde{\Omega}^{(\theta)}$, there is a unique θ -directed semi-infinite geodesic starting from this point, and conclusion (c) holds. \square

5.3. The dual environment. The remainder of Section 5 develops results that culminate in the proof of Theorem 3.17. Along the way, we prove Lemma 3.15. *Throughout this section $\theta > 0$ is fixed* and we work on the full probability event $\Omega^{(\theta)}$ of Theorem 3.2. By Remark 3.7, on this event $\mathbf{T}_{\mathbf{x}}^\theta = \mathbf{T}_{\mathbf{x}}^{\theta+} = \mathbf{T}_{\mathbf{x}}^{\theta-}$ for all $\mathbf{x} \in \mathbb{Z} \times \mathbb{R}$. To prove Theorem 3.17, we use the dual field \mathbf{X}^θ of independent Brownian motions from Theorem 3.4 and their southwest-directed semi-infinite geodesics, giving an analogue of Pimentel's dual tree [Pim16] in the exponential CGM, and adapt the argument of [Sep20].

Lemma 5.6. *For each $\theta > 0$, there exists an event $\tilde{\Omega}^{(\theta,*)}$ of probability one, on which*

$$\lim_{n \rightarrow \infty} [L_{(-n, -t_n), \mathbf{x}}(\mathbf{X}^\theta) - L_{(-n, -t_n), \mathbf{y}}(\mathbf{X}^\theta)] = \mathcal{B}^\theta(\mathbf{y}, \mathbf{x})$$

for any $\mathbf{x}, \mathbf{y} \in \mathbb{Z} \times \mathbb{R}$ and for any sequence $\{t_n\}$ satisfying $\lim_{n \rightarrow \infty} \frac{t_n}{n} = \theta$. To clarify, $L_{\mathbf{x}, \mathbf{y}}(\mathbf{X}^\theta)$ is the last-passage process on the environment \mathbf{X}^θ while \mathcal{B}^θ is the original Busemann function of the environment \mathbf{B} as in Theorem 3.2.

Proof. Recall from Theorem 3.2(vi) that

$$h_{m-1}^\theta = D(h_m^\theta, B_{m-1}), \quad X_m^\theta = R(h_m^\theta, B_{m-1}), \quad \text{and} \quad v_m^\theta = Q(h_m^\theta, B_{m-1}).$$

Further recall (3.9), which states

$$(5.14) \quad h_m^\theta = \overleftarrow{D}(h_{m-1}^\theta, X_m^\theta), \quad B_{m-1} = \overleftarrow{R}(h_{m-1}^\theta, X_m^\theta), \quad \text{and} \quad v_m^\theta = \overleftarrow{Q}(h_{m-1}^\theta, X_m^\theta).$$

Recall that $\tilde{f}(t) = -f(-t)$. Apply Lemma A.18 to deduce that

$$(5.15) \quad \tilde{h}_{m-1}^\theta = \overleftarrow{D}(\tilde{h}_m^\theta, \tilde{B}_{m-1}), \quad \tilde{X}_m^\theta = \overleftarrow{R}(\tilde{h}_m^\theta, \tilde{B}_{m-1}), \quad \text{and} \quad -\tilde{v}_m^\theta = \overleftarrow{Q}(\tilde{h}_m^\theta, \tilde{B}_{m-1}).$$

From independence and matching marginals (Part Theorem 3.2(vii) and Theorem 3.4, Parts (i) and (ii)) follows

$$(5.16) \quad (\tilde{h}_k^\theta, \tilde{B}_{k-1}, \tilde{B}_{k-2}, \tilde{B}_{k-3}, \dots) \stackrel{d}{=} (h_{m-1}^\theta, X_m^\theta, X_{m+1}^\theta, X_{m+2}^\theta, \dots) \quad \forall k, m \in \mathbb{Z}.$$

Thus, iterating mappings (5.15) backward in the index m and mappings (5.14) forward in the index m gives this equality in distribution:

$$\{\tilde{h}_{-(m-1)}^\theta, -\tilde{v}_{-(m-1)}^\theta, \tilde{X}_{-(m-1)}^\theta, \tilde{B}_{-m}\}_{m \in \mathbb{Z}} \stackrel{d}{=} \{h_{m-1}^\theta, v_m^\theta, B_{m-1}, X_m^\theta\}_{m \in \mathbb{Z}}$$

Apply Theorem 3.2(viii) to the environment $\tilde{\mathbf{X}}_-^\theta = \{\tilde{X}_{-m}^\theta\}_{m \in \mathbb{Z}}$ in the process on the left above, to deduce that there exists a full probability event $\tilde{\Omega}^{(\theta,*)}$ on which, for any $t \in \mathbb{R}$ and $m \in \mathbb{Z}$,

$$\lim_{n \rightarrow \infty} L_{(-m, -t), (n, t_n)}(\tilde{\mathbf{X}}_-^\theta) - L_{(-m, 0), (n, t_n)}(\tilde{\mathbf{X}}_-^\theta) = -\tilde{h}_{-(-m)}^\theta(-t) = h_m^\theta(t) \quad \text{and}$$

$$\lim_{n \rightarrow \infty} L_{(-m, -t), (n, t_n)}(\tilde{\mathbf{X}}_-^\theta) - L_{(-m+1, -t), (n, t_n)}(\tilde{\mathbf{X}}_-^\theta) = -\tilde{v}_{-(-m)}^\theta(-t) = v_m^\theta(t).$$

Adding together horizontal and vertical steps for the general case, the proof is complete by noting that, for any $m, k \in \mathbb{Z}$ and $s, t \in \mathbb{R}$,

$$\lim_{n \rightarrow \infty} L_{(-m, -s), (n, t_n)}(\tilde{\mathbf{X}}_-^\theta) - L_{(-k, -t), (n, t_n)}(\tilde{\mathbf{X}}_-^\theta) = \lim_{n \rightarrow \infty} L_{(-n, -t_n), (m, s)}(\mathbf{X}^\theta) - L_{(-n, -t_n), (k, t)}(\mathbf{X}^\theta)$$

because

$$\begin{aligned} L_{(-m, -s), (n, t_n)}(\tilde{\mathbf{X}}_-^\theta) &= \sup \left\{ \sum_{r=-m}^n \tilde{X}_{-r}^\theta(s_{r-1}, s_r) : -s = s_{-m-1} \leq \dots \leq s_n = t_n \right\} \\ &= \sup \left\{ \sum_{r=-m}^n X_{-r}^\theta(-s_r, -s_{r-1}) : -s = s_{-m-1} \leq \dots \leq s_n = t_n \right\} \\ &= \sup \left\{ \sum_{r=-m}^n X_{-r}^\theta(-s_r, -s_{r-1}) : -t_n = -s_n \leq -s_{n-1} \leq \dots \leq -s_{m-1} = s \right\} \\ &= \sup \left\{ \sum_{r=-n}^m X_r^\theta(\tilde{s}_{r-1}, \tilde{s}_r) : -t_n = \tilde{s}_{-n-1} \leq \tilde{s}_{-n} \leq \dots \leq \tilde{s}_m = s \right\} = L_{(-n, -t_n), (m, s)}(\mathbf{X}^\theta). \end{aligned}$$

To get the second-to-last line above, simply set $\tilde{s}_k = -s_{-k-1}$. \square




FIGURE 5.2. Regular axis (solid) and the dual axis (dashed). Dual levels are labeled with a \star .

5.4. Dual geodesics. Since Brownian motion is distributionally symmetric about the origin, there exist θ -directed dual southwest semi-infinite geodesics for the environment \mathbf{X}^θ . These are constructed in a very similar manner as the northeast geodesics in Definition 3.6. Specifically, for $(m, t) \in \mathbb{Z} \times \mathbb{R}$, let $\mathbf{T}_{(m,t)}^{\theta,\star}$ be the set of sequences $t = \tau_m^\star \geq \tau_{m-1}^\star \geq \dots$ satisfying

$$(5.17) \quad h_{r-1}^\theta(\tau_{r-1}^\star) - X_r^\theta(\tau_{r-1}^\star) = \sup_{-\infty < s \leq \tau_r^\star} \{h_{r-1}^\theta(s) - X_r^\theta(s)\} \quad \text{for each } r \leq m.$$

Define the left-most and right-most sequences similarly, by $\tau_{(m,t),r}^{\theta,L^\star}$ and $\tau_{(m,t),r}^{\theta,R^\star}$. These sequences define southwest semi-infinite paths, similar as for the northeast paths. We graphically represent southwest paths on a dual plane, where the continuous coordinate is not changed, but the discrete coordinate is shifted down by $\frac{1}{2}$. That is, for $m \in \mathbb{Z}$, denote $m^\star = m - \frac{1}{2}$, and for $\mathbf{x} = (m, t) \in \mathbb{Z} \times \mathbb{R}$, denote $\mathbf{x}^\star = (m - \frac{1}{2}, t)$. Then, for $\{\tau_r^\star\}_{r \leq m} \in \mathbf{T}_{(m,t)}^{\theta,\star}$, the backwards path consists of horizontal and vertical line segments, where τ_r^\star denotes the position of the vertical segment connecting levels $(r+1)^\star$ and r^\star . Specifically, the paths consist of the points

$$\bigcup_{r=-\infty}^m \{(u, r^\star) : u \in [\tau_{r-1}^\star, \tau_r^\star]\} \cup \bigcup_{r=-\infty}^{m-1} \{(\tau_r^\star, v) : v \in [(r-1)^\star, r^\star]\}.$$

Note that in the above, we put the continuous coordinate first to match the graphical description in \mathbb{R}^2 . Figure 5.2 shows the regular axes and the dual axes together, with a southwest dual geodesic traveling on this dual plane. Analogous results as for the original northwest semi-infinite geodesics hold, demonstrated by the following theorem.

Theorem 5.7. *Fix $\theta > 0$. Then, for every $\mathbf{x} \in \mathbb{Z} \times \mathbb{R}$, every semi-infinite path in $\mathbf{T}_{\mathbf{x}^\star}^{\theta,\star}$ is a semi-infinite geodesic for Brownian last-passage percolation with environment \mathbf{X}^θ . Specifically, the following hold.*

- (i) *On the full probability event $\Omega^{(\theta)}$ of Theorem 3.2, let $\mathbf{x} \in \mathbb{Z} \times \mathbb{R}$, and let Γ^\star be any semi-infinite path in $\mathbf{T}_{\mathbf{x}^\star}^{\theta,\star}$. Then, for any $\mathbf{y} \leq \mathbf{z} \in \mathbb{Z} \times \mathbb{R}$ with $\mathbf{y}^\star, \mathbf{z}^\star$ lying along the semi-infinite path, the energy of the portion of that path between \mathbf{y}^\star and \mathbf{z}^\star , in the environment \mathbf{X}^θ , is*

$$L_{\mathbf{y},\mathbf{z}}(\mathbf{X}^\theta) = \mathcal{B}^\theta(\mathbf{y}, \mathbf{z}),$$

and this energy is maximal among all paths between \mathbf{y} and \mathbf{z} in the environment \mathbf{X}^θ . To be clear, \mathcal{B}^θ is the original Busemann function for the environment \mathbf{B} .

- (ii) *On $\Omega^{(\theta)}$, for all $(m, t) \in \mathbb{Z} \times \mathbb{R}$ and $r \leq m$,*

$$\tau_{(m,t),r-1}^{\theta,R^\star} = \sup\{u \leq \tau_{(m,t),r}^{\theta,R^\star} : v_r^\theta(u) = 0\} \text{ and}$$

$$\tau_{(m,t),r-1}^{\theta,L^\star} = \inf\{u \leq \tau_{(m,t),r}^{\theta,L^\star} : h_r^\theta(u, \tau_{(m,t),r}^{\theta,L^\star}) = X_r^\theta(u, \tau_{(m,t),r}^{\theta,L^\star})\}$$

More specifically, if $u \leq \tau_{(m,t),r}^{\theta,L^\star}$, then $h_r^\theta(u, \tau_{(m,t),r}^{\theta,L^\star}) = X_r^\theta(u, \tau_{(m,t),r}^{\theta,L^\star})$ if and only if $u \geq \tau_{(m,t),r-1}^{\theta,L^\star}$.

- (iii) *On $\Omega^{(\theta)}$, if, for some $\mathbf{z} \in \mathbb{Z} \times \mathbb{R}$, $\mathbf{x}^\star \geq \mathbf{y}^\star$ lie along the left-most semi-infinite geodesic in $\mathbf{T}_{\mathbf{z}^\star}^{\theta,\star}$, then the portion of the path between \mathbf{x}^\star and \mathbf{y}^\star , shifted back up by $\frac{1}{2}$ to lie on integer levels, is the left-most geodesic between \mathbf{x} and \mathbf{y} in the environment \mathbf{X}^θ . The analogous statement holds for the right-most geodesics.*
- (iv) *The following distributional equality holds.*

$$\{(\tau_{(m,t),r}^{\theta,R}, \tau_{(m,t),r}^{\theta,L}) : (m, t) \in \mathbb{Z}, r \geq m\}$$

$$\stackrel{d}{=} \{(-\tau_{(-m,-t),-(r+1)}^{\theta,L\star}, -\tau_{(-m,-t),-(r+1)}^{\theta,R\star}) : (m,t) \in \mathbb{Z}, r \geq m\}$$

(v) *There exists an event of full probability, $\Omega^{(\theta,\star)} \subseteq \Omega^{(\theta)}$, on which, for every $(m,t) \in \mathbb{Z} \times \mathbb{R}$ and every sequence $\{\tau_r^{\theta,\star}\}_{r \leq m} \in \mathbf{T}_{(m,t)}^{\theta,\star}$,*

$$\lim_{n \rightarrow \infty} \frac{\tau_{-n}^{\star}}{-n} = \theta.$$

Proof. Parts (i)-(iii): On $\Omega^{(\theta)}$, let $(m,t) \in \mathbb{Z} \times \mathbb{R}$, and $\{\tau_r^{\star}\}_{r \leq m} \in \mathbf{T}_{(m,t)}^{\theta,\star}$. By (5.14) and the definitions (3.4)–(3.6), for all $u, v \in \mathbb{R}$ and $r \in \mathbb{Z}$,

$$(5.18) \quad v_r^\theta(u) = \sup_{-\infty < s \leq u} \{X_r^\theta(s, u) - h_{r-1}^\theta(s, u)\},$$

$$(5.19) \quad \begin{aligned} h_r^\theta(u, v) &= h_{r-1}^\theta(u, v) + \sup_{-\infty < s \leq v} \{X_r^\theta(s, v) - h_{r-1}^\theta(s, v)\} - \sup_{-\infty < s \leq u} \{X_r^\theta(s, u) - h_{r-1}^\theta(s, u)\} \\ &= X_r^\theta(u, v) + \sup_{-\infty < s \leq v} \{h_{r-1}^\theta(s) - X_r^\theta(s)\} - \sup_{-\infty < s \leq u} \{h_{r-1}^\theta(s) - X_r^\theta(s)\}, \text{ and} \end{aligned}$$

$$(5.20) \quad \begin{aligned} B_{r-1}(u, v) &= X_r^\theta(u, v) + \sup_{-\infty < s \leq u} \{X_r^\theta(s, u) - h_{r-1}^\theta(s, u)\} - \sup_{-\infty < s \leq v} \{X_r^\theta(s, v) - h_{r-1}^\theta(s, v)\} \\ &= h_{r-1}^\theta(u, v) + \sup_{-\infty < s \leq u} \{h_{r-1}^\theta(s) - X_r^\theta(s)\} - \sup_{-\infty < s \leq v} \{h_{r-1}^\theta(s) - X_r^\theta(s)\}. \end{aligned}$$

As τ_{r-1}^{\star} is a maximizer of $h_{r-1}^\theta(s) - X_r^\theta(s)$ over $s \in (\infty, \tau_r^{\star}]$, from (5.18), it follows that $v_r^\theta(\tau_{r-1}^{\star}) = 0$ for each $r \leq m$. By (5.19), $h_r^\theta(u, v) = X_r^\theta(u, v)$ for $u, v \in [\tau_{r-1}^{\star}, \tau_r^{\star}]$. In general, $v_r^\theta(u) \geq 0$ and $h_r^\theta(u, v) \geq X_r^\theta(u, v)$ for $u \leq v$. Then, Parts (i)-(iii) follow just as for the analogous statements in the proofs of Theorem 3.8, Parts (i)-(ii) and Lemma 5.2.

Parts (iv)-(v): By (5.16), $\{\tilde{h}_{r-1}^\theta, \tilde{X}_r^\theta\}_{r \in \mathbb{Z}}$ has the same distribution as $\{h_{-(r-1)}^\theta, B_{-r}\}_{r \in \mathbb{Z}}$. Furthermore, the left-most (right-most) maximizers of $h_{r-1}^\theta(s) - X_r^\theta(s)$ for $s \in (-\infty, t]$ are the negative of the right-most (resp. left-most) maximizers of $\tilde{X}_r^\theta(s) - \tilde{h}_{r-1}^\theta(s)$ for $s \in [-t, \infty)$, establishing Part (iv). Part (v) then follows from Theorem 3.8(v). \square

Theorem 5.8. *On the event $\Omega^{(\theta)}$, the following hold for all $s, t \in \mathbb{R}, m \in \mathbb{Z}$.*

- (i) *If $\tau_{(m,s),m}^{\theta,R} < t$, then also $\tau_{(m,s),m}^{\theta,R} < \tau_{(m+1,t),m}^{\theta,L\star}$. See Figure 5.3 for clarity.*
- (ii) *If $t \leq \tau_{(m,s),m}^{\theta,R}$, then $\tau_{(m+1,t),m}^{\theta,L\star} \leq s$. See Figure 5.4 for clarity.*
- (iii) *If $\tau_{(m,s),m}^{\theta,L} \leq t$, then also $\tau_{(m,s),m}^{\theta,L} \leq \tau_{(m+1,t),m}^{\theta,R\star}$. See Figure 5.5 for clarity.*
- (iv) *If $t < \tau_{(m,s),m}^{\theta,L}$, then $\tau_{(m,+1,t),m}^{\theta,R\star} < s$. See Figure 5.6 for clarity.*

Remark 5.9. All of the statements of Theorem 5.8 only mention the first jump time for northeast geodesics and the first time of descent for southwest geodesics. However, for any $\mathbf{x} \in \mathbb{Z} \times \mathbb{R}$ and any point \mathbf{y} along the right-most (resp. left-most) semi-infinite geodesic in $\mathbf{T}_{\mathbf{x}}^\theta$, the right-most (resp. left-most) semi-infinite geodesic in $\mathbf{T}_{\mathbf{y}}^\theta$ agrees with the remainder of the original semi-infinite geodesic started from \mathbf{x} . Thus, the results of Theorem 5.8 can be extended by induction. For example, Part (i) implies that if a left-most dual southwest geodesic starts strictly to the right and below a right-most northeast geodesic, it remains to the right and below the northeast geodesic. See Figure 5.3.

Proof. Part (i): By Lemma 5.2, if $u \geq s$, then

$$(5.21) \quad B_m(s, u) = h_m^\theta(s, u) \text{ if and only if } u \leq \tau_{(m,s),m}^{\theta,R}.$$

By Theorem 5.7(ii), if $u \leq t$, then

$$(5.22) \quad X_{m+1}^\theta(u, t) = h_{m+1}^\theta(u, t) \text{ if and only if } u \geq \tau_{(m+1,t),m}^{\theta,L\star}.$$

By (5.19) and (5.20),

$$\begin{aligned} X_{m+1}^\theta(u, v) = h_{m+1}^\theta(u, v) &\iff B_m(u, v) = h_m^\theta(u, v) \\ &\iff \sup_{-\infty \leq w \leq u} \{h_m^\theta(w) - X_{m+1}^\theta(w)\} = \sup_{-\infty \leq w \leq v} \{h_m^\theta(w) - X_{m+1}^\theta(w)\}. \end{aligned}$$

This along with (5.22) implies that if $u \leq t$,

$$(5.23) \quad B_m^\theta(u, t) = h_m^\theta(u, t) \text{ if and only if } u \geq \tau_{(m+1,t),m}^{\theta,L^*}.$$

By assumption, $\tau_{(m,s),m}^{\theta,R} < t$. Equation (5.21) and the monotonicity of Theorem 3.2(ii) imply that

$$(5.24) \quad B_m(s, t) < h_m^\theta(s, t).$$

Assume, by way of contradiction, that $\tau_{(m+1,t),m}^{\theta,L^*} \leq \tau_{(m,s),m}^{\theta,R}$. Then, there exists $u \leq \tau_{(m,s),m}^{\theta,R}$ with $u \geq s \vee \tau_{(m,t),m}^{\theta,L^*}$. Then, for such u , by (5.21) and (5.23),

$$B_r(s, u) = h_r^\theta(s, u) \quad \text{and} \quad B_r^\theta(u, t) = h_r^\theta(u, t).$$

Adding these two equations gives us a contradiction to (5.24).

Part (ii): Let $t \leq \tau_{(m,s),m}^{\theta,R}$, and assume by way of contradiction, that $\tau_{(m+1,t),m}^{\theta,L^*} > s$. This also implies that $s < t$ since $t \geq \tau_{(m+1,t),m}^{\theta,L^*}$. By Equation (5.23), $B_m(s, t) < h_m^\theta(s, t)$. However, since $s < t \leq \tau_{(m,s),m}^{\theta,R}$, Equation (5.21), implies that $B_m(s, t) = h_m^\theta(s, t)$, giving the desired contradiction.

Part (iii): Assume that $\tau_{(m,s),m}^{\theta,L} \leq t$. By Theorem 5.7(ii),

$$\tau_{(m+1,t),m}^{\theta,R^*} = \sup\{u \leq t : v_{m+1}^\theta(u) = 0\}.$$

By Lemma 5.2, $v_{m+1}^\theta(\tau_{(m,s),m}^{\theta,L}) = 0$, so the desired conclusion follows.

Part (iv): Assume that $t < \tau_{(m,s),m}^{\theta,L}$. By Lemma 5.2,

$$\tau_{(m,s),m}^{\theta,L} = \inf\{u \geq s : v_{m+1}^\theta(u) = 0\}.$$

By assumption, $\tau_{(m+1,t),m}^{\theta,R^*} \leq t < \tau_{(m,s),m}^{\theta,L}$. By Theorem 5.7(ii), $v_{m+1}^\theta(\tau_{(m+1,t),m}^{\theta,R^*}) = 0$, so $\tau_{(m+1,t),m}^{\theta,R^*} < s$. \square




FIGURE 5.3. A right-most northeast geodesic (red/thin) lies strictly above and to the left of a left-most dual southwest geodesic (blue/thick).




FIGURE 5.4. A right-most northeast geodesic (red/thin) lies weakly to the right and below a left-most dual southwest geodesic (blue/thick).




FIGURE 5.5. A left-most northeast geodesic (red/thin) lies weakly above and to the left of a right-most dual southwest geodesic (blue/thick).




FIGURE 5.6. A left-most northeast geodesic (red/thin) lies strictly to the right and below a right-most dual southwest geodesic (blue/thick).

We use these non-intersection properties to prove the following theorem. Refer to Figure 5.7 for clarity.

Theorem 5.10. *Fix $\theta > 0$. Recall the event Ω_2 of Theorem 3.8(v) and the event $\Omega^{(\theta,*)}$ of Theorem 5.7(v). On the event $\Omega^{(\theta)} \cap \Omega^{(\theta,*)} \cap \Omega_2$, if for any $\mathbf{x}, \mathbf{y} \in \mathbb{Z} \times \mathbb{R}$, the right-most semi-infinite geodesics in $\mathbf{T}_{\mathbf{x}}^\theta$ and $\mathbf{T}_{\mathbf{y}}^\theta$ are disjoint (i.e. the paths share no points), then there exists a bi-infinite, up right path defined by jump times $\cdots \leq \tau_{-1}^* \leq \tau_0^* \leq \tau_1^* \leq \cdots$ (where τ_i^* denotes the jump time from level i to level $i + 1$) that satisfies the following:*

- (i) *For any point \mathbf{x}^* along the path, the portion of that path to the south and west of \mathbf{x}^* is the left-most semi-infinite geodesic in the set $\mathbf{T}_{\mathbf{x}^*}^{\theta,*}$. Specifically, when shifted back up by $\frac{1}{2}$, the path is a bi-infinite geodesic for the environment \mathbf{X}^θ .*
- (ii) *The sequence $\{\tau_n^*\}_{n \in \mathbb{Z}}$ satisfies*

$$\lim_{n \rightarrow \infty} \frac{\tau_n^*}{n} = \theta = \lim_{n \rightarrow \infty} \frac{\tau_{-n}^*}{-n}.$$

The analogous result holds if we assume that two left-most θ -directed semi-infinite geodesics are disjoint. In this case the portion of the path to the south and west of each of its points is the right-most dual geodesic.

Proof. We follow a similar compactness argument as in the proof of Lemma 4.6 in [Sep20], appropriately modified for the semi-discrete setting of BLPP. On $\Omega^{(\theta)} \cap \Omega^{(\theta,*)} \cap \Omega_2$, let $\mathbf{x} = (m, t)$, $\mathbf{y} = (k, t)$. Assume that Γ_1 and Γ_2 are disjoint, where Γ_1 denotes the path defined by the jump times $t = \tau_{(k,t),k-1}^{\theta,R} \leq \tau_{(k,t),k}^{\theta,R} \leq \cdots$ and Γ_2 denotes the path defined by the jump times $s = \tau_{(m,s),m-1}^{\theta,R} \leq \tau_{(m,s),m}^{\theta,R} \leq \cdots$. By Theorem 3.8(v), these paths satisfy

$$(5.25) \quad \lim_{n \rightarrow \infty} \frac{\tau_{(m,s),n}^{\theta,R}}{n} = \lim_{n \rightarrow \infty} \frac{\tau_{(k,t),n}^{\theta,R}}{n} = \theta,$$

so both the space and time coordinates of these up-right paths go to ∞ . Assume, without loss of generality, that Γ_2 lies above and to the left of Γ_1 . Then, by picking t_0 large enough, there exists k_0^1 and k_0^2 such that $(k_0^1, t_0) \in \Gamma_1$ and $(k_0^2, t_0) \in \Gamma_2$. For each t_0 , there are only finitely many choices of k_0^1 and k_0^2 , so assume that k_0^1 is the largest such choice and k_0^2 is the smallest such choice. Then, $k_0^1 < k_0^2$ by the assumption that Γ_2 lies above and to the left of Γ_1 . See Figure 5.8 for clarity. For $i = 1, 2, \dots$, define $t_i = t_0 + i$. For each time i , similarly define k_i^1 and k_i^2 so that $k_i^1 < k_i^2$ and (k_i^j, t_i) lies on Γ_j for $j = 1, 2$. Recall the notation $m^* = m - \frac{1}{2}$. Then, for $i = 0, 1, 2, \dots$, there exists $k_i \in \mathbb{Z}$ with $k_i^1 < k_i^* < k_i^2$. This gives us an infinite sequence $(k_i^*, t_i)_{i \geq 0}$ such that, for each i , the point (k_i^*, t_i) , lies between the paths Γ_1 and Γ_2 (see Figure 5.8). Starting from each of these points (k_i^*, t_i) , let Γ_i^* be the left-most dual geodesic path in $\mathbf{T}_{(k_i^*, t_i)}^{\theta,*}$. Each Γ_i^* is an infinite, down-left path satisfying the limit condition of Theorem 5.7(v), so for each $j < i$, each of the paths Γ_i^* intersects the vertical line $t = t_j$. Let $k_{i,i}^* = k_i^*$, and for $j < i$, let $k_{i,j}$ be the maximal integer such that $(k_{i,j}^*, t_j)$ lies on the path Γ_i^* . Since (k_i^*, t_i) lies between Γ_1 and Γ_2 , and Γ_1, Γ_2 are right-most semi-infinite geodesics constructed from the Busemann functions, Theorem 5.8(i)–(ii) implies that for $i \geq 0$, Γ_i^* lies strictly below and to the right of Γ_2 , and weakly above and to the left of Γ_1 (see also Remark 5.9).

Since $k_j^1 < k_{i,j}^* < k_j^2$ for $0 \leq j \leq i$, there are only finitely many values of $k_{i,j}$ for each value of j . Then, there exists a subsequence k_{i_r} such that for some $N_0 \in \mathbb{Z}$ with $k_0^1 < N_0 < k_0^2$ and all ℓ , $k_{i_r,0} = N_0$. Take a further subsequence $k_{i_{r_\ell}}$ such that for some $k_1^1 < N_1^* < k_1^2$ and every element of this subsequence, $k_{i_{r_\ell},1} = N_1$. Continuing in this way, there exists a sequence (N_i, t_i) of elements of $\mathbb{Z} \times \mathbb{R}$, that (N_i^*, t_i) lies between the paths Γ_1 and Γ_2 , and such that the left-most southwest dual semi-infinite geodesic starting from (N_i^*, t_i) passes through (N_j^*, t_j) for $0 \leq j \leq i$. Since we chose the left-most paths in $\mathbf{T}_{(N_i^*, t_i)}^{\theta,*}$ at each step, the paths are consistent at all space-time points. Hence, this construction gives a bi-infinite path such that, for any point along the path, the part of the path to the southwest of that point is a semi-infinite southwest dual

geodesic. To get the last part of the theorem about asymptotic direction of the paths, we already showed the direction to the southwest. The direction to the northeast follows by (5.25) because the path lies between Γ_1 and Γ_2 .

The proof for the disjoint left geodesics is analogous, replacing the use of left-most southwest dual geodesics with right-most southwest geodesics, and now, the bi-infinite path lies weakly below and to the right of Γ_2 and strictly above and to the left of Γ_1 . \square




FIGURE 5.7. The outcome of Theorem 5.10: two disjoint semi-infinite northeast paths (red/thin) and the dual bi-infinite path (blue/thick)




FIGURE 5.8. Constructing a backwards semi-infinite path from each discrete time point. The upper red/thin path is Γ_2 and the lower red/thin path is Γ_1 .

5.5. Proof of Lemma 3.15 and Theorem 3.17.

Proof of Lemma 3.15. We follow the procedure of the proof of Theorem 4.12 in the arXiv version of [Sep18]. We take $(m, t) = \mathbf{0}$, and the general case follows analogously. We simplify further: for any sequence $\{t_k\}_{k \in \mathbb{Z}}$ satisfying

$$\lim_{n \rightarrow \infty} \frac{t_n}{n} = \theta \quad \text{and} \quad \lim_{n \rightarrow \infty} \frac{t_{-n}}{-n} = \eta,$$

set $\tilde{t}_k = -t_{-(k+1)}$. Then, $\{\tilde{t}_k\}_{k \in \mathbb{Z}}$ is a sequence satisfying

$$\lim_{n \rightarrow \infty} \frac{\tilde{t}_n}{n} = \eta \quad \text{and} \quad \lim_{n \rightarrow \infty} \frac{t_{-n}}{-n} = \theta.$$

Furthermore, a geodesic path between $(-n, t_{-n})$ and (n, t_n) for the original field of Brownian motions $\mathbf{B} = \{B_r\}_{r \in \mathbb{Z}}$, when reflected through the origin, becomes a geodesic path between $(-n, \tilde{t}_{-n})$ and (n, \tilde{t}_n) for the field of Brownian motions $\{\tilde{B}_{-r}\}_{r \in \mathbb{Z}}$. Then, without loss of generality, assume that

$$(5.26) \quad \mathbb{E}[\mathcal{B}^\eta((1, -1), \mathbf{0})] \geq \mathbb{E}[\mathcal{B}^\theta((1, -1), \mathbf{0})].$$

If a geodesic between $(-n, t_{-n})$ and (n, t_n) passes through $\mathbf{0}$, then the geodesic cannot pass through any point above and to the left of $\mathbf{0}$. Then, for any $\mathbf{x} \in \mathbb{Z}_{>0} \times \mathbb{R}_{<0}$,

$$L_{(-n, t_{-n}), \mathbf{0}} + L_{\mathbf{0}, (n, t_n)} \geq L_{(-n, t_{-n}), \mathbf{x}} + L_{\mathbf{x}, (n, t_n)}$$

Specifically, for all $k \in \mathbb{Z}_{>0}$,

$$L_{(-n, t_{-n}), \mathbf{0}} - L_{(-n, t_{-n}), (k, -k)} \geq L_{(k, -k), (n, t_n)} - L_{\mathbf{0}, (n, t_n)}.$$

For fixed $\mathbf{x}, \mathbf{y} \in \mathbb{Z} \times \mathbb{R}$, let $\tilde{\mathcal{B}}^\eta(\mathbf{x}, \mathbf{y})$ denote the almost sure limit

$$\lim_{n \rightarrow \infty} L_{(-n, t_{-n}), \mathbf{x}} - L_{(-n, -t_{-n}), \mathbf{y}}.$$

By Theorem 3.4(ii) and Lemma 5.6, the limit exists almost surely, and $\tilde{\mathcal{B}}^\theta(\mathbf{x}, \mathbf{y}) \stackrel{d}{=} \mathcal{B}^\theta(\mathbf{y}, \mathbf{x})$. Let $\Omega_{\mathbb{Z}}$ be the full probability event on which, for every $k \in \mathbb{Z}_{>0}$ and for every sequence $\{t_n\}_{n \in \mathbb{Z}}$ satisfying

$$\lim_{n \rightarrow \infty} \frac{t_n}{n} = \theta \quad \text{and} \quad \lim_{n \rightarrow \infty} \frac{t_{-n}}{-n} = \eta,$$

$$\lim_{n \rightarrow \infty} L_{(-n, t_{-n}), \mathbf{0}}(\mathbf{B}) - L_{(-n, t_{-n}), (k, -k)}(\mathbf{B}) = \tilde{\mathcal{B}}^\eta(\mathbf{0}, (k, -k)) \quad \text{and}$$

$$\lim_{n \rightarrow \infty} L_{(k, -k), (n, t_n)}(\mathbf{B}) - L_{\mathbf{0}, (n, t_n)}(\mathbf{B}) = \mathcal{B}^\theta((k, -k), \mathbf{0}).$$

Therefore, we have the following inclusion of sets

$$\left\{ \begin{array}{l} \text{There exists a sequence } \{t_n\}_{n \in \mathbb{Z}} \text{ satisfying } \lim_{n \rightarrow \infty} \frac{t_n}{n} = \theta \text{ and } \lim_{n \rightarrow \infty} \frac{t_{-n}}{-n} = \eta \\ \\ \text{such that, for all } n \in \mathbb{Z}_{>0}, \text{ some geodesic between } (-n, t_{-n}) \text{ and } (n, t_n) \text{ passes through } \mathbf{0} \end{array} \right\}$$

$$\subseteq \bigcap_{m=1}^{\infty} \left\{ \begin{array}{l} \text{There exists a sequence } \{t_n\}_{n \in \mathbb{Z}} \text{ satisfying } \lim_{n \rightarrow \infty} \frac{t_n}{n} = \theta \text{ and } \lim_{n \rightarrow \infty} \frac{t_{-n}}{-n} = \eta \text{ such that} \\ \\ \text{for all } n \in \mathbb{Z}_{>0} \text{ and } k = 1, \dots, m, L_{(-n, t_{-n}), \mathbf{0}}(\mathbf{B}) - L_{(-n, t_{-n}), (k, -k)}(\mathbf{B}) \geq L_{(k, -k), (n, t_n)}(\mathbf{B}) - L_{\mathbf{0}, (n, t_n)}(\mathbf{B}) \end{array} \right\}$$

$$\subseteq \bigcap_{m=1}^{\infty} \left\{ \tilde{\mathcal{B}}^\eta(\mathbf{0}, (k, -k)) \geq \mathcal{B}^\theta((k, -k), \mathbf{0}), \quad k = 1, \dots, m \right\} \cup \Omega_{\mathbb{Z}}^C.$$

We show that this last event has probability 0. Define the process $\{S_k\}_{k \geq 0}$ by $S_0 = 0$ and for $k \geq 1$,

$$S_k = \mathcal{B}^\eta(\mathbf{0}, (k, -k)) - \mathcal{B}^\theta((k, -k), \mathbf{0}) = \sum_{i=1}^k (\tilde{\mathcal{B}}^\eta((i-1, -i+1), (i, -i)) - \mathcal{B}^\theta((i, -i), (i-1, -i+1))).$$

The sequences

$$\{\mathcal{B}^\theta((i, -i), (i-1, -i+1))\}_{i \in \mathbb{Z}_{>0}} \quad \text{and} \quad \{\tilde{\mathcal{B}}^\eta((i-1, -i+1), (i, -i))\}_{i \in \mathbb{Z}_{>0}}$$

are independent because they are constructed from disjoint increments of the field of independent Brownian motions \mathbf{B} . Furthermore,

$$\{\mathcal{B}^\theta((i, -i), (i-1, -i+1))\}_{i \in \mathbb{Z}_{>0}} = \{v_i^\theta(-i) + h_{i-1}^\theta(-i, -i+1)\}_{i \in \mathbb{Z}_{>0}}$$

is an i.i.d. collection of random variables by Theorem 3.4(iv). By reflection applied to the Busemann functions $\tilde{\mathcal{B}}^\eta$, the process $\{S_k\}_{k \geq 0}$ has independent increments. By Theorem 3.4, the increments of this random walk are independent sums of normal and exponential random variables and therefore have finite

first and second moment. Let μ be the mean and σ^2 the variance of S_1 . Then, $\mu \leq 0$ by the assumption (5.26). Thus, for $m \geq 1$,

$$\begin{aligned} \mathbb{P}(\tilde{\mathcal{B}}^n(\mathbf{0}, (k, -k)) \geq \mathcal{B}^\theta((k, -k), \mathbf{0}), \quad k = 1, \dots, m) &= \mathbb{P}(S_k \geq 0, \quad k = 1, \dots, m) \\ &= \mathbb{P}\left(\frac{S_{\lfloor um \rfloor} - um\mu}{\sqrt{m\sigma^2}} \geq -\frac{um\mu}{\sqrt{m\sigma^2}} \text{ for } u \in [0, 1]\right) \leq \mathbb{P}\left(\inf_{u \in [0, 1]} \frac{S_{\lfloor um \rfloor} - um\mu}{\sqrt{m\sigma^2}} \geq 0\right) \xrightarrow{m \rightarrow \infty} 0. \end{aligned}$$

The convergence in the last step holds because

$$\inf_{u \in [0, 1]} \frac{S_{\lfloor um \rfloor} - um\mu}{\sqrt{m\sigma^2}} \xrightarrow{m \rightarrow \infty} \inf_{u \in [0, 1]} B(u)$$

where the convergence holds in distribution by Donsker's Theorem, and B is a standard Brownian motion. \square

Proof of Theorem 3.17. Define

$$\widehat{\Omega}^{(\theta)} = \widetilde{\Omega}^{(\theta)} \cap \Omega^{(\theta, \star)} \cap \Omega^{(\theta, \theta)}(\mathbf{X}^\theta),$$

where $\widetilde{\Omega}^{(\theta)}$ is the event of Theorem 3.12, $\Omega^{(\theta, \star)}$ is the event of Theorem 5.7(v), and $\Omega^{(\theta, \theta)}(\mathbf{X}^\theta)$ is the event of Theorem 3.16, applied to the random environment \mathbf{X}^θ of i.i.d Brownian motions (Lemma 3.4(ii)).

By the construction of the $\mathbf{T}_\mathbf{x}^\theta$ in terms of the variational formula (Definition 3.6), if for some $\mathbf{x}, \mathbf{y} \in \mathbb{Z} \times \mathbb{R}$, the right-most semi-infinite geodesics in $\mathbf{T}_\mathbf{x}^\theta$ and $\mathbf{T}_\mathbf{y}^\theta$ ever intersect, they agree above and to the right of the point of intersection. The same is true of left-most geodesics. Because $\widetilde{\Omega}^{(\theta)} \subseteq \Omega^{(\theta)} \cap \Omega_2$ (Theorem 3.12),

$$\widehat{\Omega}^{(\theta)} \subseteq \Omega^{(\theta)} \cap \Omega^{(\theta, \star)} \cap \Omega_2.$$

If, by way of contradiction, for some $\omega \in \widehat{\Omega}^{(\theta)}$, there exists \mathbf{x} and $\mathbf{y} \in \mathbb{Z} \times \mathbb{R}$ such that the right-most geodesics in $\mathbf{T}_\mathbf{x}^\theta$ and $\mathbf{T}_\mathbf{y}^\theta$ are disjoint, then by Theorem 5.10, there exists a bi-infinite geodesic for the environment \mathbf{X}^θ that is defined by jump times $\{\tau_r\}_{r \in \mathbb{Z}}$ and satisfies

$$\lim_{n \rightarrow \infty} \frac{\tau_n}{n} = \theta = \lim_{n \rightarrow \infty} \frac{\tau_{-n}}{-n}.$$

This is a contradiction since $\omega \in \Omega^{(\theta, \theta)}(\mathbf{X}^\theta)$. Therefore, for all $\omega \in \widehat{\Omega}^{(\theta)}$, whenever $(m, s), (k, t) \in \mathbb{Z} \times \mathbb{R}$, the right-most geodesics in $\mathbf{T}_{(m, s)}^\theta$ and $\mathbf{T}_{(k, t)}^\theta$ coalesce. The same is true by replacing ‘‘right’’ with ‘‘left.’’ In other words, for all sufficiently large r ,

$$(5.27) \quad \tau_{(m, s), r}^{\theta, R} = \tau_{(k, t), r}^{\theta, R} \quad \text{and} \quad \tau_{(m, s), r}^{\theta, L} = \tau_{(k, t), r}^{\theta, L}.$$

Now, let $s = s_{m-1} \leq s_m \leq \dots$ and $t = t_{k-1} \leq t_k \leq \dots$ be any sequences defining θ -directed semi-infinite geodesics starting from (m, s) and (k, t) , respectively. We show that these semi-infinite geodesics coalesce. Without loss of generality, assume that $m \leq k$ and $s_{k-1} < t_{k-1} = t$. The other cases are handled similarly. Then, by Theorem 3.10(ii) and Corollary 3.14, for all $r \geq k$,

$$\tau_{(k, s_{k-1}), r}^{\theta, L} \leq s_r \leq \tau_{(k, s_{k-1}), r}^{\theta, R} \leq \tau_{(k, t), r}^{\theta, L} \leq t_r \leq \tau_{(k, t), r}^{\theta, R}.$$

Then, by (5.27), these inequalities are all equalities for all sufficiently large r , and the geodesics coalesce. \square

APPENDIX A. TECHNICAL LEMMAS AND THEOREMS

A.1. Deterministic facts about continuous functions.

Lemma A.1. *Let $\eta^1, \eta^2 : \mathbb{R} \rightarrow \mathbb{R}$ be continuous functions satisfying*

$$\limsup_{t \rightarrow \infty} \eta^i(t) = -\infty \quad \text{for } i = 1, 2.$$

Further, assume that $\eta^1(s, u) \leq \eta^2(s, u)$ for all $s < u$. For each $t \in \mathbb{R}$ and $i = 1, 2$, let $s_t^{i, L}$ and $s_t^{i, R}$ be the left-most and right-most maximizers of η^i on the set $[t, \infty)$. Then,

$$s_t^{1, L} \leq s_t^{2, L} \quad \text{and} \quad s_t^{1, R} \leq s_t^{2, R}.$$

Proof. Since $s_t^{2, L}$ is a maximizer for η^2 on $[t, \infty)$, $\eta^2(s_t^{2, L}) \geq \eta^2(s)$, or equivalently, $\eta^2(s_t^{2, L}, s) \leq 0$ for all $s \geq t$. Then, for all $s \geq s_t^{2, L} \geq t$, the hypothesis of the lemma gives

$$\eta^1(s_t^{2, L}, s) \leq \eta^2(s_t^{2, L}, s) \leq 0.$$

Thus, the left-most maximum of η^1 on the larger set $[t, \infty)$ can be no larger than $s_t^{2, L}$.

For right-most maximizers, the lemma follows by similar reasoning: for $s > s_t^{2, R}$,

$$\eta^1(s_t^{2, R}, s) \leq \eta^2(s_t^{2, R}, s) < 0,$$

so $\eta^1(s_t^{2, R}) > \eta^1(s)$ for all $s > s_t^{2, R}$, and no maximizer of $\eta^1(s)$ over $s \in [t, \infty)$ can be larger than $s_t^{2, R}$. \square

Lemma A.2. *Let $S \subseteq \mathbb{R}^n$, and let $f_n : S \rightarrow \mathbb{R}$ be a sequence of continuous functions, converging uniformly to the function $f : S \rightarrow \mathbb{R}$. Assume that there exists a sequence $\{c_n\}$, of maximizers of f_n , converging to some $c \in S$. Then, c is a maximizer of f .*

Proof. $f_n(c_n) \geq f_n(x)$ for all $x \in S$, so it suffices to show that $f_n(c_n) \rightarrow f(c)$. This follows from the uniform convergence of f_n to f , the continuity of f , and

$$|f_n(c_n) - f(c)| \leq |f_n(c_n) - f(c_n)| + |f(c_n) - f(c)|. \quad \square$$

Lemma A.3. *Let S_n for $n \geq 0$ be subsets of some set $\tilde{S} \subseteq \mathbb{R}^n$, on which the function $f : \tilde{S} \rightarrow \mathbb{R}$ is continuous. Assume that each point $x \in S_0$ is the limit of a sequence $\{x_n\}$, where $x_n \in S_n$ for each n . Assume that $\{c_n\}$ is a sequence of maximizers of f on S_n . Assume further that c_n converges to some $c \in S_0$. Then, c is a maximizer of f on S_0 .*

Proof. For each $x_0 \in S_0$, write $x_0 = \lim_{n \rightarrow \infty} x_n$, where $x_n \in S_n$ for each n . Then, $f(c_n) \geq f(x_n)$ for all $n \geq 1$, and the result follows by taking limits. \square

Lemma A.4. *Let $\mathbf{X} = \{X_m\}_{m \in \mathbb{Z}}$, where each $X_m : \mathbb{R} \rightarrow \mathbb{R}$ is a continuous function. Let $0 < s < t < T < u$ and $m \leq n$. Then,*

$$X_m(s, t) \leq L_{(m,s),(n,u)}(\mathbf{X}) - L_{(m,t),(n,u)}(\mathbf{X}) \leq L_{(m,s),(n,T)}(\mathbf{X}) - L_{(m,t),(n,T)}(\mathbf{X})$$

Similarly, let $0 < s < t < u < \infty$ and $m < n$. Then,

$$0 \leq L_{(m,s),(n,t)}(\mathbf{X}) - L_{(m+1,s),(n,t)}(\mathbf{X}) \leq L_{(m,s),(n,u)}(\mathbf{X}) - L_{(m+1,s),(n,u)}(\mathbf{X}).$$

Remark A.5. This is a deterministic statement. The only necessary ingredient is the continuity of the X_m so that each of the last-passage times is finite and has a sequence of maximizing times.

Proof. This proof follows a standard paths-crossing argument. For example, Lemma 4.6 in [ARAS20] follows the same procedure. We prove the first statement, and the second is proven similarly. By definition of last-passage time,

$$X_m(s, t) + L_{(m,t),(n,u)} \leq L_{(m,s),(n,u)}.$$

Since $0 < s < t < T < u$, the geodesic between (m, s) and (n, u) must cross the geodesic between (m, t) and (n, T) . Let $\mathbf{z} \in \mathbb{Z} \times \mathbb{R}$ be a point of intersection. Then,

$$L_{(m,s),(n,u)} = L_{(m,s),\mathbf{z}} + L_{\mathbf{z},(n,u)} \quad \text{and} \quad L_{(m,t),(n,T)} = L_{(m,t),\mathbf{z}} + L_{\mathbf{z},(n,T)},$$

so

$$L_{(m,s),(n,u)} - L_{(m,t),(n,u)} \leq L_{(m,s),\mathbf{z}} + L_{\mathbf{z},(n,u)} - (L_{(m,t),\mathbf{z}} + L_{\mathbf{z},(n,u)}) = L_{(m,s),\mathbf{z}} - L_{(m,t),\mathbf{z}}$$

$$= L_{(m,s),\mathbf{z}} + L_{\mathbf{z},(n,T)} - (L_{(m,t),\mathbf{z}} + L_{\mathbf{z},(n,T)}) \leq L_{(m,s),(n,T)} - L_{(m,t),(n,T)}. \quad \square$$

A.2. Probabilistic results. The following is a classical result that is often used in this paper.

Lemma A.6 ([BS02], Equation (1) on pg 251). *For a standard Brownian motion B and $\lambda > 0$,*

$$\sup_{0 \leq s < \infty} \{\sqrt{2}B(s) - \lambda s\} \sim \text{Exp}(\lambda).$$

Theorem A.7. *Let B be a standard Brownian motion, and for $t > 0$, let*

$$\begin{aligned} D(t) &:= \sup_{0 \leq s < \infty} \{\sqrt{2}B(s) - \lambda s\} - \sup_{t \leq s < \infty} \{\sqrt{2}B(s) - \lambda s\} \\ &= \left(\sup_{0 \leq s \leq t} \{\sqrt{2}B(t, s) + \lambda(t - s)\} - \sup_{t \leq s < \infty} \{\sqrt{2}B(t, s) + \lambda(t - s)\} \right)^+. \end{aligned}$$

Then, for all $z \geq 0$,

$$(A.1) \quad \mathbb{P}(D(t) \leq z) = \Phi\left(\frac{z - \lambda t}{\sqrt{2t}}\right) + e^{\lambda z} \left((1 + \lambda z + \lambda^2 t) \Phi\left(-\frac{z + \lambda t}{\sqrt{2t}}\right) - \lambda \sqrt{\frac{t}{\pi}} e^{-\frac{(z + \lambda t)^2}{4t}} \right).$$

Proof. For $x \geq 0$, by time reversal and [BS02], Equation 1.2.4 on page 251,

$$\begin{aligned} \mathbb{P}\left(\sup_{0 \leq s \leq t} \{\sqrt{2}B(t, s) + \lambda(t - s)\} \leq x\right) &= \mathbb{P}\left(\sup_{0 \leq s \leq t} \{\sqrt{2}B(t, t - s) - \lambda(t - s - t)\} \leq x\right) \\ &= \mathbb{P}\left(\sup_{0 \leq s \leq t} \{\sqrt{2}B(s) + \lambda s\} \leq x\right) = \Phi\left(\frac{x - \lambda t}{\sqrt{2t}}\right) - e^{\lambda x} \Phi\left(\frac{-x - \lambda t}{\sqrt{2t}}\right). \end{aligned}$$

By Lemma A.6,

$$\sup_{t \leq s < \infty} \{\sqrt{2}B(t, s) + \lambda(t - s)\} \sim \text{Exp}(\lambda).$$

The conclusion of the theorem follows by a simple, but tedious convolution. It suffices to show the $t = 1$ case, and the general case follows by Brownian scaling.

$$\begin{aligned} \mathbb{P}(D(1) \leq z) &= \int_{-\infty}^0 \left(\Phi\left(\frac{z-y-\lambda}{\sqrt{2}}\right) - e^{\lambda(z-y)} \Phi\left(\frac{-z+y-\lambda}{\sqrt{2}}\right) \right) \lambda e^{\lambda y} dy \\ &= \int_{-\infty}^0 \int_{-\infty}^{\frac{z-y-\lambda}{\sqrt{2}}} \frac{\lambda}{\sqrt{2\pi}} e^{-x^2/2} e^{\lambda y} dx dy - e^{\lambda z} \int_{-\infty}^0 \int_{-\infty}^{\frac{-z+y-\lambda}{\sqrt{2}}} \frac{\lambda}{\sqrt{2\pi}} e^{-x^2/2} dx dy. \end{aligned}$$

We now use Fubini's Theorem to switch the order of integration. This results in

$$\begin{aligned} &= \int_{-\infty}^{\frac{z-\lambda}{\sqrt{2}}} \int_{-\infty}^0 \frac{\lambda}{\sqrt{2\pi}} e^{-x^2/2} e^{\lambda y} dy dx + \int_{\frac{z-\lambda}{\sqrt{2}}}^{\infty} \int_{-\infty}^{z-\lambda-\sqrt{2}x} \frac{\lambda}{\sqrt{2\pi}} e^{-x^2/2} e^{\lambda y} dy dx \\ &\quad - e^{\lambda z} \int_{-\infty}^{-\frac{z+\lambda}{\sqrt{2}}} \int_{\sqrt{2}x+z+\lambda}^0 \frac{\lambda}{\sqrt{2\pi}} e^{-x^2/2} dy dx \\ &= \int_{-\infty}^{\frac{z-\lambda}{\sqrt{2}}} \frac{1}{\sqrt{2\pi}} e^{-x^2/2} dx + \int_{\frac{z-\lambda}{\sqrt{2}}}^{\infty} \frac{1}{\sqrt{2\pi}} e^{-x^2/2} e^{z\lambda - \sqrt{2}\lambda x - \lambda^2} dx + e^{\lambda z} \int_{-\infty}^{-\frac{z+\lambda}{\sqrt{2}}} (\sqrt{2}x + z + \lambda) \frac{\lambda}{\sqrt{2\pi}} e^{-x^2/2} dx \\ &= \Phi\left(\frac{z-\lambda}{\sqrt{2}}\right) + e^{\lambda z} \int_{\frac{z-\lambda}{\sqrt{2}}}^{\infty} \frac{1}{\sqrt{2\pi}} e^{-\frac{(x+\sqrt{2}\lambda)^2}{2}} dx + e^{\lambda z} \left(\int_{-\infty}^{-\frac{z+\lambda}{\sqrt{2}}} \frac{\lambda}{\sqrt{\pi}} x e^{-x^2/2} dx + (\lambda z + \lambda^2) \Phi\left(-\frac{z+\lambda}{\sqrt{2}}\right) \right) \\ &= \Phi\left(\frac{z-\lambda}{\sqrt{2}}\right) + e^{\lambda z} \int_{\frac{z+\lambda}{\sqrt{2}}}^{\infty} \frac{1}{\sqrt{2\pi}} e^{-u^2/2} du + e^{\lambda z} \left(-\frac{\lambda}{\sqrt{\pi}} e^{-\frac{(z+\lambda)^2}{4}} + (\lambda z + \lambda^2) \Phi\left(-\frac{z+\lambda}{\sqrt{2}}\right) \right) \\ &= \Phi\left(\frac{z-\lambda}{\sqrt{2}}\right) + e^{\lambda z} \left((1 + \lambda z + \lambda^2) \Phi\left(-\frac{z+\lambda}{\sqrt{2}}\right) - \frac{\lambda}{\sqrt{\pi}} e^{-\frac{(z+\lambda)^2}{4}} \right). \quad \square \end{aligned}$$

Theorem A.8. *Let B be a standard Brownian motion and let $\lambda > 0$. Let T be the unique maximizer of $\sqrt{2}B(t) - \lambda t$ for $t \in [0, \infty)$. Then, for $t \geq 0$,*

$$\mathbb{P}(T > t) = (2 + \lambda^2 t) \Phi\left(-\lambda \sqrt{\frac{t}{2}}\right) - \lambda \sqrt{\frac{t}{\pi}} e^{-\frac{\lambda^2 t}{4}}.$$

Remark A.9. Theorem A.8 should be credited to Norros and Salminen, who computed the Laplace transform of this random variable in Proposition 3.9 of [SN01]. One can also obtain by integrating the probability density of the time of maximum of Brownian motion with drift on the interval $[0, t]$ found in [Buf03], Equation (1.3), and then taking $t \rightarrow \infty$. See also the discussion after Equation (1.3) equation in [Buf03] for more historical details on this formula. Theorem A.7 will be used in subsequent papers, so we use it to give our own proof of Theorem A.8.

Proof. Note that $T > t$ if and only if $D(t) = 0$ in the sense of Theorem A.7. Thus, the Lemma follows by setting $z = 0$ in the formula (A.1). \square

Theorem A.10. *Let X be a two-sided Brownian motion with strictly negative drift. Let*

$$M = \{t \in \mathbb{R} : X(t) = \sup_{t \leq s < \infty} X(s)\}.$$

Furthermore, we define

$$M^U = \left\{ t \in M : X(t) > X(s) \text{ for all } s > t \right\}$$

to be the set of elements of M that are unique maximizers of $X(s)$ over $s \in [t, \infty)$ and define

$$M^N = M \setminus M^U$$

to be the set of elements of M that are non-unique maximizers of $X(s)$ over $s \in [t, \infty)$. Then, there exists an event of probability one, on which the following hold.

- (i) M is a closed set.
- (ii) M^N is a countably infinite set.
- (iii) For all $\hat{t} \in M^U$ and $\varepsilon > 0$, there exists $t \in M^N$ satisfying $\hat{t} < t < \hat{t} + \varepsilon$. For all $t \in M^N$ and $\varepsilon > 0$, there exists $\hat{t} \in M^U$ satisfying $t - \varepsilon < \hat{t} < t$.
- (iv) For all $t \in M^N$ and $\varepsilon > 0$, there exists $t^* \in M^N$ with $t - \varepsilon < t^* < t$. For each $t \in M^N$, there exists $\delta > 0$ such that $M \cap (t, t + \delta) = \emptyset$.

Proof. By the $n = m$ case of Lemma 2.3, for each fixed $t \in \mathbb{R}$, $X(s)$ is almost surely uniquely maximized for $s \in [t, \infty)$. By Theorem A.8, this maximizer is almost surely strictly greater than t . Let $\Omega_{\mathbb{Q}}$ be the full probability event on which,

$$\lim_{s \rightarrow -\infty} X(s) = \infty \quad \text{and} \quad \lim_{s \rightarrow \infty} X(s) = -\infty,$$

and such that, for every $q \in \mathbb{Q}$, there is a unique maximizer of $X(s)$ over $s \in [q, \infty)$ that is strictly larger than q .

Part (i): This follows from Lemma A.3.

Part (ii): The set M is nonempty because, for any $s \in \mathbb{R}$, any maximizer t of $X(u)$ over $[s, \infty)$ lies in M . Then, the fact that M^N is infinite follows from Part (iv). We show that M^N is at most countably infinite and then prove Parts (iii) and (iv). To do this, we first show that for each $\omega \in \Omega_{\mathbb{Q}}$, there exists no $t \in \mathbb{R}$ such that $X(s)$ has three maximizers over $s \in [t, \infty)$. If, on the contrary, such a t exists, at least two of the maximizers must be greater than t , and therefore, these two maximizers are also maximizers of $X(s)$ over $s \in [q, \infty)$ for some $q \in \mathbb{Q}$. Thus, on $\Omega_{\mathbb{Q}}$, for each $t \in M^N$, there exists a unique $\hat{t} \in M^U$ with $\hat{t} > t$ and $X(t) = X(\hat{t})$. Therefore, M^N is at most countable because each element $t \in M^N$ is associated with the nonempty open interval (t, \hat{t}) , whose intersection with M is the empty set.

Part (iii): Let $\omega \in \Omega_{\mathbb{Q}}$, $\hat{t} \in M^U$ and $\varepsilon > 0$, and let $q \in \mathbb{Q}$ satisfy $\hat{t} < q < \hat{t} + \varepsilon$. Then, there is a unique maximizer, $s > q$ of $X(u)$ over $u \in [q, \infty)$. By assumption, \hat{t} uniquely maximizes $X(u)$ over the larger set $[\hat{t}, \infty)$, so $X(q) < X(s) < X(\hat{t})$. By the intermediate value theorem, there exists a point u with $\hat{t} < u < q$ and $X(u) = X(s)$. Then, set

$$t = \max\{u \in (\hat{t}, s) : X(u) = X(s)\}.$$

Then, $t < q < s$, and

$$X(t) = X(s) = \sup_{t \leq u < \infty} X(u)$$

because $X(u) < X(s)$ for all $t < u < q$. Therefore, $\hat{t} < t < q < \hat{t} + \varepsilon$, and $t \in M^N$.

Now, let $t \in M^N$ and $\varepsilon > 0$. Let $q \in \mathbb{Q}$ satisfy $t - \varepsilon < q < t$. Then, there is a unique maximizer \hat{t} of $X(s)$ over $s \in [q, \infty)$, so $X(\hat{t}) > X(s)$ for all $s \in [q, \infty) \setminus \{\hat{t}\}$. Specifically, $\hat{t} \in M^U$ and $X(\hat{t}) > X(t)$, so since t maximizes $X(s)$ over $s \in [t, \infty)$, \hat{t} cannot be greater than or equal to t . Therefore, $t - \varepsilon < q < \hat{t} < t$.

Part (iv): The first statement follows immediately from Part (iii). The second statement was shown in the proof of Part (ii). \square

Remark A.11. Originally discovered by Taylor [Tay55], the zero set of Brownian motion almost surely has Hausdorff dimension $\frac{1}{2}$. Parts (i) and (iii) of Theorem A.10 imply that M is the supporting set of the Lebesgue-Stieltjes measure defined by the nondecreasing function $t \mapsto -\sup_{t \leq s < \infty} X(s)$. Using this fact, Taylor's proof (see also Theorem 4.24 in [MP10]) can be modified to show that the set M almost surely has Hausdorff dimension $\frac{1}{2}$. This will be used in future work when studying the competition interface for BLPP.

A.3. The Brownian queue and stationary last-passage process. This section discusses the Brownian queue in the formulation of [OY01]. Let A and S be two independent, two-sided Brownian motions and let $\lambda > 0$. For $s < t$, $A(s, t)$ represents the arrivals to the queue in the time interval $(s, t]$, and $\lambda(t - s) - S(s, t)$ is the amount of service available in $(s, t]$. For $t \in \mathbb{R}$, set

$$q(t) = \sup_{-\infty < s \leq t} \{A(s, t) + S(s, t) - \lambda(t - s)\} \quad \text{and} \quad d(t) = A(t) + q(0) - q(t).$$

In queuing terms, $q(t)$ is the length of the queue at time t , and for $s < t$, $d(s, t)$ is the number of departures from the queue in the interval $(s, t]$. These processes are not integer-valued, but are viewed as heavy-traffic limits. We also define $e(t) = S(t) + q(0) - q(t)$. The following is due to O'Connell and Yor [OY01]. Without the statements for the process e the theorem is a special case of a more general result previously shown by Harrison and Williams [HW90].

Theorem A.12 ([OY01], Theorem 4). *The processes d and e are independent, two-sided Brownian motions. Furthermore, for each $t \in \mathbb{R}$, $\{d(s, t), e(s, t) : -\infty < s \leq t\}$ is independent of $\{q(u) : u \geq t\}$.*




FIGURE A.1. Independence structure for stationary BLPP

We reformulate Theorem A.12 in terms of the queuing mappings of (3.1)–(3.3) and (3.4)–(3.6).

Theorem A.13. *Let Y be a two-sided Brownian motion with drift $\lambda > 0$, independent of the two-sided Brownian motion C (with no drift). Then, $\overleftarrow{D}(Y, C)$ is a two-sided Brownian motion with drift λ , independent of the two-sided Brownian motion $\overleftarrow{R}(Y, C)$. Furthermore, for all $t \in \mathbb{R}$, $\{(\overleftarrow{D}(Y, C)(s, t), \overleftarrow{R}(Y, C)(s, t)) : -\infty < s \leq t\}$ is independent of $\{\overleftarrow{Q}(Y, C)(u) : u \geq t\}$.*

Proof. Let $A(t) = \lambda t - Y(t)$ and $S(t) = C(t)$, two independent two-sided Brownian motions. Then

$$q(t) = \sup_{-\infty < s \leq t} \{A(s, t) + S(s, t) - \lambda(t - s)\} = \sup_{-\infty < s \leq t} \{C(s, t) - Y(s, t)\} = \overleftarrow{Q}(Y, C)(t).$$

Next observe that

$$\begin{aligned} \overleftarrow{D}(Y, C)(t) &= Y(t) + \overleftarrow{Q}(Y, C)(t) - \overleftarrow{Q}(Y, C)(0) = -A(t) + \lambda t + q(t) - q(0) = \lambda t - d(t), \\ \text{and } \overleftarrow{R}(Y, C)(t) &= C(t) + \overleftarrow{Q}(Y, C)(0) - \overleftarrow{Q}(Y, C)(t) = S(t) + q(0) - q(t) = e(t). \end{aligned}$$

The result follows from Theorem A.12. \square

Fix a parameter $\lambda > 0$. Given an environment of Brownian motions $\mathbf{B} = \{B_m\}_{m \in \mathbb{Z}}$, set

$$Y_0^\lambda(t) = -B_0(t) + \lambda t.$$

For $m > 0$, recalling definitions (3.4)–(3.6), set

$$(A.2) \quad q_m^\lambda(t) := \overleftarrow{Q}(Y_{m-1}^\lambda, B_m)(t), \quad Y_m^\lambda(t) := \overleftarrow{D}(Y_{m-1}^\lambda, B_m)(t), \quad \text{and } W_{m-1}^\lambda(t) := \overleftarrow{R}(Y_{m-1}^\lambda, B_m)(t).$$

The increment-stationary BLPP is constructed as follows: For $(n, t) \in \mathbb{Z}_{>0} \times \mathbb{R}$, set

$$L_{(n,t)}^\lambda = \sup_{-\infty < s \leq t} \{Y_0^\lambda(s) + L_{(1,s),(n,t)}(\mathbf{B})\}.$$

One can check inductively, as in Section 4 of [OY01], that for $m \geq 1$ and $s, t \in \mathbb{R}$,

$$q_m^\lambda(t) = L_{(m,t)}^\lambda - L_{(m-1,t)}^\lambda, \quad \text{and } Y_m^\lambda(s, t) = L_{(m,t)}^\lambda - L_{(m,s)}^\lambda.$$

The stationarity of these increments comes from the next theorem that follows from Theorem A.13 and induction. It is the zero-temperature analogue of Theorem 3.3 in [SV10] and Theorem 2.11 in [ARAS20]. Figure A.1 demonstrates the independence structure, and we give credit to [ARAS20] for a very similar picture.

Theorem A.14. *Let $Y_m^\lambda, q_m^\lambda, W_m^\lambda$ be as constructed in (A.2). Then, the following hold.*

- (i) *For all $m > 0$ and $t \in \mathbb{R}$, $q_m^\lambda(t) \sim \text{Exp}(\lambda)$. $\{W_m^\lambda\}_{m \geq 0}$ is a field of independent two-sided Brownian motions. $\{Y_m^\lambda\}_{m \geq 0}$ is a field of (non-independent) two-sided Brownian motions with drift λ .*
- (ii) *Let $-\infty < t_n \leq t_{n-1} \leq \dots \leq t_1 < \infty$. Then, the following are mutually independent:*

$$\{W_0^\lambda(u, t_1) : u \leq t_1\}, \{Y_0^\lambda(t_1, u) : u \geq t_1\},$$

$$q_r^\lambda(t_r), \{B_r(t_r, u) : u \geq t_r\}, \{W_{r-1}^\lambda(u, t_r) : u \leq t_r\}, \text{ and } \{Y_r^\lambda(t_{r+1}, u) : t_{r+1} \leq u \leq t_r\} \text{ for } 1 \leq r \leq n-1,$$

$$\text{and } q_n^\lambda(t_n), \{B_n(t_n, u) : u \geq t_n\}, \text{ and } \{Y_n^\lambda(u, t_n) : u \leq t_n\}.$$

Proof. To prove Part (i), begin with the assumption:

$$(Y_0^\lambda, B_1, \dots, B_n) \quad \text{are mutually independent.}$$

By Theorem A.13, $Y_1^\lambda = \overleftarrow{D}(Y_0, B_1)$ is a two-sided Brownian motion with drift λ , independent of the two-sided Brownian motion $W_0^\lambda = \overleftarrow{R}(Y_0, B_1)$. Hence

$$(W_0^\lambda, Y_1^\lambda, B_2, \dots, B_n) \quad \text{are mutually independent.}$$

Inductively, assume that Y_r^λ is a two-sided Brownian motion with drift λ and

$$(W_0, \dots, W_{r-1}, Y_r^\lambda, B_{r+1}, \dots, B_n) \quad \text{are mutually independent.}$$

Apply Theorem A.13 to $Y_{r+1}^\lambda = \overleftarrow{D}(Y_r^\lambda, B_{r+1})$ and $W_r^\lambda = \overleftarrow{R}(Y_r^\lambda, B_{r+1})$ to continue the induction. Part (i) is proved. The proof of Part (ii) is the proof of Theorem 3.3 in [SV10]. \square

The proof of Theorem 2.5 is achieved through the coupling of q_r^λ , Y_r^λ , and W_r^λ . The following theorem is the key.

Theorem A.15 (Zero-temperature analogue of Theorem 3.1 in [ARAS20]). *Fix real numbers $t > s$, $\lambda > 0$, and $\gamma > \lambda^{-2} > \delta > 0$. Then, with probability one,*

$$(A.3) \quad \limsup_{n \rightarrow \infty} [L_{(0,t),(n,n\delta)}(\mathbf{W}^\lambda) - L_{(0,s),(n,n\delta)}(\mathbf{W}^\lambda)] \leq B_0(s, t) - \lambda(t - s) = -Y_0^\lambda(s, t) \\ \leq \liminf_{n \rightarrow \infty} [L_{(0,t),(n,n\gamma)}(\mathbf{W}^\lambda) - L_{(0,s),(n,n\gamma)}(\mathbf{W}^\lambda)],$$

and

$$(A.4) \quad \limsup_{n \rightarrow \infty} [L_{(0,t),(n,n\delta)}(\mathbf{W}^\lambda) - L_{(1,t),(n,n\delta)}(\mathbf{W}^\lambda)] \\ \leq q_1^\lambda(t) \leq \liminf_{n \rightarrow \infty} [L_{(0,t),(n,n\gamma)}(\mathbf{W}^\lambda) - L_{(1,t),(n,n\gamma)}(\mathbf{W}^\lambda)].$$

Here, $L_{\mathbf{x}, \mathbf{y}}(\mathbf{W}^\lambda)$ is the last passage time between points $\mathbf{x} \leq \mathbf{y} \in \mathbb{Z}_{>0} \times \mathbb{R}$ when $\mathbf{W}^\lambda := \{W_m^\lambda\}_{m \geq 0}$ is the random environment. (W_m^λ need not be defined for $m < 0$ since all points $(m, t) \in \mathbb{Z} \times \mathbb{R}$ in the above expression satisfy $m \geq 0$). While Theorem A.15 is not explicitly stated in [ARAS20], all necessary lemmas are provided in Section 4 of that paper.

Lemma A.16. *Let $\mathcal{A} \subseteq \mathbb{R}$ be any countable set. Consider the space $\mathbb{R}^{\mathcal{A}}$ of functions $\mathcal{A} \rightarrow \mathbb{R}$, equipped with the Borel product σ -algebra. Fix $\theta > 0$. Then,*

$$\{h_0^\theta(t) : t \in \mathcal{A}\} \stackrel{d}{=} \{-B_0(t) + \frac{1}{\sqrt{\theta}}t : t \in \mathcal{A}\} \quad \text{and} \quad \{v_1^\theta(t) : t \in \mathcal{A}\} \stackrel{d}{=} \{q_1^{\frac{1}{\sqrt{\theta}}}(t) : t \in \mathcal{A}\}.$$

Proof. This proof is a multivariate extension of the ‘‘Proof of Theorem 2.5, assuming Theorem 3.1’’ on page 1937 of [ARAS20]. This method originated in the setting of the log-gamma polymer in [GRASY15]. From definition of h_0^θ (Equation (2.6)), $h_0^\theta(0) = 0 = B_0(0)$. Hence, to prove the first statement, it is sufficient to show that

$$\{h_0^\theta(s, t) : s, t \in \mathcal{A}, s < t\} \stackrel{d}{=} \{-B_0(s, t) + \frac{1}{\sqrt{\theta}}(t - s) : s, t \in \mathcal{A}, s < t\}.$$

Theorem 2.5 establishes that, for fixed $s, t \in \mathbb{R}$,

$$h_0^\theta(s, t) = \lim_{n \rightarrow \infty} L_{(0,s),(n,n\theta)}(\mathbf{B}) - L_{(0,t),(n,n\theta)} \text{ a.s.}$$

By Theorem A.14, $\{W_m^\lambda\}_{m \geq 0}$ is a field of independent, two-sided Brownian motions. By equality of distribution between $\{W_m^\lambda\}_{m \geq 0}$ and $\{B_m\}_{m \geq 0}$, the first inequality of (A.3) implies that if $\{s_1, \dots, s_k\}$ and $\{t_1, \dots, t_k\}$ are finite subsets of \mathcal{A} with $s_i < t_i$ for all i , and $x_1, \dots, x_k \in \overline{\mathbb{R}}$, then whenever $\theta < \lambda^{-2}$, or equivalently, $\lambda < \frac{1}{\sqrt{\theta}}$,

$$\mathbb{P}(-h_0^\theta(s_i, t_i) \geq x_i, \quad 1 \leq i \leq k) \leq \mathbb{P}(B_0(s_i, t_i) - \lambda(t_i - s_i) \geq x_i, \quad 1 \leq i \leq k).$$

Since $\lambda > 0$ and $s_i < t_i$ for all i , the right-hand side decreases as λ increases, so taking $\lambda \nearrow \frac{1}{\sqrt{\theta}}$ gives us

$$\mathbb{P}(-h_0^\theta(s_i, t_i) \geq x_i, \quad 1 \leq i \leq k) \leq \mathbb{P}\left(B_0(s_i, t_i) - \frac{1}{\sqrt{\theta}}(t_i - s_i) \geq x_i, \quad 1 \leq i \leq k\right).$$

The inequality on the right of (A.3) with $\lambda \searrow \frac{1}{\sqrt{\theta}}$ establishes the reverse inequality.

The same argument works for v_1^θ , utilizing this monotonicity for $\gamma < \delta$:

$$q_1^\gamma(t) = \overleftarrow{Q}(Y, B_1)(t) = \sup_{-\infty < s \leq t} \{B_1(s, t) + B_0(s, t) - \gamma(t - s)\}$$

$$\geq \sup_{-\infty < s \leq t} \{B_1(s, t) + B_0(s, t) - \delta(t - s)\} \geq q_1^\delta(t). \quad \square$$

A.4. Time reversal.

Theorem A.17. *Let Z, B, Y, C be continuous functions satisfying $Z(0) = B(0) = Y(0) = C(0) = 0$ and*

$$\lim_{t \rightarrow \pm\infty} (B(t) - Z(t)) = \lim_{t \rightarrow \pm\infty} (C(t) - Y(t)) = \mp\infty.$$

Then,

$$Y = D(Z, B) \text{ and } C = R(Z, B) \quad \text{if and only if} \quad Z = \overleftarrow{D}(Y, C) \text{ and } B = \overleftarrow{R}(Y, C),$$

The equalities above denote equality as functions of t . If either of the above equivalent conditions are satisfied, then also $Q(Z, B) = \overleftarrow{Q}(Y, C)$.

The following lemmas will help to prove Theorem A.17.

Lemma A.18. *Let $Z, B : \mathbb{R} \rightarrow \mathbb{R}$ be continuous functions satisfying $X(0) = B(0) = 0$ and*

$$\lim_{t \rightarrow \pm\infty} (B(t) - X(t)) = \mp\infty.$$

For a function $f : \mathbb{R} \rightarrow \mathbb{R}$, recall that we define $\tilde{f} : \mathbb{R} \rightarrow \mathbb{R}$ by $\tilde{f}(t) = -f(-t)$. Then, for all $t \in \mathbb{R}$,

$$Q(Z, B)(-t) = \overleftarrow{Q}(\tilde{Z}, \tilde{B})(t), \quad -D(Z, B)(-t) = \overleftarrow{D}(\tilde{Z}, \tilde{B})(t), \quad \text{and} \quad -R(Z, B)(-t) = \overleftarrow{R}(\tilde{Z}, \tilde{B})(t).$$

Proof. This is a routine check, using the definitions. \square

The following is a well-known fact, but is often stated without proof, so we include full justification for the sake of completeness.

Lemma A.19 ([Pit75], Equation (1.4)). *Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be a continuous function such that*

$$\lim_{t \rightarrow \pm\infty} f(t) = \pm\infty.$$

Set $F(t) = \sup_{-\infty < s \leq t} f(s)$. Then,

$$(A.5) \quad \inf_{t \leq s < \infty} (2F(s) - f(s)) = F(t).$$

Proof. The left-hand side of (A.5) is

$$\inf_{t \leq s < \infty} (2F(s) - f(s)) = \inf_{t \leq s < \infty} \left(2 \sup_{-\infty < u \leq s} f(u) - f(s) \right).$$

For all $s \geq t$, $\sup_{-\infty < u \leq s} f(u)$ is greater than or equal to both $f(s)$ and $\sup_{-\infty < u \leq t} f(u)$. Therefore,

$$2 \sup_{-\infty < u \leq s} f(u) - f(s) \geq \sup_{-\infty < u \leq t} f(u) + f(s) - f(s) = F(t).$$

This establishes one direction of Equation (A.5). To show the other direction, we show that there exists $s \geq t$ such that

$$2F(s) - f(s) = F(t).$$

Note that $f(t) \leq F(t)$ and that $\lim_{s \rightarrow \infty} f(s) = \infty$ by assumption. Hence, by continuity of f , $f(s) = F(t)$ for some $s \geq t$. Let

$$s^* = \inf\{s \geq t : f(s) = F(t)\}.$$

Then, $F(s^*) = F(t)$. Therefore,

$$2F(s^*) - f(s^*) = 2F(t) - F(t) = F(t). \quad \square$$

Proof of Theorem A.17. Assume that Z, B, Y, C satisfy the conditions of the Theorem. First assume that $Z = \overleftarrow{D}(Y, C)$ and $B = \overleftarrow{R}(Y, C)$. By definitions (3.4)–(3.6),

$$\begin{aligned} Z(s) &= Y(s) + \sup_{-\infty < u \leq s} \{C(u, s) - Y(u, s)\} - \sup_{-\infty < u \leq 0} \{C(u, 0) - Y(u, 0)\} \\ &= C(s) + \sup_{-\infty < u \leq s} \{Y(u) - C(u)\} - \sup_{-\infty < u \leq 0} \{Y(u) - C(u)\}, \quad \text{and} \end{aligned}$$

$$\begin{aligned} B(s) &= C(s) + \sup_{-\infty < u \leq 0} \{C(u, 0) - Y(u, 0)\} - \sup_{-\infty < u \leq s} \{C(u, s) - Y(u, s)\} \\ &= Y(s) + \sup_{-\infty < u \leq 0} \{Y(u) - C(u)\} - \sup_{-\infty < u \leq s} \{Y(u) - C(u)\}. \end{aligned}$$

Then,

$$\begin{aligned}
 D(Z, B)(t) &= Z(t) + Q(Z, B)(0) - Q(Z, B)(t) \\
 &= Z(t) + \sup_{0 \leq s < \infty} \{B(0, s) - Z(0, s)\} - \sup_{t \leq s < \infty} \{B(t, s) - Z(t, s)\} \\
 &= B(t) + \sup_{0 \leq s < \infty} \{B(s) - Z(s)\} - \sup_{t \leq s < \infty} \{B(s) - Z(s)\} \\
 &= Y(t) + \sup_{-\infty < s \leq 0} \{Y(s) - C(s)\} - \sup_{-\infty < s \leq t} \{Y(s) - C(s)\} \\
 &\quad + \sup_{0 \leq s < \infty} \{Y(s) - C(s) - 2 \sup_{-\infty < u \leq s} \{Y(u) - C(u)\}\} \\
 &\quad - \sup_{t \leq s < \infty} \{Y(s) - C(s) - 2 \sup_{-\infty < u \leq s} \{Y(u) - C(u)\}\}.
 \end{aligned}$$

To show that this equals $Y(t)$, it is therefore sufficient to show that for $t \in \mathbb{R}$,

$$\sup_{-\infty < s \leq t} \{Y(s) - C(s)\} = \inf_{t \leq s < \infty} \left\{ 2 \sup_{-\infty < u \leq s} \{Y(u) - C(u)\} - (Y(s) - C(s)) \right\},$$

which follows from Lemma A.19. The proof that $R(Z, B)(t) = C(t)$ follows by the same reasoning. The converse then follows by the previous case and Lemma A.18: if $Y = D(Z, B)$ and $C = R(Z, B)$, $\tilde{Y} = \overleftarrow{D}(\tilde{Z}, \tilde{B})$ and $\tilde{C} = \overleftarrow{R}(\tilde{Z}, \tilde{B})$, so $\tilde{Z} = D(\tilde{Y}, \tilde{C})$ and $\tilde{B} = R(\tilde{Y}, \tilde{C})$. Hence $Z = \overleftarrow{D}(Y, C)$ and $B = \overleftarrow{R}(Y, C)$.

We finish by showing that $Q(Z, B) = \overleftarrow{Q}(Y, C)$ whenever the two equivalent conditions of the theorem are met. For all $t \in \mathbb{R}$,

$$Y(t) = D(Z, B)(t) = Z(t) + Q(Z, B)(0) - Q(Z, B)(t), \text{ and}$$

$$Z(t) = \overleftarrow{D}(Y, C) = Y(t) + \overleftarrow{Q}(Y, C)(t) - \overleftarrow{Q}(Y, C)(0).$$

Putting these two equations together,

$$Q(Z, B)(t) - \overleftarrow{Q}(Y, C)(t) = Q(Z, B)(0) - \overleftarrow{Q}(Z, B)(0).$$

This is true for all $t \in \mathbb{R}$, so $Q(Z, B)(t) - \overleftarrow{Q}(Y, C)(t)$ must be equal to some constant. Recall that

$$Q(Z, B)(t) = \sup_{t \leq s < \infty} \{B(t, s) - Z(t, s)\}, \quad \text{and} \quad \overleftarrow{Q}(Y, C)(t) = \sup_{-\infty < s \leq t} \{C(s, t) - Y(s, t)\}.$$

By the limit conditions of the theorem, maximizers exist for each of the supremums above, so $Q(Z, B)$ and $\overleftarrow{Q}(Y, C)$ both achieve a minimum value of 0. Hence, the constant must be 0. \square

REFERENCES

- [AH16] Daniel Ahlberg and Christopher Hoffman. Random coalescing geodesics in first-passage percolation. *arXiv e-prints*, page arXiv:1609.02447, 2016.
- [ARAS20] Tom Alberts, Firas Rassoul-Agha, and Mackenzie Simper. Busemann functions and semi-infinite O’Connell-Yor polymers. *Bernoulli*, 26(3):1927–1955, 2020.
- [Bar01] Yu. Baryshnikov. GUEs and queues. *Probab. Theory Related Fields*, 119(2):256–274, 2001.
- [BBS20] Márton Balázs, Ofer Busani, and Timo Seppäläinen. Non-existence of bi-infinite geodesics in the exponential corner growth model. *Forum Math. Sigma*, 8:Paper No. e46, 34, 2020.
- [BBS21] Márton Balázs, Ofer Busani, and Timo Seppäläinen. Local stationarity of exponential last passage percolation. *Probab. Theory Related Fields*, 2021. To appear.
- [BF20] Ofer Busani and Patrik Ferrari. Universality of the geodesic tree in last passage percolation. *arXiv e-prints*, page arXiv:2008.07844, 2020.
- [BGH19] Erik Bates, Shirshendu Ganguly, and Alan Hammond. Hausdorff dimensions for shared endpoints of disjoint geodesics in the directed landscape. *arXiv e-prints*, page arXiv:1912.04164, 2019.
- [BGH21] Riddhipratim Basu, Shirshendu Ganguly, and Alan Hammond. Fractal geometry of Airy₂ processes coupled via the Airy sheet. *Ann. Probab.*, 49(1):485–505, 2021.
- [BHS18] Riddhipratim Basu, Christopher Hoffman, and Allan Sly. Nonexistence of bigeodesics in integrable models of last passage percolation. *arXiv e-prints*, page arXiv:1811.04908, 2018.
- [BKS03] Itai Benjamini, Gil Kalai, and Oded Schramm. First passage percolation has sublinear distance variance. *Ann. Probab.*, 31(4):1970–1978, 2003.
- [BS02] Andrei N. Borodin and Paavo Salminen. *Handbook of Brownian motion—facts and formulae*. Probability and its Applications. Birkhäuser Verlag, Basel, second edition, 2002.

- [BSS19] Riddhipratim Basu, Sourav Sarkar, and Allan Sly. Coalescence of geodesics in exactly solvable models of last passage percolation. *J. Math. Phys.*, 60(9):093301, 22, 2019.
- [Buf03] Emmanuel Buffet. On the time of the maximum of Brownian motion with drift. *J. Appl. Math. Stochastic Anal.*, 16(3):201–207, 2003.
- [CH14] Ivan Corwin and Alan Hammond. Brownian Gibbs property for Airy line ensembles. *Invent. Math.*, 195(2):441–508, 2014.
- [CHHM21] Ivan Corwin, Alan Hammond, Milind Hegde, and Konstantin Matetski. Exceptional times when the KPZ fixed point violates Johansson’s conjecture on maximizer uniqueness. *arXiv e-prints*, page arXiv:2101.04205, 2021.
- [DH14] Michael Damron and Jack Hanson. Busemann functions and infinite geodesics in two-dimensional first-passage percolation. *Comm. Math. Phys.*, 325(3):917–963, 2014.
- [DH17] Michael Damron and Jack Hanson. Bigeodesics in first-passage percolation. *Comm. Math. Phys.*, 349(2):753–776, 2017.
- [DNV19] Duncan Dauvergne, Mihai Nica, and Bálint Virág. Uniform convergence to the Airy line ensemble. *arXiv e-prints*, page arXiv:1907.10160, 2019.
- [DOV18] Duncan Dauvergne, Janosch Ortmann, and Bálint Virág. The directed landscape. *arXiv e-prints*, page arXiv:1812.00309, 2018.
- [FP05] Pablo A. Ferrari and Leandro P. R. Pimentel. Competition interfaces and second class particles. *Ann. Probab.*, 33(4):1235–1254, 2005.
- [FS20] Wai-Tong (Louis) Fan and Timo Seppäläinen. Joint distribution of Busemann functions for the exactly solvable corner growth model. *Probability and Mathematical Physics*, 1(1):55–100, 2020.
- [GRAS17a] Nicos Georgiou, Firas Rassoul-Agha, and Timo Seppäläinen. Geodesics and the competition interface for the corner growth model. *Probab. Theory Related Fields*, 169(1-2):223–255, 2017.
- [GRAS17b] Nicos Georgiou, Firas Rassoul-Agha, and Timo Seppäläinen. Stationary cocycles and Busemann functions for the corner growth model. *Probab. Theory Related Fields*, 169(1-2):177–222, 2017.
- [GRASY15] Nicos Georgiou, Firas Rassoul-Agha, Timo Seppäläinen, and Atilla Yilmaz. Ratios of partition functions for the log-gamma polymer. *Ann. Probab.*, 43(5):2282–2331, 2015.
- [GTW01] Janko Gravner, Craig A. Tracy, and Harold Widom. Limit theorems for height fluctuations in a class of discrete space and time growth models. *J. Statist. Phys.*, 102(5-6):1085–1132, 2001.
- [GW91] Peter W. Glynn and Ward Whitt. Departures from many queues in series. *Ann. Appl. Probab.*, 1(4):546–572, 1991.
- [Ham19] Alan Hammond. A patchwork quilt sewn from Brownian fabric: regularity of polymer weight profiles in Brownian last passage percolation. *Forum Math. Pi*, 7:e2, 69, 2019.
- [Har85] J. Michael Harrison. *Brownian motion and stochastic flow systems*. Wiley Series in Probability and Mathematical Statistics: Probability and Mathematical Statistics. John Wiley & Sons, Inc., New York, 1985.
- [HN01] C. Douglas Howard and Charles M. Newman. Geodesics and spanning trees for Euclidean first-passage percolation. *Ann. Probab.*, 29(2):577–623, 2001.
- [Hof08] Christopher Hoffman. Geodesics in first passage percolation. *Ann. Appl. Probab.*, 18(5):1944–1969, 2008.
- [HW90] J. M. Harrison and R. J. Williams. On the quasireversibility of a multiclass Brownian service station. *Ann. Probab.*, 18(3):1249–1268, 1990.
- [HW92] J. M. Harrison and R. J. Williams. Brownian models of feedforward queueing networks: quasireversibility and product form solutions. *Ann. Appl. Probab.*, 2(2):263–293, 1992.
- [JRS19] Christopher Janjigian, Firas Rassoul-Agha, and Timo Seppäläinen. Geometry of geodesics through Busemann measures in directed last-passage percolation. *arXiv e-prints*, page arXiv:1908.09040, 2019.
- [LN96] Cristina Licea and Charles M. Newman. Geodesics in two-dimensional first-passage percolation. *Ann. Probab.*, 24(1):399–410, 1996.
- [MP10] Peter Mörters and Yuval Peres. *Brownian motion*, volume 30 of *Cambridge Series in Statistical and Probabilistic Mathematics*. Cambridge University Press, Cambridge, 2010. With an appendix by Oded Schramm and Wendelin Werner.
- [MQR16] Konstantin Matetski, Jeremy Quastel, and Daniel Remenik. The KPZ fixed point. *arXiv e-prints*, page arXiv:1701.00018, 2016.
- [New95] Charles M. Newman. A surface view of first-passage percolation. In *Proceedings of the International Congress of Mathematicians, Vol. 1, 2 (Zürich, 1994)*, pages 1017–1023. Birkhäuser, Basel, 1995.
- [OY01] Neil O’Connell and Marc Yor. Brownian analogues of Burke’s theorem. *Stochastic Process. Appl.*, 96(2):285–304, 2001.
- [OY02] Neil O’Connell and Marc Yor. A representation for non-colliding random walks. *Electron. Comm. Probab.*, 7:1–12, 2002.
- [Pim16] Leandro P. R. Pimentel. Duality between coalescence times and exit points in last-passage percolation models. *Ann. Probab.*, 44(5):3187–3206, 2016.
- [Pit75] J. W. Pitman. One-dimensional Brownian motion and the three-dimensional Bessel process. *Advances in Appl. Probability*, 7(3):511–526, 1975.
- [QS20] Jeremy Quastel and Sourav Sarkar. Convergence of exclusion processes and KPZ equation to the KPZ fixed point. *arXiv e-prints*, page arXiv:2008.06584, 2020.
- [Sep18] Timo Seppäläinen. The corner growth model with exponential weights. In *Random growth models*, volume 75 of *Proc. Sympos. Appl. Math.*, pages 133–201. Amer. Math. Soc., Providence, RI, 2018.
- [Sep20] Timo Seppäläinen. Existence, uniqueness and coalescence of directed planar geodesics: proof via the increment-stationary growth process. *Ann. Inst. Henri Poincaré Probab. Stat.*, 56(3):1775–1791, 2020.
- [SN01] Paavo Salminen and Ilkka Norros. On busy periods of the unbounded Brownian storage. *Queueing Syst.*, 39(4):317–333 (2002), 2001.
- [SS20] Timo Seppäläinen and Xiao Shen. Coalescence estimates for the corner growth model with exponential weights. *Electron. J. Probab.*, 25:Paper No. 85, 31, 2020.

- [SV10] Timo Seppäläinen and Benedek Valkó. Bounds for scaling exponents for a $1 + 1$ dimensional directed polymer in a Brownian environment. *ALEA Lat. Am. J. Probab. Math. Stat.*, 7:451–476, 2010.
- [Tay55] S. J. Taylor. The α -dimensional measure of the graph and set of zeros of a Brownian path. *Proc. Cambridge Philos. Soc.*, 51:265–274, 1955.
- [Vír20] Bálint Vírág. The heat and the landscape I. *arXiv e-prints*, page arXiv:2008.07241, 2020.
- [WW98] Jan Wehr and Jung Woo. Absence of geodesics in first-passage percolation on a half-plane. *Ann. Probab.*, 26(1):358–367, 1998.

TIMO SEPPÄLÄINEN, UNIVERSITY OF WISCONSIN-MADISON, MATHEMATICS DEPARTMENT, VAN VLECK HALL, 480 LINCOLN DR., MADISON WI 53706-1388, USA.

Email address: `seppalai@math.wisc.edu`

EVAN SORENSEN, UNIVERSITY OF WISCONSIN-MADISON, MATHEMATICS DEPARTMENT, VAN VLECK HALL, 480 LINCOLN DR., MADISON WI 53706-1388, USA.

Email address: `elsorensen@wisc.edu`