

On the law of the iterated logarithm under the sub-linear expectations

LI-XIN ZHANG¹

School of Mathematical Sciences, Zhejiang University, Hangzhou 310027

(Email:stazlx@zju.edu.cn)

Abstract: In this paper, we establish some general forms of the law of the iterated logarithm for independent random variables in a sub-linear expectation space, where the random variables are not necessarily identically distributed. Exponential inequalities for the maximum sum of independent random variables and Kolmogorov's converse exponential inequalities are established as tools for showing the law of the iterated logarithm. As an application, the sufficient and necessary conditions of the law of iterated logarithm for independent and identically distributed random variables under the sub-linear expectation are obtained.

Keywords: sub-linear expectation, capacity, Kolmogorov's exponential inequality, laws of the iterated logarithm

AMS 2020 subject classifications: 60F15, 60F05

1 Introduction and notations.

Let $\{X_n; n \geq 1\}$ be a sequence of independent random variables on a probability space $(\Omega, \mathcal{F}, \mathbb{P})$. Denote $S_n = \sum_{i=1}^n X_i$, $s_n^2 = \sum_{i=1}^n \mathbb{E}X_i^2$, $\log x = \ln \max(e, x)$, where \mathbb{E} is the expectation with respect to \mathbb{P} . The almost sure limit behavior of $\{S_n/\sqrt{2s_n^2 \log \log s_n^2}; n \geq 1\}$ has been studied extensively. It is known, under some conditions, that

$$\mathbb{P} \left(\limsup_{n \rightarrow \infty} \frac{|S_n|}{\sqrt{2s_n^2 \log \log s_n^2}} = 1 \right) = 1. \quad (1.1)$$

This is the "law of the the iterated logarithm" (LIL). In his famous paper, Wittmann (1985) established a general theorem for LIL which states that (1.1) holds if the following conditions are fulfilled:

$$\mathbb{E}X_n = 0 \text{ and } \mathbb{E}X_n^2 < \infty, \quad n \geq 1, \quad (1.2)$$

$$\sum_{n=1}^{\infty} \frac{\mathbb{E}|X_n|^p}{(2s_n^2 \log \log s_n^2)^{p/2}} < \infty \text{ for some } 2 < p \leq 3, \quad (1.3)$$

¹This work was Supported by grants from the NSF of China (Grant No.11731012,12031005), Ten Thousands Talents Plan of Zhejiang Province (Grant No. 2018R52042) and the Fundamental Research Funds for the Central Universities

[or $\mathbb{E}X_n^3 = 0$, $n \geq 1$, and (1.3) holds for some $3 < p \leq 4$],

$$\lim_{n \rightarrow \infty} s_n^2 = \infty \quad \text{and} \quad \limsup_{n \rightarrow \infty} \frac{s_{n+1}^2}{s_n^2} < \infty. \quad (1.4)$$

According to Wittmann, the classical result of Hartman and Wintner (1941) is just a corollary of his theorem. That is, if $\{X_n; n \geq 1\}$ is a sequence of independent and identically distributed (i.i.d.) random variables, then

$$\mathbb{P} \left(\limsup_{n \rightarrow \infty} \frac{|S_n|}{\sqrt{2n \log \log n}} = (\mathbb{E}X_1^2)^{1/2} \right) = 1 \quad (1.5)$$

if

$$\mathbb{E}X_1 = 0 \quad \text{and} \quad \mathbb{E}X_1^2 < \infty. \quad (1.6)$$

Wittmann (1987) showed that his theorem also holds when $p > 3$. Chen (1993) extended Wittmann's theorem to the case of random variables taking their values in a Banach space and weakened the condition (1.3) to that for every $\epsilon > 0$ there exists $p > 2$ such that

$$\sum_{n=1}^{\infty} \frac{\mathbb{E}|X_n|^p I\{|X_n| \geq \epsilon \sqrt{s_n^2 / \log \log s_n^2}\}}{(2s_n^2 \log \log s_n^2)^{p/2}} < \infty. \quad (1.7)$$

In this paper, we consider the random variables in a sub-linear expectation space. Let $\{X_n; n \geq 1\}$ be sequence of independent random variables in a sub-linear expectation space $(\Omega, \mathcal{H}, \widehat{\mathbb{E}})$ with a related upper capacity \mathbb{V} . Chen and Hu (2014) showed that, if $\{X_n; n \geq 1\}$ is a sequence of i.i.d. random variables, then

$$\mathbb{V} \left(\limsup_{n \rightarrow \infty} \frac{|S_n|}{\sqrt{2n \log \log n}} > (\widehat{\mathbb{E}}X_1^2)^{1/2} \right) = 0 \quad (1.8)$$

if

$$\widehat{\mathbb{E}}[X_1] = \widehat{\mathbb{E}}[-X_1], \quad (1.9)$$

$$X_1, X_2, \dots, \text{ are bounded random variables.} \quad (1.10)$$

Zhang (2016) showed that (1.8) holds if (1.9), and

$$\lim_{c \rightarrow \infty} \widehat{\mathbb{E}}[(X_1^2 - c)^+] = 0, \quad (1.11)$$

$$\int_0^{\infty} \mathbb{V}(|X_1|^2 / \log \log |X_1| \geq x) dx < \infty. \quad (1.12)$$

It is obvious that (1.11) and (1.12) are much weaker than (1.10), and together with (1.8) are very close to Hartman and Wintner's condition (1.6). Zhang also showed that (1.12) is necessary for (1.8) to hold. Nevertheless two important questions remained unanswered:

1. Is (1.11) a sharpest condition? It is known that Hartman and Wintner's condition (1.6) is also necessary for (1.6) to hold (cf. Strassen (1966)). What are the sufficient and necessary conditions for (1.8) to hold? It should be noted that, unlike in the classical probability space, $\widehat{\mathbb{E}}[X_1^2] < \infty$ and (1.11) are not equivalent and neither of them implies (1.12).
2. Does Wittmann's theorem also hold under the sub-linear expectation?

A big difficulty for showing the necessity of a kind of the condition (1.11) for (1.8) is that the symmetry argument is not valid under the sub-linear expectation. As for Wittmann's LIL, beside we have not enough powerful exponential inequalities, a difficulty is that we can not use the truncation argument under the sub-linear expectation as freely as under the classical expectation because, if a random variable X is partitioned to $X_1 + X_2$, the sub-linear expectation $\widehat{\mathbb{E}}[X]$ is no longer $\widehat{\mathbb{E}}[X_1] + \widehat{\mathbb{E}}[X_2]$. The purpose of this paper is to establish LIL for independent random variables under the sub-linear expectation, where the random variables are not necessarily identically-distributed. As a corollary, we obtain the sufficient and necessary conditions of the LIL for i.i.d. random variables. The theorems on the LIL are given in Section 5. Before that, some notations under the sub-linear expectation are given in the next section, the main tools are established in Section 3, including exponential inequalities for the maximum sum of independent random variables and Kolmogorov's converse exponential inequality, and some properties of the capacities are given in Section 4 where as a corollary, it is showed that the G -capacity is not continuous, and a Broel-Cantelli lemma is established for a capacity which is not necessarily continuous. The proofs of the LILs are given in the last section. The limit theorems are established without any assumption on the continuity of the capacities.

2 Basic settings

We use the framework and notations of Peng (2008, 2019). Let (Ω, \mathcal{F}) be a given measurable space and let \mathcal{H} be a linear space of real functions defined on (Ω, \mathcal{F}) such that if $X_1, \dots, X_n \in \mathcal{H}$ then $\varphi(X_1, \dots, X_n) \in \mathcal{H}$ for each $\varphi \in C_{l,Lip}(\mathbb{R}_n)$, where $C_{l,Lip}(\mathbb{R}_n)$ denotes the linear space of (local Lipschitz) functions φ satisfying

$$|\varphi(\mathbf{x}) - \varphi(\mathbf{y})| \leq C(1 + |\mathbf{x}|^m + |\mathbf{y}|^m)|\mathbf{x} - \mathbf{y}|, \quad \forall \mathbf{x}, \mathbf{y} \in \mathbb{R}_n,$$

for some $C > 0, m \in \mathbb{N}$ depending on φ .

\mathcal{H} is considered as a space of “random variables”. In this case we denote $X \in \mathcal{H}$. We also denote $C_{b,Lip}(\mathbb{R}_n)$ the space of bounded Lipschitz functions.

Definition 2.1 A sub-linear expectation $\widehat{\mathbb{E}}$ on \mathcal{H} is a function $\widehat{\mathbb{E}} : \mathcal{H} \rightarrow \overline{\mathbb{R}}$ satisfying the following properties: for all $X, Y \in \mathcal{H}$, we have

- (a) *Monotonicity:* If $X \geq Y$ then $\widehat{\mathbb{E}}[X] \geq \widehat{\mathbb{E}}[Y]$;
- (b) *Constant preserving:* $\widehat{\mathbb{E}}[c] = c$;
- (c) *Sub-additivity:* $\widehat{\mathbb{E}}[X+Y] \leq \widehat{\mathbb{E}}[X] + \widehat{\mathbb{E}}[Y]$ whenever $\widehat{\mathbb{E}}[X] + \widehat{\mathbb{E}}[Y]$ is not of the form $+\infty - \infty$ or $-\infty + \infty$;
- (d) *Positive homogeneity:* $\widehat{\mathbb{E}}[\lambda X] = \lambda \widehat{\mathbb{E}}[X]$, $\lambda \geq 0$.

Here $\overline{\mathbb{R}} = [-\infty, \infty]$. The triple $(\Omega, \mathcal{H}, \widehat{\mathbb{E}})$ is called a sub-linear expectation space. Give a sub-linear expectation $\widehat{\mathbb{E}}$, let us denote the conjugate expectation $\widehat{\mathcal{E}}$ of $\widehat{\mathbb{E}}$ by

$$\widehat{\mathcal{E}}[X] := -\widehat{\mathbb{E}}[-X], \quad \forall X \in \mathcal{H}.$$

From the definition, it is easily shown that $\widehat{\mathcal{E}}[X] \leq \widehat{\mathbb{E}}[X]$, $\widehat{\mathbb{E}}[X+c] = \widehat{\mathbb{E}}[X] + c$ and $\widehat{\mathbb{E}}[X-Y] \geq \widehat{\mathbb{E}}[X] - \widehat{\mathbb{E}}[Y]$ for all $X, Y \in \mathcal{H}$ with $\widehat{\mathbb{E}}[Y]$ being finite. Further, if $\widehat{\mathbb{E}}[|X|]$ is finite, then $\widehat{\mathcal{E}}[X]$ and $\widehat{\mathbb{E}}[X]$ are both finite. By Theorem 1.2.1 of Peng (2019), there exists a family of finite additive linear expectations $E_\theta : \mathcal{H} \rightarrow \overline{\mathbb{R}}$ indexed by $\theta \in \Theta$, such that

$$\widehat{\mathbb{E}}[X] = \max_{\theta \in \Theta} E_\theta[X] \quad \text{for } X \in \mathcal{H}. \quad (2.1)$$

Moreover, for each $X \in \mathcal{H}$, there exists $\theta_X \in \Theta$ such that $\widehat{\mathbb{E}}[X] = E_{\theta_X}[X]$ if $\widehat{\mathbb{E}}[X]$ is finite.

Definition 2.2 (See Peng (2008, 2019))

- (i) (Identical distribution) Let \mathbf{X}_1 and \mathbf{X}_2 be two n -dimensional random vectors defined respectively in sub-linear expectation spaces $(\Omega_1, \mathcal{H}_1, \widehat{\mathbb{E}}_1)$ and $(\Omega_2, \mathcal{H}_2, \widehat{\mathbb{E}}_2)$. They are called identically distributed, denoted by $\mathbf{X}_1 \stackrel{d}{=} \mathbf{X}_2$ if

$$\widehat{\mathbb{E}}_1[\varphi(\mathbf{X}_1)] = \widehat{\mathbb{E}}_2[\varphi(\mathbf{X}_2)], \quad \forall \varphi \in C_{b,Lip}(\mathbb{R}_n).$$

A sequence $\{X_n; n \geq 1\}$ of random variables is said to be identically distributed if $X_i \stackrel{d}{=} X_1$ for each $i \geq 1$.

- (ii) (Independence) In a sub-linear expectation space $(\Omega, \mathcal{H}, \widehat{\mathbb{E}})$, a random vector $\mathbf{Y} = (Y_1, \dots, Y_n)$, $Y_i \in \mathcal{H}$ is said to be independent to another random vector $\mathbf{X} = (X_1, \dots, X_m)$, $X_i \in \mathcal{H}$ under $\widehat{\mathbb{E}}$ if for each test function $\varphi \in C_{l,Lip}(\mathbb{R}_m \times \mathbb{R}_n)$ we have $\widehat{\mathbb{E}}[\varphi(\mathbf{X}, \mathbf{Y})] = \widehat{\mathbb{E}}[\widehat{\mathbb{E}}[\varphi(\mathbf{x}, \mathbf{Y})] |_{\mathbf{x}=\mathbf{X}}]$, whenever $\overline{\varphi}(\mathbf{x}) := \widehat{\mathbb{E}}[|\varphi(\mathbf{x}, \mathbf{Y})|] < \infty$ for all \mathbf{x} and $\widehat{\mathbb{E}}[|\overline{\varphi}(\mathbf{X})|] < \infty$.

(iii) (Independent random variables) A sequence of random variables $\{X_n; n \geq 1\}$ is said to be independent, if X_{i+1} is independent to (X_1, \dots, X_i) for each $i \geq 1$.

It is easily seen that, if $\{X_1, \dots, X_n\}$ are independent, then $\widehat{\mathbb{E}}[\sum_{i=1}^n X_i] = \sum_{i=1}^n \widehat{\mathbb{E}}[X_i]$.

Next, we consider the capacities corresponding to the sub-linear expectations. Let $\mathcal{G} \subset \mathcal{F}$. A function $V : \mathcal{G} \rightarrow [0, 1]$ is called a capacity if

$$V(\emptyset) = 0, V(\Omega) = 1 \text{ and } V(A) \leq V(B) \quad \forall A \subset B, A, B \in \mathcal{G}.$$

It is called to be sub-additive if $V(A \cup B) \leq V(A) + V(B)$ for all $A, B \in \mathcal{G}$ with $A \cup B \in \mathcal{G}$.

Let $(\Omega, \mathcal{H}, \widehat{\mathbb{E}})$ be a sub-linear expectation space. We denote $(\mathbb{V}, \mathcal{V})$ be a pair of capacities with the properties that

$$\widehat{\mathbb{E}}[f] \leq \mathbb{V}(A) \leq \widehat{\mathbb{E}}[g] \quad \text{if } f \leq I_A \leq g, f, g \in \mathcal{H} \text{ and } A \in \mathcal{F}, \quad (2.2)$$

\mathbb{V} is sub-additive

and $\mathcal{V}(A) := 1 - \mathbb{V}(A^c)$, $A \in \mathcal{F}$. It is obvious that

$$\mathcal{V}(A \cup B) \leq \mathcal{V}(A) + \mathbb{V}(B). \quad (2.3)$$

We call \mathbb{V} and \mathcal{V} the upper and the lower capacity, respectively. In general, we can choose \mathbb{V} as

$$\mathbb{V}(A) := \inf\{\widehat{\mathbb{E}}[\xi] : I_A \leq \xi, \xi \in \mathcal{H}\}, \quad \forall A \in \mathcal{F}, \quad (2.4)$$

To distinguish this capacity with others, we denote it by $\widehat{\mathbb{V}}$, and $\widehat{\mathcal{V}}(A) = 1 - \widehat{\mathbb{V}}(A)$. $\widehat{\mathbb{V}}$ is the largest capacity satisfying (2.2).

When there exists a family of probability measure on (Ω, \mathcal{F}) such that

$$\widehat{\mathbb{E}}[X] = \sup_{P \in \mathcal{P}} P[X] =: \sup_{P \in \mathcal{P}} \int X dP, \quad (2.5)$$

\mathbb{V} can be defined as

$$\mathbb{V}(A) = \sup_{P \in \mathcal{P}} P(A). \quad (2.6)$$

Also, we define the Choquet integrals/expectations $(C_{\mathbb{V}}, C_{\mathcal{V}})$ by

$$C_V[X] = \int_0^\infty V(X \geq t) dt + \int_{-\infty}^0 [V(X \geq t) - 1] dt$$

with V being replaced by \mathbb{V} and \mathcal{V} respectively. If \mathbb{V}_1 and \mathbb{V}_2 are two capacities having the property (2.2), then for any random variable $X \in \mathcal{H}$,

$$\mathbb{V}_1(X \geq x + \epsilon) \leq \mathbb{V}_2(X \geq x) \leq \mathbb{V}_1(X \geq x - \epsilon) \quad \text{for all } \epsilon > 0 \text{ and } x. \quad (2.7)$$

In fact, let $f, g \in C_{b,Lip}(R)$ such that $I\{y \geq x + \epsilon\} \leq f(y) \leq I\{y \geq x\} \leq g(y) \leq I\{y \geq x - \epsilon\}$.

Then

$$\mathbb{V}_1(X \geq x + \epsilon) \leq \widehat{\mathbb{E}}[f(X)] \leq \mathbb{V}_2(X \geq x) \leq \widehat{\mathbb{E}}[g(X)] \leq \mathbb{V}_1(X \geq x - \epsilon).$$

It follows from (2.7) that

$$\mathbb{V}_1(X \geq x) = \mathbb{V}_2(X \geq x) \quad \text{for all but except countable many } x$$

and so

$$C_{\mathbb{V}_1}(X) = C_{\mathbb{V}_2}(X). \quad (2.8)$$

Finally, for real numbers x and y , denote $x \vee y = \max(x, y)$, $x \wedge y = \min(x, y)$, $x^+ = \max(0, x)$, $x^- = \max(0, -x)$ and $\log x = \ln \max(e, x)$. For a random variable X , because $XI\{|X| \leq c\}$ may be not in \mathcal{H} , we will truncate it in the form $(-c) \vee X \wedge c$ denoted by $X^{(c)}$.

3 Exponential inequalities

Exponential inequalities and Kolmogorov's converse exponential inequality are basic tools for established the LIL. In this section, we give the exponential inequalities under both the upper capacity \mathbb{V} and the lower capacity \mathcal{V} and Kolmogorov's converse exponential inequalities under the upper capacity \mathbb{V} . The next lemma gives the Kolmogorov type exponential inequalities for maximum sums of independent random variables.

Lemma 3.1 *Let $\{X_1, \dots, X_n\}$ be a sequence of independent random variables in the sub-linear expectation space $(\Omega, \mathcal{H}, \widehat{\mathbb{E}})$. Set $S_n = \sum_{i=1}^n X_i$, $B_n^2 = \sum_{i=1}^n \widehat{\mathbb{E}}[X_i^2]$, $b_n^2 = \sum_{i=1}^n \widehat{\mathcal{E}}[X_i^2]$, and $A_n(p, y) = \sum_{i=1}^n \widehat{\mathbb{E}}[(X_i^+ \wedge y)^p]$, $p \geq 2$. Denote*

$$B_{n,y}^2 = \sum_{i=1}^n \widehat{\mathbb{E}}[(X_i \wedge y)^2], \quad b_{n,y}^2 = \sum_{i=1}^n \widehat{\mathcal{E}}[(X_i \wedge y)^2], \quad y > 0.$$

(I) For all $x, y > 0$,

$$\begin{aligned} & \mathbb{V}\left(\max_{k \leq n}(S_k - \widehat{\mathbb{E}}[S_k]) \geq x\right) \quad \left(\text{resp. } \mathcal{V}\left(\max_{k \leq n}(S_k - \widehat{\mathcal{E}}[S_k]) \geq x\right)\right) \\ & \leq \mathbb{V}\left(\max_{k \leq n} X_k > y\right) + \exp\left\{-\frac{x^2}{2(xy + B_{n,y}^2)}\left(1 + \frac{2}{3} \ln\left(1 + \frac{xy}{B_{n,y}^2}\right)\right)\right\}; \end{aligned} \quad (3.1)$$

and for all $p \geq 2$, $x, y > 0$ and $0 < \delta \leq 1$,

$$\begin{aligned} & \mathbb{V}\left(\max_{k \leq n}(S_k - \widehat{\mathbb{E}}[S_k]) \geq x\right) \quad \left(\text{resp. } \mathcal{V}\left(\max_{k \leq n}(S_k - \widehat{\mathcal{E}}[S_k]) \geq x\right)\right) \\ & \leq \mathbb{V}\left(\max_{i \leq n} X_i > y\right) + 2 \exp\{p^p\} \left\{\frac{A_n(p, y)}{y^p}\right\}^{\frac{\delta x}{10y}} + \exp\left\{-\frac{x^2}{2(1 + \delta)B_{n,y}^2}\right\}; \end{aligned} \quad (3.2)$$

(II) For all $x, y > 0$,

$$\begin{aligned} & \mathcal{V}\left(\max_{k \leq n}(S_k - \widehat{\mathbb{E}}[S_k]) \geq x\right) \\ & \leq \mathbb{V}\left(\max_{k \leq n} X_k > y\right) + \exp\left\{-\frac{x^2}{2(xy + b_{n,y}^2)}\left(1 + \frac{2}{3} \ln\left(1 + \frac{xy}{b_{n,y}^2}\right)\right)\right\}; \end{aligned} \quad (3.3)$$

and for all $p \geq 2$, $x, y > 0$ and $0 < \delta \leq 1$,

$$\begin{aligned} & \mathcal{V}\left(\max_{k \leq n}(S_k - \widehat{\mathbb{E}}[S_k]) \geq x\right) \\ & \leq \mathbb{V}\left(\max_{i \leq n} X_i > y\right) + 2 \exp\{p^p\} \left\{\frac{A_n(p, y)}{y^p}\right\}^{\frac{\delta x}{10y}} + \exp\left\{-\frac{x^2}{2(1 + \delta)b_{n,y}^2}\right\}. \end{aligned} \quad (3.4)$$

Further, the upper bounds in (3.2) and (3.4) can be replaced respectively, by

$$\begin{aligned} & C_p \delta^{-p} \frac{1}{x^p} \sum_{i=1}^n \widehat{\mathbb{E}}[(X_i^+)^p] + \exp\left\{-\frac{x^2}{2(1 + \delta)B_n^2}\right\}, \\ & C_p \delta^{-p} \frac{1}{x^p} \sum_{i=1}^n \widehat{\mathbb{E}}[(X_i^+)^p] + \exp\left\{-\frac{x^2}{2(1 + \delta)b_n^2}\right\}. \end{aligned} \quad (3.5)$$

Remark 3.1 (3.2) and (3.4) are Fuk and Nagaev (1971)'s type inequalities.

Proof. The upper bound in (3.1) for $\mathbb{V}(S_n - \widehat{\mathbb{E}}[S_n] \geq x)$ and $\mathcal{V}(S_n - \widehat{\mathcal{E}}[S_n] \geq x)$ are derived by Zhang (2016). Here, we consider the maximum sums. First, we give the proof of (3.3) and (3.4). Let $Y_k = X_k \wedge y$, $T_n = \sum_{i=1}^n (Y_i - \widehat{\mathbb{E}}[X_i])$. Then $X_k - Y_k = (X_k - y)^+ \geq 0$ and $\widehat{\mathbb{E}}[Y_k] \leq \widehat{\mathbb{E}}[X_k]$. From the fact $\mathcal{V}(A \cup B) \leq \mathcal{V}(A) + \mathbb{V}(B)$ it follows that

$$\mathcal{V}\left(\max_{k \leq n}(S_k - \widehat{\mathbb{E}}[S_k]) \geq x\right) \leq \mathbb{V}\left(\max_{k \leq n} X_k > y\right) + \mathcal{V}\left(\max_{k \leq n} T_k \geq x\right),$$

and for any $t > 0$, $\varphi(x) =: e^{t(x \wedge y)}$ is a bounded non-decreasing function and belongs to $C_{b,Lip}(\mathbb{R})$ since $0 \leq \varphi'(x) \leq te^{ty}$. From

$$e^{tY_k} = 1 + tY_k + \frac{e^{tY_k} - 1 - tY_k}{Y_k^2} Y_k^2 \leq 1 + tY_k + \frac{e^{ty} - 1 - ty}{y^2} Y_k^2$$

and the fact $\widehat{\mathcal{E}}[X + Y] \leq \widehat{\mathbb{E}}[X] + \widehat{\mathcal{E}}[Y]$, we have

$$\begin{aligned} \widehat{\mathcal{E}}[e^{tY_k}] & \leq 1 + t\widehat{\mathbb{E}}[X_k] + \frac{e^{ty} - 1 - ty}{y^2} \widehat{\mathcal{E}}[Y_k^2] \leq \exp\left\{t\widehat{\mathbb{E}}[X_k] + \frac{e^{ty} - 1 - ty}{y^2} \widehat{\mathcal{E}}[Y_k^2]\right\}, \\ \widehat{\mathcal{E}}[e^{t(T_k - T_{k-1})}] & \leq \exp\left\{\frac{e^{ty} - 1 - ty}{y^2} \widehat{\mathcal{E}}[Y_k^2]\right\}. \end{aligned}$$

Write

$$U_0 = 1, \quad U_k = \exp\left\{-\frac{e^{ty} - 1 - ty}{y^2} b_{k,y}^2\right\} e^{tT_k}, \quad k = 1, \dots, k_n.$$

Then

$$\begin{aligned} & \widehat{\mathcal{E}}[U_k - U_{k-1} | X_1, \dots, X_{k-1}] \\ &= U_{k-1} \widehat{\mathcal{E}} \left[\exp \left\{ -\frac{e^{ty} - 1 - ty}{y^2} \widehat{\mathcal{E}}[Y_k^2] \right\} e^{t(T_k - T_{k-1})} - 1 \right] \leq 0. \end{aligned} \quad (3.6)$$

For any $\alpha > 0$ and given $\beta \in (0, \alpha)$, let $f(x)$ be a continuous function with bounded derivation such that $I\{x \leq \alpha - \beta\} \leq f(x) \leq I\{x < \alpha\}$. Define $f_0 = 1$, $f_k = f(U_1) \cdots f(U_k)$.

Then

$$\begin{aligned} & f_0 U_0 + \sum_{k=1}^n f_{k-1} (U_k - U_{k-1}) = f_n U_n + \sum_{k=1}^n f_{k-1} (1 - f(U_k)) U_k \\ & \geq f_n U_n + \sum_{k=1}^n f_{k-1} (1 - f(U_k)) (\alpha - \beta) = (\alpha - \beta) (1 - f_n) + f_n U_n \\ & \geq (\alpha - \beta) I\{\max_{k \leq n} U_k \geq \alpha\}. \end{aligned}$$

By the independence,

$$\begin{aligned} & \widehat{\mathcal{E}} \left[f_0 U_0 + \sum_{k=1}^n f_{k-1} (U_k - U_{k-1}) \right] \\ &= \widehat{\mathcal{E}} \left[f_0 U_0 + \sum_{k=1}^{n-1} f_{k-1} (U_k - U_{k-1}) + f_{n-1} \widehat{\mathcal{E}}[U_n - U_{n-1} | X_1, \dots, X_{n-1}] \right] \\ & \leq \widehat{\mathcal{E}} \left[f_0 U_0 + \sum_{k=1}^{n-1} f_{k-1} (U_k - U_{k-1}) \right] \leq \cdots \leq \widehat{\mathcal{E}}[f_0 U_0]. \end{aligned}$$

It follows that

$$(\alpha - \beta) \mathcal{V}(\max_{k \leq n} U_k \geq \alpha) \leq \widehat{\mathcal{E}}[f_0 U_0] = \widehat{\mathcal{E}}[U_0].$$

By letting $\beta \rightarrow 0$, we have

$$\mathcal{V} \left(\max_{k \leq n} U_k \geq \alpha \right) \leq \frac{\widehat{\mathcal{E}}[U_0]}{\alpha} = \frac{1}{\alpha}. \quad (3.7)$$

Note

$$\exp \left\{ t \max_{k \leq n} T_k \right\} \leq \max_{k \leq n} U_k \exp \left\{ \frac{e^{ty} - 1 - ty}{y^2} b_{n,y}^2 \right\}.$$

Hence by (3.7),

$$\begin{aligned} \mathcal{V} \left(\max_{k \leq n} T_k \geq x \right) & \leq \mathcal{V} \left(\max_{k \leq n} U_k \geq \exp \left\{ tx - \frac{e^{ty} - 1 - ty}{y^2} b_{n,y}^2 \right\} \right) \\ & \leq \exp \left\{ -tx + \frac{e^{ty} - 1 - ty}{y^2} b_{n,y}^2 \right\}. \end{aligned}$$

Choosing $t = \frac{1}{y} \ln \left(1 + \frac{xy}{b_{n,y}^2} \right)$ yields

$$\mathcal{V} \left(\max_{k \leq n} T_k \geq x \right) \leq \exp \left\{ \frac{x}{y} - \frac{x}{y} \left(\frac{b_{n,y}^2}{xy} + 1 \right) \ln \left(1 + \frac{xy}{b_{n,y}^2} \right) \right\}. \quad (3.8)$$

Applying the elementary inequality

$$\ln(1+t) \geq \frac{t}{1+t} + \frac{t^2}{2(1+t)^2} \left(1 + \frac{2}{3} \ln(1+t)\right)$$

yields

$$\left(\frac{b_{n,y}}{xy} + 1\right) \ln\left(1 + \frac{xy}{b_{n,y}}\right) \geq 1 + \frac{xy}{2(xy + b_{n,y})} \left(1 + \frac{2}{3} \ln\left(1 + \frac{xy}{b_{n,y}}\right)\right).$$

(3.3) is proved.

Next we show (3.4). If $xy \leq \delta b_{n,y}^2$, then

$$\frac{x^2}{2(xy + b_{n,y})} \left(1 + \frac{2}{3} \ln\left(1 + \frac{xy}{b_{n,y}}\right)\right) \geq \frac{x^2}{2(1+\delta)b_{n,y}^2}.$$

If $xy \geq \delta b_{n,y}^2$, then

$$\frac{x^2}{2(xy + b_{n,y}^2)} \left(1 + \frac{2}{3} \ln\left(1 + \frac{xy}{b_{n,y}}\right)\right) \geq \frac{x}{2(1+1/\delta)y}.$$

It follows that

$$\mathbb{V}(T_n \geq x) \leq \exp\left\{-\frac{x^2}{2(1+\delta)b_{n,y}^2}\right\} + \exp\left\{-\frac{x}{2(1+1/\delta)y}\right\} \quad (3.9)$$

by (3.8). For $z > 0$, let

$$\beta(z) = \beta_p(z) = \frac{1}{z^p} \sum_{k=1}^n \widehat{\mathbb{E}}[(X_k^+ \wedge z)^p],$$

and choose

$$\rho = 1 \wedge \frac{1}{(1+\delta) \ln(1/\beta(z))}, \quad y = \rho z.$$

Note $b_{n,y}^2 \leq b_{n,z}^2$. Then by (3.9),

$$\begin{aligned} \mathbb{V}(S_n \geq (1+\delta)x) &\leq \mathbb{V}(T_n \geq x) + \mathbb{V}\left(\sum_{i=1}^n (X_i - y)^+ \geq \delta x\right) \\ &\leq \exp\left\{-\frac{x^2}{2(1+\delta)b_{n,z}^2}\right\} + (\beta(z))^{\delta x/2z} + \mathbb{V}(\max_{i \leq n} X_i > z) + \mathbb{V}\left(\sum_{i=1}^n (X_i - \rho z)^+ \wedge z \geq \delta x\right). \end{aligned}$$

On the other hand, for $t > 0$,

$$\begin{aligned} \mathbb{V}\left(\sum_{i=1}^n (X_i - \rho z)^+ \wedge z \geq \delta x\right) &= \mathbb{V}\left(\sum_{i=1}^n \left[\left(\frac{X_i}{z} - \rho\right)^+ \wedge 1\right] \geq \frac{\delta x}{z}\right) \\ &\leq e^{-t \frac{\delta x}{z}} \widehat{\mathbb{E}} \exp\left\{t \sum_{i=1}^n \left[\left(\frac{X_i}{z} - \rho\right)^+ \wedge 1\right]\right\} = e^{-t \frac{\delta x}{z}} \prod_{i=1}^n \widehat{\mathbb{E}} \exp\left\{t \left[\left(\frac{X_i}{z} - \rho\right)^+ \wedge 1\right]\right\} \\ &\leq e^{-t \frac{\delta x}{z}} \prod_{i=1}^n [1 + e^t \mathbb{V}(X_i \geq \rho z)] \leq \exp\left\{-t \frac{\delta x}{z} + e^t \frac{\beta(z)}{\rho^p}\right\}. \end{aligned}$$

By taking the minimum over $t \geq 0$, it follows that

$$\mathbb{V}\left(\sum_{i=1}^n (X_i - \rho z)^+ \wedge z \geq \delta x\right) \leq \exp\left\{\frac{\delta x}{z}\left(1 - \ln \frac{\delta x}{z} + \ln \frac{\beta(z)}{\rho^p}\right)\right\}.$$

Assume $\beta(z) < 1$. When $\rho = 1/[(1 + \delta) \ln \beta(z)]$, by the fact $\sqrt{x}(\ln \frac{1}{x})^p \leq (2pe^{-1})^p$ we have

$$\begin{aligned} & \frac{\delta x}{z}\left(1 - \ln \frac{\delta x}{z} + \ln \frac{\beta(z)}{\rho^p}\right) \\ &= \frac{\delta x}{z}\left(1 - \ln \frac{\delta x}{z} + \ln \beta(z) + p \ln \ln \frac{1}{\beta(z)} + p \ln(1 + \delta)\right) \\ &\leq \frac{\delta x}{z}\left(-\ln \frac{\delta x}{z} + 1 + p \ln(2(1 + \delta)pe^{-1})\right) + \frac{\delta x}{2z} \ln \beta(z) \\ &\leq (2(1 + \delta)pe^{-1})^p + \frac{\delta x}{2z} \ln \beta(z). \end{aligned}$$

The last inequality is because of $\max_{x \geq 0}\{x(C + 1 - \ln x)\} = e^C$. When $\rho = 1$,

$$\frac{\delta x}{z}\left(1 - \ln \frac{\delta x}{z} + \ln \frac{\beta(z)}{\rho^p}\right) \leq 1 + \frac{\delta x}{z} \ln \beta(z) \leq 1 + \frac{\delta x}{2z} \ln \beta(z).$$

It follows that

$$\begin{aligned} \mathbb{V}(S_n \geq (1 + \delta)x) &\leq \exp\left\{-\frac{x^2}{2(1 + \delta)b_{n,z}^2}\right\} + \mathbb{V}(\max_{i \leq n} X_i > z) \\ &\quad + 2 \exp\{(2(1 + \delta)pe^{-1})^p\} (\beta(z))^{\delta x/2z}. \end{aligned}$$

Therefore,

$$\begin{aligned} \mathbb{V}(S_n \geq x) &\leq \exp\left\{-\frac{x^2}{2(1 + \delta)^3 b_{n,z}^2}\right\} + \mathbb{V}(\max_{i \leq n} X_i > z) \\ &\quad + 2 \exp\{(2(1 + \delta)pe^{-1})^p\} (\beta(z))^{\frac{\delta x}{2(1 + \delta)z}}. \end{aligned}$$

For $0 < \delta' \leq 1$, let $\delta = \sqrt[3]{1 + \delta'} - 1$. Then $\frac{\delta}{1 + \delta} \geq \frac{\delta'}{5}$, $2(1 + \delta) < e$. It follows that

$$\mathbb{V}(S_n \geq x) \leq \exp\left\{-\frac{x^2}{2(1 + \delta')b_{n,z}^2}\right\} + \mathbb{V}(\max_{i \leq n} X_i > z) + 2 \exp\{p^p\} (\beta(z))^{\frac{\delta' x}{10z}}.$$

If $\beta(z) \geq 1$, then the above inequality is obvious. (3.4) is proved.

For (I), it is sufficient to note that

$$\widehat{\mathcal{E}}[e^{tY_k}] \leq 1 + t\widehat{\mathcal{E}}[X_k] + \frac{e^{ty} - 1 - ty\widehat{\mathbb{E}}[Y_k^2]}{y^2},$$

$$\widehat{\mathcal{E}}[e^{t(Y_k - \widehat{\mathcal{E}}[X_k])}] \leq \exp\left\{\frac{e^{ty} - 1 - ty\widehat{\mathbb{E}}[Y_k^2]}{y^2}\right\}$$

and

$$\widehat{\mathbb{E}}[e^{tY_k}] \leq 1 + t\widehat{\mathbb{E}}[X_k] + \frac{e^{ty} - 1 - ty\widehat{\mathbb{E}}[Y_k^2]}{y^2},$$

$$\widehat{\mathbb{E}}[e^{t(Y_k - \widehat{\mathbb{E}}[X_k])}] \leq \exp \left\{ \frac{e^{ty} - 1 - ty \widehat{\mathbb{E}}[Y_k^2]}{y^2} \right\}.$$

For (3.5), it is sufficient to choose $y = \delta x/10$ and note that $b_{n,y}^2 \leq b_n^2$, $B_{n,y}^2 \leq B_n^2$,

$$\mathbb{V}(\max_{i \leq n} X_i > y) \leq \frac{A_n(p, y)}{y^p}, \quad y > 0. \quad \square$$

The following lemma is a kind of analog of Kolmogorov's converse exponential inequality.

Lemma 3.2 *Let $\{X_{n,i}; i = 1, \dots, k_n\}$ be an array of independent random variables in the sub-linear expectation space $(\Omega, \mathcal{H}, \widehat{\mathbb{E}})$ with $s_n^2 = \sum_{i=1}^{k_n} \widehat{\mathbb{E}}[X_{n,i}^2]$. Let $\{x_n\}$ be a sequence of positive numbers with $x_n \rightarrow \infty$ and $x_n/s_n \rightarrow 0$. Suppose*

$$\frac{\sum_{i=1}^{k_n} |\widehat{\mathbb{E}}[X_{n,i}]|}{s_n x_n} \rightarrow 0, \quad \frac{\sum_{i=1}^{k_n} |\widehat{\mathcal{E}}[X_{n,i}]|}{s_n x_n} \rightarrow 0$$

and there exists a positive number α such that

$$|X_{n,i}| \leq \alpha \frac{s_n}{x_n}, \quad i = 1, \dots, k_n.$$

Then for any $\gamma > 0$, there exists a positive constant $\pi(\gamma)$ (small enough) such that

$$\liminf_{n \rightarrow \infty} x_n^{-2} \ln \mathbb{V} \left(\sum_{i=1}^{k_n} X_{n,i} \geq z s_n x_n \right) \geq -\frac{z^2}{2} (1 + \gamma) \text{ for all } 0 < z\alpha \leq \pi(\gamma). \quad (3.10)$$

Proof. We use an analogue argument of Stout (1974), cf. Petrov (1995, Page 241-243).

First, it is easily seen that

$$\frac{1}{s_n^2} \sum_{i=1}^{k_n} \left| \widehat{\mathbb{E}}[(X_{n,i} - \widehat{\mathbb{E}}[X_{n,i}])^2] - \widehat{\mathbb{E}}[X_{n,i}^2] \right| \leq \frac{3\alpha}{s_n x_n} \sum_{i=1}^{k_n} |\widehat{\mathbb{E}}[X_{n,i}]| \rightarrow 0.$$

Without loss of generality, we can assume that $\widehat{\mathbb{E}}[X_{n,i}] = 0$, $i = 1, \dots, k_n$. For otherwise, we consider $X_{n,i} - \widehat{\mathbb{E}}[X_{n,i}]$ instead.

Let $S_n = \sum_{i=1}^{k_n} X_{n,i}$ and $q_n(y) = \mathbb{V}(S_n \geq y s_n x_n)$. Then by (3.1),

$$q_n(y) \leq \exp \left\{ -\frac{y^2 x_n^2}{2(y\alpha + 1)} \right\}. \quad (3.11)$$

For any $t > 0$ with $t\alpha < 1/32$, we have

$$\begin{aligned} \exp\{t X_{n,i} \frac{x_n}{s_n}\} &\geq 1 + t X_{n,i} \frac{x_n}{s_n} + \frac{t^2}{2} X_{n,i}^2 \frac{x_n^2}{s_n^2} \left(1 - \sum_{i=3}^{\infty} \frac{2(t\alpha)^{i-2}}{i!} \right) \\ &\geq 1 + t X_{n,i} \frac{x_n}{s_n} + \frac{t^2}{2} X_{n,i}^2 \frac{x_n^2}{s_n^2} (1 - t\alpha/2). \end{aligned}$$

Then from the fact $\widehat{\mathbb{E}}[X + Y] \geq \widehat{\mathcal{E}}[X] + \widehat{\mathbb{E}}[Y]$ it follows that

$$\widehat{\mathbb{E}} \left[\exp\{t X_{n,i} \frac{x_n}{s_n}\} \right] \geq 1 + t \frac{x_n}{s_n} \widehat{\mathcal{E}}[X_{n,i}] + \frac{t^2}{2} \frac{x_n^2}{s_n^2} (1 - t\alpha/2) \widehat{\mathbb{E}}[X_{n,i}^2].$$

Applying $\ln(1+x) \geq x - x^2$ ($x \geq -1/4$) yields

$$\begin{aligned} \ln \widehat{\mathbb{E}} \left[\exp\left\{tX_{n,i} \frac{x_n}{s_n}\right\} \right] &\geq t \frac{x_n}{s_n} \widehat{\mathcal{E}}[X_{n,i}] + \frac{t^2 x_n^2}{2 s_n^2} (1 - t\alpha/2) \widehat{\mathbb{E}}[X_{n,i}^2] \\ &\quad - \left(t \frac{x_n}{s_n} |\widehat{\mathcal{E}}[X_{n,i}]| (t\alpha + (t\alpha)^2) + \frac{t^2 x_n^2}{2 s_n^2} \widehat{\mathbb{E}}[X_{n,i}^2] (1 - t\alpha/2)^2 \frac{(t\alpha)^2}{2} \right). \end{aligned}$$

It follows that

$$\begin{aligned} x_n^{-2} \ln \widehat{\mathbb{E}} \left[\exp\left\{tS_n \frac{x_n}{s_n}\right\} \right] &\geq -t \frac{\sum_{i=1}^{k_n} |\widehat{\mathcal{E}}[X_{n,i}]|}{s_n x_n} (1+2) + \frac{t^2}{2} (1-t\alpha) \\ &\rightarrow \frac{t^2}{2} (1-t\alpha). \end{aligned}$$

Note

$$\widehat{\mathbb{E}} \left[\exp\left\{tS_n \frac{x_n}{s_n}\right\} \right] \leq C_V \left(\exp\left\{tS_n \frac{x_n}{s_n}\right\} \right) = \int_{-\infty}^{\infty} \mathbb{V} \left(\exp\left\{tS_n \frac{x_n}{s_n}\right\} > y \right) dy$$

It follows that

$$\liminf_{n \rightarrow \infty} x_n^{-2} \ln C_V \left(\exp\left\{tS_n \frac{x_n}{s_n}\right\} \right) \geq \frac{t^2}{2} (1-t\alpha) \quad \text{for all } 0 < t\alpha < 1/32. \quad (3.12)$$

Now, for $\delta < 1/4$, let $t = z/(1-\delta)$. Then

$$\begin{aligned} C_V \left(\exp\left\{tS_n \frac{x_n}{s_n}\right\} \right) &= \int_{-\infty}^{\infty} t x_n^2 e^{t x_n^2 y} q_n(y) dy \\ &= \left(\int_{-\infty}^0 + \int_0^{t(1-\delta)} + \int_{t(1-\delta)}^{t(1+\delta)} + \int_{t(1+\delta)}^{8t} + \int_{8t}^{\infty} \right) t x_n^2 e^{t x_n^2 y} q_n(y) dy \\ &=: I_1 + I_2 + I_3 + I_4 + I_5. \end{aligned} \quad (3.13)$$

It is obvious that

$$I_3 \leq 2t^2 x_n^2 \exp\{t^2 x_n^2 (1+\delta)\} q_n(t(1-\delta)) = 2t^2 x_n^2 \exp\{t^2 x_n^2 (1+\delta)\} q_n(z) \quad (3.14)$$

and

$$I_1 \leq \int_{-\infty}^0 t x_n^2 e^{t x_n^2 y} dy \leq 1. \quad (3.15)$$

Assume that $8t\alpha \leq 1$. by (3.11), if $y\alpha \geq 1$, then

$$e^{t x_n^2 y} q_n(y) \leq \exp \left\{ t x_n^2 y - \frac{y x_n^2}{4\alpha} \right\} \leq e^{-t x_n^2 y},$$

and, if $8t \leq y \leq 1/\alpha$,

$$e^{t x_n^2 y} q_n(y) \leq \exp \left\{ t x_n^2 y - \frac{y^2 x_n^2}{4} \right\} \leq e^{-t x_n^2 y}.$$

It follows that

$$I_5 \leq \int_{8t}^{\infty} t x_n^2 e^{-t x_n^2 y} dy \leq 1, \quad 8t\alpha \leq 1. \quad (3.16)$$

Now, consider I_2 and I_4 . Choose a positive constant β . Then if $y\alpha \leq \beta < \delta$, then

$$q_n(y) \leq \exp \left\{ -\frac{y^2 x_n^2}{2(1+\beta)} \right\}, \quad \text{if } y \leq 8t \text{ and } 8t\alpha \leq \beta.$$

Let $\psi(y) = ty - \frac{y^2}{2(1+\beta)}$. Thus we arrive the inequality

$$I_2 + I_4 \leq t x_n^2 \int_D e^{\psi(y) x_n^2} dy \quad \text{with } D = (0, t(1-\delta)) \cup (t(1+\delta), 8t).$$

The function $\psi(y)$ has a maximum at the point $y = t(1+\beta)$ which lies in the interval $(t(1-\delta), t(1+\delta))$. Therefore,

$$\begin{aligned} \sup_{y \in D} \psi(y) &= \max\{\psi(t(1-\delta)), \psi(t(1+\delta))\} \\ &= \frac{t^2}{2} \left(1 - \delta^2 + (1+\delta)^2 \frac{\beta}{1+\beta} \right) \leq \frac{t^2}{2} \left(1 - \frac{\delta^2}{2} - \frac{\delta^2}{2} \frac{\beta}{1+\beta} \right), \end{aligned}$$

if $\beta = \delta^2/(2(1+\delta)^2)$. It follows that

$$I_2 + I_4 \leq 8t^2 x_n^2 \exp \left\{ \max_{\in D} \psi(y) x_n^2 \right\} \leq \frac{1}{4} \exp \left\{ \frac{t^2 x_n^2}{2} \left(1 - \frac{\delta^2}{2} \right) \right\} \quad (3.17)$$

for n large enough if $8t\alpha \leq \beta = \delta^2/(2(1+\delta)^2)$. On the other hand, if $t\alpha \leq \beta = \delta^2/2(1+\delta)^2$, it follows from (3.12) that

$$C_{\mathbb{V}} \left(\exp \left\{ t S_n \frac{x_n}{s_n} \right\} \right) \geq \exp \left\{ \frac{t^2 x_n^2}{2} \left(1 - \frac{\delta^2}{2} \right) \right\} \geq 8 \quad (3.18)$$

for n large enough. It follows from (3.14)-(3.18) that, for $0 < t\alpha \leq \delta^2/(16(1+\delta)^2)$,

$$I_1 + I_2 + I_4 + I_5 \leq \frac{1}{2} C_{\mathbb{V}} \left(\exp \left\{ t S_n \frac{x_n}{s_n} \right\} \right), \quad (3.19)$$

and therefore,

$$2t^2 x_n^2 \exp \{ t^2 x_n^2 (1+\delta) \} q_n(z) \geq I_3 \geq \frac{1}{2} C_{\mathbb{V}} \left(\exp \left\{ t S_n \frac{x_n}{s_n} \right\} \right) \geq \frac{1}{2} \exp \left\{ \frac{t^2 x_n^2}{2} \left(1 - \frac{\delta^2}{2} \right) \right\} \quad (3.20)$$

when n is large enough. It follows that

$$\begin{aligned} \liminf_{n \rightarrow \infty} x_n^{-2} \ln q_n(z) &\geq \frac{t^2}{2} \left(1 - \frac{\delta^2}{2} \right) - t^2(1+\delta) \\ &> -\frac{z^2}{2} \frac{(1+\delta)^2}{(1-\delta)^2}, \quad \text{if } 0 < z\alpha < \delta^2/(16(1+\delta)^2), \delta < 1/4. \end{aligned}$$

At last, for every $\gamma > 0$, choose $0 < \delta < 1/4$ such that $\frac{(1+\delta)^2}{(1-\delta)^2} \leq 1 + \gamma$. Then (3.10) holds with $\pi(\gamma) = \delta^2/(16(1+\delta)^2)$. The proof is completed. \square

We conjecture that for the lower capacity \mathcal{V} , we have an analogue Kolmogorov's converse exponential inequality.

Conjecture Let $\{X_{n,i}; i = 1, \dots, k_n\}$ be an array of independent random variables in the sub-linear expectation space $(\Omega, \mathcal{H}, \widehat{\mathbb{E}})$ with $\underline{\sigma}_n^2 = \sum_{i=1}^{k_n} \widehat{\mathcal{E}}[X_{n,i}^2]$. Let x_n be a sequence of positive numbers with $x_n \rightarrow \infty$ and $x_n/\underline{\sigma}_n \rightarrow 0$. Suppose

$$\frac{\sum_{i=1}^{k_n} |\widehat{\mathbb{E}}[X_{n,i}]|}{\underline{\sigma}_n x_n} \rightarrow 0, \quad \frac{\sum_{i=1}^{k_n} |\widehat{\mathcal{E}}[X_{n,i}]|}{\underline{\sigma}_n x_n} \rightarrow 0$$

and there exists a positive number α such that

$$|X_{n,i}| \leq \alpha \underline{\sigma}_n / x_n, \quad i = 1, \dots, k_n.$$

Then for any $\gamma > 0$, there exists a positive constant $\pi(\gamma)$ (small enough) such that

$$\liminf_{n \rightarrow \infty} x_n^{-2} \ln \mathcal{V} \left(\sum_{i=1}^{k_n} X_{n,i} \geq z \underline{\sigma}_n x_n \right) \geq -\frac{z^2}{2} (1 + \gamma) \text{ for all } 0 < z\alpha \leq \pi(\gamma). \quad (3.21)$$

It seems that it is not an easy task to obtain the lower bound of the tail capacity under \mathcal{V} . Recently, Peng, Yang and Yao (2020) and Peng and Zhou (2020) studied the tail behavior of the G-normal distribution through analyzing a nonlinear heat equation. Let $\xi \sim N(0, [\underline{\sigma}^2, \overline{\sigma}])$ in sub-linear expectation space $(\widetilde{\Omega}, \widetilde{\mathcal{H}}, \widetilde{\mathbb{E}})$. By Corollary 1 of Peng and Zhou (2020) we have

$$\begin{aligned} \widetilde{\mathbb{V}}(\xi > x) &= \frac{2}{\underline{\sigma} + \overline{\sigma}} \int_x^\infty (\phi(z/\overline{\sigma})I\{z \geq 0\} + \phi(z/\underline{\sigma})I\{z < 0\}) dz \\ &= \begin{cases} \frac{2\overline{\sigma}}{\underline{\sigma} + \overline{\sigma}} \left[1 - \Phi\left(\frac{x}{\overline{\sigma}}\right) \right], & x \geq 0, \\ 1 - \frac{2\overline{\sigma}}{\underline{\sigma} + \overline{\sigma}} \Phi\left(\frac{x}{\underline{\sigma}}\right), & x \leq 0, \end{cases} \end{aligned}$$

where $\Phi(x)$ and $\phi(x)$ are respectively the distribution function and density of a standard normal random variables in a probability space. Hence, by the fact that $-\xi \stackrel{d}{=} \xi$,

$$\begin{aligned} \widetilde{\mathbb{V}}(\xi \geq x) &= \widetilde{\mathbb{V}}(\xi \leq -x) = 1 - \widetilde{\mathbb{V}}(\xi > -x) \\ &= \begin{cases} \frac{2\overline{\sigma}}{\underline{\sigma} + \overline{\sigma}} \left[1 - \Phi\left(\frac{x}{\underline{\sigma}}\right) \right], & x \geq 0, \\ 1 - \frac{2\overline{\sigma}}{\underline{\sigma} + \overline{\sigma}} \Phi\left(\frac{x}{\overline{\sigma}}\right), & x \leq 0. \end{cases} \end{aligned} \quad (3.22)$$

From (3.22) and the central limit theorem, we can derive a lower bound of an exponential inequality under \mathcal{V} for independent and identically distributed random variables.

Lemma 3.3 Suppose $\{X_{ni}; i = 1, \dots, k_n\}$ is an array of independent and identically distributed random variables in the sub-linear expectation space $(\Omega, \mathcal{H}, \widehat{\mathbb{E}})$ with

$$\widehat{\mathbb{E}}[X_{n1}^2] \rightarrow \overline{\sigma}^2 < \infty, \quad \widehat{\mathcal{E}}[X_{n1}^2] \rightarrow \underline{\sigma}^2 > 0.$$

Let $\{x_n\}$ be a sequence of positive numbers such that $x_n \rightarrow \infty$, $x_n/\sqrt{k_n} \rightarrow 0$. Assume

$$\widehat{\mathbb{E}}[(X_{n1}^2 - \epsilon k_n/x_n^2)^+] \rightarrow 0 \text{ for all } \epsilon > 0,$$

and

$$\frac{\sum_{i=1}^{k_n} (|\widehat{\mathbb{E}}[X_{ni}]| + |\widehat{\mathcal{E}}[X_{ni}]|)}{x_n \sqrt{k_n}} = \frac{\sqrt{k_n} (|\widehat{\mathbb{E}}[X_{n1}]| + |\widehat{\mathcal{E}}[X_{n1}]|)}{x_n} \rightarrow 0.$$

Denote $S_n = \sum_{i=1}^{k_n} X_{ni}$. Then for any $z > 0$,

$$\liminf_{n \rightarrow \infty} x_n^{-2} \ln \mathcal{V} \left(S_n \geq z \underline{\sigma} x_n \sqrt{k_n} \right) \geq -\frac{z^2}{2}. \quad (3.23)$$

Proof. Denote $S_{n,0} = 0$, $S_{n,k} = \sum_{i=1}^k X_{ni}$. For $t > 2$, let

$$N = [k_n t^2 / x_n^2], \quad m = [x_n^2 / t^2]; \quad r = \sqrt{k_n} x_n / (tm).$$

Then $mN \leq k_n$, $r \sim \sqrt{N}$ and

$$\begin{aligned} & \left\{ \frac{S_n}{\underline{\sigma} x_n \sqrt{k_n}} \geq z \right\} \supset \left\{ \frac{S_{n,Nm}}{\underline{\sigma} x_n \sqrt{k_n}} \geq z + \epsilon/2 \right\} \cap \left\{ \left| \frac{S_n - S_{n,Nm}}{\underline{\sigma} x_n \sqrt{k_n}} \right| \leq \epsilon/2 \right\} \\ & = \left\{ \frac{S_{n,Nm}}{r \underline{\sigma}} \geq tm(z + \epsilon/2) \right\} \cap \left\{ \left| \frac{S_n - S_{n,Nm}}{\underline{\sigma} x_n \sqrt{k_n}} \right| \leq \epsilon/2 \right\} \\ & \supset \bigcap_{i=1}^m \left\{ \frac{S_{n,Ni} - S_{n,N(i-1)}}{tr \underline{\sigma}} \geq z + \epsilon/2 \right\} \cap \left\{ \left| \frac{S_n - S_{n,Nm}}{\underline{\sigma} x_n \sqrt{k_n}} \right| \leq \epsilon/2 \right\}. \end{aligned}$$

For given $z > 0$ and $\epsilon > 0$. Let $f, g \in C_{b,Lip}(R)$ such that $I\{x \geq z + \epsilon/2\} \geq f(x) \geq I\{x \geq z + \epsilon\}$ and $I\{|x| \leq \epsilon/2\} \geq g(x) \geq I\{|x| \leq \epsilon/4\}$. It follows that

$$I \left\{ \frac{S_n}{\underline{\sigma} x_n \sqrt{k_n}} \geq z + \epsilon \right\} \geq \prod_{i=1}^m f \left(\frac{S_{n,Ni} - S_{n,N(i-1)}}{tr \underline{\sigma}} \right) g \left(\frac{S_n - S_{n,Nm}}{\underline{\sigma} x_n \sqrt{k_n}} \right).$$

Note that $\{S_{n,Ni} - S_{n,N(i-1)}, i = 1, \dots, m, S_n - S_{n,Nm}\}$ are independent under $\widehat{\mathbb{E}}$ (and $\widehat{\mathcal{E}}$).

By (2.2), we have

$$\mathcal{V} \left(\frac{S_n}{\underline{\sigma} x_n \sqrt{k_n}} \geq z + \epsilon \right) \geq \left(\widehat{\mathcal{E}} \left[f \left(\frac{S_{n,N}}{tr \underline{\sigma}} \right) \right] \right)^m \widehat{\mathcal{E}} \left[g \left(\frac{S_n - S_{n,Nm}}{\underline{\sigma} x_n \sqrt{k_n}} \right) \right].$$

Note

$$\frac{\sum_{i=1}^N \widehat{\mathbb{E}}[(X_{ni}^2 - \epsilon N)^+]}{N} \rightarrow 0, \quad \frac{\sum_{i=1}^N (|\widehat{\mathbb{E}}[X_{ni}]| + |\widehat{\mathcal{E}}[X_{ni}]|)}{\sqrt{N}} \rightarrow 0.$$

By applying the Lindeberg limit theorem of Zhang (2021), we have

$$\lim_{n \rightarrow \infty} \widehat{\mathbb{E}} \left[\varphi \left(\frac{S_{n,N}}{r} \right) \right] = \lim_{n \rightarrow \infty} \widehat{\mathbb{E}} \left[\varphi \left(\frac{S_{n,N}}{\sqrt{N}} \right) \right] = \widetilde{\mathbb{E}}[\varphi(\xi)], \quad \text{for all } \varphi \in C_{b,Lip}(R),$$

where $\xi \sim N(0, [\underline{\sigma}^2, \overline{\sigma}^2])$ under $\widetilde{\mathbb{E}}$. It follows that

$$\begin{aligned} & \lim_{n \rightarrow \infty} \widehat{\mathcal{E}} \left[f \left(\frac{S_{n,N}}{tr \underline{\sigma}} \right) \right] = \widetilde{\mathcal{E}} \left[f \left(\frac{\xi}{t \underline{\sigma}} \right) \right] \\ & \geq \widetilde{\mathcal{V}} \left(\xi \geq t(z + \epsilon) \underline{\sigma} \right) = \frac{2 \underline{\sigma}}{\underline{\sigma} + \overline{\sigma}} \left[1 - \Phi \left(tz(1 + \epsilon) \right) \right], \end{aligned}$$

by (3.22). On the other hand,

$$1 - \widehat{\mathcal{E}} \left[g \left(\frac{S_n - S_{n,Nm}}{\underline{\sigma} x_n \sqrt{k_n}} \right) \right] \leq \mathbb{V} \left(\frac{|S_{n,k_n} - Nm|}{\underline{\sigma} x_n \sqrt{k_n}} \geq \epsilon/4 \right) \leq \frac{C \widehat{\mathbb{E}}[X_{n1}^2] k_n - Nm}{\epsilon^2 \underline{\sigma}^2 k_n x_n^2} \rightarrow 0.$$

It follows that

$$\begin{aligned} & \liminf_{n \rightarrow \infty} x_n^{-2} \ln \mathcal{V} \left(\frac{S_n}{\underline{\sigma} x_n \sqrt{k_n}} \geq z \right) \\ & \geq \liminf_{t \rightarrow \infty} \liminf_{n \rightarrow \infty} t^{-2} m^{-1} \ln \mathcal{V} \left(\frac{S_n}{\underline{\sigma} x_n \sqrt{k_n}} \geq z + \epsilon \right) \\ & \geq \liminf_{t \rightarrow \infty} \liminf_{n \rightarrow \infty} t^{-2} \ln \widehat{\mathcal{E}} \left[f \left(\frac{S_{n,N}}{tr \underline{\sigma}} \right) \right] = \liminf_{t \rightarrow \infty} t^{-2} \ln \widetilde{\mathcal{E}} \left[f \left(\frac{\xi}{t \underline{\sigma}} \right) \right] \\ & \geq \liminf_{t \rightarrow \infty} t^{-2} \ln \left[1 - \Phi \left(tz(1 + \epsilon) \right) \right] = -\frac{(z(1 + \epsilon))^2}{2}. \end{aligned}$$

The proof is completed. \square

4 Properties of the Capacities

Before we give the laws of the iterated logarithm, we need some more notations and properties about capacities.

Definition 4.1 (I) A function $V : \mathcal{F} \rightarrow [0, 1]$ is called to be countably sub-additive if

$$V \left(\bigcup_{n=1}^{\infty} A_n \right) \leq \sum_{n=1}^{\infty} V(A_n) \quad \forall A_n \in \mathcal{F}.$$

(II) A capacity $V : \mathcal{F} \rightarrow [0, 1]$ is called to be continuous from below if it satisfies that $V(A_n) \uparrow V(A)$ whenever $A_n \uparrow A$, where $A_n, A \in \mathcal{F}$, and, it is called to be continuous from above if it satisfies that $V(A_n) \downarrow V(A)$ whenever $A_n \downarrow A$, where $A_n, A \in \mathcal{F}$.

It is obvious that the continuity from above with the sub-additivity implies the continuity from below, and the continuity from the below with the sub-additivity implies the countable sub-additivity. So, we call a sub-additive capacity to be continuous if it is continuous from above. Also, if V is a capacity continuous from above, then

$$V \left(\bigcap_{i=1}^{\infty} A_i \right) = 1 \text{ for events } \{A_n\} \text{ with } A_n \supset A_{n+1} \text{ and } V(A_n) = 1, n = 1, 2, \dots \quad (4.1)$$

It is obvious that the lower capacity \mathcal{V} has the property (4.1) when the upper one \mathbb{V} is countably sub-additive.

The following lemma is the Borel-Cantelli Lemma and its converse under capacities.

Lemma 4.1 (i) Let $\{A_n, n \geq 1\}$ be a sequence of events in \mathcal{F} . Suppose $\sum_{n=1}^{\infty} V(A_n) < \infty$.

Then

$$\lim_{N \geq n \rightarrow \infty} V\left(\bigcup_{i=n}^N A_i\right) = 0.$$

If V is a countably sub-additive capacity, then

$$V(A_n \text{ i.o.}) = 0, \quad \text{where } \{A_n \text{ i.o.}\} = \bigcap_{n=1}^{\infty} \bigcup_{i=n}^{\infty} A_i. \quad (4.2)$$

(ii) Suppose $\{\xi_n; n \geq 1\}$ is a sequence of independent random variables in $(\Omega, \mathcal{H}, \widehat{\mathbb{E}})$. Suppose $\sum_{n=1}^{\infty} \mathbb{V}(\{\xi_n \geq 1 + \epsilon\}) = \infty$ for some $\epsilon > 0$. Then

$$\mathbb{V}\left(\bigcup_{m=n}^{\infty} \{\xi_m \geq 1\}\right) \geq \mathbb{V}\left(\bigcup_{m=n}^N \{\xi_m \geq 1\}\right) \rightarrow 1 \text{ as } N \rightarrow \infty, \quad (4.3)$$

and

$$\mathbb{V}(\{\xi_n \geq 1\} \text{ i.o.}) = 1 \quad \text{if } \mathbb{V} \text{ has the property (4.1).}$$

(iii) Suppose $\{\xi_n; n \geq 1\}$ is a sequence of independent random variables in $(\Omega, \mathcal{H}, \widehat{\mathbb{E}})$ with a countably sub-additive \mathbb{V} . Then

$$\mathcal{V}(\{\xi_n \geq 1\} \text{ i.o.}) = 1 \quad \text{if } \sum_{n=1}^{\infty} \mathcal{V}(\{\xi_n \geq 1 + \epsilon\}) = \infty \text{ for some } \epsilon > 0. \quad (4.4)$$

(iv) Suppose $\{\xi_n; n \geq 1\}$ is a sequence of independent random variables in $(\Omega, \mathcal{H}, \widehat{\mathbb{E}})$. Suppose $\sum_{n=1}^{\infty} \mathcal{V}(\{\xi_n \geq 1 - \epsilon\}) < \infty$ for some $\epsilon > 0$. Then

$$\lim_{N \geq n \rightarrow \infty} \mathcal{V}\left(\bigcup_{m=n}^N \{\xi_m \geq 1\}\right) \rightarrow 0, \quad (4.5)$$

and

$$\mathcal{V}(\{\xi_n \geq 1\} \text{ i.o.}) = 0 \quad \text{if } \mathbb{V} \text{ is continuous.} \quad (4.6)$$

Lemma 4.1 (i) (resp. (iv)) is the direct part of the Borel-Cantelli Lemma for \mathbb{V} (resp. \mathcal{V}).

(ii) or (iii) are the the converse ones.

Proof. (i) is trivial. For (ii), denote $A_n = \{\xi_n \geq 1\}$. Let $g(x)$ be a Lipschitz function

with $I\{x \geq 1 + \epsilon\} \leq g(x) \leq I\{x \geq 1\}$. Then

$$\begin{aligned}
\mathcal{V} \left(\bigcap_{i=n}^{\infty} A_i^c \right) &\leq \mathcal{V} \left(\bigcap_{i=n}^N A_i^c \right) \leq \widehat{\mathcal{E}} \left[\prod_{i=n}^N (1 - g(\xi_i)) \right] \\
&= \prod_{i=n}^N \widehat{\mathcal{E}} [(1 - g(\xi_i))] = \prod_{i=n}^N (1 - \widehat{\mathbb{E}}[g(\xi_i)]) \\
&\leq \exp \left\{ - \sum_{i=n}^N \widehat{\mathbb{E}}[g(\xi_i)] \right\} \leq \exp \left\{ - \sum_{i=n}^N \mathbb{V}(\xi_i \geq 1 + \epsilon) \right\} \\
&\rightarrow \exp \left\{ - \sum_{i=n}^{\infty} \mathbb{V}(\xi_i \geq 1 + \epsilon) \right\} = 0 \quad \text{if} \quad \sum_{n=1}^{\infty} \mathbb{V}(\{\xi_n \geq 1 + \epsilon\}) = \infty.
\end{aligned} \tag{4.7}$$

That is $\mathbb{V}(\bigcup_{i=n}^{\infty} A_i) = 1$ and $\mathbb{V}(\bigcup_{i=n}^N A_i) \rightarrow 1$ as $N \rightarrow \infty$.

For (iii), similar to (4.7) we have

$$\mathbb{V} \left(\bigcap_{i=n}^{\infty} A_i^c \right) \leq \exp \left\{ - \sum_{i=n}^{\infty} \mathcal{V}(\xi_i \geq 1 + \epsilon) \right\} = 0 \quad \text{if} \quad \sum_{n=1}^{\infty} \mathcal{V}(\{\xi_n \geq 1 + \epsilon\}) = \infty.$$

It follows from the countable sub-additivity of \mathbb{V} that

$$\mathbb{V}(\{A_n \text{ i.o.}\}^c) \leq \sum_{n=1}^{\infty} \mathbb{V} \left(\bigcap_{i=n}^{\infty} A_i^c \right) = 0.$$

Therefore, $\mathcal{V}(A_n \text{ i.o.}) = 1$.

For (iv), we let $g(x)$ be a Lipschitz function with $I\{x \geq 1\} \leq g(x) \leq I\{x \geq 1 - \epsilon\}$. Suppose $\sum_{n=1}^{\infty} \mathcal{V}(\xi_i \geq 1 - \epsilon) < \infty$. Then

$$\begin{aligned}
\mathbb{V} \left(\bigcap_{i=n}^N A_i^c \right) &\geq \widehat{\mathbb{E}} \left[\prod_{i=n}^N (1 - g(\xi_i)) \right] = \prod_{i=n}^N \widehat{\mathbb{E}} [(1 - g(\xi_i))] = \prod_{i=n}^N (1 - \widehat{\mathcal{E}}[g(\xi_i)]) \\
&\geq \prod_{i=n}^N (1 - \mathcal{V}(\xi_i \geq 1 - \epsilon)) \geq \exp \left\{ -2 \sum_{i=n}^N \mathcal{V}(\xi_i \geq 1 - \epsilon) \right\}
\end{aligned}$$

for N, n large enough. The last inequality is computed from the fact that $1 - x \geq e^{-2x}$ for all $x \leq 1/2$. (4.5) is proved. If \mathbb{V} is continuous, then

$$\mathbb{V} \left(\bigcup_{n=1}^{\infty} \bigcap_{i=n}^{\infty} A_i^c \right) \geq \lim_{n \rightarrow \infty} \lim_{N \rightarrow \infty} \mathbb{V} \left(\bigcap_{i=n}^N A_i^c \right) \geq \lim_{n \rightarrow \infty} \exp \left\{ -2 \sum_{i=n}^{\infty} \mathcal{V}(\xi_i \geq 1 - \epsilon) \right\} = 1.$$

Therefore, the proof is completed. \square .

Proposition 4.1 *Let $(\Omega, \mathcal{H}, \widehat{\mathbb{E}})$ be a sub-linear expectation space with a sequence of independent and identically distributed random variables $\{X_n; n \geq 1\}$. Consider the subspace*

$$\widetilde{\mathcal{H}} = \left\{ Y = \varphi(X_1, X_2, \dots, X_n) : \varphi \in C_{l, Lip}(\mathbb{R}_n), \widehat{\mathbb{E}}[(|Y| - c)^+] \rightarrow 0, n \geq 1 \right\}. \tag{4.8}$$

If \mathbb{V} is continuous, then $\widehat{\mathbb{E}}$ is linear on $\widetilde{\mathcal{H}}$.

Proof. It is sufficient to show that

$$\widehat{\mathbb{E}}[Y] = \widehat{\mathbb{E}}[-Y] \quad \text{for all } Y \in \widetilde{\mathcal{H}}. \quad (4.9)$$

Without loss of generality, assume $Y = \varphi(X_1)$ and $|Y| \leq c$. Denote $Y_n = \varphi(X_n)$. Then $\{Y_n; n \geq 1\}$ is a sequence of independent and identically distributed random variables with $|Y_n| \leq c$. By (3.1),

$$\mathcal{V} \left(\frac{S_m}{m} \leq \widehat{\mathbb{E}}[Y] - \epsilon \right) = \mathcal{V} \left(\sum_{i=1}^m (-Y_i + \widehat{\mathbb{E}}[Y_i]) \geq \epsilon m \right) \leq \exp \left\{ -\frac{\epsilon^2 m^2}{2(\epsilon m c + c^2 m)} \right\} \rightarrow 0.$$

Hence

$$\mathbb{V} \left(\frac{S_m}{m} > \widehat{\mathbb{E}}[Y] - \epsilon \right) \rightarrow 1 \quad \text{for all } \epsilon > 0. \quad (4.10)$$

On the other hand, let $I(k) = \{2^k + 1, \dots, 2^{k+1}\}$. By (3.1) again,

$$\mathcal{V} \left(\max_{n \in I(k)} \sum_{j \in I(k), j \leq n} (Y_j - \widehat{\mathcal{E}}[Y_j]) \geq 2^{k+1} \epsilon \right) \leq \exp \left\{ -\frac{\epsilon^2 2^{2(k+1)}}{2(\epsilon 2^{k+1} c + c^2 2^k)} \right\} \leq \exp \left\{ -\frac{\epsilon^2}{c^2} 2^k \right\}.$$

Let $T_n = \sum_{j=1}^n (Y_j - \widehat{\mathcal{E}}[Y_j])$. Note the independence of the random variables. By Lemma 4.1 (iv), it follows that

$$\lim_{L \geq l \rightarrow \infty} \mathbb{V} \left(\max_{l \leq k \leq L} \max_{n \in I(k)} \frac{T_n - T_{2^k}}{2^{k+1}} < \epsilon \right) = 1 \quad \text{for all } \epsilon > 0,$$

which implies

$$\lim_{N \geq n \rightarrow \infty} \mathbb{V} \left(\max_{n \leq l \leq N} \frac{T_l}{l} < \epsilon \right) = 1 \quad \text{for all } \epsilon > 0.$$

That is

$$\lim_{N \geq n \rightarrow \infty} \mathbb{V} \left(\max_{n \leq l \leq N} \frac{S_l}{l} \leq \widehat{\mathcal{E}}[Y] + \epsilon \right) = 1 \quad \text{for all } \epsilon > 0. \quad (4.11)$$

Let f and g be two Lipschitz functions with $I\{x \leq \epsilon\} \geq f(x) \geq I\{x \leq \epsilon/2\}$ and $I\{x \geq -\epsilon\} \geq g(x) \geq I\{x \geq -\epsilon/2\}$. By the independence of the random variables, it follows from (4.11) that

$$\begin{aligned} & \lim_{N \geq n \rightarrow \infty} \widehat{\mathbb{E}} \left[f \left(\max_{n \leq l \leq N} \frac{S_l - S_m}{l} - \widehat{\mathbb{E}}[Y] \right) \cdot g \left(\frac{S_m}{m} - \widehat{\mathbb{E}}[Y] \right) \right] \\ & \geq \lim_{N \geq n \rightarrow \infty} \widehat{\mathbb{E}} \left[f \left(\max_{n \leq l \leq N} \frac{S_l - S_m}{l} - \widehat{\mathcal{E}}[Y] \right) \right] \cdot \widehat{\mathbb{E}} \left[g \left(\frac{S_m}{m} - \widehat{\mathbb{E}}[Y] \right) \right] \\ & \geq \lim_{N \geq n \rightarrow \infty} \mathbb{V} \left(\max_{n \leq l \leq N} \frac{S_l - S_m}{l} \leq \widehat{\mathcal{E}}[Y] + \epsilon/2 \right) \cdot \mathbb{V} \left(\frac{S_m}{m} \geq \widehat{\mathbb{E}}[Y] - \epsilon/2 \right) \\ & \geq \lim_{N \geq n \rightarrow \infty} \mathbb{V} \left(\max_{n \leq l \leq N} \frac{S_l}{l} \leq \widehat{\mathcal{E}}[Y] + \epsilon/3 \right) \cdot \mathbb{V} \left(\frac{S_m}{m} \geq \widehat{\mathbb{E}}[Y] - \epsilon/2 \right) \\ & = \mathbb{V} \left(\frac{S_m}{m} \geq \widehat{\mathbb{E}}[Y] - \epsilon/2 \right). \end{aligned} \quad (4.12)$$

Up to now, we have not used the continuity or the property (4.1) of \mathbb{V} . Now, note

$$\left\{ \limsup_{n \rightarrow \infty} \frac{S_n}{n} \leq \widehat{\mathcal{E}}[Y] + 2\epsilon \right\} \supset \bigcup_{n=1}^{\infty} \bigcap_{N=n}^{\infty} \left\{ \max_{n \leq l \leq N} \frac{S_l - S_m}{l} \leq \widehat{\mathcal{E}}[Y] + \epsilon \right\}.$$

By the continuity of \mathbb{V} , it follows from (4.12) that

$$\begin{aligned} & \mathbb{V} \left(\limsup_{n \rightarrow \infty} \frac{S_n}{n} \leq \widehat{\mathcal{E}}[Y] + 2\epsilon \text{ and } \frac{S_m}{m} \geq \widehat{\mathbb{E}}[Y] - \epsilon \right) \\ & \geq \lim_{n \rightarrow \infty} \lim_{N \rightarrow \infty} \mathbb{V} \left(\max_{n \leq l \leq N} \frac{S_l - S_m}{l} \leq \widehat{\mathcal{E}}[Y] + \epsilon \text{ and } \frac{S_m}{m} \geq \widehat{\mathbb{E}}[Y] - \epsilon \right) \\ & \geq \lim_{N \geq n \rightarrow \infty} \widehat{\mathbb{E}} \left[f \left(\max_{n \leq l \leq N} \frac{S_l - S_m}{l} - \widehat{\mathcal{E}}[Y] \right) \cdot g \left(\frac{S_m}{m} - \widehat{\mathbb{E}}[Y] \right) \right] \\ & \geq \mathbb{V} \left(\frac{S_m}{m} \geq \widehat{\mathbb{E}}[Y] - \epsilon/2 \right). \end{aligned} \tag{4.13}$$

By letting $m \rightarrow \infty$, it follows from (4.10) that

$$\begin{aligned} & \mathbb{V} \left(\limsup_{n \rightarrow \infty} \frac{S_n}{n} \leq \widehat{\mathcal{E}}[Y] + 2\epsilon \text{ and } \limsup_{m \rightarrow \infty} \frac{S_m}{m} \geq \widehat{\mathbb{E}}[Y] - \epsilon \right) \\ & \geq \limsup_{m \rightarrow \infty} \mathbb{V} \left(\limsup_{n \rightarrow \infty} \frac{S_n}{n} \leq \widehat{\mathcal{E}}[Y] + \epsilon \text{ and } \frac{S_m}{m} \geq \widehat{\mathbb{E}}[Y] - \epsilon \right) = 1. \end{aligned}$$

Therefore, $\widehat{\mathbb{E}}[Y] - \epsilon < \widehat{\mathcal{E}}[Y] + 2\epsilon$ for every $\epsilon > 0$. Hence, (4.9) is verified and the proof is completed. \square .

Let $B(t)$ be a G -Brownian motion. Denote $X_n = \sqrt{n(n+1)} \left(B(1 - 1/(n+1)) - B(1 - 1/n) \right)$. Then X_1, X_2, \dots is a sequence of independent and identically distributed G -normal random variables. Applying Proposition 4.1, we have the following corollary.

Corollary 4.1 *The G -capacity \hat{c} as defined in Section 6.3 of Peng (2019) is not continuous unless $B(t)$ is a classical Brownian motion in a probability space.*

According to Proposition 4.1, the continuity of a sub-additive capacity is a very stringent condition. It is needed to avoid it. Because the Borel-Cantelli lemma (Lemma 4.1 (i)) is needed when the strong limit theorems as the LIL are considered, we usually assume that the capacity \mathbb{V} is countably sub-additive. Such a condition is satisfied when $\widehat{\mathbb{E}}$ can be presented in the form of (2.5) (cf. Chapters 3 and 6 of Peng (2019)). To make the converse part of the Borel-Cantelli lemma (Lemma 4.1 (ii)) valid, it would be reasonable to assume (4.1) instead of the continuity of \mathbb{V} . Unfortunately, the following proposition tells us that (4.1) is also a stringent condition.

Proposition 4.2 *Let $(\Omega, \mathcal{H}, \widehat{\mathbb{E}})$ be a sub-linear expectation space with a sequence of independent and identically distributed random variables $\{X_n; n \geq 1\}$. Consider the subspace $\widetilde{\mathcal{H}}$ defined as in (4.8). Suppose one of the following conditions is satisfied.*

(a) \mathbb{V} is defined as (2.4);

(b) Suppose Ω is a complete separable metric space, each element $X(\omega)$ in \mathcal{H} is a continuous function on Ω , and, the sub-linear expectation $\widehat{\mathbb{E}}$ and the capacity \mathbb{V} are defined as (2.5) and (2.6), i.e.

$$\widehat{\mathbb{E}}[X] = \max_{P \in \mathcal{P}} P[X], \quad \mathbb{V}(A) = \max_{P \in \mathcal{P}} P(A),$$

where \mathcal{P} is a weakly compact family of probability measures on $(\Omega, \mathcal{B}(\Omega))$.

If \mathbb{V} has the property that

$$\mathbb{V}\left(\bigcap_{i=1}^{\infty} A_i\right) > 0 \text{ for events } \{A_n\} \text{ with } A_n \supset A_{n+1} \text{ and } \mathbb{V}(A_n) = 1, n = 1, 2, \dots, \quad (4.14)$$

then $\widehat{\mathbb{E}}$ is linear on $\widetilde{\mathcal{H}}$.

Proof. Let Y, Y_1, Y_2, \dots , be defined as in the proof of Proposition 4.1. It is sufficient to show that $\widehat{\mathbb{E}}[Y] = \widehat{\mathcal{E}}[Y]$. With the same arguments above, (4.10) and (4.12) remain also true. Write $\mathbf{Y} = (Y_1, Y_2, \dots)$.

(a) We first assume that \mathbb{V} is defined as in (2.4). Note the expression (2.1). Consider the linear expectation E_θ on $\mathcal{H}_{1:d} = \{\varphi(Y_1, \dots, Y_d) : \varphi \in C_{b,Lip}(R^d)\}$. Note $|Y_i| \leq c, i = 1, 2, \dots$. It is obvious that, if $C_{l,Lip}(R^d) \ni \varphi_n \downarrow 0$, then

$$|\varphi_n(Y_1, \dots, Y_d)| \leq \sup_{|y_i| \leq c, i=1, \dots, d} |\varphi_n(y_1, \dots, y_d)| \rightarrow 0.$$

So, for each $\theta \in \Theta$, $E_\theta[\varphi_n(Y_1, \dots, Y_d)] \leq \widehat{\mathbb{E}}[\varphi_n(Y_1, \dots, Y_d)] \downarrow 0$. Hence, similar to Lemma 6.2.2 of Peng (2019), by Daniell-Stone's theorem, there is a unique probability P_θ^d on $\sigma(Y_1, \dots, Y_d)$ such that

$$E_\theta[\varphi(Y_1, \dots, Y_d)] = P_\theta^d[\varphi(Y_1, \dots, Y_d)], \quad \forall \varphi \in C_{b,Lip}(R^d), d \geq 1.$$

Then by Kolmogorov's consistent theorem, there exists a unique probability measure P_θ on $\sigma(\mathbf{Y})$ such that

$$P_\theta[\varphi(Y_1, \dots, Y_d)] = P_\theta^d[\varphi(Y_1, \dots, Y_d)] = E_\theta[\varphi(Y_1, \dots, Y_d)], \quad \forall \varphi \in C_{b,Lip}(R^d), d \geq 1. \quad (4.15)$$

It is obvious that the family $\{P_\theta(\mathbf{Y} \in \cdot) : \theta \in \Theta\}$ is tight on R^d because of the boundedness of Y_1, \dots, Y_d , and therefore it is relatively weakly compact. Hence, $\{P_\theta(\mathbf{Y} \in \cdot) : \theta \in \Theta\}$ is relatively weakly compact on R^∞ (cf. Billingsley (1968, Page38)). That is, on the sigma field $\sigma(\mathbf{Y})$, $\{P_\theta : \theta \in \Theta\}$ is relatively weakly compact, and so has a weakly compact closure $\mathcal{Q} = cl(\{P_\theta : \theta \in \Theta\})$. Hence

$$\widehat{\mathbb{E}}[\varphi(Y_1, \dots, Y_d)] = \max_{\theta \in \Theta} P_\theta[\varphi(Y_1, \dots, Y_n)] = \max_{Q \in \mathcal{Q}} Q[\varphi(Y_1, \dots, Y_d)], \quad \forall \varphi \in C_{b,Lip}(R^d),$$

$d \geq 1$. Denote

$$\tilde{V}(\mathbf{Y} \in A) = \sup_{Q \in \mathcal{Q}} Q(\mathbf{Y} \in A), \quad A \in \mathcal{B}(R^\infty).$$

For any $A \in \mathcal{B}(R^d)$ with $f \leq I_A \leq g$, $f, g \in C_{b,Lip}(R^d)$, we have

$$Q[f(Y_1, \dots, Y_d)] \leq Q(\mathbf{Y} \in A) \leq Q[g(Y_1, \dots, Y_d)].$$

Taking the supremum over $Q \in \mathcal{Q}$ yields

$$\widehat{\mathbb{E}}[f(Y_1, \dots, Y_d)] \leq \tilde{V}(\mathbf{Y} \in A) \leq \widehat{\mathbb{E}}[g(Y_1, \dots, Y_d)]. \quad (4.16)$$

Note the weak compactness of \mathcal{Q} . By Lemma 6.1.12 of Peng (2019), for any sequence of closed sets $F_n \downarrow F$, we have $\tilde{V}(\mathbf{Y} \in F_n) \downarrow \tilde{V}(\mathbf{Y} \in F)$. Now, $\{\max_{n \leq l \leq N} \frac{S_l - S_m}{l} \leq \widehat{\mathcal{E}}[Y] + \epsilon$ and $\frac{S_m}{m} \geq \widehat{\mathbb{E}}[Y] - \epsilon\}_{N=n}^\infty$ is a decreasing sequence of closed sets of (Y_1, Y_2, \dots) .

Hence

$$\begin{aligned} & \tilde{V} \left(\max_{l \geq n} \frac{S_l - S_m}{l} \leq \widehat{\mathcal{E}}[Y] + \epsilon \text{ and } \frac{S_m}{m} \geq \widehat{\mathbb{E}}[Y] - \epsilon \right) \\ &= \lim_{N \rightarrow \infty} \tilde{V} \left(\max_{n \leq l \leq N} \frac{S_l - S_m}{l} \leq \widehat{\mathcal{E}}[Y] + \epsilon \text{ and } \frac{S_m}{m} \geq \widehat{\mathbb{E}}[Y] - \epsilon \right) \\ &\geq \lim_{N \rightarrow \infty} \widehat{\mathbb{E}} \left[f \left(\max_{n \leq l \leq N} \frac{S_l - S_m}{l} - \widehat{\mathcal{E}}[Y] \right) \cdot g \left(\frac{S_m}{m} - \widehat{\mathbb{E}}[Y] \right) \right], \end{aligned}$$

where the inequality is due to (4.16). It follows from (4.12) that

$$\begin{aligned} & \tilde{V} \left(\limsup_{n \rightarrow \infty} \frac{S_n}{n} \leq \widehat{\mathcal{E}}[Y] + 2\epsilon \text{ and } \frac{S_m}{m} \geq \widehat{\mathbb{E}}[Y] - \epsilon \right) \\ &\geq \lim_{N \geq n \rightarrow \infty} \widehat{\mathbb{E}} \left[f \left(\max_{n \leq l \leq N} \frac{S_l - S_m}{l} - \widehat{\mathcal{E}}[Y] \right) \cdot g \left(\frac{S_m}{m} - \widehat{\mathbb{E}}[Y] \right) \right] \\ &\geq \mathbb{V} \left(\frac{S_m}{m} \geq \widehat{\mathbb{E}}[Y] - \epsilon/2 \right) \rightarrow 1 \text{ as } m \rightarrow \infty \end{aligned}$$

by (4.10). Note

$$\left\{ \limsup_{n \rightarrow \infty} \frac{S_n}{n} \leq \widehat{\mathcal{E}}[Y] + 2\epsilon \right\} \supset \bigcup_{n=1}^{\infty} \left\{ \max_{l \geq n} \frac{S_l - S_m}{l} \leq \widehat{\mathcal{E}}[Y] + \epsilon \right\}.$$

Therefore,

$$\tilde{V} \left(\limsup_{n \rightarrow \infty} \frac{S_n}{n} \leq \widehat{\mathcal{E}}[Y] + 2\epsilon \text{ and } \bigcup_{m=l}^{\infty} \left\{ \frac{S_m}{m} \geq \widehat{\mathbb{E}}[Y] - \epsilon \right\} \right) = 1 \text{ for all } l \geq 1. \quad (4.17)$$

Write

$$A_l = \left\{ \limsup_{n \rightarrow \infty} \frac{S_n}{n} \leq \widehat{\mathcal{E}}[Y] + 2\epsilon \text{ and } \bigcup_{m=l}^{\infty} \left\{ \frac{S_m}{m} \geq \widehat{\mathbb{E}}[Y] - \epsilon \right\} \right\}.$$

Next, we show that

$$\tilde{V}(A_l) \leq \mathbb{V}(A_l). \quad (4.18)$$

By the definition (2.4) of $\mathbb{V}(A)$, for any $\delta > 0$ there exists a random variable $Z \in \mathcal{H}$ with $0 \leq Z \leq 1$ such that

$$I_{A_l} \leq Z, \quad \widehat{\mathbb{E}}[Z] \leq \mathbb{V}(A_l) + \delta.$$

Consider the sequence Z, Y_1, Y_2, \dots . With the same argument as above, for each $\theta \in \Theta$, there exists a unique probability measure \overline{P}_θ on $\sigma(Z, \mathbf{Y})$ such that

$$E_\theta[\phi(Z, Y_1, \dots, Y_d)] = \overline{P}_\theta[\phi(Z, Y_1, \dots, Y_d)], \quad \forall \phi \in C_{l, Lip}(R^{1+d}), d \geq 1.$$

In particular, by the above equality and (4.15),

$$\overline{P}_\theta[\varphi(Y_1, \dots, Y_d)] = E_\theta[\varphi(Y_1, \dots, Y_d)] = P_\theta[\varphi(Y_1, \dots, Y_d)], \quad \forall \varphi \in C_{l, Lip}(R^d), d \geq 1.$$

By the consistency, $\overline{P}_\theta(\mathbf{Y} \in A^\infty) = P_\theta(\mathbf{Y} \in A^\infty)$ for any Borel set A^∞ of (y_1, y_2, \dots) . On the other hand, for the same reason, the family of probability measures $\{\overline{P}_\theta; \theta \in \Theta\}$ is relatively weakly compact on the sigma field $\sigma(Z, \mathbf{Y})$, and has a weakly compact closure $\overline{\mathcal{Q}}$ such that

$$\widehat{\mathbb{E}}[\varphi(Z, Y_1, \dots, Y_n)] = \max_{\overline{Q} \in \overline{\mathcal{Q}}} \overline{Q}[\varphi(Z, Y_1, \dots, Y_n)], \quad \varphi \in C_{l, Lip}(R^{1+d}), d \geq 1. \quad (4.19)$$

Now, let $Q \in \mathcal{Q}$ such that $\tilde{V}(A_l) - \delta \leq Q(A_l)$. Then there exists a sequence $P_{\theta_i} \implies Q$. For the sequence $\{\overline{P}_{\theta_i}\}$, there exists a subsequence $\overline{P}_{\theta'_i} \implies \overline{Q} \in \overline{\mathcal{Q}}$. Therefore, from $\overline{P}_\theta(\mathbf{Y} \in A^\infty) = P_\theta(\mathbf{Y} \in A^\infty)$ it follows that $\overline{Q}(\mathbf{Y} \in A^\infty) = Q(\mathbf{Y} \in A^\infty)$. Hence, by (4.19),

$$\tilde{V}(A_l) - \delta < Q(A_l) = \overline{Q}(A_l) \leq \overline{Q}[Z] \leq \widehat{\mathbb{E}}[Z] \leq \mathbb{V}(A_l) + \delta.$$

(4.18) is proved. According to (4.18) and (4.17), we have

$$\mathbb{V}(A_l) = 1 \quad \text{for all } l.$$

Therefore, by the property (4.14) we have

$$\mathbb{V} \left(\limsup_{n \rightarrow \infty} \frac{S_n}{n} \leq \widehat{\mathcal{E}}[Y] + 2\epsilon \text{ and } \limsup_{m \rightarrow \infty} \frac{S_m}{m} \geq \widehat{\mathbb{E}}[Y] - \epsilon \right) = \mathbb{V} \left(\bigcap_{l=1}^{\infty} A_l \right) > 0. \quad (4.20)$$

It follows that $\widehat{\mathbb{E}}[Y] - \epsilon \leq \widehat{\mathcal{E}}[Y] + 2\epsilon$ for all $\epsilon > 0$. Therefore, $\widehat{\mathbb{E}}[Y] = \widehat{\mathcal{E}}[Y]$.

(b) Note the weakly compactness of \mathcal{P} . By Lemma 6.1.12 of Peng (2019) again, for any sequence of closed sets $F_n \downarrow F$, we have $\mathbb{V}(F_n) \downarrow \mathbb{V}(F)$. Since, $\left\{ \max_{n \leq l \leq N} \frac{S_l - S_m}{l} \leq \widehat{\mathcal{E}}[Y] + \epsilon \text{ and } \frac{S_m}{m} \geq \widehat{\mathbb{E}}[Y] - \epsilon \right\}_{N=n}^\infty$ is a decreasing sequence of closed sets in Ω . Hence

$$\begin{aligned} & \mathbb{V} \left(\max_{l \geq n} \frac{S_l - S_m}{l} \leq \widehat{\mathcal{E}}[Y] + \epsilon \text{ and } \frac{S_m}{m} \geq \widehat{\mathbb{E}}[Y] - \epsilon \right) \\ &= \lim_{N \rightarrow \infty} \mathbb{V} \left(\max_{n \leq l \leq N} \frac{S_l - S_m}{l} \leq \widehat{\mathcal{E}}[Y] + \epsilon \text{ and } \frac{S_m}{m} \geq \widehat{\mathbb{E}}[Y] - \epsilon \right). \end{aligned}$$

(4.17) remains true for \mathbb{V} . It follows that (4.20) holds by the property (4.14). The proof is completed. \square

Because a capacity \mathbb{V} may be not countably sub-additive and may have not the property (4.1) so that the Borel-Cantelli lemma is not valid, we consider a countably sub-additive extension \mathbb{V}^* which defined by

$$\mathbb{V}^*(A) = \inf \left\{ \sum_{n=1}^{\infty} \widehat{\mathbb{V}}(A_n) : A \subset \bigcup_{n=1}^{\infty} A_n \right\}, \quad \mathcal{V}^*(A) = 1 - \mathbb{V}^*(A^c), \quad A \in \mathcal{F}, \quad (4.21)$$

where $\widehat{\mathbb{V}}$ is defined as in (2.4). As shown in Zhang (2016), \mathbb{V}^* is countably sub-additive, $\mathbb{V}^*(A) \leq \widehat{\mathbb{V}}(A)$ and, $\mathbb{V}^* = \widehat{\mathbb{V}}$ when $\widehat{\mathbb{V}}$ is countably sub-additive. It is shown by Zhang (2016) that, if V is also a countably sub-additive capacity satisfying $V(A) \leq \widehat{\mathbb{E}}[g]$ whenever $I_A \leq g \in \mathcal{H}$, then $V(A) \leq \mathbb{V}^*(A)$. So, if $\widehat{\mathbb{E}}$ has the form (2.5), then

$$\sup_{P \in \mathcal{P}} P(A) \leq \mathbb{V}^*(A).$$

The next lemma tells us that \mathbb{V}^* also has the property (2.2).

Lemma 4.2 *The countably sub-additive capacity \mathbb{V}^* has the property (2.2).*

Proof. Let $f, g \in \mathcal{H}$ with $f \leq I_A \leq g$. The inequality $\mathbb{V}^*(A) \leq \widehat{\mathbb{E}}[g]$ is obvious by the definition. Suppose $A \subset \bigcup_{i=1}^{\infty} A_i$, and $\mathbb{V}^*(A) \geq \sum_{i=1}^{\infty} \widehat{\mathbb{V}}(A_i) - \epsilon/2$. For each i , by the definition (2.4), there exists $g_i \in \mathcal{H}$ such that $I_{A_i} \leq g_i$ and $\widehat{\mathbb{V}}(A_i) \geq \widehat{\mathbb{E}}[g_i] - \epsilon/2^{i+1}$. Without loss of generality, we can assume $0 \leq g_i \leq 1$. It follows that

$$f \leq I_A \leq \sum_{i=1}^{\infty} g_i \quad \text{and} \quad \mathbb{V}^*(A) \geq \sum_{i=1}^{\infty} \widehat{\mathbb{E}}[g_i] - \epsilon.$$

It is sufficient to show that

$$\widehat{\mathbb{E}}[f] \leq \sum_{i=1}^{\infty} \widehat{\mathbb{E}}[g_i]. \quad (4.22)$$

Without loss of generality, we can assume that $0 \leq f \leq 1$. Note the expression (2.1) and the boundedness of f, g_1, g_2, \dots . By the same arguments in the proof of Proposition 4.2 (a), for each $\theta \in \Theta$, there exists a probability measure P_θ on $\sigma(f, g_1, g_2, \dots)$ such that

$$P_\theta[\varphi(f, g_1, \dots, g_d)] = E_\theta[\varphi(f, g_1, \dots, g_d)] \quad \text{for all } \varphi \in C_{l,Lip}(R^{d+1}), \quad d \geq 1.$$

By the countably additivity of the linear mathematical expectation,

$$E_\theta[f] = P_\theta[f] \leq P_\theta\left[\sum_{i=1}^{\infty} g_i\right] = \sum_{i=1}^{\infty} P_\theta[g_i] = \sum_{i=1}^{\infty} E_\theta[g_i].$$

By taking the supremum over $\theta \in \Theta$, we obtain (4.22). \square

Note that \mathbb{V}^* is countably sub-additive and has the property (2.2). (i) and (iii) of Lemma 4.1 are valid for \mathbb{V}^* . With a similar argument as in Proposition 4.2, we can show that \mathbb{V}^* neither is continuous nor has the property (4.14) unless $\widehat{\mathbb{E}}$ is linear. However, the following lemma tells us that the converse part of the Borel-Cantelli lemma for \mathbb{V}^* and the direct part of the Borel-Cantelli lemma for \mathcal{V}^* are still valid even without the assumption (4.1) and the continuity.

Lemma 4.3 *Let $\widehat{\mathbb{V}}$ be defined as in (2.4) and \mathbb{V}^* be defined as in (4.21).*

(i) *Let $\{\xi_n; n \geq 1\}$ be a sequence of independent random variables in $(\Omega, \mathcal{H}, \widehat{\mathbb{E}})$. Suppose $\sum_{n=1}^{\infty} \mathbb{V}^*(\xi_n \geq 1 + \epsilon) = \infty$ or $\sum_{n=1}^{\infty} \widehat{\mathbb{V}}(\xi_n \geq 1 + \epsilon) = \infty$ for some $\epsilon > 0$. Then*

$$\widehat{\mathbb{V}}(\xi_n \geq 1 \text{ i.o.}) = \mathbb{V}^*(\xi_n \geq 1 \text{ i.o.}) = 1.$$

More generally, suppose \mathbb{N}_1 and \mathbb{N}_2 are disjoint subsets of \mathbb{N} , $\sum_{n \in \mathbb{N}_j} \mathbb{V}^(\xi_n \geq 1 + \epsilon) = \infty$ or $\sum_{n \in \mathbb{N}_j} \widehat{\mathbb{V}}(\xi_n \geq 1 + \epsilon) = \infty$ for some $\epsilon > 0$, $j = 1, 2$. Then*

$$\begin{aligned} & \widehat{\mathbb{V}}\left(\{\xi_n \geq 1, n \in \mathbb{N}_1 \text{ i.o.}\} \cap \{\xi_n \geq 1, n \in \mathbb{N}_2 \text{ i.o.}\}\right) \\ &= \mathbb{V}^*\left(\{\xi_n \geq 1, n \in \mathbb{N}_1 \text{ i.o.}\} \cap \{\xi_n \geq 1, n \in \mathbb{N}_2 \text{ i.o.}\}\right) = 1. \end{aligned}$$

(ii) *Let $\{\xi_n; n \geq 1\}$ be a sequence of independent random variables in $(\Omega, \mathcal{H}, \widehat{\mathbb{E}})$. Suppose $\sum_{n=1}^{\infty} \mathcal{V}^*(\xi_n \geq 1 - \epsilon) < \infty$ or $\sum_{n=1}^{\infty} \widehat{\mathcal{V}}(\xi_n \geq 1 - \epsilon) < \infty$ for some $\epsilon > 0$. Then*

$$\widehat{\mathcal{V}}(\xi_n \geq 1 \text{ i.o.}) = \mathcal{V}^*(\xi_n \geq 1 \text{ i.o.}) = 0. \quad (4.23)$$

Proof. Without loss of generality, we assume $0 \leq \xi_i \leq 2$, for otherwise we can replace it by $0 \vee \xi_i \wedge 2$.

(i) Let $g \in C_{b,Lip}(R)$ such that $I\{x \geq 1\} \geq g(x) \geq I\{x \geq 1 + \epsilon\}$. Then

$$\sum_{i \in \mathbb{N}_j} \widehat{\mathbb{E}}[g(\xi_i)] = \infty, \quad j = 1, 2.$$

By the expression (2.1), for each i there exists $\theta_i \in \Theta$ such that

$$E_{\theta_i}[g(\xi_i)] = \widehat{\mathbb{E}}[g(\xi_i)].$$

For the linear operator E_{θ_i} , by noting the boundedness of ξ_i , we have $E_{\theta_i}[\varphi_n(\xi_i)] \searrow 0$ whenever $C_{l,Lip}(R) \ni \varphi_n \searrow 0$. By Daniell-Stone's theorem, there exists a probability P_{θ_i} on $\sigma(\xi_i)$ such that

$$P_{\theta_i}[\varphi(\xi_i)] = E_{\theta_i}[\varphi(\xi_i)] \leq \widehat{\mathbb{E}}[\varphi(\xi_i)], \quad \forall \varphi \in C_{l,Lip}(R). \quad (4.24)$$

Now, we introduce a product probability measure on $\sigma(\xi_1, \xi_2, \dots)$ defined by

$$P = P_{\theta_1} \big|_{\sigma(\xi_1)} \times P_{\theta_2} \big|_{\sigma(\xi_2)} \times \dots.$$

Then, under the probability measure P , ξ_1, ξ_2, \dots is a sequence of independent random variables and,

$$\sum_{i \in \mathbb{N}_j} P(\xi_i \geq 1) \geq \sum_{i \in \mathbb{N}_j} P[g(\xi_i)] = \sum_{i \in \mathbb{N}_j} P_{\theta_i}[g(\xi_i)] = \sum_{i \in \mathbb{N}_j} E_{\theta_i}[g(\xi_i)] = \infty.$$

So, by the Borell-Cantelli lemma for a probability measure,

$$P(\xi_n, n \in \mathbb{N}_j \text{ i.o.}) = 1, \quad j = 1, 2.$$

Note that $\{\xi_n \geq 1, n \in \mathbb{N}_1 \text{ i.o.}\}$ and $\{\xi_n \geq 1, n \in \mathbb{N}_2 \text{ i.o.}\}$ are two independent events under P . It follows that

$$\begin{aligned} & P\left(\{\xi_n \geq 1, n \in \mathbb{N}_1 \text{ i.o.}\} \cap \{\xi_n \geq 1, n \in \mathbb{N}_2 \text{ i.o.}\}\right) \\ &= P(\xi_n \geq 1, n \in \mathbb{N}_1 \text{ i.o.}) \cdot P(\xi_n \geq 1, n \in \mathbb{N}_2 \text{ i.o.}) = 1. \end{aligned}$$

Next, we show that

$$\mathcal{V}^*(B) \leq P(B) \leq \mathbb{V}^*(B) \quad \text{for all } B \in \sigma(\xi_1, \xi_2, \dots). \quad (4.25)$$

Note that for every $\varphi(x_1, \dots, x_d) \in C_{l, Lip}(R^d)$,

$$P[\varphi(x_1, \dots, x_{d-1}, \xi_d)] = P_{\theta_d}[\varphi(x_1, \dots, x_{d-1}, \xi_d)] \leq \widehat{\mathbb{E}}[\varphi(x_1, \dots, x_{d-1}, \xi_d)]$$

and

$$\begin{aligned} & P[\varphi(x_1, \dots, \xi_{d-1}, \xi_d)] = P_{\theta_{d-1}} \left[P_{\theta_d}[\varphi(x_1, \dots, x_{d-1}, \xi_d)] \big|_{x_{d-1}=\xi_{d-1}} \right] \\ & \leq P_{\theta_{d-1}} \left[\widehat{\mathbb{E}}[\varphi(x_1, \dots, x_{d-1}, \xi_d)] \big|_{x_{d-1}=\xi_{d-1}} \right] \leq \widehat{\mathbb{E}} \left[\widehat{\mathbb{E}}[\varphi(x_1, \dots, x_{d-1}, \xi_d)] \big|_{x_{d-1}=\xi_{d-1}} \right] \\ & = \widehat{\mathbb{E}}[\varphi(x_1, \dots, \xi_{d-1}, \xi_d)]. \end{aligned}$$

By induction, we conclude that

$$P[\varphi(\xi_1, \dots, \xi_d)] \leq \widehat{\mathbb{E}}[\varphi(\xi_1, \dots, \xi_d)], \quad \text{for all } \varphi \in C_{l, Lip}(R^d), \quad d \geq 1.$$

Consider a real linear subspace of \mathcal{H} defined as

$$\mathcal{L} = \{\varphi(\xi_1, \dots, \xi_d) : \varphi \in C_{l, Lip}(R^d), d \geq 1\}.$$

Then the linear expectation P on \mathcal{L} is dominated by $\widehat{\mathbb{E}}$. By the Hahn-Banach theorem, there exists a finite additive linear expectation E defined on \mathcal{L} such that $E = P$ on \mathcal{L} and $E \leq \widehat{\mathbb{E}}$ on \mathcal{H} .

By the definition of \mathbb{V}^* , there exists a sequence $0 \leq Z_i \leq 1$ in \mathcal{H} such that

$$I_B \leq \sum_{i=1}^{\infty} Z_i \quad \text{and} \quad \mathbb{V}^*(B) \geq \sum_{i=1}^{\infty} \widehat{\mathbb{E}}[Z_i] - \epsilon.$$

Note that $\xi_1, Z_1, \xi_2, Z_2, \dots$ are bounded random variables. By the same arguments in the proof of Proposition 4.2 (a), there exists a probability measure Q on $\sigma(\xi_1, Z_1, \xi_2, Z_2, \dots)$ such that

$$Q[\varphi(Z_1, \dots, Z_p, \xi_1, \dots, \xi_d)] = E[\varphi(Z_1, \dots, Z_p, \xi_1, \dots, \xi_d)] \quad \text{for all } \varphi \in C_{l,Lip}(R^{p+d}),$$

$p, d \geq 1$. By the consistency, $Q|_{\sigma(\xi_1, \xi_2, \dots)} = P$. It follows that

$$\begin{aligned} P(B) &= Q(B) \leq Q\left[\sum_{i=1}^{\infty} Z_i\right] = \sum_{i=1}^{\infty} Q[Z_i] \\ &= \sum_{i=1}^{\infty} E[Z_i] \leq \sum_{i=1}^{\infty} \widehat{\mathbb{E}}[Z_i] \leq \mathbb{V}^*(B) + \epsilon. \end{aligned}$$

By the arbitrariness of $\epsilon > 0$, we have $\mathbb{V}^*(B) \geq P(B)$. Further, $P(B) = 1 - P(B^c) \geq 1 - \mathbb{V}^*(B^c) = \mathcal{V}^*(B)$. So, (4.25) is proved.

Now, by letting $B = \{\xi_n \geq 1, n \in \mathbb{N}_1 \text{ i.o.}\} \cap \{\xi_n \geq 1, n \in \mathbb{N}_2 \text{ i.o.}\}$, we have $\mathbb{V}^*(B) \geq P(B) = 1$. Note $\widehat{\mathbb{V}}(B) \geq \mathbb{V}^*(B)$. The proof of (i) is completed.

(ii) Let $g \in C_{b,Lip}(R)$ such that $I\{x \geq 1\} \leq g(x) \leq I\{x \geq 1 - \epsilon\}$. Then

$$\sum_{i=1}^{\infty} \widehat{\mathcal{E}}[g(\xi_i)] < \infty.$$

By the expression (2.1), for each i there exists $\theta_i \in \Theta$ such that

$$E_{\theta_i}[-g(\xi_i)] = \widehat{\mathbb{E}}[-g(\xi_i)] \quad \text{i.e.} \quad E_{\theta_i}[g(\xi_i)] = \widehat{\mathcal{E}}[g(\xi_i)].$$

Through E_{θ_i} , $i = 1, 2, \dots$, we define the probability measure P as before. Under P , $\{\xi_n\}$ are independent random variables. Then

$$\sum_{i=1}^{\infty} P(\xi_i \geq 1) \leq \sum_{i=1}^{\infty} P[g(\xi_i)] = \sum_{i=1}^{\infty} E_{\theta_i}[g(\xi_i)] = \sum_{i=1}^{\infty} \widehat{\mathcal{E}}[g(\xi_i)] < \infty,$$

where the first equality is due to (4.24). It follows that $P(\xi_n \geq 1 \text{ i.o.}) = 0$. By (4.25), $\widehat{\mathcal{V}}(\xi_n \geq 1 \text{ i.o.}) \leq \mathcal{V}^*(\xi_n \geq 1 \text{ i.o.}) \leq P(\xi_n \geq 1 \text{ i.o.}) = 0$. The proof is now completed. \square .

Remark 4.1 We conjecture that Lemma 4.3 remains true for the capacity \mathbb{V} defined as in Proposition 4.2 (b). This is true when \mathcal{P} is rich enough, for example, if \mathcal{P} contains all probabilities measures P on $(\Omega, \mathcal{B}(\Omega))$ for which $P \leq \widehat{\mathbb{E}}$.

5 The law of the iterated logarithm

We state the results for the general LIL. The first two theorems describe Wittmann's LIL for independent random variables which are not-necessarily identically distributed. Let $\{X_n; n \geq 1\}$ be a sequence of independent random variables in a sub-linear expectation space $(\Omega, \mathcal{H}, \widehat{\mathbb{E}})$. Denote $s_n^2 = \sum_{k=1}^n \widehat{\mathbb{E}}[X_k^2]$, $t_n = \sqrt{2 \log \log s_n^2}$, $a_n = s_n t_n$,

$$\Gamma_n(p, \alpha) = \widehat{\mathbb{E}}\left[\left(\left(|X_n| - \alpha s_n/t_n\right)^+\right)^p\right], \bar{\Gamma}_n(p, \alpha) = \widehat{\mathbb{E}}\left[\left(\left(|X_n| \wedge a_n - \alpha s_n/t_n\right)^+\right)^p\right]$$

and

$$\Lambda_n(p, \alpha) = \sum_{j=1}^n \widehat{\mathbb{E}}\left[\left(\left(|X_j| - \alpha s_n/t_n\right)^+\right)^p\right], \bar{\Lambda}_n(p, \alpha) = \sum_{j=1}^n \widehat{\mathbb{E}}\left[\left(\left(|X_j| \wedge a_n - \alpha s_n/t_n\right)^+\right)^p\right].$$

Theorem 5.1 *Let $\{X_n; n \geq 1\}$ be a sequence of independent random variables in the sub-linear expectation space $(\Omega, \mathcal{H}, \widehat{\mathbb{E}})$ with $s_n^2 \rightarrow \infty$. Suppose*

$$\sum_{n=1}^{\infty} \mathbb{V}(|X_n| \geq \epsilon a_n) < \infty \text{ for all } \epsilon > 0, \quad (5.1)$$

and, for every $\alpha > 0$ there exist some $p \geq 2$ and $d \geq 0$ such that

$$\sum_{n=1}^{\infty} \frac{\bar{\Gamma}_n(p, \alpha)}{a_n^p} \left(\frac{\bar{\Lambda}_n(p, \alpha)}{a_n^p}\right)^d < \infty. \quad (5.2)$$

Then, for every $\epsilon > 0$,

$$\lim_{N \geq n \rightarrow \infty} \mathbb{V}\left(\inf_{n \leq m \leq N} \frac{S_m - \widehat{\mathcal{E}}[S_m]}{a_m} < -1 - \epsilon \text{ and } \sup_{n \leq m \leq N} \frac{S_m - \widehat{\mathbb{E}}[S_m]}{a_m} > 1 + \epsilon\right) = 0. \quad (5.3)$$

In particular, if \mathbb{V}^* is defined as in (4.21), then

$$\mathbb{V}^*\left(\liminf_{n \rightarrow \infty} \frac{S_n - \widehat{\mathcal{E}}[S_n]}{a_n} < -1 \text{ and } \limsup_{n \rightarrow \infty} \frac{S_n - \widehat{\mathbb{E}}[S_n]}{a_n} > 1\right) = 0. \quad (5.4)$$

Theorem 5.2 *Let $\{X_n; n \geq 1\}$ be a sequence of independent random variables in the sub-linear expectation space $(\Omega, \mathcal{H}, \widehat{\mathbb{E}})$. Suppose (5.1) and, for every $\alpha > 0$ there exist some $p \geq 2$ and $d \geq 0$ such that*

$$\sum_{n=1}^{\infty} \frac{\Gamma_n(p, \alpha)}{a_n^p} \left(\frac{\Lambda_n(p, \alpha)}{a_n^p}\right)^d < \infty, \quad (5.5)$$

and further,

$$\sum_{n=1}^{\infty} s_n^{-2} (\log s_n^2)^{\delta-1} \widehat{\mathbb{E}}[X_n^2] = \infty \text{ for all } \delta > 0, \quad (5.6)$$

$$\frac{\sum_{j=1}^n |\widehat{\mathbb{E}}[X_j]| + \sum_{j=1}^n |\widehat{\mathcal{E}}[X_j]|}{a_n} \rightarrow 0. \quad (5.7)$$

Then, for every $\epsilon > 0$,

$$\lim_{N \geq n \rightarrow \infty} \mathbb{V} \left(\max_{n \leq m \leq N} \frac{|S_m|}{a_m} > 1 + \epsilon \right) = 0, \quad (5.8)$$

$$\lim_{n \rightarrow \infty} \mathbb{V} \left(\max_{m \geq n} \frac{-S_m}{a_m} \geq 1 - \epsilon \right) = \lim_{n \rightarrow \infty} \mathbb{V} \left(\max_{m \geq n} \frac{S_m}{a_m} \geq 1 - \epsilon \right) = 1. \quad (5.9)$$

Further, if \mathbb{V}^* is defined as in (4.21), then

$$\mathbb{V}^* \left(\limsup_{n \rightarrow \infty} \frac{|S_n|}{a_n} > 1 \right) = 0 \quad (5.10)$$

and

$$\mathbb{V}^* \left(\limsup_{n \rightarrow \infty} \frac{S_n}{a_n} = 1 \text{ and } \liminf_{n \rightarrow \infty} \frac{S_n}{a_n} = -1 \right) = 1. \quad (5.11)$$

The following are some remarks on the conditions.

Remark 5.1 When $d = 0$, (5.5) and (5.2) are

$$\sum_{n=1}^{\infty} \frac{\widehat{\mathbb{E}} \left[\left((|X_n| - \alpha s_n / t_n)^+ \right)^p \right]}{a_n^p} < \infty \quad (5.12)$$

and

$$\sum_{n=1}^{\infty} \frac{\widehat{\mathbb{E}} \left[\left((|X_n| \wedge a_n - \alpha s_n / t_n)^+ \right)^p \right]}{a_n^p} < \infty, \quad (5.13)$$

respectively. Further, if n is large enough such that $\epsilon a_n / 2 > \alpha s_n / t_n$, then

$$\frac{\epsilon}{2} I\{|X_n| \geq \epsilon a_n\} \leq \frac{(|X_n| \wedge a_n - \alpha s_n / t_n)^+}{a_n} \leq \frac{(|X_n| - \alpha s_n / t_n)^+}{a_n}.$$

Therefore, (5.12) implies (5.13), and (5.13) implies (5.1). It follows that, if (5.12) or (5.13) is satisfied, then (5.1) can be removed.

By Theorem 5.2 and Remark 5.1, we have the following corollary.

Corollary 5.1 Let $\{X_n; n \geq 1\}$ be a sequence of independent random variables in the sub-linear expectation space $(\Omega, \mathcal{H}, \widehat{\mathbb{E}})$. Suppose (5.6) and (5.7) hold, and for every $\alpha > 0$, there exists $p \geq 2$ such that (5.12) holds. Then (5.8)-(5.11) hold.

Remark 5.2 As shown by Wittmann, (5.6) is implied by

$$s_n^2 = \sum_{i=1}^n \widehat{\mathbb{E}}[X_i^2] \rightarrow \infty \quad \text{and} \quad \limsup_{n \rightarrow \infty} \frac{s_{n+1}}{s_n} < \infty. \quad (5.14)$$

In fact,

$$\begin{aligned} & \sum_{n=3}^{\infty} s_n^{-2} (\log s_n^2)^{\delta-1} \widehat{\mathbb{E}}[X_n^2] \\ &= \sum_{n=3}^{\infty} \int_{s_{n-1}^2}^{s_n^2} s_n^{-2} (\log s_n^2)^{\delta-1} dx \geq c \sum_{n=3}^{\infty} \int_{s_{n-1}^2}^{s_n^2} s_{n-1}^{-2} (\log s_{n-1}^2)^{\delta-1} dx \\ &\geq c \sum_{n=3}^{\infty} \int_{s_{n-1}^2}^{s_n^2} x (\log x)^{\delta-1} dx = c \int_{s_2^2}^{\infty} x (\log x)^{\delta-1} dx = \infty. \end{aligned}$$

Remark 5.3 If $\{X_n; n \geq 1\}$ satisfies Kolomogrov's (1929) condition as

$$|X_n| \leq \alpha_n \frac{s_n}{t_n}, \quad n = 1, 2, \dots \text{ and } \alpha_n \rightarrow 0, \quad (5.15)$$

then the conditions (5.12) (and therefore (5.13)) and (5.14) are satisfied.

Hence, we have the following corollary.

Corollary 5.2 (Kolomogrov's LIL) Let $\{X_n; n \geq 1\}$ be a sequence of independent random variables in the sub-linear expectation space $(\Omega, \mathcal{H}, \widehat{\mathbb{E}})$. Suppose (5.7) and (5.15) hold. Then (5.8)-(5.11) hold.

Remark 5.4 It is obvious that (5.5) implies (5.2). If we have an additional condition $s_n^{-2} \sum_{i=1}^n \widehat{\mathbb{E}}[X_j^2 \wedge a_j] \rightarrow 1$, then (5.5) can be weakened to (5.2). Also, it is obvious that

$$\bar{\Lambda}_n(p, \alpha) \leq \bar{\Lambda}_n(2, \alpha) \leq \frac{\sum_{j=1}^n \widehat{\mathbb{E}}[X_j^2]}{a_n^2} \leq \frac{1}{2 \log \log s_n^2},$$

and therefore, (5.2) is satisfied if

$$\sum_{n=1}^{\infty} \frac{\bar{\Gamma}_n(p, \alpha)}{s_n^{p/2} (\log \log s_n^2)^{d'}} < \infty \text{ for some } d' > 0. \quad (5.16)$$

As for the condition (5.5), when $p = 2$ it is just (i) of Theorem 4.1 of Wittmann (1987). Hence, Theorem 5.2 has extended Wittmann's LIL even in the classical case.

Note $\Gamma_n(2, 2\alpha)/a_n^2 \leq \frac{\Gamma_n(p, \alpha)}{s_n^{p/2}} \alpha^{2-p} t_n^{2p-2}$, $\frac{\Lambda_n(2, 2\alpha)}{a_n^2} \leq t_n^{-2}$. Hence, if for every $\alpha > 0$, there exist constants $p \geq 2$ and $d' > 0$ such that

$$\sum_{n=1}^{\infty} \frac{\Gamma(p, \alpha)}{s_n^{p/2} (\log \log s_n^2)^{d'}} < \infty, \quad (5.17)$$

then (5.2) is satisfied with $p = 2$ and every $\alpha > 0$.

Next, we consider the i.i.d. case. For a random variable X , we denote

$$\check{\mathbb{E}}[X] = \lim_{c \rightarrow \infty} \widehat{\mathbb{E}}[(-c) \vee (X \wedge c)]$$

if the limit exists. It can be verified that $\check{\mathbb{E}}[X]$ exists if $C_{\mathbb{V}}(|X|) < \infty$ or $\check{\mathbb{E}}[|X|^{1+\epsilon}] < \infty$, $\check{\mathbb{E}}[|X|] \leq C_{\mathbb{V}}(|X|)$. Further, $\check{\mathbb{E}}[X] = \widehat{\mathbb{E}}[X]$ if $\widehat{\mathbb{E}}[|X| - c]^+ \rightarrow 0$ as $c \rightarrow \infty$.

The following two theorems on the LIL for a sequence of independent and identically distributed random variables are corollaries of Theorems 5.1 and 5.2.

Theorem 5.3 Let $\{Y_n; n \geq 1\}$ be independent and identical distributed random variables in the sub-linear expectation space $(\Omega, \mathcal{H}, \widehat{\mathbb{E}})$, and \mathbb{V}^* be defined as in (4.21). Suppose

$$C_{\mathbb{V}} \left[\frac{Y_1^2}{\log \log |Y_1|} \right] < \infty, \quad (5.18)$$

Denote $\bar{\sigma}_2^2 = \check{\mathbb{E}}[(Y_1 - \check{\mathbb{E}}[Y_1])^2]$ and $\bar{\sigma}_1^2 = \check{\mathbb{E}}[(Y_1 + \check{\mathbb{E}}[-Y_1])^2]$ (finite or infinite). Then

$$\mathbb{V}^* \left(\liminf_{n \rightarrow \infty} \frac{\sum_{i=1}^n (Y_i + \check{\mathbb{E}}[-Y_i])}{\sqrt{2n \log \log n}} < -\bar{\sigma}_1 \text{ or } \limsup_{n \rightarrow \infty} \frac{\sum_{i=1}^n (Y_i - \check{\mathbb{E}}[Y_i])}{\sqrt{2n \log \log n}} > \bar{\sigma}_2 \right) = 0. \quad (5.19)$$

Theorem 5.4 Let $\{Y_n; n \geq 1\}$ be independent and identical distributed random variables in the sub-linear expectation space $(\Omega, \mathcal{H}, \widehat{\mathbb{E}})$, and \mathbb{V}^* be defined as in (4.21). Denote $\bar{\sigma}^2 = \check{\mathbb{E}}[Y_1^2]$ (finite or infinite).

(a) Suppose (5.18) and

$$\check{\mathbb{E}}[Y_1] = \check{\mathbb{E}}[-Y_1] = 0. \quad (5.20)$$

Then

$$\mathbb{V}^* \left(\limsup_{n \rightarrow \infty} \frac{|\sum_{i=1}^n Y_i|}{\sqrt{2n \log \log n}} > \bar{\sigma} \right) = 0, \quad (5.21)$$

$$\mathbb{V}^* \left(\limsup_{n \rightarrow \infty} \frac{\sum_{i=1}^n Y_i}{\sqrt{2n \log \log n}} = \bar{\sigma} \text{ and } \liminf_{n \rightarrow \infty} \frac{\sum_{i=1}^n Y_i}{\sqrt{2n \log \log n}} = -\bar{\sigma} \right) = 1, \quad (5.22)$$

$$\mathbb{V}^* \left(C \left\{ \frac{\sum_{i=1}^n Y_i}{\sqrt{2n \log \log n}} \right\} = [-\bar{\sigma}, \bar{\sigma}] \right) = 1, \quad (5.23)$$

where $C\{x_n\}$ denote the cluster set of a sequence of $\{x_n\}$ in \mathbb{R} .

(b) Suppose

$$\mathbb{V}^* \left(\limsup_{n \rightarrow \infty} \frac{|\sum_{i=1}^n Y_i|}{\sqrt{2n \log \log n}} = +\infty \right) < 1. \quad (5.24)$$

Then (5.18) and (5.20) hold, and

$$\bar{\sigma}^2 = \lim_{c \rightarrow \infty} \widehat{\mathbb{E}}[Y_1^2 \wedge c] < \infty. \quad (5.25)$$

Remark 5.5 Theorem 5.4 removed the continuity of \mathbb{V}^* and countably sub-additivity of $\widehat{\mathbb{E}}$ in Theorem 3.11 (b) of Zhang (2016). It tells us that (5.18), (5.20) and (5.25) are the sufficient and necessary conditions for Hartman and Wintner's type LIL under the sub-linear expectation.

Comparing with the LIL, the necessary and necessary conditions of the central limit theorem for independent and identically distributed random variables are shown to be (5.20), (5.25) and $\mathbb{V}(|X_1| \geq x) = o(x^{-2})$ as $x \rightarrow \infty$ by Zhang (2020).

The following Theorem gives the result under the lower capacity \mathcal{V} .

Theorem 5.5 Let $\{Y_n; n \geq 1\}$ be independent and identical distributed random variables in the sub-linear expectation space $(\Omega, \mathcal{H}, \widehat{\mathbb{E}})$, and the related capacity \mathcal{V} be countably sub-additive. Denote $\bar{\sigma}^2 = \check{\mathbb{E}}[Y_1^2]$, $\underline{\sigma}^2 = \lim_{c \rightarrow \infty} \widehat{\mathbb{E}}[Y_1^2 \wedge c]$, $T_n = \sum_{i=1}^n Y_i$ and $d_n = \sqrt{2n \log \log n}$. Suppose (5.18), (5.20) and (5.25) are satisfied. Then

$$\mathcal{V} \left(\underline{\sigma} \leq \limsup_{n \rightarrow \infty} \frac{T_n}{d_n} \leq \bar{\sigma} \right) = 1,$$

$$\mathcal{V} \left(-\bar{\sigma} \leq \liminf_{n \rightarrow \infty} \frac{T_n}{d_n} \leq -\underline{\sigma} \right) = 1$$

and

$$\mathcal{V} \left([-\bar{\sigma}, \bar{\sigma}] \supset C \left\{ \frac{T_n}{d_n} \right\} = \left[\liminf_{n \rightarrow \infty} \frac{T_n}{d_n}, \limsup_{n \rightarrow \infty} \frac{T_n}{d_n} \right] \supset [-\underline{\sigma}, \underline{\sigma}] \right) = 1.$$

In particular, the above equalities holds for \mathcal{V}^* which is defined as in (4.21).

Theorem 5.5 removes the continuity of \mathbb{V} in Corollary 3.13 of Zhang (2016) so that it is consistent with Theorem 1 of Chen and Hu (2014) where the random variables are assumed to be bounded. The condition that $\lim_{c \rightarrow \infty} \widehat{\mathbb{E}}[|Y_1|^2 - c]^+ = 0$ in Corollary 3.13 of Zhang (2016) is now weakened to (5.25). It should be noted that (5.25), $\widehat{\mathbb{E}}[|Y_1|^2] < \infty$ and $\lim_{c \rightarrow \infty} \widehat{\mathbb{E}}[|Y_1|^2 - c]^+ = 0$ are not equivalent under the sub-linear expectation, and (5.25) is the weakest among them. Also, they does not imply (5.18).

6 Proofs of the laws of the iterated logarithm

In this section, we give the proofs of the theorems in Section 5.

Proof of Theorem 5.1. By Wittmann (1985, Lemma 3.3), for any $\lambda > 1$, there exists a sequence $\{n_k\} \subset \mathbb{N}$ with

$$\lambda a_{n_k} \leq a_{n_{k+1}} \leq \lambda^3 a_{n_{k+1}}. \quad (6.1)$$

It can be checked that

$$\lambda s_{n_k}^2 \leq s_{n_{k+1}}^2 \leq \lambda^6 s_{n_{k+1}}^2, \quad \lambda^{1/2} \frac{s_{n_k}}{t_{n_k}} \leq \frac{s_{n_{k+1}}}{t_{n_{k+1}}} \leq \lambda^3 \frac{s_{n_{k+1}}}{t_{n_{k+1}}} \quad (6.2)$$

and $\log s_{n_{k+1}}^2 \sim \frac{1}{2} \log a_{n_{k+1}} \geq ck$. Hence

$$\sum_{k=1}^{\infty} \exp \left\{ -\frac{(1+\epsilon)t_{n_{k+1}}^2}{2} \right\} = \sum_{k=1}^{\infty} \exp \left\{ -(1+\epsilon) \log \log s_{n_{k+1}}^2 \right\} < \infty \quad \text{for all } \epsilon > 0. \quad (6.3)$$

We write $I(k)$ to denote the set $\{n_k + 1, \dots, n_{k+1}\}$. Denote $b_j = \alpha \lambda^3 s_j / t_j$, where $0 < \alpha < 1/10$ is a constant and to be specified. Denote

$$\bar{\Lambda}_{n_k, n_{k+1}}(p, \alpha) = \sum_{j \in I(k)} \widehat{\mathbb{E}}[((|X_j| \wedge a_{n_{k+1}} - \alpha s_{n_{k+1}} / t_{n_{k+1}})^+)^p].$$

It follows from (5.2) and (6.1) that (cf. the arguments of Wittmann (1987, page 526))

$$\sum_{k=1}^{\infty} \left(\frac{\bar{\Lambda}_{n_k, n_{k+1}}(p, \lambda^3 \alpha)}{a_{n_{k+1}}^p} \right)^{d+1} < \infty \quad (6.4)$$

Let

$$\mathbb{N}_1 = \left\{ k \in \mathbb{N}; \frac{\bar{\Lambda}_{n_k, n_{k+1}}(p, \lambda^3 \alpha)}{a_{n_{k+1}}^p} \leq t_{n_{k+1}}^{-2p} \right\}. \quad (6.5)$$

It follows from (6.4) that

$$\sum_{k \in \mathbb{N} \setminus \mathbb{N}_1} t_{n_{k+1}}^{-2p(d+1)} < \infty. \quad (6.6)$$

We consider the sequences $\{X_j; j \in \mathbb{N} \setminus \mathbb{N}_1\}$ and $\{X_j; j \in \mathbb{N}_1\}$, respectively. Let \bar{X}_j be X_j if $j \in I(k)$ and $k \in \mathbb{N} \setminus \mathbb{N}_1$, and 0 for otherwise. Denote $\tilde{X}_j = X_j - \bar{X}_j$.

First, we consider $\{X_j; j \in \mathbb{N} \setminus \mathbb{N}_1\}$. Denote $\bar{S}_n = \sum_{j=1}^n (\bar{X}_j - \widehat{\mathbb{E}}[\bar{X}_j])$. Then $\bar{S}_n = \sum_{j \in I(k), k \in \mathbb{N} \setminus \mathbb{N}_1, j \leq n} (X_j - \widehat{\mathbb{E}}[X_j])$. Let $x = \epsilon a_{n_{k+1}}$ and $y = \epsilon' a_{n_{k+1}}$, where $\epsilon' > 0$ is chosen such that $x/(10y) \geq d+1$. By the inequality (3.2) (with $\delta = 1$) in Lemma 3.1,

$$\begin{aligned} & \mathbb{V} \left(\max_{n \in I(k)} \frac{\sum_{j=n_k+1}^n (X_j - \widehat{\mathbb{E}}[X_j])}{a_n} \geq \epsilon \lambda^3 \right) \\ & \leq \mathbb{V} \left(\max_{n \in I(k)} \sum_{j=n_k+1}^n (X_j - \widehat{\mathbb{E}}[X_j]) \geq \epsilon \lambda^3 a_{n_{k+1}} \right) \\ & \leq \mathbb{V} \left(\max_{n \in I(k)} \sum_{j=n_k+1}^n (X_j - \widehat{\mathbb{E}}[X_j]) \geq \epsilon a_{n_{k+1}} \right) \\ & \leq \exp \left\{ -\frac{\epsilon^2}{2(1+1)} t_{n_{k+1}}^2 \right\} + \sum_{j \in I(k)} \mathbb{V}(|X_j| \geq \epsilon' a_{n_{k+1}}) \\ & \quad + C \left(\frac{\sum_{j \in I(k)} \widehat{\mathbb{E}}[|X_j|^{2p} \wedge a_{n_{k+1}}^{2p}]}{a_{n_{k+1}}^{2p}} \right)^{d+1}. \end{aligned}$$

For $j \in I(k)$, we have

$$\begin{aligned} \frac{\widehat{\mathbb{E}}[|X_j|^{2p} \wedge a_{n_{k+1}}^{2p}]}{a_{n_{k+1}}^{2p}} & \leq \frac{\widehat{\mathbb{E}}[|X_j|^{2p} \wedge b_{n_{k+1}}^{2p}]}{a_{n_{k+1}}^{2p}} + \frac{\widehat{\mathbb{E}}[((|X_j| \wedge a_{n_{k+1}} - b_{n_{k+1}})^+)^{2p}]}{a_{n_{k+1}}^{2p}} \\ & \leq C \frac{\widehat{\mathbb{E}}[X_j^2]}{s_{n_{k+1}}^2 t_{n_{k+1}}^{4p-2}} + \frac{\widehat{\mathbb{E}}[((|X_j| \wedge a_{n_{k+1}} - b_{n_{k+1}})^+)^p]}{a_{n_{k+1}}^p}. \end{aligned}$$

It follows that

$$\sum_{j \in I(k)} \frac{\widehat{\mathbb{E}}[|X_j|^{2p} \wedge a_{n_{k+1}}^{2p}]}{a_{n_{k+1}}^{2p}} \leq C t_{n_{k+1}}^{-2p} + \frac{\bar{\Lambda}_{n_k, n_{k+1}}(p, \lambda^3 \alpha)}{a_{n_{k+1}}^{2p}}.$$

Therefore, for $k \notin \mathbb{N}_1$,

$$\begin{aligned} & \mathbb{V} \left(\max_{n \in I(k)} \frac{\sum_{j=n_k+1}^n (X_j - \widehat{\mathbb{E}}[X_j])}{a_n} \geq \epsilon \lambda^3 \right) \\ & \leq C t_{n_{k+1}}^{-2p(d+1)} + C \left(\frac{\bar{\Lambda}_{n_k, n_{k+1}}(p, \lambda^3 \alpha)}{a_{n_{k+1}}^p} \right)^{d+1} + \sum_{j \in I(k)} \mathbb{V}(|X_j| \geq \epsilon' a_j). \end{aligned}$$

Hence, it follows from (5.1), (6.4) and (6.6) that

$$\sum_{k \in \mathbb{N} \setminus \mathbb{N}_1} \mathbb{V} \left(\max_{n \in I(k)} \frac{\sum_{j=n_k+1}^n (X_j - \widehat{\mathbb{E}}[X_j])}{a_n} \geq \epsilon \right) < \infty \text{ for all } \epsilon > 0.$$

That is,

$$\sum_{k=1}^{\infty} \mathbb{V} \left(\max_{n \in I(k)} \frac{\bar{S}_n - \bar{S}_{n_k}}{a_n} \geq \epsilon \right) < \infty \text{ for all } \epsilon > 0. \quad (6.7)$$

It follows from (6.7) that

$$\begin{aligned} & \mathbb{V} \left(\sup_{K \leq k \leq M} \max_{n \in I(k)} \frac{\bar{S}_n - \bar{S}_{n_k}}{a_n} > \epsilon \right) \\ & \leq \sum_{k=K}^{\infty} \mathbb{V} \left(\max_{n \in I(k)} \frac{\bar{S}_n - \bar{S}_{n_k}}{a_n} \geq \epsilon \right) \rightarrow 0 \text{ as } M \geq K \rightarrow \infty \text{ for all } \epsilon > 0, \end{aligned}$$

which implies

$$\lim_{N \geq n \rightarrow \infty} \mathbb{V} \left(\sup_{n \leq m \leq N} \frac{\bar{S}_m}{a_m} > \epsilon \right) = 0 \text{ for all } \epsilon > 0. \quad (6.8)$$

Next, consider $\{X_j; j \in \mathbb{N}_1\}$. We use the truncation method. Denote

$$Z_j = \tilde{X}_j \wedge (2b_{n_{k+1}}), j \in I(k), k \geq 0,$$

and $\tilde{S}_n = \sum_{j=1}^n (Z_j - \widehat{\mathbb{E}}[\tilde{X}_j])$. Then

$$S_n - \widehat{\mathbb{E}}[S_n] = \bar{S}_n + \tilde{S}_n + \sum_{j=1}^n (\tilde{X}_j - Z_j). \quad (6.9)$$

Note that $\tilde{X}_j - Z_j = 0$ when $j \in I(k)$ and $k \in \mathbb{N}_1$, $= (X_j - 2b_{n_{k+1}})^+ \geq 0$ for otherwise. It is easily seen that

$$\begin{aligned} \widehat{\mathbb{E}}[(X_j \wedge a_{n_{k+1}} - 2b_{n_{k+1}})^+] & \leq \widehat{\mathbb{E}}[((X_{n_{k+1}} \wedge a_{n_{k+1}} - b_{n_{k+1}})^+)^p] b_{n_{k+1}}^{1-p} \\ \widehat{\mathbb{E}}[((X_j \wedge a_{n_{k+1}} - 2b_{n_{k+1}})^+)^2] & \leq \widehat{\mathbb{E}}[((X_j \wedge a_{n_{k+1}} - b_{n_{k+1}})^+)^p] b_{n_{k+1}}^{2-p}. \end{aligned}$$

It follows that for $k \in \mathbb{N}_1$,

$$\frac{\sum_{j \in I(k)} \widehat{\mathbb{E}}[\tilde{X}_j \wedge a_{n_{k+1}} - Z_j]}{a_{n_{k+1}}} \leq \frac{\bar{A}_{n_k, n_{k+1}}(p, \alpha \lambda^3)}{a_{n_{k+1}}^p} \alpha^{1-p} t_{n_{k+1}}^{2p-2} \leq \alpha^{1-p} t_{n_{k+1}}^{-2} \rightarrow 0, \quad (6.10)$$

$$\frac{\sum_{j \in I(k)} \widehat{\mathbb{E}}[(\tilde{X}_j \wedge a_{n_{k+1}} - Z_j)^2]}{s_{n_{k+1}}^2} \leq \frac{\bar{A}_{n_k, n_{k+1}}(p, \alpha \lambda^3)}{a_{n_{k+1}}^p} \alpha^{2-p} t_{n_{k+1}}^{2p-2} \leq \alpha^{2-p} t_{n_{k+1}}^{-2} \rightarrow 0. \quad (6.11)$$

Let $x = \epsilon a_{n_{k+1}}/2$ and $y = \epsilon' a_{n_{k+1}}$, where $0 < \epsilon' < 1$ is chosen such that $x/(10y) \geq d + 1$.

By the inequality (3.2) (with $\delta = 1$) in Lemma 3.1, we have that for $k \in \mathbb{N}_1$ large enough,

$$\begin{aligned}
& \mathbb{V}\left(\sum_{j \in I(k)} |\tilde{X}_j - Z_j| \geq \epsilon a_{n_{k+1}}\right) \\
& \leq \mathbb{V}\left(\sum_{j \in I(k)} |\tilde{X}_j \wedge a_{n_{k+1}} - Z_j| \geq \epsilon a_{n_{k+1}}\right) + \sum_{j \in I(k)} \mathbb{V}(X_j > a_{n_{k+1}}) \\
& \leq \mathbb{V}\left(\sum_{j \in I(k)} (|\tilde{X}_j \wedge a_{n_{k+1}} - Z_j| - \widehat{\mathbb{E}}[|\tilde{X}_j \wedge a_{n_{k+1}} - Z_j|]) \geq \epsilon a_{n_{k+1}}/2\right) + \sum_{j \in I(k)} \mathbb{V}(X_j > a_j) \\
& \leq \exp\left\{-\frac{\epsilon^2/4}{2(1+1)} \frac{a_{n_{k+1}}^2}{\sum_{j \in I(k)} \widehat{\mathbb{E}}[(\tilde{X}_j \wedge a_j - Z_j)^2]}\right\} \\
& \quad + C \left(\sum_{j \in I(k)} \frac{\widehat{\mathbb{E}}[(|X_j| \wedge a_{n_{k+1}} - b_{n_{k+1}})^+)^p]}{a_{n_{k+1}}^p}\right)^{d+1} + 2 \sum_{j \in I(k)} \mathbb{V}(X_j > \epsilon' a_{n_{k+1}}) \\
& \leq \exp\{-2t_{n_{k+1}}^2\} + C \left(\frac{\bar{\Lambda}_{n_k, n_{k+1}}(p, \lambda^3 \alpha)}{a_{n_{k+1}}^p}\right)^{d+1} + 2 \sum_{j \in I(k)} \mathbb{V}(X_j > \epsilon' a_j). \tag{6.12}
\end{aligned}$$

Note $\tilde{X}_j - Z_j = 0$ when $j \in I(k)$ and $k \in \mathbb{N} \setminus \mathbb{N}_1$. It follows from (6.12), (6.3) and (6.4) and (5.1) that

$$\sum_{k=1}^{\infty} \mathbb{V}\left(\sum_{j \in I(k)} |\tilde{X}_j - Z_j| \geq \epsilon a_{n_{k+1}}\right) = \sum_{k \in \mathbb{N}_1} \mathbb{V}\left(\sum_{j \in I(k)} |\tilde{X}_j - Z_j| \geq \epsilon a_{n_{k+1}}\right) < \infty.$$

Thus,

$$\lim_{M \geq K \rightarrow \infty} \mathbb{V}\left(\sup_{K \leq k \leq M} \frac{\sum_{j \in I(k)} |\tilde{X}_j - Z_j|}{a_{n_{k+1}}} > \epsilon\right) = 0 \text{ for all } \epsilon > 0,$$

which implies

$$\lim_{N \geq n \rightarrow \infty} \mathbb{V}\left(\sup_{n \leq m \leq N} \frac{\sum_{j=1}^m |\tilde{X}_j - Z_j|}{a_m} > \epsilon\right) = 0 \text{ for all } \epsilon > 0. \tag{6.13}$$

At last, we consider Z_j . For $0 < \epsilon < 1/2$, choose $\alpha > 0$ such that $8\alpha\lambda^3 < \epsilon$. Note that $\widehat{\mathbb{E}}[Z_j] \leq \widehat{\mathbb{E}}[\tilde{X}_j]$ and $\sum_{j=1}^n \widehat{\mathbb{E}}[Z_j^2] \leq s_n^2$. Let $y_k = 2b_{n_{k+1}}$, $x_k = (1 + \epsilon)a_{n_{k+1}}$. Applying (3.1) in Lemma 3.1 yields

$$\begin{aligned}
& \mathbb{V}\left(\max_{n \in I(k)} \frac{\tilde{S}_n}{a_n} \geq (1 + \epsilon)\lambda^3\right) \leq \mathbb{V}\left(\max_{n \in I(k)} \tilde{S}_n \geq (1 + \epsilon)a_{n_{k+1}}\right) \\
& \leq \exp\left\{-\frac{x_k^2}{2(x_k y_k + \sum_{j=1}^{n_{k+1}} \widehat{\mathbb{E}}[Z_j^2])}\right\} \leq \exp\left\{-\frac{x_k^2}{2(x_k y_k + s_{n_{k+1}}^2)}\right\} \\
& = \exp\left\{-\frac{(1 + \epsilon)^2 t_{n_{k+1}}^2}{2(1 + 8\alpha\lambda^3)}\right\} \leq \exp\{-(1 + \epsilon) \log \log s_{n_{k+1}}^2\},
\end{aligned}$$

when k is large enough. It follows (6.3) that

$$\sum_{k=1}^{\infty} \mathbb{V}\left(\max_{n \in I(k)} \frac{\tilde{S}_n}{a_n} \geq (1 + \epsilon)\lambda^3\right) < \infty \quad \forall \epsilon > 0,$$

which implies

$$\lim_{N \geq n \rightarrow \infty} \mathbb{V} \left(\sup_{n \leq m \leq N} \frac{\tilde{S}_m}{a_m} > (1 + \epsilon)\lambda^3 \right) = 0. \quad (6.14)$$

By combing (6.9), (6.8), (6.13) and (6.14), it follows that

$$\lim_{N \geq n \rightarrow \infty} \mathbb{V} \left(\sup_{n \leq m \leq N} \frac{S_m - \widehat{\mathbb{E}}[S_m]}{a_m} > (1 + \epsilon)\lambda^3 + \epsilon \right) = 0 \text{ for all } \epsilon > 0 \text{ and } \lambda > 1.$$

Therefore,

$$\lim_{N \geq n \rightarrow \infty} \mathbb{V} \left(\sup_{n \leq m \leq N} \frac{S_m - \widehat{\mathbb{E}}[S_m]}{a_m} > 1 + \epsilon \right) = 0 \text{ for all } \epsilon > 0.$$

For $-X_j$ s, we have the same result. The proof (5.3) is now completed.

For (5.4), let $\epsilon_k = 1/2^k$. Note that \mathbb{V}^* satisfies (2.4). By (2.7), (5.1) is equivalent to that it holds for \mathbb{V}^* . So, (5.3) holds for \mathbb{V}^* . Then, there is a sequence of $n_k \uparrow \infty$ such that $\mathbb{V}^*(B_k) \leq \epsilon_k$ with

$$B_k = \left\{ \inf_{n_k \leq m \leq n_{k+1}} \frac{S_m - \widehat{\mathcal{E}}[S_m]}{a_m} < 1 + \epsilon_k \text{ and } \sup_{n_k \leq m \leq n_{k+1}} \frac{S_m - \widehat{\mathbb{E}}[S_m]}{a_m} > 1 + \epsilon_k \right\}.$$

Note $\sum_{k=1}^{\infty} \mathbb{V}^*(B_k) < \infty$. By the countable sub-additivity of \mathbb{V}^* , we have

$$\mathbb{V}^*(B_k \text{ i.o.}) = 0,$$

which implies (5.4). \square

Proof of Theorem 5.2. (5.8) follows from Theorem 5.1. Now, we consider (5.9). Let $\lambda > 1$ be large enough. Let $\{n_k = n_k(\lambda)\} \subset \mathbb{N}$ satisfy (6.1), and denote $I(k) = I(k, \lambda) = \{n_k + 1, \dots, n_{k+1}\}$. Then (6.2) is satisfied. Denote $b_j = \alpha \lambda^3 s_j / t_j$, where $0 < \alpha < 1/(10\lambda^3)$ is a constant and to be specified. Redefine

$$Z_j = (-2b_{n_{k+1}}) \vee X_j \wedge (2b_{n_{k+1}}), \quad j \in I(k), \quad k \geq 0.$$

Let $B_k^2 = \sum_{j \in I(k)} \widehat{\mathbb{E}}[X_j^2]$. Then $B_k^2 = s_{n_{k+1}}^2 - s_{n_k}^2 \geq (1 - 1/\lambda)s_{n_{k+1}}^2$, $\log s_{n_k}^2 \geq ck$. From the condition (5.6), it follows that

$$\sum_{k=1}^{\infty} (\log s_{n_k}^2)^{\delta-1} = \infty \text{ for all } \delta > 0 \quad (6.15)$$

by Lemma 2.3 of Wittmann (1987). Similar to (6.4), it follows from (5.5) and (6.2) that

$$\sum_{k=1}^{\infty} \left(\frac{\Lambda_{n_k, n_{k+1}}(p, \lambda^3 \alpha)}{a_{n_{k+1}}^p} \right)^{d+1} < \infty, \quad (6.16)$$

where

$$\Lambda_{n_k, n_{k+1}}(p, \alpha) = \sum_{j \in I(k)} \widehat{\mathbb{E}}[(|X_j| - \alpha s_{n_{k+1}}/t_{n_{k+1}})^+]^p.$$

Let

$$\mathbb{N}_1 = \left\{ k \in \mathbb{N}; \frac{\Lambda_{n_k, n_{k+1}}(p, \lambda^3 \alpha)}{a_{n_{k+1}}^p} \leq t_{n_{k+1}}^{-2p} \right\}. \quad (6.17)$$

It follows from (6.16) that (6.6) holds. By (5.7), we have

$$\frac{\sum_{j \in I(k)} |\widehat{\mathbb{E}}[X_j]|}{a_{n_{k+1}}} + \frac{\sum_{j \in I(k)} |\widehat{\mathcal{E}}[X_j]|}{a_{n_{k+1}}} \rightarrow 0. \quad (6.18)$$

Note

$$\begin{aligned} \widehat{\mathbb{E}}|X_j - Z_j| &\leq \widehat{\mathbb{E}}[(|X_j| - b_{n_{k+1}})^+]^p b_{n_{k+1}}^{1-p}, \\ \widehat{\mathbb{E}}|X_j - Z_j|^2 + \widehat{\mathbb{E}}|X_j^2 - Z_j^2| &\leq 2\widehat{\mathbb{E}}[(|X_j| - b_{n_{k+1}})^+]^p b_{n_{k+1}}^{2-p}. \end{aligned}$$

Similar to (6.10) and (6.11), it follows that for $k \in \mathbb{N}_1$,

$$\frac{\sum_{j \in I(k)} \widehat{\mathbb{E}}|X_j - Z_j|}{a_{n_{k+1}}} \leq \alpha^{1-p} t_{n_{k+1}}^{-2} \rightarrow 0, \quad (6.19)$$

$$\frac{\sum_{j \in I(k)} (\widehat{\mathbb{E}}|X_j - Z_j|^2 + \widehat{\mathbb{E}}|X_j^2 - Z_j^2|)}{s_{n_{k+1}}^2} \leq 2\alpha^{2-p} t_{n_{k+1}}^{-2} \rightarrow 0. \quad (6.20)$$

Thus, similar to (6.12), by Lemma 3.1 we have that for $k \in \mathbb{N}_1$ large enough,

$$\begin{aligned} &\mathbb{V}\left(\sum_{j \in I(k)} |X_j - Z_j| \geq \epsilon a_{n_{k+1}}\right) \\ &\leq \exp\{-2t_{n_{k+1}}^2\} + C \left(\frac{\Lambda_{n_k, n_{k+1}}(p, \lambda^3 \alpha)}{a_{n_{k+1}}^p}\right)^{d+1} + \sum_{j \in I(k)} \mathbb{V}(|X_j| \geq \epsilon' a_j). \end{aligned}$$

It follows that

$$\sum_{k \in \mathbb{N}_1} \mathbb{V}\left(\sum_{j \in I(k)} |X_j - Z_j| \geq \epsilon a_{n_{k+1}}\right) < \infty \quad \text{for all } \epsilon > 0. \quad (6.21)$$

Next, we consider Z_j . Let $\widetilde{B}_k^2 = \sum_{j \in I(k)} \widehat{\mathbb{E}}[Z_j^2]$. It follows from (6.2) and (6.20) that

$$\widetilde{B}_k^2 \sim B_k^2 \geq (1 - 1/\lambda) s_{n_{k+1}}^2, \quad k \in \mathbb{N}_1.$$

Without loss of generality, we assume that

$$\frac{s_{n_{k+1}}}{\widetilde{B}_k} \leq \frac{\lambda}{\lambda - 1}, \quad k \in \mathbb{N}_1.$$

It follows from (6.18) and (6.19) that

$$\frac{\sum_{j \in I(k)} |\widehat{\mathbb{E}}[Z_j]|}{a_{n_{k+1}}} + \frac{\sum_{j \in I(k)} |\widehat{\mathcal{E}}[Z_j]|}{a_{n_{k+1}}} \rightarrow 0, \quad \mathbb{N}_1 \ni k \rightarrow \infty.$$

Further,

$$|Z_j| \leq 2b_{n_{k+1}} = 2\alpha\lambda^3 \frac{s_{n_{k+1}}}{t_{n_{k+1}}} \leq 2\alpha\lambda^3 \frac{\lambda}{\lambda-1} \frac{\tilde{B}_k}{t_{n_{k+1}}}, \quad j \in I(k)$$

for k large enough. For every $\epsilon > 0$, let $\gamma = \epsilon/2$ and $\pi(\gamma)$ be the constant defined as in Lemma (3.2). Choose α such that $2\alpha \frac{\lambda^4}{\lambda-1} < \pi(\gamma)$. By Lemma 3.2, we have that for $k \in \mathbb{N}_1$ large enough,

$$\begin{aligned} & \mathbb{V}\left(\sum_{j \in I(k)} Z_j \geq (1-\epsilon)(1-1/\lambda)a_{n_{k+1}}\right) \geq \mathbb{V}\left(\sum_{j \in I(k)} Z_j \geq (1-2\epsilon)\tilde{B}_k t_{n_{k+1}}\right) \\ & \geq \exp\left\{-\frac{(1-\epsilon)^2 t_{n_{k+1}}^2}{2}(1+\epsilon)\right\} \geq \exp\left\{-\frac{(1-\epsilon^2)t_{n_{k+1}}^2}{2}\right\}. \end{aligned} \quad (6.22)$$

It follows from (6.21), (6.22), (6.15) and (6.6) that

$$\begin{aligned} & \sum_{k \in \mathbb{N}_1} \mathbb{V}\left(\sum_{j \in I(k)} X_j \geq (1-2\epsilon)(1-1/\lambda)a_{n_{k+1}}\right) \\ & \geq \sum_{k \in \mathbb{N}_1} \mathbb{V}\left(\sum_{j \in I(k)} Z_j \geq (1-\epsilon)(1-1/\lambda)a_{n_{k+1}}\right) - C \\ & \geq \sum_k \exp\left\{-\frac{(1-\epsilon^2)t_{n_{k+1}}^2}{2}\right\} - \sum_{k \in \mathbb{N} \setminus \mathbb{N}_1} \exp\left\{-\frac{(1-\epsilon^2)t_{n_{k+1}}^2}{2}\right\} - C \\ & \geq \sum_k (\log s_{n_{k+1}}^2)^{\epsilon^2-1} - c \sum_{k \in \mathbb{N} \setminus \mathbb{N}_1} t_{n_{k+1}}^{-2p(d+1)} - C = \infty. \end{aligned}$$

Hence

$$\sum_{k=1}^{\infty} \mathbb{V}\left(\sum_{j \in I(k)} X_j \geq (1-2\epsilon)(1-1/\lambda)a_{n_{k+1}}\right) = \infty. \quad (6.23)$$

Note the the independence of $\sum_{j \in I(k)} X_j$, $k = 1, 2, \dots$, by Lemma 4.1 (ii) it follows that

$$\mathbb{V}\left(\max_{K \leq k \leq M} \frac{\sum_{j \in I(k)} X_j}{a_{n_{k+1}}} \geq (1-3\epsilon)(1-1/\lambda)\right) \rightarrow 1 \text{ as } M \rightarrow \infty \text{ for all } K. \quad (6.24)$$

On the other hand,

$$\frac{|S_{n_k}|}{a_{n_{k+1}}} \leq \frac{|S_{n_k}|}{a_{n_k}} \cdot \frac{a_{n_k}}{a_{n_{k+1}}} \leq \frac{|S_{n_k}|}{a_{n_k}} \frac{1}{\lambda} \quad (6.25)$$

for k large enough. It follows that for K large enough,

$$\begin{aligned} & \mathbb{V}\left(\max_{k \geq K} \max_{n \in I_k} \frac{S_n}{a_n} \geq (1-3\epsilon)(1-1/\lambda) - (1+\epsilon)/\lambda\right) \\ & \geq \mathbb{V}\left(\max_{K \leq k \leq M} \frac{S_{n_{k+1}}}{a_{n_{k+1}}} \geq (1-3\epsilon)(1-1/\lambda) - (1+\epsilon)/\lambda\right) \\ & \geq \mathbb{V}\left(\max_{K \leq k \leq M} \frac{\sum_{j \in I(k)} X_j}{a_{n_{k+1}}} \geq (1-3\epsilon)(1-1/\lambda)\right) - \mathbb{V}\left(\max_{K \leq k \leq M} \frac{|S_{n_k}|}{a_{n_k}} \geq 1+\epsilon\right) \\ & \rightarrow 1 \text{ as } M \rightarrow \infty \text{ and then } K \rightarrow \infty, \end{aligned}$$

by (5.8) and (6.24). By the arbitrariness of $\epsilon > 0$ being small enough and $\lambda > 1$ being large enough. We obtain

$$\lim_{n \rightarrow \infty} \mathbb{V} \left(\max_{m \geq n} \frac{S_m}{a_m} \geq 1 - \epsilon \right) = 1 \text{ for all } \epsilon > 0.$$

For $-X_j$, we have the same conclusion. (5.9) is proved.

At last, we show (5.10)-(5.11). As in Theorem 5.1, by the countable sub-additivity of \mathbb{V}^* , (5.8) implies (5.10).

For (5.11), note that (6.23) holds for \mathbb{V}^* . Further, note $\log s_{n_{k+1}}^2 \sim \log s_{n_k}^2$. We have

$$\sum_{k:k \text{ even}} \left(\text{resp. } \sum_{k:k \text{ odd}} \right) (\log s_{n_k}^2)^{\delta-1} \geq c \sum_{k=1}^{\infty} (\log s_{n_k}^2)^{\delta-1} = \infty.$$

It follows that (6.23) remains true if the summation is taken over even numbers or odd numbers. So,

$$\sum_{k:k \text{ even}} \left(\text{resp. } \sum_{k:k \text{ odd}} \right) \mathbb{V}^* \left(\sum_{j \in I(k)} X_j \geq (1 - 2/\lambda) a_{n_{k+1}} \right) = \infty. \quad (6.26)$$

For $-X_j$, we have a same equality as above. Note that $n_k = n_k(\lambda)$ and $I(k) = I(k, \lambda)$ depend on $\lambda > 1$, and $n_k(\lambda)$ satisfies (6.1). We denote $Z_k(\lambda) = \frac{\sum_{j \in I(k)} X_j}{a_{n_{k+1}}}$. Now, suppose $\lambda_0 > 1$ is large enough and $\lambda_0 \leq \lambda_i \nearrow \infty$. For λ_1 , choose $\underline{k}_1 = 0$ and \bar{k}_1 such that

$$\sum_{k \in J(1), k \text{ odd}} \mathbb{V}^* \left(Z_k(\lambda_1) \geq 1 - 2/\lambda_1 \right) > 1$$

and

$$\sum_{k \in J(1), k \text{ even}} \mathbb{V}^* \left(-Z_k(\lambda_1) \geq 1 - 2/\lambda_1 \right) > 1$$

where $J(1) = \{\underline{k}_1 + 1, \dots, \bar{k}_1\}$. For λ_2 , we first choose $\underline{k}_2 \geq \bar{k}_1$ such that $n_{\underline{k}_2}(\lambda_2) \geq n_{\bar{k}_1 + 1}(\lambda_1)$.

And then choose \bar{k}_2 with $J(2) = \{\underline{k}_2 + 1, \dots, \bar{k}_2\}$ such that

$$\sum_{k \in J(2), k \text{ odd}} \mathbb{V}^* \left(Z_k(\lambda_2) \geq 1 - 2/\lambda_2 \right) > 1$$

and

$$\sum_{k \in J(2), k \text{ even}} \mathbb{V}^* \left(-Z_k(\lambda_2) \geq 1 - 2/\lambda_2 \right) > 1$$

Inductively,

$$\sum_{k \in J(i), k \text{ odd}} \mathbb{V}^* \left(Z_k(\lambda_i) \geq 1 - 2/\lambda_i \right) > 1$$

and

$$\sum_{k \in J(i), k \text{ even}} \mathbb{V}^* \left(-Z_k(\lambda_i) \geq 1 - 2/\lambda_i \right) > 1,$$

where $J(i) = \{\underline{k}_i + 1, \dots, \bar{k}_i\}$ and $\underline{k}_i \geq \bar{k}_{i-1}$, $n_{\underline{k}_i}(\lambda_i) \geq n_{\bar{k}_{i-1}+1}(\lambda_{i-1})$. Let $\gamma_k = \lambda_i$ if $k \in J(i)$. Denote $\tilde{\mathbb{N}}_1 = \{k : k \in J(i), k \text{ is odd}\}$ and $\tilde{\mathbb{N}}_2 = \{k : k \in J(i), k \text{ is even}\}$. Then $\tilde{\mathbb{N}}_1$ and $\tilde{\mathbb{N}}_2$ are disjoint subsets of $\tilde{\mathbb{N}}$ with

$$\sum_{k \in \tilde{\mathbb{N}}_1} \mathbb{V}^* \left(\gamma_k [Z_k(\gamma_k) - 1] \geq -2 \right) = \infty$$

and

$$\sum_{k \in \tilde{\mathbb{N}}_2} \mathbb{V}^* \left(\gamma_k [-Z_k(\gamma_k) - 1] \geq -2 \right) = \infty$$

Let $A_k = \{\gamma_k [Z_k(\gamma_k) - 1] \geq -3\}$ if $k \in \tilde{\mathbb{N}}_1$, and $B_k = \{\gamma_k [-Z_k(\gamma_k) - 1] \geq -3\}$ if $k \in \tilde{\mathbb{N}}_2$. By the independence of $Z_k(\gamma_k)$, $k \in J(i)$, $i = 1, 2, \dots$, and Lemma 4.3 (i),

$$\mathbb{V}^* \left(\{A_k, k \in \tilde{\mathbb{N}}_1 \text{ i.o.}\} \cap \{B_k, k \in \tilde{\mathbb{N}}_2 \text{ i.o.}\} \right) = 1. \quad (6.27)$$

Write $A = \{A_k, k \in \tilde{\mathbb{N}}_1 \text{ i.o.}\} \cap \{B_k, k \in \tilde{\mathbb{N}}_2 \text{ i.o.}\}$ and $C = \{\limsup_{n \rightarrow \infty} \frac{|S_n|}{a_n} > 1\}$. Note for $k \in J(i)$,

$$\begin{aligned} \frac{S_{n_{k+1}(\gamma_k)}}{a_{n_{k+1}(\gamma_k)}} &\geq \frac{\sum_{j \in I(k, \gamma_k)} X_j}{a_{n_{k+1}(\gamma_k)}} - \frac{|S_{n_k(\gamma_k)}|}{a_{n_{k+1}(\gamma_k)}} \geq Z_k(\gamma_k) - \frac{1}{\gamma_k} \frac{|S_{n_k(\gamma_k)}|}{a_{n_k(\gamma_k)}}, \\ \frac{-S_{n_{k+1}(\gamma_k)}}{a_{n_{k+1}(\gamma_k)}} &\geq \frac{-\sum_{j \in I(k, \gamma_k)} X_j}{a_{n_{k+1}(\gamma_k)}} - \frac{|S_{n_k(\gamma_k)}|}{a_{n_{k+1}(\gamma_k)}} \geq -Z_k(\gamma_k) - \frac{1}{\gamma_k} \frac{|S_{n_k(\gamma_k)}|}{a_{n_k(\gamma_k)}} \end{aligned}$$

by (6.1), and $\gamma_k \rightarrow \infty$. So, on AC^c we have

$$\limsup_{n \rightarrow \infty} \frac{S_n}{a_n} \geq 1 - 4 \limsup_{k \rightarrow \infty} \frac{1}{\gamma_k} = 1 \quad \text{and} \quad \limsup_{n \rightarrow \infty} \frac{-S_n}{a_n} \geq 1 - 4 \limsup_{k \rightarrow \infty} \frac{1}{\gamma_k} = 1.$$

It follows that

$$\mathbb{V}^* \left(\limsup_{n \rightarrow \infty} \frac{S_n}{a_n} \geq 1 \text{ and } \limsup_{n \rightarrow \infty} \frac{-S_n}{a_n} \geq 1 \right) \geq \mathbb{V}^*(AB^c) \geq \mathbb{V}^*(A) - \mathbb{V}^*(B) = 1 - 0 = 1,$$

by (5.10) and (6.27). Hence (5.11) is proved by (5.10). \square

For proving Theorems 5.3 and 5.4 for independent and identically distributed random variables, we need more two lemmas.

Lemma 6.1 *Suppose $X \in \mathcal{H}$.*

(i) *For any $\delta > 0$,*

$$\sum_{n=1}^{\infty} \mathbb{V}(|X| \geq \delta \sqrt{n \log \log n}) < \infty \iff C_V \left[\frac{X^2}{\log \log |X|} \right] < \infty.$$

(ii) *If $C_V \left[\frac{X^2}{\log \log |X|} \right] < \infty$, then for any $\delta > 0$ and $p > 2$,*

$$\sum_{n=1}^{\infty} \frac{\widehat{\mathbb{E}}[(|X| \wedge (\delta \sqrt{n \log \log n}))^p]}{(n \log \log n)^{p/2}} < \infty.$$

(iii) $C_V \left[\frac{X^2}{\log \log |X|} \right] < \infty$, then for any $\delta > 0$,

$$\widehat{\mathbb{E}}[X^2 \wedge (2\delta n \log \log n)] = o(\log \log n)$$

and

$$\check{\mathbb{E}}[(|X| - \delta\sqrt{2n \log \log n})^+] = o(\sqrt{\log \log n/n}). \quad \square$$

Proof. The proof of (i) and (ii) can be found in Zhang (2016). For (iii), we denote $d_n = \sqrt{2n \log \log n}$. Let $f(x)$ be the inverse function of $g(x) = \sqrt{2x \log \log x}$ ($x > 0$). Then $cx^2/\log \log |x| \leq f(|x|) \leq Cx^2/\log \log |x|$. It follows that $\int_0^\infty \mathbb{V}(f(|X|) \geq y) dy < \infty$. Hence

$$\begin{aligned} \frac{\widehat{\mathbb{E}}[(|X| \wedge \delta d_n)^2]}{\log \log n} &\leq \frac{C_V((|X| \wedge \delta d_n)^2)}{\log \log n} = \frac{1}{\log \log n} \int_0^{(\delta d_n)^2} \mathbb{V}(|X|^2 > x) dx \\ &= \frac{1}{\log \log n} 2 \int_0^{\delta d_n} x \mathbb{V}(|X| > x) dx = \frac{2}{\log \log n} \int_0^{g(\delta d_n)} g(y) \mathbb{V}(|X| > g(y)) dg(y) \\ &\leq 4 \int_0^{C_\delta n} \frac{\log \log y}{\log \log n} \mathbb{V}(f(|X|) > y) dy \rightarrow 0 \end{aligned}$$

and

$$\begin{aligned} \widehat{\mathbb{E}}[(|X| - \delta d_n)^+] &\leq C_V((|X| - \delta d_n)^+) \leq \int_{\delta d_n}^\infty \mathbb{V}(|X| \geq x) dx \\ &= \int_{g(\delta d_n)}^\infty \mathbb{V}(|X| \geq g(y)) dg(y) \leq 2\sqrt{2} \int_{c_\delta n}^\infty \sqrt{\log \log y/y} \mathbb{V}(f(|X|) \geq y) dy \\ &\leq 2\sqrt{2} \sqrt{\log \log n/n} \int_{c_\delta n}^\infty \mathbb{V}(f(|X|) \geq y) dy = o(\sqrt{\log \log n/n}). \quad \square \end{aligned}$$

Lemma 6.2 Let $\{Y_n; n \geq 1\}$ be a sequence of independent and identical distributed random variables in the sub-linear expectation space $(\Omega, \mathcal{H}, \widehat{\mathbb{E}})$ with $C_V \left[\frac{Y_1^2}{\log \log |Y_1|} \right] < \infty$. Then

$$\mathbb{V} \left(\frac{\sum_{i=1}^n (Y_i - \check{\mathbb{E}}[Y_1])}{\sqrt{2n \log \log n}} \geq \epsilon \right) \rightarrow 0 \quad \text{for all } \epsilon > 0, \quad (6.28)$$

$$\mathbb{V} \left(\frac{\sum_{i=1}^n (-Y_i + \check{\mathbb{E}}[Y_1])}{\sqrt{2n \log \log n}} \geq \epsilon \right) \rightarrow 0 \quad \text{for all } \epsilon > 0. \quad (6.29)$$

Proof. For a random variable Y , we denote $Y^{(c)} = (-c) \vee Y \wedge c$. Denote $d_n = \sqrt{2 \log \log n}$.

Then by applying (3.2) and (3.5) with $p = 2$, we obtain

$$\mathbb{V} \left(\sum_{i=1}^n (Y_i^{(d_n)} - n\widehat{\mathbb{E}}[Y_1^{(d_n)}]) \geq \epsilon d_n \right) \leq C \frac{n\widehat{\mathbb{E}}[(|Y_1| \wedge d_n)^2]}{\epsilon^2 d_n^2} \rightarrow 0$$

and

$$\frac{n|\check{\mathbb{E}}[Y_1] - \widehat{\mathbb{E}}[Y_1^{(d_n)}]|}{d_n} = \frac{n\check{\mathbb{E}}[(|Y_1| - d_n)^+]}{d_n} \rightarrow 0$$

by Lemma 6.1 (iii). On the other hand,

$$\mathbb{V} \left(Y_i^{(d_n)} \neq Y_i \text{ for some } i = 1, \dots, n \right) \leq n\mathbb{V}(|Y_1| \geq d_n) \rightarrow 0$$

by Lemma 6.1 (i). Therefore, (6.28) holds. The proof of (6.29) is similar. \square

Proof of Theorems 5.3 and 5.4. If $\bar{\sigma}^2 = 0$, then $|\check{\mathbb{E}}[\pm Y_1]| \leq \check{\mathbb{E}}[|Y_1|] \leq (\check{\mathbb{E}}[Y_1^2])^{1/2} = 0$, $\bar{\sigma}_1^2 = \bar{\sigma}_2^2 = \bar{\sigma}^2 = 0$, and for any $\epsilon > 0$, $\mathbb{V}(|Y_1| \geq \epsilon) \leq \widehat{\mathbb{E}}[Y_1^2 \wedge \epsilon^2]/\epsilon^2 = 0$. By the countable subadditivity of \mathbb{V}^* , it follows that $\mathbb{V}^*(|Y_n| > 0) \leq \sum_{j=1}^{\infty} \mathbb{V}^*(|Y_n| > 1/j) \leq \sum_{j=1}^{\infty} l^2 \widehat{\mathbb{E}}[Y_1^2 \wedge j^2] = 0$. Hence $\mathbb{V}^*(|Y_n| \neq 0 \text{ for some } n) = 0$. And then, (5.18), (5.19) in Theorem 5.2 and (5.20)-(5.22) in Theorem 5.3 hold automatically. Therefore, without loss of generality, we assume $0 < \bar{\sigma} \leq \infty$.

We first suppose that (5.18) is satisfied. By (2.8), (5.18) is equivalent to that it holds for \mathbb{V}^* . Let $d_n = \sqrt{2n \log \log n}$ and $X_n = (-d_n) \vee Y_n \wedge d_n$. Denote $S_n = \sum_{i=1}^n X_i$, $s_n^2 = \sum_{i=1}^n \widehat{\mathbb{E}}[X_i^2]$, $t_n = \sqrt{2 \log \log s_n}$ and $a_n = t_n s_n$. Then $n \leq C s_n^2$, $d_n \leq C a_n$. Note that

$$\begin{aligned} \sum_{n=1}^{\infty} \mathbb{V}^*(Y_n \neq X_n) &= \sum_{n=1}^{\infty} \mathbb{V}^*(|Y_1| > d_n) < \infty, \\ |\check{\mathbb{E}}[Y_n] - \widehat{\mathbb{E}}[X_n]| + |\check{\mathbb{E}}[-Y_n] - \widehat{\mathbb{E}}[-X_n]| \\ &\leq 2\check{\mathbb{E}}[|Y_n - X_n|] \leq 2\check{\mathbb{E}}[(|Y_1| - d_n)^+] = o\left(\sqrt{\log \log n}/\sqrt{n}\right) \end{aligned}$$

by Lemma 6.1. It follows that

$$\frac{\sum_{i=1}^n |\check{\mathbb{E}}[Y_i] - \widehat{\mathbb{E}}[X_i]| + |\check{\mathbb{E}}[-Y_i] - \widehat{\mathbb{E}}[-X_i]|}{d_n} \rightarrow 0, \quad (6.30)$$

$$\sum_{n=1}^{\infty} \frac{\widehat{\mathbb{E}}[|X_n|^p]}{a_n^p} \leq c \sum_{n=1}^{\infty} \frac{\widehat{\mathbb{E}}[|Y_1|^p \wedge d_n^p]}{d_n^p} < \infty \quad (\text{by Lemma 6.1}), \quad (6.31)$$

$$\mathbb{V}^* \left(\lim_{n \rightarrow \infty} \frac{\sum_{i=1}^n (X_i - Y_i)}{d_n} \neq 0 \right) \leq \mathbb{V}^*(Y_n \neq X_n \text{ i.o.}) = 0. \quad (6.32)$$

Moreover,

$$\frac{\widehat{\mathbb{E}}[X_{n+1}^2]}{s_n^2} \leq \frac{C_0 d_n C_{\mathbb{V}}(|Y_1|)}{n} \rightarrow 0.$$

which implies

$$s_n^2 \rightarrow \infty \quad \text{and} \quad \frac{s_{n+1}^2}{s_n^2} \rightarrow 1, \quad \frac{a_{n+1}}{a_n} \rightarrow 1. \quad (6.33)$$

We first show that

$$\mathbb{V}^* \left(\limsup_{n \rightarrow \infty} \frac{\sum_{i=1}^n (Y_i - \check{\mathbb{E}}[Y_i])}{\sqrt{2n \log \log n}} > \bar{\sigma}_2 \right) = 0. \quad (6.34)$$

Without loss of generality, assume $\check{\mathbb{E}}[Y_1] = 0$. It follows from (6.31) and (6.33) that the conditions (5.12) in Remark 5.1 and (5.14) in Remark 5.2 are satisfied. By Theorem 5.1 and Remarks 5.2, 5.1, we have

$$\mathbb{V}^* \left(\limsup_{n \rightarrow \infty} \frac{\sum_{i=1}^n (X_i - \check{\mathbb{E}}[X_i])}{s_n t_n} > 1 \right) = 0,$$

which, together with (6.30) and (6.32), implies

$$\mathbb{V}^* \left(\limsup_{n \rightarrow \infty} \frac{\sum_{i=1}^n (Y_i - \check{\mathbb{E}}[Y_i])}{s_n t_n} > 1 \right) = 0.$$

Note

$$\lim_{n \rightarrow \infty} \frac{s_n^2}{n} = \bar{\sigma}^2 \quad (\text{finite and infinite}).$$

(6.34) is proved. For $-X_j$ we have similar conclusion, and therefore, Theorem 5.3 is proved.

Next, we turn to the proof of Theorem 5.4. For the part (a), besides (5.18) we assume further (5.20), i.e., $\check{\mathbb{E}}[Y_n] = \check{\mathbb{E}}[-Y_n] = 0$. It follows from (6.30) that the condition (5.7) in Theorem 5.2 is also satisfied. Then (5.21) and (5.22) are implied by (5.10) and (5.11), respectively. (5.23) is a direct corollary of (5.22).

Now, we consider the part (b). Suppose

$$C_{\mathbb{V}} \left[\frac{Y_1^2}{\log \log |Y_1|} \right] = \infty.$$

Then

$$C_{\mathbb{V}^*} \left[\frac{Y_1^2}{\log \log |Y_1|} \right] = \infty.$$

By Lemma 6.1,

$$\sum_{n=1}^{\infty} \mathbb{V}^*(|Y_n| \geq 2M d_n) \geq \sum_{n=1}^{\infty} \mathbb{V}^*(|Y_1| \geq 3M d_n) = \infty \quad \text{for all } M > 0.$$

It follows that there exists a sequence $M_n \nearrow \infty$ such that

$$\sum_{n=1}^{\infty} \mathbb{V}^*(|Y_n| \geq 2M_n d_n) = \infty.$$

By Lemma 4.3 (i), it follows that

$$\mathbb{V}^* \left(\limsup_{n \rightarrow \infty} \frac{|Y_n|}{M_n d_n} \geq 1 \right) \geq \mathbb{V}^*(|Y_n| \geq M_n d_n \text{ i.o.}) = 1.$$

Note $|Y_m| \leq |\sum_{i=1}^m Y_i| + |\sum_{i=1}^{m-1} Y_i|$. It follows that

$$\mathbb{V}^* \left(\limsup_{n \rightarrow \infty} \frac{|\sum_{i=1}^n Y_i|}{M_n d_n} \geq \frac{1}{2} \right) = 1.$$

It follows that

$$\mathbb{V}^* \left(\limsup_{n \rightarrow \infty} \frac{|\sum_{i=1}^n Y_i|}{d_n} = +\infty \right) \geq \mathbb{V}^* \left(\limsup_{n \rightarrow \infty} \frac{|\sum_{i=1}^n Y_i|}{M_n d_n} \geq \frac{1}{2} \right) = 1,$$

which contradicts (5.24). It follows that (5.18) holds. Under (5.18), $\check{\mathbb{E}}[Y_1]$ and $\check{\mathbb{E}}[-Y_1]$ exist and are finite. Let $\{n_k\} \subset \mathbb{N}$ such that $n_k \nearrow \infty$ and $n_{k-1}/n_k \rightarrow 0$. From (5.24) and the fact

$$\limsup_{k \rightarrow \infty} \frac{|\sum_{i=n_{k-1}+1}^{n_k} Y_i|}{a_{n_k}} \leq 2 \limsup_{n \rightarrow \infty} \frac{|\sum_{i=1}^n Y_i|}{a_n},$$

it follows that

$$\mathbb{V}^* \left(\limsup_{k \rightarrow \infty} \frac{|\sum_{i=n_{k-1}+1}^{n_k} Y_i|}{a_{n_k}} = \infty \right) := \tau < 1.$$

On the other hand, by Lemma 6.2, for every $\epsilon > 0$

$$\mathbb{V}^* \left(\frac{\sum_{i=1}^n Y_i - n\check{\mathbb{E}}[Y_1]}{d_n} \geq \epsilon \right) \rightarrow 0,$$

$$\mathbb{V}^* \left(\frac{\sum_{i=1}^n Y_i - n\check{\mathbb{E}}[Y_1]}{d_n} \geq -\epsilon \right) = 1 - \mathcal{V}^* \left(\frac{\sum_{i=1}^n (-Y_i + \check{\mathbb{E}}[Y_1])}{d_n} > \epsilon \right) \rightarrow 1$$

for every $\epsilon > 0$. Then

$$\mathbb{V}^* \left(\frac{\sum_{i=n_{k-1}+1}^{n_k} (Y_i - \check{\mathbb{E}}[Y_1])}{d_{n_k}} \geq -\epsilon \right) \rightarrow 1 \quad \text{for all } \epsilon > 0.$$

Note the independence of the random variables. By Lemma 4.3(i),

$$\mathbb{V}^* \left(\limsup_{k \rightarrow \infty} \frac{\sum_{i=n_{k-1}+1}^{n_k} (Y_i - \check{\mathbb{E}}[Y_1])}{d_{n_k}} \geq -2\epsilon \right) = 1 \quad \text{for all } \epsilon > 0.$$

It follows that

$$\mathbb{V}^* \left(\limsup_{k \rightarrow \infty} \frac{\sum_{i=n_{k-1}+1}^{n_k} (Y_i - \check{\mathbb{E}}[Y_1])}{d_{n_k}} \geq -2\epsilon, \quad \limsup_{k \rightarrow \infty} \frac{|\sum_{i=n_{k-1}+1}^{n_k} Y_i|}{a_{n_k}} < \infty \right) \geq 1 - \tau > 0.$$

Therefore,

$$\liminf_{k \rightarrow \infty} \frac{(n_k - n_{k-1})\check{\mathbb{E}}[Y_1]}{d_{n_k}} < \infty.$$

It follows that $\check{\mathbb{E}}[Y_1] \leq 0$. Similarly, $\check{\mathbb{E}}[-Y_1] \leq 0$. From the fact $\check{\mathbb{E}}[Y_1] + \check{\mathbb{E}}[-Y_1] \geq 0$, it follows that (5.20) holds.

Under (5.18) and (5.20), we still have (5.22) which contradicts (5.24) if $\bar{\sigma} = \infty$. Hence (5.25) holds. \square

Proof of Theorem 5.5. Note the countable sub-additivity of \mathbb{V} . We have $\mathbb{V}(A) \leq \mathbb{V}^*(A)$. By (5.21),

$$\mathbb{V} \left(\limsup_{n \rightarrow \infty} \frac{|T_n|}{d_n} > \bar{\sigma} \right) = 0.$$

It is sufficient to show that

$$\mathcal{V} \left(\limsup_{n \rightarrow \infty} \frac{T_n}{d_n} \geq \underline{\sigma} \right) = 1.$$

Note that \mathcal{V} has the property (4.1) since \mathbb{V} is countably sub-additive. It is sufficient to show that

$$\mathcal{V} \left(\limsup_{n \rightarrow \infty} \frac{T_n}{d_n} \geq \underline{\sigma} - \epsilon \right) = 1, \quad \forall \epsilon > 0,$$

because

$$\mathcal{V} \left(\limsup_{n \rightarrow \infty} \frac{T_n}{d_n} \geq \underline{\sigma} \right) = \mathcal{V} \left(\bigcap_{l=1}^{\infty} \left\{ \limsup_{n \rightarrow \infty} \frac{T_n}{d_n} \geq \underline{\sigma} - \frac{1}{l} \right\} \right).$$

When $\underline{\sigma} = 0$, the conclusion is trivial because \mathcal{V} has the property (4.1) and

$$\mathcal{V} \left(\bigcup_{n=m}^{\infty} \frac{T_n}{d_n} \geq -\epsilon \right) \geq \mathcal{V} \left(\frac{T_n}{d_n} \geq -\epsilon \right) = 1 - \mathbb{V} \left(\frac{-T_n}{d_n} > \epsilon \right) \rightarrow 1, \text{ for all } m,$$

by Lemma 6.2.

Suppose $\underline{\sigma} > 0$. Let $\lambda > 1$ be large enough. Denote $n_k = \lfloor \lambda^k \rfloor$ and $I(k) = \{n_k + 1, \dots, n_{k+1}\}$. Note $n_k/n_{k+1} \rightarrow 1/\lambda$, $d_{n_k}/d_{n_{k+1}} \rightarrow 1/\sqrt{\lambda}$,

$$\frac{T_{n_{k+1}}}{d_{n_{k+1}}} = \frac{T_{n_{k+1}} - T_{n_k}}{\sqrt{2(n_{k+1} - n_k) \log \log n_{k+1}}} \sqrt{1 - \frac{n_k}{n_{k+1}}} + \frac{T_{n_k}}{d_{n_k}} \frac{d_{n_k}}{d_{n_{k+1}}}$$

and (5.21). It is sufficient to show that,

$$\mathcal{V} \left(\limsup_{k \rightarrow \infty} \frac{T_{n_{k+1}} - T_{n_k}}{\sqrt{2(n_{k+1} - n_k) \log \log n_{k+1}}} \geq \underline{\sigma} - \epsilon \right) = 1, \quad \forall \epsilon > 0. \quad (6.35)$$

Denote $t_j = \sqrt{2 \log \log j}$ and $b_j = \alpha_j \sqrt{j} / \sqrt{2 \log \log j}$, where $\alpha_j \rightarrow 0$ is specified such that $\alpha_j \rightarrow 0$ and $\alpha_j^{1-p} t_j^{-2} \rightarrow 0$. Define

$$Z_j = (-2b_{n_{k+1}}) \vee Y_j \wedge 2b_{n_{k+1}}, \quad j \in I(k), \quad k \geq 0.$$

By Lemma 6.1 (ii), we have

$$\sum_{k=1}^{\infty} \frac{\Lambda_{n_k, n_{k+1}}(p)}{d_{n_{k+1}}^p} < \infty, \quad (6.36)$$

where

$$\Lambda_{n_k, n_{k+1}}(p) = \sum_{j \in I(k)} \widehat{\mathbb{E}}[(|X_j| \wedge d_{n_{k+1}})^p].$$

Let

$$\mathbb{N}_1 = \left\{ k \in \mathbb{N}; \frac{\Lambda_{n_k, n_{k+1}}(p)}{d_{n_{k+1}}^p} \leq t_{n_{k+1}}^{-2p} \right\}. \quad (6.37)$$

Similar to (6.10) and (6.11), we have for $k \in \mathbb{N}_1$,

$$\begin{aligned} & \frac{\sum_{j \in I(k)} \widehat{\mathbb{E}}[(|-d_{n_{k+1}}| \vee Y_j \wedge d_{n_{k+1}} - Z_j)^p]}{d_{n_{k+1}}^p} \\ & \leq \frac{\Lambda_{n_k, n_{k+1}}(p)}{d_{n_{k+1}}^p} \alpha_{n_{k+1}}^{1-p} t_{n_{k+1}}^{2p-2} \leq \alpha_{n_{k+1}}^{1-p} t_{n_{k+1}}^{-2} \rightarrow 0, \end{aligned} \quad (6.38)$$

$$\begin{aligned} & \frac{\sum_{j \in I(k)} \widehat{\mathbb{E}}[(|-d_{n_{k+1}}| \vee Y_j \wedge a_{n_{k+1}} - Z_j)^2]}{n_{k+1}} \\ & \leq \frac{\Lambda_{n_k, n_{k+1}}(p)}{d_{n_{k+1}}^p} \alpha_{n_{k+1}}^{2-p} t_{n_{k+1}}^{2p-2} \leq \alpha_{n_{k+1}}^{2-p} t_{n_{k+1}}^{-2} \rightarrow 0, \end{aligned} \quad (6.39)$$

by noting $\alpha_j \rightarrow 0$ such that $\alpha_j^{1-p} t_j^{-2} \rightarrow 0$. Similar to (6.12), we have that for $k \in \mathbb{N}_1$ large enough,

$$\begin{aligned} & \mathbb{V}\left(\sum_{j \in I(k)} |Y_j - Z_j| \geq \epsilon d_{n_{k+1}}\right) \\ & \leq \mathbb{V}\left(\sum_{j \in I(k)} |(-d_{n_{k+1}}) \vee Y_j \wedge d_{n_{k+1}} - Z_j| \geq \epsilon d_{n_{k+1}}\right) + \sum_{j \in I(k)} \mathbb{V}(|X_j| > d_{n_{k+1}}) \\ & \leq \exp\{-2t_{n_{k+1}}^2\} + C \frac{\Lambda_{n_k, n_{k+1}}(p)}{d_{n_{k+1}}^p} + 2 \sum_{j \in I(k)} \mathbb{V}(X_j > \epsilon' d_j). \end{aligned}$$

It follows that

$$\sum_{k \in \mathbb{N}_1} \mathbb{V}\left(\sum_{j \in I(k)} |Y_j - Z_j| \geq \epsilon d_{n_{k+1}}\right) < \infty. \quad (6.40)$$

Next, we apply Lemma 3.3 to the array $\{Z_j; j \in I(k)\}$ of independent and identically random variables, $k \in \mathbb{N}_1$. By (6.39), we have $\widehat{\mathbb{E}}[Z_j^2] \sim \widehat{\mathbb{E}}[Y_1^2 \wedge d_{n_{k+1}}^2] \rightarrow \bar{\sigma}^2$ and $\widehat{\mathcal{E}}[Z_j^2] \sim \widehat{\mathcal{E}}[Y_1^2 \wedge d_{n_{k+1}}^2] \rightarrow \underline{\sigma}^2$. By (6.38), Lemma 6.1 (iii) and the fact that $\check{\mathbb{E}}[Y_1] = \check{\mathbb{E}}[-Y_1] = 0$, we have

$$\frac{\sum_{j \in I(k)} (|\widehat{\mathbb{E}}[Z_j]| + |\widehat{\mathcal{E}}[Z_j]|)}{\sqrt{n_{k+1} - n_k} t_{n_{k+1}}} \rightarrow 0.$$

Note

$$|Z_j| \leq 2b_{n_{k+1}} = o\left(\sqrt{n_{k+1} - n_k}/t_{n_{k+1}}\right), \quad j \in I(k).$$

Applying Lemma 3.3 with $k_n = n_{k+1} - n_k$ and $y_n = t_{n_{k+1}}$ yields

$$\begin{aligned} & \mathcal{V}\left(\frac{\sum_{j \in I(k)} Z_j}{\sqrt{2(n_{k+1} - n_k) \log \log n_{k+1}}} \geq \underline{\sigma}(1 - \epsilon)\right) \\ & \geq \exp\{-(1 - \epsilon) \log \log n_{k+1}\} \geq ck^{-(1-\epsilon)}, \end{aligned} \quad (6.41)$$

for $k \in \mathbb{N}_1$ large enough. Note

$$\sum_{k \notin \mathbb{N}_1} \exp\{-(1 - \epsilon) \log \log n_{k+1}\} \leq C \sum_{k \notin \mathbb{N}_1} t_{n_{k+1}}^{-2p} < \infty, \quad (6.42)$$

by (6.36) and (6.37). From (6.40), (6.41) and (6.42), we conclude that

$$\sum_{k \in \mathbb{N}_1} \mathcal{V}\left(\frac{\sum_{j \in I(k)} Y_j}{\sqrt{2(n_{k+1} - n_k) \log \log n_{k+1}}} \geq \underline{\sigma}(1 - \epsilon)\right) = \infty.$$

Hence,

$$\sum_{k=1}^{\infty} \mathcal{V}\left(\frac{T_{n_{k+1}} - T_{n_k}}{\sqrt{2(n_{k+1} - n_k) \log \log n_{k+1}}} \geq \underline{\sigma}(1 - \epsilon)\right) = \infty, \quad \forall \epsilon > 0.$$

Note the independence of random variables. By Lemma 4.1 (iii), we have

$$\mathcal{V}\left(\frac{T_{n_k} - T_{n_{k-1}}}{\sqrt{2(n_k - n_{k-1}) \log \log n_k}} \geq \underline{\sigma}(1 - \epsilon) - \epsilon \text{ i.o.}\right) = 1, \quad \forall \epsilon > 0.$$

(6.35) is proved. \square

References

- [1] Billingsley, P. (1968) *Convergence of Probability Measures*, Wiley, New York.
- [2] Chen, Z. J. and Hu, F. (2014). A law of the iterated logarithm for sublinear expectations. *Journal of Financial Engineering*, **1**(2). arXiv: 1103.2965v2[math.PR].
- [3] Chen, X. (1993). On the law of the iterated logarithm for independent Banach space valued random variables. *Ann. Probab.*, **21**(4):1991-2011.
- [4] Fuk, D. K. and Nagaev, S. V. (1971). Probability inequalities for sums of independent random variables. *Teor. Veroyatn. Primen.*, **16**(4): 660-675.
- [5] Hartman, P. and Wintner, A. (1941). On the law of iterated logarithm. *Amer. J. Math.*, **63**:169-176.
- [6] Kolmogorov, A. (1929). Uber das Gesetz des iterierten Logarithmus. *Math. Ann.*, **101**: 126-135.
- [7] Peng, S. (2008). A new central limit theorem under sublinear expectations. ArXiv:0803.2656v1.
- [8] Peng S. (2009). Survey on normal distributions, central limit theorem, Brownian motion and the related stochastic calculus under sublinear expectations. *Sci. China Ser. A*, **52**(7): 1391-1411
- [9] Peng, S. (2019). *Nonlinear Expectations and Stochastic Calculus under Uncertainty with Robust CLT and G-Brownian Motion*. Springer.
- [10] Peng, S., Yang, S. Z. and Yao, J. F. (2020). Improving Value-at-Risk prediction under model uncertainty. arXiv:1805.03890 [math.PR] (2020).
- [11] Peng, S. and Zhou, Q.(2020). A hypothesis-testing perspective on the G-normal distribution theory. *Stat. & Probab. Lett.*, **156**. <https://doi.org/10.1016/j.spl.2019.108623>.
- [12] Petrov, V. V. (1995). *Limit Theorem of Probability Theory- Sequences of independent Random Variables*. Clarendon Press·Oxford.
- [13] Stout, W. F. (1974). *Almost sure convergence*. Academic Press, New York.
- [14] Strassen, V. (1966). A converse of the law of the iterated logarithm. *Z. Wahrsch. Verw. Geb.*, **4**: 265-268.

- [15] Wittmann, R. (1985). A general law of iterated logarithm. *Z. Wahrsch. Verw. Gebiete*, **68**: 521-543.
- [16] Wittmann, R. (1987). Sufficient moment and truncated moment conditions for the law of iterated logarithm. *Probab. Theory Related Fields*, **75**: 509-530.
- [17] Zhang, L.-X. (2016). Exponential inequalities under the sub-linear expectations with applications to laws of the iterated logarithm. *Sci. China Math.*, **59**(12), 2503-2526.
- [18] Zhang, L.-X. (2020). The convergence of the sums of independent random variables under the sub-linear expectations, *Acta Mathematica Sinica, English Series*, **36** (3), 224-244.
- [19] Zhang, L.-X. (2021). Lindeberg's central limit theorems for martingale like sequences under sub-linear expectations, *Sci. China Math.*, to appear. <https://doi.org/10.1007/s11425-018-9556-7>