

ANTI-SELF-DUAL CONNECTIONS OVER THE 5D HEISENBERG GROUP AND THE TWISTOR METHOD

GUANGZHEN REN AND WEI WANG

ABSTRACT. In this paper, we introduce notions of α -planes in 5D complex Heisenberg group and the twistor space as the moduli space of all α -planes. So we can define an anti-self-dual (ASD) connection as a connection flat over all α -planes. This geometric approach allows us to establish Penrose-Ward correspondence between ASD connections over 5D complex Heisenberg group and a class of holomorphic vector bundles on the twistor space. By Atiyah-Ward ansatz, we also construct a family of ASD connections on 5D complex Heisenberg group. When restricted to 5D real Heisenberg group, the flat model of 5D contact manifolds, an ASD connection satisfies the horizontal part of the contact instanton equation introduced by physicists.

1. INTRODUCTION

Since 3D Chern-Simons theory is very successful, people are interested in its generalization to higher odd dimension. When studying 5D Chern-Simons theory, physicists introduced the supersymmetric Yang-Mills equation on 5D contact manifolds (c.f. [6] [8]). In particular, they introduced the *contact instanton equation* on 5D contact manifolds as

$$(1) \quad \mathcal{F} = \pm \iota_T * \mathcal{F},$$

where \mathcal{F} is the curvature for a connection, T is the Reeb vector field and ι_T is the contraction with T . Hosomichi-Seong-Terashima [6] constructed a 5D $\mathcal{N} = 1$ supersymmetric Yang-Mills theory on the five sphere, and showed the fields in a vector multiplet localized to contact instantons by using the localization technique. Källén-Zabzine [8] localized the path integral of Chern-Simons theory on circle fibrations over 4D symplectic manifolds to contact instantons. On the other hand, Itoh [7] has constructed the CR twistor space over 5D contact manifolds by using differential geometry about two decades ago. Wolf [17] used the twistor method to study the contact instanton equation on contact manifolds.

In the classical flat case, by complexifying 4D Minkowski space as \mathbb{C}^4 , ones can construct twistor space by using complex geometry method (cf. [2] [10] [16]). If we denote a point of \mathbb{C}^4 by

$$\mathbf{y} = \begin{pmatrix} y_{00'} & y_{01'} \\ y_{10'} & y_{11'} \end{pmatrix},$$

Key words and phrases. the twistor method; 5D Heisenberg group; anti-self-dual connection; the contact instanton equation; Penrose-Ward correspondence; Atiyah-Ward ansatz; ASD Yang-Mills equation.

The first author is partially supported by National Nature Science Foundation in China (No. 11801508, 11801523, 11971425); The second author is partially supported by National Nature Science Foundation in China (No. 11971425).

an α -plane in \mathbb{C}^4 is the set of all \mathbf{y} satisfying

$$(2) \quad \begin{pmatrix} y_{00'} & y_{01'} \\ y_{10'} & y_{11'} \end{pmatrix} \begin{pmatrix} \pi_{0'} \\ \pi_{1'} \end{pmatrix} = \begin{pmatrix} \omega_0 \\ \omega_1 \end{pmatrix}$$

for fixed $0 \neq (\pi_{0'}, \pi_{1'}) \in \mathbb{C}^2$ and $w_0, w_1 \in \mathbb{C}$. The moduli space of all α -planes is the twistor space \mathcal{P}_0 , which is an open subset of $\mathbb{C}P^3$. Then we have the double fibration

$$\begin{array}{ccc} & \mathbb{C}^4 \times \mathbb{C}P^1 & \\ \eta \swarrow & & \searrow \tau \\ \mathcal{P}_0 & & \mathbb{C}^4, \end{array}$$

based on which there exists the Penrose correspondence between the solution of massless field equations and the first cohomology groups of certain line bundles over \mathcal{P}_0 , and Penrose-Ward correspondence between the solutions of ASD Yang-Mills equation and holomorphic bundles over \mathcal{P}_0 , trivial over the complex projective line $\hat{x} = \eta \circ \tau^{-1}(x)$ for any $x \in \mathbb{C}^4$. In this paper, we consider the simplest 5D contact manifold, 5D real Heisenberg group, and complexify it as \mathbb{C}^5 . We will generalize this theory to 5D Heisenberg group.

5D complex *Heisenberg group* \mathcal{H} is $\mathbb{C}^5 := \{(\mathbf{y}, t) | \mathbf{y} \in \mathbb{C}^4, t \in \mathbb{C}\}$ with the multiplication given by

$$(3) \quad (\mathbf{y}, t) \circ (\mathbf{y}', t') = (\mathbf{y} + \mathbf{y}', t + t' + B(\mathbf{y}, \mathbf{y}')),$$

where $B(\mathbf{y}, \mathbf{y}') = y_{00'}y'_{11'} - y_{01'}y'_{10'} + y_{10'}y'_{01'} - y_{11'}y'_{00'}$. We have left invariant vector fields on \mathcal{H} :

$$(4) \quad \begin{aligned} V_{00'} &:= \frac{\partial}{\partial y_{00'}} - y_{11'}T, & V_{01'} &:= \frac{\partial}{\partial y_{01'}} + y_{10'}T, \\ V_{10'} &:= \frac{\partial}{\partial y_{10'}} - y_{01'}T, & V_{11'} &:= \frac{\partial}{\partial y_{11'}} + y_{00'}T, & T &:= \frac{\partial}{\partial t}. \end{aligned}$$

It is easy to see that

$$(5) \quad [V_{00'}, V_{11'}] = [V_{10'}, V_{01'}] = 2T,$$

and all other brackets vanish. Consequently, for fixed $0 \neq (\pi_{0'}, \pi_{1'}) \in \mathbb{C}^2$, if denote

$$(6) \quad V_A := \pi_{0'}V_{A0'} - \pi_{1'}V_{A1'}, \quad A = 0, 1,$$

we have

$$[V_0, V_1] = 0.$$

Namely, $\text{span}\{V_0, V_1\}$ is an abelian Lie subalgebra and an integrable distribution for fixed $0 \neq (\pi_{0'}, \pi_{1'}) \in \mathbb{C}^2$. Their integral surfaces are hyperplanes (cf. (25)), which we also call α -planes. The *twistor space* \mathcal{P} is the moduli space of all α -planes, which is a 4D complex manifold. We have the double fibration over 5D complex Heisenberg group as follows

$$(7) \quad \begin{array}{ccc} & \mathcal{F} = \mathbb{C}^5 \times \mathbb{C}P^1 & \\ \eta \swarrow & & \searrow \tau \\ \mathcal{P} & & \mathcal{H} \cong \mathbb{C}^5. \end{array}$$

A connection is called *anti-self-dual* (briefly ASD) if it is flat over any α -plane. Let $\Phi = \Phi_{00'}\theta^{00'} + \Phi_{10'}\theta^{10'} + \Phi_{01'}\theta^{01'} + \Phi_{11'}\theta^{11'} + \Phi_T\theta$ be a \mathfrak{g} -valued connection form on \mathcal{H} , where $\{\theta^{AA'}, \theta\}$ are 1-forms dual to $\{V_{AA'}, T\}$. Φ is ASD if and only if it satisfies the *ASD Yang-Mills equation*

$$(8) \quad \begin{cases} V_{00'}(\Phi_{10'}) - V_{10'}(\Phi_{00'}) + [\Phi_{00'}, \Phi_{10'}] = 0, \\ V_{01'}(\Phi_{10'}) + V_{00'}(\Phi_{11'}) - V_{10'}(\Phi_{01'}) - V_{11'}(\Phi_{00'}) + [\Phi_{00'}, \Phi_{11'}] + [\Phi_{01'}, \Phi_{10'}] = 0, \\ V_{01'}(\Phi_{11'}) - V_{11'}(\Phi_{01'}) + [\Phi_{01'}, \Phi_{11'}] = 0. \end{cases}$$

An open subset U of \mathcal{H} is called *elementary* if for every α -plane $\tilde{Z} = \tau \circ \eta^{-1}(Z)$ ($Z \in \mathcal{P}$) the intersection $\tilde{Z} \cap U$ is connected and simply connected. Then we have Penrose-Ward correspondence.

Theorem 1.1. *Let U be an elementary open set in \mathcal{H} . There is a one-to-one correspondence between*

- (1) *gauge equivalence classes of ASD connections with the gauge group $\mathrm{GL}(n, \mathbb{C})$ over U ;*
- (2) *holomorphic vector bundles $E' \rightarrow \hat{U} = \eta \circ \tau^{-1}(U)$ such that $E'|_{\hat{x}}$ is trivial, where $\hat{x} := \eta \circ \tau^{-1}(x)$ for each $x \in U$.*

Define the *Sub-Laplacian*

$$\Delta_b := V_{00'}V_{11'} - V_{10'}V_{01'},$$

and *partial exterior differential operators* d_0 and d_1 by

$$(9) \quad d_0 f := V_{00'}f \cdot \theta^{00'} + V_{10'}f \cdot \theta^{10'}, \quad d_1 f := V_{01'}f \cdot \theta^{01'} + V_{11'}f \cdot \theta^{11'}$$

for a function $f \in C^\infty(U, \mathbb{C})$. By using Atiyah-Ward ansatz, we can construct a family of ASD connections.

Theorem 1.2. (1) *If φ satisfies*

$$(10) \quad \Delta_b \varphi = 0,$$

then the connection form

$$(11) \quad \Phi = \left[\begin{array}{cc} \frac{1}{2}(d_0 \ln \varphi - d_1 \ln \varphi) & V_{01'}(\ln \varphi)\theta^{00'} + V_{11'}(\ln \varphi)\theta^{10'} \\ V_{00'}(\ln \varphi)\theta^{01'} + V_{10'}(\ln \varphi)\theta^{11'} & -\frac{1}{2}(d_0 \ln \varphi - d_1 \ln \varphi) \end{array} \right] + \Phi_T \theta$$

is ASD.

(2) *In particular,*

$$(12) \quad \varphi := \frac{1}{\|\mathbf{y}\|^4 - t^2}, \quad \text{where} \quad \|\mathbf{y}\|^2 = \det \begin{bmatrix} y_{00'} & y_{01'} \\ y_{10'} & y_{11'} \end{bmatrix},$$

is the solution to (10) on $\mathcal{H} \setminus \{\|\mathbf{y}\|^4 = t^2\}$.

Now we consider 5D real Heisenberg group $\mathcal{H}^{\mathbb{R}} \cong \mathbb{R}^5$ with multiplication given by

$$(13) \quad (\mathbf{y}, s) \circ (\mathbf{y}', s') = (\mathbf{y} + \mathbf{y}', s + s' + \langle \mathbf{y}, \mathbf{y}' \rangle),$$

where $\langle \mathbf{y}, \mathbf{y}' \rangle = 2(y_1 y'_2 - y_2 y'_1 - y_3 y'_4 + y_4 y'_3)$, $\mathbf{y}, \mathbf{y}' \in \mathbb{R}^4$ and $s, s' \in \mathbb{R}$. By the real imbedding $\mathbb{R}^5 \rightarrow \mathbb{C}^5$ given by

$$(14) \quad \begin{bmatrix} y_{00'} & y_{01'} \\ y_{10'} & y_{11'} \end{bmatrix} := \begin{bmatrix} y_1 + \mathbf{i}y_2 & -y_3 + \mathbf{i}y_4 \\ y_3 + \mathbf{i}y_4 & y_1 - \mathbf{i}y_2 \end{bmatrix}, \quad t = -\mathbf{i}s,$$

$\mathcal{H}^{\mathbb{R}}$ is a subgroup of \mathcal{H} . It is a flat model of 5D contact manifolds.

Recall the contact instanton equation on 5D contact manifolds [6] [8]. For a connection ∇ , let us consider its Yang-Mills action

$$YM(\nabla) = - \int_{\mathcal{H}^{\mathbb{R}}} \text{Tr}(F \wedge *F),$$

where F is the curvature of ∇ and $*$ is the Hodge star over $\mathcal{H}^{\mathbb{R}}$. F_H and F_V are horizontal and vertical part of F respectively, which satisfy $\iota_T F_H = 0$ and $\iota_T F_V \neq 0$ respectively. As $F_H \wedge *F_V = 0$ and $F_V \wedge *F_H = 0$, we have

$$YM(\nabla) = - \int_{\mathcal{H}^{\mathbb{R}}} \text{Tr}(F_H \wedge *F_H + F_V \wedge *F_V).$$

Take F_H^+ and F_H^- as horizontal self-dual and anti-self-dual parts of F , which satisfy $\iota_T *F_H^+ = F_H^+$ and $\iota_T *F_H^- = -F_H^-$ respectively. $F_H^+ = 0$ or $F_H^- = 0$ together with $F_V = 0$ are critical points of $YM(\nabla)$. The anti-self-dual contact instanton equation [6] [8] is

$$(15) \quad F_H^+ = 0 \quad \text{and} \quad F_V = 0,$$

while the self-dual one is

$$F_H^- = 0 \quad \text{and} \quad F_V = 0.$$

The ASD equations (8) restricted to $\mathcal{H}^{\mathbb{R}}$ is exactly $F_H^+ = 0$. So it satisfies (15) without $F_V = 0$. If φ in (11) is replaced by

$$\varphi^{\mathbb{R}} = \frac{1}{|y|^4 + s^2}, \quad \text{where } |y| = (y_1^2 + y_2^2 + y_3^2 + y_4^2)^{\frac{1}{2}},$$

we get ASD connection forms on 5D real Heisenberg group $\mathcal{H}^{\mathbb{R}}$.

Baston and Easteod [1] generalize the twistor theory to a general setting based on the double fibration

$$(16) \quad \begin{array}{ccc} & G/(P \cap Q) & \\ \eta \swarrow & & \searrow \tau \\ G/Q & & G/P, \end{array}$$

where G is a complex semisimple Lie group, P, Q are its parabolic subgroups. If we take $G = \text{SO}(6, \mathbb{C})$ and suitable subgroups P and Q , by using the method in [13], we can also write down local coordinate charts of the above homogeneous spaces and the mapping η and τ in terms of local coordinates to obtain (7).

The construction of this paper is based on the fact that V_0 and V_1 in (6) span an abelian subalgebra. Its real version plays a very important role in developing a theory of quaternionic Monge-Ampère operator in [14] and tangential k -Cauchy-Fueter complex [12] over the Heisenberg group.

In Section 2, we introduce the twistor transform for 5D complex Heisenberg group and derive the ASD Yang-Mills equation. In Section 3, we give Penrose-Ward correspondence between ASD connections and holomorphic vector bundles over the twistor space, which are trivial over a class of projective lines in the twistor space, and construct a family of ASD connections by using Atiyah-Ward ansatz. In Section 4, by the real imbedding of $\mathcal{H}^{\mathbb{R}}$ into \mathcal{H} , we find that the ASD Yang-Mills equation coincides with the horizontal part of the contact instanton

equation. In Appendix, by constructing local coordinate charts of homogeneous spaces in the double fibration (16) with $G = \text{SO}(6, \mathbb{C})$, we reproduce the basic ingredients of twistor method for 5-D complex Heisenberg group.

2. THE TWISTOR TRANSFORM ON 5D COMPLEX HEISENBERG GROUP

2.1. α -planes and the twistor space of 5D complex Heisenberg group. Define a symmetric product $\langle \cdot, \cdot \rangle$ on \mathbb{C}^2 by

$$(17) \quad \langle \mathbf{w}, \tilde{\mathbf{w}} \rangle := w_1 \tilde{w}_2 + \tilde{w}_1 w_2$$

for $\mathbf{w} = (w_1, w_2)$, $\tilde{\mathbf{w}} = (\tilde{w}_1, \tilde{w}_2) \in \mathbb{C}^2$. If we denote $\mathbf{y} = (\mathbf{y}_{0'}, \mathbf{y}_{1'})$ with $\mathbf{y}_{0'} = (y_{00'}, y_{10'})$ and $\mathbf{y}_{1'} = (y_{01'}, y_{11'})$, the multiplication (3) of the Heisenberg group can be also written as

$$(18) \quad (\mathbf{y}_{0'}, \mathbf{y}_{1'}, t) \circ (\tilde{\mathbf{y}}_{0'}, \tilde{\mathbf{y}}_{1'}, \tilde{t}) = (\mathbf{y}_{0'} + \tilde{\mathbf{y}}_{0'}, \mathbf{y}_{1'} + \tilde{\mathbf{y}}_{1'}, t + \tilde{t} + \langle \mathbf{y}_{0'}, \tilde{\mathbf{y}}_{1'} \rangle - \langle \mathbf{y}_{1'}, \tilde{\mathbf{y}}_{0'} \rangle).$$

Recall the *left translation*: for fixed $(\mathbf{y}', t') \in \mathcal{H}$,

$$(19) \quad \tau_{(\mathbf{y}', t')} : (\mathbf{y}, t) \mapsto (\mathbf{y}', t') \cdot (\mathbf{y}, t), \quad (\mathbf{y}, t) \in \mathcal{H}$$

and the *dilation*:

$$(20) \quad \delta_r : (\mathbf{y}, t) \mapsto (r\mathbf{y}, r^2 t),$$

on the Heisenberg group. A vector field V over \mathcal{H} is called *left invariant* if for any $(\mathbf{y}', t') \in \mathcal{H}$, we have

$$\tau_{(\mathbf{y}', t')}^* V = V,$$

where $\tau_{(\mathbf{y}', t')}$ is the left translation in (19). Define

$$(21) \quad (V_{AA'} f)(\mathbf{y}, t) := \left. \frac{d}{ds} f((\mathbf{y}, t)(se_{AA'})) \right|_{s=0}, \quad (Tf)(\mathbf{y}, t) := \left. \frac{d}{ds} f((\mathbf{y}, t)(se_0)) \right|_{s=0}$$

for $A = 0, 1, A' = 0', 1'$, where $e_{AA'}$ is a vector in \mathbb{C}^5 with all entries vanishing except for the (AA') -entry to be 1, and $e_0 = (0, 0, 0, 0, 1)$. For example,

$$(22) \quad \begin{aligned} V_{00'} f &:= \left. \frac{d}{ds} f((\mathbf{y}, t)(se_{00'})) \right|_{s=0} = \left. \frac{d}{ds} f((\mathbf{y}, t)(s, 0, 0, 0, 0)) \right|_{s=0} \\ &= \left. \frac{d}{ds} f(y_{00'} + s, y_{10'}, y_{01'}, y_{11'}, t - sy_{11'}) \right|_{s=0} \\ &= \left(\frac{\partial}{\partial y_{00'}} - y_{11'} \frac{\partial}{\partial t} \right) f. \end{aligned}$$

We can describe α -planes, the integral surfaces of V_0 and V_1 , explicitly as follows. $\mathbb{C}^5 \times \mathbb{C}P^1$ is the complex manifold with two coordinate charts $\mathbb{C}^5 \times \mathbb{C}$ and $\mathbb{C}^5 \times \mathbb{C}$, glued by the mapping $\kappa : \mathbb{C}^5 \times \mathbb{C} \setminus \{0\} \rightarrow \mathbb{C}^5 \times \mathbb{C} \setminus \{0\}$ given by

$$(23) \quad (\mathbf{y}, \zeta) \mapsto (\mathbf{y}, \zeta^{-1}).$$

Then if we use the nonhomogeneous coordinates, $\tau : \mathbb{C}^5 \times \mathbb{C}P^1 \rightarrow \mathbb{C}^5$ is given by $(\mathbf{y}, \zeta) \rightarrow \mathbf{y}$ and $(\mathbf{y}, \tilde{\zeta}) \rightarrow \mathbf{y}$, and the vector field V_A in (6) can be rewritten as

$$V_A = \pi_{1'} V_A^\zeta = \pi_{0'} \tilde{V}_A^{\tilde{\zeta}},$$

where $\zeta = \frac{\pi_{0'}}{\pi_{1'}}$, $\tilde{\zeta} = \frac{\pi_{1'}}{\pi_{0'}}$ and

$$V_A^\zeta = \zeta V_{A0'} - V_{A1'}, \quad \tilde{V}_A^{\tilde{\zeta}} = V_{A0'} - \tilde{\zeta} V_{A1'}.$$

Let us check that the integral surfaces of V_A^ζ and $\tilde{V}_A^{\tilde{\zeta}}$ lifted to $\mathbb{C}^5 \times \mathbb{C}P^1$ by τ are the fiber of the mapping $\eta : \mathbb{C}^5 \times \mathbb{C} \rightarrow \mathbb{C}^4$ and $\tilde{\eta} : \mathbb{C}^5 \times \mathbb{C} \rightarrow \mathbb{C}^4$, respectively.

Proposition 2.1. *Let $\eta : \mathbb{C}^5 \times \mathbb{C} \rightarrow W \cong \mathbb{C}^4$ be the mapping given by*

$$(24) \quad \omega = \eta(\mathbf{y}, t, \zeta) = \begin{pmatrix} \eta_0(\mathbf{y}, t, \zeta) \\ \eta_1(\mathbf{y}, t, \zeta) \\ \eta_2(\mathbf{y}, t, \zeta) \\ \eta_3(\mathbf{y}, t, \zeta) \end{pmatrix} = \begin{pmatrix} y_{00'} + \zeta y_{01'} \\ y_{10'} + \zeta y_{11'} \\ t - \langle \mathbf{y}_{0'} + \zeta \mathbf{y}_{1'}, \mathbf{y}_{1'} \rangle \\ \zeta \end{pmatrix} \in W.$$

Then $\tau \circ \eta^{-1}(w)$ is a 2-D plane parameterized as

$$(25) \quad \begin{cases} y_{01'} = s_0, \\ y_{11'} = s_1, \\ y_{00'} = \omega_0 - \zeta s_0, \\ y_{10'} = \omega_1 - \zeta s_1, \\ t = \omega_2 + s_1 \omega_0 + s_0 \omega_1, \end{cases}$$

with parameters $s_0, s_1 \in \mathbb{C}$. V_0^ζ and V_1^ζ are tangential to this plane, and so it is an α -plane.

Proof. It is direct to see that $V_{AA'}(y_{BB'}) = \delta_{AB} \delta_{A'B'}$ by the expression of $V_{AA'}$'s in (4). So we have

$$V_A^\zeta(\eta_j(\mathbf{y}, t, \zeta)) = 0, \quad j = 0, 1.$$

Noting that

$$(26) \quad \langle \mathbf{y}_{0'} + \zeta \mathbf{y}_{1'}, \mathbf{y}_{1'} \rangle = y_{00'} y_{11'} + y_{10'} y_{01'} + 2\zeta y_{01'} y_{11'},$$

we have

$$\begin{aligned} V_0^\zeta(\eta_2(\mathbf{y}, t, \zeta)) &= \zeta(-y_{11'} - y_{11'}) - (y_{10'} - y_{10'} - 2\zeta y_{11'}) = 0, \\ V_1^\zeta(\eta_2(\mathbf{y}, t, \zeta)) &= \zeta(-y_{01'} - y_{01'}) - (y_{00'} - y_{00'} - 2\zeta y_{01'}) = 0. \end{aligned}$$

Thus V_0^ζ and V_1^ζ are tangential to each fiber of η . Note that for a fixed point $\omega = (\omega_0, \omega_1, \omega_2, \zeta) \in W$, $\eta^{-1}(w)$ in $\mathbb{C}^5 \times \mathbb{C}$ has fixed last coordinate ζ . So $\tau \circ \eta^{-1}(w)$ is the plane determined by

$$(27) \quad \eta_0(\mathbf{y}, t, \zeta) = \omega_0, \quad \eta_1(\mathbf{y}, t, \zeta) = \omega_1, \quad \eta_2(\mathbf{y}, t, \zeta) = \omega_2.$$

The solutions of linear equations $\begin{cases} y_{00'} + \zeta y_{01'} = \omega_0 \\ y_{10'} + \zeta y_{11'} = \omega_1 \end{cases}$ are given by $y_{01'} = s_0$, $y_{00'} = \omega_0 - s_0 \zeta$, $y_{11'} = s_1$, $y_{10'} = \omega_1 - s_1 \zeta$ with parameters $s_0, s_1 \in \mathbb{C}$. Then t is given by the last equation in (25) by the third equation of (24). \square

On the other hand, if $\pi_{0'} \neq 0$, integral surfaces of $\tilde{V}_0^{\tilde{\zeta}}$ and $\tilde{V}_1^{\tilde{\zeta}}$ are fibers of the mapping $\tilde{\eta} : \mathbb{C}^5 \times \mathbb{C} \rightarrow \tilde{W} \cong \mathbb{C}^4$ given by

$$(28) \quad \tilde{\omega} = \tilde{\eta}(\mathbf{y}, t, \tilde{\zeta}) = \begin{pmatrix} \tilde{\eta}_0(\mathbf{y}, t, \tilde{\zeta}) \\ \tilde{\eta}_1(\mathbf{y}, t, \tilde{\zeta}) \\ \tilde{\eta}_2(\mathbf{y}, t, \tilde{\zeta}) \\ \tilde{\eta}_3(\mathbf{y}, t, \tilde{\zeta}) \end{pmatrix} = \begin{pmatrix} \tilde{\zeta} y_{00'} + y_{01'} \\ \tilde{\zeta} y_{10'} + y_{11'} \\ t + \langle \tilde{\zeta} \mathbf{y}_{0'} + \mathbf{y}_{1'}, \mathbf{y}_{0'} \rangle \\ \tilde{\zeta} \end{pmatrix} \in \tilde{W},$$

and for $\tilde{\omega} = (\tilde{w}_0, \tilde{w}_1, \tilde{w}_2, \tilde{\zeta}) \in \tilde{W}$, the α -plane $\tau \circ \tilde{\eta}^{-1}(\tilde{\omega})$ is given by

$$(29) \quad \begin{cases} y_{01'} = \tilde{s}_0 \\ y_{11'} = \tilde{s}_1 \\ y_{00'} = \tilde{\omega}_0 - \tilde{\zeta}\tilde{s}_0 \\ y_{10'} = \tilde{\omega}_1 - \tilde{\zeta}\tilde{s}_1 \\ t = \tilde{\omega}_2 + \tilde{s}_1\tilde{\omega}_0 + \tilde{s}_0\tilde{\omega}_1, \end{cases}$$

with parameters $\tilde{s}_0, \tilde{s}_1 \in \mathbb{C}$. If (\mathbf{y}, t, ζ) satisfies (27) with $\omega_0, \omega_1, \omega_2, \zeta = \tilde{\zeta}^{-1} \in \mathbb{C}$, then we have

$$(30) \quad \tilde{\eta}_A(\mathbf{y}, t, \tilde{\zeta}) = \zeta^{-1}\omega_A, \quad A = 0, 1,$$

and

$$(31) \quad \begin{aligned} \tilde{\eta}_2(\mathbf{y}, t, \tilde{\zeta}) &= t + \langle \zeta^{-1}\mathbf{y}_{0'} + \mathbf{y}_{1'}, \mathbf{y}_{0'} \rangle \\ &= t + y_{01'}y_{10'} + y_{11'}y_{00'} + 2\zeta^{-1}y_{00'}y_{10'} \\ &= t - \langle \mathbf{y}_{0'} + \zeta\mathbf{y}_{1'}, \mathbf{y}_{1'} \rangle + 2\zeta^{-1}(y_{00'} + \zeta y_{01'})(y_{10'} + \zeta y_{11'}) \\ &= \omega_2 + 2\zeta^{-1}\omega_0\omega_1, \end{aligned}$$

by (26). So $\kappa(\mathbf{y}, t, \zeta) = (\mathbf{y}, t, \tilde{\zeta})$ maps a fiber of η over $(\omega_0, \omega_1, \omega_2, \zeta)$ to a fiber of $\tilde{\eta}$ over $(\tilde{\omega}_0, \tilde{\omega}_1, \tilde{\omega}_2, \tilde{\zeta})$ with the mapping

$$(32) \quad \begin{aligned} \Phi: W \setminus \{\zeta = 0\} &\rightarrow \tilde{W} \setminus \{\tilde{\zeta} = 0\} \\ (w_0, w_1, w_2, \zeta) &\mapsto (\tilde{\omega}_0, \tilde{\omega}_1, \tilde{\omega}_2, \tilde{\zeta}) = (\zeta^{-1}w_0, \zeta^{-1}w_1, w_2 + 2\zeta^{-1}w_1w_2, \zeta^{-1}), \end{aligned}$$

which glues W and \tilde{W} to get a complex manifold \mathcal{P} . It is the moduli space of all α -planes, which is our twistor space.

Moreover, we have the commutative diagram

$$(33) \quad \begin{array}{ccc} \mathbb{C}^5 \times \mathbb{C} & \xrightarrow{\kappa} & \mathbb{C}^5 \times \mathbb{C} \\ \downarrow \eta & & \downarrow \tilde{\eta} \\ W & \xrightarrow{\Phi} & \tilde{W} \end{array}$$

In fact, for $(\mathbf{y}_{0'}, \mathbf{y}_{1'}, \zeta) \in \mathbb{C}^5 \times \mathbb{C}$,

$$(34) \quad \begin{aligned} \Phi \circ \eta(\mathbf{y}_{0'}, \mathbf{y}_{1'}, \zeta) &= \Phi(\mathbf{y}_{0'} + \zeta\mathbf{y}_{1'}, t - \langle \mathbf{y}_{0'} + \zeta\mathbf{y}_{1'}, \mathbf{y}_{1'} \rangle, \zeta) \\ &= (\zeta^{-1}\mathbf{y}_{0'} + \mathbf{y}_{1'}, t + \langle \zeta^{-1}\mathbf{y}_{0'} + \mathbf{y}_{1'}, \mathbf{y}_{0'} \rangle, \zeta^{-1}) \\ &= \tilde{\eta} \circ \kappa(\mathbf{y}_{0'}, \mathbf{y}_{1'}, \zeta), \end{aligned}$$

by (31). Thus η and $\tilde{\eta}$ are glued to give the mapping $\mathbb{C}^5 \times \mathbb{C}P^1 \rightarrow \mathcal{P}$ in the double fibration (7).

2.2. ASD Equation. Let 1-forms $\{\theta^{AA'}, \theta\}$ be dual to left invariant vector fields $\{V_{AA'}, T\}$ in (4) on \mathcal{H} , i.e. $\theta^{AA'}(V_{BB'}) = \delta_{AB}\delta_{A'B'}$, $\theta^{AA'}(T) = 0$, $\theta(V_{BB'}) = 0$, $\theta(T) = 1$, where $A, B = 0, 1$ and $A', B' = 0', 1'$. Then $du = \sum_{A, A'} V_{AA'}u \cdot \theta^{AA'} + Tu \cdot \theta$ for a function u on \mathcal{H} . By the expression of $V_{AA'}$ in (4), we get that $\theta^{AA'} = dy_{AA'}$, $\theta = dt + y_{11'}dy_{00'} + y_{01'}dy_{10'} - y_{10'}dy_{01'} - y_{00'}dy_{11'}$. Exterior differentiation gives us $d\theta^{AA'} = 0$ ($A = 0, 1, A' = 0', 1'$) and $d\theta = -2\theta^{00'} \wedge \theta^{11'} - 2\theta^{10'} \wedge \theta^{01'}$.

The curvature of the connection form Φ is $F = (d + \Phi)^2 = d\Phi + \Phi \wedge \Phi$ given by

$$(35) \quad \begin{aligned} F(X, Y) &= d\Phi(X, Y) + \Phi \wedge \Phi(X, Y) \\ &= X(\Phi(Y)) - Y(\Phi(X)) - \Phi([X, Y]) + \Phi(X)\Phi(Y) - \Phi(Y)\Phi(X) \\ &= X\Phi_Y - Y\Phi_X - \Phi_{[X, Y]} + [\Phi_X, \Phi_Y]. \end{aligned}$$

Here we use the notation $\Phi_X := \Phi(X)$ for a vector field X on \mathcal{H} . Define the \mathfrak{g} -valued differential operators associated to the connection form Φ

$$(36) \quad \nabla_A = \nabla_{A1'} - \zeta \nabla_{A0'} := (V_{A1'} + \Phi_{A1'}) - \zeta(V_{A0'} + \Phi_{A0'}),$$

$A = 0, 1$, for fixed $\zeta \in \mathbb{C}$. A connection on \mathcal{H} is *ASD* if its curvature vanishes along each α -surface, i.e.

$$(37) \quad F(V_0^\zeta, V_1^\zeta) = 0.$$

The ASD condition (37) is equivalent to

$$\zeta^2 F(V_{00'}, V_{10'}) - \zeta(F(V_{00'}, V_{11'}) + F(V_{01'}, V_{10'})) + F(V_{01'}, V_{11'}) = 0.$$

Comparing the coefficients of ζ^2 , ζ^1 and ζ^0 , we get

$$(38) \quad F(V_{00'}, V_{10'}) = 0, \quad F(V_{00'}, V_{11'}) + F(V_{01'}, V_{10'}) = 0, \quad F(V_{01'}, V_{11'}) = 0,$$

which is equivalent to (8) by (35). Here $\Phi_{[V_{00'}, V_{11'}]} + \Phi_{[V_{01'}, V_{10'}]} = 0$ by the brackets in (5).

3. PENROSE-WARD CORRESPONDENCE AND ATIYAH-WARD ANSÄTZ ON 5D HEISENBERG GROUP

3.1. Proof of Theorem 1.1. The proof is similar to the classical case (cf. [16]). The objects in (1) and (2) are regarded as being specified modulo the usual equivalence relations. Given a $GL(n, \mathbb{C})$ vector bundle V with ASD connection ∇ over $U \subseteq \mathbb{C}^5$, to construct a vector bundle E over \hat{U} , we assign a copy of the vector space \mathbb{C}^n to each point Z of \hat{U} with

$$(39) \quad E_Z = \{\psi : \nabla\psi|_{\tilde{Z} \cap U} = 0\}$$

over Z , where $\tilde{Z} = \tau \circ \eta^{-1}(Z)$. As ∇ is an ASD connection, for fixed $\zeta \in \mathbb{C}$, the integrability condition $F(V_0, V_1) = 0$ implies the existence of solutions to the equations $(V_A + \Phi_A)h = 0$, $A = 0, 1$ on connected and simply connected domain $\tilde{Z} \cap U$. So we have $E_Z \neq \emptyset$. Since the whole procedure is holomorphic, we have constructed a holomorphic vector bundle E over \hat{U} . By construction (39), the bundle E is trivial when restricted to the projective line \hat{x} for a point x in U . This is because a vector $\psi \in V_x$ at x determines a parallel field ψ on each α -plane through x , and hence determines a point in E_Z for every point Z on the line \hat{x} in \hat{U} . Namely, each $\psi \in V_x \cong \mathbb{C}^n$ determines a holomorphic section of $E|_{\hat{x}}$. Therefore n linearly independent ψ 's in V_x give us n linearly independent sections of $E|_{\hat{x}}$. So $E|_{\hat{x}}$ is trivial. This completes the proof of the first part of the theorem.

Conversely, let E' be a holomorphic rank- n vector bundle over \hat{U} , such that $E'|_{\hat{x}}$ is trivial for all $x \in U$. We have to construct a connection form Φ on U . Since \hat{U} is covered by two charts $\hat{U} \cap W$ and $\hat{U} \cap \tilde{W}$ in (32), the vector bundle E' consists of two parts, which are $E'_{\hat{U} \cap W} \cong (\hat{U} \cap W) \times \mathbb{C}^n$

and $E'_{\hat{U} \cap \widetilde{W}} \cong (\hat{U} \cap \widetilde{W}) \times \mathbb{C}^n$ glued by a holomorphic $n \times n$ transition matrix F on the intersection $\hat{U} \cap W \cap \widetilde{W}$. The transition relation is

$$\widetilde{\xi} = F\xi,$$

where $\widetilde{\xi}$ and ξ are column n -vectors whose components serve as coordinates on the fibers of E' above $\hat{U} \cap W$ and $\hat{U} \cap \widetilde{W}$, respectively. Consider the pull-back η^*E' , which is a bundle over $\mathcal{F}_U = U \times \mathbb{C}P^1$, where η is given by (24) and (28). For a point $Z \in \hat{U}$, the restriction of η^*E' to $\eta^{-1}(Z) \in \mathcal{F}_U$ is a product bundle. We define a bundle $E \rightarrow U$ by $E_x = \Gamma(\hat{x}, E')$, where Γ denotes the space of holomorphic sections. And we have $\eta^*E' = \tau^*E$. Recall that the tangential space of leaves of the projection $\eta : \mathcal{F}_U \rightarrow \hat{U}$ are spanned by the vector fields V_0 and V_1 on \mathcal{F} . We define a partial connection ∇ that allows us to differentiate the sections of η^*E' along the fibers, where we have $\nabla_{V_A}s = V_A s$ in the trivialization $\eta^*E'|_{\eta^{-1}(Z)} = \eta^{-1}(Z) \times E'_Z$. The sections for which $\nabla_{V_A}s = V_A s$ vanish are the pull-backs to \mathcal{F} of local sections of E' . We now pick a local trivialization of E over open subset U . This determines a local trivialization of η^*E' , in which

$$\nabla_A = V_A + \Phi_A, \quad A = 0, 1,$$

for some matrix-valued functions Φ in ζ and (\mathbf{y}, t) . Note that $\eta^*E'|_{U \times O_0} \cong U \times O_0 \times \mathbb{C}^n$ and $\eta^*E'|_{U \times O_1} \cong U \times O_1 \times \mathbb{C}^n$, where $O_0 \cong \mathbb{C}$ and $O_1 \cong \mathbb{C}$ are a covering of $\mathbb{C}P^1$. Their transition function G over the intersection $U \times (\mathbb{C} \setminus \{0\})$ is given by

$$G(\mathbf{y}, t, \zeta) = F(\mathbf{y}_{0'} + \zeta \mathbf{y}_{1'}, t - \langle \mathbf{y}_{0'} + \zeta \mathbf{y}_{1'}, \mathbf{y}_{1'} \rangle, \zeta).$$

Now let us find the holomorphic sections of $E'|_{\hat{x}}$. Find non singular $n \times n$ matrices f and \widetilde{f} , with f holomorphic over $W \cap \hat{x}$ and \widetilde{f} holomorphic over $\widetilde{W} \cap \hat{x}$, such that $F = \widetilde{f}^{-1}f$ is valid on $W \cap \widetilde{W} \cap \hat{x}$. Since E'_x is trivial, by the *Birkhoff factorization*, such matrices f and \widetilde{f} must exist. Each section of E'_x is then given by $\xi = f^{-1}\psi$, $\widetilde{\xi} = \widetilde{f}^{-1}\psi$, where ψ is a constant n -vector, i.e. $\psi \in \mathbb{C}^n$. Now we identify \widetilde{f} and f with their pulling back to $U \times \mathbb{C}P^1$. So we have that

$$\begin{aligned} 0 &= \nabla_A \psi = (-V_{A1'} + \zeta V_{A0'})\psi + (-\Phi_{A1'} + \zeta \Phi_{A0'})\psi \\ (40) \quad &= (-V_{A1'}f + \zeta V_{A0'}f)\xi + (-\Phi_{A1'} + \zeta \Phi_{A0'})\psi \\ &= -(V_{A1'}f \cdot f^{-1} + \Phi_{A1'})\psi + \zeta(V_{A0'}f \cdot f^{-1} + \Phi_{A0'})\psi. \end{aligned}$$

So we have $\Phi_{AA'} = -V_{AA'}f \cdot f^{-1}$. Moreover, since V_A is tangential to the fiber of η , we have

$$\begin{aligned} (41) \quad 0 &= V_A G = V_A(\widetilde{f}^{-1}f) = -\widetilde{f}^{-1}V_A \widetilde{f} \cdot \widetilde{f}^{-1}f + \widetilde{f}^{-1}V_A f \\ &= \widetilde{f}^{-1}(V_A f \cdot f^{-1} - V_A \widetilde{f} \cdot \widetilde{f}^{-1})f. \end{aligned}$$

We have $V_A f \cdot f^{-1} = V_A \widetilde{f} \cdot \widetilde{f}^{-1}$. By Liouville's theorem, both sides must be of the form $-\Phi_{A1'} + \zeta \Phi_{A0'}$ for $A = 0, 1$, where $\Phi_{A0'}$ and $\Phi_{A1'}$ are matrix-valued functions over \mathbb{C}^5 . Therefore

$$V_{A1'}f \cdot f^{-1} - \zeta V_{A0'}f \cdot f^{-1} = -\Phi_{A1'} + \zeta \Phi_{A0'},$$

i.e.

$$(42) \quad (V_{A1'} + \Phi_{A1'})f - \zeta(V_{A0'} + \Phi_{A0'})f = 0.$$

Thus f is the solution to

$$\nabla_{A1'}f - \zeta \nabla_{A0'}f = 0,$$

$A = 0, 1$. Consequently, we have $F(V_0, V_1) = 0$, i.e. ∇ is ASD connection. Similarly, \tilde{f} is solution to

$$(43) \quad (V_{A0'} + \Phi_{A0'})\tilde{f} - \tilde{\zeta}(V_{A1'} + \Phi_{A1'})\tilde{f} = 0,$$

i.e. $\nabla_{A0'}\tilde{f} - \tilde{\zeta}\nabla_{A1'}\tilde{f} = 0$, where $A = 0, 1$. From (42) and (43), we have

$$(44) \quad \Phi_{A1'} = -V_{A1'}f \cdot f^{-1}|_{\zeta=0}, \quad \Phi_{A0'} = -V_{A0'}\tilde{f} \cdot \tilde{f}^{-1}|_{\tilde{\zeta}=0},$$

where $A = 0, 1$. So the connection Φ has the form

$$(45) \quad \Phi = -d_1f \cdot f^{-1}|_{\zeta=0} - d_0\tilde{f} \cdot \tilde{f}^{-1}|_{\tilde{\zeta}=0} + \Phi_T\theta,$$

which is ASD. \square

3.2. Atiyah-Ward ansatz For $GL(2, \mathbb{C})$ ASD connection on 5D Heisenberg group.

By the construction in the proof of Theorem 1.1, we get a $GL(2, \mathbb{C})$ ASD connection if we find a transition matrix for a rank-2 bundle over \hat{U} , which is trivial along the projective line $\hat{x} = \eta \circ \tau^{-1}(x)$ for each $x \in U \subseteq \mathcal{H}$. Namely, to find a 2×2 matrix

$$G(\mathbf{y}, t, \zeta) = \begin{pmatrix} \zeta & \gamma(\mathbf{y}, t, \zeta) \\ 0 & \zeta^{-1} \end{pmatrix}$$

defined over $U \times (\mathbb{C} \setminus \{0\})$ (the intersection of two coordinate charts of $\mathbb{C}^5 \times \mathbb{C}P^1$), and is constant along the fibers of η and trivial over $\hat{\mathbf{x}}$ for each $\mathbf{x} = (\mathbf{y}, t) \in U \subseteq \mathbb{C}^5$. It defines a function \hat{G} on $\hat{U} \cap W \cap \hat{W}$

$$G(\mathbf{y}, t, \zeta) = \hat{G}(\mathbf{y}_{0'} + \zeta\mathbf{y}_{1'}, t - \langle \mathbf{y}_{0'} + \zeta\mathbf{y}_{1'}, \mathbf{y}_{1'} \rangle, \zeta).$$

This is called *Atiyah-Ward Ansatz*.

In terms of Laurent series of ζ , we write

$$(46) \quad \gamma = \sum_{i=-\infty}^{+\infty} \gamma_{-i}\zeta^i = \gamma_{-} + \gamma_0 + \gamma_{+},$$

where $\gamma_{-} := \sum_{i=1}^{i=\infty} \gamma_i\zeta^{-i}$, $\gamma_{+} := \sum_{i=1}^{i=\infty} \gamma_{-i}\zeta^i$. Since γ need to be constant along each fiber of η , we have $V_A\gamma \equiv 0$, $A = 0, 1$, i.e.

$$(47) \quad V_{A1'}\gamma_i = V_{A0'}\gamma_{i+1}, \quad A = 0, 1, \quad i = \dots, -1, 0, 1, \dots.$$

Lemma 3.1. *Suppose that $U \subseteq \mathbb{C}^5$ is elementary and the one form $\omega = g_0\theta^{00'} + g_1\theta^{10'}$ (or $\omega = g_0\theta^{01'} + g_1\theta^{11'}$) with $g_0, g_1 \in \mathcal{O}(U)$ is d_0 -closed (or d_1 -closed), i.e. $d_0\omega = 0$ (or $d_1\omega = 0$). Then there exists a function $f \in \mathcal{O}(U)$, such that $d_0f = \omega$ (or $d_1f = \omega$).*

Proof. By definition, $d_0\omega = 0$ is equivalent to

$$(48) \quad V_{10'}g_0 = V_{00'}g_1,$$

and $d_0f = \omega$ is equivalent to

$$(49) \quad V_{00'}f = g_0 \quad \text{and} \quad V_{10'}f = g_1.$$

If we take coordinate transformation $\Psi : \mathbb{C}^5 \rightarrow \mathbb{C}^5$ given by

$$(y_{00'}, y_{10'}, y_{01'}, y_{11'}, t) := \Psi(z_0, z_1, z_2, z_3, z_4) = (z_0, z_1, z_2, z_3, z_4 - z_0z_3 - z_1z_2),$$

we have

$$\Psi_* \frac{\partial}{\partial z_0} = V_{00'}, \quad \Psi_* \frac{\partial}{\partial z_1} = V_{10'},$$

by expression of $V_{AA'}$ in (4). Take $G_A = g_A \circ \Psi$, $A = 0, 1$, and $F = f \circ \Psi$. Then by pulling back, we need to solve

$$(50) \quad \frac{\partial F}{\partial z_0} = G_0, \quad \frac{\partial F}{\partial z_1} = G_1.$$

Under the condition $\frac{\partial G_0}{\partial z_1} = \frac{\partial G_1}{\partial z_0}$, it follows from Poincaré's lemma that (50) has a solution. Therefore, $f = F \circ \Psi^{-1}$ is the solution to (49). \square

Lemma 3.2. *Suppose γ_0 satisfy the equation (10), i.e. $\Delta_b \gamma_0 = 0$, then (47) is solvable.*

Proof. Inductively, for fixed $i = 0, -1, \dots$, assuming that there exists γ_i and γ_{i-1} satisfying

$$(51) \quad \begin{cases} V_{01'} \gamma_{i-1} = V_{00'} \gamma_i, \\ V_{11'} \gamma_{i-1} = V_{10'} \gamma_i, \end{cases}$$

we need to find γ_{i-2} such that

$$(52) \quad \begin{cases} V_{01'} \gamma_{i-2} = V_{00'} \gamma_{i-1}, \\ V_{11'} \gamma_{i-2} = V_{10'} \gamma_{i-1}. \end{cases}$$

Denote

$$\Lambda_{i-1} := V_{00'} \gamma_{i-1} \theta^{01'} + V_{10'} \gamma_{i-1} \theta^{11'}.$$

Then Λ_{i-1} is d_1 -closed, since

$$\begin{aligned} d_1 \Lambda_{i-1} &= (V_{01'} V_{10'} - V_{11'} V_{00'}) \gamma_{i-1} \theta^{01'} \wedge \theta^{11'} = (V_{10'} V_{01'} - V_{00'} V_{11'}) \gamma_{i-1} \theta^{01'} \wedge \theta^{11'} \\ &= (V_{10'} V_{00'} - V_{00'} V_{10'}) \gamma_i \theta^{01'} \wedge \theta^{11'} = 0, \end{aligned}$$

by using $[V_{00'}, V_{11'}] = [V_{10'}, V_{01'}] = T$ in (5) in the second identity, (51) and $[V_{00'}, V_{10'}] = 0$ in the last identity. It follows from Lemma 3.1 that there exists γ_{i-2} such that $d_1 \gamma_{i-2} = \Lambda_{i-1}$, i.e. equation (52) is satisfied. So solving the equations (47) for $i = \dots, -1, 0$, is reduced to the solve the equation $d_1 \Lambda_0 = 0$ by induction. But

$$\begin{aligned} d_1 \Lambda_0 &= (V_{01'} V_{10'} - V_{11'} V_{00'}) \gamma_0 \theta^{01'} \wedge \theta^{11'} \\ &= (V_{10'} V_{01'} - V_{00'} V_{11'}) \gamma_0 \theta^{01'} \wedge \theta^{11'} \\ &= -\Delta_b \gamma_0 \theta^{01'} \wedge \theta^{11'} = 0. \end{aligned}$$

Hence, (47) is solvable for $i = \dots, -1, 0$.

On the other hand, for $i = 0, 1, \dots$, if there exists γ_i, γ_{i+1} such that

$$(53) \quad \begin{cases} V_{01'} \gamma_i = V_{00'} \gamma_{i+1}, \\ V_{11'} \gamma_i = V_{10'} \gamma_{i+1}, \end{cases}$$

we need to show

$$(54) \quad \begin{cases} V_{00'} \gamma_{i+2} = V_{01'} \gamma_{i+1}, \\ V_{10'} \gamma_{i+2} = V_{11'} \gamma_{i+1} \end{cases}$$

has a solution γ_{i+2} . Denote

$$\tilde{\Lambda}_{i+1} := V_{01'} \gamma_{i+1} \theta^{00'} + V_{11'} \gamma_{i+1} \theta^{10'}.$$

As above Λ_{i+1} is d_0 -closed, since

$$\begin{aligned} d_0 \tilde{\Lambda}_{i+1} &= (V_{00'} V_{11'} - V_{10'} V_{01'}) \gamma_{i+1} \theta^{00'} \wedge \theta^{10'} = (V_{11'} V_{00'} - V_{01'} V_{10'}) \gamma_{i+1} \theta^{00'} \wedge \theta^{10'} \\ &= (V_{11'} V_{01'} - V_{01'} V_{11'}) \gamma_i \theta^{00'} \wedge \theta^{10'} = 0, \end{aligned}$$

by using $[V_{00'}, V_{11'}] = [V_{10'}, V_{01'}] = T$ in the second identity, (53) and $[V_{11'}, V_{01'}] = 0$ in the third identity. By Lemma 3.1 again, there exists γ_{i+2} such that $d_0 \gamma_{i+2} = \tilde{\Lambda}_{i+1}$, i.e. the equation (54) is solvable. The equation (47), for $i = 0, 1, \dots$, is also reduced to the equation

$$0 = d_0 \tilde{\Lambda}_0 = (V_{00'} V_{11'} - V_{10'} V_{01'}) \gamma_0 \theta^{00'} \wedge \theta^{10'} = \Delta_b \gamma_0 \theta^{00'} \wedge \theta^{10'}.$$

It holds since $\Delta_b \gamma_0 = 0$. The lemma is proved. \square

Proof of Theorem 1.2. (1) As in the classical case (cf. e.g. [10, Example 10.1.2]), we take the Birkhoff decomposition

$$(55) \quad G(\mathbf{y}, t, \zeta) = \begin{pmatrix} \zeta & \gamma(\mathbf{y}, t, \zeta) \\ 0 & \zeta^{-1} \end{pmatrix} = \tilde{f}^{-1} f,$$

with

$$(56) \quad \begin{aligned} f &= \frac{1}{\sqrt{\varphi}} \begin{pmatrix} \zeta & \varphi + \gamma_+ \\ -1 & -\zeta^{-1} \gamma_+ \end{pmatrix} \in \mathcal{O}(U \times \mathbb{C}), \\ \tilde{f} &= \frac{1}{\sqrt{\varphi}} \begin{pmatrix} 1 & -\zeta \gamma_- \\ -\zeta^{-1} & \varphi + \gamma_- \end{pmatrix} \in \mathcal{O}(U \times \mathbb{C}), \end{aligned}$$

where $U \times \mathbb{C}$ and $U \times \mathbb{C}$ are local coordinate charts of $U \times \mathbb{C}P^1$. Their inverses are

$$(57) \quad \begin{aligned} f^{-1} &= \sqrt{\varphi} \begin{pmatrix} -\zeta^{-1} \varphi^{-1} \gamma_+ & -1 - \gamma_+ \varphi^{-1} \\ \varphi^{-1} & \zeta \varphi^{-1} \end{pmatrix}, \\ \tilde{f}^{-1} &= \sqrt{\varphi} \begin{pmatrix} 1 + \varphi^{-1} \gamma_- & \zeta \gamma_- \varphi^{-1} \\ \varphi^{-1} \zeta^{-1} & \varphi^{-1} \end{pmatrix}, \end{aligned}$$

respectively. It is direct to check (55) by \tilde{f}^{-1} and f in (56)-(57), if γ_{\pm} satisfy (46) and $\gamma_0 = \varphi$. Then

$$(58) \quad h := f|_{\zeta=0} = \frac{1}{\sqrt{\varphi}} \begin{pmatrix} 0 & \varphi \\ -1 & -\gamma_{-1} \end{pmatrix}, \quad \tilde{h} := \tilde{f}|_{\zeta=0} = \frac{1}{\sqrt{\varphi}} \begin{pmatrix} 1 & -\gamma_1 \\ 0 & \varphi \end{pmatrix},$$

By (9) and (47), we have $d_1 \gamma_{-1} = V_{00'} \varphi \cdot \theta^{01'} + V_{10'} \varphi \cdot \theta^{11'}$ and $d_0 \gamma_1 = V_{01'} \varphi \cdot \theta^{00'} + V_{11'} \varphi \cdot \theta^{10'}$, and

$$(59) \quad h^{-1} = \sqrt{\varphi} \begin{pmatrix} -\varphi^{-1} \gamma_{-1} & -1 \\ \varphi^{-1} & 0 \end{pmatrix}, \quad \tilde{h}^{-1} = \sqrt{\varphi} \begin{pmatrix} 1 & \gamma_1 \varphi^{-1} \\ 0 & \varphi^{-1} \end{pmatrix}.$$

It is direct to check that

$$(60) \quad \begin{aligned} \Phi &= -d_1 h \cdot h^{-1} - d_0 \tilde{h} \cdot \tilde{h}^{-1} + \Phi_T \theta \\ &= \left[\begin{array}{cc} \frac{1}{2}(-d_1 \ln \varphi + d_0 \ln \varphi) & V_{01'}(\ln \varphi) \theta^{00'} + V_{11'}(\ln \varphi) \theta^{10'} \\ V_{00'}(\ln \varphi) \theta^{01'} + V_{10'}(\ln \varphi) \theta^{11'} & \frac{1}{2}(d_1 \ln \varphi - d_0 \ln \varphi) \end{array} \right] + \Phi_T \theta, \end{aligned}$$

which is ASD by (45) and Theorem 1.1.

(2) Noting that $V_{AA'}(y_{BB'}) = \delta_{AB} \delta_{A'B'}$ and $\|\mathbf{y}\|^2 = y_{00'} y_{11'} - y_{10'} y_{01'}$, we have

$$V_{00'} \|\mathbf{y}\|^2 = y_{11'}, \quad V_{01'} \|\mathbf{y}\|^2 = -y_{10'}, \quad V_{10'} \|\mathbf{y}\|^2 = -y_{01'}, \quad V_{11'} \|\mathbf{y}\|^2 = y_{00'}.$$

It's direct to check that

$$\begin{aligned} V_{00'}(\|\mathbf{y}\|^4 - t^2) &= 2y_{11'}(\|\mathbf{y}\|^2 + t), \\ V_{01'}(\|\mathbf{y}\|^4 - t^2) &= -2y_{10'}(\|\mathbf{y}\|^2 + t), \\ V_{10'}(\|\mathbf{y}\|^4 - t^2) &= 2y_{01'}(-\|\mathbf{y}\|^2 + t), \\ V_{11'}(\|\mathbf{y}\|^4 - t^2) &= 2y_{00'}(\|\mathbf{y}\|^2 - t), \end{aligned}$$

and

$$\begin{aligned} V_{10'}V_{01'}(\|\mathbf{y}\|^4 - t^2) &= -2\|\mathbf{y}\|^2 - 2t + 4y_{10'}y_{01'}, \\ V_{00'}V_{11'}(\|\mathbf{y}\|^4 - t^2) &= 2\|\mathbf{y}\|^2 - 2t + 4y_{00'}y_{11'}. \end{aligned}$$

Consequently, for $\varphi = \frac{1}{\|\mathbf{y}\|^4 - t^2}$, we have

$$\begin{aligned} V_{10'}V_{01'}(\varphi) &= -\frac{V_{10'}V_{01'}(\|\mathbf{y}\|^4 - t^2)}{(\|\mathbf{y}\|^4 - t^2)^2} + 2\frac{V_{01'}(\|\mathbf{y}\|^4 - t^2)V_{10'}(\|\mathbf{y}\|^4 - t^2)}{(\|\mathbf{y}\|^4 - t^2)^3} \\ &= \frac{2y_{00'}y_{11'} + 2y_{10'}y_{01'} + 2t}{(\|\mathbf{y}\|^4 - t^2)^2}, \\ V_{00'}V_{11'}(\varphi) &= -\frac{V_{00'}V_{11'}(\|\mathbf{y}\|^4 - t^2)}{(\|\mathbf{y}\|^4 - t^2)^2} + 2\frac{V_{11'}(\|\mathbf{y}\|^4 - t^2)V_{00'}(\|\mathbf{y}\|^4 - t^2)}{(\|\mathbf{y}\|^4 - t^2)^3} \\ &= \frac{2y_{00'}y_{11'} + 2y_{10'}y_{01'} + 2t}{(\|\mathbf{y}\|^4 - t^2)^2}. \end{aligned}$$

Therefore $V_{10'}V_{01'}(\varphi) = V_{00'}V_{11'}(\varphi)$, i.e. φ is the solution to (10). \square

4. THE REAL CASE

4.1. SD and ASD horizontal form on real Heisenberg group $\mathcal{H}^{\mathbb{R}}$. The real Heisenberg group is a flat model of contact manifolds. Let us compare the ASD connection restricted to the real 5D Heisenberg group with contact instantons on 5D contact manifolds introduced by physicists (cf.[8] [6] [17]). Take a metric on $\mathcal{H}^{\mathbb{R}}$ as $g = dy_1^2 + dy_2^2 + dy_3^2 + dy_4^2 + dt^2$. The relevant volume element on $\mathcal{H}^{\mathbb{R}}$ is $dV := dy_1 \wedge dy_2 \wedge dy_3 \wedge dy_4 \wedge dt$. For brevity, we denote $dy_5 := dt$. The *Hodge star* $*$ with respect to the metric g on $\mathcal{H}^{\mathbb{R}}$ is given by $* : \Omega^k(\mathcal{H}^{\mathbb{R}}) \rightarrow \Omega^{5-k}(\mathcal{H}^{\mathbb{R}})$, $w \mapsto *w$, such that

$$(61) \quad w \wedge *w = dV.$$

Namely, we have $*(dy_{i_1} \wedge \cdots \wedge dy_{i_k}) = \varepsilon_{i_1 \cdots i_k j_1 \cdots j_{5-k}} dy_{j_1} \wedge \cdots \wedge dy_{j_{5-k}}$, where $\varepsilon_{i_1 \cdots i_k j_1 \cdots j_{5-k}}$ is the permutation index from $\{i_1, \cdots, i_k, j_1, \cdots, j_{5-k}\}$ to $\{1, \cdots, 5\}$. Denote by $\Omega_H^2(\mathcal{H}^{\mathbb{R}})$ the space of horizontal two forms on $\mathcal{H}^{\mathbb{R}}$, i.e. $\iota_T \omega = 0$ for $\omega \in \Omega_H^2(\mathcal{H}^{\mathbb{R}})$, and $\Omega_V^2(\mathcal{H}^{\mathbb{R}})$ the space of vertical two forms on $\mathcal{H}^{\mathbb{R}}$. We have the decomposition

$$(62) \quad \Omega^2(\mathcal{H}^{\mathbb{R}}) = \Omega_H^2(\mathcal{H}^{\mathbb{R}}) \oplus \Omega_V^2(\mathcal{H}^{\mathbb{R}}), \quad \omega = \omega_H + \omega_V,$$

where $\omega_H = \iota_T(dt \wedge \omega)$ and $\omega_V = dt \wedge (\iota_T \omega)$. Denote by $\Omega_H^{2+}(\mathcal{H}^{\mathbb{R}})$ ($\Omega_H^{2-}(\mathcal{H}^{\mathbb{R}})$) the space of all horizontal (anti-)self-dual spaces of two forms on \mathcal{H} , of which elements satisfy

$$(63) \quad \iota_T * \omega = \omega \quad (\iota_T * \omega = -\omega).$$

Then we have the decomposition

$$(64) \quad \Omega_H^2(\mathcal{H}^{\mathbb{R}}) = \Omega_H^{2+}(\mathcal{H}^{\mathbb{R}}) \oplus \Omega_H^{2-}(\mathcal{H}^{\mathbb{R}}) \quad \omega_H = \omega_H^+ + \omega_H^-,$$

where $\omega_H^+ = \frac{1}{2}(1 + \iota_T^*)\omega_H \in \Omega_H^{2+}(\mathcal{H}^{\mathbb{R}})$ and $\omega_H^- = \frac{1}{2}(1 - \iota_T^*)\omega_H \in \Omega_H^{2-}(\mathcal{H}^{\mathbb{R}})$. In fact, noting that $\iota_T * \iota_T^* = id$, we have $\iota_T * \omega_H^+ = \frac{1}{2}\iota_T * (1 + \iota_T^*)\omega_H = \frac{1}{2}(\iota_T * + 1)\omega_H = \omega_H^+$, i.e. $\omega_H^+ \in \Omega_H^{2+}(\mathcal{H}^{\mathbb{R}})$. And $\iota_T * \omega_H^- = \frac{1}{2}\iota_T * (1 - \iota_T^*)\omega_H = \frac{1}{2}(\iota_T * - 1)\omega_H = -\omega_H^-$, i.e. $\omega_H^- \in \Omega_H^{2-}(\mathcal{H}^{\mathbb{R}})$.

4.2. Contact instantons on $\mathcal{H}^{\mathbb{R}}$. The double fibration over real Heisenberg group $\mathcal{H}^{\mathbb{R}}$ becomes

$$(65) \quad \begin{array}{ccc} & \mathcal{F}^{\mathbb{R}} & \\ \eta \swarrow & & \searrow \tau \\ \mathcal{P}^{\mathbb{R}} & & \mathcal{H}^{\mathbb{R}} \end{array}$$

where $\mathcal{F}^{\mathbb{R}} = \mathcal{H}^{\mathbb{R}} \times \mathbb{C}P^1$ and $\mathcal{P}^{\mathbb{R}} = \eta(\mathcal{H}^{\mathbb{R}} \times \mathbb{C}P^1)$.

Corollary 4.1. *The fiber of the mapping $\eta : \mathbb{R}^5 \times \mathbb{C} \rightarrow \mathcal{P}^{\mathbb{R}}$ given by*

$$(66) \quad \begin{aligned} \omega_0 &:= \eta_0(\mathbf{y}_{0'}, \mathbf{y}_{1'}, \mathbf{t}, \zeta) = y_1 + \mathbf{i}y_2 + \zeta(-y_3 + \mathbf{i}y_4), \\ \omega_1 &:= \eta_1(\mathbf{y}_{0'}, \mathbf{y}_{1'}, \mathbf{t}, \zeta) = y_3 + \mathbf{i}y_4 + \zeta(y_1 - \mathbf{i}y_2), \\ \omega_2 &:= \eta_2(\mathbf{y}_{0'}, \mathbf{y}_{1'}, \mathbf{t}, \zeta) = -\mathbf{i}s - 2\zeta(-y_3 + \mathbf{i}y_4)(y_1 - \mathbf{i}y_2) - y_1^2 - y_2^2 + y_3^2 + y_4^2, \\ \omega_3 &:= \eta_3(\mathbf{y}_{0'}, \mathbf{y}_{1'}, \mathbf{t}, \zeta) = \zeta, \end{aligned}$$

is an abelian subgroup of dimension 4, whose tangential space is spanned by $\{V_0, V_1\}$. We denote $\hat{y} = (\omega_0, \omega_1, \omega_2, \omega_3)$.

Proposition 4.1. *For x and $y \in \mathbb{R}^5$, if $\hat{x} \cap \hat{y} \neq \emptyset$, then we have $x = y$. So $\mathcal{P}^{\mathbb{R}}$ is the trivial $\mathbb{C}P^1$ bundle over \mathbb{R}^5 .*

Proof. If we write

$$(67) \quad \begin{aligned} x &= (x_{00'}, x_{10'}, x_{01'}, x_{11'}, t_1) = (x_1 + \mathbf{i}x_2, x_3 + \mathbf{i}x_4, -x_3 + \mathbf{i}x_4, x_1 - \mathbf{i}x_2, -\mathbf{i}s_1), \\ y &= (y_{00'}, y_{10'}, y_{01'}, y_{11'}, t_2) = (y_1 + \mathbf{i}y_2, y_3 + \mathbf{i}y_4, -y_3 + \mathbf{i}y_4, y_1 - \mathbf{i}y_2, -\mathbf{i}s_2), \end{aligned}$$

by the embedding (14) of $\mathcal{H}^{\mathbb{R}}$ into \mathcal{H} , then we have

$$(68) \quad \begin{aligned} \hat{x} &= \eta \circ \tau^{-1}(x) = (x_{00'} + \zeta x_{01'}, x_{10'} + \zeta x_{11'}, -\mathbf{i}s_1 - \langle \mathbf{x}_{0'} + \zeta \mathbf{x}_{1'}, \mathbf{x}_{1'} \rangle, \zeta), \\ \hat{y} &= \eta \circ \tau^{-1}(y) = (y_{00'} + \zeta y_{01'}, y_{10'} + \zeta y_{11'}, -\mathbf{i}s_2 - \langle \mathbf{y}_{0'} + \zeta \mathbf{y}_{1'}, \mathbf{y}_{1'} \rangle, \zeta), \end{aligned}$$

by (24). If $\hat{x} \cap \hat{y} \neq \emptyset$, then there exists ζ such that equations

$$(69) \quad \begin{cases} x_{00'} + \zeta x_{01'} = y_{00'} + \zeta y_{01'}, \\ x_{10'} + \zeta x_{11'} = y_{10'} + \zeta y_{11'}, \\ -\mathbf{i}s_1 - \langle \mathbf{x}_{0'} + \zeta \mathbf{x}_{1'}, \mathbf{x}_{1'} \rangle = -\mathbf{i}s_2 - \langle \mathbf{y}_{0'} + \zeta \mathbf{y}_{1'}, \mathbf{y}_{1'} \rangle \end{cases}$$

must have a solution. So equations

$$\begin{pmatrix} x_{00'} - y_{00'} & x_{01'} - y_{01'} \\ x_{10'} - y_{10'} & x_{11'} - y_{11'} \end{pmatrix} \begin{pmatrix} a_1 \\ a_2 \end{pmatrix} = 0$$

has a nontrivial solution. Consequently, its coefficient matrix is singular, i.e.

$$\begin{aligned} 0 &= \begin{vmatrix} x_{00'} - y_{00'} & x_{01'} - y_{01'} \\ x_{10'} - y_{10'} & x_{11'} - y_{11'} \end{vmatrix} = \begin{vmatrix} x_1 - y_1 + \mathbf{i}x_2 - \mathbf{i}y_2 & -x_3 + y_3 + \mathbf{i}x_4 - \mathbf{i}y_4 \\ x_3 - y_3 + \mathbf{i}x_4 - \mathbf{i}y_4 & x_1 - y_1 - \mathbf{i}x_2 + \mathbf{i}y_2 \end{vmatrix} \\ &= (x_1 - y_1)^2 + (x_2 - y_2)^2 + (x_3 - y_3)^2 + (x_4 - y_4)^2, \end{aligned}$$

i.e., $x_k = y_k$ for $k = 1, \dots, 4$. So $s_1 = s_2$ in (67) by the third equation of (69). This implies η in (65) is one to one, and so a diffeomorphism. Therefore, $\mathcal{P}^{\mathbb{R}}$ is a topologically trivial $\mathbb{C}P^1$ bundle over \mathbb{R}^5 . \square

Recall the real embedding (14) of $\mathcal{H}^{\mathbb{R}}$ into \mathcal{H} . Then we have the left invariant complex vector field over $\mathcal{H}^{\mathbb{R}}$

$$(70) \quad \begin{aligned} V_{00'} &= \frac{1}{2}(\partial_{y_1} - \mathbf{i}\partial_{y_2}) - \mathbf{i}(y_1 - \mathbf{i}y_2)\partial_s, & V_{01'} &= \frac{1}{2}(-\partial_{y_3} - \mathbf{i}\partial_{y_4}) + \mathbf{i}(y_3 + \mathbf{i}y_4)\partial_s, \\ V_{10'} &= \frac{1}{2}(\partial_{y_3} - \mathbf{i}\partial_{y_4}) + \mathbf{i}(y_3 - \mathbf{i}y_4)\partial_s, & V_{11'} &= \frac{1}{2}(\partial_{y_1} + \mathbf{i}\partial_{y_2}) + \mathbf{i}(y_1 + \mathbf{i}y_2)\partial_s, & T &= \mathbf{i}\partial_s, \end{aligned}$$

and the relevant dual forms becomes

$$(71) \quad \begin{aligned} \theta^{00'} &= dy_1 + \mathbf{i}dy_2, & \theta^{01'} &= -dy_3 + \mathbf{i}dy_4, & \theta^{10'} &= dy_3 + \mathbf{i}dy_4, & \theta^{11'} &= dy_1 - \mathbf{i}dy_2, \\ \theta &= -\mathbf{i}ds + 2\mathbf{i}(y_1dy_2 - y_2dy_1 + y_4dy_3 - y_3dy_4). \end{aligned}$$

Proposition 4.2. *The space of horizontal self-dual 2-forms on $\mathcal{H}^{\mathbb{R}}$ is spanned by $\{S^{0'0'}, S^{0'1'}, S^{1'1'}\}$ with*

$$(72) \quad S^{0'0'} = \theta^{00'} \wedge \theta^{10'}, \quad S^{0'1'} = \theta^{00'} \wedge \theta^{11'} - \theta^{10'} \wedge \theta^{01'}, \quad S^{1'1'} = \theta^{01'} \wedge \theta^{11'}.$$

The space of horizontal anti-self-dual 2-forms is spanned by $\{S^{00}, S^{01}, S^{11}\}$ with

$$(73) \quad S^{00} = \theta^{00'} \wedge \theta^{01'}, \quad S^{01} = \theta^{00'} \wedge \theta^{11'} + \theta^{10'} \wedge \theta^{01'}, \quad S^{11} = \theta^{10'} \wedge \theta^{11'}.$$

The curvature of the connection form $\Phi = \Phi_{AB'}\theta^{AB'} + \Phi_T\theta$ is

$$(74) \quad \begin{aligned} F &= d\Phi + \Phi \wedge \Phi \\ &= d(\Phi_{AB'}) \wedge \theta^{AB'} + d(\Phi_T) \wedge \theta + \Phi_T d\theta + \Phi_{AB'}\Phi_{CD'}\theta^{AB'} \wedge \theta^{CD'} + [\Phi_{AB'}, \Phi_T]\theta^{AB'} \wedge \theta \\ &= V_{CD'}(\Phi_{AB'})\theta^{CD'} \wedge \theta^{AB'} + T(\Phi_{AB'})\theta \wedge \theta^{AB'} + V_{AB'}(\Phi_T)\theta^{AB'} \wedge \theta + \Phi_T d\theta \\ &\quad + \Phi_{AB'}\Phi_{CD'}\theta^{AB'} \wedge \theta^{CD'} + [\Phi_{AB'}, \Phi_T]\theta^{AB'} \wedge \theta \\ &= [V_{AB'}(\Phi_{CD'}) + \Phi_{AB'}\Phi_{CD'}]\theta^{AB'} \wedge \theta^{CD'} + \Phi_T d\theta \\ &\quad + (V_{AB'}(\Phi_T) - T(\Phi_{AB'}) + [\Phi_{AB'}, \Phi_T])\theta^{AB'} \wedge \theta, \end{aligned}$$

by relabeling indices. Here we use the Einstein convention of summation over repeated indices. Then we have

$$(75) \quad F_V = \theta \wedge \iota_T F = (V_{AB'}(\Phi_T) - T(\Phi_{AB'}) + [\Phi_{AB'}, \Phi_T])\theta^{AB'} \wedge \theta,$$

and

$$\begin{aligned}
F_H &= \iota_T(\theta \wedge F) = [V_{AB'}(\Phi_{CD'}) + \Phi_{AB'}\Phi_{CD'}]\theta^{AB'} \wedge \theta^{CD'} + \Phi_T d\theta \\
&= (V_{00'}\Phi_{01'} - V_{01'}\Phi_{00'} + [\Phi_{00'}, \Phi_{01'}])\theta^{00'} \wedge \theta^{01'} \\
&\quad + (V_{00'}\Phi_{10'} - V_{10'}\Phi_{00'} + [\Phi_{00'}, \Phi_{10'}])\theta^{00'} \wedge \theta^{10'} \\
(76) \quad &\quad + (V_{00'}\Phi_{11'} - V_{11'}\Phi_{00'} + [\Phi_{00'}, \Phi_{11'}] - 2\Phi_T)\theta^{00'} \wedge \theta^{11'} \\
&\quad + (V_{01'}\Phi_{10'} - V_{10'}\Phi_{01'} + [\Phi_{01'}, \Phi_{10'}] + 2\Phi_T)\theta^{01'} \wedge \theta^{10'} \\
&\quad + (V_{01'}\Phi_{11'} - V_{11'}\Phi_{01'} + [\Phi_{01'}, \Phi_{11'}])\theta^{01'} \wedge \theta^{11'} \\
&\quad + (V_{10'}\Phi_{11'} - V_{11'}\Phi_{10'} + [\Phi_{10'}, \Phi_{11'}])\theta^{10'} \wedge \theta^{11'}.
\end{aligned}$$

So its horizontal self-dual parts is

$$\begin{aligned}
(77) \quad F_H^+ &= (V_{00'}\Phi_{10'} - V_{10'}\Phi_{00'} + [\Phi_{00'}, \Phi_{10'}])S^{0'0'} + (V_{01'}\Phi_{11'} - V_{11'}\Phi_{01'} + [\Phi_{01'}, \Phi_{11'}])S^{1'1'} \\
&\quad + \frac{1}{2}(V_{00'}\Phi_{11'} - V_{11'}\Phi_{00'} + [\Phi_{00'}, \Phi_{11'}] + V_{01'}\Phi_{10'} - V_{10'}\Phi_{01'} + [\Phi_{01'}, \Phi_{10'}])S^{0'1'}.
\end{aligned}$$

As a result, we get that

Proposition 4.3. (1) $F_H^+ = 0$ is equivalent to $F(V_0, V_1) = 0$, i.e.,

$$(78) \quad \begin{cases} V_{00'}(\Phi_{10'}) - V_{10'}(\Phi_{00'}) + [\Phi_{00'}, \Phi_{10'}] = 0, \\ V_{01'}(\Phi_{10'}) + V_{00'}(\Phi_{11'}) - V_{10'}(\Phi_{01'}) - V_{11'}(\Phi_{00'}) + [\Phi_{00'}, \Phi_{11'}] + [\Phi_{01'}, \Phi_{10'}] = 0, \\ V_{01'}(\Phi_{11'}) - V_{11'}(\Phi_{01'}) + [\Phi_{01'}, \Phi_{11'}] = 0. \end{cases}$$

(2) $F_V = 0$ is equivalent to $V_{AB'}(\Phi_T) - T(\Phi_{AB'}) + [\Phi_{AB'}, \Phi_T] = 0$.

When restricted to the real Heisenberg group $\mathcal{H}^{\mathbb{R}}$, the *Sub-Laplacian* becomes

$$\Delta_b := (X_1^2 + X_2^2 + X_3^2 + X_4^2),$$

where

$$(79) \quad X_1 = \frac{1}{2}\partial_{x_1} + x_2\partial_s, \quad X_2 = \frac{1}{2}\partial_{x_2} - x_1\partial_s, \quad X_3 = \frac{1}{2}\partial_{x_3} + x_4\partial_s, \quad X_4 = \frac{1}{2}\partial_{x_4} - x_3\partial_s.$$

Note that $[X_1, X_2] = -\partial_s$, $[X_3, X_4] = -\partial_s$. Then

$$\varphi = \frac{1}{|x|^4 + s^2}, \quad \text{where } |x| = (x_1^2 + x_2^2 + x_3^2 + x_4^2)^{\frac{1}{2}},$$

is the solution to the Sub-Laplacian equation

$$(80) \quad \Delta_b \varphi = 0.$$

Corollary 4.2. If φ is the solution to the equation (80), then the connection form Φ in (11) restricted to the real Heisenberg group $\mathcal{H}^{\mathbb{R}}$ satisfies the horizontal part of the ASD contact instanton equation $F_H^+ = 0$.

Remark 4.1. By Penrose-Ward correspondence, Wolf [17] has already characterized the solution to the horizontal part of the “self-dual” contact instanton equation on a contact manifold as follows. Let M be a 5D K -contact manifold with Cauchy-Riemann twistor space $\pi : Z \rightarrow M$ and integrable Cauchy-Riemann structure. There is a one-to-one correspondence between

(i) rank- r Cauchy-Riemann vector bundles $E_Z \rightarrow Z$ such that the restriction $E_Z|_{\pi^{-1}(p)}$ is holomorphically trivial for all $p \in M$ and

(ii) rank- r complex vector bundles $E_M \rightarrow M$ equipped with a connection ∇ and curvature $F = \nabla^2$ such that the projection on the contact distribution is $F_H \in \Omega_+^2(M, \text{End}E_M)$, that is, $F_- = 0$.

APPENDIX A. DOUBLE FIBRATION AND TWISTOR TRANSFORM

Let us use the method in [13] to write down local coordinates of the double fibration

$$(81) \quad \begin{array}{ccc} & \text{SO}(6, \mathbb{C})/R & \\ \eta \swarrow & & \searrow \tau \\ \text{SO}(6, \mathbb{C})/Q & & \text{SO}(6, \mathbb{C})/P \end{array}$$

and the mappings η and τ in term of their local coordinates, where P , Q and R are subgroups of $\text{SO}(6, \mathbb{C})$ in the following. Here the mapping η and τ are given by

$$\eta(gR) = gQ, \quad \tau(gR) = gP.$$

$\text{SO}(6, \mathbb{C})$ is the group of all matrices preserving the matrices $I = \begin{pmatrix} 0 & E_3 \\ E_3 & 0 \end{pmatrix}$, where E_3 is 3×3 identity matrix, i.e. all $X \in \text{GL}(6, \mathbb{C})$ such that $X^t I X = I$. The Lie algebra $\mathfrak{so}(6, \mathbb{C})$ of $\text{SO}(6, \mathbb{C})$ consists of all $X \in \mathfrak{gl}(6, \mathbb{C})$ such that $X^t I + I X = 0$ (cf. [4] [5]), i.e.

$$\mathfrak{so}(6, \mathbb{C}) = \left\{ \begin{bmatrix} A & B \\ C & -A^t \end{bmatrix}; B^t = -B, C^t = -C, A, B, C \in \mathfrak{gl}(3, \mathbb{C}) \right\}.$$

Consider the subalgebra of $\mathfrak{so}(6, \mathbb{C})$,

$$\mathfrak{p} = \left\{ \begin{bmatrix} * & * & * & 0 & * & * \\ * & * & * & * & 0 & * \\ 0 & 0 & * & * & * & 0 \\ 0 & 0 & 0 & * & * & 0 \\ 0 & 0 & 0 & * & * & 0 \\ 0 & 0 & 0 & * & * & * \end{bmatrix} \in \mathfrak{so}(6, \mathbb{C}) \right\}, \quad \mathfrak{h} = \left\{ \begin{bmatrix} 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ y_{00'} & y_{01'} & 0 & 0 & 0 & 0 \\ 0 & -\frac{1}{2}t & -y_{10'} & 0 & 0 & -y_{00'} \\ \frac{1}{2}t & 0 & -y_{11'} & 0 & 0 & -y_{01'} \\ y_{10'} & y_{11'} & 0 & 0 & 0 & 0 \end{bmatrix} \in \mathfrak{so}(6, \mathbb{C}) \right\}.$$

We have the decomposition $\mathfrak{so}(6, \mathbb{C}) = \mathfrak{h} \oplus \mathfrak{p}$. Similarly, we have the decomposition $\mathfrak{so}(6, \mathbb{C}) = \mathfrak{n} \oplus \mathfrak{q}$ with

$$\mathfrak{q} = \left\{ \begin{bmatrix} * & * & * & 0 & * & * \\ 0 & * & * & * & 0 & * \\ 0 & * & * & * & * & 0 \\ 0 & 0 & 0 & * & 0 & 0 \\ 0 & 0 & * & * & * & * \\ 0 & * & 0 & * & * & * \end{bmatrix} \in \mathfrak{so}(6, \mathbb{C}) \right\}, \quad \mathfrak{n} = \left\{ \begin{bmatrix} 0 & 0 & 0 & 0 & 0 & 0 \\ y_1 & 0 & 0 & 0 & 0 & 0 \\ y_2 & 0 & 0 & 0 & 0 & 0 \\ 0 & -y_3 & -y_4 & 0 & -y_1 & -y_2 \\ y_3 & 0 & 0 & 0 & 0 & 0 \\ y_4 & 0 & 0 & 0 & 0 & 0 \end{bmatrix} \in \mathfrak{so}(6, \mathbb{C}) \right\},$$

and $\mathfrak{so}(6, \mathbb{C}) = \mathfrak{m} \oplus \mathfrak{r}$, with

$$\mathfrak{r} = \mathfrak{q} \cap \mathfrak{p} = \left\{ \begin{bmatrix} * & * & * & 0 & * & * \\ 0 & * & * & * & 0 & * \\ 0 & 0 & * & * & * & 0 \\ 0 & 0 & 0 & * & 0 & 0 \\ 0 & 0 & 0 & * & * & 0 \\ 0 & 0 & 0 & * & * & * \end{bmatrix} \in \mathfrak{so}(6, \mathbb{C}) \right\}, \quad \mathfrak{m} = \left\{ \begin{bmatrix} 0 & 0 & 0 & 0 & 0 & 0 \\ * & 0 & 0 & 0 & 0 & 0 \\ * & * & 0 & 0 & 0 & 0 \\ 0 & * & * & 0 & * & * \\ * & 0 & * & 0 & 0 & * \\ * & * & 0 & 0 & 0 & 0 \end{bmatrix} \in \mathfrak{so}(6, \mathbb{C}) \right\}.$$

Take P , Q and R to be Lie groups with Lie algebra to be \mathfrak{p} , \mathfrak{q} and \mathfrak{r} , respectively.

To write down the double fibration concretely, let us recall the model case $SL(2, \mathbb{C})/B_0$ as in [13] with the parabolic subgroup

$$B_0 = \left\{ \begin{bmatrix} a & b \\ 0 & c \end{bmatrix}; a, b, c \in \mathbb{C} \text{ and } ac \neq 0 \right\}.$$

The homogeneous space $SL(2, \mathbb{C})/B_0$ is covered by two coordinate charts $\iota_1 : \mathbb{C} \rightarrow SL(2, \mathbb{C})/B_0$ given by $z \mapsto M_z B_0$ and $\iota_2 : \mathbb{C} \rightarrow SL(2, \mathbb{C})/B_0$ given by $z \mapsto W_0 M_{\bar{z}} B_0$, where $M_z = \begin{bmatrix} 1 & 0 \\ z & 1 \end{bmatrix}$, $W_0 = \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}$ is the Weyl element. Since $W_0 M_{\bar{z}} \in \begin{bmatrix} 1 & 0 \\ -\frac{1}{\bar{z}} & 1 \end{bmatrix} B_0$, the transition function of these two coordinate charts is

$$\iota_1^{-1} \circ \iota_2 : \mathbb{C} \setminus \{0\} \rightarrow \mathbb{C} \setminus \{0\}, \quad \tilde{z} \mapsto -\frac{1}{\tilde{z}}.$$

So we get $SL(2, \mathbb{C})/B_0 \cong \mathbb{C}P^1$.

For an element Y of \mathfrak{h} given by

$$Y = \begin{bmatrix} 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ y_{00'} & y_{01'} & 0 & 0 & 0 & 0 \\ 0 & -\frac{1}{2}t & -y_{10'} & 0 & 0 & -y_{00'} \\ \frac{1}{2}t & 0 & -y_{11'} & 0 & 0 & -y_{01'} \\ y_{10'} & y_{11'} & 0 & 0 & 0 & 0 \end{bmatrix},$$

it's direct to see that

$$Y^2 = \begin{bmatrix} 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ -\langle \mathbf{y}_{0'}, \mathbf{y}_{0'} \rangle & -\langle \mathbf{y}_{0'}, \mathbf{y}_{1'} \rangle & 0 & 0 & 0 & 0 \\ -\langle \mathbf{y}_{0'}, \mathbf{y}_{1'} \rangle & -\langle \mathbf{y}_{1'}, \mathbf{y}_{1'} \rangle & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{bmatrix}, \quad Y^3 = 0.$$

So we have

$$(82) \quad e^Y = \begin{bmatrix} 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ y_{00'} & y_{01'} & 1 & 0 & 0 & 0 \\ -\frac{1}{2}\langle \mathbf{y}_{0'}, \mathbf{y}_{0'} \rangle & -\frac{1}{2}t - \frac{1}{2}\langle \mathbf{y}_{0'}, \mathbf{y}_{1'} \rangle & -y_{10'} & 1 & 0 & -y_{00'} \\ \frac{1}{2}t - \frac{1}{2}\langle \mathbf{y}_{0'}, \mathbf{y}_{1'} \rangle & -\frac{1}{2}\langle \mathbf{y}_{1'}, \mathbf{y}_{1'} \rangle & -y_{11'} & 0 & 1 & -y_{01'} \\ y_{10'} & y_{11'} & 0 & 0 & 0 & 1 \end{bmatrix},$$

which we denote by $H_{(\mathbf{y}, t)}$. Then we see that $\mathcal{H} = \{H_{(\mathbf{y}, t)} | \mathbf{y} \in \mathbb{C}^4, t \in \mathbb{C}\}$ is the Lie group with Lie algebra \mathfrak{h} .

Lemma A.1. \mathcal{H} is a group, and $H : \mathcal{H} \rightarrow \mathcal{H}$, given by $(\mathbf{y}, \mathbf{t}) \mapsto H_{(\mathbf{y}, \mathbf{t})}$ is an isomorphism.

Proof. For $(\mathbf{y}, t), (\hat{\mathbf{y}}, \hat{t}) \in \mathcal{H}$, it is direct to check that

$$(83) \quad H_{(\mathbf{y}, t)} \cdot H_{(\hat{\mathbf{y}}, \hat{t})} = \begin{bmatrix} 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ y_{00'} + \hat{y}_{00'} & y_{01'} + \hat{y}_{01'} & 1 & 0 & 0 & 0 \\ \tilde{t}_{00'} & \tilde{t}_{01'} & y_{10'} + \hat{y}_{10'} & 1 & 0 & -(y_{00'} + \hat{y}_{00'}) \\ \tilde{t}_{10'} & \tilde{t}_{11'} & y_{11'} + \hat{y}_{11'} & 0 & 1 & -(y_{01'} + \hat{y}_{01'}) \\ y_{10'} + \hat{y}_{10'} & y_{11'} + \hat{y}_{11'} & 0 & 0 & 0 & 1 \end{bmatrix}$$

with

$$\tilde{t}_{00'} = -\frac{1}{2}\langle \mathbf{y}_{0'}, \mathbf{y}_{0'} \rangle - y_{10'}\hat{y}_{00'} - \frac{1}{2}\langle \hat{\mathbf{y}}_{0'}, \hat{\mathbf{y}}_{0'} \rangle - y_{00'}\hat{y}_{10'} = -\frac{1}{2}\langle \mathbf{y}_{0'} + \hat{\mathbf{y}}_{0'}, \mathbf{y}_{0'} + \hat{\mathbf{y}}_{0'} \rangle,$$

$$\begin{aligned} \tilde{t}_{10'} &= \frac{1}{2}t - \frac{1}{2}\langle \mathbf{y}_{0'}, \mathbf{y}_{1'} \rangle - \hat{y}_{00'}y_{11'} + \frac{1}{2}\hat{t} - \frac{1}{2}\langle \hat{\mathbf{y}}_{0'}, \hat{\mathbf{y}}_{1'} \rangle - y_{01'}\hat{y}_{10'} \\ &= \frac{1}{2}t + \frac{1}{2}\hat{t} - \frac{1}{2}\langle \hat{\mathbf{y}}_{0'}, \mathbf{y}_{1'} \rangle + \frac{1}{2}\langle \mathbf{y}_{0'}, \hat{\mathbf{y}}_{1'} \rangle - \frac{1}{2}\langle \mathbf{y}_{0'} + \hat{\mathbf{y}}_{0'}, \mathbf{y}_{1'} + \hat{\mathbf{y}}_{1'} \rangle, \end{aligned}$$

$$\begin{aligned} \tilde{t}_{01'} &= -\frac{1}{2}t - \frac{1}{2}\langle \mathbf{y}_{0'}, \mathbf{y}_{1'} \rangle - y_{10'}\hat{y}_{01'} - \frac{1}{2}\hat{t} - \frac{1}{2}\langle \hat{\mathbf{y}}_{0'}, \hat{\mathbf{y}}_{1'} \rangle - y_{00'}\hat{y}_{11'} \\ &= -\frac{1}{2}t - \frac{1}{2}\hat{t} + \frac{1}{2}\langle \hat{\mathbf{y}}_{0'}, \mathbf{y}_{1'} \rangle - \frac{1}{2}\langle \mathbf{y}_{0'}, \hat{\mathbf{y}}_{1'} \rangle - \frac{1}{2}\langle \mathbf{y}_{0'} + \hat{\mathbf{y}}_{0'}, \mathbf{y}_{1'} + \hat{\mathbf{y}}_{1'} \rangle, \end{aligned}$$

$$\tilde{t}_{11'} = -\frac{1}{2}\langle \mathbf{y}_{1'}, \mathbf{y}_{1'} \rangle - y_{11'}\hat{y}_{11'} - \frac{1}{2}\langle \hat{\mathbf{y}}_{1'}, \hat{\mathbf{y}}_{1'} \rangle - y_{01'}\hat{y}_{11'} = -\frac{1}{2}\langle \mathbf{y}_{1'} + \hat{\mathbf{y}}_{1'}, \mathbf{y}_{1'} + \hat{\mathbf{y}}_{1'} \rangle.$$

It follows that $H_{(\mathbf{y}, t)} \cdot H_{(\hat{\mathbf{y}}, \hat{t})} = H_{(\mathbf{y} + \hat{\mathbf{y}}, t + \hat{t} - \langle \hat{\mathbf{y}}_{0'}, \mathbf{y}_{1'} \rangle + \langle \mathbf{y}_{0'}, \hat{\mathbf{y}}_{1'} \rangle)}$, i.e. H is a group homomorphism. Obviously, it is an isomorphism. \square

Denote

$$P_\zeta := \begin{bmatrix} 1 & 0 & 0 & 0 & 0 & 0 \\ \zeta & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & -\zeta & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix} \in \text{SO}(6, \mathbb{C}).$$

In fact, it's direct to check that $P_\zeta^t I P_\zeta = I$. We consider the two coordinate charts of G/R as follows. $\varphi_1 : \mathcal{H} \times \mathbb{C} \rightarrow G/R$, $(\mathbf{y}, t, \zeta) \mapsto H_{(\mathbf{y}, t)} \cdot P_\zeta \cdot R$, and $\varphi_2 : \mathcal{H} \times \mathbb{C} \rightarrow G/R$, $(\mathbf{y}, t, \zeta) \mapsto H_{(\mathbf{y}, t)} \cdot W \cdot P_\zeta \cdot R$, where W is the element

$$W := \begin{bmatrix} 0 & 1 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix}.$$

Proposition A.1. *The transition function of these two coordinates charts is $\varphi_1^{-1} \circ \varphi_2 : \mathcal{H} \times \mathbb{C}^* \rightarrow \mathcal{H} \times \mathbb{C}^*$, $(\mathbf{y}, t, \zeta) \mapsto (\mathbf{y}, t, \zeta^{-1})$. And $\mathcal{H} \times \mathbb{C}P^1$ is biholomorphically embedded in $\text{SO}(6, \mathbb{C})/R$.*

Proof. Note that

$$(84) \quad W \cdot P_\zeta = \begin{bmatrix} \zeta & 1 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 & -\zeta & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 & 0 & 0 & 0 \\ \zeta^{-1} & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & -\zeta^{-1} & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix} \cdot \begin{bmatrix} \zeta & 1 & 0 & 0 & 0 & 0 \\ 0 & -\zeta^{-1} & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & \zeta^{-1} & 0 & 0 \\ 0 & 0 & 0 & 1 & -\zeta & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix} \\ := P_{\zeta^{-1}} \cdot A_\zeta.$$

Then

$$\varphi_2(\mathbf{y}, t, \zeta) = H_{(\mathbf{y}, t)} \cdot W \cdot P_\zeta \cdot R = H_{(\mathbf{y}, t)} \cdot P_{\zeta^{-1}} \cdot A_\zeta \cdot R = H_{(\mathbf{y}, t)} \cdot P_{\zeta^{-1}} \cdot R,$$

since $A_\zeta \in R$. So we get the transition function $\varphi_1^{-1} \circ \varphi_2(\mathbf{y}, t, \zeta) = \varphi_1^{-1}(H_{(\mathbf{y}, t)} \cdot P_{\zeta^{-1}} \cdot R) = (\mathbf{y}, t, \zeta^{-1})$. The proposition is proved. \square

Similarly, we consider two coordinate charts of G/Q . One piece is $\psi_1 : \mathbb{C}^4 \rightarrow G/Q$ given by $(\mathbf{x}, t, \zeta) \mapsto P_\zeta \cdot N_{(\mathbf{x}, t)} \cdot Q$, where $\mathbf{x} = (x_1, x_2) \in \mathbb{C}^2$ and $N_{(\mathbf{x}, t)} = H_{(\mathbf{y}, t)}$ with $\mathbf{y} = \begin{bmatrix} x_1 & 0 \\ x_2 & 0 \end{bmatrix}$. The other is $\psi_2 : \mathbb{C}^4 \rightarrow G/Q$ given by $(\mathbf{x}, t, \zeta) \mapsto W \cdot P_\zeta \cdot N_{(\mathbf{x}, t)} \cdot Q$.

Proposition A.2. *The transition function $\psi_1^{-1} \circ \psi_2 : \mathbb{C}^4 \setminus (\mathbb{C}^3 \times \{0\}) \rightarrow \mathbb{C}^4 \setminus (\mathbb{C}^3 \times \{0\})$ of these two coordinates charts is given by $(\mathbf{x}, t, \zeta) \mapsto (\zeta^{-1}\mathbf{x}, -t - \zeta^{-1}x_1x_2, \zeta^{-1})$, which coincides with (32) up to a sign.*

Proof. Note that

$$(85) \quad \psi_2(\mathbf{x}, t, \zeta) = W \cdot P_\zeta \cdot N_{(\mathbf{x}, t)} \cdot Q = P_{\zeta^{-1}} \cdot A_\zeta \cdot H_{(x_1, x_2, 0, 0, t)} \cdot A_\zeta^{-1} \cdot Q,$$

by (84) and $A_\zeta \in Q$. Note that

$$\begin{aligned} & A_\zeta \cdot H_{(x_1, x_2, 0, 0, t)} \cdot A_\zeta^{-1} \\ = & \begin{bmatrix} \zeta & 1 & 0 & 0 & 0 & 0 \\ 0 & -\zeta^{-1} & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & \zeta^{-1} & 0 & 0 \\ 0 & 0 & 0 & 1 & -\zeta & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix} \cdot \begin{bmatrix} 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ x_1 & 0 & 1 & 0 & 0 & 0 \\ -x_1x_2 & -\frac{1}{2}t & -x_2 & 1 & 0 & -x_1 \\ \frac{1}{2}t & 0 & 0 & 0 & 1 & 0 \\ x_2 & 0 & 0 & 0 & 0 & 1 \end{bmatrix} \cdot A_\zeta^{-1} \\ = & \begin{bmatrix} \zeta & 1 & 0 & 0 & 0 & 0 \\ 0 & -\zeta^{-1} & 0 & 0 & 0 & 0 \\ x_1 & 0 & 1 & 0 & 0 & 0 \\ -\zeta^{-1}x_1x_2 & -\frac{1}{2}\zeta^{-1}t & -\zeta^{-1}x_2 & \zeta^{-1} & 0 & -\zeta^{-1}x_1 \\ -x_1x_2 - \frac{1}{2}t\zeta & -\frac{1}{2}t & -x_2 & 1 & -\zeta & -x_1 \\ x_2 & 0 & 0 & 0 & 0 & 1 \end{bmatrix} \cdot \begin{bmatrix} \zeta^{-1} & 1 & 0 & 0 & 0 & 0 \\ 0 & -\zeta & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & \zeta & 0 & 0 \\ 0 & 0 & 0 & 1 & -\zeta^{-1} & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix} \end{aligned}$$

$$\begin{aligned}
 &= \begin{bmatrix} 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ \zeta^{-1}x_1 & x_1 & 1 & 0 & 0 & 0 \\ -\zeta^{-2}x_1x_2 & \frac{1}{2}t - \zeta^{-1}x_1x_2 & -\zeta^{-1}x_2 & 1 & 0 & -\zeta^{-1}x_1 \\ -\frac{1}{2}t - \zeta^{-1}x_1x_2 & -x_1x_2 & -x_2 & 0 & 1 & -x_1 \\ \zeta^{-1}x_2 & x_2 & 0 & 0 & 0 & 1 \end{bmatrix} \\
 &= H_{(\zeta^{-1}x_1, \zeta^{-1}x_2, x_1, x_2, -t)} \\
 &= H_{(\zeta^{-1}x_1, \zeta^{-1}x_2, 0, 0, -t-2\zeta^{-1}x_1x_2)} \cdot H_{(0, 0, x_1, x_2, 0)}.
 \end{aligned}$$

Substituting the above identity to (85) to get

$$\psi_2(x_1, x_2, t, \zeta) = P_{\zeta^{-1}} \cdot H_{(\zeta^{-1}x_1, \zeta^{-1}x_2, 0, 0, -t-2\zeta^{-1}x_1x_2)} \cdot Q = P_{\zeta^{-1}} \cdot N_{(\zeta^{-1}x_1, \zeta^{-1}x_2, -t-2\zeta^{-1}x_1x_2)} Q,$$

by $H_{(0, 0, x_1, x_2, 0)} \in Q$. So we have

$$\psi_1^{-1} \circ \psi_2(x_1, x_2, t, \zeta) = \psi_1^{-1}(P_{\zeta^{-1}} \cdot N_{(\zeta^{-1}x_1, \zeta^{-1}x_2, -t-2\zeta^{-1}x_1x_2)} Q) = (\zeta^{-1}x_1, \zeta^{-1}x_2, -t-2\zeta^{-1}x_1x_2, \zeta^{-1})$$

by definition. The proposition is proved. \square

Lemma A.2. $P_{\zeta}^{-1} H_{(\mathbf{y}_{0'}, \mathbf{y}_{1'}, t)} P_{\zeta} = H_{(\mathbf{y}_{0'} + \zeta \mathbf{y}_{1'}, \mathbf{y}_{1'}, t)}$.

Proof. We have

$$\begin{aligned}
 &P_{\zeta}^{-1} H_{(\mathbf{y}_{0'}, \mathbf{y}_{1'}, t)} P_{\zeta} \\
 &= \begin{bmatrix} 1 & 0 & 0 & 0 & 0 & 0 \\ -\zeta & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & \zeta & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix} \cdot \begin{bmatrix} 1 & 0 & 0 & 0 & 0 & 0 \\ y_{00'} & y_{01'} & 1 & 0 & 0 & 0 \\ -\frac{1}{2}\langle \mathbf{y}_{0'}, \mathbf{y}_{0'} \rangle & -\frac{1}{2}t - \frac{1}{2}\langle \mathbf{y}_{0'}, \mathbf{y}_{1'} \rangle & -y_{10'} & 1 & 0 & -y_{00'} \\ \frac{1}{2}t - \frac{1}{2}\langle \mathbf{y}_{0'}, \mathbf{y}_{1'} \rangle & -\frac{1}{2}\langle \mathbf{y}_{1'}, \mathbf{y}_{1'} \rangle & -y_{11'} & 0 & 1 & -y_{01'} \\ y_{10'} & y_{11'} & 0 & 0 & 0 & 1 \end{bmatrix} \cdot P_{\zeta} \\
 &= \begin{bmatrix} 1 & 0 & 0 & 0 & 0 & 0 \\ -\zeta & 1 & 0 & 0 & 0 & 0 \\ y_{00'} & y_{01'} & 1 & 0 & 0 & 0 \\ -\frac{1}{2}\langle \mathbf{y}_{0'}, \mathbf{y}_{0'} \rangle + \frac{1}{2}t\zeta - \frac{1}{2}\zeta\langle \mathbf{y}_{0'}, \mathbf{y}_{1'} \rangle & -\frac{1}{2}t - \frac{1}{2}\langle \mathbf{y}_{0'}, \mathbf{y}_{1'} \rangle - \zeta\langle \mathbf{y}_{1'}, \mathbf{y}_{1'} \rangle & -y_{10'} - \zeta y_{11'} & 1 & \zeta & -y_{00'} - \zeta y_{01'} \\ \frac{1}{2}t - \frac{1}{2}\langle \mathbf{y}_{0'}, \mathbf{y}_{1'} \rangle & -\frac{1}{2}\langle \mathbf{y}_{1'}, \mathbf{y}_{1'} \rangle & -y_{11'} & 0 & 1 & -y_{01'} \\ y_{10'} & y_{11'} & 0 & 0 & 0 & 1 \end{bmatrix} \cdot P_{\zeta} \\
 &= \begin{bmatrix} 1 & 0 & 0 & 0 & 0 & 0 \\ y_{00'} + \zeta y_{01'} & y_{01'} & 1 & 0 & 0 & 0 \\ -\frac{1}{2}\langle \mathbf{y}_{0'} + \zeta \mathbf{y}_{1'}, \mathbf{y}_{0'} + \zeta \mathbf{y}_{1'} \rangle & -\frac{1}{2}t - \frac{1}{2}\langle \mathbf{y}_{0'} + \zeta \mathbf{y}_{1'}, \mathbf{y}_{1'} \rangle & -y_{10'} - \zeta y_{11'} & 1 & 0 & -y_{00'} - \zeta y_{01'} \\ \frac{1}{2}t - \frac{1}{2}\langle \mathbf{y}_{0'} + \zeta \mathbf{y}_{1'}, \mathbf{y}_{1'} \rangle & -\frac{1}{2}\langle \mathbf{y}_{1'}, \mathbf{y}_{1'} \rangle & -y_{11'} & 0 & 1 & -y_{01'} \\ y_{10'} + \zeta y_{11'} & y_{11'} & 0 & 0 & 0 & 1 \end{bmatrix} \\
 &= H_{(\mathbf{y}_{0'} + \zeta \mathbf{y}_{1'}, \mathbf{y}_{1'}, t)}
 \end{aligned}$$

by the definition (82) of $H_{(\mathbf{y}_{0'}, \mathbf{y}_{1'}, t)}$. \square

Proposition A.3. *The mapping $\eta : \text{SO}(6, \mathbb{C})/R \longrightarrow \text{SO}(6, \mathbb{C})/Q$ in (81) is locally given by*

$$\eta(H_{(\mathbf{y}, t)} \cdot P_{\zeta} R) = P_{\zeta} \cdot N_{(y_{00'} + \zeta y_{01'}, y_{10'} + \zeta y_{11'}, t - \langle \mathbf{y}_{0'} + \zeta \mathbf{y}_{1'}, \mathbf{y}_{1'} \rangle)} Q,$$

which coincides with (24).

Proof. By (A.2), we have

$$\begin{aligned}
 \eta(H_{(\mathbf{y}, t)} \cdot P_{\zeta} R) &= H_{(\mathbf{y}_{0'}, \mathbf{y}_{1'}, t)} \cdot P_{\zeta} Q = P_{\zeta} \cdot P_{\zeta}^{-1} \cdot H_{(\mathbf{y}_{0'}, \mathbf{y}_{1'}, t)} \cdot P_{\zeta} Q = P_{\zeta} \cdot H_{(\mathbf{y}_{0'} + \zeta \mathbf{y}_{1'}, \mathbf{y}_{1'}, t)} Q \\
 &= P_{\zeta} \cdot H_{(\mathbf{y}_{0'} + \zeta \mathbf{y}_{1'}, \mathbf{0}, t - \langle \mathbf{y}_{0'} + \zeta \mathbf{y}_{1'}, \mathbf{y}_{1'} \rangle)} \cdot H_{(\mathbf{0}, \mathbf{y}_{1'}, 0)} Q = P_{\zeta} \cdot H_{(\mathbf{y}_{0'} + \zeta \mathbf{y}_{1'}, \mathbf{0}, t - \langle \mathbf{y}_{0'} + \zeta \mathbf{y}_{1'}, \mathbf{y}_{1'} \rangle)} Q \\
 &= P_{\zeta} \cdot N_{(y_{00'} + \zeta y_{01'}, y_{10'} + \zeta y_{11'}, t - \langle \mathbf{y}_{0'} + \zeta \mathbf{y}_{1'}, \mathbf{y}_{1'} \rangle)} Q
 \end{aligned}$$

by $H_{(\mathbf{0}, \mathbf{y}_{1'}, 0)} \in Q$. The result follows. \square

REFERENCES

- [1] Baston, R. and Eastwood, M., *The Penrose transform. Its interaction with representation theory*, Oxford Mathematical Monographs, The Clarendon Press, Oxford University Press, New York, 1989.
- [2] Dunajski, M., *Solitons, instantons, and twistors*, Oxford Graduate Texts in Mathematics **19**, Oxford University Press, Oxford, 2010.
- [3] Eastwood, M., Penrose, R. and Wells, R., Cohomology and massless fields, *Comm. Math. Phys.* **78** (1980), 305-351.
- [4] Fulton, W. and Harris, J., *Representation theory, A first course*, Graduate Texts in Mathematics **129**, Springer-Verlag, New York, 1991.
- [5] Hall, B., *Lie groups, Lie algebras, and representations, an elementary introduction*, Graduate Texts in Mathematics **222**, Springer-Verlag, New York, 2003.
- [6] Hosomichi, K., Seong, R.-K. and Terashima, S., Supersymmetric gauge theories on the five-sphere, *Nuclear Phys. B.* **865** (2012), 376-396.
- [7] Itoh, M., Contact metric 5-manifolds, CR twistor spaces and integrability, *J. of Math. Phys.* **43** (2002), 3783-3797.
- [8] Källén, J. and Zabzine, M., Twisted supersymmetric 5D Yang-Mills theory and contact geometry, *J. High Energy Phys.* **5** (2012), Article 125.
- [9] Manin, Y., *Gauge field theory and complex geometry, Grundlehren der Mathematischen Wissenschaften* **289**, Springer-Verlag, Berlin, second edition, 1997.
- [10] Mason, L. and Woodhouse, N., *Integrability, self-duality, and twistor theory*, London Mathematical Society Monographs (New Series) **15**, The Clarendon Press, Oxford University Press, New York, 1996.
- [11] Popov, A.D. and Martin, W., Hidden symmetries and integrable hierarchy of the $N = 4$ supersymmetric Yang-Mills equations, *Comm. Math. Phys.* **275** (2007), 685-708.
- [12] Ren, G.-Z., Shi, Y. and Wang, W., The tangential k -Cauchy-Fueter operator and k -CF functions over the Heisenberg group, *Adv. Appl. Clifford Algebr.* **30(2)** (2020), Article 20.
- [13] Wang, W., On twistor transformations and invariant differential operator of simple Lie group $G_{2(2)}$, *J. Math. Phys.* **54** (2013), 013502.
- [14] Wang, W., The quaternionic Monge-Ampère operator and plurisubharmonic functions on the Heisenberg group, *Math. Z.* (2021), <https://doi.org/10.1007/s00209-020-02608-3>.
- [15] Ward, R. S., On self-dual gauge fields, *Phys. Lett. A.* **61** (1991), 81-82.
- [16] Ward, R. S. and Wells, R., *Twistor geometry and field theory*, Cambridge Monographs on Mathematical Physics, Cambridge University Press, Cambridge, 1990.
- [17] Wolf, M., Contact manifolds, contact instantons, and twistor geometry, *J. High Energy Phys.* **7** (2012), Article 074.

DEPARTMENT OF MATHEMATICS, ZHEJIANG INTERNATIONAL STUDIES UNIVERSITY, HANGZHOU 310023, PR CHINA

Email address, G.-Z. Ren: gzren@zisu.edu.cn

DEPARTMENT OF MATHEMATICS, ZHEJIANG UNIVERSITY, ZHEJIANG 310027, PR CHINA

Email address, W. Wang: wwang@zju.edu.cn