

Convergence of series of moments on general exponential inequality

João Lita da Silva¹

*Department of Mathematics and GeoBioTec
NOVA School of Sciences and Technology
NOVA University of Lisbon
Quinta da Torre, 2829-516 Caparica, Portugal*

Vanda Lourenço²

*Department of Mathematics and CMA
NOVA School of Sciences and Technology
NOVA University of Lisbon
Quinta da Torre, 2829-516 Caparica, Portugal*

Abstract

For an array $\{X_{n,j}, 1 \leq j \leq k_n, n \geq 1\}$ of random variables and a sequence $\{c_n\}$ of positive numbers, sufficient conditions are given under which, for all $\varepsilon > 0$, $\sum_{n=1}^{\infty} c_n \mathbb{E} \left[\max_{1 \leq i \leq k_n} \left| \sum_{j=1}^i (X_{n,j} - \mathbb{E} X_{n,j} I_{\{|X_{n,j}| \leq \delta\}}) \right| - \varepsilon \right]_+^p < \infty$, where x_+ denotes the positive part of x and $p \geq 1, \delta > 0$. Our statements are announced in a general setting allowing to conclude the previous convergence for well-known dependent structures. As an application, we study complete consistency and consistency in the r th mean of cumulative sum type estimators of the change in the mean of dependent observations.

Keywords: maximum of partial sums, convergence of series of moments, dependent random variables, CUSUM-type estimators

2010 Mathematics Subject Classification: 60F15; 62F12

1 Introduction

Probability inequalities play a central role in proofs of asymptotic results for sums of random variables with maximal probability inequalities being key to the establishment of sharp rates of convergence. The main goal of this paper is to obtain the convergence of series of moments involving the maximum of partial row sums of arrays of random variables under a general exponential-type maximal inequality. Our results encompass some of those of [3],

¹*E-mail address:* jfls@fct.unl.pt; joao.lita@gmail.com (corresponding author)

²*E-mail address:* vmml@fct.unl.pt

[16] and [19] as particular cases. Furthermore, our proofs are presented in a more direct and optimised way that enables to weaken the assumptions presented in [16].

The paper is organized as follows. Section 2 presents our results and proofs. In Section 3, we consider a (general) change-point model and apply our results to a cumulative sum (CUSUM) estimator of the point of shift in the mean of dependent observations, providing sufficient conditions for its complete consistency and consistency in the r th mean. Our theoretical results are illustrated via simulation and at the light of the CUSUM estimator described in Section 3.

In what follows, $\{k_n\}$ indicates a sequence of positive integers such that $k_n \rightarrow \infty$ as $n \rightarrow \infty$, and I_A stands for the indicator random variable of an event A . We additionally write $x \wedge y$ and $x \vee y$ for $\min(x, y)$ and $\max(x, y)$, respectively. Moreover, for each $\ell > 0$, we define the function $g_\ell(x) = (x \wedge \ell) \vee (-\ell)$, which describes the truncation at level ℓ .

2 Main results

Considering an array $\{X_{n,j}, 1 \leq j \leq k_n, n \geq 1\}$ of random variables, we admit that, for each $n \geq 1$ and all $\ell, \lambda, \eta > 0$, there exist $C_1, C_2 > 0$ non-depending on n, ℓ, λ, η such that

$$\begin{aligned} & \mathbb{P} \left\{ \max_{1 \leq i \leq k_n} \left| \sum_{j=1}^i [g_\ell(X_{n,j}) - \mathbb{E} g_\ell(X_{n,j})] \right| \geq \lambda \right\} \\ & \leq \alpha_n \mathbb{P} \left\{ \max_{1 \leq j \leq k_n} |[g_\ell(X_{n,j}) - \mathbb{E} g_\ell(X_{n,j})]| > C_1 \eta \right\} + \beta_n \left\{ \frac{\sum_{j=1}^{k_n} \mathbb{E}[g_\ell(X_{n,j}) - \mathbb{E} g_\ell(X_{n,j})]^2}{C_2 \lambda \eta} \right\}^{\lambda/\eta} \end{aligned} \quad (2.1)$$

for some sequences $\{\alpha_n\}, \{\beta_n\}$ of nonnegative numbers. Let us point out that well-known inequalities are contained in (2.1), particularly, the ones listed below.

EXAMPLE Consider $s_n(\ell) := \sum_{j=1}^{k_n} \mathbb{E}[g_\ell(X_{n,j}) - \mathbb{E} g_\ell(X_{n,j})]^2 \neq 0$ for all $\ell > 0$ and $n \geq 1$.

1. The notion of m -negatively associated random variables was introduced in [9] (see Definition 2 of [9]). Let $\{X_{n,j}, 1 \leq j \leq k_n, n \geq 1\}$ be an array of *row-wise m -negatively associated random variables*, i.e. for each $n \geq 1$, the sequence $\{X_{n,j}, 1 \leq j \leq k_n\}$ of random variables is m -negatively associated. According to Property 6 of [10], for all $\ell > 0$ and each $n \geq 1$, $\{g_\ell(X_{n,j}), 1 \leq j \leq k_n\}$ is a sequence of m -negatively associated random variables, so that $\{g_\ell(X_{n,j}), 1 \leq j \leq k_n, n \geq 1\}$ is an array of row-wise m -negatively associated random variables. By Lemma 2.1 and Remark 2.1 of [20], we have, for all $x, a > 0$ and $n \geq 1$,

$$\begin{aligned} & \mathbb{P} \left\{ \max_{1 \leq i \leq k_n} \left| \sum_{j=1}^i [g_\ell(X_{n,j}) - \mathbb{E} g_\ell(X_{n,j})] \right| \geq x \right\} \\ & \leq 2m \mathbb{P} \left\{ \max_{1 \leq j \leq k_n} |[g_\ell(X_{n,j}) - \mathbb{E} g_\ell(X_{n,j})]| > a \right\} + 8m \left[1 + \frac{3xa}{2ms_n(\ell)} \right]^{-x/(12ma)} \quad (2.2) \\ & \leq 2m \mathbb{P} \left\{ \max_{1 \leq j \leq k_n} |[g_\ell(X_{n,j}) - \mathbb{E} g_\ell(X_{n,j})]| > a \right\} + 8m \left[\frac{2ms_n(\ell)}{3xa} \right]^{x/(12ma)}. \end{aligned}$$

In fact, inequality (2.2) can be directly obtained from (2.1) by choosing $\lambda = x$, $\eta = 12ma$, $\alpha_n = 2m$, $\beta_n = 8m$, with $C_1 = 1/(12m)$ and $C_2 = 1/(8m^2)$.

2. The concept of negatively superadditive dependent random variables appeared in [8] and it has been employed by some authors since then. If for each $n \geq 1$, the sequence $\{X_{n,j}, 1 \leq j \leq k_n\}$ of random variables is negatively superadditive dependent, then the array $\{X_{n,j}, 1 \leq j \leq k_n, n \geq 1\}$ is said to be *row-wise negatively superadditive dependent*. Supposing an array $\{X_{n,j}, 1 \leq j \leq k_n, n \geq 1\}$ of row-wise negatively superadditive dependent random variables and $\ell > 0$, we have that, for each $n \geq 1$, $\{g_\ell(X_{n,j}), 1 \leq j \leq k_n\}$ is still a sequence of negatively superadditive dependent random variables (see P3 in [8]), and whence $\{g_\ell(X_{n,j}), 1 \leq j \leq k_n, n \geq 1\}$ is an array of negatively superadditive dependent random variables. Thus, for every $x, y > 0$ and $n \geq 1$,

$$\begin{aligned} & \mathbb{P} \left\{ \max_{1 \leq i \leq k_n} \left| \sum_{j=1}^i [g_\ell(X_{n,j}) - \mathbb{E} g_\ell(X_{n,j})] \right| \geq x \right\} \\ & \leq 2\mathbb{P} \left\{ \max_{1 \leq j \leq k_n} |[g_\ell(X_{n,j}) - \mathbb{E} g_\ell(X_{n,j})]| > y \right\} + 8 \left[\frac{2s_n(\ell)}{3xy} \right]^{x/(12y)} \end{aligned} \quad (2.3)$$

(see Lemma 3.1 of [19]), which follows from (2.1) by taking $\lambda = x$, $\eta = 12a$, $\alpha_n = 2$, $\beta_n = 8$; in this case, $C_1 = 1/12$ and $C_2 = 1/8$.

Throughout the proofs below, given $t > 0$ and an array of random variables $\{X_{n,j}, 1 \leq j \leq k_n, n \geq 1\}$, we denote

$$\begin{aligned} X'_{n,j}(t) &:= X_{n,j} I_{\{|X_{n,j}| \leq t\}}, \\ X''_{n,j}(t) &:= t I_{\{X_{n,j} > t\}} - t I_{\{X_{n,j} < -t\}}, \\ S_{n,i}(t) &:= \sum_{j=1}^i (X_{n,j} - \mathbb{E} X_{n,j} I_{\{|X_{n,j}| \leq t\}}), \quad i = 1, 2, \dots, k_n, \\ \Gamma_n(t) &:= \bigcap_{j=1}^{k_n} \{X_{n,j} = X'_{n,j}(t)\}. \end{aligned} \quad (2.4)$$

Our first general statement is the following.

Theorem 1 *Let $\{c_n\}$ be a sequence of positive numbers, and $\{X_{n,j}, 1 \leq j \leq k_n, n \geq 1\}$ be an array of random variables satisfying (2.1) for some sequences $\{\alpha_n\}$, $\{\beta_n\}$ of nonnegative numbers. If*

(i) *for all $\lambda > 0$, $\sum_{n=1}^{\infty} c_n(1 + \alpha_n) \sum_{j=1}^{k_n} \mathbb{P}\{|X_{n,j}| > \lambda\} < \infty$,*

(ii) *there exist $\delta > 0$ and $q \geq 1$ such that*

$$\sum_{n=1}^{\infty} c_n \beta_n \left(\sum_{j=1}^{k_n} \mathbb{P}\{|X_{n,j}| > \delta\} \right)^q < \infty \quad \text{and} \quad \sum_{n=1}^{\infty} c_n \beta_n \left(\sum_{j=1}^{k_n} \mathbb{V} X_{n,j} I_{\{|X_{n,j}| \leq \delta\}} \right)^q < \infty,$$

then, for all $\varepsilon > 0$,

$$\sum_{n=1}^{\infty} c_n \mathbb{P} \left\{ \max_{1 \leq i \leq k_n} \left| \sum_{j=1}^i (X_{n,j} - \mathbb{E} X_{n,j} I_{\{|X_{n,j}| \leq \delta\}}) \right| > \varepsilon \right\} < \infty. \quad (2.5)$$

Proof. Fixing arbitrarily $\varepsilon > 0$ and choosing $\lambda = \varepsilon/2$, $\eta = \varepsilon/(2q)$ there are $C_1, C_2 > 0$ such that (2.1) holds with sequences $\{\alpha_n\}$, $\{\beta_n\}$. Considering $X'_{n,j}(\delta)$, $X''_{n,j}(\delta)$, $\Gamma_n(\delta)$ and $S_{n,i}(\delta)$, $i = 1, 2, \dots, k_n$ defined in (2.4), it follows $X'_{n,j}(\delta) + X''_{n,j}(\delta) = g_\delta(X_{n,j})$ and

$$\begin{aligned} & \sum_{n=1}^{\infty} c_n \mathbb{P} \left\{ \max_{1 \leq i \leq k_n} |S_{n,i}(\delta)| > \varepsilon \right\} \\ &= \sum_{n=1}^{\infty} c_n \mathbb{P} \left[\left\{ \max_{1 \leq i \leq k_n} |S_{n,i}(\delta)| > \varepsilon \right\} \cap \Gamma_n(\delta) \right] + \sum_{n=1}^{\infty} c_n \mathbb{P} \left[\left\{ \max_{1 \leq i \leq k_n} |S_{n,i}(\delta)| > \varepsilon \right\} \cap \Gamma_n(\delta)^c \right] \\ &\leq \sum_{n=1}^{\infty} c_n \mathbb{P} \left\{ \max_{1 \leq i \leq k_n} \left| \sum_{j=1}^i [X'_{n,j}(\delta) - \mathbb{E} X'_{n,j}(\delta)] \right| > \varepsilon \right\} + \sum_{n=1}^{\infty} c_n \sum_{j=1}^{k_n} \mathbb{P}\{|X_{n,j}| > \delta\} \\ &= \sum_{n=1}^{\infty} c_n \mathbb{P} \left\{ \max_{1 \leq i \leq k_n} \left| \sum_{j=1}^i [g_\delta(X_{n,j}) - \mathbb{E} g_\delta(X_{n,j}) + \mathbb{E} X''_{n,j}(\delta) - X''_{n,j}(\delta)] \right| > \varepsilon \right\} \\ &\quad + \sum_{n=1}^{\infty} c_n \sum_{j=1}^{k_n} \mathbb{P}\{|X_{n,j}| > \delta\} \\ &\leq \sum_{n=1}^{\infty} c_n \mathbb{P} \left\{ \max_{1 \leq i \leq k_n} \left| \sum_{j=1}^i [g_\delta(X_{n,j}) - \mathbb{E} g_\delta(X_{n,j})] \right| > \frac{\varepsilon}{2} \right\} \\ &\quad + \sum_{n=1}^{\infty} c_n \mathbb{P} \left\{ \max_{1 \leq i \leq k_n} \left| \sum_{j=1}^i [X''_{n,j}(\delta) - \mathbb{E} X''_{n,j}(\delta)] \right| > \frac{\varepsilon}{2} \right\} + \sum_{n=1}^{\infty} c_n \sum_{j=1}^{k_n} \mathbb{P}\{|X_{n,j}| > \delta\} \\ &\leq \sum_{n=1}^{\infty} c_n \mathbb{P} \left\{ \max_{1 \leq i \leq k_n} \left| \sum_{j=1}^i [g_\delta(X_{n,j}) - \mathbb{E} g_\delta(X_{n,j})] \right| > \frac{\varepsilon}{2} \right\} \\ &\quad + \sum_{n=1}^{\infty} \frac{2c_n}{\varepsilon} \mathbb{E} \left\{ \max_{1 \leq i \leq k_n} \left| \sum_{j=1}^i [X''_{n,j}(\delta) - \mathbb{E} X''_{n,j}(\delta)] \right| \right\} + \sum_{n=1}^{\infty} c_n \sum_{j=1}^{k_n} \mathbb{P}\{|X_{n,j}| > \delta\} \\ &\leq \sum_{n=1}^{\infty} c_n \mathbb{P} \left\{ \max_{1 \leq i \leq k_n} \left| \sum_{j=1}^i [g_\delta(X_{n,j}) - \mathbb{E} g_\delta(X_{n,j})] \right| > \frac{\varepsilon}{2} \right\} + \left(1 + \frac{4\delta}{\varepsilon}\right) \sum_{n=1}^{\infty} c_n \sum_{j=1}^{k_n} \mathbb{P}\{|X_{n,j}| > \delta\}. \end{aligned}$$

According to assumption (i), it suffices to prove

$$\sum_{n=1}^{\infty} c_n \mathbb{P} \left\{ \max_{1 \leq i \leq k_n} \left| \sum_{j=1}^i [g_\delta(X_{n,j}) - \mathbb{E} g_\delta(X_{n,j})] \right| > \frac{\varepsilon}{2} \right\} < \infty.$$

Letting \mathbf{N} be the set of positive integers, and

$$N = N(\delta, \varepsilon, q) = \left\{ n \in \mathbf{N} : \sum_{j=1}^{k_n} \mathbb{P} \left\{ |X_{n,j}| > \delta \wedge \frac{\varepsilon C_1}{8q} \right\} \geq \frac{C_1 \varepsilon}{16\delta q} \right\} \quad (2.6)$$

we have $\mathbf{N} = (\mathbf{N} \setminus N) \cup N$. Since (i) implies

$$\begin{aligned} & \sum_{n \in N} c_n \mathbb{P} \left\{ \max_{1 \leq i \leq k_n} \left| \sum_{j=1}^i [g_\delta(X_{n,j}) - \mathbb{E}g_\delta(X_{n,j})] \right| > \frac{\varepsilon}{2} \right\} \leq \sum_{n \in N} c_n \\ & \leq \sum_{n \in N} \frac{16\delta q}{C_1 \varepsilon} c_n \sum_{j=1}^{k_n} \mathbb{P} \left\{ |X_{n,j}| > \delta \wedge \frac{C_1 \varepsilon q}{8q} \right\} \leq \frac{16\delta q}{C_1 \varepsilon} \sum_{n=1}^{\infty} c_n \sum_{j=1}^{k_n} \mathbb{P} \left\{ |X_{n,j}| > \delta \wedge \frac{C_1 \varepsilon}{8q} \right\} < \infty, \end{aligned}$$

one only needs to show that

$$\sum_{n \in \mathbf{N} \setminus N} c_n \mathbb{P} \left\{ \max_{1 \leq i \leq k_n} \left| \sum_{j=1}^i [g_\delta(X_{n,j}) - \mathbb{E}g_\delta(X_{n,j})] \right| > \frac{\varepsilon}{2} \right\} < \infty. \quad (2.7)$$

From (2.1), we obtain

$$\begin{aligned} & \sum_{n \in \mathbf{N} \setminus N} c_n \mathbb{P} \left\{ \max_{1 \leq i \leq k_n} \left| \sum_{j=1}^i [g_\delta(X_{n,j}) - \mathbb{E}g_\delta(X_{n,j})] \right| > \frac{\varepsilon}{2} \right\} \\ & \leq \sum_{n \in \mathbf{N} \setminus N} c_n \alpha_n \mathbb{P} \left\{ \max_{1 \leq j \leq k_n} |g_\delta(X_{n,j}) - \mathbb{E}g_\delta(X_{n,j})| > \frac{C_1 \varepsilon}{2q} \right\} + \sum_{n \in \mathbf{N} \setminus N} c_n \beta_n \left[\frac{4q s_n(\delta)}{C_2 \varepsilon^2} \right]^q. \end{aligned}$$

Since for every $n \in \mathbf{N} \setminus N$,

$$\begin{aligned} & \max_{1 \leq j \leq k_n} |\mathbb{E}g_\delta(X_{n,j})| \\ & \leq \max_{1 \leq j \leq k_n} \left[\mathbb{E}|X_{n,j}| I_{\{|X_{n,j}| \leq \delta\}} + \delta \mathbb{P}\{|X_{n,j}| > \delta\} \right] \\ & \leq \max_{1 \leq j \leq k_n} \left[\mathbb{E}|X_{n,j}| I_{\{|X_{n,j}| \leq \delta \wedge \frac{C_1 \varepsilon}{8q}\}} + \mathbb{E}|X_{n,j}| I_{\{\delta \wedge \frac{C_1 \varepsilon}{8q} < |X_{n,j}| \leq \delta\}} + \delta \mathbb{P}\{|X_{n,j}| > \delta\} \right] \\ & \leq \delta \wedge \frac{C_1 \varepsilon}{8q} + \max_{1 \leq j \leq k_n} \left[\delta \mathbb{P} \left\{ |X_{n,j}| > \delta \wedge \frac{C_1 \varepsilon}{8q} \right\} + \delta \mathbb{P}\{|X_{n,j}| > \delta\} \right] \\ & \leq \frac{C_1 \varepsilon}{8q} + \max_{1 \leq j \leq k_n} \left[2\delta \mathbb{P} \left\{ |X_{n,j}| > \delta \wedge \frac{C_1 \varepsilon}{8q} \right\} \right] \\ & \leq \frac{C_1 \varepsilon}{8q} + 2\delta \sum_{j=1}^{k_n} \mathbb{P} \left\{ |X_{n,j}| > \delta \wedge \frac{C_1 \varepsilon}{8q} \right\} \\ & < \frac{C_1 \varepsilon}{4q} \end{aligned}$$

we get

$$\begin{aligned}
& \sum_{n \in \mathbf{N} \setminus N} c_n \alpha_n \mathbb{P} \left\{ \max_{1 \leq j \leq k_n} |g_\delta(X_{n,j}) - \mathbb{E}g_\delta(X_{n,j})| > \frac{C_1 \varepsilon}{2q} \right\} \\
& \leq \sum_{n \in \mathbf{N} \setminus N} c_n \alpha_n \mathbb{P} \left\{ \max_{1 \leq j \leq k_n} |g_\delta(X_{n,j})| > \frac{C_1 \varepsilon}{4q} \right\} \\
& \leq \sum_{n \in \mathbf{N} \setminus N} c_n \alpha_n \sum_{j=1}^{k_n} \mathbb{P} \left\{ |g_\delta(X_{n,j})| > \frac{C_1 \varepsilon}{4q} \right\} \\
& \leq \sum_{n \in \mathbf{N} \setminus N} c_n \alpha_n \sum_{j=1}^{k_n} \mathbb{P} \left\{ |X_{n,j}| > \frac{C_1 \varepsilon}{4q} \right\} \\
& \leq \sum_{n=1}^{\infty} c_n \alpha_n \sum_{j=1}^{k_n} \mathbb{P} \left\{ |X_{n,j}| > \frac{C_1 \varepsilon}{4q} \right\} \\
& < \infty
\end{aligned} \tag{2.8}$$

by virtue of assumption (i). According to Cauchy-Schwarz inequality (see [15], page 150) and the elementary inequality, $2uv \leq u^2 + v^2$ for all real numbers u, v , we have

$$\begin{aligned}
s_n(\delta) & := \sum_{j=1}^{k_n} \mathbb{E}[g_\delta(X_{n,j}) - \mathbb{E}g_\delta(X_{n,j})]^2 \\
& = \sum_{j=1}^{k_n} \mathbb{V}[g_\delta(X_{n,j})] \\
& = \sum_{j=1}^{k_n} \mathbb{V}X'_{n,j}(\delta) + \sum_{j=1}^{k_n} \mathbb{V}X''_{n,j}(\delta) + 2 \sum_{j=1}^{k_n} \text{Cov} [X'_{n,j}(\delta), X''_{n,j}(\delta)] \\
& \leq \sum_{j=1}^{k_n} \mathbb{V}X'_{n,j}(\delta) + \sum_{j=1}^{k_n} \mathbb{V}X''_{n,j}(\delta) + 2 \sum_{j=1}^{k_n} [\mathbb{V}X'_{n,j}(\delta)]^{1/2} [\mathbb{V}X''_{n,j}(\delta)]^{1/2} \\
& \leq 2 \sum_{j=1}^{k_n} \mathbb{V}X'_{n,j}(\delta) + 2 \sum_{j=1}^{k_n} \mathbb{V}X''_{n,j}(\delta) \\
& \leq 2 \sum_{j=1}^{k_n} \mathbb{V}X'_{n,j}(\delta) + 2\delta^2 \sum_{j=1}^{k_n} \mathbb{P}\{|X_{n,j}| > \delta\}
\end{aligned}$$

where the last inequality follows from $\mathbb{V}X''_{n,j}(\delta) \leq \mathbb{E}[X''_{n,j}(\delta)]^2 = \delta^2 \mathbb{P}\{|X_{n,j}| > \delta\}$. By assump-

tion (ii) and the inequality $(u + v)^q \leq 2^{q-1}(u^q + v^q)$ for any $u, v \geq 0, q \geq 1$, we obtain

$$\begin{aligned}
& \sum_{n \in \mathbf{N} \setminus N} c_n \beta_n \left[\frac{4q s_n(\delta)}{C_2 \varepsilon^2} \right]^q \\
& \leq \sum_{n \in \mathbf{N} \setminus N} c_n \beta_n \frac{2^{3q-1} q^q}{C_2^q \varepsilon^{2q}} \left\{ \left[2 \sum_{j=1}^{k_n} \mathbb{V} X'_{n,j}(\delta) \right]^q + \left(2\delta^2 \sum_{j=1}^{k_n} \mathbb{P}\{|X_{n,j}| > \delta\} \right)^q \right\} \\
& \leq \frac{2^{4q-1} q^q}{C_2^q \varepsilon^{2q}} \sum_{n=1}^{\infty} c_n \beta_n \left(\sum_{j=1}^{k_n} \mathbb{V} X_{n,j} I_{\{|X_{n,j}| \leq \delta\}} \right)^q + \frac{2^{4q-1} q^q \delta^{2q}}{C_2^q \varepsilon^{2q}} \sum_{n=1}^{\infty} c_n \beta_n \left(\sum_{j=1}^{k_n} \mathbb{P}\{|X_{n,j}| > \delta\} \right)^q \\
& < \infty.
\end{aligned} \tag{2.9}$$

The thesis is established. \square

Inequality (2.1) with $\{\alpha_n\}$ and $\{\beta_n\}$ bounded sequences allows us to improve assumption (ii) of Theorem 1.

Theorem 2 *Let $\{c_n\}$ be a sequence of positive numbers, and $\{X_{n,j}, 1 \leq j \leq k_n, n \geq 1\}$ be an array of random variables satisfying (2.1) for sequences $\{\alpha_n\}, \{\beta_n\}$ of nonnegative numbers such that $\alpha_n = O(1) = \beta_n$ as $n \rightarrow \infty$. If*

(i') *for all $\lambda > 0, \sum_{n=1}^{\infty} c_n \sum_{j=1}^{k_n} \mathbb{P}\{|X_{n,j}| > \lambda\} < \infty,$*

(ii') *there exist $\delta > 0$ and $q \geq 1$ such that $\sum_{n=1}^{\infty} c_n \left(\sum_{j=1}^{k_n} \mathbb{V} X_{n,j} I_{\{|X_{n,j}| \leq \delta\}} \right)^q < \infty,$*

then, for all $\varepsilon > 0,$ (2.5) holds.

Proof. The proof follows exactly the same steps of the preceding one, except (2.9). From $\alpha_n = O(1) = \beta_n, n \rightarrow \infty$ there is $C > 0$ (non-depending on n) such that $\alpha_n, \beta_n \leq C$ for all $n \geq 1$ and instead of (2.9), we have

$$\begin{aligned}
& \sum_{n \in \mathbf{N} \setminus N} c_n \beta_n \left[\frac{4q s_n(\delta)}{C_2 \varepsilon^2} \right]^q \\
& \leq \sum_{n \in \mathbf{N} \setminus N} c_n \frac{2^{3q-1} q^q C}{C_2^q \varepsilon^{2q}} \left\{ \left[2 \sum_{j=1}^{k_n} \mathbb{V} X'_{n,j}(\delta) \right]^q + \left(2\delta^2 \sum_{j=1}^{k_n} \mathbb{P}\{|X_{n,j}| > \delta\} \right)^q \right\} \\
& \leq \sum_{n=1}^{\infty} c_n \frac{2^{4q-1} q^q C}{C_2^q \varepsilon^{2q}} \left(\sum_{j=1}^{k_n} \mathbb{V} X_{n,j} I_{\{|X_{n,j}| \leq \delta\}} \right)^q + \sum_{n \in \mathbf{N} \setminus N} c_n \frac{\delta^q C_1^q C}{2C_2^q \varepsilon^q} \left(\frac{16\delta q}{C_1 \varepsilon} \sum_{j=1}^{k_n} \mathbb{P}\{|X_{n,j}| > \delta\} \right)^q \\
& \leq \frac{2^{4q-1} q^q C}{C_2^q \varepsilon^{2q}} \sum_{n=1}^{\infty} c_n \left(\sum_{j=1}^{k_n} \mathbb{V} X_{n,j} I_{\{|X_{n,j}| \leq \delta\}} \right)^q + \sum_{n \in \mathbf{N} \setminus N} c_n \frac{8q \delta^{q+1} C_1^{q-1} C}{C_2^q \varepsilon^{q+1}} \sum_{j=1}^{k_n} \mathbb{P}\{|X_{n,j}| > \delta\} \\
& \leq \frac{2^{4q-1} q^q C}{C_2^q \varepsilon^{2q}} \sum_{n=1}^{\infty} c_n \left(\sum_{j=1}^{k_n} \mathbb{V} X_{n,j} I_{\{|X_{n,j}| \leq \delta\}} \right)^q + \frac{8q \delta^{q+1} C_1^{q-1} C}{C_2^q \varepsilon^{q+1}} \sum_{n=1}^{\infty} c_n \sum_{j=1}^{k_n} \mathbb{P}\{|X_{n,j}| > \delta\}
\end{aligned}$$

$< \infty$

because for each $n \in \mathbf{N} \setminus N$, $\sum_{j=1}^{k_n} \mathbb{P}\{|X_{n,j}| > \delta\} \leq \sum_{j=1}^{k_n} \mathbb{P}\{|X_{n,j}| > \delta \wedge C_1\varepsilon/(8q)\} < C_1\varepsilon/(16\delta q)$. The proof is complete. \square

Remark Let us observe that Theorem 1 in [3] or Theorem 1 in [9] can be both obtained from Theorem 2 by using inequality (2.2) (particularly with $m = 1$ in the former). Further, by employing inequality (2.3), Theorem 3.3 of [19] is also a consequence of Theorem 2.

The next result presents sufficient conditions under which a series of moments involving the maximum of partial row sums of arrays of random variables converges.

Theorem 3 *Let $p \geq 1$, $\{c_n\}$ be a sequence of positive numbers, and $\{X_{n,j}, 1 \leq j \leq k_n, n \geq 1\}$ be an array of random variables verifying (2.1) for some sequences $\{\alpha_n\}, \{\beta_n\}$ of nonnegative numbers. If there exists a constant $\delta > 0$ such that*

(a) *for any $\lambda > 0$,*

$$\sum_{n=1}^{\infty} c_n \mathbb{P} \left\{ \max_{1 \leq i \leq k_n} \left| \sum_{j=1}^i (X_{n,j} - \mathbb{E} X_{n,j} I_{\{|X_{n,j}| \leq \delta\}}) \right| > \lambda \right\} < \infty,$$

(b) $\max_{1 \leq j \leq k_n} \mathbb{P}\{|X_{n,j}| > \delta\} = o(1)$ and $\sum_{j=1}^{k_n} \mathbb{P}\{|X_{n,j}| > \delta\} = O(1)$ as $n \rightarrow \infty$,

(c) $\sum_{n=1}^{\infty} \sum_{j=1}^{k_n} c_n (1 + \alpha_n) \int_{\delta^p}^{\infty} \mathbb{P}\{|X_{n,j}|^p > t\} dt < \infty$,

(d) *for some $q > p$, $\sum_{n=1}^{\infty} c_n \beta_n \int_{\delta^p}^{\infty} \left(\sum_{j=1}^{k_n} \mathbb{P}\{|X_{n,j}|^p > t\} dt \right)^q < \infty$,*

$$\sum_{n=1}^{\infty} c_n \beta_n \left(\sum_{j=1}^{k_n} \mathbb{V} X_{n,j} I_{\{|X_{n,j}| \leq \delta\}} \right)^q < \infty \quad \text{and} \quad \sum_{n=1}^{\infty} c_n \beta_n \left(\sum_{j=1}^{k_n} \mathbb{E} |X_{n,j}| I_{\{|X_{n,j}| > \delta\}} \right)^q < \infty,$$

then for all $\varepsilon > 0$,

$$\sum_{n=1}^{\infty} c_n \mathbb{E} \left[\max_{1 \leq i \leq k_n} \left| \sum_{j=1}^i (X_{n,j} - \mathbb{E} X_{n,j} I_{\{|X_{n,j}| \leq \delta\}}) \right| - \varepsilon \right]_+^p < \infty. \quad (2.10)$$

Proof. From assumption (b), there is a positive constant C_δ such that

$$\sum_{j=1}^{k_n} \mathbb{P}\{|X_{n,j}| > \delta\} \leq C_\delta$$

for all n . Consider $X_{n,j}^I(t^{1/p}), X_{n,j}^{II}(t^{1/p}), \Gamma_n(t^{1/p}), S_{n,i}(\delta) := \sum_{j=1}^i (X_{n,j} - \mathbb{E} X_{n,j} I_{\{|X_{n,j}| \leq \delta\}})$, $i = 1, 2, \dots, k_n$ given by (2.4), fix arbitrarily $\varepsilon > 0$ and let ρ be a positive number that will be determined later. Hence,

$$\sum_{n=1}^{\infty} c_n \mathbb{E} \left(\max_{1 \leq i \leq k_n} |S_{n,i}(\delta)| - \varepsilon \right)_+^p = \sum_{n=1}^{\infty} c_n \int_0^{\infty} \mathbb{P} \left\{ \max_{1 \leq i \leq k_n} |S_{n,i}(\delta)| > \varepsilon + u^{1/p} \right\} du$$

$$\begin{aligned}
&= (2C_\delta)^p \sum_{n=1}^{\infty} c_n \int_0^\infty \mathbb{P} \left\{ \max_{1 \leq i \leq k_n} |S_{n,i}(\delta)| > \varepsilon + 2C_\delta t^{1/p} \right\} dt \\
&\leq (2C_\delta)^p \left(\frac{\delta}{\rho} \right)^p \vee \delta^p \sum_{n=1}^{\infty} c_n \mathbb{P} \left\{ \max_{1 \leq i \leq k_n} |S_{n,i}(\delta)| > \varepsilon \right\} \\
&\quad + (2C_\delta)^p \sum_{n=1}^{\infty} c_n \int_{(\delta/\rho)^p \vee \delta^p}^\infty \mathbb{P} \left\{ \max_{1 \leq i \leq k_n} |S_{n,i}(\delta)| > 2C_\delta t^{1/p} \right\} dt
\end{aligned}$$

and by assumption (a), (2.10) holds if

$$\sum_{n=1}^{\infty} c_n \int_{(\delta/\rho)^p \vee \delta^p}^\infty \mathbb{P} \left\{ \max_{1 \leq i \leq k_n} |S_{n,i}(\delta)| > 2C_\delta t^{1/p} \right\} dt < \infty.$$

Thus,

$$\begin{aligned}
&\sum_{n=1}^{\infty} c_n \int_{(\delta/\rho)^p \vee \delta^p}^\infty \mathbb{P} \left\{ \max_{1 \leq i \leq k_n} |S_{n,i}(\delta)| > 2C_\delta t^{1/p} \right\} dt \\
&= \sum_{n=1}^{\infty} c_n \int_{(\delta/\rho)^p \vee \delta^p}^\infty \mathbb{P} \left[\left\{ \max_{1 \leq i \leq k_n} |S_{n,i}(\delta)| > 2C_\delta t^{1/p} \right\} \cap \Gamma_n(t^{1/p}) \right] dt \\
&\quad + \sum_{n=1}^{\infty} c_n \int_{(\delta/\rho)^p \vee \delta^p}^\infty \mathbb{P} \left[\left\{ \max_{1 \leq i \leq k_n} |S_{n,i}(\delta)| > 2C_\delta t^{1/p} \right\} \cap \Gamma_n(t^{1/p})^c \right] dt \\
&\leq \sum_{n=1}^{\infty} c_n \int_{(\delta/\rho)^p \vee \delta^p}^\infty \mathbb{P} \left\{ \max_{1 \leq i \leq k_n} \left| \sum_{j=1}^i \left(X_{n,j} I_{\{|X_{n,j}| \leq t^{1/p}\}} - \mathbb{E} X_{n,j} I_{\{|X_{n,j}| \leq \delta\}} \right) \right| > 2C_\delta t^{1/p} \right\} dt \\
&\quad + \sum_{n=1}^{\infty} c_n \int_{\delta^p}^\infty \sum_{j=1}^{k_n} \mathbb{P} \{ |X_{n,j}| > t^{1/p} \} dt.
\end{aligned}$$

According to (c), it suffices to prove

$$\sum_{n=1}^{\infty} c_n \int_{(\delta/\rho)^p \vee \delta^p}^\infty \mathbb{P} \left\{ \max_{1 \leq i \leq k_n} \left| \sum_{j=1}^i \left(X_{n,j} I_{\{|X_{n,j}| \leq t^{1/p}\}} - \mathbb{E} X_{n,j} I_{\{|X_{n,j}| \leq \delta\}} \right) \right| > 2C_\delta t^{1/p} \right\} dt < \infty. \tag{2.11}$$

Since

$$\begin{aligned}
\max_{1 \leq i \leq k_n} \left| \sum_{j=1}^i \mathbb{E} X_{n,j} I_{\{\delta < |X_{n,j}| \leq t^{1/p}\}} \right| &\leq \sum_{j=1}^{k_n} \mathbb{E} |X_{n,j}| I_{\{\delta < |X_{n,j}| \leq t^{1/p}\}} \\
&\leq t^{1/p} \sum_{j=1}^{k_n} \mathbb{P} \{ |X_{n,j}| > \delta \} \\
&\leq C_\delta t^{1/p}
\end{aligned}$$

and

$$\mathbb{P} \left\{ \max_{1 \leq i \leq k_n} \left| \sum_{j=1}^i \left[\mathbb{E} X_{n,j}''(t^{1/p}) - X_{n,j}''(t^{1/p}) \right] \right| > \frac{C_\delta t^{1/p}}{2} \right\}$$

$$\begin{aligned}
&\leq \frac{2}{C_\delta t^{1/p}} \mathbb{E} \left\{ \max_{1 \leq i \leq k_n} \left| \sum_{j=1}^i \left[\mathbb{E} X_{n,j}''(t^{1/p}) - X_{n,j}''(t^{1/p}) \right] \right| \right\} \\
&\leq \frac{4}{C_\delta t^{1/p}} \sum_{j=1}^{k_n} \mathbb{E} |X_{n,j}''(t^{1/p})| \\
&= \frac{4}{C_\delta} \sum_{j=1}^{k_n} \mathbb{P}\{|X_{n,j}| > t^{1/p}\}
\end{aligned}$$

we obtain, for each $t \geq (\delta/\rho)^p \vee \delta^p$,

$$\begin{aligned}
&\mathbb{P} \left\{ \max_{1 \leq i \leq k_n} \left| \sum_{j=1}^i \left(X_{n,j} I_{\{|X_{n,j}| \leq t^{1/p}\}} - \mathbb{E} X_{n,j} I_{\{|X_{n,j}| \leq \delta\}} \right) \right| > 2C_\delta t^{1/p} \right\} \\
&= \mathbb{P} \left\{ \max_{1 \leq i \leq k_n} \left| \sum_{j=1}^i \left[X'_{n,j}(t^{1/p}) - \mathbb{E} X'_{n,j}(t^{1/p}) + \mathbb{E} X_{n,j} I_{\{\delta < |X_{n,j}| \leq t^{1/p}\}} \right] \right| > 2C_\delta t^{1/p} \right\} \\
&\leq \mathbb{P} \left\{ \max_{1 \leq i \leq k_n} \left| \sum_{j=1}^i \left[X'_{n,j}(t^{1/p}) - \mathbb{E} X'_{n,j}(t^{1/p}) \right] \right| + \max_{1 \leq i \leq k_n} \left| \sum_{j=1}^i \mathbb{E} X_{n,j} I_{\{\delta < |X_{n,j}| \leq t^{1/p}\}} \right| > 2C_\delta t^{1/p} \right\} \\
&\leq \mathbb{P} \left\{ \max_{1 \leq i \leq k_n} \left| \sum_{j=1}^i \left[X'_{n,j}(t^{1/p}) - \mathbb{E} X'_{n,j}(t^{1/p}) \right] \right| > C_\delta t^{1/p} \right\} \\
&\leq \mathbb{P} \left\{ \max_{1 \leq i \leq k_n} \left| \sum_{j=1}^i \left[g_{t^{1/p}}(X_{n,j}) - \mathbb{E} g_{t^{1/p}}(X_{n,j}) \right] \right| > \frac{C_\delta t^{1/p}}{2} \right\} \\
&\quad + \mathbb{P} \left\{ \max_{1 \leq i \leq k_n} \left| \sum_{j=1}^i \left[\mathbb{E} X_{n,j}''(t^{1/p}) - X_{n,j}''(t^{1/p}) \right] \right| > \frac{C_\delta t^{1/p}}{2} \right\} \\
&\leq \mathbb{P} \left\{ \max_{1 \leq i \leq k_n} \left| \sum_{j=1}^i \left[g_{t^{1/p}}(X_{n,j}) - \mathbb{E} g_{t^{1/p}}(X_{n,j}) \right] \right| > \frac{C_\delta t^{1/p}}{2} \right\} + \frac{4}{C_\delta} \sum_{j=1}^{k_n} \mathbb{P}\{|X_{n,j}| > t^{1/p}\}
\end{aligned}$$

Thus, (2.11) holds if

$$\sum_{n=1}^{\infty} c_n \int_{(\delta/\rho)^p \vee \delta^p}^{\infty} \mathbb{P} \left\{ \max_{1 \leq i \leq k_n} \left| \sum_{j=1}^i \left[g_{t^{1/p}}(X_{n,j}) - \mathbb{E} g_{t^{1/p}}(X_{n,j}) \right] \right| > \frac{C_\delta t^{1/p}}{2} \right\} < \infty \quad (2.12)$$

and

$$\sum_{n=1}^{\infty} c_n \int_{(\delta/\rho)^p \vee \delta^p}^{\infty} \sum_{j=1}^{k_n} \mathbb{P}\{|X_{n,j}| > t^{1/p}\} < \infty. \quad (2.13)$$

Since (2.13) follows from assumption (c), it remains to show (2.12). By assumption (b), there is $n_0 = n_0(\delta, \rho)$ such that

$$\max_{1 \leq j \leq k_n} \mathbb{P}\{|X_{n,j}| > \delta\} < \frac{\rho}{2}, \quad \forall n \geq n_0 \quad (2.14)$$

and whence, for any $t \geq (\delta/\rho)^p \vee \delta^p$ and $n \geq n_0$, we have

$$\begin{aligned}
& \max_{1 \leq j \leq k_n} |\mathbb{E}g_{t^{1/p}}(X_{n,j})| \\
& \leq \max_{1 \leq j \leq k_n} \left[\mathbb{E}|X_{n,j}|I_{\{|X_{n,j}| \leq t^{1/p}\}} + t^{1/p}\mathbb{P}\{|X_{n,j}| > t^{1/p}\} \right] \\
& \leq \max_{1 \leq j \leq k_n} \left[\mathbb{E}|X_{n,j}|I_{\{|X_{n,j}| \leq \delta\}} + \mathbb{E}|X_{n,j}|I_{\{\delta < |X_{n,j}| \leq t^{1/p}\}} + t^{1/p}\mathbb{P}\{|X_{n,j}| > t^{1/p}\} \right] \\
& \leq \delta + \max_{1 \leq j \leq k_n} \left[t^{1/p}\mathbb{P}\{|X_{n,j}| > \delta\} + t^{1/p}\mathbb{P}\{|X_{n,j}| > t^{1/p}\} \right] \\
& \leq \rho t^{1/p} + \max_{1 \leq j \leq k_n} \left[2t^{1/p}\mathbb{P}\{|X_{n,j}| > \delta\} \right] \\
& < 2\rho t^{1/p}.
\end{aligned} \tag{2.15}$$

By taking $\lambda = C_\delta t^{1/p}/2$ and $\eta = C_\delta t^{1/p}/(2q)$, there are $C_1, C_2 > 0$ such that (2.1) holds with the sequences $\{\alpha_n\}, \{\beta_n\}$. Putting $\rho = C_1 C_\delta/(6q)$, estimates (2.15) and

$$\begin{aligned}
s_n(t^{1/p}) &= \sum_{j=1}^{k_n} \mathbb{E}[g_{t^{1/p}}(X_{n,j}) - \mathbb{E}g_{t^{1/p}}(X_{n,j})]^2 \\
&= \sum_{j=1}^{k_n} \mathbb{V}[g_{t^{1/p}}(X_{n,j})] \\
&\leq 2 \sum_{j=1}^{k_n} \mathbb{V}X'_{n,j}(t^{1/p}) + 2 \sum_{j=1}^{k_n} \mathbb{V}X''_{n,j}(t^{1/p}) \\
&\leq 2 \sum_{j=1}^{k_n} \mathbb{V} \left[X'_{n,j}(\delta) + X_{n,j}I_{\{\delta < |X_{n,j}| \leq t^{1/p}\}} \right] + 2 \sum_{j=1}^{k_n} \mathbb{E} \left[X''_{n,j}(t^{1/p}) \right]^2 \\
&\leq 4 \sum_{j=1}^{k_n} \mathbb{V}X'_{n,j}(\delta) + 4 \sum_{j=1}^{k_n} \mathbb{V}X_{n,j}I_{\{\delta < |X_{n,j}| \leq t^{1/p}\}} + 2t^{2/p} \sum_{j=1}^{k_n} \mathbb{P}\{|X_{n,j}| > t^{1/p}\} \\
&\leq 4 \sum_{j=1}^{k_n} \mathbb{V}X_{n,j}I_{\{|X_{n,j}| \leq \delta\}} + 4 \sum_{j=1}^{k_n} \mathbb{E}X_{n,j}^2 I_{\{\delta < |X_{n,j}| \leq t^{1/p}\}} + 2t^{2/p} \sum_{j=1}^{k_n} \mathbb{P}\{|X_{n,j}| > t^{1/p}\} \\
&\leq 4 \sum_{j=1}^{k_n} \mathbb{V}X_{n,j}I_{\{|X_{n,j}| \leq \delta\}} + 4t^{1/p} \sum_{j=1}^{k_n} \mathbb{E}|X_{n,j}|I_{\{|X_{n,j}| > \delta\}} + 2t^{2/p} \sum_{j=1}^{k_n} \mathbb{P}\{|X_{n,j}| > t^{1/p}\}
\end{aligned}$$

for all $t \geq (\delta/\rho)^p \vee \delta^p$, lead to

$$\begin{aligned}
& \mathbb{P} \left\{ \max_{1 \leq i \leq k_n} \left| \sum_{j=1}^i [g_{t^{1/p}}(X_{n,j}) - \mathbb{E}g_{t^{1/p}}(X_{n,j})] \right| > \frac{C_\delta t^{1/p}}{2} \right\} \\
& \leq \alpha_n \mathbb{P} \left\{ \max_{1 \leq j \leq k_n} |g_{t^{1/p}}(X_{n,j}) - \mathbb{E}g_{t^{1/p}}(X_{n,j})| > 3\rho t^{1/p} \right\} + \frac{\beta_n 2^{2q} q^q}{C_\delta^{2q} C_2^q} \left[\frac{s_n(t^{1/p})}{t^{2/p}} \right]^q \\
& \leq \alpha_n \mathbb{P} \left\{ \max_{1 \leq j \leq k_n} |g_{t^{1/p}}(X_{n,j})| > \rho t^{1/p} \right\} + \frac{\beta_n 2^{3q-1} q^q}{C_\delta^{2q} C_2^q} \left(\frac{4}{t^{2/p}} \sum_{j=1}^{k_n} \mathbb{V}X_{n,j}I_{\{|X_{n,j}| \leq \delta\}} \right)^q
\end{aligned}$$

$$\begin{aligned}
& + \frac{\beta_n 2^{3q-1} q^q}{C_\delta^{2q} C_2^q} \left(\frac{4}{t^{1/p}} \sum_{j=1}^{k_n} \mathbb{E}|X_{n,j}| I_{\{|X_{n,j}|>\delta\}} + 2 \sum_{j=1}^{k_n} \mathbb{P}\{|X_{n,j}| > t^{1/p}\} \right)^q \\
& \leq \alpha_n \sum_{j=1}^{k_n} \mathbb{P}\{|g_{t^{1/p}}(X_{n,j})| > \rho t^{1/p}\} + \frac{\beta_n 2^{5q-1} q^q}{C_\delta^{2q} C_2^q t^{2q/p}} \left(\sum_{j=1}^{k_n} \mathbb{V}X_{n,j} I_{\{|X_{n,j}| \leq \delta\}} \right)^q \\
& + \frac{\beta_n 2^{4q-2} q^q}{C_\delta^{2q} C_2^q} \left(\frac{4}{t^{1/p}} \sum_{j=1}^{k_n} \mathbb{E}|X_{n,j}| I_{\{|X_{n,j}|>\delta\}} \right)^q + \frac{\beta_n 2^{4q-2} q^q}{C_\delta^{2q} C_2^q} \left(2 \sum_{j=1}^{k_n} \mathbb{P}\{|X_{n,j}| > t^{1/p}\} \right)^q \\
& \leq \alpha_n \sum_{j=1}^{k_n} \mathbb{P}\{|X_{n,j}| > \rho t^{1/p}\} + \frac{\beta_n 2^{5q-1} q^q}{C_\delta^{2q} C_2^q t^{2q/p}} \left(\sum_{j=1}^{k_n} \mathbb{V}X_{n,j} I_{\{|X_{n,j}| \leq \delta\}} \right)^q \\
& + \frac{\beta_n 2^{6q-2} q^q}{C_\delta^{2q} C_2^q t^{q/p}} \left(\sum_{j=1}^{k_n} \mathbb{E}|X_{n,j}| I_{\{|X_{n,j}|>\delta\}} \right)^q + \frac{\beta_n 2^{5q-2} q^q}{C_\delta^{2q} C_2^q} \left(\sum_{j=1}^{k_n} \mathbb{P}\{|X_{n,j}| > t^{1/p}\} \right)^q
\end{aligned}$$

for every $t \geq (\delta/\rho)^p \vee \delta^p$ and n large enough. Thereby,

$$\begin{aligned}
& \sum_{n=1}^{\infty} c_n \int_{(\delta/\rho)^p \vee \delta^p}^{\infty} \mathbb{P} \left\{ \max_{1 \leq i \leq k_n} \left| \sum_{j=1}^i [g_{t^{1/p}}(X_{n,j}) - \mathbb{E}g_{t^{1/p}}(X_{n,j})] \right| > \frac{C_\delta t^{1/p}}{2} \right\} dt \\
& \leq \sum_{n=1}^{\infty} c_n \alpha_n \int_{(\delta/\rho)^p}^{\infty} \sum_{j=1}^{k_n} \mathbb{P}\{|X_{n,j}| > \rho t^{1/p}\} dt + \frac{2^{5q-1} q^q}{C_\delta^{2q} C_2^q} \sum_{n=1}^{\infty} c_n \beta_n \int_{\delta^p}^{\infty} \frac{dt}{t^{2q/p}} \left(\sum_{j=1}^{k_n} \mathbb{V}X_{n,j} I_{\{|X_{n,j}| \leq \delta\}} \right)^q \\
& + \frac{2^{6q-2} q^q}{C_\delta^{2q} C_2^q} \sum_{n=1}^{\infty} c_n \beta_n \int_{\delta^p}^{\infty} \frac{dt}{t^{q/p}} \left(\sum_{j=1}^{k_n} \mathbb{E}|X_{n,j}| I_{\{|X_{n,j}|>\delta\}} \right)^q \\
& + \frac{2^{5q-2} q^q}{C_\delta^{2q} C_2^q} \sum_{n=1}^{\infty} c_n \beta_n \int_{\delta^p}^{\infty} \left(\sum_{j=1}^{k_n} \mathbb{P}\{|X_{n,j}|^p > t\} dt \right)^q \\
& \leq \rho^{-p} \sum_{n=1}^{\infty} c_n \alpha_n \int_{\delta^p}^{\infty} \sum_{j=1}^{k_n} \mathbb{P}\{|X_{n,j}|^p > u\} du + \frac{p 2^{5q-2} q^q \delta^{p-2q}}{(2q-p) C_\delta^{2q} C_2^q} \sum_{n=1}^{\infty} c_n \beta_n \left(\sum_{j=1}^{k_n} \mathbb{V}X_{n,j} I_{\{|X_{n,j}| \leq \delta\}} \right)^q \\
& + \frac{p 2^{6q-2} q^q \delta^{p-q}}{(q-p) C_\delta^{2q} C_2^q} \sum_{n=1}^{\infty} c_n \beta_n \left(\sum_{j=1}^{k_n} \mathbb{E}|X_{n,j}| I_{\{|X_{n,j}|>\delta\}} \right)^q \\
& + \frac{2^{5q-2} q^q}{C_\delta^{2q} C_2^q} \sum_{n=1}^{\infty} c_n \beta_n \int_{\delta^p}^{\infty} \left(\sum_{j=1}^{k_n} \mathbb{P}\{|X_{n,j}|^p > t\} dt \right)^q \\
& < \infty,
\end{aligned}$$

and the proof is concluded. \square

By means of Theorems 2 and 3, we state the following corollary.

Corollary 1 Let $p \geq 1$, $\{c_n\}$ be a sequence of positive numbers, and $\{X_{n,j}, 1 \leq j \leq k_n, n \geq 1\}$ be an array of random variables satisfying (2.1) for sequences $\{\alpha_n\}, \{\beta_n\}$ of nonnegative numbers such that $\alpha_n = O(1) = \beta_n$ as $n \rightarrow \infty$. If assumption (i') holds, and there exists a constant $\delta > 0$ such that

$$(b') \sum_{j=1}^{k_n} \mathbb{P}\{|X_{n,j}| > \delta\} = o(1) \text{ as } n \rightarrow \infty,$$

$$(c') \sum_{n=1}^{\infty} \sum_{j=1}^{k_n} c_n \int_{\delta^p}^{\infty} \mathbb{P}\{|X_{n,j}|^p > t\} dt < \infty,$$

(d') for some $q > p$,

$$\sum_{n=1}^{\infty} c_n \left(\sum_{j=1}^{k_n} \mathbb{V} X_{n,j} I_{\{|X_{n,j}| \leq \delta\}} \right)^q < \infty \quad \text{and} \quad \sum_{n=1}^{\infty} c_n \left(\sum_{j=1}^{k_n} \mathbb{E}|X_{n,j}| I_{\{|X_{n,j}| > \delta\}} \right)^q < \infty,$$

then, for all $\varepsilon > 0$, (2.10) holds.

Proof. Since (b') ensures $\sum_{j=1}^{k_n} \mathbb{P}\{|X_{n,j}|^p > t\} \leq \sum_{j=1}^{k_n} \mathbb{P}\{|X_{n,j}| > \delta\} \leq 1$ for any $t \geq \delta^p$ and n large enough, assumption (c') entails

$$\sum_{n=1}^{\infty} c_n \beta_n \int_{\delta^p}^{\infty} \left(\sum_{j=1}^{k_n} \mathbb{P}\{|X_{n,j}|^p > t\} dt \right)^q \leq C \sum_{n=1}^{\infty} c_n \int_{\delta^p}^{\infty} \sum_{j=1}^{k_n} \mathbb{P}\{|X_{n,j}|^p > t\} dt < \infty$$

where C is some positive constant (non-depending on n) such that $\beta_n \leq C$ for all n . The thesis is a consequence of Theorems 2 and 3 by noting that (b') implies (b). \square

It is worthy to note that assumption (i) of Theorem 3.1 in [16] implies both conditions (i') and (c') of Corollary 1 when $\alpha_n = 2$ and $\beta_n = 8$. Indeed, the example below shows that $\int_{\delta^p}^{\infty} \mathbb{P}\{|X_{n,j}|^p > t\} dt$ can be of smaller order than $\mathbb{E}|X_{n,j}|^p I_{\{|X_{n,j}| > \delta\}}$ on the one hand, and on the other hand, $\mathbb{P}\{|X_{n,j}| > \varepsilon\}$ can also be of smaller order than $\mathbb{E}|X_{n,j}|^p I_{\{|X_{n,j}| > \varepsilon\}}$.

EXAMPLE Let $\{\xi_n, n \geq 1\}$ be a sequence of i.i.d. random variables and $X_{n,j} := \xi_j/b_n$ where $\{b_n\}$ is a sequence of positive constants satisfying $b_n \rightarrow \infty$ as $n \rightarrow \infty$. If the tail distribution of $|\xi_1|^p$ is rapidly varying with index $-\infty$ (e.g. $|\xi_1|^p$ having exponential distribution) and $\delta > 0$, then

$$\frac{\int_{\delta^p}^{\infty} \mathbb{P}\{|X_{n,j}|^p > t\} dt}{\mathbb{E}|X_{n,j}|^p I_{\{|X_{n,j}| > \delta\}}} \leq \frac{\int_{(\delta b_n)^p}^{\infty} \mathbb{P}\{|\xi_1|^p > u\} du}{(\delta b_n)^p \mathbb{P}\{|\xi_1| > \delta b_n\}} = o(1), \quad n \rightarrow \infty$$

(see [6], page 570), i.e. $\int_{\delta^p}^{\infty} \mathbb{P}\{|X_{n,j}|^p > t\} dt = o\left(\mathbb{E}|X_{n,j}|^p I_{\{|X_{n,j}| > \delta\}}\right)$, $n \rightarrow \infty$. Instead, if $\mathbb{P}\{|\xi_1|^p > t\} \sim L(t)/t$ as $t \rightarrow \infty$, where L is a slowly varying function at ∞ locally bounded in $[t_0, \infty)$, for some $t_0 \geq 0$, and such that $\int_{t_0}^{\infty} L(t)/t dt < \infty$, then, for each fixed $\varepsilon > 0$,

$$\frac{\int_{\varepsilon^p}^{\infty} \mathbb{P}\{|X_{n,j}|^p > t\} dt}{\varepsilon^p \mathbb{P}\{|X_{n,j}| > \varepsilon\}} = \frac{\int_{(\varepsilon b_n)^p}^{\infty} \mathbb{P}\{|\xi_1|^p > u\} du}{(\varepsilon b_n)^p \mathbb{P}\{|\xi_1| > \varepsilon b_n\}} \rightarrow \infty, \quad n \rightarrow \infty$$

(see [6], page 567) and whence $\mathbb{P}\{|X_{n,j}| > \varepsilon\} = o\left(\mathbb{E}|X_{n,j}|^p I_{\{|X_{n,j}| > \varepsilon\}}\right)$, $n \rightarrow \infty$.

3 Change-point estimator

In the nineties, the problem of a change in the mean of a sequence of observations received attention by many authors (see [1], [2], [5], [12] among others). Let $\{\mu_{n,k}, 1 \leq k \leq n, n \geq 1\}$ be a triangular array of real numbers and $\{X_{n,k}, 1 \leq k \leq n, n \geq 1\}$ be a triangular array of zero-mean random variables. Consider the model

$$Y_{n,k} = \begin{cases} \mu_{n,k} + X_{n,k}, & 1 \leq k \leq k^* \\ \mu_{n,k} + \Delta_n + X_{n,k}, & k^* < k \leq n \end{cases} \quad (3.1)$$

where $k^* = [n\tau^*]$ is an unknown change-point, Δ_n is the (unknown) change-amount, and $\tau^* \in (0, 1)$ is fixed (here, $[x]$ stands for the integer part of x). In [12], the authors proved the (weak) consistency of CUSUM-type estimator of the point of shift in the mean of a sequence of observations, i.e. by considering the estimators

$$\hat{k}_n := \min \left\{ k : |U_{n,k}| = \max_{1 \leq j < n} |U_{n,j}| \right\}$$

and $\hat{\tau}_n := \hat{k}_n/n$ of k^* and τ^* , respectively, with

$$U_{n,k} := \frac{(n-k)^{1-\gamma}}{n^{1-\gamma}k^\gamma} \sum_{j=1}^k Y_{n,j} - \frac{k^{1-\gamma}}{n^{1-\gamma}(n-k)^\gamma} \sum_{j=k+1}^n Y_{n,j} \quad (0 \leq \gamma < 1)$$

conditions were given for $\hat{\tau}_n$ to be weakly consistent of τ^* . The next statement gives us the complete convergence, in the sense of Hsu and Robbins [7], of $\hat{\tau}_n$, i.e. we provide conditions under which $\hat{\tau}_n$ is *completely consistent* for τ^* . We assume that $C(\cdot)$ denotes a positive generic constant depending (only) on the arguments specified within the parentheses that may assume different values at each appearance.

Theorem 4 *Suppose that in model (3.1), $\Delta_n \neq 0$ for any $n \geq 1$, and $\{X_{n,j}, 1 \leq j \leq n, n \geq 1\}$ is an array of zero-mean random variables satisfying (2.1) for sequences $\{\alpha_n\}, \{\beta_n\}$ of non-negative numbers such that $\alpha_n = O(1) = \beta_n$ as $n \rightarrow \infty$. If either (a) $\sup_{n,j} \mathbb{E}|X_{n,j}|^r < \infty$ for $1 < r \leq 2$ verifying*

$$\begin{cases} \sum_{n=1}^{\infty} |\Delta_n|^{-r} n^{1-r}, & 0 \leq \gamma < 1/r \\ \sum_{n=1}^{\infty} |\Delta_n|^{-r} n^{1-r} \log n, & \gamma = 1/r \\ \sum_{n=1}^{\infty} |\Delta_n|^{-r} n^{r(\gamma-1)}, & 1/r < \gamma < 1, \end{cases} \quad (3.2)$$

or (b) $\sup_{n,j} \mathbb{E}|X_{n,j}|^r < \infty$ for some $r > 2$ satisfying

$$\begin{cases} \sum_{n=1}^{\infty} [|\Delta_n|^{-r} + |\Delta_n|^{2(1-r)}] n^{1-r} < \infty, & 0 \leq \gamma < 1/r \\ \sum_{n=1}^{\infty} [|\Delta_n|^{-r} \log n + |\Delta_n|^{2(1-r)}] n^{1-r} < \infty, & \gamma = 1/r \\ \sum_{n=1}^{\infty} [|\Delta_n|^{-r} + |\Delta_n|^{2r(\gamma-1)}] n^{r(\gamma-1)} < \infty, & 1/r < \gamma < 1/2 \\ \sum_{n=1}^{\infty} [|\Delta_n|^{-r} n^{r/2-1} + |\Delta_n|^{2(1-r)} \log^{r-1} n] n^{1-r} < \infty, & \gamma = 1/2 \\ \sum_{n=1}^{\infty} |\Delta_n|^{-r} n^{r(\gamma-1)} < \infty, & 1/2 < \gamma < 1, \end{cases} \quad (3.3)$$

then $\hat{\tau}_n$ converges completely to τ^* . Furthermore, for all $\varepsilon > 0$, $\sum_{n=1}^{\infty} \mathbb{E}(|\hat{\tau}_n - \tau^*| - \varepsilon)_+^r < \infty$.

Proof. As in proof of Theorem 1.1 of [12] (see [12], page 391), it can be shown that

$$|\Delta_n|(1-\gamma)(\tau^*)^{-\gamma}(1-\tau^*)\min\{\tau^*, 1-\tau^*\}|\tau^* - \hat{\tau}_n| \leq 2n^{\gamma-1} \max_{1 \leq k < n} |U_{n,k} - \mathbb{E}U_{n,k}|,$$

whence

$$|\tau^* - \hat{\tau}_n| \leq C(\gamma, \tau^*) \frac{n^{\gamma-1}}{|\Delta_n|} \max_{1 \leq k < n} |U_{n,k} - \mathbb{E}U_{n,k}|. \quad (3.4)$$

By employing Abel's lemma (see [4], page 114), we have

$$\begin{aligned} & \max_{1 \leq k < n} |U_{n,k} - \mathbb{E}U_{n,k}| \\ & \leq \max_{1 \leq k < n} \frac{1}{k^\gamma} \left| \sum_{j=1}^k (Y_{n,j} - \mathbb{E}Y_{n,j}) \right| + \max_{1 \leq k < n} \frac{1}{(n-k)^\gamma} \left| \sum_{j=k+1}^n (Y_{n,j} - \mathbb{E}Y_{n,j}) \right| \\ & = \max_{1 \leq k < n} \frac{1}{k^\gamma} \left| \sum_{j=1}^k (Y_{n,j} - \mathbb{E}Y_{n,j}) \right| + \max_{1 \leq k < n} \frac{1}{k^\gamma} \left| \sum_{j=1}^k (Y_{n,n-j+1} - \mathbb{E}Y_{n,n-j+1}) \right| \\ & = \max_{1 \leq k < n} \frac{1}{k^\gamma} \left| \sum_{j=1}^k X_{n,j} \right| + \max_{1 \leq k < n} \frac{1}{k^\gamma} \left| \sum_{j=1}^k X_{n,n-j+1} \right| \\ & \leq 2 \max_{1 \leq k < n} \left| \sum_{j=1}^k \frac{X_{n,j}}{j^\gamma} \right| + 2 \max_{1 \leq k < n} \left| \sum_{j=1}^k \frac{X_{n,n-j+1}}{j^\gamma} \right| \end{aligned} \quad (3.5)$$

and it suffices to prove that, for any $\varepsilon > 0$,

$$\sum_{n=2}^{\infty} \mathbb{P} \left\{ \max_{1 \leq k < n} \left| \sum_{j=1}^k \frac{n^{\gamma-1} X_{n,j}}{|\Delta_n| j^\gamma} \right| > \varepsilon \right\} < \infty, \quad (3.6)$$

$$\sum_{n=2}^{\infty} \mathbb{P} \left\{ \max_{1 \leq k < n} \left| \sum_{j=1}^k \frac{n^{\gamma-1} X_{n,n-j+1}}{|\Delta_n| j^\gamma} \right| > \varepsilon \right\} < \infty. \quad (3.7)$$

For (3.6) notice that $\mathbb{E}X_{n,k} = 0$ and for each $s > 0$,

$$\sum_{j=1}^n j^{-s} \leq \begin{cases} C(s)n^{1-s}, & s < 1 \\ C(s) \log n, & s = 1 \\ C(s), & s > 1. \end{cases} \quad (3.8)$$

Thus,

$$\begin{aligned} \max_{1 \leq k < n} \left| \sum_{j=1}^k \mathbb{E} \left(\frac{n^{\gamma-1} X_{n,j}}{|\Delta_n| j^\gamma} I_{\left\{ \frac{n^{\gamma-1} |X_{n,j}|}{|\Delta_n| j^\gamma} \leq \delta \right\}} \right) \right| &= \max_{1 \leq k < n} \left| \sum_{j=1}^k \mathbb{E} \left(\frac{n^{\gamma-1} X_{n,j}}{|\Delta_n| j^\gamma} I_{\left\{ \frac{n^{\gamma-1} |X_{n,j}|}{|\Delta_n| j^\gamma} > \delta \right\}} \right) \right| \\ &\leq \sum_{j=1}^{n-1} \frac{n^{\gamma-1}}{|\Delta_n| j^\gamma} \mathbb{E} |X_{n,j}| I_{\left\{ \frac{n^{\gamma-1} |X_{n,j}|}{|\Delta_n| j^\gamma} > \delta \right\}} \end{aligned}$$

$$\begin{aligned}
&\leq \delta^{1-r} \sum_{j=1}^{n-1} \frac{n^{r(\gamma-1)}}{|\Delta_n|^r j^{r\gamma}} \mathbb{E}|X_{n,j}|^r \\
&\leq \delta^{1-r} \frac{n^{r(\gamma-1)}}{|\Delta_n|^r} \left(\sup_{n,j} \mathbb{E}|X_{n,j}|^r \right) \sum_{j=1}^{n-1} j^{-r\gamma} \\
&\leq \begin{cases} C(r, \delta, \gamma) |\Delta_n|^{-r} n^{1-r}, & 0 \leq \gamma < 1/r \\ C(r, \delta) |\Delta_n|^{-r} n^{1-r} \log n, & \gamma = 1/r \\ C(r, \delta, \gamma) |\Delta_n|^{-r} n^{r(\gamma-1)}, & 1/r < \gamma < 1 \end{cases}
\end{aligned}$$

and (3.2)-(3.3) imply

$$\max_{1 \leq k < n} \left| \sum_{j=1}^k \mathbb{E} \left(\frac{n^{\gamma-1} X_{n,j}}{|\Delta_n| j^\gamma} I_{\left\{ \frac{n^{\gamma-1} |X_{n,j}|}{|\Delta_n| j^\gamma} \leq \delta \right\}} \right) \right| = o(1), \quad n \rightarrow \infty. \quad (3.9)$$

According to Markov inequality and (3.8), we get

$$\begin{aligned}
\sum_{j=1}^{n-1} \mathbb{P}\{|X_{n,j}| > \varepsilon n^{1-\gamma} j^\gamma |\Delta_n|\} &\leq \sum_{j=1}^{n-1} \frac{n^{r(\gamma-1)} \mathbb{E}|X_{n,j}|^r}{\varepsilon^r |\Delta_n|^r j^{r\gamma}} \\
&\leq \frac{n^{r(\gamma-1)}}{\varepsilon^r |\Delta_n|^r} \sup_{n,j} \mathbb{E}|X_{n,j}|^r \sum_{j=1}^{n-1} j^{-r\gamma} \\
&\leq \begin{cases} C(r, \varepsilon, \gamma) |\Delta_n|^{-r} n^{1-r}, & 0 \leq \gamma < 1/r \\ C(r, \varepsilon) |\Delta_n|^{-r} n^{1-r} \log n, & \gamma = 1/r \\ C(r, \varepsilon, \gamma) |\Delta_n|^{-r} n^{r(\gamma-1)}, & 1/r < \gamma < 1 \end{cases}
\end{aligned} \quad (3.10)$$

and (3.2)-(3.3) ensure assumption (i') in Theorem 2. Moreover,

$$\begin{aligned}
\left[\sum_{j=1}^{n-1} \mathbb{V} \left(\frac{n^{\gamma-1} X_{n,j}}{|\Delta_n| j^\gamma} I_{\left\{ \frac{n^{\gamma-1} |X_{n,j}|}{|\Delta_n| j^\gamma} \leq \delta \right\}} \right) \right]^q &\leq \left[\sum_{j=1}^{n-1} \frac{n^{2(\gamma-1)}}{\Delta_n^2 j^{2\gamma}} \mathbb{E} X_{n,j}^2 I_{\left\{ \frac{n^{\gamma-1} |X_{n,j}|}{|\Delta_n| j^\gamma} \leq \delta \right\}} \right]^q \\
&\leq \delta^{q(2-r)} \left[\sum_{j=1}^{n-1} \frac{n^{r(\gamma-1)}}{|\Delta_n|^r j^{r\gamma}} \mathbb{E}|X_{n,j}|^r \right]^q \\
&\leq C(r, \delta, q) \frac{n^{qr(\gamma-1)}}{|\Delta_n|^{qr}} \left(\sup_{n,j} \mathbb{E}|X_{n,j}|^r \right)^q \left(\sum_{j=1}^{n-1} j^{-r\gamma} \right)^q \\
&\leq \begin{cases} C(r, q, \delta, \gamma) |\Delta_n|^{-qr} n^{q(1-r)}, & 0 \leq \gamma < 1/r \\ C(r, q, \delta) |\Delta_n|^{-qr} n^{q(1-r)} \log^q n, & \gamma = 1/r \\ C(r, q, \delta, \gamma) |\Delta_n|^{-qr} n^{qr(\gamma-1)}, & 1/r < \gamma < 1 \end{cases}
\end{aligned}$$

when $r \leq 2$, and

$$\left[\sum_{j=1}^{n-1} \mathbb{V} \left(\frac{n^{\gamma-1} X_{n,j}}{|\Delta_n| j^\gamma} I_{\left\{ \frac{n^{\gamma-1} |X_{n,j}|}{|\Delta_n| j^\gamma} \leq \delta \right\}} \right) \right]^q \leq \left[\sum_{j=1}^{n-1} \frac{n^{2(\gamma-1)}}{\Delta_n^2 j^{2\gamma}} \mathbb{E} X_{n,j}^2 I_{\left\{ \frac{n^{\gamma-1} |X_{n,j}|}{|\Delta_n| j^\gamma} \leq \delta \right\}} \right]^q$$

$$\begin{aligned}
&\leq \left(\sup_{n,j} \mathbb{E}|X_{n,j}|^r \right)^{2q/r} \frac{n^{2q(\gamma-1)}}{|\Delta_n|^{2q}} \left(\sum_{j=1}^{n-1} j^{-2\gamma} \right)^q \\
&\leq \begin{cases} C(q, r, \gamma) |\Delta_n|^{-2q} n^{-q}, & 0 \leq \gamma < 1/2 \\ C(q, r) |\Delta_n|^{-2q} n^{-q} \log^q n, & \gamma = 1/2 \\ C(q, r, \gamma) |\Delta_n|^{-2q} n^{2q(\gamma-1)}, & 1/2 < \gamma < 1 \end{cases}
\end{aligned}$$

whenever $r > 2$. By taking

$$q = \begin{cases} r(1 - \gamma), & 1/r < \gamma < 1/2 \\ r/2, & 1/2 < \gamma < 1 \\ r - 1, & \text{otherwise} \end{cases}$$

in the latter inequality and any $q \geq 1$ in the former inequality, it follows

$$\sum_{n=2}^{\infty} \left[\sum_{j=1}^{n-1} \mathbb{V} \left(\frac{n^{\gamma-1} X_{n,j}}{|\Delta_n| j^\gamma} I_{\left\{ \frac{n^{\gamma-1} |X_{n,j}|}{|\Delta_n| j^\gamma} \leq \delta \right\}} \right) \right]^q < \infty \quad (3.11)$$

via (3.2)-(3.3). Hence, (3.6) holds from Theorem 2 with $c_n = 1$ and (3.9). Similarly, it can be shown (3.7); the details are omitted.

For all reals numbers x, y and $p, \varepsilon > 0$, we have

$$(|x + y| - \varepsilon)_+^p \leq \max\{1, 2^{p-1}\} \left[\left(|x| - \frac{\varepsilon}{2} \right)_+^p + \left(|y| - \frac{\varepsilon}{2} \right)_+^p \right],$$

whence, from (3.4) and (3.5), it remains to prove

$$\sum_{n=2}^{\infty} \mathbb{E} \left[\max_{1 \leq k < n} \left| \sum_{j=1}^k \frac{n^{\gamma-1} X_{n,j}}{|\Delta_n| j^\gamma} \right| - \varepsilon \right]_+^r < \infty, \quad (3.12)$$

$$\sum_{n=2}^{\infty} \mathbb{P} \left[\max_{1 \leq k < n} \left| \sum_{j=1}^k \frac{n^{\gamma-1} X_{n,n-j+1}}{|\Delta_n| j^\gamma} \right| - \varepsilon \right]_+^r < \infty \quad (3.13)$$

for every $\varepsilon > 0$. In addition to (3.10), we still have

$$\sum_{j=1}^{n-1} \mathbb{P} \left\{ \frac{n^{1-\gamma} |X_{n,j}|}{|\Delta_n| j^\gamma} > \delta \right\} \leq \begin{cases} C(r, \delta, \gamma) |\Delta_n|^{-r} n^{1-r}, & 0 \leq \gamma < 1/r \\ C(r, \delta) |\Delta_n|^{-r} n^{1-r} \log n, & \gamma = 1/r \\ C(r, \delta, \gamma) |\Delta_n|^{-r} n^{r(\gamma-1)}, & 1/r < \gamma < 1, \end{cases}$$

and

$$\sum_{j=1}^{n-1} \int_{\delta^r}^{\infty} \mathbb{P} \left\{ \frac{n^{r(\gamma-1)} |X_{n,j}|^r}{|\Delta_n|^r j^{r\gamma}} > t \right\} dt \leq \sum_{j=1}^{n-1} \frac{n^{r(\gamma-1)} \mathbb{E}|X_{n,j}|^r}{|\Delta_n|^r j^{r\gamma}} I_{\left\{ \frac{n^{\gamma-1} |X_{n,j}|}{|\Delta_n| j^\gamma} > \delta \right\}}$$

$$\begin{aligned}
&\leq \frac{n^{r(\gamma-1)}}{|\Delta_n|^r} \left(\sup_{n,j} \mathbb{E}|X_{n,j}|^r \right) \sum_{j=1}^{n-1} j^{-r\gamma} \\
&\leq \begin{cases} C(r, \gamma) |\Delta_n|^{-r} n^{1-r}, & 0 \leq \gamma < 1/r \\ C(r) |\Delta_n|^{-r} n^{1-r} \log n, & \gamma = 1/r \\ C(r, \gamma) |\Delta_n|^{-r} n^{r(\gamma-1)}, & 1/r < \gamma < 1. \end{cases}
\end{aligned}$$

From (3.2)-(3.3), assumptions (b') and (c') in Corollary 1 hold, as well as (i') in Theorem 2. Since (3.11) remains valid, and

$$\begin{aligned}
\sum_{j=1}^{n-1} \mathbb{E} \left(\frac{n^{\gamma-1} |X_{n,j}|}{|\Delta_n| j^\gamma} I_{\left\{ \frac{n^{\gamma-1} |X_{n,j}|}{|\Delta_n| j^\gamma} > \delta \right\}} \right) &\leq \delta^{1-r} \sum_{j=1}^{n-1} \frac{n^{r(\gamma-1)} \mathbb{E}|X_{n,j}|^r}{|\Delta_n|^r j^{r\gamma}} \\
&\leq \frac{n^{r(\gamma-1)}}{\delta^{r-1} |\Delta_n|^r} \left(\sup_{n,j} \mathbb{E}|X_{n,j}|^r \right) \sum_{j=1}^{n-1} j^{-r\gamma} \\
&\leq \begin{cases} C(r, \delta, \gamma) |\Delta_n|^{-r} n^{1-r}, & 0 \leq \gamma < 1/r \\ C(r, \delta) |\Delta_n|^{-r} n^{1-r} \log n, & \gamma = 1/r \\ C(r, \delta, \gamma) |\Delta_n|^{-r} n^{r(\gamma-1)}, & 1/r < \gamma < 1, \end{cases}
\end{aligned}$$

assumption (d') in Corollary 1 is also met. Hence, (3.12) follows via Corollary 1. Analogously, we can demonstrate (3.13). The proof is complete. \square

Remark If $\Delta_n = O(1)$ as $n \rightarrow \infty$, assumption (3.3) can be replaced by

$$\begin{cases} \sum_{n=1}^{\infty} |\Delta_n|^{2(1-r)} n^{1-r} < \infty, & 0 \leq \gamma < 1/r \\ \sum_{n=1}^{\infty} |\Delta_n|^{2(1-r)} n^{1-r} \log n < \infty, & \gamma = 1/r \\ \sum_{n=1}^{\infty} |\Delta_n|^{2r(\gamma-1)} n^{r(\gamma-1)} < \infty, & 1/r < \gamma < 1/2 \\ \sum_{n=1}^{\infty} |\Delta_n|^{2(1-r)} n^{-r/2} < \infty, & \gamma = 1/2 \\ \sum_{n=1}^{\infty} |\Delta_n|^{-r} n^{r(\gamma-1)} < \infty, & 1/2 < \gamma < 1. \end{cases}$$

Since, for every $\varepsilon > 0$,

$$\mathbb{E}|\hat{\tau}_n - \tau^*|^r \leq 2^{r-1} \left[\mathbb{E}(|\hat{\tau}_n - \tau^*| - \varepsilon)_+^r + \varepsilon^r \right]$$

assumptions of Theorem 4 also entail that $\hat{\tau}_n$ is consistent in the r th mean for τ^* .

In [17], the authors investigated strong consistency of $\hat{\tau}_n$ by considering the model

$$Y_i = \begin{cases} X_i, & i \leq k^* \\ \delta_0 + X_i, & i > k^* \end{cases} \quad (3.14)$$

where $\{X_i, i \geq 1\}$ is either a sequence of independent random variables or a sequence having certain dependence structure. Let us emphasize that, not only model (3.1) is more general than (3.14), but also asymptotic results of [17] are no longer guaranteed in our scenario. As a matter of fact, strong convergence for arrays of random variables requires stronger conditions on (absolute) moments of the random variables (see, for instance, [13]).

3.1 Simulations

In model (3.1), we shall assume the existence of a change-point k^* and

$$(X_{n,1}, X_{n,2}, \dots, X_{n,n}) \stackrel{d}{=} \mathcal{N}(\mathbf{0}, \boldsymbol{\Sigma}_n) \quad (3.15)$$

i.e., the n -dimensional random variable $(X_{n,1}, X_{n,2}, \dots, X_{n,n})$ has (nonsingular) multivariate normal distribution with mean vector zero and covariance matrix

$$\boldsymbol{\Sigma}_n = \begin{bmatrix} 1/n + \sigma^{-n-1}/(\sigma - 1) & -\sigma^{-n-3} & \dots & -\sigma^{-2n-1} \\ -\sigma^{-n-3} & 2/n + 2\sigma^{-n-2}/(\sigma - 1) & \dots & -\sigma^{-2n-2} \\ \vdots & \vdots & \ddots & \vdots \\ -\sigma^{-2n-1} & -\sigma^{-2n-2} & \dots & 1 + n\sigma^{-2n}/(\sigma - 1) \end{bmatrix} \quad (\sigma > 1).$$

It is well-known that if (3.15) holds with $\boldsymbol{\Sigma}_n$ having non-positive off-diagonal entries then, for each $n \geq 1$, the sequence of random variables $\{X_{n,j}, 1 \leq j \leq n\}$ is negatively associated (see, for instance, Theorem 8 of [11]). Further, $X_{n,j} \stackrel{d}{=} \mathcal{N}(0, j/n + j\sigma^{-n-j}/(\sigma - 1))$ (see [18]) and

$$\mathbb{E}|X_{n,j}|^r = \frac{2^{r/2}}{\sqrt{\pi}} \Gamma\left(\frac{r+1}{2}\right) \left(\frac{j}{n} + \frac{j\sigma^{-n-j}}{\sigma - 1}\right)^{r/2} \quad (r > 0)$$

with $\Gamma(z) := \int_0^\infty t^{z-1} e^{-t} dt$, $\text{Re}(z) > 0$ denoting the gamma function.

For the purpose of the simulations, also note that, by taking $\Delta_n = n^\theta$ in (3.3), we have

$$\begin{cases} \sum_{n=1}^\infty [n^{(1-r)(2\theta+1)} + n^{1-(1+\theta)r}], & 0 \leq \gamma < 1/r \\ \sum_{n=1}^\infty [n^{(1-r)(2\theta+1)} + n^{1-(1+\theta)r} \log n], & \gamma = 1/r \\ \sum_{n=1}^\infty [n^{r(\gamma-\theta-1)} + n^{r(\gamma-1)(2\theta+1)}], & 1/r < \gamma < 1/2 \\ \sum_{n=1}^\infty [n^{(r-1)(2\theta+1)} \log^{r-1} n + n^{-r(\theta+1/2)}], & \gamma = 1/2 \\ \sum_{n=1}^\infty n^{r(\gamma-\theta-1)}, & 1/2 < \gamma < 1 \end{cases}$$

and hence, the prior five series converge if

$$\begin{cases} \theta > (2-r)/(2r-2), & 0 \leq \gamma \leq 1/r \\ \theta > 1/(2r-2r\gamma) - 1/2, & 1/r < \gamma < 1/2 \\ \theta > (2-r)/(2r), & \gamma = 1/2 \\ \theta > \gamma - 1 + 1/r, & 1/2 < \gamma < 1. \end{cases} \quad (3.16)$$

We additionally consider the following simulation parameters:

i. $\sigma = 2$, $\mu_{n,k} = 1$, $\tau^* = 0.5$ and $\gamma = 0, 0.1, 0.5, 0.7, 0.9$;

ii. $\Delta_n = n^\theta$ with

$$\theta = \begin{cases} -0.29, -0.2, -0.09, 0, 0.1 & \text{if } \gamma = 0, 0.1, 0.5 \\ -0.19, -0.1, -0.01, 0, 0.1 & \text{if } \gamma = 0.7 \\ -0.01, 0, 0.1 & \text{if } \gamma = 0.9 \end{cases}$$

where the values of θ are in agreement with (3.16);

iii. $n = 50, 100, 500, 1000, 1500, 2000, 3000, 4000$;

iv. $m = 1000$ replications.

The simulation results are displayed in Figures 1 and 2. In all cases, and as expected, one observes the convergence of the CUSUM estimator from some value of n onwards.




Figure 1: Simulation boxplots referring to parameters $\gamma = 0, 0.1, 0.5$




Figure 2: Simulation boxplots referring to parameters $\gamma = 0.7, 0.9$

Supplementary materials

The R simulation codes are available upon request.

Acknowledgements

This work is funded by national funds through the FCT - Fundação para a Ciência e a Tecnologia, I.P., under the scope of projects UIDB/04035/2020 (GeoBioTec) and UIDB/00297/2020 (Center for Mathematics and Applications)

References

- [1] J. Antoch, M. Hušková, Change-point problem and bootstrap, *J. Nonparametr. Stat.* 5 (1995) 123–144.
- [2] J. Antoch, M. Hušková, Z. Prášková, Effect of dependence on statistics for determination of change, *J. Statist. Plann. Inference* 60 (1997) 291–310.
- [3] P. Chen, T. C. Hu, X. Liu, A. Volodin, On complete convergence for arrays of row-wise negatively associated random variables, *Theory Probab. Appl.* 52(2) (2008) 323–328.
- [4] Y. S. Chow, H. Teicher, *Probability Theory: Independence, Interchangeability, Martingales*, 3rd ed. Springer-Verlag, New York, 1997.
- [5] M. Csörgő, L. Horváth, *Limit Theorems in Change-Point Analysis*, John Wiley & Sons, Chichester, 1997.
- [6] P. Embrechts, C. Klüppelberg, T. Mikosch, *Modelling Extremal Events*, Springer-Verlag, Berlin Heidelberg, 1997.
- [7] P. L. Hsu and H. Robbins, Complete convergence and the law of large numbers, *Proc. Natl. Acad. Sci. USA* 33(2) (1947) 25–31.
- [8] T. Z. Hu, Negatively superadditive dependence of random variables with applications, *Chin. J. Appl. Probab. Stat.* 16 (2000) 133–144.
- [9] T. C. Hu, C. Y. Chiang, R. L. Taylor, On complete convergence for arrays of rowwise m -negatively associated random variables, *Nonlinear Analysis* 71 (2009) e1075–e1081.
- [10] K. Joag-Dev, F. Proschan, Negative association of random variables, with applications, *Ann. Statist.* 11 (1983) 286–295.
- [11] S. Karlin, Y. Rinott, M -matrices as covariances matrices of multinormal distributions, *Linear Algebra Appl.* 52/53 (1983) 419–438.
- [12] P. Kokoszka, R. Leipus, Change-point in the mean of dependent observations, *Statist. Probab. Lett.* 40 (1998) 385–393.
- [13] T. C. Hu, F. Moricz, R. L. Taylor, Strong laws of large numbers for arrays of rowwise independent random variables, *Acta Math. Hungar.* 54(1–2) (1989) 153–162.

- [14] T. C. Hu, S. H. Sung, A. Volodin, A note on the strong laws of large numbers for random variables, *Acta Math. Hungar.* 150(2) (2016) 412–422.
- [15] N. Mukhopadhyay, *Probability and Statistical Inference*, Marcel Dekker, New York, 2000.
- [16] A. Shen, M. Xue, A. Volodin, Complete moment convergence for arrays of rowwise NSD random variables, *Stochastics* 88(4) (2016) 606–621.
- [17] Xiaoping Shi, Yuehua Wu, Baiqi Miao, Strong convergence rate of estimators of change point and its application, *Comput. Statist. Data Anal.* 53 (2009) 990–998.
- [18] Y. L. Tong, *The Multivariate Normal Distribution*, Springer-Verlag, New York, 1990.
- [19] X. Wang, X. Deng, L. Zheng, S. Hu, Complete convergence for arrays of rowwise negatively superadditive-dependent random variables and its applications, *Statistics* 48(4) (2014) 834–850.
- [20] Y. Wu, T. C. Hu, A. Volodin, Complete convergence and complete moment convergence for weighted sums of m -NA random variables, *J. Inequal. Appl.* 2015, Article number 200 (2015).