

Evaporation of four-dimensional dynamical black holes sourced by the quantum trace anomaly

Paolo Meda^{1,2,a}, Nicola Pinamonti^{3,2,b}, Simone Roncallo^{1,2,c}, Nino Zanghì^{1,2,d}

¹ *Dipartimento di Fisica, Università di Genova, Italy.*

² *Istituto Nazionale di Fisica Nucleare - Sezione di Genova, Italy.*

³ *Dipartimento di Matematica, Università di Genova, Italy.*

E-mail: ^apaolo.meda@ge.infn.it, ^bpinamont@dima.unige.it ^csimoneroncallo1996@gmail.com, ^dnino.zanghi@ge.infn.it,

Abstract. In this paper we study the evaporation of a four-dimensional spherically symmetric black hole formed during a gravitational collapse, analyzing the backreaction of a massless conformally coupled quantum scalar field by means of the semiclassical Einstein equation. We link the evaporation and the corresponding black hole mass loss to an ingoing negative energy flux at the dynamical horizon and we show how the quantum matter trace anomaly sources that flux once a suitable averaged energy condition is assumed outside the black hole horizon. This condition is satisfied by the background matter in most realistic classical models of spherically symmetric stellar collapse. Finally, as an example, we evaluate explicitly the negative ingoing flux and the corresponding rate of evaporation in case of a null radiating star described by the Vaidya spacetime.

1 Introduction

Even if a complete description of the interplay between matter and gravity at a fundamental level should be provided by a full theory of quantum gravity, semiclassical gravity and quantum field theory on curved spacetimes brought an important framework to investigate how quantum effects related to matter fields affect the geometry of the spacetime. Once a state for the quantum field is fixed, the *back-reaction* on the classical background can be studied using the *semiclassical Einstein equation*

$$G_{\mu\nu} = 8\pi \langle :T_{\mu\nu}: \rangle_{\omega} \quad (1)$$

in units convention $G = c = \hbar = 1$. In this approximation, the classical *Einstein tensor* $G_{\mu\nu}$ is equated to the expectation value of the *stress-energy tensor* $:T_{\mu\nu}:$ of a quantum theory, evaluated on a state ω . The classical background is a four-dimensional globally hyperbolic spacetime (\mathcal{M}, g) , where \mathcal{M} is a smooth manifold and g a Lorentzian metric with signature $(-, +, +, +)$. It is often argued that solutions of eq. (1) furnish approximations to models of a more fundamental theory of quantum gravity valid in the regime where $R_{\mu\nu\rho\sigma}R^{\mu\nu\rho\sigma} \ll m_P^4$, with m_P denoting the Planck mass, and when the fluctuations of the stress-energy tensor are small [KF93]. One of the most important application of the semiclassical Einstein equation consists in the analysis of the *evaporation of dynamical black holes* [Wal95, Bal84, BB89] formed by the gravitational collapse. Actually, when the back-reaction is not taken into account and when matter is described by any quantum field which is in the vacuum in the asymptotic past, it happens that a Schwarzschild black hole emits radiation at large distance. This radiation is called *Hawking radiation*

and it can be shown that its power spectrum is of thermal nature. The temperature T_H read in this spectrum is known as *Hawking temperature* and it is given in terms of the black hole mass

$$T_H = \frac{1}{8\pi k_B M}, \quad (2)$$

where k_B is the Boltzmann constant. Since the state of quantum matter is assumed to be the vacuum in the asymptotic past, the power of the radiation at future infinity is associated with an instantaneous loss of the black hole mass. In the adiabatic approximation, namely with the assumption that the spacetime is instantaneously described by the Schwarzschild metric, one obtains that the black hole is evaporating at a fixed rate proportional to M^{-2} [Haw74, Haw75, Pag76]. Actually, since the horizon radius is also proportional to the mass, the black hole is expected to shrink during the evaporation.

It would be desirable to go beyond the adiabatic approximation and to investigate black hole evaporation with a full dynamical model governed by a *local equation*. In this case, concepts like event horizon and its adiabatic changes need to be substituted with apparent horizon and its evolution during evaporation [AK02, AK03, AK04]. The main difference between an event (static) horizon and an apparent (dynamical) horizon consists in its properties as a geometric surface: whereas an event horizon is always of null type, a trapping horizon can evolve in a spacelike or timelike surface, which implies that the area of the horizon can increase or decrease, as well for the black hole mass, respectively in a formation or evaporation process. Hence, while the value of the Hawking temperature results to be constant on an event horizon, for dynamical black holes it depends on the evolution of the apparent horizon [VAC11, KPV21].

The process of evaporation can be semiclassically explained by the presence of an ingoing negative energy flux on the horizon, violating the classical null energy condition $T(k, k) \geq 0$ for all null vectors k . This flux is responsible of both the shrink of its area and of the production of an outgoing positive energy flux, asymptotically described as Hawking radiation. Actually, such a violation is not indeed surprising, since it has been proven that a very large class of pointwise energy conditions are not valid when quantum fields are involved, even in flat spacetime (the Casimir effect, for instance). See e.g. [FR97, Fla97] and the references therein. However, these conditions can often be replaced by their averaged counterparts, i.e., the *averaged energy conditions*, in which the pointwise expectation value of the stress-energy tensor is replaced by a renormalized stress-energy tensor smeared with a suitable positive test function. The validity of the quantum averaged energy conditions has been established in many contexts, see e.g. [WY91, FR95, FR96, FR03, FV03, Brow18, FK20, SV08, FKK20]. In particular, in those papers, the semiclassical Einstein equation is used together with averaged energy conditions to identify a class of physically admissible spacetimes and states. Such conditions are also used to derive singularity theorems on semiclassical grounds. For a general review about classical and quantum energy inequalities and further references about this topic, see [KS20].

The presence of Hawking radiation and the related evaporation are often ascribed to the presence of an *anomalous trace* in the stress-energy tensor of a quantum matter field near black hole horizons [CF77, DFU76, FD76, Bar81, Bar14]. In the framework of quantum fields on curved spacetimes, such an anomalous trace is a local contribution which depends only on the geometry and the linear equation of motion of the matter field. Moreover, it arises in any covariant regularization procedure, which gives origin to a *covariantly conserved* stress-energy tensor $:T_{\mu\nu}:$, while breaking the classical conformal invariance of $T_{\mu\nu}$ [Wal77, Wal78, BD84, HW04, Mor02]. Here, we shall describe the quantum matter by a *quantum massless conformally-coupled scalar field*. The expectation values of the stress-energy tensor $\langle :T_{\mu\nu}: \rangle_\omega$, and of the vacuum polarization $\langle :\phi^2: \rangle_\omega$ are not explicitly available, when the background is a generic dynamical black hole (see [Can80, How84, AHS95, AGCF20] and the references therein for an analysis of the expectation value of the stress-energy tensor and of vacuum polarization on Schwarzschild spacetime). Actually, from a mathematical point of view the computation of the expectation values of

quadratic observables like $T_{\mu\nu}$ and ϕ^2 is not straightforward and requests some prescriptions about the choice of the state ω , which must be sufficiently regular to give finite expectation values of *normal ordered fields* [Wal95,HW01,HW02,BFK96]. On the other hand, the trace-anomaly of the quantum stress-energy tensor can be evaluated explicitly and independently of the state also for dynamical black holes. For a discussion about quantum field theory on curved spacetimes and its applications like Hawking radiation, see [HW15].

In this paper we analyze the influence of a massless conformally-coupled quantum scalar field on a spherically symmetric dynamical black hole formed after a gravitational collapse, namely a solution of the (semiclassical) Einstein equation with future outer trapping horizon \mathcal{H} and fully described by the Hawking mass $m(V, U)$ and the radius $r(V, U)$ in double-null coordinates (V, U) . First, we study the dynamics of the expansion parameter θ_+ of an outgoing null geodesics congruence emanated from the horizon, with a purely local dynamical equation. The expansion parameter is directly related to the derivative of the black hole mass along the outgoing null geodesics. We then use a semiclassical Raychaudhuri equation to relate the V -derivative of that expansion with $\langle :T_{VV}: \rangle_\omega$, which describes the flux of ingoing energy, and we show that a negative flux on the horizon induces a positive variation of the expansion parameter and consequently a mass loss; in other words, the spatial area of the dynamical horizon is shrinking and the black hole is evaporating. More precisely, as recalled in eqs. (23) to (25), a negative energy flux is directly related to the derivative of the mass with respect to the advanced time v in advanced Eddington-Finkelstein coordinates (v, r) . The negative derivative of the mass (with respect to v) drives the decrease of the spatial area of the dynamical horizon, which turns to be a timelike surface. In addition, assuming the validity of the semiclassical Einstein equation and quantum matter with vanishing initial conditions in the past, we show in Theorem 3.1 that ΔM , the total variation of the mass, can be obtained applying the divergence theorem (Stokes' theorem) to a current J_r^μ given in eq. (26) on a suitable domain. We shall see in eq. (33) that ΔM is sourced by a volume integral involving the divergence of the current J_r^μ , while it is balanced by a boundary contribution proportional to $\langle :T_{UV}: \rangle_\omega$. Such a volume integral corresponds also to the outgoing positive flux emitted towards future infinity, see eq. (35).

Then, we consider the role of the trace anomaly in this balance to show that a negative mass variation always occurs under certain conditions on the matter. In the case of a massless quantum matter scalar field conformally coupled with the curvature, $\langle :T_\rho{}^\rho: \rangle_\omega$ does not depend on the quantum state ω , but it is only a local function of the curvature of the spacetime. Its explicit form is given in eq. (37). We show in Theorem 4.1 that this anomalous term induces a negative variation of the mass, and hence it sources the evaporation process of the black hole, when a certain averaged energy condition is assumed on the component $\langle :T_{UV}: \rangle_\omega$, see eq. (39). We observe that this energy condition is physically reasonable in the light of Hayward's interpretation of $\langle :T_{UV}: \rangle_\omega$ as the work done by the matter field on the system. Furthermore, it is fulfilled in case of classical matter which sources most of the physical realistic models of gravitational collapse, e.g. in those of Oppenheimer-Snyder and Lemaître-Tolman-Bondi (see e.g. [GP09] and references therein) and also when the collapsing matter is described by a classical scalar field, as shown in the works of Christodoulou [Chr86a, Chr86b, Chr91]. As an example, we consider the evaporation induced on the spacetime generated by a null radiating star, namely the Vaidya spacetime, parametrized by the time-dependent mass function $m(v)$ in coordinates (v, r) . In this case, the anomalous term takes the simple form of eq. (40) and the rate of evaporation can be evaluated explicitly, after obtaining the negative flux $\langle :T_{VV}: \rangle_\omega$ on the horizon from the conservation equation $\nabla^\mu \langle :T_{\mu\nu}: \rangle_\omega = 0$ along ingoing null geodesics (see eqs. (42) to (44)).

The paper is organised as follows. In Section 2 we recollect some geometric aspects of spherically symmetric spacetimes and apparent horizons and we recall Hayward's thermodynamic interpretation of black hole dynamics. In Section 3 we describe the semiclassical process of evaporation due to the negative

ingoing flux on the horizon and provide an equation for the variation of the mass. In [Section 4](#) we show that the quantum trace anomaly of a free massless conformally-coupled scalar field drives the evaporation assuming a certain averaged quantum energy inequality, which is also satisfied by the background matter in most realistic classical models of collapse. As an example, we compute the rate of evaporation in the Vaidya spacetime. [Section 5](#) contains the conclusions and some possible future developments. The proofs of [Theorem 3.1](#) and [Theorem 4.1](#) are collected in the [Appendix](#).

2 Spherically symmetric black holes

Each *spherically symmetric spacetime* (\mathcal{M}, g) can be always represented by the manifold $\mathcal{M} = \Gamma \times \mathbb{S}^2$, where \mathbb{S}^2 is the two-dimensional sphere of unital radius and Γ is a two-dimensional space normal to \mathbb{S}^2 , and by the metric

$$ds^2 = g_{\mu\nu} dx^\mu dx^\nu = \gamma_{ij} dx^i dx^j + r^2 d\theta^2 + r^2 \sin^2 \theta d\varphi^2, \quad \mu, \nu = 0, \dots, 3, \quad i, j = 1, 2,$$

where $r^2 \in \mathcal{C}^\infty(\Gamma)$ measures the curvature of each sphere. The two-dimensional spacetime (Γ, γ) corresponds to the quotient of \mathcal{M} with respect to the $SO(3)$ group centered at the origin $r = 0$. The invariant $\nabla_\mu r \nabla^\mu r$ defines the *Misner-Sharp energy*

$$m \doteq \frac{r}{2} (1 - \nabla_\mu r \nabla^\mu r), \quad (3)$$

which describes the energy enclosed inside the sphere of radius r (it is a special case of the *Hawking mass* for the class of spherically symmetric spacetimes) [[MS64](#), [Haw68](#)].

In order to describe an evaporating dynamical black hole, the two-dimensional normal line element $d\gamma^2$ is often represented in the *Bardeen-Vaidya metric* [[Bar81](#)]

$$d\gamma^2 = -e^{2\Psi(v,r)} C(v,r) dv^2 + 2e^{\Psi(v,r)} dv dr, \quad (4)$$

where v is the advanced time and

$$C(v,r) \doteq 1 - \frac{2m(v,r)}{r}. \quad (5)$$

In this parametrization, which corresponds to the *advanced Eddington-Finkelstein coordinates* in the vacuum case, a radial curve at constant v describes an ingoing null geodesic. Different from $C(v,r)$, $\Psi(v,r)$ is not an invariant due to the parametrization freedom $v \mapsto \tilde{v}(v)$.

Since any two-dimensional metric is locally conformally flat, we can also parametrize $d\gamma^2$ in terms of *double-null coordinates* (V, U)

$$d\gamma^2 = -2A(V, U) dV dU \quad (6)$$

with respect to the null normal directions ∂_V and ∂_U . The orientation of the spacetime can be also chosen so that $A(V, U) > 0$ and at spatial infinity $\partial_V r > 0$, $\partial_U r < 0$. The metric is invariant under any re-parametrization $U \mapsto \tilde{U}(U)$ and $V \mapsto \tilde{V}(V)$, then we can represent the future-directed null normal vector fields as

$$\ell_+ \doteq A^{-1} \partial_V, \quad \ell_- \doteq \partial_U, \quad (7)$$

which, respectively, describe the outgoing and the ingoing light rays across the spheres that foliates \mathcal{M} . In this parametrization, the vector field ℓ_+ fulfils the geodesic equation whereas ℓ_- is an auxiliary vector,

and they are normalized such that $g_{\mu\nu}\ell_+^\mu\ell_-^\nu = -1$. The local change of coordinates which relates the metrics (4) and (6) is given by

$$\begin{cases} 2AdU = e^{2\Psi(v,r)}C(v,r)dv - 2e^{\Psi(v,r)}dr, \\ dV = dv. \end{cases} \quad (8a)$$

$$\quad (8b)$$

According to the definitions given by Hayward [Hay93, Hay96, Hay98, Hay00], each sphere that foliates \mathcal{M} is defined to be *untrapped*, *marginal* or *trapped* depending on whether the dual vector $\nabla^\mu r$ is spacelike, lightlike or timelike, respectively. If $\nabla^\mu r$ is future/past-directed, then the sphere is *future/past trapped*: the past case is related to white holes, whereas the future one to black holes, where both outgoing and ingoing light rays are trapped into the surfaces. An hypersurface foliated by marginal spheres is called a *trapping horizon* and a trapping horizon is *outer*, *degenerate* or *inner* when $\nabla^2 r > 0$, $\nabla^2 r = 0$ or $\nabla^2 r < 0$, respectively. In the (V, U) foliation one considers the *expansion parameters* of the congruences of outgoing/ingoing radial null geodesics

$$\theta_+ \doteq \frac{2}{Ar}\partial_V r, \quad \theta_- \doteq \frac{2}{Ar}\partial_U r. \quad (9)$$

Then, a *trapping surface* is defined as a compact spatial two-surface with $\theta_+\theta_- \geq 0$; it is future/past when $\theta_\pm > 0$ or $\theta_\pm < 0$, respectively, and marginal when $\theta_+ = 0$. A trapping horizon is defined as an hypersurface foliated by marginal surfaces; moreover, it is future if $\theta_- < 0$ or past if $\theta_- > 0$, outer if $\partial_U\theta_+ < 0$ or inner if $\partial_U\theta_+ > 0$. Note that both the expansions are smooth functions outside $r = 0$ since \mathcal{M} is smooth. In the framework of black hole physics, an *apparent horizon* is defined to be a *future outer trapping horizon* satisfying

$$\theta_+ \stackrel{\mathcal{H}}{=} 0, \quad \theta_- \stackrel{\mathcal{H}}{<} 0, \quad \partial_U\theta_+ \stackrel{\mathcal{H}}{<} 0, \quad (10)$$

where the subscript \mathcal{H} labels the evaluation on the apparent horizon. The first two conditions capture the fact that no outgoing rays can escape from \mathcal{H} , different from the ingoing ones which are converging therein; the third condition means that the area of the outgoing congruence is increasing just outside \mathcal{H} and it is decreasing just inside \mathcal{H} . On spherically symmetric spacetimes, the apparent horizon is the three-dimensional hypersurface

$$\mathcal{H} = \{(p_\gamma, \Omega) \in \mathcal{M} : r - 2m = 0\}, \quad (11)$$

which is also a *dynamical horizon* according to the definition given in [AK02, AK03, AK04]. Hence, the *mass of the black hole* M is defined as the Misner-Sharp energy evaluated on the horizon $r_{\mathcal{H}} \doteq 2m$

$$M \doteq m(r_{\mathcal{H}}) = \frac{r_{\mathcal{H}}}{2}. \quad (12)$$

In coordinates (v, r) the apparent horizon is described by the line $r_{\mathcal{H}}(v)$ defined by $C(v, r) = 0$. Furthermore, the mass $M(v) \doteq m(v, r_{\mathcal{H}}(v))$ is fully determined by the *rate of evaporation*

$$\dot{M}(v) \doteq \partial_v M(v). \quad (13)$$

On spherically symmetric spacetimes a preferred notion of “time” exists due to the definition of the *Kodama vector* [Kod80, AV10]

$$K \doteq g^{-1}(*dr) = A^{-1}(\partial_V r \partial_U - \partial_U r \partial_V), \quad (14)$$

where $*$ is the Hodge operator in the space normal to the spheres. This vector reduces to be parallel to the timelike Killing vector ∂_t on static spherically symmetric spacetimes. Considering any covariantly

conserved tensor $T^\mu{}_\nu$, even if K is not a Killing field, it holds that $\nabla_\mu K^\mu = 0$ and thus it can be used to define covariantly conserved currents

$$\nabla_\mu (T^\mu{}_\nu K^\nu) = 0. \quad (15)$$

The Kodama vector is timelike on untrapped spheres, namely in the region outside the horizon, and becomes lightlike on a marginal sphere, and eventually it is spacelike on trapped surfaces, i.e., in the interior of the black hole. From the definition of K , one obtains also that

$$\mathcal{L}_K K_\mu = K^\nu (\nabla_\nu K_\mu - \nabla_\mu K_\nu) \stackrel{\mathcal{H}}{=} \pm \kappa K_\mu, \quad (16)$$

where \mathcal{L}_K denotes the Lie derivative along K and

$$\kappa \doteq \frac{1}{2} \gamma^{ij} \nabla_i \nabla_j r. \quad (17)$$

Hence, eq. (17) corresponds to the definition of the *surface gravity* for a dynamical black hole and reduces to the standard one in the case of a Killing vector field (for a discussion about the different definitions of κ in literature, see e.g. [VAC11]). Thus, κ represents the gravitational acceleration detected along the black hole horizon. From the definition (17), a trapping horizon is outer, degenerate or inner when κ is positive, null or negative, respectively; in particular $\kappa > 0$ along an apparent horizons like \mathcal{H} .

A thermodynamic interpretation of the evolution of the mass along apparent horizons has been given by Hayward [Hay98], who proved the *first law of black hole (thermo)dynamics* for spherically symmetric black holes,

$$\partial_k m = \frac{\kappa}{8\pi} \partial_k \mathcal{A} + w \partial_k \mathcal{V}. \quad (18)$$

Here, $\partial_k f = k \cdot \nabla f$ denotes the derivative along any vector field k tangent to the horizon, m is the mass (3), κ is the surface gravity (17), $\mathcal{A} = 4\pi r^2$ is the area, $\mathcal{V} = \frac{4}{3}\pi r^3$ the volume and $w \doteq -\frac{1}{2} \gamma^{ij} T_{ij} = A^{-1} T_{UV}$ is the *work density* done by the matter field. Furthermore, assuming the Einstein equation $w = (8\pi)^{-1} R_\theta{}^\theta$,

$$\kappa = \frac{m}{r^2} - 4\pi r w. \quad (19)$$

As in the case of static black holes, the thermodynamic interpretation of eq. (18) can be made precise only if the surface gravity is proportional to an actual temperature. This is really the case of a static black hole, where $\kappa/2\pi$, in natural units, equals the Hawking temperature of Hawking radiation observed at future infinity [Haw75, FH90, KW91, Wal01]. In the dynamical case, it is possible to show that the very same temperature can be seen in the tunnelling probability of matter across dynamical horizons. A derivation of this fact involving the WKB approximation for the one particle excitations can be found in [DNVZZ07, HDVNZ09], based on the ideas presented in [PW00]. Moreover, another derivation focusing on the properties of states for quantum fields near apparent horizons is presented in [KPV21]. The latter observation enforces the statement that κ defined in eq. (17) must be a positive quantity at least near the apparent horizon.

3 A semiclassical model of evaporation

The local dynamics of the apparent horizon \mathcal{H} can be analyzed by studying the evolution of the expansion parameter θ_+ given in eq. (9) along an outgoing null geodesic parametrized by ℓ_+ . Denoting with $d/d\lambda \doteq \ell_+^\mu \nabla_\mu$ the directional derivative along ℓ_+ , the semiclassical Einstein equation $G_{VV} = 8\pi \langle :T_{VV}: \rangle_\omega$ reads

$$\frac{d\theta_+}{d\lambda} = -\frac{1}{2} \theta_+^2 - 8\pi \langle :T_{\mu\nu}: \rangle_\omega \ell_+^\mu \ell_+^\nu, \quad (20)$$

which represents the *Raychaudhuri equation* for the null affine-parametrized outgoing geodesics congruence [Wal84] (it is sometimes referred also as the *Landau-Raychaudhuri equation*). For classical black holes, $\langle :T_{\mu\nu}: \rangle_\omega$ is replaced by $T_{\mu\nu}$, hence $d\theta_+/d\lambda \leq 0$ from the null energy condition $T_{\mu\nu}\ell_+^\mu\ell_+^\nu \geq 0$. Thus, assuming the initial condition $\theta_+(V_0) = 2/(Ar) > 0$ at the beginning of the collapse, there must exist a region where $\theta_+ = 0$ for $V > V_0$, namely that a trapped surface has formed during the gravitational collapse. On the other hand, when evaluated on the apparent horizon ($\theta_+ = 0$) and in the semiclassical case, eq. (20) reduces to

$$\frac{d\theta_+}{d\lambda} \stackrel{\mathcal{H}}{=} -8\pi \frac{\langle :T_{VV}: \rangle_\omega}{A^2}. \quad (21)$$

Hence, the presence of a *negative ingoing energy flux* $\langle :T_{VV}: \rangle_\omega$ on \mathcal{H} , violating the null energy condition, implies that $d\theta_+/d\lambda > 0$ locally on the horizon. Therefore, the trapped surface formed during the collapse tends to disappear, namely it evaporates. This process of evaporation of the horizon makes manifest as a loss of the black hole mass given in eq. (12). Given a one-dimensional portion of horizon $\delta\mathcal{H} \subseteq \pi_\gamma(\mathcal{H})$, where $\pi_\gamma : \Gamma \times_{r^2} \mathbb{S}^2 \rightarrow \Gamma$ denotes the natural projection on the first pair of coordinates, let us define the *variation of mass* of the black hole on $\delta\mathcal{H}$

$$\Delta M \doteq \int_{\delta\mathcal{H}} dm. \quad (22)$$

In coordinates (v, r) , $\delta\mathcal{H}$ is the line enclosed between two arbitrary points $(v_P, r_{\mathcal{H}}(v_P))$ and $(v_Q, r_{\mathcal{H}}(v_Q))$ in the (v, r) plane. On \mathcal{H} , the relation $dr = (1 - 2\partial_r m)^{-1} 2\partial_v m dv$ holds and eq. (21) becomes a dynamical law for the rate \dot{M} defined in eq. (13). Actually, after rescaling v so that $\Psi(v, r_{\mathcal{H}}) = 0$, both eq. (21) and the semiclassical equation $G_{vv}r^2 \stackrel{\mathcal{H}}{=} 8\pi \langle :T_{vv}: \rangle_\omega r^2$ read

$$\dot{M}(v) = \mathcal{A}_{\mathcal{H}}(v) \langle :T_{vv}: \rangle_\omega(v, r_{\mathcal{H}}(v)), \quad (23)$$

where $\mathcal{A}_{\mathcal{H}} \doteq 4\pi r_{\mathcal{H}}^2$ denotes the *area of the horizon*. Thus, evaluating the surface gravity (17) on \mathcal{H} ,

$$\Delta M = \int_{v_P}^{v_Q} \frac{\dot{M}}{4M\kappa} dv = 4\pi \int_{v_P}^{v_Q} \frac{M}{\kappa} \langle :T_{vv}: \rangle_\omega(r_{\mathcal{H}}) dv. \quad (24)$$

Since $\kappa > 0$ and $\langle :T_{vv}: \rangle_\omega \stackrel{\mathcal{H}}{=} \langle :T_{VV}: \rangle_\omega$, both the rate of evaporation (13) and the variation of the mass (22) are negative when $\langle :T_{VV}: \rangle_\omega < 0$ on the horizon. Moreover, if we consider the vector in the (v, r) plane $n_{\mathcal{H}} \doteq g^{\mu\nu}\partial_\nu \mathcal{H} \partial_\mu$ normal to the apparent horizon \mathcal{H} , we have that

$$g_{\mu\nu} n_{\mathcal{H}}^\mu n_{\mathcal{H}}^\nu \stackrel{\mathcal{H}}{=} -16M\kappa\dot{M}. \quad (25)$$

So, if $\Delta M < 0$ then $\dot{M} < 0$ and $n_{\mathcal{H}}$ is spacelike, hence \mathcal{H} is a *timelike surface*, namely it is evaporating. Finally, one can explicitly relate ΔM to the expectation value of certain components of the stress-energy tensor outside the horizon, using the *divergence theorem (Stokes' theorem)* for suitable currents on a suitable domain. To this end, let us introduce the following currents obtained contracting the expectation values of the stress-energy tensor with the gradient ∇r and with the Kodama vector (14)

$$J_r^\mu \doteq \langle :T^{\mu\nu}: \rangle_\omega \nabla_\nu r = J_1^\mu + J_2^\nu = \langle :T^{U\mu}: \rangle_\omega \partial_U r + \langle :T^{V\mu}: \rangle_\omega \partial_V r, \quad (26)$$

$$J_K^\mu \doteq \langle :T^{\mu\nu}: \rangle_\omega K_\nu = J_1^\mu - J_2^\nu = \langle :T^{U\mu}: \rangle_\omega \partial_U r - \langle :T^{V\mu}: \rangle_\omega \partial_V r. \quad (27)$$

Denoting by $\nabla \cdot J \doteq \nabla_\mu J^\mu$ the divergence of the current J , from eq. (15) we get $\nabla \cdot J_K = 0$, which implies that $\nabla \cdot J_1 = \nabla \cdot J_2$ and $\nabla \cdot J_1 = \frac{1}{2} \nabla \cdot J_r$. Moreover, we observe that on the horizon $J_2 = 0$ because

$\partial_V r = 0$. The domain D , over which the divergence theorem is applied, is obtained considering

$$S_0 \doteq \{(V, U, \theta, \varphi) \in \mathcal{M} \mid V = V_0, U = U_0\}, \quad (28)$$

$$S_1 \doteq \{(V, U, \theta, \varphi) \in \mathcal{M} \mid V = V_1, U = U_0\}, \quad (29)$$

where U_0, V_0 and $V_1 > V_0$ are chosen such that both S_0 and S_1 lie outside the apparent horizon \mathcal{H} and in such a way that $(V_1, U_2) \times \mathbb{S}^2$ is contained on \mathcal{H} for some U_2 . Consider now $\delta\mathcal{H} \times \mathbb{S}^2$ the portion of \mathcal{H} which intersects $J^+(S_0) \cap (\mathcal{M} \setminus I(S_1))$. We denote by $P_{\mathcal{H}} = (V_1, U_2)$ and $Q_{\mathcal{H}} = (V_0, U_1)$ the extreme points of $\delta\mathcal{H}$ in the (V, U) plane. The domain $D \times \mathbb{S}^2$ is then obtained considering the portion of $J^+(S_0) \cap (\mathcal{M} \setminus I(S_1))$ which lies outside the apparent horizon \mathcal{H} . If \mathcal{H} is spacelike or null,

$$D \times \mathbb{S}^2 \doteq J^+(S_0) \cap J^-(\delta\mathcal{H} \times \mathbb{S}^2), \quad (30)$$

while, if \mathcal{H} is timelike,

$$D \times \mathbb{S}^2 \doteq J^+(S_0) \cap O, \quad (31)$$

where O is the portion of $J^-(\delta\mathcal{H} \times \mathbb{S}^2)$ which lies outside the horizon. With these definitions, we denote by $\rho_0, \delta_0, \gamma \in \partial D$ the one-dimensional curves in the (V, U) plane between (V_0, U_1) and (V_0, U_0) , (V_0, U_0) and (V_1, U_0) , (V_1, U_0) and (V_1, U_2) , respectively. See Figure 1 for a representation of D .




Figure 1: Picture of the domain of integration D given a spacelike portion $\delta\mathcal{H}$ of the apparent horizon. In this case, $D \times \mathbb{S}^2 = J^+((V_0, U_0) \times \mathbb{S}^2) \cap J^-(\delta\mathcal{H} \times \mathbb{S}^2)$, with $U_2 < U_1$. The initial data are posed on the curves ρ_0 and δ_0 of the boundary ∂D .

Thus, we have the following

Theorem 3.1. *Consider a spherically symmetric spacetime parametrized in double-null coordinates (V, U) and a quantum stress-energy tensor $\langle :T_{\mu\nu}: \rangle_\omega$ which respects the spherical symmetry and which satisfies the following initial conditions at the boundary of $J^+(S_0)$, where S_0 is given in eq. (28):*

$$\langle :T_{\mu\nu}: \rangle_\omega(p) = 0, \quad \langle :T_{\mu\nu}: \rangle_\omega(p) = 0, \quad p \in \partial J^+(S_0). \quad (32)$$

Consider the domain D introduced in eq. (30) or in eq. (31), where we have denoted the disjoint components of ∂D by $\delta\mathcal{H}$, ρ_0 , δ_0 , γ and where $P_{\mathcal{H}}$ and $Q_{\mathcal{H}}$ are the extreme points of $\delta\mathcal{H}$ as explained above

(see also Figure 1). Then the variation of the mass (22) is given in terms of the current J_r defined in eq. (26) as

$$\Delta M = -2\pi \int_D \nabla \cdot J_r d\mathcal{V}_D - 4\pi \int_{U_0}^{U_2} \frac{\langle :T_{UV}: \rangle_\omega r^2}{A} (-\partial_{Ur}) dU, \quad (33)$$

where $d\mathcal{V}_D = Ar^2 dV \wedge dU$ is the volume form of the domain D and $\nabla \cdot J_r$ is the divergence of the current J_r on \mathcal{M} . Equivalently, eq. (33) can be written in the form

$$\Delta M = -4\pi \int_{U_0}^{U_2} \frac{\langle :T_{UU}: \rangle_\omega r^2}{A} \partial_V r dU - 4\pi \int_{U_0}^{U_2} \frac{\langle :T_{UV}: \rangle_\omega r^2}{A} (-\partial_{Ur}) dU. \quad (34)$$

Hence, the outgoing flux $\langle :T_{UU}: \rangle_\omega$ emitted from $\delta\mathcal{H}$ is given by

$$\int_{U_0}^{U_2} \frac{\langle :T_{UU}: \rangle_\omega r^2}{A} \partial_V r dU = \frac{1}{2} \int_D \nabla \cdot J_r d\mathcal{V}_D, \quad (35)$$

Proof. See the Appendix A.1 □

Even if the state ω used to compute the expectation values of the stress-energy tensor is not explicitly given, we have assumed implicitly that all the integrals involving $\langle :T_{\mu\nu}: \rangle_\omega$ in Theorem 3.1 are well-defined on (\mathcal{M}, g) , namely, we have assumed that the state is sufficiently regular to give finite expectation values to $\langle :T_{\mu\nu}: \rangle_\omega$ and their averages.

In view of eq. (18), i.e., the first law of black hole dynamics stated by Hayward, we can give a physical meaning to the single contributions appearing in eqs. (33) and (34). Actually, the first equation entails that the variation of the mass on the apparent horizon is sourced by the non-vanishing divergence of the radial current J_r and balanced by the semiclassical work $\langle :w: \rangle_\omega = A^{-1} \langle :T_{UV}: \rangle_\omega$ done by the quantum matter field on the geometry. Furthermore, the divergence term $\nabla \cdot J_r$ is responsible of the appearance of an *outgoing energy flux* $\langle :T_{UU}: \rangle_\omega$, which is the origin of the radiation emitted by the black hole and seen at future infinity. With this latter observation, it is natural to expect that the quantum state ω is such that

$$\int_{U_0}^{U_2} \frac{\langle :T_{UU}: \rangle_\omega r^2}{A} \partial_V r dU \geq 0, \quad (36)$$

which can be thought as an averaged energy condition along a portion of a null ingoing geodesic, smeared by the smooth positive function $\partial_V r$. In the case of matter fields and spacetimes in which $\langle :w: \rangle_\omega = 0$, from eq. (34) we have that the variation of mass is exactly equal to the outgoing energy flux emitted at infinity. We also observe that if the black hole is evaporating, $\Delta M < 0$ and hence the right-hand side of eq. (34) must be negative in that case.

4 Evaporation induced by the trace anomaly and applications to physically relevant spacetimes

For a *free massless conformally-coupled scalar field* ϕ , the non-vanishing trace of the quantum stress-energy tensor $\langle :T_{\mu\nu}: \rangle_\omega$ does not depend on the choice of the quantum state ω , but it is fixed by the geometry of the spacetime to be equal to the *quantum trace anomaly*. In four dimensions, it reads

$$\langle :T_\rho{}^\rho: \rangle_\omega = \lambda \left(C_{\alpha\beta}{}^{\gamma\delta} C_{\gamma\delta}{}^{\alpha\beta} + R_\mu{}^\nu R_\nu{}^\mu - \frac{1}{3} R^2 \right), \quad (37)$$

where

$$\lambda \doteq \frac{1}{720(4\pi^2)}, \quad (38)$$

$C_{\alpha\beta}{}^{\gamma\delta}$ is the *Weyl tensor*, $R_\mu{}^\nu$ the *Ricci tensor* and R the *Ricci scalar* [Wal78]. Using the results stated in the previous section, we shall now see how a non-vanishing trace anomaly fixes the sign of $\nabla \cdot J_r$ in eq. (33) and thus driving the evaporation, once a quantum energy condition on the matter field state holds. This result is contained in the following

Theorem 4.1. *Under the geometric assumptions of Theorem 3.1, consider a free massless, conformally coupled scalar field ϕ and its quantum anomalous trace given by eq. (37). Suppose that the quantum state ω satisfies the initial conditions stated in eq. (32) and suppose that it fulfils*

$$\int_{U_0}^{U_{\mathcal{H}}} \frac{\langle :T_{UV}: \rangle_\omega r^2}{A} f(V, U) AdU \geq 0 \quad (39)$$

for all strictly positive functions $f(V, U) \in \mathcal{C}^\infty(\mathcal{M})$, along any ingoing radial null curve connecting the initial point (V, U_0) and $(V, U_{\mathcal{H}}) \in \delta\mathcal{H}$ at fixed V . Then $\Delta M < 0$, namely evaporation occurs along $\delta\mathcal{H}$.

Proof. See the Appendix A.2 □

The hypothesis stated in eq. (39) represents a sort of *averaged energy condition* for the state ω . If the smearing function is $f(V, U) = A^{-1}(-\partial_U r)$, eq. (39) implies that the integral involving $A^{-1} \langle :T_{UV}: \rangle_\omega$ inside the eq. (34) contributes negatively to the variation of the mass. We observe that the hypotheses of Theorem 4.1 can be weakened. Actually, the proof and thus the thesis of Theorem 4.1 holds even if we assume that eq. (39) holds only for the particular function $f(V, U)$ given in eq. (57). For a classical free matter field, the initial conditions (32) are clearly compatible with $\langle :T_{UV}: \rangle_\omega = 0$. Thus, in that case hypothesis (39) holds trivially. Moreover, that condition is fully compatible with the thermodynamic interpretation given by Hayward of $\langle :w: \rangle_\omega = A^{-1} \langle :T_{UV}: \rangle_\omega$ as the work done by the quantum matter field on the system, which is usually expected to be positive.

On the other hand, the lack of control on the evolution of a quantum state ω compatible with eq. (32) prevents us to verify the desired averaged energy condition explicitly. However, we expect that such a condition is fulfilled at least in an approximate way. Namely, if we assume the background geometry outside the black hole horizon to be close to a spacetime resulting in a classical model of collapse, we expect that the quantum component $\langle :T_{UV}: \rangle_\omega$ is compatible with the Einstein tensor of that spacetime in which quantum matter propagates. Under this approximation, the condition (39) is often satisfied - even pointwise - in the most realistic spherically symmetric models of collapse, where the background matter fulfils the dominant energy condition $A^{-1}T_{UV} \geq 0$.

As examples, we can think at the *Lemaître-Tolman-Bondi* models like the *Oppenheimer-Snyder* solution, where the collapse is driven by an (in)homogeneous spherical cloud of dust at zero pressure satisfying the weak energy condition, see e.g. [GP09] and references therein. Furthermore, in the work of Christodoulou about the collapse in the case of matter described by a classical scalar field [Chr86a, Chr86b, Chr91] the collapsing matter is described by a classical massless scalar field ϕ that is invariant under rotations and having stress-energy tensor given by $T_{\mu\nu} = \partial_\mu \phi \partial_\nu \phi - \frac{1}{2} g_{\mu\nu} \partial_\rho \phi \partial^\rho \phi$. Hence, $T_{UV} = 0$ and the dominant energy condition holds again.

Actually, there are two special backgrounds fulfilling $T_{UV} = 0$ from the Einstein equation $G_{UV} = 8\pi T_{UV} = 0$, namely the *Schwarzschild* and the *Vaidya* spacetimes. The former describes a static spherically symmetric black hole, while the latter defines the geometry outside a null radiating star, see e.g. [GP09] and references therein. In Bardeen-Vaidya parametrization (4), the Schwarzschild and

the Vaidya metrics are obtained by choosing $m \in \mathbb{R}^+$ and $m = m(v)$, respectively, and by fixing $\Psi(v, r) = 0$. In these cases, the semiclassical regime where $R_{\mu\nu\rho\sigma}R^{\mu\nu\rho\sigma} \ll m_P^4$ is always satisfied for $m/m_P \gg (3/4)^{1/4} \simeq 0.93$, which holds for astrophysical masses (for a solar mass, $M_\odot/m_P \simeq 10^{38}$). Whereas in the Schwarzschild spacetime the variation of the mass on the horizon is always zero, in the Vaidya spacetime the trace anomaly is

$$\langle :T_\rho{}^\rho: \rangle_\omega = 48\lambda \frac{M(v)^2}{r^6}, \quad (40)$$

where λ is the coefficient (38). In this case the rate of evaporation can be directly computed using eq. (23) after evaluating the negative ingoing flux $\langle :T_{VV}: \rangle_\omega$ on \mathcal{H} . To obtain $\langle :T_{VV}: \rangle_\omega$, we employ the *conservation equation* $\nabla^\mu \langle :T_{\mu V}: \rangle_\omega = 0$, which reads

$$-\frac{1}{Ar^2} \partial_U (\langle :T_{VV}: \rangle_\omega r^2) - \frac{1}{r^2} \partial_V (A^{-1} \langle :T_{UV}: \rangle_\omega r^2) - 2 \langle :T_\theta{}^\theta: \rangle_\omega \frac{\partial_V r}{r} = 0.$$

Using the relation

$$\langle :T_\theta{}^\theta: \rangle_\omega = \frac{\langle :T_{UV}: \rangle_\omega}{A} + \frac{1}{2} \langle :T_\rho{}^\rho: \rangle_\omega, \quad (41)$$

and assuming the Vaidya spacetime to be a solution of the semiclassical equation $\langle :T_{UV}: \rangle_\omega = G_{UV}/8\pi = 0$, the ingoing flux fulfils the following differential equation in (v, r) coordinates

$$\partial_r (\langle :T_{VV}: \rangle_\omega r^2) = 24\lambda M(v)^2 \left(\frac{1}{r^5} - \frac{2M(v)}{r^6} \right).$$

Integrating in $(r_{\mathcal{H}}, \infty)$ and imposing the initial condition $\langle :T_{VV}: \rangle_\omega r^2 \rightarrow 0$ when $r \rightarrow \infty$, we get

$$\langle :T_{VV}: \rangle_\omega r^2 \stackrel{\mathcal{H}}{=} -24\lambda M^2 \int_{r_{\mathcal{H}}}^{+\infty} \left(\frac{1}{r^5} - \frac{2M(v)}{r^6} \right) dr = -\frac{3\lambda}{40M(v)^2}. \quad (42)$$

Hence, the rate of evaporation obtained from eq. (23) is

$$\dot{M}(v) = -\frac{3\pi\lambda}{10M(v)^2}. \quad (43)$$

Eq. (43) is an ordinary differential equation with respect to v and it can be integrated by separation of variables, yielding the evaporation law

$$M^3(v) = M_0^3 - \frac{9\pi\lambda}{10}(v - v_0), \quad (44)$$

where we have defined the total initial mass $M_0 \doteq M(v_0)$ at the initial time v_0 . Thus, the evaporation process is completed in the time interval $\Delta v \doteq v - v_0 = 10M_0^3/(9\pi\lambda)$. Moreover, according to the first law given in (18) for the Vaidya spacetime, the negative rate (43) induces also a shrink of the area of the horizon $\mathcal{A}_{\mathcal{H}}$, whose rate of variation $\dot{\mathcal{A}}_{\mathcal{H}}(v) \doteq \partial_v \mathcal{A}(v, r_{\mathcal{H}}(v))$ is governed by $\dot{M}(v) = (\kappa/8\pi)\dot{\mathcal{A}}_{\mathcal{H}}(v)$. Hence, a negative variation of the *Wald-Kodama dynamical entropy* $S_{\mathcal{H}} \doteq \mathcal{A}_{\mathcal{H}}/4$ [AH99, HMA99] holds, namely

$$\frac{dS_{\mathcal{H}}(v)}{dv} = -\frac{12\pi^2\lambda}{5M(v)}. \quad (45)$$

5 Conclusion

In this paper we have studied the process of evaporation of a four-dimensional dynamical spherically symmetric black hole, in the framework of quantum field theory on curved spacetimes and semiclassical gravity. We have shown that an evaporating black hole is always obtained when a negative ingoing flux appears on the horizon. This flux is responsible of the negative variation of the black hole mass, which was expressed in terms of the work done by the matter field and the outgoing flux emitted towards future infinity. In this model, the evaporation process is always sourced by the trace anomaly of a free massless, conformally-coupled scalar field when the quantum state, or alternatively the classical background matter, fulfils a sort of quantum averaged energy condition and once suitable vacuum-like initial conditions are assumed. Finally, we have computed the rate of evaporation explicitly in the Vaidya spacetime, finding that it is proportional to M^{-2} as in the Schwarzschild case.

A generalization of this paper can be carried out by extending our analysis to arbitrary massless conformally coupled fields, after modifying the coefficient λ inside the trace anomaly given in eq. (37). In the general case, the four-dimensional anomalous trace is given by $\langle :T_\rho{}^\rho: \rangle_\omega = b_F F + b_G G$, where $F = C_{\alpha\beta\gamma\delta} C^{\alpha\beta\gamma\delta}$ is the square of the Weyl tensor, $G = R_{\mu\nu\rho\sigma} R^{\mu\nu\rho\sigma} - 4R_{\mu\nu} R^{\mu\nu} + R^2$ is the Euler density and b_F , b_G are coefficients depending on the numbers of particles n_s of spin s . For the explicit values of b_F and b_G , see e.g. [BD84]. We argue that a generalization of the Theorem 4.1 can be obtained for arbitrary fields after choosing properly the coefficients inside $\langle :T_\rho{}^\rho: \rangle_\omega$. The generalization of the analysis presented in this paper beyond the spherically symmetric case, namely when the presence of rotation is considered, both on the Kerr and Kerr-Vaidya spacetimes [DT20] is still an open problem.

More generally, the study we have performed should be viewed as a first step towards approaching the more general issue of the evaporation in semiclassical gravity. The next step should consist in studying the back-reaction of the quantum matter field by using the semiclassical Einstein equation and restricting the class of the solutions with the averaged quantum energy conditions on $\langle :T_{\mu\nu}: \rangle_\omega$. This problem was solved in the two-dimensional case in [APR11b, APR11a], but it is still open in the four-dimensional case, because of the difficulties in controlling the state-dependent contributions in the expectation values of the stress-energy tensor $\langle :T_{\mu\nu}: \rangle_\omega$. At the same time, only the knowledge of the details of the state allows to check the validity of the energy conditions assumed in this paper and to analyze the form of the exact solution of the semiclassical equation. However, we think that a study similar to the one in [MPS20] for cosmological spacetimes may represent a possible way to solve the semiclassical Einstein equation even for the class of dynamical black holes.

A Proofs of theorems

In this appendix we collect the proofs of the theorems stated in the main text.

A.1 Proof of Theorem 3.1

Using that $T^{UV} = A^{-2}T_{UV}$ and $T^{UU} = A^{-2}T_{VV}$ we can relate the current J_1 defined in eq. (26) to the variation of the mass (22) computed along the line enclosed between $(V_0, U_1), (V_1, U_2) \in \delta\mathcal{H}$ in the (V, U) plane (see Figure 1). In (V, U) coordinates, the derivatives of the Misner-Sharp energy (3) read

$$\begin{cases} \partial_V m = \frac{4\pi r^2}{A} (\langle :T_{UV}: \rangle_\omega \partial_V r - \langle :T_{VV}: \rangle_\omega \partial_U r), \\ \partial_U m = \frac{4\pi r^2}{A} (\langle :T_{UV}: \rangle_\omega \partial_U r - \langle :T_{UU}: \rangle_\omega \partial_V r). \end{cases} \quad (46a)$$

$$\quad (46b)$$

Evaluating eqs. (46a) and (46b) on $\partial_V r \stackrel{\mathcal{H}}{=} 0$, we obtain that

$$\Delta M = 4\pi \int_{\delta\mathcal{H}} r^2 (-\partial_U r) \left(\frac{\langle :T_{VV}: \rangle_\omega}{A} dV - \frac{\langle :T_{UV}: \rangle_\omega}{A} dU \right) = 4\pi \int_{\delta\mathcal{H}} Ar^2 (J_1^V dU - J_1^U dV). \quad (47)$$

Eq. (33) can be obtained by applying the divergence theorem (Stokes' theorem) to the current J_1 on the domain $D \times \mathbb{S}^2$. Using the spherical symmetry to integrate out the angular variables (φ, θ) , we obtain that

$$-\int_D (\nabla \cdot J_1) d\mathcal{V}_D = \int_{\delta\mathcal{H}} (J_1^V Ar^2 dU - J_1^U Ar^2 dV) + \int_{\rho_0} J_1^V Ar^2 dU + \int_{\delta_0} J_1^U Ar^2 dV - \int_\gamma J_1^V Ar^2 dU.$$

With the choice of the initial conditions (32), both the integrals along ρ_0 and δ_0 vanish. By substitution of eq. (47) at the place of the integral over $\delta\mathcal{H}$, we get

$$-\int_D (\nabla \cdot J_1) d\mathcal{V}_D = \frac{\Delta M}{4\pi} - \int_{U_0}^{U_2} Ar^2 J_1^V dU.$$

Thus, eq. (33) is obtained by employing the definition $J_1 = (J_r - J_K)/2$, where J_r and J_K are given in eqs. (26) and (27), and by using that $\nabla \cdot J_1 = \frac{1}{2}\nabla \cdot J_r$, since $\nabla \cdot J_K = 0$ everywhere.

The proof of eq. (34) can be achieved by applying the divergence theorem (Stokes' theorem) to the current J_K , after observing that $\nabla \cdot J_K = 0$ and $J_K = J_1$ on the horizon. Eventually, eq. (35) is obtained by comparing eqs. (33) and (34).

A.2 Proof of Theorem 4.1

The proof consists in applying the divergence theorem (Stokes' theorem) on the domain $D \times \mathbb{S}^2$ for a current \tilde{J} which is a weighted version of J_1 given in eqs. (26) and (27). The weight is given in terms of a strictly positive function $f(V, U)$ which will be fixed later. For sake of simplicity, we shall set $8\pi G = 1$ in the semiclassical equations. Let us define

$$\tilde{J}_\mu \doteq \xi^\nu \langle :T_{\nu\mu}: \rangle_\omega, \quad \xi^\nu = f(V, U)(\partial_V)^\nu$$

and the weighted variation of the mass

$$\Delta_h M \doteq \int_{\delta\mathcal{H}} h dm \quad (48)$$

with respect to the function

$$h(V, U) \doteq \frac{f(V, U)A(V, U)}{-\partial_U r(V, U)}. \quad (49)$$

The divergence of \tilde{J} is related to the variation of the weighted mass by the following

Lemma A.1. *Under the assumptions of Theorem 3.1 in the domain D of Figure 1,*

$$\Delta_h M = -4\pi \int_D \nabla \cdot \tilde{J} d\mathcal{V}_D - 4\pi \int_{U_0}^{U_2} \frac{\langle :T_{UV}: \rangle_\omega}{A} f A r^2 dU. \quad (50)$$

Proof. Using eqs. (48) and (49), we have that

$$\Delta_h M = 4\pi \int_{\delta\mathcal{H}} r^2 \left(\tilde{J}_V dV - \tilde{J}_U dU \right).$$

Hence, the proof is accomplished applying the divergence theorem (Stokes' theorem) to the weighted current \tilde{J}_μ on the domain $D \times \mathbb{S}^2$, similarly to what already done in the proof of Theorem 3.1 for the current J_r . \square

Using the conservation equation $\nabla^\mu \langle :T_{\mu\nu}: \rangle_\omega = 0$ and the relation (41), we obtain that

$$\nabla \cdot \tilde{J} = - \left(\frac{\langle :T_{VV}: \rangle_\omega}{A} \partial_U f + \frac{\langle :T_{UV}: \rangle_\omega}{A} \partial_V f \right) + f \left(- \frac{\langle :T_{UV}: \rangle_\omega}{A} \frac{\partial_V A}{A} + 2 \frac{\langle :T_{UV}: \rangle_\omega}{A} \frac{\partial_V r}{r} + \langle :T_\rho{}^\rho: \rangle_\omega \frac{\partial_V r}{r} \right).$$

Here, $\langle :T_\rho{}^\rho: \rangle_\omega$ is a geometric quantity given in terms of the trace anomaly in eq. (37). There we can isolate a positive contribution after computing explicitly the product $C_{\alpha\beta}{}^{\gamma\delta} C_{\gamma\delta}{}^{\alpha\beta} = (R + 12\kappa/r)^2$ and the difference

$$R_\mu{}^\nu R_\nu{}^\mu - \frac{1}{3} R^2 = 2 \frac{R_{UU}}{A} \frac{R_{VV}}{A} + \frac{2}{3} \left((g^{UV} R_{UV})^2 + (R_\theta{}^\theta)^2 \right) + \frac{8A}{3r^2} R_\theta{}^\theta R_{UV}.$$

Employing the semiclassical equations $R_{UU} = \langle :T_{UU}: \rangle_\omega$, $R_{VV} = \langle :T_{VV}: \rangle_\omega$ and $G_{UV} = \langle :T_{UV}: \rangle_\omega = AR_\theta{}^\theta$, the anomaly can be rewritten as

$$\langle :T_\rho{}^\rho: \rangle_\omega = \lambda \left(\left(R + \frac{12\kappa}{r} \right)^2 + \frac{2}{3} \left((g^{UV} R_{UV})^2 + (R_\theta{}^\theta)^2 \right) + 2 \frac{\langle :T_{UU}: \rangle_\omega \langle :T_{VV}: \rangle_\omega}{A} + \frac{8A}{3r^2} R_{UV} \frac{\langle :T_{UV}: \rangle_\omega}{A} \right),$$

where the first two terms are manifestly positive. Plugging this expression inside eq. (50) yields

$$\begin{aligned} \Delta_h M &= -4\pi\lambda \int_D \left(\left(R + \frac{12\kappa}{r} \right)^2 + \frac{2}{3} (g^{UV} R_{UV})^2 + \frac{2}{3} (R_\theta{}^\theta)^2 \right) \frac{\partial_V r}{r} f d\mathcal{V}_D \\ &\quad - 4\pi \int_D \left(- \frac{\partial_V f}{f} + \partial_V \log(A^{-1} r^2) + \lambda \frac{8A}{3r^3} \partial_V r R_{UV} \right) \frac{\langle :T_{UV}: \rangle_\omega}{A} f d\mathcal{V}_D \\ &\quad - 4\pi \int_D \left(- \frac{\partial_U f}{f} + \lambda \frac{2\partial_V r \langle :T_{UU}: \rangle_\omega}{r A} \right) \frac{\langle :T_{VV}: \rangle_\omega}{A} f d\mathcal{V}_D - 4\pi \int_{U_0}^{U_2} \frac{\langle :T_{UV}: \rangle_\omega r^2}{A} f A dU. \end{aligned} \quad (51)$$

Since D is a normal domain, e.g. with respect to the V -axis for any $\mathcal{F}(V, U) \in \mathcal{C}^\infty(\mathcal{M})$, it holds that

$$\int_D \mathcal{F}(V, U) d\mathcal{V}_D = \int_{V_0}^{V_1} dV \int_{U_0}^{U_{\mathfrak{H}(V)}} \mathcal{F}(V, U) A r^2 dU, \quad (52)$$

where $U_{\mathfrak{H}}(V)$ is the solution of $2r(V, U) - m(V, U) = 0$.

Our aim is to prove now that $\Delta_h M$ is strictly negative using eq. (51). Notice that the first terms in the right-hand side gives a negative contribution to $\Delta_h M$ because $\partial_V r > 0$ outside the apparent horizon. Hence, we isolate all the integrals of this form and we tame the effects of the other choosing carefully the function f . Actually, we want to find a function $f(V, U) > 0$ such that all these unwanted terms in eq. (51) vanish. Let us define the following two functions

$$\beta(V, U) \doteq \frac{2}{r} \partial_V r \frac{\langle :T_{UU}: \rangle_\omega}{A}, \quad \alpha(V, U) \doteq \frac{8A}{3r^3} \partial_V r R_{UV} - \int \partial_V \beta(V, U) dU. \quad (53)$$

The U -derivative of f is fixed in such a way to cancel the volume integral whose integrand is proportional to $\langle :T_{VV}: \rangle_\omega$, namely it must be a solution of the equation

$$\left(-\frac{\partial_U f}{f} + \lambda \frac{2\partial_V r}{r} \frac{\langle :T_{UU}: \rangle_\omega}{A} \right) = 0.$$

Hence, we get

$$f(V, U) = f_0(V) \exp\left(\lambda \int \beta(V, U) dU\right),$$

where $f_0(V)$ is an integration constant. Plugging this function $f(V, U)$ in the contributions of eq. (51) whose integrands are proportional to $\langle :T_{UV}: \rangle_\omega$ and defining

$$\gamma_1(V) = \int_{U_0}^{U_{\mathfrak{H}}(V)} \frac{\langle :T_{UV}: \rangle_\omega}{A} \mathcal{G}(V, U) \exp\left(\lambda \int \beta(V, U) dU\right) Ar^2 dU, \quad (54)$$

$$\gamma_2(V) = \int_{U_0}^{U_{\mathfrak{H}}(V)} \frac{\langle :T_{UV}: \rangle_\omega}{A} \exp\left(\lambda \int \beta(V, U) dU\right) Ar^2 dU, \quad (55)$$

where

$$\mathcal{G}(V, U) \doteq \partial_V \log(A^{-1}r^2) + \lambda \alpha(V, U) = \partial_V \log(A^{-1}r^2) + \lambda \frac{4A}{3r^2} \frac{2}{r} \partial_V r R_{UV} - \lambda \int \partial_V \beta(V, U) dU, \quad (56)$$

we obtain that

$$\begin{aligned} & \int_D \left(-\frac{\partial_V f}{f} + \partial_V \log(A^{-1}r^2) + \lambda \frac{8A}{3r^3} \partial_V r R_{UV} \right) \frac{\langle :T_{UV}: \rangle_\omega}{A} f d\mathcal{V}_D = \\ & = \int_D \left(-\frac{\partial_V f_0}{f_0} + \partial_V \log(A^{-1}r^2) + \lambda \alpha(V, U) \right) \frac{\langle :T_{UV}: \rangle_\omega}{A} f d\mathcal{V}_D = \int_{V_0}^{V_1} f_0(V) \left(-\frac{\partial_V f_0}{f_0} \gamma_2(V) + \gamma_1(V) \right) dV. \end{aligned}$$

This integral vanishes e.g. for $f_0(V) = C \exp\left(\int \frac{\gamma_1(V)}{\gamma_2(V)} dV\right)$, where C is a strictly positive integration constant. Therefore,

$$f(V, U) = \exp\left(\int \frac{\gamma_1(V)}{\gamma_2(V)} dV + \lambda \int \beta(V, U) dU\right) \quad (57)$$

is a well-defined strictly positive smooth function in D , constructed by the functions given in eqs. (53) to (56), such that $\Delta_h M < 0$. With this choice of $f(V, U)$, $h(V, U)$ given in eq. (49) is also positive and smooth. Hence, it is bounded from below in $\delta\mathcal{H}$, so $0 > \Delta_h M \geq C' \Delta M$, where C' is another strictly positive constant, and the proof of the Theorem holds.

References

- [AV10] G. Abreu and M. Visser, “*Kodama time: Geometrically preferred foliations of spherically symmetric spacetimes,*” Phys. Rev. D **82**, 044027 (2010) [[10.1103/PhysRevD.82.044027](#)]
- [AGCF20] P. R. Anderson, S. Gholizadeh Siahmazgi, R. D. Clark, and A. Fabbri, “*Method to compute the stress-energy tensor for a quantized scalar field when a black hole forms from the collapse of a null shell,*” Phys. Rev. D **102**(12), 125035 (2020) [[10.1103/PhysRevD.102.125035](#)]
- [AHS95] P. R. Anderson, W. A. Hiscock, and D. A. Samuel, “*Stress-energy tensor of quantized scalar fields in static spherically symmetric spacetimes,*” Phys. Rev. D **51**, 4337 (1995) [[10.1103/PhysRevD.51.4337](#)]
- [AK02] A. Ashtekar and B. Krishnan, “*Dynamical horizons: Energy, angular momentum, fluxes and balance laws,*” Phys. Rev. Lett. **89**, 261101 (2002) [[10.1103/PhysRevLett.89.261101](#)]
- [AK03] A. Ashtekar and B. Krishnan, “*Dynamical horizons and their properties,*” Phys. Rev. D **68**, 104030 (2003) [[10.1103/PhysRevD.68.104030](#)]
- [AK04] A. Ashtekar and B. Krishnan, “*Isolated and dynamical horizons and their applications,*” Liv. Rev. Rel. **7**, 10 (2004) [[10.12942/lrr-2004-10](#)]
- [APR11a] A. Ashtekar, F. Pretorius, and F. M. Ramazanoglu, “*Evaporation of 2-Dimensional Black Holes,*” Phys. Rev. D **83**, 044040 (2011) [[10.1103/PhysRevD.83.044040](#)]
- [APR11b] A. Ashtekar, F. Pretorius, and F. M. Ramazanoglu, “*Surprises in the Evaporation of 2-Dimensional Black Holes,*” Phys. Rev. Lett. **106**, 161303 (2011) [[10.1103/PhysRevLett.106.161303](#)]
- [AH99] M. C. Ashworth and S. A. Hayward, “*Boundary terms and Noether current of spherical black holes,*” Phys. Rev. D **60**, 084004 (1999) [[10.1103/PhysRevD.60.084004](#)]
- [Bal84] R. Balbinot, “*Hawking radiation and the back reaction—a first approach,*” Class. Quant. Grav. **1**(5), 573 (1984) [[10.1088/0264-9381/1/5/010](#)]
- [BB89] R. Balbinot and A. Barletta, “*The backreaction and the evolution of quantum black holes,*” Class. Quant. Grav. **6**(2), 195 (1989) [[10.1088/0264-9381/6/2/013](#)]
- [Bar81] J. M. Bardeen, “*Black holes do evaporate thermally,*” Phys. Rev. Lett. **46**(6), 382 (1981) [[10.1103/PhysRevLett.46.382](#)]
- [Bar14] J. M. Bardeen, “*Black hole evaporation without an event horizon,*” (2014) [[arXiv:1406.4098 \[gr-qc\]](#)]
- [BD84] N. D. Birrell and P. C. W. Davies, *Quantum fields in curved space 7*. Cambridge University Press, 1984.
- [Brow18] P. J. Brown, C. J. Fewster, and E. A. Kontou, “*A singularity theorem for Einstein–Klein–Gordon theory,*” Gen. Rel. Grav. **50**, 121 (2018) [[10.1007/s10714-018-2446-5](#)]
- [BFK96] R. Brunetti, K. Fredenhagen, and M. Köhler, “*The microlocal spectrum condition and Wick polynomials of free fields on curved spacetimes,*” Commun. Math. Phys. **180**(3), 633 (1996) [[10.1007/s00220-003-0815-7](#)]
- [Can80] P. Candelas, “*Vacuum polarization in Schwarzschild space-time,*” Phys. Rev. D **21**, 2185 (1980) [[10.1103/PhysRevD.21.2185](#)]

- [CF77] S. M. Christensen and S. A. Fulling, “*Trace Anomalies and the Hawking Effect*,” Phys. Rev. D **15**, 2088 (1977) [[10.1103/PhysRevD.15.2088](#)]
- [Chr86a] D. Christodoulou, “*Global existence of generalized solutions of the spherically symmetric Einstein-scalar equations in the large*,” Commun. Math. Phys. **106**(4), 587 (1986) [[10.1007/BF01463398](#)]
- [Chr86b] D. Christodoulou, “*The problem of a self-gravitating scalar field*,” Commun. Math. Phys. **105**(3), 337 (1986) [[10.1007/BF01205930](#)]
- [Chr91] D. Christodoulou, “*The formation of black holes and singularities in spherically symmetric gravitational collapse*,” Commun. Pure Appl Math. **44**(3), 339 (1991) [[10.1002/cpa.3160440305](#)]
- [DT20] P. K. Dahal and D. R. Terno, “*Kerr-Vaidya black holes*,” Phys. Rev. D **102**, 124032 (2020) [[10.1103/PhysRevD.102.124032](#)]
- [DFU76] P. C. W. Davies, S. A. Fulling, and W. G. Unruh, “*Energy-momentum tensor near an evaporating black hole*,” Phys. Rev. D **13**, 2720 (1976) [[10.1103/PhysRevD.13.2720](#)]
- [DNVZZ07] R. Di Criscienzo, M. Nadalini, L. Vanzo, S. Zerbini, and G. Zoccatelli, “*On the Hawking radiation as tunneling for a class of dynamical black holes*,” Phys. Let. B **657**, 107 (2007) [[10.1016/j.physletb.2007.10.005](#)]
- [FK20] C. J. Fewster and E. A. Kontou, “*A new derivation of singularity theorems with weakened energy hypotheses*,” Class. Quant. Grav. **37**(6), 065010 (2020) [[10.1088/1361-6382/ab685b](#)]
- [FR03] C. J. Fewster and T. A. Roman, “*Null energy conditions in quantum field theory*,” Phys. Rev. D **67**, 044003 (2003) [[10.1103/PhysRevD.67.044003](#)] [Erratum: Phys.Rev.D 80, 069903 (2009)]
- [FV03] C. J. Fewster and R. Verch, “*Stability of quantum systems at three scales: Passivity, quantum weak energy inequalities and the microlocal spectrum condition*,” Commun. Math. Phys. **240**, 329 (2003) [[10.1007/s00220-003-0884-7](#)]
- [Fla97] E. E. Flanagan, “*Quantum inequalities in two-dimensional Minkowski space-time*,” Phys. Rev. D **56**, 4922 (1997) [[10.1103/PhysRevD.56.4922](#)]
- [FR95] L. H. Ford and T. A. Roman, “*Averaged energy conditions and quantum inequalities*,” Phys. Rev. D **51**, 4277 (1995) [[10.1103/PhysRevD.51.4277](#)]
- [FR96] L. H. Ford and T. A. Roman, “*Averaged energy conditions and evaporating black holes*,” Phys. Rev. D **53**, 1988 (1996) [[10.1103/PhysRevD.53.1988](#)]
- [FR97] L. H. Ford and T. A. Roman, “*Restrictions on negative energy density in flat space-time*,” Phys. Rev. D **55**, 2082 (1997) [[10.1103/PhysRevD.55.2082](#)]
- [FH90] K. Fredenhagen and R. Haag, “*On the derivation of Hawking radiation associated with the formation of a black hole*,” Commun. Math. Phys. **127**(2), 273 (1990) [[10.1007/BF02096757](#)]
- [FKK20] B. Freivogel, E. A. Kontou, and D. Krommydas, “*The return of the singularities: Applications of the smeared null energy condition*,” (2020) [[arXiv: 2012.11569 \[gr-qc\]](#)]
- [FD76] S. A. Fulling and P. C. W. Davies, “*Radiation from a moving mirror in two dimensional space-time: conformal anomaly*,” Proc. Roy. Soc. A **348**(1654), 393 (1976) [[10.1098/rspa.1976.0045](#)]
- [GP09] J. B. Griffiths and J. Podolsky, *Exact Space-Times in Einstein’s General Relativity* Cambridge Monographs on Mathematical Physics. Cambridge University Press, Cambridge, 2009. [[10.1017/CBO9780511635397](#)]

- [Haw68] S. W. Hawking, “*Gravitational radiation in an expanding universe*,” J. Math. Phys. **9**, 598 (1968) [[10.1063/1.1664615](#)]
- [Haw74] S. W. Hawking, “*Black hole explosions?*” Nature **248**(5443), 30 (1974) [[10.1038/248030a0](#)]
- [Haw75] S. W. Hawking, “*Particle creation by black holes*,” Commun. Math. Phys. **43**(3), 199 (1975) [[10.1007/BF02345020](#)]
- [HDVNZ09] S. Hayward, R. Di Criscienzo, L. Vanzo, M. Nadalini, and S. Zerbini, “*Local Hawking temperature for dynamical black holes*,” Class. Quant. Grav. **26**, 062001 (2009) [[10.1088/0264-9381/26/6/062001](#)]
- [Hay93] S. A. Hayward, “*General laws of black-hole dynamics*,” Phys. Rev. D **49**, 6467 (1994)
- [Hay96] S. A. Hayward, “*Gravitational energy in spherical symmetry*,” Phys. Rev. D **53**(6), 1938 (1996) [[10.1103/PhysRevD.53.1938](#)]
- [Hay98] S. A. Hayward, “*Unified first law of black hole dynamics and relativistic thermodynamics*,” Class. Quant. Grav. **15**, 3147 (1998) [[10.1088/0264-9381/15/10/017](#)]
- [Hay00] S. A. Hayward, “*Black holes: New horizons*,” In “*9th Marcel Grossmann Meeting on Recent Developments in Theoretical and Experimental General Relativity, Gravitation and Relativistic Field Theories (MG 9)*,” pages 568–580. 2000. [[10.1088/0264-9381/26/6/062001](#)]
- [HMA99] S. A. Hayward, S. Mukohyama, and M. C. Ashworth, “*Dynamic black-hole entropy*,” Phys. Let. A **256**(5), 347 (1999) ISSN 0375-9601. [[10.1016/S0375-9601\(99\)00225-X](#)]
- [HW01] S. Hollands and R. M. Wald, “*Local Wick polynomials and time ordered products of quantum fields in curved space-time*,” Commun. Math. Phys. **223**, 289 (2001) [[10.1007/s002200100540](#)]
- [HW02] S. Hollands and R. M. Wald, “*Existence of local covariant time ordered products of quantum fields in curved space-time*,” Commun. Math. Phys. **231**, 309 (2002) [[10.1007/s00220-002-0719-y](#)]
- [HW04] S. Hollands and R. M. Wald, “*Conservation of the stress tensor in interacting quantum field theory in curved spacetimes*,” Rev. Math. Phys. **17**, 227 (2005) [[10.1142/S0129055X05002340](#)]
- [HW15] S. Hollands and R. M. Wald, “*Quantum fields in curved spacetime*,” Physics Reports **574**, 1 (2015) [????]
- [How84] K. W. Howard, “*Vacuum T_{μ}^{ν} in Schwarzschild spacetime*,” Phys. Rev. D **30**, 2532 (1984) [[10.1103/PhysRevD.30.2532](#)]
- [KW91] B. S. Kay and R. M. Wald, “*Theorems on the uniqueness and thermal properties of stationary, nonsingular, quasifree states on spacetimes with a bifurcate Killing horizon*,” Phys. Rep. **207**(2), 49 (1991) ISSN 0370-1573. [[10.1016/0370-1573\(91\)90015-E](#)]
- [Kod80] H. Kodama, “*Conserved energy flux for the spherically symmetric system and the backreaction problem in the black hole evaporation*,” Prog. Theo. Phys. **63**(4), 1217 (1980) ISSN 0033-068X. [[10.1143/PTP.63.1217](#)]
- [KS20] E. A. Kontou and K. Sanders, “*Energy conditions in general relativity and quantum field theory*,” Class. Quant. Grav. **37**(19), 193001 (2020) [[10.1088/1361-6382/ab8fcf](#)]
- [KF93] C. I. Kuo and L. H. Ford, “*Semiclassical gravity theory and quantum fluctuations*,” Phys. Rev. D **47**, 4510 (1993) [[10.1103/PhysRevD.47.4510](#)]

- [KPV21] F. Kurpicz, N. Pinamonti, and R. Verch, “*Temperature and entropy-area relation of quantum matter near spherically symmetric outer trapping horizons,*” (2021) [[arXiv: 2102.11547 \[gr-qc\]](#)]
- [MPS20] P. Meda, N. Pinamonti, and D. Siemssen, “*Existence and uniqueness of solutions of the semi-classical einstein equation in cosmological models,*” (2020) [[arXiv: 2003.01815 \[gr-qc\]](#)]
- [MS64] C. W. Misner and D. H. Sharp, “*Relativistic equations for adiabatic, spherically symmetric gravitational collapse,*” *Phys. Rev.* **136**, B571 (1964) [[10.1103/PhysRev.136.B571](#)]
- [Mor02] V. Moretti, “*Comments on the stress-energy tensor operator in curved spacetime,*” *Commun. Math. Phys.* **232**(2), 189 (2003) [[10.1007/s00220-002-0702-7](#)]
- [Pag76] D. N. Page, “*Particle emission rates from a black hole: Massless particles from an uncharged, nonrotating hole,*” *Phys. Rev. D* **13**, 198 (1976) [[10.1103/PhysRevD.13.198](#)]
- [PW00] M. K. Parikh and F. Wilczek, “*Hawking radiation as tunneling,*” *Phys. Rev. Lett.* **85**, 5042 (2000) [[10.1103/PhysRevLett.85.5042](#)]
- [SV08] J. Schlemmer and R. Verch, “*Local thermal equilibrium states and quantum energy inequalities,*” *Ann. Henri Poincare* **9**, 945 (2008) [[10.1007/s00023-008-0380-x](#)]
- [VAC11] L. Vanzo, G. Acquaviva, and R. Di Criscienzo, “*Tunnelling methods and Hawking’s radiation: achievements and prospects,*” *Class. Quant. Grav.* **28**, 183001 (2011) [[10.1088/0264-9381/28/18/183001](#)]
- [Wal77] R. M. Wald, “*The back reaction effect in particle creation in curved spacetime,*” *Commun. Math. Phys.* **54**(1), 1 (1977) [[10.1007/bf01609833](#)]
- [Wal78] R. M. Wald, “*Trace Anomaly of a Conformally Invariant Quantum Field in Curved Space-Time,*” *Phys. Rev. D* **17**, 1477 (1978) [[10.1103/PhysRevD.17.1477](#)]
- [Wal84] R. M. Wald, *General Relativity* Chicago Univ. Pr., Chicago, USA, 1984. [[10.7208/chicago/9780226870373.001.0001](#)]
- [Wal95] R. M. Wald, *Quantum Field Theory in Curved Space-Time and Black Hole Thermodynamics* Chicago Lectures in Physics. University of Chicago Press, Chicago, IL, 1995.
- [Wal01] R. M. Wald, “*The thermodynamics of black holes,*” *Liv. Rev. Rel.* **4**, 6 (2001) [[10.12942/lrr-2001-6](#)]
- [WY91] R. M. Wald and U. Yurtsever, “*General proof of the averaged null energy condition for a massless scalar field in two-dimensional curved space-time,*” *Phys. Rev. D* **44**, 403 (1991) [[10.1103/PhysRevD.44.403](#)]