

Constraints on running vacuum models with the baryon-to-photon ratio

Hao Yu ^{a,1}, Ke Yang ^{b,2}, Jin Li ^{a,3}

^a Physics Department, Chongqing University, Chongqing 401331, China

^b School of Physical Science and Technology, Southwest University, Chongqing 400715, China

Abstract. We study the influence of running vacuum on the baryon-to-photon ratio in running vacuum models (RVMs). When there exists non-minimal coupling between photons and other matter in the expanding universe, the energy-momentum tensor of photons is no longer conserved, but the energy of photons could remain conserved. We discuss the conditions for the energy conservation of photons in RVMs. The photon number density and baryon number density, from the epoch of photon decoupling to the present day, are obtained in the context of RVMs by assuming that photons and baryons can be coupled to running vacuum, respectively. Both cases cause the baryon-to-photon ratio to no longer be a constant. However the evolution of the baryon-to-photon ratio is strictly constrained by observations. It is found that if the dynamic term of running vacuum is indeed coupled to photons or baryons, the coefficient of the dynamic term must be extremely small, which is unnatural. Therefore, our study basically rules out the possibility that running vacuum is coupled to photons or baryons in RVMs.

¹yuhaocd@cqu.edu.cn

²keyang@swu.edu.cn

³cqujinli1983@cqu.edu.cn, corresponding author

Contents

1	Introduction	1
2	Running Vacuum Models	3
3	Energy-momentum tensor conservation and energy conservation	5
3.1	Conservation in a toy universe	6
3.2	Conservation in a realistic universe	8
4	Photons in RVMs	11
5	Baryons in RVMs	13
6	The effect of running vacuum on the baryon-to-photon ratio	16
7	Discussions and conclusions	19

1 Introduction

The Big Bang theory is by far the most accepted theory explaining the birth, evolution and future of the universe. Since the measurements of the hydrogen abundance and helium abundance in the universe are exactly consistent with the predictions in the standard Big Bang nucleosynthesis (BBN) model, many cosmologists regard it as one of the strongest evidences¹ to support the Big Bang theory (see Refs. [1–6] for a review). One of the ingenuities of the Big Bang theory is that it proposes that the abundance of light elements in the universe is only related to one basic parameter, the baryon-to-photon ratio η . Therefore, the photon and baryon number densities are of interest and significant for cosmology. In the standard cosmological model (i.e., Λ CDM model), they dilute, as the universe expands, at the same dilution rate after BBN, so the present-day value of the baryon-to-photon ratio is a constant and was formed at the end of BBN [5, 7–9].

In different epochs of the universe, η can be determined independently from different perspectives. For the BBN epoch one can derive η based on the observational data on the abundance of primordial chemical elements [5–11], and for the primordial recombination epoch one can get η by analysing the anisotropy of the cosmic microwave background (CMB) radiation [12–15]. Whether these independent η 's evolving with the universe are in line with the current observed η_0 [6, 16–19], is the pivotal to judging the viability of the corresponding hypotheses and calculations in the Λ CDM model. In practice, the photon and baryon number densities are affected by many factors related to the model of the universe, especially modified gravity theories and the components of the universe. If dark matter could annihilate or decay (see Refs. [20–26] and references therein), then the baryon-to-photon ratio will deviate from the constant predicted in the Λ CDM model. But from BBN to the present, the degree of the deviation is almost negligible ($\frac{\Delta\eta}{\eta} \leq 10^{-5}$) [27, 28]. The impact of dark energy on η is mainly due to the possibility that dark energy could decay into CMB, but the decay

¹The other two evidences are the cosmic microwave background radiation and observed expansion of the universe.

rate of dark energy must be sufficiently small to be in accordance with observations [29–33]. Considering low-amplitude and large-scale fluctuations, the evolution of η may change with the region of the universe [34]. In addition, η is influenced by varying fundamental “constants” as well [35–39]. The variation of Newton’s constant could modify nucleosynthesis codes, which can be used for restraining Brans-Dicke theory and scalar-tensor theories [39, 40]. A time-varying cosmological “constant” is also able to effect the early universe and primordial nucleosynthesis [2, 41, 42]. Although most of these works do not specify the extent of the influence of varying fundamental “constants” on η , it is not hard to infer that, in these universe models, either η is a constant different from the one in the Λ CDM model, or it will inevitably evolve over time. Overall, the prediction of η is a powerful tool to verify the feasibility of various universe models.

In this work, we study the effect of varying vacuum on η in a class of cosmological models, which are so called running vacuum models (RVMs), see Refs. [43–46] and references therein. There is a long history of investigations of dynamic vacuum [47–52], but most dynamic vacuum models lack theoretical support. The proposal of RVMs is mainly attempting to connect the vacuum energy density with the quantum field theory in curved space-time [53–55]. In recent years, the research and development of RVMs have shown that running vacuum is better than a single cosmological constant in cosmology global picture [45, 46, 56, 57], observational data [51, 58–60], and alleviating tensions in cosmology [58, 61–64]. Recently, it is found that RVMs, comparing with the Λ CDM model, even possess more superior thermodynamic characteristic [65–67].

In RVMs, vacuum is actually non-minimally coupled to matter fields. According to the regular definition of the energy-momentum tensor, it can be found that $\nabla_\mu T_m^{\nu\mu} \neq 0$, where $T_m^{\mu\nu}$ is the energy-momentum tensor of all matter fields in RVMs. According to Tiberiu Harko et al. [68, 69], if the energy-momentum tensor of a matter field is not conserved, the field equation can be in comparison with the first law of thermodynamics of open systems. Therefore, $\nabla_\mu T_m^{\nu\mu} \neq 0$, at the macroscopic level, can be explained as the production (or annihilation) of corresponding particles. If photons in RVMs are coupled to vacuum, the redshift of a single photon in the expanding universe will be modified comparing with its “free expansion” in the Λ CDM model. We wonder whether such expansion of photons in RVMs could solve the problem of energy non-conservation in the Λ CDM model ² [72–74]. Furthermore, the impact of running vacuum on CMB may lead to new estimations on the size and age of the observable universe at certain epochs. Similarly, if running vacuum is coupled to baryons and the decay (or absorption) of vacuum is a baryon non-conserving process, there could be baryon production (or annihilation) even after the Hadron epoch. Since we have no any experiments and observations implying that the baryon non-conserving process can occur in the context of low energy and weak gravitational field, in order not to contradict the existing experiments and observations, the occurrence rate of running vacuum decaying into baryons (or the contrary process) must be very low. Therefore, it imposes strict constraints on the dynamic terms in the energy density of running vacuum.

Finally, according to the current η_0 [17–19], one can, in theory, derive the value of η at any epochs through assuming that vacuum is coupled to photons or baryons. However,

²Since the evolution of CMB radiation satisfies $\nabla_\mu T_r^{\nu\mu} = 0$ in the Λ CDM model (it is also known as black-body radiation, see e.g. [70, 71] and references therein), the total energy of CMB decreases with the expansion of the universe even though photons have no interaction with other matter fields. From this point of view, we have to define subtly, with only one tiny cosmological constant, the energy of space-time to guarantee energy conservation of the entire universe.

before photon decoupling (which is closely related to the end of recombination), photons are not only coupled to running vacuum, but also baryon nuclei, which makes our discussion about the impact of running vacuum on the baryon-to-photon ratio complicated. So, in this work we only discuss the evolution of the universe after photon decoupling. By estimating the value of η at the epoch of photon decoupling in RVMs and comparing it with the latest result from the anisotropy of CMB [12–15], new constraints on the dynamics term in RVMs can be obtained, which is independent on previous research on RVMs [55, 60, 63, 75].

The paper is organized as follows: Sec. 2 is devoted to the review of RVMs. Next, we analyse the energy-momentum tensor conservation and energy conservation of photons in a toy universe model and then extend our analysis to a more realistic universe in Sec. 3. In Sec. 4, we calculate the particle number density of photons in RVMs, and the similar calculations on the particle number density of baryons are presented in Sec. 5. Then, based on Secs. 4 and 5, we study the effect of running vacuum on the baryon-to-photon ratio in Sec. 6. The last part, Sec. 7, is a summary and discussion on our results and possible future work.

2 Running Vacuum Models

The background space-time of the homogeneous and isotropic universe can be described by Friedmann-Robertson-Walker metric

$$ds^2 = -dt^2 + a(t)^2(dx^2 + dy^2 + dz^2), \quad (2.1)$$

where $a(t)$ is the scale factor, the curvature of the universe has been set as zero, and we employ natural units $c = \hbar = 1$.

In the framework of RVMs, the Friedmann equations are similar to the ones in the Λ CDM model:

$$3H^2 = \kappa^2 \rho_t, \quad (2.2)$$

$$3H^2 + 2\dot{H} = -\kappa^2 p_t, \quad (2.3)$$

where $H = \dot{a}/a$ is the Hubble rate. The parameters ρ_t and p_t represent the total energy density and pressure of all components of the universe, respectively. Since the dominant component of the universe is unfixed and some of the components can be ignored at particular epochs, the specific forms of ρ_t and p_t rely on the accuracy requirement and the epoch of the universe. The basic principles and discussions of RVMs can be referred to the literature mentioned above (see Refs. [44, 46] for a review). In this work, we consider a class of well described and extensively studied RVMs, in which the vacuum energy density ρ_Λ “runs” in the form of a function of the Hubble rate based on a renormalization group equation:

$$\rho_\Lambda(H) = \frac{\Lambda(H)}{\kappa^2} = \frac{3}{\kappa^2} \left(c_0 + \nu H^2 + \alpha \frac{H^4}{H_1^2} \right). \quad (2.4)$$

Note that c_0 is a constant with the dimension of energy squared in natural units. Since the current observed value of Hubble rate is so small, to be consistent with observations, c_0 should approximately equal $\frac{1}{3}\Lambda_0$ (Λ_0 is the value of the cosmological constant in the Λ CDM model [18, 76]). The argument H_1 can be chosen as the Planck scale or at least below the Planck scale. The dimensionless coefficients ν and α have been confined to a

very narrow range by considering various cosmological phenomena. According to previous research [55, 60, 63, 75], ν and α should satisfy $|\nu| < 10^{-3}$ and $\alpha \ll 1$.

It is believed that νH^2 has been already much larger than $\alpha \frac{H^4}{H_1^2}$ after BBN [56]. In practice, the dynamic characteristics of vacuum energy is mainly manifested in νH^2 after BBN, and therefore one can neglect $\alpha \frac{H^4}{H_1^2}$ without conflicting with observations when the problem to be studied has no connection with the early universe. Since our following study does not involve inflation and prior epochs, it is reasonable to discard the subordinate term $\alpha \frac{H^4}{H_1^2}$ in Eq. (2.4). Hence it is easy to get analytical solutions of Eqs. (2.2) and (2.3) for the radiation-dominated universe by assuming that the main components of the universe are photons and vacuum energy. For such simplified universe model, Eqs. (2.2) and (2.3) are rewritten as

$$3H^2 = \kappa^2(\rho_\Lambda + \rho_r), \quad (2.5)$$

$$3H^2 + 2\dot{H} = -\kappa^2(p_\Lambda + p_r). \quad (2.6)$$

Since we have already reduced the vacuum energy density (2.4) as $\rho_\Lambda = \frac{3}{\kappa^2}(c_0 + \nu H^2)$, one can find directly from Eq. (2.5) that the energy density of photons can be written as an explicit function of the Hubble rate. And also if we express the energy density of photons as a function of the scale factor, it is found that

$$\rho_r(a) = \frac{3}{\kappa^2}(H^2 - \nu H^2 - c_0) = \rho_r(t_0)a^{-4(1-\nu)}, \quad (2.7)$$

where we have used the state equation of photons $\rho = 1/3\rho$ and $p_\Lambda = -\rho_\Lambda$. Note that the integration constant $\rho_r(t_0)$ is not necessarily equal to the currently observed power density of CMB radiation. This solution is only applicable to the radiation-dominated epoch, so the value of $\rho_r(t_0)$ should be decided by other cosmological processes in the early universe. Obviously, when $\nu = 0$, Eq. (2.7) degenerates into the formula satisfied by photons expanding freely with the universe in the Λ CDM model.

As for the matter-dominated epoch, one can assume the universe consists of dark matter, ordinary (baryonic) matter, and running vacuum. So, the Friedmann equations (2.2) and (2.3) are given as

$$3H^2 = \kappa^2(\rho_\Lambda + \rho_{dm} + \rho_b), \quad (2.8)$$

$$3H^2 + 2\dot{H} = -\kappa^2(p_\Lambda + p_{dm} + p_b). \quad (2.9)$$

For the sake of simplicity, we set the state equations of baryonic matter and dark matter as $p_b = p_{dm} = 0$. In view of the fact that baryonic matter and dark matter have the same state equation and similar energy-momentum tensor (which has only one non-zero component, i.e., energy density), we can regard them as a whole when solving the equations above. From Eqs. (2.8) and (2.9), it is clear that, the total energy density of baryonic matter and dark matter satisfies

$$\rho_{dm} + \rho_b = \frac{3}{\kappa^2}[H^2 - \nu H^2 - c_0] = [\rho_{dm}(t_0) + \rho_b(t_0)]a^{-3(1-\nu)}. \quad (2.10)$$

Here, $\rho_{dm}(t_0) + \rho_b(t_0)$ is the current total energy density of baryonic matter and dark matter. Apparently, Eq. (2.10) can also degenerate into the standard one in the Λ CDM model when $\nu = 0$.

3 Energy-momentum tensor conservation and energy conservation

In the Λ CDM model, photons in the universe travel freely after photon decoupling and the energy-momentum tensor of photons becomes conserved. Therefore, the total number of photons approximately remains as a constant, which combined with baryon number conservation after BBN leads us to deem that the baryon-to-photon ratio is unchanged after photon decoupling (some other views claim that it should be after BBN [5, 7–9]). The conservation of photon number is also based on the requirement that CMB radiation could be regarded as black-body radiation, which therefore results in non-conservation of the internal energy of photons. In the Λ CDM model, we can not explain the energy loss of photons due to the expansion of space-time [72–74] but baryons have no such problem. In this section, we analyse whether the problem can be solved in RVMs.

The internal energy and the number of photons can be expressed as

$$U_r = \rho_r V = \frac{\pi^2 k^4}{15c^3 \hbar^3} V T^4, \quad (3.1)$$

$$N_r = \frac{2k^3 \zeta(3)}{\pi^2 c^3 \hbar^3} V T^3, \quad (3.2)$$

where k is Boltzmann's constant and $\zeta(n)$ is the Riemann zeta function. In the Λ CDM model, the energy density of CMB satisfies $\rho_r \sim a(t)^{-4}$. Comparing $\rho_r \sim a(t)^{-4}$ with Eq. (3.1), it is found that the temperature of CMB is proportional to the inverse of the scale factor ($T \sim a(t)^{-1}$). Thus photon number density satisfies $n_r = N_r/V \sim T^3 \sim a(t)^{-3}$. Because of $V \sim a(t)^3$, N_r is indeed a constant and $U_r \sim T \sim a(t)^{-1}$ decreases with the expansion of the universe. Note that $\rho_r \sim a(t)^{-4}$ is derived from the conservation of the energy-momentum tensor ($\nabla_\nu T_r^{\nu\mu} = 0$). Therefore, in the Λ CDM model, the energy-momentum tensor conservation of photons ensures the conservation of photon number but leads to a decrease in the internal energy of photons. The energy lost by photons is usually considered to enter the gravitational field or background vacuum energy³.

Now let us analyse which factors are directly related to the energy loss of photons while the total number of photons remains constant ($\rho_r \sim a(t)^{-4}$). We consider a specific period of the universe from the scale factor $a(t_1)$ to $a(t_2)$ (the corresponding Hubble rate is H_1 and H_2 , respectively). Supposing that the temperature of CMB is T_1 at the moment $a(t_1)$, the number density of photons satisfies

$$n_r(t_1) = \frac{2k^3 \zeta(3)}{\pi^2 c^3 \hbar^3} T_1^3. \quad (3.3)$$

The internal energy and entropy of photons per unit volume (not per comoving volume) are, respectively,

$$\rho_r(t_1) = \frac{\pi^2 k^4}{15c^3 \hbar^3} T_1^4, \quad (3.4)$$

$$s_r(t_1) = \frac{4\pi^2 k^4}{45c^3 \hbar^3} T_1^3. \quad (3.5)$$

³For a single photon propagating in the expanding universe, both of its frequency and energy will be reduced due to the expansion of space-time. There is still controversy about the explanation on the energy loss of photons, because the definition of the energy conservation in general relativity is not complete. The energy-momentum tensor conservation is not completely equivalent to the energy conservation.

At the moment $a(t_2)$, the temperature of CMB drops to T_2 . Since the number of photons remains conserved, the number density can be written as

$$n_r(t_2) = \frac{2k^3\zeta(3)}{\pi^2c^3\hbar^3}T_2^3 = \frac{2k^3\zeta(3)}{\pi^2c^3\hbar^3}\frac{a(t_1)^3}{a(t_2)^3}T_1^3, \quad (3.6)$$

where the second equality is based on $a(t_2)^3T_2^3 = a(t_1)^3T_1^3 = \text{Const}$ (conservation of photon number). The energy lost by CMB per comoving volume is given by

$$\Delta u_r = \rho_r(t_2)a(t_2)^3 - \rho_r(t_1)a(t_1)^3 = \frac{\pi^2k^4}{15c^3\hbar^3}[T_2^4a(t_2)^3 - T_1^4a(t_1)^3]. \quad (3.7)$$

Since $a(t_2)^3T_2^3 = a(t_1)^3T_1^3$ and $T_2 < T_1$, we have $\Delta u_r < 0$, which means that as photons expand freely with the universe, the total internal energy of CMB decreases. In the Λ CDM model, we do not completely figure out where the energy flows. In addition, Eq. (3.7) manifests that if CMB stays in a state of freely expanding ($\nabla_\nu T_r^{\nu\mu} = 0$), the energy loss of CMB within a unit comoving volume is only related to its temperature gradient. Such consequence is comprehensible because photon gas per unit (comoving) volume is completely characterized by temperature [see Eqs. (3.3), (3.4) and (3.5)].

Next we study the expansion of photons in RVMs and the change in energy of photons in a unit comoving volume. We focus on the unit comoving volume because it is more practical than the unit volume. The variation of CMB within the observable universe over a period of time can be reflected in photons per unit comoving volume.

3.1 Conservation in a toy universe

We first introduce two special types of conservation, from the point of view of photons, in a toy universe model, which is only composed of photons and the aforementioned running vacuum. The toy universe model is similar to the radiation-dominated universe and it has pedagogical significance for our follow-up research on a realistic universe. The first case corresponds $\nu = 0$, and the second is a special value of ν guaranteeing that the total energy of CMB is constant. These two special situations represent two types of conservation: energy-momentum tensor conservation and energy conservation. We focus on the conditions for the energy conservation of CMB.

We first calculate the energy loss ratio of CMB for $\nu = 0$ in the context of the toy universe. The special case of $\nu = 0$ indicates that photons can “freely propagate” in the universe and the corresponding energy-momentum tensor satisfies $\nabla_\nu T_r^{\nu\mu} = 0$. Such property is similar to CMB in the Λ CDM model. Since $a(t_1)^3T_1^3 = a(t_2)^3T_2^3$, Eqs. (3.3) and (3.5) imply that the number and entropy of photons are constant. Supposing the temperature of CMB evolves from T_1 to T_2 with respective scale factors $a(t_1)$ and $a(t_2)$, $\nabla_\nu T_r^{\nu\mu} = 0$ results in $a(t_1)^3T_1^3 = a(t_2)^3T_2^3$. Therefore, the energy loss ratio of CMB (per unit comoving volume) from T_1 to T_2 is:

$$\Gamma_r = \frac{\rho_r(t_1)a(t_1)^3 - \rho_r(t_2)a(t_2)^3}{\rho_r(t_1)a(t_1)^3} = \frac{T_1 - T_2}{T_1}. \quad (3.8)$$

It can be found that if the temperature differential of CMB from $a(t_1)$ to $a(t_2)$ is large enough, the energy loss ratio could be close to 1. For example, photons started to “freely propagate” after photon decoupling in the Λ CDM model. It is generally recognized that photon decoupling occurred during recombination about 378,000 years after the Big Bang,

at a redshift of $z \sim 1100$, when the temperature of the universe was $T_1 \sim 3000$ K [6, 77]. If we take $T_2 = 2.73$ K, which is the current temperature of CMB [78–80], Γ_r is approximately equal to 1. Moreover, for a volume equal to the observable universe, the total energy loss of CMB from $T_1 \sim 3000$ K down to $T_2 = 2.73$ K is

$$\Delta U_r = \Gamma_r \frac{4\pi}{3} R_1^3 \rho_r(t_1) = \frac{4\pi}{3} \frac{T_1 - T_2}{T_1} R_1^3 * \frac{\pi^2 k^4}{15c^3 \hbar^3} T_1^4 \sim 1.6 * 10^{70} \text{ J}, \quad (3.9)$$

where $R_1 \sim 42$ million light-years is the radius of the observable universe at the end of recombination. In this work, we roughly assume that the epoch of photon decoupling is the end of recombination. Note that to get the final consequence, we have recovered the real values of all the physical quantities above.

Although $\nu = 0$ inevitably leads to the energy loss of CMB, the total energy of the toy universe can be conserved by choosing a proper c_0 . We also let $a(t_1)$ correspond to recombination and $a(t_2)$ be the current scale factor. The energy change of vacuum depends on the density of vacuum and the change in volume of the universe:

$$\Delta U_\Lambda = \frac{4\pi}{3} (R_2^3 - R_1^3) \rho_\Lambda, \quad (3.10)$$

where R_2 the radius of the observable universe and $\rho_\Lambda = \frac{3}{\kappa^2} c_0$. If $\Delta U_\Lambda = \Delta U_r$, ρ_Λ can be expressed as

$$\rho_\Lambda = \frac{T_1 - T_2}{T_1} \frac{T_2^3}{T_1^3 - T_2^3} \frac{\pi^2 k^4}{15c^3 \hbar^3} T_1^4 \sim \frac{\pi^2 k^4}{15c^3 \hbar^3} T_2^3 T_1 \sim 4.6 * 10^{-11} \text{ kg}/(\text{m} \cdot \text{s}^2), \quad (3.11)$$

which is less than the current density of vacuum (dark) energy $\rho_{\Lambda 0} \sim 6.3 * 10^{-10} \text{ kg}/(\text{m} \cdot \text{s}^2)$ [18] in the Λ CDM model. Eq. (3.11) manifests that to keep the energy gained by vacuum always equal to the energy lost by photons, vacuum can not be a constant. Regarding this, we will continue to discuss it below.

Next, let us pay attention to another case, i.e., the energy conservation of CMB. First we need to calculate the value of ν which could maintain the energy conservation of CMB. For the universe composed of photons and vacuum, the Friedmann equation is given by Eqs. (2.5) and (2.6). The energy density of photons satisfies (2.7). And the energy-momentum tensor conservation of all components leads to

$$\dot{\rho}_\Lambda + \dot{\rho}_r + 3H(p_\Lambda + p_r + \rho_\Lambda + \rho_r) = \dot{\rho}_\Lambda + \dot{\rho}_r + 4H\rho_r = 0. \quad (3.12)$$

With Eq. (2.7), we can get the value of ν directly from the point of view of photons. Since the energy conservation of photons requires

$$\rho_r(t_2)V(t_2) = \rho_r(t_1)V(t_1), \quad (3.13)$$

it is found that the parameter ν should take a value of $\frac{1}{4}$, which can be also derived from $a(t_2)^3 T_2^4 = a(t_1)^3 T_1^4$.

In order to verify whether the value of ν also guarantees the energy conservation of vacuum, we need to solve Eq. (3.17), obtain Hubble rate H as a function of $a(t)$, and then take it back to $\rho_\Lambda(H)$ to get $\rho_\Lambda(a(t))$. Finally, if $\rho_\Lambda(t_2)V(t_2) = \rho_\Lambda(t_1)V(t_1)$ is true for $\nu = \frac{1}{4}$, it proves that $\nu = \frac{1}{4}$ can indeed ensure the energy conservation of photons and vacuum,

respectively, while their total energy-momentum tensor satisfies the conservation condition (3.12). Combining Eq. (3.12) and $\rho_r(a) = \rho_r(t_0)a^{-4(1-\nu)}$, we have

$$\frac{d\rho_\Lambda}{da} + \frac{d\rho_r}{da} + 4\frac{\rho_r}{a} = \frac{d\rho_\Lambda}{da} + 4\nu\rho_r(t_0)a^{-5+4\nu} = 0. \quad (3.14)$$

The solution can be expressed as $\rho_\Lambda(a) = \frac{\nu}{1-\nu}\rho_r(t_0)a^{-4(1-\nu)} + \frac{3c_0}{1-\nu} = \frac{\nu}{1-\nu}\rho_r + \frac{3c_0}{1-\nu}$. Taking $\rho_\Lambda(a)$ into $\rho_\Lambda(t_2)V(t_2) = \rho_\Lambda(t_1)V(t_1)$ yields

$$\frac{\nu}{1-\nu}\rho_r(t_2)V(t_2) + \frac{3c_0}{1-\nu}V(t_2) = \frac{\nu}{1-\nu}\rho_r(t_1)V(t_1) + \frac{3c_0}{1-\nu}V(t_1). \quad (3.15)$$

Since $\rho_r(t_2)V(t_2) = \rho_r(t_1)V(t_1)$ and $V(t_2) \neq V(t_1)$, the premise of the above equation is that the constant term c_0 of vacuum energy is zero. Thereupon, the evolution of $\rho_\Lambda = \frac{\nu}{1-\nu}\rho_r(t_0)a^{-4(1-\nu)}$ along with the scale factor is exactly the same as the situation of photons [see Eq. (2.7)]. As far, we have proved that both of the energy conservation of photons and vacuum can be guaranteed in the toy universe model as long as $\nu = \frac{1}{4}$ and c_0 is vanishing.

Considering the temperature of CMB satisfies $a(t_2)^3T_2^4 = a(t_1)^3T_1^4$, the entropy change of CMB from recombination to the current day can be given as

$$\Delta S = \frac{4\pi^2k^4}{45c^3\hbar^3}[V(t_2)T_2^3 - V(t_1)T_1^3] > 0, \quad (3.16)$$

which conforms with the second law of thermodynamics. The entropy of the current CMB has increased by several orders of magnitude compared to its value at the end of recombination on account of $T_1 \gg T_2$.

Finally, we summarize the characteristics of CMB with regard to the two special kinds of conservation in above toy universe model. When CMB freely expands ($\nu = 0$ and so $\nabla_{\mu\nu}T_r^\mu = 0$), the corresponding photon number and entropy keep fixed. In this case, not only is the energy of CMB non-conserved, but also the total energy of the universe can not be conservative. So, in this toy universe model the energy-momentum tensor conservation of CMB inevitably causes the non-conservation of the total energy of the universe. For another special case that CMB maintains energy conservation, based on $\rho_r(t_2)V(t_2) = \rho_r(t_1)V(t_1)$, it can be found that the value of parameter ν must be $\frac{1}{4}$. The energy density of CMB evolves with the scale factor in the same way as dust. And if the constant term c_0 of vacuum energy equals zero, vacuum energy is also conserved. Therefore, $\rho_\Lambda = \frac{3}{4\kappa^2}H^2$ can guarantee both of the energy conservation of photons and vacuum. As for the rest of ν and c_0 , neither the energy-momentum tensor conservation of photons nor the energy conservation of photons can be guaranteed. And it can be seen from Eq. (3.15) that there is no other ν that keeps the total energy of the universe conserved.

Next, let us study a more realistic universe model and discuss whether RVMs could balance observational data and the energy conservation of the universe.

3.2 Conservation in a realistic universe

In this section, we study the energy-momentum tensor conservation and energy conservation of CMB in a realistic universe with running vacuum. We suppose that vacuum is only coupled to photons and the other components of universe satisfy their own energy-momentum tensor conservation and also energy conservation.

In the realistic universe model, if there is interaction ⁴ between photons and vacuum, it should start when photons emerged in the universe (the temperature of the universe was about $T_e \sim 10^{16}$ K at the end of electroweak epoch). There should also be a cutoff temperature T_c , at which photons and vacuum are decoupled. At present, we can not evaluate T_c from the first principle or existing knowledge, so we temporarily deem that T_c just need satisfy $0 < T_c < T_e$. Here, we suppose that the cutoff temperature satisfies $0 < T_c < T_0 = 2.73$ K, which indicates there is still interaction between photons and vacuum.

We have already mentioned that photons “freely propagate” after the recombination epoch. Therefore, when studying the properties of the interaction between photons and running vacuum, we only consider the period of the universe after recombination, which helps to exclude the interaction between photons and other substances. From the end of recombination to the current day, the realistic universe is no longer dominated by radiation but by dust (or baryonic matter), dark matter and vacuum energy (or dark energy). All the components of the universe are not negligible except for neutrinos, so the proper Friedmann equation should be given as

$$3H^2 = \kappa^2(\rho_\Lambda + \rho_{dm} + \rho_b + \rho_r), \quad (3.17)$$

$$3H^2 + 2\dot{H} = -\kappa^2(p_\Lambda + p_{dm} + p_b + p_r). \quad (3.18)$$

Hereafter, we just set $\kappa^2 = 1$ for simplicity. For the dust and dark matter with a vanishing pressure, energy conservation is equivalent to energy-momentum tensor conservation ($\nabla_{\mu\nu} T_{(b, dm)}^\nu = 0 \rightarrow \dot{\rho}_{(b, dm)} + 3H\rho_{(b, dm)} = 0 \rightarrow V\rho_{(b, dm)} \equiv \text{Const}$). Therefore, the total energy-momentum tensor conservation of photons and vacuum is still Eq. (3.12), which yields the energy density of photons as

$$\begin{aligned} \rho_r(a) &= \rho_{r0} a^{-4(1-\nu)} + \frac{3\nu[\rho_b(t_0) + \rho_{dm}(t_0)]}{1-4\nu} a^{-3} & (\nu \neq \frac{1}{4}) \\ \rho_r(a) &= \rho_r(t_0) a^{-3} + \frac{3[\rho_b(t_0) + \rho_{dm}(t_0)]}{4} a^{-3} \log a & (\nu = \frac{1}{4}). \end{aligned} \quad (3.19)$$

Note that $\rho_b(t_0)$ and $\rho_{dm}(t_0)$ are the current values of baryon density and dark matter density, respectively. $\rho_r(t_0)$ is the current energy density of CMB, but $\rho_{r0} \neq \rho_r(t_0)$, which is decided by $\rho_r(t_0)$:

$$\rho_{r0} = \rho_r(t_0) - \frac{3\nu[\rho_b(t_0) + \rho_{dm}(t_0)]}{1-4\nu}. \quad (3.20)$$

Combining Eqs. (3.12) and (3.19), we have

$$\frac{d\rho_\Lambda}{da} + \frac{d\rho_r}{da} + 4\frac{\rho_r}{a} = \frac{d\rho_\Lambda}{da} + 4\nu\rho_{r0}a^{-4(1-\nu)-1} + \frac{3\nu[\rho_b(t_0) + \rho_{dm}(t_0)]}{1-4\nu}a^{-4} = 0. \quad (3.21)$$

⁴The energy density of vacuum being a cosmological constant does not mean that vacuum can not interact with photons or other components of the universe. With the expansion of space-time, the total energy of vacuum within the observable universe is increasing although the energy density of vacuum is a constant. Therefore, it is entirely reasonable to believe that the increased energy of vacuum is caused by the interaction between vacuum and other substances.

The solution of vacuum density can be expressed as

$$\begin{aligned}
\rho_\Lambda(a) &= \frac{\nu}{1-\nu}\rho_{r0}a^{-4(1-\nu)} + \frac{\nu[\rho_b(t_0) + \rho_{dm}(t_0)]}{1-4\nu}a^{-3} + \frac{3}{1-\nu}c_0 \quad (\nu \neq \frac{1}{4} \text{ and } \nu \neq 1) \\
\rho_\Lambda(a) &= H_0^2 a^{-3} + \frac{1}{4}[\rho_b(t_0) + \rho_{dm}(t_0)]a^{-3} \log a + 4c_0 \quad (\nu = \frac{1}{4}) \\
\rho_\Lambda(a) &= -\frac{\rho_b(t_0) + \rho_{dm}(t_0)}{3}a^{-3} - 4\rho_{r0} \log a \quad (\nu = 1),
\end{aligned} \tag{3.22}$$

where $H_0^2 = \frac{1}{3}[\rho_b(t_0) + \rho_{dm}(t_0) + \rho_r(t_0)]$. For the solution with $\nu = 1$, we have $\rho_{r0} = -3c_0$.

With the solutions, we can study the two special types of conservation of photons in the realistic universe. When $\nu = 0$ (the energy-momentum tensor conservation of photons), RVMs revert to the Λ CDM model and photons could “freely expand” after photon decoupling. The scale factor $a(t_{re})$ of the universe at the end of recombination can be estimated by the fact that freely propagating photons satisfy $a(t)^3 T^3 = \text{Const}$. Since T_{re} is about 3000 K and $a(t_{re})^3 T_{re}^3 = a(t_0)^3 T_0^3$, according to the current radius of the observable universe ($R_0 \sim 46$ billion light-years), the radius of the universe corresponding to the observable universe at the end of recombination epoch is $R_{re} \sim 42$ million light-years. The radius is independent on the components of the universe and so is consistent with the result (see R_1 in the previous section) obtained in the toy universe model. Although the estimation on R_{re} is rough, it is still pretty much the same as the result obtained in the Λ CDM model, and such precision has no impact on our subsequent discussions. Setting the scale factor of the current universe as $a(t_0) = 1$ yields $a(t_{re}) \sim 9.1 * 10^{-4}$.

Note that when $\nu = 0$, the solution of ρ_r is consistent with the one in the Λ CDM model and also the one in the toy universe model, so we can use directly the previous calculations and results. It is shown that for the process that the temperature of CMB drops from 3000 K to 2.73 K, the energy loss of the observable universe is still Eq. (3.9). In the same process, take into account the current value of vacuum ($\rho_{\Lambda 0} \sim 6.3 * 10^{-10} \text{ kg}/(\text{m} \cdot \text{s}^2)$ [18]), the energy change of vacuum is

$$\Delta U_\Lambda = \frac{4\pi}{3} (R_0^3 - R_{re}^3) \rho_{\Lambda 0} \sim 2.2 * 10^{71} \text{ J}. \tag{3.23}$$

Comparing Eqs. (3.9) and (3.23), one can find that these two values are not accordant and the gap is more than tenfold, that is to say, the energy lost by photons is only about one-tenth of the energy obtained by vacuum. The previous toy universe model indicates that when photons expand freely, the total energy of the universe can not be maintained. It now appears that the conclusion also applies to the Λ CDM model. Moreover, since the energy-momentum tensor conservation of photons also is equivalent to photon number conservation, the entropy of CMB is also unchanging, which is about $10^{89} k$ [81, 82]. In the Λ CDM model, we can not explain why the entropy of CMB is so large [81].

In the toy universe model, we found that RVMs with $\nu = \frac{1}{4}$ and $c_0 = 0$ can conserve both of photon energy and vacuum energy, so the total energy of the universe is conserved. In the realistic universe, let us check if there are such ν and c_0 . The energy conservation of CMB signifies $a(t_2)^3 T_2^4 = a(t_1)^3 T_1^4$, with which the radius of the observable universe at the end of recombination can be obtained as $R_{re} \sim 4.1$ million light years. And then the corresponding scale factor is $a(t_{re}) \sim 8.8 * 10^{-5}$. Note that the influence of running vacuum on the scale factor is still within the acceptable range [83, 84].

Following the steps above, one may get the value of ν directly from Eq. (3.19). Taking Eqs. (3.19) and (3.22) into $\rho_r(t_{re})V(t_{re}) = \rho_r(t_0)V(t_0)$, we have

$$\begin{aligned} a(t_{re})^{-1+4\nu} &= a(t_0)^{-1+4\nu} & (\nu \neq \frac{1}{4}) \\ \log[a(t_{re})] &= \log[a(t_0)] & (\nu = \frac{1}{4}). \end{aligned} \quad (3.24)$$

Unfortunately, for the solution with $\nu \neq \frac{1}{4}$ [see Eq. (3.19)], the condition guaranteeing the energy conservation of photons is exactly $\nu = \frac{1}{4}$, while the solution with $\nu = \frac{1}{4}$ requires the scale factor is constant. Therefore, for both solutions of photon density in the realistic universe, there is no ν that could render Eq. (3.24) true due to the existence of other substances, i.e., $\rho_b(t_0) + \rho_{dm}(t_0) \neq 0$.

Although there is no ν keeping photon energy conserved, it is likely that there exists a special ν sustaining the total energy conservation of the universe, which means $[\rho_r(t_{re}) + \rho_\Lambda(t_{re})]V(t_{re}) = [\rho_r(t_0) + \rho_\Lambda(t_0)]V(t_0)$. It is found that the energy conservation of the universe is valid only when $\nu = 1$ and $\rho_{r0} = -3c_0 = 0$. Reviewing Eq. (3.19), the energy density of photons is negative in case of $\nu = 1$ and $\rho_{r0} = -3c_0 = 0$, which is obviously unacceptable.

So far, we have discussed two special couplings between running vacuum and CMB in the context of a realistic universe model. Our research shows that it is impossible for CMB to satisfy energy conservation in RVMs due to $\rho_b(t_0) + \rho_{dm}(t_0) \neq 0$. Even replacing the independent energy conservation of CMB with the energy conservation of the whole universe, it can not be achieved by choosing special values of ν and c_0 .

Next, we discuss the impact of running vacuum on the baryon-to-photon ratio. We first study the properties of photons and baryons in RVMs, respectively.

4 Photons in RVMs

Undoubtedly, CMB has crucial significance in cosmology and it is also a significant tool and indicator for studying the evolution of the universe. In different universe models, the properties of CMB may be completely different. Combining observations and reasonable inferences, one can put forward effective constraints on universe models and even eliminate fallacious universe models. In RVMs, running vacuum can influence the evolution of certain components of the universe by non-minimal coupling. If running vacuum could be coupled to CMB, after photons are decoupled from other matter, CMB is mainly controlled by the expansion of the universe (scale factor) and running vacuum.

According to the current observation of CMB and the temperature estimation of CMB at the end of recombination from the anisotropy of CMB, the effective parameter space of running vacuum may be limited to a narrow range. Considering the value of ν in RVMs has been limited to $|\nu| < 10^{-3}$ in related research [55, 60, 63, 75], our following research is only relevant to the solution for $\nu < \frac{1}{4}$ in Eqs. (3.19) and (3.22).

In order to make our discussion more clear and reasonable, let us sort out the known facts about CMB and the observable universe in advance. First of all, the current temperature of CMB ($T_0 = 2.73$ K) and the radius of the observable universe ($R_0 \sim 46$ billion light-years) are given by precise observations, which can not be violated in any model of the universe. Secondly, according to the theory of scattering describing the collision between photons and atoms (electrons), the decoupling temperature of CMB is determined around 3000 K [6, 77]. Finally, the size of the observable universe at the moment of photon decoupling is indistinct




Figure 1. Both of the upper panels are the temperature of CMB evolving with the scale factor. The under left panel is the number of photons per unit comoving volume. The under right panel is the energy density of running vacuum, which has been plotted in a dimensionless manner. For Figs. (1(a)), (1(c)), and (1(d)), seven cases of parameter ν are presented: $\nu = -10^{-3}$ (orange-solid), $\nu = -10^{-4}$ (green-solid), $\nu = -10^{-5}$ (blue-solid), $\nu = 0$ (red-solid), $\nu = 8 \times 10^{-6}$ (blue-dashed), $\nu = 4 \times 10^{-5}$ (green-dashed), and $\nu = 5.2 \times 10^{-5}$ (orange-dashed). For Fig. (1(b)), three cases of parameter ν are presented: $\nu = 10^{-3}$ (red), $\nu = 10^{-4}$ (green), $\nu = 5.3 \times 10^{-5}$ (orange). For all plots, the proportions of each component in the current universe are: 26.8% (DM), 4.9% (Baryon), 0.005% (CMB), and 68.295% (Vacuum).

(i.e., the value of the scale factor), which affects our evaluation on the age and the baryon-to-photon ratio of the universe at the epoch of photon decoupling (generally it is considered as 378,000 years in the Λ CDM model). Based on these facts, we can calculate and analyze reasonably the impact of running vacuum on CMB in RVMs. The number of photons per unit comoving volume (not unit volume) is the most concerned issue in this section.

Taking ρ_{r0} [see Eq. (3.20)] into Eq. (3.19), since the energy of photons per unit volume satisfies Eq. (3.4), the relation between the temperature of CMB and the scale factor is given as

$$\left(\rho_r(t_0) - \frac{3\nu[\rho_b(t_0) + \rho_{dm}(t_0)]}{1 - 4\nu} \right) a^{-4(1-\nu)} + \frac{3\nu[\rho_b(t_0) + \rho_{dm}(t_0)]}{1 - 4\nu} a^{-3} = \frac{\pi^2 k^4}{15c^3 \hbar^3} T^4. \quad (4.1)$$

When $T = T_0 = 2.73$ K and $a(t_0) = 1$, $\rho_r(t_0) \sim 4.0 \times 10^{-14} \text{J/m}^3$ is not related to the value of ν . When $\nu = 0$, $a(t_{re})$ is approximately given as 9.1×10^{-4} (with redshift $z \sim 1100$).

With Eq. (4.1), it is found that the sign of ν has decisive influence on the value of the scale factor [see Fig. (1(a))]. All the temperature lines in Fig. (1(a)) converge to the point ($a = 1, T = 2.73$ K), and the intersections of the temperature lines with the horizontal line $T = 3000$ K are the values of the scale factor at the end of recombination for different ν (see Table 1 for detail). When $\nu > 0$, the scale factor at the end of recombination is less than the estimated value in the Λ CDM model, and the scale factors resulted from $\nu < 0$ have exactly reverse effects. Overall, the scale factor of recombination increases with ν .

Our research shows that within the reference range of $|\nu| < 10^{-3}$, the evolution of CMB can indeed provide extra strict constraints on the the range of ν if vacuum is coupled to photons. When $\nu > 0$, we find that the upper limit of ν is about $5.25 * 10^{-5}$, and the larger ν would result in non-physical results of CMB [see Fig. (1(b))]. But when $10^{-3} < \nu < 0$, the energy density of photons is always normal.

The number of photons per unit comoving volume is given as

$$N_{rc} = \rho_r(a) a^3 = r_A \left(\frac{T_0^4 [a^{4\nu}(1 - 19024\nu) + 19020\nu a]}{a^4(1 - 4\nu)} \right)^{3/4} a^3, \quad (4.2)$$

where $r_A = \frac{2k^3\zeta(3)}{\pi^2c^3\hbar^3} \sim 2.02 * 10^7$ (K·m) $^{-3}$. For some special values of parameter ν , the evolution of N_{rc} is shown in Fig. (1(c)). Only when $\nu = 0$, N_{rc} does not evolve with the temperature change of CMB [the red-solid line in Fig. (1(c))], which is exactly what happens in the Λ CDM model. Since parameter ν does not change the temperature of CMB at present and also the energy density of photons, when $a = 1$, all cases correspond to the current number density of photons in the universe: $4.1 * 10^8$ m $^{-3}$ [see Fig. (1(c))]. We also noticed that although the number density of photons at recombination is the same (it is a function of temperature, independent of ν and the scale factor), the number of photons per unit comoving volume can differ more than two orders of magnitude due to parameter ν . Even for the case of $\nu \sim -10^{-5}$, the gap, comparing with the Λ CDM model, could be close to 15% (see more detail in Table 1).

Finally, the energy density of running vacuum is shown in Fig. (1(d)), which evolves monotonously with the scale factor (monotone decreasing for $\nu > 0$ and monotone increasing for $\nu < 0$).

Table 1. Scale factor and photon number per unit comoving volume at the end of recombination

ν	-10^{-3}	-10^{-4}	-10^{-5}	0	$8 * 10^{-6}$	$4 * 10^{-5}$	$5.2 * 10^{-5}$
$a(t_{re})$	0.00193	0.00118	0.00095	0.00091	0.00087	0.00064	0.00029
$N_{rc}(t_{re})$	$3.9 * 10^9$	$9.1 * 10^8$	$4.7 * 10^8$	$4.1 * 10^8$	$3.6 * 10^8$	$1.4 * 10^8$	$1.4 * 10^7$

5 Baryons in RVMs

Since the number of baryons in the Universe is a billion times less than the number of photons, the variation of baryon-to-photon ratio is more sensitive to the variation of the number density of baryons. In this section, we discuss the possible impact of running vacuum on baryons in the universe. Similar to the previous approach, when baryons are coupled to vacuum, we assume that the energy-momentum tensor of CMB maintains independently conservation.

For the coupling between running vacuum and baryons, the corresponding Friedmann equations are still Eqs. (3.17) and (3.18). Since the solutions for photons and running vacuum with respect to the scale factor are as with the ones in the Λ CDM model, we can immediately get the energy density of baryons being satisfied with

$$\frac{d\rho_\Lambda}{da} + \frac{d\rho_b}{da} + 3\frac{\rho_b}{a} = -\frac{3\nu}{a} \left[\frac{4}{3}\rho_r(t_0)a^{-4} + \rho_{dm}(t_0)a^{-3} + \rho_b \right] + \frac{d\rho_b}{da} + 3\frac{\rho_b}{a} = 0. \quad (5.1)$$

The solution is given as

$$\begin{aligned} \rho_b(a) &= \rho_{b0}a^{-3(1-\nu)} - \rho_{dm}(t_0)a^{-3} - \frac{4\nu\rho_r(t_0)}{1+3\nu}a^{-4} \quad (\nu \neq -\frac{1}{3}) \\ \rho_b(a) &= \rho_{b1}a^{-4} - \rho_{dm}(t_0)a^{-3} - \frac{4\rho_r(t_0)}{3}a^{-4}\log a \quad (\nu = -\frac{1}{3}). \end{aligned} \quad (5.2)$$

Note that $\rho_{b0} = \rho_b(t_0) + \rho_{dm}(t_0) + \frac{4\nu\rho_r(t_0)}{1+3\nu}$ and $\rho_{b1} = \rho_b(t_0) + \rho_{dm}(t_0)$, which insure that when $\nu = 0$, $\rho_b(a)$ could revert to the solution in the Λ CDM model. And the solution of running vacuum is

$$\begin{aligned} \rho_\Lambda(a) &= \frac{\nu\rho_r(t_0)}{1+3\nu}a^{-4} + \frac{\nu\rho_{b0}}{1-\nu}a^{-3(1-\nu)} + \frac{3}{1-\nu}c_0 \quad (\nu \neq -\frac{1}{3} \text{ and } \nu \neq 1) \\ \rho_\Lambda(a) &= -\frac{1}{4}[\rho_r(t_0) + \rho_{b1}]a^{-4} + \frac{1}{3}\rho_r(t_0)a^{-4}\log a + \frac{9}{4}c_0 \quad (\nu = -\frac{1}{3}) \\ \rho_\Lambda(a) &= 4\rho_r(t_0)a^{-4} - 3\rho_{b0}\log a \quad (\nu = 1), \end{aligned} \quad (5.3)$$

where $\rho_{b0} = -3c_0$ and the corresponding integration constant has been set to 0. With these solutions, we can study the evolution of baryons after recombination. Also since Joan Solà et al. have pointed out, with multiple observations in cosmology, $|\nu| < 10^{-3}$ [55, 60, 63, 75], we will still only discuss the impact caused by running vacuum on the number density of baryons within the scope of $|\nu| < 10^{-3}$.

Suppose there exists a baryon-generation (annihilation) process in the universe and the energy source (sink) is running vacuum⁵. According to the above analysis on the freely expanding photons, it can be known that the approximate value of the scale factor is 0.00091 at the end of recombination. The baryon density of the current universe is approximately 4.2×10^{-28} kg/m³. Therefore, the baryons density at the end of recombination can be obtained from Eq. (5.2), which is plotted in Fig. (2(a)) with five sets of ν 's. When $|\nu| < 10^{-5}$, the baryon density is almost hard to deviate from the result ($\nu = 0$) in the Λ CDM model. Even if $|\nu|$ takes the maximum and minimum values of the feasible range $|\nu| < 10^{-3}$, the deviation of the baryon density from the result in Λ CDM model is still negligible. At the end of recombination, the maximum deviation is about 15% (see Table 2). As $\nu < 0$, running vacuum presents obviously a state of negative energy, which actually should also appear in Fig. (1(d)) when $\nu < 0$ and $a(t) \ll 1$. Unlike baryons, running vacuum is sensitive to the value of ν at the end of recombination. Even a tiny deviation of ν from 0 can cause an order of magnitude difference in vacuum energy density [see Fig. (2(c))]. When $\nu = 0$, see the red-solid in Fig. (2(c)), $\rho_\Lambda = 3c_0$ is the current vacuum energy density.

⁵It seems that such a process is unlikely to exist in the late universe, but it could be possible in the early universe with high energy and strong gravitational field. Since any interaction process generally has a truncation, the known possible baryon number non-conservation processes are basically ruled out in the low-energy late universe. However we do not rule out the possibility that the interaction between running vacuum and baryons, which could be extremely weak and hardly be detected. Later we will simply estimate the incidence of the interaction in RVMs.




Figure 2. The upper left panel is the energy density of baryons. The upper right panel is the number of baryons per unit comoving volume. The under panel is the energy density of running vacuum. All plots have been plotted in a dimensionless manner. Five cases of parameter ν are presented in all plots: $\nu = -10^{-3}$ (orange-solid), $\nu = 10^{-4}$ (green-solid), $\nu = 0$ (red-solid), $\nu = 10^{-3}$ (orange-dashed), and $\nu = 10^{-4}$ (green-dashed). The current number of baryons per unit comoving volume is about 0.2. The proportions of each component in the current universe are: 26.8% (DM), 4.9% (Baryon), 0.005% (CMB), and 68.295% (Vacuum).

Table 2. Baryon density and baryon number per unit comoving volume at the end of recombination

ν	-10^{-3}	-10^{-4}	-10^{-5}	0	10^{-5}	10^{-4}	10^{-3}
$10^{-9}\rho_b(t_{re})/\rho_b(t_0)$	1.143	1.308	1.325	1.327	1.329	1.346	1.515
$n_b(t_0)^{-1}N_{bc}(t_{re})$	0.86113	0.98599	0.99860	1	1.00140	1.01404	1.14175

Now we analyse the number density of baryons, which can be expressed as

$$\rho_b(a) = n_b(a)m_0c^2, \quad (5.4)$$

where m_0 is the mass of a single baryon, $n_b(a)$ is the number density of baryons, and c is the speed of light. According to the current observations, $n_b(t_0) \sim 2.5 * 10^{-7}$ atoms \cdot cm $^{-3}$ is a relatively accurate value⁶. If m_0 is a constant, $n_b(t_{re}) = \frac{\rho_b(t_{re})}{\rho_b(t_0)}n_b(t_0)$. The number of

⁶The estimated value of $n_b(t_0)$ from BBN is in the range of $(2.4 * 10^{-7} \sim 2.7 * 10^{-7})$ atoms \cdot cm $^{-3}$ (95% CL) [17].

baryons per unit comoving volume satisfies

$$N_{bc}(a) = n_b(a)a^3 = \frac{\rho_b(a)}{\rho_b(t_0)}n_b(t_0)a^3, \quad (5.5)$$

where $\rho_b(a)$ refers to Eq. (5.2). The evolution of $N_{bc}(a)$ with the scale factor is given by Fig. (2(c)). When $\nu = 0$, it is a constant and consistent with the Λ CDM model. The value of ν has no significant impact on the estimation of the baryon number per comoving volume at the end of recombination if the Λ CDM model is regarded as a standard (see Table 2).

6 The effect of running vacuum on the baryon-to-photon ratio

From the previous sections, it is found that running vacuum can indeed influence the particle number densities of photons and baryons by assuming vacuum is non-minimally coupled to them. In this section, we estimate the possible baryon-to-photon ratio at the end of recombination in RVMs. In cosmology, the baryon-to-photon ratio at different epochs is directly related to our understanding of some processes in the earlier universe. In the Λ CDM model, the ratio is close to a constant after recombination, and it can even be traced back to the BBN epoch. However, as we saw earlier, in RVMs the particle number densities of photons and baryons could possess different modes of evolution due to running vacuum. Therefore, the baryon-to-photon ratio in RVMs is not a constant even after recombination. Since we do not figure out whether running vacuum is coupled to photons or baryons, we will study the impact of the two situations on the baryon-to-photon ratio.

It is worth noting that when $\nu < 0$, vacuum energy may be negative when the scale factor is small enough [see Figs. (1(d)) and (2(c))], which can be also speculated from the definition of vacuum energy [see Ref. (2.4)]. In general, as we discuss RVMs with $\nu < 0$, we must be very careful and insure that vacuum energy is positive in the early universe. In view of this, we only consider the baryon-to-photon ratio in the RVMs with $\nu \geq 0$.

If running vacuum is only coupled to photons, the range of ν is about $5.25 \times 10^{-5} > \nu \geq 0$ to keep the energy density of CMB normal. So, the baryon-to-photon ratio is given by

$$\eta(a) = \frac{n_b(a)}{n_r(a)} = \frac{n_b(a)a^3}{N_{rc}(a)} = \frac{N_{bc}(a)}{r_A} \left(\frac{T_0^4 [a^{4\nu}(1 - 19024\nu) + 19020\nu a]}{a^4(1 - 4\nu)} \right)^{-3/4} a^{-3}, \quad (6.1)$$

where $N_{rc}(a)$ [see Eq. (4.2)] is the number of photons per unit comoving volume and $N_{bc}(a) = n_b(a)a^3$ is the number of baryons per unit comoving volume, respectively. Since running vacuum is only coupled to photons, the total number of baryons inside the observable universe is a constant after BBN, i.e., $n_b(a)a^3 = n_b(t_0) \sim 2.5 \times 10^{-7}$ atoms·cm⁻³. From Fig. (3(a)) and Table 3, one can find that when photons are coupled to running vacuum, the baryon-to-photon ratio at the end of recombination, $\eta(a_{re})$, increases with ν , but the deviation is limited by the upper limit of ν . When $\nu = 5.2 \times 10^{-5}$, $\eta(a_{re}) \sim 1.80 \times 10^{-8}$, which is almost 30 times the result in the Λ CDM model (when $\nu = 0$, $\eta(a_{re}) \sim 6.11 \times 10^{-10}$). Some other results are also presented in Table 3.

As we know that η can be obtained independently from different epochs of the universe and here are four of the most common epochs: BBN (η^{BBN} , $z \sim 10^9$), recombination (η^{re} , $z \sim 1100$), the period of time associated with the Ly α forest ($\eta^{Ly\alpha}$, $z \sim 2 - 3$) [85], and the present epoch (η_0 , $z = 0$). If we are clear about the values of η at any two epochs of the universe, we can use them to constrain the universe models to which η is sensitive (see




Figure 3. Plots of the baryon-to-photon ratio from recombination to the present day. The upper panel is for the case of running vacuum coupled to photons [$\nu = 5.2 \times 10^{-5}$ (orange), $\nu = 4 \times 10^{-5}$ (green), $\nu = 8 \times 10^{-6}$ (blue), and $\nu = 0$ (red)]. The upper right panel is for running vacuum coupled to baryons [$\nu = 10^{-3}$ (orange), $\nu = 10^{-4}$ (green), $\nu = 10^{-5}$ (blue), and $\nu = 0$ (red)]. The current baryon-to-photon ratio is $\eta_0 \sim 6.11 \times 10^{-10}$.

Table 3. The baryon-to-photon ratio at the end of recombination with photons coupled to vacuum

ν	0	5×10^{-7}	8×10^{-6}	4×10^{-5}	5.2×10^{-5}
η_ν^{re}	6.11×10^{-10}	6.15×10^{-10}	6.92×10^{-10}	1.79×10^{-9}	1.80×10^{-8}

references mentioned in Introduction). It also applies to the RVMs in which photons are coupled to running vacuum. Considering the following aspects, it is more reasonable to use η^{re} and η_0 to constrain such RVMs:

1. First of all, η_0 must be considered because it is given directly by observations, which therefore is the most credible. Moreover, η_0 is also more accurate than other speculation results: $\eta_0 \sim 6.11 \times 10^{-10}$ [17, 18].

2. Since η_0 has been chosen, $\eta^{\text{Ly}\alpha}$ will be eliminated. The reason is that these two epochs are too close and so it is hard to inspect the influence of the universe model on η .

3. According to the latest research, the error of η^{re} from the anisotropy of CMB is small: $\eta^{re} \sim (6.090 \pm 0.060) * 10^{-10}$ [19], and the estimation result of η^{BBN} from the abundance of light elements has a large error: $\eta^{BBN} \sim (6.084 \pm 0.230) * 10^{-10}$ [19].

4. The lithium problem in BBN has been unresolved [87, 88], so the estimation of η^{BBN} is questionable. There may be new physics beyond the Standard Model.

5. During the period from BBN to recombination, the collision of photons with electrons and atomic nuclei may lead to a change in photon number density. Therefore, the change in η after BBN may not only depend on running vacuum.

Reviewing η^{re} and η_0 , we notice that $\eta_0 \sim 6.11 * 10^{-10}$ is almost in the middle of the range of $\eta^{re} \sim (6.090 \pm 0.060) * 10^{-10}$, that is to say, from the end of recombination to the present day, the value of η can be reduced or increased. We set the baryon-to-photon ratio in RVMs to equal η_0 at present, and η_ν^{re} presents the baryon-to-photon ratio in RVMs at the end of recombination. Since the error of $\eta^{re} \sim (6.090 \pm 0.060) * 10^{-10}$ is small, the dynamic term in RVMs must evolve very slowly to ensure that the value of η_ν^{re} in RVMs is not in contradiction with η^{re} obtained by the anisotropy of CMB.

We list some results of η_ν^{re} corresponding to several sets of positive ν in Table 3. It is seen that when $\nu \geq 0$, η_ν^{re} is always larger than the current η_0 and η_ν^{re} increases with the parameter ν . We also found that in order not to contradict with $\eta^{re} \sim (6.090 \pm 0.060) * 10^{-10}$, ν must be at least less than $5 * 10^{-7}$. In fact, from the perspective of baryon-to-photon ratio, it is difficult to distinguish RVMs from the Λ CDM model ($\nu = 0$) when $0 \leq \nu < 5 * 10^{-7}$. Even if $\nu = 8 * 10^{-6}$, we can find from Fig. (3(a)) that it almost overlaps with the result in the Λ CDM model. Therefore, if there exists coupling between photons and running vacuum, according to the data on the baryon-to-photon ratio, the coefficient ν of the dynamic term of vacuum must be extremely small ($0 \leq \nu < 5 * 10^{-7}$). Although we do get a new constraint on ν by considering the coupling between photons and running vacuum, the new constraint, comparing with $|\nu| < 10^{-3}$, is too strong and seems unnatural.

Next, we study the second case, that is, running vacuum is only coupled to baryons, in which the appropriate range of parameter ν is $0 \leq \nu < 10^{-3}$. The baryon-to-photon ratio is

$$\hat{\eta}(a) = \frac{N_{bc}(a)}{n_r(a)a^3} = \left[\left(\frac{317}{49} + \frac{\nu}{245(1+3\nu)} \right) a^{3\nu} - \frac{268}{49} - \frac{\nu}{245(1+3\nu)} a^{-1} \right] \frac{n_b(t_0)}{n_r(a)a^3}, \quad (6.2)$$

where $N_{bc}(a)$ is given by Eq. (5.5). Note that $\frac{n_b(t_0)}{n_r(a)a^3} = \frac{n_b(t_0)}{n_r(t_0)} \sim 6.11 * 10^{-10}$ is the current baryon-to-photon ratio for $n_b(t_0) \sim 2.5 * 10^{-7}$ atoms·cm⁻³ and $n_r(t_0) \sim 4.1 * 10^8$ m⁻³. The evolutionary trend of $\hat{\eta}(a)$ with the scale factor is shown in Fig. 3(b). As parameter ν increases, $\hat{\eta}_\nu^{re}$ will be smaller (see Table 4). Comparing these sets of data in Table 4, one finds that for $0 \leq \nu < 10^{-3}$, $\hat{\eta}_\nu^{re}$ does not change much. Compare $\hat{\eta}_\nu^{re}$ with $\eta^{re} \sim (6.090 \pm 0.060) * 10^{-10}$, the constraint range of parameter ν is reduced to $0 \leq \nu < 10^{-4}$, which is slightly stronger than $0 \leq \nu < 10^{-3}$ and seems to be a satisfactory result.

Table 4. The baryon-to-photon ratio at the end of recombination with baryons coupled to vacuum

ν	0	10^{-5}	10^{-4}	10^{-3}
$\hat{\eta}_\nu^{re}$	$6.11 * 10^{-10}$	$6.10 * 10^{-10}$	$6.02 * 10^{-10}$	$5.26 * 10^{-10}$

Unfortunately, here the baryon decay rate ⁷ is, in fact, rather fast if ν is not extreme small. One can simply make a following estimation. From recombination to the present day, the average particle number of decaying baryons per year within the observable universe can be estimated as

$$\bar{N} = \frac{N_0 - N_{re}}{\mathfrak{T}}, \quad (6.3)$$

where N_0 is total number of baryons inside the observable universe at present and N_{re} indicates the corresponding value at the end of recombination. Here, $\mathfrak{T} \sim 1.38 * 10^{10}$ years is the age of the universe (the period before recombination is negligible). If baryon decay is equally likely to occur everywhere in the universe, with such decay rate, the number of baryons decaying into running vacuum on the entire earth per year (for the present day) should be

$$\bar{N}_{\text{earth}} = \bar{N} \frac{N_{\text{earth}}}{N_0}, \quad (6.4)$$

where N_{earth} is the total number of baryons on the earth. In rough, we take the mass of the earth as $6 * 10^{24}$ kg and the average mass of baryon as $2 * 10^{-27}$ kg, so $N_{\text{earth}} \sim 10^{51}$. The total number of baryons in the current observable universe can be given by the current baryon number density $n_b(t_0) \sim 2.5 * 10^{-7}$ atoms·cm⁻³ and the current volume of the observable universe ($\sim 4 * 10^{80}$ m³). Given any value of ν , we can obtain a corresponding N_{re} and \bar{N}_{earth} . The final results are present at Table 5.

Even if $\nu = 10^{-5}$, $\bar{N}_{\text{earth}} \sim 10^{38} \sim 10^{-13} N_{\text{earth}}$ is still a huge number, in other words, the earth will lose 10^{11} kg due to the decay of baryons into running vacuum each year. Such mass is insignificant compared to the total mass of earth, but with the current observation accuracy, humans may have observed the phenomenon that baryons decay into running vacuum in the laboratory. Obviously, in order not to contradict observations and experiments (such as the half-life of a proton is at least $1.67 * 10^{34}$ years [86]), the value of ν needs to be unnaturally small.

Table 5. The baryon-to-photon ratio at the end of recombination with baryons coupled to vacuum

ν	0	10^{-5}	10^{-4}	10^{-3}
\bar{N}_{earth}	0	$\sim 10^{38}$	$\sim 10^{39}$	$\sim 10^{40}$

7 Discussions and conclusions

The energy-momentum tensor conservation of photons is not compatible with the energy conservation of photons in the Λ CDM model, and the latter is usually not satisfied because of the expansion of space-time. In RVMs, it is found that if $\rho_\Lambda = \frac{3}{4\kappa^2} H^2 + c_0$, the energy conservation of photons can be satisfied in the context of a toy universe model. And if $c_0 = 0$, the total energy of the universe can be conserved. However, these conditions can not be implemented for a realistic universe due to $\rho_b(t_0) + \rho_{dm}(t_0) \neq 0$. Therefore, for a realistic

⁷From Fig. 2(b), when $\nu \geq 0$, the number of baryons per unit comoving volume is monotonically decreasing, which means baryons can only actually annihilate into vacuum.

universe, the energy of photons (and also the universe) is not conserved even considering running vacuum.

The baryon-to-photon ratio, which is approximately a constant in the Λ CDM model, at different epochs indicates the characteristics of the cosmological processes. One can estimate the baryon-to-photon ratio of the universe at various epochs based on particle physics, which is independent on the model of the universe and also is the crucial tool to check the availability of universe models. The study on the baryon-to-photon ratio in the standard model explains the relationship between the abundance of most light elements in the universe and BBN [1–6], which is recognized by many people. However, the difference between the estimation and observation on Lithium abundance is like a dark cloud hanging over BBN and the Λ CDM model [87, 88]. A time-evolving baryon-to-photon ratio may bring about a turning point to the problem of Lithium abundance.

Since there exists non-minimal coupling between matter fields and running vacuum in RVMs, the density of certain substances will be affected by running vacuum. If the substances are photons or baryons, the baryon-to-photon ratio must evolve with running vacuum. Considering that the photon density may be affected by a variety of factors before the end of the recombination, in order to analyze and study the influence of running vacuum on the baryon-photon ratio without other distractions, we consider the period from the end of the recombination to the current universe. For photons coupling with running vacuum, the temperature of CMB at the end of recombination, still needs to meet the requirement of photon decoupling, and it is independent on the model of universe. According to the currently observed temperature of CMB, we can inversely deduce the size of the scale factor at recombination, which changes with the coefficient ν in running vacuum (Table 1). The total number of photons in the observable universe also changes with ν at recombination, see Fig. 1(c). Similarly, we obtain the baryon density, see Fig. 2(a), and baryon number density, see Fig. 2(c), for different values of ν , when running vacuum is coupled to baryons. In order to ensure that vacuum energy is positive in the early universe, the parameter ν must be equal or greater than 0. When photons are coupled to running vacuum, the upper limit of ν is approximately equal to $5.25 * 10^{-5}$, which does not appear for the coupling between baryons and vacuum.

Combining the current observations of the baryon-to-photon ratio and the corresponding value at recombination predicted by the anisotropy of CMB, we can get constraints on RVMs by considering that running vacuum is coupled to photons or baryons. When photons are coupled to running vacuum, in order to ensure that the baryon-photon ratio in RVMs does not conflict with the anisotropy of CMB, parameter ν must be abnormally small. Such a new constraint on ν is unnatural. When baryons are coupled to vacuum, we find that the value of ν just needs to be $0 \leq \nu < 10^{-4}$ to satisfy the same requirement. However, the decay rate of baryons into vacuum is too fast, unless ν is extremely small. Otherwise we should be able to observe the decay process of non-conservation of baryon number on the earth.

In summary, it can be concluded that the possibility of running vacuum coupling with photons or baryons is very low. At present, we do not figure out whether the running vacuum can be coupled with dark matter and how to determine the possibility of such coupling. It is a question worthy of further discussion. Our research on constraining RVMs is also applicable to other universe models in which the baryon-to-photon ratio may evolve over time.

Acknowledgments

This work was supported by the National Natural Science Foundation of China under Grant Nos. 11873001 and 12005174. K. Yang acknowledges the support of Natural Science Foundation of Chongqing, China under Grant No. cstc2020jcyj-msxmX0370. J. Li acknowledges the support of Natural Science Foundation of Chongqing (Grant No. cstc2018jcyjAX0767).

References

- [1] R. A. Malaney and G. J. Mathews, *Phys. Rept.* **229**, 145 (1993).
- [2] S. Sarkar, *Rept. Prog. Phys.* **59**, 1493 (1996).
- [3] D. Tytler, J. M. O’Meara, N. Suzuki, and D. Lubin, *Phys. Scripta T* **85**, 12 (2000).
- [4] B. D. Fields, P. Molaro, and S. Sarkar, *Chin. Phys. C* **38**, 339 (2014).
- [5] G. J. Mathews, M. Kusakabe, and T. Kajino, *Int. J. Mod. Phys. E* **26**, 1741001 (2017).
- [6] P. A. Zyla *et al.* [Particle Data Group], *PTEP* **2020**, 083C01 (2020).
- [7] A. M. Boesgaard and G. Steigman, *Ann. Rev. Astron. Astrophys.* **23**, 319 (1985).
- [8] G. Steigman, *Ann. Rev. Nucl. Part. Sci.* **57**, 463 (2007).
- [9] R. H. Cyburt, B. D. Fields, K. A. Olive, and T. H. Yeh, *Rev. Mod. Phys.* **88**, 015004 (2016).
- [10] S. Sarkar, [astro-ph/0205116](https://arxiv.org/abs/astro-ph/0205116) (2002).
- [11] C. J. Copi, D. N. Schramm, and M. S. Turner, *Science* **267**, 192 (1995).
- [12] J. R. Bond, R. Crittenden, R. L. Davis, G. Efstathiou, and P. J. Steinhardt, *Phys. Rev. Lett.* **72**, 13 (1994).
- [13] W. Hu and N. Sugiyama, *Phys. Rev. D* **51**, 2599 (1995).
- [14] G. Jungman, M. Kamionkowski, A. Kosowsky, and D. N. Spergel, *Phys. Rev. D* **54**, 1332 (1996).
- [15] W. Hu and S. Dodelson, *Ann. Rev. Astron. Astrophys.* **40**, 171 (2002).
- [16] M. Fukugita and P. J. E. Peebles, *Astrophys. J.* **616**, 643 (2004).
- [17] M. Tanabashi *et al.* [Particle Data Group], *Phys. Rev. D* **98**, 030001 (2018).
- [18] N. Aghanim *et al.* [Planck Collaboration], *Astron. Astrophys.* **641**, A6 (2020).
- [19] B. D. Fields, K. A. Olive, T. H. Yeh, and C. Young, *JCAP* **2003**, 010 (2020); Erratum: [*JCAP* **2011**, E02 (2020)].
- [20] P. Ullio, L. Bergstrom, J. Edsjo, and C. G. Lacey, *Phys. Rev. D* **66**, 123502 (2002).
- [21] X. L. Chen and M. Kamionkowski, *Phys. Rev. D* **70**, 043502 (2004).
- [22] N. Padmanabhan and D. P. Finkbeiner, *Phys. Rev. D* **72**, 023508 (2005).
- [23] S. Ando and E. Komatsu, *Phys. Rev. D* **73**, 023521 (2006).
- [24] G. Bertone, W. Buchmuller, L. Covi, and A. Ibarra, *JCAP* **0711**, 003 (2007).
- [25] S. Galli, F. Iocco, G. Bertone, and A. Melchiorri, *Phys. Rev. D* **80**, 023505 (2009).
- [26] S. Galli, F. Iocco, G. Bertone, and A. Melchiorri, *Phys. Rev. D* **84**, 027302 (2011).
- [27] E. O. Zavarygin and A. V. Ivanchik, *J. Phys. Conf. Ser.* **661**, 012016 (2015).
- [28] E. O. Zavarygin and A. V. Ivanchik, *Astron. Lett.* **41**, 383 (2015).

- [29] P. J. E. Peebles and B. Ratra, *Rev. Mod. Phys.* **75**, 559 (2003).
- [30] R. Opher and A. Pelinson, *Mon. Not. Roy. Astron. Soc.* **362**, 167 (2005).
- [31] R. Opher and A. Pelinson, *Braz. J. Phys.* **35**, 1206 (2005).
- [32] S. Kumar, R. C. Nunes, and S. K. Yadav, *Phys. Rev. D* **98**, 043521 (2018).
- [33] S. K. Yadav, *Mod. Phys. Lett. A* **35**, 1950358 (2019).
- [34] J. D. Barrow and R. J. Scherrer, *Phys. Rev. D* **98**, 043534 (2018).
- [35] B. A. Campbell and K. A. Olive, *Phys. Lett. B* **345**, 429 (1995).
- [36] T. Dent and M. Fairbairn, *Nucl. Phys. B* **653**, 256 (2003).
- [37] C. J. Copi, A. N. Davis, and L. M. Krauss, *Phys. Rev. Lett.* **92**, 171301 (2004).
- [38] B. Li and M. C. Chu, *Phys. Rev. D* **73**, 025004 (2006).
- [39] J. Alvey, N. Sabti, M. Escudero, and M. Fairbairn, *Eur. Phys. J. C* **80**, 148 (2020).
- [40] T. Clifton, J. D. Barrow, and R. J. Scherrer, *Phys. Rev. D* **71**, 123526 (2005).
- [41] K. Freese, F. C. Adams, J. A. Frieman, and E. Mottola, *Nucl. Phys. B* **287**, 797 (1987).
- [42] M. Birkel and S. Sarkar, *Astropart. Phys.* **6**, 197 (1997).
- [43] J. Solà, *J. Phys. Conf. Ser.* **283**, 012033 (2011).
- [44] J. Solà, *J. Phys. Conf. Ser.* **453**, 012015 (2013).
- [45] E. L. D. Perico, J. A. S. Lima, S. Basilakos, and J. Solà, *Phys. Rev. D* **88**, 063531 (2013).
- [46] J. Solà and A. Gómez-Valent, *Int. J. Mod. Phys. D* **24**, 1541003 (2015).
- [47] H. Terazawa, *Phys. Lett.* **101B**, 43 (1981).
- [48] F. G. Alvarenga and N. A. Lemos, *Gen. Rel. Grav.* **30**, 681 (1998).
- [49] X. Zhang, *Phys. Rev. D* **74**, 103505 (2006).
- [50] Y. V. Stadnik and V. V. Flambaum, *Phys. Rev. Lett.* **114**, 161301 (2015).
- [51] A. Gómez-Valent, J. Solà, and S. Basilakos, *JCAP* **1501**, 004 (2015).
- [52] Y. H. Li, J. F. Zhang, and X. Zhang, *Phys. Rev. D* **93**, 023002 (2016).
- [53] I. L. Shapiro and J. Solà, *JHEP* **0202**, 006 (2002).
- [54] I. L. Shapiro and J. Solà, *Phys. Lett. B* **682**, 105 (2009).
- [55] J. Solà, *J. Phys. A* **41**, 164066 (2008).
- [56] J. A. S. Lima, S. Basilakos, and J. Solà, *Mon. Not. Roy. Astron. Soc.* **431**, 923 (2013).
- [57] C. Q. Geng, Y. T. Hsu, L. Yin, and K. Zhang, *Chin. Phys. C* **44**, 105104 (2020).
- [58] J. Solà, A. Gómez-Valent, and J. de Cruz Pérez, *Phys. Lett. B* **774**, 317 (2017).
- [59] E. L. D. Perico and D. A. Tamayo, *JCAP* **1708**, 026 (2017).
- [60] M. Rezaei, M. Malekjani and J. Solà, *Phys. Rev. D* **100**, 023539 (2019).
- [61] J. Solà, *Int. J. Mod. Phys. A* **31**, 1630035 (2016).
- [62] C. Q. Geng, C. C. Lee, and L. Yin, *JCAP* **1708**, 032 (2017).
- [63] A. Gómez-Valent and J. Solà, *EPL* **120**, 39001 (2017).
- [64] A. Gómez-Valent and J. Solà, *Mon. Not. Roy. Astron. Soc.* **478**, 126 (2018).
- [65] M. Gonzalez-Espinoza and D. Pavón, *Mon. Not. Roy. Astron. Soc.* **484**, 2924 (2019).
- [66] J. A. S. Lima, S. Basilakos, and J. Solà, *Eur. Phys. J. C* **76**, 228 (2016).

- [67] J. Solà and H. Yu, *Gen. Rel. Grav.* **52**, 17 (2020).
- [68] T. Harko, *Phys. Rev. D* **90**, 044067 (2014).
- [69] T. Harko, F. S. N. Lobo, J. P. Mimoso, and D. Pavón, *Eur. Phys. J. C* **75**, 386 (2015).
- [70] C. Tsallis, F. C. Sa Barreto, and E. D. Loh, *Phys. Rev. E* **52**, 1447 (1995).
- [71] J. A. S. Lima, *Gen. Rel. Grav.* **29**, 805 (1997).
- [72] M. Novello, L. A. R. Oliveira, and J. M. Salim, *Class. Quant. Grav.* **7**, 51 (1990).
- [73] A. Macleod, [physics/0407077](#) (2004).
- [74] A. Macleod, [physics/0511178](#) (2005).
- [75] J. Solà, J. de Cruz Pérez, and A. Gómez-Valent, *Mon. Not. Roy. Astron. Soc.* **478**, 4357 (2018).
- [76] S. Perlmutter *et al.* [Supernova Cosmology Project Collaboration], *Astrophys. J.* **517**, 565 (1999).
- [77] B. Ryden, *Introduction to cosmology*, Addison Wesley, San Francisco, USA, (2003).
- [78] J. C. Mather, D. J. Fixsen, R. A. Shafer, C. Mosier, and D. T. Wilkinson, *Astrophys. J.* **512**, 511 (1999).
- [79] D. J. Fixsen, *Astrophys. J.* **707**, 916 (2009).
- [80] N. Aghanim *et al.* [Planck Collaboration], *Astron. Astrophys.* **641**, A5 (2020).
- [81] E. W. Kolb and M. S. Turner, *Front. Phys.* **69**, 1 (1990).
- [82] C. A. Egan and C. H. Lineweaver, *Astrophys. J.* **710**, 1825 (2010).
- [83] J. E. Felten and R. Isaacman, *Rev. Mod. Phys.* **58**, 689 (1986).
- [84] J. M. Overduin and F. I. Cooperstock, *Phys. Rev. D* **58**, 043506 (1998).
- [85] D. Kirkman, D. Tytler, N. Suzuki, J. M. O'Meara, and D. Lubin, *Astrophys. J. Suppl.* **149**, 1 (2003).
- [86] B. Bajc, J. Hisano, T. Kuwahara, and Y. Omura, *Nucl. Phys. B* **910**, 1 (2016).
- [87] R. H. Cyburt, B. D. Fields, and K. A. Olive, *JCAP* **0811**, 012 (2008).
- [88] B. D. Fields, *Ann. Rev. Nucl. Part. Sci.* **61**, 47 (2011).