

# On the Ricci curvature of homogeneous Finsler spaces with $(\alpha, \beta)$ -metrics

Sarita Rani <sup>1</sup> and Gauree Shanker\*<sup>1</sup>

<sup>1</sup>Department of Mathematics and Statistics, Central University of Punjab, Bathinda, Punjab-151 401, India.  
Email: saritas.ss92@gmail.com, grshnkr2007@gmail.com

## Abstract

The study of curvature properties of homogeneous Finsler spaces with  $(\alpha, \beta)$ -metrics is one of the central problems in Riemann-Finsler geometry. In this paper, we consider homogeneous Finsler spaces with square metric and Randers change of square metric. First, we derive the explicit formulae for Ricci curvature of homogeneous Finsler spaces with these metrics. Next, we find a necessary and sufficient condition under which a homogeneous Finsler space with either of these metrics is of vanishing  $S$ -curvature. The formulae for Ricci curvature of homogeneous Finsler spaces with square metric and Randers change of square metric having vanishing  $S$ -curvature are established. Finally, we prove that the aforesaid spaces having vanishing  $S$ -curvature and negative Ricci curvature must be Riemannian.

**Mathematics Subject Classification:** 22E60, 53C30, 53C60.

**Keywords and Phrases:** Homogeneous Finsler space, square metric, Randers change, Ricci curvature,  $S$ -curvature.

## 1 Introduction

Finsler geometry is an interesting and active area of research for both pure and applied reasons (for detail, see [1, 2, 15, 19]). One special class of Finsler spaces is homogeneous and symmetric Finsler spaces, which is an active area of research nowadays. Many authors (for detail, see [10, 14, 17, 18, 26, 29, 35]) have worked in this area. Ricci curvature, denoted by  $Ric(x, y)$  is an important entity in Riemann-Finsler geometry. It is the mathematical object that controls growth rate of volume of a metric ball in a manifold. The Ricci curvature of a Finsler space  $(M, F)$  can always be expressed in the form  $Ric(x, y) = (n - 1)\lambda(x, y)F^2$ , where  $\lambda(x, y)$  is a scalar function on  $TM$ , called Einstein scalar. If the Einstein scalar  $\lambda(x, y)$  depends only on  $x$ , then the Finsler metric  $F$  is called an Einstein metric. Einstein manifolds play an important role in Riemann-Finsler geometry. Motivated by the open problem

“Does every smooth manifold admits an Einstein Finsler metric?”

posed by Chern, so many geometers (for example, see [5, 7, 8, 13, 24, 31, 38]) have established many results on Einstein Finsler metrics.

In [27], Rezaei et al. studied some Einstein  $(\alpha, \beta)$ -metrics. Further, in [33], Wang and Shen have studied Einstein-Randers metrics on homogeneous Riemannian spaces. After that, Deng and Hu [12] extended the work done in [33] to homogeneous Finsler spaces.

In [4], Bao and Robles proved that a compact Einstein-Randers space without boundary and with

---

\* corresponding author

negative Ricci curvature must be Riemannian. The problem of negative Ricci curvature was further studied by Mo and Yu in [22].

In [39], Zhou studied Finsler spaces with  $(\alpha, \beta)$ -metrics and introduced formulae for Riemann curvature and Ricci curvature for these metrics. Further, Cheng et al. [8] studied Einstein  $(\alpha, \beta)$ -metrics. During their study, they found that the formulae given in [39] are incorrect and they provided correct version of these formulae. Based on these formulae, a formula for Ricci curvature of homogeneous  $(\alpha, \beta)$ -metrics was constructed by Yan and Deng [36].

In 1972, Matsumoto [20] introduced the concept of  $(\alpha, \beta)$ -metrics which are the generalizations of Randers metric introduced by Randers [25]. The main aim of this paper is to establish an explicit formula for Ricci-curvature of a homogeneous Finsler space with Shen's square metric, and the same for that metric having vanishing  $S$ -curvature.

The simplest non-Riemannian metrics are the Randers metrics given by  $F = \alpha + \beta$  with  $\|\beta\|_\alpha < 1$ , where  $\alpha$  is a Riemannian metric and  $\beta$  is a 1-form. Besides Randers metrics, other interesting kind of non-Riemannian metrics are square metrics. Berwald's metric, constructed by Berwald [6] in 1929 as

$$F = \frac{\left(\sqrt{(1-|x|^2)|y|^2 + \langle x, y \rangle^2} + \langle x, y \rangle\right)^2}{(1-|x|^2)^2 \sqrt{(1-|x|^2)|y|^2 + \langle x, y \rangle^2}}$$

is a classical example of square metric. Berwald's metric can be rewritten as follows:

$$F = \frac{(\alpha + \beta)^2}{\alpha}, \quad (1.1)$$

where

$$\alpha = \frac{\sqrt{(1-|x|^2)|y|^2 + \langle x, y \rangle^2}}{(1-|x|^2)^2},$$

and

$$\beta = \frac{\langle x, y \rangle}{(1-|x|^2)^2}.$$

Equation (1.1) with the condition  $\|\beta\|_\alpha < 1$  satisfies Lemma 2.1 which is a necessary and sufficient condition for  $F$  to be a Finsler metric. Therefore, the metric given by equation (1.1) is an  $(\alpha, \beta)$ -metric. An  $(\alpha, \beta)$ -metric expressed in the form (1.1) is called square metric [32]. Square metrics play an important role in Finsler geometry. The importance of square metric can be seen in papers [32, 34, 37]. Square metrics can also be expressed in the form

$$F = \frac{\left(\sqrt{(1-b^2)\alpha^2 + \beta^2} + \beta\right)^2}{(1-b^2)^2 \sqrt{(1-b^2)\alpha^2 + \beta^2}},$$

where  $b := \|\beta_x\|_\alpha$  is the length of  $\beta$  [37].

In this case,  $F = \alpha\phi\left(b^2, \frac{\beta}{\alpha}\right)$ , where  $\phi = \phi(b^2, s)$  is a smooth function, is called general  $(\alpha, \beta)$ -metric. If  $\phi = \phi(s)$  is independent of  $b^2$ , then  $F$  is called an  $(\alpha, \beta)$ -metric.

If  $(M, F)$  be an  $n$ -dimensional Finsler space and  $\beta = b_i(x)y^i$  is a 1-form on  $M$ . Then  $F \rightarrow \bar{F}$  is called Randers change if

$$\bar{F} = F + \beta \quad (1.2)$$

Above change has been introduced by Matsumoto [21], and it was named as "Randers change" by Hashiguchi and Ichijyō [16]. In the current paper, we deal with square metrics

$$F = \frac{(\alpha + \beta)^2}{\alpha} = \alpha\phi(s), \text{ where } \phi(s) = 1 + s^2 + 2s.$$

The paper is organized as follows:

In section 2, we discuss some basic definitions and results to be used in consequent sections. The explicit formulae for Ricci curvature of homogeneous Finsler spaces with square metric and Randers change of square metric have been established in section 3. In section 4, we find a necessary and sufficient condition under which a homogeneous Finsler space with either of these metrics is of vanishing  $S$ -curvature. Also, a formula for Ricci curvature of a homogeneous Finsler space with each metric having vanishing  $S$ -curvature is established. Finally, we prove that the aforesaid spaces having vanishing  $S$ -curvature and negative Ricci curvature must be Riemannian.

## 2 Preliminaries

In this section, we discuss some basic definitions and results required for consequent sections. We refer [3, 9, 11] for notations and further details.

**DEFINITION 2.1.** *An  $n$ -dimensional real vector space  $V$  is said to be a **Minkowski space** if there exists a real valued function  $F : V \rightarrow [0, \infty)$ , called **Minkowski norm**, satisfying the following conditions:*

- $F$  is smooth on  $V \setminus \{0\}$ ,
- $F$  is positively homogeneous, i.e.,  $F(\lambda v) = \lambda F(v)$ ,  $\forall \lambda > 0$ ,
- For any basis  $\{u_1, u_2, \dots, u_n\}$  of  $V$  and  $y = y^i u_i \in V$ , the Hessian matrix  $(g_{ij}) = \left( \frac{1}{2} F^2_{y^i y^j} \right)$  is positive-definite at every point of  $V \setminus \{0\}$ .

**DEFINITION 2.2.** *Let  $M$  be a connected smooth manifold. If there exists a function  $F : TM \rightarrow [0, \infty)$  which is smooth on slit tangent bundle  $TM \setminus \{0\}$  and the restriction of  $F$  to any  $T_x M$ ,  $x \in M$ , is a Minkowski norm, then  $M$  is called a **Finsler space** and  $F$  is called a **Finsler metric**.*

An  $(\alpha, \beta)$ -metric on a connected smooth manifold  $M$  is a Finsler metric  $F$  constructed from a Riemannian metric  $\alpha = \sqrt{a_{ij}(x)y^i y^j}$  and a one-form  $\beta = b_i(x)y^i$  on  $M$  and is of the form  $F = \alpha \phi \left( \frac{\beta}{\alpha} \right)$ , where  $\phi$  is a smooth function on  $M$ . Basically,  $(\alpha, \beta)$ -metrics are the generalization of Randers metrics. Many authors (for example, see [14, 17, 28, 30, 35]) have worked on  $(\alpha, \beta)$ -metrics. Let us recall following (Shen's) lemma which gives necessary and sufficient condition for a function of  $\alpha$  and  $\beta$  to be a Finsler metric:

**LEMMA 2.1.** [9] *Let  $F = \alpha \phi(s)$ ,  $s = \beta/\alpha$ , where  $\phi$  is a smooth function on an open interval  $(-b_0, b_0)$ ,  $\alpha$  is a Riemannian metric and  $\beta$  is a 1-form with  $\|\beta\|_\alpha < b_0$ . Then  $F$  is a Finsler metric if and only if the following conditions are satisfied:*

$$\phi(s) > 0, \quad \phi(s) - s\phi'(s) + (b^2 - s^2)\phi''(s) > 0, \quad \forall |s| \leq b < b_0.$$

### 2.1 Ricci curvature of an $(\alpha, \beta)$ -metric

The notion of Riemannian curvature for Riemannian spaces can be extended to Finsler spaces. For an  $n$ -dimensional Finsler space  $(M, F)$ ,  $x \in M$ ,  $y (\neq 0) \in T_x M$ , the Riemannian curvature is a linear map  $R_y : T_x M \rightarrow T_x M$  defined as

$$R_y(v) = R_k^i(y)v^k \frac{\partial}{\partial x^i}, \quad v = v^i \frac{\partial}{\partial x^i},$$

where

$$R_k^i(y) = 2 \frac{\partial G^i}{\partial x^k} - \frac{\partial^2 G^i}{\partial x^j \partial y^k} y^j + 2G^j \frac{\partial^2 G^i}{\partial y^j \partial y^k} - \frac{\partial G^i}{\partial y^j} \frac{\partial G^j}{\partial y^k},$$

and  $G^i$  are geodesic coefficients given by

$$G^i := \frac{1}{4} g^{im} \left\{ (F^2)_{x^k y^m} y^k - (F^2)_{x^m} \right\}, \quad i = 1, 2, \dots, n.$$

DEFINITION 2.3. Let  $(M, F)$  be an  $n$ -dimensional Finsler space. For  $x \in M$  and non-zero  $y \in T_x M$ , the map defined on tangent bundle  $TM$  of  $M$  as

$$\text{Ric}(y) = \text{tr}(R_y) \quad \forall y \in T_x M$$

is called **Ricci-curvature** of  $(M, F)$ .

For an  $(\alpha, \beta)$ -metric  $F = \alpha\phi(s)$ ,  $s = \frac{\beta}{\alpha}$  on a Finsler manifold  $M$ , let us take

$$\begin{aligned} r_{ij} &= \frac{b_{i;j} + b_{j;i}}{2}, & s_{ij} &= \frac{b_{i;j} - b_{j;i}}{2}, \\ r_j^i &= a^{i\ell} r_{\ell j}, & s_j^i &= a^{i\ell} s_{\ell j}, \\ r_i &= b^\ell r_{\ell i} = b_j r_i^j, & s_i &= b^\ell s_{\ell i} = b_j s_i^j, \\ r &= r_{ij} b^i b^j = b^i r_i, & r_{00} &= r_{ij} y^i y^j, \\ r_{i0} &= r_{ij} y^j, & s_{i0} &= s_{ij} y^j, \\ r_0 &= r_i y^i, & s_0 &= s_i y^i, \end{aligned}$$

where ; denotes the covariant derivative w.r.t. Levi-Civita connection on Riemannian metric  $\alpha$  and  $a^{ij} = (a_{ij})^{-1}$ ,  $b^i = a^{ij} b_j$ .

**Theorem 2.1.** [8] Let  $F$  be an  $(\alpha, \beta)$ -metric on a Finsler space  $M$  and  ${}^\alpha \text{Ric}$  be Ricci curvature of  $\alpha$ . Then Ricci curvature of  $F$  is given by  $\text{Ric} = {}^\alpha \text{Ric} + RT_l^l$ , with

$$\begin{aligned} RT_l^l &= \frac{1}{\alpha^2} \left\{ (n-1) \zeta_1 + \zeta_2 \right\} r_{00}^2 \\ &+ \frac{1}{\alpha} \left[ \left( (n-1) \zeta_3 + \zeta_4 \right) r_{00} s_0 + \left( (n-1) \zeta_5 + \zeta_6 \right) r_{00} r_0 + \left( (n-1) \zeta_7 + \zeta_8 \right) r_{00;0} \right] \\ &+ \left( (n-1) \zeta_9 + \zeta_{10} \right) s_0^2 + (r r_{00} - r_0^2) \zeta_{11} + \left( (n-1) \zeta_{12} + \zeta_{13} \right) r_0 s_0 \\ &+ \left( r_{00} r_l^l - r_{0l} r_0^l + r_{00;l} b^l - r_{0l;0} b^l \right) \zeta_{14} + \left( (n-1) \zeta_{15} + \zeta_{16} \right) r_{0l} s_0^l \\ &+ \left( (n-1) \zeta_{17} + \zeta_{18} \right) s_{0;0} + s_{0l} s_0^l \zeta_{19} + \alpha \left[ r s_0 \zeta_{20} + \left( (n-1) \zeta_{21} + \zeta_{22} \right) s_l s_0^l \right] \\ &+ \alpha \left( 3s_l r_0^l - 2s_0 r_l^l + 2r_l s_0^l - 2s_{0;l} b^l + s_{l;0} b^l \right) \zeta_{23} + s_{0;l}^l \zeta_{24} \\ &+ \alpha^2 \left( s_l s^l \zeta_{25} + s_l^i s_i^l \zeta_{26} \right), \end{aligned}$$

where

$$\begin{aligned} \zeta_1 &= 2\psi \Theta_s (B - s^2) - 2s\psi \Theta + \Theta^2 - \Theta_s, \\ \zeta_2 &= 2\psi \psi_{ss} (B - s^2)^2 - (6s\psi \psi_s + \psi_{ss}) (B - s^2) + 2s\psi_s, \\ \zeta_3 &= -4(2Q\Theta_s + Q_s \Theta) \psi (B - s^2) + 4Q\Theta_s + 2Q_s \Theta + 4Q\Theta (s\psi - \Theta) - 2\Theta_B, \\ \zeta_4 &= -4\psi (2Q\psi_{ss} + Q_s \psi_s + Q_{ss} \psi_s^2) (B - s^2)^2 \\ &+ (-4\psi^2 (Q - sQ_s) + 4Q_{ss} \psi + 2Q_s \psi_s + 4Q\psi_{ss} - 2\psi_{sB} + 20sQ\psi \psi_s) (B - s^2) \\ &+ 2\psi (Q - sQ_s) - 4\psi_s - Q_{ss} - 10sQ\psi_s, \\ \zeta_5 &= 4\psi \Theta - 2\Theta_B, \\ \zeta_6 &= 2(2\psi \psi_s - \psi_{sB}) (B - s^2) - 2\psi_s, \\ \zeta_7 &= -\Theta, \end{aligned}$$

$$\begin{aligned}
\zeta_8 &= -\psi_s (B - s^2), \\
\zeta_9 &= 8Q\psi (Q\Theta_s + Q_s\Theta) (B - s^2) + 4Q^2 (\Theta^2 - \Theta_s) + 4Q (\Theta_B - Q_s), \\
\zeta_{10} &= (4\psi^2 (2QQ_{ss} - Q_s^2) + 8Q\psi (Q\psi_{ss} + Q_s\psi_s) - 4Q^2\psi_s^2) (B - s^2)^2 \\
&\quad + (-16sQ\psi (Q\psi_s + Q_s\psi) - 4\psi (2QQ_{ss} - Q_s^2) - 4Q (Q\psi_{ss} + Q_s\psi_s) + 4Q\psi_{sB} + 4Q_s\psi_B) (B - s^2) \\
&\quad - 4s^2Q^2\psi^2 + 4(2 + 3sQ) (Q\psi_s + Q_s\psi) - 8Q^2\psi + 2QQ_{ss} - Q_s^2 + 4sQ\psi_B, \\
\zeta_{11} &= 4\psi^2 + 4\psi_B, \\
\zeta_{12} &= 4Q(-2\psi\Theta + \Theta_B), \\
\zeta_{13} &= (8\psi (Q_s\psi - Q\psi_s) + 4Q\psi_{sB} + 4Q_s\psi_B) (B - s^2) + 8sQ\psi^2 + 4Q\psi_s - 4(1 - sQ)\psi_B, \\
\zeta_{14} &= 2\psi, \\
\zeta_{15} &= 4Q\Theta, \\
\zeta_{16} &= 4(Q\psi_s - Q_s\psi) (B - s^2) + 2Q_s - 2(1 + 2sQ)\psi, \\
\zeta_{17} &= 2Q\Theta, \\
\zeta_{18} &= 2(Q\psi_s + Q_s\psi) (B - s^2) + 2sQ\psi - Q_s, \\
\zeta_{19} &= 2(1 + sQ)Q_s - 2Q^2, \\
\zeta_{20} &= -8(\psi^2 + \psi_B)Q, \\
\zeta_{21} &= -4Q^2\Theta, \\
\zeta_{22} &= 2Q\psi - 4Q^2\psi_s (B - s^2), \\
\zeta_{23} &= 2Q\psi, \\
\zeta_{24} &= 2Q, \\
\zeta_{25} &= -4Q^2\psi, \\
\zeta_{26} &= -Q^2, \\
Q &= \frac{\phi'}{\phi - s\phi'}, \\
\Theta &= \frac{\phi\phi' - s(\phi\phi'' + \phi'\phi')}{2\phi(\phi - s\phi' + (B - s^2)\phi'')}, \\
\psi &= \frac{\phi''}{2\phi(\phi - s\phi' + (B - s^2)\phi'')}, \\
B &= b^2.
\end{aligned}$$

## 2.2 Homogeneous Finsler spaces

Let  $G$  be a Lie group and  $M$ , a smooth manifold. Then a smooth map  $f : G \times M \longrightarrow M$  satisfying

$$f(g_2, f(g_1, x)) = f(g_2g_1, x), \quad \text{for all } g_1, g_2 \in G, \quad \text{and } x \in M$$

is called a smooth action of  $G$  on  $M$ .

**DEFINITION 2.4.** *Let  $M$  be a smooth manifold and  $G$ , a Lie group. If  $G$  acts smoothly on  $M$ , then  $G$  is called a **Lie transformation group** of  $M$ .*

The following theorem gives us a differentiable structure on the coset space of a Lie group.

**Theorem 2.2.** *Let  $G$  be a Lie group and  $H$ , its closed subgroup. Then there exists a unique differentiable structure on the left coset space  $G/H$  with the induced topology that turns  $G/H$  into a smooth manifold such that  $G$  is a Lie transformation group of  $G/H$ .*

DEFINITION 2.5. Let  $(M, F)$  be a connected Finsler space and  $I(M, F)$  the group of isometries of  $(M, F)$ . If the action of  $I(M, F)$  is transitive on  $M$ , then  $(M, F)$  is said to be a **homogeneous Finsler space**.

Let  $G$  be a Lie group acting transitively on a smooth manifold  $M$ . Then for  $a \in M$ , the isotropy subgroup  $G_a$  of  $G$  is a closed subgroup and by Theorem 2.2,  $G$  is a Lie transformation group of  $G/G_a$ . Further,  $G/G_a$  is diffeomorphic to  $M$ .

**Theorem 2.3.** [11] Let  $(M, F)$  be a Finsler space. Then  $G = I(M, F)$ , the group of isometries of  $M$  is a Lie transformation group of  $M$ . Let  $a \in M$  and  $I_a(M, F)$  be the isotropy subgroup of  $I(M, F)$  at  $a$ . Then  $I_a(M, F)$  is compact.

Let  $(M, F)$  be a homogeneous Finsler space, i.e.,  $G = I(M, F)$  acts transitively on  $M$ . For  $a \in M$ , let  $H = I_a(M, F)$  be a closed isotropy subgroup of  $G$  which is compact. Then  $H$  is a Lie group itself being a closed subgroup of  $G$ . Write  $M$  as the quotient space  $G/H$ .

DEFINITION 2.6. [23] Let  $\mathfrak{g}$  and  $\mathfrak{h}$  be the Lie algebras of the Lie groups  $G$  and  $H$  respectively. Then the direct sum decomposition of  $\mathfrak{g}$  as  $\mathfrak{g} = \mathfrak{h} + \mathfrak{k}$ , where  $\mathfrak{k}$  is a subspace of  $\mathfrak{g}$  such that  $Ad(h)(\mathfrak{k}) \subset \mathfrak{k} \forall h \in H$ , is called a reductive decomposition of  $\mathfrak{g}$ , and if such decomposition exists, then  $(G/H, F)$  is called reductive homogeneous space.

Therefore, we can write, any homogeneous Finsler space as a coset space of a connected Lie group with an invariant Finsler metric. Here, the Finsler metric  $F$  is viewed as  $G$  invariant Finsler metric on  $M$ .

We can identify  $\mathfrak{k}$  with the tangent space  $T_H(G/H)$  of  $G/H$  at the origin  $H$ . Let  $\left\{w_1, w_2, w_3, \dots, w_n = \frac{w}{c}\right\}$  be the orthonormal basis of  $\mathfrak{k}$ , where  $w$  is the invariant vector field of length  $c$  corresponding to the 1-form  $\beta$ , i.e.,  $\langle w, Y \rangle = \beta(Y)$ ,  $\forall Y \in \mathfrak{k}$ , and  $\langle \cdot, \cdot \rangle$  is the restriction of Riemannian metric  $\alpha$  to  $\mathfrak{k}$ .

### 3 Ricci Curvature of Homogeneous Finsler Spaces with Certain $(\alpha, \beta)$ -Metrics

$\Gamma_{ij}^m$  are the Christoffel symbols given by

$$\nabla \frac{\partial}{\partial x^i} \frac{\partial}{\partial x^j} = \Gamma_{ij}^m \frac{\partial}{\partial x^m}.$$

To compute  $\Gamma_{ij}^k$ , we need following notations:

For  $1 \leq i, j, m \leq n$ , define the structure constants of  $\mathfrak{k}$  as follows:

$$C_{ij}^m = \langle [w_i, w_j]_{\mathfrak{k}}, w_m \rangle,$$

where  $[w_i, w_j]_{\mathfrak{k}}$  denotes the projection of  $[w_i, w_j]$  to  $\mathfrak{k}$ .

Take

$$\sum_{t=1}^n C_{ij}^t C_{ml}^t = C_{ij}^t C_{ml}^t,$$

and

$$\langle [w_i, w_j]_{\mathfrak{k}}, Y \rangle = C_{ij}^0$$

for any non-zero  $Y \in \mathfrak{k}$ .

Define

$$f(i, j) = \begin{cases} 1, & \text{if } i < j, \\ 0, & \text{if } i \geq j. \end{cases}$$

With above notations, let us recall following lemmas for later use:

LEMMA 3.1. [12] At the origin  $o = eH$ , we have:

- (i)  $\Gamma_{ij}^m = f(i, j)C_{ij}^m + \langle \nabla_{\hat{w}_i} \hat{w}_j, \hat{w}_m \rangle$ ,
- (ii)  $\langle \nabla_{\hat{w}_i} \hat{w}_j, \hat{w}_m \rangle = -\frac{1}{2} \left( C_{jm}^i + C_{im}^j + C_{ij}^m \right)$ ,
- (iii)  $\langle \nabla_{\hat{w}_i} \hat{w}_j, \hat{w}_m \rangle \hat{w}_l = \frac{1}{2} \left( C_{lt}^i C_{jm}^t + C_{lt}^j C_{im}^t + C_{lt}^m C_{ij}^t + C_{jm}^t C_{li}^t + C_{im}^t C_{lj}^t + C_{ij}^t C_{lm}^t \right)$ ,  
where  $w_i, w_j, w_m, w_l \in \mathfrak{k}$ .

LEMMA 3.2. [12] At the origin  $o = eH$ , we have:

$$\begin{aligned}
 b_k &= c\delta_{nk}, \\
 s_{jk} &= \frac{c}{2}C_{jk}^n, \\
 s_k &= \frac{c^2}{2}C_{nk}^n, \\
 r_{jk} &= \frac{c}{2} \left( C_{jn}^k + C_{kn}^j \right), \\
 s_{jk;m} &= \frac{c}{2} \left\{ -C_{jk}^t C_{mt}^n + \frac{1}{2}C_{jt}^n \left( -C_{km}^t + C_{kt}^m + C_{mt}^k \right) + \frac{1}{2}C_{tk}^n \left( -C_{jm}^t + C_{jt}^m + C_{mt}^j \right) \right\}, \\
 s_{j;k} &= c \left( s_{nj;k} + \frac{c}{2}C_{tj}^n \Gamma_{nk}^t \right), \\
 b_{j;k;m} &= c \left( -\Gamma_{nt}^j \langle \nabla_{\hat{w}_m} \hat{w}_k, \hat{w}_t \rangle - \Gamma_{nk}^t \langle \nabla_{\hat{w}_m} \hat{w}_j, \hat{w}_t \rangle + C_{mn}^t \langle \nabla_{\hat{w}_t} \hat{w}_k, \hat{w}_j \rangle \right. \\
 &\quad \left. + \hat{w}_m \langle \nabla_{\hat{w}_n} \hat{w}_k, \hat{w}_j \rangle \right), \\
 r_{jk;m} &= s_{jk;m} + b_{k;j;m}.
 \end{aligned}$$

Next, we calculate the quantities used in Theorem 2.1 at the origin for a homogeneous Finsler space using Lemma 3.1 and 3.2.

$$\begin{aligned}
 r_{00} &= cC_{0n}^0, \\
 s_j &= b^l s_{lj} = c s_{nj}, \\
 s_0 &= c s_{n0} = \frac{c^2}{2} C_{n0}^n, \\
 r_j &= b^l r_{lj} = c r_{nj}, \\
 r_0 &= c r_{n0} = -\frac{c^2}{2} (C_{n0}^n + C_{nn}^0) = \frac{c^2}{2} C_{0n}^n, \\
 r_{00;0} &= c C_{0i}^0 (C_{in}^0 + C_{0n}^i), \\
 r &= r_{ij} b^i b^j = c^2 r_{nn} = 0, \\
 r_l^l &= a^{li} r_{il} = -\frac{c}{2} \delta^{li} (C_{nl}^i + C_{ni}^l) = -\frac{c}{2} (C_{nl}^l + C_{nl}^l) = c C_{ln}^l, \\
 r_0^l &= a^{li} r_{i0} = \delta^{li} r_{i0} = r_{l0}, \\
 r_{0l} r_0^l &= r_{0l} r_{l0} = \left( -\frac{c}{2} \right)^2 (C_{nl}^0 + C_{n0}^l) (C_{n0}^l + C_{nl}^0) = \frac{c^2}{4} (C_{nl}^0 + C_{n0}^l)^2, \\
 r_{00;l} b^l &= r_{00;n} b^n = c r_{00;n} = c (s_{00;n} + b_{0;0;n}) = \frac{c^2}{2} (C_{0n}^t + C_{tn}^0) (C_{0t}^n + C_{nt}^0 + C_{0n}^t),
 \end{aligned}$$

$$\begin{aligned}
 r_{0l;0}b^l &= r_{0n;0}b^n = cr_{0n;0} = c(s_{0n;0} + b_{n;0;0}) \\
 &= \frac{c^2}{4} \{2 C_{nt}^n C_{t0}^0 + (C_{t0}^n + C_{tn}^0 + C_{0n}^t) (C_{nt}^0 + C_{n0}^t)\}, \\
 s_0^l &= a^{li} s_{i0} = \delta^{li} s_{i0} = s_{l0}, \\
 r_{0l}s_0^l &= r_{0l}s_{l0} = \frac{c^2}{4} (C_{nl}^0 C_{0l}^n + C_{n0}^l C_{0l}^n), \\
 s_{0;0} &= cs_{n0;0} + \frac{c^2}{2} C_{l0}^m \Gamma_{n0}^l = \frac{c^2}{2} C_{nt}^m C_{0t}^0, \\
 s_{0l}s_0^l &= s_{0l}s_{l0} = -\frac{c^2}{4} (C_{0l}^n)^2, \\
 s_l s_0^l &= s_l s_{l0} = \frac{c^3}{4} C_{nl}^m C_{l0}^m, \\
 s_l r_0^l &= s_l r_{l0} = -\frac{c^3}{4} C_{nl}^m (C_{nl}^0 + C_{n0}^l), \\
 r_l s_0^l &= r_l s_{l0} = \frac{c^3}{4} C_{ln}^m C_{l0}^m, \\
 s_{0;l}b^l &= cs_{0;n} = c \left( cs_{n0;n} + \frac{c^2}{2} C_{t0}^m \Gamma_{nn}^t \right) = \frac{c^3}{4} C_{nt}^m (C_{0n}^t + C_{nt}^0 + C_{0t}^m), \\
 s_{l;0}b^l &= cs_{n;0} = c \left( cs_{nn;0} + \frac{c^2}{2} C_{tn}^m \Gamma_{n0}^t \right) = -\frac{c^3}{4} C_{tn}^m (C_{0t}^n + C_{nt}^0 + C_{n0}^t), \\
 s_{0;l}^l &= a^{li} s_{i0;l} = \delta^{li} s_{i0;l} = s_{l0;l} = \frac{c}{4} \{2C_{t0}^n C_{lt}^l + C_{lt}^n (C_{0l}^t + C_{0t}^l + C_{lt}^0)\}, \\
 s_l s_0^l &= s_l s_{l0} = \frac{c^4}{4} (C_{nl}^m)^2, \\
 s_l^j s_j^l &= -\frac{c^2}{4} (C_{jl}^m)^2.
 \end{aligned}$$

Now, we calculate  $\zeta_1$  to  $\zeta_6$  for square metric as follows:

$$\begin{aligned}
 \zeta_1 &= -\frac{3(10s^4 - 8s^3 - (4B + 3)s^2 + (4 + 4B)s - 1 - 2B)}{(1 - 3s^2 + 2B)^3}, \\
 \zeta_2 &= -\frac{6(15s^6 + (-25B - 2)s^4 + (4B + 8B^2 - 3)s^2 + 2B^2 + B)}{(1 - 3s^2 + 2B)^4}, \\
 \zeta_3 &= -\frac{8(-24s^4 + 15s^3 + (10 + 8B)s^2 + (-9 - 6B)s + 4B + 2)}{(1 - s)(1 - 3s^2 + 2B)^3}, \\
 \zeta_4 &= \frac{-48(B - s^2)^2(-63s^6 + 117s^5 + (36B - 36)s^4 - (30 + 60B)s^3 + (4B^2 + 28B + 25)s^2 - (12B + 3 + 12B^2)s + 2 + 8B + 8B^2)}{(1 - s)^3(1 - 3s^2 + 2B)^5} \\
 &\quad + \frac{8(111s^4 - 168s^3 + (58 - 22B)s^2 - 12Bs + 7 + 18B + 8B^2)(B - s^2)}{(1 - 3s^2 + 2B)^3(1 - s)^3} \\
 &\quad - \frac{4(39s^4 - 51s^3 + (7 - 16B)s^2 + (9 + 6B)s + 2B + 4B^2)}{(1 - 3s^2 + 2B)^2(1 - s)^3},
 \end{aligned}$$



$$\zeta_5 = \frac{8(1-2s)}{(1-3s^2+2B)^2},$$

$$\zeta_6 = \frac{12s(4B-3s^2-1)}{(1-3s^2+2B)^3},$$

$$\zeta_7 = \frac{-1+2s}{1-3s^2+2B},$$

$$\zeta_8 = \frac{-6s(B-s^2)}{(1-3s^2+2B)^2},$$

$$\zeta_9 = \frac{-16(-27s^6 + (54B+42)s^4 - (20B+7)s^3 - (28B+20+36B^2)s^2 + (9+10B+8B^2)s - 1 + 8B^3 + 8B^2)}{(1-s)^3(1-3s^2+2B)^3},$$

$$\zeta_{10} = \frac{48(45s^4 - 60s^3 + (18-12B)s^2 + (-4-8B)s + 5 + 4B^2 + 12B)(B^2 - 2s^2B + s^4)}{(1-3s^2+2B)^4(1-s)^4}$$

$$- \frac{32(60s^4 - 72s^3 + (9-21B)s^2 + (4-4B)s + 5 + 13B + 6B^2)(B-s^2)}{(1-s)^4(1-3s^2+2B)^3}$$

$$+ \frac{16(-17s^3 + 2s^2 + (4B+8)s + 1 + 2B)}{(1-s)^3(1-3s^2+2B)^2}$$

$$+ \frac{4(55s^4 - 60s^3 + (10-52B)s^2 + (32B+12)s - 5 - 4B + 12B^2)}{(1-s)^4(1-3s^2+2B)^2},$$

$$\zeta_{11} = \frac{-4}{(1-3s^2+2B)^2},$$

$$\zeta_{12} = \frac{-32(1-2s)}{(1-s)(1-3s^2+2B)^2},$$

$$\zeta_{13} = \frac{8(21s^3 - 3s^2 - (26B-5)s + 2B + 1)}{(1-s)(1-3s^2+2B)^3},$$

$$\zeta_{14} = \frac{2}{1-3s^2+2B},$$

$$\zeta_{15} = \frac{8(1-2s)}{(1-s)(1-3s^2+2B)},$$

$$\zeta_{16} = \frac{2(21s^4 - 18s^3 + (-2-22B)s^2 + (-2+20B)s + 1 + 2B)}{(1-s)^2(1-3s^2+2B)^2},$$

$$\zeta_{17} = \frac{4(1-2s)}{(1-s)(1-3s^2+2B)},$$

$$\zeta_{18} = \frac{2(15s^4 - 18s^3 + (2-14B)s^2 + (2+16B)s - 2B - 1)}{(1-s)^2(1-3s^2+2B)^2},$$

$$\zeta_{19} = \frac{-4(1-3s)}{(1-s)^3},$$

$$\begin{aligned}
 \zeta_{20} &= \frac{16}{(1-s)(1-3s^2+2B)^2}, \\
 \zeta_{21} &= \frac{-16(1-2s)}{(1-s)^2(1-3s^2+2B)}, \\
 \zeta_{22} &= \frac{4(27s^3-3s^2-(26B+1)s+1+2B)}{(1-s)^2(1-3s^2+2B)^2}, \\
 \zeta_{23} &= \frac{4}{(1-s)(1-3s^2+2B)}, \\
 \zeta_{24} &= \frac{4}{1-s}, \\
 \zeta_{25} &= \frac{-16}{(1-s)^2(1-3s^2+2B)}, \\
 \zeta_{26} &= \frac{-4}{(1-s)^2}.
 \end{aligned} \tag{3.1}$$

From above calculations, we have the following:

**Theorem 3.1.** *Let  $G/H$  be a homogeneous Finsler space with square metric  $F$ . Then, the Ricci curvature is given by*

$$\begin{aligned}
 Ric(Z) = {}^\alpha Ric(Z) &+ \frac{c^2(C_{0n}^0)^2}{\alpha^2(Z)} \left\{ (n-1)\zeta_1 + \zeta_2 \right\} + \frac{c^3 C_{0n}^0 C_{n0}^n}{2\alpha(Z)} \left\{ (n-1)(\zeta_3 - \zeta_5) + \zeta_4 - \zeta_6 \right\} \\
 &- \frac{c C_{0l}^0 (C_{nl}^0 + C_{n0}^l)}{\alpha(Z)} \left\{ (n-1)\zeta_7 + \zeta_8 \right\} + \frac{c^4 (C_{n0}^n)^2}{4} \left\{ (n-1)(\zeta_9 - \zeta_{12}) + \zeta_{10} - \zeta_{11} - \zeta_{13} \right\} \\
 &- \frac{c^2}{4} \left\{ 4C_{0n}^0 C_{nl}^l + (C_{n0}^l + C_{nl}^0) (2C_{nl}^0 + C_{0l}^n + 2C_{0n}^l) + 2C_{nl}^n C_{l0}^0 \right\} \zeta_{14} \\
 &+ \frac{c^2}{4} C_{0l}^n (C_{n0}^l + C_{nl}^0) \left\{ (n-1)\zeta_{15} + \zeta_{16} \right\} + \frac{c^2}{2} C_{nl}^n C_{0l}^0 \left\{ (n-1)\zeta_{17} + \zeta_{18} \right\} \\
 &- \frac{c^2}{4} (C_{l0}^n)^2 \zeta_{19} + \frac{c^3}{4} \alpha(Z) C_{l0}^n C_{nl}^n \left\{ (n-1)\zeta_{21} + \zeta_{22} \right\} \\
 &+ \frac{c^3}{4} \alpha(Z) \left\{ 4C_{n0}^n C_{nl}^l - C_{nl}^n (4C_{nl}^0 - C_{0l}^n) \right\} \zeta_{23} + \frac{c}{4} \alpha(Z) \left\{ 2C_{t0}^n C_{lt}^l + C_{lt}^n C_{lt}^0 \right\} \zeta_{24} \\
 &+ \frac{c^2}{4} \alpha^2(Z) \left\{ c^2 (C_{nl}^n)^2 \zeta_{25} - (C_{li}^n)^2 \zeta_{26} \right\},
 \end{aligned}$$

where  $Z$  is a non-zero vector in  $\mathfrak{k}$ , and  $\zeta_1$  to  $\zeta_{26}$  are given by (3.1).

Next,  $\zeta_1$  to  $\zeta_{26}$  for Randers change of square metric are calculated as follows:

$$\begin{aligned}
 \zeta_1 &= \frac{3(-40s^8 - 192s^7 + (16B - 235)s^6 + (72B + 96)s^5 + (235 + 90B)s^4 - (73 + 20B)s^2 + 24sB + 18B + 9)}{4(1-3s^2+2B)^3(1+3s+s^2)^2}, \\
 \zeta_2 &= \frac{-6(9s^6 + (-13B - 2)s^4 + (2B^2 + 4B - 3)s^2 + B + 2B^2)}{(1-3s^2+2B)^4},
 \end{aligned}$$

$$\begin{aligned}
 \zeta_3 &= \frac{4(2s+3)(12s^6+54s^5+(73+8B)s^4+(48B-12)s^3}{(1-s^2)(1+3s+s^2)^2(1-3s^2+2B)^3} \\
 &\quad + \frac{(-42+78B)s^2+(6+48B)s+18B+9}{(1-s^2)^2(1-3s^2+2B)^2} \frac{(B-s^2)}{(1-s^2)(1+3s+s^2)(1-3s^2+2B)^2} \\
 &\quad - \frac{2(2s+3)(-3+9s^2+4s^3)s}{(1-s^2)^2(1-3s^2+2B)^2} \\
 &\quad - \frac{(2s+3)(40s^6+180s^5+(227+16B)s^4+(96B-48)s^3}{(1+3s+s^2)^2(1-3s^2+2B)^2(1-s^2)} \\
 &\quad \quad - (138-156B)s^2+(12+96B)s+36B+27)}{(1-s^2)^2(1+3s+s^2)(1-3s^2+2B)^2}, \\
 \zeta_4 &= -24 \left( \frac{(3+2s)(18s^2+2+4B+s-3s^3+2Bs)}{(1-3s^2+2B)^4(1-s^2)} \right. \\
 &\quad \left. + \frac{12(6s+2s^3+9s^2+3)s^2}{(1-s^2)^3(1-3s^2+2B)^5} \right) (B-s^2)^2 \\
 &\quad + 4 \left( \frac{10s^3-3+27s^2+6s}{(1-s^2)^2(1-3s^2+2B)^2} + \frac{2(6s+2s^3+9s^2+3)}{(1-s^2)^3(1-3s^2+2B)} \right. \\
 &\quad \left. + \frac{6(2s+3)(14s^2+1+2B)}{(1-3s^2+2B)^3(1-s^2)} + \frac{12s}{(1-3s^2+2B)^3} \right) (B-s^2) \\
 &\quad - \frac{2(4s^3-3+9s^2)}{(1-3s^2+2B)(1-s^2)^2} - \frac{12s(2+8s^2+15s)}{(1-3s^2+2B)^2(1-s^2)} \\
 &\quad - \frac{2(6s+2s^3+9s^2+3)}{(1-s^2)^3}, \\
 \zeta_5 &= \frac{-4(-3+9s^2+4s^3)}{(1+3s+s^2)(1-3s^2+2B)^2}, \\
 \zeta_6 &= \frac{12s(4B-3s^2-1)}{(1-3s^2+2B)^3}, \\
 \zeta_7 &= \frac{-3+9s^2+4s^3}{2(1+3s+s^2)(1-3s^2+2B)}, \\
 \zeta_8 &= \frac{-6s(B-s^2)}{(1-3s^2+2B)^2}, \\
 \zeta_9 &= \frac{8(3+2s)(-3+9s^2+4s^3)(B-s^2)}{(1-s^2)^3(1-3s^2+2B)^2} \\
 &\quad - \frac{4(3+2s)^2(12s^6+54s^5+(73+8B)s^4+(48B-12)s^3}{(1-s^2)^2(1+3s+s^2)^2(1-3s^2+2B)^3} \\
 &\quad \quad + (-42+78B)s^2+(6+48B)s+18B+9)(B-s^2)}{(3+2s)^2(3-9s^2-4s^3)^2} \\
 &\quad + \frac{(3+2s)^2(3-9s^2-4s^3)^2}{(1-s^2)^2(1+3s+s^2)^2(1-3s^2+2B)^2} \\
 &\quad + \frac{2(3+2s)^2(12s^6+54s^5+(73+8B)s^4+(48B-12)s^3}{(1-s^2)^2(1+3s+s^2)^2(1-3s^2+2B)^2} \\
 &\quad \quad + (-42+78B)s^2+(6+48B)s+18B+9)}{4(3+2s)(3-9s^2-4s^3)} - \frac{8(3+2s)(1+3s+s^2)}{(1-s^2)^3},
 \end{aligned}$$

$$\begin{aligned}
 \zeta_{10} &= \left\{ \frac{16 \left( 18s + 28s^2 + 18s^3 + 3s^4 + 8 \right)}{(1-s^2)^4 (1-3s^2+2B)^2} - \frac{144s^2(3+2s)^2}{(1-3s^2+2B)^4 (1-s^2)^2} \right. \\
 &\quad \left. + \frac{48(3+2s)(-24s^5 - 45s^4 + 12s^3 + (6B+30)s^2 + (4+8B)s + 3+6B)}{(1-3s^2+2B)^4 (1-s^2)^3} \right\} (B-s^2)^2 \\
 &\quad - \left\{ \frac{8(3+2s)(-108s^5 - 207s^4 + (8B+52)s^3 + (42B+138)s^2 + (16+32B)s + 9+18B)}{(1-s^2)^3 (1-3s^2+2B)^3} \right. \\
 &\quad \left. + \frac{16 \left( 18s + 28s^2 + 18s^3 + 3s^4 + 8 \right)}{(1-s^2)^4 (1-3s^2+2B)} \right. \\
 &\quad \left. + \frac{16 \left( -15s^4 - 27s^3 + (2B+10)s^2 + (6B+21)s + 1+2B \right)}{(1-3s^2+2B)^3 (1-s^2)^2} \right\} (B-s^2) \\
 &\quad + \frac{8(2+4s^2+9s)(-9s^4 - 18s^3 + (2B+4)s^2 + (6B+12)s + 1+2B)}{(1-s^2)^3 (1-3s^2+2B)^2} \\
 &\quad - \frac{4(3+2s)(-12s^3 - 15s^2 + (6+8B)s + 6+12B)}{(1-3s^2+2B)^2 (1-s^2)^2} + \\
 &\quad \frac{418s + 28s^2 + 18s^3 + 3s^4 + 8}{(1-s^2)^4}, \\
 \zeta_{11} &= \frac{-4}{(1-3s^2+2B)^2}, \\
 \zeta_{12} &= \frac{8(3+2s)(-3+9s^2+4s^3)}{(1+3s+s^2)(1-3s^2+2B)^2(1-s^2)}, \\
 \zeta_{13} &= \frac{8 \left( 21s^4 + 27s^3 - (26B-2)s^2 - (36B-9)s + 1+2B \right)}{(1-3s^2+2B)^3(1-s^2)}, \\
 \zeta_{14} &= \frac{2}{1-3s^2+2B}, \\
 \zeta_{15} &= \frac{2(2s+3)(3-9s^2-4s^3)}{(1-s^2)(1+3s+s^2)(1-3s^2+2B)}, \\
 \zeta_{16} &= \frac{2 \left( -21s^4 - 36s^3 + (22B-4)s^2 + 36sB + 2B+1 \right)}{(1-3s^2+2B)^2(1-s^2)}, \\
 \zeta_{17} &= \frac{(2s+3)(3-9s^2-4s^3)}{(1-s^2)(1+3s+s^2)(1-3s^2+2B)}, \\
 \zeta_{18} &= \frac{-2 \left( 15s^4 + 18s^3 - (14B+4)s^2 - 18Bs + 1+2B \right)}{(1-s^2)(1-3s^2+2B)^2}, \\
 \zeta_{19} &= \frac{2 \left( -7 + 27s^2 + 24s^3 + 6s^4 \right)}{(1-s^2)^3}, \\
 \zeta_{20} &= \frac{8(2s+3)}{(1-3s^2+2B)^2(1-s^2)}, \\
 \zeta_{21} &= \frac{2(2s+3)^2(-3+9s^2+4s^3)}{(1-s^2)^2(1+3s+s^2)(1-3s^2+2B)},
 \end{aligned}$$

$$\begin{aligned}
\zeta_{22} &= \frac{-2(2s+3)(-27s^4 - 36s^3 + (26B+4)s^2 + 36Bs - 1 - 2B)}{(1-3s^2+2B)^2(1-s^2)^2}, \\
\zeta_{23} &= \frac{2(2s+3)}{(1-s^2)(1-3s^2+2B)}, \\
\zeta_{24} &= \frac{2(2s+3)}{1-s^2}, \\
\zeta_{25} &= \frac{-4(9+4s^2+12s)}{(1-s^2)^2(1-3s^2+2B)}, \\
\zeta_{26} &= \frac{-4(9+4s^2+12s)}{(1-s^2)^2}.
\end{aligned} \tag{3.2}$$

**Theorem 3.2.** *Let  $F$  be Randers change of square metric on a homogeneous Finsler space  $G/H$ . Then, the Ricci curvature is given by*

$$\begin{aligned}
Ric(Z) &= {}^\alpha Ric(Z) + \frac{c^2(C_{0n}^0)^2}{\alpha^2(Z)} \left\{ (n-1)\zeta_1 + \zeta_2 \right\} + \frac{c^3 C_{0n}^0 C_{n0}^n}{2\alpha(Z)} \left\{ (n-1)(\zeta_3 - \zeta_5) + \zeta_4 - \zeta_6 \right\} \\
&\quad - \frac{c C_{0l}^0 (C_{nl}^0 + C_{n0}^l)}{\alpha(Z)} \left\{ (n-1)\zeta_7 + \zeta_8 \right\} + \frac{c^4 (C_{n0}^n)^2}{4} \left\{ (n-1)(\zeta_9 - \zeta_{12}) + \zeta_{10} - \zeta_{11} - \zeta_{13} \right\} \\
&\quad - \frac{c^2}{4} \left\{ 4C_{0n}^0 C_{nl}^l + (C_{n0}^l + C_{nl}^0) (2C_{nl}^0 + C_{0l}^n + 2C_{0n}^l) + 2C_{nl}^n C_{l0}^0 \right\} \zeta_{14} \\
&\quad + \frac{c^2}{4} C_{0l}^n (C_{n0}^l + C_{nl}^0) \left\{ (n-1)\zeta_{15} + \zeta_{16} \right\} + \frac{c^2}{2} C_{nl}^n C_{0l}^0 \left\{ (n-1)\zeta_{17} + \zeta_{18} \right\} \\
&\quad - \frac{c^2}{4} (C_{l0}^n)^2 \zeta_{19} + \frac{c^3}{4} \alpha(Z) C_{l0}^n C_{nl}^n \left\{ (n-1)\zeta_{21} + \zeta_{22} \right\} \\
&\quad + \frac{c^3}{4} \alpha(Z) \left\{ 4C_{n0}^n C_{nl}^l - C_{nl}^n (4C_{nl}^0 - C_{0l}^n) \right\} \zeta_{23} \\
&\quad + \frac{c}{4} \alpha(Z) \left\{ 2C_{i0}^n C_{it}^l + C_{it}^n C_{it}^0 \right\} \zeta_{24} + \frac{c^2}{4} \alpha^2(Z) \left\{ c^2 (C_{nl}^n)^2 \zeta_{25} - (C_{li}^n)^2 \zeta_{26} \right\},
\end{aligned}$$

where  $Z$  is a non-zero vector in  $\mathfrak{k}$ , and  $\zeta_1$  to  $\zeta_{26}$  are given by (3.2).

## 4 Ricci Curvature of Homogeneous Finsler Spaces with Certain $(\alpha, \beta)$ -Metrics Having Vanishing $S$ -Curvature

We have already given the formula for  $S$ -curvature of homogeneous Finsler space with  $G$ -invariant square metric in [29], and with  $G$ -invariant Randers change of square metric in [26]. Here, we give the equivalent condition for these spaces to have vanishing  $S$ -curvature.

**Theorem 4.1.** *Let  $G/H$  be a compact homogeneous Finsler space and  $F$  be a  $G$ -invariant square metric on  $G/H$ . Then  $(G/H, F)$  has vanishing  $S$ -curvature if and only if  $\langle [w, Y]_{\mathfrak{k}}, Y \rangle = 0 \forall Y \in \mathfrak{k}$  and  $w \in \mathfrak{k}$  corresponds to the 1-form  $\beta$ .*

*Proof.* We know that  $F$  has vanishing  $S$ -curvature if and only if

$$r_{ij} = 0, \text{ and } s_i = 0 \quad \forall 1 \leq i, j \leq n. \tag{4.1}$$

Therefore, we have to prove that the conditions in equation (4.1) are equivalent to  $\langle [w, Y]_{\mathfrak{k}}, Y \rangle = 0 \forall Y \in \mathfrak{k}$ .

Firstly, suppose that  $r_{ij} = 0$ , and  $s_i = 0 \quad \forall 1 \leq i, j \leq n$ . Using Lemma 3.2, we have

$$\frac{c}{2} \left( C_{in}^j + C_{jn}^i \right) = 0, \quad (4.2)$$

and

$$\frac{c^2}{2} C_{ni}^n = 0. \quad (4.3)$$

For  $i = j$ , equation (4.2) gives us  $\langle [w, w_i]_{\mathfrak{k}}, w_i \rangle = 0$ .

Since  $\left\{ w_1, w_2, w_3, \dots, w_n = \frac{w}{c} \right\}$  is the orthonormal basis of  $\mathfrak{k}$ , we have

$$\langle [w, Y]_{\mathfrak{k}}, Y \rangle = 0 \quad \forall Y \in \mathfrak{k}.$$

For the second part, suppose

$$\langle [w, Y]_{\mathfrak{k}}, Y \rangle = 0 \quad \forall Y \in \mathfrak{k}.$$

Therefore,

$$\langle [w, w_i]_{\mathfrak{k}}, w_i \rangle = 0 \quad \forall 1 \leq i \leq n, \quad (4.4)$$

$$\langle [w, w_i + w_j]_{\mathfrak{k}}, w_i + w_j \rangle = 0 \quad \forall 1 \leq i, j \leq n. \quad (4.5)$$

$$\langle [w, w + w_i]_{\mathfrak{k}}, w + w_i \rangle = 0 \quad \forall 1 \leq i \leq n. \quad (4.6)$$

From (4.4) and (4.5), we get  $C_{nj}^i + C_{ni}^j = 0$ , i.e.,  $r_{ij} = 0$ .

Further, from (4.4) and (4.6), we get  $\langle [w, w_i]_{\mathfrak{k}}, w \rangle = 0$ , i.e.,  $s_i = 0$ .  $\square$

Similarly, we can prove following theorem:

**Theorem 4.2.** *Let  $G/H$  be a compact homogeneous Finsler space and  $F$  be a  $G$ -invariant Randers change of square metric on  $G/H$ . Then  $(G/H, F)$  has vanishing  $S$ -curvature if and only if  $\langle [w, Y]_{\mathfrak{k}}, Y \rangle = 0 \quad \forall Y \in \mathfrak{k}$  and  $w \in \mathfrak{k}$  corresponds to the 1-form  $\beta$ .*

**Theorem 4.3.** *Let  $G/H$  be a compact homogeneous Finsler space with  $G$ -invariant square metric  $F$ . If  $(G/H, F)$  has vanishing  $S$ -curvature, then Ricci curvature is given by*

$$Ric(Z) = Ric^\alpha(Z) - \frac{c^2}{4} (C_{i0}^n)^2 \zeta_{19} + \frac{c}{4} \alpha(Z) \left( 2 C_{i0}^n C_{lt}^l + C_{lt}^n C_{lt}^0 \right) \zeta_{24} - \frac{c^2}{4} \alpha^2(Z) (C_{ik}^n)^2 \zeta_{26},$$

where  $Z$  is a non-zero vector in  $\mathfrak{k}$ , and  $\zeta_{19}$ ,  $\zeta_{24}$ ,  $\zeta_{26}$  are same as given in (3.1).

*Proof.* Since  $(G/H, F)$  has vanishing  $S$ -curvature, by Theorem 4.1,

$$\langle [w, Y]_{\mathfrak{k}}, Y \rangle = 0 \quad \forall Y \in \mathfrak{k}.$$

Therefore,

$$\begin{aligned} C_{nl}^n &= \langle [w_n, w_l]_{\mathfrak{k}}, w_n \rangle \\ &= \frac{1}{c^2} \left\{ \langle [w, w_l]_{\mathfrak{k}}, w \rangle + \langle [w, w_l]_{\mathfrak{k}}, w_l \rangle \right\} \\ &= \frac{1}{c^2} \langle [w, w_l]_{\mathfrak{k}}, w + w_l \rangle \\ &= \frac{1}{c^2} \langle [w, w]_{\mathfrak{k}} + [w, w_l]_{\mathfrak{k}}, w + w_l \rangle \\ &= \frac{1}{c^2} \langle [w, w + w_l]_{\mathfrak{k}}, w + w_l \rangle \\ &= 0 \quad \forall 1 \leq l \leq n. \end{aligned}$$

Further, for non-zero  $X \in \mathfrak{k}$ , we have

$$\begin{aligned}
 C_{n0}^n &= \langle [w_n, X]_{\mathfrak{k}}, w_n \rangle \\
 &= \frac{1}{c^2} \{ \langle [w, X]_{\mathfrak{k}}, w \rangle + \langle [w, X]_{\mathfrak{k}}, X \rangle \} \\
 &= \frac{1}{c^2} \langle [w, X]_{\mathfrak{k}}, w + X \rangle \\
 &= \frac{1}{c^2} \langle [w, w + X]_{\mathfrak{k}}, w + X \rangle = 0, \\
 C_{0n}^0 &= \langle [X, w_n]_{\mathfrak{k}}, X \rangle = \left\langle \left[ X, \frac{w}{c} \right]_{\mathfrak{k}}, X \right\rangle = 0,
 \end{aligned}$$

and

$$\begin{aligned}
 C_{nl}^0 + C_{n0}^l &= \langle [w_n, w_l]_{\mathfrak{k}}, X \rangle + \langle [w_n, X]_{\mathfrak{k}}, w_l \rangle \\
 &= \frac{1}{c} \left\{ \langle [w, w_l]_{\mathfrak{k}}, X \rangle + \langle [w, X]_{\mathfrak{k}}, w_l \rangle \right\} \\
 &= \frac{1}{c} \left\{ \langle [w, w_l]_{\mathfrak{k}}, X \rangle + \langle [w, X]_{\mathfrak{k}}, w_l \rangle + \langle [w, w_l]_{\mathfrak{k}}, w_l \rangle + \langle [w, X]_{\mathfrak{k}}, X \rangle \right\} \\
 &= \frac{1}{c} \left\{ \langle [w, w_l]_{\mathfrak{k}}, X + w_l \rangle + \langle [w, X]_{\mathfrak{k}}, X + w_l \rangle \right\} \\
 &= \frac{1}{c} \left\{ \langle [w, w_l]_{\mathfrak{k}} + [w, X]_{\mathfrak{k}}, X + w_l \rangle \right\} \\
 &= \frac{1}{c} \langle [w, X + w_l]_{\mathfrak{k}}, X + w_l \rangle \\
 &= 0.
 \end{aligned}$$

Using the above values in Theorem 3.1, we get the required result.  $\square$

In similar manner, we can prove the following theorem:

**Theorem 4.4.** *Let  $G/H$  be a compact homogeneous Finsler space and  $F$  be a  $G$ -invariant Randers change of square metric on  $G/H$ . If  $(G/H, F)$  has vanishing  $S$ -curvature, then Ricci curvature is given by*

$$Ric(Z) = Ric^\alpha(Z) - \frac{c^2}{4} (C_{i0}^n)^2 \zeta_{19} + \frac{c}{4} \alpha(Z) \left( 2 C_{i0}^n C_{it}^l + C_{it}^n C_{it}^0 \right) \zeta_{24} - \frac{c^2}{4} \alpha^2(Z) (C_{ik}^n)^2 \zeta_{26},$$

where  $Z$  is a non-zero vector in  $\mathfrak{k}$ , and  $\zeta_{19}$ ,  $\zeta_{24}$ ,  $\zeta_{26}$  are same as given in (3.2).

**Corollary 4.5.** *Let  $G/H$  be a homogeneous Finsler space and  $F$  be a  $G$ -invariant square metric or Randers change of square metric on  $G/H$ . If  $(G/H, F)$  has vanishing  $S$ -curvature and negative Ricci curvature, then it must be Riemannian.*

## Acknowledgements

First author is very much thankful to UGC for providing financial assistance in terms of JRF fellowship vide letter with Sr. No. 2061641032 and Ref. No. 19/06/2016(i)EU-V.

## References

- [1] Amari, S. I.; Nagaoka, H., *Methods of Information Geometry*, Translations of Mathematical Monographs, AMS, 191, Oxford Univ. Press, 2000.

- [2] Antonelli, P. L.; Ingarden, R. S.; Matsumoto, M., *The Theory of Sprays and Finsler spaces with Applications in Physics and Biology*, Vol. 58, Springer Science & Business Media, 2013.
- [3] Bao, D.; Chern, S. S.; Shen, Z., *An Introduction to Riemann-Finsler Geometry*, Springer-Verlag, New York, 2000.
- [4] Bao, D.; Robles, C., Ricci and flag curvatures in Finsler geometry, *A Sampler of Riemann-Finsler Geometry*, **50** (2004), 197-259.
- [5] Bao, D.; Robles, C.; Shen, Z., Zermelo navigation on Riemannian manifolds, *Journal of Differential Geometry*, **66**(3) (2004), 377-435.
- [6] Berwald, L., Über  $n$ -dimensionalen Geometrien konstanter Krümmung, in denen die Geraden die kürzesten sind, *Mathematische Zeitschrift*, **30** (1) (1929), 449-469.
- [7] Besse, A., *Einstein Manifolds*, Springer-Verlag, 1987.
- [8] Cheng, X.; Shen, Z.; Tian, Y., A class of Einstein  $(\alpha, \beta)$ -metrics, *Israel Journal of Mathematics*, **192** (2012), 221-249.
- [9] Chern, S. S.; Shen, Z., *Riemann-Finsler Geometry*, Nankai Tracts in Mathematics, Vol. 6, World Scientific, Singapore, 2005.
- [10] Deng, S., The  $S$ -curvature of homogeneous Randers spaces, *Differential Geometry and its Applications*, **27** (2009), 75-84.
- [11] Deng, S., *Homogeneous Finsler Spaces*, Springer Monographs in Mathematics, New York, 2012.
- [12] Deng, S.; Hu, Z., Curvatures of homogeneous Randers spaces, *Advances in Mathematics*, **240** (2013), 194-226.
- [13] Deng, S.; Kertész, D. C.; Yan, Z., There are no proper Berwald-Einstein manifolds, *Publicationes Mathematicae Debrecen*, **86** (1-2) (2015), 245-249.
- [14] Deng, S.; Wang, X., The  $S$ -curvature of homogeneous  $(\alpha, \beta)$ -metrics, *Balkan Journal of Geometry and Its Applications*, **15** (2) (2010), 39-48.
- [15] Gardner, R.; Wilkens, G., A pseudo-group isomorphism between control systems and certain generalized Finsler structures, *Contemporary Mathematics*, **196** (1996), 231-244.
- [16] Hashiguchi, M.; Ichijyō, Y., Randers spaces with rectilinear geodesics, *Rep. Fac. Sci., Kagoshima Univ., (Math., Phys. & Chem.)*, **13** (1980), 33-40.
- [17] Hu, Z.; Deng, S., Homogeneous Randers spaces with isotropic  $S$ -curvature and positive flag curvature, *Mathematische Zeitschrift*, **270** (2012), 989-1009.
- [18] Huang, L., On the fundamental equations of homogeneous Finsler spaces, *Differential Geometry and its Applications*, **40** (2015), 187-208.
- [19] Kron, G., Non-Riemannian dynamics of rotating electrical machinery, *Studies in Applied Mathematics*, **13** (1934), 103-194.
- [20] Matsumoto, M., On C-reducible Finsler-spaces, *Tensor, N. S.*, **24** (1972), 29-37.
- [21] Matsumoto, M., On Finsler spaces with Randers metric and special forms of important tensors, *Journal of mathematics of Kyoto University*, **14** (3) (1974), 477-498.



- [22] Mo, X.; Yu, C. T., On the Ricci curvature of a Randers metric of isotropic  $S$ -curvature, *Acta Mathematica Sinica, English Series*, **24** (6) (2008), 911-916.
- [23] Nomizu, K., Invariant affine connections on homogeneous spaces, *Amer. J. Math.*, **76** (1954), 33-65.
- [24] Rafie-Rad, M.; Rezaei, B., On Einstein Matsumoto metrics, *Nonlinear Analysis: Real World Applications*, **13**(2) (2012), 882-886.
- [25] Randers, G., On an asymmetric metric in the four-space of general relativity, *Phys. Rev.*, **59** (1941), 195-199.
- [26] Rani, S., Shanker, G., On  $S$ -curvature of a homogeneous Finsler space with Randers changed square metric, *Facta Universitatis, Series: Mathematics and Informatics*, **35** (3) (2020), 673-691.
- [27] Rezaei, B.; Razavi, A.; Sadeghzadeh, N., On Einstein  $(\alpha, \beta)$ -metrics, *Iranian Journal of Science and Technology*, **32** (4) (2008), 403-412.
- [28] Shanker, G.; Kaur, K., Homogeneous Finsler space with infinite series  $(\alpha, \beta)$ -metric, *Applied Sciences*, **21** (2019), 220-236.
- [29] Shanker, G., Rani, S., On  $S$ -Curvature of a Homogeneous Finsler space with square metric, *International Journal of Geometric Methods in Modern Physics*, **17** (02) (2020), 2050019 (16 pages).
- [30] Shanker, G., Rani, S., Kaur, K., Dually flat Finsler spaces associated with Randers- $\beta$  change, *Journal of Rajasthan Academy of Physical Sciences*, **18** (1-2), 95-106 (2019)
- [31] Shen, Z., On projectively related Einstein metrics in Riemann-Finsler geometry, *Mathematische Annalen*, **320**(4) (2001), 625-647.
- [32] Shen, Z.; Yu, C., On Einstein square metrics, *Publicationes Mathematicae Debrecen*, **85** (3-4) (2014), 413-424.
- [33] Wang, H.; Deng, S., Some Einstein-Randers metrics on homogeneous spaces, *Nonlinear Analysis*, **72** (2010), 4407-4414.
- [34] Xia, Q., On a class of projectively flat Finsler metrics, *Differential Geometry and its Applications*, **44** (2016), 1-16.
- [35] Xu, M.; Deng, S., Homogeneous  $(\alpha, \beta)$ -spaces with positive flag curvature and vanishing  $S$ -curvature, *Nonlinear Analysis*, **127** (2015), 45-54.
- [36] Yan, Z.; Deng, S., On homogeneous Einstein  $(\alpha, \beta)$ -metrics, *Journal of Geometry and Physics*, **103** (2016), 20-36.
- [37] Yu, C.; Zhu, H., Projectively flat general  $(\alpha, \beta)$ -metrics with constant flag curvature, *Journal of Mathematical Analysis and Applications*, **429** (2) (2015), 1222-1239.
- [38] Yu, Y.; You, Y., On Einstein  $m^{th}$  root metrics, *Differential Geometry and its Applications*, **28** (3) (2010), 290-294.
- [39] Zhou, L., A local classification of a class of  $(\alpha, \beta)$ -metrics with constant flag curvature, *Differential Geometry and its Applications*, **28** (2010), 170-193.