

Specific neutral and charged black holes in $f(R)$ gravitational theory

G. G. L. Nashed*

*Centre for Theoretical Physics, The British University,
P.O. Box 43, El Sherouk City, Cairo 11837, Egypt*

Shin'ichi Nojiri†

*Department of Physics, Nagoya University, Nagoya 464-8602
ε3*

*Japan Kobayashi-Maskawa Institute for the Origin of Particles and the Universe,
Nagoya University, Nagoya 464-8602, Japan*

(Dated: August 18, 2021)

With the successes of $f(R)$ theory as a neutral modification of Einstein's general relativity (GR), we continue our study in this field and attempt to find general natural and charged black hole (BH) solutions. In the previous papers [54, 55], we applied the field equation of the $f(R)$ gravity to a spherically symmetric space-time $ds^2 = -U(r)dt^2 + \frac{dr^2}{V(r)} + r^2(d\theta^2 + \sin^2\theta d\phi^2)$ with unequal metric potentials $U(r)$ and $V(r)$ and with/without electric charge. Then we have obtained equations which include all the possible static solutions with spherical symmetry. To ensure the closed form of system of the resulting differential equations in order to obtain specific solutions, we assumed the derivative of the $f(R)$ with respect to the scalar curvature R to have a form $F_1(r) = \frac{df(R(r))}{dR(r)} \propto \frac{c}{r^n}$ but in case $n > 2$, the resulting black hole solutions with/without charge do not generate asymptotically GR BH solutions in the limit $c \rightarrow 0$ which means that the only case that can generate GR BHs is $n = 2$. In this paper, we assume another form, i.e., $F_1(r) = 1 - \frac{F_0 - (n-3)}{r^n}$ with a constant F_0 and show that we can generate asymptotically GR BH solutions for $n > 2$ but we show that the $n = 2$ case is not allowed. This form of $F_1(r)$ could be the most acceptable physical form that we can generate from it physical metric potentials that can have a well-known asymptotic form and we obtain the metric of the Einstein general relativity in the limit of $F_0 \rightarrow n - 3$. We show that the form of the electric charge depends on n and that $n \neq 2$. Our study shows that the power n is sensitive and why we should exclude the case $n = 2$ for the choice of $F_1(r)$ presented in this study. We also study the physics of these black hole solutions by calculating their thermodynamical quantities, like entropy, the Hawking temperature and Gibb's free energy, and derive the stability conditions by using geodesic deviations.

arXiv:2103.02382v5 [gr-qc] 17 Aug 2021

* nashed@bue.edu.eg

† nojiri@gravity.phys.nagoya-u.ac.jp

I. INTRODUCTION

The $f(R)$ gravity is a modified gravitational theory where the action is given in a generic form of scalar curvature, R and the $f(R)$ gravity may describe the early and late cosmological evolution. The $f(R)$ theory can describe dark energy and dark matter consistently without imposing any new material that has not yet been revealed by experiments [1–6]. Amending the Lagrangian of general relativity (GR) does not only influence the dynamical system of the universe but it can also change the dynamical system at the galactic or solar system scales. Thus, amended theories of gravity with higher-order curvature corrections provide a deeper understanding of gravity.

The Einstein-Hilbert action, which reproduces the field equation of GR, is linear to the Ricci scalar R . By changing the action to include the non-linear terms, the Ricci curvature and/or Riemann curvatures, many viable modified gravitational theories are presented by the scientific society to describe the cosmic evolution in early times. Most of those theories use a gravitational Lagrangian which contains some of the four possible second-order curvature invariants. Moreover, many models that use higher-order invariants as a function of R are introduced in the gravitational action and different $f(R)$ gravitational models are obtained [7–19]. Aside from the ability of these theories to eliminate the contributions of curvature invariants other than the Ricci scalar R , they could also prevent the Ostrogradski issue [20], which is a problem that characterizes any higher derivative theories [21].

The earliest modification of GR could be the one presented in [22]. A natural extension of GR is to include expressions such as R^n with a constant n similar to the Starobinsky model $f(R) = R + \epsilon R^2$, where ϵ is also a constant [23]. When $n < 0$, the contribution of R^n could investigate the late epoch and describe self-accelerating vacuum solutions [24–27]. Nevertheless, such solutions suffer from instabilities [28, 29] and have strong limitations from the solar system test [30]. To avoid the previously mentioned issues, scientists have considered the $f(R)$ gravitational theory, which can accommodate a wide range of phenomena. Many applications carried use the framework of the $f(R)$ gravitational theory such as gravitational wave detection [31, 32], early-time inflation [33], cosmological phases [34–36], the singularity problem [37], stability of solutions [38–40], and other different issues [41].

Many black hole (BH) solutions in the $f(R)$ theory coincide with the BH solutions of GR or differ from them. Among these solutions, the authors in [42] derived static spherically symmetric solutions and showed that the de Sitter(dS)-Schwarzschild metric is a solution to the field equations of the $f(R)$ theory. Moreover, spherically symmetric solutions are derived in [43] by involving a perfect fluid. The BH solution with/without electric charge is presented in [44]. Many analytic spherically symmetric solutions are derived in [45–52]. Analytic charged D -dimensional BH solutions are derived and discussed in [53]. Moreover, analytic spherically symmetric BH solutions with/without electric charge are derived in [54, 55]. Those BH solutions were different from the BH solutions of GR and coincide with them under certain special conditions. The study in this paper aims to generalize this special condition and derive analytic BH solutions with/without electric charge in the $f(R)$ gravitational theory.

This paper is organized as follows: In Section II, we give the building block of the $f(R)$ gravitational theory and obtain its field equations including the contribution from the Maxwell field. In Section III, we apply the charged field equations of the $f(R)$ gravity to a space-time having spherically symmetric and unequal metric potentials. We present the non-linear differential equations which are composed of five non-linear differential equations having four unknown functions, one is the electric charge, one is the derivative $f'(R)$ of the $f(R)$, and the other two are related to the metric potentials. We study special cases that give a consistent BH solution which was derived in the previous literature. Then, we study the general case and divide it into two classes: The first class is the one without charge and derive an original new BH solution assuming the first derivative of $f(R)$ concerning R to has the form $f'(R) = 1 - \frac{F_0 - (n-3)}{r^n}$ where n can be any value. The second class is the one with the electric charge, and we derive a new charged BH solution assuming that the derivative $f'(R)$ is not changed from that in the case without electric charge. In Section IV, we study the physical properties of these new BH solutions with/without charge by giving the form of the metric potentials in asymptotic form and show that they are different from GR metric potentials either the Schwarzschild or the Reissner-Nordström due to the contribution of the non-linear curvature scalar terms. We also study the scalars of the BH solution and show that its peculiarity is softer than that of GR for the case with an electric charge. In Section V, we present the stability constraints of those BH solutions by using geodesic deviation and investigate the regions of stability graphically. In Section VI, we evaluate the basic thermodynamical expressions, that is, the Hawking-temperature, entropy, quasi-local energy, heat capacity, and Gibb's free energy, related to our new BH solutions and show that the solutions are physically acceptable. In subsection VIB, we explain that the new BH solutions fulfill the first law of thermodynamics. In the final section, we discuss our derived results.

II. $f(R)$ AMENDED THEORY

If $f(R) \neq R$ then, we have a modified gravitational theory that is unlike GR. The four dimensional action of the $f(R)$ gravitational theory takes the following form [1, 25–27, 56–60],

$$\mathcal{S} := \frac{1}{2\kappa} \int d^4x \sqrt{-g} f(R) + \int d^4x \sqrt{-g} \mathcal{L}_{\text{em}}, \quad (1)$$

with g being the determinant of the metric and κ being the Newtonian gravitational constant. The Maxwell electromagnetic field Lagrangian \mathcal{L}_{em} is given by $\mathcal{L}_{\text{em}} = \frac{1}{4}F^2 \equiv F_{\mu\nu}F^{\mu\nu}$ where $F_{\mu\nu} = \partial_\mu\eta_\nu - \partial_\nu\eta_\mu$ and $\eta = \eta_\mu dx^\mu$ is the electromagnetic Maxwell gauge potential 1-form [61].

Using the variations principle of the action (1), we obtain the field equations of the $f(R)$ gravity in the following form [62],

$$R_{\mu\nu}f_R - \frac{1}{2}g_{\mu\nu}f(R) + [g_{\mu\nu}\square - \nabla_\mu\nabla_\nu]f_R = -\frac{1}{2}\kappa T_{\mu\nu}^{\text{em}}, \quad (2)$$

where \square is the d'Alembertian operator, and $f_R = \frac{df}{dR}$ and $T_{\mu\nu}^{\text{em}}$ is the energy-momentum tensor of the Maxwell field defined as

$$T_{\mu\nu}^{\text{em}} = F_{\mu\alpha}F_\nu^\alpha - \frac{1}{4}g_{\mu\nu}F^2. \quad (3)$$

Furthermore, by the variation of equation (1) with respect to the gauge potential, η_μ , we obtain

$$\partial_\nu(\sqrt{-g}F^{\mu\nu}) = 0. \quad (4)$$

Taking the trace of equations (2), we find

$$3\square f_R + Rf_R - 2f(R) = 0. \quad (5)$$

Using Eq. (5) we obtain the form of the $f(R)$ as follows,

$$f(R) = \frac{1}{2}[3\square f_R + Rf_R]. \quad (6)$$

From Eqs. (6) and (2), we obtain [63]

$$R_{\mu\nu}f_R - \frac{1}{4}g_{\mu\nu}Rf_R + \frac{1}{4}g_{\mu\nu}\square f_R - \nabla_\mu\nabla_\nu f_R + \frac{1}{2}\kappa T_{\mu\nu}^{\text{em}} = 0. \quad (7)$$

Therefore, a significant step is to test Eqs. (4) and (7) to a spherically symmetric space-time whose line element has two different unknown functions.

III. SPHERICALLY SYMMETRIC BH SOLUTIONS

The spherically symmetric line-element is assumed to be given by

$$ds^2 = -U(r)dt^2 + \frac{dr^2}{V(r)} + r^2(d\theta^2 + \sin^2\theta d\phi^2), \quad (8)$$

where $U(r)$ and $V(r)$ are two unknowns of r . For the space-time (8), the Ricci scalar is evaluated as,

$$R(r) = \frac{r^2VV''^2 - r^2UU'V' - 2r^2UVU'' - 4rU[VU' - UV'] + 4U^2(1 - V)}{2r^2U^2}, \quad (9)$$

where $U \equiv U(r)$, $V \equiv V(r)$, $U' = \frac{dU}{dr}$, $U'' = \frac{d^2U}{dr^2}$ and $V' = \frac{dV}{dr}$. Plugging Eqs. (5) and (7) with Eq. (8) and by using Eq. (9), we obtain the (t, t) , (r, r) , and (θ, θ) (or (ϕ, ϕ)) components of the $f(R)$ equation are given by

$$0 = \frac{1}{8r^2W^2} \left\{ r^2 \left[VF_1W'^2 - 3WF_1V'W' - 2WVF_1W'' - 2W^2F_1V'' - 3WVW'F_1' - 2W^2V'F_1' + 2VW^2F_1'' \right] - 4rWV[F_1W' - WF_1'] - 4W^2F_1[1 - V]' - 8Wr^2\eta'^2 \right\}, \quad (10)$$

$$0 = \frac{1}{8r^2W^2} \left\{ r^2 \left[VF_1W'^2 - 3WF_1V'W' - 2WVF_1W'' - 2W^2F_1V'' + WVW'F_1' - 2W^2V'F_1' - 6VW^2F_1'' \right] + 4rWV[F_1W' + WF_1'] - 4W^2F_1[1 - V] - 8Wr^2\eta'^2 \right\}, \quad (11)$$

$$0 = \frac{1}{8r^2W^2} \left\{ r^2 \left[3WF_1V'W' + 2WVF_1W'' + 2W^2F_1V'' - VF_1W'^2 + WVW'F_1' + 2W^2V'F_1' + 2VW^2F_1'' \right] - 4rW^2VF_1' + 4W^2F_1[1 - V] + 8Wr^2\eta'^2 \right\}. \quad (12)$$

Other components of the $f(R)$ equation vanish. The trace of the $f(R)$ equation is given by

$$0 = \frac{1}{2r^2W^2} \left\{ r^2 \left[6W^2V'F_1' - 3WF_1V'W' - 2WVF_1W'' - 2W^2F_1V'' + VF_1W'^2 + 3WVW'F_1' + 6VW^2F_1'' \right] + 4rW[3WVW'F_1' - F_1VW' - 2F_1WV'] + 4W^2F_1[1 - V] - 4r^2W^2f(r) \right\}, \quad (13)$$

where $W(r) = \frac{U(r)}{V(r)}$ and $F_1 \equiv F_1(r) = \frac{df(R(r))}{dR(r)}$, $F_1' = \frac{dF_1(r)}{dr}$, $F_1'' = \frac{d^2F_1(r)}{dr^2}$, $F_1''' = \frac{d^3F_1(r)}{dr^3}$. The Maxwell field has the following non-vanishing components:

$$\frac{\eta' [rW' - 4W] - 2rW\eta''}{2rW^2} = 0, \quad (14)$$

where η is the component of the electric field, i.e., $\eta_\alpha = (\eta(r), 0, 0, 0)$. By using the trace equation (13), Eqs. (10), (11), and (12) can be rewritten as follows,

$$0 = r^2 \left[VF_1W'^2 - 3WF_1V'W' - 2WVF_1W'' - 2W^2F_1V'' - 3WVW'F_1' - 2W^2V'F_1' + 2VW^2F_1'' \right] - 4rWV[F_1W' - WF_1'] - 4W^2F_1[1 - V] - 8Wr^2\eta'^2, \quad (15)$$

$$0 = r^2 \left[VF_1W'^2 - 3WF_1V'W' - 2WVF_1W'' - 2W^2F_1V'' + WVW'F_1' - 2W^2V'F_1' - 6VW^2F_1'' \right] + 4rWV[F_1W' + WF_1'] - 4W^2F_1[1 - V] - 8Wr^2\eta'^2, \quad (16)$$

$$0 = r^2 \left[-VF_1W'^2 + 3WF_1V'W' + 2WVF_1W'' + 2W^2F_1V'' + WVW'F_1' + 2W^2V'F_1' + 2VW^2F_1'' \right] - 4rW^2VF_1' + 4W^2F_1[1 - V] + 8Wr^2\eta'^2. \quad (17)$$

Using Eqs. (15) and (16), i.e., (15) minus (16), we obtain

$$0 = r^2 [-4WVW'F_1' + 8VW^2F_1''] - 8rWW'VF_1. \quad (18)$$

Moreover, Eqs. (15) and (17), i.e., (15) plus (17) give:

$$0 = -2r^2WVW'F_1' + 4r^2VW^2F_1'' - 4rWW'VF_1. \quad (19)$$

A careful check shows that Eqs. (18) and (19) coincide with each other. Therefore we derive two independent equations from Eqs. (15), (16), and (17). From the above calculations, it is easy to prove that Eq. (15) is equal Eq. (16) with minus sign and equal minus two times Eq. (17). Hence, Equations (15) and (19) are independent equations, which include all the possible solutions. Now, we have four unknown functions V , W , η , and F , which is the reason why we are not able to determine one function¹. In order to obtain concrete and specific solutions and investigate the physical properties of the solutions in order to show that Eqs. (15) and (19) include physically reasonable and natural solutions, we make an assumption of the form on F in the following.

¹ Note that Eq. (14) can determine the unknown η .

In our previous studies, we showed that when $W = 1$ and providing $V \neq 0$, we get from Eq. (18)

$$F_1'' = 0, \quad \text{that leads,} \quad F_1 = f_2 + f_3 r. \quad (20)$$

From Eqs. (20) and (15) we obtain

$$\begin{aligned} 0 &= r^2 [-2F_1 V'' - 2V' F_1'] + 4rV F_1' - 4F_1 [1 - V] - 8r^2 \eta'^2 \\ &= -2r^2 (f_2 + f_3 r) V'' - 2r^2 f_3 V' + 4(f_2 + 2f_3 r) V - 4(f_2 + f_3 r) - 8r^2 \eta'^2. \end{aligned} \quad (21)$$

Assuming $f_3 = 0$, Eq. (21) gives

$$0 = f_2 [r^2 V'' - 2V + 2] + 4r^2 \eta'^2. \quad (22)$$

From Eq. (22) after using Eq. (14), we obtained the following solution

$$V = 1 + \frac{V_0}{r} + V_1 r^2 - \frac{V_2}{f_2 r^2}, \quad \eta = \frac{V_2}{r}, \quad (23)$$

where V_0 , V_1 , and V_2 are integration constants. Equation (23) is the well-known Reissner-Nordström-(anti-)de Sitter space-time.

We studied the case $f_2 = 0$ in which Eq. (21) gives,

$$0 = -2r^2 f_3 r V'' - 2r^2 f_3 V' + 8f_3 r V - 4f_3 r - 8r^2 \eta'^2. \quad (24)$$

The solution of Eq. (24) together with Eq. (14) have the following solution

$$V = \frac{1}{2} + \frac{\tilde{V}_0 r^2}{20} + \frac{\tilde{V}_1}{r^2} + \frac{\tilde{V}_2}{2r^3}, \quad \eta = \frac{\sqrt{5f_3 \tilde{V}_2}}{2r}. \quad (25)$$

Here \tilde{V}_0 , \tilde{V}_1 , and \tilde{V}_2 are integration constants, again. Equation (25) corresponds to the solution derived in [64, 65].

In [64, 65], we also solved the system of differential equations (13) by assuming $F_1 = 1 + \frac{c}{r^2}$. In the present paper, we solve such a system by assuming

$$F_1 = 1 - \frac{F_0 - (n-3)}{r^n}, \quad (26)$$

where n can take any value and F_0 is a constant. Using Eq. (26) in the system (15), (16), and (17), we obtain

$$\begin{aligned} 0 &= r^2 (r^n + F_0) \left[2W^2 V'' + 2V W W'' - V W'^2 \right] + r W W' \left[3r (r^2 + F_0) V' + V (4r^n + 4F_0 - 3F_0 n) \right] \\ &\quad - 2W \left[r n F_0 W V' - 4r^{2+n} \eta'^2 + 2W r^n (V - 1) - F_0 (2 - V (n^2 - n + 2)) \right], \end{aligned} \quad (27)$$

$$\begin{aligned} 0 &= r^2 (r^n + F_0) \left[2W^2 V'' + 2V W W'' - V W'^2 \right] + r W W' \left[3r (r^2 + F_0) V' - V (4r^n + 4F_0 - 3F_0 n) \right] \\ &\quad - 2W \left[r n F_0 W V' - 4r^{2+n} \eta'^2 + W [2r^n (1 - V)] - F_0 (2 + V (n^2 + 5n - 2)) \right], \end{aligned} \quad (28)$$

$$\begin{aligned} 0 &= r^2 (r^n + F_0) \left[2W^2 V'' + 2V W W'' - V W'^2 \right] + r W W' \left[3r (r^2 + F_0) V' - n F V \right] \\ &\quad - 2W \left[r n F_0 W V' - 4r^{2+n} \eta'^2 - W [2r^n (1 - V)] + F_0 (2 + V (n^2 + 3n - 2)) \right]. \end{aligned} \quad (29)$$

We then solve the above system of differential equations in cases $\eta = 0$ and $\eta \neq 0$.

A. The case of $\eta = 0$

The analytic solution of the above system takes the following form,

$$\begin{aligned} V(r) &= \frac{r^{\frac{2(n^2+2n-2)}{n-2}}}{X_1^{\frac{2(n+1)}{n-2}}} \left[c_2 + \int \frac{X_1^{\frac{n+1}{n-2}}}{r^{\frac{7n-8}{n-2}} X_2} dr \left(c_1 - 2 \int \frac{X_1^{\frac{n+1}{n-2}} X_2}{c_3 r^{\frac{n(2n-1)}{n-2}}} dr \right) + 2 \int \frac{X_2 X_1^{\frac{n+1}{n-2}} \int \frac{X_1^{\frac{n+1}{n-2}}}{r^{\frac{7n-8}{n-2}} X_2} dr}{c_3 r^{\frac{n(2n-1)}{n-2}}} dr \right], \\ W(r) &= c_3 r^{\frac{2n(n+1)}{2-n}} X_1^{\frac{2(n+1)}{n-2}}, \quad U(r) = W(r)V(r), \quad F_1 = 1 - \frac{F_0 - (n-3)}{r^n}, \end{aligned} \quad (30)$$

where $X_1(r) = \frac{F_0(n-2)}{2} - \frac{(n-2)(n-3)}{2} + r^n$ and $X_2(r) = F_0 - n + 3 - r^n$.

As Eq. (30) shows, the case of $n = 2$ is not allowed. However this case was studied in [54] and we obtained

$$V(r) = \frac{e^{\frac{3c_1}{2r^2}}}{r} \left\{ \text{H}c_2 + \text{H}_1 r^3 c_3 + 2\text{H}_1 r^3 \int \frac{e^{-\frac{3c_1}{2r^2}} \text{H}}{r [(2c_1 \text{H}_2 - 3r^2 \text{H}) \text{H}_1 - 2c_1 \text{H} \text{H}_3]} dr \right. \\ \left. - 2\text{H} \int \frac{e^{-\frac{3c_1}{2r^2}} r^2 \text{H}_1}{(2c_1 \text{H}_2 - 3r^2 \text{H}) \text{H}_1 - 2c_1 \text{H} \text{H}_3} dr \right\}, \\ U(r) = e^{\frac{3c_1}{2r^2}}, \quad W(r) = N(r)B(r), \quad F_1 = 1 + \frac{c_1}{r^2}, \quad (31)$$

where $\text{H} = \text{HeunC} \left(\frac{3}{2}, \frac{3}{2}, 0, \frac{3}{8}, \frac{9}{8}, -\frac{c_1}{r^2} \right)$, $\text{H}_1 = \text{HeunC} \left(\frac{3}{2}, -\frac{3}{2}, 0, \frac{3}{8}, \frac{9}{8}, -\frac{c_1}{r^2} \right)$, $\text{H}_2 = \text{HeunCPrime} \left(\frac{3}{2}, \frac{3}{2}, 0, \frac{3}{8}, \frac{9}{8}, -\frac{c_1}{r^2} \right)$, and $\text{H}_3 = \text{HeunCPrime} \left(\frac{3}{2}, -\frac{3}{2}, 0, \frac{3}{8}, \frac{9}{8}, -\frac{c_1}{r^2} \right)^2$. Therefore, as Eqs. (30) and (31) show the case of $n = 2$ is defined by the function HeunC while the case of $n \neq 2$ is defined by (30). We will discuss this case in detail below.

Using Eq. (30) in (13) we derive the form of $f(r)$ as follows,

$$f(r) = \frac{12}{r^{n+2} X_1^{\frac{3n}{n-2}} X_2} \left\{ r^{\frac{2(n^2+2n-2)}{n-2}} \left[f_1^2(n^2-1) - \frac{(n^2-4)r^n}{2} f_1 - r^{2n} \right] X_2 \left\{ \int \frac{X_1^{\frac{n+1}{n-2}} X_2 \int \frac{X_1^{\frac{n+1}{n-2}}}{r^{\frac{7n-8}{n-2}} X_2} dr}{r^{\frac{n(2n-1)}{n-2}}} dr \right. \right. \\ \left. \left. - \int \frac{X_1^{\frac{n+1}{n-2}}}{r^{\frac{7n-8}{n-2}} X_2} dr \left[\int \frac{X_1^{\frac{n+1}{n-2}} X_2}{r^{\frac{n(2n-1)}{n-2}}} dr - \frac{c_1}{2} \right] \right\} \right. \\ \left. - r^{\frac{2(n^2-n+1)}{n-2}} ((F_0 - n + 3)(n+1) - r^n) X_1^{\frac{(2n-1)}{n-2}} \int \frac{X_1^{\frac{n+1}{n-2}} X_2}{3r^{\frac{n(2n-1)}{n-2}}} dr - \frac{1}{6} X_1^{\frac{3n}{n-2}} X_2^2 \right. \\ \left. + \frac{1}{2} c_2 r^{\frac{2(n^2+2n-2)}{n-2}} X_2 \left[\left\{ f_1(n^2-1) - \frac{(n^2-4)r^n}{2} \right\} f_1 - r^{2n} \right] \right. \\ \left. + \frac{1}{6} c_1 r^{\frac{2(n^2-n+1)}{n-2}} X_1^{\frac{2n-1}{n-2}} (f_1(n+1) - r^n) \right\}, \quad (32)$$

where $f_1 = F_0 - n + 3$. Using Eq. (32) in Eq. (9) we obtain the Ricci scalar in the form

$$R = - \frac{12}{r^2 X_1^{\frac{3n}{n-2}} X_2^2} \left\{ r^{\frac{2(n^2+2n-2)}{n-2}} \left\{ \frac{f_1(n-1)(n+4)}{2} + 2r^n \right\} X_2^2 \left\{ \int \frac{X_2 X_1^{\frac{n+1}{n-2}} \int \frac{X_1^{\frac{n+1}{n-2}}}{r^{\frac{7n-8}{n-2}} X_2} dr}{r^{\frac{n(2n-1)}{n-2}}} dr \right. \right. \\ \left. \left. - \left[\int \frac{X_1^{\frac{n+1}{n-2}} X_2}{r^{\frac{n(2n-1)}{n-2}}} dr - \frac{c_1}{2} \right] \int \frac{X_1^{\frac{n+1}{n-2}}}{r^{\frac{7n-8}{n-2}} X_2} dr \right\} - r^{\frac{2(n^2-n+1)}{n-2}} \left(\frac{f_1(n+4)}{2} - 2r^n \right) X_1^{\frac{(2n-1)}{n-2}} \int \frac{X_1^{\frac{n+1}{n-2}} X_2}{3r^{\frac{n(2n-1)}{n-2}}} dr \right. \\ \left. - X_2^2 \left\{ \frac{1}{3} X_1^{\frac{3n}{n-2}} - \frac{1}{2} c_2 r^{\frac{2(n^2+2n-2)}{n-2}} \left[\frac{f_1(n-1)(n+4)}{2} + 2r^n \right] \right\} \right. \\ \left. + \frac{1}{6} c_1 r^{\frac{2(n^2-n+1)}{n-2}} X_1^{\frac{2n-1}{n-2}} \left(\frac{f_1(n+4)}{2} - 2r^n \right) \right\}. \quad (33)$$

² The special function HeunC is defined as the solution of the Heun Confluent equation, that has the form

$$X''(r) - \frac{1 + \beta - (\alpha - \beta - \gamma - 2)r - r^2 \alpha}{r(r-1)} X'(r) - \frac{\alpha(1 + \beta) - \gamma - 2\eta - (1 + \gamma)\beta - r(2\delta + [2 + \gamma + \beta])}{2r(r-1)} X(r) = 0.$$

The above differential equation has the following solution: $\text{HeunC}(\alpha, \beta, \gamma, \delta, \eta, r)$. Interested readers can check [66, 67] for more details. The special function HeunCPrime is defined as the derivative of the Heun Confluent function.

B. The case of $\eta \neq 0$

Now we are going to find an analytic solution of the system (27), (28), and (29) in the case of $\eta \neq 0$ and obtain

$$\begin{aligned}
V(r) &= \frac{r^{\frac{2(n^2+2n-2)}{n-2}}}{X_1^{\frac{2(n+1)}{n-2}}} \left[c_2 + \int \frac{X_1^{\frac{n+1}{n-2}}}{r^{\frac{7n-8}{n-2}} X_2} dr \left(c_1 - 2 \int \frac{X_1^{\frac{n+1}{n-2}} X_3}{c_3 r^{\frac{n(2n-1)}{n-2}}} dr \right) + 2 \int \frac{X_3 X_1^{\frac{n+1}{n-2}} \int \frac{X_1^{\frac{n+1}{n-2}}}{r^{\frac{7n-8}{n-2}} X_2} dr}{c_3 r^{\frac{n(2n-1)}{n-2}}} dr \right], \\
W(r) &= c_3 r^{\frac{2n(n+1)}{2-n}} X_1^{\frac{2(n+1)}{n-2}}, \quad U(r) = W(r)V(r), \\
F_1 &= 1 - \frac{F_0 - (n-3)}{r^n}, \quad \eta(r) = c_4 + c_5 \int \frac{X_1^{\frac{n+1}{n-2}}}{r^{\frac{(n-1)(n+4)}{n-2}}} dr, \tag{34}
\end{aligned}$$

where $X_3 = c_3 X_2 - 2c_2^2 r^{n-2}$. Equation (34) reduces to (30) when $\eta = 0$. As Eq. (34) shows the case $n = 2$ is not allowed. However this case is studied in [55] where we obtained

$$\begin{aligned}
V(r) &= \frac{e^{\frac{3a_1}{2r^2}}}{r} \left\{ H a_2 + H_1 r^3 a_3 + 2H_1 r^3 \int \frac{e^{-\frac{3a_1}{2r^2}} H [r^2 + 13a_1]}{r(r^2 + a_1) [(2a_1 H_2 - 3r^2 H) H_1 - 2a_1 H H_3]} dr \right. \\
&\quad \left. - 2H \int \frac{e^{-\frac{3a_1}{2r^2}} r^2 H_1 [r^2 + 13a_1]}{(r^2 + a_1) [(2a_1 H_2 - 3r^2 H) H_1 - 2a_1 H H_3]} dr \right\}, \\
W(r) &= e^{-\frac{3a_1}{r^2}}, \quad U(r) = V(r)W(r), \quad \eta = a_0 + \sqrt{\pi} \operatorname{erf} \left[\frac{\sqrt{6a_1}}{2r} \right]. \tag{35}
\end{aligned}$$

Using Eq. (34) in the trace equation given by Eq. (13), we obtain the following form of $f(r)$ as

$$\begin{aligned}
f(r) &= \frac{12}{c_3 r^{n+2} X_1^{\frac{3n}{n-2}} X_2} \left\{ r^{\frac{2(n^2+2n-2)}{n-2}} \left[f_1^2 (n^2 - 1) - \frac{(n^2 - 4)r^n}{2} f_1 - r^{2n} \right] X_2 \left\{ \int \frac{X_1^{\frac{n+1}{n-2}} X_3 \int \frac{X_1^{\frac{n+1}{n-2}}}{r^{\frac{7n-8}{n-2}} X_2} dr}{r^{\frac{n(2n-1)}{n-2}}} dr \right. \right. \\
&\quad \left. \left. - \int \frac{X_1^{\frac{n+1}{n-2}}}{r^{\frac{7n-8}{n-2}} X_2} dr \left[\int \frac{X_1^{\frac{n+1}{n-2}} X_3}{r^{\frac{n(2n-1)}{n-2}}} dr - \frac{c_3 c_1}{2} \right] \right\} \right. \\
&\quad \left. - r^{\frac{2(n^2-n+1)}{n-2}} \times ((F_0 - n + 3)(n + 1) - r^n) X_1^{\frac{(2n-1)}{n-2}} \int \frac{X_1^{\frac{n+1}{n-2}} X_3}{3r^{\frac{n(2n-1)}{n-2}}} dr \right. \\
&\quad \left. - \frac{1}{6} X_1^{\frac{3n}{n-2}} X_2 X_4 + \frac{1}{2} c_2 r^{\frac{2(n^2+2n-2)}{n-2}} X_2 \left[\left\{ f_1 (n^2 - 1) - \frac{(n^2 - 4)r^n}{2} \right\} f_1 - r^{2n} \right] \right. \\
&\quad \left. + \frac{1}{6} c_1 c_3 r^{\frac{2(n^2-n+1)}{n-2}} X_1^{\frac{2n-1}{n-2}} (f_1 (n + 1) - r^n) \right\}, \tag{36}
\end{aligned}$$

where $X_4 = c_3 X_2 - c_3^2 r^{n-2}$. Using Eq. (34) in Eq. (9), we obtain the Ricci scalar in the following form

$$\begin{aligned}
R = & -\frac{12}{c_3 r^2 X_1^{\frac{3n}{n-2}} X_2^2} \left\{ r^{\frac{2(n^2+2n-2)}{n-2}} \left\{ \frac{f_1(n-1)(n+4)}{2} + 2r^n \right\} X_2^2 \left\{ \int \frac{X_3 X_1^{\frac{n+1}{n-2}}}{r^{\frac{n(2n-1)}{n-2}}} \frac{X_1^{\frac{n+1}{n-2}}}{r^{\frac{7n-8}{n-2}} X_2} dr \right. \right. \\
& - \left. \left[\int \frac{X_1^{\frac{n+1}{n-2}} X_3}{r^{\frac{n(2n-1)}{n-2}}} dr - \frac{c_1}{2} \right] \int \frac{X_1^{\frac{n+1}{n-2}}}{r^{\frac{7n-8}{n-2}} X_2} dr \right\} - r^{\frac{2(n^2-n+1)}{n-2}} \left(\frac{f_1(n+4)}{2} - 2r^n \right) X_1^{\frac{(2n-1)}{n-2}} \int \frac{X_1^{\frac{n+1}{n-2}} X_3}{3r^{\frac{n(2n-1)}{n-2}}} dr \\
& - X_2 \left\{ \frac{1}{3} X_4 X_1^{\frac{3n}{n-2}} - \frac{1}{2} c_2 c_3 r^{\frac{2(n^2+2n-2)}{n-2}} X_2 \left[\frac{f_1(n-1)(n+4)}{2} + 2r^n \right] \right\} \\
& + \left. \frac{1}{6} c_1 c_3 r^{\frac{2(n^2-n+1)}{n-2}} X_1^{\frac{2n-1}{n-2}} \left(\frac{f_1(n+4)}{2} - 2r^n \right) \right\}. \tag{37}
\end{aligned}$$

In the following section, we will study the physics beyond the BH solutions with/without charge given by Eq. (30) and (34)

IV. INHERENT PHYSICS OF THE BH SOLUTIONS (30) AND (34)

An important detail to emphasize is that when $F_0 = n - 3$, we recover the GR BH, as Eqs. (30) and (34) show. Now, we extract the inherent physics of the BH solutions (30) and (34). We therefore concentrate on the case $n = 4$ and write the metric potentials of the neutral BH solution (30) as

$$\begin{aligned}
V(r) = & \frac{c_2 r^{22}}{(F_0 - 1 + r^4)^5} + \frac{r^{22}}{(F_0 - 1 + r^4)^5} \left\{ c_1 \int \frac{(F_0 - 1 + r^4)^{\frac{5}{2}}}{r^{10} (F_0 - 1 - r^4)^5} dr \right. \\
& + 2 \int \frac{(F_0 - 1 - r^4) \left[\int \frac{(F_0 - 1 + r^4)^{\frac{5}{2}}}{r^{10} (F_0 - 1 - r^4)} dr \right] (F_0 - 1 + r^4)^{\frac{5}{2}}}{r^{14}} dr \\
& \left. - 2 \int \frac{(F_0 - 1 + r^4)^{\frac{5}{2}} (F_0 - 1 - r^4)}{r^{14}} dr \int \frac{(F_0 - 1 + r^4)^{\frac{5}{2}}}{r^{10} (F_0 - 1 - r^4)} dr \right\}, \\
W(r) = & \frac{c_3 (F_0 - 1 + r^4)^5}{r^{20}}, \quad U(r) = W(r)V(r), \quad F_1 = 1 - \frac{F_0 - 1}{r^4}. \tag{38}
\end{aligned}$$

The asymptotic forms of the metric potentials given by Eq. (38) take the following forms

$$\begin{aligned}
V(r) = & r^2 c_2 + 1 + \frac{c_1}{3r} - (F_0 - 1) \left[\frac{5c_2}{r^2} + \frac{4}{r^4} + \frac{7c_1}{6r^5} - \frac{15c_1 (F_0 - 1)}{r^6} + \dots \right], \\
U(r) = & r^2 c_2 + 1 + \frac{c_1}{3r} + (F_0 - 1) \left[\frac{1}{r^4} + \frac{c_1}{2r^5} + \frac{26(F_0 - 1)}{35r^8} + \dots \right], \tag{39}
\end{aligned}$$

where we have use $c_3 = 1$. Using Eq. (39) in Eq. (8) we obtain the line element in the form

$$\begin{aligned}
ds^2 \approx & - \left\{ r^2 \Lambda_{\text{eff}} + 1 - \frac{2m}{r} + (F_0 - 1) \left[\frac{1}{r^4} - \frac{3m}{r^5} + \frac{26(F_0 - 1)}{35r^8} \right] \right\} dt^2 \\
& + \frac{dr^2}{r^2 \Lambda_{\text{eff}} + 1 - \frac{2m}{r} - (F_0 - 1) \left[\frac{5\Lambda_{\text{eff}}}{r^2} + \frac{4}{r^4} - \frac{7m}{r^5} + \frac{90m(F_0 - 1)}{r^6} \right]} + d\Omega^2, \tag{40}
\end{aligned}$$

where $d\Omega^2 = r^2(d\theta^2 + \sin^2 \theta d\phi^2)$ is the line-element for the two dimensional unit sphere and $m = -\frac{c_1}{6}$ and $c_2 = \Lambda_{\text{eff}}$.

Equation (40) shows that the line element expresses the asymptotic anti-de Sitter (A)dS space-time and is not identical with the Schwarzschild space-time due to the contribution of the extra terms of the higher-order curvature of the $f(R)$ gravity. Equation (40) ensures what we have stated in the introduction, that is, in the $f(R)$ gravity, one can derive a space-time that is different from the Schwarzschild one and when $F_0 = 1$, i.e., $F_1 = 1$, we recover the Schwarzschild (A)dS metric [68] as usual. In conclusion, at a higher-order curvature, we can obtain a neutral space-time that is unlike the Schwarzschild solution and coincides with the Schwarzschild (A)dS at a lower order of $f(R) = R + \text{constant}$.

Now we use Eq. (39) in Eq. (9) and obtain³

$$R(r) \approx -12\Lambda_{\text{eff}} - \frac{36(F_0 - 1)}{r^6} \Rightarrow r(R) = \frac{6^{1/3} \left[(R + 12\Lambda_{\text{eff}})^5 (1 - F_0) \right]^{1/6}}{R + 12\Lambda_{\text{eff}}}, \quad (41)$$

where we have omitted the other terms in Ricci scalar to be able to write the radial coordinate r as a function of Ricci scalar, i.e., $r(R)$. From Eq. (41), we can clearly see that when $F_0 = 1$, we obtain a constant value of the Ricci scalar because when $F_1 = 1$ and $f_R = \text{const}$. The asymptote of $f(r)$ given by Eq. (32) becomes

$$f(r) \approx -6\Lambda_{\text{eff}} - \frac{36(F_0 - 1)}{r^6} + \frac{96m(F_0 - 1)}{r^7} \dots \quad (42)$$

Using Eq. (41) in (42), we obtain

$$f(R) \approx 6\Lambda_{\text{eff}} + R - \frac{12m6^{2/3}R^{7/6}}{27(1 - F_0)^{1/6}} - \frac{168m\Lambda_{\text{eff}}6^{2/3}R^{1/6}}{27(1 - F_0)^{1/6}} - \frac{168m\Lambda_{\text{eff}}^2 6^{2/3}}{27(1 - F_0)^{1/6} R^{5/6}} \dots \quad (43)$$

Equation (43) shows that $f(R)$ has ultraviolet and infrared terms.

The expression of Eq. (43) might look strange because there is a divergence in the limit of $F_0 \rightarrow 1$, which should correspond to the standard Einstein-Hilbert action with a cosmological constant. This occurs because we have used an expansion assuming that r is large in Eq. (42), but after that, if we consider the limit of $F_0 \rightarrow 1$, the scalar curvature R becomes a constant, which means that R is independent of r and therefore we cannot solve r with respect to R . Then it is natural if the expression (43), which is obtained by combining (41) and (42), becomes singular in the limit that of $F_0 \rightarrow 1$. The singularity is rather an artificial one coming from the non-commutability of the two limits that $r \rightarrow \infty$ and $F_0 \rightarrow 1$ but the behavior makes the situations ambiguous and often uncontrollable. In order to avoid this difficulty, we also use the numerical calculations.

Using Eq. (39), we obtain the invariants of solution (30) as

$$\begin{aligned} R_{\mu\nu\rho\sigma}R^{\mu\nu\rho\sigma} &= R_{\mu\nu}R^{\mu\nu} = 24\Lambda_{\text{eff}}^2 + \frac{144\Lambda_{\text{eff}}(F_0 - 1) + 48m^2}{r^6} - \frac{384\Lambda_{\text{eff}}m(F_0 - 1) + 48m^2}{r^7} - \frac{120\Lambda_{\text{eff}}^2(F_0 - 1)^2}{r^8} + \dots \\ R &= -12\Lambda_{\text{eff}} - \frac{36(F_0 - 1)}{r^6} + \frac{32m(F_0 - 1)}{r^7} + \dots, \end{aligned} \quad (44)$$

where $(R_{\mu\nu\rho\sigma}R^{\mu\nu\rho\sigma}, R_{\mu\nu}R^{\mu\nu}, R)$ are the Kretschmann scalar, the Ricci tensor square and the Ricci scalar, respectively, and all of them have a true singularity when $r = 0$. An important detail to highlight is the fact that F_0 is the main reason of the differentiation of the present study from the (A)dS Schwarzschild BH solution of GR whose invariants behave as $(R_{\mu\nu\rho\sigma}R^{\mu\nu\rho\sigma}, R_{\mu\nu}R^{\mu\nu}, R) = \left(24\Lambda^2 + \frac{48M^2}{r^6}, 36\Lambda^2, 12\Lambda\right)$. Equation (44) shows that the leading order of the scalars $(R_{\mu\nu\rho\sigma}R^{\mu\nu\rho\sigma}, R_{\mu\nu}R^{\mu\nu}, R)$ is $\left(\frac{1}{r^6}, \frac{1}{r^6}, \frac{1}{r^6}\right)$ which coincides with the form of the (A)dS Schwarzschild BH solution whose leading terms of the Kretschmann is $\left(\frac{1}{r^6}\right)$. Thus, Eq. (44) shows that the singularity of the Kretschmann coincides with the (A)dS Schwarzschild BH solution of GR.

Now we apply the same procedure that was used for the neutral case to the charged one. The analytic solution

³ The asymptote of Ricci scalar is given as

$$R(r) = -12\Lambda_{\text{eff}} - \frac{12(F_0 - 1)}{r^6} \left[3 + \frac{8m}{r} - \frac{15(F_0 - 1)\Lambda_{\text{eff}}}{r^2} + O\left(\frac{1}{r^4}\right) \dots \right].$$

(34) in the case $n = 4$ takes the form

$$\begin{aligned}
V(r) &= \frac{r^{22}}{c_3 (F_0 - 1 + r^4)^5} \left\{ c_2 c_3 + c_1 c_3 \int \frac{(F_0 - 1 + r^4)^{\frac{5}{2}}}{r^{10} (F_0 - 1 - r^4)^5} dr - 2 \int \frac{(F_0 - 1 + r^4)^{\frac{5}{2}} [c_3 (F - 1 - r^4) - 2c_5^2 r^2]}{r^{14}} dr \right. \\
&\quad \times \int \frac{(F_0 - 1 + r^4)^{\frac{5}{2}}}{r^{10} (F_0 - 1 - r^4)} dr + 2 \int \frac{[c_3 (F - 1 - r^4) - 2c_5^2 r^2] (F_0 - 1 + r^4)^{\frac{5}{2}} \left[\int \frac{(F_0 - 1 + r^4)^{\frac{5}{2}}}{r^{10} (F_0 - 1 - r^4)} dr \right]}{r^{14}} dr \left. \right\}, \\
W(r) &= \frac{c_3 (F_0 - 1 + r^4)^5}{r^{20}}, \quad U(r) = W(r)V(r), \quad F_1 = 1 - \frac{F_0 - 1}{r^4}, \quad \eta(r) = c_4 + c_5 \int \frac{(F_0 - 1 + r^4)^{\frac{5}{2}}}{r^{12}} dr. \quad (45)
\end{aligned}$$

Equation (45) reduces to Eq. (38) when the constant $c_5 = 0$ which makes the electric charge $\eta(r) = \text{const}$. To further examine this charged solution, we calculate its asymptotic form and obtain

$$\begin{aligned}
V(r) &\approx r^2 c_2 + 1 + \frac{c_1}{3r} + \frac{c_5^2}{r^2} - (F_0 - 1) \left[\frac{5c_2}{r^2} + \frac{4}{r^4} + \frac{7c_1}{6r^5} + \frac{3[c_5^2 - 5c_1(F_0 - 1)]}{r^6} + \dots \right], \\
U(r) &\approx r^2 c_2 + 1 + \frac{c_1}{3r} + \frac{c_5^2}{r^2} + (F_0 - 1) \left[\frac{1}{r^4} + \frac{c_1}{2r^5} + \frac{2c_5^2}{r^6} + \frac{26(F_0 - 1)}{35r^8} + \dots \right], \quad (46)
\end{aligned}$$

where we use $c_3 = 1$. Using Eq. (46) in Eq. (8), we obtain the line element in the following form

$$\begin{aligned}
ds^2 &\approx - \left\{ r^2 \Lambda_{\text{eff}} + 1 - \frac{2m}{r} + \frac{q^2}{r^2} + (F_0 - 1) \left[\frac{1}{r^4} - \frac{3m}{r^5} + \frac{2q^2}{r^6} + \frac{26(F_0 - 1)}{35r^8} \right] \right\} dt^2 \\
&\quad + \frac{dr^2}{r^2 \Lambda_{\text{eff}} + 1 - \frac{2m}{r} + \frac{q^2}{r^2} - (F_0 - 1) \left[\frac{5\Lambda_{\text{eff}}}{r^2} + \frac{4}{r^4} - \frac{7m}{r^5} + \frac{3[q^2 + 30m(F_0 - 1)]}{r^6} \right]} + d\Omega^2, \quad (47)
\end{aligned}$$

where $q = c_5$, Λ_{eff} and m have the same values given in the neutral case.

Equation (47) shows that the line element is asymptotic to (A)dS and is not equivalent to the Reissner-Nordström space-time in respect of the contribution of the extra terms of the higher-order curvature of the $f(R)$ gravity. Equation (47) shows clearly that in the $f(R)$ gravity, one can obtain a space-time that is different from the Reissner-Nordström space-time and when the constant $F_0 = 1$ we recover the Reissner-Nordström (A)dS metric [68]. We can summarize the results of this section by saying that at a higher order curvature we can obtain a charged space-time that is unlike Reissner-Nordström space-time and reduces to the Reissner-Nordström one (A)dS at a lower order of $f(R) = R$.

Using Eq. (46) we obtain the invariants of solution (34) as:

$$\begin{aligned}
R_{\mu\nu\rho\sigma} R^{\mu\nu\rho\sigma} &= R_{\mu\nu} R^{\mu\nu} = 24\Lambda_{\text{eff}}^2 + \frac{144\Lambda_{\text{eff}}(F_0 - 1) + 48m^2}{r^6} - \frac{384\Lambda_{\text{eff}}m(F_0 - 1) + 96mq^2}{r^7} \\
&\quad - \frac{120\Lambda_{\text{eff}}^2(F_0 - 1)^2 + 56q^2[q^2 + 30m(F_0 - 1)]}{r^8} + \dots, \\
R &= -12\Lambda_{\text{eff}} - \frac{36(F_0 - 1)}{r^6} + \frac{32m(F_0 - 1)}{r^7} + (F_0 - 1) \frac{180\Lambda_{\text{eff}}(F_0 - 1) + 60q^2}{r^8} + \dots. \quad (48)
\end{aligned}$$

Equation (48) shows that all the invariants suffer a true singularity when $r = 0$. A substantial detail to highlight is that F_0 is the origin that make the above results unlike the Reissner-Nordström (A)dS BH solution of GR whose scalars behave as $(R_{\mu\nu\rho\sigma} R^{\mu\nu\rho\sigma}, R_{\mu\nu} R^{\mu\nu}, R) = (24\Lambda^2 + \frac{48M^2}{r^6}, 36\Lambda^2 + \frac{4q^4}{r^8}, 12\Lambda)$. Equation (48) shows that the leading expression of the scalars $(R_{\mu\nu\rho\sigma} R^{\mu\nu\rho\sigma}, R_{\mu\nu} R^{\mu\nu}, R)$ is $(\frac{1}{r^6}, \frac{1}{r^6}, \frac{1}{r^6})$ which does not coincide with the form of the Reissner-Nordström (A)dS BH solution whose leading expressions of the Kretschmann and the Ricci tensor squared are $(\frac{1}{r^6}, \frac{1}{r^8})$. Thus, Eq. (48) shows that the singularity of the Ricci tensor squared is much milder than that of the Reissner-Nordström (A)dS BH solution of GR.

V. STABILITY OF THE BHS USING GEODESIC DEVIATION

The geodesic equations take the form [68],

$$\frac{d^2 x^\alpha}{d\varepsilon^2} + \left\{ \begin{array}{c} \alpha \\ \beta\rho \end{array} \right\} \frac{dx^\beta}{d\varepsilon} \frac{dx^\rho}{d\varepsilon} = 0, \quad (49)$$

where ε is the affine connection parameter. The geodesic deviation equations take the form [69, 70],

$$\frac{d^2 \epsilon^\sigma}{d\varepsilon^2} + 2 \left\{ \begin{matrix} \sigma \\ \mu\nu \end{matrix} \right\} \frac{dx^\mu}{d\varepsilon} \frac{d\epsilon^\nu}{d\varepsilon} + \left\{ \begin{matrix} \sigma \\ \mu\nu \end{matrix} \right\}_{, \rho} \frac{dx^\mu}{d\varepsilon} \frac{dx^\nu}{d\varepsilon} \epsilon^\rho = 0, \quad (50)$$

with ϵ^ρ being the four-vector deviation. Plugging Eqs. (49) and (50) into Eq. (8), we obtain

$$\frac{d^2 t}{d\varepsilon^2} = 0, \quad \frac{1}{2} U'(r) \left(\frac{dt}{d\varepsilon} \right)^2 - r \left(\frac{d\phi}{d\varepsilon} \right)^2 = 0, \quad \frac{d^2 \theta}{d\varepsilon^2} = 0, \quad \frac{d^2 \phi}{d\varepsilon^2} = 0, \quad (51)$$

and for the geodesic deviation the line-element (8) gives

$$\begin{aligned} \frac{d^2 \epsilon^1}{d\varepsilon^2} + V(r) U'(r) \frac{dt}{d\varepsilon} \frac{d\epsilon^0}{d\varepsilon} - 2rV(r) \frac{d\phi}{d\varepsilon} \frac{d\epsilon^3}{d\varepsilon} + \left[\frac{1}{2} (U'(r)V'(r) + V(r)U''(r)) \left(\frac{dt}{d\varepsilon} \right)^2 - (V(r) + rV'(r)) \left(\frac{d\phi}{d\varepsilon} \right)^2 \right] \epsilon^1 = 0, \\ \frac{d^2 \epsilon^0}{d\varepsilon^2} + \frac{V'(r)}{V(r)} \frac{dt}{d\varepsilon} \frac{d\epsilon^1}{d\varepsilon} = 0, \quad \frac{d^2 \epsilon^2}{d\varepsilon^2} + \left(\frac{d\phi}{d\varepsilon} \right)^2 \epsilon^2 = 0, \quad \frac{d^2 \epsilon^3}{d\varepsilon^2} + \frac{2}{r} \frac{d\phi}{d\varepsilon} \frac{d\epsilon^1}{d\varepsilon} = 0, \end{aligned} \quad (52)$$

where $U(r)$ and $V(r)$ are given in Eq. (39) or Eq. (46) and $'$ is the derivative with respect to the radial coordinate r . From the condition of a circular orbit, we can obtain

$$\theta = \frac{\pi}{2}, \quad \frac{d\theta}{d\varepsilon} = 0, \quad \frac{dr}{d\varepsilon} = 0. \quad (53)$$

The use of Eq. (53) in Eq. (51) leads to

$$\left(\frac{d\phi}{d\varepsilon} \right)^2 = \frac{U'(r)}{r[2U(r) - rU'(r)]}, \quad \left(\frac{dt}{d\varepsilon} \right)^2 = \frac{2}{2U(r) - rU'(r)}. \quad (54)$$

We can rewrite Eq. (52) as

$$\begin{aligned} \frac{d^2 \epsilon^1}{d\phi^2} + U(r) U'(r) \frac{dt}{d\phi} \frac{d\epsilon^0}{d\phi} - 2rV(r) \frac{d\epsilon^3}{d\phi} + \left[\frac{1}{2} [U'(r)^2 + U(r)U''(r)] \left(\frac{dt}{d\phi} \right)^2 - [U(r) + rU'(r)] \right] \zeta^1 = 0, \\ \frac{d^2 \epsilon^2}{d\phi^2} + \epsilon^2 = 0, \quad \frac{d^2 \epsilon^0}{d\phi^2} + \frac{U'(r)}{U(r)} \frac{dt}{d\phi} \frac{d\epsilon^1}{d\phi} = 0, \quad \frac{d^2 \epsilon^3}{d\phi^2} + \frac{2}{r} \frac{d\epsilon^1}{d\phi} = 0. \end{aligned} \quad (55)$$

From the second of Eq. (55), we can show that we have a simple harmonic motion, which is the stability condition of the plane $\theta = \pi/2$. The remaining equations of (55) assume the following solutions

$$\epsilon^0 = \zeta_1 e^{i\sigma\phi}, \quad \epsilon^1 = \zeta_2 e^{i\sigma\phi}, \quad \text{and} \quad \epsilon^3 = \zeta_3 e^{i\sigma\phi}, \quad (56)$$

where ζ_1, ζ_2 , and ζ_3 are constants and ω is an unknown function. Using the values of ϵ^1 , and ϵ^3 given by Eq. (56) in the fourth equation of Eq. (55), we get

$$\zeta_2 = -\frac{\zeta_3 \sigma r}{2}. \quad (57)$$

Then substituting the values of ϵ^0 , and ϵ^1 given by Eq. (56) into the third equation of Eq. (55) and using Eq. (57), we obtain

$$\zeta_1 = \frac{\zeta_3 \sqrt{2U' r^3}}{2U}. \quad (58)$$

Substituting (56), after using Eqs. (57) and (58), into the first equation of Eq. (55) we obtain the stability condition as

$$\frac{3UVU' - \sigma^2 UU' - 2rVU'^2 + rUVU''}{UU'} > 0 \Rightarrow \sigma^2 < \frac{3UVU' - 2rVU'^2 + rUVU''}{UU'}. \quad (59)$$

We depict Eq. (59) in Fig. 1 using specific values of the model. The case $q = 0$ is drawn in Fig. 1 (a), and the case $q \neq 0$ is drawn in Fig. 1 (b) for the inequality of Eq. (59). These figures exhibit the unshaded and shaded zones where the BHs are stable and not stable respectively.




Figure 1. Plot of Eq. (59) against the coordinate r for the BHs (39) and (46).

VI. THERMODYNAMICS OF THE BH SOLUTIONS (30) AND (34)

In this section, we will investigate the thermodynamical properties of the BH solutions (30) and (34)⁴. The temperature of a BH is defined as [71–77]

$$T_{(1,2)} = \frac{r_{(1,2)} - r_{(2,1)}}{4\pi r_{(1,2)}^2}, \quad (60)$$

where $r_{(1,2)}$ represents the inner and outer horizons of the space-time. The Hawking entropy of the horizons is defined as

$$\psi_{(1,2)} = \frac{1}{4} A_{(1,2)} f_R, \quad (61)$$

with $A_{(1,2)}$ being the area of the horizons. The quasi-local energy is defined as [71–74, 78, 79]

$$E_{(1,2)} = \frac{1}{4} \int \left[2f_R r_{(1,2)} + r_{(1,2)}^2 \{ F_0 (R(r_{(1,2)})) - R(r_{(1,2)}) f_R r_{(1,2)} \} \right] dr_{(1,2)}. \quad (62)$$

Finally, the Gibbs free energy is defined as [79, 80]

$$G_{(1,2)} = E_{(1,2)} - T_{(1,2)} \psi_{(1,2)}. \quad (63)$$

A. Thermodynamics of the BH (46) that has asymptotic flatness

In this subsection, we study the thermodynamics of the BH (46) which is characterized by the mass m , F , and the constant q . When $F_0 = 1$, we obtain (A)dS Reissner-Nordström BH of GR. To derive the horizons of the BH, (46),

⁴ In this study, we will not address the BH solutions (30) and (34) because one cannot easily find their explicit roots. Therefore, we will use $\Lambda_{\text{eff}} = 0$ and study these solutions up to $O\left(\frac{1}{r^4}\right)$.

we set $U(r) = 0$ [77], whose solutions give two real positive roots that have the form

$$\begin{aligned}
r_1 &= \frac{1}{6} \left(3m + 3\sqrt{\frac{q^4 + 12(F_0 - 1)}{F_1} + m^2 + \frac{2q^2}{3} + \frac{F_1}{9}} \right. \\
&\quad \left. + 3 \left\{ 2m^2 + \frac{4q^2}{3} - \frac{F_1}{9} - \frac{12(F_0 - 1)}{F_1} - \frac{q^4}{F} + \frac{2mq^2(1+m)}{\sqrt{\frac{q^4 + 12(F_0 - 1)}{F_1} + m^2 + \frac{2q^2}{3} + \frac{F_1}{9}}} \right\}^{1/2} \right), \\
r_2 &= \frac{1}{6} \left(3m + 3\sqrt{\frac{q^4 + 12(F_0 - 1)}{F_1} + m^2 + \frac{2q^2}{3} + \frac{F_1}{9}} \right. \\
&\quad \left. - 3 \left\{ 2m^2 + \frac{4q^2}{3} - \frac{F_1}{9} - \frac{12(F_0 - 1)}{F_1} - \frac{q^4}{F} + \frac{2mq^2(1+m)}{\sqrt{\frac{q^4 + 12(F_0 - 1)}{F_1} + m^2 + \frac{2q^2}{3} + \frac{F_1}{9}}} \right\}^{1/2} \right), \tag{64}
\end{aligned}$$

where

$$\begin{aligned}
F_1 &= 3 \left(6\sqrt{(1 - F_0)[18q^8 + 3m^2q^6 + 24(1 - F_0)q^4 + 108q^2m^2(1 - F_0)] + 48(1 - 3F_0 - 3F_0^2)} \right. \\
&\quad \left. - q^6 + 36q^2(F_0 - 1) + 54m^2(F_0 - 1) \right)^{1/3}.
\end{aligned}$$

The metric potentials of the BH (46) are plotted in Fig. 2 (a). Fig. 2 (a) shows that we have two positive horizons of the metric potential $U(r)$ when $F \neq 1$. The degenerate horizon of $U(r)$ occurred at $(F, m, q, r) \equiv (2, 0.77, 0.53, 1.22)$ which coincides with the Nariai BH. The behavior of the degenerate is indicated in Fig. 2 (b) which shows that $r_1 < r_d$. As we observe from Fig. 2 (b), as q decreases, we enter a parameter region where no horizon exists, and thus, the central singularity is a naked singularity. The behavior of the two horizons w.r.t. mass is drawn in Fig. 2 (c) and (d) when $F_0 = 1$ and $F \neq 1.1$. From Eq. (60) the Hawking temperature of the BH (46) is calculated and drawn in Figs. 2 (e) and (f) for $F_0 = 1$ and $F_0 = 1.1$, respectively. These figures show that $T_1 > T_2$. Figs. 2 (e) and (f) also indicate that T_1 has an increasing positive value whereas T_2 has a decreasing negative one.

From Eq. (61), the entropy of BH (46) takes the form

$$\psi_{(1,2)} = \pi r_{(1,2)}^2 \left(1 - \frac{F_0 - 1}{r_{(1,2)}^4} \right). \tag{65}$$

The plot of the entropy (65) when $F_0 = 1$ and $F_0 = 1.1$ is drawn in Figs. 2 (g) and 2 (h) which indicate an increasing value for ψ_1 and decreasing value for ψ_2 . From Eq. (62) we evaluate the local energy of BH (46) and obtain

$$E_{(1,2)} = \frac{21r_{(1,2)}^9 + (F_0 - 1)[7r_{(1,2)}^5 - 126q^2r_{(1,2)}^3 + 54r_{(1,2)}(F_0 - 1) - 126m(F_0 - 1)]}{42r_{(1,2)}^8}. \tag{66}$$

Equation (66) shows that when $F_0 = 1$, we obtain $E_{(1,2)} = \frac{r_{(1,2)}}{2}$ which is the energy of a spherically symmetric space-time. The plot of Eq. (66) when $F_0 = 1$ and $F_0 = 1.1$ is drawn in Figs. 2 (i) and 2 (j), which also indicates positive increasing values for $E_{(1,2)}$. Figs. 2 (i) and (j) also show $E_1 > E_2$. Finally, by using of Eqs. (60), (65), and (66) in (63), we evaluate Gibbs' free energies. The plot of these energies is drawn in Fig. 2 (k) and 2 (l) when $F_0 = 1$ and $F_0 = 1.1$. These figures indicate a positive increasing value for G_1 and also $G_1 > G_2$.

B. First law of thermodynamics of the BH solutions (39) and (46)

An important step for any BH solution is to check its validity of the first law of thermodynamics. Therefore, for the charged BH the Smarr formula and the differential form for the first law of thermodynamics, in the frame of the $f(R)$ gravity, can be expressed as [81, 82]

$$M = 2(T\psi - PV) + \eta Q, \quad dE = Td\psi + \eta dQ + PdV, \tag{67}$$




Figure 2. Plots of thermodynamical quantities of the BH solution Eq. (46): (a) shows the behavior of the metric potential U for $F_0 = 1$ and $F \neq 1$ when $m = 1$ and $q = 0.5$; (b) Typical behavior of the horizons, degenerate and non-horizon of the metric potential $U(r)$ given by Eq. (46) when $m = 0.77$ and $F = 2$; (c) Typical behavior of the horizons with mass given by Eq. (46) when $q = 0.05$ and $F_0 = 1$; (d) Typical behavior of the horizons with mass given by Eq. (46) when $q = 0.05$ and $F = 2$; (e) and (f) The behavior of the Hawking temperature, (60), showing T_1 has a positive decreasing value whereas T_2 has a negative increasing value when $F_0 = 1$ and $F_0 = 1.1$; (g) and (h) The behavior of the Hawking entropy, (65), which indicates $\psi_{(1,2)}$ always has positive values, ψ_1 and increases with M while ψ_2 has a positive decreasing value; (i) and (j) The behavior of quasi-local energy, (66), which indicate that $E_{(1,2)}$ has positive increasing values and also $E_1 > E_2$; (k) and (l) The behavior of Gibb's free energy which indicates the behavior of G_1 which is positive while G_2 starts with a positive value that becomes negative as M increases.

where ψ is the Hawking entropy, T is the Hawking temperature, η is the electric potential P is the radial component of the stress-energy tensor that is used as a thermodynamic pressure, i.e., $P = T_r{}^r|_{\pm}$, V is the geometric volume and $M = m$, $Q = q$. The pressure, in the context of the $f(R)$ gravity, is determined as [81]

$$P = -\frac{1}{8\pi} \left\{ \frac{F_1}{r_{(1,2)}^2} + \frac{1}{2} (f(R) - RF_1) \right\} + \frac{1}{4} \left(\frac{2F_1}{r_{(1,2)}} + F_1' \right) T. \quad (68)$$

Using Eq. (47) we get

$$M = \frac{1}{2} \left(r_{(1,2)} + \frac{q^2}{r_{(1,2)}} - \frac{(F_0 - 1)}{2r_{(1,2)}^3} \right). \quad (69)$$

By calculating the necessary components of Eq. (67) we obtain

$$\begin{aligned} P_{(1,2)} &\approx -\frac{q^2 r_{(1,2)}^2 - 3(F_0 - 1)}{6r_{(1,2)}^3}, & T_{(1,2)} &\approx \frac{r_{(1,2)}^4 - q^2 r_{(1,2)}^2 + 3(F_0 - 1)}{4\pi r_{(1,2)}^5}, \\ \psi_{(1,2)} &\approx \frac{\pi [r_{(1,2)}^4 - (F_0 - 1)]}{r_{(1,2)}^2}, & \eta_{(1,2)} &\approx \frac{q}{2r_{(1,2)}} - \frac{(F_0 - 1)q}{r_{(1,2)}^5}, & V &= \frac{4}{3}\pi r_{(1,2)}^3. \end{aligned} \quad (70)$$

Using Eq. (70) in Eq. (67), we can prove that the first law of the flat space-times (39) and (46) is verified.

VII. DISCUSSION AND CONCLUSIONS

This study focuses on deriving spherically symmetric BH solutions with/without charge in the context of the $f(R)$ modified gravity theory. The $f(R)$ gravity is known as a theory whose field equation includes fourth order derivative thereby making the derivation of the analytic solution difficult. Therefore, we use the trace equation of the $f(R)$ equation and solve it with respect to $f(R)$. Using this solution, we rewrite the charged field equations of the $f(R)$ gravity and apply them to a spherically symmetric space-time that has two unknown functions of radial coordinate. The resulting non-linear differential equations are solved under two cases: The case of vanishing the electric charge and the case of non-vanishing electric charge. In these two cases, we assume the derivative of the $F_1(r) = \frac{df(R(r))}{dR(r)} = 1 - \frac{F_0 - (n-3)}{r^n}$ where $n > 0$. We can clearly see from the previous mentioned assumption that when $F = n - 3$, we obtain $F_1(r) = 1$, which is the case of GR. In the frame of the above assumption of $F_1(r)$ we solved the field equations of the $f(R)$ gravity with/without the electric charge analytically and derived the exact form of the metric potentials and the electric charge.

To understand the physics of these BH solutions, we give the asymptotic form of the metric potentials, when $n = 4$, with/without charge. From such asymptote, we show explicitly that our BH solutions are different from GR BH solutions and coincide with them when $F_0 = 1$. Then we write the line-element of such BHs and show they asymptotically behave as (A)dS space-time. We calculate the asymptotic form of $f(R)$ and show that it contains ultraviolet and infrared terms. We calculate the invariants of these BHs with/without charge and show that the singularity of the Ricci squared tensor in the charged case is milder than that of the GR BHs. Moreover, we study the geodesic deviations of these BHs and derive the condition of stability analytically and graphically, as shown in Fig. 1.

To further examine the physics of these BHs we calculate the thermodynamical quantities such as Hawking temperature, entropy, quasi-local energy, and Gibb's free energy, and investigate their behavior analytic and graphically. Moreover, we test the first law of thermodynamics and show in detail that such BHs satisfy this law. If we apply the results of odd perturbation presented in [65] we can prove that the BHs presented in this study are stable.

In summary, we derived new BHs with/without charge in the frame of $f(R)$ and showed that their Ricci scalars are not constant. These BHs are original ones, and their originality comes from the constant that involves, F_0 . This constant comes from the assumption of the first derivative of $f(R)$, i.e., $F_1(r) = \frac{df(R(r))}{dR(r)} = 1 - \frac{F_0 - (n-3)}{r^n}$. We stress on the fact that if $F_1(r) = 1 + \frac{c}{r^n}$, $n > 2$, we get BH solutions that cannot reduce to GR BHs because c is not allow to have a zero value in these BHs. This case needs more study and will be done elsewhere.

ACKNOWLEDGMENTS

This work is supported by the JSPS Grant-in-Aid for Scientific Research (C) No. 18K03615 (S.N.).

-
- [1] S. Nojiri and S. D. Odintsov, *Phys. Rept.* **505**, 59 (2011), arXiv:1011.0544 [gr-qc].
- [2] S. Nojiri and S. D. Odintsov, *eConf C0602061*, 06 (2006), arXiv:hep-th/0601213.
- [3] E. J. Copeland, M. Sami, and S. Tsujikawa, *Int. J. Mod. Phys. D* **15**, 1753 (2006), arXiv:hep-th/0603057.
- [4] A. Aragón, P. González, E. Papantonopoulos, and Y. Vásquez, (2020), arXiv:2005.11179 [gr-qc].
- [5] T. Clifton, P. G. Ferreira, A. Padilla, and C. Skordis, *Phys. Rept.* **513**, 1 (2012), arXiv:1106.2476 [astro-ph.CO].
- [6] S. Capozziello and M. Francaviglia, *Gen. Rel. Grav.* **40**, 357 (2008), arXiv:0706.1146 [astro-ph].
- [7] M. Mohsenzadeh and E. Yusofi, *J. Theor. Appl. Phys.* **6**, 10 (2012), arXiv:1207.1803 [physics.gen-ph].
- [8] G. Cognola, E. Elizalde, S. Nojiri, S. D. Odintsov, L. Sebastiani, and S. Zerbini, *Phys. Rev.* , 046009 (2008), arXiv:0712.4017 [hep-th].
- [9] A. Awad, W. El Hanafy, G. G. L. Nashed, S. D. Odintsov, and V. K. Oikonomou, *JCAP* **1807**, 026 (2018), arXiv:1710.00682 [gr-qc].
- [10] G. Cognola, E. Elizalde, S. Nojiri, S. D. Odintsov, L. Sebastiani, and S. Zerbini, *Phys. Rev.* **D77**, 046009 (2008), arXiv:0712.4017 [hep-th].
- [11] P. Zhang, *Phys. Rev.* **D73**, 123504 (2006), arXiv:astro-ph/0511218 [astro-ph].
- [12] B. Li and J. D. Barrow, *Phys. Rev.* **D75**, 084010 (2007), arXiv:gr-qc/0701111 [gr-qc].
- [13] Y.-S. Song, H. Peiris, and W. Hu, *Phys. Rev.* **D76**, 063517 (2007), arXiv:0706.2399 [astro-ph].
- [14] S. Nojiri and S. D. Odintsov, *Phys. Rev.* **D77**, 026007 (2008), arXiv:0710.1738 [hep-th].
- [15] A. M. Awad, S. Capozziello, and G. G. L. Nashed, *Journal of High Energy Physics* **7**, 136 (2017), arXiv:1706.01773 [gr-qc].
- [16] S. Nojiri and S. D. Odintsov, *Phys. Lett.* **B657**, 238 (2007), arXiv:0707.1941 [hep-th].
- [17] S. Capozziello, C. A. Mantica, and L. G. Molinari, *Int. J. Geom. Meth. Mod. Phys.* **16**, 1950008 (2018), arXiv:1810.03204 [gr-qc].
- [18] G. G. L. Nashed, *Mod. Phys. Lett.* **A22**, 1047 (2007), arXiv:gr-qc/0609096 [gr-qc].
- [19] J. Vainio and I. Vilja, *Gen. Rel. Grav.* **49**, 99 (2017), arXiv:1603.09551 [astro-ph.CO].
- [20] M. Ostrogradsky, *Mem. Acad. St. Petersburg* **6**, 385 (1850).
- [21] R. P. Woodard, *The invisible universe: Dark matter and dark energy. Proceedings, 3rd Aegean School, Karfas, Greece, September 26-October 1, 2005*, *Lect. Notes Phys.* **720**, 403 (2007), arXiv:astro-ph/0601672 [astro-ph].
- [22] H. A. Buchdahl, *Mon. Not. Roy. Astron. Soc.* **150**, 1 (1970).
- [23] A. A. Starobinsky, *Phys. Lett.* **B91**, 99 (1980), [,771(1980)].
- [24] S. M. Carroll, M. Hoffman, and M. Trodden, *Phys. Rev. D* **68**, 023509 (2003), arXiv:astro-ph/0301273.
- [25] S. M. Carroll, V. Duvvuri, M. Trodden, and M. S. Turner, *Phys. Rev.* **D70**, 043528 (2004), arXiv:astro-ph/0306438 [astro-ph].
- [26] S. Capozziello, *Int. J. Mod. Phys. D* **11**, 483 (2002), arXiv:gr-qc/0201033 [gr-qc].
- [27] S. Capozziello, V. F. Cardone, S. Carloni, and A. Troisi, *Int. J. Mod. Phys. D* **12**, 1969 (2003), arXiv:astro-ph/0307018 [astro-ph].
- [28] M. Soussa and R. Woodard, *Gen. Rel. Grav.* **36**, 855 (2004), arXiv:astro-ph/0308114.
- [29] V. Faraoni, *Phys. Rev. D* **72**, 124005 (2005), arXiv:gr-qc/0511094.
- [30] T. Chiba, *Phys. Lett.* **B575**, 1 (2003), arXiv:astro-ph/0307338 [astro-ph].
- [31] C. Corda, *Astropart. Phys.* **30**, 209 (2008), arXiv:0812.0483 [gr-qc].
- [32] C. Corda, *Eur. Phys. J. C* **65**, 257 (2010), arXiv:1007.4077 [gr-qc].
- [33] K. Bamba and S. D. Odintsov, *JCAP* **04**, 024 (2008), arXiv:0801.0954 [astro-ph].
- [34] S. Nojiri and S. D. Odintsov, *Phys. Rev. D* **74**, 086005 (2006), arXiv:hep-th/0608008.
- [35] S. Nojiri and S. D. Odintsov, *Einstein's legacy: From the theoretical paradise to astrophysical observations*, *J. Phys. Conf. Ser.* **66**, 012005 (2007), arXiv:hep-th/0611071 [hep-th].
- [36] S. Nojiri and S. D. Odintsov, *Phys. Rev. D* **78**, 046006 (2008), arXiv:0804.3519 [hep-th].
- [37] T. Kobayashi and K.-i. Maeda, *Phys. Rev. D* **79**, 024009 (2009), arXiv:0810.5664 [astro-ph].
- [38] V. Faraoni, *Phys. Rev. D* **72**, 061501 (2005), arXiv:gr-qc/0509008.
- [39] S. Capozziello, S. Nojiri, S. D. Odintsov, and A. Troisi, *Phys. Lett.* , 135 (2006), arXiv:astro-ph/0604431 [astro-ph].
- [40] L. Amendola, R. Gannouji, D. Polarski, and S. Tsujikawa, *Phys. Rev. D* **75**, 083504 (2007), arXiv:gr-qc/0612180.
- [41] M. Akbar and R.-G. Cai, *Phys. Lett. B* **648**, 243 (2007), arXiv:gr-qc/0612089.
- [42] T. Multamaki and I. Vilja, *Phys. Rev.* **D74**, 064022 (2006), arXiv:astro-ph/0606373 [astro-ph].
- [43] T. Multamaki and I. Vilja, *Phys. Rev.* **D76**, 064021 (2007), arXiv:astro-ph/0612775 [astro-ph].
- [44] A. de la Cruz-Dombriz and D. Saez-Gomez, *Entropy* **14**, 1717 (2012), arXiv:1207.2663 [gr-qc].
- [45] S. H. Hendi, B. Eslam Panah, and S. M. Mousavi, *Gen. Rel. Grav.* **44**, 835 (2012), arXiv:1102.0089 [hep-th].
- [46] L. Sebastiani and S. Zerbini, *Eur. Phys. J. C* **71**, 1591 (2011), arXiv:1012.5230 [gr-qc].
- [47] G. G. L. Nashed, *Mod. Phys. Lett.* , 2241 (2006), arXiv:gr-qc/0401041 [gr-qc].
- [48] S. H. Hendi and D. Momeni, *Eur. Phys. J. C* **71**, 1823 (2011), arXiv:1201.0061 [gr-qc].

- [49] S. Asgari and R. Saffari, *Gen. Rel. Grav.* **44**, 737 (2012), arXiv:1104.5108 [gr-qc].
- [50] G. G. L. Nashed, *Chin. Phys. B* **19**, 020401 (2010), arXiv:0910.5124 [gr-qc].
- [51] S. G. Ghosh and S. D. Maharaj, *Eur. Phys. J. C* **75**, 7 (2015), arXiv:1410.4043 [gr-qc].
- [52] S. H. Hendi, B. Eslam Panah, and R. Saffari, *Int. J. Mod. Phys. D* **23**, 1450088 (2014), arXiv:1408.5570 [hep-th].
- [53] Z.-Y. Tang, B. Wang, and E. Papantonopoulos, (2019), arXiv:1911.06988 [gr-qc].
- [54] G. Nashed and S. Nojiri, *Phys. Rev. D* **102**, 124022 (2020), arXiv:2012.05711 [gr-qc].
- [55] G. Nashed and S. Nojiri, (2020), arXiv:2010.04701 [hep-th].
- [56] H. A. Buchdahl, *mnras* **150**, 1 (1970).
- [57] S. Nojiri and S. D. Odintsov, *Phys. Rev.* , 123512 (2003), arXiv:hep-th/0307288 [hep-th].
- [58] S. Capozziello and M. De Laurentis, *Phys. Rept.* **509**, 167 (2011), arXiv:1108.6266 [gr-qc].
- [59] G. G. L. Nashed, *Nuovo Cim. B* **117**, 521 (2002), arXiv:gr-qc/0109017.
- [60] S. Nojiri, S. D. Odintsov, and V. K. Oikonomou, *Phys. Rept.* **692**, 1 (2017), arXiv:1705.11098 [gr-qc].
- [61] S. Capozziello, P. A. Gonzalez, E. N. Saridakis, and Y. Vasquez, *JHEP* **02**, 039 (2013), arXiv:1210.1098 [hep-th].
- [62] G. Cognola, E. Elizalde, S. Nojiri, S. D. Odintsov, and S. Zerbini, *jcap* **2**, 010 (2005), hep-th/0501096.
- [63] S. Kalita and B. Mukhopadhyay, *Eur. Phys. J.* **C79**, 877 (2019), arXiv:1910.06564 [gr-qc].
- [64] G. G. L. Nashed and S. Capozziello, *Phys. Rev.* **D99**, 104018 (2019), arXiv:1902.06783 [gr-qc].
- [65] E. Elizalde, G. G. L. Nashed, S. Nojiri, and S. D. Odintsov, *Eur. Phys. J.* **C80**, 109 (2020), arXiv:2001.11357 [gr-qc].
- [66] A. Ronveaux, *Applied Mathematics and Computation* **141**, 177 (2003), advanced Special Functions and Related Topics in Differential Equations, Third Melfi Workshop, Proceedings of the Melfi School on Advanced Topics in Mathematics and Physics.
- [67] R. S. Maier, *Journal of Differential Equations* **213**, 171 (2005).
- [68] C. W. Misner, K. S. Thorne, and J. A. Wheeler, *Gravitation* (W. H. Freeman, San Francisco, 1973).
- [69] R. A. D’Inverno, *Internationale Elektronische Rundschau* (1992).
- [70] G. G. L. Nashed, *Chaos Solitons Fractals* **15**, 841 (2003), arXiv:gr-qc/0301008 [gr-qc].
- [71] A. Sheykhi, *Phys. Rev. D* **86**, 024013 (2012).
- [72] A. Sheykhi, *Eur. Phys. J.* **C69**, 265 (2010), arXiv:1012.0383 [hep-th].
- [73] S. H. Hendi, A. Sheykhi, and M. H. Dehghani, *Eur. Phys. J.* **C70**, 703 (2010), arXiv:1002.0202 [hep-th].
- [74] A. Sheykhi, M. H. Dehghani, and S. H. Hendi, *Phys. Rev. D* **81**, 084040 (2010).
- [75] T. Shirafuji and G. G. L. Nashed, *Prog. Theor. Phys.* **98**, 1355 (1997), arXiv:gr-qc/9711010 [gr-qc].
- [76] Y.-Q. Wang, Y.-X. Liu, and S.-W. Wei, *Phys. Rev.* **D99**, 064036 (2019), arXiv:1811.08795 [gr-qc].
- [77] A. Zakria and A. Afzal, (2018), arXiv:1808.04361 [hep-th].
- [78] G. Cognola, O. Gorbunova, L. Sebastiani, and S. Zerbini, *Phys. Rev. D* **84**, 023515 (2011).
- [79] Y. Zheng and R.-J. Yang, *Eur. Phys. J.* **C78**, 682 (2018), arXiv:1806.09858 [gr-qc].
- [80] W. Kim and Y. Kim, *Phys. Lett.* **B718**, 687 (2012), arXiv:1207.5318 [gr-qc].
- [81] Y. Zheng and R. Yang, *The European Physical Journal C* **78** (2018), 10.1140/epjc/s10052-018-6167-4.
- [82] O. Ökcü and E. Aydiner, *Eur. Phys. J. C* **78**, 123 (2018), arXiv:1709.06426 [gr-qc].

