

ON THE GEOMETRIC AND RIEMANNIAN STRUCTURE OF THE SPACES OF GROUP EQUIVARIANT NON-EXPANSIVE OPERATORS

PASQUALE CASCARANO, PATRIZIO FROSINI, NICOLA QUERCIOLI, AND AMIR SAKI

ABSTRACT. Group equivariant non-expansive operators have been recently proposed as basic components in topological data analysis and deep learning. In this paper we study some geometric properties of the spaces of group equivariant operators and show how a space \mathcal{F} of group equivariant non-expansive operators can be endowed with the structure of a Riemannian manifold, so making available the use of gradient descent methods for the minimization of cost functions on \mathcal{F} . As an application of this approach, we also describe a procedure to select a finite set of representative group equivariant non-expansive operators in the considered manifold.

1. INTRODUCTION

The concept of group equivariant non-expansive operator (GENEO) has been recently proposed as a versatile tool in topological data analysis and deep learning [5, 9, 10]. We recall that an operator F is called equivariant with respect to a group G if the action of G commutes with F . The study of these operators appears to be interesting for several reasons. On the one hand, the use of GENEOs allows to formalize the role of the observer in data analysis and data comparison. This is an important goal as it appears clearer and clearer that an efficient approach to these fields of research requires to examine not only the data, but also the observer that analyzes them, since his/her role is sometimes more important than the role of data themselves. On the other hand, GENEOs could be seen as architectural components for building structures in artificial intelligence. Several researchers have pointed out that the use of equivariant operators could be of great relevance in the development of machine learning [3, 7, 17], since these operators can inject pre-existing knowledge into the system and allow us to increase our control on the construction of neural networks [4]. Furthermore, it is also worth noting that invariant and non-expansive operators can be used to reduce data variability [14, 15], and that in the last years equivariant transformations have been studied for learning symmetries [2, 18]. Finally, the research concerning the approximation and representation of observers could help us in understanding the role of conflicts and contradiction in the development of intelligence [8].

This focus on spaces of group equivariant non-expansive operators stresses the need for a study of the topological and geometrical structure of these spaces, in order to simplify their exploration and use. In particular, extending the mathematical framework proposed in [5], in this paper we show how a space of group equivariant non-expansive operators can be endowed with a suitable metric that takes into account the different probability of data. With reference to such a metric, we prove that when the spaces of data are compact, the space of GENEOs is compact too, so implying that it admits arbitrarily good approximations by finite sets of operators. Furthermore, we show that we can define a suitable Riemannian structure on finite dimensional submanifolds of the space of GENEOs and use this structure to approximate those manifolds by minimizing

2020 *Mathematics Subject Classification*. Primary: 55N31; Secondary: 58D20, 58D30, 62R40, 65D18, 68T09.

a cost function via gradient descent methods. The Riemannian structure we propose is based on the comparison of the action of GENEOS on data, taking into account the probability distribution on the set of admissible signals we are considering. At the end of the paper, we describe a computational experiment, where we use our approach to select a representative set of GENEOS from a given manifold of operators.

From our perspective, endowing the spaces of operators/observers acting on the data with new topological and geometrical structures is a key point in the present research concerning the mathematical aspects of machine learning. New results in this direction could allow exploiting GENEOS as components in the realization of new kinds of neural networks, taking advantage from the dimensionality reduction offered by Topological Data Analysis [5].

The outline of the paper is as follows. In Section 2 we illustrate our mathematical setting and prove its main properties. In Section 3 we show how our framework can be used to find a good finite approximation of the considered manifold of GENEOS. In Section 4 we conclude the paper with a brief discussion.

2. MATHEMATICAL SETTING

2.1. The space of admissible signals. Let X be a set and $(V, \langle \cdot, \cdot \rangle_V)$ be a finite dimensional inner product space whose elements belong to the space \mathbb{R}_b^X of bounded functions from X to \mathbb{R} . Then, V and \mathbb{R}^n are isomorphic as vector spaces, where n is the dimension of V . Denote the norm arising from $\langle \cdot, \cdot \rangle_V$ by $\|\cdot\|_V$. For any $\varphi \in V$, let us consider its L^∞ -norm $\|\varphi\|_\infty = \sup_{x \in X} |\varphi(x)|$. Note that $\|\cdot\|_V$ and $\|\cdot\|_\infty$ are equivalent, since V is finite dimensional. Hence, there exist two real numbers $\alpha, \beta > 0$ such that $\alpha\|\cdot\|_\infty \leq \|\cdot\|_V \leq \beta\|\cdot\|_\infty$. This implies that $\|\cdot\|_V$ and $\|\cdot\|_\infty$ induce the same topology τ_V on V .

Let us now choose a Borel measure λ on V and a non-negative integrable function $f : V \rightarrow \mathbb{R}$ such that $\int_V f d\lambda = 1$. Let us define a probability Borel measure μ on V by setting $\mu(A) = \int_A f d\lambda$ for any Borel set A in V . Finally, let us assume that the support of μ is a compact subspace Φ of V . We observe that $\mu(\Phi) = \int_\Phi f d\lambda = 1$. We will call Φ the space of *admissible signals* in probability. Let $\text{Aut}_\Phi(X)$ be the set of all bijections $g : X \rightarrow X$ such that $\varphi g, \varphi g^{-1} \in \Phi$ for any $\varphi \in \Phi$. $\text{Aut}_\Phi(X)$ naturally acts on Φ by composition on the right. Let us choose a subgroup G of $\text{Aut}_\Phi(X)$. For every $g \in G$, let us consider the map $\rho_g : \Phi \rightarrow \Phi$ taking each function $\varphi \in \Phi$ to φg . Since $\|\varphi_1 g - \varphi_2 g\|_\infty = \|\varphi_1 - \varphi_2\|_\infty$, the map ρ_g is continuous. Analogously, the map $\rho_{g^{-1}} = (\rho_g)^{-1}$ is continuous too. Hence ρ_g is a homeomorphism. Therefore, if A is a Borel set in V , then $\rho_g(A)$ is a Borel set in V . We assume that the following conditions are satisfied:

- (1) $\langle \cdot, \cdot \rangle_V$ is invariant under the action of G (i.e., $\langle \rho_g(\varphi_1), \rho_g(\varphi_2) \rangle_V = \langle \varphi_1, \varphi_2 \rangle_V$ for any $\varphi_1, \varphi_2 \in V$ and $g \in G$).
- (2) f, λ (and hence μ) are invariant under the action of G (i.e., if $g \in G$ then $f(\rho_g(\varphi)) = f(\varphi)$ for any $\varphi \in V$, and $\lambda(A) = \lambda(\rho_g(A))$ for every Borel set A in V).

Condition (1) immediately implies that $\|\cdot\|_V$ is invariant under the action of G .

Under the previous assumptions, we will say that V is a (finite dimensional) *probability inner product space with equivariance group G* .

2.2. The pseudo-distance on X induced by admissible signals. In this subsection, we show that the distance on Φ , induced by $\|\cdot\|_V$, can be pulled back to a pseudo-distance on X . We recall that a pseudo-distance on a set X is a distance d without the property $d(x, y) = 0 \implies x = y$. Note that the collection of all open balls $B_X(x, r) = \{y \in X : d(x, y) < r\}$ is a basis for a topology on X , which is called the *topology induced by d* .

The following lemma prepares us to define a pseudo-distance on X .

Lemma 2.1. *Let $x_1, x_2 \in X$. Then, the map $\xi_{x_1, x_2} : \Phi \rightarrow \mathbb{R}$ defined by $\xi_{x_1, x_2}(\varphi) = |\varphi(x_1) - \varphi(x_2)|$ for any $\varphi \in \Phi$, is continuous with respect to the topology induced by the norm $\|\cdot\|_V$ on Φ , and hence it is an integrable random variable with respect to λ .*

Proof. Let $\varphi_1, \varphi_2 \in \Phi$. Then,

$$\begin{aligned} |\xi_{x_1, x_2}(\varphi_1) - \xi_{x_1, x_2}(\varphi_2)| &= \left| |\varphi_1(x_1) - \varphi_1(x_2)| - |\varphi_2(x_1) - \varphi_2(x_2)| \right| \\ &\leq |(\varphi_1(x_1) - \varphi_1(x_2)) - (\varphi_2(x_1) - \varphi_2(x_2))| \\ &= |(\varphi_1(x_1) - \varphi_2(x_1)) - (\varphi_1(x_2) - \varphi_2(x_2))| \\ &\leq |\varphi_1(x_1) - \varphi_2(x_1)| + |\varphi_1(x_2) - \varphi_2(x_2)| \\ &\leq 2\|\varphi_1 - \varphi_2\|_\infty \leq \frac{2}{\alpha}\|\varphi_1 - \varphi_2\|_V. \end{aligned}$$

Therefore, ξ_{x_1, x_2} is continuous with respect to $\|\cdot\|_V$. \square

Now, we define a pseudo-distance Δ_X on X as follows:

$$\Delta_X(x_1, x_2) = \int_{\Phi} |\varphi(x_1) - \varphi(x_2)| f(\varphi) d\lambda, \quad \forall x_1, x_2 \in X.$$

In plain words, the distance between two points x_1, x_2 is set to be the expected value of the function ξ_{x_1, x_2} .

Another relevant pseudo-metric on X is the pseudo-metric D_X defined by setting

$$D_X(x_1, x_2) := \max_{\varphi \in \Phi} |\varphi(x_1) - \varphi(x_2)|.$$

The definition of D_X is well-posed, since Φ is compact. We observe that $\Delta_X \leq D_X$, so that the topology τ_{D_X} induced by D_X is finer than the topology τ_{Δ_X} induced by Δ_X .

In the sequel, whenever not differently specified, we assume that X is equipped with the topology arising from Δ_X .

Proposition 2.2. Each function $\varphi_0 \in \Phi$ is continuous with respect to Δ_X .

Proof. After choosing an $\varepsilon > 0$, let us consider the ball

$$B_\varepsilon = B_\Phi \left(\varphi_0, \frac{\varepsilon\alpha}{4} \right) = \left\{ \varphi \in \Phi : \|\varphi_0 - \varphi\|_V \leq \frac{\varepsilon\alpha}{4} \right\}.$$

Since φ_0 is in the essential support of f , $\mu(B_\varepsilon)$ is positive. For any $\varphi \in B_\varepsilon$, we have that $\|\varphi_0 - \varphi\|_\infty \leq \frac{1}{\alpha}\|\varphi_0 - \varphi\|_V \leq \frac{\varepsilon}{4}$, and hence for every $x \in X$

$$\begin{aligned} |\varphi(x) - \varphi_0(x)| &= |(\varphi_0(x) - \varphi_0(x_0)) + (\varphi(x) - \varphi_0(x)) + (\varphi_0(x_0) - \varphi(x_0))| \\ &\geq |\varphi_0(x) - \varphi_0(x_0)| - |\varphi(x) - \varphi_0(x)| - |\varphi_0(x_0) - \varphi(x_0)| \\ &\geq |\varphi_0(x) - \varphi_0(x_0)| - 2\|\varphi_0 - \varphi\|_\infty \\ &\geq |\varphi_0(x) - \varphi_0(x_0)| - \varepsilon/2. \end{aligned}$$

This implies that for every $x \in X$

$$\begin{aligned} \Delta_X(x, x_0) &= \int_{\Phi} |\varphi(x) - \varphi(x_0)| f(\varphi) d\lambda \\ &\geq \int_{B_\varepsilon} |\varphi(x) - \varphi(x_0)| f(\varphi) d\lambda \\ &\geq \mu(B_\varepsilon)(|\varphi_0(x) - \varphi_0(x_0)| - \varepsilon/2). \end{aligned}$$

It follows that $|\varphi_0(x) - \varphi_0(x_0)| \leq \frac{\Delta_X(x, x_0)}{\mu(B_\varepsilon)} + \varepsilon/2$. Therefore, if $\Delta_X(x, x_0) \leq (\varepsilon/2)\mu(B_\varepsilon)$, then $|\varphi_0(x) - \varphi_0(x_0)| \leq \varepsilon$. \square

Proposition 2.3. Every function $\varphi \in \Phi$ is non-expansive, and hence continuous, with respect to D_X .

Proof. If $x_1, x_2 \in X$, then for any function $\varphi \in \Phi$ we have that $|\varphi(x_1) - \varphi(x_2)| \leq \max_{\varphi \in \Phi} |\varphi(x_1) - \varphi(x_2)| = D_X(x_1, x_2)$. \square

We now recall that the initial topology τ_{in} on X with respect to Φ is the coarsest topology on X such that each function φ in Φ is continuous. From Proposition 2.2 and the definition of τ_{in} , it follows that $\tau_{\text{in}} \subseteq \tau_{\Delta_X}$. We already know that $\tau_{\Delta_X} \subseteq \tau_{D_X}$. Since Φ is compact, we know that τ_{in} and τ_{D_X} are the same (see Theorem 2.1 in [5]). Hence, $\tau_{D_X} = \tau_{\Delta_X} = \tau_{\text{in}}$.

Before proceeding, we recall the following lemma (see [11]):

Lemma 2.4. *Let (P, d) be a pseudo-metric space. The following conditions are equivalent:*

- (1) P is totally bounded;
- (2) every sequence in P admits a Cauchy subsequence.

The definition of the pseudo-metric D_X on X relies on Φ . Thus, properties on Φ naturally induce properties on X , as shown in the proof of the following statement.

Proposition 2.5. X is totally bounded with respect to D_X and Δ_X .

Proof. Because of Lemma 2.4, in order to prove that X is totally bounded with respect to D_X we have just to prove that every sequence $(x_i)_{i \in \mathbb{N}}$ in X admits a Cauchy subsequence with respect to D_X . Let us consider an arbitrary sequence $(x_i)_{i \in \mathbb{N}}$ in X and an arbitrarily small $\varepsilon > 0$. Since Φ is totally bounded, we can find a finite subset $\Phi_\varepsilon = \{\varphi_1, \dots, \varphi_n\}$ of Φ , such that $\Phi = \bigcup_{i=1}^n B_\Phi(\varphi_i, \varepsilon)$, where $B_\Phi(\varphi, \varepsilon) = \{\varphi' \in \Phi : \|\varphi' - \varphi\|_\infty < \varepsilon\}$. As a consequence, if $\varphi \in \Phi$, then there exists $\varphi_{\bar{k}} \in \Phi_\varepsilon$ such that $\|\varphi - \varphi_{\bar{k}}\|_\infty < \varepsilon$. Now, we consider the real sequence $(\varphi_1(x_i))_{i \in \mathbb{N}}$ that is bounded in \mathbb{R} because all the functions in Φ are bounded. From the Bolzano-Weierstrass Theorem it follows that we can extract a Cauchy subsequence $(\varphi_1(x_{i_h}))_{h \in \mathbb{N}}$. Then we consider the sequence $(\varphi_2(x_{i_h}))_{h \in \mathbb{N}}$. Since φ_2 is bounded, we can extract a Cauchy subsequence $(\varphi_2(x_{i_{h_t}}))_{t \in \mathbb{N}}$. We can repeat the same argument for any $\varphi_k \in \Phi_\varepsilon$. Thus, we obtain a subsequence $(x_{p_j})_{j \in \mathbb{N}}$ of $(x_i)_{i \in \mathbb{N}}$, such that $(\varphi_k(x_{p_j}))_{j \in \mathbb{N}}$ is a Cauchy sequence for any $k \in \{1, \dots, n\}$. Moreover, since Φ_ε is a finite set, there exists an index \bar{j} such that for any $k \in \{1, \dots, n\}$ we have that

$$(2.1) \quad |\varphi_k(x_{p_r}) - \varphi_k(x_{p_s})| < \varepsilon, \quad \text{for all } r, s \geq \bar{j}.$$

We observe that \bar{j} does not depend on φ , but only on ε and Φ_ε .

In order to prove that $(x_{p_j})_{j \in \mathbb{N}}$ is a Cauchy sequence in X with respect to D_X , we observe that for any $r, s \in \mathbb{N}$ the following inequalities hold:

$$(2.2) \quad \begin{aligned} |\varphi(x_{p_r}) - \varphi(x_{p_s})| &= |\varphi(x_{p_r}) - \varphi_{\bar{k}}(x_{p_r}) + \varphi_{\bar{k}}(x_{p_r}) - \varphi_{\bar{k}}(x_{p_s}) + \varphi_{\bar{k}}(x_{p_s}) - \varphi(x_{p_s})| \\ &\leq |\varphi(x_{p_r}) - \varphi_{\bar{k}}(x_{p_r})| + |\varphi_{\bar{k}}(x_{p_r}) - \varphi_{\bar{k}}(x_{p_s})| + |\varphi_{\bar{k}}(x_{p_s}) - \varphi(x_{p_s})| \\ &\leq \|\varphi - \varphi_{\bar{k}}\|_\infty + |\varphi_{\bar{k}}(x_{p_r}) - \varphi_{\bar{k}}(x_{p_s})| + \|\varphi_{\bar{k}} - \varphi\|_\infty. \end{aligned}$$

It follows that $|\varphi(x_{p_r}) - \varphi(x_{p_s})| < 3\varepsilon$ for every $\varphi \in \Phi$ and every $r, s \geq \bar{j}$. Thus, $\sup_{\varphi \in \Phi} |\varphi(x_{p_r}) - \varphi(x_{p_s})| = D_X(x_{p_r}, x_{p_s}) \leq 3\varepsilon$ for any $r, s \geq \bar{j}$. Hence, the sequence $(x_{p_j})_{j \in \mathbb{N}}$ is a Cauchy sequence in X with respect to D_X . This proves that X is totally bounded with respect to D_X . Since $\Delta_X \leq D_X$, X is also totally bounded with respect to Δ_X . \square

Remark 2.6. The proof of Proposition 2.5 just relies on the assumption that Φ is totally bounded, without using the compactness of that space.

Corollary 2.7. If X is complete, then X is compact.

Proof. It follows from Proposition 2.5, by recalling that in pseudo-metric spaces a set is compact if and only if it is complete and totally bounded [11]. \square

2.3. Φ -preserving homeomorphisms. We know that the elements of $\text{Aut}_\Phi(X)$ are isometries with respect to D_X , and hence homeomorphisms with respect to D_X (cf. [5]). In the following, we show that each element of G is also an isometry, and hence a homeomorphism, with respect to Δ_X .

Proposition 2.8. Each $g \in G$ is an isometry with respect to Δ_X .

Proof. Let λ_g be the Borel measure on V defined by setting $\lambda_g(A) = \lambda(\rho_g(A))$ for any Borel set A in V . From the invariance of λ under the action of G , $\lambda_g = \lambda$. By applying a change of variable, the invariance of f under the action of each $g \in G$ implies that

$$\begin{aligned} \Delta_X(g(x_1), g(x_2)) &= \int_{\Phi} |\varphi g(x_1) - \varphi g(x_2)| f(\varphi) d\lambda \\ &= \int_{\Phi} |\varphi g(x_1) - \varphi g(x_2)| f(\varphi g) d\lambda_g \\ &= \int_{\Phi} |\varphi(x_1) - \varphi(x_2)| f(\varphi) d\lambda = \Delta_X(x_1, x_2) \end{aligned}$$

for any $x_1, x_2 \in X$. \square

Now, we turn G into a pseudo-distance space by using the measure and the norm on Φ . To do this, we need the following lemma.

Lemma 2.9. *Let $g_1, g_2 \in G$. Then, the map $\xi_{g_1, g_2} : \Phi \rightarrow \mathbb{R}$ sending each φ to $\|\varphi g_1 - \varphi g_2\|_V$ is a continuous map with respect to $\|\cdot\|_V$, and hence an integrable map with respect to λ .*

Proof. Let $\varphi_1, \varphi_2 \in \Phi$. Since $\|\cdot\|_V$ is invariant under the action of G , we have that

$$\begin{aligned} |\xi_{g_1, g_2}(\varphi_1) - \xi_{g_1, g_2}(\varphi_2)| &= | \|\varphi_1 g_1 - \varphi_1 g_2\|_V - \|\varphi_2 g_1 - \varphi_2 g_2\|_V | \\ &\leq \|(\varphi_1 g_1 - \varphi_1 g_2) - (\varphi_2 g_1 - \varphi_2 g_2)\|_V \\ &= \|(\varphi_1 g_1 - \varphi_2 g_1) - (\varphi_1 g_2 - \varphi_2 g_2)\|_V \\ &\leq \|\varphi_1 g_1 - \varphi_2 g_1\|_V + \|\varphi_1 g_2 - \varphi_2 g_2\|_V = 2\|\varphi_1 - \varphi_2\|_V. \end{aligned}$$

Therefore, ξ_{g_1, g_2} is continuous with respect to $\|\cdot\|_V$. \square

Now, we can define the following pseudo-distance on G :

$$\Delta_G(g_1, g_2) = \int_{\Phi} \|\varphi g_1 - \varphi g_2\|_V f(\varphi) d\lambda, \quad \forall g_1, g_2 \in G.$$

In [5] a different pseudo-distance D_G on G has been considered, defined by setting $D_G(g_1, g_2) := \max_{\varphi \in \Phi} \|\varphi g_1 - \varphi g_2\|_\infty$. The definition of D_G is well-posed, since Φ is compact. We observe that

$$\begin{aligned} \Delta_G(g_1, g_2) &= \int_{\Phi} \|\varphi g_1 - \varphi g_2\|_V f(\varphi) d\lambda \\ &\leq \int_{\Phi} \beta \|\varphi g_1 - \varphi g_2\|_\infty f(\varphi) d\lambda \\ &\leq \int_{\Phi} \beta D_G(g_1, g_2) f(\varphi) d\lambda \\ &= \beta D_G(g_1, g_2) \int_{\Phi} f(\varphi) d\lambda \\ &= \beta D_G(g_1, g_2) \end{aligned}$$

for every $g_1, g_2 \in G$. The inequality $\Delta_G \leq \beta D_G$ implies that the topology τ_{D_G} induced by D_G is finer than the topology τ_{Δ_G} induced by Δ_G . We recall that if (G, D_G) is complete, then it is compact [5, Theorem 3].

In the sequel, whenever not differently specified, we consider G as a pseudo-distance space (and hence a topological space) with respect to Δ_G .

Lemma 2.10. *Let $g_1, g_2, g_3 \in G$. We have that*

$$\Delta_G(g_1, g_2) = \Delta_G(g_1 g_3, g_2 g_3) = \Delta_G(g_3 g_1, g_3 g_2).$$

Proof. The first equality follows directly from the invariance of the norm $\|\cdot\|_V$ under the action of G . Now we show that $\Delta_G(g_1, g_2) = \Delta_G(g_3 g_1, g_3 g_2)$. By applying a change of variable, the invariance of λ and f under the action of G implies that

$$\begin{aligned} \Delta_G(g_3 g_1, g_3 g_2) &= \int_{\Phi} \|\varphi g_3 g_1 - \varphi g_3 g_2\|_V f(\varphi) d\lambda \\ &= \int_{\Phi} \|\varphi g_3 g_1 - \varphi g_3 g_2\|_V f(\varphi g_3) d\lambda_{g_3} \\ &= \int_{\Phi} \|\varphi g_1 - \varphi g_2\|_V f(\varphi) d\lambda \\ &= \Delta_G(g_1, g_2) \end{aligned}$$

where λ_{g_3} is the Borel measure on V defined by setting $\lambda_{g_3}(A) = \lambda(\rho_{g_3}(A))$ for any Borel set A in V . \square

Proposition 2.11. The group G is a topological group. Further, the action of G on Φ by composition on the right is continuous.

Proof. First, we show that G is a topological group. Let $\sigma : G \times G \rightarrow G$ and $\iota : G \rightarrow G$ be the composition and the inverse maps, respectively. We consider the product topology on $G \times G$. We must show that σ and ι are continuous. To show that σ is continuous, let $(g_1, g_2), (g'_1, g'_2) \in G \times G$. Using Lemma 2.10, we have that

$$\begin{aligned} \Delta_G(g_1 g_2, g'_1 g'_2) &= \Delta_G(g_1, g'_1 g'_2 g_2^{-1}) \\ &\leq \Delta_G(g_1, g'_1) + \Delta_G(g'_1, g'_1 g'_2 g_2^{-1}) \\ &= \Delta_G(g_1, g'_1) + \Delta_G(id_X, g'_2 g_2^{-1}) \\ &= \Delta_G(g_1, g'_1) + \Delta_G(g_2, g'_2). \end{aligned}$$

It follows that the composition map σ is continuous. Now, we show that ι is continuous. Consider $h_1, h_2 \in G$. We have that

$$\begin{aligned} \Delta_G(h_1^{-1}, h_2^{-1}) &= \Delta_G(h_1^{-1}h_2, h_2^{-1}h_2) \\ &= \Delta_G(h_1^{-1}h_2, id_X) \\ &= \Delta_G(h_1^{-1}h_2, h_1^{-1}h_1) \\ &= \Delta_G(h_2, h_1) \\ &= \Delta_G(h_1, h_2). \end{aligned}$$

This proves that ι is an isometry, and hence it is continuous.

Therefore, G is a topological group.

Let us now assume that $\rho : \Phi \times G \rightarrow \Phi$ is the natural action of G on Φ (i.e. $\rho(\varphi, g) = \varphi g$ for any $\varphi \in \Phi$ and $g \in G$). We have to prove that ρ is continuous, when $\Phi \times G$ is endowed with the product topology. Let $(\varphi_1, g_1), (\varphi_2, g_2) \in \Phi \times G$ and $\varepsilon > 0$. Let us define B_ε as the ball $B_\Phi(\varphi_2, \frac{\varepsilon}{4})$ in Φ with respect to $\|\cdot\|_V$. Since φ_2 is in the essential support of f , $\mu(B_\varepsilon)$ is positive. We will show that if $\|\varphi_1 - \varphi_2\|_V \leq \frac{\varepsilon}{4}$ and $\Delta_G(g_1, g_2) \leq \frac{\varepsilon}{4}\mu(B_\varepsilon)$, then $\|\rho(\varphi_1, g_1) - \rho(\varphi_2, g_2)\| \leq \varepsilon$. Since $\|\varphi - \varphi_2\|_V \leq \frac{\varepsilon}{4}$ for every $\varphi \in B_\varepsilon$, we have that

$$\begin{aligned} \Delta_G(g_1, g_2) &= \int_{\Phi} \|\varphi g_1 - \varphi g_2\|_V f(\varphi) d\lambda \\ &\geq \int_{B_\varepsilon} \|\varphi g_1 - \varphi g_2\|_V f(\varphi) d\lambda \\ &= \int_{B_\varepsilon} \|(\varphi g_1 - \varphi_2 g_1) + (\varphi_2 g_1 - \varphi_2 g_2) + (\varphi_2 g_2 - \varphi g_2)\|_V f(\varphi) d\lambda \\ &\geq \int_{B_\varepsilon} (\|\varphi_2 g_1 - \varphi_2 g_2\|_V - \|\varphi g_1 - \varphi_2 g_1\|_V - \|\varphi_2 g_2 - \varphi g_2\|_V) f(\varphi) d\lambda \\ &= \int_{B_\varepsilon} (\|\varphi_2 g_1 - \varphi_2 g_2\|_V - 2\|\varphi - \varphi_2\|_V) f(\varphi) d\lambda \\ &\geq \mu(B_\varepsilon) \left(\|\varphi_2 g_1 - \varphi_2 g_2\|_V - \frac{\varepsilon}{2} \right). \end{aligned}$$

It follows that $\|\varphi_2 g_1 - \varphi_2 g_2\|_V \leq \frac{\Delta_G(g_1, g_2)}{\mu(B_\varepsilon)} + \frac{\varepsilon}{2}$. Therefore,

$$\begin{aligned} \|\rho(\varphi_1, g_1) - \rho(\varphi_2, g_2)\|_V &= \|\varphi_1 g_1 - \varphi_2 g_2\|_V \\ &\leq \|\varphi_1 g_1 - \varphi_2 g_1\|_V + \|\varphi_2 g_1 - \varphi_2 g_2\|_V \\ &\leq \|\varphi_1 - \varphi_2\|_V + \frac{\Delta_G(g_1, g_2)}{\mu(B_\varepsilon)} + \frac{\varepsilon}{2} \\ &\leq \frac{\varepsilon}{4} + \frac{\varepsilon}{4} + \frac{\varepsilon}{2} = \varepsilon. \end{aligned}$$

Consequently, ρ is continuous. □

In order to study the compactness of G , we need the following result.

Proposition 2.12. G is totally bounded with respect to D_G and Δ_G .

Proof. Let $(g_i)_{i \in \mathbb{N}}$ and ε be a sequence in G and a positive real number, respectively. Given that Φ is totally bounded, we can find a finite subset $\Phi_\varepsilon = \{\varphi_1, \dots, \varphi_n\}$ such that if $\varphi \in \Phi$ then there exists $\varphi_{\bar{k}} \in \Phi_\varepsilon$ for which $\|\varphi_{\bar{k}} - \varphi\|_\infty < \varepsilon$.

Let us consider the sequence $(\varphi_1 g_i)_{i \in \mathbb{N}}$ in Φ . Since Φ is totally bounded, it follows that we can extract a Cauchy subsequence $(\varphi_1 g_{i_h})_{h \in \mathbb{N}}$. Then we consider the sequence $(\varphi_2 g_{i_h})_{h \in \mathbb{N}}$. Again, we can extract a Cauchy subsequence $(\varphi_2 g_{i_{h_t}})_{t \in \mathbb{N}}$. We can repeat the

same argument for any $\varphi_k \in \Phi_\varepsilon$. Thus, we are able to extract a subsequence $(g_{p_j})_{j \in \mathbb{N}}$ of $(g_i)_{i \in \mathbb{N}}$ such that $(\varphi_k g_{p_j})_{j \in \mathbb{N}}$ is a Cauchy sequence for any $k \in \{1, \dots, n\}$. For the finiteness of set Φ_ε , we can find an index \bar{j} such that for any $k \in \{1, \dots, n\}$

$$(2.3) \quad \|\varphi_k g_{p_r} - \varphi_k g_{p_s}\|_\infty < \varepsilon, \text{ for every } s, r \geq \bar{j}.$$

We observe that \bar{j} does not depend on φ , but only on ε and Φ_ε .

In order to prove that $(g_{p_j})_{j \in \mathbb{N}}$ is a Cauchy sequence, we observe that for any $r, s \in \mathbb{N}$ we have

$$(2.4) \quad \begin{aligned} \|\varphi g_{p_r} - \varphi g_{p_s}\|_\infty &\leq \|\varphi g_{p_r} - \varphi_{\bar{k}} g_{p_r}\|_\infty + \|\varphi_{\bar{k}} g_{p_r} - \varphi_{\bar{k}} g_{p_s}\|_\infty + \|\varphi_{\bar{k}} g_{p_s} - \varphi g_{p_s}\|_\infty \\ &= \|\varphi - \varphi_{\bar{k}}\|_\infty + \|\varphi_{\bar{k}} g_{p_r} - \varphi_{\bar{k}} g_{p_s}\|_\infty + \|\varphi_{\bar{k}} - \varphi\|_\infty. \end{aligned}$$

Since $\|\varphi_{\bar{k}} - \varphi\|_\infty < \varepsilon$, we get $\|\varphi g_{p_r} - \varphi g_{p_s}\|_\infty < 3\varepsilon$ for every $r, s \geq \bar{j}$. Thus, $D_G(g_{p_r}, g_{p_s}) \leq 3\varepsilon$. Hence, the sequence $(g_{p_j})_{j \in \mathbb{N}}$ is a Cauchy sequence with respect to D_G . Therefore, by Lemma 2.4, G is totally bounded with respect to D_G . Since $\Delta_G \leq \beta D_G$, G is also totally bounded with respect to Δ_G . \square

Therefore, the following statement holds, by recalling that in pseudo-metric spaces a set is compact if and only if it is complete and totally bounded [11].

Corollary 2.13. If G is complete, then G is compact.

2.4. Group equivariant operators. Let $V_1 \subseteq \mathbb{R}_b^{X_1}$ and $V_2 \subseteq \mathbb{R}_b^{X_2}$ be two finite dimensional probability inner product spaces with equivariance groups G_1 and G_2 , respectively. Let $\langle \cdot, \cdot \rangle_{V_i}$, $\|\cdot\|_{V_i}$, λ_{V_i} , f_{V_i} be the inner product, the norm, the Borel measure and the probability density function considered on V_i , for $i = 1, 2$ (all of them are G_i -invariant). Also, for $i = 1, 2$, we consider the probability Borel measure μ_{V_i} by setting $\mu_{V_i}(A_i) = \int_{A_i} f_{V_i}(\phi) d\lambda_{V_i}$ for any Borel set $A_i \subseteq V_i$. We assume that the support of μ_{V_i} is compact, and we denote it by Φ_i , for $i = 1, 2$. Moreover, we recall that since $\|\cdot\|_{V_i}$ and $\|\cdot\|_\infty$ are equivalent in V_i , there exist two real numbers $\beta_i, \alpha_i > 0$ such that

$$(2.5) \quad \alpha_i \|\cdot\|_\infty \leq \|\cdot\|_{V_i} \leq \beta_i \|\cdot\|_\infty$$

for $i = 1, 2$.

Let us consider the Bochner space $L^2(\Phi_1, V_2)$ of all square integrable maps from Φ_1 to V_2 . Explicitly:

$$(2.6) \quad L^2(\Phi_1, V_2) := \left\{ F: \Phi_1 \rightarrow V_2 : \int_{\Phi_1} \|F(\varphi)\|_{V_2}^2 f_{V_1}(\varphi) d\lambda_{V_1} \text{ exists and is finite} \right\}.$$

Now, we define an inner product on $L^2(\Phi_1, V_2)$ as follows:

$$(2.7) \quad \langle F_1, F_2 \rangle := \int_{\Phi_1} \langle F_1(\varphi), F_2(\varphi) \rangle_{V_2} f_{V_1}(\varphi) d\lambda_{V_1} \quad \forall F_1, F_2 \in L^2(\Phi_1, V_2).$$

It is well known that the space $L^2(\Phi_1, V_2)$ is a Hilbert space.

Let us select a homomorphism $T: G_1 \rightarrow G_2$. A *group equivariant operator* (or simply a GEO) from (Φ_1, G_1) to (Φ_2, G_2) is a continuous map $F: (\Phi_1, \|\cdot\|_{V_1}) \rightarrow (\Phi_2, \|\cdot\|_{V_2})$ satisfying the property $F(\varphi g) = F(\varphi)T(g)$ for any φ in Φ_1 and g in G_1 .

Remark 2.14. Since $\|\cdot\|_{V_1}$ and $\|\cdot\|_\infty$ are equivalent in V_1 and, $\|\cdot\|_{V_2}$ and $\|\cdot\|_\infty$ are equivalent in V_2 , a GEO $F: (\Phi_1, \|\cdot\|_{V_1}) \rightarrow (\Phi_2, \|\cdot\|_{V_2})$ is continuous (and hence Borel measurable) also with respect to the L^∞ -norm defined on Φ_1 and Φ_2 .

We define the following norms on the space of GEOs from (Φ_1, G_1) to (Φ_2, G_2) :

$$(1) \quad \|F\|_\infty := \max_{\varphi \in \Phi_1} \|F(\varphi)\|_\infty$$

- (2) $\|F\|_{V_2} := \max_{\varphi \in \Phi_1} \|F(\varphi)\|_{V_2}$
 (3) $\|F\|_{L^2} := \left(\int_{\Phi_1} \|F(\varphi)\|_{V_2}^2 f_{V_1}(\varphi) d\lambda_{V_1} \right)^{\frac{1}{2}}$.

Lemma 2.15. $\|\cdot\|_{\infty}$ and $\|\cdot\|_{V_2}$ induce the same topology on the space of GEOs from (Φ_1, G_1) to (Φ_2, G_2) .

Proof. We already know that for any φ in Φ_1 :

$$\alpha_2 \|F(\varphi)\|_{\infty} \leq \|F(\varphi)\|_{V_2} \leq \beta_2 \|F(\varphi)\|_{\infty}.$$

Hence, we have that:

$$\alpha_2 \max_{\varphi \in \Phi_1} \|F(\varphi)\|_{\infty} \leq \max_{\varphi \in \Phi_1} \|F(\varphi)\|_{V_2} \leq \beta_2 \max_{\varphi \in \Phi_1} \|F(\varphi)\|_{\infty}.$$

Therefore, $\|\cdot\|_{\infty}$ and $\|\cdot\|_{V_2}$ are equivalent norms, and hence they induce the same topology on the space of GEOs from (Φ_1, G_1) to (Φ_2, G_2) . \square

Lemma 2.16. *The topology induced by $\|\cdot\|_{V_2}$ on the space of GEOs from (Φ_1, G_1) to (Φ_2, G_2) is finer than the one induced by $\|\cdot\|_{L^2}$.*

Proof. Let us consider a GEO F from (Φ_1, G_1) to (Φ_2, G_2) . We have that

$$\begin{aligned} \|F\|_{L^2} &= \left(\int_{\Phi_1} \|F(\varphi)\|_{V_2}^2 f_{V_1}(\varphi) d\lambda_{V_1} \right)^{\frac{1}{2}} \\ &\leq \left(\int_{\Phi_1} \left(\max_{\varphi \in \Phi_1} \|F(\varphi)\|_{V_2}^2 \right) f_{V_1}(\varphi) d\lambda_{V_1} \right)^{\frac{1}{2}} \\ &= \|F\|_{V_2} \left(\int_{\Phi_1} f_{V_1}(\varphi) d\lambda_{V_1} \right)^{\frac{1}{2}} \\ &= \|F\|_{V_2}. \end{aligned}$$

The above inequality directly implies the statement of the lemma. \square

We can now define the concept of GENEEO.

Definition 2.17. A group equivariant non-expansive operator (in brief, GENEEO) from $((\Phi_1, \|\cdot\|_{V_1}), G_1)$ to $((\Phi_2, \|\cdot\|_{V_2}), G_2)$ is a GEO F from (Φ_1, G_1) to (Φ_2, G_2) such that

$$\|F(\varphi) - F(\varphi')\|_{V_2} \leq \|\varphi - \varphi'\|_{V_1}$$

for every $\varphi, \varphi' \in \Phi_1$.

Remark 2.18. In [5], another definition of a GENEEO was given. Indeed, in the aforementioned paper, a GENEEO is a map between two compact spaces of functions Ψ_1 and Ψ_2 , which is group equivariant with respect to a homomorphism $T : G_1 \rightarrow G_2$ and non-expansive with respect to the L^∞ -norm. In the current framework, we call the aforementioned operators L^∞ -GENEEOs. One could equip the space of L^∞ -GENEEOs with the L^∞ -norm. It was shown in [5] that the space of all L^∞ -GENEEOs from (Ψ_1, G_1) to (Ψ_2, G_2) is compact.

A key property of the space of GENEEOs in our framework is stated by the following theorem.

Theorem 2.19. *The space of GENEEOs from $((\Phi_1, \|\cdot\|_{V_1}), G_1)$ to $((\Phi_2, \|\cdot\|_{V_2}), G_2)$ is compact with respect to the norm $\|\cdot\|_{L^2}$.*

Proof. Let (F_i) be a sequence of GENEOS from $((\Phi_1, \|\cdot\|_{V_1}), G_1)$ to $((\Phi_2, \|\cdot\|_{V_2}), G_2)$. It will suffice to prove that there exists a subsequence (F_{i_j}) of (F_i) that converges in the $\|\cdot\|_{L^2}$ -topology. From (2.5) it follows that for every index i and every $\varphi, \varphi' \in \Phi_1$

$$\alpha_2 \|F_i(\varphi) - F_i(\varphi')\|_\infty \leq \|F_i(\varphi) - F_i(\varphi')\|_{V_2} \leq \|\varphi - \varphi'\|_{V_1} \leq \beta_1 \|\varphi - \varphi'\|_\infty$$

and hence

$$\|F_i(\varphi) - F_i(\varphi')\|_\infty \leq \frac{\beta_1}{\alpha_2} \|\varphi - \varphi'\|_\infty.$$

Therefore, $\left\| \frac{\alpha_2}{\beta_1} F_i(\varphi) - \frac{\alpha_2}{\beta_1} F_i(\varphi') \right\|_\infty \leq \|\varphi - \varphi'\|_\infty$, and each map $\frac{\alpha_2}{\beta_1} F_i$ is an L^∞ -GENEO from (Φ_1, G_1) to (Φ'_2, G_2) , where $\Phi'_2 = \frac{\alpha_2}{\beta_1} \Phi_2$. Since the space of L^∞ -GENEOs from (Φ_1, G_1) to (Φ'_2, G_2) is compact with respect to $\|\cdot\|_\infty$ (see Theorem 7 in [5]), we can consider the sequence $\left(\frac{\alpha_2}{\beta_1} F_i \right)$ and extract a subsequence $\left(\frac{\alpha_2}{\beta_1} F_{i_j} \right)$ that converges with respect to $\|\cdot\|_\infty$ to an L^∞ -GENEO \bar{F} from (Φ_1, G_1) to (Φ'_2, G_2) . The maps $\frac{\beta_1}{\alpha_2} \bar{F}$ and (F_{i_j}) are GEOs from (Φ_1, G_1) to (Φ_2, G_2) , and we observe that (F_{i_j}) converges to $\frac{\beta_1}{\alpha_2} \bar{F}$ with respect to $\|\cdot\|_\infty$. From Lemma 2.15, it follows that (F_{i_j}) converges to $\frac{\beta_1}{\alpha_2} \bar{F}$ with respect to $\|\cdot\|_{V_2}$.

Since $\lim_{j \rightarrow \infty} \sup_{\varphi \in \Phi_1} \left\| \frac{\beta_1}{\alpha_2} \bar{F}(\varphi) - F_{i_j}(\varphi) \right\|_{V_2} = 0$, then $\frac{\beta_1}{\alpha_2} \bar{F}(\varphi) = \lim_{j \rightarrow \infty} F_{i_j}(\varphi)$ with respect to the norm $\|\cdot\|_{V_2}$ (or, equivalently, the norm $\|\cdot\|_\infty$) for every $\varphi \in \Phi_1$. Therefore, for any pair φ_1, φ_2 in Φ_1

$$\begin{aligned} \left\| \frac{\beta_1}{\alpha_2} \bar{F}(\varphi_1) - \frac{\beta_1}{\alpha_2} \bar{F}(\varphi_2) \right\|_{V_2} &= \left\| \lim_{j \rightarrow \infty} F_{i_j}(\varphi_1) - \lim_{j \rightarrow \infty} F_{i_j}(\varphi_2) \right\|_{V_2} \\ &= \lim_{j \rightarrow \infty} \|F_{i_j}(\varphi_1) - F_{i_j}(\varphi_2)\|_{V_2} \\ &\leq \lim_{j \rightarrow \infty} \|\varphi_1 - \varphi_2\|_{V_1} \\ &= \|\varphi_1 - \varphi_2\|_{V_1}. \end{aligned}$$

This proves that $\frac{\beta_1}{\alpha_2} \bar{F}$ is a GENEEO from $(\Phi_1, \|\cdot\|_{V_1})$ to $(\Phi_2, \|\cdot\|_{V_2})$. Since (F_{i_j}) converges to $\frac{\beta_1}{\alpha_2} \bar{F}$ with respect to $\|\cdot\|_{V_2}$, Lemma 2.16 guarantees that (F_{i_j}) converges to $\frac{\beta_1}{\alpha_2} \bar{F}$ with respect to $\|\cdot\|_{L^2}$. \square

The compactness of the space of GENEOS with respect to the topology induced by the norm $\|\cdot\|_{L^2}$ guarantees that such a space can be approximated by a finite set, as stated by the following result.

Corollary 2.20. For every $\varepsilon > 0$, the space \mathcal{F}^{all} of all GENEOS from $((\Phi_1, \|\cdot\|_{V_1}), G_1)$ to $((\Phi_2, \|\cdot\|_{V_2}), G_2)$ admits a finite subset \mathcal{F} such that for every $F \in \mathcal{F}^{\text{all}}$ there exists an $F' \in \mathcal{F}$ such that $\|F - F'\|_{L^2} < \varepsilon$.

Proof. It immediately follows from Theorem 2.19, by considering a finite subcover of the open cover of \mathcal{F}^{all} whose elements are the balls of radius ε centered at points of \mathcal{F}^{all} . \square

Another important property of the space \mathcal{F}^{all} is stated by the next result.

Proposition 2.21. If Φ_2 is convex, then the set \mathcal{F}^{all} of all GENEOS from $((\Phi_1, \|\cdot\|_{V_1}), G_1)$ to $((\Phi_2, \|\cdot\|_{V_2}), G_2)$ is convex.

Proof. Let us consider $F, F' \in \mathcal{F}^{\text{all}}$ and a value $t \in [0, 1]$. We can define an operator $F_t: \Phi_1 \rightarrow \Phi_2$ by setting $F_t(\varphi) := (1 - t)F(\varphi) + tF'(\varphi)$ for any $\varphi \in \Phi_1$. Note that the

convexity of Φ_2 ensures us that F_t is well defined. First we prove that F_t is a GEO from (Φ_1, G_1) to (Φ_2, G_2) . Since F and F' are equivariant, for every $\varphi \in \Phi_1$ and every $g \in G_1$

$$F_t(\varphi g) = (1-t)F(\varphi g) + tF'(\varphi g) = (1-t)F(\varphi)T(g) + tF'(\varphi)T(g) = F_t(\varphi)T(g).$$

Since F and F' are non-expansive, F_t is non-expansive:

$$\begin{aligned} \|F_t(\varphi) - F_t(\varphi')\|_{V_2} &= \|((1-t)F(\varphi) + tF'(\varphi)) - ((1-t)F(\varphi') + tF'(\varphi'))\|_{V_2} \\ &= \|(1-t)(F(\varphi) - F(\varphi')) + t(F'(\varphi) - F'(\varphi'))\|_{V_2} \\ &\leq (1-t)\|F(\varphi) - F(\varphi')\|_{V_2} + t\|F'(\varphi) - F'(\varphi')\|_{V_2} \\ &\leq (1-t)\|\varphi - \varphi'\|_{V_1} + t\|\varphi - \varphi'\|_{V_1} \\ &= \|\varphi - \varphi'\|_{V_1}. \end{aligned}$$

Therefore F_t is a GENEIO from $((\Phi_1, \|\cdot\|_{V_1}), G_1)$ to $((\Phi_2, \|\cdot\|_{V_2}), G_2)$. \square

2.5. Submanifolds of GENEIOs. In this subsection we discuss how it is possible to define a Riemannian structure on a manifold of GENEIOs. We also briefly recall some basic definitions and results about Hilbert manifolds and Riemannian manifolds. For further details and the general theory, see [1, 12].

Since $L^2(\Phi, V_2)$ is a Hilbert space, it has a natural structure of Hilbert manifold with a single global chart given by the identity function on $L^2(\Phi, V_2)$. Each tangent space $T_p L^2(\Phi, V_2)$ at any point $p \in L^2(\Phi, V_2)$ is canonically isomorphic to $L^2(\Phi, V_2)$ itself. We can give a Riemannian structure on $L^2(\Phi, V_2)$ by defining a metric g as $g(v, w)(p) = \langle v, w \rangle$ for every $v, w \in T_p L^2(\Phi, V_2)$, where $\langle \cdot, \cdot \rangle$ is the inner product on $L^2(\Phi, V_2)$. Now, let M be a C^k -submanifold of $L^2(\Phi, V_2)$, such that each element of M is a GENEIO. Hence, M naturally inherits a Riemannian structure from $L^2(\Phi, V_2)$. In real applications we are often interested in finite dimensional manifolds. For this reason, we give an explicit definition. A subset M is called an m -dimensional C^k -submanifold of $L^2(\Phi, V_2)$, if for all $p \in M$ there exist a neighborhood U of p in $L^2(\Phi, V_2)$, an open set $O \subseteq \mathbb{R}^m$ and a C^k -map $\phi: O \rightarrow L^2(\Phi, V_2)$, such that

- (1) $\phi: O \rightarrow U \cap M$ is a homeomorphism;
- (2) the differential $D\phi(x)$ is injective for every $x \in O$.

We call ϕ a parametrization in p . In this case, we have an explicit formula for the tangent space $T_p M$ depending on ϕ :

$$T_p M = D\phi(x)\mathbb{R}^m, \quad x = \phi^{-1}(p).$$

3. AN APPLICATION OF OUR MODEL: THE USE OF THE PROPOSED FRAMEWORK IN SELECTING GENEIOs

In this section, we will assume that X is the torus $S^1 \times S^1$. The usual embedding of S^1 in \mathbb{R}^2 entails that we can endow X with the metric induced by the natural immersion in \mathbb{R}^4 . Given a positive integer n , we define the finite group G whose elements are the isometries of X that preserve the set $Y = \left\{ \left(\frac{2\pi i}{n}, \frac{2\pi j}{n} \right) : i, j \in \{0, \dots, n-1\} \right\} \subseteq S^1 \times S^1$. Moreover, we consider the function $\chi: X \rightarrow \mathbb{R}$ as in the following:

$$(3.1) \quad \chi(x, y) := \begin{cases} 1 & (x, y) \in \gamma \times \gamma \\ 0 & \text{otherwise} \end{cases}$$

where $\gamma \subset S^1$ is the arc connecting $-\frac{\pi}{n}$ and $\frac{\pi}{n}$ counterclockwise.

We assume V_1, V_2 both equal to the n^2 -dimensional vector space V whose elements are the functions $\varphi : X \rightarrow \mathbb{R}$ that can be expressed as:

$$(3.2) \quad \varphi(\alpha, \beta) := \sum_{i=0}^{n-1} \sum_{j=0}^{n-1} a_i^j \chi \left(\alpha - \frac{2\pi i}{n}, \beta - \frac{2\pi j}{n} \right),$$

where $a_i^j \in \mathbb{R}$ for every $i, j \in \{0, \dots, n-1\}$. Moreover, we consider $f_V = f_{V_1} = f_{V_2}$, $\mu_V = \mu_{V_1} = \mu_{V_2}$ and $\Phi = \Phi_1 = \Phi_2$ as the ones assumed in Section 2.4.

Our manifold M of GENEOS is the one containing every GENEOS F defined by setting:

$$(3.3) \quad F(\varphi)(\alpha, \beta) := \frac{1}{4\sqrt{m}} \left(\sum_{t=1}^m \sum_{r,s \in \{-1,1\}} u_t \varphi \left(\alpha + r \frac{k_t \pi}{n}, \beta + s \frac{h_t \pi}{n} \right) \right),$$

where m is a fixed positive integer, $u = (u_1, \dots, u_m) \in S^{m-1} \subset \mathbb{R}^m$, i.e., $\sum_{t=1}^m u_t^2 = 1$, and $k = (k_1, \dots, k_m), h = (h_1, \dots, h_m)$ are two fixed m -tuples of positive integers. Therefore, each element in M is identified by the unit vector u . We remark that the scaling factor $\frac{1}{4\sqrt{m}}$ in (3.3) ensures that the operator F is non-expansive.

We now move to a discrete setting by thinking of V as a set of 2D discrete images. Given $\varphi \in V$, we refer to a 2D discrete image as the restriction of φ to the subset $Y \subset X$, that is $\varphi|_Y : Y \rightarrow \mathbb{R}$. By abuse of notation, to improve the readability of the following lines we avoid the restriction symbol and denote by φ a generic image. More precisely, given the definitions (3.1), (3.2) and assuming Y is a discrete $n \times n$ grid in X , the generic element φ is a discrete 2D image whose (i, j) -pixel has value $\sum_{r=0}^{n-1} \sum_{s=0}^{n-1} a_i^j \chi \left(\frac{2\pi(i-r)}{n}, \frac{2\pi(j-s)}{n} \right)$. We refer to the point $\left(\frac{2\pi i}{n}, \frac{2\pi j}{n} \right) \in Y$ as (i, j) -pixel for all $i, j = 0, \dots, n-1$. Since Y is preserved by the finite group of isometries G , we assume periodic boundary conditions for each generic 2D image. Finally, we suppose the support of the density f_V is contained in the EMNIST dataset [6], therefore Φ is defined as a finite subset of the EMNIST of cardinality N .

Obviously, the cardinality of the space of GENEOS M makes unfeasible a detailed analysis of these operators. In particular, it can be desirable to restrict this space to a finite set of GENEOS which fairly approximates the whole manifold. In order to simplify the problem and to reduce the degrees of freedom we fix the vectors $k \in \mathbb{N}^m$ and $h \in \mathbb{N}^m$. Consequently, the manifold M has dimension $m-1$ and each element in M is now identified by the vector $u \in S^{m-1}$.

We now propose a procedure to extract a finite set $M_r \subset M$ of r GENEOS, where r is a positive integer. The computation of M_r must be guided by a metric which takes into account the observation that in a generic application not all the admissible signals in Φ can be considered to be equivalent. For example, in the English alphabet we can think the higher is the frequency of a letter, which is simply the amount of times it appears on average in written language, the higher is its importance. In this regard, we remark that the manifold M is endowed with the Riemannian structure defined by considering the inner product (2.7) which allows us to provide M with the norm $\|\cdot\|_{L^2}$. The latter takes into account the previous observation by weighting the contribution of each of the admissible signals in Φ through a probability density function.

The basic idea of our proposal is to cast the problem of defining M_r as a non-linear constrained optimization problem whose objective is L^2 -norm based and the constraints reflect the hypothesis $u \in S^{m-1}$. In particular, each GENEOS in M is identified through a vector $u \in S^{m-1}$, therefore seeking for r operators in M is equivalent to search for

r vectors in S^{m-1} . Then, we propose to read the task of defining M_r as the following constrained optimization problem:

$$(3.4) \quad \arg \min_{u_i \in S^{m-1}, \forall i=1 \dots r} \sum_{i=1}^r \sum_{j=i+1}^r \frac{1}{\|F_{u_i} - F_{u_j}\|_{L^2}^2}.$$

By definition (2.6), the optimization problem in (3.4) is equivalent to the following one:

$$(3.5) \quad \arg \min_{u_i \in S^{m-1}, \forall i=1 \dots r} \sum_{i=1}^r \sum_{j=i+1}^r \frac{1}{\sum_{s=1}^N f_V(\varphi_s) \|F_{u_i}(\varphi_s) - F_{u_j}(\varphi_s)\|_V^2}.$$

In our example we assume Φ is constituted by $N = 26$ different images representing the handwritten letters of the English alphabet sampled from the EMNIST dataset. We now consider φ_s as the s -th letter in Φ . Then, the probability density f_V on φ_s is defined as the frequency of the letter φ_s in the English alphabet [13]. In addition, we suppose that $f_V(\varphi) = 0$ for all $\varphi \in V \setminus \Phi$.

We stress that the objective in (3.5) is non-smooth and non-convex. A feasible (satisfying the constraints) local-minimizer is sought through the iterative *interior point* approach which is performed by using the built-in Matlab function, namely `fmincon`. The interior point approach used, at each iteration computes a feasible solution following the gradient direction of the objective [16] starting from a given feasible initial guess.

In our experiment, the iterates performed by `fmincon` are stopped when the optimality measure and the constraints tolerance are lower than 10^{-6} [16]. In addition, we have fixed a maximum number of objective function evaluations equals to 3000. We refer to M_r^k as the k -th iterate of the interior point method and, in particular, by M_r^0 we refer to the starting iteration of the optimization process. Moreover, we fix $m = 2$ and $r = 10, 20$.

In Fig. 1 we represent by a blue circular line the manifold M . In Fig 1 (a)-(c) we depict by red dots the starting iterates M_r^0 of the optimization process when $r = 10$ and $r = 20$, respectively, whereas in Fig 1 (b)-(d) the red dots represent the solutions M_{10} and M_{20} of (3.5) when M_{10}^0 and M_{20}^0 are chosen as initial guess, respectively. We observe that following the gradient flow of the objective in (3.5) the outcomes M_{10} and M_{20} present GENEOS (red points) better distributed and separated than the ones in M_{10}^0 and M_{20}^0 which have been randomly selected. Indeed, the objective has been defined to penalize mostly close (with respect to the given L^2 -metric) operators.

We now want to evaluate how much better than the random M_{10}^0 and M_{20}^0 the computed M_{10} and M_{20} approximate the manifold M . For this purpose, we consider the evaluation set $E \subset M$ made up of 100 GENEOS belonging to M randomly selected, then for every $F \in E$ we define $\eta_r(F)$ as the following ratio:

$$(3.6) \quad \eta_r(F) := \frac{\min_{F_i \in M_r} \|F - F_i\|_{L^2}^2}{\min_{F_j \in M_r^0} \|F - F_j\|_{L^2}^2}.$$

We remark that $\eta_r(F)$, according to the formula (3.6), measures the ratio of the minimum distances among the GENEOS F belonging to E and the ones in M_r and M_r^0 . Therefore, a value $\eta_r(F)$ less than one entails that $F \in E$ is closer in terms of L^2 -distance to a GENEOS in M_r , conversely $F \in E$ is closer to a GENEOS in M_r^0 . Since E is made up of 100 GENEOS we compute the mean values of $\{\eta_r(F)\}_{F \in E}$ choosing $r = 10$ and $r = 20$. In the first case the mean value is equal to 0.5833 while in the second case it is equal to 0.5442. This states that, in mean, the outcomes M_{10} and M_{20} better approximate the




FIGURE 1. Random initial guess and outcomes of the interior-point method solving the optimization problem (3.5) when $m = 2$. (A)-(B) starting feasible iterate and related solution given $r = 10$. (C)-(D) starting feasible iterate and related solution given $r = 20$.

whole manifold M if compared to their random counterparts M_{10}^0 and M_{20}^0 used as initial guesses in the iterative scheme solving (3.5) when $r = 10$ and $r = 20$, respectively.

4. DISCUSSION

In this paper we have shown how the space \mathcal{F} of all GENEOS can be endowed with a suitable geometrical structure, so allowing to prove some of the relevant properties of such a space. In particular, we have devised a metric that guarantees the compactness of \mathcal{F} under the main assumption that the set Φ of signals we are considering is compact. Such a metric differs from the one introduced in [5] and takes into account the probability of signals in Φ . Our compactness result requires new metrics on Φ and the equivariance group G and implies that \mathcal{F} can be approximated by a finite set of operators within an arbitrarily small error. Furthermore, we have shown how we can define a suitable Riemannian structure on finite dimensional submanifolds of the space of GENEOS and take advantage of this structure to build approximations of those manifolds via a gradient flow minimization. We believe that the availability of these results could be of use in reducing the complexity of selecting suitable operators in applications concerning topological data

analysis and deep learning. In the future, we would like to explore the possibility of extending our results to operators that are equivariant and non-expansive only almost everywhere on Φ , so improving the theoretical and practical use of our mathematical model. Another line of research could be reconstructing the theory in [5] by considering another metric on the space Φ instead of the one induced by the L^∞ -norm. In this way, new practical results could be compared with the ones in [5] to realize which metrics are efficient in different experiments.

ACKNOWLEDGEMENT

This research has been partially supported by INdAM-GNSAGA and INdAM-GNCS. The research of A.S. has been supported by the Institute for Research in Fundamental Sciences (IPM). The authors thank Elena Loli Piccolomini for her helpful advice. The authors would like to thank Nikolaos Chalmoukis for comments that greatly improved the manuscript. This paper is dedicated to the memory of Ivana Paoletti, Giorgio Mati and Don Piero Vannelli.

CONFLICT OF INTEREST

The authors declare that they have no conflict of interest.

AUTHORS' CONTRIBUTIONS

P.F. devised the project. P.F., N.Q. and A.S. developed the mathematical model. P.C. took care of the experiments. All authors read and approved the final manuscript. The authors of this paper have been listed in alphabetical order.

REFERENCES

1. Ralph Abraham, Jerrold E Marsden, and Tudor Ratiu, *Manifolds, tensor analysis, and applications*, vol. 75, Springer Science & Business Media, 2012.
2. Fabio Anselmi, Georgios Evangelopoulos, Lorenzo Rosasco, and Tomaso Poggio, *Symmetry-adapted representation learning*, *Pattern Recognition* **86** (2019), 201 – 208.
3. Fabio Anselmi, Lorenzo Rosasco, and Tomaso Poggio, *On invariance and selectivity in representation learning*, *Information and Inference: A Journal of the IMA* **5** (2016), no. 2, 134–158.
4. Yoshua Bengio, Aaron Courville, and Pascal Vincent, *Representation learning: A review and new perspectives*, *IEEE Trans. Pattern Anal. Mach. Intell.* **35** (2013), no. 8, 1798–1828.
5. Mattia G. Bergomi, Patrizio Frosini, Daniela Giorgi, and Nicola Quercioli, *Towards a topological-geometrical theory of group equivariant non-expansive operators for data analysis and machine learning*, *Nature Machine Intelligence* **1** (2019), 423–433.
6. Gregory Cohen, Saeed Afshar, Jonathan Tapson, and Andre Van Schaik, *Emnist: Extending mnist to handwritten letters*, 2017 International Joint Conference on Neural Networks (IJCNN), IEEE, 2017, pp. 2921–2926.
7. Taco Cohen and Max Welling, *Group equivariant convolutional networks*, International conference on machine learning, 2016, pp. 2990–2999.
8. Patrizio Frosini, *Does intelligence imply contradiction?*, *Cognitive Systems Research* **10** (2009), no. 4, 297–315.
9. ———, *Towards an observer-oriented theory of shape comparison*, Eurographics Workshop on 3D Object Retrieval (A. Ferreira, A. Giachetti, and D. Giorgi, eds.), The Eurographics Association, 2016.
10. Patrizio Frosini and Grzegorz Jabłoński, *Combining persistent homology and invariance groups for shape comparison*, *Discrete & Computational Geometry* **55** (2016), no. 2, 373–409.
11. Steven A. Gaal, *Point set topology*, Pure and Applied Mathematics, Vol. XVI, Academic Press, New York-London, 1964.
12. Wilhelm P.A. Klingenberg, *Riemannian geometry*, De Gruyter, Berlin, Boston, 2011.
13. Robert Edward Lewand, *Cryptological mathematics*, vol. 16, American Mathematical Soc., 2000.

14. Stéphane Mallat, *Group invariant scattering*, Communications on Pure and Applied Mathematics **65** (2012), no. 10, 1331–1398.
15. ———, *Understanding deep convolutional networks*, Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences **374** (2016), no. 2065, 20150203.
16. Jorge Nocedal and Stephen Wright, *Numerical optimization*, Springer Science & Business Media, 2006.
17. Daniel E Worrall, Stephan J Garbin, Daniyar Turmukhambetov, and Gabriel J Brostow, *Harmonic networks: Deep translation and rotation equivariance*, Proc. IEEE Conf. on Computer Vision and Pattern Recognition (CVPR), vol. 2, 2017.
18. Chiyuan Zhang, Stephen Voinea, Georgios Evangelopoulos, Lorenzo Rosasco, and Tomaso Poggio, *Discriminative template learning in group-convolutional networks for invariant speech representations*, INTERSPEECH-2015 (Dresden, Germany), International Speech Communication Association (ISCA), International Speech Communication Association (ISCA), 09/2015 2015.

DIPARTIMENTO DI MATEMATICA, UNIVERSITÀ DI BOLOGNA, ITALIA
Email address: pasquale.cascarano2@unibo.it

DIPARTIMENTO DI MATEMATICA, UNIVERSITÀ DI BOLOGNA, ITALIA
Email address: patrizio.frosini@unibo.it

DIPARTIMENTO DI MATEMATICA, UNIVERSITÀ DI BOLOGNA, ITALIA
Email address: nicola.quercioli2@unibo.it

SCHOOL OF MATHEMATICS, INSTITUTE FOR RESEARCH IN FUNDAMENTAL SCIENCES (IPM), IRAN
Email address: amir.saki.math@gmail.com