

A multivariate normal approximation for the Dirichlet density and some applications

Frédéric Ouimet^{a,1,*}

^aMcGill University, Montreal, Canada.

Abstract

In this short note, we prove an asymptotic expansion for the ratio of the Dirichlet density to the multivariate normal density with the same mean and covariance matrix. The expansion is then used to derive an upper bound on the total variation between the corresponding probability measures and rederive the asymptotic variance of the Dirichlet kernel estimators introduced by [Aitchison & Lauder \(1985\)](#) and studied theoretically in [Ouimet \(2020\)](#). Another potential application related to the asymptotic equivalence between the Gaussian variance regression problem and the Gaussian white noise problem is briefly mentioned but left open for future research.

Keywords: Dirichlet distribution, asymptotic statistics, expansion, normal approximation, multivariate normal, total variation, asymptotic variance, nonparametric statistics, smoothing, density estimation

2020 MSC: Primary: 62E20 Secondary: 62H10, 62H12, 62B15, 62G05, 62G07

1. Introduction

For any $d \in \mathbb{N}$ and $\mathbf{v} \in \mathbb{R}^d$, let $\|\mathbf{v}\|_1 := \sum_{i=1}^d |v_i|$ and define the d -dimensional simplex as

$$\mathcal{S}_d := \{\mathbf{v} \in [0, 1]^d : \|\mathbf{v}\|_1 \leq 1\}. \quad (1.1)$$

Given the parameters $N \in \mathbb{N}$ and $(\boldsymbol{\alpha}, \beta) \in (0, \infty)^{d+1}$, the Dirichlet($N\boldsymbol{\alpha}, N\beta$) density function is defined by

$$K_{N,\boldsymbol{\alpha},\beta}(\mathbf{x}) = \frac{\Gamma(N\|\boldsymbol{\alpha}\|_1 + N\beta)}{\Gamma(N\beta) \prod_{i=1}^d \Gamma(N\alpha_i)} \cdot (1 - \|\mathbf{x}\|_1)^{N\beta-1} \prod_{i=1}^d x_i^{N\alpha_i-1}, \quad \mathbf{x} \in \mathcal{S}_d. \quad (1.2)$$

The covariance matrix of the Dirichlet distribution is well-known to be $(N\|\boldsymbol{\alpha}\|_1 + N\beta + 1)^{-1} \Sigma_{\mathbf{r}}$, where

$$\Sigma_{\mathbf{r}} := \text{diag}(\mathbf{r}) - \mathbf{r}\mathbf{r}^\top \quad \text{and} \quad \mathbf{r} := \frac{\boldsymbol{\alpha}}{\|\boldsymbol{\alpha}\|_1 + \beta}, \quad (1.3)$$

*Corresponding author

Email address: frederic.ouimet2@mail.mcgill.ca (Frédéric Ouimet)

¹F. O. is supported by a postdoctoral fellowship from the NSERC (PDF) and the FRQNT (B3X supplement).

see e.g. (Ng *et al.*, 2011, p.39). By adapting Theorem 1 and Equation (21) in Tanabe & Sagae (1992), we also know that

$$\det(\Sigma_{\mathbf{r}}) = r_1 r_2 \dots r_{d+1} \quad \text{and} \quad (\Sigma_{\mathbf{r}}^{-1})_{ij} = \frac{1}{r_i} \mathbb{1}_{\{i=j\}} + \frac{1}{r_{d+1}}, \quad 1 \leq i, j \leq d, \quad (1.4)$$

where $r_{d+1} := 1 - \|\mathbf{r}\|_1 = \frac{\beta}{\|\boldsymbol{\alpha}\|_1 + \beta}$.

The first goal of the paper (Theorem 2.1) is to establish an asymptotic expansion for the ratio of the Dirichlet density (1.2) to the multivariate normal density with the same mean and covariances, namely:

$$\frac{\exp\left(-\frac{1}{2} \boldsymbol{\delta}_{\mathbf{x}}^{\top} \Sigma_{\mathbf{r}}^{-1} \boldsymbol{\delta}_{\mathbf{x}}\right)}{\sqrt{(2\pi)^d (N \|\boldsymbol{\alpha}\|_1 + N\beta + 1)^{-d} r_1 r_2 \dots r_{d+1}}}, \quad \mathbf{x} \in \mathbb{R}^d, \quad (1.5)$$

where

$$\boldsymbol{\delta}_{\mathbf{x}} := (\delta_{1,x_1}, \delta_{2,x_2}, \dots, \delta_{d,x_d}) \quad \text{and} \quad \delta_{i,x_i} := \frac{x_i - r_i}{(N \|\boldsymbol{\alpha}\|_1 + N\beta + 1)^{-1/2}}. \quad (1.6)$$

The second goal of the paper is to apply the asymptotic expansion to derive an upper bound on the total variation between the probability measures on \mathbb{R}^d induced by (1.2) and (1.5), and to rederive the asymptotic variance of the Dirichlet kernel estimators in the context of density estimation for compositional data. These two applications are treated in Section 3.1 and Section 3.2, respectively. There could be many other potential applications, see for example the excellent survey by Mason & Zhou (2012) on quantile coupling inequalities.

In fact, the original motivation for the present paper was the PhD thesis of Huibin Zhou (Zhou, 2004), in which a multi-resolution coupling methodology between beta and normal random variables is applied to prove the asymptotic equivalence between the Gaussian variance regression problem and the Gaussian white noise problem under Besov smoothness constraints (see also the related works Brown & Low (1996); Brown *et al.* (2002); Rohde (2004); Carter (2006, 2007); Cai & Zhou (2009); Golubev *et al.* (2010)). We believe that the main result here (Theorem 2.1) could lead to a significant simplification of the proof of (Zhou, 2004, Theorem 2.1), in analogy with the removal of the inductive part of the proof for the Le Cam distance upper bound between multinomial and multivariate normal experiments from (Carter, 2002, Theorem 1), shown in Ouimet (2021). This point is left open for future research.

Notations. Throughout the paper, the notation $u = \mathcal{O}(v)$ means that $\limsup_{N \rightarrow \infty} |u/v| < C$, where $C > 0$ is a universal constant. Whenever C might depend on some parameters, we add subscripts (for example, $u = \mathcal{O}_{\boldsymbol{\alpha}, \beta}(v)$). Also, we write

$$x_{d+1} := 1 - \|\mathbf{x}\|_1, \quad \alpha_{d+1} := \beta, \quad \text{and} \quad \varepsilon_N := \frac{1}{N(\|\boldsymbol{\alpha}\|_1 + \beta)}. \quad (1.7)$$

In particular, the definition of x_{d+1} and r_{d+1} implies that $\delta_{d+1, x_{d+1}} = -\sum_{i=1}^d \delta_{i, x_i}$.

2. Main result

First, we prove an asymptotic expansion for the ratio of the Dirichlet density to the multivariate normal density with the same mean and covariances.

Theorem 2.1. *Pick any $\eta \in (0, 1)$ and let*

$$B_\eta := \left\{ \mathbf{x} \in \mathcal{S}_d : |\delta_{i,x_i}| \leq \eta N^{1/6}, \text{ for all } i \in \{1, 2, \dots, d+1\} \right\} \quad (2.1)$$

denote the bulk of the Dirichlet distribution. Then, uniformly for $\mathbf{x} \in B_\eta$, we have

$$\begin{aligned} \log \left(\frac{K_{N,\alpha,\beta}(\mathbf{x})}{\frac{\exp\left(-\frac{1}{2}\boldsymbol{\delta}_x^\top \Sigma_r^{-1} \boldsymbol{\delta}_x\right)}{\sqrt{(2\pi)^d (1+\varepsilon_N^{-1})^{-d} r_1 r_2 \dots r_{d+1}}}} \right) &= \varepsilon_N^{1/2} \cdot \left\{ -\sum_{i=1}^{d+1} \left(\frac{\delta_{i,x_i}}{r_i}\right) + \frac{1}{3} \sum_{i=1}^{d+1} \delta_{i,x_i} \left(\frac{\delta_{i,x_i}}{r_i}\right)^2 \right\} \\ &+ \varepsilon_N \cdot \left\{ \frac{1}{2} \sum_{i=1}^{d+1} \left(\frac{\delta_{i,x_i}}{r_i}\right)^2 - \frac{1}{4} \sum_{i=1}^{d+1} \delta_{i,x_i} \left(\frac{\delta_{i,x_i}}{r_i}\right)^3 \right\} \\ &+ \mathcal{O}_{\alpha,\beta,\eta} \left(\frac{(1 + \|\boldsymbol{\delta}_x\|_1)^5}{N^{3/2}} \right), \quad \text{as } N \rightarrow \infty. \end{aligned} \quad (2.2)$$

Proof of Theorem 2.1. Using Stirling's formula,

$$\log \Gamma(z) = \frac{1}{2} \log(2\pi) + \left(z - \frac{1}{2}\right) \log z - z + \frac{1}{12z} + \mathcal{O}(z^{-3}), \quad \text{as } z \rightarrow \infty, \quad (2.3)$$

see e.g. (Abramowitz & Stegun, 1964, p.257), and taking the logarithm in (1.2), we obtain

$$\begin{aligned} \log K_{N,\alpha,\beta}(\mathbf{x}) &= \log \Gamma(N\|\boldsymbol{\alpha}\|_1 + N\beta) - \sum_{i=1}^{d+1} \log \Gamma(N\alpha_i) + \sum_{i=1}^{d+1} (N\alpha_i - 1) \log x_i \\ &= -\frac{d}{2} \log(2\pi) - \frac{d}{2} \log \varepsilon_N - \frac{1}{2} \sum_{i=1}^{d+1} \log r_i + \sum_{i=1}^{d+1} (N\alpha_i - 1) \log \left(\frac{x_i}{r_i}\right) \\ &+ \frac{\left\{1 - \sum_{i=1}^{d+1} r_i^{-1}\right\}}{12N(\|\boldsymbol{\alpha}\|_1 + \beta)} + \mathcal{O}_{\alpha,\beta}(N^{-3}). \end{aligned} \quad (2.4)$$

By writing $\frac{x_i}{r_i} = 1 + \frac{\delta_{i,x_i}}{r_i} (1 + \varepsilon_N^{-1})^{-1/2}$ in (2.4), we deduce

$$\begin{aligned} \log K_{N,\alpha,\beta}(\mathbf{x}) &= -\log \sqrt{(2\pi)^d (1 + \varepsilon_N^{-1})^{-d} r_1 r_2 \dots r_{d+1}} - \frac{d}{2} \log(1 + \varepsilon_N) \\ &+ \sum_{i=1}^{d+1} (\varepsilon_N^{-1} r_i - 1) \log \left(1 + \frac{\delta_{i,x_i}}{r_i} (1 + \varepsilon_N^{-1})^{-1/2}\right) \\ &+ \varepsilon_N \cdot \frac{1}{12} \left\{1 - \sum_{i=1}^{d+1} r_i^{-1}\right\} + \mathcal{O}_{\alpha,\beta}(N^{-3}). \end{aligned} \quad (2.5)$$

By applying the Taylor expansion

$$\log(1 + y) = y - \frac{y^2}{2} + \frac{y^3}{3} - \frac{y^4}{4} + \mathcal{O}_\eta(y^5), \quad \text{for } |y| \leq \eta, \quad (2.6)$$

and noticing that $\delta_{d+1,x_{d+1}} = -\sum_{i=1}^d \delta_{i,x_i}$, we have

$$\begin{aligned}
\log K_{N,\alpha,\beta}(\mathbf{x}) &= -\log \sqrt{(2\pi)^d (1 + \varepsilon_N^{-1})^{-d} r_1 r_2 \dots r_{d+1}} - \frac{d}{2} \{ \varepsilon_N + \mathcal{O}_{\alpha,\beta}(N^{-2}) \} \\
&\quad - (1 + \varepsilon_N^{-1})^{-1/2} \sum_{i=1}^{d+1} \frac{\delta_{i,x_i}}{r_i} + \frac{(1 + \varepsilon_N^{-1})^{-1}}{2} \sum_{i=1}^{d+1} \frac{\delta_{i,x_i}^2}{r_i^2} \\
&\quad - (1 + \varepsilon_N)^{-1} \sum_{i=1}^d \frac{\delta_{i,x_i}^2}{2} \left\{ \frac{1}{r_i} - \frac{2}{3} \cdot \frac{\delta_{i,x_i}}{r_i^2} (1 + \varepsilon_N^{-1})^{-1/2} \right. \\
&\quad \left. + \frac{1}{2} \cdot \frac{\delta_{i,x_i}^2}{r_i^3} (1 + \varepsilon_N^{-1})^{-1} + \mathcal{O}_{\alpha,\beta,\eta} \left(\frac{(1 + |\delta_{i,x_i}|^3)}{N^{3/2}} \right) \right\} \quad (2.7) \\
&\quad - (1 + \varepsilon_N)^{-1} \sum_{i,j=1}^d \frac{\delta_{i,x_i} \delta_{j,x_j}}{2} \left\{ \frac{1}{r_{d+1}} + \frac{2}{3} \cdot \sum_{\ell=1}^d \frac{\delta_{\ell,x_\ell}}{r_{d+1}^2} (1 + \varepsilon_N^{-1})^{-1/2} \right. \\
&\quad \left. + \frac{1}{2} \cdot \sum_{\ell,m=1}^d \frac{\delta_{\ell,x_\ell} \delta_{m,x_m}}{r_{d+1}^3} (1 + \varepsilon_N^{-1})^{-1} \right. \\
&\quad \left. + \mathcal{O}_{\alpha,\beta,\eta} \left(\frac{(1 + \|\delta_{\mathbf{x}}\|_1^3)}{N^{3/2}} \right) \right\} \\
&\quad + \varepsilon_N \cdot \frac{1}{12} \left\{ 1 - \sum_{i=1}^{d+1} r_i^{-1} \right\} + \mathcal{O}_{\alpha,\beta}(N^{-3}).
\end{aligned}$$

We can rewrite this as

$$\begin{aligned}
\log K_{N,\alpha,\beta}(\mathbf{x}) &= -\log \sqrt{(2\pi)^d (1 + \varepsilon_N^{-1})^{-d} r_1 r_2 \dots r_{d+1}} \\
&\quad - \varepsilon_N^{1/2} \sum_{i=1}^{d+1} \frac{\delta_{i,x_i}}{r_i} + \frac{\varepsilon_N}{2} \sum_{i=1}^{d+1} \frac{\delta_{i,x_i}^2}{r_i^2} \\
&\quad - (1 + \varepsilon_N)^{-1} \sum_{i,j=1}^d \frac{\delta_{i,x_i} \delta_{j,x_j}}{2} \{ (\Sigma_{\mathbf{r}}^{-1})_{ij} + S_{N,ij} \} \\
&\quad + \varepsilon_N \cdot \left[-\frac{d}{2} + \frac{1}{12} \left\{ 1 - \sum_{i=1}^{d+1} r_i^{-1} \right\} \right] + \mathcal{O}_{\alpha,\beta,\eta} \left(\frac{(1 + \|\delta_{\mathbf{x}}\|_1)^2}{N^{3/2}} \right). \quad (2.8)
\end{aligned}$$

where the $d \times d$ matrices $\Sigma_{\mathbf{r}}^{-1}$ and S_N have the (i, j) components:

$$(\Sigma_{\mathbf{r}}^{-1})_{ij} := \frac{1}{r_i} \mathbf{1}_{\{i=j\}} + \frac{1}{r_{d+1}}, \quad (2.9)$$

$$\begin{aligned}
S_{N,ij} &:= \frac{2}{3} \sum_{\ell=1}^d \frac{\delta_{\ell,x_\ell}}{(1 + \varepsilon_N^{-1})^{1/2}} \left\{ \frac{-1}{r_i^2} \mathbf{1}_{\{i=j=\ell\}} + \frac{1}{r_{d+1}^2} \right\} \\
&\quad + \frac{1}{2} \sum_{\ell,m=1}^d \frac{\delta_{\ell,x_\ell} \delta_{m,x_m}}{(1 + \varepsilon_N^{-1})} \left\{ \frac{1}{r_i^3} \mathbf{1}_{\{i=j=\ell=m\}} + \frac{1}{r_{d+1}^3} \right\} + \mathcal{O}_{\alpha,\beta,\eta} \left(\frac{(1 + \|\delta_{\mathbf{x}}\|_1)^3}{N^{3/2}} \right). \quad (2.10)
\end{aligned}$$

After expanding (2.8) and rearranging some terms, we get (2.2). \square

3. Applications

In this section, we present two applications of Theorem 2.1. We find an upper bound on the total variation between Dirichlet and multivariate normal distributions (Section 3.1) and we present an alternative proof for the asymptotic variance of Dirichlet kernel estimators found in Theorem 4.2 of Ouimet (2020) (Section 3.2).

3.1. Total variation bound between Dirichlet and multivariate normal distributions

Theorem 3.1. Let $\mathbb{P}_{\alpha,\beta}$ be the law of the Dirichlet($N\alpha, N\beta$) distribution, and let $\mathbb{Q}_{\alpha,\beta}$ be the law of the Normal $_d(\mathbf{r}, (1 + \varepsilon_N^{-1})^{-1} \Sigma_{\mathbf{r}})$ distribution, where recall $\Sigma_{\mathbf{r}} := \text{diag}(\mathbf{r}) - \mathbf{r}\mathbf{r}^\top$. Then, for all $(\alpha, \beta) \in (0, \infty)^{d+1}$, we have

$$\|\mathbb{P}_{\alpha,\beta} - \mathbb{Q}_{\alpha,\beta}\| = \mathcal{O}\left(\varepsilon_N^{1/2} \cdot d \sqrt{\frac{\max_{1 \leq i \leq d+1} r_i}{\min_{1 \leq i \leq d+1} r_i}}\right), \quad \text{as } N \rightarrow \infty, \quad (3.1)$$

where $\|\cdot\|$ denotes the total variation norm.

Proof. Let $\mathbf{X} \sim \mathbb{P}_{\alpha,\beta}$. By the comparison of the total variation norm with the Hellinger distance on page 726 of [Carter \(2002\)](#), we already know that

$$\|\mathbb{P}_{\alpha,\beta} - \mathbb{Q}_{\alpha,\beta}\| \leq \sqrt{2\mathbb{P}(\mathbf{X} \in B_{1/2}^c) + \mathbb{E}\left[\log\left(\frac{d\mathbb{P}_{\alpha,\beta}}{d\mathbb{Q}_{\alpha,\beta}}(\mathbf{X})\right) \mathbf{1}_{\{\mathbf{X} \in B_{1/2}\}}\right]}. \quad (3.2)$$

Then, by applying a union bound followed by large deviation bound for the beta distribution (see e.g. Theorem 2.1 of [Marchal & Arbel \(2017\)](#)), we get, for N large enough,

$$\mathbb{P}(\mathbf{X} \in B_{1/2}^c) \leq \sum_{i=1}^{d+1} \mathbb{P}\left(|\delta_{X_i}| > \frac{1}{2}N^{1/6}\right) \leq (d+1) \cdot 2 \exp\left(-\frac{1}{2}N^{1/3}\right). \quad (3.3)$$

By Theorem 2.1,

$$\begin{aligned} & \mathbb{E}\left[\log\left(\frac{d\mathbb{P}_{\alpha,\beta}}{d\mathbb{Q}_{\alpha,\beta}}(\mathbf{X})\right) \mathbf{1}_{\{\mathbf{X} \in B_{1/2}\}}\right] \\ &= \varepsilon_N^{1/2} \cdot \mathbb{E}\left[\left\{\begin{aligned} & -\sum_{i=1}^d \delta_{i,x_i} \left(\frac{1}{r_i} - \frac{1}{r_{d+1}}\right) \\ & + \frac{1}{3} \sum_{i,j,\ell=1}^d \delta_{i,x_i} \delta_{j,x_j} \delta_{\ell,x_\ell} \left(\frac{1}{r_i^2} \mathbf{1}_{\{i=j=\ell\}} - \frac{1}{r_{d+1}^2}\right) \end{aligned}\right\} \mathbf{1}_{\{\mathbf{X} \in B_{1/2}\}}\right] \\ &+ \varepsilon_N \cdot \mathcal{O}\left(\left|\sum_{i=1}^{d+1} \frac{\mathbb{E}[(X_i - r_i)^2]}{\varepsilon_N r_i^2}\right| + \left|\sum_{i=1}^{d+1} \frac{\mathbb{E}[(X_i - r_i)^4]}{\varepsilon_N^2 r_i^3}\right| + d + \sum_{i=1}^{d+1} r_i^{-1}\right) \\ &+ \mathcal{O}_{\alpha,\beta}(N^{-3/2}). \end{aligned} \quad (3.4)$$

By Lemma A.1, the second to last $\mathcal{O}(\cdot)$ term above is

$$= \mathcal{O}\left(\sum_{i=1}^{d+1} r_i^{-1}\right) = \mathcal{O}\left(\frac{d}{\min_{1 \leq i \leq d+1} r_i}\right) = \mathcal{O}\left(d^2 \frac{\max_{1 \leq i \leq d+1} r_i}{\min_{1 \leq i \leq d+1} r_i}\right). \quad (3.5)$$

(The last equality follows from $\frac{1}{d+1} \leq \max_{1 \leq i \leq d+1} r_i$, which itself is consequence of the fact that $r_i \geq 0$ and $\sum_{i=1}^{d+1} r_i = 1$.) By putting (3.5) in (3.4) and using Lemma A.2, we get

$$(3.4) = \varepsilon_N^{1/2} \cdot \left\{ \begin{aligned} & \varepsilon_N^2 (1 + \varepsilon_N^{-1})^{3/2} \cdot \sum_{i,j,\ell=1}^d \frac{\left\{ \begin{aligned} & 4r_i r_j r_\ell - 2r_i r_\ell \mathbf{1}_{\{i=j\}} - 2r_j r_\ell \mathbf{1}_{\{i=\ell\}} \\ & - 2r_i r_j \mathbf{1}_{\{j=\ell\}} + 2r_i \mathbf{1}_{\{i=j=\ell\}} \end{aligned} \right\}}{(1 + \varepsilon_N)(1 + 2\varepsilon_N)} \\ & \cdot \left\{ \frac{1}{r_i^2} \mathbf{1}_{\{i=j=\ell\}} - \frac{1}{r_{d+1}^2} \right\} \\ & + \mathcal{O}\left(\frac{d(\mathbb{P}(\mathbf{X} \in B_{1/2}^c))^{1/2}}{\min_{1 \leq i \leq d+1} r_i}\right) + \mathcal{O}\left(\frac{d^2 (\mathbb{P}(\mathbf{X} \in B_{1/2}^c))^{1/4}}{(\min_{1 \leq i \leq d+1} r_i)^2}\right) \end{aligned} \right\}$$

$$\begin{aligned}
& + \varepsilon_N \cdot \mathcal{O}\left(d^2 \frac{\max_{1 \leq i \leq d+1} r_i}{\min_{1 \leq i \leq d+1} r_i}\right) + \mathcal{O}_{\alpha, \beta}(N^{-3/2}) \\
& = \mathcal{O}\left(\varepsilon_N^{1/2} \cdot \frac{d^2 (\mathbb{P}(\mathbf{X} \in B_{1/2}^c))^{1/4}}{(\min_{1 \leq i \leq d+1} r_i)^2}\right) + \mathcal{O}\left(\varepsilon_N \cdot d^2 \frac{\max_{1 \leq i \leq d+1} r_i}{\min_{1 \leq i \leq d+1} r_i}\right). \tag{3.6}
\end{aligned}$$

Now, putting (3.3) and (3.6) together in (3.2) gives the conclusion. \square

3.2. Asymptotic variance of Dirichlet kernel estimators

Assume that we have a sequence of observations $\mathbf{X}_1, \mathbf{X}_2, \dots, \mathbf{X}_n$ that are independent and F distributed (F is unknown), with density f supported on the d -dimensional simplex \mathcal{S}_d . Then, for a given bandwidth parameter $b > 0$, let

$$\hat{f}_{n,b}(\mathbf{s}) := \frac{1}{n} \sum_{i=1}^n K_{1/b, \mathbf{s}+b, 1-\|\mathbf{s}\|_1+b}(\mathbf{X}_i), \quad \mathbf{s} \in \mathcal{S}_d, \tag{3.7}$$

be the *Dirichlet kernel estimator* for the density function f . This estimator was introduced by Aitchison & Lauder (1985) as a nonparametric method of density estimation for compositional data and its asymptotic properties were studied theoretically for the first time in Ouimet (2020). For a detailed overview of the literature on asymmetric kernel estimators, we refer the reader to Hirukawa (2018) or Section 2 in Ouimet (2020).

One interesting application of the normal approximation in Theorem 2.1 is the derivation of the asymptotic variance of $\hat{f}_{n,b}$ at each point \mathbf{s} in the interior of the simplex.

Theorem 3.2. *Assume that f is Lipschitz continuous and let $\mathbf{s} \in \text{Int}(\mathcal{S}_d)$, then*

$$\text{Var}(\hat{f}_{n,b}(\mathbf{s})) = \frac{n^{-1} b^{-d/2} f(\mathbf{s})}{\sqrt{(4\pi)^d s_1 s_2 \dots s_{d+1}}} + o_{d,\mathbf{s}}(1), \quad \text{as } n \rightarrow \infty. \tag{3.8}$$

From this result, other asymptotic expressions can be derived such as the mean squared error and the mean integrated squared error and we can also optimize the bandwidth parameter b with respect to them, see e.g. Corollary 4.3 and Theorem 4.4 in Ouimet (2020).

Proof of Theorem 3.2. Straightforward computations show that

$$\begin{aligned}
\text{Var}(\hat{f}_{n,b}(\mathbf{s})) &= n^{-1} \mathbb{E}[K_{1/b, \mathbf{s}+b, 1-\|\mathbf{s}\|_1+b}(\mathbf{X})^2] - n^{-1} (\mathbb{E}[K_{1/b, \mathbf{s}+b, 1-\|\mathbf{s}\|_1+b}(\mathbf{X})])^2 \\
&= n^{-1} \mathbb{E}[K_{1/b, \mathbf{s}+b, 1-\|\mathbf{s}\|_1+b}(\mathbf{X})^2] - \mathcal{O}(n^{-1}), \tag{3.9}
\end{aligned}$$

where

$$\begin{aligned}
& \mathbb{E}[K_{1/b, \mathbf{s}+b, 1-\|\mathbf{s}\|_1+b}(\mathbf{X})^2] \\
& \stackrel{(2.2)}{=} \int_{\mathcal{S}_d} \left(\frac{\exp\left(-\frac{1}{2} \boldsymbol{\delta}_x^\top \boldsymbol{\Sigma}_r^{-1} \boldsymbol{\delta}_x\right)}{\sqrt{(2\pi)^d (1 + \varepsilon_N^{-1})^{-d} r_1 r_2 \dots r_{d+1}}}\right)^2 f(\mathbf{x}) d\mathbf{x} + o_{d,\mathbf{s}}(1) \\
& = \frac{2^{-d/2} f(\mathbf{s})}{\sqrt{(2\pi)^d b^d r_1 r_2 \dots r_{d+1}}} \underbrace{\int_{\mathcal{S}_d} \frac{\exp\left(-\frac{1}{2} \boldsymbol{\delta}_x^\top (\frac{1}{2} \boldsymbol{\Sigma}_r)^{-1} \boldsymbol{\delta}_x\right)}{\sqrt{(2\pi)^d 2^{-d} (1 + \varepsilon_N^{-1})^{-d} r_1 r_2 \dots r_{d+1}}} d\mathbf{x}}_{= 1 + o_d(1)} + o_{d,\mathbf{s}}(1). \tag{3.10}
\end{aligned}$$

Since $r_i = (s_i + b)/(1 + b(d + 1)) = s_i + o_{d,\mathbf{s}}(1)$ for all $1 \leq i \leq d + 1$, plugging the estimate (3.10) in (3.9) gives us the conclusion. \square

A. Moments of the Dirichlet distribution

Below, we compute some of the central moments (up to four) of the Dirichlet distribution. The lemma is used to estimate the $\asymp \varepsilon_N$ errors in (3.4) of the proof of Theorem 3.1, and also as a preliminary result for the proof of Lemma A.2.

Lemma A.1. *Let $N \in \mathbb{N}$ and let $(\boldsymbol{\alpha}, \beta) \in (0, \infty)^{d+1}$ be given. If $\mathbf{X} = (X_1, X_2, \dots, X_d) \sim \text{Dirichlet}(N\boldsymbol{\alpha}, N\beta)$ according to (1.2), then, for all $i, j, \ell \in \{1, 2, \dots, d\}$,*

$$\mathbb{E}[(X_i - r_i)(X_j - r_j)] = \varepsilon_N r_i \cdot \frac{(\mathbb{1}_{\{i=j\}} - r_j)}{(1 + \varepsilon_N)}, \quad (\text{A.1})$$

$$\mathbb{E}[(X_i - r_i)(X_j - r_j)(X_\ell - r_\ell)] = \varepsilon_N^2 \cdot \frac{\left\{ \begin{array}{l} 4r_i r_j r_\ell - 2r_i r_\ell \mathbb{1}_{\{i=j\}} - 2r_j r_\ell \mathbb{1}_{\{i=\ell\}} \\ - 2r_i r_j \mathbb{1}_{\{j=\ell\}} + 2r_i \mathbb{1}_{\{i=j=\ell\}} \end{array} \right\}}{(1 + \varepsilon_N)(1 + 2\varepsilon_N)}, \quad (\text{A.2})$$

$$\mathbb{E}[(X_i - r_i)^4] = \varepsilon_N^2 r_i^2 \cdot 3(1 - r_i)^2 + \mathcal{O}_{\boldsymbol{\alpha}, \beta}(N^{-3}), \quad (\text{A.3})$$

where recall $\varepsilon_N := 1/(N\|\boldsymbol{\alpha}\|_1 + N\beta)$ and $r_i := \mathbb{E}[X_i] = \frac{\alpha_i}{\|\boldsymbol{\alpha}\|_1 + \beta}$ for all $i \in \{1, 2, \dots, d\}$.

Proof. Equation (A.1) can be found in (Ng *et al.*, 2011, p.39). Since

$$\begin{aligned} \mathbb{E}[X_i X_j] &= \begin{cases} \frac{\Gamma(N\alpha_i+1)\Gamma(N\alpha_j+1)}{\Gamma(N\alpha_i)\Gamma(N\alpha_j)} \cdot \frac{\Gamma(N\|\boldsymbol{\alpha}\|_1+N\beta)}{\Gamma(N\|\boldsymbol{\alpha}\|_1+N\beta+2)}, & \text{if } i \neq j \\ \frac{\Gamma(N\alpha_i+2)}{\Gamma(N\alpha_i)} \cdot \frac{\Gamma(N\|\boldsymbol{\alpha}\|_1+N\beta)}{\Gamma(N\|\boldsymbol{\alpha}\|_1+N\beta+2)}, & \text{if } i = j \end{cases} \\ &= \frac{\alpha_i(\alpha_j + \mathbb{1}_{\{i=j\}})N^{-1}}{(\|\boldsymbol{\alpha}\|_1 + \beta)(\|\boldsymbol{\alpha}\|_1 + \beta + N^{-1})}, \end{aligned} \quad (\text{A.4})$$

and

$$\begin{aligned} \mathbb{E}[X_i X_j X_\ell] &= \left\{ \begin{array}{ll} \frac{\Gamma(N\alpha_i+1)\Gamma(N\alpha_j+1)\Gamma(N\alpha_\ell+1)}{\Gamma(N\alpha_i)\Gamma(N\alpha_j)\Gamma(N\alpha_\ell)} \cdot \frac{\Gamma(N\|\boldsymbol{\alpha}\|_1+N\beta)}{\Gamma(N\|\boldsymbol{\alpha}\|_1+N\beta+3)}, & \text{if } i \neq j \neq \ell \neq i \\ \frac{\Gamma(N\alpha_i+2)\Gamma(N\alpha_\ell+1)}{\Gamma(N\alpha_i)\Gamma(N\alpha_\ell)} \cdot \frac{\Gamma(N\|\boldsymbol{\alpha}\|_1+N\beta)}{\Gamma(N\|\boldsymbol{\alpha}\|_1+N\beta+3)}, & \text{if } i = j \neq \ell \\ \frac{\Gamma(N\alpha_i+2)\Gamma(N\alpha_j+1)}{\Gamma(N\alpha_i)\Gamma(N\alpha_j)} \cdot \frac{\Gamma(N\|\boldsymbol{\alpha}\|_1+N\beta)}{\Gamma(N\|\boldsymbol{\alpha}\|_1+N\beta+3)}, & \text{if } i = \ell \neq j \\ \frac{\Gamma(N\alpha_j+2)\Gamma(N\alpha_i+1)}{\Gamma(N\alpha_j)\Gamma(N\alpha_i)} \cdot \frac{\Gamma(N\|\boldsymbol{\alpha}\|_1+N\beta)}{\Gamma(N\|\boldsymbol{\alpha}\|_1+N\beta+3)}, & \text{if } j = \ell \neq i \\ \frac{\Gamma(N\alpha_i+3)}{\Gamma(N\alpha_i)} \cdot \frac{\Gamma(N\|\boldsymbol{\alpha}\|_1+N\beta)}{\Gamma(N\|\boldsymbol{\alpha}\|_1+N\beta+3)}, & \text{if } i = j = \ell \end{array} \right\} \\ &= \frac{(\alpha_i + \mathbb{1}_{\{i=\ell \neq j\}})N^{-1})(\alpha_j + \mathbb{1}_{\{i=j\}})N^{-1})(\alpha_\ell + \mathbb{1}_{\{j=\ell\}}N^{-1} + \mathbb{1}_{\{i=j=\ell\}}N^{-1})}{(\|\boldsymbol{\alpha}\|_1 + \beta)(\|\boldsymbol{\alpha}\|_1 + \beta + N^{-1})(\|\boldsymbol{\alpha}\|_1 + \beta + 2N^{-1})}, \end{aligned} \quad (\text{A.5})$$

we have

$$\begin{aligned} &\mathbb{E}[(X_i - r_i)(X_j - r_j)(X_\ell - r_\ell)] \\ &= \mathbb{E}[X_i X_j X_\ell] - r_\ell \mathbb{E}[X_i X_j] - r_j \mathbb{E}[X_i X_\ell] - r_i \mathbb{E}[X_j X_\ell] + 2r_i r_j r_\ell \end{aligned}$$

$$\begin{aligned}
&= \frac{\left\{ \begin{aligned} &(\alpha_i + \mathbb{1}_{\{i=\ell \neq j\}} N^{-1})(\alpha_j + \mathbb{1}_{\{i=j\}} N^{-1})(\alpha_\ell + (\mathbb{1}_{\{j=\ell\}} + \mathbb{1}_{\{i=j=\ell\}}) N^{-1}) \cdot (\|\boldsymbol{\alpha}\|_1 + \beta)^2 \\ &-\alpha_i \alpha_\ell (\alpha_j + \mathbb{1}_{\{i=j\}} N^{-1}) \cdot (\|\boldsymbol{\alpha}\|_1 + \beta)(\|\boldsymbol{\alpha}\|_1 + \beta + 2N^{-1}) \\ &-\alpha_i \alpha_j (\alpha_\ell + \mathbb{1}_{\{i=\ell\}} N^{-1}) \cdot (\|\boldsymbol{\alpha}\|_1 + \beta)(\|\boldsymbol{\alpha}\|_1 + \beta + 2N^{-1}) \\ &-\alpha_i \alpha_j (\alpha_\ell + \mathbb{1}_{\{j=\ell\}} N^{-1}) \cdot (\|\boldsymbol{\alpha}\|_1 + \beta)(\|\boldsymbol{\alpha}\|_1 + \beta + 2N^{-1}) \\ &+ 2\alpha_i \alpha_j \alpha_\ell \cdot (\|\boldsymbol{\alpha}\|_1 + \beta + N^{-1})(\|\boldsymbol{\alpha}\|_1 + \beta + 2N^{-1}) \end{aligned} \right\}}{(\|\boldsymbol{\alpha}\|_1 + \beta)^3 (\|\boldsymbol{\alpha}\|_1 + \beta + N^{-1})(\|\boldsymbol{\alpha}\|_1 + \beta + 2N^{-1})} \\
&= N^{-2} \cdot \frac{\left\{ \begin{aligned} &4\alpha_i \alpha_j \alpha_\ell - 2\alpha_i \alpha_\ell \mathbb{1}_{\{i=j\}} (\|\boldsymbol{\alpha}\|_1 + \beta) - 2\alpha_j \alpha_\ell \mathbb{1}_{\{i=\ell\}} (\|\boldsymbol{\alpha}\|_1 + \beta) \\ &- 2\alpha_i \alpha_j \mathbb{1}_{\{j=\ell\}} (\|\boldsymbol{\alpha}\|_1 + \beta) + 2\alpha_i \mathbb{1}_{\{i=j=\ell\}} (\|\boldsymbol{\alpha}\|_1 + \beta)^2 \end{aligned} \right\}}{(\|\boldsymbol{\alpha}\|_1 + \beta)^3 (\|\boldsymbol{\alpha}\|_1 + \beta + N^{-1})(\|\boldsymbol{\alpha}\|_1 + \beta + 2N^{-1})} \\
&= N^{-2} \cdot \frac{4r_i r_j r_\ell - 2r_i r_\ell \mathbb{1}_{\{i=j\}} - 2r_j r_\ell \mathbb{1}_{\{i=\ell\}} - 2r_i r_j \mathbb{1}_{\{j=\ell\}} + 2r_i \mathbb{1}_{\{i=j=\ell\}}}{(\|\boldsymbol{\alpha}\|_1 + \beta + N^{-1})(\|\boldsymbol{\alpha}\|_1 + \beta + 2N^{-1})}, \tag{A.6}
\end{aligned}$$

which proves (A.2). Finally, trivial calculations show that

$$\begin{aligned}
\mathbb{E}[X_i^4] &= \frac{\Gamma(N\alpha_i + 4)}{\Gamma(N\alpha_i)} \cdot \frac{\Gamma(N\|\boldsymbol{\alpha}\|_1 + N\beta)}{\Gamma(N\|\boldsymbol{\alpha}\|_1 + N\beta + 4)} \\
&= \frac{N\alpha_i(N\alpha_i + 1)(N\alpha_i + 2)(N\alpha_i + 3) \cdot N^3(\|\boldsymbol{\alpha}\|_1 + \beta)^3}{N^4(\|\boldsymbol{\alpha}\|_1 + \beta)^4 \cdot (1 + \mathcal{O}_{\boldsymbol{\alpha},\beta}(N^{-1}))}. \tag{A.7}
\end{aligned}$$

We deduce

$$\begin{aligned}
\mathbb{E}[(X_i - r_i)^4] &= \mathbb{E}[X_i^4] - 4r_i \mathbb{E}[X_i^3] + 6r_i^2 \mathbb{E}[X_i^2] - 3r_i^4 \\
&= \frac{\left\{ \begin{aligned} &N\alpha_i(N\alpha_i + 1)(N\alpha_i + 2)(N\alpha_i + 3) \cdot N^3(\|\boldsymbol{\alpha}\|_1 + \beta)^3 \\ &- 4N^2\alpha_i^2(N\alpha_i + 1)(N\alpha_i + 2) \cdot N^2(\|\boldsymbol{\alpha}\|_1 + \beta)^2(N\|\boldsymbol{\alpha}\|_1 + N\beta + 3) \\ &+ 6N^3\alpha_i^3(N\alpha_i + 1) \cdot N(\|\boldsymbol{\alpha}\|_1 + \beta) \prod_{\ell=2}^3 (N\|\boldsymbol{\alpha}\|_1 + N\beta + \ell) \\ &- 3N^4\alpha_i^4 \cdot \prod_{\ell=1}^3 (N\|\boldsymbol{\alpha}\|_1 + N\beta + \ell) \end{aligned} \right\}}{N^7(\|\boldsymbol{\alpha}\|_1 + \beta)^7 \cdot (1 + \mathcal{O}_{\boldsymbol{\alpha},\beta}(N^{-1}))} \\
&= N^{-2} \cdot \frac{3\alpha_i^2(\|\boldsymbol{\alpha}\|_1 - \alpha_i + \beta)^2}{(\|\boldsymbol{\alpha}\|_1 + \beta)^6} + \mathcal{O}_{\boldsymbol{\alpha},\beta}(N^{-3}), \tag{A.8}
\end{aligned}$$

which proves (A.3). This ends the proof. \square

We can also estimate the moments of Lemma A.1 on various events. The lemma below is used to estimate the $\asymp \varepsilon_N^{1/2}$ errors in (3.4) of the proof of Theorem 3.1.

Lemma A.2. *Let $(\boldsymbol{\alpha}, \beta) \in (0, \infty)^{d+1}$ be given, and let $A \in \mathcal{B}(\mathbb{R}^d)$ be a Borel set. If $\mathbf{X} = (X_1, X_2, \dots, X_d) \sim \text{Dirichlet}(N\boldsymbol{\alpha}, N\beta)$ according to (1.2), then, for all $i, j, \ell \in \{1, 2, \dots, d\}$ and N large enough,*

$$\left| \mathbb{E}[(X_i - r_i) \mathbb{1}_{\{\mathbf{X} \in A\}}] \right| \leq \varepsilon_N^{1/2} (\mathbb{P}(\mathbf{X} \in A^c))^{1/2}, \tag{A.9}$$

$$\left| \frac{\mathbb{E}[(X_i - r_i)(X_j - r_j)(X_\ell - r_\ell) \mathbb{1}_{\{\mathbf{X} \in A\}}]}{-\varepsilon_N^2 \cdot \frac{4r_i r_j r_\ell - 2r_i r_\ell \mathbb{1}_{\{i=j\}} - 2r_j r_\ell \mathbb{1}_{\{i=\ell\}} - 2r_i r_j \mathbb{1}_{\{j=\ell\}} + 2r_i \mathbb{1}_{\{i=j=\ell\}}}{(1+\varepsilon_N)(1+2\varepsilon_N)}} \right| \leq \varepsilon_N^{3/2} (\mathbb{P}(\mathbf{X} \in A^c))^{1/4}, \tag{A.10}$$

where recall $\varepsilon_N := 1/(N\|\boldsymbol{\alpha}\|_1 + N\beta)$ and $r_i := \mathbb{E}[X_i] = \frac{\alpha_i}{\|\boldsymbol{\alpha}\|_1 + \beta}$ for all $i \in \{1, 2, \dots, d\}$.

Proof. For the bound in (A.9), note that $\mathbb{E}[X_i - r_i] = 0$. By Cauchy-Schwarz and a bound on the second moment of the beta distribution (see e.g. (A.1)), we have

$$\begin{aligned} |\mathbb{E}[(X_i - r_i)\mathbb{1}_{\{\mathbf{X} \in A\}}]| &= |\mathbb{E}[(X_i - r_i)\mathbb{1}_{\{\mathbf{X} \in A^c\}}]| \\ &\leq (\mathbb{E}[(X_i - r_i)^2])^{1/2} (\mathbb{P}(\mathbf{X} \in A^c))^{1/2} \\ &\leq \varepsilon_N^{1/2} (\mathbb{P}(\mathbf{X} \in A^c))^{1/2}. \end{aligned} \quad (\text{A.11})$$

For the bound in (A.10), Equation (A.2), Holder's inequality and a bound on the fourth moment of the beta distribution (see e.g. (A.3)) yield, for N large enough,

$$\begin{aligned} &\left| \frac{\mathbb{E}[(X_i - r_i)(X_j - r_j)(X_\ell - r_\ell)\mathbb{1}_{\{\mathbf{X} \in A\}}]}{-\varepsilon_N^2 \cdot \frac{4r_i r_j r_\ell - 2r_i r_\ell \mathbb{1}_{\{i=j\}} - 2r_j r_\ell \mathbb{1}_{\{i=\ell\}} - 2r_i r_j \mathbb{1}_{\{j=\ell\}} + 2r_i \mathbb{1}_{\{i=j=\ell\}}}{(1+\varepsilon_N)(1+2\varepsilon_N)}} \right| \\ &= \left| \mathbb{E}[(X_i - r_i)(X_j - r_j)(X_\ell - r_\ell)\mathbb{1}_{\{\mathbf{X} \in A^c\}}] \right| \\ &\leq (\mathbb{E}[(X_i - r_i)^4])^{1/4} (\mathbb{E}[(X_j - r_j)^4])^{1/4} (\mathbb{E}[(X_\ell - r_\ell)^4])^{1/4} (\mathbb{P}(\mathbf{X} \in A^c))^{1/4} \\ &\leq \varepsilon_N^{1/2} \varepsilon_N^{1/2} \varepsilon_N^{1/2} (\mathbb{P}(\mathbf{X} \in A^c))^{1/4}. \end{aligned} \quad (\text{A.12})$$

This ends the proof. \square

References

- Abramowitz, M., & Stegun, I. A. 1964. *Handbook of mathematical functions with formulas, graphs, and mathematical tables*. National Bureau of Standards Applied Mathematics Series, vol. 55. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. [MR0167642](#).
- Aitchison, J., & Lauder, I. J. 1985. Kernel density estimation for compositional data. *J. Roy. Statist. Soc. Ser. C*, **34**(2), 129–137. [doi:10.2307/2347365](#).
- Brown, L. D., & Low, M. G. 1996. Asymptotic equivalence of nonparametric regression and white noise. *Ann. Statist.*, **24**(6), 2384–2398. [MR1425958](#).
- Brown, L. D., Cai, T. T., Low, M. G., & Zhang, C.-H. 2002. Asymptotic equivalence theory for nonparametric regression with random design. *Ann. Statist.*, **30**(3), 688–707. [MR1922538](#).
- Cai, T. T., & Zhou, H. H. 2009. Asymptotic equivalence and adaptive estimation for robust nonparametric regression. *Ann. Statist.*, **37**(6A), 3204–3235. [MR2549558](#).
- Carter, A. V. 2002. Deficiency distance between multinomial and multivariate normal experiments. Dedicated to the memory of Lucien Le Cam. *Ann. Statist.*, **30**(3), 708–730. [MR1922539](#).
- Carter, A. V. 2006. A continuous Gaussian approximation to a nonparametric regression in two dimensions. *Bernoulli*, **12**(1), 143–156. [MR2202326](#).
- Carter, A. V. 2007. Asymptotic approximation of nonparametric regression experiments with unknown variances. *Ann. Statist.*, **35**(4), 1644–1673. [MR2351100](#).
- Golubev, G. K., Nussbaum, M., & Zhou, H. H. 2010. Asymptotic equivalence of spectral density estimation and Gaussian white noise. *Ann. Statist.*, **38**(1), 181–214. [MR2589320](#).
- Hirukawa, M. 2018. *Asymmetric kernel smoothing*. SpringerBriefs in Statistics. Springer, Singapore. [MR3821525](#).
- Marchal, O., & Arbel, J. 2017. On the sub-Gaussianity of the beta and Dirichlet distributions. *Electron. Commun. Probab.*, **22**, Paper No. 54, 14. [MR3718704](#).
- Mason, D. M., & Zhou, H. H. 2012. Quantile coupling inequalities and their applications. *Probab. Surv.*, **9**, 439–479. [MR3007210](#).
- Ng, K. W., Tian, G.-L., & Tang, M.-L. 2011. *Dirichlet and related distributions*. Wiley Series in Probability and Statistics. John Wiley & Sons, Ltd., Chichester. [MR2830563](#).

- Ouimet, F. 2020. Density estimation using Dirichlet kernels. *Preprint*, 1–39. [arXiv:2002.06956](#).
- Ouimet, F. 2021. A precise local limit theorem for the multinomial distribution and some applications. *J. Statist. Plann. Inference*, **215**, 218–233. [MR4249129](#).
- Rohde, A. 2004. On the asymptotic equivalence and rate of convergence of nonparametric regression and Gaussian white noise. *Statist. Decisions*, **22**(3), 235–243. [MR2125610](#).
- Tanabe, K., & Sagae, M. 1992. An exact Cholesky decomposition and the generalized inverse of the variance-covariance matrix of the multinomial distribution, with applications. *J. Roy. Statist. Soc. Ser. B*, **54**(1), 211–219. [MR1157720](#).
- Zhou, H. 2004. *Minimax estimation with thresholding and asymptotic equivalence for Gaussian variance regression*. PhD thesis, Cornell University.