

ZERO-CYCLES OVER ZERO-DIMENSIONAL CUSPS

SHOUHEI MA

ABSTRACT. We prove that all points of a toroidal compactification lying over 0-dimensional cusps are rationally equivalent in the integral Chow group for most classical modular varieties (Siegel, Hilbert, orthogonal, Hermitian, quaternionic). This gives a generalization, and even strengthening, of the Manin-Drinfeld theorem in higher dimension from the viewpoint of algebraic cycles. The same result no longer holds for Picard modular varieties, but for them we prove that the difference of any two "special" boundary points, which are dense in the boundary, is torsion.

1. INTRODUCTION

In [15], [6], Manin and Drinfeld discovered that the difference of two cusps of a congruence modular curve is torsion in its Picard group. We wish to study higher dimensional analogue of the Manin-Drinfeld theorem from the viewpoint of algebraic cycles. In [14] the Manin-Drinfeld theorem was generalized in the form of rational equivalence of two cusps in the rational Chow group of the Baily-Borel compactification of some modular varieties. The Baily-Borel compactification is however highly singular at the boundary in general, and it has been recognized that the study of Chow groups of singular varieties is more difficult than for nonsingular ones.

Toroidal compactification [1] provides an explicit desingularization of the Baily-Borel compactification when the arithmetic group is neat. Even when not smooth, its singularities are relatively mild, and its boundary is of codimension 1. It should be noticed, however, that a naive analogue of the Manin-Drinfeld theorem no longer holds for the boundary *divisors* of toroidal compactification. From the viewpoint of algebraic cycles, one of the next cycles to look at will be 0-cycles. In this paper we study some boundary 0-cycles of toroidal compactification. For classical modular varieties with tube domain model, we prove that they are rationally equivalent in the *integral* Chow group, hence define a distinguished integral Chow

2020 *Mathematics Subject Classification.* 14C25, 11F46, 11F41, 11F55.

Key words and phrases. 0-cycle, modular variety, toroidal compactification, Manin-Drinfeld theorem, Chow group.

Supported by KAKENHI 21H00971 and 20H00112.

class. The viewpoint of 0-cycles thus provides one direction in which the Manin-Drinfeld theorem can be generalized and even strengthened.

To be more specific, we consider the following series of modular varieties.

- Siegel modular variety \mathcal{D}/Γ attached to a finite-index subgroup Γ of the symplectic group $\mathrm{Sp}(\Lambda)$ of a symplectic lattice Λ of rank > 2 .
- Hilbert modular variety \mathbb{H}^n/Γ attached to a finite-index subgroup Γ of $\mathrm{SL}_2(\mathcal{O}_F)$ for a totally real number field F of degree $n > 1$.
- Orthogonal modular variety \mathcal{D}/Γ attached to a finite-index subgroup Γ of the orthogonal group $\mathrm{O}^+(\Lambda)$ of an integral quadratic form Λ of signature $(2, n)$ with $n > 2$.
- We also cover Hermitian modular varieties attached to the unitary groups $\mathrm{U}(n, n)$ and quaternionic modular varieties attached to the groups $\mathrm{SO}^*(4n)$, both with $n > 1$.

Except for the Hilbert case, \mathcal{D} is the classical irreducible Hermitian symmetric domain of type *III*, *IV*, *I_{n,n}*, *II_{2n}* respectively. According to [18], they are characterized by the existence of tube domain model. In the first two cases Γ is automatically a congruence subgroup ([17], [3], [21]).

Let X^{bb} be the Baily-Borel compactification of $Y = \mathcal{D}/\Gamma$, and X^{tor} be the toroidal compactification of Y defined by a Γ -admissible collection of fans Σ . (We suppress the dependence on Σ in the notation.) We may assume that Σ is chosen so that X^{tor} is projective ([1]). We have a natural morphism $\pi: X^{tor} \rightarrow X^{bb}$. In the Hilbert case, X^{bb} has only 0-dimensional cusps.

Our main result is the following.

Theorem 1.1. *Let Y be one of the modular varieties as above and X^{tor} be a projective toroidal compactification of Y . Let $P, Q \in X^{tor}$ be two points such that $\pi(P), \pi(Q)$ are 0-dimensional cusps (which may be distinct). Then $[P] = [Q]$ in the integral Chow group $\mathrm{CH}_0(X^{tor})$ of X^{tor} .*

Theorem 1.1 is in contrast to the case of modular curves, namely the original Manin-Drinfeld theorem. In that case, the difference of two cusps is nontrivial unless the modular curve has genus 0, and the calculation of its order in the Picard group has been the subject of many works on modular units and cuspidal divisor class groups. Thus, in this aspect, the situation gets simpler in higher dimension.

As a consequence of Theorem 1.1, we obtain a distinguished element of $\mathrm{CH}_0(X^{tor})$ that will play a fundamental role in the Chow ring of X^{tor} . In a few examples where X^{tor} is a *K3* Hilbert modular surface, this Chow class coincides with the Beauville-Voisin class [4] of the *K3* surface (Example 3.1). A similar coincidence also holds for a *K3* Picard modular surface (Example 5.4).

The proof of Theorem 1.1 is relatively simple. In the Siegel case, it proceeds as follows. We first show that $[P] - [Q]$ is torsion in $\text{CH}_0(X^{\text{tor}})$. When $\pi(P) = \pi(Q)$, this is a consequence of the structure of the boundary. When $\pi(P) \neq \pi(Q)$, we apply the Manin-Drinfeld theorem to a chain of certain modular curves in Y joining $\pi(P)$ and $\pi(Q)$. Finally, we conclude that $[P] = [Q]$ as a consequence of the Roitman theorem [19] and the Margulis normal subgroup theorem [16], two deep theorems from different branches of Mathematics. We can say that it is this last step that makes the difference between the curve case and the higher dimensional case. At the same time, this step leaves it somewhat mysterious *how* P and Q are equivalent in the integral Chow group.

The proof in the Hilbert case is similar. It is also possible to give a similar proof in the orthogonal case, but instead of such a repetition, we *reduce* the proof to the cases of Siegel 3-folds and Hilbert modular surfaces already proved. This argument enables us to prove Theorem 1.1 in the orthogonal case without assuming Γ congruence subgroup. The cases of $\text{U}(n, n)$ and $\text{SO}^*(4n)$ are similar. All classical domains for which our proof of Theorem 1.1 may work are covered.

The case excluded from the orthogonal case is $n = 2$ with Witt index 2. This is essentially the case of product of two modular curves. Theorem 1.1 does not hold for them. Indeed, their Albanese map is the product of the Abel-Jacobi maps of each curve, and so injective if both curves have genus > 0 . Thus the assumption in the orthogonal case would be best possible.

Theorem 1.1 does not hold for Picard modular varieties attached to the unitary groups $\text{U}(1, n)$. Perhaps this case could be more interesting. For simplicity we assume Γ neat (and congruence). The toroidal compactification of a Picard modular variety is canonical and has disjoint union of abelian varieties as the boundary. In general, its Albanese map does not contract the boundary to one point. Nevertheless we can prove the following analogue of Theorem 1.1. We define *special* boundary points using the structure of toroidal compactification (Definition 5.1). In each component of the boundary, such points form the set of torsion points of the abelian variety if we take one of them as the origin.

Theorem 1.2. *Let X^{tor} be the toroidal compactification of a Picard modular variety Y . Let $P, Q \in X^{\text{tor}} - Y$ be two special boundary points. Then $[P] - [Q]$ is torsion in $\text{CH}_0(X^{\text{tor}})$.*

By Theorem 1.2, the Chow class of special boundary points is uniquely determined in the rational Chow group $\text{CH}_0(X^{\text{tor}})_{\mathbb{Q}}$. By our geometric characterization of special boundary points, any intersection 0-cycle of a modular curve and a boundary divisor is contained in the span of this Chow class in $\text{CH}_0(X^{\text{tor}})_{\mathbb{Q}}$ (Corollary 5.3).

Theorem 1.2 leads to a (surviving) Chow-theoretic analogue of cuspidal divisor class groups for Picard modular varieties, as the subgroup of $\mathrm{CH}_0(X^{\mathrm{tor}})$ generated by the differences of two special boundary points (§5.4). By Theorem 1.2 and Roitman's theorem [19], to understand this group is reduced to the study of the configuration of the Albanese image of the boundary abelian varieties.

This paper is organized as follows. In §2 we prove Theorem 1.1 for Siegel modular varieties. In §3 we prove Theorem 1.1 for Hilbert modular varieties. In §4 we prove Theorem 1.1 for orthogonal modular varieties, and give an outline for the remaining cases. In §5 we prove Theorem 1.2.

2. SIEGEL MODULAR VARIETIES

In this section, after recollection in §2.1, we prove Theorem 1.1 for Siegel modular varieties in §2.2.

2.1. Siegel modular varieties. We first recall Siegel modular varieties and their Satake and toroidal compactifications. Let Λ be a free abelian group of rank $2g > 2$ equipped with a nondegenerate symplectic form $\Lambda \times \Lambda \rightarrow \mathbb{Z}$. The Hermitian symmetric domain $\mathcal{D} = \mathcal{D}_\Lambda$ attached to Λ is defined as the open locus of the Lagrangian Grassmannian consisting of g -dimensional isotropic subspaces W of $\Lambda_{\mathbb{C}}$ such that the associated Hermitian form on W is positive-definite. Let Γ be a finite-index subgroup of the symplectic group $\mathrm{Sp}(\Lambda)$. By [17], [3], Γ is a congruence subgroup. The quotient $Y = \mathcal{D}/\Gamma$ is a quasi-projective variety, called a Siegel modular variety.

The Satake compactification of Y , denoted by X^{bb} for consistency with other sections, is defined as the quotient by Γ of the union of \mathcal{D} and its rational boundary components, equipped with the Satake topology. It is normal and projective ([2]). Rational boundary components (cusps) of \mathcal{D} correspond to isotropic subspaces I of $\Lambda_{\mathbb{Q}}$. In particular, when I is maximal, i.e., $\dim I = g$, the corresponding cusp is 0-dimensional, given by the point $[I_{\mathbb{C}}]$ of the Lagrangian Grassmannian.

A toroidal compactification $X^{\mathrm{tor}} = X_{\Gamma}^{\Sigma}$ of Y ([1]) is constructed by choosing a Γ -admissible collection of fans $\Sigma = (\Sigma_I)$, one for each isotropic subspace I of $\Lambda_{\mathbb{Q}}$. Each fan Σ_I is a polyhedral cone decomposition of the cone of positive semidefinite forms with rational kernel in $\mathrm{Sym}^2 I_{\mathbb{R}}$. This defines a partial compactification $(\mathcal{D}/U(I)_{\mathbb{Z}})^{\Sigma_I}$ of $\mathcal{D}/U(I)_{\mathbb{Z}}$, where $U(I)_{\mathbb{Z}} = \Gamma \cap U(I)_{\mathbb{Q}}$ and $U(I)_{\mathbb{Q}} \simeq \mathrm{Sym}^2 I$ is the center of the unipotent part of the stabilizer $N(I)_{\mathbb{Q}}$ of I in $\mathrm{Sp}(\Lambda_{\mathbb{Q}})$. We write $\Gamma(I)_{\mathbb{Z}} = \Gamma \cap N(I)_{\mathbb{Q}}$ and $\overline{\Gamma(I)}_{\mathbb{Z}} = \Gamma(I)_{\mathbb{Z}}/U(I)_{\mathbb{Z}}$. Then X^{tor} is constructed as a gluing of Y and the quotients $(\mathcal{D}/U(I)_{\mathbb{Z}})^{\Sigma_I}/\overline{\Gamma(I)}_{\mathbb{Z}}$. X^{tor} is a compact Moishezon space with a natural morphism $\pi : X^{\mathrm{tor}} \rightarrow X^{\mathrm{bb}}$. We may choose Σ so that X^{tor} is projective ([1]).

We are interested in the case $\dim I = g$. Let $p \in X^{bb}$ be the 0-dimensional cusp corresponding to I . In this case $\mathcal{D}/U(I)_{\mathbb{Z}}$ is embedded in the algebraic torus $T_I = \text{Sym}^2 I_{\mathbb{C}}/U(I)_{\mathbb{Z}}$ by the tube domain realization of \mathcal{D} with respect to I (= isomorphism with the Siegel upper half space). The partial compactification $\mathcal{D}/U(I)_{\mathbb{Z}} \hookrightarrow (\mathcal{D}/U(I)_{\mathbb{Z}})^{\Sigma_I}$ is an open set of the torus embedding $T_I \hookrightarrow T_I^{\Sigma_I}$ defined by the fan Σ_I . If $\sigma \in \Sigma_I$ is a cone whose relative interior consists of positive-definite forms, the corresponding torus orbit $\text{orb}(\sigma) \subset T_I^{\sigma} \subset T_I^{\Sigma_I}$ is contained in $(\mathcal{D}/U(I)_{\mathbb{Z}})^{\Sigma_I}$. Then the locus of boundary points of $(\mathcal{D}/U(I)_{\mathbb{Z}})^{\Sigma_I}$ which lie over the cusp p is the union of these torus orbits ([1] p.164). This shows that

$$(2.1) \quad \pi^{-1}(p) = (\sqcup_{\sigma} \text{orb}(\sigma)) / \overline{\Gamma(I)_{\mathbb{Z}}},$$

where σ runs over cones with positive-definite relative interior, and $\sqcup_{\sigma} \text{orb}(\sigma)$ is the union of infinitely many toric varieties whose configuration is determined by Σ_I .

2.2. Proof of Theorem 1.1. We now prove Theorem 1.1 for Siegel modular varieties. Recall that the given toroidal compactification $X^{tor} = X_{\Gamma}^{\Sigma}$ is assumed to be projective. We begin with the following reduction, which will be necessary in Step 4.

Step 1. *We may assume that Γ is neat and X^{tor} is smooth and projective.*

Proof. We take a neat subgroup Γ' of Γ of finite index. Then Σ is also Γ' -admissible, so we have the toroidal compactification $X_{\Gamma'}^{\Sigma}$ of \mathcal{D}/Γ' defined by Σ . Since we have a finite morphism $X_{\Gamma'}^{\Sigma} \rightarrow X_{\Gamma}^{\Sigma}$ and $X_{\Gamma'}^{\Sigma}$ is projective, X_{Γ}^{Σ} is projective too. Next, by [1] Corollary III.7.6, we can take a Γ' -admissible subdivision Σ' of Σ such that $X_{\Gamma'}^{\Sigma'} \rightarrow X_{\Gamma'}^{\Sigma}$ is projective and $X_{\Gamma'}^{\Sigma'}$ is smooth. Then $X_{\Gamma'}^{\Sigma'}$ is also projective.

For any point of X_{Γ}^{Σ} lying over a 0-dimensional cusp of X_{Γ}^{bb} , the points in its inverse image in $X_{\Gamma'}^{\Sigma'}$ lie over 0-dimensional cusps of $X_{\Gamma'}^{bb}$. Therefore, if we could prove Theorem 1.1 for $X_{\Gamma'}^{\Sigma'}$, the assertion for X_{Γ}^{Σ} follows by pushforward. In what follows, we rewrite (Γ, Σ) as (Γ, Σ) . \square

Let P, Q be two points of X^{tor} such that $p = \pi(P)$ and $q = \pi(Q)$ are 0-dimensional cusps of X^{bb} . We first show that $[P] - [Q] \in \text{CH}_0(X^{tor})$ is torsion in two steps.

Step 2. *If $p = q$, we have $[P] = [Q]$ in $\text{CH}_0(X^{tor})$.*

Proof. By the description (2.1) of the π -fiber over p , we see that $\pi^{-1}(p)$ is a connected configuration of finitely many toric varieties. This shows that we can join P and Q by a chain of rational curves inside $\pi^{-1}(p)$. \square

Step 3. *Even when $p \neq q$, $[P] - [Q] \in \text{CH}_0(X^{tor})$ is torsion.*

Proof. We take maximal isotropic subspaces I, I' of $\Lambda_{\mathbb{Q}}$ which correspond to p, q respectively. We first consider the case when the pairing (I, I') is nondegenerate. In that case, we take splittings $I = I_1 \oplus \cdots \oplus I_g$ and $I' = I'_1 \oplus \cdots \oplus I'_g$ such that $(I_i, I'_j) = 0$ if $i \neq j$ and $(I_i, I'_i) = \mathbb{Q}$. If we put $\Lambda_i = I_i \oplus I'_i$, we have the orthogonal splitting $\Lambda_{\mathbb{Q}} = \Lambda_1 \oplus \cdots \oplus \Lambda_g$ as a symplectic space. Note that we have an isomorphism $\Lambda_i \simeq \mathbb{Q}^2$ of symplectic space which sends I_i, I'_i to $\mathbb{Q}(1, 0), \mathbb{Q}(0, 1)$ respectively. This defines an embedding of the upper half plane $\mathbb{H} = \mathcal{D}_{\mathbb{Q}^2} \subset \mathbb{P}^1$ as

$$\mathcal{D}_{\mathbb{Q}^2} \hookrightarrow \mathcal{D}_{\Lambda_1} \times \cdots \times \mathcal{D}_{\Lambda_g} \hookrightarrow \mathcal{D}_{\Lambda}, \quad V \mapsto V \oplus \cdots \oplus V.$$

Since Γ is a congruence subgroup, this induces a finite morphism $Y(N) \rightarrow Y$ from the principal congruence modular curve $Y(N) = \mathbb{H}/\Gamma(N)$ of some level N . If $X(N)$ is the compactification of $Y(N)$, this extends to a morphism $X(N) \rightarrow X^{bb}$ which sends the cusps $0, i\infty$ of $X(N)$ to $p, q \in X^{bb}$ respectively. Furthermore, since $X(N)$ is a curve, this lifts to a morphism $f : X(N) \rightarrow X^{tor}$ to the toroidal compactification. We have $[P] = [f(0)]$ and $[Q] = [f(i\infty)]$ by Step 2. By the Manin-Drinfeld theorem, $[0] - [i\infty]$ is torsion in $\text{CH}_0(X(N))$. Sending this equality by f_* , we find that

$$[P] - [Q] = [f(0)] - [f(i\infty)] = f_*([0] - [i\infty])$$

is torsion in $\text{CH}_0(X^{tor})$. This proves the assertion when the pairing (I, I') is nondegenerate.

When (I, I') is degenerate, we can find another maximal isotropic subspace I'' such that both (I, I'') and (I', I'') are nondegenerate. If we take a point $R \in X^{tor}$ lying over the I'' -cusp, we can use the above case twice to see that $[P] - [Q] = ([P] - [R]) + ([R] - [Q])$ is torsion in $\text{CH}_0(X^{tor})$. \square

Step 4. We have $[P] = [Q]$ in $\text{CH}_0(X^{tor})$.

Proof. Let $A_0(X^{tor})$ be the subgroup of $\text{CH}_0(X^{tor})$ of degree 0 cycles. By the theorem of Roitman [19], the Albanese map $A_0(X^{tor}) \rightarrow \text{Alb}(X^{tor})$ is injective on the torsion part of $A_0(X^{tor})$. On the other hand, the Albanese variety $\text{Alb}(X^{tor})$ is trivial by the following Proposition 2.1. This implies that $A_0(X^{tor})$ is torsion-free, and so $[P] - [Q] = 0$ in $\text{CH}_0(X^{tor})$. This completes the proof of Theorem 1.1 for Siegel modular varieties. \square

In the final step we used the following general consequence of the Margulis normal subgroup theorem ([16] p.4).

Proposition 2.1. *Let $\mathcal{D} = G/K$ be a Hermitian symmetric domain and $\Gamma < G$ be an arithmetic group. Assume that the Lie group G has real rank ≥ 2 and Γ is irreducible. Then the Albanese variety of a smooth projective model of \mathcal{D}/Γ is trivial.*

Proof. Let \tilde{X} be a smooth projective model of \mathcal{D}/Γ . We have a surjective homomorphism $\Gamma \twoheadrightarrow \pi_1(\tilde{X})$ (see, e.g., [20]). This induces a surjective homomorphism $\Gamma^{ab} \twoheadrightarrow H_1(\tilde{X}, \mathbb{Z})$ between their abelianizations. Then Γ^{ab} is finite by the Margulis theorem, so $H_1(\tilde{X}, \mathbb{Z})$ is finite. \square

For Siegel modular varieties, Proposition 2.1 also follows from Weisauer's theorem [23]. For orthogonal modular varieties, Proposition 2.1 is also proved in [5] when Γ is congruence, using L^2 -cohomology and Lie algebra cohomology. This can also be proved in an elementary way by induction on the dimension, using restriction to sub orthogonal modular varieties and starting from Hilbert modular surfaces.

Remark 2.2. A similar argument, using finiteness of the Mordell-Weil groups of elliptic modular surfaces [22], yields a weaker result for boundary 1-cycles in the case of Siegel 3-folds: if $Z_1, Z_2 \subset X^{tor}$ are irreducible 1-cycles such that $\pi(Z_1), \pi(Z_2)$ are 1-dimensional cusps, then $aZ_1 \sim bZ_2 + F$ for some $a, b \in \mathbb{Z}_{>0}$ and a 1-cycle F whose support is contracted by π to points.

Remark 2.3. The situation is totally different for the boundary divisors. They tend to be linearly independent already in the level of cohomology. See [10] Proposition 4.3 for the boundary divisors over cusps of corank 1. Also, subdivision of fan leads to blow-up (for Γ neat), which just adds new components to H^2 .

3. HILBERT MODULAR VARIETIES

Let F be a totally real number field of degree $n > 1$. The group $\mathrm{SL}_2(F)$ acts on \mathbb{H}^n through the natural embedding $\mathrm{SL}_2(F) \hookrightarrow \mathrm{SL}_2(\mathbb{R})^n$. Let Γ be a finite-index subgroup of $\mathrm{SL}_2(\mathcal{O}_F)$. By [21], Γ is a congruence subgroup. The quotient $Y = \mathbb{H}^n/\Gamma$ is a quasi-projective variety called a Hilbert modular variety. The Baily-Borel compactification $Y \hookrightarrow X^{bb}$ is obtained by adding finitely many points (0-dimensional cusps) corresponding to $\mathbb{P}^1(F)/\Gamma$, with the Satake topology ([2], [7]). A toroidal compactification X^{tor} of Y can be constructed by choosing a collection of suitable polyhedral cone decompositions of the cone of totally positive elements of $F \otimes_{\mathbb{Q}} \mathbb{R} \simeq \mathbb{R}^n$, one for each cusp independently ([1], [7]). We have a natural morphism $X^{tor} \rightarrow X^{bb}$. Hence $X^{tor} - Y$ is a divisor lying over the finitely many points $X^{bb} - Y$.

3.1. Proof of Theorem 1.1. The proof of Theorem 1.1 for Hilbert modular varieties is similar (and simpler) to the case of Siegel modular varieties. Instead of repetition, we just indicate that the ingredients in the argument of §2.2 have counterparts in the present case. We may assume as before that Γ is neat and X^{tor} is smooth and projective.

- The boundary divisor of X^{tor} over a cusp is again a connected configuration of toric varieties.
- The diagonal embedding $\mathbb{H} \hookrightarrow \mathbb{H}^n$ gives a congruence modular curve joining the cusps $0 = (0, \dots, 0)$ and $i\infty = (i\infty, \dots, i\infty)$. Since $SL_2(F)$ acts transitively on $\mathbb{P}^1(F) \times \mathbb{P}^1(F) \setminus \text{diagonal}$, we can translate the diagonal embedding to obtain a congruence modular curve joining any two distinct cusps.
- Proposition 2.1 applies to the Hilbert modular varieties, so the Albanese variety of X^{tor} is trivial. In fact, this follows from a more elementary argument (see [7] p.82 and p.18).

With these ingredients, Theorem 1.1 for Hilbert modular varieties can be proved by the same argument as in the case of Siegel modular varieties. \square

Example 3.1. In a few examples in $n = 2$, the desingularization of X^{tor} is a blown-up $K3$ surface, and none of the boundary curves are contracted in the minimal model X_{K3} (see [7] Chapter VII, §3 – §4). Since the boundary curves are rational, the unique class in $CH_0(X_{K3})$ given by the boundary points coincides with the Beauville-Voisin class [4] of the $K3$ surface X_{K3} . By [4], the span of the Beauville-Voisin class contains the image of the intersection product $\text{Pic}(X_{K3}) \otimes \text{Pic}(X_{K3}) \rightarrow CH_0(X_{K3})$. This provides rational equivalence between boundary 0-cycles and special 0-cycles in Y obtained as intersection of two modular curves.

3.2. Higher modular units. Let $Y \hookrightarrow X^{tor}$ be a Hilbert modular variety and its toroidal compactification as above. Theorem 1.1 leads to a higher dimensional analogue of modular units (cf. [12]) as higher Chow cycles on Y . We write $X = X^{tor}$ and $\partial X = X - Y$. We consider a part of the localization exact sequence of higher Chow groups for $Y \hookrightarrow X \leftarrow \partial X$:

$$CH_0(X, 1) \xrightarrow{i^*} CH_0(Y, 1) \xrightarrow{\delta} CH_0(\partial X) \xrightarrow{j_*} CH_0(X).$$

Here $i: Y \hookrightarrow X$ and $j: \partial X \hookrightarrow X$ are natural embeddings and δ is the connecting map. Let $p \neq q \in X^{bb}$ be two cusps. We take $P, Q \in X$ such that $\pi(P) = p$ and $\pi(Q) = q$. We can view P, Q also as 0-cycles on ∂X .

Corollary 3.2. *There exists a nonzero element $Z_{p,q} \in CH_0(Y, 1)$, uniquely determined from (p, q) up to $i^*CH_0(X, 1)$, such that*

$$\delta(Z_{p,q}) = [P] - [Q] \in CH_0(\partial X).$$

Proof. Since $j_*([P] - [Q]) = 0$ in $CH_0(X)$ by Theorem 1.1, the existence and uniqueness of $Z_{p,q}$ follow from the exactness of the localization sequence. It is independent of the choice of P and Q because $[P] = [P'] \in CH_0(\partial X)$ if $\pi(P) = \pi(P')$ (and similarly for Q). Since P and Q belong to different connected components of ∂X , we have $[P] \neq [Q]$ in $CH_0(\partial X)$ and so $Z_{p,q} \neq 0$. \square

The higher Chow cycle $Z_{p,q}$ can be thought of as an analogue of modular units. Since it is left somewhat mysterious *how* P and Q are rationally equivalent, so is $Z_{p,q}$. If $p, q, r \in X^{bb}$ are three cusps, then

$$Z_{p,q} + Z_{q,r} \equiv Z_{p,r} \pmod{i^* \text{CH}_0(X, 1)}.$$

Therefore the element $Z_{p,q} + Z_{q,r} - Z_{p,r}$ of $\text{CH}_0(Y, 1)$ provides an element of $\text{CH}_0(X, 1)$.

4. ORTHOGONAL MODULAR VARIETIES

In §4.2 we prove Theorem 1.1 for orthogonal modular varieties, after recollection in §4.1. In §4.3 we give an outline for the remaining cases (Hermitian and quaternionic modular varieties).

4.1. Orthogonal modular varieties. We begin by recalling orthogonal modular varieties and their Baily-Borel and toroidal compactifications. In this section we let Λ be a free abelian group of rank $2 + n$ equipped with a nondegenerate symmetric bilinear form $\Lambda \times \Lambda \rightarrow \mathbb{Z}$ of signature $(2, n)$. Let $Q \subset \mathbb{P}\Lambda_{\mathbb{C}}$ be the isotropic quadric defined by $(\omega, \omega) = 0$. The Hermitian symmetric domain \mathcal{D} attached to Λ is one of the two connected components of the open set of Q defined by $(\omega, \bar{\omega}) > 0$. Let $O^+(\Lambda)$ be the index ≤ 2 subgroup of the orthogonal group $O(\Lambda)$ preserving the component \mathcal{D} . Let Γ be a subgroup of $O^+(\Lambda)$ of finite index. The quotient $Y = \mathcal{D}/\Gamma$ is a quasi-projective variety.

The domain \mathcal{D} has 0-dimensional and 1-dimensional rational boundary components, corresponding to 1-dimensional and 2-dimensional isotropic subspaces I of $\Lambda_{\mathbb{Q}}$ respectively. When $\dim I = 1$, the corresponding boundary point is the point $[I_{\mathbb{C}}]$ of Q . The Baily-Borel compactification X^{bb} of Y is defined as the quotient by Γ of the union of \mathcal{D} and the rational boundary components, equipped with the Satake topology.

A toroidal compactification X^{tor} of Y is constructed by choosing a finite collection of suitable fans $\Sigma = (\Sigma_I)$, one for each Γ -equivalence class of isotropic lines I in $\Lambda_{\mathbb{Q}}$ independently. For each I , the tube domain realization (= the linear projection from the boundary point) defines an isomorphism of $\mathcal{D}/U(I)_{\mathbb{Z}}$ with an open set of the algebraic torus $T_I = U(I)_{\mathbb{C}}/U(I)_{\mathbb{Z}}$. Here $U(I)_{\mathbb{Z}} = \Gamma \cap U(I)_{\mathbb{Q}}$ and $U(I)_{\mathbb{Q}}$ is the unipotent part of the stabilizer of I in $O^+(\Lambda_{\mathbb{Q}})$. $U(I)_{\mathbb{Q}}$ is canonically isomorphic to the quadratic space $(I^{\perp}/I) \otimes I$. The fan Σ_I gives a polyhedral cone decomposition of the extended positive cone of $U(I)_{\mathbb{R}}$. It defines the partial compactification $\mathcal{D}/U(I)_{\mathbb{Z}} \hookrightarrow (\mathcal{D}/U(I)_{\mathbb{Z}})^{\Sigma_I}$ inside the torus embedding $T_I \hookrightarrow T_I^{\Sigma_I}$. If $\sigma \in \Sigma_I$ is a cone which is not an isotropic ray, its relative interior is contained in the

positive cone, and the corresponding torus orbit $\text{orb}(\sigma)$ in $T_I^{\Sigma_I}$ is also contained in $(\mathcal{D}/U(I_{\mathbb{Z}}))^{\Sigma_I}$. The union of these torus orbits $\text{orb}(\sigma)$ is the locus of boundary points of $(\mathcal{D}/U(I_{\mathbb{Z}}))^{\Sigma_I}$ which lie over the I -cusp ([1] p.164).

We also have the partial compactifications at the 1-dimensional cusps, which are canonical and require no choice. The toroidal compactification X^{tor} is then defined as a gluing of Y and natural quotients of these partial compactifications ([1]). X^{tor} is a compact Moishezon space with a morphism $\pi : X^{\text{tor}} \rightarrow X^{\text{bb}}$.

4.2. Proof of Theorem 1.1. We now prove Theorem 1.1 in the orthogonal case. As before we may assume that Γ is neat and X^{tor} is smooth and projective. Let P, Q be two points of X^{tor} such that $p = \pi(P)$, $q = \pi(Q)$ are 0-dimensional cusps, corresponding to isotropic lines I_1, I_2 of $\Lambda_{\mathbb{Q}}$ respectively. When $p = q$, we have $[P] = [Q]$ in $\text{CH}_0(X^{\text{tor}})$ because $\pi^{-1}(p)$ is a connected union of finitely many toric varieties by our description of the partial compactifications. In the general case $p \neq q$, it is possible to run a similar argument as before when Γ is a congruence subgroup. But instead of such a repetition, we give another proof: we reduce the proof to the cases of Siegel 3-folds and Hilbert modular surfaces which were already proved. We do not assume that Γ is a congruence subgroup.

We first consider the case Λ has Witt index 2. Then we can find a subspace $\Lambda'_{\mathbb{Q}} \subset \Lambda_{\mathbb{Q}}$ containing I_1, I_2 such that $\Lambda'_{\mathbb{Q}} \simeq 2U_{\mathbb{Q}} \oplus \langle -2t \rangle$ for some $t > 0$. Here $U_{\mathbb{Q}}$ is the hyperbolic plane. We have $\mathcal{D}_{\Lambda'} \subset \mathcal{D}_{\Lambda}$, and this induces a morphism $Y' = \mathcal{D}_{\Lambda'}/\Gamma' \rightarrow Y$ for some arithmetic subgroup Γ' of $\text{SO}^+(\Lambda'_{\mathbb{Q}})$. This extends to a morphism between the Baily-Borel compactifications, say $f : (X')^{\text{bb}} \rightarrow X^{\text{bb}}$, and also between toroidal compactifications, say $f : (X')^{\text{tor}} \rightarrow X^{\text{tor}}$, for a suitable choice of fans Σ' for Γ' (cf. [8]). Then I_1, I_2 give 0-dimensional cusps p', q' of $(X')^{\text{bb}}$ such that $f(p') = p$ and $f(q') = q$ respectively. Now Y' is isomorphic to a Siegel modular 3-fold. Therefore we can apply Theorem 1.1 to $(X')^{\text{tor}}$ which is already proved in §2.2. This shows that $[P'] = [Q']$ in $\text{CH}_0((X')^{\text{tor}})$ for any points $P', Q' \in (X')^{\text{tor}}$ lying over p', q' respectively. Sending this equality by f_* , we see that

$$[P] = [f(P')] = [f(Q')] = [Q]$$

in $\text{CH}_0(X^{\text{tor}})$.

In the case Λ has Witt index 1, we can instead take a subspace $\Lambda'_{\mathbb{Q}} \supset I_1, I_2$ isometric to $U_{\mathbb{Q}} \oplus K$ with K anisotropic of signature $(1, 1)$. Then $\mathcal{D}_{\Lambda'}/\Gamma'$ for a suitable $\Gamma' < \text{SO}^+(\Lambda'_{\mathbb{Q}})$ is isomorphic to a Hilbert modular surface, for which Theorem 1.1 is proved in §3.1. This finishes the proof of Theorem 1.1 for orthogonal modular varieties. \square

4.3. **Other modular varieties.** Theorem 1.1 also holds in the following cases:

- The unitary group $U(n, n)$ of a Hermitian form of signature (n, n) and Witt index n with $n > 1$ over an imaginary quadratic field. The domain \mathcal{D} is of type $I_{n,n}$ (Hermitian upper half space), and its compact dual is the Grassmannian $G(n, 2n)$.
- The quaternionic group, usually written as $SO^*(4n)$ or sometimes $Sp(2n, \mathbb{H})$, attached to a skew-Hermitian space of dimension $2n$ and Witt index n with $n > 1$ over a definite quaternion algebra over \mathbb{Q} (cf. [13]). The domain \mathcal{D} is of type II_{2n} (quaternion upper half space), and its compact dual is the orthogonal Grassmannian $OG(2n, 4n)$.

Since we are considering 0-dimensional cusps, the (skew-)Hermitian form must have maximal Witt index, and hence unique up to isometry. In both cases, 0-dimensional cusps correspond to rational maximal isotropic subspaces. The Siegel domain realization at a 0-dimensional cusp is a tube domain (Hermitian/quaternion upper half space), so the boundary locus of toroidal compactification over that cusp is a union of toric varieties. Proposition 2.1 also holds in these cases. For joining two 0-dimensional cusps, we can use (a chain of translates of) diagonal embeddings $I_{1,1} \hookrightarrow I_{n,n}$ and $II_2 \hookrightarrow II_{2n}$ respectively. In the spirit of §4.2, we can use

$$III_n \hookrightarrow I_{n,n} \hookrightarrow II_{2n}.$$

We can also use $IV_4 \simeq I_{2,2} \hookrightarrow I_{2n,2n}$ and $IV_6 \simeq II_4 \hookrightarrow II_{4n}$ when the dimension is even.

This exhausts all possible classical domains for which our proof of Theorem 1.1 may work. In general, Step 2 in §2.2 works if the unipotent radical of the stabilizer of the 0-dimensional cusp is abelian. Step 4 works if the Lie group has real rank > 1 and the arithmetic group is irreducible. It is not clear if Step 3 can be done systematically in the general framework of Shimura varieties.

5. PICARD MODULAR VARIETIES

As explained in §1, Theorem 1.1 does not hold for Picard modular varieties. Besides explicit example (e.g., [9], [11]), we can tell which part of the argument in §2.2 does not work: Step 2, as the boundary divisors are abelian varieties; Step 4, as the Albanese variety of a Picard modular variety is in general nontrivial. In this section we show that nevertheless we can formulate and prove an analogous result (Theorem 1.2).

In §5.1 we recall Picard modular varieties and their compactifications. In §5.2 we define *special* boundary points. In §5.3 we prove Theorem 1.2. In

§5.4 we explain that Theorem 1.2 leads to a 0-cycle analogue of cuspidal class groups for Picard modular varieties.

5.1. Picard modular varieties. Let K be an imaginary quadratic field. Let V be a K -linear space of dimension $n + 1$ endowed with a Hermitian form $V \times V \rightarrow K$ of signature $(1, n)$. We write $V_{\mathbb{C}} = V \otimes_K \mathbb{C} = V \otimes_{\mathbb{Q}} \mathbb{R}$. The space

$$\mathcal{D} = \{\mathbb{C}v \in \mathbb{P}V_{\mathbb{C}} \mid (v, v) > 0\}$$

is isomorphic to the unit ball in \mathbb{C}^n . Let $U(V)$ be the unitary group of V and Γ be a congruence subgroup of $U(V)$. We assume, for simplicity of exposition, that Γ is neat. The quotient $Y = \mathcal{D}/\Gamma$ is a smooth quasi-projective variety called a Picard modular variety. (It is also sometimes called a *ball quotient*, especially when Γ is not necessarily arithmetic.) Rational boundary points of \mathcal{D} correspond to isotropic K -lines I in V . Explicitly, to I corresponds the point $p_I = [I_{\mathbb{C}}]$ of $\mathbb{P}V_{\mathbb{C}}$. The Baily-Borel compactification X^{bb} of Y is obtained by adding finitely many points corresponding to the Γ -equivalence classes of these points p_I .

Let $I \subset V$ be an isotropic K -line. We write $V(I) = (I^{\perp}/I) \otimes_K \bar{I}$ and $H(I) = \mathbb{P}(V/I)_{\mathbb{C}} - \mathbb{P}(I^{\perp}/I)_{\mathbb{C}}$. Then $H(I)$ is an affine space isomorphic to $V(I)_{\mathbb{C}}$ (with no specified base point). Consider the projection

$$\phi: \mathbb{P}V_{\mathbb{C}} - p_I \rightarrow \mathbb{P}(V/I)_{\mathbb{C}}$$

from the point p_I . Each point of $\mathbb{P}(V/I)_{\mathbb{C}}$, say $[W]$, corresponds to a \mathbb{C} -plane $W \subset V_{\mathbb{C}}$ containing $I_{\mathbb{C}}$, and the ϕ -fiber over $[W]$ is the affine line $\mathbb{P}W - p_I$. When $[W] \in H(I)$, the Hermitian form on W is nondegenerate of signature $(1, 1)$. This shows that $\phi|_{\mathcal{D}}$ realizes \mathcal{D} as a fibration of upper half planes over $H(I)$, with the fiber over $[W]$ being the positive ball $\mathcal{D} \cap \mathbb{P}W$ in $\mathbb{P}W$. This is the Siegel domain realization of \mathcal{D} with respect to I .

Let $N(I)_{\mathbb{Q}} \subset U(V)$ be the stabilizer of I in $U(V)$. We write

$$W(I)_{\mathbb{Q}} = \text{Ker}(N(I)_{\mathbb{Q}} \rightarrow \text{GL}(I) \times U(I^{\perp}/I)),$$

$$U(I)_{\mathbb{Q}} = \text{Ker}(N(I)_{\mathbb{Q}} \rightarrow \text{GL}(I^{\perp})).$$

Then $U(I)_{\mathbb{Q}} \simeq \mathbb{Q}$ is isomorphic to the purely imaginary part of $I \otimes_K \bar{I} \simeq K$. Explicitly, if we choose $v \in I$ and isotropic $v' \in V$ with $(v, v') = 1$, a purely imaginary $\alpha \in K$ corresponds to the isometry sending $v' \mapsto v' + \alpha v$ and identity on I^{\perp} . The group $W(I)_{\mathbb{Q}}$ is the unipotent part of $N(I)_{\mathbb{Q}}$. It is a Heisenberg group with center $U(I)_{\mathbb{Q}}$ and $W(I)_{\mathbb{Q}}/U(I)_{\mathbb{Q}} \simeq V(I)$ as \mathbb{Q} -linear space. The group $U(I)_{\mathbb{Q}}$ acts on $H(I)$ trivially and on the fibers of $\mathcal{D} \rightarrow H(I)$ by translation. The quotient $W(I)_{\mathbb{Q}}/U(I)_{\mathbb{Q}}$ acts on the affine space $H(I)$ as the translation by $V(I)$.

Let now $\Gamma < U(V)$ be a neat congruence subgroup. We write

$$W(I)_{\mathbb{Z}} = \Gamma \cap W(I)_{\mathbb{Q}}, \quad U(I)_{\mathbb{Z}} = \Gamma \cap U(I)_{\mathbb{Q}}, \quad V(I)_{\mathbb{Z}} = W(I)_{\mathbb{Z}}/U(I)_{\mathbb{Z}}.$$

Since Γ is neat, $W(I)_{\mathbb{Z}}$ coincides with $\Gamma \cap N(I)_{\mathbb{Q}}$. $U(I)_{\mathbb{Z}} \simeq \mathbb{Z}$ is a lattice in $U(I)_{\mathbb{Q}} \simeq \mathbb{Q}$ and $V(I)_{\mathbb{Z}}$ is a full lattice in $V(I)$, namely $V(I)_{\mathbb{Z}} \otimes_{\mathbb{Z}} \mathbb{Q} = V(I)$. The quotient

$$A(I) = V(I)_{\mathbb{C}}/V(I)_{\mathbb{Z}}$$

is an abelian variety isogenous to the $(n - 1)$ -fold self product of an elliptic curve with complex multiplication by K .

The toroidal compactification X^{tor} of Y is a canonically constructed smooth projective variety with a morphism $\pi: X^{tor} \rightarrow X^{bb}$. No choice of fan is required: it is canonical. Explicitly, for each I , we consider the punctured disc bundle $\mathcal{D}/U(I)_{\mathbb{Z}}$ over $H(I)$ and take the partial compactification, say $\overline{\mathcal{D}/U(I)_{\mathbb{Z}}}$, by filling in the origins. Then X^{tor} is obtained by gluing Y and $\overline{\mathcal{D}/U(I)_{\mathbb{Z}}}/V(I)_{\mathbb{Z}}$ around each cusp p_I . By construction the boundary divisor $\Delta(I) = \pi^{-1}(p_I)$ over p_I is canonically identified with $H(I)/V(I)_{\mathbb{Z}}$. This is a torsor for the abelian variety $A(I)$ (with no specified origin).

5.2. Special boundary points. Let $I \subset V$ be an isotropic K -line. The affine space $H(I) = \mathbb{P}(V/I)_{\mathbb{C}} - \mathbb{P}(I^{\perp}/I)_{\mathbb{C}}$ is naturally defined over K . A K -rational point of $H(I)$ can be written as $[W] = [W_K \otimes_K \mathbb{C}]$ for a K -plane $W_K \subset V$ containing I . Since the Hermitian form on W_K is nondegenerate, we can write $W_K = \langle I, J \rangle$ for another isotropic K -line $J \subset V$. Hence $\mathbb{P}W$ is the line joining the two rational boundary points p_I and p_J .

If we choose one K -rational point as the base point of $H(I)$ and identify $H(I) \simeq V(I)_{\mathbb{C}}$ accordingly, the set of all K -rational points of $H(I)$ is identified with $V(I) \subset V(I)_{\mathbb{C}}$. The translation by $V(I)_{\mathbb{Z}}$ preserves this set.

Definition 5.1. A point of the boundary divisor $\Delta(I) = H(I)/V(I)_{\mathbb{Z}}$ is called a *special point* if it is the image of a K -rational point of $H(I)$.

If we choose one special point of $\Delta(I)$ as the origin and identify $\Delta(I) \simeq A(I)$ accordingly, the set of all special points of $\Delta(I)$ is identified with the set of torsion points of the abelian variety $A(I)$. For $\gamma \in \Gamma$, the induced map $\Delta(I) \rightarrow \Delta(\gamma I)$ sends special points of $\Delta(I)$ to those of $\Delta(\gamma I)$.

We have the following geometric characterization of special boundary points. We take another isotropic K -line $J \neq I \subset V$ and put $W = \langle I, J \rangle_{\mathbb{C}}$. The 1-dimensional ball $\mathcal{D} \cap \mathbb{P}W \subset \mathcal{D}$ joining p_I and p_J defines a congruence modular curve Y_W and a morphism $Y_W \rightarrow Y$. Let X_W be the compactification of Y_W . Note that I also defines a cusp of X_W . Since X_W is 1-dimensional and X^{tor} is compact, $Y_W \rightarrow Y$ extends to a morphism $X_W \rightarrow X^{tor}$. Recalling that $\mathcal{D} \cap \mathbb{P}W$ is just the ϕ -fiber over $[W] \in H(I)$, we find that the image of the I -cusp of X_W in X^{tor} is nothing but the point $[W] \in \Delta(I)$. Since every K -rational point of $H(I)$ can be written as $[\langle I, J \rangle_{\mathbb{C}}]$ for some isotropic K -line $J \subset V$, we obtain the following.

Lemma 5.2. *A point of $\Delta(I)$ is special if and only if it is the image of the I -cusp of X_W by $X_W \rightarrow X^{tor}$ for some 1-dimensional ball $\mathcal{D} \cap \mathbb{P}W$ joining p_I and another rational boundary point.*

5.3. Proof of Theorem 1.2. Now we can prove Theorem 1.2. Let P, Q be two special boundary points of X^{tor} . We want to show that $[P] - [Q]$ is torsion in $\text{CH}_0(X^{tor})$.

We first consider the case $\pi(P) = \pi(Q)$, say p_I . We take Q to be the base point of $\Delta(I)$ and identify $\Delta(I) = A(I)$ accordingly. Since $A(I)$ is isogenous to a self product of a CM elliptic curve E , there exists a homomorphism $f: E \rightarrow A(I)$ of abelian varieties such that $f(o_E) = Q$ and $f(p) = P$ for a point p of E of finite order. Since the map $E \rightarrow \text{Pic}^0(E)$, $x \mapsto [x] - [o_E]$, is an isomorphism of abelian groups, $[p] - [o_E] \in \text{Pic}^0(E)$ is torsion. Thus we find that $[P] - [Q] = f_*([p] - [o_E])$ is torsion in $\text{CH}_0(X^{tor})$.

Next let $\pi(P) \neq \pi(Q)$. Let $\pi(P) = p_I$ and $\pi(Q) = p_J$ for isotropic K -lines $I, J \subset V$ and consider the \mathbb{C} -plane $W = \langle I, J \rangle_{\mathbb{C}}$. As explained in §5.2, the 1-dimensional ball $\mathcal{D} \cap \mathbb{P}W$ defines a morphism $f: X_W \rightarrow X^{tor}$ from a compactified congruence modular curve X_W . Let q_I, q_J be the cusps of X_W defined by I, J respectively. Let $P' = f(q_I)$ and $Q' = f(q_J)$. By the Manin-Drinfeld theorem for X_W , we see that $[P'] - [Q']$ is torsion in $\text{CH}_0(X^{tor})$. By Lemma 5.2, P' and Q' are special points in $\Delta(I)$ and $\Delta(J)$ respectively. Therefore $[P] - [P']$ and $[Q] - [Q']$ are torsion by the preceding case. Thus $[P] - [Q]$ is torsion in $\text{CH}_0(X^{tor})$. Theorem 1.2 is proved. \square

By Theorem 1.2, the *rational* Chow class of special boundary points is uniquely determined. By Lemma 5.2, every intersection point of a modular curve X_W and a boundary divisor $\Delta(I)$ is a special point. Therefore

Corollary 5.3. *The intersection 0-cycle of any modular curve and any boundary divisor in the rational Chow group $\text{CH}_0(X^{tor})_{\mathbb{Q}}$ is contained in the span of the rational Chow class of special boundary points.*

This holds even in the integral Chow group $\text{CH}_0(X^{tor})$ when the first Betti number of X^{tor} happens to be zero, by the Roitman theorem [19].

Of course toroidal compactification can be constructed even when Γ is not necessarily neat. In that general case, the boundary divisor over a cusp p_I is the quotient of $H(I)/V(I)_{\mathbb{Z}}$ by the finite abelian group $(\Gamma \cap N(I)_{\mathbb{Q}})/W(I)_{\mathbb{Z}}$. Special boundary points can be defined similarly, as the image of K -rational points of $H(I)$. Obviously Theorem 1.2 still holds in this case.

Example 5.4. By Holzapfel [11], there is an example of Γ in $n = 2$ such that the minimal desingularization of X^{tor} is a $K3$ surface X_{K3} (of Picard number 20 and discriminant 3). By Roitman's theorem for X_{K3} , any two special boundary points in X_{K3} are rationally equivalent. Then the Chow class of

these points coincides with the Beauville-Voisin class of X_{K3} . This can be seen either by considering intersection cycle of modular curve and boundary curve, or from the fact that the boundary contains a rational component.

5.4. Cuspidal Chow class group. Theorem 1.2 motivates to consider the following Chow-theoretic analogue of cuspidal divisor class groups (cf. [12]) for Picard modular varieties. (A similar group is trivial for other modular varieties as in Theorem 1.1.)

Let \mathcal{C} be the subgroup of $\text{CH}_0(X^{\text{tor}})$ generated by the differences $[P] - [Q]$ of two special boundary points. By Theorem 1.2, \mathcal{C} is a torsion group. In order to describe \mathcal{C} more explicitly, let $X^{\text{tor}} - Y = \Delta_1 + \cdots + \Delta_N$ be the irreducible decomposition of the boundary. We choose a special point $P_i \in \Delta_i$ for each i and let $\alpha_i: X^{\text{tor}} \rightarrow \text{Alb}(X^{\text{tor}})$ be the Albanese map with base point P_i . Then $A_i = \alpha_i(\Delta_i)$ is a sub abelian variety of $\text{Alb}(X^{\text{tor}})$. Let $(A_i)_{\text{tor}}$ be the torsion part of A_i . By Roitman's theorem [19], the Albanese map induces an isomorphism

$$\mathcal{C} \rightarrow \langle (A_i)_{\text{tor}} + \alpha_1(P_i) \rangle_i \subset \text{Alb}(X^{\text{tor}}).$$

Therefore, to understand \mathcal{C} is essentially reduced to understanding the configuration of the shifted sub abelian varieties $A_i + \alpha_1(P_i)$ in $\text{Alb}(X^{\text{tor}})$. Note that the subgroup $\langle (A_i)_{\text{tor}} + \alpha_1(P_i) \rangle_i$ of $\text{Alb}(X^{\text{tor}})$ can also be written as $(\sum_i A_i)_{\text{tor}} + \langle \alpha_1(P_i) \rangle_i$. Thus the sub abelian variety $\sum_i A_i$ of $\text{Alb}(X^{\text{tor}})$ gives the divisible part of \mathcal{C} (isomorphic to $(\mathbb{Q}/\mathbb{Z})^a$), and the subgroup of $\text{Alb}(X^{\text{tor}})/\sum_i A_i$ generated by the torsion points $\alpha_1(P_i)$, $1 \leq i \leq N$, gives the finite abelian part of \mathcal{C} .

REFERENCES

- [1] Ash, A.; Mumford, D.; Rapoport, M.; Tai, Y. *Smooth compactifications of locally symmetric varieties*. 2nd edition. Cambridge Univ. Press, 2010.
- [2] Baily, W. L., Jr.; Borel, A. *Compactification of arithmetic quotients of bounded symmetric domains*. Ann. of Math. (2) **84** (1966), 442–528.
- [3] Bass, H.; Lazard, M.; Serre, J.-P. *Sous-groupes d'indices finis dans $\text{SL}(n, \mathbb{Z})$* . Bull. Amer. Math. Soc. **70** (1964) 385–392.
- [4] Beauville, A.; Voisin, C. *On the Chow ring of a K3 surface*. J. Alg. Geom. **13** (2004), 417–426.
- [5] Bergeron, N.; Li, Z.; Millson, J.; Moeglin, C. *The Noether-Lefschetz conjecture and generalizations*. Invent. Math. **208** (2017), no. 2, 501–552.
- [6] Drinfeld, V. G. *Two theorems on modular curves*. Funct. Anal. Appl. **7** (1973), no.2, 155–156.
- [7] van der Geer, G. *Hilbert modular surfaces*. Springer, 1988.
- [8] Harris, M. *Functorial properties of toroidal compactifications of locally symmetric varieties*. Proc. London Math. Soc. **59** (1989), 1–22.
- [9] Hirzebruch, F. *Chern numbers of algebraic surfaces: an example*. Math. Ann. **266** (1984), no. 3, 351–356.

- [10] Hoffman, J.W.; Weintraub, S. H. *The Siegel modular variety of degree two and level three*. Trans. Amer. Math. Soc. **353** (2000), 3267–3305.
- [11] Holzapfel, R.P. *Chern numbers of algebraic surfaces – Hirzebruch’s examples are Picard modular surfaces*. Math. Nachr. **126** (1986), 255–273.
- [12] Kubert, D.; Lang, S. *Modular units*. Springer, 1981.
- [13] Krieg, A. *Modular forms on half-spaces of quaternions*. Lecture Notes in Math. **1143**, Springer, 1985.
- [14] Ma, S. *Rational equivalence of cusps*. Ann. K-Theory **5** (2020), no.3, 395–410.
- [15] Manin, J. I. *Parabolic points and zeta functions of modular curves*. Math. USSR Izv. **6** (1972), no.1, 19–64.
- [16] Margulis, G. A. *Discrete subgroups of semisimple Lie groups*. Springer, 1991.
- [17] Mennicke, J. *Zur Theorie der Siegelschen Modulgruppe*. Math. Ann. **159** (1965), 115–129.
- [18] Piatetskii-Shapiro, I. *Geometry of classical domains and the theory of automorphic functions*. Gordon and Breach, 1969.
- [19] Roitman, A. A. *The torsion of the group of 0-cycles modulo rational equivalence*. Ann. of Math. (2) **111** (1980), no. 3, 553–569.
- [20] Sankaran, G. K. *Fundamental group of locally symmetric varieties*. Manuscripta Math. **90** (1996), no. 1, 39–48.
- [21] Serre, J.-P. *Le problème des groupes de congruence pour SL_2* . Ann. of Math. (2) **92** (1970), 489–527.
- [22] Shioda, T. *On elliptic modular surfaces*. J. Math. Soc. Japan **24** (1972), 20–59.
- [23] Weissauer, R. *Vektorwertige Siegelsche Modulformen kleinen Gewichtes*. J. Reine Angew. Math. **343** (1983), 184–202.

DEPARTMENT OF MATHEMATICS, TOKYO INSTITUTE OF TECHNOLOGY, TOKYO 152-8551, JAPAN
Email address: ma@math.titech.ac.jp