

Small Sample Spaces for Gaussian Processes

Toni Karvonen

The Alan Turing Institute, United Kingdom

October 7, 2021

Abstract

It is known that the membership in a given reproducing kernel Hilbert space (RKHS) of the samples of a Gaussian process X is controlled by a certain nuclear dominance condition. However, it is less clear how to identify a “small” set of functions (not necessarily a vector space) that contains the samples. This article presents a general approach for identifying such sets. We use *scaled RKHSs*, which can be viewed as a generalisation of Hilbert scales, to define the *sample support set* as the largest set which is contained in every element of full measure under the law of X in the σ -algebra induced by the collection of scaled RKHS. This potentially non-measurable set is then shown to consist of those functions that can be expanded in terms of an orthonormal basis of the RKHS of the covariance kernel of X and have their squared basis coefficients bounded away from zero and infinity, a result suggested by the Karhunen–Loève theorem.

1 Introduction

Let $K: T \times T \rightarrow \mathbb{R}$ be a positive-semidefinite kernel on a set T and consider any Gaussian process $(X(t))_{t \in T}$ with mean zero and covariance K , which we denote $(X(t))_{t \in T} \sim \mathcal{GP}(0, K)$. Let $H(K)$ be the reproducing kernel Hilbert space (RKHS) of K equipped with inner product $\langle \cdot, \cdot \rangle_K$ and norm $\|\cdot\|_K$. It is a well-known fact, apparently originating with Parzen (1963), that the samples of X are not contained in $H(K)$ if this space is infinite-dimensional. Furthermore, Driscoll (1973); Fortet (1973); and Lukić and Beder (2001) have used the zero-one law of Kallianpur (1970) to show essentially that, given another kernel R and under certain mild assumptions,

$$\mathbb{P}[X \in H(R)] = 1 \text{ if } R \gg K \quad \text{and} \quad \mathbb{P}[X \in H(R)] = 0 \text{ if } R \not\gg K,$$

where $R \gg K$ signifies that R *dominates* K (i.e., $H(K) \subset H(R)$) and moreover that the dominance is *nuclear* (see Section 4 for details). This *Driscoll’s theorem* is an exhaustive tool for verifying whether or not the samples from a Gaussian process are contained in a given RKHS. A review of the topic can be found in Chapter 4 of Gualtierotti (2015). Two questions now arise:

- How to construct a kernel R such that $R \gg K$?
- Is it possible to exploit the fact that $\mathbb{P}[X \in H(R_1) \setminus H(R_2)] = 1$ for any kernels such that $R_1 \gg K$ and $R_2 \not\gg K$ to identify in some sense the smallest set of functions which contains the samples with probability one?

Answers to questions such as these are instructive for theory and design of Gaussian process based learning (van der Vaart and van Zanten, 2011; Flaxman et al., 2016), emulation and approximation (Xu and Stein, 2017; Karvonen et al., 2020), and optimisation (Bull, 2011) methods. For simplicity we assume that the domain T is a complete separable metric space, that

the kernel K is continuous and its RKHS is separable, and that the samples of X are continuous. Although occasionally termed “rather restrictive” (Driscoll, 1973, p. 309), these continuity assumptions are satisfied by the vast majority of domains and Gaussian processes commonly used in statistics and machine learning literature (Stein, 1999; Rasmussen and Williams, 2006), such as stationary processes with Gaussian or Matérn covariance kernels. Our motivation for imposing these restrictions is that they imply that RKHSs are measurable.

First, this article presents the most flexible construction we are aware of for a kernel R such that $R \gg K$. For any orthonormal basis $\Phi = (\phi_n)_{n=1}^\infty$ of $H(K)$ the kernel can be written as $K(t, t') = \sum_{n=1}^\infty \phi_n(t)\phi_n(t')$ for all $t, t' \in T$. Given a positive sequence $A = (\alpha_n)_{n=1}^\infty$ such that $\sum_{n=1}^\infty \alpha_n \phi_n(t)^2 < \infty$ for all $t \in T$, we define the *scaled kernel*

$$K_{A,\Phi}(t, t') = \sum_{n=1}^\infty \alpha_n \phi_n(t)\phi_n(t'). \quad (1.1)$$

This is a significant generalisation of the concept of powers of kernels which has been previously used to construct RKHSs which contain the samples by Steinwart (2019). The sequence A is called a Φ -*scaling* of $H(K)$. If $\alpha_n \rightarrow \infty$ as $n \rightarrow \infty$, the corresponding *scaled RKHS*¹, $H(K_{A,\Phi})$, is a proper superset of $H(K)$. We show that convergence of the series $\sum_{n=1}^\infty \alpha_n^{-1}$ controls whether or not samples from X are contained in $H(K_{A,\Phi})$. If $\sum_{n=1}^\infty \alpha_n^{-1}$ converges slowly, we can therefore interpret $H(K_{A,\Phi})$ as a “small” RKHS which contains the samples.

Main Result I (Theorem 4.3). *Let $\Phi = (\phi_n)_{n=1}^\infty$ be an orthonormal basis of $H(K)$ and $A = (\alpha_n)_{n=1}^\infty$ a Φ -scaling of $H(K)$. If $K_{A,\Phi}$ is continuous and*

$$d_K(t, t') = \|K(\cdot, t) - K(\cdot, t')\|_K = \sqrt{K(t, t) - 2K(t, t') + K(t', t')}$$

is a metric on T , then either

$$\mathbb{P}[X \in H(K_{A,\Phi})] = 0 \quad \text{and} \quad \sum_{n=1}^\infty \frac{1}{\alpha_n} = \infty$$

or

$$\mathbb{P}[X \in H(K_{A,\Phi})] = 1 \quad \text{and} \quad \sum_{n=1}^\infty \frac{1}{\alpha_n} < \infty.$$

In Section 5, we use this result to study sample path properties of Gaussian processes defined by infinitely smooth kernels. These appear to be the first sufficiently descriptive results of their kind. For example, let $T \subset \mathbb{R}$ and $(X(t))_{t \in T} \sim \mathcal{GP}(0, K)$ for the univariate Gaussian kernel $K(t, t') = \exp(-(t - t')^2 / (2\ell^2))$ with length-scale $\ell > 0$. We explicitly construct several scaled kernels R whose RKHSs are “small” but still contain the samples of X . In Section 6, Theorem 4.3 is applied to provide an intuitive explanation for a conjecture by Xu and Stein (2017) on asymptotic behaviour of the maximum likelihood estimate of the scaling parameter of the Gaussian kernel when the data are generated by a monomial function on a uniform grid.

Secondly, we use Theorem 4.3 to construct a “small” set which functions as a nucleus for the samples in that it “almost” contains the samples. This *sample support set* is distinct from the traditional topological support of a Gaussian measure; see the discussion at the end of Section 2. Let $C(T)$ denote the set of continuous function on T .

Main Result II (Theorems 4.8 and 4.11). *Let $\Phi = (\phi_n)_{n=1}^\infty$ be an orthonormal basis of $H(K)$ and suppose there is a Φ -scaling $A = (\alpha_n)_{n=1}^\infty$ such that $\sum_{n=1}^\infty \alpha_n^{-1} < \infty$ and $K_{A,\Phi}$ is continuous. Let $\mathcal{S}(\mathfrak{R})$ be the σ -algebra generated by the collection of scaled RKHSs consisting*

¹These spaces are not to be confused with classical Hilbert scales defined via powers of a strictly positive self-adjoint operator; see Krein and Petunin (1966) and Engl et al. (1996, Section 8.4).

of continuous functions and $S_{\mathfrak{R}}(K)$ the largest subset of $C(T)$ that is contained in every $H \in \mathcal{S}(\mathfrak{R})$ such that $\mathbb{P}[X \in H] = 1$. Suppose that $d_K(t, t') = \|K(\cdot, t) - K(\cdot, t')\|_K$ is a metric on T . Then $S_{\mathfrak{R}}(K)$ consists precisely of the functions $f = \sum_{n=1}^{\infty} f_n \phi_n$ such that

$$\liminf_{n \rightarrow \infty} f_n^2 > 0 \quad \text{and} \quad \sup_{n \geq 1} f_n^2 < \infty. \quad (1.2)$$

Furthermore, for every $H \in \mathcal{S}(\mathfrak{R})$ such that $\mathbb{P}[X \in H] = 1$ there exists $F \in \mathcal{S}(\mathfrak{R})$ such that $S_{\mathfrak{R}}(K)$ is a proper subset of F and F is a proper subset of H .

The set $S_{\mathfrak{R}}(K)$ may fail to be measurable. The latter part of the above result is therefore important in demonstrating that it is possible to construct, using countably many elementary set operations of scaled RKHSs, sets which are arbitrarily close to $S_{\mathfrak{R}}(K)$ and contain the samples. At its core this is a manifestation of the classical result that there does not exist a meaningful notion of a boundary between convergent and divergent series (Knopp, 1951, § 41). The general form of the Karhunen–Loève theorem is useful in explaining the characterisation in (1.2). If $(\phi_n)_{n=1}^{\infty}$ is any orthonormal basis of $H(K)$, then the Gaussian process can be written as $X(t) = \sum_{n=1}^{\infty} \zeta_n \phi_n(t)$ for all $t \in T$, where ζ_n are independent standard normal random variables. In general the expansion converges in $L^2(\mathbb{P})$, but if almost all samples of the process are continuous, then convergence is also uniform on T with probability one (Adler, 1990, Theorem 3.8). Because $\|X(t)\|_K^2 = \sum_{n=1}^{\infty} \zeta_n^2$ and $\mathbb{E}[\zeta_n^2] = 1$ for every n , the Karhunen–Loève expansion suggests, somewhat informally speaking, that the samples can be found in the set of functions $f = \sum_{n=1}^{\infty} f_n \phi_n$ for which the sequence $(f_n^2)_{n=1}^{\infty}$ “is of constant order”.

1.1 On Measurability and Continuity

Suppose for a moment that T is an arbitrary uncountable set, K a positive-semidefinite kernel on T such that its RKHS $H(K)$ is infinite-dimensional, and $(X(t))_{t \in T} \sim \mathcal{GP}(0, K)$ a generic Gaussian process defined on a generic probability space $(\Omega, \mathcal{A}, \mathbb{P})$. Let \mathbb{R}^T be the collection of real-valued functions on T and $\tilde{\mathcal{B}}$ the σ -algebra generated by cylinder sets of the form $\{f \in \mathbb{R}^T : (f(t_1), \dots, f(t_n)) \in B^n\}$ any for $n \in \mathbb{N}$ and any Borel set $B^n \subset \mathbb{R}^n$. Let $\Phi_X(\omega) = X(\cdot, \omega)$. Then $\tilde{\mu}_X = \mathbb{P} \circ \Phi_X^{-1}$ is the law of X on the measurable space $(\mathbb{R}^T, \tilde{\mathcal{B}})$. Consequently,

$$\mathbb{P}(X \in H) = \tilde{\mu}(\{\omega \in \Omega : X(\cdot, \omega) \in H\})$$

for $H \in \tilde{\mathcal{B}}$. Let $(\mathbb{R}^T, \tilde{\mathcal{B}}_0, \tilde{\mu}_{X,0})$ be the completion of $(\mathbb{R}^T, \tilde{\mathcal{B}}, \tilde{\mu}_X)$ and $R: T \times T \rightarrow \mathbb{R}$ a positive-semidefinite kernel and $H(R)$ its RKHS. The following facts are known about measurability of $H(K)$ and $H(R)$:

- In general, $H(R) \notin \tilde{\mathcal{B}}$. For example, if T is equipped with a topology and R is continuous on $T \times T$, then $H(R) \subset C(T)$. However, no non-empty subset of $C(T)$ can be an element of $\tilde{\mathcal{B}}$.
- LePage (1973, p. 347) has proved that $H(K) \in \tilde{\mathcal{B}}_0$ and $\tilde{\mu}_{X,0}(H(K)) = 0$, a claim which originates with Parzen (1959, 1963). A less general version which requires separability and continuity has been proved by Kallianpur (1971).
- More generally, if $H(R)$ is infinite-dimensional and $R \not\gg K$, then $H(R) \in \tilde{\mathcal{B}}_0$ and $\tilde{\mu}_{X,0}(H(R)) = 0$. This result is contained in the proof of Theorem 7.3 in Lukić and Beder (2001). See also Proposition 4.5.1 in Gualtierotti (2015).
- LePage (1973, Corollary 2) has proved a dichotomy result which says that if $G \subset \mathbb{R}^T$ is an additive group and $G \in \tilde{\mathcal{B}}_0$, then either $\tilde{\mu}_{X,0}(G) = 0$ or $\tilde{\mu}_{X,0}(G) = 1$. Furthermore $\tilde{\mu}_{X,0}(G) = 1$ implies that $H(K) \subset G$. This is the most general version of the famous zero-one law of Kallianpur (1970, Theorem 2).

It appears that not much more can be said without imposing additional structure or constructing versions of X , as is done in Lukić and Beder (2001).

Suppose that (T, d_T) is a complete separable metric space, kernel K is continuous, and almost all samples of $(X(t))_{t \in T} \sim \mathcal{GP}(0, K)$ are continuous (i.e., the (\mathcal{B}_0, μ_0) -outer measure of $C(T)$ is one). Define the probability space $(C(T), \mathcal{B}, \mu_X)$ as

$$\mathcal{B} = C(T) \cap \tilde{\mathcal{B}}_0 \quad \text{and} \quad \mu_X(C(T) \cap H) = \tilde{\mu}_{X,0}(H) \quad \text{for } H \in \tilde{\mathcal{B}}_0. \quad (1.3)$$

The rest of this article is concerned with $(C(T), \mathcal{B}, \mu_X)$ and it is to be understood that $\mathbb{P}(X \in H)$ stands for $\mu_X(H)$ for any $H \in \mathcal{B}$. In this setting Driscoll (1973, p. 313) has proved that $H(R) \in \mathcal{B}$ if R is continuous and positive-definite. By using Theorem 1.1 in Fortet (1973) (or Lukić and Beder, 2001, Theorem 4.1) one can generalise this result to the case where R is continuous and positive-semidefinite; see the proof of Theorem 7.3 in Lukić and Beder (2001).

1.2 Notation and Terminology

Let F and G be two normed spaces. The space F is *continuously embedded* in G , denoted $F \hookrightarrow G$, if $F \subset G$ and there is a constant $C > 0$ such that $\|f\|_G \leq C \|f\|_F$ for every $f \in F$. If we have both $F \hookrightarrow G$ and $G \hookrightarrow F$, then the spaces are said to be *norm-equivalent*. For non-negative real sequences $(a_n)_{n=1}^\infty$ and $(b_n)_{n=1}^\infty$ we write $a_n \preceq b_n$ if there is $C > 0$ such that $a_n \leq C b_n$ for all sufficiently large n . If we have both $a_n \preceq b_n$ and $b_n \preceq a_n$, we write $a_n \asymp b_n$. If $a_n/b_n \rightarrow 1$ as $n \rightarrow \infty$, we write $a_n \sim b_n$. For two sets F and G we use $F \subsetneq G$ to indicate that F is a proper subset of G . A kernel $R: T \times T \rightarrow \mathbb{R}$ is *positive-semidefinite* if

$$\sum_{i=1}^N \sum_{j=1}^N a_i a_j R(t_i, t_j) \geq 0 \quad (1.4)$$

for any $N \geq 1$, $a_1, \dots, a_N \in \mathbb{R}$ and $t_1, \dots, t_N \in T$. In the remainder of this article positive-semidefinite kernels are simply referred to as *kernels*. If the inequality in (1.4) is strict, the kernel is said to be *positive-definite*.

1.3 Standing Assumptions

For ease of reference our standing assumptions are collected here. We assume that (i) (T, d_T) is a *complete separable metric space*; (ii) the covariance kernel $K: T \times T \rightarrow \mathbb{R}$ is *continuous* and *positive-semidefinite* on $T \times T$; (iii) the RKHS $H(K)$ induced by K is *infinite-dimensional* and *separable*²; and (iv) $(X(t))_{t \in T} \sim \mathcal{GP}(0, K)$ is a zero-mean Gaussian process on a probability space $(\Omega, \mathcal{A}, \mathbb{P})$ with *continuous paths*. The law μ_X of X is defined on the measurable space $(C(T), \mathcal{B})$ which was constructed in Section 1.1. Some of our results have natural generalisations for general second-order stochastic processes; see Lukić and Beder (2001), in particular Sections 2 and 5, and Steinwart (2019). These generalisations are not pursued here.

2 Related Work

Reproducing kernel Hilbert spaces which contain the samples of $(X(t))_{t \in T} \sim \mathcal{GP}(0, K)$ have been constructed by means of integrated kernels in Lukić (2004), convolution kernels in Cialenco et al. (2012) and Flaxman et al. (2016, Section 3.1), and, most importantly, *powers of RKHSs* (Steinwart and Scovel, 2012) in Steinwart (2019) and Kanagawa et al. (2018, Section 4). Namely, let T be a compact

²Most famously, separable RKHSs are induced by Mercer kernels, which are continuous kernels defined on compact subsets of \mathbb{R}^d (Paulsen and Raghupathi, 2016, Section 11.3).

metric space, K a continuous kernel on $T \times T$, and ν a finite and strictly positive Borel measure on T . Then the integral operator T_ν , defined for $f \in L^2(\nu)$ via

$$(T_\nu f)(t) = \int_T K(t, t') f(t') d\nu(t'), \quad (2.1)$$

has eigenvalues $(\lambda_n)_{n=1}^\infty$, which vanish as $n \rightarrow \infty$, and eigenfunctions $(\psi_n)_{n=1}^\infty \subset H(K)$ such that $(\sqrt{\lambda_n} \psi_n)_{n=1}^\infty$ is an orthonormal basis of $H(K)$. The kernel has the uniformly convergent Mercer expansion $K(t, t') = \sum_{n=1}^\infty \lambda_n \psi_n(t) \psi_n(t')$ for all $t, t' \in T$. For $\theta > 0$ the θ th power of K is defined as

$$K^{(\theta)}(t, t') = \sum_{n=1}^\infty \lambda_n^\theta \psi_n(t) \psi_n(t'), \quad (2.2)$$

which is well-defined if $\sum_{n=1}^\infty \lambda_n^\theta \psi_n(t)^2 < \infty$ for all $t \in T$. Furthermore, if $\theta_1 < \theta_2$, then $H(K^{(\theta_2)}) \subsetneq H(K^{(\theta_1)})$. It can be then proved that $\mathbb{P}[X \in H(K^{(\theta)})] = 1$ if and only if $\sum_{n=1}^\infty \lambda_n^{1-\theta} < \infty$ (Steinwart, 2019, Theorem 5.2). This construction has two significant downsides: (i) The measure ν is a *nuisance parameter*. If one is only interested in sample path properties of Gaussian processes this measure should not have an intrinsic part to play in the analysis and results. (ii) The construction is somewhat *inflexible* and *unsuitable for infinitely smooth kernels*. Because the power RKHS $H(K^{(\theta)})$ consists precisely of the functions $f = \sum_{n=1}^\infty f_n \lambda_n^{1/2} \psi_n$ such that $\sum_{n=1}^\infty f_n^2 \lambda_n^{1-\theta} < \infty$ and $\lambda_n \rightarrow 0$ as $n \rightarrow \infty$, how much larger $H(K^{(\theta)})$ is than $H(K)$, given by $\theta = 1$, is determined by rate of decay of the eigenvalues. Power RKHSs are more prescriptive when the kernel is finitely smooth and its eigenvalues have a polynomial decay (e.g., Matérn kernels) than when the kernel is infinitely smooth with exponential eigenvalue decay (e.g., Gaussian): the change from a polynomial decay condition $\sum_{n=1}^\infty f_n^2 n^{-a}$ for the coefficients $(f_n)_{n=1}^\infty$ to $\sum_{n=1}^\infty f_n^2 n^{-a(1-\theta)} < \infty$ is arguably more substantial than that from an exponential decay condition $\sum_{n=1}^\infty f_n^2 e^{-bn} < \infty$ to $\sum_{n=1}^\infty f_n^2 e^{-b(1-\theta)n} < \infty$. Indeed, as pointed out by Kanagawa et al. (2018, Section 4.4), when the kernel is Gaussian every $H(K^{(\theta)})$ with $\theta < 1$ contains the samples with probability one, which renders the concept of dubious utility in that setting. The relationship between powers of RKHSs and scaled RKHSs is discussed more at the end of Section 3. In Section 5 we demonstrate that scaled RKHSs are more useful in describing sample path properties of Gaussian processes defined by infinitely smooth kernels than powers of RKHSs.

To the best of our knowledge, the question about a “minimal” set which contains the samples with probability one has received only cursory discussion in the literature. Perhaps the most relevant digression on the topic is an observation by Steinwart (2019, pp. 369–370), given here in a somewhat applied form and without some technicalities, that the samples are contained in the set

$$\left(\bigcap_{r < s} W_2^r(T) \right) \setminus W_2^s(T) \quad (2.3)$$

with probability one if $H(K)$ is norm-equivalent to the fractional Sobolev space $W_2^{s+d/2}(T)$ for $s > 0$ on a suitable domain $T \subset \mathbb{R}^d$. In the Sobolev case the samples are therefore “ $d/2$ less smooth” than functions in the RKHS of K . Because $W_2^s(T) = \cup_{r \geq s} W_2^r(T)$, the set in (2.3) has the same form as the sample support set in (4.2). This observation is, of course, a general version of the familiar result that the sample paths of the Brownian motion, whose covariance kernel $K(t, t') = \min\{t, t'\}$ on $T = [0, 1]$ induces the Sobolev space $W_2^1([0, 1])$ with zero boundary condition at the origin, have regularity $1/2$ in the sense that they are almost surely α -Hölder continuous if and only if $\alpha < 1/2$. That is, there is $C > 0$ such that, for almost every $\omega \in \Omega$,

$$|X(t, \omega) - X(t', \omega)| \leq C |t - t'|^\alpha$$

for all $t, t' \in [0, 1]$ and any $\alpha < 1/2$. However, Lévy's modulus of continuity theorem (Mörsters and Peres, 2010, Section 1.2) improves this to

$$|X(t, \omega) - X(t', \omega)| \leq C\sqrt{h \log(1/h)}$$

when $h = |t - t'|$ is sufficiently small. Since the Sobolev space $W_2^s(T)$ consists of those functions $f: T \rightarrow \mathbb{R}$ which admit an $L^2(\mathbb{R}^d)$ -extension $f_e: \mathbb{R}^d \rightarrow \mathbb{R}$ whose Fourier transform satisfies

$$\int_{\mathbb{R}^d} (1 + \|\xi\|^2)^s |\widehat{f_e}(\xi)|^2 d\xi < \infty, \quad (2.4)$$

Lévy's modulus of continuity theorem suggests replacing the weight in (2.4) with, for example, $(1 + \|\xi\|^2)^s \log(1 + \|\xi\|)$ so that the resulting function space is a proper superset of $W_2^s(T)$ and a proper subset of $W_2^r(T)$ for every $r < s$ and hence a proper subset of the set in (2.3). Some results and discussion in Scheuerer (2010) and Karvonen et al. (2020, Section 4.2) have this flavour.

Finally, we remark that classical results about the topological support of a Gaussian measure are distinct from the results in this article. Let $C(T)$ be equipped with the standard supremum norm $\|\cdot\|_\infty$. The *topological support*, $\text{supp}_{C(T)}(\mu_X)$, of the measure μ_X is the set

$$\text{supp}_{C(T)}(\mu_X) = \{f \in C(T) : \mu_X(B(f, r)) > 0 \text{ for all } r > 0\},$$

where $B(f, r)$ is the f -centered r -ball in $(C(T), \|\cdot\|_\infty)$. It is a classical result that (Kallianpur, 1971, Theorem 3)

$$\text{supp}_{C(T)}(\mu_X) = \overline{H(K)},$$

where $\overline{H(K)}$ is the closure of $H(K)$ in $(C(T), \|\cdot\|_\infty)$. In other words, the topological support of μ_X contains every continuous function f such that, for every $\varepsilon > 0$, there exist $g_\varepsilon \in H(K)$ satisfying $\|f - g_\varepsilon\|_\infty < \varepsilon$. Now, recall that a kernel R is *universal* if $H(R)$ is dense in $C(T)$ (Steinwart and Christmann, 2008, Section 4.6). Most kernels of interest to practitioners are universal, including Gaussians, Matérns, and power series kernels. But, by definition, for a universal kernel the closure of the RKHS is the entirety of $C(T)$. Therefore

$$\text{supp}_{C(T)}(\mu_X) = \overline{H(K)} = C(T)$$

if K is a universal kernel. This result does not provide any information about the samples because we have assumed that the samples are continuous to begin with. See, for example, Bogachev (1998, Section 3.6) for further results on topological supports of Gaussian measures, including with respect to arbitrary topologies.

3 Scaled Reproducing Kernel Hilbert Spaces

Let $\Phi = (\phi_n)_{n=1}^\infty$ be any orthonormal basis of $H(K)$. Then the kernel has the pointwise convergent expansion

$$K(t, t') = \sum_{n=1}^{\infty} \phi_n(t)\phi_n(t') \quad \text{for all } t, t' \in T$$

and, by the standard characterisation of a separable Hilbert space, the RKHS consists of precisely those functions $f: T \rightarrow \mathbb{R}$ that admit an expansion

$$f = \sum_{n=1}^{\infty} f_n \phi_n \quad \text{for coefficients such that } \sum_{n=1}^{\infty} f_n^2 < \infty.$$

By the Cauchy–Schwarz inequality this expansion converges pointwise on T . For two functions $f = \sum_{n=1}^{\infty} f_n \phi_n$ and $g = \sum_{n=1}^{\infty} g_n \phi_n$ in the RKHS the inner product is $\langle f, g \rangle_K = \sum_{n=1}^{\infty} f_n g_n$.

Definition 3.1 (Scaled kernel and RKHS). We say that a positive sequence $A = (\alpha_n)_{n=1}^{\infty}$ is a Φ -scaling of $H(K)$ if

$$\sum_{n=1}^{\infty} \alpha_n \phi_n(t)^2 < \infty \quad \text{for every } t \in T. \quad (3.1)$$

The kernel

$$K_{A,\Phi}(t, t') = \sum_{n=1}^{\infty} \alpha_n \phi_n(t) \phi_n(t') \quad (3.2)$$

is called a *scaled kernel* and its RKHS $H(K_{A,\Phi})$ a *scaled RKHS*.

See Rakotomamonjy and Canu (2005); Xu and Zhang (2009); and Zhang and Zhao (2013) for prior appearances of scaled kernels under different names and not in the context of Gaussian processes.

Proposition 3.2. Let $\Phi = (\phi_n)_{n=1}^{\infty}$ be an orthonormal basis of $H(K)$ and $A = (\alpha_n)_{n=1}^{\infty}$ a Φ -scaling of $H(K)$. Then (i) the scaled kernel $K_{A,\Phi}$ is positive-semidefinite, (ii) the collection $(\sqrt{\alpha_n} \phi_n)_{n=1}^{\infty}$ is an orthonormal basis of $H(K_{A,\Phi})$, and (iii) the scaled RKHS is

$$\begin{aligned} H(K_{A,\Phi}) &= \left\{ f = \sum_{n=1}^{\infty} f_n \sqrt{\alpha_n} \phi_n : \|f\|_{K_{A,\Phi}}^2 = \sum_{n=1}^{\infty} f_n^2 < \infty \right\} \\ &= \left\{ f = \sum_{n=1}^{\infty} f_n \phi_n : \|f\|_{K_{A,\Phi}}^2 = \sum_{n=1}^{\infty} \frac{f_n^2}{\alpha_n} < \infty \right\}, \end{aligned} \quad (3.3)$$

where convergence is pointwise, and its inner product is

$$\langle f, g \rangle_{K_{A,\Phi}} = \sum_{n=1}^{\infty} \frac{f_n g_n}{\alpha_n} \quad (3.4)$$

for any $f = \sum_{n=1}^{\infty} f_n \phi_n$ and $g = \sum_{n=1}^{\infty} g_n \phi_n$ in $H(K_{A,\Phi})$.

Proof. By the Cauchy–Schwarz inequality and (3.1),

$$\sum_{n=1}^{\infty} |\alpha_n \phi_n(t) \phi_n(t')| \leq \left(\sum_{n=1}^{\infty} \alpha_n \phi_n(t)^2 \right)^{1/2} \left(\sum_{n=1}^{\infty} \alpha_n \phi_n(t')^2 \right)^{1/2}$$

for any $t, t' \in T$. This proves that the scaled kernel in (3.2) is well-defined via an absolutely convergent series. To verify that $K_{A,\Phi}$ is positive-semidefinite, note that, for any $N \geq 1$, $a_1, \dots, a_N \in \mathbb{R}$ and $t_1, \dots, t_N \in T$,

$$\sum_{i=1}^N \sum_{j=1}^N a_i a_j K_{A,\Phi}(t_i, t_j) = \sum_{n=1}^{\infty} \alpha_n \sum_{i=1}^N \sum_{j=1}^N a_i a_j \phi_n(t_i) \phi_n(t_j) = \sum_{n=1}^{\infty} \alpha_n \left(\sum_{i=1}^N a_i \phi_n(t_i) \right)^2$$

is non-negative because each α_n is positive. Because

$$\sum_{n=1}^{\infty} |f_n \sqrt{\alpha_n} \phi_n(t)| \leq \left(\sum_{n=1}^{\infty} f_n^2 \right)^{1/2} \left(\sum_{n=1}^{\infty} \alpha_n \phi_n(t)^2 \right)^{1/2} < \infty \quad \text{for every } t \in T$$

if $\sum_{n=1}^{\infty} f_n^2 < \infty$, the space defined in (3.3) and (3.4) is a Hilbert space of functions with an orthonormal basis $(\sqrt{\alpha_n} \phi_n)_{n=1}^{\infty}$. Since $K_{A,\Phi}(t, t') = \sum_{n=1}^{\infty} \alpha_n \phi_n(t) \phi_n(t')$, the scaled kernel is the unique reproducing kernel of this space (Minh, 2010, Theorem 9). \square

A scaled RKHS depends on the ordering of the orthonormal basis of $H(K)$ used in its construction. As an example, let $\Phi = (\phi_n)_{n=1}^\infty$ be an orthonormal basis of $H(K)$ and suppose that coefficients $\alpha_n = n$ define a Φ -scaling of $H(K)$. Define an orthonormal basis $\Psi = (\psi_n)_{n=1}^\infty$ by setting $\psi_{2n+1} = \phi_{2n}$ for $n \geq 0$ and interleaving the remaining ϕ_n to produce

$$\Psi = (\phi_1, \phi_3, \phi_2, \phi_5, \phi_4, \phi_6, \phi_8, \phi_7, \phi_{16}, \dots).$$

Now, the function $f = \sum_{n=0}^\infty \phi_{2n} = \sum_{n=1}^\infty f_{\Phi,n} \phi_n$ is in $H(K_{A,\Phi})$ because

$$\|f\|_{K_{A,\Phi}}^2 = \sum_{n=1}^\infty \frac{f_{\Phi,n}^2}{\alpha_n} = \sum_{n=0}^\infty \frac{1}{2^n} < \infty$$

but not in $H(K_{A,\Psi})$ because $f = \sum_{n=0}^\infty \phi_{2n} = \sum_{n=0}^\infty \psi_{2n+1} = \sum_{n=1}^\infty f_{\Psi,n} \psi_n$ and therefore

$$\|f\|_{K_{A,\Psi}}^2 = \sum_{n=1}^\infty \frac{f_{\Psi,n}^2}{\alpha_n} = \sum_{n=0}^\infty \frac{1}{2n+1} = \infty.$$

The following results compare sizes of scaled RKHSs: the faster α_n grows, the larger the RKHS $H(K_{A,\Phi})$ is. A number of additional properties between scaled RKHSs can be proved in a similar manner but are not needed in the developments of this article. See Xu and Zhang (2009, Section 6) and Zhang and Zhao (2013, Section 4) for Proposition 3.5 and some additional results.

Proposition 3.3. *Let $\Phi = (\phi_n)_{n=1}^\infty$ be an orthonormal basis of $H(K)$ and $A = (\alpha_n)_{n=1}^\infty$ and $B = (\beta_n)_{n=1}^\infty$ two Φ -scalings of $H(K)$. Then*

$$H(K_{B,\Phi}) \hookrightarrow H(K_{A,\Phi}) \quad \text{if and only if} \quad \beta_n \preceq \alpha_n.$$

In particular,

$$H(K) \hookrightarrow H(K_{A,\Phi}) \quad \text{if and only if} \quad \inf_{n \geq 1} \alpha_n > 0.$$

Proof. If $\beta_n \preceq \alpha_n$, then for any $f = \sum_{n=1}^\infty f_n \phi_n \in H(K_{B,\Phi})$ we have

$$\|f\|_{K_{A,\Phi}}^2 = \sum_{n=1}^\infty \frac{f_n^2}{\alpha_n} = \sum_{n=1}^\infty \frac{\beta_n}{\alpha_n} \frac{f_n^2}{\beta_n} \leq \|f\|_{K_{B,\Phi}}^2 \sup_{n \geq 1} \frac{\beta_n}{\alpha_n} < \infty$$

and thus $H(K_{B,\Phi}) \hookrightarrow H(K_{A,\Phi})$. Suppose then that $H(K_{B,\Phi}) \subset H(K_{A,\Phi})$ and assume to the contrary that $\sup_{n \geq 1} \alpha_n^{-1} \beta_n = \infty$ so that there is a subsequence $(n_m)_{m=1}^\infty$ such that $\alpha_{n_m}^{-1} \beta_{n_m} \geq 2^m$. Then $f = \sum_{m=1}^\infty 2^{-m/2} \sqrt{\beta_{n_m}} \phi_{n_m} \in H(K_{B,\Phi})$ since

$$\|f\|_{K_{B,\Phi}}^2 = \sum_{m=1}^\infty \left(\frac{\sqrt{\beta_{n_m}}}{2^{m/2}} \right)^2 \frac{1}{\beta_{n_m}} = \sum_{m=1}^\infty 2^{-m} = 1$$

but

$$\|f\|_{K_{A,\Phi}}^2 = \sum_{m=1}^\infty 2^{-m} \frac{\beta_{n_m}}{\alpha_{n_m}} \geq \sum_{m=1}^\infty 1 = \infty,$$

which contradicts the assumption that $H(K_{B,\Phi}) \subset H(K_{A,\Phi})$. Therefore $\sup_{n \geq 1} \alpha_n^{-1} \beta_n < \infty$. The second statement follows by setting $\beta_n = 1$ for every $n \in \mathbb{N}$ and noting that in this case $H(K_{B,\Phi}) = H(K)$. \square

Corollary 3.4. *Let $\Phi = (\phi_n)_{n=1}^\infty$ be an orthonormal basis of $H(K)$ and $A = (\alpha_n)_{n=1}^\infty$ and $B = (\beta_n)_{n=1}^\infty$ two Φ -scalings of $H(K)$. Then $H(K_{A,\Phi})$ and $H(K_{B,\Phi})$ are norm-equivalent if and only if $\alpha_n \asymp \beta_n$.*

Proposition 3.5. *Let $\Phi = (\phi_n)_{n=1}^\infty$ be an orthonormal basis of $H(K)$ and $A = (\alpha_n)_{n=1}^\infty$ a Φ -scaling of $H(K)$. Then a positive sequence $B = (\beta_n)_{n=1}^\infty$ is a Φ -scaling of $H(K)$ and $H(K_{B,\Phi}) \subsetneq H(K_{A,\Phi})$ whenever $\lim_{n \rightarrow \infty} \alpha_n \beta_n^{-1} = \infty$.*

Proof. By the assumption that $\lim_{n \rightarrow \infty} \alpha_n \beta_n^{-1} = \infty$ there is $C > 0$ such that $\beta_n \leq C \alpha_n$ for all $n \geq 1$. Therefore B is a Φ -scaling of $H(K)$. By Proposition 3.3, $H(K_{B,\Phi}) \subset H(K_{A,\Phi})$. Therefore $H(K_{B,\Phi})$ is a proper subset of $H(K_{A,\Phi})$ if $H(K_{A,\Phi})$ is not a subset of $H(K_{B,\Phi})$. Again, Proposition 3.3 gives that $H(K_{A,\Phi}) \subset H(K_{B,\Phi})$ if and only if $\alpha_n \preceq \beta_n$, which contradicts the assumption that $\lim_{n \rightarrow \infty} \alpha_n \beta_n^{-1} = \infty$. Hence $H(K_{B,\Phi})$ is a proper subset of $H(K_{A,\Phi})$. \square

Remark 3.6. Let $H(R)$ be another separable RKHS of functions on T . The RKHSs $H(K)$ and $H(R)$ are *simultaneously diagonalisable* if there exists an orthonormal basis $(\phi_n)_{n=1}^\infty$ of $H(K)$ which is an orthogonal basis of $H(R)$. That is, $(\|\phi_n\|_R^{-1} \phi_n)_{n=1}^\infty$ is an orthonormal basis of $H(R)$ and consequently $H(R) = H(K_{A,\Phi})$ for the scaling with $\alpha_n = \|\phi_n\|_R^{-2}$.

We conclude this section by demonstrating that scaled RKHSs generalise powers of RKHSs. We say that a Φ -scaling $A_\rho = (\alpha_n)_{n=1}^\infty$ of $H(K)$ is ρ -hyperharmonic if $\alpha_n = n^\rho$ for some $\rho \geq 0$. The RKHS $H(K_{A_\rho,\Phi})$ is a ρ -hyperharmonic scaled RKHS. Recall from Section 2 that if T is a compact metric space, K is continuous on $T \times T$, and ν is a finite and strictly positive Borel measure on T , then the integral operator in (2.1) has eigenfunctions $(\psi_n)_{n=1}^\infty$ and decreasing positive eigenvalues $(\lambda_n)_{n=1}^\infty$ and $\Psi = (\sqrt{\lambda_n} \psi_n)_{n=1}^\infty$ is an orthonormal basis of $H(K)$. For $\theta > 0$ the kernel $K^{(\theta)}(t, t') = \sum_{n=1}^\infty \lambda_n^\theta \psi_n(t) \psi_n(t')$ is a θ th power of K and its RKHS $H(K^{(\theta)})$ is a θ th power of $H(K)$. These objects are well-defined if $\sum_{n=1}^\infty \lambda_n^\theta \psi_n(t) < \infty$ for all $t \in T$. We immediately recognise that $K^{(\theta)}$ equals the scaled kernel $K_{A,\Psi}$ for the scaling $A = (\lambda_n^{\theta-1})_{n=1}^\infty$ because

$$K_{A,\Psi}(t, t') = \sum_{n=1}^\infty \lambda_n^{\theta-1} \lambda_n \psi_n(t) \psi_n(t') = \sum_{n=1}^\infty \lambda_n^\theta \psi_n(t) \psi_n(t') = K^{(\theta)}(t, t').$$

An important special case is when the eigenvalues have polynomial decay, $\lambda_n \asymp n^{-p}$ for some $p > 0$. For example, this holds, with $p = -2s/d$, if $T \subset \mathbb{R}^d$ and $H(K)$ is norm-equivalent to the Sobolev space $W_2^s(T)$ for $s > d/2$ (Steinwart, 2019, p. 370). If $\lambda_n \asymp n^{-p}$ and $\rho = p(1-\theta)$, the ρ -hyperharmonic scaled RKHS $H(K_{A_\rho,\Psi})$ is norm-equivalent to the θ th power RKHS $H(K^{(\theta)})$ by Corollary 3.4 because $n^\rho \lambda_n \asymp n^{-\theta p} \asymp \lambda_n^\theta$.

4 Sample Path Properties

A kernel R on T *dominates* K if $H(K) \subset H(R)$. In this case there exists (Lukić and Beder, 2001, Theorem 1.1) a unique linear operator $L: H(R) \rightarrow H(K)$, called the *dominance operator*, whose range is contained in $H(K)$ and which satisfies

$$\langle f, g \rangle_R = \langle Lf, g \rangle_K \quad \text{for all } f \in H(R) \text{ and } g \in H(K).$$

The dominance is said to be *nuclear*, denoted $R \gg K$, if $H(R)$ is separable and the operator L is nuclear, which is to say that

$$\text{tr}(L) = \sum_{n=1}^\infty \langle L\psi_n, \psi_n \rangle_R < \infty \quad (4.1)$$

for any orthonormal basis $(\psi_n)_{n=1}^\infty$ of $H(R)$.³

³A change of basis shows that $\text{tr}(L)$ does not depend on the orthonormal basis.

Let

$$d_R(t, t') = \|R(\cdot, t) - R(\cdot, t')\|_R = \sqrt{R(t, t) - 2R(t, t') + R(t', t')}.$$

This defines a pseudometric on T . If R is positive-definite, d_R is a metric. However, positive-definiteness is not necessary for d_R to be a metric. For example, the Brownian motion kernel $R(t, t') = \min\{t, t'\}$ on $T = [0, 1]$ is positive-semidefinite because $R(t, 0) = 0$ for any $t \in T$ but nevertheless yields a metric since $d_R(t, t') = \sqrt{t - 2\min\{t, t'\} + t'}$ vanishes if and only if $t = t'$. See Lukić and Beder (2001, Section 4) for more properties of d_R . An equivalent condition to d_R being a metric is to require that the mapping $t \mapsto R(\cdot, t)$ be injective.

According to the following theorem, a special case of the zero-one law of Kallianpur (Kallianpur, 1970; LePage, 1973) and a generalisation by Lukić and Beder (2001, Theorem 7.5) of an earlier result by Driscoll (1973, Theorem 3), the nuclear dominance condition determines whether or not samples of a Gaussian process $(X(t))_{t \in T} \sim \mathcal{GP}(0, K)$ lie in $H(R)$. In particular, the probability of them being in $H(R)$ is always either one or zero.

Theorem 4.1 (Driscoll–Lukić–Beder). *Let R be a continuous kernel on $T \times T$ with separable RKHS and $(X(t))_{t \in T} \sim \mathcal{GP}(0, K)$. If d_R is a metric, then either*

$$\mathbb{P}[X \in H(R)] = 0 \quad \text{and} \quad R \not\gg K$$

or

$$\mathbb{P}[X \in H(R)] = 1 \quad \text{and} \quad R \gg K.$$

Proof. Theorem 7.5 in Lukić and Beder (2001) is otherwise identical except that R is not assumed d_T -continuous and the samples of X are assumed d_R continuous. However, when R is d_T -continuous, d_T -continuity implies d_R -continuity. \square

Summability of the reciprocal scaling coefficients controls whether or not a scaled RKHS contains the sample paths.

Lemma 4.2. *Let $\Phi = (\phi_n)_{n=1}^\infty$ be an orthonormal basis of $H(K)$ and $A = (\alpha_n)_{n=1}^\infty$ a Φ -scaling of $H(K)$. Let $R = K_{A, \Phi}$. If d_K is a metric, then so is d_R .*

Proof. Because d_K is a metric,

$$d_K(t, t')^2 = K(t, t) - 2K(t, t') + K(t', t') = \sum_{n=1}^{\infty} [\phi_n(t) - \phi_n(t')]^2$$

vanishes if and only if $t = t'$. Since $d_R(t, t')^2 = \sum_{n=1}^{\infty} \alpha_n [\phi_n(t) - \phi_n(t')]^2$ and α_n are positive, we conclude that $d_R(t, t') = 0$ if and only if $t = t'$. \square

Theorem 4.3. *Let $\Phi = (\phi_n)_{n=1}^\infty$ be an orthonormal basis of $H(K)$, $A = (\alpha_n)_{n=1}^\infty$ a Φ -scaling of $H(K)$, and $(X(t))_{t \in T} \sim \mathcal{GP}(0, K)$. If $K_{A, \Phi}$ is continuous and d_K is a metric, then either*

$$\mathbb{P}[X \in H(R)] = 0 \quad \text{and} \quad \sum_{n=1}^{\infty} \frac{1}{\alpha_n} = \infty$$

or

$$\mathbb{P}[X \in H(R)] = 1 \quad \text{and} \quad \sum_{n=1}^{\infty} \frac{1}{\alpha_n} < \infty.$$

Proof. Assume first that the scaling is such that $H(K) \subset H(K_{A, \Phi})$. It is easy to verify using Proposition 3.2 that the unique dominance operator $L: H(K_{A, \Phi}) \rightarrow H(K)$ is given by

$$Lf = \sum_{n=1}^{\infty} \frac{f_n}{\alpha_n} \phi_n \quad \text{for any} \quad f = \sum_{n=1}^{\infty} f_n \phi_n \in H(K_{A, \Phi}).$$

Since $(\sqrt{\alpha_n}\phi_n)_{n=1}^\infty$ is an orthonormal basis of $H(K_{A,\Phi})$ and $L(\sqrt{\alpha_n}\phi_n) = \alpha_n^{-1/2}$, the nuclear dominance condition (4.1) is

$$\mathrm{tr}(L) = \sum_{n=1}^{\infty} \langle \sqrt{\alpha_n}L\phi_n, \sqrt{\alpha_n}\phi_n \rangle_{K_{A,\Phi}} = \sum_{n=1}^{\infty} \langle \phi_n, \phi_n \rangle_{K_{A,\Phi}} = \sum_{n=1}^{\infty} \frac{1}{\alpha_n},$$

and the claim thus follows from Theorem 4.1 since Lemma 4.2 guarantees that d_R is a metric when R is obtained by scaling K . Then, if $H(K) \not\subset H(K_{A,\Phi})$ it is trivially true that $K_{A,\Phi} \not\gg K$ and thus to complete the proof we have to show that this implies $\sum_{n=1}^{\infty} \alpha_n^{-1} = \infty$. If we had $\sum_{n=1}^{\infty} \alpha_n^{-1} < \infty$, then it would necessarily hold that $\sup_{n \geq 1} \alpha_n^{-1} < \infty$ and consequently $H(K) \subset H(K_{A,\Phi})$ by Proposition 3.3. Therefore $\sum_{n=1}^{\infty} \alpha_n^{-1} = \infty$. \square

4.1 Sample Support Sets

Theorems 4.1 and 4.3 motivate us to define the sample support set of a Gaussian process with respect to a collection of kernels as the largest set on the “boundary” between their induced RKHSs of probabilities one and zero. Let \mathfrak{R} be a collection of continuous kernels R on T for which d_R is a metric and $H(\mathfrak{R})$ the corresponding set of RKHSs. Every element of $H(\mathfrak{R})$ is a subset of $C(T)$. By the Driscoll–Lukić–Beder theorem each element of $H(\mathfrak{R})$ has μ_X -measure one or zero, depending on the nuclear dominance condition. Define the disjoint sets

$$\mathfrak{R}_1(K) = \{R \in \mathfrak{R} : R \gg K\} \quad \text{and} \quad \mathfrak{R}_0(K) = \{R \in \mathfrak{R} : R \not\gg K\}$$

which partition \mathfrak{R} . We assume that both $\mathfrak{R}_1(K)$ and $\mathfrak{R}_0(K)$ are non-empty.

Definition 4.4 (Sample support set). Let $\mathcal{S}(\mathfrak{R}) = \sigma(H(\mathfrak{R}))$ be the σ -algebra generated by $H(\mathfrak{R})$. The *sample support set*, $S_{\mathfrak{R}}(K)$, of the Gaussian process $(X(t))_{t \in T} \sim \mathcal{GP}(0, K)$ with respect to \mathfrak{R} is the largest subset of $C(T)$ such that

$$S_{\mathfrak{R}}(K) \subset H \quad \text{for every} \quad H \in \mathcal{S}(\mathfrak{R}) \quad \text{such that} \quad \mu_X(H) = 1.$$

Proposition 4.5. *It holds that*

$$S_{\mathfrak{R}}(K) = \bigcap_{R_1 \in \mathfrak{R}_1(K)} H(R_1) \setminus \bigcup_{R_0 \in \mathfrak{R}_0(K)} H(R_0). \quad (4.2)$$

Proof. Suppose that there is $f \in S_{\mathfrak{R}}(K)$ which is not contained in the set on the right-hand side of (4.2). That is, we have either that

$$f \notin \bigcap_{R_1 \in \mathfrak{R}_1(K)} H(R_1) \quad \text{or} \quad f \in \bigcup_{R_0 \in \mathfrak{R}_0(K)} H(R_0).$$

In the former case there is $R_1 \in \mathfrak{R}_1(K)$ such that $f \notin H(R_1)$. But because $\mu_X(H(R_1)) = 1$ and $S_{\mathfrak{R}}(K) \subset H(R_1)$ by definition, this violates the assumption that $f \in S_{\mathfrak{R}}(K)$. In the latter case there is $R_0 \in \mathfrak{R}_0(K)$ such that $f \in H(R_0)$. As $\mu_X(H(R_0)) = 0$, we have for any $R_1 \in \mathfrak{R}_1$ that $\mu_X(H(R_1) \setminus H(R_0)) = 1$. But since $f \notin H(R_1) \setminus H(R_0)$, the assumption that $f \in S_{\mathfrak{R}}(K)$ is again violated and we conclude that $S_{\mathfrak{R}}(K) \subset \bigcap_{R_1 \in \mathfrak{R}_1(K)} H(R_1) \setminus \bigcup_{R_0 \in \mathfrak{R}_0(K)} H(R_0)$.

Because all elements of $H(\mathfrak{R})$ are either of measure zero or one, so are those of $\mathcal{S}(\mathfrak{R})$. It is therefore clear that $\bigcap_{R_1 \in \mathfrak{R}_1(K)} H(R_1) \setminus \bigcup_{R_0 \in \mathfrak{R}_0(K)} H(R_0)$ is contained in every $H \in \mathcal{S}(\mathfrak{R})$ such that $\mu_X(H) = 1$. Consequently, $\bigcap_{R_1 \in \mathfrak{R}_1(K)} H(R_1) \setminus \bigcup_{R_0 \in \mathfrak{R}_0(K)} H(R_0) \subset S_{\mathfrak{R}}(K)$. This concludes the proof. \square

The sample support set is the largest set which is contained in every set of probability one under the law of X that can be expressed in terms of countably many elementary set operations of the RKHSs $H(R)$ for $R \in \mathfrak{R}$. If \mathfrak{R} is countable, the sample support set is in the σ -algebra \mathcal{B} , defined in (1.3), and has μ_X -measure one. However, when \mathfrak{R} is uncountable and does not contain countable subsets $\mathfrak{R}'_1(K) \subset \mathfrak{R}_1(K)$ and $\mathfrak{R}'_0(K) \subset \mathfrak{R}_0(K)$ such that

$$\bigcap_{R_1 \in \mathfrak{R}'_1(K)} H(R_1) = \bigcap_{R_1 \in \mathfrak{R}_1(K)} H(R_1) \quad \text{and} \quad \bigcup_{R_0 \in \mathfrak{R}'_0(K)} H(R_0) = \bigcup_{R_0 \in \mathfrak{R}_0(K)} H(R_0),$$

it cannot be easily determined if $S_{\mathfrak{R}}(K)$ is an element of \mathcal{B} . The above concepts assume substantially more concrete forms when \mathfrak{R} is a collection of scaled kernels.

Definition 4.6 (Approximately constant sequence). Let Σ be a collection of non-negative sequences. A non-negative sequence $(a_n)_{n=1}^{\infty}$ is said to be Σ -approximately constant if for every $(b_n)_{n=1}^{\infty} \in \Sigma$ the series $\sum_{n=1}^{\infty} b_n$ and $\sum_{n=1}^{\infty} a_n b_n$ either both converge or diverge.

For the RKHS $H(K)$ and any of its orthonormal bases $\Phi = (\phi_n)_{n=1}^{\infty}$ we let $\mathcal{R}(\Sigma, \Phi)$ denote the set of all functions $f = \sum_{n=1}^{\infty} f_n \phi_n$ such that the series converges pointwise on T and $(f_n^2)_{n=1}^{\infty}$ is a Σ -approximately constant sequence. The following theorem provides a crucial connection between sample sets with respect to scaled kernels and functions defined as orthonormal expansions with approximately constant coefficients.

Theorem 4.7. Let $\Phi = (\phi_n)_{n=1}^{\infty}$ be an orthonormal basis of $H(K)$ and Σ_{Φ} a collection of Φ -scalings of $H(K)$ such that the corresponding scaled kernels are continuous. Suppose that d_K is a metric and let $\mathfrak{R} = \{K_{A, \Phi} : A \in \Sigma_{\Phi}\}$. Then $S_{\mathfrak{R}}(K) = \mathcal{R}(\Sigma_{\Phi}, \Phi)$.

Proof. Note first that, by Lemma 4.2, d_R is a metric for every $R \in \mathfrak{R}$. Because every scaling of $H(K)$ has an orthonormal basis that is a scaled version of $(\phi_n)_{n=1}^{\infty}$, every $f \in S_{\mathfrak{R}}(K)$ can be written as $f = \sum_{n=1}^{\infty} f_n \phi_n$ for some real coefficients f_n . Let $\Sigma_1(K)$ and $\Sigma_0(K)$ stand for the collections $(\alpha_n)_{n=1}^{\infty} \in \Sigma_{\Phi}$ such that $\sum_{n=1}^{\infty} \alpha_n^{-1} < \infty$ and $\sum_{n=1}^{\infty} \alpha_n^{-1} = \infty$, respectively. Then, by Theorem 4.3, $K_{A, \Phi} \in \mathfrak{R}_1(K)$ if $A \in \Sigma_1(K)$ and $K_{A, \Phi} \in \mathfrak{R}_0(K)$ if $A \in \Sigma_0(K)$. Because, by definition, $S_{\mathfrak{R}}(K) \subset H(K_{A, \Phi})$ for any $A \in \Sigma_1(K)$ and $S_{\mathfrak{R}}(K) \cap H(K_{A, \Phi}) = \emptyset$ for any $A \in \Sigma_0(K)$ it follows that for every $f \in S_{\mathfrak{R}}(K)$ and any $(\alpha_n)_{n=1}^{\infty} \in \Sigma_{\Phi}$ we have

$$\sum_{n=1}^{\infty} \frac{f_n^2}{\alpha_n} < \infty \quad \text{and} \quad \sum_{n=1}^{\infty} \frac{1}{\alpha_n} < \infty \quad \text{or} \quad \sum_{n=1}^{\infty} \frac{f_n^2}{\alpha_n} = \infty \quad \text{and} \quad \sum_{n=1}^{\infty} \frac{1}{\alpha_n} = \infty.$$

That is, $(f_n^2)_{n=1}^{\infty}$ is a Σ_{Φ} -approximately constant sequence and thus $S_{\mathfrak{R}}(K) \subset \mathcal{R}(\Sigma_{\Phi}, \Phi)$. Conversely, if $f \in \mathcal{R}(\Sigma_{\Phi}, \Phi)$, then $f \in H(K_{A, \Phi})$ for every $A \in \Sigma_1(K)$ and $f \notin H(K_{A, \Phi})$ for every $A \in \Sigma_0(K)$. Hence $f \in S_{\mathfrak{R}}(K)$ and thus $S_{\mathfrak{R}}(K) = \mathcal{R}(\Sigma_{\Phi}, \Phi)$. \square

Next we use Theorem 4.7 to describe the sample set in more concrete terms.

4.2 Sample Support Sets for Scaled RKHSs

Let Σ be the set of all positive sequences. Then the collection of Σ -approximately constant sequences is precisely the collection of non-negative sequences $(a_n)_{n=1}^{\infty}$ such that

$$\liminf_{n \rightarrow \infty} a_n > 0 \quad \text{and} \quad \sup_{n \geq 1} a_n < \infty. \quad (4.3)$$

For suppose that there were a Σ -approximately constant sequence $(a_n)_{n=1}^{\infty}$ that violated (4.3). If $\liminf_{n \rightarrow \infty} a_n = 0$, then there is a subsequence $(a_{n_m})_{m=1}^{\infty}$ such that $a_{n_m} \leq 2^{-m}$ for all

$m \in \mathbb{N}$. Let $(b_n)_{n=1}^\infty \in \Sigma$ be a sequence such that $b_n = 2^{-n}a_n^{-1}$ for $n \notin (n_m)_{m=1}^\infty$ and $b_{n_m} = 1$ for $m \in \mathbb{N}$. Then $\sum_{n=1}^\infty b_n$ diverges but

$$\sum_{n=1}^\infty a_n b_n = \sum_{n \notin (n_m)_{m=1}^\infty} a_n b_n + \sum_{m=1}^\infty a_{n_m} b_{n_m} \leq \sum_{n \notin (n_m)_{m=1}^\infty} 2^{-n} + \sum_{m=1}^\infty 2^{-m} < \infty,$$

which contradicts the assumption that $(a_n)_{n=1}^\infty$ is a Σ -approximately constant sequence. An analogous argument with $a_{n_m} \geq 2^m$, $b_n = 2^{-n}$, and $b_{n_m} = 2^{-m}$ shows that the second condition in (4.3) cannot be violated either and thus every Σ -approximately constant sequence satisfies (4.3). It is trivial that a sequence satisfying (4.3) is Σ -approximately constant because the conditions imply the existence of positive constants $c_1 \leq c_2$ such that $c_1 \leq a_n \leq c_2$ for all sufficiently large n . This, together with Theorem 4.7, yields the following theorem which we consider the main result of this article. The proof of this theorem is somewhat more complicated because we cannot assume that every positive sequence is a scaling of $H(K)$.

Theorem 4.8. *Let $\Phi = (\phi_n)_{n=1}^\infty$ be an orthonormal basis of $H(K)$ and suppose that there is a Φ -scaling $A = (\alpha_n)_{n=1}^\infty$ of $H(K)$ such that $\sum_{n=1}^\infty \alpha_n^{-1} < \infty$ and $K_{A,\Phi}$ is continuous. Let Σ_Φ be the collection Φ -scalings of $H(K)$ such that the corresponding scaled kernels are continuous. Suppose that d_K is a metric and let $\mathfrak{R} = \{K_{A,\Phi} : A \in \Sigma_\Phi\}$. Then $S_{\mathfrak{R}}(K)$ is non-empty and consists precisely of the functions $f = \sum_{n=1}^\infty f_n \phi_n$ such that*

$$\liminf_{n \rightarrow \infty} f_n^2 > 0 \quad \text{and} \quad \sup_{n \geq 1} f_n^2 < \infty. \quad (4.4)$$

Proof. By Theorem 4.7, we know that $S_{\mathfrak{R}}(K) = \mathcal{R}(\Sigma_\Phi, \Phi)$ and the assumption that there exists a scaling $A = (\alpha_n)_{n=1}^\infty$ of $H(K)$ such that $\sum_{n=1}^\infty \alpha_n^{-1} < \infty$ implies that $\mathcal{R}(\Sigma_\Phi, \Phi)$ is non-empty. Hence we have to show that functions in $\mathcal{R}(\Sigma_\Phi, \Phi)$ satisfy (4.4). Suppose that there is a function $f = \sum_{n=1}^\infty f_n \phi_n \in \mathcal{R}(\Sigma_\Phi, \Phi)$ that violates (4.4). If $\sup_{n \geq 1} f_n^2 = \infty$, then there is a subsequence $(f_{n_m}^2)_{m=1}^\infty$ such that $f_{n_m}^2 \geq 2^m$ for all $m \in \mathbb{N}$. Define a sequence $B = (\beta_n)_{n=1}^\infty$ by setting $\beta_{n_m} = 2^{-m} \leq f_{n_m}^2$ for m such that $f_{n_m}^2 < \alpha_{n_m}$ and $\beta_n = \alpha_n$ for all other n . Then B is a Φ -scaling of $H(K)$ since $\beta_n \preceq \alpha_n$ and $\sum_{n=1}^\infty \beta_n^{-1} \leq \sum_{n=1}^\infty \alpha_n^{-1} + \sum_{m=1}^\infty 2^{-m} < \infty$. Moreover, Proposition 3.3 implies that $H(K_{B,\Phi}) \subset H(K_{A,\Phi})$ and it thus follows from the continuity of $K_{A,\Phi}$ that $K_{B,\Phi}$ is continuous, so that $B \in \Sigma_\Phi$. However,

$$\begin{aligned} \sum_{n=1}^\infty \frac{f_n^2}{\beta_n} &= \sum_{n \notin (n_m)_{m=1}^\infty} \frac{f_n^2}{\beta_n} + \sum_{\substack{m \in \mathbb{N} \\ f_{n_m}^2 \geq \alpha_{n_m}}} \frac{f_{n_m}^2}{\beta_{n_m}} + \sum_{\substack{m \in \mathbb{N} \\ f_{n_m}^2 < \alpha_{n_m}}} \frac{f_{n_m}^2}{\beta_{n_m}} \\ &= \sum_{n \notin (n_m)_{m=1}^\infty} \frac{f_n^2}{\alpha_n} + \sum_{\substack{m \in \mathbb{N} \\ f_{n_m}^2 \geq \alpha_{n_m}}} \frac{f_{n_m}^2}{\alpha_{n_m}} + \sum_{\substack{m \in \mathbb{N} \\ f_{n_m}^2 < \alpha_{n_m}}} \frac{f_{n_m}^2}{2^m} \\ &\geq \sum_{n \notin (n_m)_{m=1}^\infty} \frac{f_n^2}{\alpha_n} + \sum_{\substack{m \in \mathbb{N} \\ f_{n_m}^2 \geq \alpha_{n_m}}} 1 + \sum_{\substack{m \in \mathbb{N} \\ f_{n_m}^2 < \alpha_{n_m}}} 1 \\ &= \sum_{n \notin (n_m)_{m=1}^\infty} \frac{f_n^2}{\alpha_n} + \sum_{m=1}^\infty 1 \\ &= \infty, \end{aligned}$$

which contradicts the assumption that $(f_n^2)_{n=1}^\infty$ is Σ_Φ -approximately constant. On the other hand, if $\liminf_{n \rightarrow \infty} f_n^2 = 0$, then there is a subsequence $(f_{n_m}^2)_{m=1}^\infty$ such that $f_{n_m}^2 \leq 2^{-m}$ for all $m \in \mathbb{N}$. The sequence $B = (\beta_n)_{n=1}^\infty$ defined as $\beta_{n_m} = 1$ for $m \in \mathbb{N}$ and $\beta_n = \alpha_n f_n^2$ for

other n is a Φ -scaling of $H(K)$ and $K_{B,\Phi} \in \Sigma_\Phi$ because we have proved that $\sup_{n \geq 1} f_n^2 < \infty$. Furthermore, clearly $\sum_{n=1}^{\infty} \beta_n^{-1} = \infty$ but

$$\sum_{n=1}^{\infty} \frac{f_n^2}{\beta_n} = \sum_{n \notin (n_m)_{m=1}^{\infty}} \frac{f_n^2}{\beta_n} + \sum_{m=1}^{\infty} \frac{f_{n_m}^2}{\beta_{n_m}} \leq \sum_{n \notin (n_m)_{m=1}^{\infty}} \frac{1}{\alpha_n} + \sum_{m=1}^{\infty} 2^{-m} < \infty,$$

which again contradicts the assumption that $(f_n^2)_{n=1}^{\infty}$ is Σ_Φ -approximately constant. This completes the proof. \square

Theorem 4.11 below demonstrates that the sample support set of Theorem 4.8 cannot be expressed in terms of countably many elementary set operations of scaled RKHSs—and thus could be non-measurable. The reason for this is that there does not exist a useful notion of a “boundary” between convergent and divergent series, a result which is partially contained in Lemma 4.10. The lemma is a modified version of a result originally due to du Bois-Reymond and Hadamard (Knopp, 1951, pp. 301–302). In its proof a classical result by Dini (Knopp, 1951, p. 293) is needed.

Lemma 4.9 (Dini). *Let $(a'_n)_{n=1}^{\infty}$ be a positive sequence such that $\sum_{n=1}^{\infty} a'_n < \infty$ and set*

$$a_n = \frac{a'_n}{\left(\sum_{l=n}^{\infty} a'_l\right)^c}.$$

Then $\lim_{n \rightarrow \infty} (a'_n)^{-1} a_n = \infty$ and the series $\sum_{n=1}^{\infty} a_n$ converges if and only if $0 \leq c < 1$.

Lemma 4.10. *Suppose that $(a_{m,n})_{n=1}^{\infty}$ is a positive sequence such that $\sum_{n=1}^{\infty} a_{m,n} < \infty$ for each $m \in \mathbb{N}$. Then there is a positive sequence $(a_n)_{n=1}^{\infty}$ such that*

$$\sum_{n=1}^{\infty} a_n < \infty \quad \text{and} \quad \lim_{n \rightarrow \infty} \frac{a_n}{a_{m,n}} = \infty \quad \text{for every } m \in \mathbb{N}.$$

Proof. For each $m \in \mathbb{N}$, define the sequence $(\bar{a}_{m,n})_{n=1}^{\infty}$ by setting $\bar{a}_{m,n} = \sum_{k=1}^m a_{k,n}$, so that $\bar{a}_{m,n} \leq \bar{a}_{m+1,n}$ and $\sum_{n=1}^{\infty} \bar{a}_{m,n} < \infty$ for every m . Therefore there are n_m such that $n_m < n_{m+1}$ and $\sum_{n=n_m}^{\infty} \bar{a}_{m,n} \leq 2^{-m}$ for every $m \geq 2$. Define $a'_n = \bar{a}_{1,n}$ when $n < n_2$ and $a'_n = \bar{a}_{m,n}$ when $n_m \leq n < n_{m+1}$ for $m \geq 2$. Then

$$\sum_{n=1}^{\infty} a'_n = \sum_{n=1}^{n_2-1} \bar{a}_{1,n} + \sum_{m=2}^{\infty} \sum_{n=n_m}^{n_{m+1}-1} \bar{a}_{m,n} \leq \sum_{n=1}^{\infty} \bar{a}_{1,n} + \sum_{m=2}^{\infty} 2^{-m} < \infty$$

and, because $\bar{a}_{m,n} \leq \bar{a}_{m+1,n}$ and $\bar{a}_{m,n} \geq a_{m,n}$, we have $a'_n \geq \bar{a}_{m,n} \geq a_{m,n}$ for all m and $n \geq n_m$. Finally, selecting

$$a_n = \frac{a'_n}{\left(\sum_{l=n}^{\infty} a'_l\right)^{1/2}}$$

yields, by Lemma 4.9, a convergent series such that

$$\lim_{n \rightarrow \infty} \frac{a_n}{a_{m,n}} \leq \lim_{n \rightarrow \infty} \frac{a_n}{a'_n} = \lim_{n \rightarrow \infty} \left(\sum_{l=n}^{\infty} a'_l \right)^{-1/2} = \infty.$$

\square

Theorem 4.11. *Let $\Phi = (\phi_n)_{n=1}^{\infty}$ be an orthonormal basis of $H(K)$ and suppose that there is a Φ -scaling $A = (\alpha_n)_{n=1}^{\infty}$ of $H(K)$ such that $\sum_{n=1}^{\infty} \alpha_n^{-1} < \infty$ and $K_{A,\Phi}$ is continuous. Let Σ_Φ be the collection Φ -scalings of $H(K)$ such that the corresponding scaled kernels are continuous.*

Suppose that d_K is a metric and let $\mathfrak{K} = \{K_{A,\Phi} : A \in \Sigma_\Phi\}$. Let $A_m = (\alpha_{m,n})_{n=1}^\infty$ and $B_m = (\beta_{m,n})_{n=1}^\infty$ for $m \in \mathbb{N}$ be any elements of Σ_Φ such that

$$\sum_{n=1}^{\infty} \frac{1}{\alpha_{m,n}} < \infty \quad \text{and} \quad \sum_{n=1}^{\infty} \frac{1}{\beta_{m,n}} = \infty \quad \text{for every } m \in \mathbb{N}.$$

If $(X(t))_{t \in T} \sim \mathcal{GP}(0, K)$, then there exists $F \in \mathcal{B}$ such that $\mu_X(F) = 1$ and

$$S_{\mathfrak{K}}(K) \subsetneq F \subsetneq \bigcap_{m=1}^{\infty} H(K_{A_m, \Phi}) \setminus \bigcup_{m=1}^{\infty} H(K_{B_m, \Phi}).$$

Proof. By Lemma 4.10 there is a positive sequence $A' = (\alpha'_n)_{n=1}^\infty$ such that

$$\sum_{n=1}^{\infty} \frac{1}{\alpha'_n} < \infty \quad \text{and} \quad \lim_{n \rightarrow \infty} \frac{\alpha_{m,n}}{\alpha'_n} = \infty \quad \text{for every } m \in \mathbb{N}.$$

It follows from Proposition 3.5 that $H(K_{A', \Phi}) \subsetneq H(K_{A_m, \Phi})$ for every m and therefore $H(K_{A', \Phi}) \subset \bigcap_{m=1}^{\infty} H(K_{A_m, \Phi})$. Using Lemma 4.9 and Proposition 3.5 we can construct a Φ -scaling A such that $H(K_{A, \Phi}) \subsetneq H(K_{A', \Phi})$ and hence the set $F = H(K_{A, \Phi}) \setminus \bigcup_{m=1}^{\infty} H(K_{B_m, \Phi})$ has the desired properties by Theorem 4.3. \square

In Theorem 4.11 the set F was constructed by finding a scaled RKHS which is smaller than each $H(K_{A_m, \Phi})$ while still containing the sample set. A natural question is if we can also construct a scaled RKHS that contains each $H(K_{B_m, \Phi})$ but not the sample support set. Although Lemma 4.10 has a counterpart for divergent series, it seems difficult to guarantee that the resulting sequence is a scaling. Namely, it can be shown that if $(\beta_{m,n})_{n=1}^\infty$ in Theorem 4.11 are such that

$$\lim_{n \rightarrow \infty} \frac{\beta_{m,n}}{\beta_{m+1,n}} = 0 \quad \text{for every } m \in \mathbb{N}$$

then there exists $B = (\beta_n)_{n=1}^\infty$ such that

$$\sum_{n=1}^{\infty} \frac{1}{\beta_n} = \infty \quad \text{and} \quad \lim_{n \rightarrow \infty} \frac{\beta_{m,n}}{\beta_n} = 0 \quad \text{for every } m \in \mathbb{N}.$$

However, even though we know that there is a Φ -scaling of $H(K)$ such that $\sum_{n=1}^{\infty} \alpha_n^{-1} < \infty$, the fact that $\sum_{n=1}^{\infty} \beta_n^{-1}$ diverges does not necessarily imply that B is a Φ -scaling because *a divergent series can have terms that are arbitrarily larger than those of a convergent series* (Knopp, 1951, p. 303): there exists a positive sequence $(\gamma_n)_{n=1}^\infty$ such that

$$\sum_{n=1}^{\infty} \frac{1}{\gamma_n} < \infty \quad \text{and} \quad \liminf_{n \rightarrow \infty} \frac{\gamma_n}{\beta_n} = 0.$$

If $(\gamma_n)_{n=1}^\infty$ is a Φ -scaling of $H(K)$, we cannot directly deduce that so is B because $\beta_n \preceq \gamma_n$ does not hold.

4.3 Sample Support Sets for Monotonely Scaled RKHSs

In this section we demonstrate that the characterisation (4.4) fails if only *monotone* scalings are used. Let Σ_M be the collection of monotone Φ -scalings of $H(K)$ and \mathfrak{K}_M the collection of corresponding scaled kernels. Then the sample support set $S_{\mathfrak{K}_M}(K)$ is smaller than the sample support set $S_{\mathfrak{K}}(K)$ characterised by Theorem 4.8 because a Σ_M -approximately constant sequence can be unbounded, as long as it does not grow too rapidly, and contain arbitrarily long

sequences of zeros. Let $(g_m)_{m=0}^\infty$ be a strictly increasing sequence of positive integers and set $a_{g_m} = g_{m+1} - g_m$ for $m \geq 0$ and $a_n = 0$ otherwise. Then

$$\sum_{n=1}^{\infty} a_n b_n = \sum_{m=0}^{\infty} (g_{m+1} - g_m) b_{g_m}. \quad (4.5)$$

Schlömilch's generalisation of the Cauchy condensation test (Knopp, 1951, p. 121) implies that, under the assumption $g_{m+1} - g_m \leq C(g_m - g_{m-1})$ for some $C > 0$ and all $m \geq 1$, for any non-increasing positive sequence $(b_n)_{n=1}^\infty$ the series $\sum_{n=1}^\infty b_n$ and $\sum_{m=0}^\infty (g_{m+1} - g_m) b_{g_m}$ either both converge or diverge. That $(a_n)_{n=1}^\infty$ is Σ_M -approximately constant follows then from (4.5). The condition $g_{m+1} - g_m \leq C(g_m - g_{m-1})$, or equivalently $a_{g_m} \leq C a_{g_{m-1}}$, guarantees that a_{g_m} does not grow too fast in relation to g_m . The canonical example is obtained by setting $g_m = 2^m$ so that $a_{2^m} = 2^m$ and $C = 2$ and $\sum_{m=0}^\infty (g_{m+1} - g_m) b_{g_m} = \sum_{m=0}^\infty 2^m b_{2^m}$ converges or diverges with $\sum_{n=1}^\infty b_n$ by the standard Cauchy condensation test. Note that, because

$$\frac{1}{g_m} \sum_{n=1}^{g_m} a_n = \frac{1}{g_m} \sum_{k=0}^m (g_{k+1} - g_k) = \frac{g_{m+1} - g_0}{g_m} \leq \frac{g_{m+1}}{g_m} \leq \frac{(1+C)g_m}{g_m} = 1 + C$$

and

$$\frac{1}{g_m - 1} \sum_{n=1}^{g_m-1} a_n = \frac{1}{g_m - 1} \sum_{k=0}^{m-1} (g_{k+1} - g_k) = \frac{g_m - g_0}{g_m - 1},$$

the elements of the sequence $(a_n)_{n=1}^\infty$ are, on average, constants:

$$\liminf_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N a_n = 1 \quad \text{and} \quad \sup_{N \geq 1} \frac{1}{N} \sum_{n=1}^N a_n \leq 1 + C.$$

This can be seen as a weaker form of (4.4).

Although Σ_M contains unbounded sequences and sequences with zero lower limit, no sequence in this set can have infinity or zero as its limit. This can be shown by using results in Knopp (1951, § 41) which, given a sequence such that $\lim_{n \rightarrow \infty} a_n = \infty$, guarantee the existence of a monotone sequence $(c_n)_{n=1}^\infty$ such that $\sum_{n=1}^\infty c_n < \infty$ but $\sum_{n=1}^\infty a_n c_n = \infty$. Conversely, given a sequence such that $\lim_{n \rightarrow \infty} a_n = 0$ it is possible to construct a monotone sequence $(d_n)_{n=1}^\infty$ such that $\sum_{n=1}^\infty d_n = \infty$ but $\sum_{n=1}^\infty a_n d_n < \infty$.

5 Examples

This section contains examples of kernels to which Theorem 4.3 can be applied to construct “small” RKHSs that contain the samples and “slightly smaller” RKHSs that do not. For simplicity we let the domain T be a finite interval on the real line and occasionally index the orthonormal bases starting from zero. Note that to use our results one needs to have access to an orthonormal expansion of the kernel. Previous results based on powers of RKHSs and convolution kernels are easier to apply in this regard, but are less flexible and expressive. See Steinwart (2019) and Kanagawa et al. (2018, Section 4.4) for examples featuring powers of RKHSs (in particular for Sobolev kernels and the Gaussian kernel) and Flaxman et al. (2016, Appendices A.2 and A.3) for convolution kernel examples. Lukić (2004) has examples involving integrated kernels.

5.1 Iterated Brownian Bridge Kernels

Let $T = [0, 1]$ and consider the *iterated Brownian bridge kernel* of order $s \in \mathbb{N}$ (Cavoretto et al., 2015, Section 4.1) defined as

$$K^s(t, t') = \frac{2}{\pi^{2s}} \sum_{n=1}^{\infty} \frac{\sin(\pi n t) \sin(\pi n t')}{n^{2s}} = \sum_{n=1}^{\infty} \phi_n(t) \phi_n(t'), \quad (5.1)$$

where $\phi_n(t) = \sqrt{2}(\pi n)^{-sn} \sin(\pi n t)$. It can be shown that the kernel (5.1) has the representation

$$K^s(t, t') = (-1)^{s-1} \frac{2^{2s-1}}{(2s)!} \left[B_{2s} \left(\frac{|t-t'|}{2} \right) - B_{2s} \left(\frac{t+t'}{2} \right) \right],$$

where B_p is the Bernoulli polynomial of degree p . For $s = 1$ we obtain the Brownian bridge kernel

$$K^1(t, t') = \min\{t, t'\} - tt'.$$

For $s \geq 1$ and fixed $t' \in [0, 1]$ the translates $K^s(\cdot, t')$ are piecewise polynomials of order $2s - 1$.

Iterated Brownian bridge kernels are natural candidates for hyperharmonic scalings, as the 2ρ -hyperharmonic scaling $A_{2\rho} = (n^{2\rho})_{n=1}^{\infty}$ for $\rho \in \mathbb{N}$ such that $\rho \leq s - 1$ gives

$$K_{A_{2\rho}, \Phi}^s(t, t') = \frac{2}{\pi^{2s}} \sum_{n=1}^{\infty} \frac{\sin(\pi n t) \sin(\pi n t')}{n^{2(s-\rho)}} = \frac{1}{\pi^{2\rho}} K^{s-\rho}(t, t').$$

By Theorem 4.3, samples of $(X(t))_{t \in T} \sim \mathcal{GP}(0, K^s)$ are in the RKHS of this kernel for any such ρ because $\sum_{n=1}^{\infty} n^{-2\rho} < \infty$ whenever $\rho > 1/2$. Alternatively, using the identity $2 \int_0^1 \sin(\pi n t) \sin(\pi m t) dt = \delta_{nm}$ we see that the Mercer expansion with respect to the Lebesgue measure on $[0, 1]$ of K^s is

$$K^s(t, t') = \sum_{n=1}^{\infty} \lambda_n \psi_n(t) \psi_n(t') \quad \text{with} \quad \lambda_n = \frac{1}{(\pi n)^{2s}} \quad \text{and} \quad \psi_n(t) = \sqrt{2} \sin(\pi n t).$$

Therefore the θ th power of $H(K^s)$, with $\theta = 1 - \rho/s$, equals $H(K_{A_{2\rho}, \Phi}^s)$ as a set. In this instance the power RKHS is recovered as an instance of a scaled RKHS. We can also consider more exotic scalings, which do not correspond to power RKHSs, such as $A = (\alpha_n)_{n=1}^{\infty}$ for $\alpha_n = n \log(n+1)^2$ and $A' = (\alpha'_n)_{n=1}^{\infty}$ for $\alpha'_n = n \log(n+1)$. These yield the scaled kernels

$$K_{A, \Phi}^s(t, t') = \frac{2}{\pi^{2s}} \sum_{n=1}^{\infty} \frac{\sin(\pi n t) \sin(\pi n t')}{n^{2s-1} \log(n+1)^2} \quad \text{and} \quad K_{A', \Phi}^s(t, t') = \frac{2}{\pi^{2s}} \sum_{n=1}^{\infty} \frac{\sin(\pi n t) \sin(\pi n t')}{n^{2s-1} \log(n+1)},$$

which do not appear to have closed form expressions. Because

$$\sum_{n=1}^{\infty} \frac{1}{n \log(n+1)^2} < \infty \quad \text{but} \quad \sum_{n=1}^{\infty} \frac{1}{n \log(n+1)} = \infty,$$

the samples from $(X(t))_{t \in T} \sim \mathcal{GP}(0, K^s)$ are located in $H(K_{A, \Phi}^s)$ but not in $H(K_{A', \Phi}^s)$.

5.2 Gaussian Kernel

The ubiquitous Gaussian kernel with a length-scale parameter $\ell > 0$ is

$$K(t, t') = \exp\left(-\frac{(t-t')^2}{2\ell^2}\right) = \exp\left(-\frac{t^2}{2\ell^2}\right) \exp\left(-\frac{(t')^2}{2\ell^2}\right) \sum_{n=0}^{\infty} \frac{1}{\ell^{2n} n!} (tt')^n \quad (5.2)$$

and the functions

$$\phi_n(t) = \frac{1}{\ell^n \sqrt{n!}} t^n \exp\left(-\frac{t^2}{2\ell^2}\right) \quad \text{for } n \geq 0 \quad (5.3)$$

form an orthonormal basis of its RKHS (Steinwart et al., 2006; Minh, 2010).

The easiest way to proceed is to consider hyperharmonic scalings $A_\rho = (n^\rho)_{n=0}^\infty$ for $\rho > 0$ with the convention $0^0 = 1$. Then

$$K_{A_\rho, \Phi}(t, t') = \exp\left(-\frac{t^2 + (t')^2}{2\ell^2}\right) \left(1 + \sum_{n=1}^{\infty} \frac{n^\rho}{\ell^{2n} n!} (tt')^n\right).$$

For $\rho = 1$ we get a simple analytic expression for the scaled kernel:

$$\begin{aligned} K_{A_1, \Phi}(t, t') &= \exp\left(-\frac{t^2 + (t')^2}{2\ell^2}\right) \left(1 + \frac{tt'}{\ell^2} \sum_{n=0}^{\infty} \frac{1}{n!} \left(\frac{tt'}{\ell^2}\right)^n\right) \\ &= \exp\left(-\frac{t^2 + (t')^2}{2\ell^2}\right) \left(1 + \frac{tt'}{\ell^2} \exp\left(\frac{tt'}{\ell^2}\right)\right). \end{aligned} \quad (5.4)$$

Since the harmonic series $\sum_{n=1}^{\infty} n^{-1}$ is the prototypical example of an ‘‘almost’’ convergent series, we can informally say that the RKHS of (5.4) is only slightly too small to contain samples of the Gaussian process $(X(t))_{t \in T} \sim \mathcal{GP}(0, K)$ with covariance kernel (5.2). For a few larger values of ρ we compute that

$$\begin{aligned} K_{A_2, \Phi}(t, t') &= \exp\left(-\frac{t^2 + (t')^2}{2\ell^2}\right) [1 + (a + a^2) \exp(a)], \\ K_{A_3, \Phi}(t, t') &= \exp\left(-\frac{t^2 + (t')^2}{2\ell^2}\right) [1 + (a + 3a^2 + a^3) \exp(a)], \\ K_{A_4, \Phi}(t, t') &= \exp\left(-\frac{t^2 + (t')^2}{2\ell^2}\right) [1 + (a + 7a^2 + 6a^3 + a^4) \exp(a)], \end{aligned}$$

where $a = tt'/\ell^2$. By Theorem 4.3 the RKHSs of these kernels contain the samples since $\sum_{n=1}^{\infty} n^{-\rho} < \infty$ if and only if $\rho > 1$.

Another example can be constructed using the scaling $A = (\tau^{2n})_{n=0}^\infty$ for some $\tau > 0$. Then

$$K_{A, \Phi}(t, t') = \exp\left(-\frac{t^2 + (t')^2}{2\ell^2}\right) \sum_{n=0}^{\infty} \frac{1}{n!} \left(\frac{\tau^2 tt'}{\ell^2}\right)^n = \exp\left(-\frac{t^2 + (t')^2}{2\ell^2} + \frac{\tau^2 tt'}{\ell^2}\right).$$

The RKHS of this kernel contains the samples if and only if $\sum_{n=0}^{\infty} \tau^{-2n}$ converges, which is equivalent to $\tau > 1$. Since $K_{A, \Phi}$ equals the original kernel if $\tau = 1$, this class of scalings is not particularly expressive. Because the Mercer eigenvalues of the Gaussian kernel have an exponential decay this example is reminiscent (but more explicit) of the power RKHS construction for the Gaussian kernel in Section 4.4 of Kanagawa et al. (2018). Observe also that with the selection $\ell^2 = (1 - r^2)/r^2$ and $\tau^2 = 1/r$ for $r \in (0, 1)$ the kernel equals the scaled Mehler kernel

$$K_M^r(t, t') = \exp\left(-\frac{r^2(t^2 + (t')^2) - 2rtt'}{2(1 - r^2)}\right) = \sqrt{1 - r^2} \sum_{n=0}^{\infty} \frac{r^n}{n!} H_n(t) H_n(t'),$$

where H_n is the n th probabilists’ Hermite polynomial. The RKHS the Mehler kernel is analysed in Irgeher and Leobacher (2015).

5.3 Power Series Kernels

A *power series kernel* (Zwicky, 2009; Zwicky and Schaback, 2013) is a kernel of the form

$$K(t, t') = \sum_{n=0}^{\infty} \frac{w_n}{(n!)^2} (tt')^n \quad (5.5)$$

for positive coefficients w_n such that the series converges for all $t, t' \in T$. These kernels do not appear to have seen much use in the probabilistic setting, but are useful in functional analysis and approximation theory because, for particular choices of the coefficients, they reproduce important classical function spaces. For instance, the selection $w_n = (n!)^2$ yields the *Szegő kernel*

$$K(t, t') = \sum_{n=0}^{\infty} (tt')^n = \frac{1}{1 - tt'} \quad (5.6)$$

which converges as long as $T \subset (-1, 1)$ and whose complex extension is the reproducing kernel of the Hardy space \mathbb{H}_1^2 on the unit disc. The *exponential kernel*

$$K(t, t') = \sum_{n=0}^{\infty} \frac{1}{n!} (tt')^n = \exp(tt'), \quad (5.7)$$

obtained by setting $w_n = n!$, is a power series kernel that is convergent on the whole real line.

From the expansion in (5.5) we see that the functions

$$\phi_n(t) = \frac{\sqrt{w_n}}{n!} t^n \quad \text{for } n \geq 0$$

form an orthonormal basis of $H(K)$ and therefore a scaled kernel is

$$K_{A, \Phi}(t, t') = \sum_{n=0}^{\infty} \frac{\alpha_n w_n}{(n!)^2} (tt')^n,$$

which itself is a power series kernel. By Theorem 4.3 the samples from $(X(t))_{t \in T} \sim \mathcal{GP}(0, K)$ are in the RKHS of a different power series kernel

$$\bar{K}(t, t') = \sum_{n=0}^{\infty} \frac{\bar{w}_n}{(n!)^2} (tt')^n$$

if and only if

$$\sum_{n=0}^{\infty} \frac{w_n}{\bar{w}_n} < \infty.$$

For example, samples from a Gaussian process with the exponential covariance kernel (5.7) are in the RKHS of the Szegő kernel (5.6) if $T \subset (-1, 1)$ because

$$\sum_{n=0}^{\infty} \frac{w_n}{\bar{w}_n} = \sum_{n=0}^{\infty} \frac{n!}{(n!)^2} < \infty.$$

Its samples are *not* in the RKHS of the kernel

$$\bar{K}(t, t') = 1 + \sum_{n=1}^{\infty} \frac{n!n}{(n!)^2} (tt')^n = 1 + tt' \exp(tt'),$$

where $w_0 = 1$ and $w_n = n!n$ for $n \geq 1$, because

$$\sum_{n=0}^{\infty} \frac{w_n}{\bar{w}_n} = 1 + \sum_{n=1}^{\infty} \frac{1}{n} = \infty.$$

Note that this is essentially the same example as the one for the Gaussian kernel that involved the kernel (5.4).

6 Application to Maximum Likelihood Estimation

Gaussian processes are often used to model deterministic data-generating functions in, for example, design of computer experiments (Sacks et al., 1989) and probabilistic numerical computation (Diaconis, 1988; Cockayne et al., 2019). Conditioning the Gaussian process $(X(t))_{t \in T} \sim \mathcal{GP}(0, K)$, defined by a positive-definite kernel, on evaluations of a deterministic function $f: T \rightarrow \mathbb{R}$ at distinct points $t_1, \dots, t_N \in T$ gives rise to the conditional process

$$(X_{f,N}(t))_{t \in T} \sim \mathcal{GP}(I_{f,N}, K_{f,N})$$

whose mean and covariance functions have simple linear-algebraic expressions in terms of evaluations of the function f and the kernel K . The conditional mean is the unique minimum-norm interpolant in the RKHS of K :

$$I_{f,N} = \arg \min_{g \in H(K)} \{ \|g\|_K : g(t_i) = f(t_i) \text{ for every } i = 1, \dots, N \}.$$

The conditional covariance $K_{f,N}$ depends only on K and the points t_i and therefore lacks the data-dependence that one would hope for when quantifying one's uncertainty in $f(t)$ at new points $t \in T$. One of the simplest ways to remedy this shortcoming is to introduce a positive scaling parameter σ and employ the model $(X(t))_{t \in T} \sim \mathcal{GP}(0, \sigma^2 K)$. Then the conditional process is $(X_{f,N}(t))_{t \in T} \sim \mathcal{GP}(I_{f,N}, \sigma^2 K_{f,N})$ and the conditional covariance becomes data-dependent if the scaling parameter is selected based on the data. A popular approach is to use the maximum likelihood estimate $\hat{\sigma}_{f,N}$ which has the explicit form (Karvonen et al., 2020, Section 3.3)

$$\hat{\sigma}_{f,N} = \frac{\|I_{f,N}\|_K}{\sqrt{N}}. \quad (6.1)$$

It is then natural to ask how $\hat{\sigma}_{f,N}$ behaves for different f and different sets of points. One would expect that $\hat{\sigma}_{f,N}$ converge to a constant if f is akin to a sample from X and the points $\{t_1, \dots, t_N\}$ cover T sufficiently well.

When $H(K)$ is norm-equivalent to a Sobolev space Karvonen et al. (2020) have used results in the scattered data approximation literature to argue that the maximum likelihood estimate decays to zero if f is too regular to be a sample from X and explodes if f is too irregular. We now use Theorem 4.3 to provide a similar explanation for the behaviour of $\hat{\sigma}_{f,N}$ that was observed and conjectured by Xu and Stein (2017) when K is the Gaussian kernel (5.2). Let $T = [0, 1]$ and consider the uniform points $t_i = i/N$ for $i = 1, \dots, N$. Xu and Stein (2017) used explicit Cholesky decomposition formulae found in Loh and Kam (2000) to prove that, for certain positive constants $C_{\ell,0}$ and $C_{\ell,1}$,

$$\hat{\sigma}_{f,N}^2 \sim C_{\ell,0} N^{-1/2} \quad \text{as } N \rightarrow \infty$$

if $f \equiv 1$ and

$$\liminf_{N \rightarrow \infty} N^{-1/2} \hat{\sigma}_{f,N}^2 \geq C_{\ell,1}$$

if $f(t) = t$. Furthermore, they conjectured that

$$\hat{\sigma}_{f,N}^2 \sim C_{\ell,p} N^{p-1/2} \quad \text{as } N \rightarrow \infty$$

if $f(t) = t^p$ for any $p \geq 0$ and a positive constant $C_{\ell,p}$.⁴ In particular, their results show that $\hat{\sigma}_{f,N} \rightarrow 0$ if $f \equiv 1$ and $\hat{\sigma}_{f,N} \rightarrow \infty$ if $f(t) = t$. We now employ a technique similar to that

⁴As pointed out by Dette and Zhigljavsky (2021, Section 4.1), it appears that the constant Xu and Stein give, $C_{\ell,p} = \ell^{2p}/(\sqrt{2\pi}(p+1/2))$, is erroneous. By truncating the expansion (6.2) after N terms and reasoning that the RKHS norm of the resulting function should be asymptotically equal to $\|I_{f,N}\|_K$ one arrives, after some relatively straightforward computations, to the conjecture that the constant is $C_{\ell,p} = 2^p \ell^{2p}/(\sqrt{\pi}(p+1/2))$.

used in Minh (2010) to demonstrate that this is due to the fact that the constant function is too “regular” to be a sample from the Gaussian process, while the linear function and higher-order monomials are too “irregular”. Xu and Stein (2017, p. 142) lacked an intuitive explanation for these results.

Any monomial can be written in terms of the orthonormal basis (5.3) as

$$\begin{aligned} t^p &= \exp\left(\frac{t^2}{2\ell^2}\right) t^p \exp\left(-\frac{t^2}{2\ell^2}\right) = \sum_{n=0}^{\infty} \frac{t^{2n+p}}{2^n \ell^{2n} n!} \exp\left(-\frac{t^2}{2\ell^2}\right) \\ &= \sum_{n=0}^{\infty} \frac{\ell^p \sqrt{(2n+p)!}}{2^n n!} \phi_{2n+p}(t). \end{aligned} \quad (6.2)$$

Let $A = (\alpha_n)_{n=1}^{\infty}$ be a Φ -scaling of $H(K)$. If the p th monomial is to be an element of the scaled RKHS $H(K_{A,\Phi})$ it must therefore hold that

$$\sum_{n=0}^{\infty} \alpha_{2n+p}^{-1} \frac{(2n+p)!}{2^{2n} (n!)^2} < \infty. \quad (6.3)$$

Stirling’s formula gives that, as $n \rightarrow \infty$,

$$\begin{aligned} \frac{(2n+p)!}{2^{2n} (n!)^2} &\sim \frac{\sqrt{2\pi}(2n+p)^{2n+p+1/2} e^{-(2n+p)}}{2\pi 2^{2n} n^{2n+1} e^{-2n}} = \frac{1}{\sqrt{2\pi} e^p} \frac{(2n+p)^{2n+p+1/2}}{(2n)^{2n}} \\ &= \frac{1}{\sqrt{2\pi} e^p} \left(1 + \frac{p}{2n}\right)^{2n} \frac{(2n+p)^{p+1/2}}{n}. \end{aligned}$$

Since $\lim_{n \rightarrow \infty} \left(1 + \frac{p}{2n}\right)^{2n} = e^p$, we conclude that

$$\frac{(2n+p)!}{2^{2n} (n!)^2} \sim \frac{2^p}{\sqrt{\pi}} n^{p-1/2} \quad \text{as } n \rightarrow \infty.$$

Therefore (6.3) holds if and only if $\sum_{n=1}^{\infty} \alpha_{2n+p}^{-1} n^{p-1/2} < \infty$. If we constraint ourselves to hyperharmonic scalings this is clearly equivalent to $\alpha_n = n^\rho$ for $\rho > p + 1/2$. But because $\sum_{n=1}^{\infty} n^{-\rho} < \infty$ if and only if $\rho > 1$ it follows that (a) if $p < 1/2$, then $f(t) = t^p$ is an element of a scaled RKHS $H(K_{A,\Phi})$ such that $\mathbb{P}[X \in H(K_{A,\Phi})] = 0$ and (b) if $p \geq 1/2$, then $f(t) = t^p$ is an element of a hyperharmonic scaled RKHS $H(K_{A,\Phi})$ if and only if $\mathbb{P}[X \in H(K_{A,\Phi})] = 1$. In other words, the constant function is too close to $H(K)$, or too “regular”, to be a sample while higher-order monomials are too far from $H(K)$, or too “irregular”, and this provides some intuition behind the difference in asymptotic behaviour of maximum likelihood estimates for these functions.

Acknowledgements

The author was supported by the Lloyd’s Register Foundation Programme for Data-Centric Engineering at the Alan Turing Institute, United Kingdom. I am grateful to Motonobu Kanagawa for his patience in enduring persistent questions over the course of a conference in Sydney in July 2019 during which the main ideas underlying this article were first conceived and to Chris Oates for extensive commentary on the draft at various stages of preparation. Discussions with Anatoly Zhigljavsky led to improvements in Section 6.

References

Adler, R. J. (1990). *An Introduction to Continuity, Extrema, and Related Topics for General Gaussian Processes*. Number 12 in Lecture Notes–Monograph Series. Institute of Mathematical Statistics.

- Bogachev, V. I. (1998). *Gaussian Measures*. Number 62 in Mathematical Surveys and Monographs. American Mathematical Society.
- Bull, A. D. (2011). Convergence rates of efficient global optimization algorithms. *Journal of Machine Learning Research*, 12:2879–2904.
- Cavoretto, R., Fasshauer, G. E., and McCourt, M. (2015). An introduction to the Hilbert-Schmidt SVD using iterated Brownian bridge kernels. *Numerical Algorithms*, 68:393–422.
- Cialenco, I., Fasshauer, G. E., and Ye, Q. (2012). Approximation of stochastic partial differential equations by a kernel-based collocation method. *International Journal of Computer Mathematics*, 89(18):2543–2561.
- Cockayne, J., Oates, C., Sullivan, T., and Girolami, M. (2019). Bayesian probabilistic numerical methods. *SIAM Review*, 61(4):756–789.
- Dette, H. and Zhigljavsky, A. (2021). Reproducing kernel Hilbert spaces, polynomials and the classical moment problems. *arXiv:2101.11968v2*.
- Diaconis, P. (1988). Bayesian numerical analysis. In *Statistical decision theory and related topics IV*, volume 1, pages 163–175. Springer-Verlag New York.
- Driscoll, M. F. (1973). The reproducing kernel Hilbert space structure of the sample paths of a Gaussian process. *Zeitschrift für Wahrscheinlichkeitstheorie und Verwandte Gebiete*, 26(4):309–316.
- Engl, H. W., Hanke, M., and Neubauer, A. (1996). *Regularization of Inverse Problems*, volume 375 of *Mathematics and Its Applications*. Springer.
- Flaxman, S., Sejdinovic, D., Cunningham, J., and Filippi, S. (2016). Bayesian learning of kernel embeddings. In *Uncertainty in Artificial Intelligence*, pages 182–191.
- Fortet, R. (1973). Espaces à noyau reproduisant et lois de probabilités des fonctions aléatoires. *Annales de l’Institut Henri Poincaré, section B*, 9(1):41–58.
- Gualtierotti, A. F. (2015). *Detection of Random Signals in Dependent Gaussian Noise*. Springer.
- Irrgeher, C. and Leobacher, G. (2015). High-dimensional integration on \mathbb{R}^d , weighted Hermite spaces, and orthogonal transforms. *Journal of Complexity*, 31(2):174–205.
- Kallianpur, G. (1970). Zero-one laws for Gaussian processes. *Transactions of the American Mathematical Society*, 149:199–211.
- Kallianpur, G. (1971). Abstract Wiener processes and their reproducing kernel Hilbert spaces. *Zeitschrift für Wahrscheinlichkeitstheorie und Verwandte Gebiete*, 17:113–123.
- Kanagawa, M., Hennig, P., Sejdinovic, D., and Sriperumbudur, B. K. (2018). Gaussian processes and kernel methods: A review on connections and equivalences. *arXiv:1807.02582v1*.
- Karvonen, T., Wynne, G., Tronarp, F., Oates, C. J., and Särkkä, S. (2020). Maximum likelihood estimation and uncertainty quantification for Gaussian process approximation of deterministic functions. *SIAM/ASA Journal on Uncertainty Quantification*, 8(3):926–958.
- Knopp, K. (1951). *Theory and Application of Infinite Series*. Blackie & Son, 2nd edition.
- Krein, S. G. and Petunin, Yu. I. (1966). Scales of Banach spaces. *Russian Mathematical Surveys*, 21(2):85–159.

- LePage, R. (1973). Subgroups of paths and reproducing kernels. *The Annals of Probability*, 1(2):345–347.
- Loh, W.-L. and Kam, T.-K. (2000). Estimating structured correlation matrices in smooth Gaussian random field models. *Annals of Statistics*, 28(3):880–904.
- Lukić, M. N. (2004). Integrated Gaussian processes and their reproducing kernel Hilbert spaces. In *Stochastic Processes and Functional Analysis: A Volume of Recent Advances in Honor of M. M. Rao*, number 238 in Lecture Notes in Pure and Applied Mathematics, pages 241–263. Marcel Dekker.
- Lukić, M. N. and Beder, J. H. (2001). Stochastic processes with sample paths in reproducing kernel Hilbert spaces. *Transactions of the American Mathematical Society*, 353(10):3945–3969.
- Minh, H. Q. (2010). Some properties of Gaussian reproducing kernel Hilbert spaces and their implications for function approximation and learning theory. *Constructive Approximation*, 32(2):307–338.
- Mörters, P. and Peres, Y. (2010). *Brownian Motion*. Number 30 in Cambridge Series in Statistical and Probabilistic Mathematics. Cambridge University Press.
- Parzen, E. (1959). Statistical inference on time series by Hilbert space methods, I. Technical Report 23, Applied Mathematics and Statistics Laboratory, Stanford University.
- Parzen, E. (1963). Probability density functionals and reproducing kernel Hilbert spaces. In *Proceedings of the Symposium on Time Series Analysis*, pages 155–169. John Wiley & Sons.
- Paulsen, V. I. and Raghupathi, M. (2016). *An Introduction to the Theory of Reproducing Kernel Hilbert Spaces*. Number 152 in Cambridge Studies in Advanced Mathematics. Cambridge University Press.
- Rakotomamonjy, A. and Canu, S. (2005). Frames, reproducing kernels, regularization and learning. *Journal of Machine Learning Research*, 6:1485–1515.
- Rasmussen, C. E. and Williams, C. K. I. (2006). *Gaussian Processes for Machine Learning*. Adaptive Computation and Machine Learning. MIT Press.
- Sacks, J., Welch, W. J., M. T. J., and Wynn, H. P. (1989). Design and analysis of computer experiments. *Statistical Science*, 4(4):409–435.
- Scheuerer, M. (2010). Regularity of the sample paths of a general second order random field. *Stochastic Processes and their Applications*, 120(10):1879–1897.
- Stein, M. L. (1999). *Interpolation of Spatial Data: Some Theory for Kriging*. Springer Series in Statistics. Springer.
- Steinwart, I. (2019). Convergence types and rates in generic Karhunen-Loève expansions with applications to sample path properties. *Potential Analysis*, 51:361–395.
- Steinwart, I. and Christmann, A. (2008). *Support Vector Machines*. Information Science and Statistics. Springer.
- Steinwart, I., Hush, D., and Scovel, C. (2006). An explicit description of the reproducing kernel Hilbert spaces of Gaussian RBF kernels. *IEEE Transactions on Information Theory*, 52(10):4635–4643.
- Steinwart, I. and Scovel, C. (2012). Mercer’s theorem on general domains: On the interaction between measures, kernels, and RKHSs. *Constructive Approximation*, 35:363–417.

- van der Vaart, A. and van Zanten, H. (2011). Information rates of nonparametric Gaussian process methods. *Journal of Machine Learning Research*, 12(6):2095–2119.
- Xu, W. and Stein, M. L. (2017). Maximum likelihood estimation for a smooth Gaussian random field model. *SIAM/ASA Journal on Uncertainty Quantification*, 5(1):138–175.
- Xu, Y. and Zhang, H. (2009). Refinement of reproducing kernels. *Journal of Machine Learning Research*, 10:107–140.
- Zhang, H. and Zhao, L. (2013). On the inclusion relation of reproducing kernel Hilbert spaces. *Analysis and Applications*, 11(2):1350014.
- Zwicker, B. (2009). Power series kernels. *Constructive Approximation*, 29(1):61–84.
- Zwicker, B. and Schaback, R. (2013). Interpolation and approximation in Taylor spaces. *Journal of Approximation Theory*, 171:65–83.