

The Impossibility Region for Detecting Sparse Mixtures using the Higher Criticism

David L. Donoho, and Alon Kipnis

Department of Statistics, Stanford University, Stanford, CA 94041, USA

Abstract: Consider a multiple hypothesis testing setting involving rare/weak effects: relatively few tests, out of possibly many, deviate from their null hypothesis behavior. Summarizing the significance of each test by a P -value, we construct a global test against the null using the Higher Criticism (HC) statistics of these P -values. We calibrate the rare/weak model using parameters controlling the asymptotic distribution of non-null P -values near zero. We derive a region in the parameter space where the HC test is asymptotically powerless. Our derivation involves very different tools than previously used to show powerlessness of HC, relying on properties of the empirical processes underlying HC. In particular, our result applies to situations where HC is not asymptotically optimal, or when the asymptotically detectable region of the parameter space is unknown.

Keywords and phrases: multiple testing, empirical process, higher criticism, phase transition.

1. Background

Multiple hypothesis testing involving rare/weak effects has attracted significant interest in statistics and machine learning over the last two decades [Donoho and Jin \(2004, 2008, 2009\)](#); [Hall and Jin \(2010\)](#); [Cai et al \(2011\)](#); [Arias-Castro et al \(2011\)](#); [Cai and Wu \(2014\)](#); [Arias-Castro and Wang \(2015\)](#); [Li and Siegmund \(2015\)](#); [Moscovich et al \(2016\)](#); [Gontscharuk et al \(2016\)](#); [Kipnis \(2019\)](#); [Donoho and Kipnis \(2020\)](#). Roughly speaking, the rare/weak effect setting proposes that the non-null effect is only moderately large, and, the non-null effects are present only in a relatively small, unknown subset of the tests. Here ‘moderately large’ means that the effects are not so large that they are individually obvious; e.g. they do not individually stand out after straightforward Bonferroni correction, but they might conceivably still be detectable using more sensitive methods.

Many of the works cited above show that the Higher Criticism (HC) test of [Donoho and Jin \(2004\)](#) has interesting optimality properties under a phase diagram analysis in which the phase space quantifies how rare and how weak the non-null effects are. In this paper, we derive a region in the phase space where HC is asymptotically powerless, in the sense that the sum of its Type I and Type II errors approaches one within this region.

All previous works deriving a region where HC is powerless argue that over the given region, *all possible* multiple hypothesis tests must be asymptotically

powerless. Consequently, this derivation settles matters only in cases where HC is asymptotically fully powerful inside the complement of the derived region. In such settings, HC will have the optimal phase diagram.

Actually, because the HC statistic is a practitioner's - rather than a theoretician's - tool, it can be used in a wide variety of multiple testing situations. We have no reason to believe that it has the optimal phase diagram in general. In situations where HC does not have the optimal phase diagram, the existing strategy for establishing its region of powerlessness does not settle matters. There remains the possibility that in certain settings, a region exists where HC is powerless, and yet some more sensitive multiple testing approach is fully powerful. As an example when this situation can arise, suppose that there exists a covariate predicting the presence/absence of the rare and weak effects. A naive approach is to ignore this covariate and apply HC to P-values obtained by testing each feature independently. Arguably, a better approach is to reduce each equivalence class indicated by levels of the covariate by combining the responses within this class, and applying HC to the collection of P-values obtained from each equivalent class after this reduction. As another example, suppose that many effects exist under the alternative hypothesis, but these effects are too weak for HC to be effective. Therefore, a test based on a statistics that aggregates the individual weak effects is expected to be much more powerful than HC. In Section 3 below we provide specific cases corresponding to these two examples.

2. Problem Formulation and Statement of the Main Result

2.1. Rare/Weak Features

Consider a multiple hypothesis testing situation involving a collection of P-values π_1, \dots, π_n associated with many independent, although not necessarily identical, tests. We are interested in testing the global null

$$H_0 : \pi_i \stackrel{\text{iid}}{\sim} \text{Unif}(0, 1), \quad \forall i = 1, \dots, n, \quad (1a)$$

against an alternative of the form

$$H_1^{(n)} : \pi_i \sim (1 - \epsilon_n)\text{Unif}(0, 1) + \epsilon_n G_{n,i}, \quad \text{independently, } \forall i = 1, \dots, n. \quad (1b)$$

Here $\epsilon_n = n^{-\beta}$, $\beta \in (0, 1)$ is the rarity parameter. $G_{n,i}$ is a probability distribution stochastically dominated by $\text{Unif}(0, 1)$ that specifies the behavior of the i th test whenever this hypothesis is non-null. We assume that, for $X \sim G_{n,i}$ and $q > 0$,

$$\max_{i=1, \dots, n} (-\log(\Pr(X < n^{-q}))) \leq \log(n) (\alpha(q, r) + o(1)), \quad (2)$$

for some bivariate function $\alpha(q, r)$ that is continuous, non-negative, increasing in q and decreasing in r , and $o(1)$ indicates a sequence tending to zero uniformly

in i and $q \in [1/2, 1]$ as n goes to infinity. Note that, because $G_{n,i}$ is stochastically smaller than $\text{Unif}(0, 1)$, only the case $\alpha(q, r) \leq q$ is interesting. As it turns out, the large n behavior of our test to be defined below only depends on the bound of the near-zero "tail" behavior of $G_{n,i}$ specified by $\alpha(q, r)$ in (2). Effectively, (2) characterizes the way $G_{n,i}$ is smaller than $\text{Unif}(0, 1)$ on values scaling as n^{-q} , $q > 0$. This characterization arises in several studies of rare/weak effects considered in the past:

- (i) The normal means model of [Donoho and Jin \(2004\)](#) corresponds to:

$$\alpha(q, r) = (\sqrt{q} - \sqrt{r})^2.$$

- (ii) The two-sample normal means and the two-sample large Poisson means models of [Donoho and Kipnis \(2020\)](#) correspond to:

$$\alpha(q, r) = (\sqrt{q} - \sqrt{r/2})^2.$$

- (iii) The two-sample small Poisson means model of [Donoho and Kipnis \(2020\)](#) corresponds to:

$$\alpha(q, r) = q \frac{\log\left(\frac{2q}{r \log(2)}\right) - 1}{\log(2)} + \frac{r}{2}.$$

- (iv) The normal heteroscedastic mixture model of [Cai et al \(2011\)](#) corresponds to:

$$\alpha(q, r; \sigma^2) = (\sqrt{q} - \sqrt{r})^2 / \sigma^2, \quad \sigma^2 > 0. \quad (3)$$

More generally, (3) arises under a verity of models in which the non-null effects are on the moderate deviation scale [Kipnis \(2021\)](#).

2.2. Higher Criticism

Let

$$\text{HC}_{n,i} \equiv \sqrt{n} \frac{i/n - \pi_{(i)}}{\sqrt{\pi_{(i)}(1 - \pi_{(i)})}},$$

where $\pi_{(i)}$ is the i -th order statistic of $\{\pi_i, i = 1, \dots, n\}$. The HC statistic is:

$$\text{HC}_n^* \equiv \max_{1 \leq i \leq n\gamma_0} \text{HC}_{n,i}, \quad (4)$$

where $0 < \gamma_0 < 1$ is a tunable parameter that does not affect the asymptotic properties of HC_n^* under $H_1^{(n)}$. It would simplify certain arguments below to fix it at some value smaller than 1/2, eg. 1/10.

The HC test rejects H_0 for large values of HC_n^* .

2.3. Impossibility of Detection

We say that a sequence of tests $\{T_n\}$ based on the P-values π_1, \dots, π_n is asymptotically powerless if

$$\Pr_{H_0}(T_n > h(n)) + \Pr_{H_1^{(n)}}(T_n \leq h(n)) \rightarrow 1,$$

for any sequence $\{h(n)\}_{n \in \mathbb{N}}$.

Consider the curve $\rho(\beta)$ defined in the plane (β, r) as

$$\rho(\beta) \equiv \sup \left\{ r > 0 : \max_{q \in (0,1]} \left(\frac{1+q}{2} - \alpha(q, r) - \beta \right) < 0 \right\}. \quad (5)$$

Our main result implies that a test based on HC_n^* is asymptotically powerless when $r < \rho(\beta)$. Namely, whenever (β, r) satisfies

$$\max_{q \in (0,1]} \left(\frac{1+q}{2} - \alpha(q, r) - \beta \right) < 0.$$

Theorem 2.1. *Consider testing H_0 against $H_1^{(n)}$. Assume that $G_{n,i}$ has a continuous density f_i , and that, for some $C < \infty$, $\|f_i\|_\infty = C$ for all $i = 1, \dots, n$. HC_n^* is asymptotically powerless if $r < \rho(\beta)$.*

3. Stylized Applications

This paper's characterization of the region of powerlessness for HC would not be necessary in situations where some lower bound argument exists showing that no statistic – including HC – can be powerful over that region. The powerlessness characterization is much more useful in cases where we want to show that HC does not have the optimal phase diagram, i.e., in cases where there exists some *other* multiple testing procedure that is powerful within the region where HC is powerless. In such situations, this paper's characterisation may allow us to establish limitations on HC which some other procedures escape. We discuss two situations of this form.

3.1. Coupled rare and weak effects

Suppose that we have an array of $n \times k$ response variables $\{X_{i,j}\}_{1 \leq i \leq n, 1 \leq j \leq k}$ experiencing rare and weak effects where the noise is independent but the signal has structure: If the rare effect is present in some variable X_{i_0, j_0} , it is also present in the entire row $\{X_{i_0, j}\}_{j=1, \dots, k}$. This scenario is a stylized version of the situation mentioned earlier, when a covariate predicting the presence/absence of the rare and weak effects exists. In this scenario, rather than considering each pair (i, j) separately, one might instead combine data within each row to a

P-value *per row* i and then apply HC to the n per-row P-values associated with each row.

As a concrete model, consider the following rare and weak normal means setting,

$$\begin{aligned} H_0 : X_{i,j} &\stackrel{\text{iid}}{\sim} \mathcal{N}(0,1) \quad i = 1, \dots, n, \quad j = 1, \dots, k \\ H_1 : (X_{i,1}, \dots, X_{i,k}) &\stackrel{\text{iid}}{\sim} (1 - \epsilon_n) \mathcal{N}^k(0,1) + \epsilon_n \cdot \mathcal{N}^k(\mu_n, 1), \quad i = 1, \dots, n, \end{aligned} \quad (6)$$

where μ_n and ϵ_n are calibrated to n in the standard manner:

$$\mu_n = \sqrt{2r \log(n)}, \quad \epsilon_n = n^{-\beta}. \quad (7)$$

We compare the asymptotic power of HC in testing H_0 versus H_1 under two collections of P-values:

$$\pi_{i,k} = \Pr(X_{i,j} \geq \mathcal{N}(0,1)), \quad i = 1, \dots, n, \quad j = 1, \dots, k, \quad (8a)$$

versus

$$\bar{\pi}_i = \Pr\left(\frac{1}{\sqrt{k}} \sum_{j=1}^k X_{i,j} \geq \mathcal{N}(0,1)\right), \quad i = 1, \dots, n. \quad (8b)$$

Note that $X_{i,j} \sim (1 - \epsilon_n) \mathcal{N}(0,1) + \epsilon_n \mathcal{N}(\mu_n, 1)$ under H_1 , hence (2) holds for $X \sim \mathcal{N}(\mu_n, 1)$ with $\alpha(q, r) = (\sqrt{q} - \sqrt{r})^2$. The upper boundary for the of impossibility in (5) evaluates to

$$\tilde{\rho}(\beta) \equiv \begin{cases} \beta - 1/2 & \frac{1}{2} < \beta < \frac{3}{4}, \\ 1 - (1 - \sqrt{\beta})^2 & \frac{3}{4} \leq \beta < 1. \end{cases}$$

It is straightforward to extend the results in Section 2 to the case where the response variables are exchangeable but not necessarily independent as in (6), implying that HC of the usual P-values (8a) is asymptotically powerless whenever $r < \tilde{\rho}$. On the other hand, in view of (7) and (8b), the effect in $\bar{\pi}_i$ under H_1 is k times stronger than the effect in $\pi_{i,j}$, while the linear reduction in the sample size from $n \cdot k$ to n when moving from $\{\pi_{i,j}\}$ to $\{\bar{\pi}_i\}$ does not affect the asymptotic performance of HC. Consequently, a standard analysis of HC applied to $\{\bar{\pi}_i\}$ implies that it has full power whenever $r > \tilde{\rho}/k$ [Donoho and Jin \(2004\)](#).

In short, the region of full power in the phase diagram based on reduction to $\bar{\pi}_i$ is considerably larger than the region of full power for the usual p-values π_i , thereby enabling a comparison between a better phase diagram and a less good one.

3.2. Many weak effects but too few moderately large ones

Another situation where HC is ineffective compared to other tests is in the presence of a mixture of two types of effects: many weak effects and rare, moderately

large effects. In certain settings it can be better to aggregate all the individual responses before reducing each response to a P-value.

As a specific model, consider the multiple hypothesis testing problem

$$H_0 : X_i \stackrel{\text{iid}}{\sim} \mathcal{N}(0, 1), \quad i = 1, \dots, n, \quad (9)$$

$$H_1 : X_i \stackrel{\text{iid}}{\sim} (1 - \epsilon_n)\mathcal{N}(a_n, 1) + \epsilon_n\mathcal{N}(\mu_n, 1), \quad i = 1, \dots, n, \quad (10)$$

where ϵ_n and μ_n are calibrated as in (7). If $a_n \rightarrow 0$ and $r < \tilde{\rho}(\beta)$, then the analysis provided in this paper implies that HC is asymptotically powerless. Nevertheless, an easy calculation shows that a test that is based on $\sum_{i=1}^n X_i^2$ is asymptotically powerful provided $a_n n^{1/2} \rightarrow \infty$.

4. Proof of Theorem 2.1

4.1. Coupling

Let I be a random subset of indices in $[n] \equiv \{1, \dots, n\}$ such that $i \in I$ with probability $\epsilon_n = n^{-\beta}$, for $i = 1, \dots, n$. An equivalent way of specifying $H_1^{(n)}$ of (1b) sets

$$\pi_i \sim \begin{cases} G_{n,i} & i \in I \\ \text{Unif}(0, 1) & i \notin I. \end{cases} \quad (11)$$

Let $Q_i^{(0)}$ and $Q_i^{(1)}$ denote the realized P-values under H_0 and $H_1^{(n)}$, respectively. On a common probability space, we couple these P-values so that

$$\begin{aligned} Q_i^{(0)} &\neq Q_i^{(1)}, & i \in I, \\ Q_i^{(0)} &= Q_i^{(1)}, & i \in I^c. \end{aligned} \quad (12)$$

For $h \in \{0, 1\}$, define

$$\text{HC}_n^{(h)} = \max_{i=1, \dots, \gamma_0 n} \sqrt{n} \frac{\frac{i}{n} - Q_{(i)}^{(h)}}{\sqrt{Q_{(i)}^{(h)}(1 - Q_{(i)}^{(h)})}}.$$

For $h = 0, 1$, the joint distribution of the collection $\{Q_i^{(h)}\}_{i=1}^n$ is identical to the joint distribution of $\{\pi_i\}_{i=1}^n$ under $H_h^{(n)}$. The coupling device should make clear to the reader that the vast majority of P-values are the same actual numbers under either hypothesis; the ones that are different appear sporadically in $[n]$; they are concentrated in the index list I . We will argue that under certain conditions, the different P-values are so thinned out that their differences are not sufficient to allow the HC statistic to behave differently between the two hypotheses.

Theorem 4.1. Let $\text{HC}_n^{(0)}$ and $\text{HC}_n^{(1)}$ be defined on a common probability space where the underlying P-values $\{Q_i^{(0)}\}_{i=1}^n$ and $\{Q_i^{(1)}\}_{i=1}^n$ are coupled as in (12). Assume that $G_{n,i}$ has a continuous density f_i , and that, for some $C < \infty$, $\|f_i\|_\infty = C$ for all $i = 1, \dots, n$. Suppose that (r, β) obeys $r < \rho(\beta)$. Then, for $c > 0$,

$$\Pr\left(\text{HC}_n^{(1)} > \text{HC}_n^{(0)} + c\right) \rightarrow 0, \quad \text{as } n \rightarrow \infty.$$

Theorem 2.1 follows immediately. The proof of Theorem 4.1 requires additional notation and preliminary technical lemmas as provided below.

4.2. Preliminaries

For any $(\beta, r) \in (1/2, 1) \times (0, \infty)$,

$$\max_{q \in (0, 1]} \left(\frac{1+q}{2} - \alpha(q, r) - \beta \right) < 0 \Leftrightarrow r < \rho(\beta). \quad (13)$$

The condition:

$$1 - \alpha(1, r) - \beta < 0 \quad (14)$$

coincides with the condition $r < \rho(\beta)$ only if $q = 1$ is the maximizer of (13). Therefore, any (β, r) such that $r < \rho(\beta)$ also satisfies (14).

Denote by $M \equiv M_n \sim \text{Bin}(n, \epsilon_n)$ the RV indicating the cardinality of I . For $h \in \{0, 1\}$, denote by

$$F_n^{(h)}(t) \equiv \frac{1}{n} \sum_{i=1}^n \mathbf{1}(Q_i^{(h)} \leq t),$$

the empirical CDF of the P-values $\{Q_i^{(h)}\}_{i=1}^n$, and denote by

$$H^{(h)}(u) \equiv \frac{1}{M} \sum_{i \in I} \mathbf{1}(Q_i^{(h)} \leq u), \quad (15)$$

the empirical CDF's of P-values in the thin subset I . Define

$$\Delta_n(t) \equiv F_n^{(1)}(t) - F_n^{(0)}(t).$$

Also, set

$$\tilde{F}^{(h)}(t) \equiv \frac{M}{n} H^{(h)}(t). \quad (16)$$

Note that although $\tilde{F}^{(h)}(u)$ is scaled like the empirical CDF of n random variables, there are actually only M RVs involved. Furthermore,

$$\Delta_n(t) = \tilde{F}^{(1)}(t) - \tilde{F}^{(0)}(t). \quad (17)$$

It is crucial for our analysis that the difference CDF is normalized in some sense by n , while coupling shows that it involves only M nontrivial contributions.

Enumerate the elements of I with indices i_1, \dots, i_M :

$$\bar{Q}_j^{(h)} \equiv Q_{i_j}^{(h)}, \quad i \in I.$$

Our key assumption (1b) implies that

$$-\log(\Pr(\bar{Q}_j^{(1)} < n^{-a})) = \log(n) (\alpha(q, r) + o(1)), \quad (18)$$

where the term $o(1)$ is independent of j . Finally, consider a grid

$$\mathcal{U}_n = \{u_{k;n}\} \subset (0, 1) \quad (19)$$

with spacing n^{-3} .

In what follows, we state and prove a series of preliminary technical lemmas.

Lemma 4.2. *For $\beta \in (0, 1)$ and $\eta > 0$ such that $1 - \beta + \eta > 0$, define the event*

$$\Omega_n \equiv \{M < n^{1-\beta+\eta}\}. \quad (20)$$

There exists $n_0 \equiv n_0(\beta, \eta)$ such that

$$\Pr(\Omega_n) \leq e^{-n^{1-\beta+\eta}}, \quad n \geq n_0. \quad (21)$$

Proof of Lemma 4.2

As $M \sim \text{Bin}(n, n^{-\beta})$, by the Chernoff bound (Mitzenmacher and Upfal, 2017, Ch. 4),

$$\Pr(M \geq n^{1-\beta+\eta}) \leq \exp(-nD(n^{-\beta+\eta} \parallel n^{-\beta})), \quad (22)$$

where $D(x \parallel y) \equiv x \log(x/y) + (1-x) \log((1-x)/(1-y))$. Inspection of the first term $x \log(x/y)$ shows that, for some $c > 0$ and $n_0 = n_0(\beta, c)$ we have:

$$nD(n^{-\beta+\eta} \parallel n^{-\beta}) \geq n^{1-\beta+\eta} \log(n) \cdot c, \quad n \geq n_0 \quad (23)$$

for all $n \geq n_0(\beta, c)$. Now combine (22) with (23). \square

The next lemma expresses a key insight. Consider the relatively few P -values that are different under H_0 and H_1 ; the smallest of any of these is still much much larger than many P -values that are the same under H_0 and H_1 . Hence the “place” we will be looking for differences in the behavior of $F_n^{(h)}(x)$ will not be at the far extreme $x \approx 0$.

Lemma 4.3. *Let $\bar{Q}_- = \min\{\bar{Q}_{(1)}^{(0)}, \bar{Q}_{(1)}^{(1)}\}$ and $Q_- = \min\{Q_{(1)}^{(0)}, Q_{(1)}^{(1)}\}$. Suppose that (β, r) satisfies (14). Then, for $\delta > 0$,*

$$\Pr(n\bar{Q}_- > n^\delta) \rightarrow 1, \quad (24)$$

and

$$\Pr(Q_- < \bar{Q}_-) \rightarrow 1. \quad (25)$$

Proof of Lemma 4.3

As (β, r) satisfies (14) and since $\alpha(q, r)$ is continuous in a semi-neighborhood around $q < 1$, there exists $\delta > 0$ and $\eta > 0$ such that

$$1 - \alpha(1 - \delta, r) - \beta - \eta < 0. \quad (26)$$

From (18), we have, conditionally on M ,

$$\begin{aligned} \Pr\left(\bar{Q}_{(1)}^{(1)} \leq n^{-1+\delta} | M\right) &= 1 - \Pr(\bar{Q}_j^{(h)} > n^{-1+\delta}, j = 1, \dots, M) \\ &= 1 - \prod_{j=1}^M \left(1 - \Pr\left(\bar{Q}_j^{(1)} \leq n^{-1+\delta}\right)\right) \\ &\leq 1 - \left(1 - n^{-\alpha(1-\delta, r)+o(1)}\right)^M \leq Mn^{-\alpha(1-\delta, r)+o(1)}, \end{aligned}$$

where in the last transition we used the inequality¹ $(1+x)^M \geq 1+Mx$, $x \geq -1$; where, of course, the term $n^{-\alpha(q, r)+o(1)} < 1$ (ie. it represents a term which always belongs to $(0, 1)$). Hence unconditionally, by Lemma 4.2,

$$\begin{aligned} \Pr\left(\bar{Q}_{(1)}^{(1)} \leq n^{-1+\delta}\right) &\leq \Pr\left(M \cdot n^{-\alpha(1-\delta, r)+o(1)} \mid \Omega_n\right) + P(\Omega_n^c) \\ &\leq n^{1-\beta+\eta} \cdot n^{-\alpha(1-\delta, r)+o(1)} + e^{-n^{1-\beta+\eta/2}}. \end{aligned}$$

Our assumption $r < \rho(\beta)$ yields (26) and, so

$$\Pr\left(\bar{Q}_{(1)}^{(1)} \leq n^{-1+\delta}\right) \leq n^{-\eta/2+o(1)} + o(1) \rightarrow 0;$$

hence $\Pr\left(n\bar{Q}_{(1)}^{(1)} \geq n^\delta\right) \rightarrow 1$. Similarly, conditionally on M , $\bar{Q}_{(1)}^{(0)}$ is the minimum of M random variables $(W_j)_{j=1}^M$ (say) iid uniform over $(0, 1)$, and so:

$$\begin{aligned} \Pr\left(\bar{Q}_{(1)}^{(0)} \leq n^{\delta-1} | M\right) &\geq 1 - \Pr\{W_j > n^{\delta-1}, j = 1, \dots, M\} \\ &= 1 - (1 - n^{\delta-1})^M \\ &\leq Mn^{-1} \leq n^{-\beta}(1 + o(1)), \end{aligned}$$

Unconditionally,

$$\Pr\left(\bar{Q}_{(1)}^{(1)} \leq n^{-1+\delta}\right) \leq n^{-\beta}(1 + o(1)) + P(\Omega_n^c) \rightarrow 0,$$

using $\beta > 0$ and (by Lemma 4.2) $\Pr(\Omega_n) \rightarrow 1$. Hence $\Pr\left(n\bar{Q}_{(1)}^{(0)} \geq n^\delta\right) \rightarrow 1$ and (24) follows. For the claim in (25), note that $Q_{(0)}^{(0)}$ is the minimum of n iid RVs uniform over $(0, 1)$, hence

$$\Pr\left(Q_{(0)}^{(0)} \leq \frac{\log(n)}{n}\right) = 1 - \left(1 - \frac{\log(n)}{n}\right)^n \rightarrow 1,$$

¹Indeed, $M \cdot \log(1+x) > \log(1+Mx)$.

while (24) implies

$$\Pr\left(\bar{Q}_- \leq \frac{\log(n)}{n}\right) \rightarrow 0.$$

Consequently,

$$\begin{aligned} \Pr(Q_- < \bar{Q}_-) &\geq \Pr\left(Q_{(1)}^{(0)} \leq \frac{\log(n)}{n} < \bar{Q}_-\right) \\ &\geq 1 - \Pr\left(Q_{(1)}^{(0)} > \frac{\log(n)}{n}\right) - \Pr\left(\bar{Q}_- \leq \frac{\log(n)}{n}\right) \rightarrow 1. \end{aligned}$$

Lemma 4.4. *Under the conditions of Lemma 4.3, we may define*

$$Q_+ \triangleq \max\{Q_i^{(0)} : Q_i^{(0)} < \bar{Q}_-\}.$$

Then for $\delta_1 > 0$,

$$n^{1-\delta_1} |Q_+ - \bar{Q}_-| \rightarrow_p 0. \quad (27)$$

Furthermore, for $\delta_2 > 0$,

$$\Pr(nQ_+ > n^{\delta_2}) \rightarrow 1. \quad (28)$$

Proof of Lemma 4.4

Set $S \equiv \bar{Q}_{(1)}^{(0)} - Q_+$, so that $S \geq |\bar{Q}_- - Q_+|$. Now, Q_+ and $\bar{Q}_{(1)}^{(0)}$ correspond to a consecutive pair in $\{Q_{(1)}^{(0)}, \dots, Q_{(n)}^{(0)}\}$. Therefore, $S \sim \text{Beta}(1, n)$, i.e., $\Pr(S \leq t) = 1 - (1 - t)^n$. Consequently, for any $\epsilon > 0$,

$$\Pr(n^{1-\delta_1} |\bar{Q}_- - Q_+| < \epsilon) \leq \Pr(S < \epsilon n^{\delta_1-1}) = 1 - (1 - \epsilon n^{\delta_1-1})^n \sim 1 - e^{-\epsilon n^{\delta_1}} \rightarrow 0.$$

This proves the claim (27). Next, fix $\delta_2 > 0$. From Lemma 4.3,

$$\Pr(n\bar{Q}_- > n^{\delta_2}) \rightarrow 1.$$

Take $\delta_1 < \delta_2$. We have $\bar{Q}_i - Q_+ = o_p(1)n^{\delta_1-1} \leq o_p(1)n^{\delta_2-1} \leq o_p(1)\bar{Q}_-$, hence

$$Q_+ = (1 + o_p(1))\bar{Q}_-,$$

and

$$\Pr(nQ_+ > n^{\delta_2}) = \Pr(n\bar{Q}_-(1 + o_p(1)) > n^{\delta_2}) \rightarrow 1,$$

which implies (25). \square

Fix $t_1 < 1/2$. Let $\bar{Q}_- \equiv \min\{\bar{Q}_{(1)}^{(0)}, \bar{Q}_{(1)}^{(1)}\}$ and $Q_- = \min\{Q_{(1)}^{(0)}, Q_{(1)}^{(1)}\}$. Suppose that $Q_- < \bar{Q}_-$. Then the set $\{Q_i^{(0)} : Q_i^{(0)} < \bar{Q}_-\}$ is non-empty by Lemma 4.3. Define $Q_+ = \max\{Q_i^{(0)} : Q_i^{(0)} < \bar{Q}_-\}$.

Lemma 4.5. *We have*

$$\text{HC}^{(1)} - \text{HC}^{(0)} \leq \sup_{t \in [Q_+, t_1]} \sqrt{n} \Delta_n(t) w(t),$$

where

$$w(t) \equiv \frac{1}{\sqrt{t(1-t)}}.$$

4.2.1. *Proof of Lemma 4.5*

Because $Q_{(1)}^{(0)} < \bar{Q}_-$, it is also true that $Q_{(1)}^{(1)} < \bar{Q}_-$ and, moreover, that $Q_{(1)}^{(1)} = Q_{(1)}^{(0)} = Q_-$. In fact,

$$\left\{ Q_i^{(0)} : Q_i^{(0)} < \bar{Q}_- \right\} = \left\{ Q_i^{(1)} : Q_i^{(1)} < \bar{Q}_- \right\}, \quad (29)$$

and the set defined this way is nonempty. Also Q^+ is a jump point of both $F_n^{(0)}(t)$ and of $F_n^{(1)}(t)$.

For any left-closed interval $D = [a, b)$ or $D = [a, b]$, define

$$\text{HC}_D^{(h)} = \max_{t \in \{Q_i^{(h)}\} \cap D} \sqrt{n} \cdot \left(F_n^{(h)}(t) - t \right) w(t); \quad (30)$$

i.e. in each case we are maximizing of the jump points of $F_n^{(h)}(t)$ in D . From (29),

$$\begin{aligned} \text{HC}_{[Q_-, Q_+]}^{(0)} &= \text{HC}_{[Q_-, Q_+]}^{(1)} = \text{HC}_{[Q_{(1)}^{(0)}, Q_+]}^{(0)} \\ &= \text{HC}_{[Q_{(1)}^{(1)}, Q_+]}^{(1)} = \text{HC}_{[Q_-, Q_+]}^{(1)}. \end{aligned}$$

Now, observe that, for $h \in \{0, 1\}$,

$$\text{HC}^{(h)} = \max \left(\text{HC}_{[Q_{(1)}^{(h)}, Q_+]}^{(h)}, \text{HC}_{[Q_+, t_1]}^{(h)} \right).$$

It follows that

$$\text{HC}^{(1)} - \text{HC}^{(0)} \leq \text{HC}_{[Q_+, t_1]}^{(1)} - \text{HC}_{[Q_+, t_1]}^{(0)}.$$

For a right-open interval $D = [a, b)$ or (a, b) , define

$$\overline{\text{HC}}_D^{(h)} = \sup_{t \in D} \sqrt{n} \left(F_n^{(h)}(t) - t \right) w(t);$$

i.e. the definition is maximizing over all points in D . Note that if the left endpoint a of $D = [a, b)$ coincides with one of the jump points of $F_N^{(h)}(t)$, then

$$\text{HC}_D^{(h)} = \overline{\text{HC}}_D^{(h)}.$$

Indeed, $F_n^{(h)}(t)$ is piecewise constant with positive jumps, so that $t \rightarrow F_n^{(h)}(t) - t$ is strictly decreasing. Meanwhile, for $t \leq t_1 < 1/2$, $t \rightarrow w(t)$ is strictly decreasing. Hence, the supremum over $t \in D = [Q_i^{(h)}, t]$ always occurs at some jump point $Q_j^{(h)} \in D$. Now, for the interval $D = [Q_+, t]$, Q_+ is a jump point of both $F_N^{(0)}(t)$ and $F_N^{(1)}(t)$ – because of the set equality (29). Hence

$$\text{HC}_{[Q_+, t_1]}^{(h)} = \overline{\text{HC}}_{[Q_+, t_1]}^{(h)}, \quad h \in \{0, 1\}.$$

Finally, since $\Delta_n(t) = F_n^{(1)}(t) - F_n^{(0)}(t)$, we have

$$\overline{\text{HC}}_{[Q_+, t_1]}^{(1)} - \overline{\text{HC}}_{[Q_+, t_1]}^{(0)} \leq \sup_{t \in [Q_+, t_1]} \sqrt{n} \Delta_n(t) w(t).$$

□

Lemma 4.6. *Let $U_{(1)} \leq \dots \leq U_{(n)}$ denote uniform order statistics. Set $U_{(0)} = 0$, $U_{(n+1)} = 1$, and $S_i \equiv U_{(i+1)} - U_{(i)}$, $i = 0, \dots, n$. Let $\eta > 0$.*

$$\Pr(\min_i S_i > n^{-2-\eta}) \rightarrow 1, \quad n \rightarrow \infty.$$

also

$$\Pr(n^{2+\eta} \min_i S_i \rightarrow \infty) = 1.$$

Proof of Lemma 4.6

We invoke familiar properties of the minimum of uniform spacings [Bairamov et al \(2010\)](#), Set $t = x/(n(n+1))$; then for $x \geq 0$ we have

$$\Pr(n(n+1) \min_i S_i > x) = \Pr(\min_i S_i > t) = (1 - t(n+1))^n = \left(1 - \frac{x}{n}\right)^n \rightarrow e^{-x}. \quad (31)$$

Both conclusions follow directly. □

Lemma 4.7. *Consider the grid \mathcal{U}_n of (19). For $U_1, \dots, U_n \stackrel{\text{iid}}{\sim} \text{Unif}(0, 1)$, let $\Omega_n^{(1)}$ denote the event that there exists, for $i = 1, \dots, n$, at least one grid point situated between each consecutive pair $[U_{(i)}, U_{(i+1)}]$, $1 \leq i \leq n-1$. Then*

$$\Pr(\Omega_n^{(1)}) \rightarrow 1.$$

Proof of Lemma 4.7

Lemma 4.6 implies that the maximal spacing between each consecutive pair eventually exceeds $n^{-2-\eta}$, for each $\eta > 0$. Because the spacings in \mathcal{U}_n all equal n^{-3} , the probability that there exists at least one grid point between each consecutive pair tends to one. □

Lemma 4.8. Consider $X_i \sim P_{X_i}$, for $i = 1, \dots, n$ independently, where we assume that each P_{X_i} has a continuous density function f_{X_i} , and that $\|f_{X_i}\|_\infty = C < \infty$ for all $i = 1, \dots, n$ for some $C > 0$. Consider the grid $\mathcal{U}_n = \{u_k\}$ of (19). Set $T_i \equiv X_{(i+1)} - X_{(i)}$. Then for $\eta > 0$,

$$\Pr \left\{ n^{2+\eta} \min_i T_i > x \right\} \rightarrow 1, \quad (32)$$

and

$$\Pr(n^{2+\eta} \min_i T_i \rightarrow \infty) = 1.$$

Proof of Lemma 4.8

Let $F_X = \text{Ave}\{F_{X_i}\}_{i=1}^n$ denote the marginal distribution; it has continuous density f_X obeying $f_X \leq C$ as well. Now define RV's $U_{(i)}$ by

$$F_X(X_i) = U_i, \quad i = 1, \dots, n.$$

These are iid uniform RV's obeying

$$F_X(X_{(i)}) = U_{(i)}, \quad i = 1, \dots, n.$$

Set $X_{(0)} = 0, X_{(n+1)} = 1$ The RV's $T_i = X_{(i)} - X_{(i-1)}$, for $i = 1, \dots, n+1$ may be written as

$$T_i = \frac{U_{(i)} - U_{(i-1)}}{f_X(x_i^*)}, \quad i = 1, \dots, n+1,$$

where x_i^* is chosen by the mean value theorem to obey

$$U_{(i+1)} - U_{(i)} = F_X(X_{(i+1)}) - F_X(X_{(i)}) = f_X(x_i^*) (X_{(i+1)} - X_{(i)}).$$

Because $f_X(x_i^*) \leq C$,

$$T_i = X_{(i)} - X_{(i-1)} \geq \frac{1}{C} (U_{(i+1)} - U_{(i)}).$$

Set $S_i = U_{(i+1)} - U_{(i)}$, $i = 1, \dots, n$. Lemma 4.6 implies

$$\Pr \left\{ n^{2+\eta} \min T_i \geq x \right\} \geq \Pr \left\{ C n^{2+\eta} \min S_i \geq x \right\} \rightarrow 1. \quad (33)$$

Lemma 4.9. Let $\Omega_n^{(2)}$ denote the event that there exists, for $i = 1, \dots, n$, at least one grid point situated between each consecutive pair $[Q_{(i)}^{(1)}, Q_{(i+1)}^{(1)}]$, $1 \leq i \leq n-1$. Then $\Pr(\Omega_n^{(2)}) \rightarrow 1$

Proof of Lemma 4.9

Using Lemma 4.8 with $X_i = Q_i^{(1)}$, for $i = 1, \dots, n$, and $x = \log(n)$, there exists n_0 such that

$$\Pr \{Cn(n+1) \min T_i \geq \log(n)\} < 1/n, \quad n \geq n_0. \quad (34)$$

Consequently, the probability that the minimal spacing between each consecutive pair among $\{Q_i^{(1)}, i = 1, \dots, n\}$ is greater than $n^{-2.5}$ is tending to 1. The stated claim follows because the spacings in the grid \mathcal{U}_n all equal n^{-3} .

Lemma 4.10. *Let $F(x)$ denote the empirical CDF of $x_1, \dots, x_n \subset (0, 1)$, and let $\mathcal{U}_n = \{u_k\}$ be the grid of (19). Assume that there exists, for $i = 1, \dots, n-1$, at least one grid point situated between each consecutive pair $(x_{(i)}, x_{(i+1)})$. Then for all sufficiently large n ,*

$$\max_{i=1, \dots, n} \sqrt{n} \frac{F(x_i) - x_i}{\sqrt{x_i(1-x_i)}} - \max_{u_k \in \mathcal{U}_n} \sqrt{n} \frac{F(u_k) - u_k}{\sqrt{u_k(1-u_k)}} \leq n^{-1}.$$

Proof of Lemma 4.10

Proof. Let $u_{k(i)}$ denote the smallest value of u_k satisfying $u \in (x_{(i)}, x_{(i+1)})$ and $u \in \mathcal{U}_n$. Then

$$F_n^{(1)}(x_i) = F_n^{(1)}(u_{k(i)}), \quad i = 1, \dots, n. \quad (35)$$

Hence:

$$\begin{aligned} & \max_{i=1, \dots, n} \left(\frac{F(x_i)}{\sqrt{x_i(1-x_i)}} - \frac{F(u_{k(i)})}{\sqrt{u_{k(i)}(1-u_{k(i)})}} \right) \\ &= \max_{i=1, \dots, n} F(x_i) \left(\frac{1}{\sqrt{x_i(1-x_i)}} - \frac{1}{\sqrt{u_{k(i)}(1-u_{k(i)})}} \right) \\ &\leq \max_{i=1, \dots, n} \left| \frac{1}{\sqrt{x_i(1-x_i)}} - \frac{1}{\sqrt{u_{k(i)}(1-u_{k(i)})}} \right|. \end{aligned}$$

Also

$$\max_{u_k \in \mathcal{U}_n} \frac{F(u_k) - u_k}{\sqrt{u_k(1-u_k)}} \geq \max_{i=1, \dots, n} \frac{F(u_{k(i)}) - u_{k(i)}}{\sqrt{u_{k(i)}(1-u_{k(i)})}},$$

admittedly, generally with equality. Combining these:

$$\begin{aligned}
& \max_{i=1,\dots,n} \frac{F(x_i) - x_i}{\sqrt{x_i(1-x_i)}} - \max_{u_k \in \mathcal{U}_n} \frac{F(u_k) - u_k}{\sqrt{u_k(1-u_k)}} \\
& \leq \max_{i=1,\dots,n} \left(\frac{F(x_i) - x_i}{\sqrt{x_i(1-x_i)}} - \frac{F(u_{k(i)}) - u_{k(i)}}{\sqrt{u_{k(i)}(1-u_{k(i)})}} \right) \\
& \leq \max_{i=1,\dots,n} \left| \frac{F(x_i)}{\sqrt{x_i(1-x_i)}} - \frac{F(u_{k(i)})}{\sqrt{u_{k(i)}(1-u_{k(i)})}} \right| \\
& + \max_{i=1,\dots,n} \left| \frac{x_i}{\sqrt{x_i(1-x_i)}} - \frac{u_{k(i)}}{\sqrt{u_{k(i)}(1-u_{k(i)})}} \right| \\
& \leq \max_{i=1,\dots,n} \left| \frac{1}{\sqrt{x_i(1-x_i)}} - \frac{1}{\sqrt{u_{k(i)}(1-u_{k(i)})}} \right| \\
& + \max_{i=1,\dots,n} \left| \frac{x_i}{\sqrt{x_i(1-x_i)}} - \frac{u_{k(i)}}{\sqrt{u_{k(i)}(1-u_{k(i)})}} \right| \\
& \leq 2/n^{9/4} + 4/n^{9/4}.
\end{aligned}$$

where the last step holds for $n > n_0$ by the next lemma. Reinstating the original \sqrt{n} normalization gives the bound

$$\max_{i=1,\dots,n} \sqrt{n} \frac{F(x_i) - x_i}{\sqrt{x_i(1-x_i)}} - \max_{u_k \in \mathcal{U}_n} \sqrt{n} \frac{F(u_k) - u_k}{\sqrt{u_k(1-u_k)}} \leq n^{1/2} \cdot \frac{6}{n^{9/4}}, \quad n > n_0.$$

As soon as $n > \max(n_0, 6^{4/3})$, our main claim follows, as then $n^{-1} > 6/n^{7/4}$. \square

Lemma 4.11. *Let $(x_i)_{i=1}^n$ denote a numerical sequence taking values in $(0, 1)$, obeying*

$$n^{-3/2} < \min_i x_i \leq \max_i x_i < 1 - n^{-3/2}.$$

Let, for each i , $u_k(i)$ denote the closest grid point from our equispaced grid \mathcal{U}_n that is as large or larger. Then

$$|u_{k(i)} - x_i| \leq n^{-3}; \quad i = 1, \dots, n. \quad (36)$$

For $n > n_0$, and $i = 1, \dots, n$,

$$\left| \frac{1}{\sqrt{u_{k(i)}(1-u_{k(i)})}} - \frac{1}{\sqrt{x_i(1-x_i)}} \right| \leq 2 \cdot n^{-9/4} \quad (37)$$

$$\left| \sqrt{\frac{u_{k(i)}}{1-u_{k(i)}}} - \sqrt{\frac{x_i}{1-x_i}} \right| \leq 4 \cdot n^{-9/4} \quad (38)$$

Proof. (36) follows essentially by definition.

The function $w(x) = \frac{1}{\sqrt{x(1-x)}}$ is convex on $(0, 1)$, with derivative $w'(x) = \frac{(1/2-x)}{(x(1-x))^{3/2}}$. Now $|w'(x)| \leq (d(x, \{0, 1\})/2)^{-3/2}/2$. Since $\min_i d(x_i, \{0, 1\}) \geq n^{-3/2}$, we get

$$\max_i |w'(x_i)| \leq (n^{-3/2}/2)^{-3/2}/2 = n^{3/4} \cdot \sqrt{2}.$$

On $0 < x < u < 1/2$, w is convex decreasing, so $w(x) > w(u) > w(x) + w'(x)|u - x|$; so there, $|w(x) - w(u)| \leq (\max_i |w'(x_i)|) \cdot |u - x|$. Since $|u - x| \leq n^{-3}$ we have

$$|w(x) - w(u)| \leq \sqrt{2} \cdot n^{3/4} \cdot n^{-3} = \sqrt{2} \cdot n^{-9/4}; \quad 0 < x < u < 1/2.$$

Arguing similarly for $1/2 \leq x < u < 1$, where w is convex increasing, $w(u) - w'(u)|u - x| \leq w(x) < w(u)$. To determine $\max_{1/2 < x_i < 1} |w'(u_{k(i)})|$, use $|w'(u)| \leq (d(u, \{0, 1\})/2)^{-3/2}/2$ and notice that $\min_i d(u_{k(i)}, \{0, 1\}) \geq n^{-3/2} - n^{-3} > n^{-3/2}/2$; hence

$$\max_{1/2 < x_i < 1} |w'(u_{k(i)})| \leq (n^{-3/2}/4)^{-3/2}/2 \leq n^{3/4} \cdot 2,$$

since $4^{3/2}/2 = 2$. Combining $|w(x) - w(u)| \leq (\max_i |w'(u_{k(i)})|) \cdot |u - x|$ and $|u - x| \leq n^{-3}$, (37) follows for the case $1/2 \leq x < u < 1$. We omit the argument for the case $1/2 - n^{-1/3} \leq x \leq 1/2 \leq u \leq 1/2 + n^{-1/3}$. (37) follows.

Now consider the function $h(x) = \sqrt{\frac{x}{1-x}}$; this is increasing on $(0, 1)$, with derivative $h'(x) = x^{-1/2}(1-x)^{-3/2}/2$. Now $|h'(x)| \leq (d(x, \{0, 1\})/2)^{-3/2}$. Since $\min_i d(x_i, \{0, 1\}) \geq n^{-3/2}$, we get $\max_i |h'(x_i)| \leq n^{-3/4} \cdot 2^{3/2}$.

On $0 < x < u < 1/2$, h is concave, so $h(x) < h(u) < h(x) + h'(x)|u - x|$; so there, $|h(x) - h(u)| \leq (\max_i |h'(x_i)|) \cdot |u - x|$; since $|u - x| \leq n^{-3}$ we get

$$|h(x) - h(u)| \leq 2^{3/2} \cdot n^{3/4} \cdot n^{-3} = 4n^{-9/4}; \quad 0 < x < u < 1/2.$$

On $1/2 \leq x < u < 1$, h is convex, and $h(u) - h'(u)|u - x| \leq h(x) < h(u)$; we have the bound $|h(x) - h(u)| \leq (\max_i |h'(u_{k(i)})|) \cdot |u - x|$, which requires us to bound $\max_{1/2 < x_i < 1} |h'(u_{k(i)})|$. We notice that $\min_i d(u_{k(i)}, \{0, 1\}) \geq n^{-3/2} - n^{-3} > n^{-3/2}/2$; hence

$$\max_{1/2 < x_i < 1} |h'(u_{k(i)})| \leq (n^{-3/2}/4)^{-3/2}/2 \leq n^{3/4} \cdot 2,$$

- again since $4^{3/2}/2 = 2$. Again using $|u - x| \leq n^{-3}$, we get

$$|h(x) - h(u)| \leq 2^{3/2} \cdot n^{3/4} \cdot n^{-3} \leq 4n^{-9/4}; \quad 1/2 < x < u < 1 - n^{-3}.$$

We again omit the argument for the case $1/2 - n^{-1/3} \leq x \leq 1/2 \leq u \leq 1/2 + n^{-1/3}$. (38) follows. \square

Lemma 4.12. For any $\eta > 0$ and $a > 0$, uniformly in $u \in [n^{\eta-1}, 1/2]$,

$$\Pr \left(\sqrt{n} \left(\tilde{F}_n^{(0)}(u) - \mathbb{E} \left[\tilde{F}_n^{(0)}(u) \right] \right) > a\sqrt{u(1-u)} \right) \leq e^{-a \cdot c \cdot n^{\eta/2}} + e^{-n^{1-(\beta+1/2)/2}},$$

for some $c = c(\beta) > 0$.

4.2.2. Proof of Lemma 4.12

$$\sqrt{n} \left(\tilde{F}^{(0)}(u) - \mathbb{E} \left[\tilde{F}^{(0)}(u) \right] \right) = \sqrt{n} \frac{M}{n} \left(H^{(0)}(u) - u \right), \quad (39)$$

so, conditional on M ,

$$\Pr \left(\sqrt{n} \left(\tilde{F}^{(0)}(u) - \mathbb{E} \left[\tilde{F}^{(0)}(u) \right] \right) > a \sqrt{u(1-u)} | M \right) \quad (40)$$

$$= \Pr \left(H^{(0)}(u) - u > a \frac{\sqrt{n}}{M} \sqrt{u(1-u)} | M \right) \quad (41)$$

$$= \Pr \left(H^{(0)}(u) > (1 + \kappa)u | M \right) =: P_{n,M}, \quad (42)$$

say, where

$$\kappa := \kappa(u; n, M, a) := a \frac{\sqrt{n}}{M} \sqrt{\frac{1-u}{u}}. \quad (43)$$

Conditional on M , apply the Chernoff inequality (Mitzenmacher and Upfal, 2017, Ch. 4):

$$P_{M,n} \leq \exp \left(-M \frac{\kappa^2}{1 + \kappa} u \right). \quad (44)$$

Consider now the event $\Omega_n \equiv \{M < n^{1-\beta+\eta_1}\}$. From Lemma 4.2, for any $\eta_1 > 0$, there exists $n_0 \equiv n_0(\beta, c', \eta_1)$ such that

$$\Pr(\Omega_n) \leq c' e^{-n^{1-\beta+\eta_1}}, \quad n \geq n_0.$$

Since $\beta \in (1/2, 1)$ we may specifically choose $\eta_1 = (\beta - 1/2)/2 > 0$, yielding $1 - \beta + \eta_1 = 1 - (\beta + 1/2)/2 > 0$.

At the same time, this choice of η_1 gives us $-1/2 + \beta - \eta_1 = (\beta - 1/2)/2 (> 0, \text{ as } \beta > 1/2)$. We are assuming $u \leq 1/2$ so on the event Ω_n :

$$\kappa = a \frac{\sqrt{n}}{M} \sqrt{\frac{1-u}{u}} \geq a \cdot n^{-1/2+\beta-\eta_1} = a \cdot n^{(\beta-1/2)/2} \rightarrow \infty. \quad (45)$$

Consequently, $\frac{\kappa^2}{1+\kappa} = \kappa \cdot (1 + o(1))$ and, for large n , $M \frac{\kappa^2}{2+\kappa} \geq c'' M \cdot \kappa$ for $c'' = c''(\beta)$. Now $M \kappa u = a \sqrt{n} \sqrt{u(1-u)}$, and since $u < 1/2$,

$$M \kappa u = a \sqrt{n} \sqrt{u(1-u)} > a \sqrt{nu/2}.$$

By hypothesis, $u \geq n^{\eta-1}$, hence

$$M \cdot \frac{\kappa^2}{2 + \kappa} \cdot u \geq a \cdot c''' \cdot n^{\eta/2}, \quad (46)$$

for $n > n'''(\beta)$, where $c''' = c'''(\beta)$. Equation (44) and Ω_n together imply that for $n > n'''(\beta)$

$$P_{M,n} \leq \exp\{-a \cdot c''' \cdot n^{\eta/2}\}. \quad (47)$$

Let P_n denote the probability mentioned in the statement of the lemma; we conclude that

$$P_n \leq \Pr(\Omega_n^c) + P_{M,n} \leq e^{-n^{1-(\beta+1/2)/2}} + e^{-a \cdot c''' \cdot n^{\eta/2}}. \quad (48)$$

as advertised. \square

Lemma 4.13. *Fix $t_1 < 1/2$ and let $t_0 = n^{-1}$. For any $\epsilon > 0$, there exists $n_0 \equiv n_0(\epsilon)$ such that, if $r < \rho(\beta)$, then*

$$\sup_{t \in (t_0, t_1)} \sqrt{n} \left| \mathbb{E} \left[\tilde{F}^{(1)}(t) \right] - \mathbb{E} \left[\tilde{F}^{(0)}(t) \right] \right| w(t) \leq \epsilon,$$

for all $n \geq n_0$.

4.2.3. Proof of Lemma 4.13

We have $\mathbb{E} \left[\tilde{F}^{(0)}(t) | M \right] = tM/n$ for $t \in (0, 1)$. Consider $t_n \in (t_0, t_1)$, parameterized as $t_n = n^{-q_n}$, where $q_n \in (-\log(t_1)/\log(n), 1)$. From (18), we have

$$\mathbb{E} \left[\tilde{F}^{(1)}(t_n) | M \right] = \frac{M}{n} n^{-\alpha(q,r)(1+o_p(1))}, \quad (49)$$

as $n \rightarrow \infty$. Let $\eta > 0$. On the event $\Omega_n \equiv \{M < n^{1-\beta+\eta}\}$, using also that $q \geq \alpha(q, r)$, we get that $M/n < n^{-\beta+\eta}$; since $w(t_n) \leq 2 \cdot n^{q_n/2}$ we have (here $q = q_n$):

$$\sqrt{n} \left| \mathbb{E} \left[\tilde{F}^{(1)}(t_n) | M \right] - \mathbb{E} \left[\tilde{F}^{(0)}(t_n) | M \right] \right| w(t_n) \quad (50)$$

$$\leq \sqrt{n} \cdot \frac{M}{n} \left| n^{-\alpha(q,r)(1+o_p(1))} - n^{-q} \right| \cdot n^{q/2} \cdot 2 \quad (51)$$

$$\leq n^{1/2} \cdot n^{-\beta+\eta} \cdot n^{-\alpha(q,r)(1+o_p(1))} \cdot n^{q/2} \cdot 2 \quad (52)$$

$$= n^{-\beta+\eta+\frac{1+q}{2}-\alpha(q,r)+o_p(1)\alpha(q,r)}. \quad (53)$$

As (β, r) satisfy (13), we can choose $\eta > 0$ such that

$$\kappa \equiv \beta + \alpha(q, r) - \frac{1+q}{2} - \eta > 0.$$

Now, $\alpha(q, r)$ is continuous in q by assumption and hence bounded for $q \in [0, 1]$, so $\alpha(q, r) o_p(1) = o_p(1)$. It follows that, on the event Ω_n , for all n large enough,

$$\sup_{t \in (t_0, t_1)} \sqrt{n} \left| \mathbb{E} \left[\tilde{F}^{(1)}(t) | M \right] - \mathbb{E} \left[\tilde{F}^{(0)}(t) | M \right] \right| w(t) \quad (54)$$

$$\leq \max_{q \in [0, 1]} n^{-\beta+\eta+\frac{1+q}{2}-\alpha(q,r)+o_p(1)} \quad (55)$$

$$= n^{-\kappa+o_p(1)} \rightarrow 0. \quad (56)$$

Since $F^{(h)}(t) \in [0, 1]$,

$$|\mathbb{E} \left[\mathbb{E} \left[\tilde{F}_n^{(h)}(t) \mid M \right] \mid \Omega_n \right] - \mathbb{E} \left[\tilde{F}^{(h)}(t) \right]| \leq 2 \cdot \Pr(\Omega_n^c); \quad h \in \{0, 1\}.$$

Moreover $\sqrt{n} \cdot w(t_n) \leq n$ over the range $t_n \in (n^{-1}, t_1)$. Finally, by Lemma 4.2, $n \cdot \Pr(\Omega_n^c) \rightarrow 0$ at an exponential rate that only depends on η . \square

Lemma 4.14. *Consider a fixed $u \in (0, 1)$. Then for $a > 0$ and $\beta \in (1/2, 1)$,*

$$\Pr \left(\sqrt{n} \left(\tilde{F}_n^{(1)}(u) - \mathbb{E} \left[\tilde{F}_n^{(1)}(u) \right] \right) > a\sqrt{u(1-u)} \right) \leq e^{-a\sqrt{nu(1-u)}} + e^{-n^{1/4}}. \quad (57)$$

Proof of Lemma 4.14

Define

$$\delta = \delta(u; n, M, a) \equiv \frac{a\sqrt{n}\sqrt{u(1-u)}}{M \cdot \mathbb{E} \left[H_M^{(1)}(u) \right]}.$$

Conditioning on M ,

$$\begin{aligned} & \Pr \left(\sqrt{n} \left(\tilde{F}_n^{(1)}(u) - \mathbb{E} \left[\tilde{F}_n^{(1)}(u) \right] \right) > a\sqrt{u(1-u)} \mid M \right) \\ &= \Pr \left(H_M^{(1)}(u) - \mathbb{E} \left[H_M^{(1)}(u) \right] > \frac{a\sqrt{n}}{M} \sqrt{u(1-u)} \right) \\ &= \Pr \left(H_M^{(1)}(u) > (1 + \delta) \mathbb{E} \left[H_M^{(1)}(u) \right] \right) \leq \exp \left\{ -M \frac{\delta^2}{1 + \delta} \mathbb{E} \left[H_M^{(1)}(u) \right] \right\}, \end{aligned}$$

The last step follows by the Chernoff inequality:

$$\Pr(X \geq (1 + \delta)\mu) \leq \left(\frac{e^{-\delta}}{(1 + \delta)^{1+\delta}} \right)^\mu \leq e^{-\mu \frac{\delta^2}{1+\delta}}, \quad (58)$$

This is valid for any $\delta > 1$, where X is the sum of M independent Bernoulli random variables with $\mu = \mathbb{E}[X]$ (Mitzenmacher and Upfal, 2017, Ch. 4).

Set $\eta_1 = (\beta - 1/2)/2$. Since $\mathbb{E} \left[H_M^{(h)}(u) \right] \leq 1$, on the event $\Omega_n \equiv \{M \leq n^{1-\beta+\eta_1}\}$, we have

$$\delta \geq \frac{a \cdot \sqrt{n}}{n^{1-\beta+\eta_1}} \frac{\sqrt{(1-u)u}}{\mathbb{E} \left[H_M^{(h)}(u) \right]} \quad (59)$$

$$\geq a \cdot n^{\beta-1/2-\eta_1} \sqrt{(1-u)u}. \quad (60)$$

Since $\beta > 1/2$, our choice of η_1 gives $\beta - 1/2 - \eta_1 = (\beta - 1/2)/2 > 0$, and so $\delta \rightarrow \infty$ with increasing n . Hence $\frac{\delta^2}{1+\delta} = \delta(1 + o(1))$, and so for large n

$$M \frac{\delta^2}{1 + \delta} \mathbb{E} \left[H_M^{(1)}(u) \right] \geq \frac{1}{2} M \cdot \delta \cdot \mathbb{E} \left[H_M^{(1)}(u) \right] = \frac{a}{2} \sqrt{n} \sqrt{u(1-u)}.$$

Therefore, on Ω_n ,

$$\Pr\left(\sqrt{n}\left(\tilde{F}_n^{(1)}(u) - \mathbb{E}\left[\tilde{F}_n^{(1)}(u)\right]\right) > a\sqrt{u(1-u)} \mid M\right) \leq \exp\left\{-\frac{a}{2}\sqrt{n}\sqrt{u(1-u)}\right\}.$$

Finally, Lemma 4.2 implies that $\Pr(\Omega_n) < e^{-n^{1-\beta+\eta_1}}$. Our choice $\eta_1 = (\beta - 1/2)/2$ gives $1 - \beta + \eta_1 = 3/4 - \beta/2 > 1/4$ since $1 > \beta$, and so

$$\Pr\left(\sqrt{n}\left(\tilde{F}_n^{(1)}(u) - \mathbb{E}\left[\tilde{F}_n^{(1)}(u)\right]\right) > a\sqrt{u(1-u)}\right) \leq e^{-a\sqrt{n}\sqrt{u(1-u)}} + e^{-n^{(3-2\beta)/4}} \quad (61)$$

$$\leq e^{-a\sqrt{n}\sqrt{u(1-u)}} + e^{-n^{1/4}} \quad (62)$$

□

Lemma 4.15. Fix $0 < \eta < 1/2$. Using the grid \mathcal{U}_n of (19), define the subgrid $\mathcal{U}_n^\eta \equiv \mathcal{U}_n \cap (n^{-1+\eta}, 1/2)$. For $c > 0$, and $h \in 0, 1$,

$$\Pr\left\{\max_{u \in \mathcal{U}_n^\eta} \sqrt{n}\left(\tilde{F}^{(h)}(u) - \mathbb{E}\left[\tilde{F}^{(h)}(u)\right]\right) w(u) \geq c\right\} \rightarrow 0. \quad (63)$$

Proof of Lemma 4.15

Lemmas 4.12 and 4.14 each imply exponential inequalities of the form

$$\max_{u \in (n^{-1+\eta}, 1/2)} \Pr\left\{\sqrt{n} \frac{\tilde{F}^{(h)}(u) - \mathbb{E}\left[\tilde{F}^{(h)}(u)\right]}{\sqrt{u(1-u)}} \geq c\right\} \leq 2 \cdot e^{-(a+bc) \cdot n^\delta}, \quad (64)$$

where $a, b \geq 0$, $a + bc > 0$, $\delta = \delta(\eta) > 0$ - valid for all $n \geq n_0(c)$. Therefore, applying the union bound across the individual grid points, we get, for $h \in \{0, 1\}$,

$$\Pr\left\{\max_{u \in \mathcal{U}_n^\eta} \sqrt{n} \frac{\tilde{F}^{(h)}(u) - \mathbb{E}\left[\tilde{F}^{(h)}(u)\right]}{\sqrt{u(1-u)}} \geq c\right\} \quad (65)$$

$$\leq \sum_{u \in \mathcal{U}_n^\eta} \Pr\left\{\sqrt{n} \frac{\tilde{F}^{(h)}(u) - \mathbb{E}\left[\tilde{F}^{(h)}(u)\right]}{\sqrt{u(1-u)}} \geq c\right\} \quad (66)$$

$$\leq \text{card}(\mathcal{U}_n^\eta) \cdot 2e^{-(a+bc)n^\delta} = O(n^4 e^{-(a+bc)n^\delta}) \rightarrow 0, \quad (67)$$

since $\text{card}(\mathcal{U}_n) \leq n^4$. □

4.3. Proof of Theorem 4.1

Set $t_1 = \gamma_0 n < 1/2$ and $t_0 = \bar{Q}_-$. Fix $\delta > 0$. Denote by Ψ_n the event that between each consecutive pair $(Q_{(i)}^{(h)}, Q_{(i+1)}^{(h)})$, there exists at least one grid point.

From Lemmas 4.7 and 4.9, we have that $\Pr(\Psi_n) > 1 - \delta/2$ for all $n \geq n_1(\delta)$ for some $n_1(\delta)$. Conditioning on Ψ_n , Lemma 4.5, implies

$$\text{HC}_n^{(1)} - \text{HC}_n^{(0)} \leq \sup_{t \leq t_1} \sqrt{n} \Delta_n(t) w(t) = \sup_{t \in [t_0, t_1]} \sqrt{n} \Delta_n(t) w(t), \quad (68)$$

while Lemma 4.10 further implies that

$$\sup_{t \in (t_0, t_1)} \sqrt{n} \Delta_n(t) w(t) \leq \max_{u_k \in \mathcal{U}_n \cap (t_0, t_1)} \sqrt{n} \Delta_n(u_k) w(u_k) + 2n^{-1}. \quad (69)$$

Write

$$\begin{aligned} \Delta_n(t) &= \left(\tilde{F}^{(1)}(t) - \mathbb{E} \left[\tilde{F}^{(1)}(t) \right] \right) + \left(\mathbb{E} \left[\tilde{F}^{(1)}(t) \right] - \mathbb{E} \left[\tilde{F}^{(0)}(t) \right] \right) \\ &\quad + \left(\mathbb{E} \left[\tilde{F}^{(0)}(t) \right] - \tilde{F}^{(0)}(t) \right). \end{aligned}$$

Fix $\eta > 0$ and $\epsilon > 0$. From Lemma 4.3, $n^{\eta-1} \bar{Q}_- > 1$ with probability at least $1 - \delta/2$ for all $n \geq n_2(\eta, \delta)$. Therefore, Lemmas 4.12 and 4.14, via Lemma 4.15, imply that

$$\max_{u_k \in \mathcal{U}_n \cap (t_0, t_1)} \sqrt{n} \left(\tilde{F}^{(0)}(u_k) - \mathbb{E} \left[\tilde{F}^{(0)}(u_k) \right] \right) w(u_k) \leq \epsilon,$$

and

$$\max_{u_k \in \mathcal{U}_n \cap (t_0, t_1)} \sqrt{n} \left(\tilde{F}^{(1)}(u_k) - \mathbb{E} \left[\tilde{F}^{(1)}(u_k) \right] \right) w(u_k) \leq \epsilon,$$

for $n \geq n_3(\eta, \epsilon)$ with probability greater than $1 - \delta/2$. From Lemma 4.13 we have that

$$\sup_{t \in (t_0, t_1)} \sqrt{n} \left(\mathbb{E} \left[\tilde{F}^{(1)}(t) \right] - \mathbb{E} \left[\tilde{F}^{(0)}(t) \right] \right) w(t) \leq \epsilon,$$

for all $n \geq n_3(\epsilon)$. Putting it all together,

$$\text{HC}_n^{(1)} - \text{HC}_n^{(0)} \leq 3\epsilon + 2n^{-1}, \quad (70)$$

with probability at least $1 - \delta$ for all $n \geq n_4(\epsilon, \eta, \delta)$. As δ, η and ϵ are arbitrary, the proof is completed.

Funding

This work is supported in part by funding from the Koret Foundation and the NSF under Grant No. DMS-1816114.

References

- Arias-Castro E., Wang M. (2015) The sparse Poisson means model. *Electronic Journal of Statistics* 9(2):2170–2201
- Arias-Castro E., Candès E. J., Plan Y. (2011) Global testing under sparse alternatives: Anova, multiple comparisons and the higher criticism. *The Annals of Statistics* 39(5):2533–2556
- Bairamov I., Berred A., Stepanov A. (2010) Limit results for ordered uniform spacings. *Statistical Papers* 51(1):227
- Cai T. T., Wu Y. (2014) Optimal detection of sparse mixtures against a given null distribution. *IEEE Transactions on Information Theory* 60(4):2217–2232
- Cai T. T., Jeng J. X., Jin J. (2011) Optimal detection of heterogeneous and heteroscedastic mixtures. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)* 73(5):629–662
- Donoho D. L., Jin J. (2004) Higher criticism for detecting sparse heterogeneous mixtures. *The Annals of Statistics* 32(3):962–994
- Donoho D. L., Jin J. (2008) Higher criticism thresholding: Optimal feature selection when useful features are rare and weak. *Proceedings of the National Academy of Sciences* 105(39):14,790–14,795
- Donoho D. L., Jin J. (2009) Feature selection by higher criticism thresholding achieves the optimal phase diagram. *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences* 367(1906):4449–4470
- Donoho D. L., Kipnis A. (2020) Higher criticism to compare two large frequency tables, with sensitivity to possible rare and weak differences. (in preparation) URL <https://arxiv.org/abs/2007.01958>, [2007.01958](https://arxiv.org/abs/2007.01958)
- Gontscharuk V., Landwehr S., Finner H., et al (2016) Goodness of fit tests in terms of local levels with special emphasis on higher criticism tests. *Bernoulli* 22(3):1331–1363
- Hall P., Jin J. (2010) Innovated higher criticism for detecting sparse signals in correlated noise. *The Annals of Statistics* 38(3):1686–1732
- Kipnis A. (2019) Higher criticism for discriminating word-frequency tables and testing authorship. URL <https://arxiv.org/abs/1911.01208>, [1911.01208](https://arxiv.org/abs/1911.01208)
- Kipnis A. (2021) Sparsely log chi-squared p-values of rare and weak features. In preparation.
- Li J., Siegmund D. (2015) Higher criticism: p -values and criticism. *The Annals of Statistics* 43(3):1323–1350
- Mitzenmacher M., Upfal E. (2017) *Probability and computing: Randomization and probabilistic techniques in algorithms and data analysis*. Cambridge university press
- Moscovich A., Nadler B., Spiegelman C. (2016) On the exact berk-jones statistics and their p -value calculation. *Electronic Journal of Statistics* 10(2):2329–2354