

Currents, charges and algebras in exceptional generalised geometry

David Osten

*Institute for Theoretical and Mathematical Physics
Lomonosov Moscow State University
Lomonosovskiy Avenue, Moscow, 119991, Russia*

ostend@itmp.msu.ru

Abstract

A classical $E_{d(d)}$ -invariant Hamiltonian formulation of world-volume theories of half-BPS p -branes in type IIb and eleven-dimensional supergravity is proposed, extending known results to $d \leq 6$. It consists of a Hamiltonian, characterised by a generalised metric, and a current algebra constructed s.t. it reproduces the $E_{d(d)}$ generalised Lie derivative. $E_{d(d)}$ -covariance necessitates the introduction of so-called charges, specifying the type of p -brane and the choice of section. For $p > 2$, currents of p -branes are generically non-geometric due to the imposition of U -duality, e.g. the M5-currents contain coordinates associated to the M2-momentum.

A derivation of the $E_{d(d)}$ current algebra from a canonical Poisson structure is in general not possible. At most, one can derive a current algebra associated to para-Hermitian exceptional geometry.

The membrane in the $SL(5)$ -theory is studied in detail. It is shown, that in a generalised frame the current algebra is twisted by the generalised fluxes. As a consistency check, the double dimensional reduction from membranes in M-theory to strings in type IIa string theory is performed. Many features generalise to p -branes in $SL(p+3)$ generalised geometries that form building blocks for the $E_{d(d)}$ -currents.

Contents

1	Introduction	2
1.1	The string case	3
1.2	Summary of results	5
2	Membranes in the $SL(5)$-theory	7
2.1	The $SL(5)$ generalised Lie derivative on the membrane phase space	8
2.2	Charges and $SL(5)$ -covariance	9
2.3	Double reduction of membrane current algebra	10
2.4	Twist by generalised vielbein and the embedding tensor	11
3	Canonical and exceptional currents	13
3.1	Para-Hermitian generalised geometries	14
3.1.1	$O(d, d)$	14
3.1.2	$SL(p + 3)$ and $E_{d(d)}$	16
3.2	Charges, currents and the (generalised) Lie derivatives	17
3.3	Canonical p -form currents, their current algebra and $SL(p + 3)$	19
3.4	Exceptional currents for $d \leq 6$	20
4	Outlook	23
A	Conventions and identities	24
B	The $E_{d(d)}$ η- and ω-symbols for $d \leq 6$	25

1 Introduction

$O(d, d)$ generalised geometry and double field theory [1–13] has been very useful in the study of world-sheet string theory in features like non-abelian and Poisson-Lie T -duality or classical world-sheet integrability [14–26]. In both cases arranging the degrees of freedom $O(d, d)$ -covariantly was advantageous. For instance, a Hamiltonian formulation of strings in generic backgrounds is possible, using only input from generalised geometry: the generalised metric determines the Hamiltonian and the invariant $O(d, d)$ -metric the Poisson structure [24, 27]. In case of generalised parallelisable spaces, this formulation has been called \mathcal{E} -model and appeared first in the study of Poisson-Lie T -duality [14, 15, 28, 29].

At the same time, exceptional field theory and $E_{d(d)}$ generalised geometry [30–40] has been a powerful tool in organising compactifications of eleven-dimensional supergravity, but not necessarily in understanding the underlying world-volume theories better. Additional complications in phrasing world-volume theories in an $E_{d(d)}$ -invariant way are to be expected. That is, because U -duality acts, in general, non-perturbatively and maps different world-volume theories to each other – in contrast to the (bosonic) string, where $O(d, d)$ maps the theory to itself. For example, problems concerning U -duality of the classical membrane have been studied in [41, 42].

The problem of $E_{d(d)}$ -invariant world-volume formulations has, so far, mainly been studied at the level of construction of actions [43–48]. In all of these proposals, the introduction

of *charges*, describing the kind of object one works with, was necessary in order to achieve duality covariance.

But, success in the strive for an \mathcal{E} -model like formulation has, by now, only been achieved for the $E_{3(3)}$ -theory and partly the $E_{4(4)}$ -theory [31, 41, 42, 49–51]. Generalising [50], such a formulation might – similar to Poisson-Lie T -duality as canonical transformation of classical string theory – help to understand to which extent the recently discussed Poisson-Lie U -duality, transformations of exceptional Drinfeld algebras [50–59], can be understood as some kind of classical duality transformation of world-volume theories in exceptional field theory.

The aim of this paper is to close that gap and propose an $E_{d(d)}$ -covariant Hamiltonian formulation of world-volume theories, building on previous attempt [49, 60, 61]. Besides the typical objects of $E_{d(d)}$ generalised geometry an additional object, the above mentioned charge, that specifies the type of world-volume, needs to be introduced to ensure $E_{d(d)}$ -covariance.

Before an overview over the results is given, the string and its Hamiltonian formulation in terms of $O(d, d)$ generalised geometry is presented, in order to clarify the key points one might strive for in a generalisation to the $E_{d(d)}$ -case.

1.1 The string case

The classical dynamics of a string coupling to the NS-NS part of an arbitrary supergravity background can be described by the Hamiltonian

$$H = \frac{1}{2} \int d\sigma \mathcal{H}^{IJ}(G(x(\sigma)), B(x(\sigma))) \mathbf{E}_I(\sigma) \mathbf{E}_J(\sigma), \quad (1.1)$$

where the embedding of the string into target space $x(\sigma)$ and the canonical momentum $p(\sigma)$ are collected in $\mathbf{E}_M(\sigma) = (p_\mu(\sigma), \partial x^\mu(\sigma))$, from now called *current*. σ is the spatial world-volume coordinate, $\partial = \partial_\sigma$. \mathcal{H} is the generalised metric, encoding the σ -model couplings, metric G and the B -field, in the Hamiltonian and the indices I, J, K, \dots are raised and lowered by the $O(d, d)$ -metric η :

$$\mathcal{H} = \begin{pmatrix} G - BG^{-1}B & BG^{-1} \\ -G^{-1}B & G^{-1} \end{pmatrix}, \quad \eta = \begin{pmatrix} 0 & \mathbb{1} \\ \mathbb{1} & 0 \end{pmatrix}. \quad (1.2)$$

The Virasoro constraints take the form $\mathcal{H}^{IJ} \mathbf{E}_I \mathbf{E}_J = 0 = \eta^{IJ} \mathbf{E}_I \mathbf{E}_J$.

Current algebra and generalised Lie brackets. For a Hamiltonian description we still need to specify the Poisson brackets of the currents \mathbf{E}_I – the *current algebra*. In this article, we often discuss two types of current algebra brackets, the Dorfman current algebra

$$\{\mathbf{E}_I(\sigma), \mathbf{E}_J(\sigma')\}_D = \eta_{IJ} \partial \delta(\sigma - \sigma') \quad (1.3)$$

and the Lie current algebra

$$\{\mathbf{E}_I(\sigma), \mathbf{E}_J(\sigma')\}_L = \eta_{IJ} \frac{1}{2} (\partial - \partial') \delta(\sigma - \sigma') + \omega_{IJ} \frac{1}{2} (\partial + \partial') \delta(\sigma - \sigma') \quad (1.4)$$

where $\omega_{IJ} = \begin{pmatrix} 0 & -\mathbb{1} \\ \mathbb{1} & 0 \end{pmatrix}$. The difference between these current algebra brackets is a total world-sheet derivative term $\int d\sigma \partial(\dots)$. Depending on the boundary conditions on $x(\sigma)$ such

total derivative terms could be neglected, they correspond to boundary contributions for the open string and winding contributions for the closed string. They have been shown to be important on the quantum level [62] and on the level of action the difference between the $O(d, d)$ invariant version and the standard non-linear σ -model is a topological term [1,63,64].

As functions on the phase space we consider functionals

$$\phi[x] = - \int d\sigma \phi^I(x(\sigma)) \mathbf{E}_I(\sigma) \in (T \oplus T^*)LM. \quad (1.5)$$

The Poisson brackets of these reproduce the generalised Lie derivative of $O(d, d)$ generalised geometry for (1.3) respectively the canonical Lie bracket for (1.4):

$$\{\phi_1, \phi_2\}_D = [v_1, v_2] + \mathcal{L}_{v_1} \xi_2 - \mathbf{d}(\iota_{v_2} \xi_1), \quad \{\phi_1, \phi_2\}_L = [v_1, v_2] + \mathcal{L}_{v_1} v_2 - \mathcal{L}_{v_2} v_1 \quad (1.6)$$

with $\phi_i = v_i + \xi_i \in (T \oplus T^*)LM$. A detailed discussion can be found in [24] and will be revisited in section 3.1.

Deriving the current algebra. In a manifestly $O(d, d)$ -covariant way, the current is defined as $\mathbf{E}_I = \eta_{IJ} \partial X^J$ for extended coordinates $X^M = (x^\mu, \tilde{x}_\mu)$. The *current algebra* can be obtained *non-covariantly* by the identification $p_\mu = \partial \tilde{x}_\mu$, as above, and applying the canonical Poisson brackets

$$\{p_\mu(\sigma), x^\nu(\sigma')\} = -\delta_\mu^\nu \delta(\sigma - \sigma'), \quad \{p_\mu(\sigma), p_\nu(\sigma')\} = \{x^\mu(\sigma), x^\nu(\sigma')\} = 0. \quad (1.7)$$

The $O(d, d)$ -covariant way to derive the current algebra is as Dirac brackets [51, 65] for the constraints

$$\Phi_I = \mathbf{P}_I - \mathbf{E}_I = \mathbf{P}_I - \eta_{IJ} \partial X^J \quad (1.8)$$

on an *extended* canonical phase space with

$$\{\mathbf{P}_M(\sigma), X^N(\sigma')\} = -c \delta_M^N \delta(\sigma - \sigma'), \quad \{\mathbf{P}_M(\sigma), \mathbf{P}_N(\sigma')\} = \{X^M(\sigma), X^N(\sigma')\} = 0. \quad (1.9)$$

The Dirac procedure is necessary, because the constraints Φ_M are second class, i.e.

$$\{\Phi_M(\sigma), \Phi_N(\sigma')\} = C_{MN}(\sigma, \sigma') = -2c \eta_{MN} \partial \delta(\sigma - \sigma') \quad (1.10)$$

is invertible. The resulting Dirac brackets for the currents \mathbf{E} are

$$\begin{aligned} \{\mathbf{E}_M(\sigma), \mathbf{E}_N(\sigma')\}_{D.B.} &= - \int d\sigma'' \int d\sigma''' \{\mathbf{E}_M(\sigma), \Phi_K(\sigma'')\} (C^{-1})^{KL}(\sigma'', \sigma''') \{\Phi_L(\sigma'''), \mathbf{E}_N(\sigma')\} \\ &= \frac{c}{2} \eta_{MN} \delta \delta(\sigma - \sigma'). \end{aligned} \quad (1.11)$$

$$\{\mathbf{E}_M(\sigma), X^N(\sigma')\}_{D.B.} = -\frac{c}{2} \delta_M^N \delta(\sigma - \sigma'). \quad (1.12)$$

If one trusts such a calculation, involving inverting $\partial \delta(\sigma - \sigma')$, $\{\mathbf{E}_M(\sigma), \mathbf{E}_N(\sigma')\}_{D.B.}$ coincides with (1.3) for $c = 2$. As will be important later, the Dirac procedure naturally constructs a Lie bracket (skewsymmetric and satisfying Jacobi identity) but the calculation seems to work only *when neglecting total spatial world-sheet derivative terms*. Hence, for the string one can derive the Dorfman current algebra (1.3) as a Dirac bracket up to such terms.

\mathcal{G}	$O(d, d)$	$SL(3) \times SL(2)$	$SL(5)$	$SO(5, 5)$	$E_{6(6)}$	$SL(p + 3)$
$\dim(M)$	d	3	4	5	6	$p + 2$
object	string	miscellaneous				p -brane
\mathcal{R}_1	$2\mathbf{d}$	$(\bar{\mathbf{3}}, \mathbf{2})$	$\mathbf{10}$	$\mathbf{16}$	$\mathbf{27}$	$\frac{(\mathbf{p}+3)(\mathbf{p}+2)(\mathbf{p}+1)}{3!}$
\mathcal{R}_2	$\mathbf{1}$	$(\mathbf{3}, \mathbf{1})$	$\bar{\mathbf{5}}$	$\mathbf{10}$	$\bar{\mathbf{27}}$	$\frac{(\mathbf{p}+4)(\mathbf{p}+3)(\mathbf{p}+2)(\mathbf{p}+1)}{4!}$
\mathcal{R}_3		$(\mathbf{1}, \mathbf{2})$	$\bar{\mathbf{5}}$	$\bar{\mathbf{16}}$	$\mathbf{78}$	
\mathcal{R}_4		$(\bar{\mathbf{3}}, \mathbf{1})$	$\bar{\mathbf{10}}$	$\mathbf{45}$	$\mathbf{351}$	
\vdots		\vdots	\vdots	\vdots	\vdots	

Table 1: The relevant representations of duality/generalised geometry groups \mathcal{G} including the begin of the tensor hierachy for the $E_{d(d)}$ groups.

Generalised flux frame. A Different choice of basis for the currents $\mathbf{E}_A = E_A^I \mathbf{E}_I$ with $E_A^I \in O(d, d)$ being a generalised vielbein, i.e. $E_A^I E_B^J \mathcal{H}_{IJ} = \delta_{AB}$, leads to a diagonalisation of the Hamiltonian $H = \frac{1}{2} \int d\sigma \delta^{AB} \mathbf{E}_A(\sigma) \mathbf{E}_B(\sigma)$ and a twist of the current algebra

$$\{\mathbf{E}_A(\sigma), \mathbf{E}_B(\sigma')\}_D = \eta_{AB} \partial \delta(\sigma - \sigma') - \mathbf{F}^C_{AB}(\sigma) \delta(\sigma - \sigma') \quad (1.13)$$

by the generalised fluxes $\mathbf{F}_{ABC} = (\partial_{[A} E_{B]}^I) E_{C]I}$ with $\partial_A = E_A^M \partial_M = E_A^M (\partial_\mu, 0)$. In case of a generalised paralisable background the \mathbf{F}_{ABC} can be chosen to be constant, then this Hamiltonian formulation takes the form of an \mathcal{E} -model.

1.2 Summary of results

All of the above points for the string are generalised to some extent to objects in $E_{d(d)}$, and in part $SL(d + 1)$, generalised geometry. A duality invariant Hamiltonian formulation of the $\frac{1}{2}$ -BPS p -branes is proposed. For that we consider a generalised geometry to some duality group \mathcal{G} on a d -dimensional manifold M with coordinates x^μ , including:

- *two representations of \mathcal{G} :* \mathcal{R}_1 with indices K, L, M, \dots and \mathcal{R}_2 with indices $\mathcal{K}, \mathcal{L}, \mathcal{M}, \dots$. The \mathcal{R}_1 -representation of the generalised tangent bundle, e.g. $(T \oplus \wedge^2 T^*)M$ for the (M-section) $SL(5)$ -theory, and of the extended coordinates $X^M = (x^\mu, \dots)$ that are used in exceptional field theory. The representations to the duality groups \mathcal{G} , relevant in that paper, are collected in table 1.
- *η -symbols $\eta_{\mathcal{M}, \mathcal{KL}}$ and $\eta^{\mathcal{M}, \mathcal{KL}}$ – invariants of \mathcal{G} :* From these one can derive the Y -tensor: $Y^{\mathcal{KL}}_{MN} = \eta^{\mathcal{P}, \mathcal{KL}} \eta_{\mathcal{P}, MN}$ that defines the *generalised Lie derivative* and the section condition $\eta^{\mathcal{M}, \mathcal{KL}} \partial_K f \partial_L g = 0$ on functions f, g of the extended coordinates X^M .
- *ω -symbols:* defining a para-Hermitian generalised geometry (discussed in section 3.1) and characterising the difference of the *standard* Lie derivate on the generalised tangent bundle to the *generalised* Lie derivate. Explicite expressions for η - and ω -symbols (and their decomposition in M-theory and type IIb sections) in the conventions of this paper can be found in appendix B.
- *a generalised metric \mathcal{H}*

The fundamental objects of such a p -brane world-volume theory are currents $\mathbf{Z} \in \mathcal{R}_1$ that are spatial top-forms on the world-volumes, i.e. (spatial) world-volume p -forms:

$$\mathbf{Z}_K(\sigma) = \frac{1}{p} \eta_{N,KL} \mathcal{Q}^N \wedge dX^L. \quad (1.14)$$

$\sigma = (\sigma^1, \dots, \sigma^p)$ always denote the spatial coordinates of the p -brane world-volume, \int the integral of the p -dimensional spatial part of the world-volume and $\delta(\sigma - \sigma')$ the δ -distribution in p -dimension.

In terms of this current the proposed Hamiltonian and (reducible) set of spatial diffeomorphism constraints are:

$$H = \frac{1}{2} \int \mathbf{Z}_K \wedge \star \mathbf{Z}_L \mathcal{H}^{KL}, \quad 0 = \eta^{M,KL} \mathbf{Z}_K \wedge \star \mathbf{Z}_L. \quad (1.15)$$

The current \mathbf{Z} is characterised by a $(p-1)$ -form charge $\mathcal{Q} \in \mathcal{R}_2$ and is supposed to satisfy the fundamental current algebra

$$\{\mathbf{Z}_M(\sigma), \mathbf{Z}_N(\sigma')\}_D = \eta_{MN} \wedge d\delta(\sigma - \sigma') \quad (1.16)$$

such that the model exhibits \mathcal{G} -covariance. Indeed, when applying this to functionals $\phi = -\int \phi^K \mathbf{Z}_K$ one reproduces the generalised Lie derivative of \mathcal{G} generalised geometry, if the charge fulfils the condition

$$\mathcal{Q}^M \wedge dX^N \partial_N = \frac{1}{p} \eta^{M,NP} \eta_{K,LP} \mathcal{Q}^K \wedge dX^L \partial_N. \quad (1.17)$$

Alternatively, similar to the string, one could also consider the current algebra

$$\{\mathbf{Z}_M(\sigma), \mathbf{Z}_N(\sigma')\}_L = \mathcal{Q}^L \wedge \left(\left(\eta_{\mathcal{L},MN} + \omega_{\mathcal{L},(MN)} \right) \frac{1}{2} (d - d') + \omega_{\mathcal{L},[MN]} \frac{1}{2} (d + d') \right) \delta(\sigma - \sigma') \quad (1.18)$$

that is a Lie bracket and corresponds to the *standard* Lie derivative on the generalised tangent bundle. For $E_{d(d)}$ with $d \geq 5$ the ω -symbols are not skewsymmetric anymore and, hence, the standard and generalised Lie derivatives on the generalised tangent bundle are **not** equivalent up to total derivative terms anymore.

As a result, it will turn out that current representation for the 'Lie' current algebra (1.18) can be generically represented by a form like:

$$\mathbf{Z}_M = (p_\mu, dx^{\mu_1} \wedge \dots \wedge dx^{\mu_p}). \quad (1.19)$$

from which one can even derive (1.18) by means of the canonical Poisson brackets for p and x . In case of the p -brane in the $SL(p+3)$ theory this will be equivalent up to total world-sheet derivative terms to (1.16). Then, one can even generalise a derivation by Dirac brackets, as in the string case. This is demonstrated in section 3.3

For *exceptional currents*, in which we are really interested, such a derivation is *not* possible, and the currents itself cannot be written only in terms of canonical momenta $p(\sigma)$ and field $x(\sigma)$. Solutions to the charge condition (1.17) dictate, that current algebra representations of exceptional symmetries must contain differentials '*non-geometric*' coordinates, i.e. those that

are not part of a section. These could be identified with momentum d.o.f.s of other objects in the theory – for example for the M5-brane, the charge \mathcal{Q} and the current \mathbf{Z} seem to contain a combination of geometric (x^μ) and ‘non-geometric’ ($\tilde{x}_{\mu\nu}$) coordinates, latter are associated to the M2 momentum, i.e. schematically

$$\mathbf{Z}_M \sim (p_\mu, dx^{\mu_1} \wedge dx^{\mu_2} \wedge dx^{\mu_3} \wedge dx^{\mu_4} \wedge d\tilde{x}_{\mu_3\mu_4}, dx^{\mu_1} \wedge \dots \wedge dx^{\mu_5}). \quad (1.20)$$

This is presented also for all the other $\frac{1}{2}$ -BPS p -pranes in $d \leq 6$ in detail in section 3.4.

For the membrane in $SL(5)$, all of this is shown explicitly in section 2, revisiting partly known results in the literature. It is observed that the key point of the string generalise: derivation of the current algebra from the canonical Poisson brackets and the \mathcal{E} -model like form of the current algebra in the generalised flux frame (section 2.4). The introduction of charges in section 2.2 achieves manifest $SL(5)$ -invariance of the current algebra and together with known Hamiltonian in that case [49] motivates of the form of Hamiltonian (1.15) and the current algebra (1.16), that conjectured here to hold more generally.

2 Membranes in the $SL(5)$ -theory

Let us consider the bosonic part of the standard Polyakov-type action of an M2-brane

$$S = T \int_{\Sigma} \left(\frac{1}{2} dx^\mu \wedge \star dx^\nu G_{\mu\nu}(x) + \frac{1}{3!} dx^\mu \wedge dx^\nu \wedge dx^\rho C_{\mu\nu\rho}(x) + \frac{1}{2} \star 1 \right) \quad (2.1)$$

with tension T and coupling to the space-time metric G and a three-form gauge potential C . For the target space, we restrict ourselves to work only on the internal part of some compactification on a four-dimensional manifold, so that $\kappa, \lambda, \mu, \dots = 1, \dots, d = 4$ and work on a three-dimensional world-volume Σ with coordinates $\xi = (\tau, \sigma^\alpha)$ and a dynamical world-volume metric. In the following we integrate out latter [66] and work in the Hamiltonian formalism. It was shown that Hamiltonian and spatial diffeomorphism constraints take the following form¹ [49]:

$$H = \frac{1}{2} \int \mathbf{Z}_K \wedge \star \mathbf{Z}_L \mathcal{H}^{KL}(G, C) \quad (2.2)$$

$$0 = \eta^{\mathcal{M}, KL} \mathbf{Z}_K \wedge \star \mathbf{Z}_L \sim e^{\mathcal{M}KK' \mathcal{L}\mathcal{L}'} \mathbf{Z}_{KK'} \wedge \star \mathbf{Z}_{\mathcal{L}\mathcal{L}'}. \quad (2.3)$$

Let us explain the notation. From here on all the differential geometric expressions, like \wedge , \star , ‘d’ or integrals, are to be understood on the (two-dimensional) spatial part of the membrane world-volume. The indices $\mathcal{K}, \mathcal{L}, \mathcal{M}, \dots = 1, \dots, 5$ denote (fundamental) \mathcal{R}_2 -indices and $K, L, M, \dots \mathcal{R}_1 = \mathbf{10}$ -indices of $SL(5)$, with decompositions into the four-dimensional M-theory target space indices $\kappa, \lambda, \mu, \dots = 1, \dots, 4$ as $V^{\mathcal{M}} = \frac{1}{\sqrt{2}} V^{\mathcal{M}\mathcal{M}'} = (v^\mu, \frac{1}{\sqrt{2}} v_{\mu\mu'})$ and $V^{\mathcal{M}} = (v^\mu, v^5)$.² \mathcal{H}^{KL} is the (inverse) generalised metric of $SL(5)$ generalised geometry, parameterised for example as

$$\mathcal{H}^{KL}(G, C) = G^{-\frac{1}{3}} \begin{pmatrix} G^{\kappa\lambda} & -G^{\kappa\mu} C_{\mu\lambda\lambda'} \\ -C_{\kappa\kappa'\mu} G^{\mu\lambda} & 2G_{\kappa[\lambda} G_{\lambda']\kappa'} + C_{\kappa\kappa'\mu} G^{\mu\mu'} C_{\mu'\lambda\lambda'} \end{pmatrix}. \quad (2.4)$$

¹The a priori five diffeomorphism constraints (2.3) can indeed be reduced to two constraints, but only in that redundant form the $SL(5)$ -covariance is manifest.

²The general conventions for $E_{d(d)}$ η -symbols are collected in appendix B.

The η -symbols $\eta_{\mathcal{M},KL}$, $\eta^{M,KL}$ are nothing else than the invariant tensor of $SL(5)$ $\eta^{M,KK'\mathcal{L}\mathcal{L}'} = \epsilon^{MKK'\mathcal{L}\mathcal{L}'}$, and can be used to define the Y -tensor and the section condition:

$$Y^{KL}{}_{MN} = \eta^{P,KL} \eta_{P,MN} \quad \eta^{M,KL} \partial_K \otimes \partial_L = 0. \quad (2.5)$$

The currents $\mathbf{Z} \in \mathcal{R}_1$ are a collection of the canonical momenta $p(\sigma)$ and differentials of the fields $x(\sigma)$:

$$\mathbf{Z}_K = \frac{1}{\sqrt{2}} \mathbf{Z}_{K\mathcal{K}'} = \left(p_{\kappa} \star 1, \frac{1}{\sqrt{2}} dx^{\kappa} \wedge dx^{\kappa'} \right) \quad (2.6)$$

in the form of spatial 2-forms on the M2-brane world-volume.

2.1 The $SL(5)$ generalised Lie derivative on the membrane phase space

We revisit the study in [49] of the Poisson structure of the above \mathbf{Z}_M in the following, in order to clarify the appearance of the generalised derivative in the sense of [24] by neglecting a topological contribution. The aim is to see how the generalised Lie derivative

$$[\phi_1, \phi_2]_D^K = \phi_1^L \partial_L \phi_2^K - \phi_2^L \partial_L \phi_1^K - Y^{KL}{}_{MN} \phi_1^M \partial_L \phi_2^N \quad (2.7)$$

is encoded in the current algebra of an M2-brane. From the canonical (equal time) Poisson brackets of the $p(\sigma)$ and $x(\sigma)$ one derives the current algebra³

$$\{\mathbf{Z}_K(\sigma), \mathbf{Z}_L(\sigma')\} = dx^\mu(\sigma) \wedge \left(\frac{1}{2} (d - d') \delta(\sigma - \sigma') \epsilon_{\mu KL} + \frac{1}{2} (d + d') \delta(\sigma - \sigma') \omega_{\mu, KL} \right) \quad (2.8)$$

As in the string case we can consider the corresponding 'Dorfman' and 'Courant' brackets:

$$\{\mathbf{Z}_K(\sigma), \mathbf{Z}_L(\sigma')\}_D = \epsilon_{\mu KL} dx^\mu(\sigma) \wedge d\delta(\sigma - \sigma'), \quad (2.9)$$

$$\{\mathbf{Z}_K(\sigma), \mathbf{Z}_L(\sigma')\}_C = \epsilon_{\mu KL} dx^\mu(\sigma) \wedge \frac{1}{2} (d - d') \delta(\sigma - \sigma'), \quad (2.10)$$

the $SL(5)$ -invariant contribution to (2.8), when neglecting total spatial derivatives. These names are appropriate because, indeed, they lead to the generalised Lie derivative:

$$\{\phi_1, \phi_2\}_D = -2 \int \phi_1^L \partial_L \phi_2^K \mathbf{Z}_K \epsilon_{\mu KL} + \int \phi_1^K \partial_{\mu'} \phi_2^L dx^\mu \wedge dx^{\mu'} \quad (2.11)$$

$$\begin{aligned} \delta_v^\mu \delta_{v'}^{\mu'} dx^v \wedge dx^{v'} \partial_{\mu'} &= \frac{1}{2} \epsilon^{\mu\mu'}{}_{\nu\nu'} dx^\nu \wedge dx^{\nu'} \partial_{\mu'} = \frac{1}{4} \epsilon^{\mu\mu\nu\nu'} \mathbf{Z}_{\nu\nu'} \partial_{\mu\mu'} = \epsilon^{\mu MN} \mathbf{Z}_N \partial_M \\ \Rightarrow \{\phi_1, \phi_2\}_D^N(\sigma) &= \phi_1^M \partial_M \phi_2^N - \phi_2^M \partial_M \phi_1^N - Y^{MN}{}_{KL} \phi_1^K \partial_M \phi_2^L \end{aligned} \quad (2.12)$$

for local functionals $\phi = - \int \phi^K(\sigma) \mathbf{Z}_K(\sigma)$, by use of the section condition $\tilde{\partial}^{\mu\mu'} \phi(X) = 0$ and the canonical Poisson brackets in \mathcal{R}_1 -indices, $\{\mathbf{Z}_K(\sigma), X^L(\sigma')\} = -\delta_K^L \delta(\sigma - \sigma')$.

The second term in (2.8), including

$$\omega_{\mu, KL} = \frac{1}{2} \begin{pmatrix} 0 & -\epsilon_{\mu\kappa}{}^{\lambda\lambda'} \\ \epsilon_{\mu\lambda}{}^{\kappa\kappa'} & 0 \end{pmatrix}, \quad (2.13)$$

³using the convention and identities from appendix A

breaks the $SL(5)$ -invariance of (2.8), but makes it a Lie bracket. In fact it will be, as in the $O(d, d)$ case, simply the canonical Lie bracket on $(T \oplus \wedge^2 T^*)M$. To summarise

$$\{\phi_1, \phi_2\}_D = [v_1, v_2] + \mathcal{L}_{v_1}\xi_2 - d(\iota_{v_2}\xi_1), \quad \{\phi_1, \phi_2\}_L = [v_1, v_2] + \mathcal{L}_{v_1}\xi_2 - \mathcal{L}_{v_2}\xi_1 \quad (2.14)$$

with $\phi_i = v_i + \xi_i = -\int \phi_i^K \mathbf{Z}_K \in (T \oplus \wedge^2 T^*)M$. Again, as in the string case, the ω -term corresponds to a boundary/topological contribution, i.e. treated as a distribution

$$\omega_{\mu, KL} \int dx^\mu \wedge d\phi, \quad (2.15)$$

giving rise to wrapping contributions $\sim \int dx^\mu \wedge dx^{\mu'}$. So, the difference between Courant and Dorfman bracket is of this form of a total differential under a spatial world volume integral. This result is completely analogous to the string resp. $O(d, d)$ case [24]. There the winding contribution was equivalent to a topological term in the action, necessary for its $O(d, d)$ -invariance. This will be discussed further, also for other generalised geometries, in section 3.1.

2.2 Charges and $SL(5)$ -covariance

One can write the current algebra in a manifestly $SL(5)$ -invariant way with the use of a charge $\mathbf{q}_M \in \overline{\mathcal{R}}_2$. E.g. the *Dorfman* current bracket can be written as

$$\{\mathbf{Z}_K(\sigma), \mathbf{Z}_L(\sigma')\}_D = \epsilon_{MKL} \mathbf{q}_{M'} dX^{MM'}(\sigma) \wedge d\delta(\sigma - \sigma') \quad (2.16)$$

or with help of a 1-form valued ($SL(5)$ -invariant) 'metric', that can be used to lower the indices,

$$\eta_{KL} = \epsilon_{MKL} \mathbf{q}_{M'} dX^{MM'}, \quad \text{s.t. } d\eta = 0, \quad (2.17)$$

as

$$\{\mathbf{Z}_K(\sigma), \mathbf{Z}_L(\sigma')\}_D = \eta_{KL} \wedge d\delta(\sigma - \sigma'). \quad (2.18)$$

Acting on local functionals $\phi = -\int \phi^K \mathbf{Z}_K$ this gives

$$\{\phi_1, \phi_2\}_D = -\int \left(2\phi_{[1}^J \partial_J \phi_2^I \mathbf{Z}_I - \epsilon_{MKL} \mathbf{q}_{M'} \phi_1^K \partial_N \phi_2^L dX^{MM'} \wedge dX^N \right). \quad (2.19)$$

The generalised Lie derivative is reproduced if we identify

$$dX^{MM'} \wedge dX^{NN'} \mathbf{q}_{M'} \partial_{NN'} = \frac{3}{2} dX^{[M|M'} \wedge dX^{N|N']} \mathbf{q}_{M'} \partial_{NN'} = \epsilon^{MM'NN'K} \mathbf{Z}_K \partial_{NN'}, \quad (2.20)$$

with the consistency condition $\mathbf{q}_{[M} \partial_{N'}] = 0$, that was used in the first step. In that way the charge \mathbf{q}_M was already introduced in [46].

In the string case we had that the currents $\mathbf{E}_I = (p_i, \partial x^i)$ are related to the doubled coordinates like $\mathbf{E}_I = \eta_{IJ} \partial X^J$. This allowed to derive the (Dorfman, Courant, Lie) brackets of the doubled coordinates $X_I(\sigma)$ and study non-geometric on the string phase space [24, 27, 65]. In a similar fashion the 'extended coordinate fields' X^M of $SL(5)$ exceptional field theory are

related to the membrane currents \mathbf{Z}_M by lowering the index with η_{MN} and taking the spatial world-sheet differential, leading to the following objects and their M-theory decompositions

$$\begin{aligned}
\text{0-forms (coordinates):} \quad X^M &= \left(x^\mu, \frac{1}{\sqrt{2}} \tilde{x}_{\mu\mu'} \right) \\
\text{1-forms:} \quad X_M &= \eta_{MN} X^N = \epsilon_{\mathcal{L}MN} \mathbf{q}_{\mathcal{L}'} X^{\mathcal{L}\mathcal{L}'} dX^N, \quad \mathbf{Z}^M \equiv \frac{1}{2} dX^M \\
\text{2-forms (currents):} \quad \mathbf{Z}_M &= \frac{1}{2} \eta_{MN} \wedge dX^N = \frac{1}{2} \epsilon_{\mathcal{K}LM} \mathbf{q}_{\mathcal{K}'} dX^{\mathcal{K}\mathcal{K}'} \wedge dX^L \quad (2.21) \\
&= \left(\frac{1}{2} d\tilde{x}_{\mu\nu} \wedge dx^\nu, \frac{1}{\sqrt{2}} dx^\mu \wedge dx^{\mu'} \right),
\end{aligned}$$

when using the choice of charge $\mathbf{q}_5 = 1$ (with all other components vanishing) that corresponds to the choice of M-theory section used before.⁴

2.3 Double reduction of membrane current algebra

As a consistency check for the logic behind identifying the current algebra with objects of exceptional generalised geometry and exceptional field theory, we study the reduction of the membrane to the type IIa string. For that, we perform the usual double dimensional reduction [67]:

$$x^4(\sigma^1, \sigma^2) = \sigma^2, \quad x^\mu(\sigma^1, \sigma^2) = x^\mu(\sigma^1) \equiv x^\mu(\sigma). \quad (2.22)$$

for $\underline{\kappa}, \underline{\lambda}, \dots = 1, 2, 3$. The membrane current (2.6) becomes

$$\mathbf{Z}_{\mathcal{M}\mathcal{M}'} = \left(dx^\mu \wedge dx^{\mu'}, dx^\mu \wedge dx^4, p_\mu, p_4 \right) = \left(0, dx^\mu \wedge d\sigma^2, p_\mu, 0 \right) \quad (2.23)$$

such that with $\mathbf{Z}_{\mathcal{M}\mathcal{M}'} \rightarrow \mathbf{z}_{\mathcal{M}\mathcal{M}'}(\sigma^1) \wedge d\sigma^2$

$$\mathbf{z}_{\mathcal{M}\mathcal{M}'} = \left(\underbrace{\epsilon^{\mu\mu'\nu} \mathbf{z}_{\mu\mu'}}_{\mathbf{z}_m}, \underbrace{\mathbf{z}_{\mu 5}, \mathbf{z}_{\mu 4}, \mathbf{z}_{45}}_{\mathbf{z}_{\bar{m}}} \right) = \left(dx^\mu, p_\mu, 0, 0 \right) \quad (2.24)$$

At the same time the membrane current algebra (2.8) reduces to

$$\{ \mathbf{z}_K(\sigma), \mathbf{z}_L(\sigma') \}_L = \eta_{KL} \frac{1}{2} (d - d') \delta(\sigma - \sigma') + \frac{1}{2} \omega_{KL} (d + d') \delta(\sigma - \sigma') \quad (2.25)$$

$$\{ \mathbf{z}_K(\sigma), \mathbf{z}_L(\sigma') \}_D = \eta_{KL} d\delta(\sigma - \sigma'), \quad (2.26)$$

with $\eta_{KL} = \epsilon_{4KL}$ and $\omega_{KL} = \omega_{4,KL}$. The only non-vanishing components are:

$$\eta_{\mu\mu'v5} = \eta_{v5\mu\mu'} = \omega_{\mu\mu'v5} = -\omega_{v5\mu\mu'} = \epsilon_{\mu\mu'v} \quad (2.27)$$

⁴The last identification of \mathbf{Z}_M with $dX^K \wedge dX^L$ is equivalent to (2.20):

$$\begin{aligned}
e^{\mathcal{N}\mathcal{R}\mathcal{R}'\mathcal{K}\mathcal{K}'} \mathbf{Z}_{\mathcal{K}\mathcal{K}'} &= \frac{1}{4} \epsilon^{\mathcal{N}\mathcal{R}\mathcal{R}'\mathcal{K}\mathcal{K}'} \epsilon_{\mathcal{M}\mathcal{L}\mathcal{L}'\mathcal{K}\mathcal{K}'} \mathbf{q}_{\mathcal{M}} dX^{\mathcal{M}\mathcal{M}'} \wedge dX^{\mathcal{L}\mathcal{L}'} \\
&= \frac{1}{4} 2! 3! \delta_{[\mathcal{M}}^{\mathcal{N}} \delta_{\mathcal{L}}^{\mathcal{R}} \delta_{\mathcal{L}']}^{\mathcal{R}'} \mathbf{q}_{\mathcal{M}} dX^{\mathcal{M}\mathcal{M}'} \wedge dX^{\mathcal{L}\mathcal{L}'} = 3 \mathbf{q}_{\mathcal{M}} dX^{[\mathcal{N}]\mathcal{M}'} \wedge dX^{|\mathcal{R}\mathcal{R}']} \left(= 2 \epsilon^{\mathcal{N}\mathcal{R}\mathcal{R}'\mathcal{K}} \mathbf{Z}_{\mathcal{K}} \right)
\end{aligned}$$

and then as above in (2.20). For that reason the extra factor $\frac{1}{2}$ was needed in the definition of \mathbf{Z} in (2.21).

or in the conventions of (2.24)

$$\eta_{KL} = \begin{pmatrix} \frac{\eta_{kl}}{0} & 0 \\ 0 & 0 \end{pmatrix}, \quad \underline{\eta} = \begin{pmatrix} 0 & \mathbb{1}_3 \\ \mathbb{1}_3 & 0 \end{pmatrix}; \quad \omega_{KL} = \begin{pmatrix} \frac{\omega_{kl}}{0} & 0 \\ 0 & 0 \end{pmatrix}, \quad \underline{\omega} = \begin{pmatrix} 0 & -\mathbb{1}_3 \\ \mathbb{1}_3 & 0 \end{pmatrix}$$

which, restricted to the $k, l, \dots = (\underline{\kappa}, \underline{\kappa}), (\underline{\lambda}, \underline{\lambda}), \dots$ indices, is the canonical $O(3,3)$ metric and the components of the canonical symplectic form ω as expected from the string discussion. T -duality transformations are defined by $M_M^N \in \text{SL}(5)$ with $M_M^K M_N^L \eta_{KL} = \eta_{MN}$.

String charges and $\text{SL}(5)$ -covariance. The type IIa discussion above motivates

$$\{\mathbf{z}_K(\sigma), \mathbf{z}_L(\sigma')\} = \mathbf{q}^M \left(\eta_{M,KL} \frac{1}{2} (d - d') \delta(\sigma - \sigma') + \frac{1}{2} \omega_{M,KL} (d + d') \delta(\sigma - \sigma') \right) \quad (2.28)$$

$$\{\mathbf{z}_K(\sigma), \mathbf{z}_L(\sigma')\}_D = \mathbf{q}^M \eta_{M,KL} d\delta(\sigma - \sigma'), \quad (2.29)$$

as the $\text{SL}(5)$ string current algebra using the string charge $\mathbf{q}^M \in \mathcal{R}_2$, fulfilling $\mathbf{q}^M \partial_{\mathcal{M}\mathcal{M}'} = 0$ [46]. But it seems to hold for both the IIa and IIb section. The $\text{SL}(5)$ -invariant $M \rightarrow \text{IIa}$ reduction condition would be something like

$$\mathbf{q}^M \sigma^2 = \mathbf{q}_{M'} X^{\mathcal{M}\mathcal{M}'}. \quad (2.30)$$

Let us proceed in the same way as for the membrane current algebra, and try to reproduce the generalised Lie derivative form the Dorfman current algebra (2.29)

$$\{\phi_1, \phi_2\} = - \int \left(2\phi_{[1}^K \partial_K \phi_{2]}^L \mathbf{z}_L - \mathbf{q}^P \eta_{P,KL} \phi_1^K \partial_M \phi_2^L dX^M \right) \quad (2.31)$$

We need to make a similar identification as in the membrane case. Let us define $\eta_{KL} = \frac{1}{2} \mathbf{q}^P \epsilon_{PKL}$ in an $\text{SL}(5)$ -invariant way. Then it is natural to define

$$\mathbf{z}_K = \eta_{KL} dX^L = \eta_{M,KL} \mathbf{q} dX^L, \quad (2.32)$$

from which

$$\begin{aligned} \epsilon^{\mathcal{P}\mathcal{M}\mathcal{M}'\mathcal{N}\mathcal{N}'} \mathbf{z}_{\mathcal{N}\mathcal{N}'} &= \frac{1}{4} \epsilon^{\mathcal{P}\mathcal{M}\mathcal{M}'\mathcal{N}\mathcal{N}'} \mathbf{q}^{\mathcal{K}} \epsilon^{\mathcal{K}\mathcal{L}\mathcal{L}'\mathcal{N}\mathcal{N}'} dX^{\mathcal{L}\mathcal{L}'} = \frac{1}{4} 3! 2! \mathbf{q}^{[\mathcal{P}} dX^{\mathcal{M}\mathcal{M}']} = 3\mathbf{q}^{[\mathcal{P}} dX^{\mathcal{M}\mathcal{M}']} \\ 3\mathbf{q}^{[\mathcal{P}} dX^{\mathcal{M}\mathcal{M}']} \partial_{\mathcal{M}\mathcal{M}'} &= \mathbf{q}^{\mathcal{P}} dX^{\mathcal{M}\mathcal{M}'} \partial_{\mathcal{M}\mathcal{M}'} \end{aligned}$$

follows, making use of the charge condition $\mathbf{q}^M \partial_{\mathcal{M}\mathcal{M}'} = 0$. Then, (2.31) becomes

$$\{\phi_1, \phi_2\} = - \int \left(2\phi_{[1}^K \partial_K \phi_{2]}^N - Y^{MN}{}_{KL} \phi_1^K \partial_M \phi_2^L \right) \mathbf{z}_N \quad (2.33)$$

as wished.

2.4 Twist by generalised vielbein and the embedding tensor

In analogy to the string case [24], we aim to diagonalise the Hamiltonian and all the constraints via $\mathbf{Z}_A(\sigma) = E_A^K(\sigma) \mathbf{Z}_K(\sigma)$ in order to characterise the model via a twist of the current algebra and to bring the model into the form of an \mathcal{E} -model. E_A^K are generalised frame

fields⁵ of the SL(5)-theory. For the canonical (Lie) current algebra (2.8) we have

$$\begin{aligned} \{\mathbf{Z}_A(\sigma), \mathbf{Z}_B(\sigma')\}_L &= \frac{1}{2} \left(j^{\mathcal{C}}(\sigma) \wedge d\delta(\sigma - \sigma') - j^{\mathcal{C}}(\sigma') \wedge d'\delta(\sigma - \sigma') \right) \eta_{\mathcal{C},AB} \\ &+ dj^{\mathcal{C}} \wedge \frac{1}{2} (d + d') \delta(\sigma - \sigma') \omega_{\mathcal{C},AB}(\sigma, \sigma') \\ &- \mathbf{F}^{\mathcal{C}}_{[AB]}(\sigma) \mathbf{Z}_C(\sigma) \delta(\sigma - \sigma') \end{aligned} \quad (2.34)$$

with $j^{\mathcal{C}} = E_{\mu}^{\mathcal{C}} dx^{\mu}$. The ϵ -symbol is SL(5)-invariant, whereas $\omega_{\mu,AB}(\sigma, \sigma') = E_A^K(\sigma) E_B^L(\sigma') \omega_{\mu,KL}$ is not SL(5)-invariant, but is again necessary for (2.34) to be a Lie bracket. The twist is characterised by the SL(5) generalised fluxes [68,69] $[E_A, E_B]_D = \mathbf{F}^{\mathcal{C}}_{AB} E_C$, i.e.

$$\mathbf{F}^{\mathcal{C}}_{AB} = 2E_N^{\mathcal{C}} \partial_{[A} E_B]^{N} - Y^{CD}{}_{AE} E_N^E \partial_D E_B^N. \quad (2.35)$$

From this definition it is quite obvious to see, why they should appear generically in the current algebra that reproduces the generalised Lie derivative. The twists can also be defined for the Courant algebroid brackets

$$\{\mathbf{Z}_A(\sigma), \mathbf{Z}_B(\sigma')\}_D = j^{\mathcal{C}}(\sigma) \wedge d\delta(\sigma - \sigma') \epsilon_{\mathcal{C}AB} - \mathbf{F}^{\mathcal{C}}_{AB}(\sigma) \mathbf{Z}_C(\sigma) \delta(\sigma - \sigma') \quad (2.36)$$

$$\{\mathbf{Z}_A(\sigma), \mathbf{Z}_B(\sigma')\}_C = j^{\mathcal{C}}(\sigma) \wedge \frac{1}{2} (d - d') \delta(\sigma - \sigma') \epsilon_{\mathcal{C}AB} - \mathbf{F}^{\mathcal{C}}_{AB}(\sigma) \mathbf{Z}_C(\sigma) \delta(\sigma - \sigma'). \quad (2.37)$$

Some kind of Courant algebroid conditions will put conditions on the $\mathbf{F}^{\mathcal{C}}_{AB}$, corresponding to a (dynamical) Bianchi identity of these fluxes.

The type of finite-dimensional algebras, as for example the recently discussed exceptional Drinfeld algebras [50–59], are contained in the current algebra as the algebra of the zero modes $z_A = -\int \mathbf{Z}_A$ in case \mathbf{F} is constant:

$$\begin{aligned} \{z_A, z_B\}_D &= \int \mathbf{F}^{\mathcal{C}}_{AB}(\sigma) \mathbf{Z}_C(\sigma) = \mathbf{F}^{\mathcal{C}}_{AB} z_C \\ \{z_A, z_B\}_C &= \mathbf{F}^{\mathcal{C}}_{[AB]} z_C, \\ \{z_A, z_B\} &= \mathbf{F}^{\mathcal{C}}_{[AB]} z_C + \int \eta^{\mathcal{E},CD} \partial_D (\omega_{\mathcal{E},AB}) \mathbf{Z}_C. \end{aligned} \quad (2.38)$$

The boundary/winding contribution to the zero mode algebra ensures (similarly to [24] for the string case) that this zero mode algebra in the last line is indeed a Lie algebra. The $\{z_A, z_B\}_D$ bracket is the one appearing in the recent discussions on exceptional Drinfeld algebras.

⁵The SL(5)-vielbeine and their invariance conditions, used here, are:

$$\begin{aligned} \text{'little' vielbein:} & \quad E_{\mathcal{A}_1}^{\mathcal{M}_1} \dots E_{\mathcal{A}_5}^{\mathcal{M}_5} \epsilon^{\mathcal{A}_1 \dots \mathcal{A}_5} = \epsilon^{\mathcal{M}_1 \dots \mathcal{M}_5} \\ \text{'big' vielbein:} & \quad E_A^K = \frac{1}{2} E_{\mathcal{A}\mathcal{A}'}^{\mathcal{K}\mathcal{K}'} = E_{[\mathcal{A}}^{\mathcal{K}} E_{\mathcal{A}']}^{\mathcal{K}'}, \quad E_A^K E_B^L E_M^C E_N^D Y^{MN}{}_{KL} = Y^{CD}{}_{AB}. \end{aligned}$$

In a frame, defined by such a vielbein, also the following object, a 'little' current, appears:

$$j^{\mathcal{C}} = E^{\mathcal{C}}{}_{\mu} dx^{\mu} \quad \text{with} \quad d(\eta_{\mathcal{C},AB} j^{\mathcal{C}}) = d(\epsilon_{\mu KL} E_A^K E_B^L dx^{\mu}) = -Y^{MN}{}_{KL} \partial_M (E_A^K E_B^L) \mathbf{Z}_N$$

$\mathcal{A}, \mathcal{B}, \dots$ are the flat (generalised flux frame) \mathcal{R}_2 SL(5)-indices and $A = [\mathcal{A}\mathcal{A}']$, ... flat (generalised flux frame) \mathcal{R}_1 SL(5)-indices.

3 Canonical and exceptional currents

We aim to apply the same concepts as in section 2 to all the world-volume theories of all the $\frac{1}{2}$ -BPS objects in string and M-theory, and investigate how their current algebra can be understood to incorporate generalised geometry structures.

There are two generalisations that we will be concerned with. Of course the key interest lies in an $E_{d(d)}$ -covariant description of world-volume theories of objects in type II string and M-theories. However, some features are shared by the simpler case of p -branes in $SL(p+3)$ generalised geometry, considered in [38]. In all cases here, the Y -tensor can be defined in terms of η -symbols

$$Y^{MN}{}_{KL} = \eta^{\mathcal{P},MN} \eta_{\mathcal{P},KL} \quad (3.1)$$

\mathcal{R}_1 -indices are denoted by K, L, M, \dots and \mathcal{R}_2 -indices by $\mathcal{K}, \mathcal{L}, \mathcal{M}, \dots$, see table 1. Let us first settle our conventions for the appearing duality groups, here.

$SL(p+3)$ generalised geometry is the geometry on the generalised tangent bundle $(T \oplus \wedge^p T^*) M$. The η -symbol and the corresponding ω -symbol, that will be used in the following, are

$$\eta_{\kappa_1 \dots \kappa_{p-1}, MN} = \begin{pmatrix} 0 & \frac{\delta^{\nu_1 \dots \nu_p}}{\sqrt{p!}} \\ \frac{\delta^{\mu_1 \dots \mu_p}}{\sqrt{p!}} & 0 \end{pmatrix}, \quad \omega_{\kappa_1 \dots \kappa_{p-1}, MN} = \begin{pmatrix} 0 & -\frac{\delta^{\nu_1 \dots \nu_p}}{\sqrt{p!}} \\ \frac{\delta^{\mu_1 \dots \mu_p}}{\sqrt{p!}} & 0 \end{pmatrix}, \quad (3.2)$$

where other components are supposed to vanish and the following decompositions of \mathcal{R}_1 - and \mathcal{R}_2 -indices into spacetime indices $\kappa, \lambda, \mu = 1, \dots, d = p+2$ are made:

$$X^M = \left(x^\mu, \frac{\tilde{x}^{\mu_1 \dots \mu_p}}{\sqrt{p!}} \right) \in \mathcal{R}_1, \quad Q^{\mathcal{M}} = \left(\frac{Q^{\mu_1 \dots \mu_{p-1}}}{\sqrt{(p-1)!}}, \frac{Q^{\mu_1 \mu_2 \mu_3 \mu_4}}{\sqrt{4!}} \right) \in \mathcal{R}_2.$$

\mathcal{R}_1 and \mathcal{R}_2 correspond to the three- resp. four-fold skewsymmetric representation of $SL(p+3)$. That generalises the $p=2$ -case, that is the membrane case from section 2. The generalised Lie derivative $[\phi_1, \phi_2]_D$ takes the same form to the $O(d, d)$ and $SL(5)$ for all p :

$$[\phi_1, \phi_2]_D = [v_1, v_2] + \mathcal{L}_{v_1} \xi_2 - d(\iota_{v_2} \xi_1) \quad (3.3)$$

with $\phi_i = v_i + \xi_i \in (T \oplus \wedge^p T^*) M$.

$E_{d(d)}$ generalised geometry. The η -symbols and conventions for decomposition of the \mathcal{R}_1 - and \mathcal{R}_2 -representation $E_{d(d)}$ for $d \leq 6$ in the type IIB and M-theory section are collected in appendix B [70]. The representations of the duality groups that will be relevant are collected in table 1. Mainly the representations \mathcal{R}_1 and \mathcal{R}_2 will matter in the following, whereas the higher representations of the tensor hierarchy will not appear in the following. For $d \leq 6$ the generalised Lie derivative always takes the form (in the M-theory section):

$$[\phi_1, \phi_2]_D = [v_1, v_2] + (\mathcal{L}_{v_1} \omega_2 - d(\iota_{v_2} \omega_1)) + (\mathcal{L}_{v_1} \xi_2 - d(\iota_{v_2} \xi_1) + \omega_1 \wedge d\omega_2) \quad (3.4)$$

with $\phi_i = v_i + \omega_i + \xi_i \in (T \oplus \wedge^2 T^* \oplus \wedge^5 T^*) M$.

The general setup. The proposition of this section is a Hamiltonian formulation of any p -brane object in a duality covariant way. The obvious conjecture, generalising the membrane in $SL(5)$, for the Hamiltonian and spatial diffeomorphism constraints is

$$H = \frac{1}{2} \int \mathbf{Z}_K \wedge \star \mathbf{Z}_L \mathcal{H}^{KL}, \quad 0 = \eta^{M,KL} \mathbf{Z}_K \wedge \star \mathbf{Z}_L. \quad (3.5)$$

for the generalised metric \mathcal{H} and the η -symbols $\eta_{M,KL}$ of that theory. The currents \mathbf{Z}_M are spatial world-volume p -forms. In the following, these currents \mathbf{Z} and their algebra will be the main focus having two main questions in mind:

- How should the currents \mathbf{Z} and a currents algebra look, such that they form some kind of representation of the exceptional symmetry? I.e. how must the currents and their algebra look, such that bracket of $\phi = - \int \phi^M \mathbf{Z}_M$ will reproduce generalised Lie brackets (3.3) or (3.4)?
- Can these currents be constructed by means of the canonical Poisson structure? I.e. Will the currents be in generality derivatives of coordinate fields $x(\sigma)$ and their canonical momenta $p(\sigma)$, arranged into representations of the duality group as in (2.21) for the membrane?

3.1 Para-Hermitian generalised geometries

In order to answer the above questions we first go a step back, and reconsider why the construction of the generalised Lie derivative from the canonical Poisson structure (up to total derivative terms) worked for the string and the membrane.

In these constructions a non duality-invariant ω -symbol appeared, ensuring the Lie bracket properties of the current algebra. Let us investigate how the invariants $\eta_{M,KL}$ and the ω -symbols combine to produce a Lie bracket. This is part of so-called para-Hermitian geometry [71–76].

3.1.1 $O(d, d)$

We define the projector

$$P^{KL}{}_{MN} = \frac{1}{2} \left(Y^{KL}{}_{MN} + \Omega^{KL}{}_{MN} \right)$$

with $Y^{KL}{}_{MN} = \eta^{KL} \eta_{MN}$ and $\Omega^{KL}{}_{MN} = \eta^{KL} \omega_{MN}$ for $O(d, d)$ generalised geometry. This notation is chosen in a way that it directly generalises to exceptional generalised geometry. The projector has the following properties:

$$P^{KL}{}_{MN} = P^{LK}{}_{MN}, \quad P^{KL}{}_{MN} \partial_K \otimes \partial_L = 0 \quad (\text{section condition})$$

Using this section condition, one can derive the identities:

$$P^{KL}{}_{MN} P^{NP}{}_{RS} \partial_P = P^{KL}{}_{RS} \begin{pmatrix} 0 \\ \delta^m \end{pmatrix} \approx 0, \quad (3.6)$$

$$P^{KL}{}_{NM} P^{NP}{}_{RS} \partial_P = P^{KL}{}_{RS} \begin{pmatrix} \partial_m \\ 0 \end{pmatrix} \approx P^{KL}{}_{RS} \partial_M. \quad (3.7)$$

In comparison to the standard Courant algebroid structure, there are now two total derivative⁶ bilinear objects:

$$((\phi_1, \phi_2))^K = \frac{1}{2} Y^{KL}{}_{MN} \partial_L (\phi_1^M \phi_2^N), \quad \llbracket \phi_1, \phi_2 \rrbracket^K = \frac{1}{2} \Omega^{KL}{}_{MN} \partial_L (\phi_1^M \phi_2^N) \quad (3.8)$$

Besides the standard Dorfman bracket $[\phi_1, \phi_2]_D^K$, which is defined in terms of the Y -tensor, one can define a bracket in terms of the projector P :

$$[\phi_1, \phi_2]_L^K = \phi_1^L \partial_L \phi_2^K - \phi_2^L \partial_L \phi_1^K + 2P^{KL}{}_{MN} \phi_1^M \partial_L \phi_2^N \quad (3.9)$$

By the identities (3.6) and (3.7) this is indeed a Lie bracket and in fact

$$[\phi_1, \phi_2]_L = [v_1, v_2] + \mathcal{L}_{v_1} \xi_2 - \mathcal{L}_{v_2} \xi_1 = [\phi_{[1}, \phi_{2]}]_D + \llbracket \phi_1, \phi_2 \rrbracket \equiv [\phi_1, \phi_2]_C + \llbracket \phi_1, \phi_2 \rrbracket, \quad (3.10)$$

the standard Lie derivative on $(T \oplus T^*)M$ with $\phi_i = v_i + \xi_i \in (T \oplus T^*)M$. Also, $[\phi_1, \phi_2]_L \sim [\phi_1, \phi_2]_C$ up to total derivative terms in the $O(d, d)$ case.

To summarise: the para-Hermitian geometry connected to the pair (η, ω) defines a (standard) Lie algebroid over $TM \oplus T^*M$. Connected to that, Lie bracket by total derivative terms, are other algebraic entities with interesting properties, including the standard Courant algebroid over M .

ω -term and the section. The crucial point of the ' ω -geometry' is that it, in contrast to the standard approach to generalised geometry or double field theory, allows for a reconstruction of the section from the choice of ω and vice versa.

We choose the $P^{KL}{}_{MN}$ as the fundamental object obeying the identities (3.14) and start with the standard section $\partial_N \approx (\partial_n, 0)$. Then, as $P^{KL}{}_{MN} = \frac{1}{2} \eta^{KL} (\eta_{MN} + \omega_{MN})$, the identities (3.14) imply

$$\omega = \begin{pmatrix} 2B & -\mathbb{1} \\ \mathbb{1} & 0 \end{pmatrix} \quad (3.11)$$

for some skewsymmetric matrix B . If we took an arbitrary section $\partial'_M = M_M{}^N \partial_N$, ω transforms as $\omega' = M \cdot \omega \cdot M^T$, for $M \in O(d, d)$. Up to a (constant) B -shift, the choice of section determines the form of ω and vice versa. This B -shift symmetry is also a well-known property of a Courant algebroid.

A conceptual consequence is that one can reconstruct the Lie algebroid structure of the current algebra from the standard Courant algebroid *plus* a choice of section.

Para-Hermitian and para-Kähler geometries. The current algebra is characterised by the pair (η, ω) and could be completed to a compatible triple (η, ω, I) by $I^M{}_N = \eta^{MK} \omega_{KN}$. If I is a real structure, $I^2 = \mathbb{1}$, the geometry is called *para-Hermitian*, if $d\omega = 0$ *para-Kähler* [71, 73, 75].

⁶Contracting these with the $O(d, d)$ -currents \mathbf{E} makes it obvious that they correspond to total derivatives under the spatial world-volume integrals:

$$\begin{aligned} ((\phi_1, \phi_2)) &= \int ((\phi_1, \phi_2))^K \mathbf{E}_K = 2\eta_{MN} \int d(\phi_1^M \phi_2^N) = 2 \int d(\phi_1 \bullet \phi_2) \\ \llbracket \phi_1, \phi_2 \rrbracket &= \int \llbracket \phi_1, \phi_2 \rrbracket^K \mathbf{E}_K = 2\omega_{MN} \int d(\phi_1^M \phi_2^N) = 2 \int d(\phi_1 \circ \phi_2). \end{aligned}$$

In addition, a string model is defined by a generalised metric \mathcal{H} in the Hamiltonian formalism. Recently, Born geometry was introduced as para-Kähler geometry of the triple $(\eta, \omega, \mathcal{H})$, subject to the conditions [72]

$$\eta^{-1}\mathcal{H} = \mathcal{H}^{-1}\eta, \quad \omega^{-1}\mathcal{H} = -\mathcal{H}^{-1}\omega. \quad (3.12)$$

A central result of [72, 73] was that, following from this, there exists a frame in which all the defining structures take a canonical form:

$$\mathcal{H} = \mathbb{1}, \quad \omega = \begin{pmatrix} 0 & -\mathbb{1} \\ \mathbb{1} & 0 \end{pmatrix}, \quad \eta = \begin{pmatrix} 0 & \mathbb{1} \\ \mathbb{1} & 0 \end{pmatrix}. \quad (3.13)$$

The input that we obtain from the Hamiltonian formulation of the string is different, though. In the generalised metric formulation, in which we worked so far, – meaning canonical coordinates on the phase space and background information encoded in the Hamiltonian via the generalised metric – we get η and ω in their canonical form and $\mathcal{H}(G, B)$ in a general background dependent form. So, unless we are in flat (or toroidal) space, where we can choose $\mathcal{H} = \mathbb{1}$, the classical phase space geometry of the string σ -model is *not* described by Born geometry.

3.1.2 $\mathrm{SL}(p+3)$ and $E_{d(d)}$

In total analogy to the $O(d, d)$ case, we study the role that the ω -term plays in exceptional generalised geometry. The main question will be whether a similar relation to (3.10) holds, i.e. whether generalised and standard Lie derivatives are the same up to total derivative terms.

We use the definitions of η - and ω -symbols in (3.2) and appendix B in order to define the projector $P^{KL}{}_{MN} = \frac{1}{2}(Y^{KL}{}_{MN} + \Omega^{KL}{}_{MN}) = \frac{1}{2}\eta^{P, KL}(\eta_{P, MN} + \omega_{P, MN})$. This projector has the following properties:

$$P^{KL}{}_{MN} = P^{LK}{}_{MN}, \quad P^{KL}{}_{MN}\partial_K \otimes \partial_L = 0 \quad (\text{section condition})$$

Using this section condition one derives

$$P^{KL}{}_{MN}P^{NP}{}_{RS}\partial_L \otimes \partial_P \approx 0$$

$$P^{KL}{}_{NM}P^{NP}{}_{RS}\partial_{(L} \otimes \partial_{P)} \approx P^{KL}{}_{RS}\partial_{(L} \otimes \partial_{M)} \quad (3.14)$$

$$\text{e.g. } P^{KL}{}_{NM}P^{NP}{}_{RS} \left(\partial_L \phi_{[1}^M \right) \left(\partial_P \phi_{2]}^S \right) \approx P^{KL}{}_{RS} \left(\partial_L \phi_{[1}^P \right) \left(\partial_P \phi_{2]}^S \right) \quad (3.15)$$

As is also the case in the usual exceptional generalised geometry in identities regarding the Y -tensor [34], these identities involving the projector P are a weaker than in the string case.

Instead of taking a given section and computing these identities, one could again take the opposite route and choose the $P^{KL}{}_{MN}$ as the fundamental object obeying the identities (3.14). Then one sees that a choice of ω is equivalent to a choice of (M-theory or IIb) section up to a gauge-transformation of the three-form gauge fields.

Again, one can define a total derivative object

$$\llbracket \phi_1, \phi_2 \rrbracket^K = \frac{1}{2}\Omega^{KL}{}_{MN}\partial_L \left(\phi_1^M \phi_2^N \right) \quad (3.16)$$

and, besides the standard Dorfman brackets (3.3) and (3.4), a bracket in terms of the projector P :

$$[\phi_1, \phi_2]_L^K = \phi_1^L \partial_L \phi_2^K - \phi_2^L \partial_L \phi_1^K + 2P^{KL}{}_{MN} \phi_1^M \partial_L \phi_2^N. \quad (3.17)$$

By the identities (3.14) this is a Lie bracket. From here on, the stories slightly diverge:

- For the $SL(p+3)$ -theory, we have

$$[\phi_1, \phi_2]_L = [v_1, v_2] + \mathcal{L}_{v_1} \xi_2 - \mathcal{L}_{v_2} \xi_1 = [\phi_1, \phi_2]_C + \llbracket \phi_1, \phi_2 \rrbracket, \quad (3.18)$$

with $\phi_i = v_i + \xi_i \in (T \oplus \wedge^p T^*) M$ as in the $O(d, d)$ -case.

- For the $E_{d(d)}$ -theory, on the other hand, we have

$$[\phi_1, \phi_2]_L = [v_1, v_2] + (\mathcal{L}_{v_1} \omega_2 - \mathcal{L}_{v_2} \omega_1) + (\mathcal{L}_{v_1} \xi_2 - \mathcal{L}_{v_2} \xi_1) \quad (3.19)$$

with $\phi_i = v_i + \omega_i + \xi_i \in (T \oplus \wedge^2 T^* \oplus \wedge^5 T^*) M$. So, essentially, the para-Hermitian $E_{d(d)}$ -geometry is basically *the same* as multiple copies of para-Hermitian $SL(p+3)$ -geometry put together. But, the key difference is that the generalised Lie derivative of $E_{d(d)}$ -generalised geometry and the standard Lie derivative on the extended tangent bundle do **not** only differ by a total derivative term: $[\phi_1, \phi_2]_L \neq [\phi_1, \phi_2]_C + \llbracket \phi_1, \phi_2 \rrbracket$. The reason is, that the ‘interaction’ term $\omega_{[1} \wedge d\omega_{2]}$ in (the Courant bracket version of) (3.4) is *not* a total derivative. On the level of the ω -symbol this has also the consequence that the ω -symbols are in general *not* skewsymmetric: $\omega_{\mathcal{M}, \mathcal{K}L} \neq -\omega_{\mathcal{M}, L\mathcal{K}}$, see appendix B for the explicit expressions.

3.2 Charges, currents and the (generalised) Lie derivatives

As currents of p -branes we postulate, motivated by our investigation of the M2-brane and the type IIa and type IIb string in the $SL(5)$ -theory in section 2, spatial top-forms on the world-volumes, i.e. (spatial) world-volume p -forms:

$$\mathbf{Z}_K(\sigma) = \frac{1}{p} \eta_{\mathcal{N}, KL} \mathcal{Q}^{\mathcal{N}} \wedge dX^K. \quad (3.20)$$

For sake of simplicity of notation, $\sigma = (\sigma^1, \dots, \sigma^p)$ always denote the spatial coordinates of the p -brane world-volume, \int the integral of the p -dimensional spatial part of the world-volume and $\delta(\sigma - \sigma')$ the δ -distribution in p -dimension. The p , for which σ , \int , δ and the spatial world-volume differential d are to be understood, should be clear from context.

The currents (3.20) are characterised by a charge $\mathcal{Q} : \wedge^{p-1} \mathcal{R}_1 \rightarrow \mathcal{R}_2$, an \mathcal{R}_2 -valued $(p-1)$ -form on the extended space, parameterised by coordinates X^M . We define

$$\eta_{KL} = \eta_{\mathcal{N}, KL} \mathcal{Q}^{\mathcal{N}} = \frac{1}{(p-1)!} \eta_{\mathcal{N}, KL} q_{N_1 \dots N_{p-1}}^{\mathcal{N}} dX^{N_1} \wedge \dots \wedge dX^{N_{p-1}}$$

in term of constants $q_{1 \dots N_{p-1}}^{\mathcal{N}}$ and consequentially assume that $d\mathcal{Q} = 0$ resp. $d\eta = 0$. In such a form a charge appeared already in [45, 48].

The aim again is that the algebra of sections $\phi[X] = -\int \phi^K(X(\sigma)) \mathbf{Z}_K(\sigma)$, satisfying the section condition, is the standard exceptional generalised Lie derivative

$$\{\phi_1, \phi_2\}_D = -\int \left(\phi_1^K \partial_K \phi_2^L - \phi_2^K \partial_K \phi_1^L - Y^{KL}{}_{MN} \phi_1^M \partial_K \phi_2^N \right) \mathbf{Z}_L \quad (3.21)$$

respectively the para-Hermitian exceptional generalised Lie derivative $\{\phi_1, \phi_2\}_L$ for $P^{KL}{}_{MN}$ instead of $Y^{KL}{}_{MN}$. This will be the case if we assume $\{\mathbf{Z}_M(\sigma), X^N(\sigma')\} = \delta_M^N \delta(\sigma - \sigma')$ and the current algebra

$$\{\mathbf{Z}_M(\sigma), \mathbf{Z}_N(\sigma')\}_D = \eta_{MN} \wedge d\delta(\sigma - \sigma') \quad (3.22)$$

respectively

$$\{\mathbf{Z}_M(\sigma), \mathbf{Z}_N(\sigma')\}_L = \mathcal{Q}^L \wedge \left(\left(\eta_{L,MN} + \omega_{L,(MN)} \right) \frac{1}{2} (d - d') + \omega_{L,[MN]} \frac{1}{2} (d + d') \right) \delta(\sigma - \sigma'), \quad (3.23)$$

and if the charge \mathcal{Q}^M fulfils the *charge condition*

$$\mathcal{Q}^M \wedge dX^N \partial_N = \eta^{M,NP} \mathbf{Z}_P \partial_N. \quad (3.24)$$

In terms of the \mathcal{Q} alone it is, using (3.20), given by⁷

$$\mathcal{Q}^M \wedge dX^N \partial_N = \frac{1}{p} \eta^{M,NP} \eta_{K,LP} \mathcal{Q}^K \wedge dX^L \partial_N. \quad (3.26)$$

In order to find a solution for the charge \mathcal{Q}^M , it seems that the section condition needs to be imposed first. Only in special cases like $\text{SL}(5)$, it seems that one can solve the charge condition first and derive a section condition from a concrete charge.

Connecting to the previous results for $\text{SL}(5)$. For the 1- and 2-branes we can make contact with section 2 and [46]

$$p = 1: \quad q^M \equiv \mathbf{q}^M \in \mathcal{R}_2 \quad (3.27)$$

$$p = 2: \quad q_{M_1}^M = q_{M_1 M_2}^M \equiv \delta_{M_1 M_2}^{M M'} \mathbf{q}_{M'}, \quad \text{with } \mathbf{q}_{M'} \in \overline{\mathcal{R}}_2 = \mathcal{R}_3 \quad (3.28)$$

$$p = 3: \quad q_{M_1 M_2}^M = \epsilon^{M M_3 M_4} \epsilon_{N_1 M_1 M_3} \epsilon_{N_2 M_2 M_4} \mathbf{q}^{N_1 N_2} \quad \text{with } \mathbf{q}^{N_1 N_2} \in \bigwedge^2 \mathcal{R}_2 = \mathcal{R}_4. \quad (3.29)$$

The charge condition (3.26) implies

$$\mathbf{q}^M \partial_{M M'} = 0 \in \mathcal{R}_3, \quad \mathbf{q}_{[M_1} \partial_{M_2 M_3]} = 0 \in \mathcal{R}_4, \quad \mathbf{q}^{M_1 M_2} \partial_{M_1 M_2} = 0 \in \mathcal{R}_5, \quad (3.30)$$

using the $\text{SL}(5)$ representations \mathcal{R}_p . Motivated by that one might suggest that also for general d the charges of p -branes are elements of representations \mathcal{R}_{p+1} appearing in the tensor

⁷If multiplied by $\eta_{M,KL}$ the charge condition can be put in terms of the η_{KL} as

$$\eta_{KL} \wedge dX^M \partial_M = Y^{MN}{}_{KL} \mathbf{Z}_M \partial_N = \frac{1}{p} Y^{MN}{}_{KL} \eta_{MP} \wedge X^P \partial_N. \quad (3.25)$$

As such, the condition already appeared for the string ($p = 1$) in [46].

hierarchy, c.f. table 1, $\mathcal{Q}_{(p)} \in \mathcal{R}_{p+1}$ and there charge conditions to be elements in \mathcal{R}_{p+2} as was already done in [46]. This claim is true for the $\text{SL}(5)$ -theory as demonstrated above.

Apart for the $p = 3$ -case, D3-branes in the IIB theory, for which a similar embedding $\mathcal{Q}_{(3)} = \mathcal{R}_4 \in \wedge^2 \mathcal{R}_1 \otimes \mathcal{R}_2$ seems possible for arbitrary d , and the obvious $\mathcal{Q}_{(1)} \in \mathcal{R}_2$, there seems no necessity in our approach to the current algebra to assume that the charges or their charge condition should fit into the tensor hierarchy. We worked so far purely from the point of view of the theory in the internal space. Furthermore, the charges should exist also in cases, which are normally not considered to be part of the tensor hierarchy, as for example for the 5-brane multiplet in the $E_{6(6)}$, for which the charges then would take values in \mathcal{R}_6 . So, in the following, we will continue to consider the charges to be rather $\mathcal{Q}_{(p)} \in (\wedge^{p-1} \overline{\mathcal{R}}_1) \otimes \mathcal{R}_2$ as before, and not elements of \mathcal{R}_{p+1} .

3.3 Canonical p -form currents, their current algebra and $\text{SL}(p+3)$

The obvious generalisation of the type II string in $d = 3$ and membrane in $d = 4$ cases is the p -brane current in a $p+2$ -dimensional target space⁸

$$\mathbf{Z}_M(\sigma) = \left(p_\mu(\sigma) \star 1, \frac{1}{\sqrt{p!}} dx^{\mu_1} \wedge \dots \wedge dx^{\mu_p} \right). \quad (3.31)$$

Using the canonical Poisson structure, one obtains

$$\{\mathbf{Z}_M(\sigma), \mathbf{Z}_N(\sigma')\} = \mathcal{Q}^{\mathcal{L}} \wedge \left(\eta_{\mathcal{L},MN} \frac{1}{2} (d - d') + \omega_{\mathcal{L},MN} \frac{1}{2} (d + d') \right) \delta(\sigma - \sigma') \quad (3.32)$$

with η - and ω -symbols of the $\text{SL}(p+3)$ -group (3.2) and

$$\mathcal{Q}^{\mathcal{M}} = \left(\frac{1}{\sqrt{(p-1)!}} \mathcal{Q}^{\mu_1 \dots \mu_{p-1}}, 0 \right), \quad \mathcal{Q}^{\mu_1 \dots \mu_{p-1}} = dx^{\mu_1} \wedge \dots \wedge dx^{\mu_{p-1}}. \quad (3.33)$$

This is a solution to the charge condition (3.26) to the $\text{SL}(p+3)$ η -symbols and indeed the form of the current fits into the above system $\mathbf{Z}_M = \frac{1}{p} \eta_{\mathcal{K},LM} \mathcal{Q}^{\mathcal{K}} \wedge dX^{\mathcal{L}}$:

$$\mathbf{Z}_M = \left(\frac{1}{p!} dx^{\kappa_1} \wedge \dots \wedge dx^{\kappa_{p-1}} \wedge d\tilde{x}_{\kappa_1 \dots \kappa_{p-1} \kappa}, \frac{1}{\sqrt{p!}} dx^{\mu_1} \wedge \dots \wedge dx^{\mu_p} \right) \quad (3.34)$$

with coordinates $\tilde{x}_{\kappa_1 \dots \kappa_p}$ corresponding to the p -brane momentum (c.f. the string case – $p = \partial x$). When neglecting total derivative contributions (3.32) becomes

$$\{\mathbf{Z}_M(\sigma), \mathbf{Z}_N(\sigma')\} = \eta_{\mathcal{L},MN} \mathcal{Q}^{\mathcal{L}} \wedge d\delta(\sigma - \sigma'), \quad (3.35)$$

as wished, and gives the $\text{SL}(p+3)$ generalised Lie derivative (3.3), when applied on $\phi = -\int \phi^{\mathcal{K}} \mathbf{Z}_{\mathcal{K}}$. If, instead, one used (3.32), one reproduces the standard Lie derivative of the para-Hermitian generalised geometry on $(T \oplus \wedge^p T^*)M$ (3.18).

⁸Similar nice looking expressions can be obtained for any $d \geq p+1$, but only for $d = p+2$ the currents form a representation of an SL group, $\text{SL}(p+3)$.

Dirac brackets. As for the $O(d, d)$ string, an alternative derivation of (3.35) exists. This again employs Dirac brackets on an extended phase space with \mathcal{R}_1 momenta and coordinates

$$\begin{aligned}\{\mathbf{P}_M(\sigma), X^N(\sigma')\} &= -(p+1)\delta_M^N\delta(\sigma - \sigma') \\ \{\mathbf{P}_M(\sigma), \mathbf{P}_N(\sigma')\} &= \{X^M(\sigma), X^N(\sigma')\} = 0\end{aligned}\quad (3.36)$$

for constraints $\Phi_M = \mathbf{P}_M - \mathbf{Z}_M = \mathbf{P}_M - \frac{1}{p}\eta_{\mathcal{K}, LM}\mathcal{Q}^{\mathcal{K}} \wedge dX^L$. One finds, neglecting total world-volume derivatives,

$$\begin{aligned}A_{KL}(\sigma, \sigma') &= \{\mathbf{Z}_K(\sigma), \Phi_L(\sigma')\} = \{\mathbf{Z}_K(\sigma), \mathbf{P}_L(\sigma')\} \\ &= (p+1) \begin{pmatrix} \alpha_{\kappa\lambda} & \delta_{\kappa}^{[\lambda p} \mathcal{Q}^{\lambda_1 \dots \lambda_{p-1}]} \\ p\delta_{\lambda}^{[\kappa p} \mathcal{Q}^{\kappa_1 \dots \kappa_{p-1}]} & 0 \end{pmatrix} \wedge d\delta(\sigma - \sigma') \\ B_{KL}(\sigma, \sigma') &= \{\Phi_K(\sigma), \Phi_L(\sigma')\} = -\frac{(p+1)^2}{p}\eta_{\mathcal{M}, KL}\mathcal{Q}^{\mathcal{M}} \wedge d\delta(\sigma - \sigma')\end{aligned}$$

with $\alpha_{\kappa\lambda} = -\alpha_{\lambda\kappa}$. The Dirac brackets on (3.36) are, as striven for,

$$\begin{aligned}\{\mathbf{Z}_K(\sigma), \mathbf{Z}_L(\sigma')\}_{D.B.} &= -\int'' \int''' A_{KM}(\sigma, \sigma'')(B^{-1})^{MN}(\sigma'', \sigma''')A_{NL}(\sigma''', \sigma') \\ &= \eta_{\mathcal{M}, KL}\mathcal{Q}^{\mathcal{M}} \wedge d\delta(\sigma - \sigma'), \\ \{\mathbf{Z}_M(\sigma), X^N(\sigma')\}_{D.B.} &= -\delta_M^N\delta(\sigma - \sigma').\end{aligned}\quad (3.37)$$

3.4 Exceptional currents for $d \leq 6$

The previous setting was quite restrictive and only in part related to the exceptional duality groups that we want to geometrize. In particular, one can find solutions to the charge condition (3.26) for each $\frac{1}{2}$ -BPS object in type IIb and eleven-dimensional supergravity and construct potential currents of world-volume theories, such that their algebra respect the exceptional symmetries.

Solutions to the charge condition in $E_{d(d)}$. In the following, solutions of the charge condition and their corresponding currents $\mathbf{Z}_M = \frac{1}{p}\eta_{\mathcal{K}, LM}\mathcal{Q}^{\mathcal{K}} \wedge dX^L$ in an M-theory or type IIb section in $E_{d(d)}$ are collected.⁹ In contrast to the previously discussed $p = 1, 2$ cases in the $SL(5)$ -theory, it seems not to be possible to solve the condition without specifying the section first.

- In the **M-theory section**, we have directly from (3.24):

$$\mathbf{Z}_M = \left(Z_{\mu}, \frac{1}{\sqrt{2!}}\mathcal{Q}^{[\mu_1} \wedge dx^{\mu_2]}, \frac{1}{\sqrt{5!}}\mathcal{Q}^{[\mu_1\mu_2\mu_3\mu_4} \wedge dx^{\mu_5]} \right) \quad (3.38)$$

⁹The currents and charges for the **type IIa theory** objects – F1, D2, D4, NS5 – can be obtained dimensional reduction analogous to section 2.3:

- M2 \rightarrow F1, M5 \rightarrow D4: $x^d \equiv \sigma^p$, $x^{\underline{\mu}} = x^{\underline{\mu}}(\sigma^1, \dots, \sigma^{p-1})$, $\mu = (\underline{\mu}, d)$
- M2 \rightarrow D2, M5 \rightarrow NS5: $dx^d \equiv 0$

The point particle, $p = 0$, exists as well ($\mathcal{Q} = 0$) [45], but has no spatial world-volume and, hence, there is also no interesting current algebra defined on it.

– M2-brane:

$$\begin{aligned} Q^\mu &= q dx^\mu, & Q^{\mu_1\mu_2\mu_3\mu_4} &= 0 \\ \mathbf{Z}_M &= q \left(\frac{1}{2!} dx^\lambda \wedge d\tilde{x}_{\lambda\mu}, \frac{1}{\sqrt{2!}} dx^{\mu_1} \wedge dx^{\mu_2}, 0 \right) \end{aligned} \quad (3.39)$$

– M5-brane:

$$\begin{aligned} Q^\mu &= -\frac{q}{6} dx^\mu \wedge dx^{\mu_1} \wedge dx^{\mu_2} \wedge d\tilde{x}_{\mu_1\mu_2} \\ Q^{\mu_1\mu_2\mu_3\mu_4} &= q dx^{\mu_1} \wedge dx^{\mu_2} \wedge dx^{\mu_3} \wedge dx^{\mu_4} \\ \mathbf{Z}_M &= q \left(\frac{1}{5!} dx^{\mu_1} \wedge \dots \wedge dx^{\mu_4} \wedge d\tilde{x}_{\mu_1\dots\mu_4\mu} - \frac{1}{6 \cdot 5} dx^{\mu_1} \wedge \dots \wedge dx^{\mu_3} \wedge d\tilde{x}_{\mu_1\mu_2} \wedge d\tilde{x}_{\mu_3\mu} \right. \\ &\quad \left. \frac{1}{\sqrt{2!}} \frac{1}{6} dx^{\mu_1} \wedge dx^{\mu_2} \wedge dx^{\mu_3} \wedge dx^{\mu_4} \wedge d\tilde{x}_{\mu_3\mu_4}, \frac{1}{\sqrt{5!}} dx^{\mu_1} \wedge \dots \wedge dx^{\mu_5} \right) \end{aligned} \quad (3.40)$$

• In the type IIB section one derives from (3.24):

$$\mathbf{Z}_M = \left(Z_{\underline{\mu}}, Q_m \wedge dx^\mu, \frac{1}{\sqrt{3!}} Q^{\underline{\mu}_1\underline{\mu}_2} \wedge dx^{\underline{\mu}_3}, \frac{1}{\sqrt{5!}} Q_m^{\underline{\mu}_1\underline{\mu}_2\underline{\mu}_3\underline{\mu}_4} \wedge dx^{\underline{\mu}_5} \right) \quad (3.41)$$

– F1/D1-system resp. (p, q) -string

$$\begin{aligned} Q_m &= q_m = \begin{pmatrix} p \\ q \end{pmatrix}, & Q^{\underline{\mu}_1\underline{\mu}_2} &= 0, & Q_m^{\underline{\mu}_1\underline{\mu}_2\underline{\mu}_3\underline{\mu}_4} &= 0 \\ \mathbf{Z}_M &= q_m \left(d\tilde{x}_{\underline{\mu}}^m, dx^\mu, 0, 0 \right) \end{aligned} \quad (3.42)$$

– D3-brane:¹⁰

$$\begin{aligned} Q_m &= \frac{1}{2} q \epsilon_{mn} dx^\mu \wedge d\tilde{x}_{\underline{\mu}}^n, & Q^{\underline{\mu}_1\underline{\mu}_2} &= q dx^\mu \wedge dx^\nu, & Q_m^{\underline{\mu}_1\underline{\mu}_2\underline{\mu}_3\underline{\mu}_4} &= 0 \\ \mathbf{Z}_M &= q \left(\frac{1}{3!} \epsilon_{mn} dx^\nu \wedge d\tilde{x}_{\underline{\nu}}^m \wedge d\tilde{x}_{\underline{\mu}}^n - \frac{1}{3!} dx^{\underline{\mu}_1} \wedge dx^{\underline{\mu}_2} \wedge d\tilde{x}_{\underline{\mu}_1\underline{\mu}_2\underline{\mu}'} \right. \\ &\quad \left. \frac{1}{2} \epsilon_{mn} dx^\mu \wedge dx^\nu \wedge d\tilde{x}_{\underline{\mu}}^n, \frac{1}{\sqrt{3!}} dx^{\underline{\mu}_1} \wedge dx^{\underline{\mu}_2} \wedge dx^{\underline{\mu}_3}, 0 \right) \end{aligned} \quad (3.43)$$

– NS5/D5-brane:

$$\begin{aligned} Q_m &= \frac{1}{4!} q_l \left(3 \epsilon_{mn} dx^\nu \wedge d\tilde{x}_{\underline{\nu}}^n \wedge dx^\lambda \wedge d\tilde{x}_{\underline{\lambda}}^l - \delta_m^l dx^{\underline{\mu}_1} \wedge dx^{\underline{\mu}_2} \wedge dx^{\underline{\mu}_3} \wedge d\tilde{x}_{\underline{\mu}_1\underline{\mu}_2\underline{\mu}_3} \right) \\ Q^{\underline{\mu}_1\underline{\mu}_2} &= \frac{1}{2} q_m dx^{\underline{\mu}_1} \wedge dx^{\underline{\mu}_2} \wedge dx^{\underline{\mu}_3} \wedge d\tilde{x}_{\underline{\mu}_3}^m \\ Q_m^{\underline{\mu}_1\underline{\mu}_2\underline{\mu}_3\underline{\mu}_4} &= q_m dx^{\underline{\mu}_1} \wedge dx^{\underline{\mu}_2} \wedge dx^{\underline{\mu}_3} \wedge dx^{\underline{\mu}_4} \\ \mathbf{Z}_M &= \left(\frac{1}{5} \left[Q_m \wedge d\tilde{x}_{\underline{\mu}}^m + \frac{1}{2!} Q^{\underline{\mu}_1\underline{\mu}_2} \wedge d\tilde{x}_{\underline{\mu}_1\underline{\mu}_2\underline{\mu}} + \frac{1}{4!} Q_m^{\underline{\mu}_1\underline{\mu}_2\underline{\mu}_3\underline{\mu}_4} \wedge d\tilde{x}_{\underline{\mu}_1\underline{\mu}_2\underline{\mu}_3\underline{\mu}_4\underline{\mu}_5}^m \right], \right. \\ &\quad \left. - \frac{1}{4!} q_l dx^\mu \wedge dx^{\underline{\mu}_1} \wedge dx^{\underline{\mu}_2} \wedge \left(3 \epsilon_{mn} d\tilde{x}_{\underline{\mu}_1}^n \wedge d\tilde{x}_{\underline{\mu}_2}^l + \delta_m^l dx^{\underline{\mu}_3} \wedge d\tilde{x}_{\underline{\mu}_1\underline{\mu}_2\underline{\mu}_3} \right), \right. \\ &\quad \left. \frac{1}{\sqrt{3!}} \frac{1}{2} q_m dx^{\underline{\mu}_1} \wedge \dots \wedge dx^{\underline{\mu}_4} \wedge d\tilde{x}_{\underline{\mu}_4}^m, \frac{1}{\sqrt{5!}} q_m dx^{\underline{\mu}_1} \wedge \dots \wedge dx^{\underline{\mu}_5} \right) \end{aligned} \quad (3.44)$$

¹⁰As suggested by (3.29) there might be an embedding of $\mathcal{R}_4 = \wedge^2 \mathcal{R}_2$ of this charge solution. This is indeed the case - the only non-zero component of $\mathbf{q}^{\mathcal{M}_1\mathcal{M}_2}$ is $\mathbf{q}_{m_1m_2} = \frac{1}{2} q \epsilon_{m_1m_2}$ in the IIB section. With (3.29) one reproduces $Q^{\mathcal{M}}$ of the D3-brane.

Suggested by these charge solutions, we see the typical decomposition of \mathcal{R}_1 and \mathcal{R}_2 in the M-theory and type IIb sections

$$\begin{aligned} \mathbf{Z}_M &= \left(Z_\mu, \underbrace{\frac{\mathbf{Z}^{\mu_1\mu_2}}{\sqrt{2!}}}_{\text{M2}}, \underbrace{\frac{\mathbf{Z}^{\mu_1\mu_2\mu_3\mu_4\mu_5}}{\sqrt{5!}}}_{\text{M5}} \right), & \mathcal{Q}^M &= \left(\underbrace{Q^\mu}_{\text{M2}}, \underbrace{\frac{Q^{\mu_1\mu_2\mu_3\mu_4}}{\sqrt{4!}}}_{\text{M5}}, \underbrace{\frac{Q^{\mu_1\mu_2\mu_3\mu_4\mu_5\mu_6,\mu}}{\sqrt{6!}}}_{\text{KKM}} \right) \\ \mathbf{Z}_M &= \left(Z_\mu, \underbrace{Z_m^\mu}_{\text{F1/D1}}, \underbrace{\frac{\mathbf{Z}^{\mu_1\mu_2\mu_3}}{\sqrt{3!}}}_{\text{D3}}, \underbrace{\frac{\mathbf{Z}^{\mu_1\mu_2\mu_3\mu_4\mu_5}}{\sqrt{5!}}}_{\text{NS5/D5}} \right), & \mathcal{Q}^M &= \left(\underbrace{Q_m}_{\text{F1/D1}}, \underbrace{\frac{Q^{\mu_1\mu_2}}{\sqrt{2!}}}_{\text{D3}}, \underbrace{\frac{Q_m^{\mu_1\mu_2\mu_3\mu_4}}{\sqrt{4!}}}_{\text{NS5/D5}}, \underbrace{\frac{Q^{\mu_1\mu_2\mu_3\mu_4\mu_5,\mu}}{\sqrt{5!}}}_{\text{KKM}} \right) \end{aligned} \quad (3.45)$$

The already existing place for the Kalazu-Klein monopole (KKM) in the \mathcal{R}_2 -representation is empty in the above collected list of currents. It is simply not a solution to the charge condition (3.26) for $d \leq 6$ yet. The decomposition of the \mathbf{Z} is similar to the one of the generalised coordinates in X^M . For both sections the x^μ and $Z_\mu = p_\mu$ -components are shared in all currents.

The *non-geometric coordinates* of X^M , i.e. the ones that do not belong to a section, could be associated to the momenta of the corresponding objects – in the same decomposition as the one for the \mathbf{Z} . This is analogous to the string in $O(d, d)$ generalised geometry, where $p_\mu = d\tilde{x}_\mu$ [24, 27, 65].

The non-geometry of the currents. As already noted previously [45, 48], where σ -model constructions of such world-volume theories were considered, we observe that for the ‘higher’ ($p > 2$) world-volume not only the corresponding Q -component but also all the ‘lower’ ones are turned on – e.g. for the M5-brane (3.40) not only $Q^{\mu_1\cdots\mu_4}$ but also Q^μ . In the end, the reason for that is $E_{d(d)}$ -covariance.

In [45, 48] this was done by introducing auxiliary gauge fields. Here, instead, these components contain differentials of *non-geometric coordinates*, that are associated to the momenta of the lower dimensional branes. In the example of the NS/D5-brane (3.44), the D3-charge and current contain the differential $d\tilde{x}_{\mu_4}^m$ and so on.

Consequently, from the point of view of such a higher brane, whose phase space coordinate normally just would be x^μ and $p_\mu = Z_\mu$, the currents itself already are non-geometric as also the additional non-geometric coordinates \tilde{x} appear inevitably. Finally, such terms are logical to appear if one strives for an $E_{d(d)}$ -covariant setup: As for example the M2- and M5-brane or the D-branes are connected by duality, in general M5 currents can’t appear alone in such a setup. Another way to understand that fact is, that in the generalised Lie derivative for $d > 4$ (in the M-theory picture) a ‘coupling’ of two-forms appears: $[\omega_1, \omega_2]_D = \omega_1 \wedge d\omega_2$ for two-forms ω_i . Hence, the current algebra bracket of two M2-currents forms an M5-current, necessitating that in presence of an M5-current also M2-currents have to be present. Interestingly, the converse does not seem to be true: one can formulate an M2-brane current algebra, by consistently setting the (canonical) M5-current to zero.

Deriving the current algebra? As the generalised Lie derivative for $E_{d(d)}$ for $d \geq 5$ is not equivalent to the standard Lie derivative up to total derivatives, *one cannot hope to derive this current algebra* in a similar way to the p -brane in $SL(p+3)$ or the string in $O(d, d)$. Neither starting from canonical Poisson brackets nor a Dirac bracket approach can lead to a non-Lie bracket. The only reason, why one could make it work in the previous cases, is that there the

difference is always only a total-derivative term on the spatial world-volume, that would not contribute classically.

It is possible to derive a para-Hermitian $E_{d(d)}$ current, though. For example, for $d = 5, 6$, it could simply be realised as a 'sum' of the $p = 2$ - and $p = 5$ -currents resp. their corresponding $SL(p + 3)$ -theory

$$\mathbf{Z}_M = \mathbf{Z}_M^{(2)}(\sigma) + \mathbf{Z}_M^{(5)}(\rho) \sim \left(p_\mu^{(2)}(\sigma) + p_\mu^{(5)}(\rho), dx^{\mu_1}(\sigma) \wedge dx^{\mu_2}(\sigma), dx^{\mu_1}(\rho) \wedge \dots \wedge dx^{\mu_5}(\rho) \right)$$

in the M-theory section. They correspond to decoupled M2 and M5 world-volume theories, without exhibiting the exceptional symmetry at the classical level.

4 Outlook

There are several conceptual questions left open in this article. The main result, the $E_{d(d)}$ -covariant current algebra

$$\{\mathbf{Z}_M(\sigma), \mathbf{Z}_N(\sigma')\} = \eta_{\mathcal{L}, MN} \mathcal{Q}^{\mathcal{L}} \wedge d\delta(\sigma - \sigma') \quad (4.1)$$

of a p -brane current was postulated but could not be derived from some underlying principle, like a canonical (Poisson) structure, in general. It would be nice to change that by considering higher algebraic structures, like aspects of dg-symplectic geometry, L_∞ algebras or AKSZ constructions, see for example [77] for similar approaches in the context of exceptional generalised geometry. On a more fundamental level, one might ask whether a canonical phase space structure could ever capture the symmetries of M-theory, or whether one should take a structure like (4.1) to be the fundamental one.

The current algebra was constructed as a simple way to write down an $E_{d(d)}$ -invariant world-volume theory, motivated by the membrane and string in the $SL(5)$ -theory where it could be derived from the standard Polyakov type action, when neglecting total world-sheet derivatives. A similar clean path to an action could be possible for D-branes, following [60]. For $p > 2$ the difference between the *canonical* and the $E_{d(d)}$ -current algebra is not a total world-volume differential anymore (as in the string and the membrane case). This relates directly to discussions at the level of the action, where such topological contributions have been shown to appear as well, as difference between duality invariant and standard versions of actions [1, 62–64]. Hence, for $p > 2$ severe modification of actions in contrast to the normal Polyakov Nambu-Goto type ones are to be expected, when demanding $E_{d(d)}$ covariance. Several candidates of such $E_{d(d)}$ -invariant world-volume actions have been discussed in the literature [43–46, 48]. It would be interesting to study whether they would correspond to the Hamiltonian settings presented in this paper. This would also clarify, whether the conjectured Hamiltonian and spatial diffeomorphism

$$H = \frac{1}{2} \int \mathbf{Z}_K \wedge \star \mathbf{Z}_L \mathcal{H}^{KL}, \quad 0 = \eta^{\mathcal{M}, KL} \mathbf{Z}_K \wedge \star \mathbf{Z}_L. \quad (4.2)$$

make sense for $p > 2$ and $d > 4$ as well. The Hamiltonian formulation of membranes in a duality invariant way might be helpful in context of the recent study of exceptional Drinfeld algebras [50–59], in particular in the question whether certain transformations, e.g. polyvector shifts a.k.a. generalised Yang-Baxter deformations, of Drinfeld algebras really correspond to dualities of some world-volume theory, generalising work of [50] to generic

objects and $d > 4$. But one should not forget that the current algebra (4.1) is, in general, not a Lie algebra. Tools of conventional classical field theory, like Legendre transformations or canonical transformations, have to be reconsidered. Generalising techniques of geometric quantisation from the recently studies $O(d, d)$ case [78] could be necessary, as well.

The fact, that all the $p > 2$ exceptional currents in section 3.4 are manifestly non-geometric, is a puzzling results. Perhaps one must consider multiple interacting world-volume theories at once, e.g. M2- and M5-branes, to make sense out of that Hamiltonian theory. Potentially this might help resolving the famous membrane duality problem [41, 42]. In the string case, the study of the current algebra was feasible in order interpret backgrounds as non-commutative and non-associative ones [24, 65]. Integrating (4.1) could give a similar picture for M-theory backgrounds. The picture should be more involved, as one would expect all kinds of higher brackets to appear.

Generalisations of the present setup to $E_{7(7)}$ and $E_{8(8)}$ generalised geometry should be possible as well. The additional duality group invariants, that appear in the generalised Lie derivatives there, have to be introduced into the current algebra (4.1). Extending the analysis to include the exterior space and tensor hierarchies seems straightforward.

For the membrane in the $SL(5)$ -theory it was recently shown [79] that the solution to the classical world-volume theory can be interpreted as generalised geodesics in $SL(5)$ generalised geometry. Studying the equations of motion for the $E_{d(d)}$ - and $SL(p+3)$ -currents with the above Hamiltonian, might extend that picture to higher dimensional objects and higher space-time dimension. This might give a geometric interpretation of the charge \mathcal{Q} . So far, it only appears as a auxiliary object to ensure manifest $E_{d(d)}$ -covariance.

Acknowledgements

I thank Alex Arvanitakis, Chris Blair and Daniel Thompson for discussions at the start of this project. Also, I thank the Max-Planck-Institute for Physics Munich where a large portion of this work was performed.

A Conventions and identities

We define the generalised Kronecker symbols as follows:

$$\epsilon^{i_1 \dots i_k j_1 \dots j_{n-k}} \epsilon_{i_1 \dots i_k k_1 \dots k_{n-k}} = k! \delta_{k_1 \dots k_{n-k}}^{j_1 \dots j_{n-k}} = k! (n-k)! \delta_{[k_1 \dots k_{n-k}]}^{[j_1 \dots j_{n-k}]}$$

The standard δ -distribution $f(\sigma) = \int d\sigma' f(\sigma') \delta(\sigma - \sigma')$ behaves in a maybe unexpected way, when applied to functions on compact spaces. In particular $(\partial_1 + \partial_2) \delta(\sigma_2 - \sigma_1) \neq 0$, but instead

$$\int d\sigma_1 d\sigma_2 \phi(\sigma_1, \sigma_2) (\partial_1 + \partial_2) \delta(\sigma_2 - \sigma_1) = \int d\sigma \partial (\phi(\sigma, \sigma)). \quad (\text{A.1})$$

As discussed in the main text, this term is not vanishing in general. Strings and membranes can have non-trivial winding around a non-trivial cycle in target space. In that case, the coordinate fields $x(\sigma)$ are not smooth, such that in particular $\int d\sigma \partial x(\sigma) \neq 0$. For open world-volumes there are boundary contributions, as well.

In many calculations it is helpful to write

$$\delta(\sigma_2 - \sigma_1) = \int d\sigma \delta(\sigma - \sigma_1) \delta(\sigma - \sigma_2), \quad (\text{A.2})$$

to see for example that $(\partial_1 + \partial_2)\delta(\sigma_2 - \sigma_1) = \int d\sigma \partial \delta(\sigma - \sigma_1) \delta(\sigma - \sigma_2)$. The following distributional identities follow:

$$\begin{aligned}
& \int d\sigma f(\sigma) (\partial_1 + \dots + \partial_n) (\delta(\sigma - \sigma_1) \cdot \dots \cdot \delta(\sigma - \sigma_n)) \\
&= \int d\sigma ((\partial f(\sigma)) + (n-1)f(\sigma)\partial) (\delta(\sigma - \sigma_1) \cdot \dots \cdot \delta(\sigma - \sigma_n)) \\
& \frac{1}{2} e(\sigma_1) \cdot e^{-1}(\sigma_2) (\partial_1 - \partial_2) (\delta(\sigma - \sigma_1) \delta(\sigma - \sigma_2)) \\
&= \frac{1}{2} (\partial_1 - \partial_2) (\delta(\sigma - \sigma_1) \delta(\sigma - \sigma_2)) \mathbb{1} - ((\partial e) \cdot e^{-1})(\sigma) \delta(\sigma - \sigma_1) \delta(\sigma - \sigma_2) \\
& (f(\sigma_2)\partial_1 + f(\sigma_1)\partial_2) \delta(\sigma - \sigma_1) \delta(\sigma - \sigma_2) \\
&= (\partial f(\sigma)) \delta(\sigma - \sigma_1) \delta(\sigma - \sigma_2) + \partial(f(\sigma) \delta(\sigma - \sigma_1) \delta(\sigma - \sigma_2))
\end{aligned} \tag{A.3}$$

for arbitrary (matrix-valued) functions e and f which hold without any additional boundary terms.

The higher world-volume currents are written directly as forms (on the spatial world-volume). Typical important identities are

$$\begin{aligned}
dx^\mu(\sigma) \wedge d\delta(\sigma - \sigma') &= dx^\mu(\sigma') \wedge d\delta(\sigma - \sigma') \\
\omega(\sigma') \wedge d\delta(\sigma - \sigma') &= \omega(\sigma) \wedge d\delta(\sigma - \sigma') + (d\omega)(\sigma) \delta(\sigma - \sigma') \quad \text{for 1-forms } \omega.
\end{aligned}$$

B The $E_{d(d)}$ η - and ω -symbols for $d \leq 6$

For $d \leq 6$ the Y -tensor can be written in terms of the η -symbols:

$$Y^{KL}{}_{MN} = \eta^{P,KL} \eta_{P,MN} \tag{B.1}$$

when we choose the normalisation

$$\eta^{M,KL} \eta_{N,KL} = 2(d-1) \delta_N^M. \tag{B.2}$$

Indices K, L, M, \dots resp. $\mathcal{K}, \mathcal{L}, \mathcal{M}, \dots$ denote \mathcal{R}_1 - resp. \mathcal{R}_2 -indices of the duality group $E_{d(d)}$.

In the following, we give the η - and ω -symbols explicitly in the conventions in which contractions of the totally antisymmetric multi-indices $^{\mu_1 \dots \mu_p}$ receiving a factor $\frac{1}{p!}$. Let us note, that for $d > 4$ the ω -symbols are not skewsymmetric anymore: $\omega_{\mathcal{M},KL} \neq -\omega_{\mathcal{M},LK}$.

- M-theory decomposition with target space indices $\kappa, \lambda, \mu, \dots = 1, \dots, d$:

$$\begin{aligned}
X^{\mathcal{M}} &= \left(x^\mu, \frac{\tilde{x}^{\mu_1 \mu_2}}{\sqrt{2!}}, \frac{\tilde{x}^{\mu_1 \mu_2 \mu_3 \mu_4 \mu_5}}{\sqrt{5!}} \right) \in \mathcal{R}_1 \\
\mathcal{Q}^{\mathcal{M}} &= \left(Q^\mu, \frac{Q^{\mu_1 \mu_2 \mu_3 \mu_4}}{\sqrt{4!}}, \frac{Q^{\mu_1 \mu_2 \mu_3 \mu_4 \mu_5 \mu_6, \mu}}{\sqrt{6!}} \right) \in \mathcal{R}_2
\end{aligned}$$

$$\begin{aligned}
\eta_{\mathcal{M},KL} : \quad \eta_{\mu,KL} &= \begin{pmatrix} 0 & \frac{\delta^{\lambda_1 \lambda_2}}{\sqrt{2!}} & 0 \\ \frac{\delta^{\kappa_1 \kappa_2}}{\sqrt{2!}} & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \\
\eta_{\mu_1 \mu_2 \mu_3 \mu_4, KL} &= \begin{pmatrix} 0 & 0 & \frac{\delta^{\lambda_1 \lambda_2 \lambda_3 \lambda_4 \lambda_5}}{\sqrt{5!}} \\ 0 & \frac{\delta^{\kappa_1 \kappa_2 \lambda_1 \lambda_2}}{\sqrt{2!} \sqrt{2!}} & 0 \\ \frac{\delta^{\kappa_1 \kappa_2 \kappa_3 \kappa_4 \kappa_5}}{\sqrt{5!}} & 0 & 0 \end{pmatrix} \\
(\eta_{\mu_1 \mu_2 \mu_3 \mu_4 \mu_5 \mu_6, \mu})_{KL} &= \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & \frac{\delta^{\lambda_1 \lambda_2 \lambda_3 \lambda_4 \lambda_5 \nu} \delta^{\nu \mu}}{\sqrt{5!} \sqrt{2!}} \\ 0 & \frac{\delta^{\kappa_1 \kappa_2 \kappa_3 \kappa_4 \kappa_5 \nu} \delta^{\nu \mu}}{\sqrt{5!} \sqrt{2!}} & 0 \end{pmatrix} \\
\omega_{\mathcal{M},KL} : \quad \omega_{\mu,KL} &= \begin{pmatrix} 0 & -\frac{\delta^{\lambda_1 \lambda_2}}{\sqrt{2!}} & 0 \\ \frac{\delta^{\kappa_1 \kappa_2}}{\sqrt{2!}} & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \\
\omega_{\mu_1 \mu_2 \mu_3 \mu_4, KL} &= \begin{pmatrix} 0 & 0 & -\frac{\delta^{\lambda_1 \lambda_2 \lambda_3 \lambda_4 \lambda_5}}{\sqrt{5!}} \\ 0 & -\frac{\delta^{\kappa_1 \kappa_2 \lambda_1 \lambda_2}}{\sqrt{2!} \sqrt{2!}} & 0 \\ \frac{\delta^{\kappa_1 \kappa_2 \kappa_3 \kappa_4 \kappa_5}}{\sqrt{5!}} & 0 & 0 \end{pmatrix} \\
(\omega_{\mu_1 \mu_2 \mu_3 \mu_4 \mu_5 \mu_6, \mu})_{KL} &= \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & -\frac{\delta^{\lambda_1 \lambda_2 \lambda_3 \lambda_4 \lambda_5 \nu} \delta^{\nu \mu}}{\sqrt{5!} \sqrt{2!}} \\ 0 & -\frac{\delta^{\kappa_1 \kappa_2 \kappa_3 \kappa_4 \kappa_5 \nu} \delta^{\nu \mu}}{\sqrt{5!} \sqrt{2!}} & 0 \end{pmatrix}
\end{aligned}$$

- Type IIb decomposition with $SL(2)$ -indices $k, l, m, \dots = 1, 2$ and target-space indices $\underline{\kappa}, \underline{\lambda}, \underline{\mu}, \dots = 1, \dots, d-1$:

$$\begin{aligned}
X^M &= \left(x^\mu, \tilde{x}_\mu^m, \frac{\tilde{x}_{\mu_1 \mu_2 \mu_3}^m}{\sqrt{3!}}, \frac{\tilde{x}_{\mu_1 \mu_2 \mu_3 \mu_4 \mu_5}^m}{\sqrt{5!}} \right) \in \mathcal{R}_1 \\
Q^{\mathcal{M}} &= \left(Q_m, \frac{Q_{\mu_1 \mu_2}^m}{\sqrt{2!}}, \frac{Q_{\mu_1 \mu_2 \mu_3 \mu_4}^m}{\sqrt{4!}}, \frac{Q_{\mu_1 \mu_2 \mu_3 \mu_4 \mu_5 \mu_6}^m}{\sqrt{5!}} \right) \in \mathcal{R}_2
\end{aligned}$$

$$\begin{aligned}
\eta_{\mathcal{M},KL} : \quad (\eta^m)_{KL} &= \begin{pmatrix} 0 & \delta_l^m \delta_{\underline{\kappa}}^\lambda & 0 & 0 \\ \delta_k^m \delta_{\underline{\lambda}}^\kappa & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix} \\
(\eta_{\underline{\mu}_1 \underline{\mu}_2})_{KL} &= \begin{pmatrix} 0 & 0 & \frac{\delta_{\underline{\mu}_1 \underline{\mu}_2 \underline{\kappa}}^{\underline{\lambda}_1 \underline{\lambda}_2 \underline{\lambda}_3}}{\sqrt{3!}} & 0 \\ 0 & \epsilon_{kl} \delta_{\underline{\mu}_1 \underline{\mu}_2}^{\underline{\kappa} \underline{\lambda}} & 0 & 0 \\ \frac{\delta_{\underline{\mu}_1 \underline{\mu}_2 \underline{\lambda}}^{\underline{\lambda}_1 \underline{\lambda}_2 \underline{\lambda}_3}}{\sqrt{3!}} & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix} \\
(\eta_{\underline{\mu}_1 \underline{\mu}_2 \underline{\mu}_3 \underline{\mu}_4}^m)_{KL} &= \begin{pmatrix} 0 & 0 & 0 & \frac{\delta_l^m \delta_{\underline{\mu}_1 \underline{\mu}_2 \underline{\mu}_3 \underline{\mu}_4 \underline{\kappa}}^{\underline{\lambda}_1 \underline{\lambda}_2 \underline{\lambda}_3 \underline{\lambda}_4 \underline{\lambda}_5}}{\sqrt{5!}} \\ 0 & 0 & \frac{\delta_k^m \delta_{\underline{\mu}_1 \underline{\mu}_2 \underline{\mu}_3 \underline{\mu}_4}^{\underline{\kappa} \underline{\lambda}_1 \underline{\lambda}_2 \underline{\lambda}_3}}{\sqrt{3!}} & 0 \\ 0 & \frac{\delta_l^m \delta_{\underline{\mu}_1 \underline{\mu}_2 \underline{\mu}_3 \underline{\mu}_4}^{\underline{\lambda}_1 \underline{\lambda}_2 \underline{\lambda}_3 \underline{\kappa}}}{\sqrt{3!}} & 0 & 0 \\ \frac{\delta_k^m \delta_{\underline{\mu}_1 \underline{\mu}_2 \underline{\mu}_3 \underline{\mu}_4 \underline{\lambda}}^{\underline{\kappa}_1 \underline{\kappa}_2 \underline{\kappa}_3 \underline{\kappa}_4 \underline{\kappa}_5}}{\sqrt{5!}} & 0 & 0 & 0 \end{pmatrix} \\
(\eta_{\underline{\mu}_1 \underline{\mu}_2 \underline{\mu}_3 \underline{\mu}_4 \underline{\mu}_5}^m)_{KL} &= \begin{pmatrix} 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & \epsilon^{lk} \frac{\delta_{\underline{\mu}_1 \underline{\mu}_2 \underline{\mu}_3 \underline{\mu}_4 \underline{\mu}_5}^{\underline{\kappa} \underline{\lambda}_1 \underline{\lambda}_2 \underline{\lambda}_3 \underline{\lambda}_4 \underline{\lambda}_5}}{\sqrt{5!}} \\ 0 & 0 & \frac{1}{2} \left(\frac{\delta_{\underline{\mu}_1 \underline{\mu}_2 \underline{\mu}_3 \underline{\mu}_4 \underline{\mu}_5}^{\underline{\kappa}_1 \underline{\kappa}_2 \underline{\kappa}_3 \underline{\nu}_1 \underline{\nu}_2} \delta_{\underline{\nu}_1 \underline{\nu}_2 \underline{\mu}}^{\underline{\lambda}_1 \underline{\lambda}_2 \underline{\lambda}_3}}{2! \sqrt{3! 3!}} + (\underline{\kappa} \leftrightarrow \underline{\lambda}) \right) & 0 & 0 \\ 0 & \epsilon^{kl} \frac{\delta_{\underline{\mu}_1 \underline{\mu}_2 \underline{\mu}_3 \underline{\mu}_4 \underline{\mu}_5}^{\underline{\lambda}_1 \underline{\kappa}_1 \underline{\kappa}_2 \underline{\kappa}_3 \underline{\kappa}_4 \underline{\kappa}_5}}{\sqrt{5!}} & 0 & 0 & 0 \end{pmatrix}
\end{aligned}$$

The ω -symbols are given by the same terms as the η -symbols, but with different signs, as in the M-theory section as well:

$$\omega_{\bullet, MN} = \begin{pmatrix} 0 & - & - & - \\ + & - & - & - \\ + & - & - & 0 \\ + & - & 0 & 0 \end{pmatrix}.$$

References

- [1] A. A. Tseytlin, *Duality Symmetric Formulation of String World Sheet Dynamics*, *Phys. Lett.* **B242** (1990) 163–174.
- [2] A. A. Tseytlin, *Duality symmetric closed string theory and interacting chiral scalars*, *Nucl. Phys.* **B350** (1991) 395–440.
- [3] W. Siegel, *Manifest duality in low-energy superstrings*, in *International Conference on Strings 93 Berkeley, California, May 24-29, 1993*, pp. 353–363, 1993. hep-th/9308133.
- [4] W. Siegel, *Superspace duality in low-energy superstrings*, *Phys. Rev.* **D48** (1993) 2826–2837, [hep-th/9305073].

- [5] N. Hitchin, *Generalized Calabi-Yau manifolds*, *Quart. J. Math.* **54** (2003) 281–308, [math/0209099].
- [6] M. Gualtieri, *Generalized complex geometry*. PhD thesis, Oxford U., 2003. math/0401221.
- [7] M. Graña, R. Minasian, M. Petrini, and D. Waldram, *T-duality, Generalized Geometry and Non-Geometric Backgrounds*, *JHEP* **04** (2009) 075, [arXiv:0807.4527].
- [8] C. Hull and B. Zwiebach, *Double Field Theory*, *JHEP* **09** (2009) 099, [arXiv:0904.4664].
- [9] B. Zwiebach, *Double Field Theory, T-Duality, and Courant Brackets*, *Lect. Notes Phys.* **851** (2012) 265–291, [arXiv:1109.1782].
- [10] G. Aldazabal, D. Marqués, and C. Nuñez, *Double Field Theory: A Pedagogical Review*, *Class. Quant. Grav.* **30** (2013) 163001, [arXiv:1305.1907].
- [11] O. Hohm, D. Lüst, and B. Zwiebach, *The Spacetime of Double Field Theory: Review, Remarks, and Outlook*, *Fortsch. Phys.* **61** (2013) 926–966, [arXiv:1309.2977].
- [12] D. Geissbühler, D. Marqués, C. Nuñez, and V. Penas, *Exploring Double Field Theory*, *JHEP* **06** (2013) 101, [arXiv:1304.1472].
- [13] E. Plauschinn, *Non-geometric backgrounds in string theory*, *Phys. Rept.* **798** (2019) 1–122, [arXiv:1811.11203].
- [14] C. Klimčík and P. Ševera, *Poisson-Lie T duality and loop groups of Drinfeld doubles*, *Phys. Lett.* **B372** (1996) 65–71, [hep-th/9512040].
- [15] C. Klimčík and P. Ševera, *NonAbelian momentum winding exchange*, *Phys. Lett.* **B383** (1996) 281–286, [hep-th/9605212].
- [16] D. Osten and S. J. van Tongeren, *Abelian Yang-Baxter deformations and TsT transformations*, *Nucl. Phys.* **B915** (2017) 184–205, [arXiv:1608.08504].
- [17] F. Hassler, *Poisson-Lie T-Duality in Double Field Theory*, arXiv:1707.08624.
- [18] J.-i. Sakamoto, Y. Sakatani, and K. Yoshida, *Homogeneous Yang-Baxter deformations as generalized diffeomorphisms*, *J. Phys.* **A50** (2017), no. 41 415401, [arXiv:1705.07116].
- [19] R. Borsato and L. Wulff, *On non-abelian T-duality and deformations of supercoset string sigma-models*, *JHEP* **10** (2017) 024, [arXiv:1706.10169].
- [20] J. J. Fernandez-Melgarejo, J.-i. Sakamoto, Y. Sakatani, and K. Yoshida, *T-folds from Yang-Baxter deformations*, *JHEP* **12** (2017) 108, [arXiv:1710.06849].
- [21] J.-I. Sakamoto and Y. Sakatani, *Local β -deformations and Yang-Baxter sigma model*, *JHEP* **06** (2018) 147, [arXiv:1803.05903].
- [22] D. Lüst and D. Osten, *Generalised fluxes, Yang-Baxter deformations and the $O(d,d)$ structure of non-abelian T-duality*, *JHEP* **05** (2018) 165, [arXiv:1803.03971].
- [23] S. Demulder, F. Hassler, and D. C. Thompson, *Doubled aspects of generalised dualities and integrable deformations*, *JHEP* **02** (2019) 189, [arXiv:1810.11446].

- [24] D. Osten, *Current algebras, generalised fluxes and non-geometry*, *J. Phys. A* **53** (2020), no. 26 265402, [arXiv:1910.00029].
- [25] D. C. Thompson, *An Introduction to Generalised Dualities and their Applications to Holography and Integrability*, in *18th Hellenic School and Workshops on Elementary Particle Physics and Gravity (CORFU2018) Corfu, Corfu, Greece, August 31-September 28, 2018*, 2019. arXiv:1904.11561.
- [26] R. Borsato and S. Driezen, *Supergravity solution-generating techniques and canonical transformations of σ -models from $O(D, D)$* , arXiv:2102.04498.
- [27] C. D. A. Blair, E. Malek, and A. J. Routh, *An $O(D, D)$ invariant Hamiltonian action for the superstring*, *Class. Quant. Grav.* **31** (2014), no. 20 205011, [arXiv:1308.4829].
- [28] K. Sfetsos, *Canonical equivalence of nonisometric sigma models and Poisson-Lie T duality*, *Nucl. Phys.* **B517** (1998) 549–566, [hep-th/9710163].
- [29] C. Klimčík, *η and λ deformations as \mathcal{E} -models*, *Nucl. Phys.* **B900** (2015) 259–272, [arXiv:1508.05832].
- [30] P. Pires Pacheco and D. Waldram, *M-theory, exceptional generalised geometry and superpotentials*, *JHEP* **09** (2008) 123, [arXiv:0804.1362].
- [31] D. S. Berman and M. J. Perry, *Generalized Geometry and M theory*, *JHEP* **06** (2011) 074, [arXiv:1008.1763].
- [32] D. S. Berman, H. Godazgar, M. J. Perry, and P. West, *Duality Invariant Actions and Generalised Geometry*, *JHEP* **02** (2012) 108, [arXiv:1111.0459].
- [33] A. Coimbra, C. Strickland-Constable, and D. Waldram, *$E_{d(d)} \times \mathbb{R}^+$ generalised geometry, connections and M theory*, *JHEP* **02** (2014) 054, [arXiv:1112.3989].
- [34] D. S. Berman, M. Cederwall, A. Kleinschmidt, and D. C. Thompson, *The gauge structure of generalised diffeomorphisms*, *JHEP* **01** (2013) 064, [arXiv:1208.5884].
- [35] D. S. Berman, E. T. Musaev, D. C. Thompson, and D. C. Thompson, *Duality Invariant M-theory: Gauged supergravities and Scherk-Schwarz reductions*, *JHEP* **10** (2012) 174, [arXiv:1208.0020].
- [36] A. Coimbra, C. Strickland-Constable, and D. Waldram, *Supergravity as Generalised Geometry II: $E_{d(d)} \times \mathbb{R}^+$ and M theory*, *JHEP* **03** (2014) 019, [arXiv:1212.1586].
- [37] O. Hohm and H. Samtleben, *Exceptional Form of D=11 Supergravity*, *Phys. Rev. Lett.* **111** (2013) 231601, [arXiv:1308.1673].
- [38] K. Lee, C. Strickland-Constable, and D. Waldram, *Spheres, generalised parallelisability and consistent truncations*, *Fortsch. Phys.* **65** (2017), no. 10-11 1700048, [arXiv:1401.3360].
- [39] O. Hohm and H. Samtleben, *Consistent Kaluza-Klein Truncations via Exceptional Field Theory*, *JHEP* **01** (2015) 131, [arXiv:1410.8145].
- [40] D. S. Berman and C. D. A. Blair, *The Geometry, Branes and Applications of Exceptional Field Theory*, *Int. J. Mod. Phys. A* **35** (2020), no. 30 2030014, [arXiv:2006.09777].

- [41] M. J. Duff, *Duality Rotations in String Theory*, *Nucl. Phys.* **B335** (1990) 610.
- [42] M. J. Duff, J. X. Lu, R. Percacci, C. N. Pope, H. Samtleben, and E. Sezgin, *Membrane Duality Revisited*, *Nucl. Phys.* **B901** (2015) 1–21, [arXiv:1509.02915].
- [43] Y. Sakatani and S. Uehara, *Branes in Extended Spacetime: Brane Worldvolume Theory Based on Duality Symmetry*, *Phys. Rev. Lett.* **117** (2016), no. 19 191601, [arXiv:1607.04265].
- [44] C. D. A. Blair and E. T. Musaev, *Five-brane actions in double field theory*, *JHEP* **03** (2018) 111, [arXiv:1712.01739].
- [45] Y. Sakatani and S. Uehara, *Exceptional M-brane sigma models and η -symbols*, *PTEP* **2018** (2018), no. 3 033B05, [arXiv:1712.10316].
- [46] A. S. Arvanitakis and C. D. Blair, *The Exceptional Sigma Model*, *JHEP* **04** (2018) 064, [arXiv:1802.00442].
- [47] C. D. A. Blair, *Open exceptional strings and D-branes*, *JHEP* **07** (2019) 083, [arXiv:1904.06714].
- [48] Y. Sakatani and S. Uehara, *Born sigma model for branes in exceptional geometry*, arXiv:2004.09486.
- [49] M. Hatsuda and K. Kamimura, *SL(5) duality from canonical M2-brane*, *JHEP* **11** (2012) 001, [arXiv:1208.1232].
- [50] Y. Sakatani and S. Uehara, *Non-Abelian U-duality for membranes*, *PTEP* **2020** (2020), no. 7 073B01, [arXiv:2001.09983].
- [51] Y. Sakatani, *Extended Drinfel'd algebras and non-Abelian duality*, arXiv:2009.04454.
- [52] Y. Sakatani, *U-duality extension of Drinfel'd double*, arXiv:1911.06320.
- [53] E. Malek and D. C. Thompson, *Poisson-Lie U-duality in Exceptional Field Theory*, arXiv:1911.07833.
- [54] I. Bakhmatov, K. Gubarev, and E. T. Musaev, *Non-abelian tri-vector deformations in $d = 11$ supergravity*, *JHEP* **05** (2020) 113, [arXiv:2002.01915].
- [55] C. D. A. Blair, D. C. Thompson, and S. Zhidkova, *Exploring Exceptional Drinfeld Geometries*, *JHEP* **09** (2020) 151, [arXiv:2006.12452].
- [56] E. T. Musaev, *On non-abelian U-duality of 11D backgrounds*, arXiv:2007.01213.
- [57] E. Malek, Y. Sakatani, and D. C. Thompson, *$E_{6(6)}$ exceptional Drinfel'd algebras*, *JHEP* **01** (2021) 020, [arXiv:2007.08510].
- [58] K. Gubarev and E. T. Musaev, *Polyvector deformations in eleven-dimensional supergravity*, arXiv:2011.11424.
- [59] E. T. Musaev and Y. Sakatani, *Non-abelian U-duality at work*, arXiv:2012.13263.
- [60] M. Hatsuda and T. Kimura, *Canonical approach to Courant brackets for D-branes*, *JHEP* **06** (2012) 034, [arXiv:1203.5499].

- [61] M. Hatsuda and K. Kamimura, *M5 algebra and SO(5,5) duality*, *JHEP* **06** (2013) 095, [arXiv:1305.2258].
- [62] D. S. Berman and N. B. Copland, *The String partition function in Hull's doubled formalism*, *Phys. Lett. B* **649** (2007) 325–333, [hep-th/0701080].
- [63] C. M. Hull, *A Geometry for non-geometric string backgrounds*, *JHEP* **10** (2005) 065, [hep-th/0406102].
- [64] C. M. Hull, *Global aspects of T-duality, gauged sigma models and T-folds*, *JHEP* **10** (2007) 057, [hep-th/0604178].
- [65] C. D. A. Blair, *Non-commutativity and non-associativity of the doubled string in non-geometric backgrounds*, *JHEP* **06** (2015) 091, [arXiv:1405.2283].
- [66] E. Bergshoeff, E. Sezgin, and P. Townsend, *Supermembranes and Eleven-Dimensional Supergravity*, *Phys. Lett. B* **189** (1987) 75–78.
- [67] M. Duff, P. S. Howe, T. Inami, and K. Stelle, *Superstrings in D=10 from Supermembranes in D=11*, *Phys. Lett. B* **191** (1987) 70.
- [68] E. Malek, *U-duality in three and four dimensions*, *Int. J. Mod. Phys. A* **32** (2017), no. 27 1750169, [arXiv:1205.6403].
- [69] C. D. A. Blair and E. Malek, *Geometry and fluxes of SL(5) exceptional field theory*, *JHEP* **03** (2015) 144, [arXiv:1412.0635].
- [70] Y. Sakatani and S. Uehara, *η -symbols in exceptional field theory*, *PTEP* **2017** (2017), no. 11 113B01, [arXiv:1708.06342].
- [71] I. Vaisman, *Towards a double field theory on para-Hermitian manifolds*, *J. Math. Phys.* **54** (2013) 123507, [arXiv:1209.0152].
- [72] L. Freidel, F. J. Rudolph, and D. Svoboda, *A Unique Connection for Born Geometry*, *Commun. Math. Phys.* **372** (2019), no. 1 119–150, [arXiv:1806.05992].
- [73] D. Svoboda, *Algebroid Structures on Para-Hermitian Manifolds*, *J. Math. Phys.* **59** (2018), no. 12 122302, [arXiv:1802.08180].
- [74] V. E. Marotta and R. J. Szabo, *Para-Hermitian Geometry, Dualities and Generalized Flux Backgrounds*, arXiv:1810.03953.
- [75] F. Hassler, D. Lüst, and F. J. Rudolph, *Para-Hermitian Geometries for Poisson-Lie Symmetric σ -models*, arXiv:1905.03791.
- [76] V. E. Marotta and R. J. Szabo, *Born Sigma-Models for Para-Hermitian Manifolds and Generalized T-Duality*, arXiv:1910.09997.
- [77] A. S. Arvanitakis, *Brane Wess-Zumino terms from AKSZ and exceptional generalised geometry as an L_∞ -algebroid*, *Adv. Theor. Math. Phys.* **23** (2019), no. 5 1159–1213, [arXiv:1804.07303].

- [78] L. Alfonsi and D. S. Berman, *Double Field Theory and Geometric Quantisation*,
arXiv:2101.12155.
- [79] C. Strickland-Constable, *Classical worldvolumes as generalised geodesics*,
arXiv:2102.00555.