

ON REALIZATIONS OF THE LIE GROUPS $G_{2,\mathbf{H}}, F_{4,\mathbf{H}}, E_{6,\mathbf{H}}, E_{7,\mathbf{H}}, E_{8,\mathbf{H}}$, SECOND EDITION

TOSHIKAZU MIYASHITA

ABSTRACT. In order to define the exceptional compact Lie groups G_2, F_4, E_6, E_7, E_8 , we usually use the Cayley algebra \mathfrak{C} or its complexification \mathfrak{C}^C . In the present article, we consider replacing the Cayley algebra \mathfrak{C} with the field of quaternion numbers \mathbf{H} in the definition of the groups above, and these groups are denoted as in title above. Our aim is to determine the structure of these groups. We call ‘‘realization’’ to determine the structure of the groups.

1. INTRODUCTION

The realizations of all exceptional simple Lie groups had been completed by Ichiro Yokota and the members of his school thirty-five years ago. Until now, their study has yielded the valuable and useful results, and those are summarized in [9] and [10].

As mentioned in abstract, in order to define the exceptional compact Lie groups G_2, F_4, E_6, E_7, E_8 , we usually use the Cayley algebra \mathfrak{C} or its complexification \mathfrak{C}^C . We consider replacing \mathfrak{C} with the field of quaternion numbers \mathbf{H} in the definition of the groups above. These groups are denoted by $G_{2,\mathbf{H}}, F_{4,\mathbf{H}}, E_{6,\mathbf{H}}, E_{7,\mathbf{H}}, E_{8,\mathbf{H}}$. In the present article, our aim is to determine the structure of those groups using the results obtained and the ideas of several proofs in the exceptional Lie groups.

Here, we will sketch out the contents of this article. First, we describe the definitions of the groups $G_{2,\mathbf{H}}, F_{4,\mathbf{H}}, E_{6,\mathbf{H}}, E_{7,\mathbf{H}}, E_{8,\mathbf{H}}$. For details of the definition, see the text of the article.

$$\begin{aligned} G_{2,\mathbf{H}} &= \{\alpha \in \text{Iso}_{\mathbf{R}}(\mathbf{H}) \mid (\alpha x)(\alpha y) = \alpha(xy)\}, \\ F_{4,\mathbf{H}} &= \{\alpha \in \text{Iso}_{\mathbf{R}}(\mathfrak{J}(3, \mathbf{H})) \mid \alpha X \circ \alpha Y = \alpha(X \circ Y)\}, \\ E_{6,\mathbf{H}} &= \{\alpha \in \text{Iso}_{\mathbf{C}}(\mathfrak{J}(3, \mathbf{H}^C)) \mid \det \alpha X = \det X, \langle \alpha X, \alpha Y \rangle = \langle X, Y \rangle\}, \\ E_{7,\mathbf{H}} &= \{\alpha \in \text{Iso}_{\mathbf{C}}((\mathfrak{B}_{\mathbf{H}})^C) \mid \alpha(P \times Q)\alpha^{-1} = \alpha P \times \alpha Q, \langle \alpha P, \alpha Q \rangle = \langle P, Q \rangle\}, \\ E_{8,\mathbf{H}} &= \{\alpha \in \text{Iso}_{\mathbf{C}}((\mathfrak{e}_{8,\mathbf{H}})^C) \mid \alpha[R_1, R_2] = [\alpha R_1, \alpha R_2], \langle \alpha R_1, \alpha R_2 \rangle = \langle R_1, R_2 \rangle\}. \end{aligned}$$

In Section 2, we describe the definitions of the Cayley algebras $\mathfrak{C}, \mathfrak{C}^C$, the exceptional Jordan algebras $\mathfrak{J}(3, \mathfrak{C}), \mathfrak{J}(3, \mathfrak{C}^C)$, the Freudenthal \mathbf{C} -vector space \mathfrak{B}^C , the 248-dimensional \mathbf{C} -vector space \mathfrak{e}_8^C , the exceptional Lie groups G_2, F_4, E_6, E_7, E_8 and these complexification, and in addition, we also describe the properties related to algebras, vector spaces or the results related to the exceptional Lie groups.

In Section 3, by using the isomorphism $(G_{2,\mathbf{H}})^C \cong Sp(1, \mathbf{H}^C)$ which had already been proved by Ichiro Yokota ([4, Proposition 1.3.1]), we will determine the structure of the group $G_{2,\mathbf{H}}$.

In Sections 4, 5, since the structure of the groups $F_{4,\mathbf{H}}$ and $E_{6,\mathbf{H}}$ had already been proved ([9, Propositions 2.11.3, 3.11.3]), these results will be described as theorems only. Moreover, in order to determine the structure of $E_{8,\mathbf{H}}$ in final section, we will determine the root system and the

2010 *Mathematics Subject Classification.* 53C30, 53C35, 17B40.

Key words and phrases. exceptional Lie groups.

Dynkin diagram of the Lie algebras $(\mathfrak{f}_4^C)^{\varepsilon_1, \varepsilon_2}$ and $(\mathfrak{e}_6^C)^{\varepsilon_1, \varepsilon_2}$ of the groups $(F_4^C)^{\varepsilon_1, \varepsilon_2}$ and $(E_6^C)^{\varepsilon_1, \varepsilon_2}$, respectively.

In Section 6, by using the isomorphism $(E_7^C)^{\varepsilon_1, \varepsilon_2} \cong Spin(12, C)$ ([2, Proposition 1.1.7]), we can show the isomorphism $(E_7)^{\varepsilon_1, \varepsilon_2} \cong Spin(12)$, so that we will determine the structure of the group $E_{7, \mathbf{H}}$. Moreover, as in the previous sections, we will determine the root system and the Dynkin diagram of the Lie algebras $(\mathfrak{e}_7^C)^{\varepsilon_1, \varepsilon_2}$ of the group $(E_7^C)^{\varepsilon_1, \varepsilon_2}$.

In the final section (Section 7), first the connectedness of the group $(E_{8, \mathbf{H}})^C$ is shown, so that we can prove that the connectedness of the group $E_{8, \mathbf{H}}$ follows from the connectedness of the group $(E_{8, \mathbf{H}})^C$. Subsequently, the connectedness of the group $(E_8^C)^{\varepsilon_1, \varepsilon_2}$ is shown, moreover we can draw the Dynkin diagram of the Lie algebra $(\mathfrak{e}_8^C)^{\varepsilon_1, \varepsilon_2}$ by using the results of root systems obtained in the previous sections, so that the type of the group $(E_8^C)^{\varepsilon_1, \varepsilon_2}$ as Lie algebras is determined. Hence, together with the result of the center $z((E_8^C)^{\varepsilon_1, \varepsilon_2})$, we will determine the structure of the group $(E_8^C)^{\varepsilon_1, \varepsilon_2}$. Finally, we will determine the structure of the compact real form $(E_8)^{\varepsilon_1, \varepsilon_2}$ of $(E_8^C)^{\varepsilon_1, \varepsilon_2}$, and we will determine the structure of the group $E_{8, \mathbf{H}}$ by defining a mapping $f_8 : (E_8)^{\varepsilon_1, \varepsilon_2} \rightarrow E_{8, \mathbf{H}}$.

Our results are given as follows.

$$\begin{aligned} G_{2, \mathbf{H}} &\cong Sp(1)/\mathbf{Z}_2, \\ F_{4, \mathbf{H}} &\cong Sp(3)/\mathbf{Z}_2, \\ E_{6, \mathbf{H}} &\cong SU(6)/\mathbf{Z}_2, \\ E_{7, \mathbf{H}} &\cong Ss(12) := Spin(12)/\mathbf{Z}_2, \\ E_{8, \mathbf{H}} &\cong E_7/\mathbf{Z}_2. \end{aligned}$$

These results are in [3] except $G_{2, \mathbf{H}} \cong Sp(1)/\mathbf{Z}_2$. However, there exists no proof in there. Thus, since Ichiro Yokota had determined the structure of the groups $F_{4, \mathbf{H}}$ and $E_{6, \mathbf{H}}$ ([9]), the author have added the proofs for the remaining groups $G_{2, \mathbf{H}}$, $E_{7, \mathbf{H}}$ and $E_{8, \mathbf{H}}$.

We refer the reader to [2] and [9] for notations. The author would like to say that one of the features of this article is to prove the connectedness of various groups by elementary methods.

2. PRELIMINARIES

In this section, we will focus on the definitions of the exceptional complex and compact Lie groups of type G_2, F_4, E_6, E_7 and E_8 , the structure of those Lie algebras and several results related to them. Since the proofs are omitted, if necessary, refer to [4], [5], [6], [8] for the complex Lie groups and [9] for the compact Lie groups.

2.1. The Lie groups G_2^C and G_2 of the type G_2 . Let \mathfrak{C} and \mathfrak{C}^C be the division Cayley algebra and its complexification, where $\mathfrak{C} = \{e_0 = 1, e_1, e_2, e_3, e_4, e_5, e_6, e_7\}_{\mathbf{R}}$. In \mathfrak{C} (resp. \mathfrak{C}^C), since the multiplication and the inner product are well known, these are omitted.

The simply connected complex Lie group G_2^C and the compact Lie group G_2 are defined by

$$\begin{aligned} G_2^C &= \{\alpha \in \text{Iso}_C(\mathfrak{C}^C) \mid \alpha(xy) = (\alpha x)(\alpha y)\}, \\ G_2 &= \{\alpha \in \text{Iso}_{\mathbf{R}}(\mathfrak{C}) \mid \alpha(xy) = (\alpha x)(\alpha y)\}. \end{aligned}$$

We define an \mathbf{R} -linear transformation γ of \mathfrak{C} by

$$\gamma(m + ne_4) = m - ne_4, \quad m + ne_4 \in \mathbf{H} \oplus \mathbf{H}e_4 = \mathfrak{C}.$$

Then we have $\gamma \in G_2$ and $\gamma^2 = 1$. Hence γ induces the involutive inner automorphism $\tilde{\gamma}$ on G_2 : $\tilde{\gamma}(\alpha) = \gamma\alpha\gamma, \alpha \in G_2$. Besides, γ is naturally extended to the C -linear transformation of \mathfrak{C}^C .

We need the following well-known result later.

Theorem 2.1 ([9, Theorem 1.10.1]). *The group $(G_2)^\gamma$ is isomorphic to the group $(Sp(1) \times Sp(1))/\mathbf{Z}_2, \mathbf{Z}_2 = \{(1, 1), (-1, -1)\}$: $(G_2)^\gamma \cong (Sp(1) \times Sp(1))/\mathbf{Z}_2$.*

2.2. The Lie groups $F_4^{\mathbb{C}}$ and F_4 of the type F_4 . Let $\mathfrak{J}(3, \mathbb{C})$ and $\mathfrak{J}(3, \mathbb{C}^{\mathbb{C}})$ be the exceptional Jordan algebra and its complexification. In $\mathfrak{J}(3, \mathbb{C})$ (resp. $\mathfrak{J}(3, \mathbb{C}^{\mathbb{C}})$), the Jordan multiplication $X \circ Y$, the inner product $\langle X, Y \rangle$ and a cross multiplication $X \times Y$, called the Freudenthal multiplication, are defined by

$$X \circ Y = \frac{1}{2}(XY + YX), \quad \langle X, Y \rangle = \text{tr}(X \circ Y),$$

$$X \times Y = \frac{1}{2}(2X \circ Y - \text{tr}(X)Y - \text{tr}(Y)X + (\text{tr}(X)\text{tr}(Y) - \langle X, Y \rangle)E),$$

respectively, where E is the 3×3 unit matrix. Moreover, we define the trilinear form $\langle X, Y, Z \rangle$, the determinant $\det X$ by

$$\langle X, Y, Z \rangle = \langle X, Y \times Z \rangle, \quad \det X = \frac{1}{3}\langle X, X, X \rangle,$$

respectively, and briefly denote $\mathfrak{J}(3, \mathbb{C})$ (resp. $\mathfrak{J}(3, \mathbb{C}^{\mathbb{C}})$) by $\mathfrak{J}^{\mathbb{C}}$ (resp. \mathfrak{J}).

In \mathfrak{J} (resp. $\mathfrak{J}^{\mathbb{C}}$), we often use the following nations:

$$E_1 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad E_2 = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad E_3 = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix},$$

$$F_1(x) = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & x \\ 0 & \bar{x} & 0 \end{pmatrix}, \quad F_2(x) = \begin{pmatrix} 0 & 0 & \bar{x} \\ 0 & 0 & 0 \\ x & 0 & 0 \end{pmatrix}, \quad F_3(x) = \begin{pmatrix} 0 & x & 0 \\ \bar{x} & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}.$$

The simply connected complex Lie group $F_4^{\mathbb{C}}$ and the compact Lie group F_4 are defined by

$$F_4^{\mathbb{C}} = \{\alpha \in \text{Iso}_{\mathbb{C}}(\mathfrak{J}^{\mathbb{C}}) \mid \alpha(X \circ Y) = \alpha X \circ \alpha Y\},$$

$$F_4 = \{\alpha \in \text{Iso}_{\mathbb{R}}(\mathfrak{J}) \mid \alpha(X \circ Y) = \alpha X \circ \alpha Y\}.$$

The Lie algebra $\mathfrak{f}_4^{\mathbb{C}}$ of the group $F_4^{\mathbb{C}}$ is given by

$$\mathfrak{f}_4^{\mathbb{C}} = \{\delta \in \text{Hom}_{\mathbb{C}}(\mathfrak{J}^{\mathbb{C}}) \mid \delta(X \circ Y) = \delta X \circ Y + X \circ \delta Y\},$$

and any element δ of the Lie algebras $\mathfrak{f}_4^{\mathbb{C}}$ can be uniquely expressed by

$$\delta = (D_1, D_2, D_3) + \tilde{A}_1(a_1) + \tilde{A}_2(a_2) + \tilde{A}_3(a_3), \quad D_i \in \mathfrak{so}(8, \mathbb{C}), \quad a_k \in \mathbb{C}^{\mathbb{C}},$$

where $D_2, D_3 \in \mathfrak{so}(8, \mathbb{C})$ are uniquely determined by the Principle of triality on $\mathfrak{so}(8, \mathbb{C})$: $(D_1 x)y + x(D_2 y) = \overline{D_3(\bar{x}y)}$, $x, y \in \mathbb{C}^{\mathbb{C}}$ for a given $D_1 \in \mathfrak{so}(8, \mathbb{C})$ and $\tilde{A}_k(a_k)$ is the \mathbb{C} -linear mapping of $\mathfrak{J}^{\mathbb{C}}$.

2.3. The Lie groups $E_6^{\mathbb{C}}$ and E_6 of the type E_6 . The simply connected complex Lie group $E_6^{\mathbb{C}}$ and the compact Lie group E_6 are defined by

$$E_6^{\mathbb{C}} = \{\alpha \in \text{Iso}_{\mathbb{C}}(\mathfrak{J}^{\mathbb{C}}) \mid \det \alpha X = \det X\},$$

$$E_6 = \{\alpha \in \text{Iso}_{\mathbb{C}}(\mathfrak{J}^{\mathbb{C}}) \mid \det \alpha X = \det X, \langle \alpha X, \alpha Y \rangle = \langle X, Y \rangle\},$$

where the Hermite inner product $\langle X, Y \rangle$ is defined by $(\tau X, Y)$ using τ is the complex conjugation in $\mathfrak{J}^{\mathbb{C}}$: $\langle X, Y \rangle = (\tau X, Y)$.

Then we have naturally the inclusion $F_4^{\mathbb{C}} \hookrightarrow E_6^{\mathbb{C}}$ and $(E_6^{\mathbb{C}})_E = F_4^{\mathbb{C}}$ (resp. $F_4 \hookrightarrow E_6$ and $(E_6)_E = F_4$).

The Lie algebra $\mathfrak{e}_6^{\mathbb{C}}$ of the group $E_6^{\mathbb{C}}$ is given by

$$\mathfrak{e}_6^{\mathbb{C}} = \{\phi \in \text{Hom}_{\mathbb{C}}(\mathfrak{J}^{\mathbb{C}}) \mid (\phi X, X, X) = 0\},$$

and any element ϕ of Lie algebras $\mathfrak{e}_6^{\mathbb{C}}$ can be uniquely expressed by

$$\phi = \delta + \tilde{T}, \quad \delta \in \mathfrak{f}_4^{\mathbb{C}}, \quad T \in (\mathfrak{J}^{\mathbb{C}})_0,$$

where $(\mathfrak{J}^C)_0 = \{X \in \mathfrak{J}^C \mid \text{tr}(X) = 0\}$ and the C -linear mapping \tilde{T} of \mathfrak{J}^C is defined by $\tilde{T}X = T \circ X, X \in \mathfrak{J}^C$.

2.4. The Lie groups E_7^C and E_7 of the type E_7 . Let \mathfrak{P}^C be the 56-dimensional Freudenthal C -vector space

$$\mathfrak{P}^C = \mathfrak{J}^C \oplus \mathfrak{J}^C \oplus C \oplus C,$$

with the inner product (P, Q) , the Hermite inner product $\langle P, Q \rangle$ and the skew-symmetric inner product $\{P, Q\}$ which are defined by

$$\begin{aligned} (P, Q) &:= (X, Z) + (Y, W) + \xi\zeta + \eta\omega \\ \langle P, Q \rangle &:= \langle X, Z \rangle + \langle Y, W \rangle + (\tau\xi) + (\tau\eta)\omega, \\ \{P, Q\} &:= (X, W) - (Y, Z) + \xi\omega - \eta\zeta, \end{aligned}$$

respectively, where $P = (X, Y, \xi, \eta), Q = (Z, W, \zeta, \omega) \in \mathfrak{P}^C$. Note that $\langle P, Q \rangle = (\tau P, Q) = \{\tau\lambda P, Q\}$.

In \mathfrak{P}^C , we often use the following notations:

$$\dot{X} := (X, 0, 0, 0), \dot{Y} := (0, Y, 0, 0), \dot{1} := (0, 0, 1, 0), \underline{1} := (0, 0, 0, 1).$$

For $\phi \in \mathfrak{e}_6^C, A, B \in \mathfrak{J}^C$ and $\nu \in C$, we define a C -linear mapping $\Phi(\phi, A, B, \nu) : \mathfrak{P}^C \rightarrow \mathfrak{P}^C$ by

$$\Phi(\phi, A, B, \nu) \begin{pmatrix} X \\ Y \\ \xi \\ \eta \end{pmatrix} = \begin{pmatrix} \phi X - \frac{1}{3}\nu X + 2B \times Y + \eta A \\ 2A \times X - {}^t\phi Y + \frac{1}{3}\nu Y + \xi B \\ (A, Y) + \nu\xi \\ (B, X) - \nu\eta \end{pmatrix},$$

where ${}^t\phi \in \mathfrak{e}_6^C$ is the transpose of ϕ with respect to the inner product (X, Y) : $({}^t\phi X, Y) = (X, \phi Y)$.

For $P = (X, Y, \xi, \eta), Q = (Z, W, \zeta, \omega) \in \mathfrak{P}^C$, we define a C -linear mapping $P \times Q : \mathfrak{P}^C \rightarrow \mathfrak{P}^C$, called the Freudenthal cross operation, by

$$P \times Q := \Phi(\phi, A, B, \nu), \quad \begin{cases} \phi = -\frac{1}{2}(X \vee W + Z \vee Y) \\ A = -\frac{1}{4}(2Y \times W - \xi Z - \zeta X) \\ B = \frac{1}{4}(2X \times Z - \eta W - \omega Y) \\ \nu = \frac{1}{8}((X, W) + (Z, Y) - 3(\xi\omega + \zeta\eta)), \end{cases}$$

where $X \vee W \in \mathfrak{e}_6^C$ is defined by

$$X \vee W = [\tilde{X}, \tilde{W}] + (X \circ W - \frac{1}{3}(X, W)E)^\sim,$$

here the C -linear mappings \tilde{X}, \tilde{W} of \mathfrak{J}^C are the same ones as that in E_6^C .

The simply connected complex Lie group E_7^C and the compact Lie E_7 are defined by

$$E_7^C = \{\alpha \in \text{Iso}_C(\mathfrak{P}^C) \mid \alpha(P \times Q)\alpha^{-1} = \alpha P \times \alpha Q\},$$

$$E_7 = \{\alpha \in \text{Iso}_C(\mathfrak{P}^C) \mid \alpha(P \times Q)\alpha^{-1} = \alpha P \times \alpha Q, \langle \alpha P, \alpha Q \rangle = \langle P, Q \rangle\}.$$

The Lie algebra \mathfrak{e}_7^C of the group E_7^C and the Lie algebra \mathfrak{e}_7 of the group E_7 are given by

$$\mathfrak{e}_7^C = \{\Phi(\phi, A, B, \nu) \in \text{Hom}(\mathfrak{P}^C) \mid \phi \in \mathfrak{e}_6^C, A, B \in \mathfrak{J}^C, \nu \in C\},$$

$$\mathfrak{e}_7 = \{\Phi(\phi, A, -\tau A, \nu) \in \text{Hom}(\mathfrak{P}^C) \mid \phi \in \mathfrak{e}_6, A \in \mathfrak{J}^C, \nu \in i\mathbf{R}\}.$$

For $\alpha \in E_6^C$, the mapping $\tilde{\alpha} : \mathfrak{P}^C \rightarrow \mathfrak{P}^C$ is defined by

$$\tilde{\alpha}(X, Y, \xi, \eta) = (\alpha X, {}^t\alpha^{-1}Y, \xi, \eta),$$

then we have $\tilde{\alpha} \in E_7^C$, and so α and $\tilde{\alpha}$ will be identified. The group E_7^C contains E_6^C as a subgroup by

$$E_6^C = (E_7^C)_{\mathbf{i}, \mathbf{1}} := \{\alpha \in E_7^C \mid \alpha \mathbf{i} = \mathbf{i}, \alpha \mathbf{1} = \mathbf{1}\}.$$

Hence we have the inclusion $F_4^C \hookrightarrow E_6^C \hookrightarrow E_7^C$. For $\alpha \in E_6$, suppose $\alpha \mathbf{i} = \mathbf{i}$, we have $\alpha \mathbf{1} = \mathbf{1}$, so E_7 contains E_6 as a subgroup by

$$E_6 = (E_7)_{\mathbf{i}}.$$

Hence we have the inclusion $F_4 \hookrightarrow E_6 \hookrightarrow E_7$. Finally, note that $\alpha \in E_7^C$ leaves the skew-symmetric inner product invariant: $\{\alpha P, \alpha Q\} = \{P, Q\}$, $P, Q \in \mathfrak{P}^C$.

2.5. The Lie group E_8^C and E_8 of the type E_8 . Let \mathfrak{e}_8^C be the 248-dimensional C -vector space

$$\mathfrak{e}_8^C = \mathfrak{e}_7^C \oplus \mathfrak{P}^C \oplus \mathfrak{P}^C \oplus C \oplus C \oplus C.$$

We define a Lie bracket $[R_1, R_2]$, $R_1 = (\Phi_1, P_1, Q_1, r_1, s_1, t_1)$, $R_2 = (\Phi_2, P_2, Q_2, r_2, s_2, t_2)$, by

$$[R_1, R_2] := (\Phi, P, Q, r, s, t), \quad \begin{cases} \Phi = [\Phi_1, \Phi_2] + P_1 \times Q_2 - P_2 \times Q_1 \\ P = \Phi_1 P_2 - \Phi_2 P_1 + r_1 P_2 - r_2 P_1 + s_1 Q_2 - s_2 Q_1 \\ Q = \Phi_1 Q_2 - \Phi_2 Q_1 - r_1 Q_2 + r_2 Q_1 + t_1 P_2 - t_2 P_1 \\ r = -\frac{1}{8}\{P_1, Q_2\} + \frac{1}{8}\{P_2, Q_1\} + s_1 t_2 - s_2 t_1 \\ s = \frac{1}{4}\{P_1, P_2\} + 2r_1 s_2 - 2r_2 s_1 \\ t = -\frac{1}{4}\{Q_1, Q_2\} - 2r_1 t_2 + 2r_2 t_1. \end{cases}$$

Then the C -vector space \mathfrak{e}_8^C becomes a complex simple Lie algebra of type E_8 .

In \mathfrak{e}_8^C , we often use the following notations:

$$\begin{aligned} \Phi &= (\Phi, 0, 0, 0, 0, 0), & P^- &= (0, P, 0, 0, 0, 0), & Q_- &= (0, 0, Q, 0, 0, 0), \\ \tilde{r} &= (0, 0, 0, r, 0, 0), & s^- &= (0, 0, 0, 0, s, 0), & t_- &= (0, 0, 0, 0, 0, t). \end{aligned}$$

We define a C -linear transformation λ_ω of \mathfrak{e}_8^C by

$$\lambda_\omega(\Phi, P, Q, r, s, t) = (\lambda\Phi\lambda^{-1}, \lambda Q, -\lambda P, -r, -t, -s),$$

where a C -linear transformation λ of \mathfrak{P}^C on the right-hand side is defined by $\lambda(X, Y, \xi, \eta) = (Y, -X, \eta, -\xi)$. As in \mathfrak{J}^C , the complex conjugation in \mathfrak{e}_8^C is denoted by τ :

$$\tau(\Phi, P, Q, r, s, t) = (\tau\Phi\tau, \tau P, \tau Q, \tau r, \tau s, \tau t).$$

The simply connected complex Lie group E_8^C and the compact Lie group E_8 are defined by

$$E_8^C = \text{Aut}(\mathfrak{e}_8^C) = \{\alpha \in \text{Iso}_C(\mathfrak{e}_8^C) \mid \alpha[R, R'] = [\alpha R, \alpha R']\},$$

$$E_8 = \{\alpha \in E_8^C \mid \langle \alpha R_1, \alpha R_2 \rangle = \langle R_1, R_2 \rangle\} = (E_8^C)^{\tau\lambda_\omega},$$

where the Hermite inner product $\langle R_1, R_2 \rangle$ is defined by $(-1/15)B_8(\tau\lambda_\omega R_1, R_2)$ using the Killing form B_8 of \mathfrak{e}_8^C : $\langle R_1, R_2 \rangle = (-1/15)B_8(\tau\lambda_\omega R_1, R_2)$.

The Lie algebra \mathfrak{e}_8^C of the group E_8^C and the Lie algebra \mathfrak{e}_8 of the group E_8 are given by

$$\mathfrak{e}_8^C = \{(\Phi, P, Q, r, s, t) \mid \Phi \in \mathfrak{e}_7^C, P, Q \in \mathfrak{J}^C, r, s, t \in C\},$$

$$\mathfrak{e}_8 = \{(\Phi, P, -\tau\lambda P, r, s, -\tau s) \mid \Phi \in \mathfrak{e}_7, P \in \mathfrak{J}^C, r \in i\mathbf{R}, s \in C\}.$$

For $\alpha \in E_7^C$, the mapping $\tilde{\alpha} : \mathfrak{e}_8^C \rightarrow \mathfrak{e}_8^C$ is defined by

$$\tilde{\alpha}(\Phi, P, Q, r, s, t) = (\alpha\Phi\alpha^{-1}, \alpha P, \alpha Q, r, s, t),$$

then we have $\tilde{\alpha} \in E_8^C$, and so α and $\tilde{\alpha}$ will be identified. The group E_8^C contains E_7^C as a subgroup by

$$E_7^C = (E_8^C)_{\tilde{1}, 1^-, 1_-} \left(= \{ \alpha \in (E_8^C) \mid \alpha\tilde{1} = \tilde{1}, \alpha 1^- = 1^-, \alpha 1_- = 1_- \} \right).$$

Hence we have the inclusion $F_4^C \hookrightarrow E_6^C \hookrightarrow E_7^C \hookrightarrow E_8^C$. For $\alpha \in E_8$, suppose $\alpha 1_- = 1_-$, we have $\alpha\tilde{1} = \tilde{1}$ and $\alpha 1^- = 1^-$, so E_8 contains E_7 as a subgroup by

$$E_7 = (E_8)_{1_-}.$$

Hence we have the inclusion $F_4 \hookrightarrow E_6 \hookrightarrow E_7 \hookrightarrow E_8$.

2.6. Useful lemma.

Lemma 2.2. *For Lie groups G, G' , let a mapping $\varphi : G \rightarrow G'$ be a homomorphism of Lie groups. When G' is connected, if $\text{Ker } \varphi$ is discrete and $\dim(\mathfrak{g}) = \dim(\mathfrak{g}')$, φ is surjective.*

Proof. The proof is omitted (cf. [4, Lemma 0.6 (2)]). \square

Lemma 2.3 (E. Cartan-Ravsevskii). *Let G be a simply connected Lie group with a finite order automorphism σ of G . Then G^σ is connected.*

Proof. The proof is omitted (cf. [4, Lemma 0.7]). \square

After this, using these lemmas without permission each times, we often prove lemma, proposition or theorem.

3. THE GROUP $G_{2, \mathbf{H}}$

Let \mathbf{H} and \mathbf{H}^C be the field of quaternion number and its complexification. We consider the following groups

$$\begin{aligned} (G_{2, \mathbf{H}})^C &:= \{ \alpha \in \text{Iso}_C(\mathbf{H}^C) \mid \alpha(xy) = (\alpha x)(\alpha y) \}, \\ G_{2, \mathbf{H}} &:= \{ \alpha \in \text{Iso}_R(\mathbf{H}) \mid \alpha(xy) = (\alpha x)(\alpha y) \}, \end{aligned}$$

In addition, we define a subgroup $((G_{2, \mathbf{H}})^C)^\tau$ of $(G_{2, \mathbf{H}})^C$ by

$$((G_{2, \mathbf{H}})^C)^\tau := \{ \alpha \in (G_{2, \mathbf{H}})^C \mid \tau\alpha = \alpha\tau \}.$$

Then we have the following proposition.

Lemma 3.1. *The group $((G_{2, \mathbf{H}})^C)^\tau$ coincides with the group $G_{2, \mathbf{H}}$: $((G_{2, \mathbf{H}})^C)^\tau = G_{2, \mathbf{H}}$.*

Proof. Let $\alpha \in G_{2, \mathbf{H}}$. Then we define an action of α to $x := x_1 + ix_2 \in \mathbf{H}^C, x_i \in \mathbf{H}$ by

$$\alpha x = \alpha(x_1 + ix_2) = \alpha x_1 + i\alpha x_2 \in \mathbf{H}^C.$$

Hence, for $x = x_1 + ix_2, y = y_1 + iy_2 \in \mathbf{H}^C, x_i, y_i \in \mathbf{H}$, it follows that

$$\begin{aligned} (\alpha x)(\alpha y) &= \alpha(x_1 + ix_2)\alpha(y_1 + iy_2) = (\alpha x_1 + i\alpha x_2)(\alpha y_1 + i\alpha y_2) \\ &= ((\alpha x_1)(\alpha y_1) - (\alpha x_2)(\alpha y_2)) + i((\alpha x_1)(\alpha y_2) + (\alpha x_2)(\alpha y_1)) \\ &= (\alpha(x_1 y_1) - \alpha(x_2 y_2)) + i(\alpha(x_1 y_2) + \alpha(x_2 y_1)) \\ &= \alpha(x_1 y_1 - x_2 y_2) + i\alpha(x_1 y_2 + x_2 y_1) \\ &= \alpha(x_1 y_1 - x_2 y_2 + i(x_1 y_2 + x_2 y_1)) \\ &= \alpha((x_1 + ix_2)(y_1 + iy_2)) \\ &= \alpha(xy). \end{aligned}$$

Hence we have $\alpha \in (G_{2,\mathbf{H}})^C$. Moreover, it is easy to verify that $\tau\alpha = \alpha\tau$. Indeed, it follows that

$$\begin{aligned}\tau\alpha x &= \tau\alpha(x_1 + ix_2) = \tau(\alpha x_1 + i\alpha x_2) = \alpha x_1 - i\alpha x_2 \\ &= \alpha(x_1 - ix_2) = \alpha\tau(x_1 + ix_2) = \alpha\tau x, \quad x = x_1 + ix_2 \in \mathbf{H}^C,\end{aligned}$$

that is, $\tau\alpha = \alpha\tau$. Thus we have $\alpha \in ((G_{2,\mathbf{H}})^C)^\tau$.

Conversely, let $\beta \in ((G_{2,\mathbf{H}})^C)^\tau$. From $\tau\beta = \beta\tau$, β induces an \mathbf{R} -isomorphism of \mathbf{H} . Indeed, note that $\tau y = y, y \in \mathbf{H}$, it follows that

$$\tau(\beta y) = \beta\tau y = \beta y,$$

so that $\beta y \in \mathbf{H}$. Thus we have $\beta \in G_{2,\mathbf{H}}$. With above, the proof of this lemma is completed. \square

Here, in order to determine the structure of the group $G_{2,\mathbf{H}}$, we need the following result.

Proposition 3.2. ([4, Proposition 1.3.1]) *The group $(G_{2,\mathbf{H}})^C$ is isomorphic to the group $Sp(1, \mathbf{H}^C)/\mathbf{Z}_2, \mathbf{Z}_2 = \{1, -1\}$: $(G_{2,\mathbf{H}})^C \cong Sp(1, \mathbf{H}^C)/\mathbf{Z}_2$.*

Now, we determine the structure of the group $G_{2,\mathbf{H}}$.

Theorem 3.3. *The group $G_{2,\mathbf{H}}$ is isomorphic to the group $Sp(1)/\mathbf{Z}_2, \mathbf{Z}_2 = \{1, -1\}$: $G_{2,\mathbf{H}} \cong Sp(1)/\mathbf{Z}_2$.*

Proof. Let $Sp(1) = \{q \in \mathbf{H} \mid q\bar{q} = 1\} \subset Sp(1, \mathbf{H}^C)$. We define a mapping $\varphi_{2,\mathbf{H}} : Sp(1) \rightarrow G_{2,\mathbf{H}}$ by

$$\varphi_{2,\mathbf{H}}(q)m = qm\bar{q}, \quad m \in \mathbf{H}.$$

First, we can easily confirm that $\varphi_{2,\mathbf{H}}$ is well-defined. Indeed, it follows that

$$(\varphi_{2,\mathbf{H}}(q)m)(\varphi_{2,\mathbf{H}}(q)n) = (qm\bar{q})(qn\bar{q}) = q(mn)\bar{q} = \varphi_{2,\mathbf{H}}(q)(mn).$$

Moreover, since this mapping is the restriction of the mapping $\varphi_{2,\mathbf{H}^C} : Sp(1, \mathbf{H}^C) \rightarrow (G_{2,\mathbf{H}})^C$, it is clear that $\varphi_{2,\mathbf{H}}$ is a homomorphism.

Next, we will prove that $\varphi_{2,\mathbf{H}}$ is surjective. Let $\alpha \in G_{2,\mathbf{H}} \subset (G_{2,\mathbf{H}})^C$. There exists $q \in Sp(1, \mathbf{H}^C)$ such that $\alpha = \varphi_{2,\mathbf{H}^C}(q)$ (Proposition 3.2). In addition, α satisfies the condition $\tau\alpha = \alpha\tau$ (Lemma 3.1), that is, $\tau\varphi_{2,\mathbf{H}^C}(q)\tau = \varphi_{2,\mathbf{H}^C}(q)$, so that we have $\varphi_{2,\mathbf{H}^C}(\tau q) = \varphi_{2,\mathbf{H}^C}(q)$. Hence we have the following

$$\tau q = q \quad \text{or} \quad \tau q = -q.$$

In the former case, we obtain $q \in Sp(1)$. In the latter case, we have $q = iq', q' \in \mathbf{H}$. However, this case is impossible because of $1 = q\bar{q} = (iq')(i\bar{q}') = -q'\bar{q}' \leq 0$. The proof of surjective is completed.

Finally, we will determine $\text{Ker } \varphi_{2,\mathbf{H}}$. Since $\varphi_{2,\mathbf{H}}$ is the restriction of the mapping φ_{2,\mathbf{H}^C} , it is clear $\text{Ker } \varphi_{2,\mathbf{H}} = \text{Ker } \varphi_{2,\mathbf{H}^C} = \{1, -1\} \cong \mathbf{Z}_2$.

Therefore we have the required isomorphism

$$G_{2,\mathbf{H}} \cong Sp(1)/\mathbf{Z}_2.$$

\square

After this, in all sections, we will determine the root system and the Dynkin diagram of some Lie algebras, where some Lie algebras are defined later. Its final aim is to determine the type of the complex Lie group $(E_{8,\mathbf{H}})^C$ as Lie algebras,

4. THE GROUP $F_{4,\mathbf{H}}$ AND THE ROOT SYSTEM, THE DYNKIN DIAGRAM OF THE LIE ALGEBRA $(\mathfrak{f}_4^C)^{\varepsilon_1, \varepsilon_2}$

Let $\mathfrak{J}(3, \mathfrak{C})$ be the exceptional Jordan algebra. We consider an algebra $\mathfrak{J}(3, \mathbf{H})$ which are defined by replacing \mathfrak{C} with \mathbf{H} in $\mathfrak{J}(3, \mathfrak{C})$, and we briefly denote $\mathfrak{J}(3, \mathbf{H})$ by $\mathfrak{J}_{\mathbf{H}}$.

Then we consider the following group $F_{4,\mathbf{H}}$ which is defined by replacing \mathfrak{J} with $\mathfrak{J}_{\mathbf{H}}$ in F_4 :

$$\begin{aligned} F_{4,\mathbf{H}} &= \{\alpha \in \text{Iso}_{\mathbf{R}}(\mathfrak{J}_{\mathbf{H}}) \mid \alpha(X \circ Y) = \alpha X \circ \alpha Y\} \\ &= \{\alpha \in \text{Iso}_{\mathbf{R}}(\mathfrak{J}_{\mathbf{H}}) \mid \alpha(X \times Y) = \alpha X \times \alpha Y\}. \end{aligned}$$

This group has been investigated by Ichiro Yokota, so that the following result was obtained.

Theorem 4.1. ([9, Proposition 2.11.3]) *The group $F_{4,\mathbf{H}}$ is isomorphic to the group $Sp(3)/\mathbf{Z}_2$, $\mathbf{Z}_2 = \{E, -E\}$: $F_{4,\mathbf{H}} \cong Sp(3)/\mathbf{Z}_2$.*

Now, we move the determination of the root system and the Dynkin diagram of the Lie algebra $(\mathfrak{f}_4^C)^{\varepsilon_1, \varepsilon_2}$.

We arrange \mathbf{R} -linear transformations of \mathfrak{C} below. By using the mapping $\varphi_{G_2} : Sp(1) \times Sp(1) \rightarrow (G_2)^\gamma$,

$$\varphi_{G_2}(p, q)(m + ne_4) = qm\bar{q} + (pn\bar{q})e_4, \quad m + ne_4 \in \mathbf{H} \oplus \mathbf{H}e_4 = \mathfrak{C}$$

(Theorem 2.1), we define \mathbf{R} -linear transformations $\varepsilon_1, \varepsilon_2$ of \mathfrak{C} by

$$\varepsilon_1 := \varphi_{G_2}(e_1, 1), \quad \varepsilon_2 := \varphi_{G_2}(e_2, 1),$$

where e_1, e_2 are two of basis in $\mathbf{H} \subset \mathfrak{C}$. Besides, the \mathbf{R} -linear transformation γ of \mathfrak{C} defined in previous section is expressed by

$$\gamma = \varphi_{G_2}(-1, 1).$$

These \mathbf{R} -linear transformations of \mathfrak{C} are naturally extended to \mathbf{R} -linear transformations of \mathfrak{J} as follows:

$$\varepsilon_i \begin{pmatrix} \xi_1 & x_3 & \bar{x}_2 \\ \bar{x}_3 & \xi_2 & x_1 \\ x_2 & \bar{x}_1 & \xi_3 \end{pmatrix} = \begin{pmatrix} \xi_1 & \varepsilon_i x_3 & \bar{\varepsilon}_i \bar{x}_2 \\ \bar{\varepsilon}_i \bar{x}_3 & \xi_2 & \varepsilon_i x_1 \\ \varepsilon_i x_2 & \bar{\varepsilon}_i \bar{x}_1 & \xi_3 \end{pmatrix}, \quad i = 1, 2, \quad \gamma \begin{pmatrix} \xi_1 & x_3 & \bar{x}_2 \\ \bar{x}_3 & \xi_2 & x_1 \\ x_2 & \bar{x}_1 & \xi_3 \end{pmatrix} = \begin{pmatrix} \xi_1 & \gamma x_3 & \bar{\gamma} \bar{x}_2 \\ \bar{\gamma} \bar{x}_3 & \xi_2 & \gamma x_1 \\ \gamma x_2 & \bar{\gamma} \bar{x}_1 & \xi_3 \end{pmatrix},$$

moreover these are also extended to \mathbf{C} -linear transformations of \mathfrak{J}^C . Then we have $\varepsilon_1, \varepsilon_2, \gamma \in G_2 \subset F_4 \subset F_4^C$ and $\varepsilon_1^2 = \varepsilon_2^2 = \gamma, \varepsilon_1^4 = \varepsilon_2^4 = 1, \gamma^2 = 1$.

Now, we consider a subgroup $(F_4^C)^{\varepsilon_1, \varepsilon_2}$ of F_4^C by

$$(F_4^C)^{\varepsilon_1, \varepsilon_2} := \{\alpha \in F_4^C \mid \varepsilon_1 \alpha = \alpha \varepsilon_1, \varepsilon_2 \alpha = \alpha \varepsilon_2\}.$$

Then the group $(F_4^C)^{\varepsilon_1, \varepsilon_2}$ is isomorphic to the group $Sp(3, \mathbf{H}^C)$: $(F_4^C)^{\varepsilon_1, \varepsilon_2} \cong Sp(3, \mathbf{H}^C)$, however since this group does not use directly in this article, the proof of isomorphism is omitted.

Now, we move the determination of the root system and the Dynkin diagram of the Lie algebra $(\mathfrak{f}_4^C)^{\varepsilon_1, \varepsilon_2}$ of the group $(F_4^C)^{\varepsilon_1, \varepsilon_2}$.

We prove the following lemma needed later.

Lemma 4.2. *The Lie algebra $(\mathfrak{f}_4^C)^{\varepsilon_1, \varepsilon_2}$ of the group $(F_4^C)^{\varepsilon_1, \varepsilon_2}$ is given by*

$$(\mathfrak{f}_4^C)^{\varepsilon_1, \varepsilon_2} = \left\{ \begin{array}{l} \delta = (D_1, D_2, D_3) \\ +\tilde{A}_1(a_1) + \tilde{A}_2(a_2) + \tilde{A}_3(a_3) \in \mathfrak{f}_4^C \end{array} \left| \begin{array}{l} D_1 = d_{01}G_{01} + d_{02}G_{02} + d_{03}G_{03} \\ +d_{12}G_{12} + d_{13}G_{13} + d_{23}G_{23} \\ +d_{45}(G_{45} + G_{67}) + d_{46}(G_{46} - G_{57}) \\ +d_{47}(G_{47} + G_{56}), d_{ij} \in \mathbf{C}, \\ D_2 = \pi\kappa D_1, D_3 = \kappa\pi D_1, \\ a_k \in \mathbf{H}^C \end{array} \right. \right\}.$$

In particular, we have $\dim_{\mathbf{C}}((\mathfrak{f}_4^C)^{\varepsilon_1, \varepsilon_2}) = 9 + 4 \times 3 = 21$.

Proof. First, $\varepsilon_1, \varepsilon_2$ are expressed as elements of the group $SO(8)$ as follows:

$$\varepsilon_1 = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & -1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & -1 & 0 \end{pmatrix}, \quad \varepsilon_2 = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & -1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & -1 & 0 & 0 \end{pmatrix}.$$

For $\delta \in \mathfrak{f}_4^C$, since it follows that

$$\varepsilon_i \delta \varepsilon_i^{-1} = (\varepsilon_i D_1 \varepsilon_i^{-1}, \varepsilon_i D_2 \varepsilon_i^{-1}, \varepsilon_i D_3 \varepsilon_i^{-1}) + \tilde{A}_1(\varepsilon_i a_1) + \tilde{A}_2(\varepsilon_i a_2) + \tilde{A}_3(\varepsilon_i a_3), i = 1, 2,$$

we obtain the required result using the matrix representations of $\varepsilon_1, \varepsilon_2$ above. \square

Here, we define a Lie subalgebra \mathfrak{h}_4 of $(\mathfrak{f}_4^C)^{\varepsilon_1, \varepsilon_2}$ by

$$\mathfrak{h}_4 := \left\{ \delta_4 = (L_1, L_2, L_3) \in (\mathfrak{f}_4^C)^{\varepsilon_1, \varepsilon_2} \mid \begin{array}{l} L_1 = \lambda_0(iG_{01}) + \lambda_1(iG_{23}) + \lambda_2(i(G_{45} + G_{67})) \\ L_2 = \pi\kappa L_1, L_3 = \kappa\pi L_1, \lambda_i \in \mathbb{C} \end{array} \right\},$$

where the explicit forms of L_2, L_3 are given as follow:

$$L_2 = \frac{1}{2}(-\lambda_0 + \lambda_1 + 2\lambda_2)(iG_{01}) + \frac{1}{2}(-\lambda_0 + \lambda_1 - 2\lambda_2)(iG_{23}) + \frac{1}{2}(-\lambda_0 - \lambda_1)(i(G_{45} + G_{67})),$$

$$L_3 = \frac{1}{2}(-\lambda_0 - \lambda_1 - 2\lambda_2)(iG_{01}) + \frac{1}{2}(\lambda_0 + \lambda_1 - 2\lambda_2)(iG_{23}) + \frac{1}{2}(\lambda_0 - \lambda_1)(i(G_{45} + G_{67})).$$

Then \mathfrak{h}_4 is a Cartan subalgebra of $(\mathfrak{f}_4^C)^{\varepsilon_1, \varepsilon_2}$. Indeed, it is clear that $(\mathfrak{f}_4^C)^{\varepsilon_1, \varepsilon_2}$ is abelian. Next, by doing straightforward computation, we can confirm that $[\delta, \delta_4] \in \mathfrak{h}_4, \delta \in (\mathfrak{f}_4^C)^{\varepsilon_1, \varepsilon_2}$ implies $\delta \in \mathfrak{h}_4$ for any $\delta_4 \in \mathfrak{h}_4$.

Theorem 4.3. *The rank of the Lie algebra $(\mathfrak{f}_4^C)^{\varepsilon_1, \varepsilon_2}$ is three. The roots Δ of $(\mathfrak{f}_4^C)^{\varepsilon_1, \varepsilon_2}$ relative to \mathfrak{h}_4 are given by*

$$\Delta = \left\{ \begin{array}{l} \pm(\lambda_0 - \lambda_1), \pm(\lambda_0 + \lambda_1), \pm 2\lambda_2, \pm\lambda_0, \pm\lambda_1, \\ \pm\frac{1}{2}(-\lambda_0 - \lambda_1 - 2\lambda_2), \pm\frac{1}{2}(\lambda_0 + \lambda_1 - 2\lambda_2) \end{array} \right\}.$$

Proof. Let $\delta_4 = (L_1, L_2, L_3) \in \mathfrak{h}_4$ and $\delta_{01} := (U_{01}, \pi\kappa U_{01}, \kappa\pi U_{01}) \in (\mathfrak{f}_4^C)^{\varepsilon_1, \varepsilon_2}$ where $U_{01} := (G_{02} + G_{13}) + i(G_{03} - G_{12})$. Then it follows that

$$\begin{aligned} [L_1, U_{01}] &= [\lambda_0(iG_{01}) + \lambda_1(iG_{23}) + \lambda_2(i(G_{45} + G_{67})), (G_{02} + G_{13}) + i(G_{03} - G_{12})] \\ &= (\lambda_0 - \lambda_1)G_{02} + (\lambda_0 - \lambda_1)G_{13} + (\lambda_0 - \lambda_1)iG_{03} - (\lambda_0 - \lambda_1)iG_{12} \\ &= (\lambda_0 - \lambda_1)((G_{02} + G_{13}) + i(G_{03} - G_{12})) \\ &= (\lambda_0 - \lambda_1)U_{01}, \end{aligned}$$

that is, $[L_1, U_{01}] = (\lambda_0 - \lambda_1)U_{01}$. Note that $[L_1, U_{01}] = (\lambda_0 - \lambda_1)U_{01}$ implies

$$\begin{aligned} [L_2, \pi\kappa U_{01}] &= [\pi\kappa L_1, \pi\kappa U_{01}] = (\lambda_0 - \lambda_1)\pi\kappa U_{01}, \\ [L_3, \kappa\pi U_{01}] &= [\kappa\pi L_1, \kappa\pi U_{01}] = (\lambda_0 - \lambda_1)\kappa\pi U_{01}, \end{aligned}$$

so that we have $[\delta_4, \delta_{01}] = (\lambda_0 - \lambda_1)\delta_{01}$. Hence we see that $\lambda_0 - \lambda_1$ is a root of $(\mathfrak{f}_4^C)^{\varepsilon_1, \varepsilon_2}$ and δ_{01} is a root vector associated with its root. Similarly, for the elements of $(\mathfrak{f}_4^C)^{\varepsilon_1, \varepsilon_2}$

$$\begin{aligned} \delta_{01}^- &:= (U_{01}^-, \pi\kappa U_{01}^-, \kappa\pi U_{01}^-), U_{01}^- := i(G_{02} + G_{13}) + (G_{03} - G_{12}), \\ \varepsilon_{01} &:= (V_{01}, \pi\kappa V_{01}, \kappa\pi V_{01}), V_{01} := (G_{02} - G_{13}) - i(G_{03} + G_{12}), \\ \varepsilon_{01}^- &:= (V_{01}^-, \pi\kappa V_{01}^-, \kappa\pi V_{01}^-), V_{01}^- := i(G_{02} - G_{13}) - (G_{03} + G_{12}), \end{aligned}$$

$$\begin{aligned}\varepsilon_{23} &:= (V_{23}, \pi\kappa V_{23}, \kappa\pi V_{23}), V_{23} := (G_{46} - G_{57}) - i(G_{47} + G_{56}), \\ \varepsilon_{23}^- &:= (V_{23}^-, \pi\kappa V_{23}^-, \kappa\pi V_{23}^-), V_{23}^- := i(G_{46} - G_{57}) - (G_{47} + G_{56}),\end{aligned}$$

we have

$$\begin{aligned}[\delta_4, \delta_{01}^-] &= -(\lambda_0 - \lambda_1)\delta_{01}^-, [\delta_4, \varepsilon_{01}] = (\lambda_0 + \lambda_1)\varepsilon_{01}, [\delta_4, \varepsilon_{01}^-] = -(\lambda_0 + \lambda_1)\varepsilon_{01}^-, \\ [\delta_4, \varepsilon_{23}] &= 2\lambda_2\varepsilon_{23}, [\delta_4, \varepsilon_{23}^-] = -2\lambda_2\varepsilon_{23}^-. \end{aligned}$$

The remainders of roots and root vectors associated with these roots are obtained as follows and together with the results above:

roots	root vectors associated with roots
$-(\lambda_0 - \lambda_1)$	$(U_{01}^-, \pi\kappa U_{01}^-, \kappa\pi U_{01}^-), U_{01}^- = i(G_{02} + G_{13}) + (G_{03} - G_{12})$
$\lambda_0 + \lambda_1$	$(V_{01}, \pi\kappa V_{01}, \kappa\pi V_{01}), V_{01} = (G_{02} - G_{13}) - i(G_{03} + G_{12})$
$-(\lambda_0 + \lambda_1)$	$(V_{01}^-, \pi\kappa V_{01}^-, \kappa\pi V_{01}^-), V_{01}^- = i(G_{02} - G_{13}) - (G_{03} + G_{12})$
$2\lambda_2$	$(V_{23}, \pi\kappa V_{23}, \kappa\pi V_{23}), V_{23} = (G_{46} - G_{57}) - i(G_{47} + G_{56})$
$-2\lambda_2$	$(V_{23}^-, \pi\kappa V_{23}^-, \kappa\pi V_{23}^-), V_{23}^- = i(G_{46} - G_{57}) - (G_{47} + G_{56})$
λ_0	$\tilde{A}_1(1 - ie_1)$
$-\lambda_0$	$\tilde{A}_1(1 + ie_1)$
λ_1	$\tilde{A}_1(e_2 - ie_3)$
$-\lambda_1$	$\tilde{A}_1(e_2 + ie_3)$
$(1/2)(-\lambda_0 + \lambda_1 + 2\lambda_2)$	$\tilde{A}_2(1 - ie_1)$
$-(1/2)(-\lambda_0 + \lambda_1 + 2\lambda_2)$	$\tilde{A}_2(1 + ie_1)$
$(1/2)(-\lambda_0 + \lambda_1 - 2\lambda_2)$	$\tilde{A}_2(e_2 - ie_3)$
$-(1/2)(-\lambda_0 + \lambda_1 - 2\lambda_2)$	$\tilde{A}_2(e_2 + ie_3)$
$(1/2)(-\lambda_0 - \lambda_1 - 2\lambda_2)$	$\tilde{A}_3(1 - ie_1)$
$-(1/2)(-\lambda_0 - \lambda_1 - 2\lambda_2)$	$\tilde{A}_3(1 + ie_1)$
$(1/2)(\lambda_0 + \lambda_1 - 2\lambda_2)$	$\tilde{A}_3(e_2 - ie_3)$
$-(1/2)(\lambda_0 + \lambda_1 - 2\lambda_2)$	$\tilde{A}_3(e_2 + ie_3)$


Thus The Lie algebra $(\mathfrak{f}_4^C)^{\varepsilon_1, \varepsilon_2}$ is spanned by \mathfrak{h}_4 and the roots vector associated with roots above, the roots obtained above are all. The rank of the Lie algebra $(\mathfrak{f}_4^C)^{\varepsilon_1, \varepsilon_2}$ follows from the dimension of \mathfrak{h}_4 . \square

Subsequently, we prove the following theorem.

Theorem 4.4. *In the root system Δ of Theorem 5.3,*

$$\Pi = \{\alpha_1, \alpha_2, \alpha_3\}$$

is a fundamental root system of $(\mathfrak{f}_4^C)^{\varepsilon_1, \varepsilon_2}$, where $\alpha_1 = (1/2)(-\lambda_0 + \lambda_1 + 2\lambda_2)$, $\alpha_2 = (1/2)(-\lambda_0 - \lambda_1 - 2\lambda_2)$, $\alpha_3 = \lambda_0 + \lambda_1$. The Dynkin diagram of $(\mathfrak{f}_4^C)^{\varepsilon_1, \varepsilon_2}$ is given by



Proof. The all positive roots are expressed by $\alpha_1, \alpha_2, \alpha_3$ as follows:

$$-(\lambda_0 - \lambda_1) = 2\alpha_1 + 2\alpha_2 + \alpha_3,$$

$$\begin{aligned}
\lambda_0 + \lambda_1 &= \alpha_3, \\
-2\lambda_2 &= 3\alpha_2 + \alpha_3, \\
-\lambda_0 &= \alpha_1 + \alpha_2, \\
\lambda_1 &= \alpha_1 + \alpha_2 + \alpha_3, \\
(1/2)(-\lambda_0 + \lambda_1 + 2\lambda_2) &= \alpha_1, \\
(1/2)(-\lambda_0 + \lambda_1 - 2\lambda_2) &= \alpha_1 + 2\alpha_2 + \alpha_3, \\
(1/2)(-\lambda_0 - \lambda_1 - 2\lambda_2) &= \alpha_2, \\
(1/2)(\lambda_0 + \lambda_1 - 2\lambda_2) &= \alpha_2 + \alpha_3.
\end{aligned}$$

Hence Π is a fundamental root system of $(\mathfrak{f}_4^C)^{\varepsilon_1, \varepsilon_2}$.

Then the Killing form B_4 of \mathfrak{f}_4^C is given by $B_4(\delta, \delta') = 3\text{tr}(\delta\delta')$ ([9, Theorem 2.5.3]), so that for $\delta_4 := (L_1, L_2, L_3), \delta_4' := (L_1', L_2', L_3') \in \mathfrak{h}_4$, we have

$$B_4(\delta_4, \delta_4') = 3\text{tr}(\delta_4\delta_4') = 18(\lambda_0\lambda_0' + \lambda_1\lambda_1' + 2\lambda_2\lambda_2').$$

Indeed, it follows that

$$\begin{aligned}
B_4(\delta_4, \delta_4') &= 3\text{tr}(\delta_4\delta_4') \\
&= 3\text{tr}((L_1, L_2, L_3)(L_1' L_2' L_3')) \\
&= 3\text{tr}(L_1 L_1', L_2 L_2', L_3 L_3') \\
&= 3(\text{tr}(L_1 L_1') + \text{tr}(L_2 L_2') + \text{tr}(L_3 L_3')) \\
&= 3(3(2\lambda_0\lambda_0' + 2\lambda_1\lambda_1' + 4\lambda_2\lambda_2')) \\
&= 18(\lambda_0\lambda_0' + \lambda_1\lambda_1' + 2\lambda_2\lambda_2').
\end{aligned}$$

Now, the canonical elements $\delta_{\alpha_1}, \delta_{\alpha_2}, \delta_{\alpha_3} \in \mathfrak{h}_4$ corresponding to $\alpha_1, \alpha_2, \alpha_3$ are determined as follows:

$$\begin{aligned}
\delta_{\alpha_1} &= -\frac{1}{36}(iG_{01}) + \frac{1}{36}(iG_{23}) + \frac{1}{36}(i(G_{45} + G_{67})), \\
\delta_{\alpha_2} &= -\frac{1}{36}(iG_{01}) - \frac{1}{36}(iG_{23}) - \frac{1}{36}(i(G_{45} + G_{67})), \\
\delta_{\alpha_3} &= \frac{1}{18}(iG_{01}) + \frac{1}{18}(iG_{23}).
\end{aligned}$$

Indeed, as for δ_{α_1} , set $\delta_{\alpha_1} := (L_1, L_2, L_3) \in \mathfrak{h}_4$, $L_1 := r_0(iG_{01}) + r_1(iG_{23}) + r_2(i(G_{45} + G_{67}))$, then since δ_{α_1} satisfies $B_4(\delta_{\alpha_1}, \delta_4) = \alpha_1(\delta_4)$ for any $\delta_4 \in \mathfrak{h}_4$, it follows from

$$B_4(\delta_{\alpha_1}, \delta_4) = 18(r_0\lambda_0 + r_1\lambda_1 + 2r_2\lambda_2), \quad \alpha_1(\delta_4) = (1/2)(-\lambda_0 + \lambda_1 + 2\lambda_2)$$

that

$$r_0 = -\frac{1}{36}, \quad r_1 = \frac{1}{36}, \quad r_2 = \frac{1}{36},$$

that is, $\delta_{\alpha_1} = -\frac{1}{36}(iG_{01}) + \frac{1}{36}(iG_{23}) + \frac{1}{36}(i(G_{45} + G_{67}))$. As for the others, we have the required results by doing similar computations.

Hence we have the following

$$\begin{aligned}
(\alpha_1, \alpha_1) &= B_4(\delta_{\alpha_1}, \delta_{\alpha_1}) = 18 \left(\left(-\frac{1}{36}\right)^2 + \left(\frac{1}{36}\right)^2 + \left(\frac{1}{36}\right)^2 \right) = \frac{1}{18}, \\
(\alpha_1, \alpha_2) &= B_4(\delta_{\alpha_1}, \delta_{\alpha_2}) = 18 \left(\left(-\frac{1}{36}\right)^2 + \left(\frac{1}{36}\right) \left(-\frac{1}{36}\right) + \left(\frac{1}{36}\right) \left(-\frac{1}{36}\right) \right) = -\frac{1}{12},
\end{aligned}$$

$$\begin{aligned}
(\alpha_1, \alpha_3) &= B_4(\delta_{\alpha_1}, \delta_{\alpha_3}) = 18 \left(\left(-\frac{1}{36} \right) \left(\frac{1}{18} \right) + \left(\frac{1}{36} \right) \left(\frac{1}{18} \right) \right) = 0, \\
(\alpha_2, \alpha_2) &= B_4(\delta_{\alpha_2}, \delta_{\alpha_2}) = 18 \left(\left(-\frac{1}{36} \right)^2 + \left(-\frac{1}{36} \right)^2 + \left(-\frac{1}{36} \right)^2 \right) = \frac{1}{18}, \\
(\alpha_2, \alpha_3) &= B_4(\delta_{\alpha_2}, \delta_{\alpha_3}) = 18 \left(\left(-\frac{1}{36} \right) \left(\frac{1}{18} \right) + \left(-\frac{1}{36} \right) \left(\frac{1}{18} \right) \right) = -\frac{1}{18}, \\
(\alpha_3, \alpha_3) &= B_4(\delta_{\alpha_3}, \delta_{\alpha_3}) = 18 \left(\left(\frac{1}{18} \right)^2 + \left(\frac{1}{18} \right)^2 \right) = \frac{1}{9}.
\end{aligned}$$

Thus, using the inner products above, we have

$$\begin{aligned}
\cos \theta_{12} &= \frac{(\alpha_1, \alpha_2)}{\sqrt{(\alpha_1, \alpha_1)(\alpha_2, \alpha_2)}} = -\frac{1}{2}, & \cos \theta_{13} &= \frac{(\alpha_1, \alpha_3)}{\sqrt{(\alpha_1, \alpha_1)(\alpha_3, \alpha_3)}} = 0, \\
\cos \theta_{23} &= \frac{(\alpha_2, \alpha_3)}{\sqrt{(\alpha_2, \alpha_2)(\alpha_3, \alpha_3)}} = -\frac{1}{\sqrt{2}}, & (\alpha_2, \alpha_2) &= \frac{1}{18} < \frac{1}{9} = (\alpha_3, \alpha_3),
\end{aligned}$$

so that we can draw the Dynkin diagram. \square

5. THE GROUP $E_{6,\mathbf{H}}$ AND THE ROOT SYSTEM, THE DYNKIN DIAGRAM OF THE LIE ALGEBRA $(\mathfrak{e}_6^C)^{\varepsilon_1, \varepsilon_2}$

Let $\mathfrak{J}(3, \mathfrak{C}^C)$ be the complex exceptional Jordan algebra. As in previous section, we consider an algebra $\mathfrak{J}(3, \mathbf{H}^C)$ which are defined by replacing \mathfrak{C} with \mathbf{H} in $\mathfrak{J}(3, \mathfrak{C}^C)$, and we briefly denote $\mathfrak{J}(3, \mathbf{H}^C)$ by $(\mathfrak{J}_{\mathbf{H}})^C$.

Then we consider the following group $E_{6,\mathbf{H}}$ which is defined by replacing \mathfrak{J}^C with $(\mathfrak{J}_{\mathbf{H}})^C$ in E_6 :

$$\begin{aligned}
E_{6,\mathbf{H}} &= \{ \alpha \in \text{Iso}_C((\mathfrak{J}_{\mathbf{H}})^C) \mid \det(\alpha X) = \det X, \langle \alpha X, \alpha Y \rangle = \langle X, Y \rangle \} \\
&= \{ \alpha \in \text{Iso}_C((\mathfrak{J}_{\mathbf{H}})^C) \mid \tau \alpha \tau(X \times Y) = \alpha X \times \alpha Y, \langle \alpha X, \alpha Y \rangle = \langle X, Y \rangle \}.
\end{aligned}$$

This group also has been investigated by Ichiro Yokota, so that the following result was obtained.

Theorem 5.1. ([9, Proposition 3.11.3]) *The group $E_{6,\mathbf{H}}$ is isomorphic to the group $SU(6)/\mathbf{Z}_2$, $\mathbf{Z}_2 = \{E, -E\}$: $E_{6,\mathbf{H}} \cong SU(6)/\mathbf{Z}_2$.*

We consider a subgroup $(E_6^C)^{\varepsilon_1, \varepsilon_2}$ of E_6^C by

$$(E_6^C)^{\varepsilon_1, \varepsilon_2} := \{ \alpha \in E_6^C \mid \varepsilon_1 \alpha = \alpha \varepsilon_1, \varepsilon_2 \alpha = \alpha \varepsilon_2 \}.$$

Then the group $(E_6^C)^{\varepsilon_1, \varepsilon_2}$ has been studied by author in [2, Proposition 1.1.1], and its result was obtained as follows:

$$(E_6^C)^{\varepsilon_1, \varepsilon_2} \cong SU^*(6, \mathbf{C}^C) (\cong SU(6, \mathbf{C}^C)).$$

Now, we move the determination of the root system and the Dynkin diagram of the Lie algebra $(\mathfrak{e}_6^C)^{\varepsilon_1, \varepsilon_2}$ of the group $(E_6^C)^{\varepsilon_1, \varepsilon_2}$. After this, we will determine those by the same procedure as in $(\mathfrak{f}_4^C)^{\varepsilon_1, \varepsilon_2}$.

We prove the following lemma needed later.

Lemma 5.2. *The Lie algebra $(\mathfrak{e}_6^C)^{\varepsilon_1, \varepsilon_2}$ of the group $(E_6^C)^{\varepsilon_1, \varepsilon_2}$ is given by*

$$(\mathfrak{e}_6^C)^{\varepsilon_1, \varepsilon_2} = \left\{ \begin{array}{l} \delta = (D_1, D_2, D_3) \\ + \tilde{A}_1(a_1) + \tilde{A}_2(a_2) + \tilde{A}_3(a_3) \\ + (\tau_1 E_1 + \tau_2 E_2 + \tau_3 E_3 + F_1(t_1) \\ + F_2(t_2) + F_3(t_3))^\sim \in \mathfrak{e}_6^C \end{array} \left| \begin{array}{l} D_1 = d_{01}G_{01} + d_{02}G_{02} + d_{03}G_{03} \\ + d_{12}G_{12} + d_{13}G_{13} + d_{23}G_{23} \\ + d_{45}(G_{45} + G_{67}) + d_{46}(G_{46} - G_{57}) \\ + d_{47}(G_{47} + G_{56}), d_{ij} \in C, \\ D_2 = \pi\kappa D_1, D_3 = \kappa\pi D_1, \\ a_k, t_k \in \mathbf{H}^C, \\ \tau_k \in C, \tau_1 + \tau_2 + \tau_3 = 0 \end{array} \right. \right\}.$$

In particular, we have $\dim_C((\mathfrak{e}_6^C)^{\varepsilon_1, \varepsilon_2}) = 9 + 4 \times 3 + (3 - 1 + 4 \times 3) = 35$.

Proof. For $\phi := \delta + \tilde{T} \in \mathfrak{e}_6^C, \delta \in \mathfrak{f}_4^C, T \in (\mathfrak{Z}^C)_0 := \{T \in \mathfrak{Z}^C \mid \text{tr}(T) = 0\}$, it follows that

$$\varepsilon_i \phi \varepsilon_i^{-1} = \varepsilon_i(\delta + \tilde{T})\varepsilon_i^{-1} = \varepsilon_i \delta \varepsilon_i^{-1} + (\varepsilon_i T)^\sim, \quad i = 1, 2$$

using Lemma 4.2, so that we have the required result. \square

Here, we define a Lie subalgebra \mathfrak{h}_6 of $(\mathfrak{e}_6^C)^{\varepsilon_1, \varepsilon_2}$ by

$$\mathfrak{h}_6 := \left\{ \begin{array}{l} \phi_6 = \delta_4 + (\mu_1 E_1 + \mu_2 E_2 + \mu_3 E_3)^\sim \in (\mathfrak{e}_6^C)^{\varepsilon_1, \varepsilon_2} \end{array} \left| \begin{array}{l} \delta_4 := (L_1, L_2, L_3), \\ L_1 = \lambda_0(iG_{01}) + \lambda_1(iG_{23}) \\ + \lambda_2(i(G_{45} + G_{67})), \\ L_2 = \pi\kappa L_1, L_3 = \kappa\pi L_1, \lambda_i \in C, \\ \mu_k \in C, \mu_1 + \mu_2 + \mu_3 = 0 \end{array} \right. \right\}.$$

Then \mathfrak{h}_6 is a Cartan subalgebra of $(\mathfrak{e}_6^C)^{\varepsilon_1, \varepsilon_2}$. Indeed, it is clear that $(\mathfrak{e}_6^C)^{\varepsilon_1, \varepsilon_2}$ is abelian. Next, by doing straightforward computation, we can confirm that $[\phi, \phi_6] \in \mathfrak{h}_6, \delta \in (\mathfrak{e}_6^C)^{\varepsilon_1, \varepsilon_2}$ implies $\phi \in \mathfrak{h}_6$ for any $\phi_6 \in \mathfrak{h}_6$.

Theorem 5.3. *The rank of the Lie algebra $(\mathfrak{e}_6^C)^{\varepsilon_1, \varepsilon_2}$ is five. The roots Δ of $(\mathfrak{e}_6^C)^{\varepsilon_1, \varepsilon_2}$ relative to \mathfrak{h}_6 are given by*

$$\Delta = \left\{ \begin{array}{l} \pm(\lambda_0 - \lambda_1), \pm(\lambda_0 + \lambda_1), \pm 2\lambda_2, \\ \pm \frac{1}{2}(-2\lambda_0 + \mu_2 - \mu_3), \pm \frac{1}{2}(-2\lambda_0 - \mu_2 + \mu_3), \\ \pm \frac{1}{2}(-2\lambda_1 + \mu_2 - \mu_3), \pm \frac{1}{2}(-2\lambda_1 - \mu_2 + \mu_3), \\ \pm \frac{1}{2}(\lambda_0 - \lambda_1 - 2\lambda_2 - \mu_1 + \mu_3), \pm \frac{1}{2}(\lambda_0 - \lambda_1 - 2\lambda_2 + \mu_1 - \mu_3), \\ \pm \frac{1}{2}(\lambda_0 - \lambda_1 + 2\lambda_2 - \mu_1 + \mu_3), \pm \frac{1}{2}(\lambda_0 - \lambda_1 + 2\lambda_2 + \mu_1 - \mu_3), \\ \pm \frac{1}{2}(\lambda_0 + \lambda_1 + 2\lambda_2 + \mu_1 - \mu_2), \pm \frac{1}{2}(\lambda_0 + \lambda_1 + 2\lambda_2 - \mu_1 + \mu_2), \\ \pm \frac{1}{2}(-\lambda_0 - \lambda_1 + 2\lambda_2 + \mu_1 - \mu_2), \pm \frac{1}{2}(-\lambda_0 - \lambda_1 + 2\lambda_2 - \mu_1 + \mu_2). \end{array} \right\}.$$

Proof. First, the roots $\pm(\lambda_0 - \lambda_1), \pm(\lambda_0 + \lambda_1), \pm 2\lambda_2$ of $(\mathfrak{f}_4^C)^{\varepsilon_1, \varepsilon_2}$ are also the roots of $(\mathfrak{e}_6^C)^{\varepsilon_1, \varepsilon_2}$. Indeed, for example, let $\lambda_0 - \lambda_1$ and its associated root vector δ_{01} . Then it follows that

$$\begin{aligned} [\phi_6, \delta_{01}] &= [\delta_4 + (\tau_1 E_1 + \tau_2 E_2 + \tau_3 E_3)^\sim, \delta_{01}] \\ &= [\delta_4, \delta_{01}] + [(\tau_1 E_1 + \tau_2 E_2 + \tau_3 E_3)^\sim, \delta_{01}] \quad ([(\tau_1 E_1 + \tau_2 E_2 + \tau_3 E_3)^\sim, \delta_{01}] = 0) \\ &= (\lambda_0 - \lambda_1)(\delta_{01}). \end{aligned}$$

The similar results are obtained for the others roots above.

We will determine the remainders of roots. Let $\phi_6 \in \mathfrak{h}_6$ and $\tilde{A}_1(1 + ie_1) + \tilde{F}_1(1 + ie_1) \in (\mathfrak{f}_4^C)^{\varepsilon_1, \varepsilon_2}$. Then it follows that

$$[\phi_6, \tilde{A}_1(1 + ie_1) + \tilde{F}_1(1 + ie_1)] = [(L_1, L_2, L_3) + (\mu_1 E_1 + \mu_2 E_2 + \mu_3 E_3)^\sim, \tilde{A}_1(1 + ie_1) + \tilde{F}_1(1 + ie_1)]$$

$$\begin{aligned}
&= \tilde{A}_1(L_1(1+ie_1)) + \tilde{F}_1(L_1(1+ie_1)) \\
&\quad + (1/2)\mu_2\tilde{F}_1(1+ie_1) + (1/2)\mu_2\tilde{A}_1(1+ie_1) \\
&\quad - (1/2)\mu_3\tilde{F}_1(1+ie_1) - (1/2)\mu_3\tilde{A}_1(1+ie_1) \\
&= -\lambda_0\tilde{A}_1(1+ie_1) - \lambda_0\tilde{F}_1(L_1(1+ie_1)) \\
&\quad + (1/2)\mu_2\tilde{F}_1(1+ie_1) + (1/2)\mu_2\tilde{A}_1(1+ie_1) \\
&\quad - (1/2)\mu_3\tilde{F}_1(1+ie_1) - (1/2)\mu_3\tilde{A}_1(1+ie_1) \\
&= (1/2)(-2\lambda_0 + \mu_2 - \mu_3)(\tilde{A}_1(1+ie_1) + \tilde{F}_1(1+ie_1)),
\end{aligned}$$

that is, $[\phi_6, \tilde{A}_1(1+ie_1) + \tilde{F}_1(1+ie_1)] = (1/2)(-2\lambda_0 + \mu_2 - \mu_3)(\tilde{A}_1(1+ie_1) + \tilde{F}_1(1+ie_1))$. Hence we see that $(1/2)(-2\lambda_0 + \mu_2 - \mu_3)$ is a root of $(\mathfrak{e}_6^C)^{\varepsilon_1, \varepsilon_2}$ and $\tilde{A}_1(1+ie_1) + \tilde{F}_1(1+ie_1)$ is a root vector associated with its root. Similarly, we have that $-(1/2)(-2\lambda_0 + \mu_2 - \mu_3)$ is a root of $(\mathfrak{e}_6^C)^{\varepsilon_1, \varepsilon_2}$ and $\tilde{A}_1(1-ie_1) - \tilde{F}_1(1-ie_1)$ is a root vector associated with its root.

The remainders of roots and root vectors associated with these roots are obtained as follows:

roots	root vectors associated with roots
$(1/2)(-2\lambda_0 - \mu_2 + \mu_3)$	$\tilde{A}_1(1+ie_1) - \tilde{F}_1(1+ie_1)$
$-(1/2)(-2\lambda_0 - \mu_2 + \mu_3)$	$\tilde{A}_1(1-ie_1) + \tilde{F}_1(1-ie_1)$
$(1/2)(-2\lambda_1 + \mu_2 - \mu_3)$	$\tilde{A}_1(e_2 + ie_3) + \tilde{F}_1(e_2 + ie_3)$
$-(1/2)(-2\lambda_1 + \mu_2 - \mu_3)$	$\tilde{A}_1(e_2 - ie_3) - \tilde{F}_1(e_2 - ie_3)$
$(1/2)(-2\lambda_1 - \mu_2 + \mu_3)$	$\tilde{A}_1(e_2 + ie_3) - \tilde{F}_1(e_2 + ie_3)$
$-(1/2)(-2\lambda_1 - \mu_2 + \mu_3)$	$\tilde{A}_1(e_2 - ie_3) + \tilde{F}_1(e_2 - ie_3)$
$(1/2)(\lambda_0 - \lambda_1 - 2\lambda_2 - \mu_1 + \mu_3)$	$\tilde{A}_2(1+ie_1) + \tilde{F}_2(1+ie_1)$
$-(1/2)(\lambda_0 - \lambda_1 - 2\lambda_2 - \mu_1 + \mu_3)$	$\tilde{A}_2(1-ie_1) - \tilde{F}_2(1-ie_1)$
$(1/2)(\lambda_0 - \lambda_1 - 2\lambda_2 + \mu_1 - \mu_3)$	$\tilde{A}_2(1+ie_1) - \tilde{F}_2(1+ie_1)$
$-(1/2)(\lambda_0 - \lambda_1 - 2\lambda_2 + \mu_1 - \mu_3)$	$\tilde{A}_2(1-ie_1) + \tilde{F}_2(1-ie_1)$
$(1/2)(\lambda_0 - \lambda_1 + 2\lambda_2 - \mu_1 + \mu_3)$	$\tilde{A}_2(e_2 + ie_3) + \tilde{F}_2(e_2 + ie_3)$
$-(1/2)(\lambda_0 - \lambda_1 + 2\lambda_2 - \mu_1 + \mu_3)$	$\tilde{A}_2(e_2 - ie_3) - \tilde{F}_2(e_2 - ie_3)$
$(1/2)(\lambda_0 - \lambda_1 + 2\lambda_2 + \mu_1 - \mu_3)$	$\tilde{A}_2(e_2 + ie_3) - \tilde{F}_2(e_2 + ie_3)$
$-(1/2)(\lambda_0 - \lambda_1 + 2\lambda_2 + \mu_1 - \mu_3)$	$\tilde{A}_2(e_2 - ie_3) + \tilde{F}_2(e_2 - ie_3)$
$(1/2)(\lambda_0 + \lambda_1 + 2\lambda_2 + \mu_1 - \mu_2)$	$\tilde{A}_3(1+ie_1) + \tilde{F}_3(1+ie_1)$
$-(1/2)(\lambda_0 + \lambda_1 + 2\lambda_2 + \mu_1 - \mu_2)$	$\tilde{A}_3(1-ie_1) - \tilde{F}_3(1-ie_1)$
$(1/2)(\lambda_0 + \lambda_1 + 2\lambda_2 - \mu_1 + \mu_2)$	$\tilde{A}_3(1+ie_1) - \tilde{F}_3(1+ie_1)$
$-(1/2)(\lambda_0 + \lambda_1 + 2\lambda_2 - \mu_1 + \mu_2)$	$\tilde{A}_3(1-ie_1) + \tilde{F}_3(1-ie_1)$
$(1/2)(-\lambda_0 - \lambda_1 + 2\lambda_2 + \mu_1 - \mu_2)$	$\tilde{A}_3(e_2 + ie_3) + \tilde{F}_3(e_2 + ie_3)$
$-(1/2)(-\lambda_0 - \lambda_1 + 2\lambda_2 + \mu_1 - \mu_2)$	$\tilde{A}_3(e_2 - ie_3) - \tilde{F}_3(e_2 - ie_3)$
$(1/2)(-\lambda_0 - \lambda_1 + 2\lambda_2 - \mu_1 + \mu_2)$	$\tilde{A}_3(e_2 + ie_3) - \tilde{F}_3(e_2 + ie_3)$
$-(1/2)(-\lambda_0 - \lambda_1 + 2\lambda_2 - \mu_1 + \mu_2)$	$\tilde{A}_3(e_2 - ie_3) + \tilde{F}_3(e_2 - ie_3)$


Thus, since $(\mathfrak{e}_6^C)^{\varepsilon_1, \varepsilon_2}$ is spanned by \mathfrak{h}_6 and the root vectors associated with roots above, the roots obtained above are all. The rank of the Lie algebra $(\mathfrak{e}_6^C)^{\varepsilon_1, \varepsilon_2}$ follows from the dimension of \mathfrak{e}_6 . \square

Subsequently, we prove the following theorem.

Theorem 5.4. *In the root system Δ of Theorem 5.3*

$$\Pi = \{\alpha_1, \alpha_2, \alpha_3, \alpha_4, \alpha_5\}$$

is a fundamental root system of $(\mathfrak{e}_6^C)^{\varepsilon_1, \varepsilon_2}$, where $\alpha_1 = -(\lambda_0 - \lambda_1)$, $\alpha_2 = (1/2)(\lambda_0 - \lambda_1 - 2\lambda_2 + \mu_1 - \mu_3)$, $\alpha_3 = 2\lambda_2$, $\alpha_4 = -(1/2)(\lambda_0 + \lambda_1 + 2\lambda_2 + \mu_1 - \mu_2)$, $\alpha_5 = \lambda_0 + \lambda_1$. The Dynkin diagram of $(\mathfrak{e}_6^C)^{\varepsilon_1, \varepsilon_2}$ is given by



Proof. The all positive roots are expressed by $\alpha_1, \alpha_2, \alpha_3, \alpha_4, \alpha_5$ as follows:

$$\begin{aligned} -(\lambda_0 - \lambda_1) &= \alpha_1, \\ \lambda_0 + \lambda_1 &= \alpha_5, \\ 2\lambda_2 &= \alpha_3, \\ (1/2)(-2\lambda_0 + \mu_2 - \mu_3) &= \alpha_1 + \alpha_2 + \alpha_3 + \alpha_4, \\ -(1/2)(-2\lambda_0 - \mu_2 + \mu_3) &= \alpha_2 + \alpha_3 + \alpha_4 + \alpha_5, \\ (1/2)(-2\lambda_1 + \mu_2 - \mu_3) &= \alpha_2 + \alpha_3 + \alpha_4, \\ -(1/2)(-2\lambda_1 - \mu_2 + \mu_3) &= \alpha_1 + \alpha_2 + \alpha_3 + \alpha_4 + \alpha_5, \\ -(1/2)(\lambda_0 - \lambda_1 - 2\lambda_2 - \mu_1 + \mu_3) &= \alpha_1 + \alpha_2 + \alpha_3, \\ (1/2)(\lambda_0 - \lambda_1 - 2\lambda_2 + \mu_1 - \mu_3) &= \alpha_2, \\ -(1/2)(\lambda_0 - \lambda_1 + 2\lambda_2 - \mu_1 + \mu_3) &= \alpha_1 + \alpha_2, \\ (1/2)(\lambda_0 - \lambda_1 + 2\lambda_2 + \mu_1 - \mu_3) &= \alpha_2 + \alpha_3, \\ -(1/2)(\lambda_0 + \lambda_1 + 2\lambda_2 + \mu_1 - \mu_2) &= \alpha_4, \\ (1/2)(\lambda_0 + \lambda_1 + 2\lambda_2 - \mu_1 + \mu_2) &= \alpha_3 + \alpha_4 + \alpha_5, \\ -(1/2)(-\lambda_0 - \lambda_1 + 2\lambda_2 + \mu_1 - \mu_2) &= \alpha_4 + \alpha_5, \\ (1/2)(-\lambda_0 - \lambda_1 + 2\lambda_2 - \mu_1 + \mu_2) &= \alpha_3 + \alpha_4. \end{aligned}$$

Hence Π is a fundamental root system of $(\mathfrak{e}_6^C)^{\varepsilon_1, \varepsilon_2}$.

Then, for $\phi, \phi' \in \mathfrak{e}_6^C$, the Killing form B_6 of \mathfrak{e}_6^C is given by

$$B_6(\phi, \phi') = B_6(\delta + \tilde{T}, \delta' + \tilde{T}') = \frac{4}{3}B_4(\delta, \delta') + 12(T, T')$$

([9, Theorem 3.5.3]), so that for $\phi_6 := \delta_4 + (\mu_1 E_1 + \mu_2 E_2 + \mu_3 E_3)^\sim$, $\phi_6' := \delta_4' + (\mu_1' E_1 + \mu_2' E_2 + \mu_3' E_3)^\sim \in \mathfrak{h}_6$, we have

$$\begin{aligned} B_6(\phi_4, \phi_4') &= \frac{4}{3}B_4(\delta_4, \delta_4') + 12(\mu_1 E_1 + \mu_2 E_2 + \mu_3 E_3, \mu_1' E_1 + \mu_2' E_2 + \mu_3' E_3) \\ &= \frac{4}{3} \cdot 18(\lambda_0 \lambda_0' + \lambda_1 \lambda_1' + 2\lambda_2 \lambda_2') + 12(\mu_1 \mu_1' + \mu_2 \mu_2' + \mu_3 \mu_3') \\ &= 24(\lambda_0 \lambda_0' + \lambda_1 \lambda_1' + 2\lambda_2 \lambda_2') + 12(\mu_1 \mu_1' + \mu_2 \mu_2' + \mu_3 \mu_3'). \end{aligned}$$

Now, the canonical elements $\phi_{\alpha_1}, \phi_{\alpha_2}, \phi_{\alpha_3}, \phi_{\alpha_4}, \phi_{\alpha_5}$ corresponding to $\alpha_1, \alpha_2, \alpha_3, \alpha_4, \alpha_5$ are determined as follows:

$$\begin{aligned} \phi_{\alpha_1} &= -\frac{1}{24}(iG_{01}) + \frac{1}{24}(iG_{23}), \\ \phi_{\alpha_2} &= \frac{1}{48}(iG_{01}) - \frac{1}{48}(iG_{23}) - \frac{1}{48}(i(G_{45} + G_{67})) + \left(\frac{1}{24}E_1 - \frac{1}{24}E_3\right)^\sim, \\ \phi_{\alpha_3} &= \frac{1}{24}(i(G_{45} + G_{67})), \\ \phi_{\alpha_4} &= -\frac{1}{48}(iG_{01}) - \frac{1}{48}(iG_{23}) - \frac{1}{48}(i(G_{45} + G_{67})) + \left(-\frac{1}{24}E_1 + \frac{1}{24}E_2\right)^\sim, \end{aligned}$$

$$\phi_{\alpha_5} = \frac{1}{24}(iG_{01}) + \frac{1}{24}(iG_{23}).$$

Indeed, as for ϕ_{α_1} , set $\phi_{\alpha_1} := (L_1, L_2, L_3) + (\nu_1 E_1 + \nu_2 E_2 + \nu_3 E_3) \sim \in \mathfrak{h}_6$, $L_1 := r_0(iG_{01}) + r_1(iG_{23}) + r_2(i(G_{45} + G_{67}))$, then since ϕ_{α_1} satisfies $B_6(\phi_{\alpha_1}, \phi_6) = \alpha_1(\phi_6)$ for any $\phi_6 \in \mathfrak{h}_6$, it follows from

$$\begin{aligned} B_6(\phi_{\alpha_1}, \phi_6) &= 24(r_0\lambda_0 + r_1\lambda_1 + 2r_2\lambda_2) + 12(\nu_1\mu_1 + \nu_2\mu_2 + \nu_3\mu_3), \\ \alpha_1(\phi_6) &= -\lambda_0 + \lambda_1 \end{aligned}$$

that

$$r_0 = -\frac{1}{24}, \quad r_1 = \frac{1}{24},$$

that is, $\phi_{\alpha_1} = -\frac{1}{24}(iG_{01}) + \frac{1}{24}(iG_{23})$. As for the others, we have the required results by doing similar computations.

Hence we have the following

$$\begin{aligned} (\alpha_1, \alpha_1) &= B_6(\phi_{\alpha_1}, \phi_{\alpha_1}) = 24 \left(\left(-\frac{1}{24} \right)^2 + \left(\frac{1}{24} \right)^2 \right) = \frac{1}{12}, \\ (\alpha_1, \alpha_2) &= B_6(\phi_{\alpha_1}, \phi_{\alpha_2}) = 24 \left(\left(-\frac{1}{24} \right) \left(\frac{1}{48} \right) + \left(\frac{1}{24} \right) \left(-\frac{1}{48} \right) \right) = -\frac{1}{24}, \\ (\alpha_1, \alpha_3) &= B_6(\phi_{\alpha_1}, \phi_{\alpha_3}) = 0, \\ (\alpha_1, \alpha_4) &= B_6(\phi_{\alpha_1}, \phi_{\alpha_4}) = 24 \left(\left(-\frac{1}{24} \right) \left(-\frac{1}{48} \right) + \left(\frac{1}{24} \right) \left(-\frac{1}{48} \right) \right) = 0, \\ (\alpha_1, \alpha_5) &= B_6(\phi_{\alpha_1}, \phi_{\alpha_5}) = 24 \left(\left(-\frac{1}{24} \right) \left(\frac{1}{24} \right) + \left(\frac{1}{24} \right)^2 \right) = 0, \\ (\alpha_2, \alpha_2) &= B_6(\phi_{\alpha_2}, \phi_{\alpha_2}) = 24 \left(\left(\frac{1}{48} \right)^2 + \left(-\frac{1}{48} \right)^2 + 2 \left(-\frac{1}{48} \right)^2 \right) + 12 \left(\left(\frac{1}{24} \right)^2 + \left(-\frac{1}{24} \right)^2 \right) = \frac{1}{12}, \\ (\alpha_2, \alpha_3) &= B_6(\phi_{\alpha_2}, \phi_{\alpha_3}) = 24 \cdot 2 \left(-\frac{1}{48} \right) \left(\frac{1}{24} \right) = -\frac{1}{24}, \\ (\alpha_2, \alpha_4) &= B_6(\phi_{\alpha_2}, \phi_{\alpha_4}) = 24 \left(\left(\frac{1}{48} \right) \left(-\frac{1}{48} \right) + \left(-\frac{1}{48} \right)^2 + 2 \left(-\frac{1}{48} \right)^2 \right) + 12 \left(\frac{1}{24} \right) \left(-\frac{1}{24} \right) = 0, \\ (\alpha_2, \alpha_5) &= B_6(\phi_{\alpha_2}, \phi_{\alpha_5}) = 24 \left(\left(\frac{1}{48} \right) \left(\frac{1}{24} \right) + \left(-\frac{1}{48} \right) \left(\frac{1}{24} \right) \right) = 0, \\ (\alpha_3, \alpha_3) &= B_6(\phi_{\alpha_3}, \phi_{\alpha_3}) = 24 \cdot 2 \left(\frac{1}{24} \right)^2 = \frac{1}{12}, \\ (\alpha_3, \alpha_4) &= B_6(\phi_{\alpha_3}, \phi_{\alpha_4}) = 24 \cdot 2 \left(\frac{1}{24} \right) \left(-\frac{1}{48} \right) = -\frac{1}{24}, \\ (\alpha_3, \alpha_5) &= B_6(\phi_{\alpha_3}, \phi_{\alpha_5}) = 0, \\ (\alpha_4, \alpha_4) &= B_6(\phi_{\alpha_4}, \phi_{\alpha_4}) = 24 \left(\left(-\frac{1}{48} \right)^2 + \left(-\frac{1}{48} \right)^2 + 2 \left(-\frac{1}{48} \right)^2 \right) + 12 \left(\left(-\frac{1}{24} \right)^2 + \left(\frac{1}{24} \right)^2 \right) = \frac{1}{12}, \\ (\alpha_4, \alpha_5) &= B_6(\phi_{\alpha_4}, \phi_{\alpha_5}) = 24 \left(\left(-\frac{1}{48} \right) \left(\frac{1}{24} \right) + \left(-\frac{1}{48} \right) \left(\frac{1}{24} \right) \right) = -\frac{1}{24}, \\ (\alpha_5, \alpha_5) &= B_6(\phi_{\alpha_5}, \phi_{\alpha_5}) = 24 \left(\left(\frac{1}{24} \right)^2 + \left(\frac{1}{24} \right)^2 \right) = \frac{1}{12}. \end{aligned}$$

Thus, using the inner products above, we have

$$\begin{aligned}\cos \theta_{12} &= \frac{(\alpha_1, \alpha_2)}{\sqrt{(\alpha_1, \alpha_1)(\alpha_2, \alpha_2)}} = -\frac{1}{2}, & \cos \theta_{13} &= \frac{(\alpha_1, \alpha_3)}{\sqrt{(\alpha_1, \alpha_1)(\alpha_3, \alpha_3)}} = 0, \\ \cos \theta_{14} &= \frac{(\alpha_1, \alpha_4)}{\sqrt{(\alpha_1, \alpha_1)(\alpha_4, \alpha_4)}} = 0, & \cos \theta_{15} &= \frac{(\alpha_1, \alpha_5)}{\sqrt{(\alpha_1, \alpha_1)(\alpha_5, \alpha_5)}} = 0, \\ \cos \theta_{23} &= \frac{(\alpha_2, \alpha_3)}{\sqrt{(\alpha_2, \alpha_2)(\alpha_3, \alpha_3)}} = -\frac{1}{2}, & \cos \theta_{24} &= \frac{(\alpha_2, \alpha_4)}{\sqrt{(\alpha_2, \alpha_2)(\alpha_4, \alpha_4)}} = 0, \\ \cos \theta_{25} &= \frac{(\alpha_2, \alpha_5)}{\sqrt{(\alpha_2, \alpha_2)(\alpha_5, \alpha_5)}} = 0, & \cos \theta_{34} &= \frac{(\alpha_3, \alpha_4)}{\sqrt{(\alpha_3, \alpha_3)(\alpha_4, \alpha_4)}} = -\frac{1}{2}, \\ \cos \theta_{35} &= \frac{(\alpha_3, \alpha_5)}{\sqrt{(\alpha_3, \alpha_3)(\alpha_5, \alpha_5)}} = 0, & \cos \theta_{45} &= \frac{(\alpha_4, \alpha_5)}{\sqrt{(\alpha_4, \alpha_4)(\alpha_5, \alpha_5)}} = -\frac{1}{2}.\end{aligned}$$

so that we can draw the Dynkin diagram. \square

6. THE GROUP $E_{7,\mathbf{H}}$ AND THE ROOT SYSTEM, THE DYNKIN DIAGRAM OF THE LIE ALGEBRA $(\mathfrak{e}_{7,\mathbf{H}})^C$

Let \mathfrak{B}^C be the Freudenthal C -vector space. We consider a 32 dimensional C -vector space $(\mathfrak{B}_{\mathbf{H}})^C$ which is defined by replacing \mathfrak{C} with \mathbf{H} in \mathfrak{B}^C :

$$(\mathfrak{B}_{\mathbf{H}})^C := (\mathfrak{J}_{\mathbf{H}})^C \oplus (\mathfrak{J}_{\mathbf{H}})^C \oplus C \oplus C.$$

Then we consider the following complex Lie algebra $(\mathfrak{e}_{7,\mathbf{H}})^C$ which is given by replacing \mathfrak{C} with \mathbf{H} in the complex Lie algebra $\mathfrak{e}_{7,\mathbf{H}}^C$:

$$(\mathfrak{e}_{7,\mathbf{H}})^C = \{ \Phi(\phi, A, B, \nu) \in \text{Hom}_C((\mathfrak{B}_{\mathbf{H}})^C) \mid \phi \in (\mathfrak{e}_{6,\mathbf{H}})^C, A, B \in (\mathfrak{J}_{\mathbf{H}})^C, \nu \in C \}.$$

In particular, we have $\dim_C((\mathfrak{e}_{7,\mathbf{H}})^C) = 35 + 15 \times 2 + 1 = 66$. In $(\mathfrak{e}_{7,\mathbf{H}})^C$, we can define a Lie bracket and can prove its simplicity as in $\mathfrak{e}_{7,\mathbf{H}}^C$. In addition, we define a symmetric inner product $(\Phi_1, \Phi_2)_7$ by

$$(\Phi_1, \Phi_2)_7 := -2(\phi_1, \phi_2)_6 - 4(A_1, B_2) - 4(A_2, B_1) - \frac{8}{3}\nu_1\nu_2,$$

where $\Phi_i := \Phi(\phi_i, A_i, B_i, \nu_i) \in (\mathfrak{e}_{7,\mathbf{H}})^C, i = 1, 2$. Then the symmetric inner product $(\Phi_1, \Phi_2)_7$ is $(\mathfrak{e}_{7,\mathbf{H}})^C$ -adjoint invariant (cf. [9, Lemma 4.5.1 (1)]).

Then we prove the following theorem needed in Section 8.

Theorem 6.1. *The killing form $B_{7,\mathbf{H}}$ of $(\mathfrak{e}_{7,\mathbf{H}})^C$ is given by*

$$B_{7,\mathbf{H}}(\Phi_1, \Phi_2) = -5(\Phi_1, \Phi_2)_7.$$

Proof. Since $(\mathfrak{e}_{7,\mathbf{H}})^C$ is simple, there exist $k \in C$ such that

$$B_{7,\mathbf{H}}(\Phi_1, \Phi_2) = k(\Phi_1, \Phi_2)_7, \quad \Phi_i \in (\mathfrak{e}_{7,\mathbf{H}})^C.$$

We will determine k . Let $\Phi_1 = \Phi_2 = \Phi(0, 0, 0, 1)$. Then we have

$$(\Phi_1, \Phi_2)_7 = -\frac{8}{3}.$$

On the other hand, it follows from

$$\begin{aligned}(\text{ad}\Phi_1)(\text{ad}\Phi_2)\Phi(\phi, A, B, \nu) &= [\Phi_1, [\Phi_2, \Phi(\phi, A, B, \nu)]] \\ &= [\Phi_1, \Phi(0, \frac{2}{3}A, -\frac{2}{3}B, 0)] \\ &= \Phi(0, \frac{4}{9}A, \frac{4}{9}B, 0)\end{aligned}$$

that

$$B_{7,\mathbf{H}}(\text{ad}\Phi_1\text{ad}\Phi_2) = \text{tr}(\text{ad}\Phi_1\text{ad}\Phi_2) = \frac{4}{9} \times 15 \times 2 = \frac{40}{3}.$$

Thus we have $k = -5$.

Therefore we have

$$B_{7,\mathbf{H}}(\Phi_1, \Phi_2) = -5(\Phi_1, \Phi_2)_7.$$

□

We will study a following Lie group $E_{7,\mathbf{H}}$ which is defined by replacing \mathfrak{C} with \mathbf{H} in the compact Lie group E_7 :

$$E_{7,\mathbf{H}} := \{ \alpha \in \text{Iso}_{\mathfrak{C}}((\mathfrak{P}_{\mathbf{H}})^{\mathfrak{C}}) \mid \alpha(P \times Q)\alpha^{-1} = \alpha P \times \alpha Q, \langle \alpha P, \alpha Q \rangle = \langle P, Q \rangle \},$$

where $\langle P, Q \rangle := (\tau P, Q) = \{\tau\lambda P, Q\}$. The group $E_{7,\mathbf{H}}$ is a compact Lie group as a closed subgroup of the unitary group $U(32) = U((\mathfrak{P}_{\mathbf{H}})^{\mathfrak{C}}) = \{ \alpha \in \text{Iso}_{\mathfrak{C}}((\mathfrak{P}_{\mathbf{H}})^{\mathfrak{C}}) \mid \langle \alpha P, \alpha Q \rangle = \langle P, Q \rangle \}$.

Lemma 6.2. *The Lie algebra $\mathfrak{e}_{7,\mathbf{H}}$ of the group $E_{7,\mathbf{H}}$ is given by*

$$\mathfrak{e}_{7,\mathbf{H}} = \{ \Phi(\phi, A, -\tau A, \nu) \mid \phi \in \mathfrak{e}_{6,\mathbf{H}}, A \in (\mathfrak{J}_{\mathbf{H}})^{\mathfrak{C}}, \nu \in i\mathbf{R} \}.$$

In particular, we have $\dim(\mathfrak{e}_{7,\mathbf{H}}) = 35 + 30 + 1 = 66$.

Proof. As in $\mathfrak{e}_7^{\mathfrak{C}}$, the complex Lie algebra $(\mathfrak{e}_{7,\mathbf{H}})^{\mathfrak{C}}$ of the group $(E_{7,\mathbf{H}})^{\mathfrak{C}}$ is given by

$$(\mathfrak{e}_{7,\mathbf{H}})^{\mathfrak{C}} = \{ \Phi(\phi, A, B, \nu) \mid \phi \in (\mathfrak{e}_{6,\mathbf{H}})^{\mathfrak{C}}, A, B \in (\mathfrak{J}_{\mathbf{H}})^{\mathfrak{C}}, \nu \in \mathfrak{C} \},$$

where the group $(E_{7,\mathbf{H}})^{\mathfrak{C}}$ is defined by

$$(E_{7,\mathbf{H}})^{\mathfrak{C}} = \{ \alpha \in \text{Iso}_{\mathfrak{C}}((\mathfrak{P}_{\mathbf{H}})^{\mathfrak{C}}) \mid \alpha(P \times Q)\alpha^{-1} = \alpha P \times \alpha Q \}.$$

Hence, using $((E_{7,\mathbf{H}})^{\mathfrak{C}})^{\tau\lambda} = E_{7,\mathbf{H}}$ shown in Proposition 6.3 (1), we obtain the required result. □

The immediate aim is to prove the connectedness of $E_{7,\mathbf{H}}$. We explain its procedure. First, since we have the isomorphism $E_{6,\mathbf{H}} \cong SU(6)/\mathbf{Z}_2$ (Theorem 5.1), the connectedness of $E_{6,\mathbf{H}}$ follows from its result. After this, we will prove that the homogeneous space $E_{7,\mathbf{H}}/E_{6,\mathbf{H}}$ is homeomorphic to a space $(\mathfrak{M}_{\mathbf{H}})_1$, so that the connectedness of $E_{7,\mathbf{H}}$ is finally shown, where the space $(\mathfrak{M}_{\mathbf{H}})_1$ is defined later.

Now, we will make some preparations. First, we define a space $(\mathfrak{M}_{\mathbf{H}})^{\mathfrak{C}}$ by

$$\begin{aligned} (\mathfrak{M}_{\mathbf{H}})^{\mathfrak{C}} &:= \{ P \in (\mathfrak{P}_{\mathbf{H}})^{\mathfrak{C}} \mid P \times P = 0, P \neq 0 \} \\ &= \left\{ P = (X, Y, \xi, \eta) \mid \begin{array}{l} X \vee Y = 0, X \times X = \eta Y \\ Y \times Y = \xi X, (X, Y) = 3\xi\eta \end{array} \right\}. \end{aligned}$$

Here, for example, the following elements of $(\mathfrak{M}_{\mathbf{H}})^{\mathfrak{C}}$

$$\left(X, \frac{1}{\eta}(X \times X), \frac{1}{\eta^2}\det X, \eta \right), \left(\frac{1}{\xi}(Y \times Y), Y, \xi, \frac{1}{\xi^2}\det Y \right) \quad \dot{1} = (0, 0, 1, 0), \quad \dot{1} = (0, 0, 0, 1)$$

belong to $(\mathfrak{M}_{\mathbf{H}})^{\mathfrak{C}}$, where $\xi \neq 0, \eta \neq 0$. Obviously, the group $E_{7,\mathbf{H}}$ acts on $(\mathfrak{M}_{\mathbf{H}})^{\mathfrak{C}}$.

Here, we will study the following subgroup $(E_{7,\mathbf{H}})_1$ of $E_{7,\mathbf{H}}$:

$$(E_{7,\mathbf{H}})_1 := \{ \alpha \in E_{7,\mathbf{H}} \mid \alpha \dot{1} = \dot{1} \}.$$

Proposition 6.3. (1) *The group $E_{7,\mathbf{H}}$ coincides with the group $((E_{7,\mathbf{H}})^{\mathfrak{C}})^{\tau\lambda} \cdot E_{7,\mathbf{H}} = ((E_{7,\mathbf{H}})^{\mathfrak{C}})^{\tau\lambda}$.*

(2) *If $\alpha \in E_{7,\mathbf{H}}$ satisfies $\alpha \dot{1} = \dot{1}$, then α also satisfies $\alpha \dot{1} = \dot{1}$ and conversely.*

Proof. (1) Let $\alpha \in E_{7,\mathbf{H}}$. Then, note that $\{\alpha P, \alpha Q\} = \{P, Q\}, P, Q \in (\mathfrak{P}_{\mathbf{H}})^C$, so that it follows from

$$\langle P, \lambda Q \rangle = \langle P, Q \rangle = \langle \alpha P, \alpha Q \rangle = \langle \alpha P, \lambda \alpha Q \rangle = \langle P, {}^t \alpha \lambda \alpha Q \rangle$$

that $\lambda = {}^t \alpha \lambda \alpha$, that is, ${}^t \alpha^{-1} = \lambda \alpha \lambda^{-1}$. In addition, we have ${}^t \alpha \tau \alpha = \tau$ from $\langle \alpha P, \alpha Q \rangle = \langle P, Q \rangle$. Hence, together with the formula ${}^t \alpha^{-1} = \lambda \alpha \lambda^{-1}$ above, that is, ${}^t \alpha = \lambda \alpha^{-1} \lambda^{-1}$, we have $(\tau \lambda) \alpha = \alpha(\tau \lambda)$, so that $E_{7,\mathbf{H}} \subset ((E_{7,\mathbf{H}})^C)^{\tau \lambda}$.

Conversely, let $\beta \in ((E_{7,\mathbf{H}})^C)^{\tau \lambda}$. Then it follows from $\langle P, Q \rangle = \langle \tau \lambda P, Q \rangle$ that

$$\langle \beta P, \beta Q \rangle = \langle (\tau \lambda) \beta P, \beta Q \rangle = \langle \beta(\tau \lambda) P, \beta Q \rangle = \langle \tau \lambda P, Q \rangle = \langle P, Q \rangle.$$

Hence we have $((E_{7,\mathbf{H}})^C)^{\tau \lambda} \subset E_{7,\mathbf{H}}$. Thus we have $E_{7,\mathbf{H}} = ((E_{7,\mathbf{H}})^C)^{\tau \lambda}$.

(2) By using the result of (1) above, it follows that $\alpha \mathbb{1} = \alpha(\tau \lambda)(-\mathbb{1}) = (\tau \lambda) \alpha(-\mathbb{1}) = \tau \lambda(-\mathbb{1}) = \mathbb{1}$. \square

Then we have the following theorem.

Proposition 6.4. *The group $(E_{7,\mathbf{H}})_1$ is isomorphic to the group $E_{6,\mathbf{H}}$: $(E_{7,\mathbf{H}})_1 \cong E_{6,\mathbf{H}}$.*

In particular, the group $(E_{7,\mathbf{H}})_1$ is connected.

Proof. We associate an element $\beta \in E_{6,\mathbf{H}}$ with the element

$$\tilde{\beta} = \begin{pmatrix} \beta & 0 & 0 & 0 \\ 0 & \tau \beta \tau & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \in (E_{7,\mathbf{H}})_1.$$

Then, for $P = (X, Y, \xi, \eta), Q = (Z, W, \zeta, \omega) \in (\mathfrak{P}_{\mathbf{H}})^C$, using the formula $\beta X \vee \tau \beta \tau W = \beta(X \vee W) \beta^{-1}$, it follows that

$$\begin{aligned} \tilde{\beta} P \times \tilde{\beta} Q &= (\beta X, \tau \beta \tau Y, \xi, \eta) \times (\beta Z, \tau \beta \tau W, \zeta, \omega) \\ &= \begin{pmatrix} (-1/2)(\beta X \vee \tau \beta \tau W + \beta Z \vee \tau \beta \tau Y) \\ (-1/4)(2\tau \beta \tau Y \times \tau \beta \tau W - \xi \beta Z - \zeta \beta X) \\ (1/4)(2\beta X \times \beta Z - \eta \tau \beta \tau W - \omega \tau \beta \tau Y) \\ (-1/8)((\beta X, \tau \beta \tau W) + (\beta Z, \tau \beta \tau Y) - 3(\xi \omega + \zeta \eta)) \end{pmatrix} \\ &= \begin{pmatrix} (-1/2)(\beta(X \vee W) \beta^{-1} + \beta(Z \vee Y) \beta^{-1}) \\ (-1/4)(\beta(2Y \times W - \xi Z - \zeta X)) \\ (1/4)(\tau \beta \tau(2X \times Z - \eta W - \omega Y)) \\ (-1/8)((X, W) + (Z, Y) - 3(\xi \omega + \zeta \eta)) \end{pmatrix}. \end{aligned}$$

Here, set $P \times Q =: \Phi(\phi, A, B, \nu)$, where

$$\begin{aligned} \phi &= (-1/2)(X \vee W + Z \vee Y), \quad A = (-1/4)(2Y \times W - \xi Z - \zeta X), \\ B &= (1/4)(2X \times Z - \eta W - \omega Y), \quad \nu = (1/8)((X, W) + (Z, Y) - 3(\xi \omega + \zeta \eta)), \end{aligned}$$

then we have $\tilde{\beta} P \times \tilde{\beta} Q = \Phi(\beta \phi \beta^{-1}, \beta A, \tau \beta \tau B, \nu)$.

On the other hand, for $R := (S, T, \varsigma, \varrho) \in (\mathfrak{P}_{\mathbf{H}})^C$, using the formula $(\tau \beta \tau)^t \phi(\tau \beta^{-1} \tau) = {}^t(\beta \phi \beta^{-1})$, it follows that

$$\begin{aligned} \tilde{\beta}(P \times Q) \tilde{\beta}^{-1} R &= \tilde{\beta}(P \times Q)(\beta^{-1} S, \tau \beta^{-1} \tau T, \varsigma, \varrho) \\ &=: \tilde{\beta} \Phi(\phi, A, B, \nu)(\beta^{-1} S, \tau \beta^{-1} \tau T, \varsigma, \varrho) \quad (P \times Q =: \Phi(\phi, A, B, \nu)) \end{aligned}$$

$$\begin{aligned}
&= \beta \begin{pmatrix} \phi(\beta^{-1}S) - (1/3)\nu(\beta^{-1}S) + 2B \times (\tau\beta^{-1}\tau T) + \varrho A \\ 2A \times (\beta^{-1}S) - {}^t\phi(\tau\beta^{-1}\tau T) + (1/3)\nu(\tau\beta^{-1}\tau T) + \varsigma B \\ (A, \tau\beta^{-1}\tau T) + \nu\varsigma \\ (B, \beta^{-1}S) - \nu\varrho \end{pmatrix} \\
&= \begin{pmatrix} \beta\phi\beta^{-1}S - (1/3)\nu S + \beta(2B \times (\tau\beta^{-1}\tau T)) + \varrho\beta A \\ \tau\beta\tau(2A \times (\beta^{-1}S)) - (\tau\beta\tau){}^t\phi(\tau\beta^{-1}\tau T) + (1/3)\nu T + \varsigma{}^t\beta^{-1}B \\ (\beta A, T) + \nu\varsigma \\ (\tau\beta\tau B, S) - \nu\varrho \end{pmatrix} \\
&= \begin{pmatrix} \beta\phi\beta^{-1}S - (1/3)\nu S + 2(\tau\beta\tau)B \times T + \varrho\beta A \\ 2\beta A \times S - {}^t(\beta\phi\beta^{-1})T + (1/3)\nu T + \varsigma(\tau\beta\tau)B \\ (\beta A, T) + \nu\varsigma \\ ((\tau\beta\tau)B, S) - \nu\varrho \end{pmatrix} \\
&= \Phi(\beta\phi\beta^{-1}, \beta A, (\tau\beta\tau)B, \nu)(S, T, \varsigma, \varrho),
\end{aligned}$$

that is, $\tilde{\beta}(P \times Q)\tilde{\beta}^{-1} = \Phi(\beta\phi\beta^{-1}, \beta A, {}^t\beta^{-1}B, \nu)$.

Hence we have $\tilde{\beta}P \times \tilde{\beta}Q = \tilde{\beta}(P \times Q)\tilde{\beta}^{-1}$. Moreover, it is easy to verify that $\langle \tilde{\beta}P, \tilde{\beta}Q \rangle = \langle P, Q \rangle$. Indeed, for $P = (X, Y, \xi, \eta), Q = (Z, W, \zeta, \omega) \in (\mathfrak{P}_{\mathbf{H}})^C$, note that $\tau\beta\tau \in E_{7, \mathbf{H}}$, it follows that

$$\begin{aligned}
\langle \tilde{\beta}P, \tilde{\beta}Q \rangle &= \langle (\beta X, (\tau\beta\tau)Y, \xi, \eta), (\beta Z, (\tau\beta\tau)W, \zeta, \omega) \rangle \\
&= \langle \beta X, \beta Z \rangle + \langle (\tau\beta\tau)Y, (\tau\beta\tau)W \rangle + (\tau\xi)\eta + (\tau\eta)\omega \\
&= \langle X, Z \rangle + \langle Y, W \rangle + (\tau\xi)\eta + (\tau\eta)\omega \\
&= \langle P, Q \rangle,
\end{aligned}$$

so that we have $\tilde{\beta} \in E_{7, \mathbf{H}}$. Moreover, since it is trivial that $\tilde{\beta}\mathbf{1} = \mathbf{1}$, we obtain $\tilde{\beta} \in (E_{7, \mathbf{H}})_1$.

Conversely, let $\alpha \in (E_{7, \mathbf{H}})_1$. Then α is of the form

$$\alpha = \begin{pmatrix} \beta & \varepsilon & 0 & 0 \\ \delta & \beta' & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \quad \beta, \beta', \delta, \varepsilon \in \text{Hom}_C((\mathfrak{J}_{\mathbf{H}})^C).$$

Indeed, set

$$\alpha = \begin{pmatrix} \beta & \varepsilon & C & A \\ \delta & \beta' & B & D \\ c & a & \nu & \lambda \\ b & d & \kappa & \mu \end{pmatrix}, \quad \begin{array}{l} \beta, \beta', \delta, \varepsilon \in \text{Hom}_C((\mathfrak{J}_{\mathbf{H}})^C), \\ a, b, c, d \in \text{Hom}_C((\mathfrak{J}_{\mathbf{H}})^C, C), \\ A, B, C, D \in (\mathfrak{J}_{\mathbf{H}})^C, \\ \nu, \mu, \lambda, \kappa \in C. \end{array}$$

Then, since $\alpha\mathbf{1} = \mathbf{1}$ implies $\alpha\mathbf{1} = \mathbf{1}$ (Lemma 6.2(2)), it follows that

$$\begin{aligned}
\mathbf{1} &= \alpha\mathbf{1} = \dot{C} + \dot{B} + \dot{\nu} + \kappa \\
\mathbf{1} &= \alpha\mathbf{1} = \dot{A} + \dot{D} + \dot{\lambda} + \mu.
\end{aligned}$$

Hence we have $A = B = C = D = \mathbf{O}$ (\mathbf{O} is zero matrix), $\kappa = \lambda = 0$ and $\nu = \mu = 1$. Moreover, it follows from

$$\{\alpha\dot{X}, \alpha\mathbf{1}\} = \{\alpha\dot{X}, \alpha\mathbf{1}\} = 0, \quad \{\alpha\dot{X}, \alpha\mathbf{1}\} = \{\alpha\dot{X}, \alpha\mathbf{1}\} = 0$$

that

$$\begin{aligned}
0 &= \{\alpha\dot{X}, \alpha\mathbf{1}\} = \{(\beta X, \delta X, c(X), b(X)), (0, 0, 1, 0)\} = -b(X), \\
0 &= \{\alpha\dot{X}, \alpha\mathbf{1}\} = \{(\beta X, \delta X, c(X), b(X)), (0, 0, 0, 1)\} = c(X), \\
0 &= \{\alpha\mathbf{1}, \alpha\mathbf{1}\} = \{(\varepsilon X, \beta' X, a(X), d(X)), (0, 0, 1, 0)\} = -d(X), \\
0 &= \{\alpha\mathbf{1}, \alpha\mathbf{1}\} = \{(\varepsilon X, \beta' X, a(X), d(X)), (0, 0, 0, 1)\} = a(X),
\end{aligned}$$

that is, $-b(X) = c(X) = -d(X) = a(X) = 0$ hold for all $X \in (\mathfrak{J}_{\mathbf{H}})^C$. Hence we have $a = b = c = d = 0$. With above, α is of the form above.

The group $(E_{7,\mathbf{H}})_1$ act on the space $(\mathfrak{M}_{\mathbf{H}})^C$, obviously. Hence, for $(X, \frac{1}{\eta}(X \times X), \frac{1}{\eta^2} \det X, \eta) \in (\mathfrak{M}_{\mathbf{H}})^C, \eta \neq 0$, since it follows that

$$(*) \cdots \alpha(X, \frac{1}{\eta}(X \times X), \frac{1}{\eta^2} \det X, \eta) = (\beta X + \frac{1}{\eta} \varepsilon(X \times X), \delta X + \frac{1}{\eta} \beta'(X \times X), \frac{1}{\eta^2} \det X, \eta) \in (\mathfrak{M}_{\mathbf{H}})^C,$$

the following formula

$$(\beta X + \frac{1}{\eta} \varepsilon(X \times X)) \times (\beta X + \frac{1}{\eta} \varepsilon(X \times X)) = \eta(\delta X + \frac{1}{\eta} \beta'(X \times X))$$

holds for all $\eta \in C, \eta \neq 0$. Hence the formula above holds for $\eta = 1, -1, 2, -2, 1/4$, so that we have that $\delta X = 0$ holds for all $X \in (\mathfrak{J}_{\mathbf{H}})^C$. Thus we obtain $\delta = 0$. Moreover, for $(\frac{1}{\xi}(Y \times Y), Y, \xi, \frac{1}{\xi^2} \det Y) \in (\mathfrak{M}_{\mathbf{H}})^C, \xi \neq 0$, by doing a similar computation above, we have that the formula $\beta'Y \times \beta'Y = \beta(Y \times Y) + \xi \varepsilon(Y)$ holds for all $\xi \in C, \xi \neq 0$. Hence the formula above holds for $\xi = 1, -1$, so that we have that $\varepsilon(Y) = 0$ holds for all $Y \in (\mathfrak{J}_{\mathbf{H}})^C$. Thus we obtain $\varepsilon = 0$. Therefore α is of the form

$$\alpha = \begin{pmatrix} \beta & 0 & 0 & 0 \\ 0 & \beta' & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \quad \beta, \beta' \in \text{Hom}_C((\mathfrak{J}_{\mathbf{H}})^C).$$

Again, in the formula (*) above, set $\eta = 1$. Then the condition $\alpha(X, X \times X, \det X, 1) \in (\mathfrak{M}_{\mathbf{H}})^C$ implies

$$\begin{cases} \beta X \times \beta X = \beta'(X \times X) \\ (\beta X, \beta'(X \times X)) = 3 \det X. \end{cases}$$

Hence it follows from two formulas above that

$$3 \det(\beta X) = (\beta X, \beta X \times \beta X) = (\beta X, \beta'(X \times X)) = 3 \det X,$$

that is, $\det(\beta X) = \det X$. Moreover, $\langle \alpha \dot{X}, \alpha \dot{X}' \rangle = \langle \dot{X}, \dot{X}' \rangle$ implies $\langle \beta X, \beta X' \rangle = \langle X, X' \rangle$. Thus we have $\beta \in E_{6,\mathbf{H}}$.

Next, as for β' , using the formula $\beta X \times \beta X = \beta'(X \times X)$, it follows from $\beta \in E_{6,\mathbf{H}}$ that

$$\beta'(X \times X) = \beta X \times \beta X = \tau \beta \tau(X \times X),$$

that is $\beta'(X \times X) = \tau \beta \tau(X \times X) \cdots (**)$.

Therefore we can obtain

$$\beta' = \tau \beta \tau.$$

Indeed, set $X \times X$ instead of X in the formula (**), then we have the following formula

$$(\det X) \beta' X = (\det X) \tau \beta \tau X.$$

In the case where $\det X \neq 0$, we have $\beta' X = \tau \beta \tau X$ for all $X \in (\mathfrak{J}_{\mathbf{H}})^C$, that is $\beta' = \tau \beta \tau$. In the case $\det X = 0$. We consider a space $S := \{X \in (\mathfrak{J}_{\mathbf{H}})^C \mid \det X \neq 0\}$. Then it follows from $\bar{S} = (\mathfrak{J}_{\mathbf{H}})^C$ that S is dense in $(\mathfrak{J}_{\mathbf{H}})^C$, and since $\beta', \tau \beta \tau \in \text{Hom}_C((\mathfrak{J}_{\mathbf{H}})^C)$ are continuous mappings and agree on S , these also agree on $(\mathfrak{J}_{\mathbf{H}})^C$. Namely, $\beta' = \tau \beta \tau$ holds even if $\det X = 0$.

Finally, the connectedness of the group $(E_{7,\mathbf{H}})_1$ follows from the connectedness of the group $E_{6,\mathbf{H}}$.

Consequently, the proof of this proposition is completed. \square

Here, we prove the useful lemma in order to prove Theorem 6.6 below.

Lemma 6.5. *Any element $P \in (\mathfrak{M}_{\mathbf{H}})^C$ can be transformed to a diagonal form by some element $\alpha \in (E_{7,\mathbf{H}})_0$:*

$$\alpha P = (X, Y, \xi, \eta), \quad X, Y: \text{diagonal forms}, \xi > 0.$$

Proof. Before proving this lemma, we prove the following claim.

Claim. *Any element $P \in (\mathfrak{J}_{\mathbf{H}})^C$ can be transformed to a diagonal form by some element $\beta \in E_{6,\mathbf{H}}$:*

$$\beta X = \begin{pmatrix} \xi_1 & 0 & 0 \\ 0 & \xi_2 & 0 \\ 0 & 0 & \xi_3 \end{pmatrix}, \quad \xi_i \in C.$$

Proof. The proof of this claim can be proved in exactly the same way as the proof of [9, Proposition 3.8.2], so its proof is omitted. \square

We start to prove the proof and will first show that P can be transformed to a diagonal form with $\xi \neq 0$.

Let $P = (X, Y, \xi, \eta) \in (\mathfrak{M}_{\mathbf{H}})^C$.

Case (i) where $P = (X, Y, \xi, \eta), \xi \neq 0$. Then we choose $\beta \in E_{6,\mathbf{H}}$ such that $\tau\beta\tau Y$ is a diagonal form (Claim). Hence since it follows from $X = (1/\xi)(Y \times Y)$ that

$$\beta X = \frac{1}{\xi}\beta(Y \times Y) = \frac{1}{\xi}(\tau\beta\tau Y \times \tau\beta\tau Y),$$

βX is also a diagonal form. Thus P is a diagonal form.

Case (ii) where $P = (X, Y, 0, \eta), Y \neq 0$. As in the Case (i), we choose $\beta \in E_{6,\mathbf{H}}$ such that $\tau\beta\tau Y$ is a diagonal form:

$$\tau\beta\tau Y := \text{diag}(\eta_1, \eta_2, \eta_3), \quad \eta_i \in C.$$

Hence, from $Y \neq 0$, some η_i is non-zero: $\eta_i \neq 0$. For this $\eta_i \in C$, we choose $\alpha_i(a) := \exp(\Phi(0, aE_i, -\tau aE_i, 0)) \in (E_{7,\mathbf{H}})_0, a \in C, a \neq 0$ (Lemma 6.2). The action of $\alpha_i(a)$ to $(\mathfrak{J}_{\mathbf{H}})^C$ is given as follows:

$$\alpha_i(a)(X, Y, \xi, \eta) = \begin{pmatrix} (1 + (\cos |a| - 1)p_i)X - 2\tau a \frac{\sin |a|}{|a|} E_i \times Y + \eta a \frac{\sin |a|}{|a|} E_i \\ 2a \frac{\sin |a|}{|a|} E_i \times X + (1 + (\cos |a| - 1)p_i)Y - \xi \tau a \frac{\sin |a|}{|a|} E_i \\ (a \frac{\sin |a|}{|a|} E_i, Y) + \xi \cos |a| \\ (-\tau a \frac{\sin |a|}{|a|} E_i, X) + \eta \cos |a| \end{pmatrix},$$

where $p_i : (\mathfrak{J}_{\mathbf{H}})^C \rightarrow (\mathfrak{J}_{\mathbf{H}})^C$ is defined by $p_i(X) = (X, E_i)E_i + 4E_i \times (E_i \times X)$ (cf. [4, Lemma 4.3.4]).

Apply $\alpha_i(-\pi/2)$ on $\beta P = (\beta X, \tau\beta\tau Y, 0, \eta)$, then we have the following

$$\begin{aligned} \alpha_i\left(-\frac{\pi}{2}\right)(\beta P) &= \alpha_i\left(-\frac{\pi}{2}\right)(\beta X, \tau\beta\tau Y, 0, \eta) \\ &= \begin{pmatrix} (1 - p_i)(\beta X) + 2E_i \times (\tau\beta\tau Y) - \eta E_i \\ -2E_i \times (\beta X) + (1 - p_i)(\tau\beta\tau Y) \\ (-E_i, \tau\beta\tau Y) \\ (E_i, \beta X) \end{pmatrix} \end{aligned}$$

$$= \begin{pmatrix} * \\ * \\ -\eta_i \\ * \end{pmatrix}, \quad \eta_i \neq 0.$$

Hence this case reduces to Case (i).

Case (iii) where $P = (X, 0, 0, \eta)$, $X \neq 0$. We choose $\beta \in E_{6,\mathbf{H}}$ such that $\beta X = \text{diag}(\xi_1, \xi_2, \xi_3)$, $\xi_i \in \mathbb{C}$. Then some ξ_i is non-zero from $\beta X \neq 0$: $\xi_i \neq 0$. Apply $\alpha_{i+1}(-\pi/2)$ on $\beta P = (\beta X, 0, 0, \eta)$, then we have the following

$$\begin{aligned} \alpha_{i+1}\left(-\frac{\pi}{2}\right)(\beta P) &= \alpha_{i+1}\left(-\frac{\pi}{2}\right)(\beta X, 0, 0, \eta) \\ &= \alpha_{i+1}\left(-\frac{\pi}{2}\right)(\xi_i E_i + \xi_{i+1} E_{i+1} + \xi_{i+2} E_{i+2}, 0, 0, \eta) \\ &= ((1 - p_{i+1})(\xi_i E_i + \xi_{i+1} E_{i+1} + \xi_{i+2} E_{i+2}) - \eta E_{i+1}, \\ &\quad - 2E_{i+1} \times (\xi_i E_i + \xi_{i+1} E_{i+1} + \xi_{i+2} E_{i+2}), 0, (E_{i+1}, \xi_i E_i + \xi_{i+1} E_{i+1} + \xi_{i+2} E_{i+2})) \\ &= (*, -\xi_i E_{i+2} - \xi_{i+2} E_i, 0, *), \quad \xi_i \neq 0. \end{aligned}$$

Hence this case reduces to Case (ii).

Case (iv) where $P = (0, 0, 0, \eta)$, $\eta \neq 0$. We choose $\alpha_1(a) \in (E_{7,\mathbf{H}})_0$, set $a = -\pi/2$. Apply $\alpha_1(-\pi/2)$ on P , then we have the following

$$\begin{aligned} \alpha_1\left(-\frac{\pi}{2}\right)P &= \alpha_1\left(-\frac{\pi}{2}\right)(0, 0, 0, \eta) \\ &= (\eta E_1, 0, 0, 0), \quad \eta \neq 0. \end{aligned}$$

Hence this case reduces to Case (iii).

With above, it was shown that P can be transformed to a diagonal form with $\xi \neq 0$. Finally, for $\theta \in U(1) := \{\theta \in \mathbb{C} \mid (\tau\theta)\theta = 1\}$, we define a mapping $\phi : (\mathfrak{P}_{\mathbf{H}})^{\mathbb{C}} \rightarrow (\mathfrak{P}_{\mathbf{H}})^{\mathbb{C}}$ by

$$\phi(\theta)(X, Y, \xi, \eta) = (\theta^{-1}X, \theta Y, \theta^3 \xi, \theta^{-3} \eta).$$

Then we have $\phi(\theta) \in E_{7,\mathbf{H}}$. For a given $P = (X, Y, \xi, \eta)$ transformed to a diagonal form with $\xi \neq 0$, since we can choose some $\theta \in U(1)$ such that $\xi\theta^3 > 0$, we obtain $P \in (\mathfrak{M}_{\mathbf{H}})^{\mathbb{C}}$ of the required form. The proof is completed. \square

We define a space $((\mathfrak{M}_{\mathbf{H}})^{\mathbb{C}})_1$ by

$$(\mathfrak{M}_{\mathbf{H}}^{\mathbb{C}})_1 := \{P \in (\mathfrak{M}_{\mathbf{H}})^{\mathbb{C}} \mid \langle P, P \rangle = 1\}.$$

Then we have the following theorem.

Theorem 6.6. *The homogeneous space $E_{7,\mathbf{H}}/E_{6,\mathbf{H}}$ is homeomorphic to the space $((\mathfrak{M}_{\mathbf{H}})^{\mathbb{C}})_1$: $E_{7,\mathbf{H}}/E_{6,\mathbf{H}} \simeq ((\mathfrak{M}_{\mathbf{H}})^{\mathbb{C}})_1$.*

In particular, the group $E_{7,\mathbf{H}}$ is connected.

Proof. For $\alpha \in E_{7,\mathbf{H}}$ and $P \in ((\mathfrak{M}_{\mathbf{H}})^{\mathbb{C}})_1$, we have $\alpha P \in ((\mathfrak{M}_{\mathbf{H}})^{\mathbb{C}})_1$. Hence $E_{7,\mathbf{H}}$ acts on $((\mathfrak{M}_{\mathbf{H}})^{\mathbb{C}})_1$. We will prove that the group $(E_7)_0$ acts transitively on $((\mathfrak{M}_{\mathbf{H}})^{\mathbb{C}})_1$. In order to prove this, it is sufficient to show that any element $P \in ((\mathfrak{M}_{\mathbf{H}})^{\mathbb{C}})_1$ can be transformed to $(0, 0, 1, 0) \in ((\mathfrak{M}_{\mathbf{H}})^{\mathbb{C}})_1$ by some $\alpha \in (E_{7,\mathbf{H}})_0$. Now, $P \in ((\mathfrak{M}_{\mathbf{H}})^{\mathbb{C}})_1$ can be transformed to a diagonal form by $\alpha \in (E_{7,\mathbf{H}})_0$:

$$\alpha P = \left(\frac{1}{\xi} \begin{pmatrix} \eta_2 \eta_3 & 0 & 0 \\ 0 & \eta_3 \eta_1 & 0 \\ 0 & 0 & \eta_1 \eta_2 \end{pmatrix}, \begin{pmatrix} \eta_1 & 0 & 0 \\ 0 & \eta_2 & 0 \\ 0 & 0 & \eta_3 \end{pmatrix}, \xi, \frac{1}{\xi^2} \eta_1 \eta_2 \eta_3\right), \quad \xi > 0$$

(Lemma 6.5). From the condition $\langle \alpha P, \alpha P \rangle = \langle P, P \rangle = 1$, we have

$$\frac{1}{\xi^2} (|\eta_2 \eta_3|^2 + |\eta_3 \eta_1|^2 + |\eta_1 \eta_2|^2) + (|\eta_1|^2 + |\eta_2|^2 + |\eta_3|^2) + \xi^2 + \frac{1}{\xi^4} |\eta_1 \eta_2 \eta_3|^2 = 1,$$

that is,

$$\left(1 + \frac{|\eta_1|^2}{\xi^2}\right)\left(1 + \frac{|\eta_2|^2}{\xi^2}\right)\left(1 + \frac{|\eta_3|^2}{\xi^2}\right) = \frac{1}{\xi^2}. \quad (*)$$

We choose $r_1, r_2, r_3 \in \mathbf{R}$, $0 \leq r_i < \pi/2$ such that $\tan r_i = |\eta_i|/\xi$, $i = 1, 2, 3$, then (*) becomes

$$\xi = \cos r_1 \cos r_2 \cos r_3.$$

By setting $a_i := (\eta_i/|\eta_i|)r_i$, $i = 1, 2, 3$ (if $\eta_i = 0$, then a_i means 0), we have

$$r_i = |a_i|, \quad \eta_i = \frac{1}{|a_i|} \frac{\eta_i}{|\eta_i|} r_i \frac{|\eta_i|}{\xi} \xi = \frac{a_i}{|a_i|} \tan r_i \cos r_1 \cos r_2 \cos r_3.$$

Hence we see that αP is of the form

$$\alpha P = \begin{pmatrix} \cos |a_1| a_2 \frac{\sin |a_2|}{|a_2|} a_3 \frac{\sin |a_3|}{|a_3|} & 0 & 0 \\ 0 & a_1 \frac{\sin |a_1|}{|a_1|} \cos |a_2| a_3 \frac{\sin |a_3|}{|a_3|} & 0 \\ 0 & 0 & a_1 \frac{\sin |a_1|}{|a_1|} a_2 \frac{\sin |a_2|}{|a_2|} \cos |a_3| \end{pmatrix} = \begin{pmatrix} a_1 \frac{\sin |a_1|}{|a_1|} \cos |a_2| \cos |a_3| & 0 & 0 \\ 0 & \cos |a_1| a_2 \frac{\sin |a_2|}{|a_2|} \cos |a_3| & 0 \\ 0 & 0 & \cos |a_1| \cos |a_2| a_3 \frac{\sin |a_3|}{|a_3|} \end{pmatrix}$$

Thus, using $\alpha_i(a)$ defined in the proof of Lemma 6.5, αP is expressed by

$$\alpha P = \alpha_3(a_3) \alpha_2(a_2) \alpha_1(a_1) (0, 0, 1, 0),$$

that is,

$$\alpha_1(a_1)^{-1} \alpha_2(a_2)^{-1} \alpha_3(a_3)^{-1} (\alpha P) = (0, 0, 1, 0).$$

This shows the transitivity of $(E_{7,\mathbf{H}})_0$. Since we have $((\mathfrak{M}_{\mathbf{H}})^C)_1 = (E_{7,\mathbf{H}})_0(0, 0, 1, 0)$, $((\mathfrak{M}_{\mathbf{H}})^C)_1$ is connected. The group $E_{7,\mathbf{H}}$ acts transitively on $((\mathfrak{M}_{\mathbf{H}})^C)_1$ and the isotropy subgroup of $E_{7,\mathbf{H}}$ at $\dot{1} = (0, 0, 1, 0) \in ((\mathfrak{M}_{\mathbf{H}})^C)_1$ is $E_{6,\mathbf{H}}$ (Proposition 6.3). Therefore we have the required homeomorphism

$$E_{7,\mathbf{H}}/E_{6,\mathbf{H}} \simeq ((\mathfrak{M}_{\mathbf{H}})^C)_1.$$

Finally, the connectedness of $E_{7,\mathbf{H}}$ follows from the connectedness of $((\mathfrak{M}_{\mathbf{H}})^C)_1$ and $E_{6,\mathbf{H}}$. \square

Here, we prove the following lemma.

Lemma 6.7. *The group $(E_{7,\mathbf{H}})^C$ is an algebraic subgroup of the general linear group $GL(32, \mathbf{C}) = \text{Iso}_{\mathbf{C}}((\mathfrak{P}_{\mathbf{H}})^C)$ and satisfies the condition that $\alpha \in (E_{7,\mathbf{H}})^C$ implies $\alpha^* = (\tau\lambda)\alpha^{-1}(\lambda^{-1}\tau) \in (E_{7,\mathbf{H}})^C$, where α^* is the conjugate transpose of α with respect to the Hermite inner product $\langle P, Q \rangle: \langle \alpha^* P, Q \rangle = \langle P, \alpha Q \rangle$.*

Proof. First, it is easy to verify that $\alpha^* = (\tau\lambda)\alpha^{-1}(\lambda^{-1}\tau)$. Indeed, for any $P, Q \in (\mathfrak{P}_{\mathbf{H}})^C$, it follows from

$$\begin{aligned} \langle \alpha^*P, Q \rangle &= \langle P, \alpha Q \rangle = \{\tau\lambda P, \alpha Q\} = \{\alpha\alpha^{-1}\tau\lambda P, \alpha Q\} \\ &= \{\alpha^{-1}\tau\lambda P, Q\} = \{(\tau\lambda)(\lambda^{-1}\tau)\alpha^{-1}\tau\lambda P, Q\} \\ &= \langle (\lambda^{-1}\tau)\alpha^{-1}(\tau\lambda)P, Q \rangle \end{aligned}$$

that $\alpha^* = (\lambda^{-1}\tau)\alpha^{-1}(\tau\lambda)$, that is, $\alpha^* = (\tau\lambda)\alpha^{-1}(\lambda^{-1}\tau)$. Subsequently, we will show $\alpha^* \in (E_{7,\mathbf{H}})^C$. It is clear that $\alpha^* \in \text{Iso}_C((\mathfrak{P}_{\mathbf{H}})^C)$ and it follows from $\tau P \times \tau Q = \tau(P \times Q)\tau$ and $\lambda \in (E_{7,\mathbf{H}})^C$ that

$$\begin{aligned} \alpha^*P \times \alpha^*Q &= (\tau\lambda)\alpha^{-1}(\lambda^{-1}\tau)P \times (\tau\lambda)\alpha^{-1}(\lambda^{-1}\tau)Q \\ &= \tau(\lambda\alpha^{-1}(\lambda^{-1}\tau)P \times \lambda\alpha^{-1}(\lambda^{-1}\tau)Q)\tau \\ &= \tau\lambda(\alpha^{-1}(\lambda^{-1}\tau)P \times \alpha^{-1}(\lambda^{-1}\tau)Q)\lambda^{-1}\tau \\ &= \tau\lambda\alpha^{-1}((\lambda^{-1}\tau)P \times (\lambda^{-1}\tau)Q)\alpha\lambda^{-1}\tau \\ &= (\tau\lambda)\alpha^{-1}(\lambda\tau)(P \times Q)(\tau\lambda^{-1})\alpha(\lambda^{-1}\tau) \\ &= \alpha^*(P \times Q)\alpha^{*-1}. \end{aligned}$$

Hence we have $\alpha^* \in (E_{7,\mathbf{H}})^C$.

Finally, the group $(E_{7,\mathbf{H}})^C$ is defined by the algebraic relation $\alpha P \times \alpha Q = \alpha(P \times Q)\alpha^{-1}$, so that $(E_{7,\mathbf{H}})^C$ is algebraic. \square

Let $O((\mathfrak{P}_{\mathbf{H}})^C)$ be the orthogonal subgroup $GL(32, C) = \text{Iso}_C((\mathfrak{P}_{\mathbf{H}})^C)$:

$$O(32, C) = O((\mathfrak{P}_{\mathbf{H}})^C) := \{\alpha \in \text{Iso}_C((\mathfrak{P}_{\mathbf{H}})^C) \mid \langle \alpha P, \alpha Q \rangle = \langle P, Q \rangle\}.$$

Then we have the following

$$\begin{aligned} (E_{7,\mathbf{H}})^C \cap O((\mathfrak{P}_{\mathbf{H}})^C) &= \{\alpha \in \text{Iso}_C((\mathfrak{P}_{\mathbf{H}})^C) \mid \alpha P \times \alpha Q = \alpha(P \times Q)\alpha^{-1}, \langle \alpha P, \alpha Q \rangle = \langle P, Q \rangle\} \\ &= E_{7,\mathbf{H}}. \end{aligned}$$

Using Chevalley's lemma ([1, Lemma 2]), we have a homeomorphism

$$\begin{aligned} (E_{7,\mathbf{H}})^C &\simeq ((E_{7,\mathbf{H}})^C \cap O((\mathfrak{P}_{\mathbf{H}})^C)) \times \mathbf{R}^d \\ &= E_{7,\mathbf{H}} \times \mathbf{R}^d, \end{aligned}$$

where the dimension d of the Euclidian part is computed by Lemma 6.2 as follows.

$$d = \dim((E_{7,\mathbf{H}})^C) - \dim(E_{7,\mathbf{H}}) = 66 \times 2 - 66 = 66.$$

With above, we have the following theorem needed in the next section.

Theorem 6.8. *The group $(E_{7,\mathbf{H}})^C$ is homeomorphic to the topological product of the group $E_{7,\mathbf{H}}$ and a 66-dimensional Euclidian space \mathbf{R}^{66} :*

$$(E_{7,\mathbf{H}})^C \simeq E_{7,\mathbf{H}} \times \mathbf{R}^{66}.$$

In particular, the group $(E_{7,\mathbf{H}})^C$ is connected.

Proof. The first half has already been shown above. The connectedness of the group $(E_{7,\mathbf{H}})^C$ follows from the connectedness of the group $E_{7,\mathbf{H}}$ (Theorem 6.6). \square

Using the C -linear transformations $\varepsilon_1, \varepsilon_2$ defined in previous section and the semi-linear transformation $\tau\lambda$ of $(\mathfrak{P}_{\mathbf{H}})^C$, we define a 12-dimensional \mathbf{R} -vector space V^{12} by

$$\begin{aligned} V^{12} &:= \{P \in \mathfrak{P}^C \mid \varepsilon_1 P = -iP, \varepsilon_2(\tau\lambda)P = P\} \\ &= \{P = (X, -\varepsilon_2\tau X, 0, 0) \mid X = F_1(x_1) + F_2(x_2) + F_3(x_3), x_i \in (\mathbf{H}^C e_4)_{\varepsilon_1}\} \end{aligned}$$

$$= \left\{ P = \left(\begin{array}{ccc} 0 & x_3 & \overline{x_2} \\ \overline{x_3} & 0 & x_1 \\ x_2 & \overline{x_1} & 0 \end{array} \right), \left(\begin{array}{ccc} 0 & -\varepsilon_2 \tau x_3 & -\overline{\varepsilon_2 \tau x_2} \\ -\overline{\varepsilon_2 \tau x_3} & 0 & -\varepsilon_2 \tau x_1 \\ -\varepsilon_2 \tau x_2 & -\overline{\varepsilon_2 \tau x_1} & 0 \end{array} \right), 0, 0 \right\} \left| x_i \in (\mathbf{H}^C e_4)_{\varepsilon_1} \right\}$$

with the norm

$$(P, P)_{\varepsilon_2} := \frac{1}{8} \{P, \varepsilon_2 P\},$$

where $(\mathbf{H}^C e_4)_{\varepsilon_1} := \{x \in \mathfrak{C}^C \mid x = p(e_4 + ie_5) + q(e_6 - ie_7), p, q \in \mathfrak{C}\}$. Let $x = p(e_4 + ie_5) + q(e_6 - ie_7) \in (\mathbf{H}^C e_4)_{\varepsilon_1}$, then note that $-\varepsilon_2 \tau x = -\tau q(e_4 + ie_5) + \tau p(e_6 - ie_7)$. Hence, for $P \in V^{12}$, the explicit form of $(P, P)_{\varepsilon_2}$ as $x_i := p_i(e_4 + ie_5) + q_i(e_6 - ie_7), i = 1, 2, 3$ is given by

$$(P, P)_{\varepsilon_2} = (p_1(\tau p_1) + q_1(\tau q_1)) + (p_2(\tau p_2) + q_2(\tau q_2)) + (p_3(\tau p_3) + q_3(\tau q_3)).$$

Next, we move on to study a group $(E_7)^{\varepsilon_1, \varepsilon_2}$ defined below.

We define a subgroup $(E_7)^{\varepsilon_1, \varepsilon_2}$ of E_7 by

$$(E_7)^{\varepsilon_1, \varepsilon_2} := \{\alpha \in E_7 \mid \varepsilon_1 \alpha = \alpha \varepsilon_1, \varepsilon_2 \alpha = \alpha \varepsilon_2\},$$

Then we have the following lemma and proposition.

Lemma 6.9. *The Lie algebra $(\mathfrak{e}_7)^{\varepsilon_1, \varepsilon_2}$ of the group $(E_7)^{\varepsilon_1, \varepsilon_2}$ is given by*

$$(\mathfrak{e}_7)^{\varepsilon_1, \varepsilon_2} = \left\{ \Phi(\phi, A, -\tau A, \nu) \in \mathfrak{e}_7 \left| \begin{array}{l} \phi = (D_1, D_2, D_3) + \tilde{A}_1(a_1) + \tilde{A}_2(a_2) + \tilde{A}_3(a_3) + i\tilde{T} \\ D_1 = d_{01}G_{01} + d_{02}G_{02} + d_{03}G_{03} + d_{12}G_{12} + d_{13}G_{13} \\ \quad + d_{23}G_{23} + d_{45}(G_{45} + G_{67}) + d_{46}(G_{46} - G_{57}) \\ \quad + d_{47}(G_{47} + G_{56}), d_{ij} \in \mathbf{R}, \\ D_2 = \pi \kappa D_1, D_3 = \kappa \pi D_1, \\ a_i \in \mathbf{H}, T \in (\mathfrak{J}_{\mathbf{H}})_0, \\ A \in (\mathfrak{J}_{\mathbf{H}})^C. \\ \nu \in i\mathbf{R} \end{array} \right. \right\}.$$

In particular, we have $\dim((\mathfrak{e}_7)^{\varepsilon_1, \varepsilon_2}) = (9 + 4 \times 3 + 15 - 1) + 30 + 1 = 66$.

Proof. Using the result of [9, Theorem 4.3.4], by straightforward computation, we can obtain the required result. \square

Proposition 6.10. *The group $(E_7)^{\varepsilon_1, \varepsilon_2}$ is connected.*

Proof. First, we will confirm that $\tau\lambda$ commutes with $\varepsilon_i, i = 1, 2$, respectively. Indeed, it follows that

$$\begin{aligned} (\tau\lambda)\varepsilon_i(X, Y, \xi, \eta) &= (\tau\lambda)(\varepsilon_i X, \varepsilon_i Y, \xi, \eta), (X, Y, \xi, \eta) \in \mathfrak{P}^C \\ &= (\varepsilon_i \tau Y, -\varepsilon_i \tau X, \tau \eta, -\tau \xi) \\ &= \varepsilon_i(\tau\lambda)(X, Y, \xi, \eta), \end{aligned}$$

that is, $(\tau\lambda)\varepsilon_i = \varepsilon_i(\tau\lambda)$. Hence we have

$$((E_7^C)^{\varepsilon_1, \varepsilon_2})^{\tau\lambda} = ((E_7^C)^{\tau\lambda})^{\varepsilon_1, \varepsilon_2} = (E_7)^{\varepsilon_1, \varepsilon_2}.$$

Thus, since $(E_7^C)^{\varepsilon_1, \varepsilon_2} (\cong Spin(12, C))$ is simply connected ([2, Proposition 1.1.7]), the group $(E_7)^{\varepsilon_1, \varepsilon_2} = ((E_7^C)^{\varepsilon_1, \varepsilon_2})^{\tau\lambda}$ is connected. \square

Now, we will construct the spinor group $Spin(12)$ in E_7 .

Proposition 6.11. *The group $(E_7)^{\varepsilon_1, \varepsilon_2}$ is isomorphic to the group $Spin(12):(E_7)^{\varepsilon_1, \varepsilon_2} \cong Spin(12)$.*

Proof. Let the orthogonal group $O(12) = O(V^{12}) = \{\beta \in \text{Iso}_{\mathbf{R}}(V^{12}) \mid (\beta P, \beta Q)_{\varepsilon_2} = (P, Q)_{\varepsilon_2}\}$. We consider the restriction $\alpha|_{V^{12}}$ of $\alpha \in (E_7)^{\varepsilon_1, \varepsilon_2}$ to V^{12} , then since $(E_7)^{\varepsilon_1, \varepsilon_2}$ acts on V^{12} , we see $\alpha|_{V^{12}} \in O(12)$. Moreover, since $(E_7)^{\varepsilon_1, \varepsilon_2}$ is connected (Proposition 6.10), we can define a homomorphism $p : (E_7)^{\varepsilon_1, \varepsilon_2} \rightarrow SO(V^{12}) = SO(12)$ by

$$p(\alpha) = \alpha|_{V^{12}}.$$

We will determine $\text{Ker } p$. However, since p is the restriction of the mapping π defined in [2, Proposition 1.1.7], we easily obtain $\text{Ker } p = \text{Ker } \pi = \{1, -\gamma\} \cong \mathbf{Z}_2$.

Finally, we will prove that p is surjective. Since the group $(E_7)^{\varepsilon_1, \varepsilon_2}$ is connected (Proposition 6.10) and $\text{Ker } p$ is discrete, together with $\dim((e_7)^{\varepsilon_1, \varepsilon_2}) = 66 = \dim(\mathfrak{so}(12))$ (Lemma 6.7), we see that p is surjective. Thus we have the isomorphism

$$(E_7)^{\varepsilon_1, \varepsilon_2} / \mathbf{Z}_2 \cong SO(12).$$

Therefore the group $(E_7)^{\varepsilon_1, \varepsilon_2}$ is isomorphic to $Spin(12)$ as the universal covering group of $SO(12)$, that is,

$$(E_7)^{\varepsilon_1, \varepsilon_2} \cong Spin(12).$$

□

Let $\alpha \in (E_7)^{\varepsilon_1, \varepsilon_2}$. Then, as in $(E_7^C)^{\varepsilon_1, \varepsilon_2}$, for $P \in (\mathfrak{P}_{\mathbf{H}})^C$, we have $\alpha P \in (\mathfrak{P}_{\mathbf{H}})^C$ because of $(E_7)^{\varepsilon_1, \varepsilon_2} \subset (E_7)^\gamma$. Hence α induces a C -linear isomorphism $\alpha : (\mathfrak{P}_{\mathbf{H}})^C \rightarrow (\mathfrak{P}_{\mathbf{H}})^C$.

Now, we will determine the structure of the group $E_{7,\mathbf{H}}$.

Theorem 6.12. *The group $E_{7,\mathbf{H}}$ is isomorphic to the semi-spinor group $Ss(12)$: $E_{7,\mathbf{H}} \cong Ss(12)$.*

Proof. Let $Spin(12)$ as the group $(E_7)^{\varepsilon_1, \varepsilon_2}$ (Proposition 6.11). Then we define a mapping $f_7 : (E_7)^{\varepsilon_1, \varepsilon_2} \rightarrow E_{7,\mathbf{H}}$ by

$$f_7(\alpha) = \alpha|_{(\mathfrak{P}_{\mathbf{H}})^C}.$$

As mentioned above, since $\alpha \in (E_7)^{\varepsilon_1, \varepsilon_2}$ induces the C -linear isomorphism $\alpha : (\mathfrak{P}_{\mathbf{H}})^C \rightarrow (\mathfrak{P}_{\mathbf{H}})^C$, we easily see $\alpha|_{(\mathfrak{P}_{\mathbf{H}})^C} \in E_{7,\mathbf{H}}$. Hence f_7 is well-defined. Moreover, it is clear that f_7 is a homomorphism.

Next, we will determine $\text{Ker } f_7$. In order to determine it, we will prove the following claim.

Claim 1. *The kernel of the differential mapping $f_{7*} : (e_7)^{\varepsilon_1, \varepsilon_2} \rightarrow e_{7,\mathbf{H}}$ of f_7 is trivial: $\text{Ker } f_{7*} = \{0\}$.*

Proof. Let $\Phi := \Phi(\phi, A, -\tau A, \nu) \in \text{Ker } f_{7*}$. Then Φ satisfies the condition

$$\Phi P = 0 \text{ for all } P \in (\mathfrak{P}_{\mathbf{H}})^C.$$

Set $P := (0, 0, 0, 1)$, then it follows from $0 = \Phi P = (A, 0, 0, -\nu)$ that $A = 0$ and $\nu = 0$. Hence Φ is of the form $\Phi(\phi, 0, 0, 0)$: $\Phi = \Phi(\phi, 0, 0, 0)$. In addition, set $P := (F_1(x), 0, 0, 0)$, $x \in \mathbf{H}^C$, then as $\phi = (D_1, D_2, D_3) + \tilde{A}_1(a_1) + \tilde{A}_2(a_2) + \tilde{A}_3(a_3) + (\tau_1 E_1 + \tau_2 E_2 + \tau_3 E_3 + F_1(t_1) + F_2(t_2) + F_3(t_3))^\sim$, it follows from

$$\begin{aligned} 0 = \Phi P &= (\phi F_1(x), 0, 0, 0) \\ &= (((a_1, x) + (t_1, x))E_2 + ((a_1, x) - (t_1, x))E_3 \\ &\quad + F_1(D_1 x - (1/2)\tau_1 x) + (1/2)F_2(\overline{a_3 x} + \overline{t_3 x}) + (1/2)F_3(-\overline{x a_2} + \overline{x t_2}), 0, 0, 0) \end{aligned}$$

that

$$\begin{aligned} (a_1, x) + (t_1, x) &= (a_1, x) - (t_1, x) = 0, \quad D_1 x - (1/2)\tau_1 x = 0, \\ \overline{a_3 x} + \overline{t_3 x} &= 0, \quad -\overline{x a_2} + \overline{x t_2} = 0 \text{ for all } x \in (\mathfrak{P}_{\mathbf{H}})^C. \end{aligned}$$

Moreover, set $P := (F_2(x), 0, 0, 0), (F_3(x), 0, 0, 0), x \in \mathbf{H}^C$, then similarly it follows from $\Phi P = 0$ that

$$\begin{cases} (a_2, x) + (t_2, x) = (a_2, x) - (t_2, x) = 0, & D_2x - (1/2)\tau_2x = 0, \\ \overline{a_1x} + \overline{t_1x} = 0, & -\overline{xa_3} + \overline{xt_3} = 0, \\ (a_3, x) + (t_3, x) = (a_3, x) - (t_3, x) = 0, & D_3x - (1/2)\tau_3x = 0, \\ \overline{a_2x} + \overline{t_2x} = 0, & -\overline{xa_1} + \overline{xt_1} = 0 \end{cases}$$

for all $x \in (\mathfrak{P}_{\mathbf{H}})^C$, respectively.

Hence we have

$$(*) \begin{cases} a_1 = a_2 = a_3 = 0, & t_1 = t_2 = t_3 = 0, \\ D_1x - (1/2)\tau_1x = 0, & D_2x - (1/2)\tau_2x = 0, & D_3x - (1/2)\tau_3x = 0. \end{cases}$$

Finally, set $P := (0, F_k(x), 0, 0), x \in \mathbf{H}^C$, then as results different from (*), we have

$$D_1x + (1/2)\tau_1x = 0, \quad D_2x + (1/2)\tau_2x = 0, \quad D_3x + (1/2)\tau_3x = 0$$

Therefore, together with (*), we have

$$D_1 = D_2 = D_3 = 0, \quad \tau_1 = \tau_2 = \tau_3 = 0.$$

With above, we obtain the required result. \square

Claim 2. The center $z((E_7)^{\varepsilon_1, \varepsilon_2})$ of the group $(E_7)^{\varepsilon_1, \varepsilon_2}$ coincides with the center $z((E_7^C)^{\varepsilon_1, \varepsilon_2})$ of the group $(E_7^C)^{\varepsilon_1, \varepsilon_2}$: $z((E_7)^{\varepsilon_1, \varepsilon_2}) = \{1, \gamma\} \times \{1, -\gamma\} \cong \mathbf{Z}_2 \times \mathbf{Z}_2$.

Proof. First, from Proposition 6.11 we can confirm

$$z((E_7)^{\varepsilon_1, \varepsilon_2}) \cong z(\text{Spin}(12)) \cong \mathbf{Z}_2 \times \mathbf{Z}_2,$$

and moreover from [2, Proposition 1.1.6] and $(E_7)^{\varepsilon_1, \varepsilon_2} \subset (E_7^C)^{\varepsilon_1, \varepsilon_2}$ we have

$$\begin{aligned} z((E_7)^{\varepsilon_1, \varepsilon_2}) \supset z((E_7^C)^{\varepsilon_1, \varepsilon_2}) &= \{1, \gamma\} \times \{1, -\gamma\} \\ &\cong \mathbf{Z}_2 \times \mathbf{Z}_2. \end{aligned}$$

Hence we obtain the required result. \square

Here, the group $(E_7)^{\varepsilon_1, \varepsilon_2}$ is connected (Proposition 6.10) and $\text{Ker } f_7$ is discrete (Claim 1), so that together with Claim 2, we have

$$\text{Ker } f_7 \subset z((E_7)^{\varepsilon_1, \varepsilon_2}) = \{1, \gamma\} \times \{1, -\gamma\}.$$

However, since $-1, -\gamma$ do not leave $(\mathfrak{P}_{\mathbf{H}})^C$, we see that $-1, -\gamma \notin \text{Ker } f_7$. Thus we have

$$\text{Ker } f_7 = \{1, \gamma\} = \mathbf{Z}_2.$$

Finally, we will prove that f_7 is surjective. Since the group $E_{7, \mathbf{H}}$ is connected (Theorem 6.6) and $\text{Ker } f_7$ is discrete, together with $\dim((\mathfrak{e}_7)^{\varepsilon_1, \varepsilon_2}) = 66 = \dim(\mathfrak{e}_{7, \mathbf{H}})$ (Lemmmas 6.7, 6.2), f_7 is surjective.

Therefore, from Proposition 6.11, we have the isomorphism

$$E_{7, \mathbf{H}} \cong \text{Spin}(12)/\mathbf{Z}_2,$$

that is, we have the required isomorphism

$$E_{7, \mathbf{H}} \cong \text{Ss}(12).$$

\square

We consider a subgroup $(E_7^C)^{\varepsilon_1, \varepsilon_2}$ of E_7^C by

$$(E_7^C)^{\varepsilon_1, \varepsilon_2} := \{\alpha \in E_7^C \mid \varepsilon_1 \alpha = \alpha \varepsilon_1, \varepsilon_2 \alpha = \alpha \varepsilon_2\}.$$

Then the group $(E_7^C)^{\varepsilon_1, \varepsilon_2}$ has been studied by author in [2, Proposition 1.1.7], and its result was obtained as follows:

$$(E_7^C)^{\varepsilon_1, \varepsilon_2} \cong Spin(12, C).$$

Now, we move the determination of the root system and the Dynkin diagram of the Lie algebra $(\mathfrak{e}_7^C)^{\varepsilon_1, \varepsilon_2}$ of the group $(E_7^C)^{\varepsilon_1, \varepsilon_2}$. After this, we will determine those by the same procedure as in $(\mathfrak{f}_4^C)^{\varepsilon_1, \varepsilon_2}$.

Then we need the following lemma used later.

Lemma 6.13. ([2, Lemma 1.1.2]) *The Lie algebra $(\mathfrak{e}_7^C)^{\varepsilon_1, \varepsilon_2}$ of the group $(E_7^C)^{\varepsilon_1, \varepsilon_2}$ is given by*

$$(\mathfrak{e}_7^C)^{\varepsilon_1, \varepsilon_2} = \left\{ \Phi(\phi, A, B, \nu) \in \mathfrak{e}_7^C \left| \begin{array}{l} \phi = (D_1, D_2, D_3) \\ +\tilde{A}_1(a_1) + \tilde{A}_2(a_2) + \tilde{A}_3(a_3) \\ +(\tau_1 E_1 + \tau_2 E_2 + \tau_3 E_3 + F_1(t_1) \\ +F_2(t_2) + F_3(t_3))^\sim, \\ D_1 = d_{01}G_{01} + d_{02}G_{02} + d_{03}G_{03} \\ +d_{12}G_{12} + d_{13}G_{13} + d_{23}G_{23} \\ +d_{45}(G_{45} + G_{67}) + d_{46}(G_{46} - G_{57}) \\ +d_{47}(G_{47} + G_{56}), d_{ij} \in C, \\ D_2 = \pi \kappa D_1, D_3 = \kappa \pi D_1, \\ a_k, t_k \in \mathbf{H}^C, \\ \tau_k \in C, \tau_1 + \tau_2 + \tau_3 = 0, \\ A, B \in (\mathfrak{J}_{\mathbf{H}})^C, \\ \nu \in C \end{array} \right. \right\}.$$

In particular, we have $\dim_C((\mathfrak{e}_7^C)^{\varepsilon_1, \varepsilon_2}) = (9 + 4 \times 3 + (3 - 1) + 4 \times 3) + 15 \times 2 + 1 = 66$.

Here, we define a Lie subalgebra \mathfrak{h}_7 of $(\mathfrak{e}_7^C)^{\varepsilon_1, \varepsilon_2}$ by

$$\mathfrak{h}_7 := \left\{ \Phi_7 = \Phi(\phi, 0, 0, \mu) \in (\mathfrak{e}_7^C)^{\varepsilon_1, \varepsilon_2} \left| \begin{array}{l} \phi = \delta + (\mu_1 E_1 + \mu_2 E_2 + \mu_3 E_3)^\sim \\ \delta := (L_1, L_2, L_3), \\ L_1 = \lambda_0(iG_{01}) + \lambda_1(iG_{23}) \\ +\lambda_2(i(G_{45} + G_{67})), \\ L_2 = \pi \kappa L_1, L_3 = \kappa \pi L_1, \lambda_i \in C, \\ \mu_k \in C, \mu_1 + \mu_2 + \mu_3 = 0, \\ \mu \in C \end{array} \right. \right\}.$$

Then \mathfrak{h}_7 is a Cartan subalgebra of $(\mathfrak{e}_7^C)^{\varepsilon_1, \varepsilon_2}$. Indeed, it is clear that $(\mathfrak{e}_7^C)^{\varepsilon_1, \varepsilon_2}$ is abelian. Next, by doing straightforward computation, we can confirm that $[\Phi, \Phi_7] \in \mathfrak{h}_7, \Phi \in (\mathfrak{e}_7^C)^{\varepsilon_1, \varepsilon_2}$ implies $\Phi \in \mathfrak{h}_7$ for any $\Phi_7 \in \mathfrak{h}_7$.

Theorem 6.14. *The rank of the Lie algebra $(\mathfrak{e}_7^C)^{\varepsilon_1, \varepsilon_2}$ is six. The roots Δ of $(\mathfrak{e}_7^C)^{\varepsilon_1, \varepsilon_2}$ relative to \mathfrak{h}_7 are given by*

$$\Delta = \left\{ \begin{array}{l} \pm(\lambda_0 - \lambda_1), \pm(\lambda_0 + \lambda_1), \pm 2\lambda_2, \\ \pm \frac{1}{2}(-2\lambda_0 + \mu_2 - \mu_3), \pm \frac{1}{2}(-2\lambda_0 - \mu_2 + \mu_3), \\ \pm \frac{1}{2}(-2\lambda_1 + \mu_2 - \mu_3), \pm \frac{1}{2}(-2\lambda_1 - \mu_2 + \mu_3), \\ \pm \frac{1}{2}(\lambda_0 - \lambda_1 - 2\lambda_2 - \mu_1 + \mu_3), \pm \frac{1}{2}(\lambda_0 - \lambda_1 - 2\lambda_2 + \mu_1 - \mu_3), \\ \pm \frac{1}{2}(\lambda_0 - \lambda_1 + 2\lambda_2 - \mu_1 + \mu_3), \pm \frac{1}{2}(\lambda_0 - \lambda_1 + 2\lambda_2 + \mu_1 - \mu_3), \\ \pm \frac{1}{2}(\lambda_0 + \lambda_1 + 2\lambda_2 + \mu_1 - \mu_2), \pm \frac{1}{2}(\lambda_0 + \lambda_1 + 2\lambda_2 - \mu_1 + \mu_2), \\ \pm \frac{1}{2}(-\lambda_0 - \lambda_1 + 2\lambda_2 + \mu_1 - \mu_2), \pm \frac{1}{2}(-\lambda_0 - \lambda_1 + 2\lambda_2 - \mu_1 + \mu_2), \\ \pm(\mu_1 + \frac{2}{3}\mu), \pm(\mu_2 + \frac{2}{3}\mu), \pm(\mu_3 + \frac{2}{3}\mu), \\ \pm(-\lambda_0 - \frac{1}{2}\mu_1 + \frac{2}{3}\mu), \pm(\lambda_0 - \frac{1}{2}\mu_1 + \frac{2}{3}\mu), \pm(-\lambda_1 - \frac{1}{2}\mu_1 + \frac{2}{3}\mu), \pm(\lambda_1 - \frac{1}{2}\mu_1 + \frac{2}{3}\mu), \\ \pm(-\frac{1}{2}(-\lambda_0 + \lambda_1 + 2\lambda_2) - \frac{1}{2}\mu_2 + \frac{2}{3}\mu), \pm(\frac{1}{2}(-\lambda_0 + \lambda_1 + 2\lambda_2) - \frac{1}{2}\mu_2 + \frac{2}{3}\mu), \\ \pm(-\frac{1}{2}(-\lambda_0 + \lambda_1 - 2\lambda_2) - \frac{1}{2}\mu_2 + \frac{2}{3}\mu), \pm(\frac{1}{2}(-\lambda_0 + \lambda_1 - 2\lambda_2) - \frac{1}{2}\mu_2 + \frac{2}{3}\mu), \\ \pm(-\frac{1}{2}(-\lambda_0 - \lambda_1 - 2\lambda_2) - \frac{1}{2}\mu_3 + \frac{2}{3}\mu), \pm(\frac{1}{2}(-\lambda_0 - \lambda_1 - 2\lambda_2) - \frac{1}{2}\mu_3 + \frac{2}{3}\mu), \\ \pm(-\frac{1}{2}(\lambda_0 + \lambda_1 - 2\lambda_2) - \frac{1}{2}\mu_3 + \frac{2}{3}\mu), \pm(\frac{1}{2}(\lambda_0 + \lambda_1 - 2\lambda_2) - \frac{1}{2}\mu_3 + \frac{2}{3}\mu) \end{array} \right\}.$$

Proof. The roots of $(\mathfrak{e}_6^C)^{\varepsilon_1, \varepsilon_2}$ is also the roots of $(\mathfrak{e}_7^C)^{\varepsilon_1, \varepsilon_2}$. Indeed, let the root α of $(\mathfrak{e}_6^C)^{\varepsilon_1, \varepsilon_2}$ and its associated root vector $V \in (\mathfrak{e}_6^C)^{\varepsilon_1, \varepsilon_2} \subset (\mathfrak{e}_7^C)^{\varepsilon_1, \varepsilon_2}$. Then it follows that

$$\begin{aligned} [\Phi_7, V] &= [\Phi(\phi, 0, 0, \mu), \Phi(V, 0, 0, 0)] = \Phi([\phi, V], 0, 0, 0) \\ &= \Phi(\alpha(\phi)V, 0, 0, 0) = \alpha(\phi)\Phi(V, 0, 0, 0) \\ &= \alpha(\Phi_7)V. \end{aligned}$$

We will determine the remainders of roots and give a few examples for that. First, let $\Phi_7 = \Phi(\phi, 0, 0, \nu) \in \mathfrak{h}_7$ and $\Phi(0, E_1, 0, 0) \in (\mathfrak{e}_7^C)^{\varepsilon_1, \varepsilon_2}$. Then it follows that

$$\begin{aligned} [\Phi_7, \Phi(0, E_1, 0, 0)] &= [\Phi(\phi, 0, 0, \mu), \Phi(0, E_1, 0, 0)] \\ &= \Phi(0, (\phi + (2/3)\mu)E_1, 0, 0) \\ &= \Phi(0, (\mu_1 + (2/3)\mu)E_1, 0, 0) \\ &= (\mu_1 + (2/3)\mu)\Phi(0, E_1, 0, 0), \end{aligned}$$

that is, $[\Phi_7, \Phi(0, E_1, 0, 0)] = (\mu_1 + (2/3)\mu)\Phi(0, E_1, 0, 0)$. Hence we see that $\mu_1 + (2/3)\mu$ is a root and $\Phi(0, E_1, 0, 0)$ is an associated root vector. Next, let $\Phi(0, F_1(1+ie_1), 0, 0) \in (\mathfrak{e}_7^C)^{\varepsilon_1, \varepsilon_2}$. Then it follows that

$$\begin{aligned} [\Phi, \Phi(0, F_1(1+ie_1), 0, 0)] &= [\Phi(\phi, 0, 0, \mu), \Phi(0, F_1(1), 0, 0)] \\ &= \Phi(0, (\phi + (2/3)\mu)F_1(1+ie_1), 0, 0) \\ &= \Phi(0, \phi F_1(1+ie_1) + (2/3)\mu F_1(1+ie_1), 0, 0) \\ &= \Phi(0, (-\lambda_0 + (1/2)(\mu_2 + \mu_3))F_1(1+ie_1) + (2/3)\mu F_1(1+ie_1), 0, 0) \\ &= (-\lambda_0 + (1/2)(\tau_2 + \tau_3) + (2/3)\mu)\Phi(0, F_1(1+ie_1), 0, 0), \end{aligned}$$

that is, $[\Phi_7, \Phi(0, F_1(1 + ie_1), 0, 0)] = (-\lambda_0 - (1/2)\mu_1 + (2/3)\mu)\Phi(0, F_1(1 + ie_1), 0, 0)$. Hence we see that $-\lambda_0 - (1/2)\mu_1 + (2/3)\mu$ is a root and $\Phi(0, F_1(1 + ie_1), 0, 0)$ is an associated root vector.

Together with two cases above, the remainders of roots and root vectors associated with these roots are obtained as follows:

roots	root vectors associated with roots
$\mu_1 + (2/3)\mu$	$\Phi(0, E_1, 0, 0)$
$-(\mu_1 + (2/3)\mu)$	$\Phi(0, 0, E_1, 0)$
$\mu_2 + (2/3)\mu$	$\Phi(0, E_2, 0, 0)$
$-(\mu_2 + (2/3)\mu)$	$\Phi(0, 0, E_2, 0)$
$\mu_3 + (2/3)\mu$	$\Phi(0, E_3, 0, 0)$
$-(\mu_3 + (2/3)\mu)$	$\Phi(0, 0, E_3, 0)$
$-\lambda_0 - (1/2)\mu_1 + (2/3)\mu$	$\Phi(0, F_1(1 + ie_1), 0, 0)$
$-(-\lambda_0 - (1/2)\mu_1 + (2/3)\mu)$	$\Phi(0, 0, F_1(1 - ie_1), 0)$
$\lambda_0 - (1/2)\mu_1 + (2/3)\mu$	$\Phi(0, F_1(1 - ie_1), 0, 0)$
$-(\lambda_0 - (1/2)\mu_1 + (2/3)\mu)$	$\Phi(0, 0, F_1(1 + ie_1), 0)$
$-\lambda_1 - (1/2)\mu_1 + (2/3)\mu$	$\Phi(0, F_1(e_2 + ie_3), 0, 0)$
$-(-\lambda_1 - (1/2)\mu_1 + (2/3)\mu)$	$\Phi(0, 0, F_1(e_2 - ie_3), 0)$
$\lambda_1 - (1/2)\mu_1 + (2/3)\mu$	$\Phi(0, F_1(e_2 - ie_3), 0, 0)$
$-(\lambda_1 - (1/2)\mu_1 + (2/3)\mu)$	$\Phi(0, 0, F_1(e_2 + ie_3), 0)$
$(-1/2)(-\lambda_0 + \lambda_1 + 2\lambda_2) - (1/2)\mu_2 + (2/3)\mu$	$\Phi(0, F_2(1 + ie_1), 0, 0)$
$-((-1/2)(-\lambda_0 + \lambda_1 + 2\lambda_2) - (1/2)\mu_2 + (2/3)\mu)$	$\Phi(0, 0, F_2(1 - ie_1), 0)$
$(1/2)(-\lambda_0 + \lambda_1 + 2\lambda_2) - (1/2)\mu_2 + (2/3)\mu$	$\Phi(0, F_2(1 - ie_1), 0, 0)$
$-((1/2)(-\lambda_0 + \lambda_1 + 2\lambda_2) - (1/2)\mu_2 + (2/3)\mu)$	$\Phi(0, 0, F_2(1 + ie_1), 0)$
$(-1/2)(-\lambda_0 + \lambda_1 - 2\lambda_2) - (1/2)\mu_2 + (2/3)\mu$	$\Phi(0, F_2(e_2 + ie_3), 0, 0)$
$-((-1/2)(-\lambda_0 + \lambda_1 - 2\lambda_2) - (1/2)\mu_2 + (2/3)\mu)$	$\Phi(0, 0, F_2(e_2 - ie_3), 0)$
$(1/2)(-\lambda_0 + \lambda_1 - 2\lambda_2) - (1/2)\mu_2 + (2/3)\mu$	$\Phi(0, F_2(e_2 - ie_3), 0, 0)$
$-((1/2)(-\lambda_0 + \lambda_1 - 2\lambda_2) - (1/2)\mu_2 + (2/3)\mu)$	$\Phi(0, 0, F_2(e_2 + ie_3), 0)$
$(-1/2)(-\lambda_0 - \lambda_1 - 2\lambda_2) - (1/2)\mu_3 + (2/3)\mu$	$\Phi(0, F_3(1 + ie_1), 0, 0)$
$-((-1/2)(-\lambda_0 - \lambda_1 - 2\lambda_2) - (1/2)\mu_3 + (2/3)\mu)$	$\Phi(0, 0, F_3(1 - ie_1), 0)$
$(1/2)(-\lambda_0 - \lambda_1 - 2\lambda_2) - (1/2)\mu_3 + (2/3)\mu$	$\Phi(0, F_3(1 - ie_1), 0, 0)$
$-((1/2)(-\lambda_0 - \lambda_1 - 2\lambda_2) - (1/2)\mu_3 + (2/3)\mu)$	$\Phi(0, 0, F_3(1 + ie_1), 0)$
$(-1/2)(\lambda_0 + \lambda_1 - 2\lambda_2) - (1/2)\mu_3 + (2/3)\mu$	$\Phi(0, F_3(e_2 + ie_3), 0, 0)$
$-((-1/2)(\lambda_0 + \lambda_1 - 2\lambda_2) - (1/2)\mu_3 + (2/3)\mu)$	$\Phi(0, 0, F_3(e_2 - ie_3), 0)$
$(1/2)(\lambda_0 + \lambda_1 - 2\lambda_2) - (1/2)\mu_3 + (2/3)\mu$	$\Phi(0, F_3(e_2 - ie_3), 0, 0)$
$-((1/2)(\lambda_0 + \lambda_1 - 2\lambda_2) - (1/2)\mu_3 + (2/3)\mu)$	$\Phi(0, 0, F_3(e_2 + ie_3), 0)$


Thus, since $(\mathfrak{e}_7^C)^{\varepsilon_1, \varepsilon_2}$ is spanned by \mathfrak{h}_7 and the root vectors associated with roots above, the roots obtained above are all. The rank of the Lie algebra $(\mathfrak{e}_7^C)^{\varepsilon_1, \varepsilon_2}$ follows from the dimension of \mathfrak{e}_7 . \square

Subsequently, we prove the following theorem.

Theorem 6.15. *In the root system Δ of Theorem 6.14*

$$\Pi = \{\alpha_1, \alpha_2, \alpha_3, \alpha_4, \alpha_5, \alpha_6\}$$

is a fundamental root system of $(\mathfrak{e}_7^C)^{\varepsilon_1, \varepsilon_2}$, where $\alpha_1 = -(\lambda_0 - \lambda_1)$, $\alpha_2 = (1/2)(\lambda_0 - \lambda_1 - 2\lambda_2 + \mu_1 - \mu_3)$, $\alpha_3 = 2\lambda_2$, $\alpha_4 = -(1/2)(\lambda_0 + \lambda_1 + 2\lambda_2 + \mu_1 - \mu_2)$, $\alpha_5 = \lambda_0 + \lambda_1$, $\alpha_6 = -(\mu_2 + (2/3)\mu)$. The Dynkin diagram of $(\mathfrak{e}_7^C)^{\varepsilon_1, \varepsilon_2}$ is given by



Proof. The all positive roots are expressed by $\alpha_1, \alpha_2, \alpha_3, \alpha_4, \alpha_5, \alpha_6$ as follows:

$$\begin{aligned}
-(\lambda_0 - \lambda_1) &= \alpha_1, \\
\lambda_0 + \lambda_1 &= \alpha_5, \\
2\lambda_2 &= \alpha_3, \\
(1/2)(-2\lambda_0 + \mu_2 - \mu_3) &= \alpha_1 + \alpha_2 + \alpha_3 + \alpha_4, \\
-(1/2)(-2\lambda_0 - \mu_2 + \mu_3) &= \alpha_2 + \alpha_3 + \alpha_4 + \alpha_5, \\
(1/2)(-2\lambda_1 + \mu_2 - \mu_3) &= \alpha_2 + \alpha_3 + \alpha_4, \\
-(1/2)(-2\lambda_1 - \mu_2 + \mu_3) &= \alpha_1 + \alpha_2 + \alpha_3 + \alpha_4 + \alpha_5, \\
-(1/2)(\lambda_0 - \lambda_1 - 2\lambda_2 - \mu_1 + \mu_3) &= \alpha_1 + \alpha_2 + \alpha_3, \\
(1/2)(\lambda_0 - \lambda_1 - 2\lambda_2 + \mu_1 - \mu_3) &= \alpha_2, \\
-(1/2)(\lambda_0 - \lambda_1 + 2\lambda_2 - \mu_1 + \mu_3) &= \alpha_1 + \alpha_2, \\
(1/2)(\lambda_0 - \lambda_1 + 2\lambda_2 + \mu_1 - \mu_3) &= \alpha_2 + \alpha_3, \\
-(1/2)(\lambda_0 + \lambda_1 + 2\lambda_2 + \mu_1 - \mu_2) &= \alpha_4, \\
(1/2)(\lambda_0 + \lambda_1 + 2\lambda_2 - \mu_1 + \mu_2) &= \alpha_3 + \alpha_4 + \alpha_5, \\
-(1/2)(-\lambda_0 - \lambda_1 + 2\lambda_2 + \mu_1 - \mu_2) &= \alpha_4 + \alpha_5, \\
(1/2)(-\lambda_0 - \lambda_1 + 2\lambda_2 - \mu_1 + \mu_2) &= \alpha_3 + \alpha_4 \\
-(\mu_1 + (2/3)\mu) &= \alpha_3 + 2\alpha_4 + \alpha_5 + \alpha_6, \\
-(\mu_2 + (2/3)\mu) &= \alpha_6, \\
-(\mu_3 + (2/3)\mu) &= \alpha_1 + 2\alpha_2 + 2\alpha_3 + 2\alpha_4 + \alpha_5 + \alpha_6, \\
-(-\lambda_0 - (1/2)\mu_1 + (2/3)\mu) &= \alpha_2 + \alpha_3 + \alpha_4 + \alpha_5 + \alpha_6, \\
-(\lambda_0 - (1/2)\mu_1 + (2/3)\mu) &= \alpha_1 + \alpha_2 + \alpha_3 + \alpha_4 + \alpha_6, \\
-(-\lambda_1 - (1/2)\mu_1 + (2/3)\mu) &= \alpha_1 + \alpha_2 + \alpha_3 + \alpha_4 + \alpha_5 + \alpha_6, \\
-(\lambda_1 - (1/2)\mu_1 + (2/3)\mu) &= \alpha_2 + \alpha_3 + \alpha_4 + \alpha_6, \\
-((-1/2)(-\lambda_0 + \lambda_1 + 2\lambda_2) - (1/2)\mu_2 + (2/3)\mu) &= \alpha_1 + \alpha_2 + 2\alpha_3 + 2\alpha_4 + \alpha_5 + \alpha_6, \\
-((1/2)(-\lambda_0 + \lambda_1 + 2\lambda_2) - (1/2)\mu_2 + (2/3)\mu) &= \alpha_2 + \alpha_3 + 2\alpha_4 + \alpha_5 + \alpha_6, \\
-((-1/2)(-\lambda_0 + \lambda_1 - 2\lambda_2) - (1/2)\mu_2 + (2/3)\mu) &= \alpha_1\alpha_2 + \alpha_3 + 2\alpha_4 + \alpha_5 + \alpha_6, \\
-((1/2)(-\lambda_0 + \lambda_1 - 2\lambda_2) - (1/2)\mu_2 + (2/3)\mu) &= \alpha_2 + 2\alpha_3 + 2\alpha_4 + \alpha_5 + \alpha_6, \\
-((-1/2)(-\lambda_0 - \lambda_1 - 2\lambda_2) - (1/2)\mu_3 + (2/3)\mu) &= \alpha_4 + \alpha_6, \\
-((1/2)(-\lambda_0 - \lambda_1 - 2\lambda_2) - (1/2)\mu_3 + (2/3)\mu) &= \alpha_3 + \alpha_4 + \alpha_5 + \alpha_6, \\
-((-1/2)(\lambda_0 + \lambda_1 - 2\lambda_2) - (1/2)\mu_3 + (2/3)\mu) &= \alpha_4 + \alpha_5 + \alpha_6, \\
-((1/2)(\lambda_0 + \lambda_1 - 2\lambda_2) - (1/2)\mu_3 + (2/3)\mu) &= \alpha_3 + \alpha_4 + \alpha_6.
\end{aligned}$$

Hence Π is a fundamental root system of $(\mathfrak{e}_7^C)^{\varepsilon_1, \varepsilon_2}$.

Then, for $\Phi, \Phi' \in \mathfrak{e}_7^C$, the Killing form B_7 of \mathfrak{e}_7^C is given by

$$B_7(\Phi, \Phi') = B_7(\Phi(\phi, A, B, \nu), \Phi(\phi, A, B, \nu)) = \frac{3}{2}B_6(\phi, \phi') + 36(A, B') + 36(A', B) + 24\nu\nu'$$

([9, Theorem 4.5.2]), so that for $\Phi_7 := \Phi(\phi, 0, 0, \mu), \Phi_7' := \Phi(\phi', 0, 0, \mu') \in \mathfrak{h}_7$, we have

$$\begin{aligned} B_7(\Phi_7, \Phi_7') &= \frac{3}{2}B_6(\phi, \phi') + 24\mu\mu' \\ &= \frac{3}{2}(24(\lambda_0\lambda_0' + \lambda_1\lambda_1' + 2\lambda_2\lambda_2') + 12(\mu_1\mu_1' + \mu_2\mu_2' + \mu_3\mu_3')) + 24\mu\mu' \\ &= 36(\lambda_0\lambda_0' + \lambda_1\lambda_1' + 2\lambda_2\lambda_2') + 18(\mu_1\mu_1' + \mu_2\mu_2' + \mu_3\mu_3') + 24\mu\mu', \end{aligned}$$

where $\phi := \lambda_0(iG_{01}) + \lambda_1(iG_{23}) + \lambda_2(i(G_{45} + G_{67})) + (\mu_1E_1 + \mu_2E_2 + \mu_3E_3)^\sim$, $\phi' = \lambda_0'(iG_{01}) + \lambda_1'(iG_{23}) + \lambda_2'(i(G_{45} + G_{67})) + (\mu_1'E_1 + \mu_2'E_2 + \mu_3'E_3)^\sim$. Now, the canonical elements $\Phi_{\alpha_1}, \Phi_{\alpha_2}, \Phi_{\alpha_3}, \Phi_{\alpha_4}, \Phi_{\alpha_5}, \Phi_{\alpha_6}$ corresponding to $\alpha_1, \alpha_2, \alpha_3, \alpha_4, \alpha_5, \alpha_6$ are determined as follows:

$$\begin{aligned} \Phi_{\alpha_1} &= \Phi\left(-\frac{1}{36}(iG_{01}) + \frac{1}{36}(iG_{23}), 0, 0, 0\right), \\ \Phi_{\alpha_2} &= \Phi\left(\frac{1}{72}(iG_{01}) - \frac{1}{72}(iG_{23}) - \frac{1}{72}(i(G_{45} + G_{67})) + \left(\frac{1}{36}E_1 - \frac{1}{36}E_3\right)^\sim, 0, 0, 0\right), \\ \Phi_{\alpha_3} &= \Phi\left(\frac{1}{36}(i(G_{45} + G_{67})), 0, 0, 0\right), \\ \Phi_{\alpha_4} &= \Phi\left(-\frac{1}{72}(iG_{01}) - \frac{1}{72}(iG_{23}) - \frac{1}{72}(i(G_{45} + G_{67})) + \left(-\frac{1}{36}E_1 + \frac{1}{36}E_2\right)^\sim, 0, 0, 0\right), \\ \Phi_{\alpha_5} &= \Phi\left(\frac{1}{36}(iG_{01}) + \frac{1}{36}(iG_{23}), 0, 0, 0\right), \\ \Phi_{\alpha_6} &= \Phi\left(\frac{1}{54}(E_1 - 2E_2 + E_3)^\sim, 0, 0, \frac{1}{36}\right). \end{aligned}$$

Hence we have the following

$$\begin{aligned} (\alpha_1, \alpha_1) &= B_7(\Phi_{\alpha_1}, \Phi_{\alpha_1}) = 36 \left(\left(-\frac{1}{36}\right)^2 + \left(\frac{1}{36}\right)^2 \right) = \frac{1}{18}, \\ (\alpha_1, \alpha_2) &= B_7(\Phi_{\alpha_1}, \Phi_{\alpha_2}) = 36 \left(\left(-\frac{1}{36}\right) \left(\frac{1}{72}\right) + \left(\frac{1}{36}\right) \left(-\frac{1}{72}\right) \right) = -\frac{1}{36}, \\ (\alpha_1, \alpha_3) &= B_7(\Phi_{\alpha_1}, \Phi_{\alpha_3}) = 0, \\ (\alpha_1, \alpha_4) &= B_7(\Phi_{\alpha_1}, \Phi_{\alpha_4}) = 36 \left(\left(-\frac{1}{36}\right) \left(-\frac{1}{72}\right) + \left(\frac{1}{36}\right) \left(-\frac{1}{72}\right) \right) = 0, \\ (\alpha_1, \alpha_5) &= B_7(\Phi_{\alpha_1}, \Phi_{\alpha_5}) = 36 \left(\left(-\frac{1}{36}\right) \left(\frac{1}{36}\right) + \left(\frac{1}{36}\right)^2 \right) = 0, \\ (\alpha_1, \alpha_6) &= B_7(\Phi_{\alpha_1}, \Phi_{\alpha_6}) = 0, \\ (\alpha_2, \alpha_2) &= B_7(\Phi_{\alpha_2}, \Phi_{\alpha_2}) = 36 \left(\left(\frac{1}{72}\right)^2 + \left(-\frac{1}{72}\right)^2 + 2 \left(-\frac{1}{72}\right)^2 \right) + 18 \left(\left(\frac{1}{36}\right)^2 + \left(-\frac{1}{36}\right)^2 \right) = \frac{1}{18}, \\ (\alpha_2, \alpha_3) &= B_7(\Phi_{\alpha_2}, \Phi_{\alpha_3}) = 36 \cdot 2 \left(-\frac{1}{72} \right) \left(\frac{1}{36} \right) = -\frac{1}{36}, \\ (\alpha_2, \alpha_4) &= B_6(\Phi_{\alpha_2}, \Phi_{\alpha_4}) = 36 \left(\left(\frac{1}{72}\right) \left(-\frac{1}{72}\right) + \left(-\frac{1}{72}\right)^2 + 2 \left(-\frac{1}{72}\right)^2 \right) + 18 \left(\frac{1}{36} \right) \left(-\frac{1}{36} \right) = 0, \\ (\alpha_2, \alpha_5) &= B_6(\Phi_{\alpha_2}, \Phi_{\alpha_5}) = 36 \left(\left(\frac{1}{72}\right) \left(\frac{1}{36}\right) + \left(-\frac{1}{72}\right) \left(\frac{1}{36}\right) \right) = 0, \end{aligned}$$

$$\begin{aligned}
(\alpha_2, \alpha_6) &= B_7(\Phi_{\alpha_2}, \Phi_{\alpha_6}) = 18 \left(\left(\frac{1}{36} \right) \left(\frac{1}{54} \right) + \left(-\frac{1}{36} \right) \left(\frac{1}{54} \right) \right) = 0, \\
(\alpha_3, \alpha_3) &= B_7(\Phi_{\alpha_3}, \Phi_{\alpha_3}) = 36 \cdot 2 \left(\frac{1}{36} \right)^2 = \frac{1}{18}, \\
(\alpha_3, \alpha_4) &= B_7(\Phi_{\alpha_3}, \Phi_{\alpha_4}) = 36 \cdot 2 \left(\frac{1}{36} \right) \left(-\frac{1}{72} \right) = -\frac{1}{36}, \\
(\alpha_3, \alpha_5) &= B_7(\Phi_{\alpha_3}, \Phi_{\alpha_5}) = 0, \\
(\alpha_3, \alpha_6) &= B_7(\Phi_{\alpha_3}, \Phi_{\alpha_6}) = 0, \\
(\alpha_4, \alpha_4) &= B_7(\Phi_{\alpha_4}, \Phi_{\alpha_4}) = 36 \left(\left(-\frac{1}{72} \right)^2 + \left(-\frac{1}{72} \right)^2 + 2 \left(-\frac{1}{72} \right)^2 \right) + 18 \left(\left(-\frac{1}{36} \right)^2 + \left(\frac{1}{36} \right)^2 \right) = \frac{1}{18}, \\
(\alpha_4, \alpha_5) &= B_7(\Phi_{\alpha_4}, \Phi_{\alpha_5}) = 36 \left(\left(-\frac{1}{72} \right) \left(\frac{1}{36} \right) + \left(-\frac{1}{72} \right) \left(\frac{1}{36} \right) \right) = -\frac{1}{36}, \\
(\alpha_4, \alpha_6) &= B_7(\Phi_{\alpha_4}, \Phi_{\alpha_6}) = 18 \left(\left(-\frac{1}{36} \right) \left(\frac{1}{54} \right) + \left(\frac{1}{36} \right) \left(-\frac{2}{54} \right) \right) = -\frac{1}{36}, \\
(\alpha_5, \alpha_5) &= B_7(\Phi_{\alpha_5}, \Phi_{\alpha_5}) = 36 \left(\left(\frac{1}{36} \right)^2 + \left(\frac{1}{36} \right)^2 \right) = \frac{1}{18}, \\
(\alpha_5, \alpha_6) &= B_7(\Phi_{\alpha_5}, \Phi_{\alpha_6}) = 0, \\
(\alpha_6, \alpha_6) &= B_7(\Phi_{\alpha_6}, \Phi_{\alpha_6}) = 18 \left(\left(\frac{1}{54} \right)^2 + \left(-\frac{2}{54} \right)^2 + \left(\frac{1}{54} \right)^2 \right) + 24 \left(\frac{1}{36} \right)^2 = \frac{1}{18}.
\end{aligned}$$

Thus, using the inner products above, we have

$$\begin{aligned}
\cos \theta_{12} &= \frac{(\alpha_1, \alpha_2)}{\sqrt{(\alpha_1, \alpha_1)(\alpha_2, \alpha_2)}} = -\frac{1}{2}, & \cos \theta_{13} &= \frac{(\alpha_1, \alpha_3)}{\sqrt{(\alpha_1, \alpha_1)(\alpha_3, \alpha_3)}} = 0, \\
\cos \theta_{14} &= \frac{(\alpha_1, \alpha_4)}{\sqrt{(\alpha_1, \alpha_1)(\alpha_4, \alpha_4)}} = 0, & \cos \theta_{15} &= \frac{(\alpha_1, \alpha_5)}{\sqrt{(\alpha_1, \alpha_1)(\alpha_5, \alpha_5)}} = 0, \\
\cos \theta_{16} &= \frac{(\alpha_1, \alpha_6)}{\sqrt{(\alpha_1, \alpha_1)(\alpha_6, \alpha_6)}} = 0, & \cos \theta_{23} &= \frac{(\alpha_2, \alpha_3)}{\sqrt{(\alpha_2, \alpha_2)(\alpha_3, \alpha_3)}} = -\frac{1}{2}, \\
\cos \theta_{24} &= \frac{(\alpha_2, \alpha_4)}{\sqrt{(\alpha_2, \alpha_2)(\alpha_4, \alpha_4)}} = 0, & \cos \theta_{25} &= \frac{(\alpha_2, \alpha_5)}{\sqrt{(\alpha_2, \alpha_2)(\alpha_5, \alpha_5)}} = 0, \\
\cos \theta_{26} &= \frac{(\alpha_2, \alpha_6)}{\sqrt{(\alpha_2, \alpha_2)(\alpha_6, \alpha_6)}} = 0, & \cos \theta_{34} &= \frac{(\alpha_3, \alpha_4)}{\sqrt{(\alpha_3, \alpha_3)(\alpha_4, \alpha_4)}} = -\frac{1}{2}, \\
\cos \theta_{35} &= \frac{(\alpha_3, \alpha_5)}{\sqrt{(\alpha_3, \alpha_3)(\alpha_5, \alpha_5)}} = 0, & \cos \theta_{36} &= \frac{(\alpha_3, \alpha_6)}{\sqrt{(\alpha_3, \alpha_3)(\alpha_6, \alpha_6)}} = 0, \\
\cos \theta_{45} &= \frac{(\alpha_4, \alpha_5)}{\sqrt{(\alpha_4, \alpha_4)(\alpha_5, \alpha_5)}} = -\frac{1}{2}, & \cos \theta_{46} &= \frac{(\alpha_4, \alpha_6)}{\sqrt{(\alpha_4, \alpha_4)(\alpha_6, \alpha_6)}} = -\frac{1}{2}, \\
\cos \theta_{56} &= \frac{(\alpha_5, \alpha_6)}{\sqrt{(\alpha_5, \alpha_5)(\alpha_6, \alpha_6)}} = 0.
\end{aligned}$$

so that we can draw the required Dynkin diagram. \square

7. THE GROUP $E_{8,\mathbf{H}}$ AND THE ROOT SYSTEM, THE DYNKIN DIAGRAM OF THE LIE ALGEBRA $(\mathfrak{e}_8^C)^{\varepsilon_1, \varepsilon_2}$

We consider the following complex Lie algebra $(\mathfrak{e}_{8,\mathbf{H}})^C$ which is given by replacing \mathfrak{C} with \mathbf{H} in the complex Lie algebra \mathfrak{e}_8^C :

$$\begin{aligned} (\mathfrak{e}_{8,\mathbf{H}})^C &= (\mathfrak{e}_{7,\mathbf{H}})^C \oplus (\mathfrak{P}_{\mathbf{H}})^C \oplus (\mathfrak{P}_{\mathbf{H}})^C \oplus C \oplus C \oplus C \\ &= \left\{ R = (\Phi, P, Q, r, s, t) \mid \begin{array}{l} \Phi \in (\mathfrak{e}_{7,\mathbf{H}})^C, P, Q \in (\mathfrak{P}_{\mathbf{H}})^C, \\ r, s, t \in C \end{array} \right\} \end{aligned}$$

In particular, we have $\dim_C((\mathfrak{e}_{8,\mathbf{H}})^C) = 66 + 32 \times 2 + 3 = 133$. In $(\mathfrak{e}_{8,\mathbf{H}})^C$, we can define a Lie bracket and can prove its simplicity as in \mathfrak{e}_8^C . In addition, we define a symmetric inner product $(R_1, R_2)_8$ by

$$(R_1, R_2)_8 := (\Phi_1, \Phi_2)_7 - \{Q_1, P_2\} + \{P_1, Q_2\} - 8r_1r_2 - 4t_1s_2 - 4s_1t_2,$$

where $R_i := (\Phi_i, P_i, Q_i, r_i, s_i, t_i) \in (\mathfrak{e}_{8,\mathbf{H}})^C, i = 1, 2$. Then $(\mathfrak{e}_{8,\mathbf{H}})^C$ leaves the symmetric inner product $(R_1, R_2)_8$ invariant (cf. [9, Lemma 5.3.1]).

Then we have the following theorem.

Theorem 7.1. *The Killing form $B_{8,\mathbf{H}}$ of $(\mathfrak{e}_{8,\mathbf{H}})^C$ is given by*

$$\begin{aligned} B_{8,\mathbf{H}}(R_1, R_2) &= -9(R_1, R_2)_8 \\ &= -9(\Phi_1, \Phi_2)_7 + 9\{Q_1, P_2\} - 9\{P_1, Q_2\} + 72r_1r_2 + 36t_1s_2 + 36s_1t_2 \\ &= \frac{9}{5}B_{7,\mathbf{H}}(\Phi_1, \Phi_2) + 9\{Q_1, P_2\} - 9\{P_1, Q_2\} + 72r_1r_2 + 36t_1s_2 + 36s_1t_2. \end{aligned}$$

Proof. Since $(\mathfrak{e}_{8,\mathbf{H}})^C$ is simple, there exists $k \in C$ such that

$$B_{8,\mathbf{H}}(R_1, R_2) = k(R_1, R_2)_8, \quad R_i \in (\mathfrak{e}_{8,\mathbf{H}})^C.$$

We will determine k . Let $R_1 = R_2 = (0, 0, 0, 1, 0, 0) = \tilde{1}$. Then we have

$$(\tilde{1}, \tilde{1})_8 = -8.$$

On the other hand, it follows from

$$\begin{aligned} (\text{ad } \tilde{1})(\text{ad } \tilde{1})R &= [\tilde{1}, [\tilde{1}, (\Phi, P, Q, r, s, t)]], \quad R \in (\mathfrak{e}_{8,\mathbf{H}})^C \\ &= [\tilde{1}, (0, P, -Q, 0, 2s, -2t)] \\ &= (0, P, Q, 0, 4s, 4t) \end{aligned}$$

that

$$B_{8,\mathbf{H}}(\tilde{1}, \tilde{1}) = \text{tr}((\text{ad } \tilde{1})(\text{ad } \tilde{1})) = 32 \times 2 + 4 \times 2 = 72.$$

Thus we have $k = -9$.

Therefore we have

$$B_{8,\mathbf{H}}(R_1, R_2) = -9(R_1, R_2)_8$$

and together with Theorem 6.1, the remainders of formulas can be easily obtained. \square

Now, we consider a Lie group $E_{8,\mathbf{H}}$ which is defined by

$$E_{8,\mathbf{H}} = \{ \alpha \in \text{Iso}_C((\mathfrak{e}_{8,\mathbf{H}})^C) \mid \alpha[R_1, R_2] = [\alpha R_1, \alpha R_2], \langle \alpha R_1, \alpha R_2 \rangle = \langle R_1, R_2 \rangle \},$$

where $\langle R_1, R_2 \rangle := (-1/15)B_{8,\mathbf{H}}((\tau\lambda_\omega)R_1, R_2)$. The group $E_{8,\mathbf{H}}$ is a compact Lie group as a closed subgroup of the unitary group $U(133) = U((\mathfrak{e}_{8,\mathbf{H}})^C) = \{ \alpha \in \text{Iso}_C((\mathfrak{e}_{8,\mathbf{H}})^C) \mid \langle \alpha R_1, \alpha R_2 \rangle = \langle R_1, R_2 \rangle \}$. Moreover, $E_{8,\mathbf{H}}$ is nothing but the group which is replaced \mathfrak{C} with \mathbf{H} in the definition of the compact Lie group of type E_8 (Subsection 3.5). The main aim of this section is to determine the structure of the group $E_{8,\mathbf{H}}$.

We give the explicit form of the Lie algebra $\mathfrak{e}_{8,\mathbf{H}}$ of the group $E_{8,\mathbf{H}}$ as lemma.

Lemma 7.2. *The Lie algebra $\mathfrak{e}_{8,\mathbf{H}}$ of the group $E_{8,\mathbf{H}}$ is given by*

$$\mathfrak{e}_{8,\mathbf{H}} = \{R = (\Phi, P, -\tau\lambda P, r, s, -\tau s) \mid \Phi \in \mathfrak{e}_{7,\mathbf{H}}, P \in (\mathfrak{P}_{\mathbf{H}})^C, r \in i\mathbf{R}, s \in C\}.$$

In particular, we have $\dim(\mathfrak{e}_{8,\mathbf{H}}) = 66 + 32 \times 2 + 1 + 2 = 133$.

Proof. We can confirm $((\mathfrak{e}_{8,\mathbf{H}})^C)^{\tau\lambda\omega} = \mathfrak{e}_{8,\mathbf{H}}$ as that in $(\mathfrak{e}_8^C)^{\tau\lambda\omega} = \mathfrak{e}_8$. For $R = (\Phi, P, Q, r, s, t) \in (\mathfrak{e}_{8,\mathbf{H}})^C$, we have the required result by getting R satisfying $(\tau\lambda\omega)R = R$. \square

First, we will study the complexification $(E_{8,\mathbf{H}})^C$ of $E_{8,\mathbf{H}}$:

$$\begin{aligned} (E_{8,\mathbf{H}})^C &= \text{Aut}((\mathfrak{e}_{8,\mathbf{H}})^C) \\ &= \{\alpha \in \text{Iso}_C((\mathfrak{e}_{8,\mathbf{H}})^C) \mid \alpha[R_1, R_2] = [\alpha R_1, \alpha R_2]\}. \end{aligned}$$

The immediate aim is to prove the connectedness of the group $(E_{8,\mathbf{H}})^C$. In order to prove it, we will use the manner used in [4].

First, we consider a subgroup $((E_{8,\mathbf{H}})^C)_{\bar{1},1^-,1_-}$ of $(E_{8,\mathbf{H}})^C$:

$$((E_{8,\mathbf{H}})^C)_{\bar{1},1^-,1_-} = \{\alpha \in (E_{8,\mathbf{H}})^C \mid \alpha\bar{1} = \bar{1}, \alpha 1^- = 1^-, \alpha 1_- = 1_-\}.$$

Then we have the following proposition.

Proposition 7.3. *The group $((E_{8,\mathbf{H}})^C)_{\bar{1},1^-,1_-}$ is isomorphic to the group $(E_{7,\mathbf{H}})^C$: $((E_{8,\mathbf{H}})^C)_{\bar{1},1^-,1_-} \cong (E_{7,\mathbf{R}})^C$.*

Proof. This proposition can be proved as in the proof of [9, Proposition 5.7.1] by replacing \mathfrak{C} with \mathbf{H} . \square

Subsequently, we consider a subgroup $((E_{8,\mathbf{H}})^C)_{1_-}$ of $(E_{8,\mathbf{H}})^C$:

$$((E_{8,\mathbf{H}})^C)_{1_-} = \{\alpha \in (E_{8,\mathbf{H}})^C \mid \alpha 1_- = 1_-\}.$$

Then we prove the following lemma.

Lemma 7.4. *The Lie algebra $((\mathfrak{e}_{8,\mathbf{H}})^C)_{1_-}$ of the group $((E_{8,\mathbf{H}})^C)_{1_-}$ is given by*

$$\begin{aligned} ((\mathfrak{e}_{8,\mathbf{H}})^C)_{1_-} &= \{R \in (\mathfrak{e}_{8,\mathbf{H}})^C \mid [R, 1_-] = 0\} \\ &= \{(\Phi, 0, Q, 0, 0, t) \in (\mathfrak{e}_{8,\mathbf{H}})^C \mid \Phi \in (\mathfrak{e}_{7,\mathbf{H}})^C, Q \in (\mathfrak{P}_{\mathbf{H}})^C, t \in C\}. \end{aligned}$$

In particular, we have $\dim_C(((\mathfrak{e}_{8,\mathbf{H}})^C)_{1_-}) = 66 + 32 + 1 = 99$.

Proof. By the straightforward computation, we can easily obtain the required results. \square

We prove the following proposition needed in the proof of connectedness.

Proposition 7.5. *The group $((E_{8,\mathbf{H}})^C)_{1_-}$ is a semi-direct product of groups $\exp(\text{ad}(((\mathfrak{P}_{\mathbf{H}})^C)_- \oplus C_-))$ and $(E_{7,\mathbf{H}})^C$:*

$$((E_{8,\mathbf{H}})^C)_{1_-} = \exp(\text{ad}(((\mathfrak{P}_{\mathbf{H}})^C)_- \oplus C_-)) \rtimes (E_{7,\mathbf{H}})^C.$$

In particular, the group $((E_{8,\mathbf{H}})^C)_{1_-}$ is connected.

Proof. Let $((\mathfrak{P}_{\mathbf{H}})^C)_- \oplus C_- = \{(0, 0, L, 0, 0, v) \mid L \in (\mathfrak{P}_{\mathbf{H}})^C, v \in C\}$ be a Lie subalgebra of the Lie algebra $((\mathfrak{e}_{8,\mathbf{H}})^C)_{1_-}$ (Lemma 7.4). Since it follows from $[L_-, v_-] = 0$ that $\text{ad}(L_-)$ commutes with $\text{ad}(v_-)$, we have $\exp(\text{ad}(L_- + v_-)) = \exp(\text{ad}(L_-)) \exp(\text{ad}(v_-))$. Hence the group $\exp(\text{ad}(((\mathfrak{P}_{\mathbf{H}})^C)_- \oplus C_-))$ is the connected subgroup of the group $((E_{8,\mathbf{H}})^C)_{1_-}$.

Now, let $\alpha \in ((E_{8,\mathbf{H}})^C)_{1_-}$ and set $\alpha\bar{1} := (\Phi_1, P_1, Q_1, r_1, s_1, t_1), \alpha 1^- := (\Phi_2, P_2, Q_2, r_2, s_2, t_2)$. Then, it follows from

$$[\alpha\bar{1}, 1_-] = \alpha[\bar{1}, 1_-] = -2\alpha 1_- = -21_- = (0, 0, 0, 0, 0, -2),$$

$$[\alpha\tilde{1}, 1_-] = [(\Phi_1, P_1, Q_1, r_1, s_1, t_1), (0, 0, 0, 0, 0, 1)] = (0, 0, -P_1, s_1, 0, -2r_1)$$

that $P_1 = 0, r_1 = 1, s_1 = 0$. Similarly, from $[\alpha 1^-, 1_-] = \alpha[1^-, 1_-] = \alpha\tilde{1}$, we have $\Phi_1 = 0, P_2 = -Q_1, s_2 = r_1, s_1 = 0, t_1 = -2r_2$. Hence we have

$$\alpha\tilde{1} = (0, 0, Q_1, 1, 0, t_1), \quad \alpha 1^- = (\Phi_2, -Q_1, Q_2, -\frac{t_1}{2}, 1, t_2).$$

Moreover, from $[\alpha\tilde{1}, \alpha 1^-] = \alpha[\tilde{1}, 1^-] = 2\alpha 1^-$, we obtain the following

$$\Phi_2 = \frac{1}{2}Q_1 \times Q_1, \quad Q_2 = -\frac{t}{2}Q_1 - \frac{1}{3}\Phi_2 Q_1, \quad t_2 = -\frac{t_1^2}{4} - \frac{1}{16}\{Q_1, Q_2\},$$

that is,

$$\alpha 1^- = (\frac{1}{2}Q_1 \times Q_1, -Q_1, -\frac{t}{2}Q_1 - \frac{1}{6}(Q_1 \times Q_1)Q_1, -\frac{t_1}{2}, 1, -\frac{t_1^2}{4} + \frac{1}{96}\{Q_1, (Q_1 \times Q_1)Q_1\}).$$

On the other hand, for $\exp(\text{ad}((v/2)_-)) \exp(\text{ad}(L_-)) \in \exp(\text{ad}(((\mathfrak{P}_{\mathbf{H}})^C)_- \oplus C_-))$, we have

$$\begin{aligned} & \exp\left(\text{ad}\left(\left(\frac{v}{2}\right)_-\right)\right) \exp(\text{ad}(L_-)) 1^- \\ &= \left(\frac{1}{2}L \times L, -L, -\frac{t}{2}L - \frac{1}{6}(L \times L)L, -\frac{v}{2}, 1, -\frac{v^2}{4} + \frac{1}{96}\{L, (L \times L)L\}\right), \end{aligned}$$

so that set $\delta := \exp(\text{ad}((t_1/2)_-)) \exp(\text{ad}(Q_{1-}))$, then we see

$$(\delta^{-1}\alpha)1^- = 1^-.$$

In addition, we can confirm

$$(\delta^{-1}\alpha)\tilde{1} = \tilde{1}, \quad (\delta^{-1}\alpha)1_- = 1_-.$$

Indeed, it follows that

$$\begin{aligned} & \exp\left(\text{ad}\left(\left(\frac{v}{2}\right)_-\right)\right) \exp(\text{ad}(L_-))\tilde{1} = \exp\left(\text{ad}\left(\left(\frac{v}{2}\right)_-\right)\right)(0, 0, L, 1, 0, 0) = (0, 0, L, 1, 0, v), \\ & \exp\left(\text{ad}\left(\left(\frac{v}{2}\right)_-\right)\right) \exp(\text{ad}(L_-))1_- = \exp\left(\text{ad}\left(\left(\frac{v}{2}\right)_-\right)\right)1_- = 1_-, \end{aligned}$$

so that we have $(\delta^{-1}\alpha)\tilde{1} = \delta^{-1}(0, 0, P, 1, 0, t) = \tilde{1}$, $(\delta^{-1}\alpha)1_- = \delta^{-1}1_- = 1_-$.

Hence, $\delta^{-1}\alpha \in ((E_{8,\mathbf{H}})^C)_{\tilde{1}, 1^-, 1_-} \cong (E_{7,\mathbf{H}})^C$ follows from Proposition 7.3, so that we obtain

$$((E_{8,\mathbf{H}})^C)_{1_-} = \exp(\text{ad}(((\mathfrak{P}_{\mathbf{H}})^C)_- \oplus C_-))(E_{7,\mathbf{H}})^C.$$

Furthermore, for $\beta \in (E_{7,\mathbf{H}})^C$, it is easy to verify that

$$\beta(\exp(\text{ad}(L_-)))\beta^{-1} = \exp(\text{ad}(\beta L_-)), \quad \beta((\exp(\text{ad}(v_-)))\beta^{-1} = \exp(\text{ad}(v_-)).$$

Indeed, for $(\Phi, P, Q, r, s, t) \in (\mathfrak{e}_{8,\mathbf{H}})^C$, by doing simple computation, we have the following

$$\begin{aligned} \beta \text{ad}(L_-)\beta^{-1}(\Phi, P, Q, r, s, t) &= \beta[L_-, \beta^{-1}(\Phi, P, Q, r, s, t)] \\ &= [\beta L_-, \beta\beta^{-1}(\Phi, P, Q, r, s, t)] \quad (\beta \in (E_{7,\mathbf{H}})^C \subset (E_{8,\mathbf{H}})^C) \\ &= [\beta L_-, (\Phi, P, Q, r, s, t)] \\ &= \text{ad}(\beta L_-)(\Phi, P, Q, r, s, t), \end{aligned}$$

that is, $\beta \text{ad}(L_-)\beta^{-1} = \text{ad}(\beta L_-)$. It is clear that the latter formula holds.

Hence we have

$$\beta(\exp(\text{ad}(L_-)))\beta^{-1} = \beta\left(\sum_{n=0}^{\infty} \frac{1}{n!} \text{ad}(L_-)^n\right)\beta^{-1}$$

$$\begin{aligned}
&= \sum_{n=0}^{\infty} \frac{1}{n!} (\beta \text{ad}(L_-) \beta^{-1})^n \quad (\beta \text{ad}(L_-) \beta^{-1} = \text{ad}(\beta L_-)) \\
&= \sum_{n=0}^{\infty} \frac{1}{n!} (\text{ad}(\beta L_-))^n \\
&= \exp(\text{ad}(\beta L_-)),
\end{aligned}$$

that is, $\beta(\exp(\text{ad}(L_-)))\beta^{-1} = \exp(\text{ad}(\beta L_-))$. By the argument similar to above, we also see that $\beta((\exp(\text{ad}(v_-)))\beta^{-1} = \exp(\text{ad}(v_-))$. Thus, for $\alpha \in ((E_{8,\mathbf{H}})^C)_{1_-}$, it follows from $\alpha = \exp(N_- + u_-)\beta \in \exp(\text{ad}(((\mathfrak{P}_{\mathbf{H}})^C)_- \oplus C_-)) \oplus (E_{7,\mathbf{H}})^C = ((E_{8,\mathbf{H}})^C)_{1_-}$ that

$$\begin{aligned}
&\alpha \exp(\text{ad}(L_- + v_-))\alpha^{-1} \\
&= (\exp(N_- + u_-)\beta) \exp(\text{ad}(L_- + v_-)) (\exp(N_- + u_-)\beta)^{-1} \\
&= (\exp(\text{ad}(N_-)) \exp(\text{ad}(u_-))\beta) (\exp(\text{ad}(L_-)) \exp(\text{ad}(v_-))) (\exp(\text{ad}(N_-)) \exp(\text{ad}(u_-))\beta)^{-1} \\
&= (\exp(\text{ad}(N_-)) \exp(\text{ad}(u_-))\beta) (\exp(\text{ad}(L_-)) \exp(\text{ad}(v_-))) (\beta^{-1} \exp(\text{ad}(u_-))^{-1} \exp(\text{ad}(N_-))^{-1}) \\
&= (\exp(\text{ad}(N_-)) \exp(\text{ad}(u_-))) (\exp(\text{ad}(\beta L_-)) \exp(\text{ad}(v_-))) (\exp(\text{ad}(u_-))^{-1} \exp(\text{ad}(N_-))^{-1}) \\
&= (\exp(\text{ad}(N_-)) \exp(\text{ad}(\beta L_-)) \exp(\text{ad}(N_-))^{-1}) (\exp(\text{ad}(u_-)) \exp(\text{ad}(v_-)) \exp(\text{ad}(u_-))^{-1}) \\
&\in \exp(\text{ad}(((\mathfrak{P}_{\mathbf{H}})^C)_- \oplus C_-)) = \exp(\text{ad}(((\mathfrak{P}_{\mathbf{H}})^C)_- \oplus C_-)).
\end{aligned}$$

This shows that $\exp(\text{ad}(((\mathfrak{P}_{\mathbf{H}})^C)_- \oplus C_-)) = \exp(\text{ad}(((\mathfrak{P}_{\mathbf{H}})^C)_-)) \exp(\text{ad}(C_-))$ is a normal subgroup of the group $((E_{8,\mathbf{H}})^C)_{1_-}$. In addition, we have a split exact sequence

$$1 \rightarrow \exp(\text{ad}(((\mathfrak{P}_{\mathbf{H}})^C)_- \oplus C_-)) \xrightarrow{j} ((E_{8,\mathbf{H}})^C)_{1_-} \begin{array}{c} \xrightarrow{p} \\ \xleftarrow{s} \end{array} (E_{7,\mathbf{H}})^C \rightarrow 1.$$

Indeed, first we define a mapping j by $j(\delta) = \delta$. Then it is clear that j is a injective homomorphism, and subsequently, define a mapping p by $p(\alpha) := p(\delta\beta) = \beta$. Then, since it follows that

$$\begin{aligned}
p(\alpha_1\alpha_2) &= p((\delta_1\beta_1)(\delta_2\beta_2)) \\
&= p(\delta_1(\beta_1\delta_2\beta_1^{-1})\beta_1\beta_2) \quad (\beta_1\delta_2\beta_1^{-1} =: \delta_2' \in \exp(\text{ad}(((\mathfrak{P}_{\mathbf{H}})^C)_- \oplus C_-))) \\
&= p((\delta_1\delta_2')(\beta_1\beta_2)) \\
&= \beta_1\beta_2 \\
&= p(\alpha_1)p(\alpha_2),
\end{aligned}$$

p is a homomorphism. Subsequently, let $\beta \in (E_{7,\mathbf{H}})^C$. We choose $\delta \in \exp(\text{ad}(((\mathfrak{P}_{\mathbf{H}})^C)_- \oplus C_-))$, then there exists $\alpha \in ((E_{8,\mathbf{H}})^C)_{1_-}$ such that $\alpha = \delta\beta$. This implies that p is surjective. Finally, we define a mapping s by $s(\beta) = \beta$. Then it is clear that s is also a injective homomorphism and we have

$$ps(\beta) = p(\beta) = \beta,$$

that is, $ps = 1$. With above, the short sequence is a split exact sequence.

Hence the group $((E_{8,\mathbf{H}})^C)_{1_-}$ is a semi-direct product of $\exp(\text{ad}(((\mathfrak{P}_{\mathbf{H}})^C)_- \oplus C_-))$ and $(E_{7,\mathbf{H}})^C$:

$$((E_{8,\mathbf{H}})^C)_{1_-} = \exp(\text{ad}(((\mathfrak{P}_{\mathbf{H}})^C)_- \oplus C_-)) \rtimes (E_{7,\mathbf{H}})^C.$$

Finally, the connectedness of $((E_{8,\mathbf{H}})^C)_{1_-}$ follows from the connectedness of $\exp(\text{ad}(((\mathfrak{P}_{\mathbf{H}})^C)_- \oplus C_-))$ and $(E_{7,\mathbf{H}})^C$ (Theorem 6.8). \square

Before proving the connectedness of the group $(E_{8,\mathbf{H}})^C$, we will make some preparations. For $R \in (\mathfrak{e}_{8,\mathbf{H}})^C$, we define a C -linear mapping $R \times R : (\mathfrak{e}_{8,\mathbf{H}})^C \rightarrow (\mathfrak{e}_{8,\mathbf{H}})^C$ by

$$(R \times R)R_1 = [R, [R, R_1]] + \frac{1}{18}B_{8,\mathbf{H}}(R, R_1)R, \quad R_1 \in (\mathfrak{e}_{8,\mathbf{H}})^C,$$

where $B_{8,\mathbf{H}}$ is the Killing form of $(\mathfrak{e}_{8,\mathbf{H}})^C$ (Theorem 7.1). Using this mapping, we define a space $\mathfrak{W}_{\mathbf{H}}$ by

$$(\mathfrak{W}_{\mathbf{H}})^C = \{R \in (\mathfrak{e}_{8,\mathbf{H}})^C \mid R \times R = 0, R \neq 0\}.$$

Lemma 7.6. *For $R = (\Phi, P, Q, r, s, t) \in (\mathfrak{e}_{8,\mathbf{H}})^C$, $R \neq 0$ belongs to $(\mathfrak{e}_{8,\mathbf{H}})^C$ if and only if R satisfies the following conditions*

- (1) $2s\Phi - P \times P = 0$
- (2) $2t\Phi + Q \times Q = 0$
- (3) $2r\Phi + P \times Q = 0$
- (4) $\Phi P - 3rP - 3sQ = 0$
- (5) $\Phi Q + 3rQ - 3tP = 0$
- (6) $\{P, Q\} - 16(st + r^2) = 0$
- (7) $2(\Phi P \times Q_1 + 2P \times \Phi Q_1 - rP \times Q_1 - sQ \times Q_1) - \{P, Q_1\}\Phi = 0$
- (8) $2(\Phi Q \times P_1 + 2Q \times \Phi P_1 + rQ \times P_1 - tP \times P_1) - \{Q, P_1\}\Phi = 0$
- (9) $8((P \times Q_1)Q - stQ_1 - r^2Q_1 - \Phi^2Q_1 + 2r\Phi Q_1) + 5\{P, Q_1\}Q - 2\{Q, Q_1\}P = 0$
- (10) $8((Q \times P_1)P + stP_1 + r^2P_1 + \Phi^2P_1 + 2r\Phi P_1) + 5\{Q, P_1\}P - 2\{P, P_1\}Q = 0$
- (11) $10(\text{ad } \Phi)^2\Phi_1 + Q \times \Phi_1P - P \times \Phi_1Q + B_{7,\mathbf{H}}(\Phi, \Phi_1)\Phi = 0$
- (12) $10(\Phi_1\Phi P - 2\Phi\Phi_1P - r\Phi_1P - s\Phi_1Q) + B_{7,\mathbf{H}}(\Phi, \Phi_1)P = 0$
- (13) $10(\Phi_1\Phi Q - 2\Phi\Phi_1Q + r\Phi_1Q - t\Phi_1P) + B_{7,\mathbf{H}}(\Phi, \Phi_1)Q = 0,$

for all $\Phi_1 \in (\mathfrak{e}_{7,\mathbf{H}})^C, P_1, Q_1 \in (\mathfrak{F}_{\mathbf{H}})^C$, where $B_{7,\mathbf{H}}$ is the Killing form of the Lie algebra $(\mathfrak{e}_{7,\mathbf{H}})^C$.

Proof. Note that $(\Phi, P \times Q)_7 = \{\Phi P, Q\}$ holds for $\Phi \in (\mathfrak{e}_{7,\mathbf{H}})^C, P, Q \in (\mathfrak{F}_{\mathbf{H}})^C$. For $R = (\Phi, P, Q, r, s, t) \in (\mathfrak{e}_{8,\mathbf{H}})^C$, by doing simple computation of $(R \times R)R_1 = 0$ for all $R_1 = (\Phi_1, P_1, Q_1, r_1, s_1, t_1) \in (\mathfrak{e}_{8,\mathbf{H}})^C$, we have the required relational formulas above. \square

Proposition 7.7. *The group $((E_{8,\mathbf{H}})^C)_0$ acts on $(\mathfrak{W}_{\mathbf{H}})^C$ transitively.*

Proof. Since $\alpha \in (E_{8,\mathbf{H}})^C$ leaves the Killing form $B_{8,\mathbf{H}}$ invariant : $B_{8,\mathbf{H}}(\alpha R, \alpha R') = B_{8,\mathbf{H}}(R, R')$, $R, R' \in (\mathfrak{e}_{8,\mathbf{H}})^C$, the group $(E_{8,\mathbf{H}})^C$ acts on $(\mathfrak{W}_{\mathbf{H}})^C$. Indeed, let $R \in (\mathfrak{W}_{\mathbf{H}})^C$. Then, for all $R_1 \in (\mathfrak{e}_{8,\mathbf{H}})^C$, it follows that

$$\begin{aligned} (\alpha R \times \alpha R)R_1 &= [\alpha R, [\alpha R, \alpha R_1]] + (1/18)B_{8,\mathbf{H}}(\alpha R, R_1)\alpha R \\ &= \alpha[[R, [R, \alpha^{-1}R_1]] + (1/18)B_{8,\mathbf{H}}(R, \alpha^{-1}R_1)\alpha R \\ &= \alpha((R \times R)\alpha^{-1}R_1 \\ &= 0. \end{aligned}$$

Hence the group $(E_{8,\mathbf{H}})^C$ acts on $(\mathfrak{W}_{\mathbf{H}})^C$, that is, $\alpha R \in (E_{8,\mathbf{H}})^C$. We will show that this action is transitive. First, for all $R_1 \in (\mathfrak{e}_{8,\mathbf{H}})^C$, it follows from

$$\begin{aligned} (1_- \times 1_-)R_1 &= [1_-, [1_-, (\Phi_1, P_1, Q_1, r_1, s_1, t_1)]] + (1/18)B_{8,\mathbf{H}}(1_-, R_1)1_- \\ &= [1_-, (0, 0, P_1, -s_1, 0, 2r_1)] + 2s_11_- \\ &= (0, 0, 0, 0, -2s_1) + 2s_11_- \\ &= 0 \end{aligned}$$

that $1_- \in (\mathfrak{W}_{\mathbf{H}})^C$. Then, any element $R \in (\mathfrak{W}_{\mathbf{H}})^C$ can be transformed to $1_- \in (\mathfrak{W}_{\mathbf{H}})^C$ by some $\alpha \in ((E_{8,\mathbf{H}})^C)_0$. We will prove this below.

Case (i) where $R = (\Phi, P, Q, r, s, t), t \neq 0$.

From Lemma 7.6 (2),(5) and (6), we see

$$\Phi = -\frac{1}{2t}Q \times Q, \quad P = \frac{r}{t}Q - \frac{1}{6t^2}(Q \times Q)Q, \quad s = -\frac{r^2}{t} + \frac{1}{96t^3}\{Q, (Q \times Q)Q\}.$$

Now, let $\Theta := \text{ad}(0, P_1, 0, r_1, s_1, 0) \in \text{ad}((\mathfrak{e}_{8, \mathbf{H}})^C)$, $r_1 \neq 0$, then we compute $\Theta^n 1_-$:

$$\begin{aligned} & \Theta^n 1_- \\ &= \begin{pmatrix} ((-2)^{n-1} + (-1)^n)r_1^{n-2}P_1 \times P_1 \\ \left((-2)^{n-1} - \frac{1 + (-1)^{n-1}}{2} \right) r_1^{n-2}s_1P_1 + \left(\frac{1 - (-2)^n}{6} + \frac{(-1)^n}{2} \right) r_1^{n-3}(P_1 \times P_1)P_1 \\ ((-2)^n + (-1)^{n-1})r_1^{n-1}P_1 \\ (-2)^{n-1}r_1^{n-1}s_1 \\ -\left((-2)^{n-2} + 2^{n-2} \right) r_1^{n-2}s_1^2 + \frac{2^{n-2} + (-2)^{n-2} - (-1)^{n-1} - 1}{24} r_1^{n-4}\{P_1, (P_1 \times P_1)P_1\} \\ (-2)^n r_1^n \end{pmatrix}. \end{aligned}$$

Hence, by doing straightforward computation, we have

$$\begin{aligned} & (\exp \Theta) 1_- = \left(\sum_{n=0}^{\infty} \frac{1}{n!} \Theta^n \right) 1_- \\ &= \begin{pmatrix} -\frac{1}{2r_1^2}(e^{-2r_1} - 2e^{-r_1} + 1)P_1 \times P_1 \\ \frac{s_1}{2r_1^2}(-e^{-2r_1} - e^{r_1} + e^{-r_1} + 1)P_1 + \frac{1}{6r_1^3}(-e^{-2r_1} + e^{r_1} + 3e^{-r_1} - 3)(P_1 \times P_1)P_1 \\ \frac{1}{r_1}(e^{-2r_1} - e^{-r_1})P_1 \\ \frac{s_1}{2r_1}(1 - e^{-2r_1}) \\ -\frac{s_1^2}{4r_1^2}(e^{-2r_1} + e^{2r_1} - 2) + \frac{1}{96r_1^4}(e^{2r_1} + e^{-2r_1} - 4e^{r_1} - 4e^{-r_1} + 6)\{P_1, (P_1 \times P_1)P_1\} \\ e^{-2r_1} \end{pmatrix}. \end{aligned}$$

In the case where $t \neq 1$. For a given $R = (\Phi, P, Q, r, s, t)$, $t \neq 0$, we can choose $P_1 \in (\mathfrak{B}_{\mathbf{H}})^C$, $r_1, s_1 \in C$ such that

$$\frac{1}{r_1}(e^{-2r_1} - e^{-r_1})P_1 = Q, \quad \frac{s_1}{2r_1}(1 - e^{-2r_1}) = r, \quad e^{-2r_1} = t.$$

Indeed, we choose some one value of \sqrt{t} satisfying $(\sqrt{t})^2 = t$ and some one value of $\log t$ for $t \in C$, respectively. Then because of $t - t^2 \neq 0$, $t - 1 \neq 0$, we can get

$$P_1 = \frac{(\sqrt{t} + t) \log t}{2(t - t^2)}Q, \quad r_1 = -\frac{\log t}{2}, \quad s_1 = \frac{\log t}{t - 1}r.$$

Hence, by using these P_1, r_1, s_1 , we obtain

$$(\exp \Theta) 1_- = \begin{pmatrix} -\frac{1}{2t}Q \times Q \\ \frac{r}{t}Q - \frac{1}{6t^2}(Q \times Q)Q \\ Q \\ r \\ -\frac{r^2}{t} + \frac{1}{96t^3}\{Q, (Q \times Q)Q\} \\ t \end{pmatrix} = R.$$

In the case where $t = 1$. Then, a given $R = (\Phi, P, Q, r, s, t), t \neq 0$ is of the form

$$\Phi = -\frac{1}{2}Q \times Q, \quad P = rQ - \frac{1}{6}(Q \times Q)Q, \quad s = -r^2 + \frac{1}{96}\{Q, (Q \times Q)Q\},$$

so let $\Theta := \text{ad}(0, P_1, 0, 0, s_1, 0) \in \text{ad}((\mathfrak{e}_{8,\mathbf{H}})^C)$, we have

$$(\exp \Theta)1_- = \left(\sum_{n=0}^{\infty} \frac{1}{n!} \Theta^n \right) 1_- = \begin{pmatrix} -\frac{1}{2}P_1 \times P_1 \\ -s_1P_1 + \frac{1}{6}(P_1 \times P_1)P_1 \\ -P_1 \\ s_1 \\ -s_1^2 + \frac{1}{96}\{P_1, (P_1 \times P_1)P_1\} \\ 1 \end{pmatrix}.$$

Here, as in the case $t \neq 1$, we choose $P_1 \in (\mathfrak{J}_{\mathbf{H}})^C, s_1 \in C$ such that $-P_1 = Q, s_1 = r$. Hence, by using these P_1, s_1 , we obtain

$$(\exp \Theta)1_- = \begin{pmatrix} -\frac{1}{2}Q \times Q \\ rQ - \frac{1}{6}(Q \times Q)Q \\ Q \\ r \\ -r^2 + \frac{1}{96}\{Q, (Q \times Q)Q\} \\ 1 \end{pmatrix} = R.$$

Thus R is transformed to 1_- by $(\exp \Theta)^{-1} \in ((E_{8,\mathbf{H}})^C)_0$.

Case (ii) where $R = (\Phi, P, Q, r, s, 0), s \neq 0$.

Let $\Theta := \text{ad}(0, 0, 0, 0, \pi/2, -\pi/2) \in \text{ad}((\mathfrak{e}_{8,\mathbf{H}})^C)$. Then we have

$$(\exp \Theta)R = (\Phi, Q, -P, -r, 0, -s), \quad -s \neq 0.$$

Hence this case is reduced to Case (i).

Case (iii) where $R = (\Phi, P, Q, r, 0, 0), r \neq 0$.

Again, from Lemma 7.6 (2),(5) and (6), we have

$$Q \times Q = 0, \quad \Phi Q = -3rQ, \quad \{P, Q\} = 16r^2.$$

Then, let $\Theta := \text{ad}(0, Q, 0, 0, 0, 0) \in \text{ad}((\mathfrak{e}_{8,\mathbf{H}})^C)$, we have

$$(\exp \Theta)R = (\Phi, P + 2rQ, Q, r, -4r^2, 0), \quad -4r^2 \neq 0.$$

Hence this case is reduced to Case (ii).

Case (iv) where $R = (\Phi, P, Q, 0, 0, 0), Q \neq 0$.

We can choose $P_1 \in (\mathfrak{P}_{\mathbf{H}})^C$ such that $\{P_1, Q\} \neq 0$. Indeed, $\{P_1, Q\} = 0$ for all $P_1 \in (\mathfrak{P}_{\mathbf{H}})^C$ implies $Q = 0$, so that there exists $P_1 \in (\mathfrak{P}_{\mathbf{H}})^C$ such that $\{P_1, Q\} \neq 0$.

Now, let $\Theta := \text{ad}(0, P_1, 0, 0, 0, 0) \in \text{ad}((\mathfrak{e}_{8, \mathbf{H}})^C)$, we have

$$(\exp \Theta)R = \begin{pmatrix} \Phi + P_1 \times Q \\ P - \Phi P_1 + \frac{1}{2}(P_1 \times Q)P_1 + \frac{1}{16}\{P_1, Q\}P_1 \\ Q \\ -\frac{1}{8}\{P_1, Q\} \\ \frac{1}{4}\{P_1, P\} + \frac{1}{8}\{P_1, -\Phi P_1\} + \frac{1}{24}\{P_1, (P_1 \times Q)P_1\} \\ 0 \end{pmatrix}, \quad -\frac{1}{8}\{P_1, Q\} \neq 0.$$

Hence this case is reduced to Case (iii).

Case (v) where $R = (\Phi, P, 0, 0, 0, 0)$, $P \neq 0$.

As in Case (iv), we choose $Q_1 \in (\mathfrak{B}_{\mathbf{H}})^C$ such that $\{P, Q_1\} \neq 0$. Then, let $\Theta := \text{ad}(0, 0, Q_1, 0, 0, 0) \in \text{ad}((\mathfrak{e}_{8, \mathbf{H}})^C)$, we have

$$(\exp \Theta)R = \begin{pmatrix} \Phi - P \times Q_1 \\ P \\ -\Phi Q_1 + \frac{1}{2}(P \times Q_1)Q_1 + \frac{1}{16}\{P, Q_1\}Q_1 \\ \frac{1}{8}\{P, Q_1\} \\ 0 \\ -\frac{1}{8}\{Q_1, -\Phi Q_1\} - \frac{1}{24}\{Q_1, -(P \times Q_1)Q_1\} \end{pmatrix}, \quad \frac{1}{8}\{P, Q_1\} \neq 0.$$

Hence this case is also reduced to Case (iii).

Case (vi) where $R = (\Phi, 0, 0, 0, 0, 0)$, $\Phi \neq 0$. We can choose $P_1 \in (\mathfrak{B}_{\mathbf{H}})^C$ such that $\Phi P_1 \neq 0$. Indeed, $\Phi P_1 = 0$ for all $P_1 \in (\mathfrak{B}_{\mathbf{H}})^C$ implies $\Phi = 0$, so that there exists $P_1 \in (\mathfrak{B}_{\mathbf{H}})^C$ such that $\Phi P_1 \neq 0$. Then, let $\Theta := \text{ad}(0, P_1, 0, 0, 0, 0) \in \text{ad}((\mathfrak{e}_{8, \mathbf{H}})^C)$, we have

$$(\exp \text{ad}(0, P_1, 0, 0, 0, 0))R = \left(\Phi, -\Phi P_1, 0, 0, \frac{1}{8}\{\Phi P_1, P_1\}, 0 \right), \quad -\Phi P_1 \neq 0.$$

Hence this case is reduced to Case (v).

With above, the proof of this proposition is completed. \square

Now, we will prove the theorem as the immediate aim .

Theorem 7.8. *The homogeneous space $(E_{8, \mathbf{H}})^C / ((E_{8, \mathbf{H}})^C)_{1-}$ is homeomorphic to the space $(\mathfrak{B}_{\mathbf{H}})^C$: $(E_{8, \mathbf{H}})^C / ((E_{8, \mathbf{H}})^C)_{1-} \simeq (\mathfrak{B}_{\mathbf{H}})^C$.*

In particular, the group $(E_{8, \mathbf{H}})^C$ is connected.

Proof. Since the group $(E_{8, \mathbf{H}})^C$ acts on the space $(\mathfrak{B}_{\mathbf{H}})^C$ transitively (Proposition 7.7), the former half of this theorem is proved. The latter half was shown as follows. The connectedness of the group $(E_{8, \mathbf{H}})^C$ follows from the connectedness of the group $((E_{8, \mathbf{H}})^C)_{1-}$ and the space $(\mathfrak{B}_{\mathbf{H}})^C = ((E_{8, \mathbf{H}})^C)_{01-}$ (Propositions 7.5, 7.7). \square

In order to give a polar decomposition of the group $(E_{8, \mathbf{H}})^C$, we prove the following lemma.

Lemma 7.9. *The group $(E_{8, \mathbf{H}})^C$ is an algebraic subgroup of the general linear group $GL(133, \mathbb{C}) = \text{Iso}_{\mathbb{C}}((\mathfrak{t}_{8, \mathbf{H}})^C)$ and satisfies the condition that $\alpha \in (E_{8, \mathbf{H}})^C$ implies $\alpha^* = (\tau \lambda_{\omega}) \alpha^{-1} (\lambda_{\omega} \tau) \in (E_{8, \mathbf{H}})^C$, where α^* is the transpose of α with respect to the Hermite inner product $\langle R_1, R_2 \rangle$: $\langle \alpha^* R_1, R_2 \rangle = \langle R_1, \alpha R_2 \rangle$.*

Proof. First, it is easy to verify that $\alpha^* = (\tau\lambda_\omega)\alpha^{-1}(\lambda_\omega\tau)$. Indeed, it follows from

$$\begin{aligned} \langle \alpha^* R_1, R_2 \rangle &= \langle R_1, \alpha R_2 \rangle = -\frac{1}{15} B_{8,\mathbf{H}}((\tau\lambda_\omega)R_1, \alpha R_2) \\ &= -\frac{1}{15} B_{8,\mathbf{H}}(\alpha(\alpha^{-1}(\tau\lambda_\omega)R_1), \alpha R_2) \\ &= -\frac{1}{15} B_{8,\mathbf{H}}(\alpha^{-1}(\tau\lambda_\omega)R_1, R_2) \\ &= -\frac{1}{15} B_{8,\mathbf{H}}((\tau\lambda_\omega)((\lambda_\omega\tau)\alpha^{-1}(\tau\lambda_\omega)R_1), R_2) \\ &= \langle (\lambda_\omega\tau)\alpha^{-1}(\tau\lambda_\omega)R_1, R_2 \rangle \end{aligned}$$

that $\alpha^* = (\lambda_\omega\tau)\alpha^{-1}(\tau\lambda_\omega)$, that is, $\alpha^* = (\tau\lambda_\omega)\alpha^{-1}(\lambda_\omega\tau)$.

Subsequently, we will show $\alpha^* \in (E_{8,\mathbf{H}})^C$. It is clear that $\alpha^* \in \text{Iso}_C((\mathfrak{e}_{8,\mathbf{H}})^C)$ and note that $[\tau R_1, \tau R_2] = \tau[R_1, R_2]$, $R_i \in (\mathfrak{e}_{8,\mathbf{H}})^C$ and $\lambda_\omega \in (E_{8,\mathbf{H}})^C$, it follows that

$$\begin{aligned} [\alpha^* R_1, \alpha^* R_2] &= [(\tau\lambda_\omega)\alpha^{-1}(\lambda_\omega\tau)R_1, (\tau\lambda_\omega)\alpha^{-1}(\lambda_\omega\tau)R_2] \\ &= \tau[\lambda_\omega\alpha^{-1}(\lambda_\omega\tau)R_1, \lambda_\omega\alpha^{-1}(\lambda_\omega\tau)R_2] \\ &= (\tau\lambda_\omega)[\alpha^{-1}(\lambda_\omega\tau)R_1, \alpha^{-1}(\lambda_\omega\tau)R_2] \\ &= (\tau\lambda_\omega)\alpha^{-1}[(\lambda_\omega\tau)R_1, (\lambda_\omega\tau)R_2] \\ &= (\tau\lambda_\omega)\alpha^{-1}(\lambda_\omega\tau)[R_1, R_2] \\ &= \alpha^*[R_1, R_2]. \end{aligned}$$

Hence we have $\alpha^* \in (E_{8,\mathbf{H}})^C$.

Finally, the group $(E_{8,\mathbf{H}})^C$ is defined by the algebraic relation $\alpha[R_1, R_2] = [\alpha R_1, \alpha R_2]$, so that $(E_{8,\mathbf{H}})^C$ is algebraic. \square

Let $O((\mathfrak{e}_{8,\mathbf{H}})^C)$ be the orthogonal subgroup $GL(133, C) = \text{Iso}_C((\mathfrak{e}_{8,\mathbf{H}})^C)$:

$$O(133, C) = O((\mathfrak{e}_{8,\mathbf{H}})^C) = \{\alpha \in \text{Iso}_C((\mathfrak{e}_{8,\mathbf{H}})^C) \mid \langle \alpha R_1, \alpha R_2 \rangle = \langle R_1, R_2 \rangle\}.$$

Then we have the following

$$\begin{aligned} ((E_{8,\mathbf{H}})^C \cap O((\mathfrak{e}_{8,\mathbf{H}})^C)) &= \left\{ \alpha \in \text{Iso}_C((\mathfrak{e}_{8,\mathbf{H}})^C) \mid \begin{array}{l} \alpha[R_1, R_2] = [\alpha R_1, \alpha R_2], \\ \langle \alpha R_1, \alpha R_2 \rangle = \langle R_1, R_2 \rangle \end{array} \right\} \\ &= E_{8,\mathbf{H}}. \end{aligned}$$

Using Chevally's lemma([1, Lemma 2]), from Lemma 7.9 we have a homeomorphism

$$\begin{aligned} (E_{8,\mathbf{H}})^C &\simeq ((E_{8,\mathbf{H}})^C \cap O((\mathfrak{e}_{8,\mathbf{H}})^C)) \times \mathbf{R}^d \\ &= E_{8,\mathbf{H}} \times \mathbf{R}^d, \end{aligned}$$

where the dimension d of the Euclidian part is computed from Lemma 7.2 as follows:

$$d = \dim((E_{8,\mathbf{H}})^C) - \dim(E_{8,\mathbf{H}}) = 133 \times 2 - 133 = 133.$$

With above, we have the following theorem.

Theorem 7.10. *The group $(E_{8,\mathbf{H}})^C$ is homeomorphic to the topological product of the group $E_{8,\mathbf{H}}$ and a 133-dimensional Euclidian space \mathbf{R}^{133} :*

$$(E_{8,\mathbf{H}})^C \simeq E_{8,\mathbf{H}} \times \mathbf{R}^{133}.$$

In particular, the group $E_{8,\mathbf{H}}$ is connected.

Proof. The former half has been already proved above. As for the latter half, the connectedness of the group $E_{8,\mathbf{H}}$ follows from the connectedness of the group $(E_{8,\mathbf{H}})^C$ (Theorem 7.8) \square

Next, we move on to study a group $(E_8^C)^{\varepsilon_1, \varepsilon_2}$ defined below.
We define a subgroup $(E_8^C)^{\varepsilon_1, \varepsilon_2}$ of E_8 by

$$(E_8^C)^{\varepsilon_1, \varepsilon_2} := \{\alpha \in E_8 \mid \varepsilon_1 \alpha = \alpha \varepsilon_1, \varepsilon_2 \alpha = \alpha \varepsilon_2\}.$$

The immediate aim is to prove the connectedness of the group $(E_8^C)^{\varepsilon_1, \varepsilon_2}$ by the same procedure as the proof of the connectedness of the group $(E_{8, \mathbf{H}})^C$.

First, we consider a subgroup $((E_8^C)^{\varepsilon_1, \varepsilon_2})_{\tilde{1}, 1^-, 1_-}$ of $(E_8^C)^{\varepsilon_1, \varepsilon_2}$:

$$((E_8^C)^{\varepsilon_1, \varepsilon_2})_{\tilde{1}, 1^-, 1_-} = \{\alpha \in (E_8^C)^{\varepsilon_1, \varepsilon_2} \mid \alpha \tilde{1} = \tilde{1}, \alpha 1^- = 1^-, \alpha 1_- = 1_-\}.$$

Then we have the following proposition.

Proposition 7.11. *The group $((E_8^C)^{\varepsilon_1, \varepsilon_2})_{\tilde{1}, 1^-, 1_-}$ is isomorphic to the group $(E_7^C)^{\varepsilon_1, \varepsilon_2}$:
 $((E_8^C)^{\varepsilon_1, \varepsilon_2})_{\tilde{1}, 1^-, 1_-} \cong (E_7^C)^{\varepsilon_1, \varepsilon_2}$.*

Proof. Since $\varepsilon_i, i = 1, 2$ satisfies $\varepsilon_i \tilde{1} = \tilde{1}, \varepsilon_i 1^- = 1^-, \varepsilon_i 1_- = 1_-$, ε_i induces an automorphism of the group $(E_8^C)_{\tilde{1}, 1^-, 1_-}$. Hence, note that $(E_8^C)_{\tilde{1}, 1^-, 1_-} \cong E_7^C$ ([9, Proposition 5.7.1]), we have

$$((E_8^C)^{\varepsilon_1, \varepsilon_2})_{\tilde{1}, 1^-, 1_-} = ((E_8^C)_{\tilde{1}, 1^-, 1_-})^{\varepsilon_1, \varepsilon_2} \cong (E_7^C)^{\varepsilon_1, \varepsilon_2}.$$

□

Subsequently, we consider a subgroup $((E_8^C)^{\varepsilon_1, \varepsilon_2})_{1_-}$ of $(E_8^C)^{\varepsilon_1, \varepsilon_2}$:

$$((E_8^C)^{\varepsilon_1, \varepsilon_2})_{1_-} = \{\alpha \in (E_8^C)^{\varepsilon_1, \varepsilon_2} \mid \alpha 1_- = 1_-\}.$$

Then we prove the following lemma.

Lemma 7.12. (1) *The Lie algebra $(\mathfrak{e}_8^C)^{\varepsilon_1, \varepsilon_2}$ of the group $(E_8^C)^{\varepsilon_1, \varepsilon_2}$ is given by*

$$(\mathfrak{e}_8^C)^{\varepsilon_1, \varepsilon_2} = \{R = (\Phi, P, Q, r, s, t) \mid \Phi \in (\mathfrak{e}_7^C)^{\varepsilon_1, \varepsilon_2}, P, Q \in (\mathfrak{P}_{\mathbf{H}})^C, r, s, t \in C\}.$$

In particular, we have $\dim_{\mathbb{C}}((\mathfrak{e}_8^C)^{\varepsilon_1, \varepsilon_2}) = 66 + 32 \times 2 + 3 = 133$.

(2) *The Lie algebra $((\mathfrak{e}_8^C)^{\varepsilon_1, \varepsilon_2})_{1_-}$ of the group $((E_8^C)^{\varepsilon_1, \varepsilon_2})_{1_-}$ is given by*

$$\begin{aligned} ((\mathfrak{e}_8^C)^{\varepsilon_1, \varepsilon_2})_{1_-} &= \{R \in (\mathfrak{e}_8^C)^{\varepsilon_1, \varepsilon_2} \mid [R, 1_-] = 0\} \\ &= \{(\Phi, 0, Q, 0, 0, t) \in (\mathfrak{e}_8^C)^{\varepsilon_1, \varepsilon_2} \mid \Phi \in (\mathfrak{e}_7^C)^{\varepsilon_1, \varepsilon_2}, Q \in (\mathfrak{P}_{\mathbf{H}})^C, t \in C\}. \end{aligned}$$

In particular, we have $\dim_{\mathbb{C}}(((\mathfrak{e}_8^C)^{\varepsilon_1, \varepsilon_2})_{1_-}) = 66 + 32 + 1 = 99$.

Proof. By the straightforward computation, we can easily obtain the required results. □

We prove the following proposition needed in the proof of connectedness.

Proposition 7.13. *The group $((E_8^C)^{\varepsilon_1, \varepsilon_2})_{1_-}$ is a semi-direct product of groups $\exp(\text{ad}(((\mathfrak{P}_{\mathbf{H}})^C)_- \oplus C_-))$ and $(E_7^C)^{\varepsilon_1, \varepsilon_2}$:*

$$((E_8^C)^{\varepsilon_1, \varepsilon_2})_{1_-} = \exp(\text{ad}(((\mathfrak{P}_{\mathbf{H}})^C)_- \oplus C_-)) \rtimes (E_7^C)^{\varepsilon_1, \varepsilon_2}.$$

In particular, the group $((E_8^C)^{\varepsilon_1, \varepsilon_2})_{1_-}$ is connected.

Proof. Using Proposition 7.11 and Lemma 7.12, as in the proof of Proposition 7.5, this proposition is proved by replacing $(E_{8, \mathbf{H}})^C, (E_{7, \mathbf{H}})^C, (\mathfrak{e}_{8, \mathbf{H}})^C$ of the proof of Proposition 7.5 with $(E_8^C)^{\varepsilon_1, \varepsilon_2}, (E_7^C)^{\varepsilon_1, \varepsilon_2}, (\mathfrak{e}_8^C)^{\varepsilon_1, \varepsilon_2}$, respectively. Note that the connectedness of the group $(E_7^C)^{\varepsilon_1, \varepsilon_2}$ follows from $(E_7^C)^{\varepsilon_1, \varepsilon_2} \cong \text{Spin}(12, C)$ ([2, Proposition 1.1.7]).

□

Before proving the connectedness of the group $(E_8^C)^{\varepsilon_1, \varepsilon_2}$, we will make some preparations. For $R \in \mathfrak{e}_8^C$, we define a C -linear mapping $R \times R : \mathfrak{e}_8^C \rightarrow \mathfrak{e}_8^C$ by

$$(R \times R)R_1 = [R, [R, R_1]] + \frac{1}{30}B_8(R, R_1)R, \quad R_1 \in \mathfrak{e}_8^C,$$

where B_8 is the Killing form of \mathfrak{e}_8^C ([9, Theorem 5.3.2]). Using this mapping, we define a space $(\mathfrak{W}^C)_{\varepsilon_1, \varepsilon_2}$ by

$$(\mathfrak{W}^C)_{\varepsilon_1, \varepsilon_2} = \{R \in \mathfrak{e}_8^C \mid R \times R = 0, R \neq 0, \varepsilon_1 R = R, \varepsilon_2 R = R\},$$

where the action of $\varepsilon_i, i = 1, 2$ to \mathfrak{e}_8^C is given by

$$\varepsilon_i(\Phi, P, Q, r, s, t) = ((\text{ad} \varepsilon_i)\Phi, \varepsilon_i P, \varepsilon_i Q, r, s, t), \quad (\Phi, P, Q, r, s, t) \in \mathfrak{e}_8^C,$$

and note that P, Q -part of $R = (\Phi, P, Q, r, s, t) \in (\mathfrak{W}^C)_{\varepsilon_1, \varepsilon_2}$ belong to $(\mathfrak{P}_{\mathbf{H}})^C : P, Q \in (\mathfrak{P}_{\mathbf{H}})^C$.

Lemma 7.14. *For $R = (\Phi, P, Q, r, s, t) \in \mathfrak{e}_8^C$, $R \neq 0$ belongs to $(\mathfrak{W}^C)_{\varepsilon_1, \varepsilon_2}$ if and only if R satisfies the following conditions*

- (1) $2s\Phi - P \times P = 0$ (2) $2t\Phi + Q \times Q = 0$ (3) $2r\Phi + P \times Q = 0$
- (4) $\Phi P - 3rP - 3sQ = 0$ (5) $\Phi Q + 3rQ - 3tP = 0$ (6) $\{P, Q\} - 16(st + r^2) = 0$
- (7) $2(\Phi P \times Q_1 + 2P \times \Phi Q_1 - rP \times Q_1 - sQ \times Q_1) - \{P, Q_1\}\Phi = 0$
- (8) $2(\Phi Q \times P_1 + 2Q \times \Phi P_1 + rQ \times P_1 - tP \times P_1) - \{Q, P_1\}\Phi = 0$
- (9) $8((P \times Q_1)Q - stQ_1 - r^2Q_1 - \Phi^2Q_1 + 2r\Phi Q_1) + 5\{P, Q_1\}Q - 2\{Q, Q_1\}P = 0$
- (10) $8((Q \times P_1)P + stP_1 + r^2P_1 + \Phi^2P_1 + 2r\Phi P_1) + 5\{Q, P_1\}P - 2\{P, P_1\}Q = 0$
- (11) $18(\text{ad} \Phi)^2\Phi_1 + Q \times \Phi_1 P - P \times \Phi_1 Q + B_7(\Phi, \Phi_1)\Phi = 0$
- (12) $18(\Phi_1\Phi P - 2\Phi\Phi_1 P - r\Phi_1 P - s\Phi_1 Q) + B_7(\Phi, \Phi_1)P = 0$
- (13) $18(\Phi_1\Phi Q - 2\Phi\Phi_1 Q + r\Phi_1 Q - t\Phi_1 P) + B_7(\Phi, \Phi_1)Q = 0,$

for all $\Phi_1 \in (\mathfrak{e}_7^C)^{\varepsilon_1, \varepsilon_2} P_1, Q_1 \in (\mathfrak{P}_{\mathbf{H}})^C$, where B_7 is the Killing form of the Lie algebra \mathfrak{e}_7^C .

Proof. Note that $(\Phi, P \times Q)_7 = \{\Phi P, Q\}$ holds for $\Phi \in (\mathfrak{e}_7^C)^{\varepsilon_1, \varepsilon_2}, P, Q \in (\mathfrak{P}_{\mathbf{H}})^C$. For $R = (\Phi, P, Q, r, s, t) \in (\mathfrak{W}^C)_{\varepsilon_1, \varepsilon_2}$, by doing simple computation of $(R \times R)R_1 = 0$ for all $R_1 = (\Phi_1, P_1, Q_1, r_1, s_1, t_1) \in (\mathfrak{W}^C)_{\varepsilon_1, \varepsilon_2}$, we have the required relational formulas above. \square

Proposition 7.15. *The group $((E_8^C)^{\varepsilon_1, \varepsilon_2})_0$ acts on $(\mathfrak{W}^C)_{\varepsilon_1, \varepsilon_2}$ transitively.*

Proof. Using Lemmas 7.12 (1), 7.14, as in the proof of Proposition 7.7, this proposition is proved by replacing $(E_{8,\mathbf{H}})^C, (\mathfrak{e}_{8,\mathbf{H}})^C$ of the proof of Proposition 7.7 with $(E_8^C)^{\varepsilon_1, \varepsilon_2}, (\mathfrak{e}_8^C)^{\varepsilon_1, \varepsilon_2}$, respectively. \square

Now, we will prove the theorem as the immediate aim .

Theorem 7.16. *The homogeneous space $(E_8^C)^{\varepsilon_1, \varepsilon_2} / ((E_8^C)^{\varepsilon_1, \varepsilon_2})_{1-}$ is homeomorphic to the space $(\mathfrak{W}^C)_{\varepsilon_1, \varepsilon_2} : (E_8^C)^{\varepsilon_1, \varepsilon_2} / ((E_8^C)^{\varepsilon_1, \varepsilon_2})_{1-} \simeq (\mathfrak{W}^C)_{\varepsilon_1, \varepsilon_2}$.*

In particular, the group $(E_8^C)^{\varepsilon_1, \varepsilon_2}$ is connected.

Proof. Since the group $(E_8^C)^{\varepsilon_1, \varepsilon_2}$ acts on the space $(\mathfrak{W}^C)_{\varepsilon_1, \varepsilon_2}$ transitively (Proposition 7.15), the former half of this theorem is proved.

The latter half was shown as follows. The connectedness of the group $(E_8^C)^{\varepsilon_1, \varepsilon_2}$ follows from the connectedness of the group $((E_8^C)^{\varepsilon_1, \varepsilon_2})_{1-}$ and the space $(\mathfrak{W}^C)_{\varepsilon_1, \varepsilon_2} = ((E_8^C)^{\varepsilon_1, \varepsilon_2})_{01-}$ (Propositions 7.13, 7.15). \square

In order to determine the type of the group $(E_8^C)^{\varepsilon_1, \varepsilon_2}$ as Lie algebras, we move the determination of the root system and the Dynkin diagram of the Lie algebra $(\mathfrak{e}_8^C)^{\varepsilon_1, \varepsilon_2}$.

Here, we define a Lie subalgebra \mathfrak{h}_8 of $(\mathfrak{e}_8^C)^{\varepsilon_1, \varepsilon_2}$ by

$$\mathfrak{h}_8 := \left\{ R_8 = (\Phi, 0, 0, w, 0, 0) \left| \begin{array}{l} \Phi := \Phi(\phi, 0, 0, \mu) \in \mathfrak{h}_7, \\ \phi = \delta + (\mu_1 E_1 + \mu_2 E_2 + \mu_3 E_3)^\sim \\ \delta := (L_1, L_2, L_3), \\ L_1 = \lambda_0(iG_{01}) + \lambda_1(iG_{23}) + \lambda_2(i(G_{45} + G_{67})), \\ L_2 = \pi\kappa L_1, L_3 = \kappa\pi L_1, \lambda_i \in C, \\ \mu_k \in C, \mu_1 + \mu_2 + \mu_3 = 0, \mu \in C, \\ w \in C \end{array} \right. \right\}.$$

Then \mathfrak{h}_8 is a Cartan subalgebra of $(\mathfrak{e}_8^C)^{\varepsilon_1, \varepsilon_2}$. Indeed, it is clear that $(\mathfrak{e}_8^C)^{\varepsilon_1, \varepsilon_2}$ is abelian. Next, by doing straightforward computation, we can confirm that $[R, R_8] \in \mathfrak{h}_8, R \in (\mathfrak{e}_8^C)^{\varepsilon_1, \varepsilon_2}$ implies $R \in \mathfrak{h}_8$ for any $R_8 \in \mathfrak{h}_8$.

Theorem 7.17. *The rank of the Lie algebra $(\mathfrak{e}_8^C)^{\varepsilon_1, \varepsilon_2}$ is seven. The roots Δ of $(\mathfrak{e}_8^C)^{\varepsilon_1, \varepsilon_2}$ relative to \mathfrak{h}_8 are given by*

$$\Delta = \left\{ \begin{array}{l} \pm(\lambda_0 - \lambda_1), \pm(\lambda_0 + \lambda_1), \pm 2\lambda_2, \pm(1/2)(-2\lambda_0 + \mu_2 - \mu_3), \\ \pm(1/2)(-2\lambda_0 - \mu_2 + \mu_3), \pm(1/2)(-2\lambda_1 + \mu_2 - \mu_3), \pm(1/2)(-2\lambda_1 - \mu_2 + \mu_3), \\ \pm(1/2)(\lambda_0 - \lambda_1 - 2\lambda_2 - \mu_1 + \mu_3), \pm(1/2)(\lambda_0 - \lambda_1 - 2\lambda_2 + \mu_1 - \mu_3), \\ \pm(1/2)(\lambda_0 - \lambda_1 + 2\lambda_2 - \mu_1 + \mu_3), \pm(1/2)(\lambda_0 - \lambda_1 + 2\lambda_2 + \mu_1 - \mu_3), \\ \pm(1/2)(\lambda_0 + \lambda_1 + 2\lambda_2 + \mu_1 - \mu_2), \pm(1/2)(\lambda_0 + \lambda_1 + 2\lambda_2 - \mu_1 + \mu_2), \\ \pm(1/2)(-\lambda_0 - \lambda_1 + 2\lambda_2 + \mu_1 - \mu_2), \pm(1/2)(-\lambda_0 - \lambda_1 + 2\lambda_2 - \mu_1 + \mu_2), \\ \pm(\mu_1 + (2/3)\mu), \pm(\mu_2 + (2/3)\mu), \pm(\mu_3 + (2/3)\mu), \pm(-\lambda_0 - (1/2)\mu_1 + (2/3)\mu), \\ \pm(\lambda_0 - (1/2)\mu_1 + (2/3)\mu), \pm(-\lambda_1 - (1/2)\mu_1 + (2/3)\mu), \pm(\lambda_1 - (1/2)\mu_1 + (2/3)\mu), \\ \pm(-(1/2)(-\lambda_0 + \lambda_1 + 2\lambda_2) - (1/2)\mu_2 + (2/3)\mu), \\ \pm((1/2)(-\lambda_0 + \lambda_1 + 2\lambda_2) - (1/2)\mu_2 + (2/3)\mu), \\ \pm(-(1/2)(-\lambda_0 + \lambda_1 - 2\lambda_2) - (1/2)\mu_2 + (2/3)\mu), \\ \pm((1/2)(-\lambda_0 + \lambda_1 - 2\lambda_2) - (1/2)\mu_2 + (2/3)\mu), \\ \pm(-(1/2)(-\lambda_0 - \lambda_1 - 2\lambda_2) - (1/2)\mu_3 + (2/3)\mu), \\ \pm((1/2)(-\lambda_0 - \lambda_1 - 2\lambda_2) - (1/2)\mu_3 + (2/3)\mu), \\ \pm(-(1/2)(\lambda_0 + \lambda_1 - 2\lambda_2) - (1/2)\mu_3 + (2/3)\mu), \\ \pm((1/2)(\lambda_0 + \lambda_1 - 2\lambda_2) - (1/2)\mu_3 + (2/3)\mu), \\ \pm(\mu_1 - (1/3)\mu + w), \pm(\mu_2 - (1/3)\mu + w), \pm(\mu_3 - (1/3)\mu + w), \\ \pm(-\lambda_0 - (1/2)\mu_1 - (1/3)\mu + w), \pm(\lambda_0 - (1/2)\mu_1 - (1/3)\mu + w), \\ \pm(-\lambda_1 - (1/2)\mu_1 - (1/3)\mu + w), \pm(\lambda_1 - (1/2)\mu_1 - (1/3)\mu + w), \\ \pm((-1/2)(-\lambda_0 + \lambda_1 + 2\lambda_2) - (1/2)\mu_2 - (1/3)\mu + w), \\ \pm((1/2)(-\lambda_0 + \lambda_1 + 2\lambda_2) - (1/2)\mu_2 - (1/3)\mu + w), \\ \pm((-1/2)(-\lambda_0 + \lambda_1 - 2\lambda_2) - (1/2)\mu_2 - (1/3)\mu + w), \\ \pm((1/2)(-\lambda_0 + \lambda_1 - 2\lambda_2) - (1/2)\mu_2 - (1/3)\mu + w), \\ \pm((-1/2)(-\lambda_0 - \lambda_1 - 2\lambda_2) - (1/2)\mu_3 - (1/3)v + w), \\ \pm((1/2)(-\lambda_0 - \lambda_1 - 2\lambda_2) - (1/2)\mu_3 - (1/3)v + w), \end{array} \right\}.$$

$$\Delta = \left\{ \begin{array}{l} \pm((-1/2)(\lambda_0 + \lambda_1 - 2\lambda_2) - (1/2)\mu_3 - (1/3)\nu + w), \\ \pm((1/2)(\lambda_0 + \lambda_1 - 2\lambda_2) - (1/2)\mu_3 - (1/3)\nu + w), \\ \pm(-\mu_1 + (1/3)\mu + w), \pm(-\mu_2 + (1/3)\mu + w), \pm(-\mu_3 + (1/3)\mu + w), \\ \pm(-\lambda_0 + (1/2)\mu_1 + (1/3)\mu + w), \pm(\lambda_0 + (1/2)\mu_1 + (1/3)\mu + w), \\ \pm(-\lambda_1 + (1/2)\mu_1 + (1/3)\mu + w), \pm(\lambda_1 + (1/2)\mu_1 + (1/3)\mu + w), \\ \pm((-1/2)(-\lambda_0 + \lambda_1 + 2\lambda_2) + (1/2)\mu_2 + (1/3)\mu + w), \\ \pm((1/2)(-\lambda_0 + \lambda_1 + 2\lambda_2) + (1/2)\mu_2 + (1/3)\mu + w), \\ \pm((-1/2)(-\lambda_0 + \lambda_1 - 2\lambda_2) + (1/2)\mu_2 + (1/3)\mu + w), \\ \pm((1/2)(-\lambda_0 + \lambda_1 - 2\lambda_2) + (1/2)\mu_2 + (1/3)\mu + w), \\ \pm((-1/2)(-\lambda_0 - \lambda_1 - 2\lambda_2) + (1/2)\mu_3 + (1/3)\nu + w), \\ \pm((1/2)(-\lambda_0 - \lambda_1 - 2\lambda_2) + (1/2)\mu_3 + (1/3)\nu + w), \\ \pm((-1/2)(\lambda_0 + \lambda_1 - 2\lambda_2) + (1/2)\mu_3 + (1/3)\nu + w), \\ \pm((1/2)(\lambda_0 + \lambda_1 - 2\lambda_2) + (1/2)\mu_3 + (1/3)\nu + w), \\ \pm(\mu + w), \pm(-\mu + w), \pm(2w) \end{array} \right.$$

Proof. The roots of $(\mathfrak{e}_7^C)^{\varepsilon_1, \varepsilon_2}$ are also the roots of $(\mathfrak{e}_8^C)^{\varepsilon_1, \varepsilon_2}$. Indeed, let the root α of $(\mathfrak{e}_7^C)^{\varepsilon_1, \varepsilon_2}$ and its associated root vector $\Phi_s \in (\mathfrak{e}_7^C)^{\varepsilon_1, \varepsilon_2} \subset (\mathfrak{e}_8^C)^{\varepsilon_1, \varepsilon_2}$. Then, for $R_8 \in \mathfrak{h}_8$, we have

$$\begin{aligned} [R_8, \Phi_s] &= [(\Phi, 0, 0, w, 0, 0)(\Phi_s, 0, 0, 0, 0, 0)] \\ &= ([\Phi, \Phi_s], 0, 0, 0, 0, 0) \\ &= (\alpha(\Phi)\Phi_s, 0, 0, 0, 0, 0) \\ &= \alpha(\Phi)(\Phi_s, 0, 0, 0, 0, 0) \\ &= \alpha(R_8)\Phi_s. \end{aligned}$$

We will determine the remainders of roots. We will show a few examples. First, let $R_8 = (\Phi, 0, 0, w, 0, 0) \in \mathfrak{h}_8$ and $R_{\dot{E}_1}^- := (0, \dot{E}_1, 0, 0, 0, 0) \in (\mathfrak{e}_8^C)^{\varepsilon_1, \varepsilon_2}$, where $\dot{E}_1 := (E_1, 0, 0, 0) \in (\mathfrak{P}_{\mathbf{H}})^C$. Then it follows that

$$\begin{aligned} [R_8, R_{\dot{E}_1}^-] &= [(\Phi, 0, 0, w, 0, 0), (0, \dot{E}_1, 0, 0, 0, 0)] \\ &= (0, \Phi\dot{E}_1 + r\dot{E}_1, 0, 0, 0, 0) \\ &= (0, \Phi(\phi, 0, 0, \mu)(E_1, 0, 0, 0) + w(E_1, 0, 0, 0), 0, 0, 0, 0) \\ &= (0, (\phi - (1/3)\mu + w)E_1, 0, 0, 0, 0), (\phi = \tilde{T}_0) \\ &= (0, ((\mu_1 - (1/3)\mu + w)E_1, 0, 0, 0), 0, 0, 0, 0) \\ &= (\mu_1 - (1/3)\mu + w)(0, \dot{E}_1, 0, 0, 0, 0) \\ &= (\mu_1 - (1/3)\mu + w)R_{\dot{E}_1}^-, \end{aligned}$$

that is, $[R_8, R_{\dot{E}_1}^-] = (\mu_1 - (1/3)\mu + w)R_{\dot{E}_1}^-$. Hence we see that $\mu_1 - (1/3)\mu + w$ is a root and $(0, \dot{E}_1, 0, 0, 0, 0)$ is an associated root vector. Next, let $R_{\dot{F}_1(1+ie_1)}^- := (0, \dot{F}_1(1+ie_1), 0, 0, 0, 0) \in (\mathfrak{e}_8^C)^{\varepsilon_1, \varepsilon_2}$, where $\dot{F}_1(1+ie_1) := (F_1(1+ie_1), 0, 0, 0) \in \mathfrak{P}^C$. Then it follows that

$$\begin{aligned} [R_8, R_{\dot{F}_1(1+ie_1)}^-] &= [(\Phi, 0, 0, w, 0, 0), (0, \dot{F}_1(1+ie_1), 0, 0, 0, 0)] \\ &= (0, \Phi\dot{F}_1(1+ie_1) + r\dot{F}_1(1+ie_1), 0, 0, 0, 0) \\ &= (0, \Phi(\phi, 0, 0, \mu)(F_1(1+ie_1), 0, 0, 0) + r(F_1(1+ie_1), 0, 0, 0), 0, 0, 0, 0) \\ &= (0, (\phi - (1/3)\mu + w)F_1(1+ie_1), 0, 0, 0, 0), (\phi = (L_1, L_2, L_3) + \tilde{T}_0) \\ &= (0, ((-\lambda_0 + (1/2)(\mu_2 + \mu_3) - (1/3)\mu + w)F_1(1+ie_1), 0, 0, 0), 0, 0, 0, 0) \end{aligned}$$

$$\begin{aligned}
&= (0, (-\lambda_0 - (1/2)\mu_1 - (1/3)\mu + w)\dot{F}_1(1 + ie_1), 0, 0, 0, 0) \\
&= (-\lambda_0 - (1/2)\mu_1 - (1/3)\mu + w)R^-_{\dot{F}_1(1+ie_1)},
\end{aligned}$$

that is, $[R_8, R^-_{\dot{F}_1(1+ie_1)}] = (-\lambda_0 - (1/2)\mu_1 - (1/3)\mu + w)R^-_{\dot{F}_1(1+ie_1)}$. Hence we see that $-\lambda_0 - (1/2)\mu_1 - (1/3)\mu + w$ is a root and $(0, \dot{F}_1(1 + ie_1), 0, 0, 0, 0)$ is its associated root vector. All of roots and these associated root vectors except ones of the Lie algebra $(\mathfrak{e}_7^C)^{\varepsilon_1, \varepsilon_2}$ above are obtained as follows:

roots	associated root vectors
$\mu_1 - (1/3)\mu + w$	$(0, \dot{E}_1, 0, 0, 0, 0)$
$-(\mu_1 - (1/3)\mu + w)$	$(0, 0, \dot{E}_1, 0, 0, 0)$
$\mu_2 - (1/3)\mu + w$	$(0, \dot{E}_2, 0, 0, 0, 0)$
$-(\mu_2 - (1/3)\mu + w)$	$(0, 0, \dot{E}_2, 0, 0, 0)$
$\mu_3 - (1/3)\mu + w$	$(0, \dot{E}_3, 0, 0, 0, 0)$
$-(\mu_3 - (1/3)\mu + w)$	$(0, 0, \dot{E}_3, 0, 0, 0)$
$-\lambda_0 - (1/2)\mu_1 - (1/3)\mu + w$	$(0, \dot{F}_1(1 + ie_1), 0, 0, 0, 0)$
$-(-\lambda_0 - (1/2)\mu_1 - (1/3)\mu + w)$	$(0, 0, \dot{F}_1(1 - ie_1), 0, 0, 0)$
$\lambda_0 - (1/2)\mu_1 - (1/3)\mu + w$	$(0, \dot{F}_1(1 - ie_1), 0, 0, 0, 0)$
$-(\lambda_0 - (1/2)\mu_1 - (1/3)\mu + w)$	$(0, 0, \dot{F}_1(1 + ie_1), 0, 0, 0)$
$-\lambda_1 - (1/2)\mu_1 - (1/3)\mu + w$	$(0, \dot{F}_1(e_2 + ie_3), 0, 0, 0, 0)$
$-(-\lambda_0 - (1/2)\mu_1 - (1/3)\mu + w)$	$(0, 0, \dot{F}_1(e_2 - ie_3), 0, 0, 0)$
$\lambda_1 - (1/2)\mu_1 - (1/3)\mu + w$	$(0, \dot{F}_1(e_2 - ie_3), 0, 0, 0, 0)$
$-(\lambda_1 - (1/2)\mu_1 - (1/3)\mu + w)$	$(0, 0, \dot{F}_1(e_2 + ie_3), 0, 0, 0)$
$(-1/2)(-\lambda_0 + \lambda_1 + 2\lambda_2) - (1/2)\mu_2 - (1/3)\mu + w$	$(0, \dot{F}_2(1 + ie_1), 0, 0, 0, 0)$
$-((-1/2)(-\lambda_0 + \lambda_1 + 2\lambda_2) - (1/2)\mu_2 - (1/3)\mu + w)$	$(0, 0, \dot{F}_2(1 - ie_1), 0, 0, 0)$
$(1/2)(-\lambda_0 + \lambda_1 + 2\lambda_2) - (1/2)\mu_2 - (1/3)\mu + w$	$(0, \dot{F}_2(1 - ie_1), 0, 0, 0, 0)$
$-((-1/2)(-\lambda_0 + \lambda_1 + 2\lambda_2) - (1/2)\mu_2 - (1/3)\mu + w)$	$(0, 0, \dot{F}_2(1 + ie_1), 0, 0, 0)$
$(-1/2)(-\lambda_0 + \lambda_1 - 2\lambda_2) - (1/2)\mu_2 - (1/3)\mu + w$	$(0, \dot{F}_2(e_2 + ie_3), 0, 0, 0, 0)$
$-((-1/2)(-\lambda_0 + \lambda_1 - 2\lambda_2) - (1/2)\mu_2 - (1/3)\mu + w)$	$(0, 0, \dot{F}_2(e_2 - ie_3), 0, 0, 0)$
$(1/2)(-\lambda_0 + \lambda_1 - 2\lambda_2) - (1/2)\mu_2 - (1/3)\mu + w$	$(0, \dot{F}_2(e_2 - ie_3), 0, 0, 0, 0)$
$-((-1/2)(-\lambda_0 + \lambda_1 - 2\lambda_2) - (1/2)\mu_2 - (1/3)\mu + w)$	$(0, 0, \dot{F}_2(e_2 + ie_3), 0, 0, 0)$
$(-1/2)(-\lambda_0 - \lambda_1 - 2\lambda_2) - (1/2)\mu_3 - (1/3)\nu + w$	$(0, \dot{F}_3(1 + ie_1), 0, 0, 0, 0)$
$-((-1/2)(-\lambda_0 - \lambda_1 - 2\lambda_2) - (1/2)\mu_3 - (1/3)\nu + w)$	$(0, 0, \dot{F}_3(1 - ie_1), 0, 0, 0)$
$(1/2)(-\lambda_0 - \lambda_1 - 2\lambda_2) - (1/2)\mu_3 - (1/3)\nu + w$	$(0, \dot{F}_3(1 - ie_1), 0, 0, 0, 0)$
$-((1/2)(-\lambda_0 - \lambda_1 - 2\lambda_2) - (1/2)\mu_3 - (1/3)\nu + w)$	$(0, 0, \dot{F}_3(1 + ie_1), 0, 0, 0)$
$(-1/2)(\lambda_0 + \lambda_1 - 2\lambda_2) - (1/2)\mu_3 - (1/3)\nu + w$	$(0, \dot{F}_3(e_2 + ie_3), 0, 0, 0, 0)$
$-((-1/2)(-\lambda_0 - \lambda_1 - 2\lambda_2) - (1/2)\mu_3 - (1/3)\nu + w)$	$(0, 0, \dot{F}_3(e_2 - ie_3), 0, 0, 0)$
$(1/2)(\lambda_0 + \lambda_1 - 2\lambda_2) - (1/2)\mu_3 - (1/3)\nu + w$	$(0, \dot{F}_3(e_2 - ie_3), 0, 0, 0, 0)$
$-((1/2)(\lambda_0 + \lambda_1 - 2\lambda_2) - (1/2)\mu_3 - (1/3)\nu + w)$	$(0, 0, \dot{F}_3(e_2 + ie_3), 0, 0, 0)$
$-\mu_1 + (1/3)\mu + w$	$(0, \dot{E}_1, 0, 0, 0, 0)$
$-(-\mu_1 + (1/3)\mu + w)$	$(0, 0, \dot{E}_1, 0, 0, 0)$
$-\mu_2 + (1/3)\mu + w$	$(0, \dot{E}_2, 0, 0, 0, 0)$
$-(-\mu_2 + (1/3)\mu + w)$	$(0, 0, \dot{E}_2, 0, 0, 0)$
$-\mu_3 + (1/3)\mu + w$	$(0, \dot{E}_3, 0, 0, 0, 0)$
$-(-\mu_3 + (1/3)\mu + w)$	$(0, 0, \dot{E}_3, 0, 0, 0)$

$$\begin{array}{ll}
-\lambda_0 + (1/2)\mu_1 + (1/3)\mu + w & (0, \dot{F}_1(1 + ie_1), 0, 0, 0, 0) \\
-(-\lambda_0 + (1/2)\mu_1 + (1/3)\mu + w) & (0, 0, \dot{F}_1(1 - ie_1), 0, 0, 0) \\
\lambda_0 + (1/2)\mu_1 + (1/3)\mu + w & (0, \dot{F}_1(1 - ie_1), 0, 0, 0, 0) \\
-(\lambda_0 + (1/2)\mu_1 + (1/3)\mu + w) & (0, 0, \dot{F}_1(1 + ie_1), 0, 0, 0) \\
-\lambda_1 + (1/2)\mu_1 + (1/3)\mu + w & (0, \dot{F}_1(e_2 + ie_3), 0, 0, 0, 0) \\
-(-\lambda_0 + (1/2)\mu_1 + (1/3)\mu + w) & (0, 0, \dot{F}_1(e_2 - ie_3), 0, 0, 0) \\
\lambda_1 + (1/2)\mu_1 + (1/3)\mu + w & (0, \dot{F}_1(e_2 - ie_3), 0, 0, 0, 0) \\
-(\lambda_1 + (1/2)\mu_1 + (1/3)\mu + w) & (0, 0, \dot{F}_1(e_2 + ie_3), 0, 0, 0) \\
(-1/2)(-\lambda_0 + \lambda_1 + 2\lambda_2) + (1/2)\mu_2 + (1/3)\mu + w & (0, \dot{F}_2(1 + ie_1), 0, 0, 0, 0) \\
-((-1/2)(-\lambda_0 + \lambda_1 + 2\lambda_2) + (1/2)\mu_2 + (1/3)\mu + w) & (0, 0, \dot{F}_2(1 - ie_1), 0, 0, 0) \\
(1/2)(-\lambda_0 + \lambda_1 + 2\lambda_2) + (1/2)\mu_2 + (1/3)\mu + w & (0, \dot{F}_2(1 - ie_1), 0, 0, 0, 0) \\
-((1/2)(-\lambda_0 + \lambda_1 + 2\lambda_2) + (1/2)\mu_2 + (1/3)\mu + w) & (0, 0, \dot{F}_2(1 + ie_1), 0, 0, 0) \\
(-1/2)(-\lambda_0 + \lambda_1 - 2\lambda_2) + (1/2)\mu_2 + (1/3)\mu + w & (0, \dot{F}_2(e_2 + ie_3), 0, 0, 0, 0) \\
-((-1/2)(-\lambda_0 + \lambda_1 - 2\lambda_2) + (1/2)\mu_2 + (1/3)\mu + w) & (0, 0, \dot{F}_2(e_2 - ie_3), 0, 0, 0) \\
(1/2)(-\lambda_0 + \lambda_1 - 2\lambda_2) + (1/2)\mu_2 + (1/3)\mu + w & (0, \dot{F}_2(e_2 - ie_3), 0, 0, 0, 0) \\
-((1/2)(-\lambda_0 + \lambda_1 - 2\lambda_2) + (1/2)\mu_2 + (1/3)\mu + w) & (0, 0, \dot{F}_2(e_2 + ie_3), 0, 0, 0) \\
(-1/2)(-\lambda_0 - \lambda_1 - 2\lambda_2) + (1/2)\mu_3 + (1/3)\mu + w & (0, \dot{F}_3(1 + ie_1), 0, 0, 0, 0) \\
-((-1/2)(-\lambda_0 - \lambda_1 - 2\lambda_2) + (1/2)\mu_3 + (1/3)\mu + w) & (0, 0, \dot{F}_3(1 - ie_1), 0, 0, 0) \\
(1/2)(-\lambda_0 - \lambda_1 - 2\lambda_2) + (1/2)\mu_3 + (1/3)\mu + w & (0, \dot{F}_3(1 - ie_1), 0, 0, 0, 0) \\
-((1/2)(-\lambda_0 - \lambda_1 - 2\lambda_2) + (1/2)\mu_3 + (1/3)\mu + w) & (0, 0, \dot{F}_3(1 + ie_1), 0, 0, 0) \\
(-1/2)(\lambda_0 + \lambda_1 - 2\lambda_2) + (1/2)\mu_3 + (1/3)\mu + w & (0, \dot{F}_3(e_2 + ie_3), 0, 0, 0, 0) \\
-((-1/2)(\lambda_0 + \lambda_1 - 2\lambda_2) + (1/2)\mu_3 + (1/3)\mu + w) & (0, 0, \dot{F}_3(e_2 - ie_3), 0, 0, 0) \\
(1/2)(\lambda_0 + \lambda_1 - 2\lambda_2) + (1/2)\mu_3 + (1/3)\mu + w & (0, \dot{F}_3(e_2 - ie_3), 0, 0, 0, 0) \\
-((1/2)(\lambda_0 + \lambda_1 - 2\lambda_2) + (1/2)\mu_3 + (1/3)\mu + w) & (0, 0, \dot{F}_3(e_2 + ie_3), 0, 0, 0) \\
\mu + w & (0, \dot{1}, 0, 0, 0, 0) \\
-(\mu + w) & (0, 0, \dot{1}, 0, 0, 0) \\
-\mu + w & (0, \dot{1}, 0, 0, 0, 0) \\
-(-\mu + w) & (0, 0, \dot{1}, 0, 0, 0) \\
2w & (0, 0, 0, 0, \dot{1}, 0) \\
-2w & (0, 0, 0, 0, 0, \dot{1})
\end{array}$$

where $\dot{F}_k(x) := (F_k(x), 0, 0, 0)$, $F_k(x) := (0, F_k(x), 0, 0)$, $\dot{1} := (0, 0, 1, 0)$, $\dot{1} := (0, 0, 0, 1)$ in \mathfrak{B}^C .


Thus, since $(\mathfrak{e}_8^C)^{\varepsilon_1, \varepsilon_2}$ is spanned by \mathfrak{h}_8 and associated root vectors above, the roots obtained above are all. \square

Subsequently, we prove the following theorem.

Theorem 7.18. *In the root system Δ of Theorem 7.17*

$$\Pi = \{\alpha_1, \alpha_2, \alpha_3, \alpha_4, \alpha_5, \alpha_6, \alpha_7\}$$

is a fundamental root system of $(\mathfrak{e}_8^C)^{\varepsilon_1, \varepsilon_2}$, where $\alpha_1 = \mu + w$, $\alpha_2 = -2w$, $\alpha_3 = \mu_1 - (1/3)\mu + w$, $\alpha_4 = -(1/2)(\lambda_0 + \lambda_1 + 2\lambda_2 + \mu_1 - \mu_2)$, $\alpha_5 = -(1/2)(-2\lambda_0 + \mu_2 - \mu_3)$, $\alpha_6 = -(\lambda_0 - \lambda_1)$, $\alpha_7 = 2\lambda_2$. The Dynkin diagram of $(\mathfrak{e}_8^C)^{\varepsilon_1, \varepsilon_2}$ is given by



In particular, we have that the type of the group $(E_8^C)^{\varepsilon_1, \varepsilon_2}$ as Lie algebras is E_7 .

Proof. The all positive roots are expressed by $\alpha_1, \alpha_2, \alpha_3, \alpha_4, \alpha_5, \alpha_6$ as follows:

$$\begin{aligned}
-(\lambda_0 - \lambda_1) &= \alpha_6, \\
-(\lambda_0 + \lambda_1) &= \alpha_1 + 2\alpha_2 + 3\alpha_3 + 4\alpha_4 + 2\alpha_5 + \alpha_6 + 2\alpha_7, \\
2\lambda_2 &= \alpha_7, \\
-(1/2)(-2\lambda_0 + \mu_2 - \mu_3) &= \alpha_5, \\
(1/2)(-2\lambda_0 - \mu_2 + \mu_3) &= \alpha_1 + 2\alpha_2 + 3\alpha_3 + 4\alpha_4 + 3\alpha_5 + 2\alpha_6 + 2\alpha_7, \\
-(1/2)(-2\lambda_1 + \mu_2 - \mu_3) &= \alpha_5 + \alpha_6, \\
(1/2)(-2\lambda_1 - \mu_2 + \mu_3) &= \alpha_1 + 2\alpha_2 + 3\alpha_3 + 4\alpha_4 + 3\alpha_5 + \alpha_6 + 2\alpha_7, \\
(1/2)(\lambda_0 - \lambda_1 - 2\lambda_2 - \mu_1 + \mu_3) &= \alpha_4 + \alpha_5, \\
-(1/2)(\lambda_0 - \lambda_1 - 2\lambda_2 + \mu_1 - \mu_3) &= \alpha_4 + \alpha_5 + \alpha_6 + \alpha_7, \\
(1/2)(\lambda_0 - \lambda_1 + 2\lambda_2 - \mu_1 + \mu_3) &= \alpha_4 + \alpha_5 + \alpha_7, \\
-(1/2)(\lambda_0 - \lambda_1 + 2\lambda_2 + \mu_1 - \mu_3) &= \alpha_4 + \alpha_5 + \alpha_6, \\
-(1/2)(\lambda_0 + \lambda_1 + 2\lambda_2 + \mu_1 - \mu_2) &= \alpha_4, \\
-(1/2)(\lambda_0 + \lambda_1 + 2\lambda_2 - \mu_1 + \mu_2) &= \alpha_1 + 2\alpha_2 + 3\alpha_3 + 3\alpha_4 + 2\alpha_5 + \alpha_6 + \alpha_7, \\
(1/2)(-\lambda_0 - \lambda_1 + 2\lambda_2 + \mu_1 - \mu_2) &= \alpha_1 + 2\alpha_2 + 3\alpha_3 + 3\alpha_4 + 2\alpha_5 + \alpha_6 + 2\alpha_7, \\
(1/2)(-\lambda_0 - \lambda_1 + 2\lambda_2 - \mu_1 + \mu_2) &= \alpha_4 + \alpha_7 \\
\mu_1 + (2/3)\mu &= \alpha_1 + \alpha_2 + \alpha_3, \\
-(\mu_2 + (2/3)\mu) &= \alpha_2 + 2\alpha_3 + 2\alpha_4 + 2\alpha_5 + \alpha_6 + \alpha_7, \\
\mu_3 + (2/3)\mu &= \alpha_1 + \alpha_2 + \alpha_3 + 2\alpha_4 + 2\alpha_5 + \alpha_6 + \alpha_7, \\
-\lambda_0 - (1/2)\mu_1 + (2/3)\mu &= \alpha_1 + \alpha_2 + \alpha_3 + 2\alpha_4 + \alpha_5 + \alpha_6 + \alpha_7, \\
-(\lambda_0 - (1/2)\mu_1 + (2/3)\mu) &= \alpha_2 + 2\alpha_3 + 2\alpha_4 + \alpha_5 + \alpha_6 + \alpha_7, \\
-\lambda_1 - (1/2)\mu_1 + (2/3)\mu &= \alpha_1 + \alpha_2 + \alpha_3 + 2\alpha_4 + \alpha_5 + \alpha_7, \\
-(\lambda_1 - (1/2)\mu_1 + (2/3)\mu) &= \alpha_2 + 2\alpha_3 + \alpha_4 + \alpha_5 + \alpha_7, \\
-(1/2)(-\lambda_0 + \lambda_1 + 2\lambda_2) - (1/2)\mu_2 + (2/3)\mu &= \alpha_1 + \alpha_2 + \alpha_3 + \alpha_4 + \alpha_5, \\
(1/2)(-\lambda_0 + \lambda_1 + 2\lambda_2) - (1/2)\mu_2 + (2/3)\mu &= \alpha_1 + \alpha_2 + \alpha_3 + \alpha_4 + \alpha_5 + \alpha_6 + \alpha_7, \\
-(1/2)(-\lambda_0 + \lambda_1 - 2\lambda_2) - (1/2)\mu_2 + (2/3)\mu &= \alpha_1 + \alpha_2 + \alpha_3 + \alpha_4 + \alpha_5 + \alpha_7, \\
(1/2)(-\lambda_0 + \lambda_1 - 2\lambda_2) - (1/2)\mu_2 + (2/3)\mu &= \alpha_1 + \alpha_2 + \alpha_3 + \alpha_4 + \alpha_5 + \alpha_6, \\
-((-1/2)(-\lambda_0 - \lambda_1 - 2\lambda_2) - (1/2)\mu_3 + (2/3)\mu) &= \alpha_2 + 2\alpha_3 + \alpha_4 + 2\alpha_5 + \alpha_6 + \alpha_7, \\
(1/2)(-\lambda_0 - \lambda_1 - 2\lambda_2) - (1/2)\mu_3 + (2/3)\mu &= \alpha_1 + \alpha_2 + \alpha_3 + \alpha_4, \\
-(1/2)(\lambda_0 + \lambda_1 - 2\lambda_2) - (1/2)\mu_3 + (2/3)\mu &= \alpha_1 + \alpha_2 + \alpha_3 + \alpha_4 + \alpha_7, \\
-((1/2)(\lambda_0 + \lambda_1 - 2\lambda_2) - (1/2)\mu_3 + (2/3)\mu) &= \alpha_2 + 2\alpha_3 + \alpha_4 + 2\alpha_5 + \alpha_6 + 2\alpha_7, \\
\mu_1 - (1/3)\mu + w &= \alpha_3, \\
-(\mu_2 - (1/3)\mu + w) &= \alpha_1 + 2\alpha_2 + 2\alpha_3 + 2\alpha_4 + 2\alpha_5 + \alpha_6 + \alpha_7, \\
\mu_3 - (1/3)\mu + w &= \alpha_3 + 2\alpha_4 + 2\alpha_5 + \alpha_6 + \alpha_7, \\
-\lambda_0 - (1/2)\mu_1 - (1/3)\mu + w &= \alpha_3 + 2\alpha_4 + \alpha_5 + \alpha_6 + \alpha_7,
\end{aligned}$$

$$\begin{aligned}
& -(\lambda_0 - (1/2)\mu_1 - (1/3)\mu + w) = \alpha_1 + 2\alpha_2 + 2\alpha_3 + 2\alpha_4 + \alpha_5 + \alpha_6 + \alpha_7, \\
& -\lambda_1 - (1/2)\mu_1 - (1/3)\mu + w = \alpha_3 + 2\alpha_4 + \alpha_5 + \alpha_7, \\
& -(\lambda_1 - (1/2)\mu_1 - (1/3)\mu + w) = \alpha_1 + 2\alpha_2 + 2\alpha_3 + 2\alpha_4 + \alpha_5 + \alpha_7, \\
& -(1/2)(-\lambda_0 + \lambda_1 + 2\lambda_2) - (1/2)\mu_2 - (1/3)\mu + w = \alpha_3 + \alpha_4 + \alpha_5, \\
& (1/2)(-\lambda_0 + \lambda_1 + 2\lambda_2) - (1/2)\mu_2 - (1/3)\mu + w = \alpha_3 + \alpha_4 + \alpha_5 + \alpha_6 + \alpha_7, \\
& -(1/2)(-\lambda_0 + \lambda_1 - 2\lambda_2) - (1/2)\mu_2 - (1/3)\mu + w = \alpha_3 + \alpha_4 + \alpha_5 + \alpha_7, \\
& (1/2)(-\lambda_0 + \lambda_1 + 2\lambda_2) - (1/2)\mu_2 - (1/3)\mu + w = \alpha_3 + \alpha_4 + \alpha_5 + \alpha_6 + \alpha_7, \\
& -((-1/2)(-\lambda_0 - \lambda_1 - 2\lambda_2) - (1/2)\mu_3 - (1/3)\mu + w) = \alpha_1 + 2\alpha_2 + 2\alpha_3 + 3\alpha_4 + 2\alpha_5 + \alpha_6 + 2\alpha_7, \\
& (1/2)(-\lambda_0 - \lambda_1 - 2\lambda_2) - (1/2)\mu_3 - (1/3)\mu + w = \alpha_3 + \alpha_4, \\
& (-1/2)(\lambda_0 + \lambda_1 - 2\lambda_2) - (1/2)\mu_3 - (1/3)\mu + w = \alpha_3 + \alpha_4 + \alpha_7, \\
& -((1/2)(\lambda_0 + \lambda_1 - 2\lambda_2) - (1/2)\mu_3 - (1/3)\mu + w) = \alpha_1 + 2\alpha_2 + 2\alpha_3 + 3\alpha_4 + 2\alpha_5 + \alpha_6 + 2\alpha_7, \\
& \quad -(-\mu_1 + (1/3)\mu + w) = \alpha_2 + \alpha_3, \\
& \quad \quad -\mu_2 + (1/3)\mu + w = \alpha_1 + \alpha_2 + 2\alpha_3 + 2\alpha_4 + 2\alpha_5 + \alpha_6 + \alpha_7, \\
& \quad \quad -(-\mu_3 + (1/3)\mu + w) = \alpha_2 + \alpha_3 + 2\alpha_4 + 2\alpha_5 + \alpha_6 + \alpha_7, \\
& \quad \quad -\lambda_0 + (1/2)\mu_1 + (1/3)\mu + w = \alpha_1 + \alpha_2 + 2\alpha_3 + 2\alpha_4 + \alpha_5 + \alpha_6 + \alpha_7, \\
& \quad \quad -(\lambda_0 + (1/2)\mu_1 + (1/3)\mu + w) = \alpha_2 + \alpha_3 + 2\alpha_4 + \alpha_5 + \alpha_6 + \alpha_7, \\
& \quad \quad -\lambda_1 + (1/2)\mu_1 + (1/3)\mu + w = \alpha_1 + \alpha_2 + 2\alpha_3 + 2\alpha_4 + \alpha_5 + \alpha_7, \\
& \quad \quad -(\lambda_1 + (1/2)\mu_1 + (1/3)\mu + w) = \alpha_2 + \alpha_3 + 2\alpha_4 + \alpha_5 + \alpha_7, \\
& -((-1/2)(-\lambda_0 + \lambda_1 + 2\lambda_2) + (1/2)\mu_2 + (1/3)\mu + w) = \alpha_2 + \alpha_3 + \alpha_4 + \alpha_5 + \alpha_6 + \alpha_7, \\
& -((1/2)(-\lambda_0 + \lambda_1 + 2\lambda_2) + (1/2)\mu_2 + (1/3)\mu + w) = \alpha_2 + \alpha_3 + \alpha_4 + \alpha_5, \\
& -((-1/2)(-\lambda_0 + \lambda_1 - 2\lambda_2) + (1/2)\mu_2 + (1/3)\mu + w) = \alpha_2 + \alpha_3 + \alpha_4 + \alpha_5 + \alpha_6, \\
& -((1/2)(-\lambda_0 + \lambda_1 - 2\lambda_2) + (1/2)\mu_2 + (1/3)\mu + w) = \alpha_2 + \alpha_3 + \alpha_4 + \alpha_5 + \alpha_7, \\
& -((-1/2)(-\lambda_0 - \lambda_1 - 2\lambda_2) + (1/2)\mu_3 + (1/3)\mu + w) = \alpha_2 + \alpha_3 + \alpha_4, \\
& \quad (1/2)(-\lambda_0 - \lambda_1 - 2\lambda_2) + (1/2)\mu_3 + (1/3)\mu + w = \alpha_1 + \alpha_2 + 2\alpha_3 + 3\alpha_4 + 2\alpha_5 + \alpha_6 + \alpha_7, \\
& \quad (-1/2)(\lambda_0 + \lambda_1 - 2\lambda_2) + (1/2)\mu_3 + (1/3)\mu + w = \alpha_1 + \alpha_2 + 2\alpha_3 + 3\alpha_4 + 2\alpha_5 + \alpha_6 + 2\alpha_7, \\
& -((1/2)(\lambda_0 + \lambda_1 - 2\lambda_2) + (1/2)\mu_3 + (1/3)\mu + w) = \alpha_2 + \alpha_3 + \alpha_4 + \alpha_7, \\
& \quad \quad \mu + w = \alpha_1, \\
& \quad \quad -(-\mu + w) = \alpha_1 + \alpha_2, \\
& \quad \quad -2w = \alpha_2.
\end{aligned}$$

Hence Π is a fundamental root system of $(\mathfrak{e}_8^C)^{\varepsilon_1, \varepsilon_2}$.

Then, for $R, R' \in \mathfrak{e}_8^C$, the Killing form B_8 of \mathfrak{e}_8^C is given by

$$\begin{aligned}
B_8(R, R') &= B_8((\Phi, P, Q, r, s, t), (\Phi', P', Q', r', s', t')) \\
&= \frac{5}{3}B_7(\Phi, \Phi') + 15\{Q, P'\} - 15\{P, Q'\} + 120rr' + 60ts' + 60st'
\end{aligned}$$

([9, Theorem 5.3.2]), so that for $R_8 := (\Phi, 0, 0, w, 0, 0)$, $R_8' := (\Phi', 0, 0, w', 0, 0) \in \mathfrak{h}_8$, we have

$$\begin{aligned}
B_8(R_7, R_8') &= \frac{5}{3}B_7(\Phi, \Phi') + 120ww' \\
&= \frac{5}{3}(36(\lambda_0\lambda_0' + \lambda_1\lambda_1' + 2\lambda_2\lambda_2') + 18(\mu_1\mu_1' + \mu_2\mu_2' + \mu_3\mu_3') + 24\mu\mu') + 120ww' \\
&= 60(\lambda_0\lambda_0' + \lambda_1\lambda_1' + 2\lambda_2\lambda_2') + 30(\mu_1\mu_1' + \mu_2\mu_2' + \mu_3\mu_3') + 40\mu\mu' + 120ww',
\end{aligned}$$

where $\Phi := \Phi(\lambda_0(iG_{01}) + \lambda_1(iG_{23}) + \lambda_2(i(G_{45} + G_{67}) + (\mu_1 E_1 + \mu_2 E_2 + \mu_3 E_3)^\sim, 0, 0, \mu, \Phi' := \Phi(\lambda_0'(iG_{01}) + \lambda_1'(iG_{23}) + \lambda_2'(i(G_{45} + G_{67}) + (\mu_1' E_1 + \mu_2' E_2 + \mu_3' E_3)^\sim, 0, 0, \mu')$.

Now, the canonical elements $R_{\alpha_1}, R_{\alpha_2}, R_{\alpha_3}, R_{\alpha_4}, R_{\alpha_5}, R_{\alpha_6}, R_{\alpha_7}$ corresponding to $\alpha_1, \alpha_2, \alpha_3, \alpha_4, \alpha_5, \alpha_6, \alpha_7$ are determined as follows:

$$R_{\alpha_1} = (\Phi(0, 0, 0, \frac{1}{40}), 0, 0, \frac{1}{120}, 0, 0, 0),$$

$$R_{\alpha_2} = (0, 0, 0, -\frac{1}{60}, 0, 0),$$

$$R_{\alpha_3} = (\Phi((\frac{1}{45}E_1 - \frac{1}{90}E_2 - \frac{1}{90}E_3)^\sim, 0, 0, -\frac{1}{120}), 0, 0, \frac{1}{120}, 0, 0),$$

$$R_{\alpha_4} = (\Phi(-\frac{1}{120}(iG_{01}) - \frac{1}{120}(iG_{23}) - \frac{1}{120}(i(G_{45} + G_{67})) + (-\frac{1}{60}E_1 + \frac{1}{60}E_2)^\sim, 0, 0, 0), 0, 0, 0, 0, 0),$$

$$R_{\alpha_5} = (\Phi(\frac{1}{60}(iG_{01}) + (-\frac{1}{60}E_2 + \frac{1}{60}E_3)^\sim, 0, 0, 0), 0, 0, 0, 0, 0),$$

$$R_{\alpha_6} = (\Phi(-\frac{1}{60}(iG_{01}) + \frac{1}{60}(iG_{23}), 0, 0, 0), 0, 0, 0, 0, 0),$$

$$R_{\alpha_7} = (\Phi(-\frac{1}{60}(i(G_{45} + G_{67})), 0, 0, 0), 0, 0, 0, 0, 0).$$

Hence we have the following

$$(\alpha_1, \alpha_1) = B_8(R_{\alpha_1}, R_{\alpha_1}) = 40 \cdot \left(-\frac{1}{40}\right)^2 + 120 \cdot \left(\frac{1}{120}\right)^2 = \frac{1}{30},$$

$$(\alpha_1, \alpha_2) = B_8(R_{\alpha_1}, R_{\alpha_2}) = 120 \cdot \left(\frac{1}{120}\right) \left(-\frac{1}{60}\right) = -\frac{1}{60},$$

$$(\alpha_1, \alpha_3) = (\alpha_1, \alpha_4) = (\alpha_1, \alpha_5) = (\alpha_1, \alpha_6) = (\alpha_1, \alpha_7) = 0,$$

$$(\alpha_2, \alpha_2) = B_8(R_{\alpha_2}, R_{\alpha_2}) = 120 \cdot \left(-\frac{1}{60}\right)^2 = \frac{1}{30},$$

$$(\alpha_2, \alpha_3) = B_8(R_{\alpha_2}, R_{\alpha_3}) = 120 \cdot \left(-\frac{1}{60}\right) \left(\frac{1}{120}\right) = -\frac{1}{60},$$

$$(\alpha_2, \alpha_4) = (\alpha_2, \alpha_5) = (\alpha_2, \alpha_6) = (\alpha_2, \alpha_7) = 0,$$

$$(\alpha_3, \alpha_3) = B_8(R_{\alpha_3}, R_{\alpha_3}) = 30 \cdot \left(\left(\frac{1}{45}\right)^2 + \left(-\frac{1}{90}\right)^2 + \left(-\frac{1}{90}\right)^2\right) + 40 \cdot \left(-\frac{1}{120}\right)^2 + 120 \cdot \left(\frac{1}{120}\right)^2 = \frac{1}{30},$$

$$(\alpha_3, \alpha_4) = B_8(R_{\alpha_3}, R_{\alpha_4}) = 30 \cdot \left(\left(\frac{1}{45}\right) \left(-\frac{1}{60}\right) + \left(-\frac{1}{90}\right) \left(\frac{1}{60}\right)\right) = -\frac{1}{60},$$

$$(\alpha_3, \alpha_5) = (\alpha_3, \alpha_6) = (\alpha_3, \alpha_7) = 0,$$

$$(\alpha_4, \alpha_4) = B_8(R_{\alpha_4}, R_{\alpha_4}) = 60 \cdot \left(\left(-\frac{1}{120}\right)^2 + \left(-\frac{1}{120}\right)^2 + 2 \left(-\frac{1}{120}\right)^2\right) + 30 \cdot \left(\left(-\frac{1}{60}\right)^2 + \left(\frac{1}{60}\right)^2\right) = \frac{1}{30},$$

$$(\alpha_4, \alpha_5) = B_8(R_{\alpha_4}, R_{\alpha_5}) = 60 \cdot \left(-\frac{1}{120}\right) \left(\frac{1}{60}\right) + 30 \cdot \left(\frac{1}{60}\right) \left(-\frac{1}{60}\right) = -\frac{1}{60},$$

$$(\alpha_4, \alpha_6) = 0,$$

$$(\alpha_4, \alpha_7) = 60 \cdot 2 \cdot \left(-\frac{1}{120}\right) \left(\frac{1}{60}\right) = -\frac{1}{60},$$

$$(\alpha_5, \alpha_5) = B_8(R_{\alpha_5}, R_{\alpha_5}) = 60 \cdot \left(\frac{1}{60}\right)^2 + 30 \cdot \left(\left(-\frac{1}{60}\right)^2 + \left(\frac{1}{60}\right)^2\right) = \frac{1}{30},$$

$$(\alpha_5, \alpha_6) = B_8(R_{\alpha_5}, R_{\alpha_6}) = 60 \cdot \left(\frac{1}{60}\right) \left(-\frac{1}{60}\right) = -\frac{1}{60},$$

$$(\alpha_5, \alpha_7) = 0,$$

$$(\alpha_6, \alpha_6) = B_8(R_{\alpha_6}, R_{\alpha_6}) = 60 \cdot \left(\left(-\frac{1}{60}\right)^2 + \left(\frac{1}{60}\right)^2\right) = \frac{1}{30},$$

$$(\alpha_6, \alpha_7) = 0,$$

$$(\alpha_7, \alpha_7) = B_8(R_{\alpha_7}, R_{\alpha_7}) = 60 \cdot 2 \cdot \left(\frac{1}{60}\right)^2 = \frac{1}{30}.$$

Thus, using the inner product above, we have

$$\begin{aligned} \cos \theta_{12} &= \frac{(\alpha_1, \alpha_2)}{\sqrt{(\alpha_1, \alpha_1)(\alpha_2, \alpha_2)}} = -\frac{1}{2}, & \cos \theta_{13} &= \frac{(\alpha_1, \alpha_3)}{\sqrt{(\alpha_1, \alpha_1)(\alpha_3, \alpha_3)}} = 0, \\ \cos \theta_{14} &= \frac{(\alpha_1, \alpha_4)}{\sqrt{(\alpha_1, \alpha_1)(\alpha_4, \alpha_4)}} = 0, & \cos \theta_{15} &= \frac{(\alpha_1, \alpha_5)}{\sqrt{(\alpha_1, \alpha_1)(\alpha_5, \alpha_5)}} = 0, \\ \cos \theta_{16} &= \frac{(\alpha_1, \alpha_6)}{\sqrt{(\alpha_1, \alpha_1)(\alpha_6, \alpha_6)}} = 0, & \cos \theta_{17} &= \frac{(\alpha_1, \alpha_7)}{\sqrt{(\alpha_1, \alpha_1)(\alpha_7, \alpha_7)}} = 0, \\ \cos \theta_{23} &= \frac{(\alpha_2, \alpha_3)}{\sqrt{(\alpha_2, \alpha_2)(\alpha_3, \alpha_3)}} = -\frac{1}{2}, & \cos \theta_{24} &= \frac{(\alpha_2, \alpha_4)}{\sqrt{(\alpha_2, \alpha_2)(\alpha_4, \alpha_4)}} = 0, \\ \cos \theta_{25} &= \frac{(\alpha_2, \alpha_5)}{\sqrt{(\alpha_2, \alpha_2)(\alpha_5, \alpha_5)}} = 0, & \cos \theta_{26} &= \frac{(\alpha_2, \alpha_6)}{\sqrt{(\alpha_2, \alpha_2)(\alpha_6, \alpha_6)}} = 0, \\ \cos \theta_{27} &= \frac{(\alpha_2, \alpha_7)}{\sqrt{(\alpha_2, \alpha_2)(\alpha_7, \alpha_7)}} = 0, & \cos \theta_{34} &= \frac{(\alpha_3, \alpha_4)}{\sqrt{(\alpha_3, \alpha_3)(\alpha_4, \alpha_4)}} = -\frac{1}{2}, \\ \cos \theta_{35} &= \frac{(\alpha_3, \alpha_5)}{\sqrt{(\alpha_3, \alpha_3)(\alpha_5, \alpha_5)}} = 0, & \cos \theta_{36} &= \frac{(\alpha_3, \alpha_6)}{\sqrt{(\alpha_3, \alpha_3)(\alpha_6, \alpha_6)}} = 0, \\ \cos \theta_{37} &= \frac{(\alpha_3, \alpha_7)}{\sqrt{(\alpha_3, \alpha_3)(\alpha_7, \alpha_7)}} = 0, & \cos \theta_{45} &= \frac{(\alpha_4, \alpha_5)}{\sqrt{(\alpha_4, \alpha_4)(\alpha_5, \alpha_5)}} = -\frac{1}{2}, \\ \cos \theta_{46} &= \frac{(\alpha_4, \alpha_6)}{\sqrt{(\alpha_4, \alpha_4)(\alpha_6, \alpha_6)}} = 0, & \cos \theta_{47} &= \frac{(\alpha_4, \alpha_7)}{\sqrt{(\alpha_4, \alpha_4)(\alpha_7, \alpha_7)}} = -\frac{1}{2}, \\ \cos \theta_{56} &= \frac{(\alpha_5, \alpha_6)}{\sqrt{(\alpha_5, \alpha_5)(\alpha_6, \alpha_6)}} = -\frac{1}{2}, & \cos \theta_{57} &= \frac{(\alpha_5, \alpha_7)}{\sqrt{(\alpha_5, \alpha_5)(\alpha_7, \alpha_7)}} = 0, \\ \cos \theta_{67} &= \frac{(\alpha_6, \alpha_7)}{\sqrt{(\alpha_6, \alpha_6)(\alpha_7, \alpha_7)}} = 0. \end{aligned}$$

so that we can draw the required Dynkin diagram.

Therefore we have that the type of the group $(E_8^C)^{\varepsilon_1, \varepsilon_2}$ as Lie algebras is E_7 . \square

In order to determine the structure of the group $(E_8^C)^{\varepsilon_1, \varepsilon_2}$, we determine the center $z((E_8^C)^{\varepsilon_1, \varepsilon_2})$ of $(E_8^C)^{\varepsilon_1, \varepsilon_2}$.

Proposition 7.19. *The center $z((E_8^C)^{\varepsilon_1, \varepsilon_2})$ of the group $(E_8^C)^{\varepsilon_1, \varepsilon_2}$ is isomorphic to a cyclic group of order two :*

$$z((E_8^C)^{\varepsilon_1, \varepsilon_2}) = \{1, \gamma\} \cong \mathbf{Z}_2.$$

Proof. Let $\alpha \in z((E_8^C)^{\varepsilon_1, \varepsilon_2})$. Then, since $1_- \in (\mathfrak{B}^C)_{\varepsilon_1, \varepsilon_2}$, we can set $\alpha 1_- := (\Phi, P, Q, r, s, t) \in (\mathfrak{B}^C)_{\varepsilon_1, \varepsilon_2}$. For $\beta \in (E_7^C)^{\varepsilon_1, \varepsilon_2} \subset (E_8^C)^{\varepsilon_1, \varepsilon_2}$, we see

$$\beta(\alpha 1_-) = \alpha(\beta 1_-) = \alpha 1_- (= (\Phi, P, Q, r, s, t)),$$

on the other hand, we see

$$\beta(\alpha 1_-) = \beta(\Phi, P, Q, r, s, t) = (\beta\Phi\beta^{-1}, \beta P, \beta Q, r, s, t).$$

Hence we have

$$\begin{cases} \beta\Phi\beta^{-1} = \Phi, \\ \beta P = P, \\ \beta Q = Q. \end{cases} \quad \text{for all } \beta \in (E_7^C)^{\varepsilon_1, \varepsilon_2}$$

Here, let the element $-1 \in (E_7^C)^{\varepsilon_1, \varepsilon_2}$ as β . Then we have $P = Q = 0$. Moreover, let the element $\omega 1 \in (E_6^C)^{\varepsilon_1, \varepsilon_2} \subset (E_7^C)^{\varepsilon_1, \varepsilon_2}$ as β , where $\omega \in C, \omega^3 = 1, \omega \neq 1$. Then, for $\Phi := \Phi(\phi, A, B, \nu)$, we have $A = B = 0$. Hence $\alpha 1_-$ is of the form $(\Phi(\phi, 0, 0, \nu), 0, 0, r, s, t)$: $\alpha 1_- = (\Phi(\phi, 0, 0, \nu), 0, 0, r, s, t)$, moreover we see $t \neq 0$ for $\alpha 1_- = (\Phi(\phi, 0, 0, \nu), 0, 0, r, s, t)$.

Indeed, assume $t = 0$, then we have $r = 0$ from Lemma 7.14 (6). Hence $\alpha 1_-$ is of the form $((\Phi(\phi, 0, 0, \nu), 0, 0, 0, s, 0))$. In addition, $\Phi(\phi, 0, 0, \nu)$ satisfies the condition $\Phi Q' = 0$ for all $Q' \in (\mathfrak{P}_{\mathbf{H}})^C$ from Lemma 7.14 (5), so set $Q' := (0, 0, 0, 1)$, then we have $\nu = 0$. Moreover set $Q' := (E, 0, 0, 0)$ and $Q' := (F_k(x), 0, 0, 0), x \in \mathbf{H}^C, k = 1, 2$, then we have $T = 0$ and $D_i = 0, i = 1, 2, 3, a_i = 0$, where $\phi := (D_1, D_2, D_3) + \tilde{A}_1(a_1) + \tilde{A}_2(a_2) + \tilde{A}_3(a_3) + \tilde{T} \in (\mathfrak{e}_6^C)^{\varepsilon_1, \varepsilon_2}$, that is, $\phi = 0$. Thus $\alpha 1_-$ is of the form $s 1^-$: $\alpha 1_- = s 1^-, s \neq 0$. Let the mapping $\varphi : SU(2) \rightarrow E_8 \subset E_8^C$ which is defined in [9, Theorem 5.7.4] as φ_3 . Then, for $A_1 := \begin{pmatrix} 0 & i \\ i & 0 \end{pmatrix} \in SU(2) = \{A \in M(2, C) \mid (\tau^t A)A = E, \det A = 1\}$, we choose the element $\varphi(A_1) = \exp((\pi/2)\text{ad}(0, 0, 0, 0, i, i)) \in (E_8^C)^{\varepsilon_1, \varepsilon_2}$ (Lemma 7.12 (1)). It follows that

$$\varphi(A_1)(\alpha 1_-) = \alpha(\varphi(A_1)1_-) = \alpha 1^-,$$

on the other hand, it follows that

$$\varphi(A_1)(\alpha 1_-) = \varphi(A_1)(s 1^-) = s 1_-.$$

Thus we have $\alpha 1^- = s 1_-$.

In addition, from $B_8(\alpha 1_-, \alpha 1^-) = B_8(1_-, 1^-)$, we have $s^2 = 1$. Moreover, it follows that

$$[\alpha 1_-, \alpha 1^-] = [s 1^-, s 1_-] = s^2[1^-, 1_-] = s^2 \tilde{1},$$

on the other hand, it follows that

$$[\alpha 1_-, \alpha 1^-] = \alpha[1_-, 1^-] = \alpha(-\tilde{1}) = -\alpha \tilde{1}.$$

Thus we have $\alpha \tilde{1} = -\tilde{1}$ from $s^2 = 1$. Consequently, we obtain the following

$$\alpha \tilde{1} = -\tilde{1}, \quad \alpha 1^- = s 1_-, \quad \alpha 1_- = s 1^-$$

with $s^2 = 1$, that is,

$$(i) \begin{cases} \alpha \tilde{1} = -\tilde{1} \\ \alpha 1^- = 1_- \\ \alpha 1_- = 1^- \end{cases}, \quad (ii) \begin{cases} \alpha \tilde{1} = -\tilde{1} \\ \alpha 1^- = -1_- \\ \alpha 1_- = 1^- \end{cases}, \quad (iii) \begin{cases} \alpha \tilde{1} = -\tilde{1} \\ \alpha 1^- = 1_- \\ \alpha 1_- = -1^- \end{cases}, \quad (iv) \begin{cases} \alpha \tilde{1} = -\tilde{1} \\ \alpha 1^- = -1_- \\ \alpha 1_- = -1^- \end{cases}.$$

Case (i). Again let the mapping φ be used above. Then, for $A_2 := \begin{pmatrix} \frac{i}{\sqrt{2}} & \frac{i}{\sqrt{2}} \\ i & -\frac{i}{\sqrt{2}} \end{pmatrix} \in SU(2)$,

we choose the element $\varphi(A_2) = \exp((\pi/2)\text{ad}(0, 0, 0, -i/\sqrt{2}, i/\sqrt{2}, i/\sqrt{2})) \in (E_8^C)^{\varepsilon_1, \varepsilon_2}$ (Lemma 7.12(1)). It follows from $\varphi(A_1)\alpha = \alpha\varphi(A_1)$ that

$$\varphi(A_2)(\alpha \tilde{1}) = \alpha\varphi(A_1)\tilde{1} = \alpha(0, 0, 0, 0, 1, 1) = 1^- + 1_-,$$

on the other hand, it follows that

$$\varphi(A_2)(\alpha\tilde{1}) = \varphi(A_1)(-\tilde{1}) = (0, 0, 0, 0, -1, -1) = -(1^- + 1_-).$$

Hence this result is contradiction, so that this case is impossible.

Case (ii). It follows that

$$[\alpha 1^-, \alpha 1_-] = [-1_-, 1^-] = -[1_-, 1^-] = -(2\tilde{1}) = 2\tilde{1},$$

on the other hand, it follows that

$$[\alpha 1^-, \alpha 1_-] = \alpha[1_-, 1^-] = \alpha[1_-, 1^-] = \alpha(2\tilde{1}) = 2\alpha\tilde{1}$$

Hence this results are contrary to $\alpha\tilde{1} = -\tilde{1}$, so that this case is also impossible.

Case (iii). The same result as in Case (ii) is obtained, so that this case is also impossible.

Case (iv). As in Case (i), let the mapping φ . Then, for $A_3 := \begin{pmatrix} \frac{1}{\sqrt{2}} & \frac{i}{\sqrt{2}} \\ \frac{i}{\sqrt{2}} & \frac{1}{\sqrt{2}} \end{pmatrix} \in SU(2)$,

we choose the element $\varphi(A_3) = \exp((\pi/4)\text{ad}(0, 0, 0, 0, i, i)) \in (E_8^C)^{\varepsilon_1, \varepsilon_2}$ (Lemma 7.12 (1)). It follows that

$$\varphi(A_3)(\alpha\tilde{1}) = \alpha\varphi(A_3)\tilde{1} = \alpha(0, 0, 0, 0, -i, i) = (-i)1^- + i1_-,$$

on the other hand, it follows from $\varphi(A_3)\alpha = \alpha\varphi(A_3)$ that

$$\varphi(A_3)(\alpha\tilde{1}) = \varphi(A_3)(-\tilde{1}) = -(0, 0, 0, 0, -i, i) = i1^- + (-i)1_-.$$

Hence this result is contradiction, so that this case is impossible.

With above, the assumption as $t = 0$ is unreasonable. Thus we obtain

$$\alpha 1_- = (\Phi, 0, 0, r, s, t), \quad t \neq 0.$$

Here, from Lemma 7.14 (2), we have

$$\Phi = -\frac{1}{2t}Q \times Q,$$

so that we have $\Phi = 0$ from $Q = 0$. Therefore $\alpha 1_-$ is of the form

$$\alpha 1_- = (0, 0, 0, r, s, t), \quad t \neq 0.$$

Now, let the mapping φ . Then, for $A_4 := \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} \in SU(2)$, we choose the elements $\varphi(A_4) = \exp((\pi/2)\text{ad}(0, 0, 0, 0, 1, -1)) \in (E_8^C)^{\varepsilon_1, \varepsilon_2}$ (Lemma 7.12(1)). Using $\varphi(A_1)$ defined above together, it follows that

$$\varphi(A_1)\varphi(A_4)(\alpha 1_-) = \alpha(\varphi(A_1)\varphi(A_4)1_-) = \alpha(-1_-) = -\alpha 1_- = (0, 0, 0, -r, -s, -t),$$

on the other hand, it follows that

$$\begin{aligned} \varphi(A_1)\varphi(A_4)(\alpha 1_-) &= \varphi(A_1)\varphi(A_4)(0, 0, 0, r, s, t) = \varphi(A_1)(0, 0, 0, -r, -t, -s) \\ &= (0, 0, 0, r - s, -t). \end{aligned}$$

Hence we have $r = 0$, so $\alpha 1_- = (0, 0, 0, 0, s, t), t \neq 0$. Here, since $\alpha 1_- \in \mathfrak{B}_{\varepsilon_1, \varepsilon_2}$ again, we see $st = 0$ from Lemma 7.14 (6). Thus we have $s = 0$ because of $t \neq 0$. Namely, we have

$$\alpha 1_- = t1_-, \quad t \neq 0.$$

Again, choose the element $\varphi(A_1)$. Then it follows that

$$\varphi(A_1)(\alpha 1_-) = \alpha\varphi(A_1)1_- = \alpha 1^-,$$

on the other hand, it follows that

$$\varphi(A_1)(\alpha 1_-) = \varphi(A_1)(t 1_-) = t 1^-,$$

that is, $\alpha 1^- = t 1^-$, $t \neq 0$.

Here, using the Killing form B_8 , it follows that

$$B_8(\alpha 1^-, \alpha 1_-) = B_8(1^-, 1_-) = 60,$$

on the other hand, it follows that

$$B_8(\alpha 1^-, \alpha 1_-) = B_8(t 1^-, t 1_-) = 60t^2.$$

Hence we have $t^2 = 1$,

so that it follows from $\alpha 1_- = t 1_-$, $\alpha 1^- = t 1^-$, $t \neq 0$ that

$$\begin{aligned} [\alpha 1^-, \alpha 1_-] &= \alpha [1^-, 1_-] = \alpha \tilde{1}, \\ [\alpha 1^-, \alpha 1_-] &= [t 1^-, t 1_-] = t^2 [1^-, 1_-] = t^2 \tilde{1} = \tilde{1}, \end{aligned}$$

that is, $\alpha \tilde{1} = \tilde{1}$.

Thus, from $t^2 = 1$, we have the following

$$\begin{cases} \alpha \tilde{1} = \tilde{1} \\ \alpha 1^- = 1^- \\ \alpha 1_- = 1_- \end{cases} \quad \text{or} \quad \begin{cases} \alpha \tilde{1} = \tilde{1} \\ \alpha 1^- = -1^- \\ \alpha 1_- = -1_- \end{cases}.$$

In the latter case, we choose the element $\varphi(A_2) \in (E_8^C)^{\varepsilon_1, \varepsilon_2}$ again, then it follows that

$$\varphi(A_2)(\alpha 1^-) = \alpha \varphi(A_2) 1^- = \alpha (1/2)(-\tilde{1} - 1^- + 1_-) = (1/2)(-\tilde{1} + 1^- - 1_-),$$

on the other hand, it follows that

$$\varphi(A_2)(\alpha 1^-) = \varphi(A_2)(-1^-) = -(1/2)(-\tilde{1} - 1^- + 1_-) = (1/2)(\tilde{1} + 1^- - 1_-).$$

Hence this result is contradiction, so that the latter case is impossible.

In the former case, note that $((E_8^C)^{\varepsilon_1, \varepsilon_2})_{\tilde{1}, 1^-, 1_-} = ((E_8^C)_{\tilde{1}, 1^-, 1_-})^{\varepsilon_1, \varepsilon_2} \cong (E_7^C)^{\varepsilon_1, \varepsilon_2}$, we see $\alpha \in (E_7^C)^{\varepsilon_1, \varepsilon_2}$.

Thus we obtain $\alpha \in z((E_7^C)^{\varepsilon_1, \varepsilon_2}) = \{1, \gamma, -1, -\gamma\} = \{1, \gamma\} \times \{1, -\gamma\} \cong \mathbf{Z}_2 \times \mathbf{Z}_2$ ([2, Proposition 1.1.6]). However, since it is easy to verify that $-1, -\gamma \notin E_8^C$, we have $z((E_8^C)^{\varepsilon_1, \varepsilon_2}) \subset \{1, \gamma\}$ and vice versa.

Therefore we have the required result

$$z((E_8^C)^{\varepsilon_1, \varepsilon_2}) = \{1, \gamma\} \cong \mathbf{Z}_2.$$

□

As for the group $(E_8^C)^{\varepsilon_1, \varepsilon_2}$, the following is a summary of what we have obtained up to this point in this section.

- The type of the group $(E_8^C)^{\varepsilon_1, \varepsilon_2}$ as Lie algebras is E_7 .
- The group $(E_8^C)^{\varepsilon_1, \varepsilon_2}$ is connected.
- The center $z((E_8^C)^{\varepsilon_1, \varepsilon_2})$ is isomorphic to a cyclic group of order two: $z((E_8^C)^{\varepsilon_1, \varepsilon_2}) \cong \mathbf{Z}_2$.

Now, we determine the structure of the group $(E_8^C)^{\varepsilon_1, \varepsilon_2}$.

Theorem 7.20. *The group $(E_8^C)^{\varepsilon_1, \varepsilon_2}$ is isomorphic to the group E_7^C : $(E_8^C)^{\varepsilon_1, \varepsilon_2} \cong E_7^C$.*

Proof. Since the group $(E_8^C)^{\varepsilon_1, \varepsilon_2}$ is connected (Theorem 7.16) and the type of the group $(E_8^C)^{\varepsilon_1, \varepsilon_2}$ as Lie algebras is E_7 (Theorem 7.18), note that $z(E_7^C) = \{1, -1\} \cong \mathbf{Z}_2$, we have

$$(E_8^C)^{\varepsilon_1, \varepsilon_2} \cong E_7^C \quad \text{or} \quad (E_8^C)^{\varepsilon_1, \varepsilon_2} \cong E_7^C / \mathbf{Z}_2.$$

Then, since the center $z((E_8^C)^{\varepsilon_1, \varepsilon_2})$ is isomorphic to a cyclic group of order two (Proposition 7.19), we have the required isomorphism

$$(E_8^C)^{\varepsilon_1, \varepsilon_2} \cong E_7^C.$$

□

We consider a real form $(E_8)^{\varepsilon_1, \varepsilon_2}$ of the group $(E_8^C)^{\varepsilon_1, \varepsilon_2}$:

$$(E_8)^{\varepsilon_1, \varepsilon_2} = \{ \alpha \in (E_8^C)^{\varepsilon_1, \varepsilon_2} \mid \langle \alpha R_1, \alpha R_2 \rangle = \langle R_1, R_2 \rangle \},$$

where the Hermite inner product $\langle R_1, R_2 \rangle$ is defined by $(-1/15)B_8(\tau\lambda_\omega R_1, R_2)$: $\langle R_1, R_2 \rangle = (-1/15)B_8(\tau\lambda_\omega R_1, R_2)$. Then, since $\alpha \in (E_8)^{\varepsilon_1, \varepsilon_2}$ commutes with $\varepsilon_i, i = 1, 2$, α induces a C -linear isomorphism of $(\mathfrak{e}_8^C)^{\varepsilon_1, \varepsilon_2}$. Hence the group $(E_8)^{\varepsilon_1, \varepsilon_2}$ is a compact Lie group as the closed subgroup of the unitary group $U(133) = \{ \alpha \in \text{Iso}_C((\mathfrak{e}_8^C)^{\varepsilon_1, \varepsilon_2}) \mid \langle \alpha R_1, \alpha R_2 \rangle = \langle R_1, R_2 \rangle \}$.

Then we have the following lemma.

Lemma 7.21. *The Lie algebra $(\mathfrak{e}_8)^{\varepsilon_1, \varepsilon_2}$ of the group $(E_8)^{\varepsilon_1, \varepsilon_2}$ is given by*

$$(\mathfrak{e}_8)^{\varepsilon_1, \varepsilon_2} = \{ R = (\Phi, P, -\tau\lambda P, r, s, -\tau s) \mid \Phi \in (\mathfrak{e}_7)^{\varepsilon_1, \varepsilon_2}, P \in (\mathfrak{P}_{\mathbf{H}})^C, r \in i\mathbf{R}, s \in C \},$$

where $(\mathfrak{e}_7)^{\varepsilon_1, \varepsilon_2}$ is defined by

$$(\mathfrak{e}_7)^{\varepsilon_1, \varepsilon_2} = \left\{ \Phi(\phi, A, -\tau A, \nu) \left| \begin{array}{l} \phi = (D_1, D_2, D_3) \\ +\tilde{A}_1(a_1) + \tilde{A}_2(a_2) + \tilde{A}_3(a_3) \\ +i(\tau_1 E_1 + \tau_2 E_2 + \tau_3 E_3 + F_1(t_1) \\ +F_2(t_2) + F_3(t_3))^\sim, \\ D_1 = d_{01}G_{01} + d_{02}G_{02} + d_{03}G_{03} \\ +d_{12}G_{12} + d_{13}G_{13} + d_{23}G_{23} \\ +d_{45}(G_{45} + G_{67}) + d_{46}(G_{46} - G_{57}) \\ +d_{47}(G_{47} + G_{56}), d_{ij} \in \mathbf{R}, \\ D_2 = \pi\kappa D_1, D_3 = \kappa\pi D_1, \\ a_k, t_k \in \mathbf{H}, \\ \tau_k \in \mathbf{R}, \tau_1 + \tau_2 + \tau_3 = 0, \\ A \in (\mathfrak{J}_{\mathbf{H}})^C, \nu \in i\mathbf{R} \end{array} \right. \right\}.$$

In particular, we have $\dim((\mathfrak{e}_8)^{\varepsilon_1, \varepsilon_2}) = 66 + 64 + 3 = 133$.

Proof. Note that $(\mathfrak{e}_8)^{\varepsilon_1, \varepsilon_2} = ((\mathfrak{e}_8^C)^{\varepsilon_1, \varepsilon_2})^{\tau\lambda_\omega}$, using the result of Lemma 7.12(1), we can obtain the required result by straightforward computations. □

We determine the structure of the group $(E_8)^{\varepsilon_1, \varepsilon_2}$.

Theorem 7.22. *The group $(E_8)^{\varepsilon_1, \varepsilon_2}$ is isomorphic to the group E_7 : $(E_8)^{\varepsilon_1, \varepsilon_2} \cong E_7$.*

Proof. First, it is easy to verify that $z((E_8^C)^{\varepsilon_1, \varepsilon_2}) \subset z((E_8)^{\varepsilon_1, \varepsilon_2})$. Indeed, let $\delta \in z((E_8^C)^{\varepsilon_1, \varepsilon_2})$. Then, since $(E_8)^{\varepsilon_1, \varepsilon_2} \subset (E_8^C)^{\varepsilon_1, \varepsilon_2}$, δ commutes with any elements $\beta \in (E_8)^{\varepsilon_1, \varepsilon_2}$. Hence we have $\delta \in z((E_8)^{\varepsilon_1, \varepsilon_2})$, that is, $z((E_8^C)^{\varepsilon_1, \varepsilon_2}) \subset z((E_8)^{\varepsilon_1, \varepsilon_2})$. Here, from $(E_8^C)^{\varepsilon_1, \varepsilon_2} \cong E_7^C$ (Theorem 7.20), the group $(E_8^C)^{\varepsilon_1, \varepsilon_2}$ is simply connected. Moreover, it follows from $\varepsilon_i(\tau\lambda_\omega) = (\tau\lambda_\omega)\varepsilon_i, i = 1, 2$ that

$$(E_8)^{\varepsilon_1, \varepsilon_2} = ((E_8^C)^{\tau\lambda_\omega})^{\varepsilon_1, \varepsilon_2} = ((E_8^C)^{\varepsilon_1, \varepsilon_2})^{\tau\lambda_\omega}.$$

Hence the group $(E_8)^{\varepsilon_1, \varepsilon_2}$ is connected. In addition, since the group $(E_8)^{\varepsilon_1, \varepsilon_2}$ is a real form of the group $(E_8^C)^{\varepsilon_1, \varepsilon_2}$, the type of $(E_8)^{\varepsilon_1, \varepsilon_2}$ as Lie algebras is E_7 . Thus, together with the fact that the group $(E_8)^{\varepsilon_1, \varepsilon_2}$ is compact, we have the following

$$(E_8)^{\varepsilon_1, \varepsilon_2} \cong E_7 \quad \text{or} \quad (E_8)^{\varepsilon_1, \varepsilon_2} \cong E_7/\mathbf{Z}_2.$$

In the latter case, since the center $z(E_7/\mathbf{Z}_2)$ of the group E_7/\mathbf{Z}_2 is trivial, the center $z((E_8)^{\varepsilon_1, \varepsilon_2})$ of the group $(E_8)^{\varepsilon_1, \varepsilon_2}$ is also trivial. However, it follows from $z((E_8^C)^{\varepsilon_1, \varepsilon_2}) \subset z((E_8)^{\varepsilon_1, \varepsilon_2})$ as mentioned above and $z((E_8^C)^{\varepsilon_1, \varepsilon_2}) = \{1, \gamma\}$ (Proposition 7.19), we see $\gamma \in z((E_8)^{\varepsilon_1, \varepsilon_2})$. This is contradiction, so that this case is impossible.

Therefore we have the required isomorphism

$$(E_8)^{\varepsilon_1, \varepsilon_2} \cong E_7.$$

□

We consider a subgroup $((F_4)^{\varepsilon_1, \varepsilon_2})_{E_1, E_2, E_3}$ of the group $(F_4)^{\varepsilon_1, \varepsilon_2}$:

$$((F_4)^{\varepsilon_1, \varepsilon_2})_{E_1, E_2, E_3} = \{\alpha \in (F_4)^{\varepsilon_1, \varepsilon_2} \mid \alpha E_1 = E_1, \alpha E_2 = E_2, \alpha E_3 = E_3\}.$$

Then, in order to prove the main theorem, we need the following proposition.

Proposition 7.23. *The group $((F_4)^{\varepsilon_1, \varepsilon_2})_{E_1, E_2, E_3}$ is isomorphic to the group $Sp(1) \times Sp(1) \times Sp(1)$: $((F_4)^{\varepsilon_1, \varepsilon_2})_{E_1, E_2, E_3} \cong Sp(1) \times Sp(1) \times Sp(1)$.*

Proof. We define a mapping $\varphi_{4, \varepsilon_i} : Sp(1) \times Sp(1) \times Sp(1) \rightarrow ((F_4)^{\varepsilon_1, \varepsilon_2})_{E_1, E_2, E_3}$ by

$$\varphi_{4, \varepsilon_i}(p_1, p_2, p_3)(M + \mathbf{a}) = A_3 M A_3^* + \mathbf{a} A_3^*, \quad M + \mathbf{a} \in \mathfrak{J}_H \oplus \mathbf{H}^3 = \mathfrak{J},$$

where $A_3 := \text{diag}(p_1, p_2, p_3) \in Sp(3)$. Note that $\varphi_{4, \varepsilon_i}$ is the restriction of the mapping $\varphi_4 : Sp(1) \times Sp(3) \rightarrow (F_4)^\gamma$ ([9, Theorem 2.11.2]), that is, $\varphi_{4, \varepsilon_i}(p_1, p_2, p_3) = \varphi_4(1, A_3)$.

First, we will prove that $\varphi_{4, \varepsilon_i}$ is well-defined and a homomorphism. However, since $\varphi_{4, \varepsilon_i}$ is the restriction of the mapping φ_4 , it is clear.

Next, we will prove that $\varphi_{4, \varepsilon_i}$ is surjective. Let $\alpha \in ((F_4)^{\varepsilon_1, \varepsilon_2})_{E_1, E_2, E_3}$. Then, since $\varepsilon_i^2 = \gamma, i = 1, 2$, we have $\alpha \in ((F_4)^\gamma)_{E_1, E_2, E_3} \subset (F_4)^\gamma$. Hence there exist $p \in Sp(1)$ and $A \in Sp(3)$ such that $\alpha = \varphi_4(p, A)$ ([9, Theorem 2.11.2]). First, it follows from $\alpha E_i = E_i, i = 1, 2, 3$ that

$$E_i = \alpha E_i = \varphi_4(p, A) E_i = A E_i A^*,$$

that is, $E_i = A E_i A^*, i = 1, 2, 3$. Set $A := \begin{pmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{pmatrix} \in Sp(3)$, then we have the following

$$\begin{aligned} a_{11} \overline{a_{11}} &= 1, a_{21} \overline{a_{21}} = a_{31} \overline{a_{31}} = 0 \quad (E_1 = A E_1 A^*), \\ a_{22} \overline{a_{22}} &= 1, a_{12} \overline{a_{12}} = a_{32} \overline{a_{32}} = 0 \quad (E_2 = A E_2 A^*), \\ a_{33} \overline{a_{33}} &= 1, a_{13} \overline{a_{13}} = a_{23} \overline{a_{23}} = 0 \quad (E_3 = A E_3 A^*). \end{aligned}$$

Hence we see $a_{11}, a_{22}, a_{33} \in Sp(1)$ and $a_{ij} = 0, i \neq j$, so that A is of the form

$$\text{diag}(p_1, p_2, p_3)(= A_3), \quad p_i \in Sp(1), i = 1, 2, 3.$$

Moreover, it follows from $\varepsilon_i \alpha = \alpha \varepsilon_i, i = 1, 2$ that

$$\begin{aligned} \varepsilon_i \alpha \varepsilon_i^{-1}(M + \mathbf{a}) &= \varepsilon_i \varphi_4(p, A_3)(M + (-e_i) \mathbf{a}) = \varepsilon_i (A_3 M A_3^* + p(-e_i \mathbf{a}) A_3^*) \\ &= A_3 M A_3^* + e_i (p(-e_i \mathbf{a}) A_3^*) = A_3 M A_3^* + (-e_i p e_i) \mathbf{a} A_3^* \\ &= \varphi_4(-e_i p e_i, A_3)(M + \mathbf{a}), \quad M + \mathbf{a} \in \mathfrak{J}_H \oplus \mathbf{H}^3 = \mathfrak{J}, \end{aligned}$$

that is, $\varphi_4(-e_i p e_i, A_3) = \varphi_4(p, A_3)$.

Hence we have the following

$$\begin{cases} -e_i p e_i = p \\ A_3 = A_3 \end{cases} \quad \text{or} \quad \begin{cases} -e_i p e_i = -p \\ A_3 = -A_3. \end{cases}$$

The latter case is impossible because of $A_3 \neq \mathbf{O}$ (\mathbf{O} is zero matrix). In the former case, from the first condition, we have $p \in \mathbf{R}$, and together with $p \in Sp(1)$, we see

$$p = 1 \quad \text{or} \quad p = -1.$$

Thus we have

$$\alpha = \varphi_4(1, A_3) \quad \text{or} \quad \alpha = \varphi_4(-1, A_3).$$

However, as for $\alpha = \varphi_4(-1, A_3)$, it follows from $\varphi_4(-1, -E) = 1$ that

$$\alpha = \varphi_4(-1, A_3) \cdot 1 = \varphi_4(-1, A_3)\varphi_4(-1, -E) = \varphi_4(1, -A_3), \quad -A_3 \in Sp(3).$$

With above, there exist $A_3 = \text{diag}(p_1, p_2, p_3)$, $p_i \in Sp(1), i = 1, 2, 3$ such that

$$\alpha = \varphi_4(1, A_3) = \varphi_{4, \varepsilon_i}(p_1, p_2, p_3).$$

The proof of surjective is completed.

Finally, we will determine $\text{Ker } \varphi_{4, \varepsilon_i}$. Since $\varphi_{4, \varepsilon_i}$ is the restriction of the mapping φ_4 , it is clear $\text{Ker } \varphi_{4, \varepsilon_i} = \{E\}$.

Therefore we have the required isomorphism

$$((F_4)^{\varepsilon_1, \varepsilon_2})_{E_1, E_2, E_3} \cong Sp(1) \times Sp(1) \times Sp(1).$$

□

Now, we prove the main theorem in this section.

Theorem 7.24. *The group $E_{8,\mathbf{H}}$ is isomorphic to the group E_7/\mathbf{Z}_2 , $\mathbf{Z}_2 = \{1, \gamma\}$: $E_{8,\mathbf{H}} \cong E_7/\mathbf{Z}_2$.*

Proof. Let E_7 as the group $(E_8)^{\varepsilon_1, \varepsilon_2}$ (Theorem 7.22). then we define a mapping $g_8 : (E_8)^{\varepsilon_1, \varepsilon_2} \rightarrow E_{8,\mathbf{H}}$ by

$$g_8(\alpha) = \alpha|_{(\mathfrak{e}_{8,\mathbf{H}})^C}.$$

First, we will prove that g_8 is well-defined. Let $\alpha \in (E_8)^{\varepsilon_1, \varepsilon_2}$, then we have $\alpha R \in (\mathfrak{e}_{8,\mathbf{H}})^C$ for $R \in (\mathfrak{e}_{8,\mathbf{H}})^C$. Indeed, any elements $R \in (\mathfrak{e}_{8,\mathbf{H}})^C$ satisfies $\gamma R = R$. Hence it follows from $(E_8)^{\varepsilon_1, \varepsilon_2} \subset (E_8)^\gamma$ that

$$\alpha R = \alpha(\gamma R) = \gamma(\alpha R), \quad R \in (\mathfrak{e}_{8,\mathbf{H}})^C,$$

that is, $\alpha R \in (\mathfrak{e}_{8,\mathbf{H}})^C$. Thus α induces a C -linear isomorphism of $(\mathfrak{e}_{8,\mathbf{H}})^C$, so that we have $g_8(\alpha) \in E_{8,\mathbf{H}}$, that is, g_8 is well-defined. It is clear that g_8 is a homomorphism.

Next we will determine $\text{Ker } g_8$. Let $\alpha \in \text{Ker } g_8$. For $1_- = (0, 0, 0, 0, 0, 1) \in (\mathfrak{e}_{8,\mathbf{H}})^C$, we have

$$\alpha 1_- = 1_-.$$

Hence we see $\alpha \in ((E_8)^{\varepsilon_1, \varepsilon_2})_{1_-} = ((E_8)_{1_-})^{\varepsilon_1, \varepsilon_2} \cong (E_7)^{\varepsilon_1, \varepsilon_2}$ (Subsection 2.5). Moreover, for $R^-_{\dot{1}} := (0, \dot{1}, 0, 0, 0, 0) \in (\mathfrak{e}_{8,\mathbf{H}})^C$, $\dot{1} = (0, 0, 1, 0) \in \mathfrak{P}^C$, we have

$$\alpha \dot{1} = \dot{1}.$$

Hence we see $\alpha \in ((E_7)^{\varepsilon_1, \varepsilon_2})_{\dot{1}} \subset ((E_7)_{\dot{1}})^{\varepsilon_1, \varepsilon_2} \cong (E_6)^{\varepsilon_1, \varepsilon_2}$ (Subsection 2.4). In addition, for $R^-_{\dot{E}} := (0, \dot{E}, 0, 0, 0, 0) \in (\mathfrak{e}_{8,\mathbf{H}})^C$, $\dot{E} = (E, 0, 0, 0) \in \mathfrak{P}^C$, we have

$$\alpha \dot{E} = \dot{E}.$$

Hence we see $\alpha \in ((E_6)^{\varepsilon_1, \varepsilon_2})_{\dot{E}} \subset ((E_6)_{\dot{E}})^{\varepsilon_1, \varepsilon_2} \cong (F_4)^{\varepsilon_1, \varepsilon_2}$ (Subsection 2.3). Moreover, for $R^-_{\dot{E}_i} := (0, \dot{E}_i, 0, 0, 0, 0) \in (\mathfrak{e}_{8,\mathbf{H}})^C$, $\dot{E}_i = (E_i, 0, 0, 0) \in \mathfrak{P}^C, i = 1, 2, 3$, we have

$$\alpha \dot{E}_i = \dot{E}_i, \quad i = 1, 2, 3.$$

Hence we see $\alpha \in ((F_4)^{\varepsilon_1, \varepsilon_2})_{E_1, E_2, E_3}$, so that there exists $(p_1, p_2, p_3) \in Sp(1) \times Sp(1) \times Sp(1)$ such that $\alpha = \varphi_{4, \varepsilon_i}(p_1, p_2, p_3)$ (Proposition 7.23).

Subsequently, for $R^-_{\dot{F}_k(1)} = (0, \dot{F}_k(1), 0, 0, 0, 0) \in (\mathfrak{e}_{8,\mathbf{H}})^C$, $\dot{F}_k(1) = (F_k(1), 0, 0, 0), k = 1, 2, 3$, it follows from $\alpha R^-_{\dot{F}_k(1)} = R^-_{\dot{F}_k(1)}$ that

$$\begin{aligned} R^-_{\dot{F}_k(1)} &= \alpha R^-_{\dot{F}_k(1)} = \varphi_{4, \varepsilon_i}(p_1, p_2, p_3) R^-_{\dot{F}_k(1)} = (0, \varphi_{4, \varepsilon_i}(p_1, p_2, p_3) \dot{F}_k(1), 0, 0, 0, 0) \\ &= (0, \text{diag}(p_1, p_2, p_3) \dot{F}_k(1) \text{diag}(p_1, p_2, p_3)^*, 0, 0, 0, 0). \end{aligned}$$

Hence we have $p_2\overline{p_3} = p_3\overline{p_1} = p_1\overline{p_2} = 1$, and together with $p_i \in Sp(1)$, we see $p_1 = p_2 = p_3 := p$. Thus α is of the form

$$\alpha = \varphi_{4, \varepsilon_i}(p, p, p), \quad p \in Sp(1).$$

Moreover, for $R^-_{\dot{F}_1(e_i)} = (0, \dot{F}_1(e_i), 0, 0, 0, 0) \in (\mathfrak{e}_{8, \mathbf{H}})^C$, $\dot{F}_1(e_i) = (F_1(e_i), 0, 0, 0), i = 1, 2$, it follows from $\alpha R^-_{\dot{F}_1(e_i)} = R^-_{\dot{F}_1(e_i)}$ that

$$\begin{aligned} R^-_{\dot{F}_1(e_i)} &= \alpha R^-_{\dot{F}_1(e_i)} = \varphi_{4, \varepsilon_i}(p, p, p) R^-_{\dot{F}_1(e_i)} = (0, \varphi_{4, \varepsilon_i}(p, p, p) \dot{F}_1(e_i), 0, 0, 0, 0) \\ &= (0, \text{diag}(p, p, p) \dot{F}_1(e_i) \text{diag}(p, p, p)^*, 0, 0, 0, 0). \end{aligned}$$

Hence we have $p\varepsilon_i\overline{p} = 1, i = 1, 2$, and together with $p \in Sp(1)$, we see $p^2 = 1, p \in \mathbf{R}$, that is, $p = 1$ or $p = -1$, so that we have $\alpha = 1$ when $p = 1$ or $\alpha = \gamma$ when $p = -1$. With above, we see $\text{Ker } g_8 \subset \{1, \gamma\}$ and vice versa. Thus we obtain

$$\text{Ker } g_8 = \{1, \gamma\} \cong \mathbf{Z}_2.$$

Finally, we will prove that g_8 is surjective. Since the group $E_{8, \mathbf{H}}$ is connected (Theorem 7.10) and $\text{Ker } g_8$ is discrete, and together with $\dim((\mathfrak{e}_8)^{\varepsilon_1, \varepsilon_2}) = 133 = \dim(\mathfrak{e}_{8, \mathbf{H}})$ (Lemmas 7.2, 7.21), g_8 is surjective.

Therefore, from Theorem 7.22, we have the required isomorphism

$$E_{8, \mathbf{H}} \cong E_7/\mathbf{Z}_2.$$

□

REFERENCES

- [1] C. CHEVALLEY, Theory of Lie groups I, Princeton Univ. Press, 1946.
- [2] T. MIYASHITA AND I. YOKOTA, 3-graded decompositions of exceptional Lie algebra \mathfrak{g} and group realizations of $\mathfrak{g}_{ev}, \mathfrak{g}_0$ and \mathfrak{g}_{ed} Part II, $G = E_7$, Case 1, J. Math. Kyoto Univ., 46-2(2006), 383-413.
- [3] I. YOKOTA, Non-symmetry Freudenthal's magic square, J. Fac. Sci. Shinshu Univ. 20(1985), 13-13.
- [4] I. YOKOTA, Realizations of involutive automorphisms σ and G^σ of exceptional linear Lie groups G , Part I, $G = G_2, F_4$, and E_6 , Tsukuba J. Math. 14(1990), 185-223.
- [5] I. YOKOTA, Realizations of involutive automorphisms σ and G^σ of exceptional linear Lie groups G , Part II, $G = E_7$, Tsukuba J. Math. 14(1990), 378-404.
- [6] I. YOKOTA, Realizations of involutive automorphisms σ and G^σ of exceptional linear Lie groups G , Part III, $G = E_8$, Tsukuba J. Math. 15(1991), 301-314.
- [7] I. YOKOTA, 2-graded decompositions of exceptional Lie algebra \mathfrak{g} and group realizations of $\mathfrak{g}_{ev}, \mathfrak{g}_0$ Part I, $G = G_2, F_4, E_6$, Japanese J. Math., 24-2(1998), 275-296.
- [8] I. YOKOTA, 3-graded decompositions of exceptional Lie algebra \mathfrak{g} and group realizations of $\mathfrak{g}_{ev}, \mathfrak{g}_0$ and \mathfrak{g}_{ed} Part I, $G = G_2, F_4, E_6$, J. Math. Kyoto Univ., 41-3(2001), 449-474.
- [9] I. YOKOTA, Exceptional Lie groups, arXiv:math/0902.0431v1[mathDG](2009).
- [10] I. YOKOTA, Exceptional linear simple Lie groups (in Japanese), Gendai-sugakusya, Kyoto, 1992.

1365-3 BESSHO ONSEN
 UEDA CITY
 NAGANO PREF. 386-1431
 JAPAN
 Email address: anarchybin@gmail.com