

Universal upper bound for the entropy of superconducting vortices and the Nernst effect

M. C. Diamantini,¹ C. A. Trugenberger,² and V. M. Vinokur^{3,4}

¹*NiPS Laboratory, INFN and Dipartimento di Fisica e Geologia,
University of Perugia, via A. Pascoli, I-06100 Perugia, Italy*

²*SwissScientific Technologies SA, rue du Rhone 59, CH-1204 Geneva, Switzerland*

³*Terra Quantum AG, St. Gallerstrasse 16A, CH-9400 Rorschach, Switzerland*

⁴*City College of the City University of New York, 160 Convent Ave, New York, NY 10031, USA*

We show that the entropy per vortex per layer in superconductors in external magnetic fields is bounded by the universal value $k_B \ln 2$, which explains puzzling results of recent experiments on the Nernst effect.

The Nernst effect [1, 2], the emergence of a transverse electric field produced by a longitudinal thermal gradient in the presence of a magnetic field, is currently attracting a lot of attention. The Nernst signal has a maximum as a function of temperature and magnetic field and a tail into the normal state. Recent experiments [3] revealed that, in the fluctuation region, the Nernst effect has the magnitude expected by theory. However, the peak amplitude of the Nernst signal is unexpectedly about the same in different superconductors despite the broad variance in basic superconducting parameters, and corresponds to the universal value of an entropy per vortex per layer $\approx k_B \ln 2$ [3].

Here we demonstrate that this remarkable finding is an immediate consequence of the dynamical symmetry of a superconducting ground state in an external magnetic field under the algebra $W_{1+\infty}$ of quantum area preserving the diffeomorphisms [4–6], see [7] for a review. The uniform ground states with a gap – like the Cooper pair condensate in superconductors – are incompressible in the limit of large gap, where density waves are suppressed. Their dynamical symmetry, classifying all possible configurations at the classical level, or ground states at the quantum level, is that of volume-preserving diffeomorphisms. In the presence of the applied external magnetic field the dynamical symmetry reduces to area-preserving diffeomorphisms in the plane orthogonal to the magnetic field.

This reduced case is particularly interesting since the algebra of area-preserving diffeomorphisms is a well-known extension of the 2D conformal algebra [8]. At the classical level, area-preserving diffeomorphisms are canonical transformations of a two-dimensional phase space. Normalizing all lengths by the magnetic length $\ell = \sqrt{\hbar c / e B}$, where c is the light velocity, $-e$ is the electron charge, and B is the magnetic field, and endowing the plane with coordinates z and \bar{z} with a Poisson bracket

$$\{f, g\} = i(\partial f \bar{\partial} g - \bar{\partial} f \partial g), \quad (1)$$

one describes the area-preserving diffeomorphisms $\delta z = \{\mathcal{L}, z\}$ and $\delta \bar{z} = \{\mathcal{L}, \bar{z}\}$ by generating functions $\mathcal{L}(z, \bar{z})$. The basis of generators $\mathcal{L}_{n,m} = z^n \bar{z}^m$ satisfies the classical w_∞ algebra [7]

$$\{\mathcal{L}_{n,m}, \mathcal{L}_{k,l}\} = -i(mk - nl) \mathcal{L}_{n+k-1, m+l-1}. \quad (2)$$

The operators \mathcal{L}_{nm} with $n > 0$ and $m > 0$ form two closed subalgebras, the two chiral sectors of the classical w_∞ algebra,

related by complex conjugation. Generators with both n and m negative, are descendants obtained as products of primary generators in the two fundamental chiral sectors. Finally, generators with $n = 0$ and $m = 0$ form two Abelian subalgebras.

The quantum version of this infinite-dimensional algebra is obtained by the usual substitution of Poisson brackets with quantum commutators: $i\{\cdot, \cdot\} \rightarrow [\cdot, \cdot]$. Let us denote the quantum version of $\mathcal{L}_{i-n,i}$ by V_n^i and let us first discuss one single chiral sector of the quantum algebra by restricting to positive values of i . This gives the algebra W_∞ ,

$$[V_n^i, V_m^j] = (jn - im) V_{n+m}^{i+j-1} + q(i, j, n, m) V_{n+m}^{i+j-3} + \dots + \delta^{ij} \delta_{n+m,0} c d(i, n), \quad (3)$$

where the structure constants q and d are polynomials of their arguments and the dots denote a finite number of similar terms involving the operators $V_{n+m}^{i+j-1-2k}$. The first term on the r.h.s. of Eq. (3) is the classical term (2). The remaining terms are quantum operator corrections, with the exception of the last c-number term, which represents a quantum anomaly with central charge c . All the quantum operator corrections are uniquely determined by the closure of the algebra; only the central charge c is a free parameter. If we admit also the value $i = 0$ we obtain the full algebra $W_{1+\infty}$, including the Abelian subalgebra quantum generators V_n^0 . The quantization of the full classical algebra w_∞ , involving generators in the two chiral sectors, the two Abelian subalgebras, and all their products, is then obtained as the direct product of two copies $W_{1+\infty}$ and $\overline{W}_{1+\infty}$ of opposite chirality. If the chirality breaking external magnetic field is applied, only one of the two chiral sectors survive and the dynamical symmetry reduces to $W_{1+\infty}$.

This infinite algebra has been applied to classify incompressible quantum fluids corresponding to the observed hierarchy of quantum Hall plateaus [9–11]. Not going into the details of the involved representation theory of the infinite-dimensional algebra [12], we focus on the point crucial for present purposes, namely that the two generators $V_{-1}^0 = z$ and $V_1^1 = \bar{z}$ do not commute

$$[z, \bar{z}] = [V_{-1}^0, V_1^1] = 1. \quad (4)$$

Incompressibility leads to a phase-space reduction, the coordinate space itself becomes a phase space and the two coordinates do not commute. In the presence of a magnetic field this is tantamount to the well-known non-commutativity of

the magnetic translations or of the guiding centres of charged particles in the first Landau level. Finite magnetic translations form nothing else than the group of area-preserving diffeomorphisms on the torus. This, however is just a special case. The non-commutativity follows generically from the dynamical symmetry of an incompressible quantum fluid, independently of a magnetic field [13].

As a consequence of this non-commutativity, there is a fundamental cell A_{fund} in real space, not in phase space, which is the smallest area which can accommodate a quantum degree of freedom. In superconductors, the fundamental excitations are vortices carrying one flux quantum $\Phi_0 = \pi\hbar c/e$. Vortices carrying two or more flux quanta are simply the superimposed fundamental excitations. The number of available configurations for maximally squeezed vortices is thus 2^N with $N = A/A_{\text{fund}}$ the number of fundamental cells fitting in the sample. The maximum entropy of a vortex per layer is thus the universal quantity $k_B \ln 2$, in a full accord with the Nernst effect measurements [3]. This bound is simply a universal consequence of the incompressibility of a gapped, uniform ground state in a magnetic field and is common to all possible superconductors, independently of all the details of their material parameters.

Finally, let us note that this result can be interpreted in terms of information theory. A maximally squeezed vortex is a fundamental bit. Therefore, a vortex “erased” from the sample by the transverse Nernst current takes away exactly the entropy

$k_B \ln 2$ according to the Landauer bound formula [14].

Acknowledgements

The work by V.M.V. was supported by Terra Quantum and, partially, by the NSF grant DMR-1809188 M.C.D. thanks CERN, where she completed this work, for kind hospitality.

-
- [1] K. Behnia and H. Aubin, Rep. Progr. Phys. **79**, 046502 (2016).
 - [2] A. A. Varlamov, A. Galda, and A. Glatz, Rev. Mod. Phys. **90**, 015009 (2018).
 - [3] C. W. Rischau *et al.*, Phys. Rev. Lett. **126**, 077001 (2021).
 - [4] I. Bakas, Phys. Lett. B **228**, 57 (1989).
 - [5] I. Bakas, Comm. Math. Phys. **134**, 487 (1990).
 - [6] C. N. Pope, X. Shen, and J. L. Romans, Nucl. Phys. B **339** (1990).
 - [7] X. Shen, Int. J. Mod. Phys. **A7**, 6953 (1990).
 - [8] P. Francesco, P. Mathieu & D. Sènèchal, Conformal field theory. Springer-Verlag, New York (1997).
 - [9] A. Cappelli, C. A. Trugenberger, and G. Zemba, Nucl. Phys. B **396**, 465 (1993).
 - [10] A. Cappelli, C. A. Trugenberger, and G. Zemba, Phys. Rev. Lett. **71**, 1969 (1993).
 - [11] A. Cappelli, C. A. Trugenberger, and G. Zemba, Nucl. Phys. B **448**, 470 (1995).
 - [12] V. Kac and A. Radul, Comm. Math. Phys. **157**, 429 (1993).
 - [13] M. C. Diamantini and P. Sodano, Phys. Rev. B **82**, 144515 (2010).
 - [14] R. Landauer, IBM J. Res. Develop. **5**, 183 (1961).