

BRANCHING FORMULA FOR q -TODA FUNCTIONS OF TYPE B

AYUMU HOSHINO, YUSUKE OHKUBO, AND JUN'ICHI SHIRAISHI

ABSTRACT. We present a proof of the explicit formula for the asymptotically free eigenfunctions of the B_N q -Toda operator which was conjectured by the first and third authors. This formula can be regarded as a branching formula from the B_N q -Toda eigenfunction restricted to the A_{N-1} q -Toda eigenfunctions. The proof is given by a contiguity relation of the A_{N-1} q -Toda eigenfunctions and a recursion relation of the branching coefficients.

1. INTRODUCTION

Let $f^{A_{N-1}\text{Toda}}(x|s|q)$ and $f^{B_N\text{Toda}}(x|s|q)$ be the asymptotically free eigenfunctions of the A_{N-1} and B_N q -Toda operators, respectively (Definition 2.2, Definition 2.6). Here, q is a generic parameter, and $x = (x_1, \dots, x_N)$ is an N -tuple of variables. We introduce an N -tuple of continuous parameters (or indeterminates) $s = (s_1, \dots, s_N)$, while the ordinary q -Toda functions contain a weight as a set of discrete parameters. A combinatorial explicit formula is known for the asymptotically free eigenfunctions of Macdonald's difference operator of type A [1, 2, 3], and the one of the A_{N-1} q -Toda functions $f^{A_{N-1}\text{Toda}}(x|s|q)$ can be given by taking a certain limit ($t \rightarrow 0$) of that formula. The aim of this paper is to prove the following explicit formula for $f^{B_N\text{Toda}}(x|s|q)$ in terms of $f^{A_{N-1}\text{Toda}}(x|s|q)$ that was conjectured in [4].

Theorem. 2.7. *The B_N q -Toda function $f^{B_N\text{Toda}}(x|s|q)$ is of the form*

$$\begin{aligned} & f^{B_N\text{Toda}}(x_1, \dots, x_N | s_1, \dots, s_N | q) \\ &= \sum_{\theta=(\theta_1, \dots, \theta_N) \in \mathbb{Z}_{\geq 0}^N} e_{\theta}^{B_N/A_{N-1}}(s|q) \cdot \prod_{i=1}^N x_i^{-\theta_i} \cdot f^{A_{N-1}\text{Toda}}(x_1, \dots, x_N | q^{-\theta_1} s_1, \dots, q^{-\theta_N} s_N | q), \end{aligned} \tag{1.1}$$

where we have set

$$\begin{aligned} e_{\theta}^{B_N/A_{N-1}}(s|q) &:= \prod_{k=1}^N \frac{q^{(N-k+1)\theta_k}}{(q; q)_{\theta_k} (q/s_k^2; q)_{\theta_k}} \\ &\quad \times \prod_{1 \leq i < j \leq N} \frac{1}{(qs_j/s_i; q)_{\theta_i} (q^{\theta_j - \theta_i} qs_i/s_j; q)_{\theta_i} (q/s_i s_j; q)_{\theta_i} (q/s_i s_j; q)_{\theta_j}}, \end{aligned} \tag{1.2}$$

$$(a; q)_n := \frac{(a; q)_{\infty}}{(q^n a; q)_{\infty}}, \text{ and } (a; q)_{\infty} := \prod_{k=1}^{\infty} (1 - q^{k-1} a).$$

The q -Toda system has been studied in the connection with representation theory of the quantum groups. In particular, the eigenfunctions of the q -Toda operators can be constructed by Whittaker functions in the Verma module [5, 6] and expressed via fermionic formulas [7]. Moreover, the q -Toda functions are closely related to characters of Demazure modules [8, 9, 10] and the equivariant K-theory of Laumon spaces [11, 12].

The main result (1.1) can be regarded as a branching rule for the B_N q -Toda function restricted to the A_{N-1} q -Toda eigenfunctions. The proof is given by direct calculation, in which we give a contiguity relation of $f^{A_{N-1}\text{Toda}}(x|s|q)$ (Proposition 3.1). It is an interesting problem to find similar branching formulas for q -Toda functions of other types.

This paper is organized as follows. In Section 2, we recall the definitions of the q -Toda functions and state the main theorem. The proof is given in Section 3.

2. A_{N-1} AND B_N q -TODA FUNCTIONS

First, we recall the asymptotically free eigenfunctions for A_{N-1} q -Toda operator. Let q be a generic parameter and let $s = (s_1, \dots, s_N)$ be an N -tuple of indeterminates. Set

$$\Lambda_{\mathbb{Q}(s,q)}^{A_{N-1}} = \mathbb{Q}(s,q)[[x_2/x_1, \dots, x_N/x_{N-1}]]. \quad (2.1)$$

Definition 2.1. Let $x = (x_1, \dots, x_N)$. The q -Toda operator $\mathbb{D}^{A_{N-1}\text{Toda}}(x|s|q)$ of type A acting on $\Lambda_{\mathbb{Q}(s,q)}^{A_{N-1}}$ is defined to be

$$\mathbb{D}^{A_{N-1}\text{Toda}}(x|s|q) = \sum_{i=1}^{N-1} s_i(1 - x_{i+1}/x_i)T_{q,x_i} + s_N T_{q,x_N}. \quad (2.2)$$

Here, T_{q,x_i} is the difference operator defined by

$$T_{q,x_i}f(x_1, \dots, x_N) = f(x_1, \dots, qx_i, \dots, x_N). \quad (2.3)$$

The eigenfunctions of $\mathbb{D}^{A_{N-1}\text{Toda}}(x|s|q)$ is given as follows. We use the notation in [4].

Definition 2.2 ([8]). Set

$$f^{A_{N-1}\text{Toda}}(x|s|q) = \sum_{\theta \in \mathbb{M}^{(N)}} c_N^{\text{Toda}}(\theta; s; q) \prod_{1 \leq i < j \leq N} (x_j/x_i)^{\theta_{i,j}}. \quad (2.4)$$

Here, $\mathbb{M}^{(N)} = \{\theta = (\theta_{ij})_{i,j=1}^N \mid \theta_{ij} \in \mathbb{Z}_{\geq 0}, \theta_{kl} = 0 \text{ if } k \geq l\}$ is the set of $N \times N$ strictly upper triangular matrices with non-negative integer entries, and the coefficients $c_N^{\text{Toda}}(\theta; s; q)$ are defined by

$$\begin{aligned} c_N^{\text{Toda}}(\theta; s; q) & \\ &= \prod_{k=2}^N \prod_{1 \leq i < j \leq k-1} \frac{1}{(q^{\sum_{a=k+1}^N (\theta_{i,a} - \theta_{j+1,a})} q s_{j+1}/s_i; q)_{\theta_{i,k}}} \frac{q^{\theta_{i,k}}}{(q^{\theta_{j,k} - \theta_{i,k} - \sum_{a=k+1}^N (\theta_{i,a} - \theta_{j,a})} q s_i/s_j; q)_{\theta_{i,k}}}. \end{aligned} \quad (2.5)$$

Fact 2.3 ([8, 4]). We have

$$\mathbb{D}^{A_{N-1}\text{Toda}}(x|s|q) f^{A_{N-1}\text{Toda}}(x|s|q) = \sum_{i=1}^N s_i f^{A_{N-1}\text{Toda}}(x|s|q). \quad (2.6)$$

This formula was originally proved in [8]. A combinatorial explicit formula was given for the asymptotically free eigenfunctions of the Macdonald operator in [1, 2, 3], and the formula $f^{A_{N-1}\text{Toda}}$ can also be directly obtained by taking a certain limit $t \rightarrow 0$ of that combinatorial formula [4]. As for the limit from the Macdonald functions to the Toda functions, see also [8, 9, 10].

Notation 2.4. We introduce

$$d_N^{\text{Toda}}((\theta_{i,n})_{1 \leq i \leq N-1}; (s_i)_{1 \leq i \leq N}; q) := \frac{c_N^{\text{Toda}}((\theta_{i,j})_{1 \leq i < j \leq N}; (s_i)_{1 \leq i \leq N} | q)}{c_{N-1}^{\text{Toda}}((\theta_{i,j})_{1 \leq i < j \leq N-1}; (q^{-\theta_{i,n}} s_i)_{1 \leq i \leq N-1} | q)} \quad (N \geq 2). \quad (2.7)$$

Then, d_N^{Toda} is of the form

$$\begin{aligned} d_N^{\text{Toda}}((\theta_i)_{1 \leq i \leq N-1}; (s_i)_{1 \leq i \leq N}; q) & \\ &= \prod_{i=1}^{N-1} \frac{1}{(q; q)_{\theta_i}} \frac{q^{\theta_i}}{(q s_N/s_i; q)_{\theta_i}} \prod_{1 \leq i < j \leq N-1} \frac{1}{(q s_j/s_i; q)_{\theta_i}} \frac{q^{\theta_i}}{(q^{\theta_j - \theta_i + 1} s_i/s_j; q)_{\theta_i}}, \end{aligned} \quad (2.8)$$

and the A_{N-1} q -Toda function can be expressed as

$$\begin{aligned} f^{A_{N-1}\text{Toda}}(x|s|q) & \\ &= \sum_{\theta = (\theta_1, \dots, \theta_{N-1}) \in \mathbb{Z}_{\geq 0}^{N-1}} d_N^{\text{Toda}}(\theta; s; q) \prod_{i=1}^{N-1} (x_N/x_i)^{\theta_i} \cdot f^{A_{N-2}\text{Toda}}(x | (q^{-\theta_i} s_i)_{1 \leq i \leq N-1} | q). \end{aligned} \quad (2.9)$$

Although (2.9) follows from the case of the Macdonald functions, we can also prove (2.9) in a similar manner to Section 3.

Now, we turn to the case of type B. Set

$$\Lambda_{\mathbb{Q}(s,q)}^{B_N} = \mathbb{Q}(s,q)[[x_2/x_1, \dots, x_N/x_{N-1}, 1/x_N]]. \quad (2.10)$$

Definition 2.5. Define the B_N q -Toda operator $D^{B_N \text{Toda}}(x|s|q)$ acting on $\Lambda_{\mathbb{Q}(s,q)}^{B_N}$ by

$$\begin{aligned} D^{B_N \text{Toda}}(x|s|q) &= \sum_{i=1}^{N-1} s_i(1 - x_{i+1}/x_i)T_{q,x_i} + s_N(1 - 1/x_N)T_{q,x_N} \\ &\quad + s_1^{-1}T_{q,x_1}^{-1} + \sum_{i=2}^N s_i^{-1}(1 - x_i/x_{i-1})T_{q,x_i}^{-1}. \end{aligned} \quad (2.11)$$

This operator can be obtained by the limit of the B_N Macdonald operator [4]. As for the description of the q -Toda operators by the quantum groups, see [5, 6, 7].

Definition 2.6. The asymptotically free eigenfunction $f^{B_N \text{Toda}}(x|s|q) \in \Lambda_{\mathbb{Q}(s,q)}^{B_N}$ of the B_N q -Toda operator is defined by

$$D^{B_N \text{Toda}}(x|s|q)f^{B_N \text{Toda}}(x|s|q) = \sum_{i=1}^N (s_i + s_i^{-1})f^{B_N \text{Toda}}(x|s|q), \quad (2.12)$$

$$\left[f^{B_N \text{Toda}}(x|s|q) \right]_{x,1} = 1. \quad (2.13)$$

Here, $[\]_{x,1}$ means the constant term with respect to x_i 's.

Note that $f^{B_N \text{Toda}}(x|s|q)$ is uniquely determined. We obtain an explicit formula for the B_N q -Toda function $f^{B_N \text{Toda}}(x|s|q)$ in terms of the A_{N-1} q -Toda functions $f^{A_{N-1} \text{Toda}}(x|s|q)$.

Theorem 2.7. The B_N q -Toda function $f^{B_N \text{Toda}}(x|s|q)$ satisfies the branching formula

$$\begin{aligned} &f^{B_N \text{Toda}}(x_1, \dots, x_N|s_1, \dots, s_N|q) \\ &= \sum_{\theta=(\theta_1, \dots, \theta_N) \in \mathbb{Z}_{\geq 0}^N} e_{\theta}^{B_N/A_{N-1}}(s|q) \cdot \prod_{i=1}^N x_i^{-\theta_i} \cdot f^{A_{N-1} \text{Toda}}(x_1, \dots, x_N|q^{-\theta_1}s_1, \dots, q^{-\theta_N}s_N|q), \end{aligned} \quad (2.14)$$

where we have set

$$\begin{aligned} e_{\theta}^{B_N/A_{N-1}}(s|q) &:= \prod_{k=1}^N \frac{q^{(N-k+1)\theta_k}}{(q; q)_{\theta_k} (q/s_k^2; q)_{\theta_k}} \\ &\quad \times \prod_{1 \leq i < j \leq N} \frac{1}{(qs_j/s_i; q)_{\theta_i} (q^{\theta_j - \theta_i} qs_i/s_j; q)_{\theta_i} (q/s_i s_j; q)_{\theta_i} (q/s_i s_j; q)_{\theta_j}}, \end{aligned} \quad (2.15)$$

$(a; q)_n := \frac{(a; q)_{\infty}}{(q^n a; q)_{\infty}}$, and $(a; q)_{\infty} := \prod_{k=1}^{\infty} (1 - q^{k-1}a)$. Note that the constant term of $f^{A_{N-1} \text{Toda}}$ is 1. Hence, the constant term of (2.14) is also 1.

This formula was conjectured in [4]. The proof is given in the next subsection.

Remark 2.8. The region of convergence of $f^{A_{N-1} \text{Toda}}$ can be derived from the case of the Macdonald functions (Proposition 6.1 in [2]) by taking the limit $t \rightarrow 0$. It is an interesting problem to consider the convergence of the formula (2.14).

3. PROOF OF THEOREM 2.7

In this section, we prove Theorem 2.7. First we give the following relation of the q -Toda functions of type A.

Proposition 3.1. *The q -Toda functions of type A satisfy the contiguity relation*

$$\begin{aligned} & f^{A_{N-1}\text{Toda}}(x_1, \dots, x_{N-1}, qx_N | s | q) \\ &= \sum_{k=1}^N (-1)^{N-k} \frac{q^{N-k} \prod_{i=k+1}^{N-1} s_i/s_k}{\prod_{i=k+1}^N (1 - s_i/s_k)(1 - qs_i/s_k)} (x_N/x_k) f^{A_{N-1}\text{Toda}}(x_1, \dots, x_N | q^{-\varepsilon_k} \cdot s | q). \end{aligned} \quad (3.1)$$

Here, we used the notation

$$q^{\pm \varepsilon_i} \cdot s = (s_1, \dots, s_{i-1}, q^{\pm 1} s_i, s_{i+1}, \dots, s_N). \quad (3.2)$$

Proof. First, we show the following equation of the rational functions of a_i and s_i :

$$\prod_{i=1}^{N-1} a_i = \sum_{k=1}^N (s_k/s_N) \frac{\prod_{i=1}^{N-1} (1 - a_i s_k/s_i)}{\prod_{\substack{1 \leq i \leq N \\ i \neq k}} (1 - s_k/s_i)}. \quad (3.3)$$

Regarding s_i 's in the RHS as complex variables, we set

$$F(s) := \sum_{k=1}^N (s_k/s_N) \frac{\prod_{i=1}^{N-1} (1 - a_i s_k/s_i)}{\prod_{\substack{1 \leq i \leq N \\ i \neq k}} (1 - s_k/s_i)} = \sum_{k=1}^N \frac{\prod_{i=1}^{N-1} (s_i - a_i s_k)}{\prod_{\substack{1 \leq i \leq N \\ i \neq k}} (s_i - s_k)}. \quad (3.4)$$

For any $\ell = 1, \dots, N$, the residue at $s_\ell = s_{\ell'}$ ($\ell' \neq \ell$) is

$$\begin{aligned} \text{Res}_{s_\ell = s_{\ell'}} F(s) &= \lim_{s_\ell \rightarrow s_{\ell'}} F(s)(s_\ell - s_{\ell'}) \\ &= \lim_{s_\ell \rightarrow s_{\ell'}} \left(-\frac{\prod_{i=1}^{N-1} (s_i - a_i s_\ell)}{\prod_{i \neq \ell, \ell'} (s_i - s_\ell)} + \frac{\prod_{i=1}^{N-1} (s_i - a_i s_{\ell'})}{\prod_{i \neq \ell, \ell'} (s_i - s_{\ell'})} \right) \\ &= 0. \end{aligned} \quad (3.5)$$

Hence $F(s)$ is regular on the whole complex plane with respect to each s_ℓ , and it is clear that $F(s)$ is bounded. This indicates that $F(s)$ is a constant function. By the specialization $s_i = a_{i-1}^{-1} a_{i-2}^{-1} \cdots a_1^{-1} s_1$ ($i = 2, \dots, N$), we have

$$\begin{aligned} F(s) &= \sum_{k=1}^N (a_{N-1} a_{N-2} \cdots a_k) \frac{\prod_{i=1}^{N-1} \left(1 - \frac{a_i a_{i-1} \cdots a_1}{a_{k-1} a_{k-2} \cdots a_1} \right)}{\prod_{\substack{1 \leq i \leq N \\ i \neq k}} \left(1 - \frac{a_{i-1} a_{i-2} \cdots a_1}{a_{k-1} a_{k-2} \cdots a_1} \right)} \\ &= a_{N-1} a_{N-2} \cdots a_1. \end{aligned} \quad (3.6)$$

This gives (3.3).

Substituting $a_i = q^{\theta_i}$ into (3.3) yields

$$\begin{aligned} \prod_{i=1}^{N-1} q^{\theta_i} &= \sum_{k=1}^{N-1} (-1)^{N-k} \frac{q^{N-k} \prod_{i=k+1}^{N-1} s_i/s_k}{\prod_{i=k+1}^N (1 - s_i/s_k)(1 - qs_i/s_k)} \frac{d_N^{\text{Toda}}(\theta_1, \dots, \theta_k - 1, \dots, \theta_{N-1} | q^{-\varepsilon_k} \cdot s)}{d_N^{\text{Toda}}(\theta_1, \dots, \theta_{N-1} | s)} \\ &\quad + \frac{d_N^{\text{Toda}}(\theta_1, \dots, \theta_{N-1} | q^{-\varepsilon_N} \cdot s)}{d_N^{\text{Toda}}(\theta_1, \dots, \theta_{N-1} | s)}. \end{aligned} \quad (3.7)$$

By (2.9) and (3.7), we obtain the formula (3.1). \square

Proposition 3.2. *The branching coefficients $e_\theta^{B_N/A_{N-1}}(s|q)$ satisfy the recursion relation*

$$\sum_{i=1}^N \left((1 - q^{-\theta_i}) s_i + (1 - q^{\theta_i}) s_i^{-1} \right) e_\theta^{B_N/A_{N-1}}(s|q) \quad (3.8)$$

$$= \sum_{k=1}^N s_N \frac{(-1)^{N-k+1} q^{-\theta_N + \delta_{k,N}} q^{N-k} \prod_{i=k+1}^{N-1} (q^{-\theta_i + \theta_{k-1}} s_i / s_k)}{\prod_{i=k+1}^N (1 - q^{-\theta_i + \theta_{k-1}} s_i / s_k) (1 - q q^{-\theta_i + \theta_{k-1}} s_i / s_k)} e_{(\theta_1, \dots, \theta_{k-1}, \dots, \theta_N)}^{B_N/A_{N-1}}(s|q).$$

Proof. By substituting (2.15) into (3.8), it can be shown that (3.8) is equivalent to

$$\sum_{i=1}^N \left((1 - q^{-\theta_i}) s_i + (1 - q^{\theta_i}) s_i^{-1} \right) = - \sum_{k=1}^N q^{-\theta_k} s_k \frac{\prod_{i=1}^N (1 - q^{\theta_k} s_i / s_k) (1 - q^{\theta_k} / s_i s_k)}{\prod_{i \neq k} (1 - q^{\theta_k - \theta_i} s_i / s_k) (1 - q^{\theta_k + \theta_i} / s_i s_k)}. \quad (3.9)$$

By replacing q^{θ_i} with generic parameters Q_i and shifting s_i to $Q_i s_i$, the equation (3.9) becomes

$$\sum_{i=1}^N \left((1 - Q_i) s_i + (1 - Q_i^{-1}) s_i^{-1} \right) = \sum_{k=1}^N s_k \frac{\prod_{i=1}^N (1 - Q_i s_i / s_k) (1 - Q_i^{-1} / s_i s_k)}{\prod_{i \neq k} (1 - s_i / s_k) (1 - 1 / s_i s_k)}. \quad (3.10)$$

The proof is completed by showing this equation. Regarding s_i 's as complex variables, we define the function

$$F(s) := \sum_{k=1}^N s_k \frac{\prod_{i=1}^N (1 - Q_i s_i / s_k) (1 - Q_i^{-1} / s_i s_k)}{\prod_{i \neq k} (1 - s_i / s_k) (1 - 1 / s_i s_k)}. \quad (3.11)$$

A direct calculation shows that the residue at $s_\ell = s_{\ell'}^{\pm 1}$ ($\ell \neq \ell'$) is

$$\operatorname{Res}_{s_\ell = s_{\ell'}^{\pm 1}} F(s) = \lim_{s_\ell \rightarrow s_{\ell'}^{\pm 1}} F(s) (s_\ell - s_{\ell'}) = 0 \quad (3.12)$$

and these singularities are removable. Hence, $F(s)$ is a regular with respect to each variable s_ℓ ($\ell = 1, \dots, n$) on the complex plane except for the origin 0 (and ∞). Therefore, for arbitrary ℓ , the function $F(s)$ can be given by the Laurent series on $0 < |s_\ell| < \infty$

$$F(s) = \sum_{i \in \mathbb{Z}} C_i s_\ell^i, \quad (3.13)$$

where C_i is a function of $s_1, \dots, s_{\ell-1}, s_{\ell+1}, \dots, s_N$. Since the orders of the poles at $s_\ell = 0$ and $s_\ell = \infty$ are at most 1, we have $C_i = 0$ ($i < -1$ or $i > 1$). It can be shown that the residues at $s_\ell = 0$ and $s_\ell = \infty$ are

$$C_{-1} = \operatorname{Res}_{s_\ell=0} F(s) = 1 - Q_\ell^{-1}, \quad (3.14)$$

$$C_1 = \operatorname{Res}_{s_\ell=\infty} F(s) = 1 - Q_\ell. \quad (3.15)$$

Therefore, with a constant \tilde{C}_0 independent of s_i 's, we can write

$$F(s) = \sum_{i=1}^N \left((1 - Q_i) s_i + (1 - Q_i^{-1}) s_i^{-1} \right) + \tilde{C}_0. \quad (3.16)$$

Furthermore, we obtain

$$\begin{aligned} \tilde{C}_0 &= F(\sqrt{Q_1}^{-1}, \sqrt{Q_2}^{-1}, \dots, \sqrt{Q_N}^{-1}) \\ &= \sum_{k=1}^N \sqrt{Q_k}^{-1} \frac{\prod_{i=1}^N (1 - \sqrt{Q_i Q_k}) (1 - \sqrt{Q_k / Q_i})}{\prod_{i \neq k} (1 - \sqrt{Q_k / Q_i}) (1 - \sqrt{Q_i Q_k})} \\ &= 0. \end{aligned} \quad (3.17)$$

This gives (3.10). \square

Proof of Theorem 2.7. The action of $D^{B_N \text{Toda}}(x|s|q)$ on the right hand side of (2.14) gives

$$\begin{aligned} &D^{B_N \text{Toda}}(x|s|q) \text{ (RHS of (2.14))} \\ &= \sum_{\theta \in \mathbb{Z}_{\geq 0}^N} e_\theta^{B_N/A_{N-1}}(s|q) \prod_{i=1}^N x_i^{-\theta_i} \cdot \left\{ D^{A_{N-1} \text{Toda}}(x|s|q) - q^{-\theta_N} s_N / x_N T_{q, x_N} \right\} \end{aligned}$$

$$\begin{aligned}
& + \mathbb{D}^{A_{N-1}\text{Toda}}((x_{N-i+1}^{-1})_{i=1}^N | (q^{\theta_{N-i+1}} s_{N-i+1}^{-1})_{i=1}^N | q) \Big\} f^{A_{N-1}\text{Toda}}(x | (q^{-\theta_i} s_i)_{i=1}^N | q) \\
& = \sum_{\theta \in \mathbb{Z}_{\geq 0}^N} e_{\theta}^{B_N/A_{N-1}}(s|q) \prod_{i=1}^N x_i^{-\theta_i} \cdot \left\{ \sum_{i=1}^N q^{-\theta_i} s_i + \sum_{i=1}^N q^{\theta_i} s_i^{-1} - q^{-\theta_N} s_N / x_N T_{q, x_N} \right\} \\
& \quad \times f^{A_{N-1}\text{Toda}}(x | (q^{-\theta_i} s_i) | q).
\end{aligned} \tag{3.18}$$

Here, we used Fact 2.3 and the symmetry

$$f^{A_{N-1}\text{Toda}}(x|s|q) = f^{A_{N-1}\text{Toda}}((x_{N-i+1}^{-1})_{1 \leq i \leq N} | (s_{N-i+1}^{-1})_{1 \leq i \leq N} | q). \tag{3.19}$$

By Proposition 3.1, we have

$$\begin{aligned}
& \mathbb{D}^{B_N\text{Toda}}(x|s|q) \text{ (RHS of (2.14))} \\
& = \sum_{\theta \in \mathbb{Z}_{\geq 0}^N} e_{\theta}^{B_N/A_{N-1}}(s|q) \prod_{i=1}^N x_i^{-\theta_i} \cdot \left\{ \sum_{i=1}^N (q^{-\theta_i} s_i + q^{\theta_i} s_i^{-1}) f^{A_{N-1}\text{Toda}}(x | (q^{-\theta_l} s_l) | q) \right. \\
& \quad - q^{-\theta_N} \sum_{k=1}^N (-1)^{N-k} (s_N / x_k) \frac{q^{N-k} \prod_{i=k+1}^{N-1} (q^{-\theta_i + \theta_k} s_i / s_k)}{\prod_{i=k+1}^N (1 - q^{-\theta_i + \theta_k} s_i / s_k) (1 - q q^{-\theta_i + \theta_k} s_i / s_k)} \\
& \quad \left. \times f^{A_{N-1}\text{Toda}}(x | q^{-\varepsilon_k} \cdot (q^{-\theta_l} s_l)_{1 \leq l \leq N} | q) \right\} \\
& = \sum_{\theta \in \mathbb{Z}_{\geq 0}^N} \prod_{i=1}^N x_i^{-\theta_i} \cdot \left\{ \sum_{i=1}^N (q^{-\theta_i} s_i + q^{\theta_i} s_i^{-1}) e_{\theta}^{B_N/A_{N-1}}(s|q) \right. \\
& \quad + \sum_{k=1}^N s_N \frac{(-1)^{N-k+1} q^{N-k-\theta_N + \delta_{k,N}} \prod_{i=k+1}^{N-1} (q^{-\theta_i + \theta_k - 1} s_i / s_k)}{\prod_{i=k+1}^N (1 - q^{-\theta_i + \theta_k - 1} s_i / s_k) (1 - q q^{-\theta_i + \theta_k - 1} s_i / s_k)} e_{(\theta_1, \dots, \theta_k - 1, \dots, \theta_N)}^{B_N/A_{N-1}}(s|q) \Big\} \\
& \quad \times f^{A_{N-1}\text{Toda}}(x | (q^{-\theta_l} s_l)_{1 \leq l \leq N} | q),
\end{aligned} \tag{3.20}$$

where we have used that $e_{\theta}^{B_N/A_{N-1}} = 0$ if $\theta_j = -1$ for some j . Proposition 3.2 shows that this is equal to $\sum_{i=1}^N (s_i + s_i^{-1}) \cdot$ (RHS of (2.14)). This completes the proof. \square

ACKNOWLEDGMENT

The authors would like to thank B. Feigin, M. Fukuda, M. Noumi, and L. Rybnikov for valuable discussions. The researches of J.S. and H.A. are partially supported by JSPS KAKENHI (J.S: 19K03512, H.A: 19K03530). Y.O. is partially supported by Grant-in-Aid for JSPS Research Fellow (18J00754).

REFERENCES

- [1] J. Shiraishi, "A conjecture about raising operators for Macdonald polynomials," *Lett. Math. Phys.* **73** no. 1, (2005) 71–81 [arXiv:math/0503727](#) [[math.QA](#)].
- [2] M. Noumi and J. Shiraishi, "A direct approach to the bispectral problem for the Ruijsenaars-Macdonald q-difference operators," [arXiv:1206.5364](#) [[math.QA](#)].
- [3] A. Braverman, M. Finkelberg, and J. Shiraishi, "Macdonald polynomials, Laumon spaces and perverse coherent sheaves," *Perspectives in Representation Theory, Contemp. Math* **610** (2014) 23–41.
- [4] A. Hoshino and J. Shiraishi, "Branching Rules for Koornwinder Polynomials with One Column Diagrams and Matrix Inversions," *SIGMA* **16** (2020) 084 [arXiv:2002.02148](#) [[math.QA](#)].
- [5] P. Etingof, "Whittaker functions on quantum groups and q-deformed Toda operators," [arXiv:math/9901053](#) [[math.QA](#)].
- [6] A. Sevostyanov, "Quantum deformation of Whittaker modules and the Toda lattice," *Duke Math. J.* **105** no. 2, (2000) 211–238 [arXiv:math/9905128](#) [[math.QA](#)].
- [7] B. Feigin, E. Feigin, M. Jimbo, T. Miwa, and E. Mukhin, "Fermionic formulas for eigenfunctions of the difference Toda Hamiltonian," *Lett. Math. Phys.* **88** no. 1-3, (2009) 39 [arXiv:0812.2306](#) [[math.QA](#)].
- [8] A. Gerasimov, D. Lebedev, and S. Oblezin, "On q-Deformed $\mathfrak{gl}_{\ell+1}$ -Whittaker Function I," *Commun. Math. Phys.* **294** no. 1, (2010) 97–119 [arXiv:0803.0145](#) [[math.RT](#)].
- [9] A. Gerasimov, D. Lebedev, and S. Oblezin, "On q-Deformed $\mathfrak{gl}_{\ell+1}$ -Whittaker Function II," *Commun. Math. Phys.* **294** no. 1, (2010) 121–143 [arXiv:0803.0970](#) [[math.RT](#)].

- [10] A. Gerasimov, D. Lebedev, and S. Oblezin, “On q -Deformed $\mathfrak{gl}_{\ell+1}$ -Whittaker Function III,” *Lett. Math. Phys.* **97** no. 1, (2011) 1–24 [arXiv:0805.3754](#) [[math.RT](#)].
- [11] A. Givental and Y.-P. Lee, “Quantum K-theory on flag manifolds, finite-difference Toda lattices and quantum groups,” *Inventiones mathematicae* **151** no. 1, (2003) 193–219 [arXiv:math/0108105](#) [[math.AG](#)].
- [12] A. Braverman and M. Finkelberg, “Finite difference quantum Toda lattice via equivariant K-theory,” *Transform. Groups* **10** no. 3-4, (2005) 363–386 [arXiv:math/0503456](#).

AH: HIROSHIMA INSTITUTE OF TECHNOLOGY, 2-1-1 MIYAKE, HIROSHIMA 731-5193, JAPAN
Email address: a.hoshino.c3@it-hiroshima.ac.jp

YO: DAICHI UNIVERSITY OF PHARMACY, 22-1 TAMAGAWA-CHO, MINAMI-KU, FUKUOKA 815-8511, JAPAN
Email address: yusuke.ohkubo.math@gmail.com

JS: GRADUATE SCHOOL OF MATHEMATICAL SCIENCES, THE UNIVERSITY OF TOKYO, KOMABA, TOKYO 153-8914, JAPAN
Email address: shiraish@ms.u-tokyo.ac.jp