

Moments of Markovian growth-collapse processes

Nicolas Privault

Division of Mathematical Sciences
School of Physical and Mathematical Sciences
Nanyang Technological University
21 Nanyang Link
Singapore 637371

January 5, 2022

Abstract

We apply general moment identities for Poisson stochastic integrals with random integrands to the computation of the moments of Markovian growth-collapse processes. This extends existing formulas for mean and variance available in the literature to closed form moments expressions of all orders. In comparison with other methods based on differential equations, our approach yields polynomial expressions in the time parameter. We also treat the case of the associated embedded chain.

Key words: Growth-collapse processes, Poisson shot noise, uniform cut-off, stochastic integrals with jumps, moments, cumulants.

Mathematics Subject Classification (2010): 60J27; 60G55; 60J22.

1 Introduction

Markovian growth-collapse processes, see [Eliazar and Klafter \(2004\)](#), are piecewise-deterministic Markov processes ([Davis \(1984\)](#)), that grow in between random jump times at which they may randomly crash. Growth-collapse processes are used in e.g. earth sciences and physics, and they have also been recently applied to the study of crypto-currencies, see [Frolkova and Mandjes \(2019\)](#).

The computation of moments of growth-collapse processes has been the object of several approaches, see [Boxma et al. \(2006\)](#) for the use of conditional distributions for the computa-

tion of mean and variance, and [Daw and Pender \(2020\)](#) for moment expressions of all orders using the solution of differential equations by matrix exponentials.

In this paper, we apply general moment identities written as sums over partitions for Poisson stochastic integrals with random integrands, see [Privault \(2009; 2012a;b; 2016\)](#), to the computation of the moments of growth-collapse processes. In particular, we obtain closed-form moment expressions which are polynomial in time in the case of uniformly distributed cut-off rates.

Let $(N_t)_{t \in \mathbb{R}_+}$ denote a standard Poisson process with intensity λ on \mathbb{R}_+ , and consider a process $(Y_t)_{t \in \mathbb{R}_+}$ of the form

$$Y_t = \int_0^t f(N_{s-}) dN_s, \quad t \geq 0.$$

As the left limit $f(N_{s-})$ is predictable with respect to the filtration $(\mathcal{F}_s)_{s \in \mathbb{R}_+}$ generated by $(N_s)_{s \in \mathbb{R}_+}$ the mean of Y_t can be computed from the smoothing lemma, see e.g. Theorem 9.2.1 in [Brémaud \(1999\)](#), as

$$\mathbb{E}[Y_t] = \lambda \mathbb{E} \left[\int_0^t f(N_{s-}) ds \right] = \lambda \int_0^t \mathbb{E}[f(N_{s-})] ds, \quad t \geq 0.$$

This calculation does not apply however to the process $f(N_s)$ which is only adapted and not predictable with respect to the filtration $(\mathcal{F}_s)_{s \in \mathbb{R}_+}$. In this case we may apply the Slivnyak-Mecke formula, see [Slivnyak \(1962\)](#), [Mecke \(1967\)](#), to obtain

$$\mathbb{E}[Y_t] = \lambda \mathbb{E} \left[\int_0^t \varepsilon_s^+ f(N_s) ds \right] = \lambda \mathbb{E} \left[\int_0^t f(1 + N_s) ds \right], \quad t \geq 0,$$

where ε_s^+ denotes the operator that adds one jump at the location $s \geq 0$ to the Poisson process path.

In order to compute higher order moments we will apply a nonlinear extension of the Slivnyak-Mecke identity, see Proposition 2.1 below, which allows us to express the moments of Poisson stochastic integrals as a sum of multiple integrals with respect to the intensity of the Poisson process over partitions. In Section 3 we consider the computation of moments of jump processes of the form

$$Y_t = \sum_{k=1}^{N_t} g(T_k, k, N_t) = \int_0^t g(s, N_s, N_t) dN_s,$$

where $(T_k)_{k \geq 1}$ denotes the sequence of jump times of the Poisson process $(N_t)_{t \in \mathbb{R}_+}$, see Proposition 3.1 and its Corollary 3.2. Those identities are then specialized in Section 4 to the case of uniform cut-off distributions, for processes of the form

$$Y_t = \sum_{k=1}^{N_t} f_k(T_k)(1 - U_k) \prod_{l=k+1}^{N_t} U_l, \quad t \in \mathbb{R}_+,$$

where $(U_k)_{k \geq 1}$ is an i.i.d. uniform sequence on $[0, 1]$, independent of the standard Poisson process $(N_t)_{t \in \mathbb{R}_+}$, see Corollary 4.1.

In particular, in Proposition 5.1 we obtain the closed form polynomial expression

$$\mathbb{E}[(X_t)^n] = \frac{(n+1)!}{\lambda^n} \sum_{k=0}^n (-1)^k (k+1)^{n-1} \binom{n}{k} e^{-k\lambda t/(k+1)}, \quad t \in \mathbb{R}_+, \quad (1.1)$$

for the moments of all orders $n \geq 0$ of the growth-collapse process

$$X_t = t - \sum_{k=1}^{N_t} T_k(1 - U_k) \prod_{l=k+1}^{N_t} U_l, \quad t \in \mathbb{R}_+,$$

where $(U_k)_{k \geq 1}$ is an i.i.d. uniform sequence on $[0, 1]$. This result extends Theorems 4 and 5 as well as Corollary 1 of Boxma et al. (2006) from mean and variance to higher moments of all orders, and provides a closed form alternative to Corollary 4 in Daw and Pender (2020) which uses matrix exponentials. The expression (1.1) immediately yields the asymptotic moments

$$\lim_{t \rightarrow \infty} \mathbb{E}[(X_t)^n] = \frac{(n+1)!}{\lambda^n}, \quad n \geq 1,$$

which recover the gamma stationary distribution of $(X_t)_{t \in \mathbb{R}_+}$ with shape parameter 2, see Theorem 3 in Boxma et al. (2006).

Finally, in Section 6 we show that our approach can also be applied to discrete-time embedded processes of the form

$$Y(m) = \sum_{k=1}^m g(T_k, k, m) = \int_0^{T_m} g(s, N_s, m) dN_s, \quad m \geq 1,$$

see Corollaries 6.2-6.3, and to the embedded growth-collapse chain

$$X(m) = T_m - \sum_{k=1}^m T_k(1 - U_k) \prod_{l=k+1}^m U_l, \quad m \geq 1,$$

where $(U_k)_{k \geq 1}$ is an i.i.d. uniform sequence on $[0, 1]$, see Corollaries 6.4-6.5. This recovers Theorem 7 stated for mean and variance in Boxma et al. (2006), and provides moment expressions of all orders.

We proceed as follows. In Section 2 we review the derivation of moment identities for stochastic integrals using sums over partitions, and in Section 3 we apply them to the moments of jump processes driven by a Poisson process. Those expressions are then specialized as closed form polynomial identities in Section 4 in the case of uniform cut-off distributions. The moments of growth-collapse processes are considered in Section 5, and the case of embedded chains is treated in Section 6.

2 Moment identities for Poisson stochastic integrals

In this section we review the computation of moments of Poisson stochastic integrals with random integrands using sums over partitions, see Proposition 3.1 in Privault (2012a). Consider a Poisson process $(N_t)_{t \in \mathbb{R}_+}$ constructed as $N_t = \omega([0, t])$, where $\omega(ds)$ is a Poisson random measure of intensity $\lambda(ds)$, with sequence $(T_k)_{k \geq 1}$ of jump times. For any $s_1, \dots, s_k \in \mathbb{R}_+$, we let $\epsilon_{s_1}^+ \cdots \epsilon_{s_k}^+$ denote the operator

$$(\epsilon_{s_1}^+ \cdots \epsilon_{s_k}^+ F)(\omega) = F(\omega \cup \{s_1, \dots, s_k\})$$

acting on random variables F by addition of points at locations s_1, \dots, s_k to the point process $\omega(dx)$. For example, if F takes the form $F = f(N_{t_1}, \dots, N_{t_n})$, then we have

$$\epsilon_{s_1}^+ \cdots \epsilon_{s_k}^+ F = f(N_{t_1} + \#\{k : s_k \leq t_1\}, \dots, N_{t_n} + \#\{k : s_k \leq t_n\}).$$

The following moment identity, see Proposition 3.1 in Privault (2012a) and Theorem 1 in Privault (2016), uses sums over partitions $\{\pi_1, \dots, \pi_k\}$ of $\{1, \dots, n\}$, and applies to random integrands $u : \mathbb{R}_+ \times \Omega \rightarrow \mathbb{R}$.

Proposition 2.1 *Let $(u_s(\omega))_{s \in \mathbb{R}_+}$ denote a stochastic process indexed by $s \in \mathbb{R}_+$. For any $n \geq 1$, we have*

$$\begin{aligned} & \mathbb{E} \left[\left(\int_0^T u_s dN_s \right)^n \right] \\ &= \sum_{k=1}^n \sum_{\pi_1 \cup \dots \cup \pi_k = \{1, \dots, n\}} \mathbb{E} \left[\int_0^t \cdots \int_0^t \epsilon_{s_1}^+ \cdots \epsilon_{s_k}^+ (u^{|\pi_1|}(s_1, \omega) \cdots u^{|\pi_k|}(s_k, \omega)) \lambda(ds_1) \cdots \lambda(ds_k) \right], \end{aligned} \tag{2.1}$$

where the power $|\pi_i|$ denotes the cardinality of the subset π_i and the above sum runs over all partitions π_1, \dots, π_k of $\{1, \dots, n\}$.

In the sequel we will frequently use the equivalent combinatorial expressions

$$\begin{aligned} \sum_{k=1}^n \sum_{\pi_1 \cup \dots \cup \pi_k = \{1, \dots, n\}} f_k(|\pi_1|, \dots, |\pi_k|) &= \sum_{k=1}^n \frac{n!}{k!} \sum_{\substack{p_1 + \dots + p_k = n \\ p_1, \dots, p_k \geq 1}} \frac{f_k(p_1, \dots, p_k)}{p_1! \cdots p_k!} \\ &= \sum_{k=1}^n \frac{n!}{k!} \sum_{q_0=0 < q_1 < \dots < q_k = n} \frac{f_k(q_1 - q_0, \dots, q_k - q_{k-1})}{(q_1 - q_0)! \cdots (q_k - q_{k-1})!} \end{aligned}$$

for f_k a function on \mathbb{N}^k , $k = 1, \dots, n$. In particular, for $x_1, \dots, x_n \in \mathbb{R}$ and $f_k(p_1, \dots, p_k) = x_{p_1} \cdots x_{p_k}$ this yields the Bell polynomial of order $n \geq 1$ as

$$\begin{aligned} B_n(x_1, \dots, x_n) &= \sum_{k=1}^n \frac{n!}{k!} \sum_{\substack{p_1 + \dots + p_k = n \\ p_1 \geq 1, \dots, p_k \geq 1}} \frac{x_{p_1} \cdots x_{p_k}}{p_1! \cdots p_k!} \\ &= \sum_{k=0}^n \sum_{\pi_1 \cup \dots \cup \pi_k = \{1, \dots, n\}} x_{|\pi_1|} \cdots x_{|\pi_k|}. \end{aligned}$$

We will also use the relation $\mathbb{E}[X^n] = B_n(\kappa_X^{(1)}, \dots, \kappa_X^{(n)})$ between the moments $\mathbb{E}[X^n]$ and the cumulants $\kappa_X^{(n)}$ of a random variable X , and the inversion relation

$$\kappa_n^X = \sum_{k=1}^n (k-1)! (-1)^{k-1} \sum_{\pi_1 \cup \dots \cup \pi_k = \{1, \dots, n\}} \mathbb{E}[X^{|\pi_1|}] \cdots \mathbb{E}[X^{|\pi_k|}], \quad n \geq 1, \quad (2.2)$$

see Theorem 1 of [Lukacs \(1955\)](#), or [Leonov and Shiryaev \(1959\)](#).

Shot noise processes

Before moving to the setting of Markovian growth-collapse processes, we use the case of Poisson shot noise processes as an illustration for the result of [Proposition 2.1](#). Consider a shot noise process $(S_t)_{t \in \mathbb{R}_+}$ of the form

$$S_t = \sum_{k=1}^{N_t} J_k g(T_k, t) = \sum_{k=1}^{N_t} J_{N_{T_k}} g(T_k, t) = \int_0^t J_{N_s} g(s, t) dN_s, \quad t \in \mathbb{R}_+,$$

where $(J_k)_{k \geq 0}$ is a sequence of i.i.d. random variables admitting moments of all orders, and $g(\cdot, \cdot)$ is a sufficiently integrable deterministic function. The next proposition provides a closed form expression for the moments of shot noise processes using standard Bell polynomials, see also [Corollary 2](#) in [Daw and Pender \(2020\)](#) for another expression using matrix exponentials in case $\lambda(ds) = \lambda ds$ for some rate $\lambda > 0$, and $g(s, t) = e^{-\beta(t-s)}$ for some $\beta > 0$.

Proposition 2.2 For any $n \geq 1$, we have

$$\mathbb{E}[S_t^n] = B_n \left(\mathbb{E}[J_1] \int_0^t g(s, t) ds, \dots, \mathbb{E}[J_1^n] \int_0^t g^n(s, t) ds \right),$$

where B_n is the Bell polynomial of order $n \geq 1$.

Proof. Taking $u_s(\omega) := J_{N_s} g(s, t)$, by (2.1) we have

$$\begin{aligned} \mathbb{E}[S_t^n] &= \mathbb{E} \left[\left(\sum_{k=1}^{N_t} J_{N_{T_k}} g(T_k, t) \right)^n \right] \\ &= \mathbb{E} \left[\left(\sum_{k=1}^{N_t} u_{T_k}(\omega) \right)^n \right] \\ &= \sum_{l=1}^n \sum_{\pi_1 \cup \dots \cup \pi_l = \{1, \dots, n\}} \mathbb{E} \left[\int_0^t \dots \int_0^t \epsilon_{s_1}^+ \dots \epsilon_{s_l}^+ (u^{|\pi_1|}(s_1, \omega) \dots u^{|\pi_l|}(s_l, \omega)) \lambda(ds_1) \dots \lambda(ds_l) \right] \\ &= \sum_{l=1}^n \sum_{\pi_1 \cup \dots \cup \pi_l = \{1, \dots, n\}} \int_0^t \dots \int_0^t g^{|\pi_1|}(s_1, t) \dots g^{|\pi_l|}(s_l, t) \mathbb{E}[\epsilon_{s_1}^+ \dots \epsilon_{s_l}^+ (J_{N_{s_1}}^{|\pi_1|} \dots J_{N_{s_l}}^{|\pi_l|})] \lambda(ds_1) \dots \lambda(ds_l) \\ &= \sum_{l=1}^n \sum_{\pi_1 \cup \dots \cup \pi_l = \{1, \dots, n\}} \int_0^t \dots \int_0^t g^{|\pi_1|}(s_1, t) \dots g^{|\pi_l|}(s_l, t) \mathbb{E}[J_{N_{s_1}}^{|\pi_1|}] \mathbb{E}[J_{1+N_{s_1}}^{|\pi_1|}] \dots \mathbb{E}[J_{l-1+N_{s_l}}^{|\pi_l|}] \lambda(ds_1) \dots \lambda(ds_l) \\ &= \sum_{l=1}^n \sum_{\pi_1 \cup \dots \cup \pi_l = \{1, \dots, n\}} \mathbb{E}[J_1^{|\pi_1|}] \dots \mathbb{E}[J_1^{|\pi_l|}] \int_0^t \dots \int_0^t g^{|\pi_1|}(s_1, t) \dots g^{|\pi_l|}(s_l, t) \lambda(ds_1) \dots \lambda(ds_l). \end{aligned}$$

□

We note that Proposition 2.2 is consistent with the Lévy-Khintchine formula for compound Poisson processes, as the Faà di Bruno formula, see e.g. § 2 of Lukacs (1955), yields

$$\begin{aligned} \mathbb{E}[e^{\alpha S_t}] &= \sum_{n \geq 0} \frac{\alpha^n}{n!} \mathbb{E}[S_t^n] \\ &= \sum_{n \geq 0} \frac{\alpha^n}{n!} B_n \left(\mathbb{E}[J_1] \int_0^t g(s, t) ds, \dots, \mathbb{E}[J_1^n] \int_0^t g^n(s, t) ds \right) \\ &= \exp \left(\sum_{n \geq 1} \frac{\alpha^n}{n!} \mathbb{E}[J_1^n] \int_0^t g^n(s, t) ds \right) \\ &= \exp \left(\int_0^t (e^{\alpha g(s, t) J_1} - 1) ds \right), \end{aligned}$$

which recovers the cumulants of S_t from the moments of J_1 as

$$\kappa_{S_t}^{(n)} = \mathbb{E}[J_1^n] \int_0^t g^n(s, t) ds, \quad n \geq 1.$$

3 Moments of jump processes

From now on we assume that $(N_t)_{t \in \mathbb{R}_+}$ is a standard Poisson process with intensity $\lambda > 0$, and in this section we consider jump processes built as the anticipating Poisson integrals

$$Y_t = \sum_{k=1}^{N_t} g(T_k, k, N_t) = \int_0^t g(s, N_s, N_t) dN_s, \quad t \in \mathbb{R}_+. \quad (3.1)$$

Proposition 3.1 *Let $(Y_t)_{t \in \mathbb{R}_+}$ be defined as in (3.1). For all $n \geq 1$, we have*

$$\mathbb{E}[(Y_t)^n] = \sum_{k=1}^n \lambda^k \sum_{\pi_1 \cup \dots \cup \pi_k = \{1, \dots, n\}} \int_0^t \dots \int_0^t \mathbb{E} \left[\prod_{l=1}^k g^{|\pi_l|}(s_l, l + N_{s_l}, k + N_t) \right] ds_1 \dots ds_n.$$

Proof. We have

$$\begin{aligned} \mathbb{E}[(Y_t)^n] &= \mathbb{E} \left[\left(\int_0^t g(s, N_s, N_t) dN_s \right)^n \right] \\ &= \sum_{k=1}^n \lambda^k \sum_{\pi_1 \cup \dots \cup \pi_k = \{1, \dots, n\}} \int_0^t \dots \int_0^t \mathbb{E} \left[\epsilon_{s_1}^+ \dots \epsilon_{s_n}^+ \prod_{l=1}^k g^{|\pi_l|}(s_l, N_{s_l}, N_t) \right] ds_1 \dots ds_n \\ &= \sum_{k=1}^n \lambda^k \sum_{\pi_1 \cup \dots \cup \pi_k = \{1, \dots, n\}} \int_0^t \dots \int_0^t \mathbb{E} \left[\prod_{l=1}^k g^{|\pi_l|}(s_l, l + N_{s_l}, k + N_t) \right] ds_1 \dots ds_n, \end{aligned}$$

where the sum runs over all partitions π_1, \dots, π_k of $\{1, \dots, n\}$. □

Next, we specialize Proposition 3.1 to the case where $g(s, k, n)$ takes the form

$$g(s, k, n) = f_k(s) W_k^n,$$

where

$$W_k^n = (1 - Z_k) \prod_{l=k+1}^n Z_l, \quad 1 \leq k \leq n,$$

and $(Z_k)_{k \geq 1}$ is an i.i.d. random sequence independent of the Poisson process $(N_t)_{t \in \mathbb{R}_+}$, with moment sequence $m_n = \mathbb{E}[Z^n]$, $n \geq 0$, i.e. we have

$$Y_t = \sum_{k=1}^{N_t} f_k(T_k) (1 - Z_k) \prod_{l=k+1}^{N_t} Z_l, \quad t \in \mathbb{R}_+. \quad (3.2)$$

Corollary 3.2 *Let $(Y_t)_{t \in \mathbb{R}_+}$ be defined as in (3.2). For all $n \geq 1$ we have*

$$\mathbb{E}[(Y_t)^n] = n! e^{\lambda t(m_n - 1)} \sum_{k=1}^n \lambda^k \sum_{q_0=0 < q_1 < \dots < q_k=n} 1$$

$$\int_0^t \int_0^{s_k} \cdots \int_0^{s_2} \prod_{l=1}^k \left(\frac{f_l^{q_l - q_{l-1}}(s_l)}{(q_l - q_{l-1})!} C_{q_{l-1}, q_l - q_{l-1}} e^{\lambda s_l (m_{q_{l-1}} - m_{q_l})} \right) ds_1 \cdots ds_k,$$

$t \in \mathbb{R}_+$, where

$$C_{p,q} := \mathbb{E}[(1-Z)^p Z^q] = \sum_{k=0}^p \binom{p}{k} (-1)^k m_{q+k}, \quad p, q \geq 0.$$

Proof. By Proposition 3.1, for all $n \geq 1$ we have

$$\begin{aligned} \mathbb{E}[(Y_t)^n] &= n! \sum_{k=1}^n \lambda^k \sum_{\substack{p_1 + \cdots + p_k = n \\ p_1, \dots, p_k \geq 1}} \int_0^t \int_0^{s_k} \cdots \int_0^{s_2} \frac{f^{p_1}(s_1) \cdots f^{p_k}(s_k)}{p_1! \cdots p_k!} \mathbb{E} \left[\epsilon_{s_1}^+ \cdots \epsilon_{s_n}^+ (W_{N_{s_1}}^{N_t})^{p_1} \cdots (W_{N_{s_k}}^{N_t})^{p_k} \right] ds_1 \cdots ds_k. \end{aligned}$$

For $p_1 + \cdots + p_k = n$ and $0 \leq s_1 < \cdots < s_k \leq s_{k+1} := t$, we have

$$\begin{aligned} \mathbb{E} \left[\epsilon_{s_1}^+ \cdots \epsilon_{s_k}^+ \left((W_{N_{s_1}}^{N_t})^{p_1} \cdots (W_{N_{s_k}}^{N_t})^{p_k} \right) \right] &= \mathbb{E} \left[(W_{1+N_{s_1}}^{k+N_t})^{p_1} \cdots (W_{k+N_{s_k}}^{k+N_t})^{p_k} \right] \\ &= \mathbb{E} \left[\left((1 - Z_{1+N_{s_1}}) \prod_{l=2+N_{s_1}}^{k+N_t} Z_l \right)^{p_1} \cdots \left((1 - Z_{k+N_{s_k}}) \prod_{l=k+1+N_{s_k}}^{k+N_t} Z_l \right)^{p_k} \right] \\ &= \mathbb{E} \left[\prod_{l=1}^k \left((1 - Z_{l+N_{s_l}})^{p_l} (Z_{l+N_{s_l}})^{p_1 + \cdots + p_{l-1}} \prod_{p=1+N_{s_l}}^{N_{s_{l+1}}} (Z_{l+p})^{p_1 + \cdots + p_l} \right) \right] \\ &= \mathbb{E} \left[\prod_{l=1}^k (C_{p_1 + \cdots + p_{l-1}, p_l} (m_{p_1 + \cdots + p_l})^{N_{s_{l+1}} - N_{s_l}}) \right] \\ &= \prod_{l=1}^k (C_{p_1 + \cdots + p_{l-1}, p_l} e^{\lambda (s_{l+1} - s_l) (m_{p_1 + \cdots + p_l} - 1)}) \\ &= e^{\lambda t (m_n - 1)} \prod_{l=1}^k (C_{p_1 + \cdots + p_{l-1}, p_l} e^{\lambda s_l (m_{p_1 + \cdots + p_{l-1}} - m_{p_1 + \cdots + p_l})}). \end{aligned}$$

□

4 Uniform cut-offs

In this section we assume that Z_k is uniform on $[0, 1]$, $k \geq 1$. In this case we have $m_n = 1/(n+1)$ and $C_{p,q}$ is given by the beta function as $C_{p,q} = p!q!/(p+q+1)!$, hence

$$\prod_{l=1}^k C_{p_1 + \cdots + p_{l-1}, p_l} = \prod_{l=1}^k \frac{(p_1 + \cdots + p_{l-1})! p_l!}{(p_1 + \cdots + p_l + 1)!} = \frac{1}{n!} \prod_{l=1}^k \frac{p_l!}{p_1 + \cdots + p_l + 1},$$

which yields the next consequence of Corollary 3.2.

Corollary 4.1 *Let $(Y_t)_{t \in \mathbb{R}_+}$ be defined as in (3.2), where $(Z_k)_{k \geq 1}$ is an i.i.d. uniform sequence on $[0, 1]$. For any $n \geq 1$, we have*

$$\mathbb{E}[(Y_t)^n] = e^{-n\lambda t/(n+1)} \sum_{k=1}^n \lambda^k \sum_{q_0=0 < q_1 < \dots < q_k=n} \int_0^t \int_0^{s_k} \dots \int_0^{s_2} \prod_{l=1}^k \frac{f_l^{q_l - q_{l-1}}(s_l) e^{\lambda s_l (1/q_{l-1} - 1/q_l)}}{1 + q_l} ds_1 \dots ds_k, \quad (4.1)$$

$t \in \mathbb{R}_+$.

When $f(s) = s$, $s \in \mathbb{R}_+$, the relation

$$\int_0^{s_2} s_1^n e^{\alpha s_1} ds_1 = 1 - e^{\alpha s_2} \sum_{k=0}^n (-1)^k \frac{(\alpha s_2)^k}{k!}, \quad s_2 \in \mathbb{R}_+, \quad n \geq 0, \quad (4.2)$$

can be used to compute the integrals in (4.1) by induction.

First moment

For $n = 1$ we have

$$\mathbb{E}[Y_t] = \frac{\lambda}{2} e^{-\lambda t/2} \int_0^t s_1 e^{\lambda s_1/2} ds_1 = t - \frac{2}{\lambda} (1 - e^{-\lambda t/2}), \quad (4.3)$$

which is consistent with Theorem 4 in Boxma et al. (2006), with a shorter proof, see Figure 1.

Second moment

For $n = 2$ we have

$$\begin{aligned} \mathbb{E}[(Y_t)^2] &= \frac{\lambda}{3} e^{-2\lambda t/3} \int_0^t s_1^2 e^{2\lambda s_1/3} ds_1 + \frac{\lambda^2}{6} e^{-2\lambda t/3} \int_0^t s_2 e^{\lambda s_2/6} \int_0^{s_2} s_1 e^{\lambda s_1/2} ds_1 ds_2 \\ &= \frac{18}{\lambda^2} e^{-2\lambda t/3} + \frac{4}{\lambda} e^{-\lambda t/2} t - \frac{24}{\lambda^2} e^{-\lambda t/2} + t^2 + \frac{6}{\lambda^2} - \frac{4}{\lambda} t, \end{aligned} \quad (4.4)$$

hence

$$\kappa^{(2)}(t) = \text{Var}[Y_t] = \mathbb{E}[Y_t^2] - (\mathbb{E}[Y_t])^2 = \frac{2}{\lambda^2} (9e^{-2\lambda t/3} - 2e^{-\lambda t} - 8e^{-\lambda t/2} + 1),$$

which recovers Theorem 5 in Boxma et al. (2006) with a shorter proof, see Figure 1. Figures 1 to 7 are plotted with 10 million Monte Carlo samples and $\lambda = 2$.




Figure 1: Mean and variance given by (4.3) and (4.4).

The subsequent integrals for higher order moments can be evaluated using Mathematica or based on the recurrence relation (4.2).

Third moment

For $n = 3$ we have

$$\begin{aligned}
\mathbb{E}[(Y_t)^3] &= \frac{\lambda}{4} e^{-3\lambda t/4} \int_0^t s_1^3 e^{3\lambda s_1/4} ds_1 + \frac{\lambda^2}{8} e^{-3\lambda t/4} \int_0^t s_2^2 e^{\lambda s_2/4} \int_0^{s_2} s_1 e^{\lambda s_1/2} ds_1 ds_2 \\
&+ \frac{\lambda^2}{12} e^{-3\lambda t/4} \int_0^t s_2 e^{\lambda s_2/12} \int_0^{s_2} s_1^2 e^{2\lambda s_1/3} ds_1 ds_2 \\
&+ \frac{\lambda^3}{24} e^{-3\lambda t/4} \int_0^t s_3 e^{\lambda s_3/12} \int_0^{s_3} s_2 e^{\lambda s_2/6} \int_0^{s_2} s_1 e^{\lambda s_1/2} ds_1 ds_2 ds_3 \\
&= \frac{e^{-3\lambda t/4}}{\lambda^3} (384 + 54e^{\lambda t/12}(\lambda t - 12) + 6e^{\lambda t/4}(48 + \lambda t(\lambda t - 12)) + e^{3\lambda t/4}(\lambda t(18 + \lambda t(\lambda t - 6)) - 24)),
\end{aligned} \tag{4.5}$$

see Figure 2 below.

Fourth moment

For $n = 4$ we have

$$\begin{aligned}
\mathbb{E}[(Y_t)^4] &= \frac{\lambda}{5} e^{-4\lambda t/5} \int_0^t s_1^4 e^{4\lambda s_1/5} ds_1 + \frac{\lambda^2}{10} e^{-4\lambda t/5} \int_0^t s_2^3 e^{3\lambda s_2/10} \int_0^{s_2} s_1 e^{\lambda s_1/2} ds_1 ds_2 \\
&+ \frac{\lambda^2}{20} e^{-4\lambda t/5} \int_0^t s_2 e^{\lambda s_2/20} \int_0^{s_2} s_1^3 e^{3\lambda s_1/4} ds_1 ds_2 + \frac{\lambda^2}{15} e^{-4\lambda t/5} \int_0^t s_2^2 e^{2\lambda s_2/15} \int_0^{s_2} s_1^2 e^{2\lambda s_1/3} ds_1 ds_2 \\
&+ \frac{\lambda^3}{30} e^{-4\lambda t/5} \int_0^t s_3^2 e^{2\lambda s_3/15} \int_0^{s_3} s_2 e^{\lambda s_2/6} \int_0^{s_2} s_1 e^{\lambda s_1/2} ds_1 ds_2 ds_3
\end{aligned}$$

$$\begin{aligned}
& + \frac{\lambda^3}{40} e^{-4\lambda t/5} \int_0^t s_3 e^{\lambda s_3/20} \int_0^{s_3} s_2^2 e^{\lambda s_2/4} \int_0^{s_2} s_1 e^{\lambda s_1/2} ds_1 ds_2 ds_3 \\
& + \frac{\lambda^3}{60} e^{-4\lambda t/5} \int_0^t s_3 e^{\lambda s_1/20} \int_0^{s_3} s_2 e^{\lambda s_1/12} \int_0^{s_2} s_1^2 e^{2\lambda s_1/3} ds_1 ds_2 ds_3 \\
& + \frac{\lambda^4}{120} e^{-4\lambda t/5} \int_0^t s_4 e^{\lambda s_4/20} \int_0^{s_4} s_3 e^{\lambda s_3/12} \int_0^{s_3} s_2 e^{\lambda s_2/6} \int_0^{s_2} s_1 e^{\lambda s_1/2} ds_1 ds_2 ds_3 ds_4 \\
& = \frac{e^{-4\lambda t/5}}{\lambda^4} (15000 + 1536e^{\lambda t/20}(\lambda t - 20) + 108e^{2\lambda t/15}(180 + \lambda t(\lambda t - 24)) \\
& \quad + 8e^{3\lambda t/10}(\lambda t(144 + \lambda t(\lambda t - 18)) - 480) + e^{4\lambda t/5}(120 + \lambda t(\lambda t(36 + \lambda t(\lambda t - 8)) - 96))),
\end{aligned} \tag{4.6}$$

see Figure 3 below.

5 Moments of growth-collapse processes

In this section we consider the growth-collapse process $(X_t)_{t \in \mathbb{R}_+}$ of Boxma et al. (2006), defined as $X_t := t - Y_t$, i.e.

$$X_t = t - \sum_{k=1}^{N_t} g(T_k, k, N_t) = t - \int_0^t g(s, N_s, N_t) dN_s, \quad t \in \mathbb{R}_+.$$

The moments of X_t can be recovered from (4.1) and the binomial recursion

$$\mathbb{E}[(X_t)^n] = \mathbb{E}[(t - Y_t)^n] = (-1)^n \left(\mathbb{E}[(Y_t)^n] - \sum_{k=0}^{n-1} \binom{n}{k} t^{n-k} (-1)^k \mathbb{E}[(X_t)^k] \right). \tag{5.1}$$

The next proposition extends Theorems 4 and 5 as well as Corollary 1 of Boxma et al. (2006) from mean and variance to moments of all orders, see also Corollary 4 in Daw and Pender (2020) for an expression using matrix exponentials. It is also consistent with Theorem 3 of Boxma et al. (2006) which states that the stationary distribution of the Markovian growth-collapse process $(X_t)_{t \in \mathbb{R}_+}$ is the gamma distribution $\Gamma(2, \lambda)$ with shape parameter 2 and scaling parameter λ , and cumulants $\kappa^{(n)}(\infty) = 2(n-1)!/\lambda^n$, $n \geq 1$.

Proposition 5.1 *The moments of the growth-collapse process*

$$X_t = t - \sum_{k=1}^{N_t} f_k(T_k)(1 - U_k) \prod_{l=k+1}^{N_t} U_l, \quad t \in \mathbb{R}_+,$$

with uniform cut-offs $(U_k)_{k \geq 1}$ on $[0, 1]$, are given by

$$\mathbb{E}[(X_t)^n] = \frac{(n+1)!}{\lambda^n} \sum_{k=0}^n (-1)^k (k+1)^{n-1} \binom{n}{k} e^{-k\lambda t/(k+1)}, \quad n \geq 0, \quad t \in \mathbb{R}_+.$$

As a consequence, the asymptotic moments of $(X_t)_{t \in \mathbb{R}_+}$ are given by

$$\lim_{t \rightarrow \infty} \mathbb{E}[(X_t)^n] = \frac{(n+1)!}{\lambda^n}, \quad n \geq 1.$$

Before proving Proposition 5.1, we recover the first moments and cumulants of X_t from the expressions (4.3)-(4.6) and the identity (5.1). We find

$$\mathbb{E}[X_t] = -\mathbb{E}[Y_t] + t = \frac{\lambda}{2} e^{-\lambda t/2} \int_0^t s_1 e^{\lambda s_1/2} ds_1 = \frac{2}{\lambda} (-e^{-\lambda t/2} + 1) \quad (5.2)$$

and

$$\mathbb{E}[(X_t)^2] = \mathbb{E}[(Y_t)^2] - t^2 + 2t\mathbb{E}[X_t] = \frac{3!}{\lambda^2} (3e^{-2\lambda t/3} - 4e^{-\lambda t/2} + 1), \quad (5.3)$$


see Theorems 4 and 5 of Boxma et al. (2006). Next, from (4.5) we have

$$\begin{aligned} \mathbb{E}[(X_t)^3] &= -\mathbb{E}[(Y_t)^3] + t^3 - 3t^2\mathbb{E}[X_t] + 3t\mathbb{E}[X_t^2] \\ &= \frac{4!}{\lambda^3} (-16e^{-3\lambda t/4} + 27e^{-2\lambda t/3} - 12e^{-\lambda t/2} + 1), \end{aligned} \quad (5.4)$$

and therefore the third cumulant of X_t is given by (2.2) as

$$\kappa^{(3)}(t) = 2 \frac{2!}{\lambda^3} (-27e^{-7\lambda t/6} + 96e^{-3\lambda t/4} - 135e^{-2\lambda t/3} + 4e^{-3\lambda t/2} + 24e^{-\lambda t} + 39e^{-\lambda t/2} - 1), \quad (5.5)$$

see Figure 2.



(a) Third cumulant $\kappa^{(3)}(t)$.

(b) Skewness $\kappa^{(3)}(t)/(\kappa^{(2)}(t))^{3/2}$.

Figure 2: Third cumulant (5.5) and skewness.

From (4.6) and (5.2)-(5.5) we have


$$\mathbb{E}[(X_t)^4] = \mathbb{E}[(Y_t)^4] - t^4 + 4t^3\mathbb{E}[X_t] - 6t^2\mathbb{E}[(X_t)^2] + 4t\mathbb{E}[X_t^3]$$

$$= \frac{5!}{\lambda^4} (125e^{-4\lambda t/5} - 256e^{-3\lambda t/4} + 162e^{-2\lambda t/3} - 32e^{-\lambda t/2} + 1), \quad (5.6)$$

and therefore the fourth cumulant of X_t is given by (2.2) as

$$\begin{aligned} \kappa^{(4)}(t) = & 2 \frac{3!}{\lambda^4} (-8e^{-2\lambda t} + 504e^{-7\lambda t/6} + 1250e^{-4\lambda t/5} - 256e^{-5\lambda t/4} - 2304e^{-3\lambda t/4} \\ & + 72e^{-5\lambda t/3} - 81e^{-4\lambda t/3} + 1206e^{-2\lambda t/3} - 64e^{-3\lambda t/2} - 168e^{-\lambda t} - 152e^{-\lambda t/2} + 1), \end{aligned} \quad (5.7)$$

see Figure 3.



(a) Fourth cumulant $\kappa^{(4)}(t)$.

(b) Excess kurtosis $\kappa^{(4)}(t)/(\kappa^{(2)}(t))^2$.

Figure 3: Fourth cumulant (5.7) and excess kurtosis.

Proof of Proposition 5.1. Using the infinitesimal generator of the Markov process $(X_t)_{t \in \mathbb{R}_+}$ it can be shown, see § 3 of Boxma et al. (2006) and § 3.5 of Daw and Pender (2020), that the moments $\mathbb{E}[(X_t)^n]$ satisfy the differential equation

$$\frac{d}{dt} \mathbb{E}[(X_t)^n] = n \mathbb{E}[(X_t)^{n-1}] - \frac{\lambda n}{n+1} \mathbb{E}[(X_t)^n].$$

Based on the intuition gained from (5.2)-(5.6), we search for a solution of the form

$$\mathbb{E}[(X_t)^n] = \frac{(n+1)!}{\lambda^n} \sum_{k=0}^n a_{k,n} e^{-k\lambda t/(k+1)},$$

which, by identification of terms, yields the recurrence relation

$$a_{k,n} = \frac{n(k+1)}{n-k} a_{k,n-1}, \quad 0 \leq k < n,$$

hence

$$a_{k,n} = (k+1)^{n-k} \binom{n}{k} a_{k,k}, \quad 0 \leq k < n.$$

In addition, the initial condition $t = 0$ requires

$$\sum_{k=0}^n a_{k,n} = 0,$$

hence

$$\sum_{k=0}^n (k+1)^{n-k} \binom{n}{k} a_{k,k} = 0,$$

which is solved by taking $a_{k,k} = (-1)^k (k+1)^{k-1}$, due to the combinatorial relation

$$S(n, n+1) = \sum_{k=0}^n (-1)^{n-k} (k+1)^n \binom{n+1}{k+1} = (n+1) \sum_{k=0}^n (k+1)^{n-1} \binom{n}{k} (-1)^k = 0,$$

which follows from the vanishing of the Stirling numbers of the second kind $S(n, n+1)$, see e.g. page 824 of [Abramowitz and Stegun \(1972\)](#). \square

6 Embedded growth-collapse chain

In this section we show that Proposition 2.1 can also be used to compute the moments of all orders of the embedded chain

$$Y(m) = Y_{T_m} = \sum_{k=1}^m g(T_k, k, m) = \int_0^\infty g(s, N_s, m) \mathbf{1}_{[0, T_m]}(s) dN_s, \quad m \geq 1. \quad (6.1)$$

Proposition 6.1 *Let $(Y(m))_{m \geq 1}$ be of the form (6.1). For any $n, m \geq 1$ we have*

$$\mathbb{E}[(Y(m))^n] = n! \sum_{k=1}^n \lambda^k \sum_{0=q_0 < q_1 < \dots < q_{k-1} < q_k = n} \mathbb{E} \left[\int_0^\infty \int_0^{s_k} \dots \int_0^{s_2} \mathbf{1}_{\{N_{s_k}^- \leq m-k\}} \prod_{l=1}^k \frac{(g(s_l, l + N_{s_l}, m))^{q_l - q_{l-1}}}{(q_l - q_{l-1})!} ds_1 \dots ds_k \right].$$

Proof. By Proposition 2.1 and the identity $\{s \leq T_m\} = \{N_{s^-} < m\}$, $s > 0$, for all $n \geq 1$ we have

$$\begin{aligned} \mathbb{E} \left[\left(\sum_{k=1}^m g(T_k, k, m) \right)^n \right] &= \mathbb{E} \left[\left(\int_0^\infty g(s, N_s, m) \mathbf{1}_{[0, T_m]}(s) dN_s \right)^n \right] \\ &= \sum_{k=1}^n k! \lambda^k \sum_{\pi_1 \cup \dots \cup \pi_k = \{1, \dots, n\}} \mathbb{E} \left[\int_0^\infty \int_0^{s_k} \dots \int_0^{s_2} \epsilon_{s_1}^+ \dots \epsilon_{s_k}^+ \prod_{l=1}^k (g(s_l, N_{s_l}, m) \mathbf{1}_{\{N_{s_l}^- < m\}})^{|\pi_l|} ds_1 \dots ds_k \right] \\ &= \sum_{k=1}^n k! \lambda^k \sum_{\pi_1 \cup \dots \cup \pi_k = \{1, \dots, n\}} \mathbb{E} \left[\int_0^\infty \int_0^{s_k} \dots \int_0^{s_2} \prod_{l=1}^k (g(s_l, l + N_{s_l}, m) \mathbf{1}_{\{l-1 + N_{s_l}^- < m\}})^{|\pi_l|} ds_1 \dots ds_k \right] \end{aligned}$$

$$= \sum_{k=1}^n k! \lambda^k \sum_{\pi_1 \cup \dots \cup \pi_k = \{1, \dots, n\}} \mathbb{E} \left[\int_0^\infty \int_0^{s_k} \dots \int_0^{s_2} \mathbf{1}_{\{N_{s_k}^- \leq m-k\}} \prod_{l=1}^k (g(s_l, l + N_{s_l}, m))^{\lvert \pi_l \rvert} ds_1 \dots ds_k \right].$$

□

Next, we specialize Proposition 6.1 to the case where $g(s, k, m)$ takes the form

$$g(s, k, m) = f_k(s)(1 - Z_k) \prod_{l=k+1}^m Z_l, \quad (6.2)$$

where $(Z_k)_{k \geq 1}$ is an i.i.d. random sequence independent of the Poisson process $(N_t)_{t \in \mathbb{R}_+}$, with moment sequence $m_n = \mathbb{E}[Z^n]$, $n \geq 0$.

Corollary 6.2 *Let $(Y(m))_{m \geq 1}$ be defined as in (6.2). For any $n, m \geq 1$ we have*

$$\begin{aligned} \mathbb{E}[(Y(m))^n] &= n! \sum_{k=1}^n \sum_{i=0}^{m-k} \frac{\lambda^{k+i}}{i!} m_n^{m-i-k} \sum_{0=q_0 < q_1 < \dots < q_{k-1} < q_k = n} \\ &\prod_{l=1}^k \frac{C_{q_{l-1}, q_l - q_{l-1}}}{(q_l - q_{l-1})!} \int_0^\infty e^{-\lambda s_k} \int_0^{s_k} \dots \int_0^{s_2} \left(s_1 + \sum_{l=1}^{k-1} m_{p_1 + \dots + p_l}(s_{l+1} - s_l) \right)^i \prod_{l=1}^k f_{l,m}^{q_l - q_{l-1}}(s_l) ds_1 \dots ds_k. \end{aligned} \quad (6.3)$$

Proof. Using the fact that the jumps of $(N_s)_{s \in [0, s_k]}$ are uniformly distributed on $[0, s_k]$ given that $N_{s_k} = i$, by Proposition 6.1 we have

$$\begin{aligned} \mathbb{E} \left[\left(\sum_{k=1}^m g(T_k, k, m) \right)^n \right] &= \sum_{k=1}^n k! \lambda^k \sum_{\pi_1 \cup \dots \cup \pi_k = \{1, \dots, n\}} \\ &\mathbb{E} \left[\int_0^\infty \int_0^{s_k} \dots \int_0^{s_2} \mathbf{1}_{\{N_{s_k}^- \leq m-k\}} \prod_{l=1}^k \left(f_{l,m}(s_l)(1 - Z_{l+N_{s_l}}) \prod_{j=l+1+N_{s_l}}^m Z_j \right)^{\lvert \pi_l \rvert} ds_1 \dots ds_k \right] \\ &= \sum_{k=1}^n k! \lambda^k \sum_{\pi_1 \cup \dots \cup \pi_k = \{1, \dots, n\}} \\ &\mathbb{E} \left[\int_0^\infty \int_0^{s_k} \dots \int_0^{s_2} \mathbf{1}_{\{N_{s_k}^- \leq m-k\}} \prod_{l=1}^k (f_{l,m}^{p_l}(s_l) C_{p_1 + \dots + p_{l-1}, p_l}(m_{p_1 + \dots + p_l})^{N_{s_{l+1}} - N_{s_l}}) ds_1 \dots ds_k \right] \\ &= n! \sum_{k=1}^n \frac{\lambda^k}{p_1! \dots p_k!} \sum_{i=0}^{m-k} \mathbb{P}(N_{s_k}^- = i) \sum_{\substack{p_1 + \dots + p_k = n \\ p_1, \dots, p_k \geq 1}} \\ &\int_0^\infty \int_0^{s_k} \dots \int_0^{s_2} \prod_{l=1}^k (f_{l,m}^{p_l}(s_l) C_{p_1 + \dots + p_{l-1}, p_l}) \left(\frac{s_1 + \sum_{l=1}^{k-1} m_{p_1 + \dots + p_l}(s_{l+1} - s_l)}{s_k} \right)^i m_n^{m-i-k} ds_1 \dots ds_k \end{aligned}$$

$$\begin{aligned}
&= n! \sum_{k=1}^n \lambda^{k+i} \sum_{\substack{p_1+\dots+p_k=n \\ p_1, \dots, p_k \geq 1}} \sum_{i=0}^{m-k} \frac{m_n^{m-i-k}}{i!} \prod_{l=1}^k \frac{C_{p_1+\dots+p_{l-1}, p_l}}{p_l!} \\
&\int_0^\infty e^{-\lambda s_k} \int_0^{s_k} \cdots \int_0^{s_2} \prod_{l=1}^k f_{l,m}^{p_l}(s_l) \left(s_1 + \sum_{l=1}^{k-1} m_{p_1+\dots+p_l}(s_{l+1} - s_l) \right)^i ds_1 \cdots ds_k.
\end{aligned}$$

□

Uniform cut-offs

The next result specializes Corollary 6.2 to the case of embedded growth processes with uniform cut-offs.

Corollary 6.3 *Let $(Y(m))_{m \geq 1}$ be defined as in (6.2), with $(Z_k)_{k \geq 1}$ an i.i.d. uniform random sequence. For any $n, m \geq 1$ we have*

$$\begin{aligned}
\mathbb{E}[(Y(m))^n] &= \sum_{k=1}^n \sum_{i=0}^{m-k} \frac{\lambda^{k+i}}{i!(n+1)^{m-i-k}} \sum_{0=q_0 < q_1 < \dots < q_{k-1} < q_k = n} \prod_{l=1}^k \frac{1}{1+q_l} \\
&\int_0^\infty e^{-\lambda s_k} \int_0^{s_k} \cdots \int_0^{s_2} \left(s_1 + \sum_{l=1}^{k-1} \frac{s_{l+1} - s_l}{1+q_l} \right)^i \prod_{l=1}^k f_{l,m}^{q_l - q_{l-1}}(s_l) ds_1 \cdots ds_k.
\end{aligned} \tag{6.4}$$

Proof. By Corollary 6.2 we have

$$\begin{aligned}
\mathbb{E} \left[\left(\sum_{k=1}^m g(T_k, k, m) \right)^n \right] &= n! \sum_{k=1}^n \lambda^{k+i} \sum_{\substack{p_1+\dots+p_k=n \\ p_1, \dots, p_k \geq 1}} \sum_{i=0}^{m-k} \frac{m_n^{m-i-k}}{i!} \prod_{l=1}^k \frac{C_{p_1+\dots+p_{l-1}, p_l}}{p_l!} \\
&\int_0^\infty e^{-\lambda s_k} \int_0^{s_k} \cdots \int_0^{s_2} \prod_{l=1}^k f_{l,m}^{p_l}(s_l) \left(s_1 + \sum_{l=1}^{k-1} m_{p_1+\dots+p_l}(s_{l+1} - s_l) \right)^i ds_1 \cdots ds_k \\
&= n! \sum_{k=1}^n \lambda^{k+i} \sum_{\substack{p_1+\dots+p_k=n \\ p_1, \dots, p_k \geq 1}} \sum_{i=0}^{m-k} \frac{m_n^{m-i-k}}{i!} \prod_{l=1}^k \frac{(p_1 + \dots + p_{l-1})!}{(p_1 + \dots + p_l + 1)!} \\
&\int_0^\infty e^{-\lambda s_k} \int_0^{s_k} \cdots \int_0^{s_2} \prod_{l=1}^k f_{l,m}^{p_l}(s_l) \left(s_1 + \sum_{l=1}^{k-1} m_{p_1+\dots+p_l}(s_{l+1} - s_l) \right)^i ds_1 \cdots ds_k.
\end{aligned}$$

□

The first moments of $Y(m)$ can be computed in closed form when $f_{l,m}(s) = s$, which corresponds to

$$Y(m) = Y_{T_m} = \sum_{k=1}^m T_k(1 - U_k) \prod_{l=k+1}^m U_l, \quad m \geq 1,$$

where $(U_k)_{k \geq 1}$ is a uniform random sequence on $[0, 1]$.

First moment

For $n = 1$, Corollary 6.3 yields

$$\mathbb{E}[Y(m)] = \frac{1}{\lambda} \sum_{i=0}^{m-1} \frac{1}{i!2^{m-i-1}} \int_0^\infty e^{-s_1} s_1^i \frac{s_1}{2} ds_1 = \frac{2^{-m} + m - 1}{\lambda},$$

which is consistent with Theorem 7 in Boxma et al. (2006).

Second moment

For $n = 2$, we find


$$\begin{aligned} \mathbb{E}[(Y(m))^2] &= \frac{1}{\lambda^2} \sum_{i=0}^{m-1} \frac{1}{i!3^{m-1-i}} \int_0^\infty e^{-s_1} s_1^i \frac{s_1^2}{3} ds_1 + \frac{1}{2\lambda^2} \sum_{i=0}^{m-2} \frac{1}{i!3^{m-2-i}} \int_0^\infty e^{-s_2} \int_0^{s_2} \left(\frac{s_1 + s_2}{2}\right)^i s_1 \frac{s_2}{3} ds_1 ds_2 \\ &= \frac{1}{\lambda^2} \left(\frac{2}{3^m} + \frac{m-1}{2^{m-1}} - m + m^2 \right). \end{aligned}$$

Third moment


For $n = 3$, we have

$$\begin{aligned} \mathbb{E}[Y(m)^3] &= \frac{1}{\lambda^3} \sum_{i=0}^{m-1} \frac{1}{i!4^{m-1-i}} \int_0^\infty e^{-s_1} \frac{s_1^{i+3}}{4} ds_1 \\ &\quad + \frac{1}{\lambda^3} \sum_{i=0}^{m-2} \frac{1}{i!4^{m-2-i}} \int_0^\infty e^{-s_2} \int_0^{s_2} \left(\left(\frac{s_1 + s_2}{2}\right)^i \frac{s_1 s_2^2}{2 \cdot 4} + \left(\frac{2s_1 + s_2}{3}\right)^i \frac{s_1^2 s_2}{3 \cdot 4} \right) ds_1 ds_2 \\ &\quad + \frac{1}{\lambda^3} \sum_{i=0}^{m-3} \frac{1}{i!4^{m-3-i}} \int_0^\infty e^{-s_3} \int_0^{s_3} \int_0^{s_2} \left(\frac{3s_1 + s_2 + 2s_3}{6}\right)^i \frac{s_1 s_2 s_3}{2 \cdot 3 \cdot 4} ds_1 ds_2 ds_3. \quad (6.5) \end{aligned}$$

Although the last partial summation (6.5) does not have a polynomial expression, it can easily be estimated numerically, see Figure 4.



(a) Third cumulant of $Y(m)$.



(b) Skewness $\kappa^{(3)}(m)/(\kappa^{(2)}(m))^{3/2}$.

Figure 4: Third cumulant and skewness of $Y(m)$.

Fourth moment

For $n = 4$, we find

$$\begin{aligned}
\mathbb{E}[Y(m)^4] &= \frac{1}{\lambda^4} \sum_{i=0}^{m-1} \frac{1}{i!5^{m-1-i}} \int_0^\infty e^{-s_1} \frac{s_1^{i+4}}{5} ds_1 \\
&+ \frac{1}{\lambda^4} \sum_{i=0}^{m-2} \frac{1}{i!5^{m-2-i}} \int_0^\infty e^{-s_2} \int_0^{s_2} \left(\left(\frac{s_1 + s_2}{2} \right)^i \frac{s_1 s_2^3}{2 \cdot 5} + \left(\frac{3s_1 + s_2}{4} \right)^i \frac{s_1^3 s_2}{4 \cdot 5} + \left(\frac{2s_1 + s_2}{3} \right)^i \frac{s_1^2 s_2^2}{3 \cdot 5} \right) ds_1 ds_2 \\
&+ \frac{1}{\lambda^4} \sum_{i=0}^{m-3} \frac{1}{i!5^{m-3-i}} \int_0^\infty e^{-s_3} \int_0^{s_3} \int_0^{s_2} \\
&\left(\left(\frac{8s_1 + s_2 + 3s_3}{12} \right)^i \frac{s_1^2 s_2 s_3}{3 \cdot 4 \cdot 5} + \left(\frac{4s_1 + 2s_2 + 2s_3}{8} \right)^i \frac{s_1 s_2^2 s_3}{2 \cdot 4 \cdot 5} + \left(\frac{3s_1 + s_2 + 2s_3}{6} \right)^i \frac{s_1 s_2 s_3^2}{2 \cdot 3 \cdot 5} \right) ds_1 ds_2 ds_3 \\
&+ \frac{1}{\lambda^4} \sum_{i=0}^{m-4} \frac{1}{i!5^{m-4-i}} \int_0^\infty e^{-s_4} \int_0^{s_4} \int_0^{s_3} \int_0^{s_2} \left(\frac{6s_1 + 2s_2 + s_3 + 3s_4}{12} \right)^i \prod_{l=1}^4 \frac{s_l}{2 \cdot 3 \cdot 4 \cdot 5} ds_1 ds_2 ds_3 ds_4,
\end{aligned}$$

see Figure 5.




Figure 5: Fourth cumulant and excess kurtosis of $Y(m)$.

Compensated embedded chain

Finally, we consider the compensated embedded chain

$$\begin{aligned}
X(m) &:= Y_{T_m} - f_m(T_m) \\
&= -f_m(T_m) + \sum_{k=1}^m g(T_k, k, m) \\
&= -f_m(T_m) + \sum_{k=1}^m f_k(s)(1 - Z_k) \prod_{l=k+1}^n Z_l, \quad m \geq 1,
\end{aligned} \tag{6.6}$$

where $(Z_k)_{k \geq 1}$ is an i.i.d. random sequence independent with moment sequence $m_n = \mathbb{E}[Z^n]$, $n \geq 0$. This process can be obtained from (6.1) by replacing $g(s, m, m) = f_m(s)(1 - Z_m)$ with $g(s, m, m) = f_m(s)(1 - Z_m) - f_m(s) = -f_m(s)Z_m$ in (6.1). In Corollary 6.2, this amounts to modifying last term of $i = m - k$ or $N_{s_k} = m - k$ in (6.3), by changing the last term $C_{q_{k-1}, q_k - q_{k-1}}$ of order $l = k$ in the product $\prod_{l=1}^k C_{q_{l-1}, q_l - q_{l-1}}$ into $(-1)^{q_k - q_{k-1}} m_{q_k} = (-1)^{q_k - q_{k-1}} m_n$, yielding the next corollary.

Corollary 6.4 *Let $(X(m))_{m \geq 1}$ be defined as in (6.6). For any $n, m \geq 1$ we have*

$$\begin{aligned}
\mathbb{E}[(X(m))^n] &= n! \sum_{k=1}^n \sum_{i=0}^{m-k-1} \frac{\lambda^{k+i}}{i!} m_n^{m-i-k} \sum_{0=q_0 < q_1 < \dots < q_{k-1} < q_k = n} \\
&\int_0^\infty e^{-\lambda s_k} \int_0^{s_k} \dots \int_0^{s_2} \left(s_1 + \sum_{l=1}^{k-1} m_{q_l} (s_{l+1} - s_l) \right)^i \prod_{l=1}^k \frac{C_{q_{l-1}, q_l - q_{l-1}} f_{l,m}^{q_l - q_{l-1}}(s_l)}{(q_l - q_{l-1})!} ds_1 \dots ds_k
\end{aligned}$$

$$\begin{aligned}
& + n! \sum_{k=1}^{\min(n,m)} \frac{\lambda^m}{(m-k)!} \sum_{0=q_0 < q_1 < \dots < q_{k-1} < q_k = n} (-1)^{q_k - q_{k-1}} m_{q_k} \\
& \int_0^\infty e^{-\lambda s_k} f_{k,m}^{n-q_{k-1}}(s_k) \int_0^{s_k} \dots \int_0^{s_2} \left(s_1 + \sum_{l=1}^{k-1} m_{q_l} (s_{l+1} - s_l) \right)^{m-k} \prod_{l=1}^{k-1} \frac{C_{q_{l-1}, q_l - q_{l-1}} f_{l,m}^{q_l - q_{l-1}}(s_l)}{(q_l - q_{l-1})!} ds_1 \dots ds_k.
\end{aligned}$$

Similarly, according to Corollary 6.3, computing the moments of

$$X(m) = T_m - \sum_{k=1}^m T_k (1 - U_k) \prod_{l=k+1}^m U_l, \quad (6.7)$$

with $(U_k)_{k \geq 1}$ an i.i.d. uniform random sequence, means multiplying the product $\prod_{l=1}^k \frac{1}{1+q_l}$ for $i = m - k$ in (6.4) by

$$\frac{(-1)^{q_k - q_{k-1}} m_{q_k}}{C_{q_{k-1}, q_k - q_{k-1}}} = \frac{(-1)^{q_k - q_{k-1}} (1 + q_k)!}{(1 + q_k) q_{k-1}! (q_k - q_{k-1})!} = (-1)^{q_k - q_{k-1}} \binom{n}{q_{k-1}},$$

as is done in the next corollary.


Corollary 6.5 *Let $(X(m))_{m \geq 1}$ be defined as in (6.7), with $(U_k)_{k \geq 1}$ an i.i.d. uniform random sequence. For any $n, m \geq 1$ we have*

$$\begin{aligned}
\mathbb{E}[(Y(m))^n] &= \sum_{k=1}^n \sum_{i=0}^{m-k-1} \frac{\lambda^{k+i}}{i!(n+1)^{m-i-k}} \sum_{0=q_0 < q_1 < \dots < q_{k-1} < q_k = n} \\
& \int_0^\infty e^{-\lambda s_k} \int_0^{s_k} \dots \int_0^{s_2} \left(s_1 + \sum_{l=1}^{k-1} \frac{s_{l+1} - s_l}{1 + q_l} \right)^i \prod_{l=1}^k \frac{f_{l,m}^{q_l - q_{l-1}}(s_l)}{1 + q_l} ds_1 \dots ds_k \\
& + \sum_{k=1}^{\min(n,m)} \frac{\lambda^m}{(m-k)!} \sum_{0=q_0 < q_1 < \dots < q_{k-1} < q_k = n} (-1)^{q_k - q_{k-1}} \binom{n}{q_{k-1}} \\
& \int_0^\infty e^{-\lambda s_k} \int_0^{s_k} \dots \int_0^{s_2} \left(s_1 + \sum_{l=1}^{k-1} \frac{s_{l+1} - s_l}{1 + q_l} \right)^{m-k} \prod_{l=1}^k \frac{f_{l,m}^{q_l - q_{l-1}}(s_l)}{1 + q_l} ds_1 \dots ds_k.
\end{aligned}$$


When $f_{l,m}(s) = s$, Corollary 6.5 shows that the second moment reads

$$\mathbb{E}[(X(m))^2] = \frac{1}{\lambda^2} \left(2 - 4 \left(\frac{1}{2} \right)^m + 2 \left(\frac{1}{3} \right)^m \right)$$

which recovers Theorem 7 in Boxma et al. (2006).




(a) Third cumulant of $X(m)$.




(b) Skewness $\kappa^{(3)}(m)/(\kappa^{(2)}(m))^{3/2}$.

Figure 6: Third cumulant and skewness of $X(m)$.

Higher order moments of $X(m)$ can be obtained from Corollary 6.5 using Mathematica, and are plotted with $f_{i,m}(s) = s$ at the orders 3 and 4 in Figures 6-7, along with Monte Carlo simulations used for confirmation.



(a) Fourth cumulant of $X(m)$.



(b) Excess kurtosis $\kappa^{(4)}(m)/(\kappa^{(2)}(m))^2$.

Figure 7: Fourth cumulant and kurtosis of $X(m)$.

References

- M. Abramowitz and I.A. Stegun. *Handbook of mathematical functions with formulas, graphs, and mathematical tables*, volume 55. Dover Publications, New York, 1972. 9th Edition.
- O. Boxma, D. Perry, W. Stajde, and S. Zacks. A Markovian growth-collapse model. *J. Appl. Probab.*, 38: 221–243, 2006.
- P. Brémaud. *Markov chains*, volume 31 of *Texts in Applied Mathematics*. Springer-Verlag, New York, 1999.
- M.H.A. Davis. Piecewise-deterministic Markov processes: a general class of non-diffusion stochastic models. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 46:353–388, 1984.

- A. Daw and J. Pender. Matrix calculations for moments of Markov processes. Preprint arXiv:1909.03320, 2020.
- I. Eliazar and J. Klafter. A growth-collapse model: Lévy inflow, geometric crashes, and generalized Ornstein-Uhlenbeck dynamics. *Phys. A*, 334:1–21, 2004.
- M. Frolkova and M. Mandjes. A bitcoin-inspired infinite-server model with a random fluid limit. *Stochastic Models*, 35(1):1–32, 2019.
- V.P. Leonov and A.N. Shiryaev. On a method of calculation of semi-invariants. *Theory Probab. Appl.*, 4: 319–329, 1959.
- E. Lukacs. Applications of Faà di Bruno’s formula in mathematical statistics. *Amer. Math. Monthly*, 62: 340–348, 1955.
- J. Mecke. Stationäre zufällige Masse auf lokalkompakten Abelschen Gruppen. *Z. Wahrscheinlichkeitstheorie Verw. Geb.*, 9:36–58, 1967.
- N. Privault. Moment identities for Poisson-Skorohod integrals and application to measure invariance. *C. R. Math. Acad. Sci. Paris*, 347:1071–1074, 2009.
- N. Privault. Moments of Poisson stochastic integrals with random integrands. *Probability and Mathematical Statistics*, 32(2):227–239, 2012a.
- N. Privault. Invariance of Poisson measures under random transformations. *Ann. Inst. H. Poincaré Probab. Statist.*, 48(4):947–972, 2012b.
- N. Privault. Combinatorics of Poisson stochastic integrals with random integrands. In G. Peccati and M. Reitzner, editors, *Stochastic Analysis for Poisson Point Processes: Malliavin Calculus, Wiener-Itô Chaos Expansions and Stochastic Geometry*, volume 7 of *Bocconi & Springer Series*, pages 37–80. Springer, Berlin, 2016.
- I.M. Slivnyak. Some properties of stationary flows of homogeneous random events. *Theory Probab. Appl.*, 7 (3):336–341, 1962.