

Column randomization and almost-isometric embeddings

Shahar Mendelson *

January 5, 2022

Abstract

The matrix $A : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is (δ, k) -regular if for any k -sparse vector x ,

$$\left| \|Ax\|_2^2 - \|x\|_2^2 \right| \leq \delta\sqrt{k}\|x\|_2^2.$$

We show that if A is (δ, k) -regular for $1 \leq k \leq 1/\delta^2$, then by multiplying the columns of A by independent random signs, the resulting random ensemble A_ε acts on an arbitrary subset $T \subset \mathbb{R}^n$ (almost) as if it were gaussian, and with the optimal probability estimate: if $\ell_*(T)$ is the gaussian mean-width of T and $d_T = \sup_{t \in T} \|t\|_2$, then with probability at least $1 - 2 \exp(-c(\ell_*(T)/d_T)^2)$,

$$\sup_{t \in T} \left| \|A_\varepsilon t\|_2^2 - \|t\|_2^2 \right| \leq C (\Lambda d_T \delta \ell_*(T) + (\delta \ell_*(T))^2),$$

where $\Lambda = \max\{1, \delta^2 \log(n\delta^2)\}$. This estimate is optimal for $0 < \delta \leq 1/\sqrt{\log n}$.

1 Introduction

Linear operators that act in an almost-isometric way on subsets of \mathbb{R}^n are of obvious importance. Although approximations of isometries are the only operators that almost preserve the Euclidean norm of any point in \mathbb{R}^n , one may consider a more flexible alternative: a random ensemble of operators Γ such that, for any fixed $T \subset \mathbb{R}^n$, with high probability, Γ “acts well” on every element of T . Such random ensembles have been studied extensively over the years, following the path paved by the celebrated work of Johnson and Lindenstrauss in [5]. Here we formulate the Johnson-Lindenstrauss Lemma in one of its gaussian versions:

Theorem 1.1. *There exist absolute constants c_0 and c_1 such that the following holds. Let $1 \leq m \leq n$ and set $\Gamma : \mathbb{R}^n \rightarrow \mathbb{R}^m$ to be a random matrix whose entries are independent, standard gaussian random variables. Let $T \subset S^{n-1}$ be of cardinality at most $\exp(c_0 m)$. Then for $m^{-1/2} \sqrt{\log |T|} < \rho < 1$, with probability at least $1 - 2 \exp(-c_1 \rho^2 m)$, for every $t \in T$,*

$$\left| \left\| m^{-1/2} \Gamma t \right\|_2^2 - 1 \right| \leq \rho.$$

The scope of Theorem 1.1 can be extended to more general random ensembles than the gaussian one, e.g., to a random matrix whose rows are iid copies of a centred random vector

*Centre for Mathematics and its Applications, The Australian National University. Email: shahar.mendelson@anu.edu.au

that exhibits suitable decay properties (see, e.g. [4, 8]). It is far more challenging to construct a random ensemble that, on the one hand, satisfies a version of Theorem 1.1, and on the other is based on “few random bits” or is constructed using a heavy-tailed random vector.

A significant breakthrough towards more general “Johnson-Lindenstrauss transforms” came in [7], where it was shown that a matrix that satisfies a suitable version of the *restricted isometry property*, can be converted to the wanted random ensemble by multiplying its columns by random signs. More accurately, let $\varepsilon_1, \dots, \varepsilon_n$ be independent, symmetric $\{-1, 1\}$ -valued random variables. Set $D_\varepsilon = \text{diag}(\varepsilon_1, \dots, \varepsilon_n)$ and for a matrix $A : \mathbb{R}^n \rightarrow \mathbb{R}^m$ define

$$A_\varepsilon = AD_\varepsilon.$$

From here on we denote by Σ_k the subset of S^{n-1} consisting of vectors that are supported on at most k coordinates.

Definition 1.2. *A matrix $A : \mathbb{R}^n \rightarrow \mathbb{R}^m$ satisfies the restricted isometry property of order k and level $\delta \in (0, 1)$ if*

$$\sup_{x \in \Sigma_k} \left| \|Ax\|_2^2 - 1 \right| \leq \delta.$$

Theorem 1.3. [7] *There are absolute constants c_0 and c_1 such that the following holds. Let $\lambda > 0$ and $\rho \in (0, 1)$. Consider $T \subset \mathbb{R}^n$ and let $k \geq c \log(e|T|/\lambda)$. If A satisfies the restricted isometry property of order k and at level $\delta < \rho/4$, then with probability at least $1 - \lambda$, for every $t \in T$,*

$$(1 - \rho)\|t\|_2^2 \leq \|A_\varepsilon t\|_2^2 \leq (1 + \rho)\|t\|_2^2.$$

While Theorem 1.3 does not recover the probability estimate from Theorem 1.1, it does imply at the constant probability level that A_ε is an almost isometry in the random ensemble sense: if A is a matrix that $1 \pm \delta$ -preserves the norms of vectors that are $c \log |T|$ sparse, then a typical realization of the random ensemble A_ε , $1 \pm c'\delta$ preserves the norms of all the elements in T .

Various extensions of Theorem 1.1 that hold for arbitrary subsets of \mathbb{R}^n have been studied over the years. In such extensions the “complexity parameter” $\log |T|$ is replaced by more suitable counterparts. A rather general version of Theorem 1.1 follows from a functional Bernstein inequality (see, e.g., [3, 8, 2]), and to formulate that inequality in the gaussian case we require the following definition.

Definition 1.4. *Let g_1, \dots, g_n be independent, standard gaussian random variables. For $T \subset \mathbb{R}^n$ set*

$$\ell_*(T) = \mathbb{E} \sup_{t \in T} \left| \sum_{i=1}^n g_i t_i \right| \quad \text{and} \quad d_T = \sup_{t \in T} \|t\|_2.$$

Let

$$\left(\frac{\ell_*(T)}{d_T} \right)^2$$

be the critical dimension of the set T .

The critical dimension appears naturally when studying the geometry of convex sets—for example, in the context of the Dvoretzky-Milman Theorem (see [1] and references therein for

more details). It is the natural alternative to $\log |T|$ —which was suitable for finite subsets of sphere S^{n-1} .

Let $G = (g_i)_{i=1}^n$ be the standard gaussian random vector in \mathbb{R}^n , set G_1, \dots, G_m to be independent copies of G and put

$$\Gamma = \sum_{i=1}^m \langle G_i, \cdot \rangle e_i$$

to be the random ensemble used in Theorem 1.1.

Theorem 1.5. *There exist absolute constants c_0, c_1 and C such that the following holds. If $T \subset \mathbb{R}^n$ and $u \geq c_0$ then with probability at least*

$$1 - 2 \exp \left(-c_1 u^2 \left(\frac{\ell_*(T)}{d_T} \right)^2 \right),$$

for every $t \in T$,

$$\left| \left\| m^{-1/2} \Gamma t \right\|_2^2 - \|t\|_2^2 \right| \leq C \left(u d_T \frac{\ell_*(T)}{\sqrt{m}} + u^2 \left(\frac{\ell_*(T)}{\sqrt{m}} \right)^2 \right). \quad (1.1)$$

One may use Theorem 1.5 to ensure that the uniform error in (1.1) is at most $\max\{\rho, \rho^2\} d_T^2$. Indeed, if

$$\frac{\ell_*(T)/d_T}{\sqrt{m}} \sim \rho,$$

then with probability at least $1 - 2 \exp(-c_3 \rho^2 m)$,

$$\sup_{t \in T} \left| \left\| m^{-1/2} \Gamma t \right\|_2^2 - \|t\|_2^2 \right| \leq \max\{\rho, \rho^2\} d_T^2, \quad (1.2)$$

which is a natural counterpart of Theorem 1.1 once $\log |T|$ is replaced by $(\ell_*(T)/d_T)^2$.

As it happens, a version of Theorem 1.3 that is analogous to (1.2) was proved in [9], using the notion of a multi-level RIP.

Definition 1.6. *Let $L = \lceil \log_2 n \rceil$. For $\delta > 0$ and $s \geq 1$ the matrix A satisfies a multi-scale RIP with distortion δ and sparsity s if, for every $1 \leq \ell \leq L$ and every $x \in \Sigma_{2^\ell s}$, one has*

$$\left| \|Ax\|_2^2 - \|x\|_2^2 \right| \leq \max \left\{ 2^{\ell/2} \delta, 2^\ell \delta^2 \right\}.$$

Definition 1.6 implies that if $k \geq s$ then

$$\sup_{x \in \Sigma_k} \left| \|Ax\|_2^2 - \|x\|_2^2 \right| \leq \max\{\sqrt{k} \delta, k \delta^2\}.$$

Example 1.7. *Let $\Gamma : \mathbb{R}^n \rightarrow \mathbb{R}^m$ be a gaussian matrix as above and set $A = m^{-1/2} \Gamma$. It is standard to verify (using, for example, Theorem 1.5 and a well-known estimate on $\ell_*(\Sigma_k)$) that with probability at least $1 - 2 \exp(-ck \log(en/k))$,*

$$\sup_{x \in \Sigma_k} \left| \|Ax\|_2^2 - 1 \right| \leq C \sqrt{\frac{k \log(en/k)}{m}}.$$

By the union bound over k it follows that with a nontrivial probability, A satisfies a multi-scale RIP with $s = 1$ and $\delta \sim m^{-1/2} \sqrt{\log(en)}$. Observe that the second term in the multi-scale RIP—, namely $k \delta^2$, is not needed here.

Remark 1.8. *Example 1.7 gives a good intuition on the role δ has in well-behaved situations: it should scale (roughly) like $1/\sqrt{m}$, where m is the number of rows of the matrix A .*

The following theorem is the starting point of this note: an estimate on the error a typical realization of the random ensemble $A_\varepsilon = AD_\varepsilon$ has when acting on an arbitrary $T \subset \mathbb{R}^n$, given that A satisfies an appropriate multi-scale RIP.

Theorem 1.9. *[9] There are absolute constants c and C such that the following holds. Let $\eta, \rho > 0$ and $A : \mathbb{R}^n \rightarrow \mathbb{R}^m$ that satisfies a multi-scale RIP with sparsity level $s = c(1 + \eta)$ and distortion*

$$\delta = C \frac{\rho d_T}{\max\{\ell_*(T), d_T\}}. \quad (1.3)$$

Then for $T \subset \mathbb{R}^n$, with probability at least $1 - \eta$,

$$\sup_{t \in T} \left| \|A_\varepsilon t\|_2^2 - \|t\|_2^2 \right| \leq \max\{\rho^2, \rho\} d_T^2. \quad (1.4)$$

To put Theorem 1.9 in some context, if the belief is that A_ε should exhibit the same behaviour as the gaussian matrix $m^{-1/2}\Gamma$, then (keeping in mind that δ should scale like $1/\sqrt{m}$), “a gaussian behaviour” as in Theorem 1.5 is that with high probability,

$$\sup_{t \in T} \left| \|A_\varepsilon t\|_2^2 - \|t\|_2^2 \right| \leq c (\delta d_T \ell_*(T) + (\delta \ell_*(T))^2).$$

Observe that $\ell_*(T) \gtrsim d_T$, implying by (1.3) that $\rho \sim \delta \ell_*(T)/d_T$. Hence, the error in (1.4) in terms of δ is indeed

$$\sim d_T \delta \ell_*(T) + (\delta \ell_*(T))^2.$$

However, despite the “gaussian error”, the probability estimate in Theorem 1.9 is far weaker than in Theorem 1.5—it is just at the constant level.

Our main result is that using a modified, seemingly less restrictive version of the multi-scale RIP, A_ε acts on T as if it were a gaussian operator: achieving the same distortion and probability estimate as in Theorem 1.5.

Definition 1.10. *Let $A : \mathbb{R}^n \rightarrow \mathbb{R}^m$ be a matrix. For $\delta > 0$ let $1 \leq k^* \leq n$ be the largest such that for every $1 \leq k \leq k_*$, A is a (δ, k) regular; that is, for every $1 \leq k \leq k_*$*

$$\sup_{x \in \Sigma_k} \left| \|Ax\|_2^2 - \|x\|_2^2 \right| \leq \delta \sqrt{k}.$$

Theorem 1.11. *There exist absolute constants c and C such that the following holds. Let $\delta > 0$ and set $\Lambda = \max\{1, \delta^2 \log(n\delta^2)\}$. If $k_* \geq 1/\delta^2$, $T \subset \mathbb{R}^n$ and $u \geq 1$, then with probability at least*

$$1 - 2 \exp \left(-cu^2 \left(\frac{\ell_*(T)}{d_T} \right)^2 \right),$$

we have that

$$\sup_{t \in T} \left| \|A_\varepsilon t\|_2^2 - \|t\|_2^2 \right| \leq Cu^2 (\Lambda \cdot d_T \delta \ell_*(T) + (\delta \ell_*(T))^2). \quad (1.5)$$

Remark 1.12. *The sub-optimality in Theorem 1.11 lies in the factor Λ —in the range where δ is relatively large: at least $1/\sqrt{\log n}$. For $\delta \leq 1/\sqrt{\log n}$ we have that $\Lambda = 1$ and Theorem 1.11 recovers the functional Bernstein inequality for $u \sim 1$; that holds despite the fact that A_ε is based only on n “random bits”.*

Moreover, for the error in (1.5) to have a chance of being a nontrivial two-sided estimate, i.e., that for some $0 < \rho < 1$ and every $t \in T$,

$$\| \|A_\varepsilon t\|_2^2 - \|t\|_2^2 \| \leq \rho d_T^2,$$

δ has to be smaller than $\sim d_T/\ell_(T)$. In particular, if the critical dimension of T , $(\ell_*(T)/d_T)^2$ is at least $\log n$, a choice of $\delta \leq (d_T/\ell_*(T))$ leads to $\Lambda = 1$ and thus to an optimal outcome in Theorem 1.11.*

Theorem 1.11 clearly improves the probability estimate from Theorem 1.9. The other (virtual) improvement is that the matrix A need only be (δ, k) -regular for $k \leq 1/\delta^2$, and the way A acts on Σ_k for $k > 1/\delta^2$ is of no importance. The reason for calling that improvement “virtual” is the following observation:

Lemma 1.13. *If $k^* \geq 1/\delta^2$ then for any $1 \leq s \leq n$,*

$$\sup_{x \in \Sigma_s} \left| \|Ax\|_2^2 - 1 \right| \leq 4 \max\{\delta\sqrt{s}, \delta^2 s\}.$$

In other words, the second term in the multi-scale RIP condition follows automatically from the first one and the fact that k^* is sufficiently large.

Proof. Let $x \in \Sigma_s$ for $s \geq k_*$, and let $(J_i)_{i=1}^\ell$ be a decomposition of the support of x to coordinate blocks of cardinality $k_*/2$. Set $y_i = P_{J_i}x$, that is, the projection of x onto $\text{span}\{e_m : m \in J_i\}$ and write $x = \sum_{i=1}^\ell y_i$. Note that $\ell \leq 4s/k_*$ and that

$$\|Ax\|_2^2 = \left\| \sum_{i=1}^\ell Ay_i \right\|_2^2 = \sum_{i=1}^\ell \|Ay_i\|_2^2 + \sum_{i \neq j} \langle Ay_i, Ay_j \rangle.$$

The vectors y_i are orthogonal and so $\|x\|_2^2 = \sum_{i=1}^\ell \|y_i\|_2^2$. Therefore,

$$\left| \|Ax\|_2^2 - \|x\|_2^2 \right| \leq \sum_{i=1}^\ell \left| \|Ay_i\|_2^2 - \|y_i\|_2^2 \right| + \left| \sum_{i \neq j} \langle Ay_i, Ay_j \rangle \right|.$$

For the first term, as each y_i is supported on at most $k_*/2$ coordinates, it follows from the regularity condition that

$$\sum_{i=1}^\ell \left| \|Ay_i\|_2^2 - \|y_i\|_2^2 \right| \leq \delta\sqrt{k_*/2} \sum_{i=1}^\ell \|y_i\|_2^2 = \delta\sqrt{k_*/2} \|x\|_2^2 \leq \delta\sqrt{s}.$$

As for the second term, since y_i and y_j are orthogonal, $\|y_i + y_j\|_2 = \|y_i - y_j\|_2$ and

$$\begin{aligned} \langle Ay_i, Ay_j \rangle &= \frac{1}{4} (\|A(y_i + y_j)\|_2^2 - \|A(y_i - y_j)\|_2^2) \\ &= \frac{1}{4} (\|A(y_i + y_j)\|_2^2 - \|y_i + y_j\|_2^2) - \frac{1}{4} (\|y_i - y_j\|_2^2 - \|A(y_i - y_j)\|_2^2). \end{aligned}$$

Thus, by the regularity of A and as $|\text{supp}(y_i \pm y_j)| \leq k_*$,

$$\begin{aligned} |\langle Ay_i, Ay_j \rangle| &\leq \frac{1}{4} \left(\delta \sqrt{k_*} \|y_i + y_j\|_2^2 + \delta \sqrt{k_*} \|y_i - y_j\|_2^2 \right) \\ &\leq \frac{1}{2} \delta \sqrt{k_*} (\|y_i\|_2^2 + \|y_j\|_2^2). \end{aligned}$$

Taking the sum over all pairs $i \neq j$, $i, j \leq \ell$, each factor $\|y_i\|_2^2$ appears at most 2ℓ times, and $2\ell \leq 8s/k_*$. Hence, using that $1/\delta^2 \leq k_*$

$$\sum_{i \neq j} |\langle Ay_i, Ay_j \rangle| \leq \frac{1}{2} \delta \sqrt{k_*} \cdot \frac{8s}{k_*} \sum_{i=1}^{\ell} \|y_i\|_2^2 \leq 2\delta \frac{s}{\sqrt{k_*}} \|x\|_2^2 \leq 4\delta^2 s.$$

■

Clearly, Theorem 1.11 implies a suitable version of Theorem 1.9.

Corollary 1.14. *There exist absolute constants c and c_1 such that the following holds. Let A be as above, set $T \subset \mathbb{R}^n$ and $0 < \delta < 1/\sqrt{\log n}$. Let $\rho = c\delta \ell_*(T)/d_T$. Then with probability at least $1 - 2\exp(-c_1\rho^2/\delta^2)$,*

$$\sup_{t \in T} \left| \|A_\varepsilon t\|_2^2 - \|t\|_2^2 \right| \leq \rho d_T^2.$$

Remark 1.15. *Recalling the intuition that $m \sim 1/\delta^2$, the outcome of Corollary 1.14 coincides with the estimate in (1.2).*

In Section 3 we present one simple application of Theorem 1.11. We show that column randomization of a typical realization of a Bernoulli circulant matrix (complete or partial) exhibits an almost gaussian behaviour (conditioned on the generating vector). In particular, only $2n$ random bits (n from the generating Bernoulli vector and n from the column randomization) are required if one wishes to create a random ensemble that is, effectively, an almost isometry.

The proof of Theorem 1.11 is based on a chaining argument. For more information on the *generic chaining* mechanism, see Talagrand's treasured manuscript [10]. We only require relatively basic notions from generic chaining theory, as well as the celebrated *majorizing measures theorem*.

Definition 1.16. *Let $T \subset \mathbb{R}^n$. A collection of subsets of T , $(T_s)_{s \geq 0}$, is an admissible sequence if $|T_0| = 1$ and for $s \geq 1$, $|T_s| \leq 2^{2^s}$. For every $t \in T$ denote by $\pi_s t$ a nearest point to t in T_s with respect to the Euclidean distance. Set $\Delta_s t = \pi_{s+1} t - \pi_s t$ for $s \geq 1$ and let $\Delta_0 t = \pi_0 t$.*

The γ_2 functional with respect to the ℓ_2 metric is defined by

$$\gamma_2(T, \|\cdot\|_2) = \inf_{(T_s)} \sup_{t \in T} \sum_{s \geq 0} 2^{s/2} \|\Delta_s t\|_2,$$

where the infimum is taken with respect to all admissible sequences of T .

An application of Talagrand's *majorizing measures theorem* to the gaussian process $t \rightarrow \sum_{i=1}^n g_i t_i$ shows that $\gamma_2(T, \|\cdot\|_2)$ and $\ell_*(T)$ are equivalent:

Theorem 1.17. *There are absolute constants c and C such that for every $T \subset \mathbb{R}^n$,*

$$c\gamma_2(T, \|\cdot\|_2) \leq \ell_*(T) \leq C\gamma_2(T, \|\cdot\|_2).$$

The proof of Theorem 1.17 can be found, for example, in [10].

2 Proof of Theorem 1.11

We begin the proof with a word about notation: throughout, absolute constants, that is, positive numbers that are independent of all the parameters involved in the problem, are denoted by c , c_1 , C , etc. Their value may change from line to line.

As noted previously, the proof is based on a chaining argument. Let $(T_s)_{s \geq 0}$ be an optimal admissible sequence of T . Set s_0 to satisfy that 2^{s_0} is the critical dimension of T , i.e.,

$$2^{s_0} = \left(\frac{\ell_*(T)}{d_T} \right)^2$$

(without loss of generality we may assume that equality holds). Let $s_0 \leq s_1$ to be named in what follows and observe that

$$\|A_\varepsilon t\|_2^2 = \|A_\varepsilon(t - \pi_{s_1}t) + A_\varepsilon \pi_{s_1}t\|_2^2 = \|A_\varepsilon(t - \pi_{s_1}t)\|_2^2 + 2\langle A_\varepsilon(t - \pi_{s_1}t), A_\varepsilon \pi_{s_1}t \rangle + \|A_\varepsilon \pi_{s_1}t\|_2^2.$$

Writing $t - \pi_{s_1}t = \sum_{s \geq s_1} \Delta_s t$, it follows that

$$\|A_\varepsilon(t - \pi_{s_1}t)\|_2 \leq \sum_{s \geq s_1} \|A_\varepsilon \Delta_s t\|_2, \quad (2.1)$$

and

$$|\langle A_\varepsilon(t - \pi_{s_1}t), A_\varepsilon \pi_{s_1}t \rangle| \leq \|A_\varepsilon(t - \pi_{s_1}t)\|_2 \cdot \|A_\varepsilon \pi_{s_1}t\|_2;$$

Therefore, setting

$$\Psi^2 = \sup_{t \in T} \left| \|A_\varepsilon \pi_{s_1}t\|_2^2 - \|\pi_{s_1}t\|_2^2 \right| \quad \text{and} \quad \Phi = \sum_{s \geq s_1} \|A_\varepsilon \Delta_s t\|_2$$

we have that for every $t \in T$,

$$\|A_\varepsilon \pi_{s_1}t\|_2 \leq \sqrt{\Psi^2 + d_T^2}$$

and

$$|\langle A_\varepsilon(t - \pi_{s_1}t), A_\varepsilon \pi_{s_1}t \rangle| \leq \Phi \cdot \sqrt{\Psi^2 + d_T^2}. \quad (2.2)$$

Hence,

$$\sup_{t \in T} \left| \|A_\varepsilon t\|_2^2 - \|t\|_2^2 \right| \leq \Psi^2 + 2\Phi \cdot \sqrt{\Psi^2 + d_T^2} + \Phi^2 + \sup_{t \in T} \left| \|\pi_{s_1}t\|_2^2 - \|t\|_2^2 \right|. \quad (2.3)$$

To estimate the final term, note that for every $t \in T$,

$$\begin{aligned} \left| \|t\|_2^2 - \|\pi_{s_1}t\|_2^2 \right| &\leq \|t - \pi_{s_1}t\|_2^2 + 2|\langle t - \pi_{s_1}t, \pi_{s_1}t \rangle| \leq \|t - \pi_{s_1}t\|_2^2 + 2\|t - \pi_{s_1}t\|_2 \cdot \|\pi_{s_1}t\|_2 \\ &\leq \left(\sum_{s \geq s_1} \|\Delta_s t\|_2 \right)^2 + 2d_T \sum_{s \geq s_1} \|\Delta_s t\|_2. \end{aligned}$$

By the definition of the γ_2 functional and the majorizing measures theorem, for every integer s and every $t \in T$,

$$\|\Delta_s t\|_2 \leq 2^{-s/2} \gamma_2(T, \|\cdot\|_2) \leq c_1 2^{-s/2} \ell_*(T).$$

Thus,

$$\sum_{s \geq s_1} \|\Delta_s t\|_2 \leq c_1 \ell_*(T) \sum_{s \geq s_1} 2^{-s/2} \leq c_2 2^{-s_1/2} \ell_*(T),$$

and

$$\left| \|\pi_{s_1} t\|_2^2 - \|t\|_2^2 \right| \leq c_3 \left(2^{-s_1} \ell_*^2(T) + d_T 2^{-s_1/2} \ell_*(T) \right). \quad (2.4)$$

Equation (2.4) and the wanted estimate in Theorem 1.11 hint on the identity of 2^{s_1} : it should be larger than $1/\delta^2$. Recalling that $s_0 \leq s_1$ and that $2^{s_0} = (\ell_*(T)/d_T)^2$, set

$$2^{s_1} = \max \left\{ \frac{1}{\delta^2}, \left(\frac{\ell_*(T)}{d_T} \right)^2, \log(e(1+n\delta^2)) \right\}.$$

The reason behind the choice of the third term will become clear in what follows.

With that choice of s_1 ,

$$\sup_{t \in T} \left| \|\pi_{s_1} t\|_2^2 - \|t\|_2^2 \right| \leq c_3 \left(\delta^2 \ell_*^2(T) + d_T \delta \ell_*(T) \right), \quad (2.5)$$

and the nontrivial part of the proof is to control Φ and Ψ with high probability that would lead to the wanted estimate on (2.3).

2.1 A decoupling argument

For every $t \in \mathbb{R}^n$,

$$\|A_\varepsilon t\|_2^2 = \sum_{i,j} \langle Ae_i, Ae_j \rangle \varepsilon_i \varepsilon_j t_i t_j = \sum_{i=1}^n \|Ae_i\|_2^2 t_i^2 + \sum_{i \neq j} \langle Ae_i, Ae_j \rangle \varepsilon_i \varepsilon_j t_i t_j. \quad (2.6)$$

By the assumption that A is $(\delta, 1)$ -regular,

$$\max_{1 \leq i \leq n} \left| \|Ae_i\|_2^2 - 1 \right| \leq \delta,$$

and noting that $d_T \leq c \ell_*(T)$,

$$\left| \sum_{i=1}^n \|Ae_i\|_2^2 t_i^2 - \|t\|_2^2 \right| = \left| \sum_{i=1}^n (\|Ae_i\|_2^2 - 1) t_i^2 \right| \leq \delta \|t\|_2^2 \leq \delta d_T^2 \leq c d_T \delta \ell_*(T).$$

Next, we turn to the ‘‘off-diagonal’’ term in (2.6). For $t \in \mathbb{R}^n$ let

$$Z_t = \sum_{i \neq j} \varepsilon_i \varepsilon_j t_i t_j \langle Ae_i, Ae_j \rangle$$

and let us obtain high probability estimates on $\sup_{u \in U} |Z_u|$ for various sets U .

The first step in that direction is decoupling: let η_1, \dots, η_n be independent $\{0, 1\}$ -valued random variables with mean $1/2$. Set $I = \{i : \eta_i = 1\}$ and observe that for every $(\varepsilon_i)_{i=1}^n$,

$$\sup_{u \in U} |Z_u| \leq 4 \sup_{u \in U} \mathbb{E}_\eta \left| \left\langle A \left(\sum_{i \in I} \varepsilon_i u_i e_i \right), A \left(\sum_{j \in I^c} \varepsilon_j u_j e_j \right) \right\rangle \right|. \quad (2.7)$$

Indeed, for every $(\varepsilon_i)_{i=1}^n$,

$$\begin{aligned} \sup_{u \in U} \left| \sum_{i \neq j} \langle Ae_i, Ae_j \rangle \varepsilon_i \varepsilon_j u_i u_j \right| &= 4 \sup_{u \in U} \left| \sum_{i \neq j} \langle Ae_i, Ae_j \rangle \mathbb{E}_\eta (1 - \eta_i) \eta_j \cdot \varepsilon_i \varepsilon_j u_i u_j \right| \\ &\leq 4 \sup_{u \in U} \mathbb{E}_\eta \left| \sum_{i \in I, j \in I^c} \langle Ae_i, Ae_j \rangle \varepsilon_i \varepsilon_j u_i u_j \right|; \end{aligned}$$

and for every $u \in \mathbb{R}^n$,

$$\sum_{i \in I, j \in I^c} \langle Ae_i, Ae_j \rangle \varepsilon_i \varepsilon_j u_i u_j = \left\langle A \left(\sum_{j \in I^c} \varepsilon_j u_j e_j \right), A \left(\sum_{i \in I} \varepsilon_i u_i e_i \right) \right\rangle.$$

Equation (2.7) naturally leads to the following definition:

Definition 2.1. For $v \in \mathbb{R}^n$ and $I \subset \{1, \dots, n\}$, set

$$W_{v,I} = A^* A \left(\sum_{i \in I} \varepsilon_i v_i e_i \right). \quad (2.8)$$

Recall that $\pi_{s+1}t$ is the nearest point to t in T_{s+1} and $\Delta_s t = \pi_{s+1}t - \pi_s t$.

Lemma 2.2. For every t and every $(\varepsilon_i)_{i=1}^n$,

$$\begin{aligned} |Z_{\pi_{s_1}t}| &\leq 4 \sum_{s=s_0}^{s_1-1} \left(\mathbb{E}_\eta \left| \sum_{j \in I^c} \varepsilon_j (\Delta_s t)_j (W_{\pi_{s+1}t,I})_j \right| + \mathbb{E}_\eta \left| \sum_{i \in I} \varepsilon_i (\Delta_s t)_i (W_{\pi_s t, I^c})_i \right| \right) \\ &\quad + 4 \mathbb{E}_\eta \left| \sum_{j \in I^c} \varepsilon_j (\pi_{s_0} t)_j (W_{\pi_{s_0}t,I})_j \right|. \end{aligned} \quad (2.9)$$

Proof. Fix an integer s . With the decoupling argument in mind, fix $I \subset \{1, \dots, n\}$ and observe that

$$\begin{aligned} &\left\langle A \left(\sum_{i \in I} \varepsilon_i (\pi_{s+1}t)_i e_i \right), A \left(\sum_{j \in I^c} \varepsilon_j (\pi_{s+1}t)_j e_j \right) \right\rangle \\ &= \left\langle A \left(\sum_{i \in I} \varepsilon_i (\pi_{s+1}t)_i e_i \right), A \left(\sum_{j \in I^c} \varepsilon_j (\Delta_s t)_j e_j \right) \right\rangle + \left\langle A \left(\sum_{i \in I} \varepsilon_i (\pi_{s+1}t)_i e_i \right), A \left(\sum_{j \in I^c} \varepsilon_j (\pi_s t)_j e_j \right) \right\rangle \\ &= \left\langle A \left(\sum_{i \in I} \varepsilon_i (\pi_{s+1}t)_i e_i \right), A \left(\sum_{j \in I^c} \varepsilon_j (\Delta_s t)_j e_j \right) \right\rangle + \left\langle A \left(\sum_{i \in I} \varepsilon_i (\Delta_s t)_i e_i \right), A \left(\sum_{j \in I^c} \varepsilon_j (\pi_s t)_j e_j \right) \right\rangle \\ &\quad + \left\langle A \left(\sum_{i \in I} \varepsilon_i (\pi_s t)_i e_i \right), A \left(\sum_{j \in I^c} \varepsilon_j (\pi_s t)_j e_j \right) \right\rangle. \end{aligned}$$

Moreover,

$$\begin{aligned} \left\langle A \left(\sum_{i \in I} \varepsilon_i (\pi_{s+1}t)_i e_i \right), A \left(\sum_{j \in I^c} \varepsilon_j (\Delta_s t)_j e_j \right) \right\rangle &= \left\langle A^* A \left(\sum_{i \in I} \varepsilon_i (\pi_{s+1}t)_i e_i \right), \left(\sum_{j \in I^c} \varepsilon_j (\Delta_s t)_j e_j \right) \right\rangle \\ &= \sum_{j \in I^c} \varepsilon_j (\Delta_s t)_j (W_{\pi_{s+1}t,I})_j. \end{aligned}$$

and

$$\left\langle A\left(\sum_{i \in I} \varepsilon_i(\Delta_s t)_i e_i\right), A\left(\sum_{j \in I^c} \varepsilon_j(\pi_{s t})_j e_j\right) \right\rangle = \sum_{i \in I} \varepsilon_i(\Delta_s t)_i (W_{\pi_{s t}, I^c})_i.$$

Combining these observations, for every $t \in T$ and $(\varepsilon_i)_{i=1}^n$,

$$\begin{aligned} \frac{1}{4} Z_{\pi_{s_1} t} &= \mathbb{E}_\eta \left\langle A\left(\sum_{i \in I} \varepsilon_i(\pi_{s_1} t)_i e_i\right), A\left(\sum_{j \in I^c} \varepsilon_j(\pi_{s_1} t)_j e_j\right) \right\rangle \\ &= \mathbb{E}_\eta \sum_{s=s_0}^{s_1-1} \sum_{i \in I} \varepsilon_i(\Delta_s t)_i (W_{\pi_{s t}, I^c})_i \\ &\quad + \mathbb{E}_\eta \sum_{s=s_0}^{s_1-1} \sum_{j \in I^c} \varepsilon_j(\Delta_s t)_j (W_{\pi_{s+1} t, I})_j \\ &\quad + \mathbb{E}_\eta \sum_{j \in I^c} \varepsilon_j(\pi_{s_0} t)_j (W_{\pi_{s_0} t, I})_j, \end{aligned}$$

from which the claim follows immediately. \blacksquare

As part of the decoupling argument and to deal with the introduction of the random variables $(\eta_i)_{i=1}^n$ in (2.7), we will use the following elementary fact:

Lemma 2.3. *Let f be a function of $(\varepsilon_i)_{i=1}^n$ and $(\eta_i)_{i=1}^n$. If $\mathbb{E}_\eta \mathbb{E}_\varepsilon |f|^q \leq \kappa^q$ then with $(\varepsilon_i)_{i=1}^n$ -probability at least $1 - \exp(-q)$,*

$$\mathbb{E}_\eta (|f| \mid (\varepsilon_i)_{i=1}^n) \leq e\kappa. \quad (2.10)$$

Proof. For a nonnegative function h we have that point-wise, $\mathbb{1}_{\{h \geq t\}} \leq h^q/t^q$. Let $h = \mathbb{E}_\eta (|f| \mid (\varepsilon_i)_{i=1}^n)$ and note that

$$Pr_\varepsilon (\mathbb{E}_\eta (|f| \mid (\varepsilon_i)_{i=1}^n) \geq u\kappa) = \mathbb{E}_\varepsilon \mathbb{1}_{\{h \geq u\kappa\}} \leq (u\kappa)^{-q} (\mathbb{E}_\varepsilon h^q).$$

By Jensen's inequality followed by Fubini's Theorem,

$$\mathbb{E}_\varepsilon h^q = \mathbb{E}_\varepsilon (\mathbb{E}_\eta (|f| \mid (\varepsilon_i)_{i=1}^n)^q) \leq \mathbb{E}_\varepsilon \mathbb{E}_\eta |f|^q \leq \kappa^q,$$

and setting $u = e$ proves (2.10). \blacksquare

We shall use Lemma 2.3 in situations where we actually have more information—namely that for any $(\eta_i)_{i=1}^n$, $\mathbb{E}_\varepsilon (|f|^q \mid (\eta_i)_{i=1}^n) \leq \kappa^q$ for a well chosen κ . As a result,

$$\mathbb{E}_\eta (|f| \mid (\varepsilon_i)_{i=1}^n) \leq e\kappa \quad \text{with probability at least } 1 - \exp(-q).$$

Taking into account Lemma 2.2 and Lemma 2.3, it follows that if one wishes to estimate $\sup_{t \in T} |Z_{\pi_{s_1} t}|$ using a chaining argument, it suffices to obtain, for every $I \subset \{1, \dots, n\}$, bounds on moments of random variables of the form

$$\sum_{j \in I^c} \varepsilon_j(\Delta_s t)_j (W_{\pi_{s+1} t, I})_j, \quad \sum_{i \in I} \varepsilon_i(\Delta_s t)_i (W_{\pi_{s t}, I^c})_i, \quad \text{and} \quad \sum_{j \in I^c} \varepsilon_j(\pi_{s_0} t)_j (W_{\pi_{s_0} t, I})_j, \quad (2.11)$$

as that results in high probability estimates on each of the terms in (2.9). And as there are at most $2^{2^{s+3}}$ random variables involved in this chaining argument at the s -stage, the required moment in (2.11) is $q \sim 2^s$ for the first two terms and $q \sim 2^{s_0}$ for the third one.

2.2 Preliminary estimates

For $J \subset \{1, \dots, n\}$ let $P_J x = \sum_{j \in J} x_j e_j$ be the projection of x onto $\text{span}(e_j)_{j \in J}$. The key lemma in the proof of Theorem 1.11 is:

Lemma 2.4. *There exists an absolute constant c such that the following holds. Let $I \subset \{1, \dots, n\}$ and $W_{v,I}$ be as in (2.8). Set $J \subset I^c$ such that $|J| \leq k_*$. Then for $q \geq |J|$,*

$$(\mathbb{E} \|P_J W_{v,I}\|_2^q)^{1/q} \leq c\sqrt{q}\delta \|v\|_2.$$

Proof. Let S^J be the Euclidean unit sphere in the subspace $\text{span}(e_j)_{j \in J}$ and let U be a maximal $1/10$ -separated subset of S^J . By a volumetric estimate (see, e.g. [1]), there is an absolute constant c_0 such that $|U| \leq \exp(c_0|J|)$. Moreover, a standard approximation argument shows that for every $y \in \mathbb{R}^n$,

$$\|P_J y\|_2 = \sup_{x \in S^J} \langle y, x \rangle \leq c_1 \max_{x \in U} \langle y, x \rangle,$$

where c_1 is a suitable absolute constant. Therefore,

$$\sup_{x \in S^J} \langle W_{v,I}, x \rangle \leq c_1 \max_{x \in U} \langle W_{v,I}, x \rangle,$$

and it suffices to control, with high probability,

$$\max_{x \in U} \left\langle \sum_{i \in I} \varepsilon_i v_i e_i, A^* A x \right\rangle = \max_{x \in U} \sum_{i \in I} \varepsilon_i v_i (A^* A x)_i.$$

Fix $x \in U$, recall that A is (δ, k) -regular for $1 \leq k \leq k_*$ and we first explore the case $1 \leq q \leq k_*/2$.

Denote by $I' \subset I$ the set of indices corresponding to the q largest values of $|(A^* A x)_i|$, $i \in I$. If $|I| \leq q$ then set $I' = I$.

It is straightforward to verify (e.g., using Höfdding's inequality) that there is an absolute constant c_2 such that

$$\begin{aligned} \left\| \sum_{i \in I} \varepsilon_i v_i (A^* A x)_i \right\|_{L_q} &\leq \sum_{i \in I'} |v_i (A^* A x)_i| + c_2 \sqrt{q} \left(\sum_{i \in I \setminus I'} v_i^2 (A^* A x)_i^2 \right)^{1/2} \\ &\leq \|v\|_2 \|P_{I'}(A^* A x)\|_2 + c_2 \sqrt{q} \cdot \frac{\|P_{I'}(A^* A x)\|_2}{\sqrt{q}} \|v\|_2 \\ &\leq c_3 \|v\|_2 \|P_{I'}(A^* A x)\|_2, \end{aligned}$$

where we used that fact that for $i \in I \setminus I'$,

$$|(A^* A x)_i| \leq \frac{\|P_{I'}(A^* A x)\|_2}{\sqrt{q}}.$$

Therefore,

$$\left\| \sum_{i \in I} \varepsilon_i v_i (A^* A x)_i \right\|_{L_q} \leq c_3 \|v\|_2 \max_{I' \subset I, |I'|=q} \sup_{z \in S^{I'}} \langle A^* A x, z \rangle.$$

Note that x is supported in $J \subset I^c$, while each ‘legal’ z is supported in a subset of I ; in particular, x and z are orthogonal, implying that for every such z ,

$$\|x + z\|_2 = \|x - z\|_2 \leq 2 \quad \text{and} \quad |\text{supp}(x + z)|, |\text{supp}(x - z)| \leq 2q.$$

Thus, by the $(\delta, 2q)$ -regularity of A (as $2q \leq k_*$),

$$\begin{aligned} |\langle A^* Ax, z \rangle| &= |\langle Az, Ax \rangle| = \frac{1}{4} \left| \|A(x + z)\|_2^2 - \|A(x - z)\|_2^2 \right| \\ &= \frac{1}{4} \left| (\|A(x + z)\|_2^2 - \|x + z\|_2^2) - (\|A(x - z)\|_2^2 - \|x - z\|_2^2) \right| \\ &\leq \frac{\delta}{4} \sqrt{2q} \cdot \max\{\|x + z\|_2^2, \|x - z\|_2^2\} \leq \delta \sqrt{q}, \end{aligned}$$

and it follows that

$$\left\| \sum_{i \in I} \varepsilon_i v_i (A^* Ax)_i \right\|_{L_q} \leq c_4 \|v\|_2 \delta \sqrt{q}.$$

Turning to the case $q \geq k_*/2$, let I' be the set of indices corresponding to the $k_*/2$ largest coordinates of $(|(A^* Ax)_i|)_{i \in I}$. Therefore,

$$\begin{aligned} \left\| \sum_{i \in I} \varepsilon_i v_i (A^* Ax)_i \right\|_{L_q} &\leq \sum_{i \in I'} v_i (A^* Ax)_i + c_2 \sqrt{q} \left(\sum_{i \in I \setminus I'} v_i^2 (A^* Ax)_i^2 \right)^{1/2} \\ &\leq \|v\|_2 \cdot \|P_{I'} A^* Ax\|_2 \left(1 + c_5 \sqrt{\frac{q}{k_*}} \right), \end{aligned} \quad (2.12)$$

using that for $i \in I \setminus I'$,

$$|(A^* Ax)_i| \leq \frac{\|P_{I'} A^* Ax\|_2}{\sqrt{k_*/2}}.$$

Recall that $|\text{supp}(x)| = |J| \leq k_*$ and that $|I'| = k_*/2$. The same argument used previously shows that

$$\|P_{I'} A^* Ax\|_2 \leq c_6 \delta \sqrt{k_*};$$

hence,

$$\left\| \sum_{i \in I} \varepsilon_i v_i (A^* Ax)_i \right\|_{L_q} \leq c_7 \|v\|_2 \delta \sqrt{q},$$

and the estimate holds for each $q \geq |J|$ for that fixed x .

Setting $u \geq 1$, it follows from Chebychev’s inequality that with probability at least $1 - 2 \exp(-c_8 u^2 q)$,

$$\left| \sum_{i \in I} \varepsilon_i v_i (A^* Ax)_i \right| \leq c_9 u \|v\|_2 \delta \sqrt{q},$$

and by the union bound, recalling that $q \geq |J|$, the same estimate holds uniformly for every $x \in U$, provided that $u \geq c_{10}$. Thus, with probability at least $1 - 2 \exp(-cu^2 q)$,

$$\|P_J W_{v,I}\|_2 \leq c' \max_{x \in U} \left| \sum_{i \in I} \varepsilon_i v_i (A^* Ax)_i \right| \leq C u \|v\|_2 \delta \sqrt{q},$$

and the wanted estimate follows from tail integration. ■

The next observation deals with more refined estimates on random variables of the form

$$X_{a,b} = \sum_{i \in I^c} \varepsilon_i a_i b_i.$$

Once again, we use the fact that for any $J \subset I^c$

$$\|X_{a,b}\|_{L_q} \leq \sum_{j \in J} |a_j b_j| + c\sqrt{q} \left(\sum_{j \in I^c \setminus J} a_j^2 b_j^2 \right)^{1/2}. \quad (2.13)$$

As one might have guessed, the choice of b will be vectors of the form $W_{\pi_{st}, I}$. These are random vectors that are independent of the Bernoulli random variables involved in the definition of X . At the same time, a will be deterministic.

Without loss of generality assume that $a_i, b_i \geq 0$ for every i . Let J_1 be the set of indices corresponding to the k_1 largest coordinates of $(a_i)_{i \in I^c}$, J_2 is the set corresponding to the following k_2 largest coordinates, and so on. The choice of k_1, k_2, \dots , will be specified in what follows.

Note that for any $\ell > 1$,

$$\|P_{J_\ell} a\|_\infty \leq \frac{\|P_{J_{\ell-1}} a\|_2}{\sqrt{|J_{\ell-1}|}} = \frac{\|P_{J_{\ell-1}} a\|_2}{\sqrt{k_{\ell-1}}}.$$

Set $J = J_1$, and by (2.13),

$$\begin{aligned} \|X_{a,b}\|_{L_q} &\leq \sum_{j \in J_1} |a_j b_j| + c\sqrt{q} \left(\sum_{\ell \geq 1} \sum_{j \in J_{\ell+1}} a_j^2 b_j^2 \right)^{1/2} \\ &\leq \|P_{J_1} a\|_2 \|P_{J_1} b\|_2 + c\sqrt{q} \left(\sum_{\ell \geq 1} \|P_{J_{\ell+1}} a\|_\infty^2 \|P_{J_{\ell+1}} b\|_2^2 \right)^{1/2} \\ &\leq \|P_{J_1} a\|_2 \|P_{J_1} b\|_2 + c\sqrt{q} \left(\sum_{\ell \geq 1} \frac{\|P_{J_\ell} a\|_2^2}{|J_\ell|} \|P_{J_{\ell+1}} b\|_2^2 \right)^{1/2} \\ &\leq \|P_{J_1} a\|_2 \|P_{J_1} b\|_2 + c\sqrt{q} \max_{\ell \geq 2} \frac{\|P_{J_\ell} b\|_2}{\sqrt{k_{\ell-1}}} \cdot \left(\sum_{\ell \geq 2} \|P_{J_\ell} a\|_2^2 \right)^{1/2} \\ &\leq \|a\|_2 \left(\|P_{J_1} b\|_2 + c\sqrt{q} \max_{\ell \geq 2} \frac{\|P_{J_\ell} b\|_2}{\sqrt{k_{\ell-1}}} \right). \end{aligned} \quad (2.14)$$

And, in the case where $|J_\ell| = k$ for every ℓ , it follows that

$$\|X_{a,b}\|_{L_q} \leq \|a\|_2 \left(\|P_{J_1} b\|_2 + c\sqrt{\frac{q}{k}} \max_{\ell \geq 2} \|P_{J_\ell} b\|_2 \right). \quad (2.15)$$

2.3 Estimating Φ

Recall that

$$\Phi = \sup_{t \in T} \sum_{s \geq s_1} \|A_\varepsilon \Delta_s t\|_2$$

and that for every $t \in \mathbb{R}^n$,

$$Z_t = \sum_{i \neq j} \varepsilon_i \varepsilon_j t_i t_j \langle A e_i, A e_j \rangle.$$

Theorem 2.5. *There are absolute constants c and C such that for $u > 1$, with probability at least $1 - 2\exp(-cu^22^{s_1})$,*

$$\Phi^2 \leq Cu^2\delta^2\ell_*^2(T).$$

Proof. Let $s \geq s_1$, and as noted previously, for every $t \in T$,

$$\begin{aligned} \|A_\varepsilon \Delta_{st}\|_2^2 &= \sum_{i=1}^n \|Ae_i\|_2^2 (\Delta_{st})_i^2 + \sum_{i \neq j} \langle Ae_i, Ae_j \rangle \varepsilon_i \varepsilon_j (\Delta_{st})_i (\Delta_{st})_j \\ &\leq (1 + \delta) \|\Delta_{st}\|_2^2 + \left| \sum_{i \neq j} \langle Ae_i, Ae_j \rangle \varepsilon_i \varepsilon_j (\Delta_{st})_i (\Delta_{st})_j \right| \\ &\leq 2\|\Delta_{st}\|_2^2 + |Z_{\Delta_{st}}|. \end{aligned}$$

Following the decoupling argument, one may fix $I \subset \{1, \dots, n\}$. The core of the argument is to obtain satisfactory estimates on moments of the random variables

$$V_{\Delta_{st}, I} = \sum_{j \in I^c} \varepsilon_j (\Delta_{st})_j (W_{\Delta_{st}, I})_j$$

for that (arbitrary) fixed choice of I . And, since $|T_s| = 2^{2^s}$, for a uniform estimate that holds for every random variable of the form $V_{\Delta_{st}, I}$, $t \in T$, it is enough to control the q -th moment of each $V_{\Delta_{st}, I}$ for $q \sim 2^s$.

With that in mind, denote by \mathbb{E}_{I^c} the expectation taken with respect to $(\varepsilon_i)_{i \in I^c}$, and set \mathbb{E}_I the expectation taken with respect to $(\varepsilon_i)_{i \in I}$.

Let us apply (2.15), with the choice

$$k = 1/\delta^2 \leq k_*, \quad a = \Delta_{st} \quad \text{and} \quad b = W_{\Delta_{st}, I}.$$

Thus, for every $(\varepsilon_i)_{i \in I}$,

$$(\mathbb{E}_{I^c} |V_{\Delta_{st}, I}|^q)^{1/q} \leq \|\Delta_{st}\|_2 \left(\|P_{J_1} W_{\Delta_{st}, I}\|_2 + c_0 \delta \sqrt{q} \cdot \max_{\ell \geq 2} \|P_{J_\ell} W_{\Delta_{st}, I}\|_2 \right).$$

The sets J_ℓ are all of cardinality $1/\delta^2$ and so there are at most $\max\{\delta^2 n, 1\}$ of them. By Lemma 2.4, for each one of the sets $J_\ell \subset I^c$ and $q \geq |J_\ell| = 1/\delta^2$,

$$(\mathbb{E}_I \|P_{J_\ell} W_{\Delta_{st}, I}\|_2^q)^{1/q} \leq c_1 \delta \sqrt{q} \|\Delta_{st}\|_2,$$

implying that

$$\begin{aligned} \left(\mathbb{E}_I \max_{\ell} \|P_{J_\ell} W_{\Delta_{st}, I}\|_2^q \right)^{1/q} &\leq \left(\sum_{\ell} \mathbb{E}_I \|P_{J_\ell} W_{\Delta_{st}, I}\|_2^q \right)^{1/q} \\ &\leq \ell^{1/q} \cdot c_1 \delta \sqrt{q} \|\Delta_{st}\|_2 \leq c_2 \delta \sqrt{q} \|\Delta_{st}\|_2 \end{aligned} \quad (2.16)$$

as long as $\max\{\delta^2 n, 1\} \leq \exp(q)$. In particular, since $2^{s_1} \geq \log(e(1 + n\delta^2))$, it suffices that $q \geq 2^{s_1}$ to ensure that (2.16) holds.

Hence, for every $I \subset \{1, \dots, n\}$,

$$\begin{aligned}
(\mathbb{E}|V_{\Delta_s t, I}|^q)^{1/q} &= (\mathbb{E}_I \mathbb{E}_{I^c} |V_{\Delta_s t, I}|^q)^{1/q} \\
&\leq \|\Delta_s t\|_2 \left(\mathbb{E}_I \left(\|P_{J_1} W_{\Delta_s t, I}\|_2 + c_0 \delta \sqrt{q} \cdot \max_{\ell \geq 2} \|P_{J_\ell} W_{\Delta_s t, I}\|_2 \right)^q \right)^{1/q} \\
&\leq \|\Delta_s t\|_2 \cdot 2 \left((\mathbb{E}_I \|P_{J_1} W_{\Delta_s t, I}\|_2^q)^{1/q} + c_0 \delta \sqrt{q} \left(\mathbb{E}_I \max_{\ell \geq 2} \|P_{J_\ell} W_{\Delta_s t, I}\|_2^q \right)^{1/q} \right) \\
&\leq c_3 \|\Delta_s t\|_2^2 (\delta \sqrt{q} + \delta^2 q) \\
&\leq c_4 \delta^2 q \|\Delta_s t\|_2^2,
\end{aligned}$$

because $\delta^2 q \geq \delta^2 2^{s_1} \geq 1$.

By Jensen's inequality, for every $t \in T$,

$$\begin{aligned}
(\mathbb{E}|Z_{\Delta_s t}|^q)^{1/q} &\leq 4 \left(\mathbb{E}_\eta \mathbb{E}_\varepsilon \left| \sum_{i \in I^c} \varepsilon_i (\Delta_s t)_i (W_{\Delta_s t, I})_i \right|^q \right)^{1/q} \\
&= 4 (\mathbb{E}_\eta \mathbb{E}_\varepsilon |V_{\Delta_s t, I}|^q)^{1/q} \leq c_5 q \delta^2 \|\Delta_s t\|_2^2.
\end{aligned}$$

Setting $q = u2^s$ for $u \geq 1$ and $s \geq s_1$, Chebychev's inequality implies that with $(\varepsilon_i)_{i=1}^n$ -probability at least $1 - 2 \exp(-c_6 u^2 2^s)$,

$$\begin{aligned}
\|A_\varepsilon \Delta_s t\|_2^2 &= 2 \|\Delta_s t\|_2^2 + |Z_{\Delta_s t}|^2 \leq 2 \|\Delta_s t\|_2^2 + c_7 u^2 2^s \delta^2 \|\Delta_s t\|_2^2 \\
&\leq c_8 u^2 \delta^2 2^s \|\Delta_s t\|_2^2.
\end{aligned}$$

By the union bound, this estimate holds for every $t \in T$ and $s \geq s_1$. Thus, there are absolute constants c and C such that with probability at least $1 - 2 \exp(-cu^2 2^{s_1})$,

$$\sum_{s \geq s_1} \|A_\varepsilon \Delta_s t\|_2 \leq Cu \delta \sum_{s \geq s_1} 2^{s/2} \|\Delta_s t\|_2 \leq Cu \delta \ell_*(T),$$

as claimed. ■

2.4 Estimating Ψ

Next, recall that

$$\Psi^2 = \sup_{t \in T} \left| \|A_\varepsilon \pi_{s_1} t\|_2^2 - \|\pi_{s_1} t\|_2^2 \right|.$$

Expanding as in (2.6), the ‘‘diagonal term’’ is at most $\delta^2 d_T^2$, and one has to deal with the ‘‘off-diagonal’’ term

$$\sup_{t \in T} |Z_{\pi_{s_1} t}|.$$

As observed in Lemma 2.2 combined with Lemma 2.3, it suffices to obtain, for every fixed $I \subset \{1, \dots, n\}$, estimates on the moments of

$$\sum_{j \in I^c} \varepsilon_j (\Delta_s t)_j (W_{\pi_{s+1} t, I})_j \quad \sum_{i \in I} \varepsilon_i (\Delta_s t)_i (W_{\pi_s t, I^c})_i \quad \text{and} \quad \sum_{j \in I^c} \varepsilon_j (\pi_{s_0} t)_j (W_{\pi_{s_0} t, I})_j. \quad (2.17)$$

Remark 2.6. We shall assume throughout that $(\ell_*(T)/d_T)^2 \leq 1/\delta^2$; the required modifications when the reverse inequality holds are straightforward and are therefore omitted.

We begin with a standard observation:

Lemma 2.7. *There is an absolute constant c such that the following holds. Let $(X_\ell)_{\ell \geq 0}$ be nonnegative random variables, let $q \geq 1$ and set $q_\ell = q^{2^\ell}$. If there is some κ such that for every ℓ , $\|X_\ell\|_{L_{q_\ell}} \leq \kappa$, then*

$$\|\max_\ell X_\ell\|_{L_q} \leq c\kappa$$

Proof. By Chebychev's inequality, for every $\ell \geq 1$ and $u \geq 2$,

$$\Pr(|X_\ell| \geq u\kappa) \leq \frac{\mathbb{E}|X_\ell|^{q_\ell}}{(u\kappa)^{q_\ell}} \leq u^{-q_\ell}.$$

Therefore,

$$\Pr(\exists \ell \geq 2 : |X_\ell| \geq u\kappa) \leq \sum_{\ell \geq 2} u^{-q^{2^\ell}} \leq c_1 u^{-4q},$$

implying that

$$\mathbb{E} \max_{\ell \geq 2} |X_\ell|^q \leq \int_0^\infty qu^{q-1} \Pr\left(\max_{\ell \geq 1} |X_\ell| > u\right) du \leq (c_1 \kappa)^q.$$

Hence,

$$\|\max_\ell X_\ell\|_{L_q} \leq \|X_0\|_{L_q} + \|X_1\|_{L_q} + \|\max_{\ell \geq 2} X_\ell\|_{L_q} \leq c_2 \kappa. \quad \blacksquare$$

The analysis is split into two cases.

Case I: $\log(e(1 + \delta^2 n)) \leq 1/\delta^2$.

In this case, $2^{s_1} = 1/\delta^2$. For every $s_0 \leq s \leq s_1$ we invoke (2.14) for sets J_ℓ of cardinality $2^{s+\ell}$ when $s + \ell \leq s_1$, and of cardinality $1/\delta^2$ when $s + \ell > s_1$.

Set $a = \Delta_{s,t}$ and $b = W_{\pi_{s+1}t, I}$; the treatment when $b = W_{\pi_{s,t}, I^c}$ is identical and is omitted. Let $q \geq 2^s$ and put $q_\ell = q^{2^\ell}$ if $s + \ell \leq s_1$. Finally, set $p \geq 1/\delta^2$.

Consider ℓ such that $s + \ell \leq s_1$ and observe that $q_\ell \geq |J_\ell| = 2^{s+\ell}$. By Lemma 2.4,

$$|J_{\ell-1}|^{-1/2} \left(\mathbb{E}\|P_{J_\ell} W_{\pi_{s+1}t, I}\|_2^{q_\ell}\right)^{1/q_\ell} \leq c_1 \delta \|\pi_{s+1}t\|_2 \cdot \sqrt{\frac{q^{2^\ell}}{2^{s+\ell}}} \leq c_1 \sqrt{\frac{q}{2^s}} \delta d_T. \quad (2.18)$$

Therefore, by Lemma 2.7,

$$\left(\mathbb{E} \max_{\{\ell: s+\ell \leq s_1\}} \left(|J_{\ell-1}|^{-1/2} \|P_{J_\ell} W_{\pi_{s+1}t, I}\|_2\right)^q\right)^{1/q} \leq c_2 \sqrt{\frac{q}{2^s}} \delta d_T.$$

Next, if $s + \ell > s_1$ then $p \geq |J_\ell| = 1/\delta^2$ and by Lemma 2.4,

$$|J_{\ell-1}|^{-1/2} \left(\mathbb{E}\|P_{J_\ell} W_{\pi_{s+1}t, I}\|_2^p\right)^{1/p} \leq c_3 \delta \|\pi_{s+1}t\|_2 \cdot \sqrt{\frac{p}{1/\delta^2}} \leq c_3 \sqrt{p\delta^2} \cdot \delta d_T. \quad (2.19)$$

There are at most $n\delta^2$ sets J_ℓ in that range, and because $n\delta^2 \leq \exp(p)$, it is evident that

$$\left(\mathbb{E} \max_{\{\ell: s+\ell > s_1\}} \left(|J_{\ell-1}|^{-1/2} \|P_{J_\ell} W_{\pi_{s+1}t, I}\|_2 \right)^p \right)^{1/p} \leq c_4 \sqrt{p\delta^2} \cdot \delta d_T;$$

therefore, as $q \leq p$,

$$\left(\mathbb{E} \max_{\{\ell: s+\ell > s_1\}} \left(|J_{\ell-1}|^{-1/2} \|P_{J_\ell} W_{\pi_{s+1}t, I}\|_2 \right)^q \right)^{1/q} \leq c_4 \sqrt{p\delta^2} \delta d_T;$$

Thus, for every fixed $I \subset \{1, \dots, n\}$,

$$\left(\mathbb{E} \max_{\ell} \left(|J_{\ell-1}|^{-1/2} \|P_{J_\ell} W_{\pi_{s+1}t, I}\|_2 \right)^q \right)^{1/q} \leq c_5 \max \left\{ \sqrt{\frac{q}{2^s}}, \sqrt{p\delta^2} \right\} \cdot \delta d_T. \quad (2.20)$$

Now, by (2.15),

$$\begin{aligned} & \left(\mathbb{E}_{I^c} \left| \sum_{j \in I^c} \varepsilon_j(\Delta_s t)_j (W_{\pi_{s+1}t, I})_j \right|^q \right)^{1/q} \\ & \leq \|\Delta_s t\|_2 \left(\|P_{J_1} W_{\pi_{s+1}t, I}\|_2 + c_6 \sqrt{q} \max_{\ell \geq 2} \frac{\|P_{J_{\ell+1}} W_{\pi_{s+1}t, I}\|_2}{\sqrt{|J_\ell|}} \right), \end{aligned} \quad (2.21)$$

which, combined with (2.18) and (2.20), implies that

$$\left(\mathbb{E} \left| \sum_{j \in I^c} \varepsilon_j(\Delta_s t)_j (W_{\pi_{s+1}t, I})_j \right|^q \right)^{1/q} \leq c_7 \sqrt{q} \|\Delta_s t\|_2 \cdot \left(1 + \max \left\{ \sqrt{\frac{q}{2^s}}, \sqrt{p\delta^2} \right\} \right) \cdot \delta d_T. \quad (2.22)$$

Clearly, there are at most 2^{2s+3} random variables as in (2.22). With that in mind, set $u \geq 1$ and let $q = u^{2s}$, $p = u/\delta^2$. By Lemma 2.2 and Lemma 2.3 followed by the union bound, we have that with probability at least $1 - 2 \exp(-c_8 u^2 2^s)$, for every $t \in T$,

$$\mathbb{E}_\eta \left| \sum_{j \in I^c} \varepsilon_j(\Delta_s t)_j W_{\pi_{s+1}t, I} \right| \leq c_9 u^2 2^{s/2} \|\Delta_s t\|_2 \cdot \delta d_T. \quad (2.23)$$

And, by the union bound and recalling that $2^{s_0} = (\ell_*(T)/d_T)^2$, (2.23) holds uniformly for every $t \in T$ and $s_0 \leq s \leq s_1$ with probability at least

$$1 - 2 \exp \left(-c_{10} u^2 \left(\frac{\ell_*(T)}{d_T} \right)^2 \right).$$

An identical argument shows that with probability at least

$$1 - 2 \exp(-c_{11} u^2 2^{s_0}) = 1 - 2 \exp(-c_{11} u^2 (\ell_*(T)/d_T)^2),$$

for every $t \in T$,

$$\mathbb{E}_\eta \left| \sum_{i \in I^c} \varepsilon_i(\pi_{s_0} t)_i (W_{\pi_{s_0}t, I})_i \right| \leq c_{12} u^2 2^{s_0/2} d_T \cdot \delta d_T \leq c_{13} d_T \delta \ell_*(T).$$

Finally, invoking Lemma 2.2, we have that with probability at least $1 - 2 \exp(-cu^2 (\ell_*(T)/d_T)^2)$, for every $t \in T_{s_1}$,

$$|Z_t| \leq C u^2 d_T \delta \ell_*(T),$$

as required. ■

Case II: $\log(e(1 + \delta^2 n)) > 1/\delta^2$.

The necessary modifications are minor and we shall only sketch them. In this range, $2^{s_1} = \log(e(1 + \delta^2 n))$, and the problem is that for each vector $\Delta_s t$, the number of blocks J_ℓ of cardinality $1/\delta^2$ —namely, $n\delta^2$, is larger than $\exp(2^s)$ when $s \leq s_1$. Therefore, setting $|J_\ell| = 1/\delta^2$ for every ℓ , the uniform estimate on

$$\max_{\ell} \frac{\|P_{J_{\ell+1}} W_{\pi_{s+1}t, I}\|_2}{\sqrt{|J_\ell|}} = \delta \max_{\ell} \|P_{J_\ell} W_{\pi_{s+1}t, I}\|_2$$

can be obtained by bounding $(\mathbb{E}\|P_{J_\ell} W_{\pi_{s+1}t, I}\|_2^q)^{1/q}$ for $q \geq \log(\delta^2 n)$. Indeed, by Lemma 2.4, for every $I \subset \{1, \dots, n\}$ and every $t \in T$, we have that

$$\delta \left(\mathbb{E} \max_{\ell} \|P_{J_\ell} W_{\pi_{s+1}t, I}\|_2^q \right)^{1/q} \leq \delta^2 \sqrt{q} d_T.$$

The rest of the argument is identical to the previous one and is omitted. ■

Proof of Theorem 1.11. Using the estimates we established, it follows that for $u \geq c_0$, with probability at least

$$1 - 2 \exp \left(-c_1 u^2 \left(\frac{\ell_*(T)}{d_T} \right)^2 \right),$$

$$\Phi \leq C u \delta \ell_*(T) \quad \text{and} \quad \Psi^2 \leq C u^2 d_T \delta \ell_*(T) \left(1 + \delta \sqrt{\log(e(1 + n\delta^2))} \right),$$

noting that $\delta d_T^2 \leq c_2 d_T \delta \ell_*(T)$. Since

$$\sup_{t \in T} \left| \|A_\varepsilon t\|_2^2 - \|t\|_2^2 \right| \leq \Psi^2 + 2\Phi \sqrt{\Psi^2 + d_T^2} + \Phi^2 + C' \left((\delta \ell_*(T))^2 + d_T \delta \ell_*(T) \right)$$

the claim follows from a straightforward computation, by separating to the cases $\Phi \leq \Psi$ and $\Psi \leq \Phi$. ■

3 Application - A circulant Bernoulli matrix

Let $(\varepsilon_i)_{i=1}^n$ and $(\varepsilon'_i)_{i=1}^n$ be independent Bernoulli vectors. Set $\xi = (\varepsilon'_i)_{i=1}^n$ and put $D_\varepsilon = \text{diag}(\varepsilon_1, \dots, \varepsilon_n)$. Let Γ be the circulant matrix generated by the random vector ξ ; that is, Γ is the matrix whose j -th row is the shifted vector $\tau_j \xi$, where for every $v \in \mathbb{R}^n$, $\tau_j v = (v_{j-i})_{i=1}^n$.

Fix $I \subset \{1, \dots, n\}$ of cardinality m and let

$$A = \sqrt{\frac{1}{m}} P_I \Gamma : \mathbb{R}^n \rightarrow \mathbb{R}^m$$

to be the normalized partial circulant matrix. It follows from Theorem 3.1 in [6] and the estimates in Section 4 there that for a typical realization of ξ , the matrix A is (δ, k) -regular for $\delta \sim m^{-1/2} \log^2 n$:

Theorem 3.1. [6] *There exist absolute constants c and C such that the following holds. For $x > 0$ with probability at least*

$$1 - 2 \exp \left(-c \min \left\{ x^2 \frac{m}{k}, x \sqrt{\frac{m}{k}} \log^2 n \right\} \right)$$

we have that

$$\sup_{t \in \Sigma_k} \left| \|At\|_2^2 - 1 \right| \leq C(1+x) \sqrt{\frac{k}{m}} \log^2 n.$$

By Theorem 3.1 and the union bound for $1 \leq k \leq 1/\delta^2$, there is an event Ω with probability at least $1 - 2 \exp(-c' \log^4 n)$ with respect to $(\varepsilon'_i)_{i=1}^n$, on which, for every $k \leq 1/\delta^2$,

$$\sup_{t \in \Sigma_k} \left| \|At\|_2^2 - 1 \right| \leq \delta \sqrt{k}.$$

This verifies the assumption needed in Theorem 1.11 on the event Ω . Now fix $(\varepsilon'_i)_{i=1}^n \in \Omega$ and let A be the resulting partial circulant matrix. Set $A_\varepsilon = AD_\varepsilon$ and let $T \subset \mathbb{R}^n$. By Theorem 1.11, with probability at least

$$1 - 2 \exp(-c'(\ell_*(T)/d_T)^2)$$

with respect to $(\varepsilon_i)_{i=1}^n$, we have that

$$\sup_{t \in T} \left| \|A_\varepsilon t\|_2^2 - \|t\|_2^2 \right| \leq C' \left(\Lambda d_T \frac{\ell_*(T)}{\sqrt{m}} \cdot \log^2 n + \left(\frac{\ell_*(T)}{\sqrt{m}} \right)^2 \cdot \log^4 n \right)$$

where

$$\Lambda \leq c'' \max \left\{ 1, \frac{\log^5 n}{m} \right\}.$$

Thus, a random matrix generated by $2n$ independent random signs is a good embedding of an arbitrary subset of \mathbb{R}^n with the same accuracy (up to logarithmic factors) as a gaussian matrix. Moreover, conditioned on the choice of the circulant matrix A , the probability estimate coincides with the estimate in the gaussian case.

References

- [1] Shiri Artstein-Avidan, Apostolos Giannopoulos, and Vitali D. Milman. *Asymptotic geometric analysis. Part I*, volume 202 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, 2015.
- [2] Withold Bednorz. Concentration via chaining method and its applications. *arXiv:1405.0676*.
- [3] Sjoerd Dirksen. Tail bounds via generic chaining. *Electron. J. Probab.*, 20:no. 53, 29, 2015.
- [4] Sjoerd Dirksen. Dimensionality reduction with subgaussian matrices: a unified theory. *Found. Comput. Math.*, 16(5):1367–1396, 2016.

- [5] William B. Johnson and Joram Lindenstrauss. Extensions of Lipschitz mappings into a Hilbert space. In *Conference in modern analysis and probability (New Haven, Conn., 1982)*, volume 26 of *Contemp. Math.*, pages 189–206. Amer. Math. Soc., Providence, RI, 1984.
- [6] Felix Krahmer, Shahar Mendelson, and Holger Rauhut. Suprema of chaos processes and the restricted isometry property. *Comm. Pure Appl. Math.*, 67(11):1877–1904, 2014.
- [7] Felix Krahmer and Rachel Ward. New and improved Johnson-Lindenstrauss embeddings via the restricted isometry property. *SIAM J. Math. Anal.*, 43(3):1269–1281, 2011.
- [8] Shahar Mendelson. Upper bounds on product and multiplier empirical processes. *Stochastic Process. Appl.*, 126(12):3652–3680, 2016.
- [9] Samet Oymak, Benjamin Recht, and Mahdi Soltanolkotabi. Isometric sketching of any set via the restricted isometry property. *Inf. Inference*, 7(4):707–726, 2018.
- [10] Michel Talagrand. *Upper and lower bounds for stochastic processes*. Springer, 2014.