

Low-mass inelastic dark matter direct detection via the Migdal effect

Nicole F. Bell,^{1,*} James B. Dent,^{2,†} Bhaskar Dutta,^{3,‡}
Sumit Ghosh,^{3,§} Jason Kumar,^{4,¶} and Jayden L. Newstead^{1,**}

¹*ARC Centre of Excellence for Dark Matter Particle Physics,*

School of Physics, The University of Melbourne, Victoria 3010, Australia

²*Department of Physics, Sam Houston State University, Huntsville, TX 77341, USA*

³*Mitchell Institute for Fundamental Physics and Astronomy, Department of Physics and Astronomy,
Texas A&M University, College Station, TX 77843, USA*

⁴*Department of Physics and Astronomy, University of Hawaii, Honolulu, HI 96822, USA*

We consider searches for the inelastic scattering of low-mass dark matter at direct detection experiments, using the Migdal effect. We find that there are degeneracies between the dark matter mass and the mass splitting that are difficult to break. Using XENON1T data we set bounds on a previously unexplored region of the inelastic dark matter parameter space. For the case of exothermic scattering, we find that the Migdal effect allows xenon-based detectors to have sensitivity to dark matter with $\mathcal{O}(\text{MeV})$ mass, far beyond what can be obtained with nuclear recoils.

I. INTRODUCTION

There has been significant recent interest in methods of detecting low-mass dark matter at direct detection experiments [1–30]. The main difficulty to be overcome is that, for low-mass dark matter, the recoil energy deposited in the detector is typically small relative to threshold values needed for detection. For the case where the dominant interaction is with nucleons, this problem is exacerbated by the fact that nuclear recoils are more difficult to detect than electron recoils. Some particularly promising detection strategies involve new analysis techniques, rather than new detector technologies. A particularly useful strategy, which has been the subject of several recent studies, is the Migdal effect [18, 23, 31–37]. These studies have largely focused on elastic nuclear scattering. However, inelastic dark matter scattering (iDM) is a generic feature of many classes of dark matter models [38–55]. Here we discuss the Migdal effect in the context of inelastic dark matter scattering.

If dark matter scatters off a nucleus, then electrons may be released via the Migdal effect. Essentially, the electron cloud is boosted relative to the scattered nucleus, which can result in electron emission. Because many direct detection experiments are more sensitive to energetic electrons than to recoiling nuclei, these Migdal electrons can provide the leading channel for the direct detection of low-mass dark matter.

The spectrum of energy deposition in the detector can be altered if dark matter-nucleus scattering is inelastic. Indeed, the Migdal effect itself can be thought of as a

type of inelasticity in the DM-nucleus scattering, as the Migdal electrons carry away energy, but negligible momentum. Dark matter-nucleus scattering can exhibit inelasticity in two other ways: by exciting a low-lying nuclear state, or by changing the dark matter particle mass. Here we consider the latter case, by assuming the dark matter particle emerging from the scattering process has a different mass than the incoming dark particle. If the outgoing mass is assumed to be larger (smaller) than the incoming mass, the scattering is said to be endothermic (exothermic).

Although inelastic dark matter scattering has been studied in-depth in the context of models to explain the DAMA excess [38, 42, 56–58], it is in fact a generic feature of some classes of dark matter models. While we will not explore particular inelastic dark matter models for the purpose of this paper, we provide the following as an illustrative example. Inelastic scattering mediated by a dark photon with a vector coupling to the dark matter is generic in any model where dark matter is only charged under spontaneously broken continuous symmetries. The reason is that a gauge boson can only have a vector coupling to a complex degree of freedom. But if all of the continuous symmetries under which the dark matter is charged are spontaneously broken, then the dark matter is generically expected to split into two real degrees of freedom. Since one cannot form a vector current with a single real degree of freedom, the dark photon must instead couple to an off-diagonal vector current, yielding inelastic scattering.

The relevant parameters of the dark matter model include the mass splitting, δ , as well as the mass m_χ and dark matter-nucleon cross section $\sigma_{\chi n}$. We will demonstrate that there are degeneracies among these parameters that cannot easily be broken with the data from direct detection experiments alone. We will also show that, in the case of exothermic scattering, the Migdal effect provides a unique opportunity to probe very low-mass dark matter.

* n.bell@unimelb.edu.au

† jbdent@shsu.edu

‡ dutta@physics.tamu.edu

§ ghosh@tamu.edu

¶ jkumar@hawaii.edu

** jnewstead@unimelb.edu.au

The plan of this paper is as follows. In Section II, we derive the electron recoil spectrum arising from the Migdal effect in the case of inelastic dark matter scattering. In Section III we present our results and comment on the distinguishability of the various scenarios under consideration. In Section IV, we conclude with a discussion of our results and future avenues.

II. THE MIGDAL EFFECT WITH INELASTIC SCATTERING

We focus on electron ionization in liquid xenon as a result of DM-nucleus scattering, through the Migdal effect. As noted in [59], bremsstrahlung and electron excitation are expected to be subleading effects in xenon-based detectors. Similarly, we will follow the isolated-atom approximation (see [34]), which is expected to be a good approximation for xenon at the relevant momentum transfer.

The non-relativistic DM-atomic scattering process is $\chi A \rightarrow \chi' A$. In the center-of-mass frame, we find

$$\frac{1}{2}\mu v^2 = E_{\chi'} + E_A + \delta, \quad (1)$$

where $\mu = m_\chi m_A / (m_\chi + m_A)$ is the reduced mass of the initial $\chi - A$ system, v is the relative velocity of the two initial particles, and $E_{\chi'}$ and E_A are the kinetic energies of the outgoing dark particle and the atom, respectively. Here, $\delta = E_e + m_\chi - m_{\chi'}$ represents the amount of the initial kinetic energy lost to inelastic effects, including the excitation of Migdal electrons and the mass splitting between χ and χ' . Note that we have not necessarily assumed $m_\chi < m_{\chi'}$; as such, δ may be negative.

In the frame of the detector, we then can express the atomic recoil energy as

$$E_R = \frac{\mu^2}{m_A} v^2 \left[1 - \frac{\delta}{\mu v^2} - \sqrt{1 - \frac{2\delta}{\mu v^2} \cos \theta_{cm}} \right], \quad (2)$$

where θ_{cm} is the dark matter scattering angle in center-of-mass frame, and where we have made the approximation $m_{\chi'} m_A / (m_{\chi'} + m_A) \sim \mu$.

For velocity-independent spin-independent scattering, the scattering matrix element is independent of θ_{cm} . If the DM interacts with the nucleus through s -channel exchange of a mediator with mass m_ϕ , then the squared matrix element depends on E_R as $\propto m_\phi^4 / (m_\phi^2 + 2m_A E_R)^2$, yielding a differential atomic scattering rate which scales as

$$\frac{d^2 R}{dE_R dv} \propto \frac{m_\phi^4}{(m_\phi^2 + 2m_A E_R)^2} \frac{m_A}{2\mu^2 v^2} [vf(v)] F^2(q), \quad (3)$$

where $f(v)$ is dark matter velocity distribution and $F(q)$ is the nuclear form factor. Henceforth, we will assume a contact interaction.

The electromagnetic energy spectrum is then given by

$$\frac{d^3 R}{dE_e dE_R dv} = \frac{d^2 R}{dE_R dv} \times \frac{1}{2\pi} \sum_{n,\ell} \frac{d}{dE_e} p_{qe}^c(n, \ell \rightarrow E_e), \quad (4)$$

where p_{qe}^c is the ionization probability (values for xenon can be found in [34]). Here we include the rates for Migdal electrons originating from the $n = 3, 4, 5$ shells. To obtain the deposited energy spectrum we must sum the contributions from the nuclear recoil, E_R , the ejected electron E_e and the atomic de-excitation energy E_{nl} :

$$E_{\text{det}} = \mathcal{L} E_R + E_e + E_{nl} \quad (5)$$

where we have included a quenching factor, \mathcal{L} , on the nuclear recoil energy. We take a constant value of $\mathcal{L} = 0.15$ and note that the error this introduces would be a subdominant to the uncertainties in the atomic ionization probabilities. The detected energy spectrum is then obtained by integrating Eq. 4 over the atomic recoil energies and enforcing energy conservation. As noted in [60], the impulse approximation used in the calculation of the ionization probabilities will break down if the collision time scale (set by the inverse of the recoil energy E_R^{-1}) is greater than the time taken for the atom to traverse its potential (set by the inverse of the phonon frequency ω_{ph}^{-1}). In a fluid we will take this cutoff as the time taken for an atom to traverse the average inter-atomic distance at the sound speed. For xenon at 170K that is $t_1 = a/v_s \approx 10^{-12}$ s, therefore we conservatively set the cutoff at $E_R > 100t_1^{-1} \approx 50$ meV. This limit has an increasingly large effect at lower dark matter mass and target thresholds. At sufficiently low mass the maximum recoil energy falls below this cutoff, setting a minimum mass for which the impulse approximation is valid. For elastic scattering we restrict ourselves to $m_\chi \gtrsim 0.02$ GeV. For exothermic (endothermic) scattering the limits are dependent on the mass splitting, for example at -10 keV (10 keV) we set $m_\chi \gtrsim 0.6$ MeV (0.4 GeV).

III.1. The Migdal event rate and its degeneracies

In Figure 1 we plot the differential event rate of nuclear (solid black) and Migdal (solid blue) events as a function of the detected energy (in keV_{ee}), for $m_\chi = 2$ GeV and $\sigma_{SI} = 10^{-40}$ cm². Note, the nuclear recoil energies would be below threshold for XENON1T.

The shape of the Migdal electron recoil spectrum is largely characterized by the minimum energy (below which no events are seen), the maximum recoil energy, and the peak of the spectrum. The minimum energy is determined by the energy released when a higher-shell electron de-excites to fill the orbital vacated by a Migdal electron. This minimum energy is thus determined by the energy levels of the target nucleus, and is largely independent of the dark matter parameters.




FIG. 1. Rate of Migdal events for elastic (blue) and inelastic endothermic (red) and exothermic (green) DM scattering, for parameters that illustrate the $m_\chi - \delta$ degeneracy. Also shown is the elastic nuclear recoil rate (black). The vertical gray line corresponds to the threshold of the XENON1T S2-only analysis. The cross sections of the elastic, exothermic and endothermic curves are $(1, 0.65, 3.5) \times 10^{-40} \text{ cm}^2$, respectively.




FIG. 2. Rate of Migdal events for elastic (blue), endothermic (red) and exothermic (green) DM scattering. The vertical gray line corresponds to the threshold of the XENON1T S2-only analysis. The cross sections of the elastic, exothermic and endothermic curves are $(1, 0.0084, 28) \times 10^{-40} \text{ cm}^2$, respectively.

The maximum energy of the Migdal electrons is given by

$$E_{max} = \frac{1}{2} \mu v_{max}^2 - \delta, \quad (6)$$

$$\sim \frac{1}{2} m_\chi v_{max}^2 - \delta.$$

where v_{max} is the maximum DM-nucleus relative speed, and we have assumed $m_\chi \ll m_A$. This amounts to the situation in which, in center-of-mass frame, all of the kinetic energy is absorbed by the various forms of inelasticity (the boost to laboratory frame shifts E_{max} by a negligible amount). We see that this maximum is nearly independent of the target material, and of the initial orbital of the Migdal electron. If v_{max} is taken to be the maximum speed with respect to the Earth of a particle at the galactic escape speed, then there is a degeneracy between m_χ and δ in the determination of E_{max} . In practice, the probability of a Migdal electron falls with both the atomic recoil energy and the electron energy and thus the rate falls steeply toward E_{max} .

The peak of the Migdal electron spectrum (for any shell) occurs just above the minimum energy, with an amplitude which is proportional to the DM-nucleus scattering cross section. We thus see that there should be a one-parameter family of models for which the Migdal electron energy spectrum is nearly identical. In Figure 1, we illustrate the approximate degeneracy between the mass and the mass splitting by plotting best-fit scenarios with $\delta = -4 \text{ keV}$ and $\delta = 4 \text{ keV}$. Note that the shift in the mass of the incoming particle is roughly what one would expect from Eq. 6. In Figure 2 we make a similar plot with the elastic case being $m_\chi = 0.5 \text{ GeV}$. The best-fit rate is found by minimizing the χ^2 of the integrated rate in 5 log-spaced bins above the XENON1T threshold, i.e. in the range $E_{det} = 0.186 - 2 \text{ keV}$ ($E_{det} = 0.186 - 1.5 \text{ keV}$ for the $m_\chi = 0.5 \text{ GeV}$ case). We find that while the rates are almost degenerate above threshold, we see that a opportunity to resolve the degeneracy exists below threshold. Therefore, future experiments with $\mathcal{O}(10 \text{ eV})$ thresholds will not suffer greatly from this degeneracy.

II.II. Decay of the upscattered DM particle

If dark matter scatters endothermically, it is possible for the heavier state produced by this interaction to decay back to the lighter state. If this decay proceeds through the production of photons and occurs within the detector, then these decay photons can mimic an electron recoil [55]. In the case where dark matter is non-relativistic, then this signal is essentially monoenergetic, with $E = \delta$. A key feature which distinguishes this electron recoil signal from that produced by the Migdal electrons is that every DM-nucleus endothermic scatter would produce decay photons, while only a small fraction of such scatters can produce Migdal electrons.

This implies that although one might potentially see signals from both Migdal electrons and decay photons,

one is unlikely to see them simultaneously. If the excited DM state decays promptly within the detector, and if the energy of the decay photon exceeds threshold, then the decay photon signal will swamp the Migdal electron signal (if the decay photon signal is below threshold, then one would only see the Migdal electron signal). On the other hand, if the excited DM state is relatively long-lived, then the initial scatter and subsequent decay will not both occur in the detector. Instead, one will see either the Migdal electrons from the initial scatter (with decay occurring outside the detector), or one will see the monoenergetic decay photon signal (with the initial scatter having occurred outside the detector). The relative rates of Migdal events and decay photon events will depend on the lifetime of the excited states, but these are essentially separate signals.

III. BOUNDS AND SENSITIVITIES

We set upper limits on the dark matter-nucleon inelastic cross section using the S2-only data set from the XENON1T experiment [61]. This single channel analysis does not discriminate nuclear and electronic recoils and thus it can be used to place bounds on the cross section from both nuclear recoils and the Migdal effect. We perform a single bin analysis by integrating the total event rate in the range $E = 0.186 - 3.8 \text{ keV}_{ee}$, taking into account the energy dependent efficiency. The total number of events observed in the 22 tonne-day exposure was 61, while 23.4 were expected from background. Using this data we can place an upper limit (at 90% confidence) of 49 expected events coming from dark matter. We use this upper limit to place bounds on the cross section for both nuclear and Migdal scattering. Additionally, we project the sensitivity of LZ [62, 63] assuming a 1000 day exposure of the 5.6 tonne volume. Using a region of interest of $E = 0.5 - 4 \text{ keV}_{ee}$, an expected background of $2 \times 10^{-5} \text{ /kg/day/keV}$ [63] and background uncertainty of 15%, we place an upper limit of 80 expected events from dark matter. The resulting bounds are shown in Fig. 3 for elastic, exothermic, and endothermic interactions. The exo/endothermic cases are calculated for a benchmark mass-splitting of $\pm 10 \text{ keV}$. These mass splittings demonstrate the extremal behaviour of the Migdal bounds in the inelastic parameter space, from becoming essentially mass independent, to becoming no better than the nuclear recoil bounds.

For nuclear scattering, exo/endothermic inelasticity only shifts the bounds by $\sim 1 \text{ GeV}$. As mentioned previously, the maximum energy that can go into the Migdal electrons is approximately proportional to the dark matter kinetic energy, less the mass-splitting energy (Eq. 6). This drives the loss in sensitivity of the elastic and endothermic interaction curves, while the exothermic rate remains above threshold for very small dark matter masses. Essentially, even for very small dark matter mass and kinetic energy, the energy yielded by exother-




FIG. 3. Bounds on the inelastic cross section from XENON1T on nuclear recoils (solid), XENON1T with the Migdal effect (dashed) and the projected LZ sensitivity with the Migdal effect (dotted). These are each plotted for for exothermic (green), elastic (blue) and endothermic (red) interactions, where $\delta = -10, 0, 10 \text{ keV}$ respectively. We restrict our bounds to $m_\chi \geq 0.6 \text{ MeV}$, where we can be sure the impulse approximation is valid.

mic scattering is sufficient to yield Migdal electrons above threshold. For very small m_χ , we estimate that LZ’s sensitivity will not exceed limits from XENON1T. This is due to our estimate for LZ’s threshold, which is larger than that of XENON1T’s S2-only analysis.¹

It is interesting to note that, for $m_\chi \ll \text{GeV}$, XENON1T is not able to detect exothermic nuclear recoils even for $\delta = -10 \text{ keV}$. Even though a significant amount of energy is released in exothermic scattering, for such light dark matter, the vast majority of this energy is transferred to the outgoing DM or the Migdal electrons, but not to nuclear recoils.

IV. CONCLUSION

Inelastic nuclear scattering is a feature which arises in many classes of dark matter models. We have considered the impact of this feature on searches for the scattering of low-mass dark matter using the Migdal effect. We have found that there is an irreducible degeneracy in the Migdal electron energy spectrum between DM mass and the inelastic splitting. The direct detection data alone is

¹ We note that LZ may also perform an S2-only analysis, improving their low-mass sensitivity. However, the achievable threshold, background and exposure of such an analysis is unknown.

thus insufficient to reconstruct the dark matter model. We also find that, in the case of exothermic scattering, the Migdal effect provides sensitivity to dark matter masses as low as 1 MeV, providing a new approach to searching for very low-mass inelastic dark sector models. Although we have focused on xenon-based detectors, these results should generalize to other materials.

Other than reducing the threshold, one potential way to break the degeneracy between mass and mass splitting is to search for photons produced by decay of the heavier dark state. This may occur at a location displaced from the original scatter, which itself need not even be within

the detector volume. The energy of these decay photons would determine the mass splitting, breaking the model degeneracy.

ACKNOWLEDGMENTS

The work of BD and SG are supported in part by the DOE Grant No. DE-SC0010813. The work of JK is supported in part by DOE grant DE-SC0010504. NFB and JLN are supported in part by the Australian Research Council. JBD acknowledges support from the National Science Foundation under Grant No. NSF PHY-1820801.

-
- [1] R. Essig, J. Mardon, and T. Volansky, *Phys. Rev. D* **85**, 076007 (2012), arXiv:1108.5383 [hep-ph].
- [2] R. Essig, A. Manalaysay, J. Mardon, P. Sorensen, and T. Volansky, *Phys. Rev. Lett.* **109**, 021301 (2012), arXiv:1206.2644 [astro-ph.CO].
- [3] P. W. Graham, D. E. Kaplan, S. Rajendran, and M. T. Walters, *Phys. Dark Univ.* **1**, 32 (2012), arXiv:1203.2531 [hep-ph].
- [4] H. An, M. Pospelov, J. Pradler, and A. Ritz, *Phys. Lett. B* **747**, 331 (2015), arXiv:1412.8378 [hep-ph].
- [5] R. Essig, M. Fernandez-Serra, J. Mardon, A. Soto, T. Volansky, and T.-T. Yu, *JHEP* **05**, 046 (2016), arXiv:1509.01598 [hep-ph].
- [6] Y. Hochberg, Y. Zhao, and K. M. Zurek, *Phys. Rev. Lett.* **116**, 011301 (2016), arXiv:1504.07237 [hep-ph].
- [7] S. Derenzo, R. Essig, A. Massari, A. Soto, and T.-T. Yu, *Phys. Rev. D* **96**, 016026 (2017), arXiv:1607.01009 [hep-ph].
- [8] I. M. Bloch, R. Essig, K. Tobioka, T. Volansky, and T.-T. Yu, *JHEP* **06**, 087 (2017), arXiv:1608.02123 [hep-ph].
- [9] Y. Hochberg, Y. Kahn, M. Lisanti, C. G. Tully, and K. M. Zurek, *Phys. Lett. B* **772**, 239 (2017), arXiv:1606.08849 [hep-ph].
- [10] Y. Hochberg, T. Lin, and K. M. Zurek, *Phys. Rev. D* **94**, 015019 (2016), arXiv:1604.06800 [hep-ph].
- [11] C. Kouvaris and J. Pradler, *Phys. Rev. Lett.* **118**, 031803 (2017), arXiv:1607.01789 [hep-ph].
- [12] R. Essig, T. Volansky, and T.-T. Yu, *Phys. Rev. D* **96**, 043017 (2017), arXiv:1703.00910 [hep-ph].
- [13] R. Budnik, O. Chesnovsky, O. Slone, and T. Volansky, *Phys. Lett. B* **782**, 242 (2018), arXiv:1705.03016 [hep-ph].
- [14] P. C. Bunting, G. Gratta, T. Melia, and S. Rajendran, *Phys. Rev. D* **95**, 095001 (2017), arXiv:1701.06566 [hep-ph].
- [15] S. Knapen, T. Lin, M. Pyle, and K. M. Zurek, *Phys. Lett. B* **785**, 386 (2018), arXiv:1712.06598 [hep-ph].
- [16] Y. Hochberg, Y. Kahn, M. Lisanti, K. M. Zurek, A. G. Grushin, R. Ilan, S. M. Griffin, Z.-F. Liu, S. F. Weber, and J. B. Neaton, *Phys. Rev. D* **97**, 015004 (2018), arXiv:1708.08929 [hep-ph].
- [17] S. A. Hertel, A. Biekert, J. Lin, V. Velan, and D. N. McKinsey, *Phys. Rev. D* **100**, 092007 (2019), arXiv:1810.06283 [physics.ins-det].
- [18] M. J. Dolan, F. Kahlhoefer, and C. McCabe, *Phys. Rev. Lett.* **121**, 101801 (2018), arXiv:1711.09906 [hep-ph].
- [19] T. Bringmann and M. Pospelov, *Phys. Rev. Lett.* **122**, 171801 (2019), arXiv:1810.10543 [hep-ph].
- [20] T. Emken, R. Essig, C. Kouvaris, and M. Sholapurkar, *JCAP* **09**, 070 (2019), arXiv:1905.06348 [hep-ph].
- [21] R. Essig, J. Pérez-Ríos, H. Ramani, and O. Slone, *Phys. Rev. Research* **1**, 033105 (2019), arXiv:1907.07682 [hep-ph].
- [22] Y. Ema, F. Sala, and R. Sato, *Phys. Rev. Lett.* **122**, 181802 (2019), arXiv:1811.00520 [hep-ph].
- [23] N. F. Bell, J. B. Dent, J. L. Newstead, S. Sabharwal, and T. J. Weiler, *Phys. Rev. D* **101**, 015012 (2020), arXiv:1905.00046 [hep-ph].
- [24] T. Trickle, Z. Zhang, and K. M. Zurek, *Phys. Rev. Lett.* **124**, 201801 (2020), arXiv:1905.13744 [hep-ph].
- [25] T. Trickle, Z. Zhang, K. M. Zurek, K. Inzani, and S. Griffin, *JHEP* **03**, 036 (2020), arXiv:1910.08092 [hep-ph].
- [26] S. M. Griffin, K. Inzani, T. Trickle, Z. Zhang, and K. M. Zurek, *Phys. Rev. D* **101**, 055004 (2020), arXiv:1910.10716 [hep-ph].
- [27] D. Baxter, Y. Kahn, and G. Krnjaic, *Phys. Rev. D* **101**, 076014 (2020), arXiv:1908.00012 [hep-ph].
- [28] N. A. Kurinsky, T. C. Yu, Y. Hochberg, and B. Cabrera, *Phys. Rev. D* **99**, 123005 (2019), arXiv:1901.07569 [hep-ex].
- [29] R. Catena, T. Emken, N. A. Spaldin, and W. Tarantino, *Phys. Rev. Res.* **2**, 033195 (2020), arXiv:1912.08204 [hep-ph].
- [30] S. M. Griffin, Y. Hochberg, K. Inzani, N. Kurinsky, T. Lin, and T. C. Yu, (2020), arXiv:2008.08560 [hep-ph].
- [31] A. Migdal, *J.Phys.(USSR)* **4**, 449 (1941).
- [32] J. D. Vergados and H. Ejiri, *Phys. Lett. B* **606**, 313 (2005), arXiv:hep-ph/0401151.
- [33] R. Bernabei *et al.*, *Int. J. Mod. Phys. A* **22**, 3155 (2007), arXiv:0706.1421 [astro-ph].
- [34] M. Ibe, W. Nakano, Y. Shoji, and K. Suzuki, *JHEP* **03**, 194 (2018), arXiv:1707.07258 [hep-ph].
- [35] R. Essig, J. Pradler, M. Sholapurkar, and T.-T. Yu, *Phys. Rev. Lett.* **124**, 021801 (2020), arXiv:1908.10881 [hep-ph].
- [36] C. P. Liu, C.-P. Wu, H.-C. Chi, and J.-W. Chen, *Phys. Rev. D* **102**, 121303 (2020), arXiv:2007.10965 [hep-ph].
- [37] G. Grilli di Cortona, A. Messina, and S. Piacentini, *JHEP* **11**, 034 (2020), arXiv:2006.02453 [hep-ph].
- [38] D. Tucker-Smith and N. Weiner, *Phys. Rev. D* **64**, 043502

- (2001), arXiv:hep-ph/0101138.
- [39] D. Tucker-Smith and N. Weiner, *Phys. Rev. D* **72**, 063509 (2005), arXiv:hep-ph/0402065.
- [40] D. P. Finkbeiner and N. Weiner, *Phys. Rev. D* **76**, 083519 (2007), arXiv:astro-ph/0702587.
- [41] C. Arina and N. Fornengo, *JHEP* **11**, 029 (2007), arXiv:0709.4477 [hep-ph].
- [42] S. Chang, G. D. Kribs, D. Tucker-Smith, and N. Weiner, *Phys. Rev. D* **79**, 043513 (2009), arXiv:0807.2250 [hep-ph].
- [43] Y. Cui, D. E. Morrissey, D. Poland, and L. Randall, *JHEP* **05**, 076 (2009), arXiv:0901.0557 [hep-ph].
- [44] P. J. Fox, G. D. Kribs, and T. M. Tait, *Phys. Rev. D* **83**, 034007 (2011), arXiv:1011.1910 [hep-ph].
- [45] T. Lin and D. P. Finkbeiner, *Phys. Rev. D* **83**, 083510 (2011), arXiv:1011.3052 [astro-ph.CO].
- [46] A. De Simone, V. Sanz, and H. P. Sato, *Phys. Rev. Lett.* **105**, 121802 (2010), arXiv:1004.1567 [hep-ph].
- [47] H. An, P. Dev, Y. Cai, and R. Mohapatra, *Phys. Rev. Lett.* **108**, 081806 (2012), arXiv:1110.1366 [hep-ph].
- [48] M. Pospelov, N. Weiner, and I. Yavin, *Phys. Rev. D* **89**, 055008 (2014), arXiv:1312.1363 [hep-ph].
- [49] D. P. Finkbeiner and N. Weiner, *Phys. Rev. D* **94**, 083002 (2016), arXiv:1402.6671 [hep-ph].
- [50] K. R. Dienes, J. Kumar, B. Thomas, and D. Yaylali, *Phys. Rev. Lett.* **114**, 051301 (2015), arXiv:1406.4868 [hep-ph].
- [51] G. Barello, S. Chang, and C. A. Newby, *Phys. Rev. D* **90**, 094027 (2014), arXiv:1409.0536 [hep-ph].
- [52] J. Bramante, P. J. Fox, G. D. Kribs, and A. Martin, *Phys. Rev. D* **94**, 115026 (2016), arXiv:1608.02662 [hep-ph].
- [53] N. F. Bell, G. Busoni, and S. Robles, *JCAP* **09**, 018 (2018), arXiv:1807.02840 [hep-ph].
- [54] J. R. Jordan, Y. Kahn, G. Krnjaic, M. Moschella, and J. Spitz, *Phys. Rev. D* **98**, 075020 (2018), arXiv:1806.05185 [hep-ph].
- [55] N. F. Bell, J. B. Dent, B. Dutta, S. Ghosh, J. Kumar, and J. L. Newstead, *Phys. Rev. Lett.* **125**, 161803 (2020), arXiv:2006.12461 [hep-ph].
- [56] D. Tucker-Smith and N. Weiner, *Nucl. Phys. B Proc. Suppl.* **124**, 197 (2003), arXiv:astro-ph/0208403.
- [57] D. P. Finkbeiner, T. Lin, and N. Weiner, *Phys. Rev. D* **80**, 115008 (2009), arXiv:0906.0002 [astro-ph.CO].
- [58] S. Kang, S. Scopel, and G. Tomar, *Phys. Rev. D* **99**, 103019 (2019), arXiv:1902.09121 [hep-ph].
- [59] E. Aprile *et al.* (XENON), *Phys. Rev. Lett.* **123**, 241803 (2019), arXiv:1907.12771 [hep-ex].
- [60] S. Knapen, J. Kozaczk, and T. Lin, (2020), arXiv:2011.09496 [hep-ph].
- [61] E. Aprile *et al.* (XENON), *Phys. Rev. Lett.* **123**, 251801 (2019), arXiv:1907.11485 [hep-ex].
- [62] D. Akerib *et al.* (LZ), *Nucl. Instrum. Meth. A* **953**, 163047 (2020), arXiv:1910.09124 [physics.ins-det].
- [63] D. S. Akerib *et al.* (LZ), (2021), arXiv:2102.11740 [hep-ex].