

The First Chiral Homology Group

Jethro van Ekeren *

Reimundo Heluani †

Abstract

We study the first chiral homology group of elliptic curves with coefficients in vacuum insertions of a conformal vertex algebra V . We find finiteness conditions on V guaranteeing that these homologies are finite dimensional, generalizing the C_2 -cofinite, or quasi-lisse condition in the degree 0 case. We determine explicitly the flat connections that these homologies acquire under smooth variation of the elliptic curve, as insertions of the conformal vector and the Weierstrass ζ function. We construct linear functionals associated to self-extensions of V -modules and prove their convergence under said finiteness conditions. These linear functionals turn out to be degree 1 analogs of the n -point functions in the degree 0 case. As a corollary we prove the vanishing of the first chiral homology group of an elliptic curve with values in several rational vertex algebras, including affine \mathfrak{sl}_2 at non-negative integral level, the $(2, 2k + 1)$ -minimal models and arbitrary simple affine vertex algebras at level 1. Of independent interest, we prove a Fourier space version of the Borchers formula.

1 Introduction

1.1. Let k be a field, A an associative k -algebra and $\rho_M : A \rightarrow \text{End}_k(M)$ a finite dimensional A -module. The linear functional

$$\varphi_M^0 : A \rightarrow k, \quad a \mapsto \text{tr}_M \rho_M(a), \quad (1.1)$$

defines a class in the dual of the zeroth Hochschild homology of A :

$$\text{HH}_0(A)^* = (A/[A, A])^*.$$

If A is finite dimensional and semisimple over k , then every class in $\text{HH}_0(A)^*$ arises as a linear combination of such linear functionals. In fact, as M runs over a set of representatives of the isomorphism classes of irreducible A -modules, the φ_M^0 provide a basis of $\text{HH}_0(A)^*$.

1.2. Let

$$0 \rightarrow M \rightarrow E \rightarrow M \rightarrow 0, \quad (1.2)$$

be a self-extension of finite dimensional A -modules. We choose a k -linear splitting of (1.2) so that the representation $\rho_E : A \rightarrow \text{End}_k(E)$ may be written in block matrix form as

$$\rho_E(a) = \begin{pmatrix} \rho_M(a) & \sigma(a) \\ 0 & \rho_M(a) \end{pmatrix}$$

where $\sigma : A \rightarrow \text{End}_k(M)$ is an A -derivation. The linear functional

$$\varphi_E^1 : A \otimes A \rightarrow k, \quad a \otimes b \mapsto \text{tr}_M \sigma(a) \rho_M(b),$$

defines a class in the dual $\text{HH}_1(A)^*$ of the first Hochschild homology group. This class is independent of the choice of splitting of (1.2).

*Instituto de Matemática e Estatística, Universidade Federal Fluminense, Niterói RJ, Brazil.

jethrovanekerem@gmail.com

†Instituto de Matemática Pura e Aplicada, Rio de Janeiro, RJ, Brazil

heluani@potuz.net

1.3. In this article we study a *chiral* or *vertex* analog of the construction in 1.1–1.2 with the *genus 1 chiral homology* of Beilinson and Drinfeld [1] in place of Hochschild homology above. The degree zero case is due to Zhu [2]. Let V be a conformal vertex algebra of central charge c and M an admissible V -module (that is, a positive energy V -module which decomposes into finite dimensional Jordan blocks for the action of L_0). If V is C_2 -cofinite, or more generally quasi-lisse (see [3]), then the linear functional

$$\varphi_M^0 : V \rightarrow k, \quad a \mapsto \text{tr}_M Y^M(e^{2\pi iz} L_0 a, e^{2\pi iz}) e^{2\pi i\tau(L_0 - c/24)} \quad (1.3)$$

converges to a holomorphic function of $\tau \in \mathbb{H}$ the upper half complex plane. It defines a class of the dual of the zeroth *chiral homology group of V on the universal elliptic curve*, also known as the *genus 1 conformal block of V* . If in addition V is rational then the classes φ_M^0 , where M runs over a set of representatives of the isomorphism classes of irreducible V -modules, constitute a basis for the space of conformal blocks in genus 1. In this case, the space of conformal blocks is finite dimensional. This is the chiral analog of 1.1.

1.4. Let V be a conformal vertex algebra, $\tau \in \mathbb{H}$ the upper-half plane, $q = e^{2\pi i\tau}$, and E_τ the complex elliptic curve defined as the quotient of \mathbb{C} by the lattice $\mathbb{Z} + \mathbb{Z}\tau$ or equivalently as the quotient of \mathbb{C}^* by the action of \mathbb{Z} given by $n \mapsto q^n$. In [1] Beilinson and Drinfeld define a complex of vector spaces $C_\bullet(E_\tau, V)$ whose homology $H_\bullet^{\text{ch}}(E_\tau, V)$ is known as the *chiral homology* of E_τ with coefficients in V . As E_τ varies in the moduli space of elliptic curves, the spaces $H_k^{\text{ch}}(E_\tau, V)^*$ define bundles with flat connections. For $k = 0$ the space of flat sections of this bundle coincides with the space of conformal blocks in genus 1 as defined by Zhu. In this article we detail the construction of the $k = 1$ case, leading to the notion of *degree 1 conformal block* of V in genus 1.

1.5. Let M be an admissible V -module and

$$0 \rightarrow M \rightarrow E \rightarrow M \rightarrow 0$$

a self-extension of M . Upon choosing a splitting as vector spaces, we may write the action $Y^E(\cdot, z)$ of V on E in block matrix form as

$$Y^E(a, z) = \begin{pmatrix} Y^M(a, z) & \psi(a, z) \\ 0 & Y^M(a, z) \end{pmatrix}.$$

The vertex analog of the construction described in 1.2 is as follows.

Theorem 10.23. *Let \mathcal{F} be the space of meromorphic functions $f(z)$ which are bi-periodic with possible poles at $\mathbb{Z} + \tau\mathbb{Z}$. Define the following linear functional on $V \otimes V \otimes \mathcal{F}$*

$$a \otimes b \otimes f \mapsto \frac{1}{2\pi i} \int_C dz f(e^{2\pi i(z-w)}) \text{tr}_M \psi(e^{2\pi iz} L_0 a, e^{2\pi iz}) Y^M(e^{2\pi iw} L_0 b, e^{2\pi iw}) e^{2\pi i\tau(L_0 - c/24)}, \quad (1.4)$$

where C is a small contour of z around 0 not containing the point w . Then if $\text{res}_t f(t)Y[a, t]b = 0$ and under certain finiteness conditions on V (described below), this linear functional converges to a holomorphic function of τ in the upper half plane (it is independent of w), and it determines a flat section of the dual of the first chiral homology group of V in genus 1. This class is independent of the chosen splitting of E .

Notice that in contrast to the situation of 1.2, our linear functional takes as input an elliptic function f in addition to two elements in the algebra V .

1.6. We now describe the *finiteness conditions* alluded to in Theorem 10.23. The role they play in connection with the first chiral cohomology group of V in genus 1 is analogous to that played in degree zero by the C_2 -cofiniteness condition or more generally by the quasi-lisse condition. In order to describe them, we recall from [4] that any vertex algebra carries a decreasing filtration $\{F_p V\}$ known as the *Li filtration*. The associated graded $A = \text{gr}_F V$ with respect to this filtration is a \mathbb{Z}_+ -graded commutative algebra with a derivation of degree 1 which is generated (as a differential algebra) by its degree 0 component $R_V = A^0$. The spectrum of the commutative algebra A is known as the *singular support* of V [5]. The subalgebra R_V in fact carries the natural structure of a Poisson algebra, and its spectrum is known as the *associated scheme* of V .

The vertex algebra V is said to be C_2 -cofinite (or lisse) if $\dim R_V < \infty$, and is said to be quasi-lisse if its associated variety consists of a finite number of symplectic leaves. If V is quasi-lisse then in particular the zeroth Poisson homology of its associated scheme is finite dimensional, that is,

$$\mathrm{HP}_0(R_V) = \frac{R_V}{\{R_V, R_V\}}, \quad (1.5)$$

is finite dimensional. (We briefly recall the construction of Poisson homology of Poisson algebras in Section 3.)

It follows from the work of Zhu [2] that for a C_2 -cofinite conformal vertex algebra V the zeroth chiral homology of E_τ with coefficients in V is finite dimensional. In fact, as was observed in [3], it is sufficient that V be quasi-lisse or more generally that $\dim \mathrm{HP}_0(R_V) < \infty$. And indeed we have

$$\dim H_0^{\mathrm{ch}}(E_\tau, V) \leq \dim \mathrm{HP}_0(R_V).$$

See Proposition 10.7 below. In this work we prove finite dimensionality of the first chiral homology group of E_τ under two finiteness hypotheses on V , the first being a natural degree 1 analog of (1.5), namely

$$\dim \mathrm{HP}_1(R_V) < \infty. \quad (1.6)$$

To describe the second finiteness hypothesis we recall JR_V the \mathbb{Z}_+ -graded commutative algebra with a derivation of degree 1, freely generated as a differential algebra by R_V . Its spectrum is known as the *arc space* of the associated scheme $\mathrm{Spec} R_V$. By the universal property of the arc space construction, there is a surjection of differential algebras

$$JR_V \twoheadrightarrow A \quad (1.7)$$

where, we recall, $A = \mathrm{gr}_F V$. The kernel of the surjection (1.7) is a differential ideal of JR_V , and we say that the vertex algebra V is *classically free* if (1.7) is an isomorphism. We may now state our result on finiteness of the first chiral homology group.

Theorem 10.9. *Let E_τ be an elliptic curve and V a vertex algebra that satisfies (1) the condition (1.6) and (2) that the kernel of the surjection (1.7) is finitely generated as a differential ideal. Then $\dim H_1^{\mathrm{ch}}(E_\tau, V) < \infty$.*

1.7. Here is another description of the condition on the morphism (1.7). The algebra A comes equipped with a derivation of degree 1. This derivation induces a morphism of A -modules $d : \Omega_A^1 \rightarrow A$, where Ω_A^1 is the module of Kähler differentials. This morphism can be extended by the Leibniz rule to a differential d of degree -1 of the free (super) commutative algebra $K_\bullet(A) = \wedge^\bullet \Omega_A^1$. This is the *Koszul complex* of the differential algebra A . We let $\mathrm{HK}_i(A)$ be the i^{th} homology of the complex $K_\bullet(A)$. It was shown in [6] that $\mathrm{HK}_1(A) = 0$ if and only if V is classically free, and in Proposition 3.3 below that $\dim \mathrm{HK}_1(A) < \infty$ if and only if the kernel of the morphism (1.7) is finitely generated as a differential ideal.

1.8. Stronger results can be obtained for V classically free. In [6] a relationship was established between the *nodal curve* limit $\tau \rightarrow i\infty$ of the chiral homology of E_τ , and the Hochschild homology of the Zhu algebra $\mathrm{Zhu}(V)$ of V . In the case that V is classically free this takes the particularly simple form

$$\lim_{\tau \rightarrow i\infty} H_1^{\mathrm{ch}}(E_\tau, V) \cong \mathrm{HH}_1(\mathrm{Zhu}(V)).$$

See Proposition 10.19 below. In Section 10 we exploit this result to obtain

Theorem 10.21. *Let V be a classically free vertex algebra and E_τ a smooth elliptic curve. If $\mathrm{Zhu}(V)$ is semisimple, in particular if V is rational, then $H_1^{\mathrm{ch}}(E_\tau, V) = 0$.*

Corollary 12.2. *The chiral homology $H_1^{\mathrm{ch}}(E_\tau, V)$ vanishes in the following cases:*

- (a) $V = \mathrm{Vir}_{2,2s+1}$ is a boundary Virasoro minimal model,
- (b) $V = V_k(\mathfrak{sl}_2)$ is the simple affine vertex algebra at level $k \in \mathbb{Z}_+$,
- (c) $V = V_1(\mathfrak{g})$ is the simple affine vertex algebra at level 1, for \mathfrak{g} a simple Lie algebra.

1.9. In order to prove convergence of the linear functional (1.3) Zhu showed that the right hand side, regarded as a formal power series, satisfies an ordinary differential equation in $q = e^{2\pi i\tau}$ with a regular singularity at $q = 0$. He then used results from the classical theory of ordinary differential equations, in particular the Frobenius-Fuchs theory, to guarantee the convergence of such a formal solution.

We follow the same technique in this article. We write down an explicit ordinary differential equation that our functionals (1.4) satisfy (see (5.48)). This differential equation arises as the connection that the bundles of chiral homology (or their duals) acquire as E_τ moves in the moduli space of elliptic curves. We then show that this differential equation has a regular singularity at $q = 0$.

1.10. In [2], the space of conformal blocks is defined as a system of compatible n -point functions, the trace functional (1.3) being a special case of such for $n = 1$. It is proven that one can express an $n+1$ point function as a linear combination of n -point functions. Convergence of the $n = 1$ linear functionals and this ‘‘insertion formula’’ guarantees convergence of the n -point functions. Conversely, a suitable integral of an $n + 1$ point function recovers an n -point function. Concretely, the n -point functions are given by

$$a_1 \otimes \cdots \otimes a_n \mapsto \mathrm{tr}_M Y^M(e^{2\pi iz_1 L_0} a_1, e^{2\pi iz_1}) \dots Y^M(e^{2\pi iz_n L_0} a_n, e^{2\pi iz_n}) q^{L_0 - c/24}.$$

The following algebro-geometric counterpart was given in [1]. For each collection of points t_1, \dots, t_n on the elliptic curve E_τ the authors define a complex whose homology, denoted here $H_\bullet^{ch}(V^{\otimes n}, t_1, \dots, t_n)$ represents *chiral homology of E_τ with supports on the modules V inserted at the points t_i* . The authors moreover prove [1, (4.4.2.1)] that these homologies are isomorphic for all n . These statements remain true for arbitrary (not necessarily elliptic) curves and collections of points. In the degree 0 case this was carried out explicitly in [7], recovering the result of Zhu in genus 1.

We here define explicit complexes computing the homologies $H_\bullet^{ch}(V^{\otimes n}, t_1, \dots, t_n)$ and construct natural classes, the analog of the n -point functions, associated to self extensions of modules as in 1.5. The $n = 1$ version of this construction is just (1.4). These classes are linear functionals

$$V^{\otimes n+1} \otimes \mathcal{F}_{n+1} \rightarrow \mathcal{F}_n,$$

where \mathcal{F}_n denotes the space of meromorphic functions $f(t_1, \dots, t_n)$, bi-periodic with possible poles at the diagonals $t_i = t_j$ modulo $\mathbb{Z} + \tau\mathbb{Z}$. We show that $n + 1$ point functions can be described in terms of n point functions (see Proposition 9.18), thus guaranteeing their convergence. Similarly we show that an integral of an $n + 1$ -point function is given by n -point functions (see (6.7)). Concretely these linear functionals are given by

$$a_0 \otimes \cdots \otimes a_n \otimes f(z_0, \dots, z_n) \mapsto \frac{1}{2\pi i} \int_C dz_0 f(e^{2\pi iz_0}, \dots, e^{2\pi iz_n}) \mathrm{tr}_M \psi(e^{2\pi iz_0 L_0} a_0, e^{2\pi iz_0}) \times \\ \times Y^M(e^{2\pi iz_1 L_0} a_1, e^{2\pi iz_1}) \dots Y^M(e^{2\pi iz_n L_0} a_n, e^{2\pi iz_n}) q^{L_0 - c/24},$$

where C is a small contour around 0 not containing any of the points z_1, \dots, z_n . The main result is that this formal power series converges uniformly and its limit is an elliptic function in \mathcal{F}_n (see Theorem 11.1).

1.11. Instrumental in Zhu’s construction is the identification of E_τ with the quotient $\mathbb{C}/(\mathbb{Z} + \tau\mathbb{Z})$ via the exponential map. Indeed the relationship between conformal blocks as defined by correlation functions, and as chiral homology, requires the introduction of an auxiliary vertex algebra structure associated with the exponential change of coordinates. For a conformal vertex algebra with state-field correspondence $Y(\cdot, z)$, Zhu constructs an isomorphic vertex algebra structure on V with state-field correspondence $Y[\cdot, z]$, defined by

$$Y[a, z] = Y(e^{2\pi i L_0} a, e^{2\pi iz} - 1). \quad (1.8)$$

This vertex algebra structure has already appeared above in the statement of Theorem 10.23.

A new feature that arises in our study of chiral homology in degree greater than 0, is the appearance of elliptic functions in the description of chains and not only of the differentials. This forces us to study these functions and their Fourier expansions. Recall that for a vertex algebra V and a rational function $f(s, t)$ with possible poles at $s = 0$, $t = 0$ and $s = t$ the following Borchers identity holds

$$\mathrm{res}_s \mathrm{res}_t Y(a, s) Y(b, t) c_{i_{s,t}} f(s, t) - \mathrm{res}_t \mathrm{res}_s Y(b, t) Y(a, s) c_{i_{t,s}} f(t, s) =$$

$$\text{res}_t Y(\text{res}_{s-t} i_{t,s-t} f(s,t) Y(a, s-t)b, t) c$$

where $i_{s,t} : \mathbb{C}(s,t) \rightarrow \mathbb{C}((s))((t))$ is the canonical embedding and $\text{res}_s : V((s)) \rightarrow V$ is the *residue map* (coefficient of s^{-1}). When dealing with the exponentiated coordinates $z = e^{2\pi i s}$, $w = e^{2\pi i t}$ it is useful to use the following modified vertex operators

$$X(a, z) = z^{-1} Y(z^{L_0} a, z).$$

Let $f(t)$ be a meromorphic function of a complex variable t which is periodic, i.e., $f(t+1) = f(t)$. Let $\tau \in \mathbb{H}$ and suppose f is analytic in the domain $-\Im\tau < \Im t < \Im\tau$ except possibly at $t = 0$ where it may have a pole. Put $q = e^{2\pi i \tau}$ and let $F_+(z)$ (resp. $F_-(z)$) be the Fourier expansion of f , in the domain $|q| < |z| < 1$ (resp. $1 < |z| < |q|^{-1}$). If f is *quasi-periodic* with period τ , that is, f satisfies $f(t+\tau) = f(t) - \alpha$ for all $t \in \mathbb{C}$ for some constant $\alpha \in \mathbb{C}$, then we prove

Theorem 7.3. *The following identity holds*

$$2\pi i X(a_{[f]} b, w) = \text{res}_z F_- \left(\frac{z}{w} \right) X(a, z) X(b, w) - \text{res}_z F_+ \left(\frac{z}{w} \right) X(b, w) X(a, z) \quad (1.9)$$

as an equality in $\text{Hom}(V, V((w)))$, where

$$a_{[f]} b = \text{res}_t f(t) Y[a, t] b.$$

Notice that both vertex algebra structures $Y(a, z)$ and $Y[a, z]$ of V appear in (1.9), as well as both f and its Fourier expansions on different domains appear.

1.12. This article is a self-contained introduction to chiral homology in degree 1 for vertex algebras, aimed at the representation theorist. No background on algebraic geometry nor chiral algebras is required. It is our aim to provide a bridge between the abstract world of chiral homology on arbitrary curves and the linear algebra world in the presence of coordinates. The translation between these two was carried out in [6], in this article we focus on techniques to compute these homologies. In doing so, we have had to combine tools from different areas of mathematics, the analytical proof of the Borcherds identity in Fourier coordinates (1.9), Zhu algebras and their Hochschild homology, the Li filtration and the higher Poisson homology of the associated scheme, arc spaces and classical-freeness of vertex algebras. It would be interesting to see if these techniques have geometrical counterparts. Some of these tools make sense immediately in the geometric side, for example, Li's *increasing standard filtration* makes sense as a filtration of chiral algebras. Others, like Zhu's theory, do not yet seem to be incorporated into the theory of chiral algebras.

This article continues with the program detailed in [6, §1.6]. In particular we cover item a) described there confirming that certain extensions of modules produce degree 1 chiral homology classes by extending the result from the limit as $q \rightarrow 0$ using the flat connection on the moduli space of elliptic curves. The other points of the cited program include chiral homology in higher degrees and in higher genera.

The complexes described in Section 5 admit obvious generalizations to higher degrees, and so do the finiteness conditions on $\text{HK}_\bullet(A)$ and $\text{HP}_\bullet(R_V)$. By increasing the number of insertion points, namely computing $H_k^{\text{ch}}(V^{\otimes n})$ with $n \geq k$, the higher derived functors of global sections do not play any role, as the varieties involved become affine. We expect the same techniques used this article to apply to compute higher degree chiral homology of elliptic curves.

Regarding chiral homology in higher genus. Recently, it was proved in a series of papers by Damiolini, Gibney and Tarasca [8, 9] that the degree 0 chiral homology of a C_2 cofinite vertex algebra V is finite dimensional in arbitrary curves, not only genus 1. Their approach is to describe the chiral homology on genus g by understanding the structure in the boundary of the moduli space, thus reducing to lower genera. It would be interesting to see if this approach can be generalized to higher chiral homology, and in particular if the conditions on the Poisson homology of R_V and the Koszul complex $K_\bullet(A)$ described above, continue to play the same role in arbitrary genus.

1.13 Algebraic-geometric considerations. We collect here some observations that may help the reader acquainted with [1]. This subsection is not needed anywhere in the text.

Let X be a smooth algebraic curve of genus g , V a conformal vertex algebra and $\mathcal{A} = \mathcal{A}_V$ the corresponding chiral algebra on X . There are different flavours of chiral homology that one may compute.

First, there is simply $H_{\bullet}^{\text{ch}}(X, \mathcal{A})$, the *chiral homology of X with coefficients in \mathcal{A}* . When X is an elliptic curve $E_{\tau} = \mathbb{C}/\mathbb{Z} + \tau\mathbb{Z}$, these homologies are denoted by $H_i^{\text{ch}}(E_{\tau}, V)$ below, for $i = 0, 1$.

For a collection of points $\{x_i\}_{i=1}^n \subset X$ we have the *chiral homology with supports* $H_{\bullet}^{\text{ch}}(X, \{\mathcal{A}_{x_i}\})$, that is, the chiral homology with coefficients in the collection of \mathcal{A} -modules \mathcal{A}_{x_i} , supported at the points x_i . For every n and every collection $\{x_i\}$ these homologies are isomorphic. When $X = E_{\tau}$ and x_i is the image of $t_i \in \mathbb{C}$. These homologies are denoted $H_i^{\text{ch}}(V^{\otimes n}, t_1, \dots, t_n, \tau)$ below.

The construction works in families, that is, when X/S is a curve over a base S , and $\{x_i\}$ is a collection of S -valued points. One obtains a twisted \mathcal{D}_S -module, whose fiber at $s \in S$ equals $H_{\bullet}^{\text{ch}}(X_s, \{\mathcal{A}_{x_i(s)}\})$. Applying this to $S = \mathcal{M}_{g,n}$, the moduli space of n -marked, genus g curves and X is the universal curve, the chiral homologies give rise to twisted \mathcal{D} -modules, or quasi-coherent sheaves with projectively flat connections on $\mathcal{M}_{g,n}$, denote these sheaves by $\mathcal{H}_{\bullet}^{\text{ch}}(V^{\otimes n})$. In the case when $g = 1$, the global sections of this \mathcal{D} -module on $\mathcal{M}_{1,n}$ are denoted $H_i^{\text{ch}}(V^{\otimes n})$ for $i = 0, 1$ below. Its zeroth de Rham homology (coinvariants by the action of vector fields) is denoted by $H_i^{\text{ch}}(V^{\otimes n})_{\nabla}$ below. Similarly global sections of the dual quasi-coherent sheaf $\mathcal{H}^{\text{ch}}(V^{\otimes n})^*$ are denoted $H_i^{\text{ch}}(V^{\otimes n})^*$ below, and their flat sections, or equivalently its zeroth de Rham cohomology, or the \mathcal{D} -module push-forward to the point, are denoted by $H_i^{\text{ch}}(V^{\otimes n})^{*\nabla}$.

The moduli space of marked curves fibers over the moduli space of curves $\pi : \mathcal{M}_{g,n} \rightarrow \mathcal{M}_g$ by forgetting the points (in the case $g = 1$ we do remember one point, the polarization of the elliptic curve). Instead of pushing forward $\mathcal{H}_{\bullet}^{\text{ch}}(V^{\otimes n})^*$ down to the point, we may consider π_* of it. This is the relative 0 de Rham cohomology and corresponds to taking flat sections with respect to the action of the vertical vector fields, or the relative connection corresponding to moving the points $\{x_i\}$ in the curve $X \in \mathcal{M}_g$. The global sections of $\pi_* \mathcal{H}_{\bullet}^{\text{ch}}(V^{\otimes n})^*$, when $g = 1$, are denoted $H_i^{\text{ch}}(V^{\otimes n})^{*\text{tr}}$ below.

We can fix X and move the points $\{x_i\}$, obtaining a \mathcal{D} -module on X^n , which is symmetric, namely it is a pullback of a \mathcal{D} -module on $\text{Sym}^n X$, the universal degree n -divisor on X . Its fiber at the point $\{x_i\} \in X^n \setminus \Delta$ (where $\Delta \subset X^n$ is the diagonal divisor) being $H_{\bullet}^{\text{ch}}(X, \{A_{x_i}\})$. Denote this \mathcal{D}_{X^n} -module by $\mathcal{H}_{\bullet}^{\text{ch}}(X, V^{\otimes n})$.

Here is another construction $\mathcal{H}_{\bullet}^{\text{ch}}(X, V^{\otimes n})$. When viewed as a Lie algebra on the Ran space $\text{Ran}(X)$ of X , the chiral algebra \mathcal{A} can have modules supported not only at X but at higher powers $\Delta^{(n)} : X^n \rightarrow \text{Ran}(X)$. We can consider for example the image of $\Delta_*^{(n)} \mathcal{A}^{\boxtimes n}$ as a module over \mathcal{A} . The chiral homology with coefficients in $\Delta_*^{(n)} \mathcal{A}^{\boxtimes n}$ gives rise to a \mathcal{D} -module on $\text{Sym}^n X$, its pullback to X^n coincides with $\mathcal{H}_{\bullet}^{\text{ch}}(X, V^{\otimes n})$. The construction of the complex C_{\bullet}^n below, follows this pattern in the case when X is the universal elliptic curve and we restrict to homologies in degrees 0 and 1.

1.14. The structure of this paper is as follows. In section 2 we recall the definition of vertex algebras and their modules. We recall the exponential change of coordinate formula. In section 3 we recall the definition of the Poisson homology of a Poisson algebra and prove the relation between the finite generation of the kernel of (1.7) and the first homology of the Koszul complex. In section 4 we define several spaces of meromorphic elliptic functions that we will use throughout. We study their Fourier series expansions and the failure of ellipticity when differentiating with respect to the modular parameter. In section 5 we define the complexes computing the chiral homologies and their duals in genus 1. We give explicit proofs that these complexes are in fact complexes. We define the homology with supports and study their \mathcal{D} -module structure when moving these points. We describe explicitly the flat connection with respect to the modular parameter, and the corresponding modular differential equation that flat sections satisfy. In section 6 we define the notion of *degree 1 conformal block on the torus* and prove its modular invariance. This is a generalization of the notion of n -point functions in the degree 0 case, and consists of a coherent system of flat sections of the dual of chiral homology with coefficients in $V^{\otimes n}$ for all $n \geq 1$. In section 7 we prove the main technical tool of the article, a version of Borchers identity on Fourier variables (Theorem 7.3). In section 8 we study self extensions of modules and their associated derivations. We prove a version of the above Borchers formula for these derivations. Section 9 is the main technical section of the article. We define the linear functionals associated to self extensions as formal power series. We prove that they satisfy the formal differential equation corresponding to the flat connection and we prove an *insertion formula*. In section 10 we prove the finite generation of chiral homology under the finiteness conditions on V explained above. We use this to prove convergence of the trace functional of section 9 for $n = 1$. We prove here the vanishing of chiral homology of classically free rational vertex algebras. In section 11 we prove the convergence of the n -point functions of section 9 for arbitrary n . We show that their limit is a degree 1 conformal block on the torus. In section 12 we

give some examples of vanishing of chiral homology and some non-trivial classes obtained by the above construction. In section 13 we summarize some of the results obtained.

1.15 Acknowledgements. This work has been carried out over an extended period of time. We have benefited from discussion with many mathematicians. We would like to thank Drazen Adamovic, Tomoyuki Arakawa, David Ben-Zvi, John Francis, Dennis Gaitsgory, Deniz Kus, Antun Milas, Sam Raskin and Pavel Safronov for enlightening discussions. RH was supported by CNPq Grant Number 305688/2019-7. JvE was supported by CNPq Grant Number 303806/2017-6 and by a grant from the Serrapilheira Institute (grant number Serra – 1912-31433).

2 Preliminaries on Vertex Algebras

In this section we collect the basic definitions and notations on vertex algebras. We start by describing the notation for Laurent expansions of multi-variable functions. We give the definition of vertex algebra in 2.4, and of modules in 2.6. We recall the exponential change of coordinates in 2.8.

2.1. Let $f(z, w) \in \mathbb{C}[[z, w]][z^{-1}, w^{-1}, (z-w)^{-1}]$. We denote by $i_{z,w}f(z, w) \in \mathbb{C}((z))((w))$ and $i_{w,z}f(z, w) \in \mathbb{C}((w))((z))$ the expansions of $f(z, w)$ in the domains $|z| > |w|$ and $|w| > |z|$ respectively. More generally, for $f(z_1, \dots, z_n) \in \mathbb{C}[[z_1, \dots, z_n]][z_i^{-1}, (z_i - z_j)^{-1}]_{1 \leq i \neq j \leq n}$ we denote by

$$i_{z_1, \dots, z_n}f(z_1, \dots, z_n) \in \mathbb{C}((z_1)) \dots ((z_n)),$$

the expansion of $f(z_1, \dots, z_n)$ in the domain $|z_1| > \dots > |z_n|$. This expansion may be done iteratively; first expand all instances of $(z_i - z_n)^n$ in positive powers of z_n , obtaining a Laurent series in z_n whose coefficients are functions of z_1, \dots, z_{n-1} , then repeat with z_{n-1} , etc.

Sometimes it will be required to expand $f(z_1, \dots, z_n)$ as above in powers of $z_i - z_j$. We will only need such expansions in the domain $|z_i| > |z_i - z_j|$, where $1 \leq i < j \leq n$. To perform such an expansion means to replace all instances of z_i^{-k} , where $k \geq 0$, by $((z_i - z_j) + z_j)^{-k}$ and expand in positive powers of $z_i - z_j$. Therefore such an expansion is a series in $z_1, \dots, \hat{z}_i, \dots, z_n, z_i - z_j$, where we have omitted z_i from the list.

2.2 Example. For $f = f(z, w, t)$ the expansion

$$i_{w, w-t, z-w}f(z, w, t) \in \mathbb{C}((w))((w-t))((z-w)),$$

is computed as follows. We take the Laurent series expansion $f(z, w, t) = \sum_k f_k(w, t)(z-w)^k$ of f around $z-w=0$. Then we take the Laurent series expansion $f_k(w, t) = \sum_\ell f_{\ell, k}(w)(w-t)^\ell$ of each f_k around $w-t=0$. Finally we write

$$i_{w, w-t, z-w}f(z, w, t) = \sum_{\ell, k} f_{\ell, k}(w)(w-t)^\ell (z-w)^k.$$

2.3. For $f(z, w) \in \mathbb{C}[[z, z^{-1}, w, w^{-1}]]$ we write $\text{res}_z dzf(z, w) \in \mathbb{C}[[w, w^{-1}]]$ for its coefficient of z^{-1} . For $f(z, w) \in \mathbb{C}[[z, w]][z^{-1}, w^{-1}, (z-w)^{-1}]$ the notation $\text{res}_z dzf(z, w)$ is ambiguous. We must choose an embedding $\mathbb{C}[[z, w]][z^{-1}, w^{-1}, (z-w)^{-1}] \subset \mathbb{C}[[z, z^{-1}, w, w^{-1}]]$ in order to compute this residue. For example

$$\text{res}_z dz i_{z,w} \frac{1}{z-w} = 1 \neq 0 = \text{res}_z dz i_{w,z} \frac{1}{z-w}.$$

By $\text{res}_{z=w} dzf(z, w)$ we mean the coefficient of $(z-w)^{-1}$ in the expansion of $f(z, w) = f((z-w)+w, w)$ in the domain $|w| > |z-w|$.

Notice that for any Laurent series $f(t) \in \mathbb{C}((t))$ we have

$$\text{res}_t dt \frac{d}{dt} f(t) = 0, \quad \text{res}_t dt \left(t \frac{d}{dt} + 1 \right) f(t) = 0. \quad (2.1)$$

2.4. Throughout this article $(V, \mathbf{1}, \omega, Y(\cdot, z))$ will denote a conformal vertex algebra (or vertex operator algebra). In particular $V = \bigoplus_{n \in \mathbb{Z}_+} V_n$ is a \mathbb{Z}_+ -graded complex vector space V together with two distinguished vectors $\mathbf{1}$ and ω and a bilinear map

$$V \otimes V \rightarrow V((z)), \quad a \otimes b \mapsto Y(a, z)b = \sum_{n \in \mathbb{Z}} z^{-1-n} a_{(n)} b, \quad a_{(n)} b = 0 \quad \forall n \gg 0,$$

satisfying the following axioms

- a) $Y(\mathbf{1}, z) = \text{Id}_V$.
- b) $a_{(n-1)} \mathbf{1} = \delta_{n,0} a$ for all $a \in V$ and $n \geq 0$.
- c) For any $f(z, w) \in \mathbb{C}[[z, w]][z^{-1}, w^{-1}, (z-w)^{-1}]$ and $a, b, c \in V$, the following Borcherds identity holds

$$\begin{aligned} \text{res}_z dz Y(a, z) Y(b, w) c i_{z,w} f(z, w) - \text{res}_z dz Y(b, w) Y(a, z) c i_{w,z} f(z, w) \\ = Y(\text{res}_{z=w} dz f(z, w) Y(a, z-w) b, w) c. \end{aligned} \quad (2.2)$$

- d) Put $L(z) = \sum_{n \in \mathbb{Z}} L_n z^{-n-2} = Y(\omega, z)$. Then

$$[L_m, L_n] = (m-n)L_{m+n} + \frac{c}{12}(m^3 - m)\delta_{m,-n} \text{Id}_V m$$

where $c \in \mathbb{C}$ is a constant called the *central charge* of V .

- e) $L_{-1} a = a_{(-2)} \mathbf{1}$ for every $a \in V$.
- f) For each $n \in \mathbb{Z}_+$ the restriction of L_0 to V_n is $n \text{Id}_{V_n}$ and V_n is finite dimensional.

2.5. A vertex algebra V satisfies the following skew-symmetry relation:

$$Y(a, z)b = e^{zL_{-1}} Y(b, -z)a,$$

multiplying by z^k and taking residues we get

$$a_{(k)} b = -(-1)^k \sum_{j \geq 0} \frac{(-L_{-1})^j}{j!} b_{(k+j)} a. \quad (2.3)$$

2.6. For us a *module* over the conformal vertex algebra V will be what is normally called a positive energy module or a \mathbb{N} -gradable weak module in the literature. This is a vector space M equipped with a bilinear operation

$$V \otimes M \rightarrow M((z)), \quad a \otimes m \mapsto Y^M(a, z)m = \sum_{n \in \mathbb{Z}} z^{-n-1} a_{(n)}^M m, \quad a_{(n)}^M m = 0 \quad \forall n \gg 0,$$

satisfying the following analog of the Borcherds identity for $a, b \in V, m \in M$ and $f(z, w) \in \mathbb{C}[[z, w]][z^{-1}, w^{-1}, (z-w)^{-1}]$:

$$\begin{aligned} \text{res}_z dz Y^M(a, z) Y^M(b, w) m i_{z,w} f(z, w) - \text{res}_z dz Y^M(b, w) Y^M(a, z) m i_{w,z} f(z, w) = \\ Y^M(\text{res}_{z=w} dz f(z, w) Y(a, z-w) b, w) m. \end{aligned} \quad (2.4)$$

We put $L^M(z) = Y^M(\omega, z)$ and require that L_0^M act with finite dimensional generalized eigenspaces, and that the real part of its spectrum be bounded below. That is $M = \bigoplus_{\lambda \in \mathbb{C}} M_\lambda$ with $(L_0^M|_{M_\lambda} - \lambda)^n = 0$ for $n \gg 0$. Below we will often drop the superscript M on the operators whenever no confusion should arise.

2.7. Let $\tau \in \mathbb{H}$ and $q = e^{2\pi i\tau}$. For a module M the operator $q^{L_0} = e^{2\pi i\tau L_0}$ is well-defined since the generalized eigenspaces of L_0 are finite dimensional by hypothesis. In Section 9 we will deal with formal sums like $\text{tr}_M q^{L_0}$ by which we mean, in this context, the expression

$$\sum_{\lambda} q^{\lambda} \sum_{n \geq 0} \frac{(2\pi i\tau)^n}{n!} (L_0 - \lambda)^n,$$

where q is a formal variable. Convergence of such series under appropriate hypotheses is then treated in Sections 10 and 11.

2.8. For a vertex operator algebra $(V, \mathbf{1}, \omega, Y(\cdot, z))$ we will consider another isomorphic vertex algebra $(V, \mathbf{1}, \tilde{\omega}, Y[\cdot, z])$ introduced by Zhu [2] where $\tilde{\omega} = (2\pi i)^2(\omega - \frac{c}{24}\mathbf{1})$ and

$$Y[a, z] = Y(e^{2\pi iz L_0} a, e^{2\pi iz} - 1).$$

We have for homogeneous $a \in V$ and $m \in \mathbb{Z}$

$$a_{[m]} = (2\pi i)^{-m-1} \text{res}_z (Y(a, z) (\log(1+z))^m (1+z)^{\deg a - 1}). \quad (2.5)$$

2.9. For a Laurent series $f(t) = \sum_{n \geq N} f_n t^n$, $a, b \in V$ we will use the notation

$$a_{(f)} b = \text{res}_t f(t) Y(a, t) b = \sum_{n \geq N} f_n a_{(n)} b, \quad a_{[f]} b = \text{res}_t f(t) Y[a, t] b = \sum_{n \geq N} f_n a_{[n]} b.$$

2.10 Remark. There is an ambiguous situation in that for a integer number n , $a_{(n)}$ could mean the n -product as defined in 2.4 or the f -product for the constant Laurent series $f(t) = nt^0$. We will always mean the n -th product by this notation and believe that no confusion should arise.

By integration by parts and noting that the translation operator for $(V, \mathbf{1}, \tilde{\omega}, Y[\cdot, z])$ is given by $(2\pi i)(L_0 + L_{-1})$ these operations satisfy

$$(L_{-1}a)_{(f)} b = -a_{(f')} b \quad (2.6)$$

and

$$(2\pi i)((L_0 + L_{-1})a)_{[f]} b = -a_{[f']} b. \quad (2.7)$$

We have the following

2.11 Lemma. *Let $f \in \mathbb{C}((x))$. Then*

$$a_{(f(x))} b + b_{(f(-x))} a = \sum_{j \geq 0} \frac{(-1)^j}{(j+1)!} T^{j+1} (a_{(x^{j+1} f(x))} b), \quad \text{for all } a, b \in V. \quad (2.8)$$

2.12. Let V be a vertex algebra. We denote by $V_{(-2)}V$ the subspace of V spanned by elements of the form $a_{(-2)}b$ for $a, b \in V$. The C_2 -algebra $R_V = V/V_{(-2)}V$ carries the natural structure of Poisson algebra with commutative product $a \cdot b = a_{(-1)}b$ and Poisson bracket $\{a, b\} = a_{(0)}b$. The vertex algebra V is said to be C_2 -cofinite if R_V is finite dimensional.

3 Homology

In this section we recall basic homological constructions that will be used in Section 10 to codify finiteness conditions on the vertex algebra V guaranteeing finite dimensionality of its chiral homology. We recall the construction of the *Koszul complex of a differential algebra* and the *arc algebra* of a commutative algebra. We describe a relation between the first homology group of the Koszul complex of a graded differential algebra and finite generation of a certain ideal in an associated arc algebra. We then recall the definition of Poisson homology of a Poisson algebra and give an explicit description of the differentials in low degree.

3.1. Let k be a field and A a commutative k -algebra. Denote by $\Omega_A = \Omega_{A/k}$ the A -module of Kähler differentials on A . It is isomorphic as a k -vector space to the Hochschild homology group $\mathrm{HH}_1(A)$. A derivation T of A induces a morphism

$$\iota_T : \Omega_A \rightarrow A,$$

of A -modules, which in turn extends uniquely to a differential d on the Koszul complex $\wedge^\bullet \Omega_A$. We refer to this as the Koszul complex of the differential algebra (A, T) and denote it by $K_\bullet(A)$. Explicitly, the differentials of $K_\bullet(A)$ in low degree are

$$d_1 : \Omega_A \rightarrow A, \quad a \, dx \mapsto a \cdot (Tx), \quad (3.1)$$

$$d_2 : \wedge^2 \Omega_A \rightarrow \Omega_A, \quad a \, dx \wedge dy \mapsto a \cdot (Tx) \, dy - a \cdot (Ty) \, dx. \quad (3.2)$$

We denote the homology groups of this complex by $\mathrm{HK}_i(A)$ for $i \in \mathbb{Z}_+$.

3.2. Let R be a commutative k -algebra. The arc algebra of R is the differential algebra (JR, ∂) characterized by the universal property that any morphism of algebras $R \rightarrow A$ from R to a differential algebra (A, T) , lifts uniquely to a morphism of differential algebras $(JR, \partial) \rightarrow (A, T)$ making the following diagram commute

$$\begin{array}{ccc} & & (JR, \partial) \\ & \nearrow & \downarrow \\ R & \longrightarrow & (A, T). \end{array} \quad (3.3)$$

The arc algebra is naturally \mathbb{Z}_+ -graded with $JR^0 = R$, and ∂ has degree $+1$.

3.3 Proposition. Let $A = \bigoplus_{n \in \mathbb{Z}_+} A^n$ be a graded differential k -algebra with derivation T of degree $+1$. Assume A^0 is of finite type as a k -algebra, and that A is generated by A^0 as a differential algebra. The canonical map $\pi : JA^0 \rightarrow A$ is a surjection and we write I for its kernel, which is a differential ideal of $J = JA^0$. Then

- a) $\mathrm{HK}_1(A) = 0$ if and only if π is an isomorphism, and
- b) in general there exists an injection

$$\mathrm{HK}_1(A) \rightarrow I/(J \cdot TI).$$

Proof. Part a) is [6, Thm. 13.7]. Part b) is proved by a diagram chase. Indeed we have a morphism of complexes

$$\begin{array}{ccccccc} \wedge^2 \Omega_J & \longrightarrow & \Omega_J & \longrightarrow & J & \longrightarrow & 0 \\ \downarrow & & \downarrow & & \downarrow & & \\ \wedge^2 \Omega_A & \longrightarrow & \Omega_A & \longrightarrow & A & \longrightarrow & 0, \end{array}$$

where the upper row is exact by a). Let $\omega = \sum_i a_i \, db_i \in \Omega_A$ be a 1-cocycle, and let \tilde{a}_i and \tilde{b}_i be lifts of a_i and b_i to J , and write $\tilde{\omega} = \sum_i \tilde{a}_i \, d\tilde{b}_i \in \Omega_J$. Clearly

$$d\tilde{\omega} = \sum_i \tilde{a}_i \cdot T\tilde{b}_i \in I,$$

since ω is closed. We claim that $\omega \mapsto d\tilde{\omega}$ defines a morphism

$$\delta : \mathrm{HK}_1(A) \rightarrow I/(J \cdot TI).$$

Indeed if $\omega' - \omega = d\theta$ where $\theta = \sum_i a_i \, db_i \wedge dc_i$, then

$$\tilde{\omega}' - \tilde{\omega} - d \sum_i \tilde{a}_i \, d\tilde{b}_i \wedge d\tilde{c}_i \in I \, dJ + J \, dI,$$

and so

$$\delta(\omega') - \delta(\omega) \in I \cdot (TJ) + J \cdot (TI) \subset J \cdot (TI).$$

Obviously δ is injective: $\delta(\omega) \in J \cdot TI$ implies ω possesses a lift $\tilde{\omega}$ of the form $\sum_i \tilde{a}_i d(\tilde{x}_i)$ which implies $\omega = 0$. \square

3.4. We briefly recall the definition of Poisson homology groups. These were introduced in [10] in a geometric context, the presentation here follows the algebraic formulation of [11]. First we recall the notion of Lie-Rinehart pair. Let k be a field (though the constructions work with k a commutative ring in general) and \mathcal{O} a commutative k -algebra. Then a Lie-Rinehart pair, or a \mathcal{O} -Lie algebroid consists of an \mathcal{O} -module \mathcal{L} together with a k Lie algebra structure on \mathcal{L} and an action $\omega : \mathcal{L} \rightarrow \text{Der}(\mathcal{O})$ of \mathcal{L} on \mathcal{O} by derivations. These data satisfy the conditions:

$$\omega(fx)(g) = f\omega(x)(g) \quad \text{and} \quad [x, fy] = f[x, y] + (\omega(x)f)y$$

for all $f, g \in \mathcal{O}$ and all $x, y \in \mathcal{L}$. Given a \mathcal{O} -Lie algebroid \mathcal{L} , we have its Chevalley-Eilenberg complex

$$C(\mathcal{L})_{>0} = \wedge_{\mathcal{O}}^{>0} \mathcal{L}. \quad (3.4)$$

The differential is constructed as follows. In degree 2 we let

$$d(x \otimes y) = [x, y],$$

and extend d by the Leibniz rule to be a derivation of degree -1 of the super-commutative algebra $C(\mathcal{L})$.

Let A be a Poisson algebra over k . The A -module $\Omega_{A/k} = \text{HH}_1(A)$ of Kähler differentials on A carries the structure of an A -Lie algebroid. The k -Lie algebra bracket is defined by

$$[f dx, g dy] = f\{x, b\} dy + g\{f, y\} dx + fg d\{x, y\}, \quad f, g, x, y \in A,$$

and the action is given by

$$\omega(f dx)(g) = f\{x, g\}.$$

The Eilenberg-Chevalley complex defined in (3.4) is augmented, by putting A in degree 0 and letting the differential in degree 1, be given by

$$d_1 : \Omega_{A/k} \rightarrow A, \quad a dx \mapsto \{a, x\}, \quad (3.5)$$

We let $C(\Omega_A^1)$ be this augmented complex. Its homology is known as the *Poisson homology* $\text{HP}_\bullet(-)$ of A :

$$\text{HP}_i(A) = H^i C(\Omega_A^1).$$

We record here the differential in degree 2 that will be used below:

$$d_2 : \wedge^2 \Omega_{A/k} \rightarrow \Omega_{A/k}, \quad a dx \wedge dy \mapsto \{a, x\} dy - \{a, y\} dx - a d(\{x, y\}). \quad (3.6)$$

4 Elliptic Functions

In this section we recall some material about elliptic functions. We adopt the conventions of [2]. We recall the Eisenstein series in 4.1 and the Weierstrass elliptic functions and their Fourier series expansions in 4.2. We introduce the rings of modular forms in 4.5, Jacobi forms in 4.6 and meromorphic elliptic functions in 4.8. We analyze domains of convergence of Fourier series expansions in 4.12 and define a derivation with respect to the modular parameter in 4.14.

4.1. For $k \geq 1$ we consider the Eisenstein series

$$G_{2k}(q) = 2\xi(2k) + \frac{2(2\pi i)^{2k}}{(2k-1)!} \sum_{n=1}^{\infty} \sigma_{2k-1}(n) q^n, \quad (4.1)$$

as a formal power series in the variable q . Here

$$\xi(2k) = \sum_{n \geq 1} n^{-2k} \quad \text{and} \quad \sigma_k(n) = \sum_{d|n} d^k.$$

This series converges uniformly and absolutely in the domain $|q| < 1$, and thereby defines a function $g_{2k}(\tau) = G_{2k}(e^{2\pi i \tau})$ analytic in the upper half complex plane \mathbb{H} . For $k \geq 2$ the function $g_{2k}(\tau)$ is a holomorphic modular form of weight $2k$. The function $g_2(\tau)$ transforms under the action of $\text{SL}_2(\mathbb{Z})$ as

$$g_2\left(\frac{a\tau + b}{c\tau + d}\right) = (c\tau + d)^2 g_2(\tau) - 2\pi i c(c\tau + d). \quad (4.2)$$

4.2. For $k \geq 1$ we introduce the formal Laurent series

$$\wp_k(t) = \frac{1}{t^k} + (-1)^k \sum_{n \geq 1} \binom{2n+1}{k-1} G_{2n+2}(q) t^{2n+2-k}, \quad (4.3)$$

as an element of $\mathbb{C}[[q]]((t))$. These series are related to each other by

$$\wp_{k+1}(t) = -\frac{1}{k} \frac{d}{dt} \wp_k(t). \quad (4.4)$$

Putting $q = e^{2\pi i \tau}$ in $\wp_k(t)$ yields a Laurent series in t which converges uniformly and absolutely in a neighborhood of $t = 0$. The sum can be analytically continued to a meromorphic function $\wp_k(t, \tau)$ having poles at $t = m + n\tau$ for $m, n \in \mathbb{Z}$. For $k \geq 2$ this meromorphic function is doubly periodic:

$$\wp_k(t + m + n\tau, \tau) = \wp_k(t, \tau), \quad m, n \in \mathbb{Z},$$

while for $k = 1$ one has

$$\wp_1(t + m + n\tau, \tau) = \wp_1(t, \tau) + (m + n\tau)g_2(\tau) - 2\pi i n. \quad (4.5)$$

It is convenient to introduce the functions

$$\zeta(t, \tau) = \wp_1(t, \tau) - g_2(\tau)t$$

which, in light of (4.5), satisfies

$$\zeta(t + m + n\tau, \tau) = \zeta(t, \tau) - 2\pi i n. \quad (4.6)$$

It will also be convenient to introduce the normalized functions

$$\bar{\wp}_k(t, \tau) = \begin{cases} \zeta(t, \tau) + \pi i & k = 1 \\ \wp_2(t, \tau) + g_2(\tau) & k = 2 \\ \wp_k(t, \tau) & k \geq 3. \end{cases}$$

They satisfy the same differential equation (4.4). We will often drop τ from the notation denoting $\wp_k(t)$ instead of $\wp_k(t, \tau)$ when no confusion should arise.

We also consider the formal series $P_k(z, q) \in \mathbb{C}[[q]][[z, z^{-1}]]$:

$$P_k(z, q) = \frac{(2\pi i)^k}{(k-1)!} \sum_{n=1}^{\infty} \left(\frac{n^{k-1} z^n}{1-q^n} + \frac{(-1)^k n^{k-1} z^{-n} q^n}{1-q^n} \right), \quad (4.7)$$

(where $(1-q^n)^{-1}$ stands for the geometric series $\sum_{j \geq 0} q^{nj}$). These series are related to each other by

$$P_{k+1}(z, q) = \frac{2\pi i}{k} z \frac{d}{dz} P_k(z, q). \quad (4.8)$$

The series $P_k(z, q)$ converges uniformly and absolutely in every closed subset of the domain $|q| < |z| < 1$, and can be analytically continued to a meromorphic function of $z \in \mathbb{C} \setminus \{0\}$ with poles at $z = q^n$ for $n \in \mathbb{Z}$. The series $P_k(z, q)$ should be thought of as the Fourier series of the function $\wp_k(t, \tau)$. More precisely one has

$$P_k(e^{2\pi it}, e^{2\pi i\tau}) = (-1)^k \overline{\wp}_k(t, \tau), \quad (4.9)$$

whenever $0 < \Im(t) < \Im(\tau)$. The series $P_k(qz, q)$ converges uniformly and absolutely in every closed subset of the domain $1 < |z| < |q|^{-1}$. For $k \geq 2$ the analytic continuation of the sum coincides with that of $P_k(z, q)$. On the other hand, by (4.6), the analytic continuation of $P_1(qz, q)$ coincides with that of $P_1(z, q) + 2\pi i$.

Purely formal analogues of these statements also hold, namely

$$P_1(z, q) - P_1(zq, q) + 2\pi i = 2\pi i \sum_{n \in \mathbb{Z}} z^n \quad (4.10)$$

and

$$P_k(z, q) - P_k(zq, q) = \frac{(2\pi i)^k}{(k-1)!} \sum_{n \in \mathbb{Z}} n^{k-1} z^n \quad \text{for } k \geq 2, \quad (4.11)$$

as formal power series.

4.3 Lemma. *The series $P_k(z, q)$ satisfy*

$$P_k(z^{-1}, q) = (-1)^k P_k(zq, q). \quad (4.12)$$

4.4. As in 2.9, for $a, b \in V$ we have the following elements of $V[G_4(q), G_6(q)] \subset V[[q]]$:

$$a_{[\wp_k]} b = a_{[k]} b + (-1)^k \sum_{n \geq 1} \binom{2n+1}{k-1} G_{2n+2}(q) a_{[2n+2-k]} b.$$

4.5. We will consider the graded ring of modular forms \mathbb{M}_* which is generated by $G_4(q)$ and $G_6(q)$. We will also need the ring of quasi-modular forms \mathbb{QM}_* generated by $G_2(q)$, $G_4(q)$ and $G_6(q)$. Both of these rings are Noetherian, indeed they are polynomial rings. We may consider these rings as subrings of $\mathbb{C}[[q]]$.

4.6. For each $n \geq 2$ we consider the following graded rings $\mathbb{J}_*^n = \bigoplus_{k \geq 0} \mathbb{J}_k^n$ of weak Jacobi modular forms of index 0. By definition \mathbb{J}_k^n consists of functions $f(t_1, \dots, t_n, \tau)$ satisfying

$$\begin{aligned} f(t_1, \dots, t_i + m + l\tau, \dots, t_n, \tau) &= f(t_1, \dots, t_n, \tau), & 1 \leq i \leq n, & \quad m, l \in \mathbb{Z}, \\ f\left(\frac{t_1}{c\tau + d}, \dots, \frac{t_n}{c\tau + d}, \frac{a\tau + b}{c\tau + d}\right) &= (c\tau + d)^k f(t_1, \dots, t_n, \tau), & ad - cb = 1, & \quad a, b, c, d \in \mathbb{Z}. \end{aligned} \quad (4.13)$$

The functions $f(t_1, \dots, t_n, \tau)$ are meromorphic in $\mathbb{C}^n \times \mathbb{H}$ with possible poles at $t_i - t_j = m + l\tau$, $i \neq j$, $m, l \in \mathbb{Z}$.

For every pair of non negative integers $n \geq m$ we have an inclusion $\mathbb{J}_*^m \rightarrow \mathbb{J}_*^n$ given by $f \mapsto \tilde{f}$ where $\tilde{f}(t_1, \dots, t_n; \tau) = f(t_{n-m+1}, \dots, t_n; \tau)$. In this way \mathbb{J}_*^n is an algebra over \mathbb{J}_*^m , and in particular all \mathbb{J}_*^n are algebras over \mathbb{M}_* .

Notice that $\mathbb{J}_*^0 = \mathbb{J}_*^1 = \mathbb{M}_*$. The description of \mathbb{J}_*^2 is as follows: the function $\wp_k(t_1 - t_2, \tau)$ is a weak Jacobi form of index 0 and weight k for $k \geq 2$, and \mathbb{J}_*^2 consists of linear combinations of $1, \wp_k(t_1 - t_2, \tau)$, $k \geq 2$ with coefficients in \mathbb{M}_* . It follows from [12, Prop. 2.8] that the ring \mathbb{J}_*^2 is isomorphic to the polynomial ring $\mathbb{M}[\wp_2, \wp_3, \wp_4]$.

Since every function $f(t_1, t_2, \tau)$ in \mathbb{J}_*^2 can be written in the form $f(t_1 - t_2, \tau)$ for some f , we shall often abuse notation and write $f(t, \tau) \in \mathbb{J}_*^2$, it being implicit that $f(t_1, t_2, \tau) = f(t_1 - t_2, \tau)$.

4.7. For $n \geq 3$ an inductive description of \mathbb{J}_*^n can be given. Any $f(t_1, \dots, t_n, \tau) \in \mathbb{J}_*^n$ can be written as a linear combination of the form [6]

$$\begin{aligned} f(t_1, \dots, t_n, \tau) &= g(t_2, \dots, t_n, \tau) \\ &+ \sum_{i=3}^n g_i(t_2, \dots, t_n, \tau) (\wp_1(t_{i-1} - t_1, \tau) - \wp_1(t_i - t_1, \tau) + \wp_1(t_i - t_{i-1}, \tau)) \\ &+ \sum_{i=2}^n \sum_{m \geq 2} \wp_m(t_i - t_1, \tau) g_{im}(t_2, \dots, t_n, \tau), \end{aligned}$$

where the functions $g(t_2, \dots, t_n, \tau)$, $g_i(t_2, \dots, t_n, \tau)$ for $i = 3, \dots, n$, and $g_{im}(t_2, \dots, t_n, \tau)$ for $i = 1, \dots, n$ and $m \geq 2$, are elements of \mathbb{J}_*^{n-1} . Inductively one obtains an expression for $f(t_1, \dots, t_n, \tau) \in \mathbb{J}_*^n$ as a linear combination of products of $\wp_m(t_i - t_j, \tau)$ for $1 \leq j < i \leq n$, and the functions

$$\zeta(t_i, t_{j-1}, t_j, \tau) = \wp_1(t_{j-1} - t_i, \tau) - \wp_1(t_j - t_i, \tau) + \wp_1(t_j - t_{j-1}, \tau), \quad 1 \leq i < j - 1 \leq n.$$

These are weak Jacobi forms in three variables of index 0 and weight 1. The function $\wp_1(t, \tau)$ is not a weak Jacobi modular form, as we have seen in 4.2 it is not periodic, however the combination $\zeta(t_1, t_2, t_3, \tau)$ is a Jacobi form.

4.8. For each n we shall denote by \mathcal{F}_n the ring of meromorphic functions $f(t_1, \dots, t_n, \tau)$ on $\mathbb{C}^n \times \mathbb{H}$ such that

$$f(t_1, \dots, t_i + m + l\tau, \dots, t_n, \tau) = f(t_1, \dots, t_n, \tau), \quad m, l \in \mathbb{Z},$$

and f is allowed to have poles at $t_i - t_j = m + l\tau$, $1 \leq i \neq j \leq n$, $m, l \in \mathbb{Z}$. We have $\mathbb{J}_*^n \subset \mathcal{F}_n$. Notice also that a similar recursive description as in 4.7 is valid for \mathcal{F}_n . The difference being in the first step of the induction as $\mathcal{F}_0 = \mathcal{F}_1$ consists of holomorphic functions of τ as opposed to modular forms. In particular, functions in $f(t_0, t_1, \tau) \in \mathcal{F}_2$ are in fact functions of one variable $f = f(t_0 - t_1, \tau)$. More generally we have

4.9 Lemma. Let $f(t_0, \dots, t_n, \tau) \in \mathcal{F}_{n+1}$, then

$$\sum_{i=0}^n \frac{\partial}{\partial t_i} f(t_0, \dots, t_n, \tau) = 0.$$

4.10 Remark. Notice that in the definition of $\zeta(t_1, t_2, t_3, \tau)$ we may replace $\wp_1(t, \tau)$ by $\overline{\wp}_1(t, \tau)$ without effect. It follows that the Fourier series expansion of $\zeta(t_1, t_2, t_3, \tau)$ in the domain $|qz_1| < |z_2| < |z_3| < |z_1|$ is given by

$$-P_1\left(\frac{z_2}{z_1}, q\right) + P_1\left(\frac{z_3}{z_1}, q\right) - P_1\left(\frac{z_3}{z_2}, q\right) - \pi i.$$

The following is standard

4.11 Lemma. Let $f(t_1, \dots, t_n, \tau) \in \mathbb{J}_p^n$, then for each $1 \leq i < j \leq n$ the Laurent series expansion

$$f(t_1, \dots, t_n, \tau) = \sum_{k \geq N} f_k(t_1, \dots, \hat{t}_i, \dots, t_n, \tau) (t_i - t_j)^k,$$

where the variable \hat{t}_i is missing in the coefficient $f_k \in \mathbb{J}_{p+k}^{n-1}$. These coefficients are given as an integral

$$f_k(t_1, \dots, \hat{t}_i, \dots, t_n, \tau) = \oint_{C_j} dt_i \frac{f(t_1, \dots, t_n, \tau)}{(t_i - t_j)^{k+1}},$$

where the contour C_j is a small circle in the t_i -plane around t_j and not containing any other point $t_l = t_l$ for $l \neq j$.

4.12. Being periodic, any element $f(t_1, \dots, t_n, \tau) \in \mathcal{F}^n$ admits a Fourier series. That is a formal power series

$$F(z_1, \dots, z_n, q) = \sum_{m_1, \dots, m_n \in \mathbb{Z}} F_{m_1, \dots, m_n}(q) z_1^{m_1} \dots z_n^{m_n}, \quad (4.14)$$

where each coefficient $F_{m_1, \dots, m_n}(q) \in \mathbb{C}[[q]]$. This series converges uniformly and absolutely in any closed subset of the domain $|qz_1| < |z_n| < \dots < |z_1|$ and it can be analytically continued to a meromorphic function with poles at $z_i = z_j q^m$, $i \neq j$, $m \in \mathbb{Z}$. In its domain of convergence we have

$$F(e^{2\pi i t_1}, \dots, e^{2\pi i t_n}, e^{2\pi i \tau}) = f(t_1, \dots, t_n, \tau).$$

Notice that for each $1 \leq i \leq n$ the series $F(z_1, \dots, z_{i-1}, qz_i, \dots, qz_n, q)$ converges absolutely and uniformly in any closed subset of the domain

$$|qz_i| < |z_{i-1}| < \dots < |z_1| < |z_n| < \dots < |z_i|,$$

and can be analytically extended to the same meromorphic function as $F(z_1, \dots, z_n, q)$. That is, we have ‘‘cyclically’’ shifted the order of the modulus of the variables in the domain of expansion. In the same vein, we will use the fact that

$$F(z_1, \dots, z_n, q) = F(qz_1, \dots, qz_n, q), \quad (4.15)$$

which follows since both are Fourier expansions of the same elliptic function on the same domain. The Fourier expansions of the functions in \mathbb{J}_*^2 have been described in 4.2.

4.13. For $f(t, \tau) \in \mathbb{J}_k^2$ the function

$$\left((2\pi i) \frac{d}{d\tau} - \zeta(t, \tau) \frac{d}{dt} \right) f(t, \tau), \quad (4.16)$$

is periodic in t with periods 1, τ . It is not, however, a Jacobi form. Indeed if we write

$$D = (2\pi i) \frac{d}{d\tau} - \zeta(t, \tau) \frac{d}{dt},$$

then it follows from [12, 2.15] that

$$(Df) \left(\frac{t}{c\tau + d}, \frac{a\tau + b}{c\tau + d} \right) = (c\tau + d)^{k+2} (Df)(t, \tau) + 2\pi i k c (c\tau + d)^{k+1} f(t, \tau). \quad (4.17)$$

Rather $Df \in \mathbb{J}_*^2 \otimes_{\mathbb{M}_*} \mathbb{Q}\mathbb{M}_*$. If $F(z, q)$ is the Fourier series of $f(t, \tau)$ in the domain $|q| < |z| < 1$

$$\left[(2\pi i)^2 q \frac{d}{dq} + (P_1(z, q) + \pi i) (2\pi i) z \frac{d}{dz} \right] F(z, q), \quad (4.18)$$

converges absolutely in the domain $|q| < |z| < 1$ and is the Fourier series of Df .

4.14 Lemma. Let $f = f(t_0, \dots, t_n, \tau) \in \mathcal{F}_{n+1}$. Define Df by

$$Df(t_0, \dots, t_n, \tau) = 2\pi i \frac{\partial f}{\partial \tau}(t_0, \dots, t_n, \tau) + \sum_{i=0}^{n-1} \zeta(t_n - t_i) \frac{\partial f}{\partial t_i}(t_0, \dots, t_n, \tau). \quad (4.19)$$

Then $Df \in \mathcal{F}_{n+1}$.

Proof. Fix $0 \leq j \leq n$, we take the total derivative with respect to τ of

$$f(t_0, \dots, t_j + \tau, \dots, t_n, \tau) - f(t_0, \dots, t_j, \dots, t_n, \tau) = 0,$$

obtaining

$$\frac{\partial f}{\partial \tau}(t_0, \dots, t_j + \tau, \dots, t_n, \tau) - \frac{\partial f}{\partial \tau}(t_0, \dots, t_j, \dots, t_n, \tau) = -\frac{\partial f}{\partial t_j}(t_0, \dots, t_j + \tau, \dots, t_n, \tau).$$

Recall that $\zeta(t + \tau, \tau) = \zeta(t, \tau) - 2\pi i$, by (4.6). So if we put

$$\widehat{f}(t_0, \dots, t_n, \tau) = \sum_{i=0}^{n-1} \zeta(t_n - t_i, \tau) \frac{\partial f}{\partial t_i}(t_0, \dots, t_n, \tau),$$

then for $0 \leq j \leq n-1$ we have

$$\begin{aligned} \widehat{f}(t_0, \dots, t_j + \tau, \dots, t_n, \tau) - \widehat{f}(t_0, \dots, t_j, \dots, t_n, \tau) \\ = 2\pi i \frac{\partial f}{\partial t_j}(t_0, \dots, t_j + \tau, \dots, t_n, \tau). \end{aligned}$$

In the case $j = n$ we have

$$\begin{aligned} \widehat{f}(t_0, \dots, t_n + \tau, \tau) &= \sum_{i=0}^{n-1} (\zeta(t_n - t_i) - 2\pi i) \frac{\partial f}{\partial t_i}(t_0, \dots, t_n, \tau) \\ &= \widehat{f}(t_0, \dots, t_n, \tau) - 2\pi i \sum_{i=0}^{n-1} \frac{\partial f}{\partial t_i}(t_0, \dots, t_n, \tau) = \widehat{f}(t_0, \dots, t_n, \tau) + 2\pi i \frac{\partial f}{\partial t_n}(t_0, \dots, t_n, \tau), \end{aligned}$$

where we used Lemma 4.9 in the last equation. It follows that

$$\left[2\pi i \frac{\partial}{\partial \tau} + \sum_{i=0}^{n-1} \zeta(t_n - t_i, \tau) \frac{\partial}{\partial t_i} \right] f(t_0, \dots, t_i, \dots, t_n, \tau),$$

is τ -periodic. It is obviously also 1-periodic. Hence $Df \in \mathcal{F}_{n+1}$ as desired. \square

4.15 Corollary. *Let $f \in \mathcal{F}_n$ and let $F(z_1, \dots, z_n, q)$ be its Fourier series in the domain $|qz_1| < |z_n| < \dots < |z_2| < |z_1|$. Then the series*

$$\left((2\pi i)^2 q \frac{d}{dq} - \sum_{i=1}^{n-1} \left(P_1 \left(\frac{z_n}{z_i}, q \right) + \pi i \right) 2\pi i z_i \frac{d}{dz_i} \right) F(z_1, \dots, z_n, q), \quad (4.20)$$

converges absolutely in the same domain, and is the Fourier series of $Df \in \mathcal{F}_n$.

4.16. Lemma 4.14 can be interpreted in the following way. Consider the semidirect product $\mathrm{SL}(2, \mathbb{Z}) \times \mathbb{Z}^{2n}$, where \mathbb{Z}^{2n} consists of n -copies of the standard $\mathrm{SL}(2, \mathbb{Z})$ representation. The coarse moduli space of n -marked elliptic curves is the quotient of $\mathbb{H} \times \mathbb{C}^n$ by the action of this group given as follows.

$$\begin{aligned} (k_1, l_1, \dots, k_n, l_n) \cdot (\tau, t_1, \dots, t_n) &= (\tau, t_1 + k_1 + l_1 \tau, \dots, t_n + k_n + l_n \tau), \\ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \cdot (\tau, t_1, \dots, t_n) &= \left(\frac{a\tau + b}{c\tau + d}, \frac{t_1}{c\tau + d}, \dots, \frac{t_n}{c\tau + d} \right). \end{aligned}$$

The vector fields $\frac{\partial}{\partial t_i}$, $i = 1, \dots, n-1$ are well defined in this quotient, but the vector field $\frac{\partial}{\partial \tau}$ is not. Instead, the vector field

$$D_\tau = 2\pi i \frac{\partial}{\partial \tau} + \sum_{i=1}^{n-1} \zeta(t_n - t_i, \tau) \frac{\partial}{\partial t_i}, \quad (4.21)$$

is well defined in the open locus where the points are distinct $t_i - t_j \notin \mathbb{Z} + \tau\mathbb{Z}$. Notice that the non-trivial commutators among these vector fields are

$$\left[D_\tau, \frac{\partial}{\partial t_i} \right] = -\overline{\varphi}_2(t_n - t_i) \frac{\partial}{\partial t_i}, \quad i = 1, \dots, n-1. \quad (4.22)$$

5 First chiral homology groups

In this section we define complexes computing chiral homology and their duals in genus 1. In 5.1–5.7 we define the complex computing chiral homology of the universal elliptic curve. In 5.8–5.14 we define the complex that computes chiral homology on the universal elliptic curve with n marked points. In 5.15 we study the connection obtained by moving the marked points. In 5.18–5.22 we describe the connection obtained as the base elliptic curve moves in the moduli space. We study the dual spaces of chiral homology in 5.24 and their flat sections with respect to variation of the marked points in 5.25. Finally we define flat sections of the dual of chiral homology in 5.26.

5.1. Let V be a vertex algebra. We define a complex

$$V^{\otimes 3} \otimes \mathbb{J}_*^3 \xrightarrow{d_2} V^{\otimes 2} \otimes \mathbb{J}_*^2 \xrightarrow{d_1} V \otimes \mathbb{J}_*^1,$$

as follows. For $f(t_1, t_2, \tau) = f(t_1 - t_2, \tau) \in \mathbb{J}_*^2$ we put

$$d_1(a \otimes b \otimes f(t_1, t_2)) = a_{(f)}b, \quad (5.1)$$

where the (f) -product is as defined in 2.9. In this expression, and in the sequel, we omit explicit reference to τ and we abide by the conventions of Section 2.1 and Example 2.2.

Now we let $f(t_1, t_2, t_3) \in \mathbb{J}_*^3$ and we consider the Laurent series expansions

$$f(t_1, t_2, t_3) = \sum_k f_{12k}(t_2, t_3)(t_1 - t_2)^k = \sum_k f_{13k}(t_2, t_3)(t_1 - t_3)^k = \sum_k f_{23k}(t_1, t_3)(t_2 - t_3)^k, \quad (5.2)$$

where $f_{ijk} \in \mathbb{J}_*^2$ and the sum runs over $k \geq N_{ij}$ for some fixed $N_{ij} \in \mathbb{Z}$. For instance when $i = 1, j = 2$ we have

$$f(t_1, t_2, t_3) = f(t_2 + (t_1 - t_2), t_2, t_3) = \sum_{k \geq N_{12}} f_{12k}(t_2, t_3)(t_1 - t_2)^k.$$

We now put

$$\begin{aligned} d_2(a \otimes b \otimes c \otimes f(t_1, t_2, t_3)) &= \sum_k a \otimes b_{(k)}c \otimes f_{23k}(t_1, t_2) \\ &\quad - \sum_k b \otimes a_{(k)}c \otimes f_{13k}(t_1, t_2) - \sum_k a_{(k)}b \otimes c \otimes f_{12k}(t_1, t_2). \end{aligned} \quad (5.3)$$

The following lemma is a straightforward consequence of the Borchers identity.

5.2 Lemma. $d_1 \circ d_2 = 0$.

Proof. Let $f(t_1, t_2, t_3) \in \mathbb{J}_*^3$. In the notation of Example 2.2 we have

$$\sum_k (a_{(k)}b)_{(f_{12k})}c = \text{res}_{t_2-t_3} \text{res}_{t_1-t_2} Y\left(Y(a, t_1 - t_2)b, t_2 - t_3\right) c i_{t_2, t_2-t_3, t_1-t_2} f(t_1, t_2, t_3),$$

and similar expressions for the other terms of (5.3). Thus

$$\begin{aligned} &d_1(d_2(a \otimes b \otimes c \otimes f(t_1, t_2, t_3))) \\ &= \text{res}_{t_1-t_3} \text{res}_{t_2-t_3} Y(a, t_1 - t_3)Y(b, t_2 - t_3)c i_{t_1, t_1-t_3, t_2-t_3} f(t_1, t_2, t_3) \\ &\quad - \text{res}_{t_2-t_3} \text{res}_{t_1-t_3} Y(b, t_2 - t_3)Y(a, t_1 - t_3)c i_{t_2, t_2-t_3, t_1-t_3} f(t_1, t_2, t_3) \\ &\quad - \text{res}_{t_2-t_3} \text{res}_{t_1-t_2} Y(Y(a, t_1 - t_2)b, t_2 - t_3)c i_{t_2, t_2-t_3, t_1-t_2} f(t_1, t_2, t_3). \end{aligned} \quad (5.4)$$

The lemma follows from Borchers identity (2.2) with $z = t_1 - t_3$ and $w = t_2 - t_3$. \square

5.3 Lemma. For any $a, b \in V$ and $f(t_1 - t_2) \in \mathbb{J}_*^2$ we have

$$\begin{aligned} d_1(L_{-1}a \otimes b \otimes f + a \otimes b \otimes f') &= 0, \\ d_1(a \otimes L_{-1}b \otimes f - a \otimes b \otimes f') &= L_{-1}d_1(a \otimes b \otimes f). \end{aligned} \quad (5.5)$$

Proof. This is simply an application of (2.6). □

5.4. We now define $C_0 = V \otimes \mathbb{J}_*^1$ and

$$C_1 = V^{\otimes 2} \otimes \mathbb{J}_*^2 / \left\langle L_{-1}a \otimes b \otimes f + a \otimes b \otimes f' \right\rangle, \quad (5.6)$$

and we define C_2 to be the quotient of $V^{\otimes 3} \otimes \mathbb{J}_*^3$ by the subspace spanned by the terms

$$\begin{aligned} & L_{-1}a \otimes b \otimes c \otimes f + a \otimes b \otimes c \otimes (\partial f / \partial t_1), \\ & a \otimes L_{-1}b \otimes c \otimes f + a \otimes b \otimes c \otimes (\partial f / \partial t_2), \\ & a \otimes b \otimes c \otimes f(t_1, t_2, t_3) + b \otimes a \otimes c \otimes f(t_2, t_1, t_3). \end{aligned}$$

The differentials d_1 and d_2 descend to define a complex C_\bullet as we shall see below. Let us denote by π the quotient map $V^{\otimes 2} \otimes \mathbb{J}_*^2 \rightarrow C_1$.

5.5 Lemma. *For any $a, b, c \in V$ and $f \in \mathbb{J}_*^3$ we have*

$$\begin{aligned} \pi \circ d_2 \left(L_{-1}a \otimes b \otimes c \otimes f(t_1, t_2, t_3) + a \otimes b \otimes c \otimes \frac{d}{dt_1} f(t_1, t_2, t_3) \right) &= 0, \\ \pi \circ d_2 \left(a \otimes L_{-1}b \otimes c \otimes f(t_1, t_2, t_3) + a \otimes b \otimes c \otimes \frac{d}{dt_2} f(t_1, t_2, t_3) \right) &= 0, \\ d_2 \left(a \otimes b \otimes L_{-1}c \otimes f(t_1, t_2, t_3) + a \otimes b \otimes c \otimes \frac{d}{dt_3} f(t_1, t_2, t_3) \right) &= \\ & \left(L_{-1}^{(2)} + \frac{d}{dt_2} \right) d_2 \left(a \otimes b \otimes c \otimes f(t_1, t_2, t_3) \right), \end{aligned} \quad (5.7)$$

where $L_{-1}^{(2)}$ is the operator L_{-1} applied to the second factor of $V \otimes V \otimes \mathbb{J}_*^2$.

Proof. If we write $g = \partial f / \partial t_1$ then the coefficients of the Laurent series expansions of f and g are related by

$$g_{23k} = f'_{23k}, \quad g_{13, k-1} = k f_{13k} \quad \text{and} \quad g_{12, k-1} = k f_{12k}.$$

Therefore

$$\begin{aligned} d_2 \left(a \otimes b \otimes c \otimes \frac{d}{dt_1} f(t_1, t_2, t_3) \right) &= \sum_k a \otimes b_{(k)} c \otimes f'_{23k} \\ & \quad - \sum_k (k+1) b \otimes a_{(k)} c \otimes f_{13, k+1} - \sum_k (k+1) a_{(k)} b \otimes c \otimes f_{12, k+1}. \end{aligned} \quad (5.8)$$

On the other hand, replacing a by $L_{-1}a$ in (5.3) and applying π , we have

$$\begin{aligned} \pi \circ d_2 \left(L_{-1}a \otimes b \otimes c \otimes f(t_1, t_2, t_3) \right) &= \sum_k L_{-1}a \otimes b_{(k)} c \otimes f_{23k} \\ & \quad - \sum_k b \otimes (L_{-1}a)_{(k)} c \otimes f_{13k} - \sum_k (L_{-1}a)_{(k)} b \otimes c \otimes f_{12k} = \\ & \quad - \sum_k a \otimes b_{(k)} c \otimes f'_{23k} + \sum_k k b \otimes a_{(k-1)} c \otimes f_{13k} + \sum_k k a_{(k-1)} b \otimes c \otimes f_{12k}. \end{aligned} \quad (5.9)$$

Thus the first equation of (5.7) is proved. The second equation is proved in the same way. In order to prove the third equation we consider $h = \partial f / \partial t_3$, then the coefficients of the Laurent series expansions of f and h are related by

$$h_{12k} = -f'_{12k}, \quad h_{13k} = -(f'_{13k} + (k+1)f_{13, k+1}), \quad h_{23k} = -(f'_{23k} + (k+1)f_{23, k+1}).$$

Therefore

$$\begin{aligned}
d_2 \left(a \otimes b \otimes c \otimes \frac{d}{dt_3} f(t_1, t_2, t_3) \right) &= - \sum_k a \otimes b_{(k)} c \otimes (f'_{23k} + (k+1)f_{23,k+1}) \\
&\quad + \sum_k b \otimes a_{(k)} c \otimes (f'_{13k} + (k+1)f_{13,k+1}) + \sum_k a_{(k)} b \otimes c \otimes f'_{12k} \\
&= - \sum_k a \otimes b_{(k)} c \otimes f'_{23k} + \sum_k b \otimes a_{(k)} c \otimes f'_{13k} + \sum_k a_{(k)} b \otimes c \otimes f'_{12k} \\
&\quad + \sum_k a \otimes (L_{-1}b)_{(k)} c \otimes f_{23k} - \sum_k b \otimes (L_{-1}a)_{(k)} c \otimes f_{13k}.
\end{aligned} \tag{5.10}$$

The sum of the first three terms equals $\frac{d}{dt_2} d_2(a \otimes b \otimes c \otimes f(t_1, t_2, t_3))$. On the other hand replacing c by $L_{-1}c$ in (5.3) we have

$$\begin{aligned}
d_2 \left(a \otimes b \otimes L_{-1}c \otimes f(t_1, t_2, t_3) \right) &= \sum_k a \otimes b_{(k)} L_{-1}c \otimes f_{23k} \\
&\quad - \sum_k b \otimes a_{(k)} L_{-1}c \otimes f_{13k} - \sum_k a_{(k)} b \otimes L_{-1}c \otimes f_{12k} \\
&= \sum_k a \otimes L_{-1}(b_{(k)}c) \otimes f_{23k} - \sum_k a \otimes (L_{-1}b)_{(k)} c \otimes f_{23k} \\
&\quad - \sum_k b \otimes L_{-1}(a_{(k)}c) \otimes f_{13k} + \sum_k b \otimes (L_{-1}a)_{(k)} c \otimes f_{13k} - \sum_k a_{(k)} b \otimes L_{-1}c \otimes f_{12k}.
\end{aligned} \tag{5.11}$$

The sum of the first, third and last term of the right hand side equal $L_{-1}^{(2)}(d_2(a \otimes b \otimes c \otimes f))$, the remaining terms cancel the last two terms of (5.10). \square

5.6 Lemma. *We have the identity*

$$\pi \circ d_2 \left(a \otimes b \otimes c \otimes f(t_1, t_2, t_3) \right) = -\pi \circ d_2 \left(b \otimes a \otimes c \otimes f(t_2, t_1, t_3) \right). \tag{5.12}$$

Proof. Let $g(t_1, t_2, t_3) = f(t_2, t_1, t_3)$. Then the Laurent expansions of g and of f are related by

$$f_{23k} = g_{13k}, \quad f_{13k} = g_{23k}, \quad f_{12k} = \sum_{j \geq 0} (-1)^{k+j} g_{12,k-j}^{(j)}.$$

The last one is obtained from

$$g(t_2, t_1, t_3) = \sum_j f_{12j}(t_2, t_3)(t_1 - t_2)^j = \sum_j (-1)^j g_{12j}(t_1, t_3)(t_1 - t_2)^j,$$

after multiplying by $(t_1 - t_2)^{-k-1}$ and taking residues. By (5.3) we have

$$d_2 \left(b \otimes a \otimes c \otimes g(t_1, t_2, t_3, \tau) \right) = \sum_k b \otimes a_{(k)} c \otimes f_{13k} - \sum_k a \otimes b_{(k)} c \otimes f_{23k} - \sum_k b_{(k)} a \otimes c \otimes g_{12k}. \tag{5.13}$$

We use the skew-symmetry identity (2.3) to write

$$\begin{aligned}
\pi \sum_k b_{(k)} a \otimes c \otimes g_{12k} &= -\pi \sum_k \sum_{j \geq 0} (-1)^k \frac{(-L_{-1})^j}{j!} a_{(k+j)} b \otimes c \otimes g_{12k} = \\
&= -\pi \sum_k \sum_{j \geq 0} (-1)^k a_{(k+j)} b \otimes c \otimes g_{12k}^{(j)} = -\pi \sum_k a_{(k)} b \otimes c \otimes f_{12k}.
\end{aligned}$$

This proves the Lemma. \square

5.7. Due to Lemmas 5.5 and 5.6 we may define the following complex

$$C_2 \xrightarrow{d_2} C_1 \xrightarrow{d_1} C_0 \rightarrow 0. \quad (5.14)$$

Notice that this is a complex of $\mathbb{M}_*[\partial]$ -modules, where ∂ acts by $L_{-1}^{(k+1)} + \frac{d}{dt_{k+1}}$ on C_k .

For $i = 0, 1$ the homology $H_i(C_\bullet)$ is called the i^{th} *chiral homology group* of the vertex algebra V in genus 1. We will denote these groups by $H_i^{\text{ch}}(V)$, leaving the τ dependence implicit. They correspond to the chiral homology groups of the universal elliptic curve with coefficients in the chiral algebra associated to V as defined in [1] and explained in [6]. $H_0^{\text{ch}}(V)$ is also known as the space of *coinvariants* of V in genus 1 and its dual is closely related to the space of *conformal blocks* of V in genus one as defined and studied in [2]. We will comment more on this below when we study the connection on this complex.

5.8 Homology with supports. More generally we may define, for fixed $n \geq 1$, a complex

$$V^{\otimes(n+2)} \otimes \mathbb{J}_*^{n+2} \xrightarrow{d_2} V^{\otimes(n+1)} \otimes \mathbb{J}_*^{n+1} \xrightarrow{d_1} V^{\otimes n} \otimes \mathbb{J}_*^n \rightarrow 0, \quad (5.15)$$

as follows. Let $f(t_0, \dots, t_n) \in \mathbb{J}_*^{n+1}$ and let $1 \leq i \leq n$. We consider the Laurent series expansion

$$f(t_0, \dots, t_n) = \sum_k f_{ik}(t_1, \dots, t_n)(t_0 - t_i)^k.$$

By Lemma 4.11 we have $f_{ik} \in \mathbb{J}_*^n$ for each $i = 1, \dots, n$ and for all integers k , and we define

$$d_1(a \otimes a_1 \otimes \dots \otimes a_n \otimes f(t_0, \dots, t_n)) = \sum_{i=1}^n \sum_k a_1 \otimes \dots \otimes (a_{(k)} a_i) \otimes \dots \otimes a_n \otimes f_{ik}. \quad (5.16)$$

For $f(u, v, t_1, \dots, t_n) \in \mathbb{J}_*^{n+2}$ we will consider similarly the following expansions for $i = 1, \dots, n$:

$$\begin{aligned} f(u, v, t_1, \dots, t_n) &= \sum_k f_{0ik}(v, t_1, \dots, t_n)(u - t_i)^k, \\ &= \sum_k f_{1ik}(u, t_1, \dots, t_n)(v - t_i)^k = \sum_k f_{00k}(v, t_1, \dots, t_n)(u - v)^k. \end{aligned} \quad (5.17)$$

We define

$$\begin{aligned} d_2(a \otimes b \otimes a_1 \otimes \dots \otimes a_n \otimes f) &= \sum_{i=1}^n \sum_k a \otimes a_1 \otimes \dots \otimes (b_{(k)} a_i) \otimes \dots \otimes a_n \otimes f_{1ik}(t_0, \dots, t_n) \\ &\quad - \sum_{i=1}^n \sum_k b \otimes a_1 \otimes \dots \otimes (a_{(k)} a_i) \otimes \dots \otimes a_n \otimes f_{0ik}(t_0, \dots, t_n) \\ &\quad - \sum_k a_{(k)} b \otimes a_1 \otimes \dots \otimes a_n \otimes f_{00k}(t_0, \dots, t_n). \end{aligned} \quad (5.18)$$

5.9 Lemma. $d_1 \circ d_2 = 0$

Proof. The proof of this lemma is similar to that of Lemma 5.2. Let $f(u, v, t_1, \dots, t_n) \in \mathbb{J}_*^{n+2}$. We have

$$\begin{aligned} &d_1(d_2(a \otimes b \otimes a_1 \otimes \dots \otimes a_n \otimes f(u, v, t_1, \dots, t_n))) \\ &= \sum_{i=1}^n \sum_{j=1}^{i-1} \text{res}_{u-t_j} \text{res}_{v-t_i} a_1 \otimes \dots \otimes Y(a, u - t_j) a_j \otimes \dots \otimes Y(b, v - t_i) a_i \otimes \dots \otimes a_n \otimes i_{u, u-t_j} i_{v, v-t_i} f \\ &\quad + \sum_{i=1}^n \text{res}_{u-t_i} \text{res}_{v-t_i} a_1 \otimes \dots \otimes Y(a, u - t_i) Y(b, v - t_i) a_i \otimes \dots \otimes a_n \otimes i_{t_i, u-t_i, v-t_i} f \\ &\quad + \sum_{i=1}^n \sum_{j=i+1}^n \text{res}_{u-t_j} \text{res}_{v-t_i} a_1 \otimes \dots \otimes Y(b, v - t_i) a_i \otimes \dots \otimes Y(a, u - t_j) a_j \otimes \dots \otimes a_n \otimes i_{u, u-t_j} i_{v, v-t_i} f \end{aligned}$$

$$\begin{aligned}
& - \sum_{i=1}^n \sum_{j=1}^{i-1} \text{res}_{v-t_j} \text{res}_{u-t_i} a_1 \otimes \cdots \otimes Y(b, v-t_j) a_j \otimes \cdots \otimes Y(a, u-t_i) a_i \otimes \cdots \otimes a_n \otimes i_{u, u-t_i} i_{v, v-t_j} f \\
& - \sum_{i=1}^n \text{res}_{v-t_i} \text{res}_{u-t_i} a_1 \otimes \cdots \otimes Y(b, v-t_i) Y(a, u-t_i) a_i \otimes \cdots \otimes a_n \otimes i_{t_i, v-t_i, u-t_i} f \tag{5.19} \\
& - \sum_{i=1}^n \sum_{j=i+1}^n \text{res}_{v-t_j} \text{res}_{v-t_i} a_1 \otimes \cdots \otimes Y(a, u-t_i) a_i \otimes \cdots \otimes Y(b, v-t_j) a_j \otimes \cdots \otimes a_n \otimes i_{u, u-t_j} i_{v, v-t_i} f \\
& - \sum_{i=1}^n \text{res}_{v-t_i} \text{res}_{u-v} a_1 \otimes \cdots \otimes Y(Y(a, u-v) b, v-t_i) a_i \otimes \cdots \otimes a_n \otimes i_{t_i, v-t_i, u-v} f \\
& = \sum_{i=1}^n \text{res}_{u-t_i} \text{res}_{v-t_i} a_1 \otimes \cdots \otimes Y(a, u-t_i) Y(b, v-t_i) a_i \otimes \cdots \otimes a_n \otimes i_{t_i, u-t_i, v-t_i} f \\
& - \sum_{i=1}^n \text{res}_{v-t_i} \text{res}_{u-t_i} a_1 \otimes \cdots \otimes Y(b, v-t_i) Y(a, u-t_i) a_i \otimes \cdots \otimes a_n \otimes i_{t_i, v-t_i, u-t_i} f \\
& - \sum_{i=1}^n \text{res}_{v-t_i} \text{res}_{u-v} a_1 \otimes \cdots \otimes Y(Y(a, u-v) b, v-t_i) a_i \otimes \cdots \otimes a_n \otimes i_{t_i, v-t_i, u-v} f,
\end{aligned}$$

and this vanishes because of Borchers identity (2.2). \square

From now on, it will be convenient to denote by $L_{-1}^{(i)}$ the operator $1 \otimes \cdots \otimes L_{-1} \otimes \cdots \otimes 1$ acting on $a_1 \otimes \cdots \otimes a_n \in V^{\otimes n}$ or $a_0 \otimes a_1 \otimes \cdots \otimes a_n \in V^{\otimes(n+1)}$ or $a \otimes b \otimes a_1 \otimes \cdots \otimes a_n \in V^{\otimes(n+2)}$ with L_{-1} acting in the component labelled 'i'.

5.10 Lemma. For $a_0, a_1, \dots, a_n \in V$ and $f(t_0, t_1, \dots, t_n) \in \mathbb{J}_*^{n+1}$ we have

$$d_1 \left(\left[L_{-1}^{(0)} + \frac{\partial}{\partial t_0} \right] a_0 \otimes \cdots \otimes a_n \otimes f(t_0, \dots, t_n) \right) = 0, \tag{5.20}$$

and for $j = 1, 2, \dots, n$ we have

$$d_1 \left(\left[L_{-1}^{(j)} + \frac{\partial}{\partial t_j} \right] a_0 \otimes \cdots \otimes a_n \otimes f(t_0, \dots, t_n) \right) = \left(L_{-1}^{(j)} + \frac{\partial}{\partial t_j} \right) d_1 (a_0 \otimes \cdots \otimes a_n \otimes f(t_0, \dots, t_n)). \tag{5.21}$$

Proof. Recalling the Laurent series expansion $f = \sum_k f_{0jk}(t_1, \dots, t_n)(t_0 - t_j)^k$ and the definition of d_1 , we have

$$\begin{aligned}
d_1 (L_{-1} a_0 \otimes \cdots \otimes a_n \otimes f(t_0, \dots, t_n)) &= \sum_{j=1}^n \sum_k a_1 \otimes \cdots \otimes ((L_{-1} a_0)_{(k)} a_j) \otimes \cdots \otimes a_n \otimes f_{jk} \\
&= - \sum_{j=1}^n \sum_k a_1 \otimes \cdots \otimes (a_{0(k-1)} a_j) \otimes \cdots \otimes a_n \otimes k f_{jk} \\
&= -d_1 \left(a_0 \otimes \cdots \otimes a_n \otimes \frac{\partial f}{\partial t_0}(t_0, \dots, t_n) \right),
\end{aligned}$$

and the first claim follows. Similarly the second claim follows from the equality

$$\begin{aligned}
& \sum_k a_1 \otimes \cdots \otimes (a_{0(k)} L_{-1} a_j) \otimes \cdots \otimes a_n \otimes f_{0jk}(t_0 - t_j)^k \\
&= \sum_k a_1 \otimes \cdots \otimes L_{-1}(a_{0(k)} a_j) \otimes \cdots \otimes a_n \otimes f_{0jk}(t_0 - t_j)^k \\
&\quad - \sum_k a_1 \otimes \cdots \otimes (a_{0(k-1)} a_j) \otimes \cdots \otimes a_n \otimes k f_{0jk}(t_0 - t_j)^k.
\end{aligned}$$

\square

We define \tilde{C}_0 to be $V^{\otimes n} \otimes \mathbb{J}_*^n$ and \tilde{C}_1 to be the quotient of $V^{\otimes(n+1)} \otimes \mathbb{J}_*^{n+1}$ by the linear span of the terms

$$\left(L_{-1}^{(0)} + \frac{\partial}{\partial t_0} \right) a_0 \otimes \cdots \otimes a_n \otimes f(t_0, \dots, t_n).$$

We denote by π the corresponding quotient map $V^{\otimes(n+1)} \otimes \mathbb{J}_*^{n+1} \rightarrow \tilde{C}_1$. We notice that $d_1 : \tilde{C}_1 \rightarrow \tilde{C}_0$ is well defined by (5.20) and it is a map of $k[\partial_1, \dots, \partial_n]$ -modules by (5.21). In addition, recalling that we view \mathbb{J}_*^{n+1} as a \mathbb{J}_*^n -module as in 4.6 we see that the map d_1 is \mathbb{J}_*^n -linear. These two actions make the complex $\tilde{C}_1 \rightarrow \tilde{C}_0$ a complex of D -modules. The following lemma is proved in the same way as above.

5.11 Lemma. *Let $a, b, a_1 \dots a_n \in V$ and $f(u, v, t_1, \dots, t_n) \in \mathbb{J}_*^{n+2}$. Then*

$$\pi \circ d_2 (L_{-1}a \otimes b \otimes a_1 \otimes \cdots \otimes a_n \otimes f) + \pi \circ d_2 \left(a \otimes b \otimes a_1 \otimes \cdots \otimes a_n \otimes \frac{d}{du} f(u, v, t_1, \dots, t_n) \right) = 0 \quad (5.22)$$

$$\pi \circ d_2 (a \otimes L_{-1}b \otimes a_1 \otimes \cdots \otimes a_n \otimes f) + \pi \circ d_2 \left(a \otimes b \otimes a_1 \otimes \cdots \otimes a_n \otimes \frac{d}{dv} f(u, v, t_1, \dots, t_n) \right) = 0 \quad (5.23)$$

$$\begin{aligned} & d_2 \left(\left(L_{-1}^{(j)} + \frac{d}{dt_j} \right) a \otimes b \otimes a_1 \otimes \cdots \otimes a_n \otimes f(u, v, t_1, \dots, t_n) \right) \\ &= \left(L_{-1}^{(j)} + \frac{d}{dt_j} \right) d_2 (a \otimes b \otimes a_1 \otimes \cdots \otimes a_n \otimes f(u, v, t_1, \dots, t_n)), \quad 1 \leq j \leq n. \end{aligned} \quad (5.24)$$

We also have as in Lemma 5.6:

5.12 Lemma. *Let $a, b, a_1, \dots, a_n \in V$ and $f(u, v, t_1, \dots, t_n) \in \mathbb{J}_*^{n+2}$. Then*

$$\pi d_2 (a \otimes b \otimes a_1 \otimes \cdots \otimes a_n \otimes f(u, v, t_1, \dots, t_n)) + \pi d_2 (b \otimes a \otimes a_1 \otimes \cdots \otimes a_n \otimes f(v, u, t_1, \dots, t_n)) = 0.$$

5.13. The complex (5.15) carries an action of the symmetric group S_n . Indeed for $\sigma \in S_n$ a permutation we have

$$\begin{aligned} \sigma(a_1 \otimes \cdots \otimes a_n \otimes f(t_1, \dots, t_n)) &= a_{\sigma(1)} \otimes \cdots \otimes a_{\sigma(n)} \otimes f(v, t_{\sigma(1)}, \dots, t_{\sigma(n)}), \\ \sigma(a \otimes a_1 \otimes \cdots \otimes a_n \otimes f(v, t_1, \dots, t_n)) &= a \otimes a_{\sigma(1)} \otimes \cdots \otimes a_{\sigma(n)} \otimes f(v, t_{\sigma(1)}, \dots, t_{\sigma(n)}), \\ \sigma(a \otimes b \otimes a_1 \otimes \cdots \otimes a_n \otimes f(u, v, t_1, \dots, t_n)) &= a \otimes b \otimes a_{\sigma(1)} \otimes \cdots \otimes a_{\sigma(n)} \otimes f(u, v, t_{\sigma(1)}, \dots, t_{\sigma(n)}). \end{aligned}$$

It is immediate that the differentials d_1 and d_2 commute with this action, making the complex (5.15) a complex in the category of S_n -modules.

5.14 Definition of chiral homology. Let V be a vertex algebra and $n \geq 1$.

a) Let C_0^n be $V^{\otimes n} \otimes \mathbb{J}_*^n$.

b) Let C_1^n be the quotient of $V^{\otimes(n+1)} \otimes \mathbb{J}_*^{n+1}$ by the linear span of the terms

$$L_{-1}a_0 \otimes \cdots \otimes a_n \otimes f(t_0, \dots, t_n) + a_0 \otimes \cdots \otimes a_n \otimes \frac{d}{dt_0} f(t_0, \dots, t_n). \quad (5.25)$$

c) Let C_2^n be the quotient of $V^{\otimes(n+2)} \otimes \mathbb{J}_*^{n+2}$ by the linear span of the terms

$$\begin{aligned} & a \otimes b \otimes a_1 \otimes \cdots \otimes a_n \otimes f(u, v, t_1, \dots, t_n) + b \otimes a \otimes a_1 \otimes \cdots \otimes a_n \otimes f(v, u, t_1, \dots, t_n), \\ & L_{-1}a \otimes b \otimes a_1 \otimes \cdots \otimes a_n \otimes f(u, v, t_1, \dots, t_n) + a \otimes b \otimes a_1 \otimes \cdots \otimes a_n \otimes \frac{d}{du} f(u, v, t_1, \dots, t_n). \end{aligned} \quad (5.26)$$

Let

$$C_2^m \xrightarrow{d_2} C_1^n \xrightarrow{d_1} C_0^m \rightarrow 0, \quad (5.27)$$

be the complex with differentials d_1 and d_2 as in (5.16) and (5.18) above. For $i = 0, 1$ we define i^{th} *chiral homology* of V in genus 1, with coefficients in $V^{\otimes n}$ to be the homology $H_i(C_\bullet^n)$. We denote this by $H_i^{\text{ch}}(V^{\otimes n})$.

5.15 Moving the points. The complex C_\bullet^n is a complex of \mathbb{J}_*^n -modules and a complex of $k[\partial_1, \dots, \partial_n]$ -modules. The action of \mathbb{J}_*^n is by multiplication on the right and the action of ∂_j is given by

$$\begin{aligned} & \left(L_{-1}^{(j)} + \frac{d}{dt_j} \right) a_1 \otimes \cdots \otimes a_n \otimes f(t_1, \dots, t_n), \quad \text{in degree 0,} \\ & \left(L_{-1}^{(j)} + \frac{d}{dt_j} \right) a_0 \otimes a_1 \otimes \cdots \otimes a_n \otimes f(t_0, t_1, \dots, t_n), \quad \text{in degree 1,} \\ & \left(L_{-1}^{(j)} + \frac{d}{dt_j} \right) a \otimes b \otimes a_1 \otimes \cdots \otimes a_n \otimes f(u, v, t_1, \dots, t_n), \quad \text{in degree 2,} \end{aligned} \quad (5.28)$$

where, in each case, $j = 1, \dots, n$. These two actions are compatible making C_\bullet^n a complex of D -modules. We let $H_i^{\text{ch}}(V^{\otimes n})_{\text{tr}}$ be the coinvariants by this action, namely the quotient of $H_i^{\text{ch}}(V^{\otimes n})$ by elements of the form (5.28).

5.16. In the definition of $H_i^{\text{ch}}(V^{\otimes n})_{\text{tr}}$ given in 5.15 above, the same result is obtained if we quotient the terms (5.28) for $j = 1, \dots, n-1$ rather than $j = 1, \dots, n$. Indeed let us write $\tilde{H}_i^{\text{ch}}(V^{\otimes n})_{\text{tr}}$ for the quotient of $H_i^{\text{ch}}(V^{\otimes n})$ by (5.28) for $j = 1, \dots, n-1$. We have obvious surjections $\pi_i : \tilde{H}_i^{\text{ch}}(V^{\otimes n})_{\text{tr}} \rightarrow H_i^{\text{ch}}(V^{\otimes n})_{\text{tr}}$.

Let $\alpha = a_1 \otimes \cdots \otimes a_n \otimes f(t_1, \dots, t_n, \tau) \in C_0^n$. Put $g(t_0, \dots, t_n, \tau) = f(t_1, \dots, t_n, \tau)$ so that g is independent of t_0 . Put $\beta = \omega \otimes a_1 \otimes \cdots \otimes a_n \otimes g \in C_1^n$. We have

$$d_1 \beta = \sum_{i=1}^n a_1 \otimes \cdots \otimes L_{-1} a_i \otimes \cdots \otimes a_n \otimes f(t_1, \dots, t_n, \tau) \in C_0^n.$$

Therefore in $\tilde{H}_0^{\text{ch}}(V^{\otimes n})_{\text{tr}}$ we have

$$\begin{aligned} a_1 \otimes \cdots \otimes a_{n-1} \otimes L_{-1} a_n \otimes f &= - \sum_{i=1}^{n-1} a_1 \otimes \cdots \otimes L_{-1} a_i \otimes \cdots \otimes a_n \otimes f = \\ & \sum_{i=1}^{n-1} a_1 \otimes \cdots \otimes a_n \otimes \frac{\partial}{\partial t_i} f = -a_1 \otimes \cdots \otimes a_n \otimes \frac{\partial}{\partial t_n} f, \end{aligned}$$

where in the last equality we used Lemma 4.9. Equivalently

$$\left(L_{-1}^{(n)} + \frac{\partial}{\partial t_n} \right) \alpha = 0.$$

It follows that π_0 is an isomorphism.

Similarly let $\alpha = a_0 \otimes \cdots \otimes a_{n+1} \otimes f(t_0, \dots, t_n, \tau) \in C_1^n$, be a cycle, that is $d_1 \alpha = 0$. Put $g(u, v, t_1, \dots, t_n, \tau) = f(v, t_1, \dots, t_n, \tau)$, so that g is independent of u and let $\beta = \omega \otimes a_0 \otimes a_1 \otimes \cdots \otimes a_n \otimes g \in C_2^n$. We have

$$d_2 \beta = - \sum_{i=0}^{n-1} a_0 \otimes \cdots \otimes L_{-1} a_i \otimes \cdots \otimes a_n \otimes f(t_0, \dots, t_n, \tau) \in C_1^n,$$

where we have used that $d_1 \alpha = 0$ so that the first term in the right hand side of (5.18) vanishes. Therefore in $\tilde{H}_1^{\text{ch}}(V^{\otimes n})_{\text{tr}}$ it holds

$$a_0 \otimes \cdots \otimes a_{n-1} \otimes L_{-1} a_n \otimes f = - \sum_{i=0}^{n-1} a_0 \otimes \cdots \otimes L_{-1} a_i \otimes \cdots \otimes a_n \otimes f =$$

$$\sum_{i=0}^{n-1} a_0 \otimes \cdots \otimes a_n \otimes \frac{\partial}{\partial t_i} f = -a_0 \otimes \cdots \otimes a_n \otimes \frac{\partial}{\partial t_n} f.$$

It follows that π_1 is an isomorphism. In the case of $n = 1$ we obtain in particular

$$H_i^{\text{ch}}(V) \cong H_i^{\text{ch}}(V)_{\text{tr}}, \quad i = 0, 1. \quad (5.29)$$

5.17 Remark. The homology groups $H_i^{\text{ch}}(V) \cong H_i^{\text{ch}}(V)_{\text{tr}}$ recover the chiral homology (of degree $i = 0, 1$) of the torus with coefficients in the translation invariant chiral algebra \mathcal{A}_V associated with the conformal vertex algebra V [1, 6]. To compare the construction above with that given in [6, Proposition 7.13] we temporarily denote by \mathbb{J}_*^n the ring of meromorphic elliptic functions $f(t_1, \dots, t_n, \tau)$ with Jacobi dependence on the modular parameter τ and with possible poles at $t_i = t_j$ for $i \neq j$ and $t_i = 0$ for $i = 1, \dots, n$. Due to the group structure of the elliptic curve $E_\tau = \mathbb{C}/\mathbb{Z} + \mathbb{Z}\tau$ there is no essential difference between \mathbb{J}_*^n and \mathbb{J}_*^{n+1} , indeed the map $g \mapsto f$ defined by $f(t, \dots, t_n) = g(t_1 - t, \dots, t_n - t)$ is a ring isomorphism $\mathbb{J}_*^n \rightarrow \mathbb{J}_*^{n+1}$.

We specialize equations (5.16) and (5.18) to the case of $n = 1$ and we identify $f(t_0, t_1) = g(t_0 - t_1) \in \mathbb{J}_*^2$ with $g(t_0) \in \mathbb{J}_*^1$ and $f(u, v, t_1) = g(u - t_1, v - t_1) \in \mathbb{J}_*^3$ with $g(u, v) \in \mathbb{J}_*^2$ as above. With these identifications the differentials may be rewritten as:

$$d_1(f(t_0) \cdot a \otimes a_1) = a_{(f)} a_1, \quad (5.30)$$

and

$$\begin{aligned} d_2(f(u, v) \cdot a \otimes b \otimes a_1) &= \text{res}_v f(u, v) a \otimes b(v) a_1 - \text{res}_{v-u} f(v, u) a(v-u) b \otimes a_1 \\ &\quad - \text{res}_v f(v, u) b \otimes a(v) a_1. \end{aligned} \quad (5.31)$$

In [6, Proposition 7.13] we have written down a complex A^\bullet which computes chiral homology $H_i^{\text{ch}}(E_\tau, \mathcal{A}_V)$ for $i = 0, 1$ for the elliptic curve $E_\tau = \mathbb{C}/\mathbb{Z} + \mathbb{Z}\tau$ with coefficients in the translation invariant chiral algebra \mathcal{A}_V over E_τ associated with the conformal vertex algebra V . Comparison of A^\bullet with $C_\bullet^{n=1}$, particularly with equations (5.30) and (5.31), reveals that

$$H_i^{\text{ch}}(V) \cong H_i^{\text{ch}}(X, \mathcal{A}_V),$$

for $i = 0, 1$. In fact there are two small differences between A^\bullet and $C_\bullet^{n=1}$: the first is that coinvariants by the action of S_2 are omitted from the definition of A^\bullet in [6], but this does not alter the homology groups in degrees 0 and 1. The second difference is conventional, with A^\bullet being a cohomological complex concentrated in non positive degrees and C_\bullet^n a homological complex concentrated in non negative degrees.

5.18 Connections. We extend scalars in the complex C_\bullet^n from the ring \mathbb{J}_*^n of Jacobi forms to the ring \mathcal{F}_n of meromorphic elliptic functions. Recall from 4.6 that we have chosen a \mathbb{J}_*^n -module structure on \mathbb{J}_*^{n+k} and therefore on \mathcal{F}_{n+k} . We consider $C_k^n \otimes_{\mathbb{J}_*^n} \mathcal{F}_{n+k}$. In addition to the action of $\frac{\partial}{\partial t_i}$ for $i = 1, \dots, n$, these complexes carry a canonical action of the vector field D_τ of (4.21). Define $\nabla_\tau : C_0^n \otimes_{\mathbb{J}_*^n} \mathcal{F}_n \rightarrow C_0^n \otimes_{\mathbb{J}_*^n} \mathcal{F}_n$ by

$$\begin{aligned} \nabla_\tau(a_1 \otimes \cdots \otimes a_n \otimes f) &= a_1 \otimes \cdots \otimes a_n \otimes Df \\ &\quad + a_1 \otimes \cdots \otimes a_{n-1} \otimes (\omega_{(-1)} a_n) \otimes f \\ &\quad - \sum_{k \geq 1} a_1 \otimes \cdots \otimes a_{n-1} \otimes (\omega_{(2k-1)} a_n) \otimes g_{2k}(\tau) f \\ &\quad - \sum_{i=1}^{n-1} \sum_{m \geq 0} a_1 \otimes \cdots \otimes (\omega_{(m+1)} a_i) \otimes \cdots \otimes a_n \otimes \bar{\varphi}_{m+2}(t_n - t_i) f. \end{aligned} \quad (5.32)$$

where $Df = \left(2\pi i \frac{\partial}{\partial \tau} + \sum_{i=1}^{n-1} \zeta(t_n - t_i) \frac{\partial}{\partial t_i} \right) f$ is as defined in Lemma 4.14. Every function appearing on the right hand side of (5.32) is elliptic therefore $\nabla_\tau \in \text{End}(C_0^n \otimes_{\mathbb{J}_*^n} \mathcal{F}_n)$ is well defined.

In the algebro-geometric setting, the chiral homology acquires the structure of a twisted D -module, indeed a vector bundle with projectively flat connection, over the moduli space of curves with marked points. In our setting this connection is actually flat and it corresponds to the following

Lemma. *The operators ∇_τ and $L_{-1}^{(j)} + \frac{d}{dt_j}$ satisfy, for $1 \leq j \leq n-1$, the same commutation relations as in (4.22):*

$$\left[\left(L_{-1}^{(j)} + \frac{d}{dt_j} \right), \nabla_\tau \right] = \bar{\wp}_2(t_n - t_j) \left(L_{-1}^{(j)} + \frac{d}{dt_j} \right). \quad (5.33)$$

Proof. Follows easily from $\left[\frac{d}{dt_j}, D \right] = \bar{\wp}_2(t_n - t_j) \frac{d}{dt_j}$ and $[L_{-1}, L_m] = -(m+1)L_{m-1}$ for $m \geq 0$. \square

We let $H^{(0)}$ denote the endomorphism of C_0^n defined by

$$H^{(0)}(a_1 \otimes \cdots \otimes a_n \otimes f(t_1, \dots, t_n, \tau)) = d_1 \left(\omega \otimes a_1 \otimes \cdots \otimes a_n \otimes f(t_1, \dots, t_n, \tau) \zeta(t_n - t_0) \right), \quad (5.34)$$

here abusing notation since $f(t_1, \dots, t_n, \tau) \zeta(t_0 - t_n) \notin \mathcal{F}_{n+1}$ in general. A direct calculation shows that

$$\nabla_\tau = 2\pi i \frac{\partial}{\partial \tau} + \sum_{i=1}^{n-1} \zeta(t_n - t_i) \left[L_{-1}^{(i)} + \frac{\partial}{\partial t_i} \right] - H^{(0)}.$$

The non-ellipticity of ζ and of $\partial f / \partial \tau$ cancel due to (4.16). In the right hand side of (5.32) the terms ζ and $\partial f / \partial \tau$ are combined in the operator D .

5.19. Similarly we define ∇_τ on $C_1^n \otimes_{\mathbb{J}_*^n} \mathcal{F}_{n+1}$ by

$$\begin{aligned} \nabla_\tau(a_0 \otimes \cdots \otimes a_n \otimes f) &= a_0 \otimes \cdots \otimes a_n \otimes Df(t_0, \dots, t_n, \tau) \\ &\quad + a_0 \otimes \cdots \otimes \omega_{(-1)} a_n \otimes f - \sum_{k \geq 1} a_0 \otimes \cdots \otimes \omega_{(2k-1)} a_n \otimes g_{2k}(\tau) f \\ &\quad + a_0 \otimes \cdots \otimes a_n \otimes \bar{\wp}_2(t_n - t_0) f \\ &\quad - \sum_{i=0}^{n-1} \sum_{m \geq 0} a_0 \otimes a_1 \otimes \cdots \otimes \omega_{(m+1)} a_i \otimes \cdots \otimes a_n \otimes \bar{\wp}_{m+2}(t_n - t_i) f \\ &\quad - \sum_{i=1}^n \sum_k \omega \otimes a_1 \otimes \cdots \otimes a_{0(k)} a_i \otimes \cdots \otimes a_n \otimes f_{0ik}(t_1, \dots, t_n, \tau) \zeta(t_0 - t_n, \tau). \end{aligned} \quad (5.35)$$

When $a_0 \otimes \cdots \otimes a_n \otimes f$ is a cycle (that is if it lies in the kernel of d_1) then the last term of (5.35) vanishes. The remaining terms are manifestly elliptic so that the image of ∇_τ is indeed in $C_1^n \otimes_{\mathbb{J}_*^n} \mathcal{F}_{n+1}$. For elements of $\ker d_1$ we have, in the same way as in (5.33) the commutation relations (4.22):

$$\left[\left(L_{-1}^{(j)} + \frac{d}{dt_j} \right), \nabla_\tau \right] = \bar{\wp}_2(t_n - t_j) \left(L_{-1}^{(j)} + \frac{d}{dt_j} \right). \quad (5.36)$$

for $1 \leq j \leq n-1$. Meanwhile for $j=0$ we have

$$\left[\left(L_{-1}^{(0)} + \frac{d}{dt_0} \right), \nabla_\tau \right] = \left(L_{-1}^{(0)} + \frac{d}{dt_0} \right) \bar{\wp}_2(t_n - t_0). \quad (5.37)$$

It follows that ∇_τ descends to the quotient by elements of the form (5.25).

In general we can write

$$\nabla_\tau = 2\pi i \frac{\partial}{\partial \tau} + \sum_{i=0}^{n-1} \zeta(t_n - t_i) \left[L_{-1}^{(i)} + \frac{\partial}{\partial t_i} \right] - H^{(1)} + \bar{\wp}_2(t_n - t_0), \quad (5.38)$$

where $H^{(1)}$ is defined by

$$H^{(1)}(a_0 \otimes \cdots \otimes a_n \otimes f(t_0, \dots, t_n, \tau)) = d_2 \left(\omega \otimes a_0 \otimes \cdots \otimes a_n \otimes f(t_0, t_1, \dots, t_n, \tau) \zeta(t - t_n, \tau) \right), \quad (5.39)$$

where on the right hand side the function $f \cdot \zeta$ is considered as a function on the $n + 3$ variables $t, t_0, t_1, \dots, t_n, \tau$. Once again the non-ellipticity of $\partial f / \partial \tau$ is corrected by the terms containing ζ , making the elliptic function Df appear in the right hand side of (5.35).

Notice that by commuting $\zeta(t_n - t_0)$ with $L_{-1}^0 + \frac{d}{dt_0}$ in (5.38), it is seen that ∇_τ has the following simple form in the quotient C_1^n :

$$\nabla_\tau = 2\pi i \frac{\partial}{\partial \tau} + \sum_{i=1}^{n-1} \zeta(t_n - t_i) \left[L_{-1}^{(i)} + \frac{\partial}{\partial t_i} \right] - H^{(1)} \quad (5.40)$$

5.20. We record here some particular cases of the operators ∇_τ when $n = 1$. Equation (5.32) reduces to

$$\nabla_\tau(a \otimes f) = a \otimes \frac{d}{d\tau} f + \omega_{(-1)} a \otimes f - \sum_{k \geq 1} \omega_{(2k-1)} a \otimes g_{2k}(\tau) f. \quad (5.41)$$

Similarly, equation (5.35), applied to a cycle $a \otimes b \otimes f$ reduces to

$$\begin{aligned} \nabla_\tau(a \otimes b \otimes f(t_1 - t_0, \tau)) &= a \otimes b \otimes Df(t_1 - t_0, \tau) + a \otimes \omega_{(-1)} b \otimes f(t_1 - t_0, \tau) \\ &\quad - \sum_{k \geq 1} a \otimes \omega_{(2k-1)} b \otimes g_{2k}(\tau) f(t_1 - t_0, \tau) \\ &\quad + a \otimes b \otimes \bar{\varphi}_2(t_1 - t_0) f(t_1 - t_0, \tau) \\ &\quad - \sum_{m \geq 0} \omega_{(m+1)} a \otimes b \otimes \bar{\varphi}_{m+2}(t_1 - t_0) f(t_1 - t_0, \tau). \end{aligned} \quad (5.42)$$

5.21 Proposition. *The connection operator induces well defined endomorphisms of $H_k^{ch}(V^{\otimes n})$ for $k = 0, 1$.*

Proof. We want to show that ∇_τ preserves the spaces of cycles and boundaries. In degree 0 it is therefore enough to prove that ∇_τ preserves boundaries. It is instructive to do the case $n = 1$ first, the proof for general n follows the same pattern. So let $\alpha = a \otimes b \otimes f \in C_1^{n-1}$. We have

$$d_1 \alpha = a_{(f)} b = \sum_k a_{(k)} b \otimes f_k, \quad \text{where} \quad f = \sum_k f_k(t_0 - t_1)^k,$$

and

$$\nabla_\tau d_1 \alpha = \sum_k a_{(k)} b \otimes \frac{d}{d\tau} f_k + \sum_k \omega_{(-1)} a_{(k)} b \otimes f_k - \sum_{j \geq 1} \sum_k \omega_{(2j-1)} a_{(k)} b \otimes g_{2j}(\tau) f_k.$$

Using Borchers identity (2.2) we express the right hand side as

$$\sum_k a_{(k)} b \otimes \frac{d}{d\tau} f_k + \sum_k a_{(k)} \omega_{(\zeta)} b \otimes f_k + (L_{-1} a)_{(\zeta f)} b + (\omega_{(1)} a)_{(\bar{\varphi}_2 f)} b + \sum_{m \geq 2} (\omega_{(m)} a)_{(\varphi_{m+1} f)} b$$

The third term equals $a_{(\bar{\varphi}_2 f)} b - a_{(\zeta f')} b$ by (2.6). Collecting we obtain

$$\nabla_\tau d_1 \alpha = d_1 \left(a \otimes b \otimes Df + a \otimes \omega_{(\zeta)} b \otimes f + a \otimes b \otimes \bar{\varphi}_2 f + \sum_{m \geq 1} \omega_{(m)} a \otimes b \otimes \bar{\varphi}_{m+1} f \right). \quad (5.43)$$

Therefore ∇_τ preserves homology in degree 0 in the case $n = 1$. The general case is proved in the same fashion by an application of Borchers identity. Let $\alpha = a \otimes a_1 \cdots \otimes a_n \otimes f(t_0, \dots, t_n) \in C_1^n$. We have

$$d_1 \alpha = \sum_{i=1}^n \sum_k a_1 \otimes \cdots \otimes a_{(k)} a_i \otimes \cdots \otimes a_n \otimes f_{0ik}(t_1, \dots, t_n).$$

We compute therefore

$$\begin{aligned}
\nabla_\tau d_1 \alpha &= \sum_{i=1}^n \sum_k a_1 \otimes \cdots \otimes a_{(k)} a_i \otimes \cdots \otimes a_n \otimes Df_{0ik} + \sum_{i=1}^{n-1} \sum_k a_1 \otimes \cdots \otimes a_{(k)} a_i \otimes \cdots \otimes \omega_{(\zeta)} a_n \otimes f_{0ik} \\
&\quad + \sum_k a_1 \otimes \cdots \otimes a_{n-1} \otimes \omega_{(\zeta)} a_{(k)} a_n \otimes f_{0nk} \\
&\quad - \sum_{i=1}^n \sum_k \sum_{j=1}^{i-1} \sum_{m \geq 1} a_1 \otimes \cdots \otimes \omega_{(m)} a_j \otimes \cdots \otimes a_{(k)} a_i \otimes \cdots \otimes a_n \otimes \bar{\varphi}_{m+1}(t_n - t_j) f_{0ik} \\
&\quad - \sum_{i=1}^{n-1} \sum_k \sum_{m \geq 1} a_1 \otimes \cdots \otimes \omega_{(m)} a_{(k)} a_i \otimes \cdots \otimes a_n \otimes \bar{\varphi}_{m+1}(t_n - t_i) f_{0ik} \\
&\quad - \sum_{i=1}^{n-2} \sum_k \sum_{j=i+1}^{n-1} \sum_{m \geq 1} a_1 \otimes \cdots \otimes a_{(k)} a_i \otimes \cdots \otimes \omega_{(m)} a_j \otimes \cdots \otimes a_n \otimes \bar{\varphi}_{m+1}(t_n - t_j) f_{0ik}.
\end{aligned}$$

We add and subtract the terms with $m = 0$ as in the last three sums and use Borchers identity in the third and fifth summand on the right hand side to obtain

$$\begin{aligned}
\nabla_\tau d_1 \alpha &= d_1 \left(a \otimes a_1 \otimes \cdots \otimes a_n \otimes Df + a \otimes a_1 \otimes \cdots \otimes a_{n-1} \otimes \omega_{(\zeta)} a_n \otimes f \right. \\
&\quad \left. - \sum_{j=1}^{n-1} \sum_{m \geq 1} a \otimes a_1 \otimes \cdots \otimes \omega_{(m)} a_j \otimes \cdots \otimes a_n \otimes \bar{\varphi}_{m+1}(t_n - t_j) f \right. \\
&\quad \left. + \sum_{m \geq 1} \omega_{(m)} a \otimes a_1 \otimes \cdots \otimes a_n \otimes \bar{\varphi}_{m+1}(t_n - t_0) f \right).
\end{aligned}$$

This proves that ∇_τ preserves homology in degree 0. The degree 1 case is proved in the same way. \square

5.22 Remark. In order to define ∇_τ , it is enough to extend scalars by adding $g_2(\tau)$ instead of all holomorphic functions, ie. the connection ∇_τ is well defined on $C_\bullet^n \otimes_{\mathbb{M}_*} \mathbb{Q}\mathbb{M}_* \subset C_\bullet^n \otimes_{\mathbb{J}_*^n} \mathcal{F}_*$. We can therefore consider the complexes

$$C_2^n \otimes_{\mathbb{M}_*} \mathbb{Q}\mathbb{M}_* \xrightarrow{d_2} C_1^n \otimes_{\mathbb{M}} \mathbb{Q}\mathbb{M}_* \xrightarrow{d_1} C_0^n \otimes_{\mathbb{M}} \mathbb{Q}\mathbb{M}_* \rightarrow 0, \quad (5.44)$$

and the ∇_τ - and $k[\partial_1, \dots, \partial_n]$ -coinvariants of the corresponding homology. We will denote these coinvariants by $H_i^{\text{ch}}(V^{\otimes n})_\nabla$.

Now let a_0, \dots, a_n be of homogeneous conformal weight and let $f \in \mathbb{J}_k^n$. Let $\alpha = a_0 \otimes \cdots \otimes a_n \otimes f \in C_1^n$ and put $\deg \alpha = k + \sum_{i=0}^n \deg a_i$. This defines a grading on C_1^n that we may call the *conformal grading*. We let $(C_1^n)_p$ consist of elements of conformal degree p . Suppose $d_1 \alpha = 0$. Then every element in (5.35) lies in $(C_1^n)_{\deg \alpha + 2}$ except

$$a_0 \otimes \cdots \otimes a_n \otimes \left(Df - \sum_{i=0}^n \deg a_i g_2(\tau) f \right).$$

Let $g(t_0, \dots, t_n, \tau) = D(f) - \sum_{i=0}^n \deg a_i g_2(\tau) f$. It follows from (4.17) and (4.2) that g is not modular but rather satisfies

$$g\left(\frac{t_0}{c\tau + d}, \dots, \frac{t_n}{c\tau + d}, \frac{a\tau + b}{c\tau + d}\right) = (c\tau + d)^{k+2} g(t_0, \dots, t_n, \tau) + 2\pi ic \deg \alpha (c\tau + d)^{k+1} f(t_0, \dots, t_n, \tau). \quad (5.45)$$

5.23 Specializing the curve. We can restrict the complex C_\bullet^n (or rather its extension of scalars to \mathcal{F}_n) by fixing the points $t_1, \dots, t_n \in \mathbb{C}$ with $t_i \neq t_j$ for $i \neq j$ and $\tau \in \mathbb{H}$. The homology of the resulting complex is denoted by $H_i^{\text{ch}}(V^{\otimes n}, t_1, \dots, t_n, \tau)$ and it corresponds to the chiral homology of the elliptic curve $E_\tau = \mathbb{C}/\mathbb{Z} + \tau\mathbb{Z}$ with insertions of the vacuum module V at the points t_1, \dots, t_n , as defined in [1, §4.2.19]. Explicitly, C_0^n specializes to $V^{\otimes n}$, C_1^n specializes to the quotient of $V^{\otimes n+1} \otimes \Gamma_1$, where Γ_1 consists of meromorphic functions $f(t_0)$, biperiodic with periods 1 and τ , with possible poles at $t_0 = t_i + \mathbb{Z} + \tau\mathbb{Z}$, $i = 1, \dots, n$, by the relation (5.25). An similarly C_2^n specializes to the quotient

of $V^{\otimes n+2} \otimes \Gamma_2$, where Γ_2 consists of meromorphic functions $f(u, v)$, bi-periodic, with possible poles at $u, v = t_i + \mathbb{Z} + \tau\mathbb{Z}$, $i = 1, \dots, n$, and $u = v + \mathbb{Z} + \tau\mathbb{Z}$, by elements of the form (5.26).

The chiral homology $H_i^{\text{ch}}(V^{\otimes n})$ induces a vector bundle with a flat connection over the moduli space of n -marked elliptic curves, its fiber over the curve E_τ with markings t_1, \dots, t_n is $H_i^{\text{ch}}(V^{\otimes n}, t_1, \dots, t_n, \tau)$. We remark that in the general case the connection is not flat but rather projectively flat, but in the genus 1 case (where there is only one modular parameter) this connection is actually flat as we saw in (5.33) and (5.36).

In the case $n = 1$, as we saw in 5.16 we have an isomorphism $H_i^{\text{ch}}(V) \simeq H_i^{\text{ch}}(V)_{\text{tr}}$ therefore $H_i^{\text{ch}}(V, t_1, \tau)$ does not depend on t_1 , we denote this homology by $H_i^{\text{ch}}(E_\tau, V)$.

5.24. We will be interested in the dual vector spaces $H_\bullet^{\text{ch}}(V^{\otimes n})^*$ and their flat sections as we vary the points and the modular parameter. Recall that we have fixed a \mathbb{J}_*^n -module structure on \mathbb{J}_*^m for all $m \geq n$ in 4.6 and that the complexes C_\bullet^n are complexes of \mathbb{J}_*^n -modules. Let $Z_i^n \subset C_i^n$ consist of cycles, that is $Z_i^n = \ker d_i$. We will consider linear functionals

$$S_i^n \in \text{Hom}_{\mathbb{J}_*^n}(Z_i^n, \mathcal{F}_n), \quad S_i^n(d_{i+1}\alpha) = 0, \quad \forall \alpha \in C_{i+1}^n. \quad (5.46)$$

Thus, S_i^n gives a vector in the dual to chiral homology $H_i^{\text{ch}}(V^{\otimes n})^*$.

We remark that these are not cohomology classes of the dual complex of C_\bullet^n as these linear functionals are only defined on cycles and not on all chains.

5.25 Moving the points. Recall that the complex C_\bullet^n is a complex of $k[\partial_1, \dots, \partial_n]$ -modules. \mathcal{F}_n is a $k[\partial_1, \dots, \partial_n]$ -module where ∂_i acts as $\frac{\partial}{\partial t_i}$. Hence $\text{Hom}_{\mathbb{J}_*^n}(Z_i^n, \mathcal{F}_n)$ is a $k[\partial_1, \dots, \partial_n]$ -module where the action of ∂_j is given by

$$\left(\partial_j S_i^n\right)(\alpha) = \frac{\partial}{\partial t_j} S_i^n(\alpha) - S_i^n\left(\left[L_{-1}^{(j)} + \frac{\partial}{\partial t_j}\right]\alpha\right), \quad S_i^n \in \text{Hom}_{\mathbb{J}_*^n}(Z_i^n, \mathcal{F}_n), \quad \alpha \in Z_i^n. \quad (5.47)$$

We denote by $H_i^{\text{ch}}(V^{\otimes n})^{*\text{tr}}$ the invariants of this action, namely the subspace consisting on classes S_i^n such that the right hand side of (5.47) vanishes for every $1 \leq j \leq n$.

5.26 Definition. A flat section of the dual of chiral homology, of degree $i = 0, 1$ is the cohomology class of a linear functional $S_i^n \in H_i^{\text{ch}}(V^{\otimes n})^{*\text{tr}}$ which in addition is flat with respect to the dual connection ∇_τ , that is

$$D_\tau S_i^n(\alpha) = S_i^n(\nabla_\tau \alpha), \quad \text{for all } \alpha \in Z_i^n, \quad (5.48)$$

where D_τ is the derivation defined in (4.21). We will denote by $H_i^{\text{ch}}(V^{\otimes n})^{*\nabla}$ the group of such flat sections.

5.27 Remark. In degree 0 the space of flat sections of the dual of chiral homology agrees with the space of conformal blocks of V in genus 1 as defined by Zhu [2]. It is clear that we also have $H_0^{\text{ch}}(V^{\otimes n})^{*\nabla} = (H_0^{\text{ch}}(V^{\otimes n})_{\nabla})^*$.

6 Degree 1 conformal blocks on the torus

In this section we define the degree 1 analog of the notion of conformal block as in [2]. It consists of a compatible system of flat sections of the dual of the first chiral homology with supports on n points, for every n . We prove the modular invariance of the degree 1 conformal block in 6.7.

6.1 Definition. Let $(V, \mathbf{1}, \omega, Y(\cdot, z))$ be a vertex algebra. For each $n \geq 1$, consider the complex C_\bullet^n computing the chiral homology with coefficients in $V^{\otimes n}$. We let $Z_1^n \subset C_1^n$ be the set of cycles, that is the kernel of d_1 . A system of maps $S_1 = \{S_1^n\}_{n \geq 1}$

$$S_1^n : Z_1^n \rightarrow \mathcal{F}_n, \quad (a_0, \dots, a_n, f) \mapsto S_1^n(a_0, \dots, a_n, f; t_1, \dots, t_n, \tau),$$

is said to satisfy the *genus-one property* if the following properties are satisfied.

- a) S_1^n defines a flat section of the dual of the first chiral homology in the torus: $S_1^n \in H_1^{\text{ch}}(V^{\otimes n})^{*\nabla}$.

b) For $a_1, \dots, a_n \in V$ and $f \in \mathbb{J}_*^{n+1}$:

$$S_1^n(\mathbf{1}, a_1, \dots, a_n, f; t_1, \dots, t_n, \tau) = 0 \quad (6.1)$$

c) Let $\alpha = a_0 \otimes \dots \otimes a_n \otimes f \in Z_1^n$ and let $\sigma \in S_n$. Recall the action of the symmetric group on C_\bullet^n defined in 5.13. We have

$$S_1^n(\sigma(\alpha); t_{\sigma(1)}, \dots, t_{\sigma(n)}) = S_1^n(\alpha; t_1, \dots, t_n). \quad (6.2)$$

d) Let $b, a_0, \dots, a_n \in V$ and $f = f(u, v, t_1, \dots, t_n, \tau) \in \mathbb{J}_*^{n+2}$. We can view

$$v := a_0 \otimes b \otimes a_1 \otimes \dots \otimes a_n \otimes f \in C_1^{n+1}. \quad (6.3)$$

Then if v is a cycle, that is $d_1 v = 0$, using the Laurent expansions of f as in (5.17). We have

$$\begin{aligned} S_1^{n+1}(a_0, b, a_1, \dots, a_n, f; v, t_1, \dots, t_n, \tau) = & \\ & \sum_{i=1}^n \sum_k S_1^n(a_0, a_1, \dots, b_{(k)} a_i, \dots, a_n, f_{1ik}; t_1, \dots, t_n, \tau) \zeta(v - t_i) \\ & + \sum_{i=1}^n \sum_k S_1^n(a_0, a_1, \dots, b_{(k)} a_i, \dots, a_n, f_{1ik-1}; t_1, \dots, t_n, \tau) (\wp_2(v - t_i) + g_2(\tau)) \\ & + \sum_{i=1}^n \sum_k \sum_{m \geq 0} S_1^n(a_0, a_1, \dots, b_{(k)} a_i, \dots, a_n, f_{1ik-m-2}; t_1, \dots, t_n, \tau) \wp_{m+3}(v - t_i) \\ & - \sum_{i=1}^{n-1} \sum_k S_1^n(a_0, a_1, \dots, b_{(k)} a_i, \dots, a_n, f_{1ik}; t_1, \dots, t_n, \tau) \zeta(t_n - t_i) \\ & - \sum_{i=1}^{n-1} \sum_k S_1^n(a_0, a_1, \dots, b_{(k)} a_i, \dots, a_n, f_{1ik-1}; t_1, \dots, t_n, \tau) (\wp_2(t_n - t_i) + g_2(\tau)) \\ & - \sum_{i=1}^{n-1} \sum_k \sum_{m \geq 0} S_1^n(a_0, a_1, \dots, b_{(k)} a_i, \dots, a_n, f_{1ik-2}; t_1, \dots, t_n, \tau) \wp_{m+3}(t_n - t_i) \\ & + \sum_k S_1^n(a_0, a_1, \dots, b_{(k)} a_n, f_{1nk+1}; t_1, \dots, t_n, \tau) \\ & - \sum_k \sum_{j \geq 1} S_1^n(a_0, a_1, \dots, b_{(k)} a_n, f_{1n,k+1-2j}; t_1, \dots, t_n, \tau) g_{2j}(\tau). \quad (6.4) \end{aligned}$$

6.2. We prove now that the right hand side of (6.4) is well defined. Consider

$$\alpha_s = a_0 \otimes b \otimes a_1 \otimes \dots \otimes a_n \otimes g_s(u, v, t_1, \dots, t_n),$$

where $g_s(u, v, t_1, \dots, t_n) = f(u, v, t_1, \dots, t_n) (\zeta(v - t_n) + \zeta(s - v))$ is a family of meromorphic functions parametrized by s . The functions g_s are not in \mathcal{F}_{n+2} since they have poles at $v = s$ in addition to the diagonal divisors, and are not elliptic with respect to the variable t_n . As such α cannot be viewed as an s -family of elements in C_2^n . However, we may formally define $\beta_s = d_2 \alpha_s$ using (5.18). It follows in the same way as in Lemma 5.9 that $d_1 \beta_s = 0$. Note however that β_s is not a cycle in C_1^n since the functions involved have poles at $t_i = s$ for $i = 1, \dots, n$ in addition to the diagonal divisor (there is no pole at $t_0 = s$ due to our assumption that v defined by (6.3) is a cycle). Equation (6.4) is equivalent to

$$S_1^{n+1}(a_0, b, a_1, \dots, a_n, f; v, t_1, \dots, t_n, \tau) = S_1^n(\beta_s; t_1, \dots, t_n, \tau)|_{s=v}. \quad (6.5)$$

However, the s -dependent part of β_s can be written as a linear combination of elements of the form $\gamma \otimes h \cdot \wp_k(t_i - s)$ where $\gamma \in V^{\otimes(n+1)}$, $h \in \mathcal{F}_{n+1}$, $k \geq 1$ and $1 \leq i \leq n$. It follows by the \mathcal{F}_n -linearity

of the differential that each coefficient of $\wp_k(t_i - s)$ is a cycle in C_1^n and therefore in the domain of S_1^n . Thus, the right hand side of (6.5) is well defined.

We now prove that the right hand side of (6.4) is an elliptic function in \mathcal{F}_{n+1} . Indeed every term except the first and the fourth are manifestly elliptic. Notice that the sum of these two terms is also elliptic for the variables t_i , $1 \leq i \leq n-1$. Denote by $S = S(v, t_1, \dots, t_n)$ the right hand side of (6.4). It follows that

$$\begin{aligned} S(v, t_1, \dots, t_n + \tau) - S(v, t_1, \dots, t_n) &= -S(v + \tau, t_1, \dots, t_n) + S(v, t_1, \dots, t_n) = \\ &= 2\pi i \sum_{i=1}^n \sum_k S_1^n \left(a_0, a_1, \dots, b_{(k)} a_i, \dots, a_n, f_{1ik}(t_0, \dots, t_n, \tau) \right). \end{aligned} \quad (6.6)$$

Consider the vector v defined by (6.3) but viewed as a vector in C_2^n . We compute

$$d_2 v = \sum_{i=1}^n \sum_k a_0 \otimes \dots \otimes b_{(k)} a_i \otimes \dots \otimes a_n \otimes f_{1ik}(t_0, \dots, t_n, \tau),$$

where we used that $d_1 v = 0$ when viewed as an element of C_1^{n+1} . Thus, the right hand side of (6.6) equals $2\pi i S_1^n(d_2 v) = 0$.

6.3. Given such a system, for each $a_0, \dots, a_n \in V$ and $f \in \mathbb{J}_*^{n+1}$, and a fixed $\tau \in \mathbb{H}$, we may view

$$S_1^n(a_0, \dots, a_n, f; t_1, \dots, t_n, \tau),$$

as an n -variable function on the complex torus $\mathbb{C}/\mathbb{Z} + \mathbb{Z}\tau$. These are the higher analogs of the *correlation functions* on the torus defined by Zhu in [2] in degree 0.

By definition it is clear that if $\{S_1^n\}$ and $\{S_1^m\}$ satisfy the genus-one property then so does $\{\alpha S_1^n + S_1^m\}$ for any $\alpha \in \mathbb{C}$.

Definition. The linear space of families $S_1 = \{S_1^n\}$ satisfying the genus one property is called the space of *degree 1 conformal blocks on the torus*.

6.4 Remark. Notice that all the terms in the right hand side of (6.4) involving $g_2(\tau)$ cancel each-other and can be therefore omitted from the identity.

6.5. Let $f(t_0, t, t_1, \dots, t_n, \tau) \in \mathbb{J}_*^{n+2}$ and consider its Laurent expansion near $t - t_1$:

$$f = \sum_k f_k(t_0, t_1, \dots, t_n, \tau)(t - t_1)^k.$$

It follows from (6.4) that the following holds

$$\int_C S_1^{n+1}(a_0, a, a_1, \dots, a_n, f; t, t_1, \dots, t_n, \tau) dt = \sum_k S_1^n(a_0, a_{(k)} a_1, a_2, \dots, a_n, f_k; t_1, \dots, t_n, \tau), \quad (6.7)$$

where C is a small contour around t_1 , not including any of the points t_i , $i \neq 1$. This is the degree 1 analog of [2, (4.1.4)]. To prove (6.7) notice that from the right hand side of (6.4) only the $i = 1$ summand of the first term has a non-trivial residue at $s = t_1$.

6.6. Consider (6.4) with $b = \omega$ and $f = f(t_0, \dots, t_n, \tau) \in \mathbb{J}_*^{n+1}$, that is, independent of s . It follows from (5.48) that

$$\begin{aligned} S_1^{n+1}(a_0, \omega, a_1, \dots, a_n, f; t_1, \dots, t_n) &= \left(2\pi i \frac{d}{d\tau} - \sum_{i=1}^{n-1} \zeta(s - t_i) \frac{d}{dt_i} \right) S_1^n(a_0, \dots, a_n, f; t_1, \dots, t_n, \tau) \\ &\quad - S_1^n \left(a_0, \dots, a_n, \left(2\pi i \frac{d}{d\tau} - \sum_{i=0}^{n-1} \zeta(s - t_i) \frac{d}{dt_i} \right) f; t_1, \dots, t_n, \tau \right) \\ &\quad - S_1^n \left(a_0, \dots, a_n, (\wp_2(s - t_0) + g_2(\tau)f); t_1, \dots, t_n, \tau \right). \end{aligned} \quad (6.8)$$

This is a degree 1 analog of [2, (4.1.5)].

6.7 Theorem. Let $\{S_1^n\}$ satisfy the genus-one property and let

$$\sigma = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in SL(2, \mathbb{Z}).$$

Define $\{\sigma(S_1^n)\}$ as follows. For $a_i \in V$, $i = 0, \dots, n$ of conformal weight Δ_i and $f \in \mathbb{J}_k^{n+1}$. Put

$$\sigma(S_1^n)(a_0, \dots, a_n, f; t_1, \dots, t_n, \tau) = (c\tau + d)^{-k - \sum_{i=0}^n \Delta_i} S_1^n \left(a_0, \dots, a_n, f; \frac{t_1}{c\tau + d}, \dots, \frac{t_n}{c\tau + d}, \frac{a\tau + b}{c\tau + d} \right). \quad (6.9)$$

Then $\{\sigma(S_1^n)\}$ satisfies the genus-one property.

Proof. It is clear that $\sigma(S_1^n)$ satisfies (6.2). Let now $\alpha = a_0 \otimes \dots \otimes a_n \otimes f \in Z_1^n$ with $f \in \mathbb{J}_k^{n+1}$ and a_0, \dots, a_n of homogeneous conformal weight. We put $\deg \alpha = k + \sum_{i=0}^n \deg a_i$.

$$\begin{aligned} \sigma(S_1^n)(L_{-1}a_0, \dots, a_n, f; t_1, \dots, t_n, \tau) &= (c\tau + d)^{-\deg \alpha - 1} S_1^n \left(L_{-1}a_0, \dots, a_n, f; \frac{t_1}{c\tau + d}, \dots, \frac{t_n}{c\tau + d}, \frac{a\tau + b}{c\tau + d} \right) \\ &= -(c\tau + d)^{-\deg \alpha - 1} S_1^n \left(a_0, \dots, a_n, \frac{\partial}{\partial t_0} f; \frac{t_1}{c\tau + d}, \dots, \frac{t_n}{c\tau + d}, \frac{a\tau + b}{c\tau + d} \right) \\ &= -\sigma(S_1^n) \left(a_0, \dots, a_n, \frac{\partial}{\partial t_0} f; t_1, \dots, t_n, \tau \right), \end{aligned}$$

so that $\sigma(S_1^n)$ vanishes on elements as in (5.25) and it is therefore well defined on Z_1^n .

Next, we check that $\sigma(S_1^n)$ vanishes on coboundaries. Let $a, b, a_1, \dots, a_n \in V$ be homogeneous elements, and $f = f(u, v, t_1, \dots, t_n, \tau) \in \mathbb{J}_p^{n+2}$. Let $\alpha = a \otimes b \otimes a_1 \otimes \dots \otimes a_n \otimes f \in C_2^n$. We have

$$\begin{aligned} \sigma(S_1^n)(d_2\alpha; t_1, \dots, t_n, \tau) &= \sum_{i=1}^n \sum_k \sigma(S_1^n)(a, a_1, \dots, b_{(k)}a_i, \dots, a_n, f_{1ik}; t_1, \dots, t_n, \tau) \\ &\quad - \sum_{i=1}^n \sum_k \sigma(S_1^n)(b, a_1, \dots, a_{(k)}a_i, \dots, a_n, f_{0ik}; t_1, \dots, t_n, \tau) \\ &\quad - \sum_k \sigma(S_1^n)(a_{(k)}b, a_1, \dots, a_n, f_{00k}; t_1, \dots, t_n, \tau). \end{aligned} \quad (6.10)$$

Noting that $f_{1ik} \in \mathbb{J}_{p+k}^{n+1}$ and $\deg b_{(k)}a_i = \deg b + \deg a_i - k - 1$ we have that the first term on the right hand side of (6.10) equals

$$(c\tau + d)^{1-p-\deg b-\deg a-\sum_{i=1}^n \deg a_i} S_1^n \left(a, a_1, \dots, b_{(k)}a_i, \dots, a_n, f_{1ik}; \frac{t_1}{c\tau + d}, \dots, \frac{t_n}{c\tau + d}, \frac{a\tau + b}{c\tau + d} \right).$$

The other two terms are treated similarly. It follows that

$$\sigma(S_1^n)(d\alpha; t_1, \dots, t_n, \tau) = (c\tau + d)^{1-p-\deg b-\deg a-\sum_{i=1}^n \deg a_i} S_1^n \left(d_2\alpha; \frac{t_1}{c\tau + d}, \dots, \frac{t_n}{c\tau + d}, \frac{a\tau + b}{c\tau + d} \right) = 0,$$

therefore $\sigma(S_1^n)$ defines a vector in $H_1^{\text{ch}}(V^{\otimes n}, t_1, \dots, t_n)^*$.

We now check that $\sigma(S_1^n)$ vanishes on elements as in (5.28). We have for $\alpha = a_0 \otimes \dots \otimes a_n \otimes f \in Z_1^n$ with $f \in \mathbb{J}_k^{n+1}$

$$\sigma(S_1^n)(a_0, \dots, L_{-1}a_i, \dots, a_n, f; t_1, \dots, t_n, \tau)$$

$$\begin{aligned}
&= (c\tau + d)^{-\deg \alpha - 1} S_1^n \left(a_0, \dots, L_{-1} a_i, \dots, a_n, f; \frac{t_1}{c\tau + d}, \dots, \frac{t_n}{c\tau + d}, \frac{a\tau + b}{c\tau + d} \right) \\
&= (c\tau + d)^{-\deg \alpha - 1} \frac{\partial}{\partial t_i} S_1^n \left(a_0, \dots, a_n, f; \frac{t_1}{c\tau + d}, \dots, \frac{t_n}{c\tau + d}, \frac{a\tau + b}{c\tau + d} \right) \\
&\quad - (c\tau + d)^{-\deg \alpha - 1} S_1^n \left(a_0, \dots, a_n, \frac{\partial}{\partial t_i} f; \frac{t_1}{c\tau + d}, \dots, \frac{t_n}{c\tau + d}, \frac{a\tau + b}{c\tau + d} \right) \\
&= \frac{\partial}{\partial t_i} \sigma(S)_1^n (a_0, \dots, a_n, f; t_1, \dots, t_n, \tau) \\
&\quad - \sigma(S)_1^n \left(a_0, \dots, a_n, \frac{\partial}{\partial t_i} f; t_1, \dots, t_n, \tau \right).
\end{aligned}$$

So that $\sigma(S)_1^n$ gives a well defined vector on $H_1^{\text{ch}}(V^{\otimes n})^{*\text{tr}}$.

We now check that $\sigma(S)_1^n$ satisfies (5.48). Let $a_0, \dots, a_n \in V$ be homogeneous elements and $f = f(t_0, \dots, t_n, \tau) \in \mathbb{J}_k^{n+1}$. Put $\alpha = a_0 \otimes \dots \otimes a_n \otimes f \in Z_1^n$. We have

$$\begin{aligned}
\frac{d}{d\tau} \sigma(S)_1^n(\alpha; t_1, \dots, t_n, \tau) &= \frac{d}{d\tau} (c\tau + d)^{-\deg \alpha} S_1^n \left(\alpha; \frac{t_1}{c\tau + d}, \dots, \frac{t_n}{c\tau + d}, \frac{a\tau + b}{c\tau + d} \right) \\
&= -c \deg \alpha (c\tau + d)^{-1 - \deg \alpha} S_1^n \left(\alpha; \frac{t_1}{c\tau + d}, \dots, \frac{t_n}{c\tau + d}, \frac{a\tau + b}{c\tau + d} \right) \\
&\quad - c (c\tau + d)^{-2 - \deg \alpha} \sum_{i=1}^n \left(\frac{\partial}{\partial t_i} S_1^n \right) \left(\alpha; \frac{t_1}{c\tau + d}, \dots, \frac{t_n}{c\tau + d}, \frac{a\tau + b}{c\tau + d} \right) \\
&\quad + (c\tau + d)^{-2 - \deg \alpha} \left(\frac{\partial}{\partial \tau} S_1^n \right) \left(\alpha; \frac{t_1}{c\tau + d}, \dots, \frac{t_n}{c\tau + d}, \frac{a\tau + b}{c\tau + d} \right).
\end{aligned} \tag{6.11}$$

The second term on the right hand side vanishes because of Lemma 4.9. The first term equals

$$-\frac{c \deg \alpha}{c\tau + d} \sigma(S)_1^n(\alpha; t_1, \dots, t_n, \tau). \tag{6.12}$$

Finally using that S_1^n satisfies (5.48), the third term equals

$$\frac{1}{2\pi i} (c\tau + d)^{-2 - \deg \alpha} S_1^n \left(\nabla_\tau \alpha; \frac{t_1}{c\tau + d}, \dots, \frac{t_n}{c\tau + d}, \frac{a\tau + b}{c\tau + d} \right),$$

It follows from Remark 5.22 and (5.45) that this term equals

$$\frac{1}{2\pi i} \sigma(S)_1^n(\nabla_\tau \alpha; t_1, \dots, t_n, \tau) + \frac{c \deg \alpha}{c\tau + d} \sigma(S)_1^n(\alpha; t_1, \dots, t_n, \tau).$$

The second term cancels (6.12) and the first term together with the left hand side of (6.11) show that $\sigma(S)_1^n$ is a flat section in $H_1^{\text{ch}}(V^{\otimes n})^{*\nabla}$

It is only left to prove that $\sigma(S)_1^n$ satisfies the insertion formula (6.4). We let a_0, \dots, a_n, b be homogeneous elements in V and $f = f(u, v, t_1, \dots, t_n, \tau) \in \mathbb{J}_p^{n+2}$. We define v as in (6.3) and assume it is a cycle, that is $d_1 v = 0$. We want to evaluate

$$\begin{aligned}
&\sigma(S)_1^{n+1}(a_0, b, a_1, \dots, a_n, f; v, t_1, \dots, t_n, \tau) = \\
&\quad (c\tau + d)^{-k - \deg b - \sum_{i=0}^n \deg a_i} S_1^{n+1} \left(a_0, b, a_1, \dots, a_n, f; \frac{v}{c\tau + d}, \frac{t_1}{c\tau + d}, \dots, \frac{t_n}{c\tau + d}, \frac{a\tau + b}{c\tau + d} \right).
\end{aligned}$$

We now notice that in the right hand side of (6.4) function appearing transforms manifestly as a Jacobi form except the first and the fourth terms involving a ζ function. The sum of these two terms is however manifestly modular and the Theorem follows by simply applying (6.4) for S_1^n and the definition of $\sigma(S)_1^n$. \square

7 Modified Vertex Operators

In this section we prove a version of Borcherds identity adapted to elliptic curves via the exponential map. This identity (7.3) involves the two different vertex algebra structures $Y(\cdot, z)$ and $Y[\cdot, z]$ on V , as well as the Laurent and Fourier expansions of an elliptic function in the coordinates t and $z = e^{2\pi it}$.

7.1. We introduce the following modified vertex operators

$$X(a, z) = z^{-1}Y(z^{L_0}a, z) = \sum_{n \in \mathbb{Z}} a_n z^{-n-1}.$$

7.2 Lemma. *The modified vertex operators $(X(\cdot, z))$ satisfy the following identities:*

$$X((L_{-1} + L_0)a, z) = z \frac{d}{dz} X(a, z) + X(a, z), \quad (7.1)$$

and

$$q^{L_0} X(a, z) = qX(a, qz)q^{L_0}. \quad (7.2)$$

7.3 Theorem. *Let $f(t)$ be a meromorphic function of a complex variable t which is periodic, i.e., $f(t+1) = f(t)$. Let $\tau \in \mathbb{H}$ and suppose f is analytic in the domain $-\Im\tau < \Im t < \Im\tau$ except possibly at $t = 0$ where it may have a pole. Put $q = e^{2\pi i\tau}$, $z = e^{2\pi it}$ and let $F_+(z)$ (resp. $F_-(z)$) be the Fourier expansion of f as in 4.12, in the domain $|q| < |z| < 1$ (resp. $1 < |z| < |q|^{-1}$). If either*

- $f(t)$ is regular at $t = 0$, or
- f satisfies $f(t + \tau) = f(t) - \alpha$ for all $t \in \mathbb{C}$ for some constant $\alpha \in \mathbb{C}$,

then the following identity holds

$$2\pi i X(a_{[f]}b, w) = \operatorname{res}_z F_- \left(\frac{z}{w} \right) X(a, z)X(b, w) - \operatorname{res}_z F_+ \left(\frac{z}{w} \right) X(b, w)X(a, z), \quad (7.3)$$

as an equality in $\operatorname{Hom}(V, V((w)))$.

Notice that two different vertex algebra structures of V are in play here: the usual vertex algebra structure in the form of $X(\cdot, z)$, and also the $Y[\cdot, z]$ vertex algebra structure of 2.8 is used in the left hand side of (7.3).

Proof. It is enough to prove the statement for $a, b \in V$ of homogeneous conformal weight. We assume this throughout. We also fix $\tau \in \mathbb{H}$ and write $q = e^{2\pi i\tau}$. Our hypotheses on $f(t)$ imply that it admits a Laurent expansion near $t = 0$, namely

$$f(t) = \sum_{n \geq N} f_n t^n,$$

so the left hand side of (7.3) becomes

$$X(\operatorname{res}_t f(t)Y[a, t]b dt, w) = \sum_{n \geq N} f_n X(a_{[n]}b, w) \in \operatorname{Hom}(V, V((w))). \quad (7.4)$$

Since $f(t+1) = f(t)$, the function f has Fourier series expansions

$$F_{\pm}(z) = \sum_{n \in \mathbb{Z}} F_n^{\pm} z^n, \quad \text{where } F_n^{\pm} = \int_{t_0}^{t_0+1} f(t) e^{-2\pi i n t} dt.$$

Here t_0 is chosen so that $0 < \pm \Im t_0 < \Im\tau$ and the integral is taken along the path, parallel to the real axis in the t -plane, connecting t_0 to $t_0 + 1$.

The series $F_+(z)$ converges uniformly on closed subsets of the domain $|q| < |z| < 1$ and the series $F_-(z)$ converges uniformly on closed subsets of the domain $1 < |z| < |q|^{-1}$. In their corresponding domains of convergence we have

$$F_{\pm}(e^{2\pi it}) = f(t).$$

If $f(t)$ is regular at $t = 0$ then $F_n^- - F_n^+ = 0$ for all $n \in \mathbb{Z}$ and if $f(t + \tau) = f(t) - \alpha$ then

$$F_n^- - F_n^+ = \alpha \quad \text{for all } n \in \mathbb{Z}. \quad (7.5)$$

To simplify the presentation we agree to put $\alpha = 0$ in case $f(t)$ is regular at $t = 0$, so that (7.5) holds in all cases.

$$\begin{aligned} \operatorname{res}_t f(t) Y[a, t] b dt &= \operatorname{res}_t f(t) e^{2\pi i \deg(a)t} Y(a, e^{2\pi it} - 1) b dt \\ &= \sum_{n \in \mathbb{Z}} a_{(n)} b \operatorname{res}_t f(t) e^{2\pi i \deg(a)t} (e^{2\pi it} - 1)^{-n-1} dt. \end{aligned} \quad (7.6)$$

Since $f(t)$ is analytic with possible pole at $t = 0$, we have

$$\operatorname{res}_t f(t) e^{2\pi i \deg(a)t} (e^{2\pi it} - 1)^{-n-1} dt = \frac{1}{2\pi i} \oint f(t) e^{2\pi i \deg(a)t} (e^{2\pi it} - 1)^{-n-1} dt, \quad (7.7)$$

where the integral is taken along an anticlockwise contour containing $t = 0$ and no other pole of $f(t)$. One can always find a suitable contour of the form: t_0 to $t_0 + 1$ to $t_0 + 1 + \tau$ to $t_0 + \tau$ back to t_0 . We denote the lower and upper segments of this contour by C_1 and $-C_2$ respectively. The integrals on the lateral segments cancel because $f(t + 1) = f(t)$. For $n \leq -1$ we have

$$\begin{aligned} \oint dt f(t) e^{2\pi i \deg(a)t} (e^{2\pi it} - 1)^{-n-1} &= \sum_{j \geq 0} (-1)^j \binom{-n-1}{j} \oint dt f(t) e^{2\pi i t(\deg(a) - n - 1 - j)} \\ &= \sum_{j \geq 0} (-1)^j \binom{-n-1}{j} \left(\int_{C_1} dt f(t) e^{2\pi i t(\deg(a) - n - 1 - j)} - \int_{C_2} dt f(t) e^{2\pi i t(\deg(a) - n - 1 - j)} \right) \\ &= \sum_{j \geq 0} (-1)^j \binom{-n-1}{j} \left(F_{n+1+j-\deg(a)}^- - F_{n+1+j-\deg(a)}^+ \right). \end{aligned} \quad (7.8)$$

By (7.5) the right hand side of (7.8) equals $\alpha \delta_{n,-1}$. For $n \geq 0$ we have the following convergent series expansions

$$(e^{2\pi it} - 1)^{-1-n} = \begin{cases} (-1)^{n+1} \sum_{j \geq 0} \binom{n+j}{j} e^{2\pi i j t}, & t \in \mathbb{H}, \\ \sum_{j \geq n+1} \binom{j-1}{n} e^{-2\pi i j t}, & -t \in \mathbb{H}, \end{cases} \quad (7.9)$$

and hence

$$\begin{aligned} \oint f(t) e^{2\pi i \deg(a)t} (e^{2\pi it} - 1)^{-n-1} dt &= \sum_{j \geq n+1} \binom{j-1}{n} \int_{C_1} e^{2\pi i t(\deg(a) - j)} f(t) \\ &\quad + \sum_{j \geq 0} (-1)^n \binom{n+j}{j} \int_{C_2} f(t) e^{2\pi i t(\deg(a) + j)} dt \\ &= \sum_{j \geq n+1} \binom{j-1}{n} F_{j-\deg(a)}^- + \sum_{j \geq 0} (-1)^n \binom{n+j}{j} F_{-j-\deg(a)}^+. \end{aligned} \quad (7.10)$$

Collecting (7.8) and (7.10) yields

$$\begin{aligned} 2\pi i X(a_{[f]} b, w) &= \alpha X(a_{(-1)} b, w) \\ &\quad + \sum_{n \geq 0} \left(\sum_{j \geq n+1} \binom{j-1}{n} F_{j-\deg(a)}^- + \sum_{j \geq 0} (-1)^n \binom{n+j}{j} F_{-j-\deg(a)}^+ \right) w^{\deg(a) + \deg(b) - n - 2} Y(a_{(n)} b, w). \end{aligned} \quad (7.11)$$

We now study the right hand side of (7.3):

$$\begin{aligned} \operatorname{res}_z F_- \left(\frac{z}{w} \right) X(a, z) X(b, w) &= \sum_{n \in \mathbb{Z}} F_n^- w^{-n + \deg(b) - 1} a_n Y(b, w) \\ &= \sum_{n \in \mathbb{Z}} F_{n+1 - \deg(a)}^- w^{\deg_a + \deg_b - n - 2} a_{(n)} Y(b, w) \end{aligned} \quad (7.12)$$

Similarly we obtain

$$\operatorname{res}_z F_+ \left(\frac{z}{w} \right) X(b, w) X(a, z) = \sum_{n \in \mathbb{Z}} F_{n+1 - \deg(a)}^+ w^{\deg(a) + \deg(b) - n - 2} Y(b, w) a_{(n)} \quad (7.13)$$

Subtracting (7.13) from (7.12) and using

$$[a_{(n)}, Y(b, w)] = \sum_{j \geq 0} \binom{n}{j} w^{n-j} Y(a_{(j)} b, w),$$

we obtain

$$\begin{aligned} \operatorname{res}_z F_- \left(\frac{z}{w} \right) X(a, z) X(b, w) - \operatorname{res}_z F_+ \left(\frac{z}{w} \right) X(b, w) X(a, z) &= \sum_{n < 0} F_{n+1 - \deg(a)}^- w^{\deg(a) + \deg(b) - n - 2} a_{(n)} Y(b, w) \\ &+ \sum_{n \geq 0} F_{n+1 - \deg(a)}^- \left[\sum_{j \geq 0} \binom{n}{j} w^{\deg(a) + \deg(b) - j - 2} Y(a_{(j)} b, w) + w^{\deg_a + \deg_b - n - 2} Y(b, w) a_{(n)} \right] \\ &- \sum_{n \geq 0} F_{n+1 - \deg(a)}^+ w^{\deg(a) + \deg(b) - n - 2} Y(b, w) a_{(n)} \\ &+ \sum_{n < 0} F_{n+1 - \deg(a)}^+ \left[\sum_{j \geq 0} \binom{n}{j} w^{\deg(a) + \deg(b) - j - 2} Y(a_{(j)} b, w) - w^{\deg(a) + \deg(b) - n - 2} a_{(n)} Y(b, w) \right] \\ &= \sum_{n, j \geq 0} F_{n+1 - \deg(a)}^- \binom{n}{j} w^{\deg(a) + \deg(b) - j - 2} Y(a_{(j)} b, w) \\ &+ \sum_{n < 0, j \geq 0} F_{n+1 - \deg(a)}^+ \binom{n}{j} w^{\deg(a) + \deg(b) - j - 2} Y(a_{(j)} b, w) \\ &+ \sum_{n < 0} \left(F_{n+1 - \deg(a)}^- - F_{n+1 - \deg(a)}^+ \right) a_{(n)} Y(b, w) w^{\deg(a) + \deg(b) - n - 2} \\ &+ \sum_{n \geq 0} \left(F_{n+1 - \deg(a)}^- - F_{n+1 - \deg(a)}^+ \right) w^{\deg(a) + \deg(b) - n - 2} Y(b, w) a_{(n)}. \end{aligned} \quad (7.14)$$

The sum of the final two terms of (7.14) is

$$\begin{aligned} \alpha \sum_{n < 0} a_{(n)} Y(b, w) w^{\deg(a) + \deg(b) - n - 2} + \alpha \sum_{n \geq 0} w^{\deg(a) + \deg(b) - n - 2} Y(b, w) a_{(n)} \\ = \alpha w^{\deg(a) + \deg(b) - 1} Y(a_{(-1)} b, w) = \alpha X(a_{(-1)} b, w). \end{aligned} \quad (7.15)$$

We use the identity $\binom{-n-1}{j} = (-1)^j \binom{n+j}{j}$ to rewrite the first two terms of the right hand side of (7.14) as

$$\sum_{j \geq 0} w^{\deg(a) + \deg(b) - j - 2} Y(a_{(j)} b, w) \left[\sum_{n \geq j+1} F_{n - \deg(a)}^- \binom{n-1}{j} + \sum_{n \geq 0} F_{-n - \deg(a)}^+ (-1)^j \binom{n+j}{j} \right]. \quad (7.16)$$

Comparing (7.15) and (7.16) with (7.11) yields the desired result. \square

We only use a limited number of cases of Theorem 7.3, summarized in:

7.4 Corollary. For $a, b \in V$ the modified vertex operators satisfy

$$X(a_{[0]}b, w) = \frac{1}{2\pi i} (\operatorname{res}_z X(a, z)X(b, w) - \operatorname{res}_z X(b, w)X(a, z)). \quad (7.17)$$

$$\begin{aligned} X(a_{[\varphi_1]}b, w) - G_2(q)X(a_{[1]}b, w) + \pi i X(a_{[0]}, b, w) = \\ - \frac{1}{2\pi i} \operatorname{res}_z \left(P_1\left(\frac{zq}{w}, q\right) - 2\pi i \right) X(a, z)X(b, w) + \frac{1}{2\pi i} \operatorname{res}_z P_1\left(\frac{z}{w}, q\right) X(b, w)X(a, z). \\ X(a_{[\varphi_2]}b, w) + G_2(q)X(a_{[0]}, b, w) = \frac{1}{2\pi i} \operatorname{res}_z P_2\left(\frac{zq}{w}, q\right) X(a, z)X(b, w) \\ - \frac{1}{2\pi i} \operatorname{res}_z P_2\left(\frac{z}{w}, q\right) X(b, w)X(a, z). \\ X(a_{[\varphi_k]}b, w) = \frac{(-1)^k}{2\pi i} \operatorname{res}_z P_k\left(\frac{zq}{w}, q\right) X(a, z)X(b, w) \\ - \frac{(-1)^k}{2\pi i} \operatorname{res}_z P_k\left(\frac{z}{w}, q\right) X(b, w)X(a, z) \quad k \geq 3. \end{aligned} \quad (7.18)$$

Proof. Putting $f(t) = 1$ in Theorem 7.3 immediately yields (7.17). By (4.9) and subsequent remarks, it follows that the Fourier expansions of $-\varphi_1(t, \tau) + g_2(\tau)t - \pi i$ in the domains $|q| < |z| < 1$, resp. $1 < |z| < |q|^{-1}$, are $P_1(z, q)$, resp. $P_1(qz, q) - 2\pi i$. The first equation of (7.18) now follows from Theorem 7.3. The proofs of the remaining equations are similar. \square

Finally we will need the following slight generalization of the identities proved above.


7.5 Proposition. For each $a, b, c \in V$ the following identity of elements of $V[[u, u^{-1}, w, w^{-1}, q]]$ holds:

$$\begin{aligned} \sum_{m \geq 0} (-1)^m \binom{k+m-1}{m} P_{k+m}\left(\frac{u}{w}, q\right) X(a_{[m]}b, w) c = \frac{1}{2\pi i} \operatorname{res}_z P_k\left(\frac{u}{z}, q\right) X(a, z)X(b, w)c \\ - \frac{1}{2\pi i} \operatorname{res}_z P_k\left(\frac{u}{z}, q\right) X(b, w)X(a, z)c. \end{aligned} \quad (7.19)$$

Proof. Put $z = e^{2\pi i t}$, $w = e^{2\pi i v}$, $u = e^{2\pi i s}$ and for each $k \geq 1$ and fixed v, s, τ , consider the function

$$f(t, \tau) = (-1)^k \overline{\varphi}_k(t + v - s, \tau).$$

We apply Theorem 7.3 with the above function. This function is periodic $f(t+1) = f(t)$ and analytic except at $t = s - v + m + n\tau$, $m, n \in \mathbb{Z}$, where it has a pole. This function does not satisfy the conditions for Theorem 7.3 unless $s - v = 0$. This function is analytic on an appropriate strip containing $t = 0$. Without loss of generality we may assume $0 < \Im(s - v) < \tau$. In the following picture we can see shaded the domain where the function $f(t)$ is analytic:



That is, $f(t)$ is analytic in the strip $\Im(s - v - \tau) < \Im t < \Im(s - v)$. The proof of Theorem 7.3 is adapted verbatim by replacing the contour of integration there by a contour contained in this strip (depicted in dashed in the above picture around the point 0). Recall from (4.9) that the series $P_k(\frac{u}{zw}, q)$ converges to $f(t)$ in the domain $|q| < |\frac{u}{zw}| < 1$, that is precisely the shaded strip. It follows that the right hand side of (7.3) is given by the right hand side of (7.19) times $2\pi i$.

We can compute the left hand side of (7.3) by Laurent expansion.

$$\begin{aligned} X(a_{[f]}b, w) &= \text{res}_t f(t)X(Y[a, t]b, w) = \sum_{m \geq 0} \frac{1}{m!} \frac{d^m f}{dt^m} \Big|_{t=0} t^m X(Y[a, t]b, w) \\ &= \sum_{m \geq 0} \frac{1}{m!} \frac{d^m f}{dt^m} \Big|_{t=0} X(a_{[m]}b, w). \end{aligned} \quad (7.20)$$

Using (4.4) and (4.9) we see that the series $(-1)^m \binom{k+m-1}{m} P_{k+m}(\frac{u}{w}, q)$ converges uniformly in the domain $|q| < |u/w| < 1$, to $\frac{1}{m!} \frac{d^m f}{dt^m} \Big|_{t=0}$. Therefore the left hand side of (7.3) reads

$$\sum_{m \geq 0} (-1)^m \binom{k+m-1}{m} P_{k+m}\left(\frac{u}{w}, q\right) X(a_{[m]}b, w).$$

Notice that if we consider the situation when $-\tau < \Im(s - v) < 0$, then the function $f(t)$ would be analytic in the strip $\Im(s - v) < \Im t < \Im(s - v + \tau)$. Repeating the same argument we would obtain the equation:

$$\begin{aligned} \sum_{m \geq 0} \binom{k+m-1}{m} P_{k+m}\left(\frac{w}{u}, q\right) X(a_{[m]}b, w) c &= \frac{1}{2\pi i} \text{res}_z P_k\left(\frac{z}{u}, q\right) X(a, z)X(b, w)c \\ &\quad - \frac{1}{2\pi i} \text{res}_z P_k\left(\frac{z}{u}, q\right) X(b, w)X(a, z)c, \end{aligned} \quad (7.21)$$

which is equivalent to (7.19) by (4.12) after replacing u by uq^{-1} . □

8 Derivations and Self-Extensions

In this section we study self extensions of a module M of a vertex algebra V and the corresponding $\text{End}(M)$ -valued derivation of V . We state an analogous to the Borcherds formula of Theorem 7.3 in this setting.

8.1. Let V be a vertex algebra and M be a V -module. An $\text{End}(M)$ -valued derivation of V is a bilinear map $\Psi : V \otimes M \rightarrow M((z))$ such that for every $f \in \mathbb{C}[[z, w]][[z^{-1}, w^{-1}, (z-w)^{-1}]]$ and every $a, b \in V$, $m \in M$ we have

$$\begin{aligned} \text{res}_{z-w} \Psi(Y(a, z-w)b, w) m f(z, w) &= \text{res}_z (Y(a, z)\Psi(b, w) + \Psi(a, z)Y(b, w)) m i_{z,w} f(z, w) \\ &\quad - \text{res}_z (\Psi(b, w)Y(a, z) + Y(b, w)\Psi(a, z)) m i_{w,z} f(z, w). \end{aligned}$$

8.2 Example. Let M be a V -module and E a self extension of V -modules, i.e.,

$$0 \rightarrow M \rightarrow E \rightarrow M \rightarrow 0.$$

Writing $Y^E(\cdot, z) : V \otimes E \rightarrow E((z))$ for the V -module structure of E and choosing a splitting of the sequence as vector spaces, allows us to write in matrix form

$$Y^E(a, z) = \begin{pmatrix} Y^M(a, z) & \Psi(a, z) \\ 0 & Y^M(a, z) \end{pmatrix}. \quad (8.1)$$

Then (2.4) for the V -module E implies that Ψ is an $\text{End}(M)$ -valued derivation. Conversely, given an $\text{End}(M)$ -valued derivation Ψ , we define $Y^E(a, z)$ on $M \oplus M$ by (8.1) and obtain a self extension of M .

8.3. As in Section 7.1 we introduce the modified derivation $\sigma(a, z) = z^{-1}\Psi(z^{L_0}a, z)$ and we write

$$\Psi(a, z) = \sum_{n \in \mathbb{Z}} \Psi(a)_{(n)} z^{-1-n} \quad \text{and} \quad \sigma(a, z) = \sum_{n \in \mathbb{Z}} \sigma(a)_n z^{-n-1}.$$

In the same way as in Theorem 7.3, Corollary 7.4 and Proposition 7.5 we prove

8.4 Proposition. *Let M be a V -module, $X(\cdot, z) = X^M(\cdot, z)$ the modified vertex operators. Let $\Psi(\cdot, z)$ be a derivation and $\sigma(\cdot, z)$ the corresponding modified operator as in 8.3. For $a, b \in V$ we have:*

$$\sigma(a_{[0]}b, w) = \frac{1}{2\pi i} \operatorname{res}_z \left(\sigma(a, z)X(b, w) + X(a, z)\sigma(b, w) \right) - \frac{1}{2\pi i} \operatorname{res}_z \left(\sigma(b, w)X(a, z) + X(b, w)\sigma(a, z) \right) \quad (8.2)$$

$$\begin{aligned} \sigma(a_{[\varphi_1]}b, w) - G_2(q)\sigma(a_{[1]}b, w) + \pi i\sigma(a_{[0]}b, w) \\ = -\frac{1}{2\pi i} \operatorname{res}_z \left(P_1 \left(\frac{zq}{w}, q \right) - 2\pi i \right) \left(\sigma(a, z)X(b, w) + X(a, z)\sigma(b, w) \right) \\ + \frac{1}{2\pi i} \operatorname{res}_z P_1 \left(\frac{z}{w}, q \right) \left(\sigma(b, w)X(a, z) + X(b, w)\sigma(a, z) \right), \end{aligned} \quad (8.3)$$

$$\begin{aligned} \sigma(a_{[\varphi_2]}b, w) + G_2(q)X(a_{[0]}b, w) = \frac{1}{2\pi i} \operatorname{res}_z P_2 \left(\frac{zq}{w}, q \right) \left(\sigma(a, z)X(b, w) + X(a, z)\sigma(b, w) \right) \\ - \frac{1}{2\pi i} \operatorname{res}_z P_2 \left(\frac{z}{w}, q \right) \left(\sigma(b, w)X(a, z) + X(b, w)\sigma(a, z) \right), \end{aligned} \quad (8.4)$$

$$\begin{aligned} \sigma(a_{[\varphi_k]}b, w) = \frac{(-1)^k}{2\pi i} \operatorname{res}_z P_k \left(\frac{zq}{w}, q \right) \left(\sigma(a, z)X(b, w) + X(a, z)\sigma(b, w) \right) \\ - \frac{(-1)^k}{2\pi i} \operatorname{res}_z P_k \left(\frac{z}{w}, q \right) \left(\sigma(b, w)X(a, z) + X(b, w)\sigma(a, z) \right), \quad k \geq 3 \end{aligned} \quad (8.5)$$

$$\begin{aligned} \sum_{m \geq 0} (-1)^m \binom{k+m-1}{m} P_{k+m} \left(\frac{u}{w}, q \right) \sigma(a_{[m]}b, w) c = \\ \frac{1}{2\pi i} \operatorname{res}_z P_k \left(\frac{u}{z}, q \right) \left(\sigma(a, z)X(b, w) + X(a, z)\sigma(b, w) \right) c \\ - \frac{1}{2\pi i} \operatorname{res}_z P_k \left(\frac{u}{z}, q \right) \left(X(b, w)\sigma(a, z) + \sigma(b, w)X(a, z) \right) c. \end{aligned} \quad (8.6)$$

9 Higher Trace Functions

In this section we define and study n -point linear functionals associated with a self extension of a V -module M . These functionals are formal power series, defined as graded traces of products of an $\operatorname{End}(M)$ -valued derivation and n vertex operators acting on M . We start with the $n = 1$ case in 9.1–9.9 and then treat the general case in 9.10–9.18. In Lemma 9.4 we prove that these linear functionals satisfy the formal differential equations (5.47) which correspond to moving the marked points. We show that the functionals vanish on boundaries in Proposition 9.7 in the $n = 1$ case and in Proposition 9.11 in the general case. We show that they are formal solutions of the modular differential equation (5.48) in Theorems 9.9 and 9.14. We prove a recursive formula expressing an $(n + 1)$ -point linear functional in terms of n -point linear functionals in 9.18, which we refer to as the *insertion formula*. The motif in all proofs is similar: one uses Theorem 7.3 to obtain two terms involving different Fourier expansions of the same elliptic function, inside of the trace functional. Then one commutes one of these two terms using the cyclic property of the trace and the commutation relation (7.2), of the modified vertex operators with L_0 . We often exploit the fact that most of the Fourier series appearing are expansions of elliptic functions, with the exception of P_1 which is an expansion of φ_1 . One needs to keep track of the different domains of convergence of the involved Fourier transforms.

In this section we fix a vertex algebra V , its module M and an $\operatorname{End}(M)$ -valued derivation Ψ as in 8.3. We let $X(\cdot, z)$ and $\sigma(\cdot, z)$ be the associated modified operators.

9.1. Recall the ring \mathcal{F}_{n+1} of meromorphic elliptic functions as defined in 4.8, and its subring \mathbb{J}_*^{n+1} of weak Jacobi forms of index 0. For elements $a_0, \dots, a_n \in V$ and $f(t_0, \dots, t_n, \tau) \in \mathcal{F}_{n+1}$, we let $F(z_0, \dots, z_n, q)$ be the Fourier expansion of f in the domain $|qz_0| < |z_n| < \dots < |z_1| < |z_0|$, as in 4.12, and we define

$$F_1^n(a_0, \dots, a_n, F(z_0, \dots, z_n, q)) = \text{res}_{z_0} z_1 \dots z_n F(z_0, \dots, z_n, q) \text{tr}_M \sigma(a_0, z_0) X(a_1, z_1) \dots X(a_n, z_n) q^{L_0}. \quad (9.1)$$

We will prove below (see Theorem 10.23) that, under certain finiteness conditions on V , this series converges in the domain $|qz_1| < |z_n| < \dots < |z_1|$, to an element of \mathcal{F}_n , which we denote $F_1^n(a_0, \dots, a_n, f)$.

9.2 Remark. In this section we will make extensive use of Fourier expansions of elliptic functions in different domains. To avoid notational clutter, we will adhere to the following conventions. Whenever we write a Fourier series expansion $F(z_0, \dots, z_n, q)$ of an elliptic function $f(t_0, \dots, t_n, \tau)$ in $n+1$ variables, followed by $n+1$ field insertions in a given order, it is tacitly understood that F is the expansion whose domain of convergence is given by the order the variables in the product of fields as it is in (9.1). Occasionally other domains of expansion play a role. We write $F_{>}(z_0, \dots, z_n, q)$ to mean the Fourier expansion in the domain given by decreasing order of variable modulus, in the same order as specified by the field insertion. For example, in the expression

$$F_{>}(z_0, z_1, z_2, q) X(b, z_1) \sigma(a, z_0) X(c, z_2),$$

The series $F_{>}(z_0, z_1, z_2, q)$ converges in the domain $|qz_1| < |z_2| < |z_0| < |z_1|$.

When in need of a more verbose notation, we will write a subindex with the explicit order of the expansion. Hence, in the example above we would have

$$F_{z_1, z_0, z_2}(z_0, z_1, z_2, q) = F_{>}(z_0, z_1, z_2, q).$$

9.3 Remark. The right hand side of (9.1) is well defined for any formal series $F(z_0, \dots, z_n, q)$ not necessarily the Fourier expansion of an elliptic function. We will denote the corresponding functional also with the left hand side of (9.1), noting the points where the argument is not elliptic.

9.4 Lemma. *The series F_1^n satisfies*

$$\begin{aligned} z_i \frac{d}{dz_i} F_1^n(a_0, \dots, a_n, F(z_0, \dots, z_n, q)) &= F_1^n(a_0, \dots, (L_{-1} + L_0)a_i, \dots, a_n, F(z_0, \dots, z_n, q)) \\ &\quad + F_1^n(a_0, \dots, a_n, z_i \frac{d}{dz_i} F(z_0, \dots, z_n, q)), \quad 1 \leq i \leq n, \end{aligned} \quad (9.2)$$

$$F_1^n((L_{-1} + L_0)a_0, \dots, a_n, F(z_0, \dots, z_n, q)) + F_1^n(a_0, \dots, a_n, z_0 \frac{d}{dz_0} F(z_0, \dots, z_n, q)) = 0. \quad (9.3)$$

Proof. Follows directly from (2.1) and (7.1). □

9.5. For simplicity we first describe the situation in the case $n=1$. Let $a, b \in V$ and $f(t_0, t_1, \tau) \in \mathcal{F}_2$. We have

$$a_{[f]}b = \text{res}_{t_1-t_0} dt_1 f(t_1 - t_0) Y[a, t_1 - t_0] b \in V \otimes \mathcal{F}_1.$$

If $f(t_0, t_1, \tau) \in \mathbb{J}_*^2$ then $a_{[f]}b$ is a linear combination of the \wp_k -products defined in (4.4), and we have in fact $a_{[f]}b \in V \otimes \mathbb{QM}_*$.

Let $f(t_0, t_1, \tau) \in \mathcal{F}_2$ with Fourier series $F(z_0, z_1, q)$ convergent in the domain $|qz_0| < |z_1| < |z_0|$. Abusing notation, we have $f(t_0, t_1, \tau) = f(t_1 - t_0, \tau)$, so that $F(z_0, z_1, q) = F\left(\frac{z_1}{z_0}, q\right)$ and

$$F_1^1(a, b, F(z_0, z_1, q)) = \text{res}_{z_0} z_1 F\left(\frac{z_1}{z_0}, q\right) \text{tr}_M \sigma(a, z_0) X(b, z_1) q^{L_0}.$$

We will prove in Corollary 9.8 below that it is independent of z_1 . We now prove a proposition which plays an important technical role in what follows. This is a degree 1 analog of [2, 4.3.3]. Recall that we have introduced $\zeta(t, \tau) = \wp_1(t, \tau) - g_2(\tau)t$ and that $P_1(z, q)$ is the expansion, in the domain $|q| < |z| < 1$, of $-\zeta(t, \tau) - \pi i$.

9.6 Proposition. Let $a, b, c \in V$ and let $f(t_0, t_1, \tau) = f(t_1 - t_0, \tau) \in \mathcal{F}_2$ with Fourier expansion $F\left(\frac{z_1}{z_0}, q\right)$ convergent in the domain $|qz_0| < |z_1| < |z_0|$. Then

$$\begin{aligned}
& -\operatorname{res}_{z_0} z_1 F\left(\frac{z_1}{z_0}, q\right) \operatorname{tr}_M a_0 \sigma(b, z_0) X(c, z_1) q^{L_0} \\
&= \operatorname{res}_{z_0} z_1 P_1\left(\frac{z_1}{z_0}, q\right) \operatorname{tr}_M \sigma(a, z_0) X(b_{[f]}c, z_1) q^{L_0} - \operatorname{res}_{z_0} z_1 F\left(\frac{z_1}{z_0}, q\right) \operatorname{tr}_M \sigma(b, z_0) X(a_{[\zeta]}c, z_1) q^{L_0} \\
&\quad + \pi i \operatorname{res}_{z_0} z_1 F\left(\frac{z_1}{z_0}, q\right) \operatorname{tr}_M \sigma(b, z_0) X(a_{[0]}c, z_1) q^{L_0} \\
&\quad - \sum_{m \geq 0} (-1)^m \operatorname{res}_{z_0} z_1 P_{m+1}\left(\frac{z_1}{z_0}, q\right) F\left(\frac{z_1}{z_0}, q\right) \operatorname{tr}_M \sigma(a_{[m]}b, z_0) X(c, z_1) q^{L_0}.
\end{aligned} \tag{9.4}$$

Proof. Recall from 4.12 that $F\left(\frac{z_1 q}{u}, q\right)$ is the Fourier expansion of f in the domain $|z_1 q| < |u| < |z_1|$. Using Theorem 7.4 we obtain

$$\begin{aligned}
& \operatorname{res}_{z_0} z_1 P_1\left(\frac{z_1}{z_0}, q\right) \operatorname{tr}_M \sigma(a, z_0) X(b_{[f]}c, z_1) q^{L_0} \\
&= \frac{1}{2\pi i} \operatorname{res}_{z_0} z_1 P_1\left(\frac{z_1}{z_0}, q\right) \operatorname{tr}_M \sigma(a, z_0) \operatorname{res}_u X(b, u) X(c, z_1) F\left(\frac{z_1}{u}, q\right) q^{L_0} \\
&\quad - \frac{1}{2\pi i} \operatorname{res}_{z_0} z_1 P_1\left(\frac{z_1}{z_0}, q\right) \operatorname{tr}_M \sigma(a, z_0) \operatorname{res}_u X(c, z_1) X(b, u) F\left(\frac{z_1 q}{u}, q\right) q^{L_0} \\
&= \frac{1}{2\pi i} \operatorname{res}_{z_0} \operatorname{res}_u z_1 P_1\left(\frac{z_1}{u}, q\right) F\left(\frac{z_1}{z_0}, q\right) \operatorname{tr}_M \sigma(a, u) X(b, z_0) X(c, z_1) q^{L_0} \\
&\quad - \frac{1}{2\pi i} \operatorname{res}_{z_0} \operatorname{res}_u z_1 P_1\left(\frac{z_1}{u}, q\right) F\left(\frac{z_1 q}{z_0}, q\right) \operatorname{tr}_M \sigma(a, u) X(c, z_1) X(b, z_0) q^{L_0}.
\end{aligned} \tag{9.5}$$

Similarly, using that $P_1(zq, q)$ converges in the domain $1 < |z| < |q^{-1}|$ to $-\zeta(t, \tau) + \pi i$, while $P_1(z, q)$ converges in the domain $|q| < |z| < 1$ to $-\zeta(t, \tau) - \pi i$ we obtain:

$$\begin{aligned}
& -\operatorname{res}_{z_0} z_1 F\left(\frac{z_1}{z_0}, q\right) \operatorname{tr}_M \sigma(b, z_0) X(a_{[\zeta]}c, z_1) q^{L_0} + \pi i \operatorname{res}_{z_0} z_1 F\left(\frac{z_1}{z_0}, q\right) \operatorname{tr}_M \sigma(b, z_0) X(a_{[0]}c, z_1) q^{L_0} \\
&= \frac{1}{2\pi i} \operatorname{res}_{z_0} \operatorname{res}_u z_1 F\left(\frac{z_1}{z_0}, q\right) P_1\left(\frac{uq}{z_1}, q\right) \operatorname{tr}_M \sigma(b, z_0) X(a, u) X(c, z_1) q^{L_0} \\
&\quad - \frac{1}{2\pi i} \operatorname{res}_{z_0} \operatorname{res}_u z_1 F\left(\frac{z_1}{z_0}, q\right) \left(P_1\left(\frac{u}{z_1}, q\right) + 2\pi i\right) \operatorname{tr}_M \sigma(b, z_0) X(c, z_1) X(a, u) q^{L_0}.
\end{aligned} \tag{9.6}$$

And by (8.6) we obtain

$$\begin{aligned}
& - \sum_{m \geq 0} (-1)^m \operatorname{res}_{z_0} z_1 P_{m+1}\left(\frac{z_1}{z_0}, q\right) F\left(\frac{z_1}{z_0}, q\right) \operatorname{tr}_M \sigma(a_{[m]}b, z_0) X(c, z_1) q^{L_0} \\
&= - \frac{1}{2\pi i} \operatorname{res}_{z_0} z_1 F\left(\frac{z_1}{z_0}, q\right) \operatorname{tr}_M \operatorname{res}_u P_1\left(\frac{z_1}{u}, q\right) \sigma(a, u) X(b, z_0) X(c, z_1) q^{L_0} \\
&\quad - \frac{1}{2\pi i} \operatorname{res}_{z_0} z_1 F\left(\frac{z_1}{z_0}, q\right) \operatorname{tr}_M \operatorname{res}_u P_1\left(\frac{z_1}{u}, q\right) X(a, u) \sigma(b, z_0) X(c, z_1) q^{L_0} \\
&\quad + \frac{1}{2\pi i} \operatorname{res}_{z_0} z_1 F\left(\frac{z_1}{z_0}, q\right) \operatorname{tr}_M \operatorname{res}_u P_1\left(\frac{z_1}{u}, q\right) \sigma(b, z_0) X(a, u) X(c, z_1) q^{L_0} \\
&\quad + \frac{1}{2\pi i} \operatorname{res}_{z_0} z_1 F\left(\frac{z_1}{z_0}, q\right) \operatorname{tr}_M \operatorname{res}_u P_1\left(\frac{z_1}{u}, q\right) X(b, z_0) \sigma(a, u) X(c, z_1) q^{L_0}.
\end{aligned} \tag{9.7}$$

The first term of the right hand side of (9.7) cancels the first term of the right hand side of (9.5). Using (4.12), the third term of the right hand side of (9.7) cancels the first term of the right hand side of (9.6).

In the fourth term of the right hand side of (9.7) we commute $X(b, z_0)$ with q^{L_0} using (7.2) and that the trace is a symmetric functional, so this term cancels the second term of the right hand side of (9.5). In the second term of the right hand side of (9.7) we commute $X(a, u)$ with q^{L_0} (using (7.2) again) to obtain

$$\begin{aligned} & -\frac{1}{2\pi i} \operatorname{res}_{z_0} z_1 F\left(\frac{z_1}{z_0}, q\right) \operatorname{tr}_M \operatorname{res}_u P_1\left(\frac{z_1}{u}, q\right) \sigma(b, z_0) X(c, z_1) q^{L_0} X(a, u) \\ &= -\frac{1}{2\pi i} \operatorname{res}_{z_0} z_1 F\left(\frac{z_1}{z_0}, q\right) \operatorname{tr}_M \operatorname{res}_u P_1\left(\frac{z_1}{u}, q\right) \sigma(b, z_0) X(c, z_1) X(a, qu) q q^{L_0} \\ &= -\frac{1}{2\pi i} \operatorname{res}_{z_0} z_1 F\left(\frac{z_1}{z_0}, q\right) \operatorname{tr}_M \operatorname{res}_u P_1\left(\frac{z_1 q}{u}, q\right) \sigma(b, z_0) X(c, z_1) X(a, u) q^{L_0}. \end{aligned}$$

Now, using $P_1\left(\frac{z_1 q}{u}, q\right) = -P_1\left(\frac{u}{z_1}, q\right)$, this term plus the second term on the right hand side of (9.6) yields

$$\begin{aligned} & -\operatorname{res}_{z_0} \operatorname{res}_u z_1 F\left(\frac{z_1}{z_0}, q\right) \operatorname{tr}_M \sigma(b, z_0) X(c, z_1) X(a, u) q^{L_0} \\ &= -\operatorname{res}_{z_0} z_1 F\left(\frac{z_1}{z_0}, q\right) \operatorname{tr}_M a_0 \sigma(b, z_0) X(c, z_1) q^{L_0}. \end{aligned}$$

This proves the proposition. \square

Recall the complex C_\bullet defined in 5.7, computing the chiral homology in genus 1 of the vertex algebra $(Y, \mathbf{1}, \tilde{\omega}, Y[\cdot, z])$. To lighten the burden of notation we write $a \otimes b \otimes f$ for a typical element of $\ker d_1$, implicitly we shall always mean finite sums of such terms. The following proposition shows that F_1^1 vanishes on coboundaries.

9.7 Proposition. *Let $a, b, c \in V$ and let $f = f(t_0, t_1, t_2, \tau) \in \mathcal{F}_3$. Let $F(z_0, z_1, z_2, q)$ be the Fourier expansion of f in the domain $|qz_0| < |z_2| < |z_1| < |z_0|$. Let f_{ijk} be the coefficients of the Laurent expansion of f near $t_i - t_j = 0$, that is $f = \sum_k f_{ijk}(t_i - t_j)^k$ with $f_{ijk} \in \mathcal{F}_2$. Let $F_{ijk}(z, q)$ be the Fourier expansion of f_{ijk} in the domain $|q| < |z| < 1$. Recall the definition of $d_2(a \otimes b \otimes c \otimes f)$ from (5.3). Then the following identity holds*

$$\begin{aligned} \sum_k F_1^1\left(a, b_{[k]}c, F_{12k}\left(\frac{z_1}{z_0}, q\right)\right) - \sum_k F_1^1\left(b, a_{[k]}c, F_{02k}\left(\frac{z_1}{z_0}, q\right)\right) \\ - \sum_k F_1^1\left(a_{[k]}b, c, F_{01k}\left(\frac{z_1}{z_0}, q\right)\right) = 0. \end{aligned} \quad (9.8)$$

Proof. Using Theorem 7.3 and the convention of Remark 9.2, the first term of (9.8) can be written as

$$\begin{aligned} & \frac{1}{2\pi i} \operatorname{res}_u \operatorname{res}_{z_0} z_1 F_{>}(u, z_0, z_1) \operatorname{tr}_M \sigma(a, u) X(b, z_0) X(c, z_1) q^{L_0} \\ & - \frac{1}{2\pi i} \operatorname{res}_u \operatorname{res}_{z_0} z_1 F_{>}(u, z_0, z_1) \operatorname{tr}_M \sigma(a, u) X(c, z_1) X(b, z_0) q^{L_0}. \end{aligned} \quad (9.9)$$

Similarly the second and third term are respectively written as

$$\begin{aligned} & -\frac{1}{2\pi i} \operatorname{res}_{z_0} \operatorname{res}_u z_1 F_{>}(u, z_0, z_1) \operatorname{tr}_M \sigma(b, z_0) X(a, u) X(c, z_1) q^{L_0} \\ & + \frac{1}{2\pi i} \operatorname{res}_{z_0} \operatorname{res}_u z_1 F_{>}(u, z_0, z_1) \operatorname{tr}_M \sigma(b, z_0) X(c, z_1) X(a, u) q^{L_0}. \end{aligned} \quad (9.10)$$

$$\begin{aligned} & -\frac{1}{2\pi i} \operatorname{res}_{z_0} \operatorname{res}_u z_1 F_{>}(u, z_0, z_1) \operatorname{tr}_M (\sigma(a, u) X(b, z_0) + X(a, u) \sigma(b, z_0)) X(c, z_1) q^{L_0} \\ & + \frac{1}{2\pi i} \operatorname{res}_{z_0} \operatorname{res}_u z_1 F_{>}(u, z_0, z_1) \operatorname{tr}_M (X(b, z_0) \sigma(a, u) + \sigma(b, z_0) X(a, u)) X(c, z_1) q^{L_0}. \end{aligned} \quad (9.11)$$

The first term of (9.11) cancels the first term of (9.9) and the fourth term of (9.11) cancels the first term of (9.10). Commuting $X(b, z_0)$ with q^{L_0} using (7.2) the third term of (9.11) can be written as

$$\frac{1}{2\pi i} \operatorname{res}_{z_0} \operatorname{res}_u F_{z_0, u, z_1}(u, q^{-1}z_0, z_1, q) \sigma(a, u) X(c, z_1) X(b, z_0) q^{L_0}.$$

Noting that

$$F_{z_0, u, z_1}(u, q^{-1}z_0, z_1, q) = F_{u, z_1, z_0}(u, z_0, z_1, q),$$

since they are two Fourier expansions of the same function in the same domain, we see that this term cancels the second term of (9.9). Finally by the same reasoning the second term of (9.11) cancels the second term of (9.10) after commuting $X(a, u)$ with q^{L_0} using (7.2). \square

9.8 Corollary. *Let $a \otimes b \otimes f \in \ker d_1$. The linear functional F_1^1 satisfies:*

$$z_1 \frac{d}{dz_1} F_1^1 \left(a, b, F \left(\frac{z_1}{z_0}, q \right) \right) = 0. \quad (9.12)$$

Proof. Apply proposition 9.7 to $\tilde{\omega} \otimes a \otimes b \otimes \frac{F(z_1, z_2, q)}{2\pi i}$. We have

$$d_2 \left(\tilde{\omega} \otimes a \otimes b \otimes \frac{f(t_1, t_2, \tau)}{2\pi i} \right) = \tilde{\omega} \otimes a_{[f]} b \otimes \frac{1}{2\pi i} - \tilde{\omega}_{[0]} a \otimes b \otimes \frac{f(t_1, t_2, \tau)}{2\pi i} - a \otimes \tilde{\omega}_{[0]} b \otimes \frac{f(t_1, t_2, \tau)}{2\pi i}.$$

The first term in the right hand side vanishes by our assumption. Using that $\tilde{\omega}_{[0]} = 2\pi i(L_0 + L_{-1})$ obtain

$$F_1^1 \left((L_0 + L_{-1})a, b, F \left(\frac{z_1}{z_0}, q \right) \right) + F_1^1 \left(a, (L_0 + L_{-1})b, F \left(\frac{z_1}{z_0}, q \right) \right) = 0. \quad (9.13)$$

On the other hand, it follows from Lemma 9.4 that the left hand side of (9.13) equals

$$z_1 \frac{d}{dz_1} F_1^1 \left(a, b, F \left(\frac{z_1}{z_0}, q \right) \right).$$

\square

9.9 Theorem. *Let $b, c \in V$ and let $F(z_0, z_1, q) = F \left(\frac{z_1}{z_0}, q \right)$ be the Fourier series of $f(t_0, t_1, \tau) \in \mathcal{F}_2$ in the domain $|qz_0| < |z_1| < |z_0|$. Recall the differential operator $D = (2\pi i) \frac{d}{d\tau} + \zeta(t_1 - t_0, \tau) \frac{d}{dt_0}$ of (4.19). Let $G \left(\frac{z_1}{z_0}, q \right)$ be the Fourier series of $g(t_0, t_1, \tau) = Df(t_0, t_1, \tau) \in \mathcal{F}_2$. Then the following differential equation holds*

$$\begin{aligned} (2\pi i)^2 \left[q \frac{d}{dq} - \frac{c}{24} \right] F_1^1 \left(b, c, F \left(\frac{z_1}{z_0}, q \right) \right) &= F_1^1 \left(b, \tilde{\omega}_{[\zeta]} c, F \left(\frac{z_1}{z_0}, q \right) \right) \\ &+ F_1^1 \left(b, c, P_2 \left(\frac{z_1}{z_0}, q \right) F \left(\frac{z_1}{z_0}, q \right) \right) + F_1^1 \left(b, c, G \left(\frac{z_1}{z_0}, q \right) \right) \\ &- \sum_{k \geq 0} (-1)^k F_1^1 \left(\tilde{\omega}_{[k+1]} b, c, P_{k+2} \left(\frac{z_1}{z_0}, q \right) F \left(\frac{z_1}{z_0}, q \right) \right) - F_1^1 \left(\tilde{\omega}_{[f]} c, P_1 \left(\frac{z_1}{z_0}, q \right) \right). \end{aligned} \quad (9.14)$$

Notice that the argument of F_1^1 in the last term on the right hand side is not the Fourier expansion of an elliptic function. This term however vanishes if $b \otimes c \otimes f \in \ker d_1$.

Proof. We have from (4.20):

$$G \left(\frac{z_1}{z_0}, q \right) = (2\pi i)^2 q \frac{d}{dq} F \left(\frac{z_1}{z_0}, q \right) - 2\pi i \left[P_1 \left(\frac{z_1}{z_0}, q \right) + \pi i \right] z_0 \frac{d}{dz_0} F \left(\frac{z_1}{z_0}, q \right).$$

Putting $a = \tilde{\omega} = (2\pi i)^2(\omega - \frac{c}{24}\mathbf{1})$ in Proposition 9.6 yields

$$\begin{aligned}
& -\operatorname{res}_{z_0} z_1 F\left(\frac{z_1}{z_0}, q\right) \operatorname{tr}_M \tilde{\omega}_0 \sigma(b, z_0) X(c, z_1) q^{L_0} \\
&= \operatorname{res}_{z_0} z_1 P_1\left(\frac{z_1}{z_0}, q\right) \operatorname{tr}_M \sigma(\tilde{\omega}, z_0) X(b_{[f]}c, z_1) q^{L_0} - \operatorname{res}_{z_0} z_1 F\left(\frac{z_1}{z_0}, q\right) \operatorname{tr}_M \sigma(b, z_0) X(\tilde{\omega}_{[c]}c, z_1) q^{L_0} \\
&\quad + \pi i \operatorname{res}_{z_0} z_1 F\left(\frac{z_1}{z_0}, q\right) \operatorname{tr}_M \sigma(b, z_0) X(\tilde{\omega}_{[0]}c, z_1) q^{L_0} \\
&\quad - \sum_{k \geq 0} (-1)^k \operatorname{res}_{z_0} z_1 P_{k+1}\left(\frac{z_1}{z_0}, q\right) F\left(\frac{z_1}{z_0}, q\right) \operatorname{tr}_M \sigma(\tilde{\omega}_{[k]}b, z_0) X(c, z_1) q^{L_0}.
\end{aligned} \tag{9.15}$$

The fourth term equals

$$\begin{aligned}
& \sum_{k \geq 0} (-1)^k F_1^1\left(\tilde{\omega}_{[k+1]}b, c, P_{k+2}\left(\frac{z_1}{z_0}, q\right) F\left(\frac{z_1}{z_0}, q\right)\right) \\
& - \operatorname{res}_{z_0} z_1 P_1\left(\frac{z_1}{z_0}, q\right) F\left(\frac{z_1}{z_0}, q\right) \operatorname{tr}_M \sigma(\tilde{\omega}_{[0]}b, z_0) X(c, z_1) q^{L_0}.
\end{aligned} \tag{9.16}$$

Recalling that $\tilde{\omega}_{[0]} = (2\pi i)(L_0 + L_{-1})$, and using Lemma 9.4, we expand the second term of (9.16) as

$$\begin{aligned}
& - (2\pi i) \operatorname{res}_{z_0} z_1 P_1\left(\frac{z_1}{z_0}, q\right) F\left(\frac{z_1}{z_0}, q\right) \operatorname{tr}_M \sigma((L_0 + L_{-1})b, z_0) X(c, z_1) q^{L_0} \\
&= (2\pi i) \operatorname{res}_{z_0} z_1 z_0 \frac{d}{dz_0} \left[P_1\left(\frac{z_1}{z_0}, q\right) F\left(\frac{z_1}{z_0}, q\right) \right] \sigma(b, z_0) X(c, z_1) q^{L_0}.
\end{aligned}$$

Using (4.8) this last expression equals:

$$\begin{aligned}
& = -F_1^1\left(b, c, P_2\left(\frac{z_1}{z_0}, q\right) F\left(\frac{z_1}{z_0}, q\right)\right) \\
& \quad + (2\pi i) \operatorname{res}_{z_0} z_1 P_1\left(\frac{z_1}{z_0}, q\right) z_0 \frac{d}{dz_0} \left(F\left(\frac{z_1}{z_0}, q\right) \right) \sigma(b, z_0) X(c, z_1) q^{L_0}.
\end{aligned} \tag{9.17}$$

We rewrite the second term of the right hand side of (9.17) in terms of the function G , obtaining

$$\begin{aligned}
& -F_1^1\left(b, c, G\left(\frac{z_1}{z_0}, q\right)\right) - (2\pi i)(\pi i) F_1^1\left(b, c, z_0 \frac{d}{dz_0} F\left(\frac{z_1}{z_0}, q\right)\right) \\
& \quad + (2\pi i)^2 \operatorname{res}_{z_0} z_1 \left[q \frac{d}{dq} F\left(\frac{z_1}{z_0}, q\right) \right] \operatorname{tr}_M \sigma(b, z_0) X(c, z_1) q^{L_0}.
\end{aligned} \tag{9.18}$$

Next we use $\tilde{\omega}_{(\deg \tilde{\omega} - 1)} = (2\pi i)^2(L_0 - \frac{c}{24})$ to see that:

$$\begin{aligned}
& (2\pi i)^2 \left[q \frac{d}{dq} - \frac{c}{24} \right] F_1^1\left(b, c, F\left(\frac{z_1}{z_0}, q\right)\right) \\
& \quad = \operatorname{res}_{z_0} z_1 F\left(\frac{z_1}{z_0}, q\right) \operatorname{tr}_M \tilde{\omega}_{(\deg \tilde{\omega} - 1)} \sigma(b, z_0) X(c, z_1) q^{L_0} \\
& \quad \quad + (2\pi i)^2 \operatorname{res}_{z_0} z_1 \left[q \frac{d}{dq} F\left(\frac{z_1}{z_0}, q\right) \right] \operatorname{tr}_M \sigma(b, z_0) X(c, z_1) q^{L_0}.
\end{aligned} \tag{9.19}$$

Finally we use (9.2) and Corollary 9.8 to obtain

$$F_1^1\left(b, \tilde{\omega}_{[0]}c, F\left(\frac{z_1}{z_0}, q\right)\right) = 2\pi i F_1^1\left(b, c, z_0 \frac{d}{dz_0} F\left(\frac{z_1}{z_0}, q\right)\right).$$

Putting all this together gives the claimed differential equation. \square

We now proceed to analyze the general $n \geq 1$ situation. The proofs are essentially the same as above with a little bit of notational clutter. Care must be taken when considering the domain of expansions of the respective Fourier series. The following proposition expresses the trace of the zero mode as a linear combination of n -point functions, this is the analog of [2, Prop 4.3.3] and a generalization of Proposition 9.6.

9.10 Proposition. *Let $a, a_0, \dots, a_n \in V$, $n \geq 1$ and $f(t_0, \dots, t_n, \tau) \in \mathcal{F}_{n+1}$. Let $F(z_0, \dots, z_n, q)$ be the Fourier expansion of f in the domain $|qz_0| < |z_n| < \dots < |z_0|$. For each $1 \leq i \leq n$ consider the Laurent expansion of f near $t_0 = t_i$, that is*

$$f(t_0, \dots, t_n, \tau) = \sum_k f_{0ik}(t_1, \dots, t_n, \tau)(t_0 - t_i)^k.$$

Let $F_{0ik}(z_1, \dots, z_n, q)$ be the Fourier expansion of f_{0ik} in the domain $|qz_1| < |z_n| < \dots < |z_1|$. The following holds

$$\begin{aligned} & + \sum_{i=0}^{n-1} \sum_{m \geq 0} (-1)^m F_1^n \left(a_0, \dots, a_{[m]} a_i, \dots, a_n, F(z_0, \dots, z_n, q) P_{m+1} \left(\frac{z_n}{z_i}, q \right) \right) \\ & \quad + F_1^n \left(a_0, \dots, a_{n-1}, a_{[\zeta(t) - \pi i]} a_n, F(z_0, \dots, z_n, q) \right) \\ & - \sum_{i=1}^n \sum_k F_1^n \left(a, a_1, \dots, a_{0[k]} a_i, \dots, a_n, F_{0ik}(z_1, \dots, z_n, q) P_1 \left(\frac{z_n}{z_0}, q \right) \right) = \\ & \text{res}_{z_0} z_1 \dots z_n F(z_0, \dots, z_n, q) \text{tr}_M a(\deg a - 1) \sigma(a_0, z_0) X(a_1, z_1) \dots X(a_n, z_n) q^{L_0}. \end{aligned} \tag{9.20}$$

Proof. Using (8.6) we express the term $i = 0$ in the first summand of (9.20) as

$$\begin{aligned} & \frac{1}{2\pi i} \text{res}_{z_0} \text{res}_u z_1 \dots z_n F(z_0, \dots, z_n, q) P_1 \left(\frac{z_n}{u}, q \right) \text{tr}_M \sigma(a, u) X(a_0, z_0) \dots X(a_n, z_n) q^{L_0} \\ & + \frac{1}{2\pi i} \text{res}_{z_0} \text{res}_u z_1 \dots z_n F(z_0, \dots, z_n, q) P_1 \left(\frac{z_n}{u}, q \right) \text{tr}_M X(a, u) \sigma(a_0, z_0) X(a_1, z_1) \dots X(a_n, z_n) q^{L_0} \\ & - \frac{1}{2\pi i} \text{res}_{z_0} \text{res}_u z_1 \dots z_n F(z_0, \dots, z_n, q) P_1 \left(\frac{z_n}{u}, q \right) \text{tr}_M X(a_0, z_0) \sigma(a, u) X(a_1, z_1) \dots X(a_n, z_n) q^{L_0} \\ & - \frac{1}{2\pi i} \text{res}_{z_0} \text{res}_u z_1 \dots z_n F(z_0, \dots, z_n, q) P_1 \left(\frac{z_n}{u}, q \right) \text{tr}_M \sigma(a_0, z_0) X(a, u) X(a_1, z_1) \dots X(a_n, z_n) q^{L_0}. \end{aligned} \tag{9.21}$$

The remaining terms corresponding to $1 \leq i \leq n - 1$ can be written as follows using Proposition 7.5

$$\begin{aligned} & \frac{1}{2\pi i} \sum_{i=1}^{n-1} \text{res}_{z_0} \text{res}_u z_1 \dots z_n F(z_0, \dots, z_n, q) P_1 \left(\frac{z_n}{u}, q \right) \\ & \quad \text{tr}_M \sigma(a_0, z_0) X(a_1, z_1) \dots X(a_{i-1}, z_{i-1}) X(a, u) X(a_i, z_i) \dots X(a_n, z_n) q^{L_0} \\ & - \frac{1}{2\pi i} \sum_{i=2}^{n-1} \text{res}_{z_0} \text{res}_u z_1 \dots z_n F(z_0, \dots, z_n, q) P_1 \left(\frac{z_n}{u}, q \right) \\ & \quad \text{tr}_M \sigma(a_0, z_0) X(a_1, z_1) \dots X(a_i, z_i) X(a, u) X(a_{i+1}, z_{i+1}) \dots X(a_n, z_n) q^{L_0}. \end{aligned} \tag{9.22}$$

This is a telescoping sum and the only two terms that survive are

$$\begin{aligned} & \frac{1}{2\pi i} \text{res}_{z_0} \text{res}_u z_1 \dots z_n F(z_0, \dots, z_n, q) P_1 \left(\frac{z_n}{u}, q \right) \text{tr}_M \sigma(a_0, z_0) X(a, u) X(a_1, z_1) \dots X(a_n, z_n) q^{L_0} \\ & - \frac{1}{2\pi i} \text{res}_{z_0} \text{res}_u z_1 \dots z_n F(z_0, \dots, z_n, q) P_1 \left(\frac{z_n}{u}, q \right) \\ & \quad \text{tr}_M \sigma(a_0, z_0) X(a_1, z_1) \dots X(a_{n-1}, z_{n-1}) X(a, u) X(a_n, z_n) q^{L_0}. \end{aligned} \tag{9.23}$$

The second term using Proposition 7.4 is written as

$$\begin{aligned}
F_1^n \left(a_0, \dots, a_{n-1}, a_{[\zeta(t) - \pi i]} a_n, F(z_0, \dots, z_n, q) \right) = \\
\frac{1}{2\pi i} \operatorname{res}_{z_0} \operatorname{res}_u z_1 \dots z_n F(z_0, \dots, z_n, q) P_1 \left(\frac{z_n}{u}, q \right) \\
\operatorname{tr}_M \sigma(a_0, z_0) X(a_1, z_1) \dots X(a_{n-1}, z_{n-1}) X(a, u) X(a_n, z_n) q^{L_0} \\
- \frac{1}{2\pi i} \operatorname{res}_{z_0} \operatorname{res}_u z_1 \dots z_n F(z_0, \dots, z_n, q) \left(P_1 \left(\frac{z_n q}{u}, q \right) - 2\pi i \right) \\
\operatorname{tr}_M \sigma(a_0, z_0) X(a_1, z_1) \dots X(a_n, z_n) X(a, u) q^{L_0}.
\end{aligned} \tag{9.24}$$

We see that the first two terms of (9.20) add up to

$$\begin{aligned}
\frac{1}{2\pi i} \operatorname{res}_{z_0} \operatorname{res}_u z_1 \dots z_n F(z_0, \dots, z_n, q) P_1 \left(\frac{z_n}{u}, q \right) \operatorname{tr}_M \sigma(a, u) X(a_0, z_0) \dots X(a_n, z_n) q^{L_0} \\
+ \frac{1}{2\pi i} \operatorname{res}_{z_0} \operatorname{res}_u z_1 \dots z_n F(z_0, \dots, z_n, q) P_1 \left(\frac{z_n}{u}, q \right) \operatorname{tr}_M X(a, u) \sigma(a_0, z_0) X(a_1, z_1) \dots X(a_n, z_n) q^{L_0} \\
- \frac{1}{2\pi i} \operatorname{res}_{z_0} \operatorname{res}_u z_1 \dots z_n F(z_0, \dots, z_n, q) P_1 \left(\frac{z_n}{u}, q \right) \operatorname{tr}_M X(a_0, z_0) \sigma(a, u) X(a_1, z_1) \dots X(a_n, z_n) q^{L_0} \\
- \frac{1}{2\pi i} \sum_{i=1}^n \operatorname{res}_{z_0} \operatorname{res}_u z_1 \dots z_n F(z_0, \dots, z_n, q) \left(P_1 \left(\frac{z_n q}{u}, q \right) - 2\pi i \right) \\
\operatorname{tr}_M \sigma(a_0, z_0) \dots X(a_n, z_n) X(a, u) q^{L_0}.
\end{aligned} \tag{9.25}$$

In order to compute the third term of (9.20), we recall from 4.12 that $F(q^{-1}z_0, z_1, \dots, z_n, q)$ is the Fourier series of $f(t_0, \dots, t_n, \tau)$ convergent in the domain $|z_0| < |z_n| < \dots < |z_1| < |q^{-1}z_0|$. Using Proposition 7.3 we see that this term is a telescoping sum and the two surviving terms are

$$\begin{aligned}
- \frac{1}{2\pi i} \operatorname{res}_{z_0} \operatorname{res}_u z_1 \dots z_n F(z_0, \dots, z_n, q) P_1 \left(\frac{z_n}{u} \right) \operatorname{tr}_M \sigma(a, u) X(a_0, z_0) X(a_2, z_2) \dots X(a_n, z_n) q^{L_0} \\
+ \frac{1}{2\pi i} \operatorname{res}_{z_0} \operatorname{res}_u z_1 \dots z_n F(q^{-1}z_0, \dots, z_n, q) P_1 \left(\frac{z_n}{u} \right) \operatorname{tr}_M \sigma(a, u) X(a_1, z_1) \dots X(a_n, z_n) X(a_0, z_0) q^{L_0}.
\end{aligned} \tag{9.26}$$

The first term of (9.26) cancels the first term in (9.25). Commuting $X(a_0, z_0)$ with q^{L_0} using (7.2) in the third term of (9.25) that term cancels the second term of (9.26). Finally commuting $X(a, u)$ with q^{L_0} in the second term of (9.25) we see that this term plus the fourth term add up to

$$\operatorname{res}_{z_0} \operatorname{res}_u z_1 \dots z_n F(z_0, \dots, z_n, q) \operatorname{tr}_M \sigma(a_0, z_0) X(a_1, z_1) \dots X(a_n, z_n) X(a, u) q^{L_0},$$

proving the proposition. \square

Recall the complex C_1^n of 5.14 computing the chiral homology of the vertex algebra $(V, Y[\cdot, z], \tilde{\omega}, \mathbf{1})$ with coefficients in $V^{\otimes n}$ in genus 1. The following proposition shows that F_1^n vanishes on boundaries.

9.11 Proposition. *Let $a, b, a_1, \dots, a_n \in V$ and $f = f(u, v, t_1, \dots, t_n, \tau) \in \mathcal{F}_{n+2}$. Recall the definition of d_2 in (5.18). We let f_{0ik}, f_{1ik} and f_{00k} be the coefficients of the Laurent expansions of f about $u-t_i, v-t_i$ and $u-v$ respectively as in (5.17). Let $F_{0ik}(z_0, \dots, z_n, q)$ and $F_{1ik}(z_0, \dots, z_n, q)$ and $F_{00k}(z_0, \dots, z_n, q)$ be their respective Fourier expansions, convergent in the domain*

$$|z_0 q| < |z_n| < \dots < |z_1| < |z_0|.$$

The linear functional F_1^n satisfies:

$$\begin{aligned}
\sum_{i=1}^n \sum_k F_1^n \left(a, a_1, \dots, b_{[k]} a_i, \dots, a_n, F_{1ik} \right) - \sum_{i=1}^n \sum_k F_1^n \left(b, a_1, \dots, a_{[k]} a_i, \dots, a_n, F_{0ik} \right) \\
- \sum_k F_1^n \left(a_{[k]} b, a_1, \dots, a_n, F_{00k} \right) = 0.
\end{aligned} \tag{9.27}$$

Proof. Using Theorem 7.3 and the convention of Remark 9.2, the first term of (9.27) can be written as

$$\begin{aligned} & \frac{1}{2\pi i} \sum_{i=1}^n \operatorname{res}_{z_0} \operatorname{res}_u F_{>}(z_0, u, z_1, \dots, z_n, q) z_1 \dots z_n \operatorname{tr}_M \sigma(a, z_0) X(a_1, z_1) \dots X(b, u) X(a_i, z_i) \dots X(a_n, z_n) q^{L_0} \\ & - \frac{1}{2\pi i} \sum_{i=1}^n \operatorname{res}_{z_0} \operatorname{res}_u F_{>}(z_0, u, z_1, \dots, z_n, q) z_1 \dots z_n \operatorname{tr}_M \sigma(a, z_0) X(a_1, z_1) \dots X(a_i, z_i) X(b, u) \dots X(a_n, z_n) q^{L_0}. \end{aligned}$$

This is a telescoping sum and the only surviving terms after relabeling $u \leftrightarrow z_0$ are:

$$\begin{aligned} & \frac{1}{2\pi i} \operatorname{res}_{z_0} \operatorname{res}_u z_1 \dots z_n F_{>}(u, z_0, z_1, \dots, z_n, q) \operatorname{tr}_M \sigma(a, u) X(b, z_0) X(a_1, z_1) \dots X(a_n, z_n) q^{L_0} \\ & - \frac{1}{2\pi i} \operatorname{res}_{z_0} \operatorname{res}_u z_1 \dots z_n F_{>}(u, z_0, z_1, \dots, z_n, q) \operatorname{tr}_M \sigma(a, u) X(a_1, z_1) \dots X(a_n, z_n) X(b, z_0) q^{L_0}. \end{aligned} \quad (9.28)$$

Similarly, the second term can be written as:

$$\begin{aligned} & - \frac{1}{2\pi i} \operatorname{res}_{z_0} \operatorname{res}_u z_1 \dots z_n F_{>}(u, z_0, z_1, \dots, z_n, q) \operatorname{tr}_M \sigma(b, z_0) X(a, u) X(a_1, z_1) \dots X(a_n, z_n) q^{L_0} \\ & + \frac{1}{2\pi i} \operatorname{res}_{z_0} \operatorname{res}_u z_1 \dots z_n F_{>}(u, z_0, \dots, z_n, q) \operatorname{tr}_M \sigma(b, z_0) X(a_1, z_1) \dots X(a_n, z_n) X(a, u) q^{L_0}. \end{aligned} \quad (9.29)$$

Finally the third term is written using Proposition 8.4 as

$$\begin{aligned} & - \frac{1}{2\pi i} \operatorname{res}_{z_0} \operatorname{res}_u z_1 \dots z_n F_{>}(u, z_0, \dots, z_n, q) \operatorname{tr}_M \left(\sigma(a, u) X(b, z_0) + X(a, u) \sigma(b, z_0) \right) X(a_1, z_1) \dots X(a_n, z_n) q^{L_0} \\ & + \frac{1}{2\pi i} \operatorname{res}_{z_0} \operatorname{res}_u z_1 \dots z_n F_{>}(u, z_0, \dots, z_n, q) \operatorname{tr}_M \left(\sigma(b, z_0) X(a, u) + X(b, z_0) \sigma(a, u) \right) X(a_1, z_1) \dots X(a_n, z_n) q^{L_0}. \end{aligned} \quad (9.30)$$

The first summands of each term in (9.30) cancel the first summands of (9.28) and (9.29). The second term of (9.30) after commuting with q^{L_0} using (7.2) equals

$$- \frac{1}{2\pi i} \operatorname{res}_{z_0} \operatorname{res}_u F_{u, z_0, \dots, z_n}(q^{-1}u, z_0, \dots, z_n, q) \sigma(b, z_0) X(a_1, z_1) \dots X(a_n, z_n) X(a, u) q^{L_0}.$$

Noting that

$$F_{u, z_0, \dots, z_n}(q^{-1}u, z_0, \dots, z_n, q) = F_{z_0, \dots, z_n, u}(u, z_0, \dots, z_n, q),$$

this term cancels the second term of (9.29). Similarly the fourth term of (9.30) cancels the second term of (9.28). \square

9.12 Lemma. Let $f(t_0, \dots, t_n, \tau) \in \widehat{\mathcal{F}}_{n+1}$ and consider its Laurent expansion

$$f = \sum_k f_{0ik}(t_1, \dots, t_n, \tau) (t_0 - t_i)^k.$$

The series F_1^n satisfies

$$\begin{aligned} & 2\pi i \sum_{i=1}^n z_i \frac{d}{dz_i} F_1^n(a_0, \dots, a_n, F(z_0, \dots, z_n, q)) = \\ & \sum_{i=1}^n \sum_k F_1^n(\tilde{\omega}, a_1, \dots, a_{0[k]} a_i, \dots, a_n, F_{0ik}(z_1, \dots, z_n, q)) \end{aligned}$$

Proof. From (9.2) the left hand side equals:

$$\begin{aligned} & 2\pi i \sum_{i=1}^n F_1^n(a_0, a_1, \dots, (L_{-1} + L_0) a_i, \dots, a_n, F(z_0, \dots, z_n, q)) \\ & + 2\pi i F_1^n \left(a_0, \dots, a_n, \sum_{i=1}^n z_i \frac{d}{dz_i} F(z_0, \dots, z_n, q) \right). \end{aligned}$$

Using (9.3), Lemma 4.9 and that $\tilde{\omega}_{[0]} = (2\pi i)(L_{-1} + L_0)$, this can be written as

$$\sum_{i=0}^n F_1^n \left(a_0, a_1, \dots, \tilde{\omega}_{[0]} a_i, \dots, a_n, F(z_0, \dots, z_n, q) \right)$$

The lemma follows from Proposition 9.11 with $a = \tilde{\omega}$ and $b = a_0$. \square

9.13. Let $a_0, \dots, a_n \in V$ and $F(z_0, \dots, z_n, q)$ be the Fourier series of $f(t_0, \dots, t_n, \tau) \in \mathcal{F}_{n+1}$ in the domain $|qz_0| < |z_n| < \dots < |z_1| < |z_0|$. Recall the definition of $Df \in \mathcal{F}_{n+1}$ in 4.13. Let \overline{D} be the differential operator D in the exponential variables q, z_i as in (4.20), that is

$$\overline{D}F = \left((2\pi i)^2 q \frac{d}{dq} - \sum_{i=0}^{n-1} \left(P_1 \left(\frac{z_n}{z_i}, q \right) + \pi i \right) 2\pi i z_i \frac{d}{dz_i} \right) F(z_1, \dots, z_n, q),$$

is the Fourier series of Df convergent in the domain $|qz_0| < |z_n| < \dots < |z_1| < |z_0|$ as in (4.20). For any formal series $S(z_1, \dots, z_n, q)$ on $n+1$ variables z_1, \dots, z_n, q , we will denote by $\overline{D}S$ the differential operator above with the sum starting from $i=1$.

9.14 Theorem. Let $a_0, \dots, a_n \in V$ and $f = f(t_0, \dots, t_n, \tau) \in \mathcal{F}_{n+1}$. The following differential equation holds

$$\begin{aligned} & \left(\overline{D} - (2\pi i)^2 \frac{c}{24} \right) F_1^n(a_0, \dots, a_n, F(z_0, \dots, z_n, q)) = \\ & F_1^n(a_0, \dots, a_{n-1}, \tilde{\omega}_{[\zeta]} a_n, F(z_0, \dots, z_n, q)) + F_1^n(a_0, \dots, a_n, \overline{D}F(z_0, \dots, z_n, q)) \\ & \quad + F_1^n \left(a_0, \dots, a_n, F(z_0, \dots, z_n, q) P_2 \left(\frac{z_n}{z_0}, q \right) \right) \\ & - \sum_{i=0}^{n-1} \sum_{m \geq 0} (-1)^m F_1^n \left(a_0, \dots, \tilde{\omega}_{[m+1]} a_i, \dots, a_n, F(z_0, \dots, z_n, q) P_{m+2} \left(\frac{z_n}{z_i}, q \right) \right) \\ & - \sum_{i=1}^n \sum_k F_1^n \left(\tilde{\omega}, a_1, \dots, a_{0[k]} a_i, \dots, a_n, F_{0ik}(z_1, \dots, z_n, q) \left(P_1 \left(\frac{z_n}{z_0}, q \right) + \pi i \right) \right). \end{aligned} \tag{9.31}$$

The last term vanishes if $a_0 \otimes \dots \otimes a_n \otimes f \in \ker d_1$.

Proof. Consider Proposition 9.10 with $a = \tilde{\omega} = (2\pi i)^2 \left(\omega - \frac{c}{24} \mathbf{1} \right)$, (9.20) reads

$$\begin{aligned} & \sum_{i=0}^{n-1} \sum_{m \geq 0} (-1)^m F_1^n \left(a_0, \dots, \tilde{\omega}_{[m]} a_i, \dots, a_n, F(z_0, \dots, z_n, q) P_{m+1} \left(\frac{z_n}{z_i}, q \right) \right) \\ & \quad + F_1^n(a_0, \dots, a_{n-1}, \tilde{\omega}_{[\zeta(t) - \pi i]} a_n, F(z_0, \dots, z_n, q)) \\ & - \sum_{i=1}^n \sum_k F_1^n \left(\tilde{\omega}, a_1, \dots, a_{0[k]} a_i, \dots, a_n, F_{0ik}(z_1, \dots, z_n, q) P_1 \left(\frac{z_n}{z_0}, q \right) \right) = \\ & \text{res}_{z_0} z_1 \dots z_n F(z_0, \dots, z_n, q) \text{tr}_M \tilde{\omega}_{(\deg \tilde{\omega} - 1)} \sigma(a_0, z_0) X(a_1, z_1) \dots X(a_n, z_n) q^{L_0} \end{aligned} \tag{9.32}$$

Using that $\tilde{\omega}_0 = (2\pi i)^2 (L_0 - \frac{c}{24})$, the right hand side equals

$$\begin{aligned} & (2\pi i)^2 \left(q \frac{d}{dq} - \frac{c}{24} \right) F_1^n(a_0, \dots, a_n, F(z_0, \dots, z_n, q)) \\ & - (2\pi i)^2 \text{res}_{z_0} z_1 \dots z_n \left(q \frac{d}{dq} F(z_0, \dots, z_n, q) \right) \text{tr}_M \sigma(a_0, z_0) X(a_1, z_1) \dots X(a_n, z_n) q^{L_0}. \end{aligned} \tag{9.33}$$

Using that $\tilde{\omega}_{[0]} = (2\pi i)(L_0 + L_{-1})$ and (9.3), the summand $i = m = 0$ of the first term in (9.32) equals

$$\begin{aligned} & - 2\pi i F_1^n \left(a_0, \dots, a_n, P_1 \left(\frac{z_n}{z_0}, q \right) z_0 \frac{d}{dz_0} F(z_0, \dots, z_n, q) \right) \\ & \quad + F_1^n \left(a_0, \dots, a_n, F(z_0, \dots, z_n, q) P_2 \left(\frac{z_n}{z_0}, q \right) \right). \end{aligned} \tag{9.34}$$

Similarly using (9.2) the summands $m = 0$ and $1 \leq i \leq n - 1$ in the first term of (9.32) equal:

$$\begin{aligned}
& 2\pi i \sum_{i=1}^{n-1} z_i \frac{d}{dz_i} F_1^n \left(a_0, \dots, a_n, F(z_0, \dots, z_n, q) P_1 \left(\frac{z_n}{z_i}, q \right) \right) \\
& - 2\pi i \sum_{i=1}^{n-1} F_1^n \left(a_0, \dots, a_n, P_1 \left(\frac{z_n}{z_i}, q \right) z_i \frac{d}{dz_i} F(z_0, \dots, z_n, q) \right) \\
& + \sum_{i=1}^{n-1} F_1^n \left(a_0, \dots, a_n, P_2 \left(\frac{z_n}{z_i}, q \right) F(z_0, \dots, z_n, q) \right) = \\
& \sum_{i=1}^{n-1} P_1 \left(\frac{z_n}{z_i}, q \right) 2\pi i z_i \frac{d}{dz_i} F_1^n (a_0, \dots, a_n, F(z_0, \dots, z_n, q)) \\
& - 2\pi i \sum_{i=1}^{n-1} F_1^n \left(a_0, \dots, a_n, P_1 \left(\frac{z_n}{z_i}, q \right) z_i \frac{d}{dz_i} F(z_0, \dots, z_n, q) \right).
\end{aligned} \tag{9.35}$$

We see that the second term of the right hand side of (9.35) plus the first term of (9.34) minus the second term of (9.33) equal

$$\begin{aligned}
& F_1^n \left(a_0, \dots, a_n, \overline{D}F(z_0, \dots, z_n, q) \right) + \frac{(2\pi i)^2}{2} \sum_{i=0}^{n-1} F_1^n \left(a_0, \dots, a_n, z_i \frac{d}{dz_i} F(z_0, \dots, z_n, q) \right) \\
& = F_1^n \left(a_0, \dots, a_n, \overline{D}F(z_0, \dots, z_n, q) \right) - \frac{(2\pi i)^2}{2} F_1^n \left(a_0, \dots, a_n, z_n \frac{d}{dz_n} F(z_0, \dots, z_n, q) \right),
\end{aligned} \tag{9.36}$$

where we used Lemma 4.9 in the last equation. We use Lemma 9.4 to express

$$\begin{aligned}
F_1^n \left(a_0, \dots, a_{n-1}, \tilde{\omega}_{[-\pi i]} a_n, F(z_0, \dots, z_n, q) \right) &= \frac{(2\pi i)^2}{2} F_1^n \left(a_0, \dots, a_n, z_n \frac{d}{dz_n} F(z_0, \dots, z_n, q) \right) \\
&- \frac{(2\pi i)^2}{2} z_n \frac{d}{dz_n} F_1^n \left(a_0, \dots, a_n, F(z_0, \dots, z_n, q) \right).
\end{aligned} \tag{9.37}$$

The first term cancels the second term of (9.36) and the second term can be written as

$$\begin{aligned}
& \frac{(2\pi i)^2}{2} \sum_{i=1}^{n-1} z_i \frac{d}{dz_i} F_1^n \left(a_0, \dots, a_n, F(z_0, \dots, z_n, q) \right) \\
& - \pi i \sum_{i=1}^n \sum_k F_1^n \left(\tilde{\omega}, a_1, \dots, a_{0[k]} a_i, \dots, a_n, F_{0ik}(z_1, \dots, z_n, q) \right),
\end{aligned}$$

using Lemma 9.12. Replacing in (9.32) the Theorem follows. \square

9.15. Let $1 \leq j \leq n + 1$ and $f = f(t_0, \dots, t_{j-1}, s, t_j, \dots, t_n, \tau) \in \mathcal{F}_{n+2}$. We let

$$h(v, s, t_0, \dots, t_n, \tau) = f(t_0, \dots, t_{j-1}, v, t_j, \dots, t_n, \tau) (\zeta(s - v, \tau) - \pi i).$$

We will need the Laurent expansions of h and f in powers of $v - t_i$:

$$\begin{aligned}
f(t_0, \dots, v, t_j, \dots, t_n, \tau) &= \sum_k f_{vik}(t_0, \dots, t_n, \tau) (v - t_i)^k, \\
h(v, s, t_0, \dots, t_n, \tau) &= \sum_k h_{vik}(s, t_0, \dots, t_n, \tau) (v - t_i)^k, \\
h_{vik}(s, t_0, \dots, t_n, \tau) &= f_{vik}(t_0, \dots, t_n, \tau) (\zeta(s - t_i, \tau) - \pi i) \\
&+ f_{vi, k-1}(t_0, \dots, t_n, \tau) (\wp_2(s - t_i, \tau) + g_2(\tau)) \\
&+ \sum_{m \geq 0} f_{vi, k-m-2}(t_0, \dots, t_n, \tau) \wp_{m+3}(s - t_i, \tau).
\end{aligned}$$

Notice that $F(z_0, \dots, u, z_j, \dots, z_n, q)P_1\left(\frac{u}{w}, q\right)$ is the Fourier expansion of h in the domain $|qz_0| < |z_n| < \dots < |z_j| < |u| < |w| < |z_{j-1}| < \dots < |z_0|$. We let $F_{vik}(z_0, \dots, z_n, q)$ be the Fourier expansion of $f_{vik}(t_0, \dots, t_m, \tau)$, in the domain $|qz_0| < |z_n| < \dots < |z_0|$. Similarly, we consider the Laurent expansion of f in powers of $t_0 - t_i$ for $1 \leq i \leq n$:

$$f(t_0, \dots, t_{j-1}, v, t_j, \dots, t_n, \tau) = \sum_k f_{0ik}(t_1, \dots, t_{j-1}, v, t_j, \dots, t_n, \tau)(t_0 - t_i)^k,$$

And let $F_{0ik}(z_1, \dots, z_{j-1}, u, z_j, \dots, z_n, q)$ be its Fourier expansion convergent in the domain $|qz_1| < |z_n| < \dots < |z_j| < |u| < |z_{j-1}| < \dots < |z_1|$.

The following Proposition expresses an $n+1$ -point functional in terms of n -point functionals and the trace of the zero mode. This is the analog of [2, Prop. 4.3.4] in the degree zero case.

9.16 Proposition. *Let $a, a_0, \dots, a_n \in V$, $1 \leq j \leq n+1$, and $f = f(t_0, \dots, t_{j-1}, s, t_j, \dots, t_n, \tau) \in \mathcal{F}_{n+2}$. Let $F(z_0, \dots, z_{j-1}, w, z_j, \dots, z_n, q)$ be the Fourier expansion of f in the domain*

$$|qz_0| < |z_n| < \dots < |z_j| < |w| < |z_{j-1}| < \dots < |z_1| < |z_0|,$$

Then

$$\begin{aligned} & F_1^{n+1}\left(a_0, \dots, a_{j-1}, a, a_j, \dots, a_n, F(z_0, \dots, z_{j-1}, w, z_j, \dots, z_n, q)\right) \\ &= \text{res}_{z_0} \text{res}_u z_1 \dots, z_n F_{>}(z_0, \dots, z_{j-1}, u, z_j, \dots, z_n, q) \text{tr}_M X(a, u) \sigma(a_0, z_0) X(a_1, z_1) \dots X(a_n, z_n) q^{L_0} \\ &+ \sum_{i=0}^{j-1} \sum_k \sum_{m \geq 0} F_1^n\left(a_0, \dots, a_{[k]} a_i, \dots, a_n, F_{vi, k-m}(z_0, \dots, z_n, q) \left(P_{m+1}\left(\frac{z_i q}{w}, q\right) - 2\pi i \delta_{m,0}\right)\right) \\ &+ \sum_{i=j}^n \sum_k \sum_{m \geq 0} F_1^n\left(a_0, \dots, a_{[k]} a_i, \dots, a_n, F_{vi, k-m}(z_0, \dots, z_n, q) P_{m+1}\left(\frac{z_i}{w}, q\right)\right) \\ &- \sum_{i=1}^n \sum_k F_1^n\left(a, a_1, \dots, a_{[k]} a_i, \dots, a_n, F_{0ik, z_0, \dots, z_n}(z_1, \dots, z_{j-1}, z_0, z_j, \dots, z_n, q) \left(P_1\left(\frac{z_0 q}{w}, q\right) - 2\pi i\right)\right). \end{aligned} \quad (9.38)$$

Where we have used the notation of 9.15 on the right hand side.

Proof. It follows from (4.10) that

$$w^{-1} P_1\left(\frac{u}{w}, q\right) - w^{-1} \left(P_1\left(\frac{uq}{w}, q\right) - 2\pi i\right) = 2\pi i \delta(u, w) = 2\pi i \sum_{n \in \mathbb{Z}} w^{-n-1} u^n. \quad (9.39)$$

Consider

$$\begin{aligned} & F_1^{n+1}\left(a_0, \dots, a_{j-1}, a, a_j, \dots, a_n, F(z_0, \dots, z_{j-1}, w, z_j, \dots, z_n, q)\right) \\ &= \text{res}_{z_0} w z_1 \dots, z_n F(z_0, \dots, z_{j-1}, w, z_j, \dots, z_n, q) \text{tr}_M \sigma(a_0, z_0) \dots X(a, w) X(a_j, z_j) \dots X(a_n, z_n) q^{L_0} \\ &= \frac{1}{2\pi i} \text{res}_{z_0} \text{res}_u z_1 \dots, z_n F(z_0, \dots, z_{j-1}, u, z_j, \dots, z_n, q) P_1\left(\frac{u}{w}, q\right) \\ &\quad \text{tr}_M \sigma(a_0, z_0) \dots X(a_{j-1}, z_{j-1}) X(a, u) X(a_j, z_j) \dots X(a_n, z_n) q^{L_0} \\ &- \frac{1}{2\pi i} \text{res}_{z_0} \text{res}_u z_1 \dots, z_n F(z_0, \dots, z_{j-1}, u, z_j, \dots, z_n, q) \left(P_1\left(\frac{uq}{w}, q\right) - 2\pi i\right) \\ &\quad \text{tr}_M \sigma(a_0, z_0) \dots X(a_{j-1}, z_{j-1}) X(a, u) X(a_j, z_j) \dots X(a_n, z_n) q^{L_0}. \end{aligned} \quad (9.40)$$

Using (7.21) we can express the first term on the right hand side of (9.40) as

$$\begin{aligned}
& \frac{1}{2\pi i} \operatorname{res}_{z_0} \operatorname{res}_u z_1 \dots, z_n F(z_0, \dots, z_{j-1}, u, z_j, \dots, z_n, q) P_1 \left(\frac{u}{w}, q \right) \\
& \quad \operatorname{tr}_M \sigma(a_0, z_0) \dots X(a_{j-1}, z_{j-1}) X(a, u) X(a_j, z_j) \dots X(a_n, z_n) q^{L_0} \\
& = \frac{1}{2\pi i} \operatorname{res}_{z_0} \operatorname{res}_u z_1 \dots, z_n F_{>}(z_0, \dots, z_{j-1}, u, z_j, \dots, z_n, q) P_1 \left(\frac{u}{w}, q \right) \\
& \quad \operatorname{tr}_M \sigma(a_0, z_0) \dots X(a_j, z_j) X(a, u) X(a_{j+1}, z_{j+1}) \dots X(a_n, z_n) q^{L_0} \\
& \quad + \sum_k \sum_{m \geq 0} \operatorname{res}_{z_0} z_1 \dots, z_n F_{vj, k-m, >}(z_0, \dots, z_n, q) P_{m+1} \left(\frac{z_j}{w}, q \right) \\
& \quad \operatorname{tr}_M \sigma(a_0, z_0) \dots X(a_{[k]} a_j, z_j) \dots X(a_n, z_n) q^{L_0}.
\end{aligned} \tag{9.41}$$

Repeating this argument we can express this term as

$$\begin{aligned}
& \frac{1}{2\pi i} \operatorname{res}_{z_0} \operatorname{res}_u z_1 \dots, z_n F(z_0, \dots, z_{j-1}, u, z_j, \dots, z_n, q) P_1 \left(\frac{u}{w}, q \right) \\
& \quad \operatorname{tr}_M \sigma(a_0, z_0) \dots X(a_{j-1}, z_{j-1}) X(a, u) X(a_j, z_j) \dots X(a_n, z_n) q^{L_0} \\
& = \frac{1}{2\pi i} \operatorname{res}_{z_0} \operatorname{res}_u z_1 \dots, z_n F_{>}(z_0, \dots, z_{j-1}, u, z_j, \dots, z_n, q) P_1 \left(\frac{u}{w}, q \right) \\
& \quad \operatorname{tr}_M \sigma(a_0, z_0) \dots X(a_n, z_n) X(a, u) q^{L_0} \\
& \quad + \sum_{i=j}^n \sum_k \sum_{m \geq 0} \operatorname{res}_{z_0} z_1 \dots, z_n F_{vi, k-m, >}(z_0, \dots, z_n, q) P_{m+1} \left(\frac{z_i}{w}, q \right) \\
& \quad \operatorname{tr}_M \sigma(a_0, z_0) \dots X(a_{[k]} a_i, z_i) \dots X(a_n, z_n) q^{L_0}.
\end{aligned} \tag{9.42}$$

Similarly, using (7.21) and repeating the same procedure we can express the second term on the right hand side of (9.40) as

$$\begin{aligned}
& - \frac{1}{2\pi i} \operatorname{res}_{z_0} \operatorname{res}_u z_1 \dots, z_n F(z_0, \dots, z_{j-1}, u, z_j, \dots, z_n, q) \left(P_1 \left(\frac{uq}{w}, q \right) - 2\pi i \right) \\
& \quad \operatorname{tr}_M \sigma(a_0, z_0) \dots X(a_{j-1}, z_{j-1}) X(a, u) X(a_j, z_j) \dots X(a_n, z_n) q^{L_0} \\
& = - \frac{1}{2\pi i} \operatorname{res}_{z_0} \operatorname{res}_u z_1 \dots z_n F_{>}(z_0, \dots, z_{j-1}, u, z_j, \dots, z_n, q) \left(P_1 \left(\frac{uq}{w}, q \right) - 2\pi i \right) \\
& \quad \operatorname{tr}_M \sigma(a_0, z_0) X(a, u) X(a_1, z_1) \dots X(a_n, z_n) q^{L_0} \\
& \quad + \sum_{i=1}^{j-1} \sum_k \sum_{m \geq 0} \operatorname{res}_{z_0} z_1 \dots z_n F_{vi, k-m, >}(z_0, \dots, z_n, q) \left(P_{m+1} \left(\frac{z_i q}{w}, q \right) - 2\pi i \delta_{m,0} \right) \\
& \quad \operatorname{tr}_M \sigma(a_0, z_0) \dots X(a_{[k]} a_i, z_i) \dots X(a_n, z_n) q^{L_0}.
\end{aligned} \tag{9.43}$$

Using Proposition 8.4, the first term on the right hand side of (9.43) can be written as

$$\begin{aligned}
& - \frac{1}{2\pi i} \operatorname{res}_{z_0} \operatorname{res}_u z_1 \dots z_n F_{>}(z_0, \dots, z_{j-1}, u, z_j, \dots, z_n, q) \left(P_1 \left(\frac{uq}{w}, q \right) - 2\pi i \right) \\
& \quad \operatorname{tr}_M \sigma(a_0, z_0) X(a, u) X(a_1, z_1) \dots X(a_n, z_n) q^{L_0} \\
& = - \frac{1}{2\pi i} \operatorname{res}_{z_0} \operatorname{res}_u z_1 \dots z_n F_{>}(z_0, \dots, u, z_j, \dots, z_n, q) \left(P_1 \left(\frac{uq}{w}, q \right) - 2\pi i \right) \\
& \quad \operatorname{tr}_M \left(X(a, u) \sigma(a_0, z_0) + \sigma(a, u) X(a_0, z_0) \right) X(a_1, z_1) \dots X(a_n, z_n) q^{L_0} \\
& \quad + \frac{1}{2\pi i} \operatorname{res}_{z_0} \operatorname{res}_u z_1 \dots z_n F_{>}(z_0, \dots, u, z_j, \dots, z_n, q) \left(P_1 \left(\frac{uq}{w}, q \right) - 2\pi i \right) \\
& \quad \operatorname{tr}_M X(a_0, z_0) \sigma(a, u) X(a_1, z_1) \dots X(a_n, z_n) q^{L_0} \\
& \quad + \sum_k \sum_{m \geq 0} \operatorname{res}_{z_0} z_1 \dots z_n F_{v0, k-m, >}(z_0, \dots, z_n, q) \left(P_{m+1} \left(\frac{z_0 q}{w}, q \right) - 2\pi i \delta_{m,0} \right) \\
& \quad \operatorname{tr}_M \sigma(a_{[k]} a_0, z_0) X(a_1, z_1) \dots X(a_n, z_n) q^{L_0}.
\end{aligned} \tag{9.44}$$

Commuting $X(a, u)$ with q^{L_0} using (7.2) in the first term of (9.44) we cancel the first term in (9.42) and remains

$$\text{res}_{z_0} \text{res}_u z_1 \dots z_n F_{>}(z_0, \dots, u, z_j, \dots, z_n, q) \text{tr}_M \sigma(a_0, z_0) X(a_1, z_1) \dots X(a_n, z_n) X(a, u) q^{L_0}.$$

Commuting $X(a_0, z_0)$ with q^{L_0} in the second term, we can write this term as

$$\begin{aligned} & \frac{1}{2\pi i} \text{res}_{z_0} \text{res}_u z_1 \dots z_n F_{>}(z_0, \dots, u, z_j, \dots, z_n, q) \left(P_1 \left(\frac{uq}{w}, q \right) - 2\pi i \right) \\ & \quad \text{tr}_M X(a_0, z_0) \sigma(a, u) X(a_1, z_1) \dots X(a_n, z_n) q^{L_0} \\ & = \frac{1}{2\pi i} \text{res}_{z_0} \text{res}_u z_1 \dots z_n F_{z_0, u, z_1, \dots, z_n}(q^{-1}z_0, z_1, \dots, u, z_j, \dots, z_n, q) \left(P_1 \left(\frac{uq}{w}, q \right) - 2\pi i \right) \\ & \quad \text{tr}_M \sigma(a, u) X(a_1, z_1) \dots X(a_n, z_n) X(a_0, z_0) q^{L_0}. \end{aligned}$$

Noting that $F_{z_0, u, z_1, \dots, z_n}(q^{-1}z_0, \dots, u, z_j, \dots, z_n, q) = F_{u, z_1, \dots, z_n, z_0}(z_0, \dots, u, z_j, \dots, z_n)$ this last expression equals

$$\begin{aligned} & \frac{1}{2\pi i} \text{res}_{z_0} \text{res}_u z_1 \dots z_n F_{>}(z_0, \dots, u, z_j, \dots, z_n, q) \left(P_1 \left(\frac{uq}{w}, q \right) - 2\pi i \right) \\ & \quad \text{tr}_M \sigma(a, u) X(a_1, z_1) \dots X(a_n, z_n) X(a_0, z_0) q^{L_0}. \end{aligned}$$

Repeated use of Proposition 7.3 expresses this term as

$$\begin{aligned} & \frac{1}{2\pi i} \text{res}_{z_0} \text{res}_u z_1 \dots z_n F_{>}(z_0, \dots, u, z_j, \dots, z_n, q) \left(P_1 \left(\frac{uq}{w}, q \right) - 2\pi i \right) \\ & \quad \text{tr}_M \sigma(a, u) X(a_0, z_0) X(a_1, z_1) \dots X(a_n, z_n) q^{L_0} \\ & - \sum_{i=1}^n \sum_k \text{res}_u z_1 \dots z_n F_{0ik, >}(z_1, \dots, u, z_j, \dots, z_n, q) \left(P_1 \left(\frac{uq}{w}, q \right) - 2\pi i \right) \\ & \quad \text{tr}_M \sigma(a, u) X(a_1, z_1) \dots X(a_{0[k]} a_i, z_i) \dots X(a_n, z_n) q^{L_0}. \end{aligned} \tag{9.45}$$

The second term is the last term of (9.38) and the first term cancels the second summand on the first term in (9.44), proving the Proposition. \square

We will need the following generalization of Proposition 9.10:

9.17 Proposition. *Let $a, a_0, \dots, a_n \in V$, $n \geq 1$, $1 \leq j \leq n+1$ and $f(t_0, \dots, t_{j-1}, s, t_j, \dots, t_n, \tau) \in \mathcal{F}_{n+2}$. With the notation of 9.15, the following holds:*

$$\begin{aligned} & + \sum_{i=0}^{n-1} \sum_k \sum_{m \geq 0} (-1)^m F_1^n \left(a_0, \dots, a_{[k]} a_i, \dots, a_n, F_{vi, k-m}(z_0, \dots, z_n, q) P_{m+1} \left(\frac{z_n}{z_i}, q \right) \right) \\ & \quad + \sum_k F_1^n \left(a_0, \dots, a_{[k]} a_n, F_{vn, k+1}(z_0, \dots, z_n, q) \right) \\ & \quad - \pi i \sum_k F_1^n \left(a_0, \dots, a_{[k]} a_n, F_{vn, k}(z_0, \dots, z_n, q) \right) \\ & \quad - \sum_k \sum_{j \geq 1} F_1^n \left(a_0, \dots, a_{[k]} a_n, G_{2j}(q) F_{vn, k+1-2j}(z_0, \dots, z_n, q) \right) \\ & - \sum_{i=1}^n \sum_k F_1^n \left(a, a_1, \dots, a_{0[k]} a_i, \dots, a_n, F_{0ik, z_0, \dots, z_n}(z_1, \dots, z_{j-1}, z_0, z_j, \dots, z_n, q) P_1 \left(\frac{z_n}{z_0}, q \right) \right) = \\ & \text{res}_{z_0} \text{res}_u z_1 \dots z_n F_{>}(z_0, \dots, z_{j-1}, u, z_{j+1}, \dots, z_n, q) \text{tr}_M X(a, u) \sigma(a_0, z_0) X(a_1, z_1) \dots X(a_n, z_n) q^{L_0}. \end{aligned} \tag{9.46}$$

Proof. Using (8.6) we express the term $i = 0$ in the first summand of (9.46) as

$$\begin{aligned}
& \frac{1}{2\pi i} \operatorname{res}_{z_0} \operatorname{res}_u z_1 \dots z_n F_{>}(z_0, \dots, z_{j-1}, u, z_j, \dots, z_n, q) P_1\left(\frac{z_n}{u}, q\right) \operatorname{tr}_M \sigma(a, u) X(a_0, z_0) \dots X(a_n, z_n) q^{L_0} \\
& + \frac{1}{2\pi i} \operatorname{res}_{z_0} \operatorname{res}_u z_1 \dots z_n F_{>}(z_0, \dots, z_{j-1}, u, z_j, \dots, z_n, q) P_1\left(\frac{z_n}{u}, q\right) \\
& \quad \operatorname{tr}_M X(a, u) \sigma(a_0, z_0) X(a_1, z_1) \dots X(a_n, z_n) q^{L_0} \\
& - \frac{1}{2\pi i} \operatorname{res}_{z_0} \operatorname{res}_u z_1 \dots z_n F_{>}(z_0, \dots, z_{j-1}, u, z_j, \dots, z_n, q) P_1\left(\frac{z_n}{u}, q\right) \\
& \quad \operatorname{tr}_M X(a_0, z_0) \sigma(a, u) X(a_1, z_1) \dots X(a_n, z_n) q^{L_0} \\
& - \frac{1}{2\pi i} \operatorname{res}_{z_0} \operatorname{res}_u z_1 \dots z_n F_{>}(z_0, \dots, z_{j-1}, u, z_j, \dots, z_n, q) P_1\left(\frac{z_n}{u}, q\right) \\
& \quad \operatorname{tr}_M \sigma(a_0, z_0) X(a, u) X(a_1, z_1) \dots X(a_n, z_n) q^{L_0}.
\end{aligned} \tag{9.47}$$

The remaining terms corresponding to $1 \leq i \leq n-1$ can be written as follows using Proposition 7.5

$$\begin{aligned}
& \frac{1}{2\pi i} \sum_{i=1}^{n-1} \operatorname{res}_{z_0} \operatorname{res}_u z_1 \dots z_n F_{>}(z_0, \dots, z_{j-1}, u, z_j, \dots, z_n, q) P_1\left(\frac{z_n}{u}, q\right) \\
& \quad \operatorname{tr}_M \sigma(a_0, z_0) X(a_1, z_1) \dots X(a_{i-1}, z_{i-1}) X(a, u) X(a_i, z_i) \dots X(a_n, z_n) q^{L_0} \\
& - \frac{1}{2\pi i} \sum_{i=2}^{n-1} \operatorname{res}_{z_0} \operatorname{res}_u z_1 \dots z_n F_{>}(z_0, \dots, z_{j-1}, u, z_j, \dots, z_n, q) P_1\left(\frac{z_n}{u}, q\right) \\
& \quad \operatorname{tr}_M \sigma(a_0, z_0) X(a_1, z_1) \dots X(a_i, z_i) X(a, u) X(a_{i+1}, z_{i+1}) \dots X(a_n, z_n) q^{L_0}.
\end{aligned} \tag{9.48}$$

This is a telescoping sum and the only two terms that survive are

$$\begin{aligned}
& \frac{1}{2\pi i} \operatorname{res}_{z_0} \operatorname{res}_u z_1 \dots z_n F_{>}(z_0, \dots, z_{j-1}, u, z_j, \dots, z_n, q) P_1\left(\frac{z_n}{u}, q\right) \operatorname{tr}_M \sigma(a_0, z_0) X(a, u) X(a_1, z_1) \dots X(a_n, z_n) q^{L_0} \\
& - \frac{1}{2\pi i} \operatorname{res}_{z_0} \operatorname{res}_u z_1 \dots z_n F_{>}(z_0, \dots, z_{j-1}, u, z_j, \dots, z_n, q) P_1\left(\frac{z_n}{u}, q\right) \\
& \quad \operatorname{tr}_M \sigma(a_0, z_0) X(a_1, z_1) \dots X(a_{n-1}, z_{n-1}) X(a, u) X(a_n, z_n) q^{L_0}.
\end{aligned} \tag{9.49}$$

The sum of the second, third and fourth terms of (9.46) can be written as follows using Proposition 7.4:

$$\begin{aligned}
& \frac{1}{2\pi i} \operatorname{res}_{z_0} \operatorname{res}_u z_1 \dots z_n F_{>}(z_0, \dots, z_{j-1}, u, z_j, \dots, z_n, q) P_1\left(\frac{z_n}{u}, q\right) \\
& \quad \operatorname{tr}_M \sigma(a_0, z_0) X(a_1, z_1) \dots X(a_{n-1}, z_{n-1}) X(a, u) X(a_n, z_n) q^{L_0} \\
& - \frac{1}{2\pi i} \operatorname{res}_{z_0} \operatorname{res}_u z_1 \dots z_n F_{>}(z_0, \dots, z_{j-1}, u, z_j, \dots, z_n, q) \left(P_1\left(\frac{z_n q}{u}, q\right) - 2\pi i \right) \\
& \quad \operatorname{tr}_M \sigma(a_0, z_0) X(a_1, z_1) \dots X(a_n, z_n) X(a, u) q^{L_0}.
\end{aligned} \tag{9.50}$$

We see that the first four terms of (9.46) add up to

$$\begin{aligned}
& \frac{1}{2\pi i} \operatorname{res}_{z_0} \operatorname{res}_u z_1 \dots z_n F_{>}(z_0, \dots, z_{j-1}, u, z_j, \dots, z_n, q) P_1\left(\frac{z_n}{u}, q\right) \\
& \quad \operatorname{tr}_M \sigma(a, u) X(a_0, z_0) \dots X(a_n, z_n) q^{L_0} \\
& + \frac{1}{2\pi i} \operatorname{res}_{z_0} \operatorname{res}_u z_1 \dots z_n F_{>}(z_0, \dots, z_{j-1}, u, z_j, \dots, z_n, q) P_1\left(\frac{z_n}{u}, q\right) \\
& \quad \operatorname{tr}_M X(a, u) \sigma(a_0, z_0) X(a_1, z_1) \dots X(a_n, z_n) q^{L_0} \\
& - \frac{1}{2\pi i} \operatorname{res}_{z_0} \operatorname{res}_u z_1 \dots z_n F_{>}(z_0, \dots, z_{j-1}, u, z_j, \dots, z_n, q) P_1\left(\frac{z_n}{u}, q\right) \\
& \quad \operatorname{tr}_M X(a_0, z_0) \sigma(a, u) X(a_1, z_1) \dots X(a_n, z_n) q^{L_0} \\
& - \frac{1}{2\pi i} \sum_{i=1}^n \operatorname{res}_{z_0} \operatorname{res}_u z_1 \dots z_n F_{>}(z_0, \dots, z_{j-1}, u, z_j, \dots, z_n, q) \left(P_1\left(\frac{z_n q}{u}, q\right) - 2\pi i \right) \\
& \quad \operatorname{tr}_M \sigma(a_0, z_0) \dots X(a_n, z_n) X(a, u) q^{L_0}.
\end{aligned} \tag{9.51}$$

In order to compute the last term of the left hand side of (9.46) we use Proposition 7.3 to express this term as a telescoping sum and the two surviving terms are

$$\begin{aligned}
& -\frac{1}{2\pi i} \operatorname{res}_{z_0} \operatorname{res}_u z_1 \dots z_n F_{>}(z_0, \dots, z_{j-1}, u, z_j, \dots, z_n, q) P_1 \left(\frac{z_n}{u} \right) \\
& \quad \operatorname{tr}_M \sigma(a, u) X(a_0, z_0) X(a_2, z_2) \dots X(a_n, z_n) q^{L_0} \\
& + \frac{1}{2\pi i} \operatorname{res}_{z_0} \operatorname{res}_u z_1 \dots z_n F_{>}(z_0, \dots, z_{j-1}, u, z_j, \dots, z_n, q) P_1 \left(\frac{z_n}{u} \right) \\
& \quad \operatorname{tr}_M \sigma(a, u) X(a_1, z_1) \dots X(a_n, z_n) X(a_0, z_0) q^{L_0}.
\end{aligned} \tag{9.52}$$

The first term of (9.52) cancels the first term in (9.51). Commuting $X(a_0, z_0)$ with q^{L_0} using (7.2) in the third term of (9.51) we cancel the second term of (9.52). Finally commuting $X(a, u)$ with q^{L_0} in the second term of (9.51) we see that this term plus the fourth term add up to

$$\operatorname{res}_{z_0} \operatorname{res}_u z_1 \dots z_n F_{>}(z_0, \dots, z_{j-1}, u, z_j, \dots, z_n, q) \operatorname{tr}_M \sigma(a_0, z_0) X(a_1, z_1) \dots X(a_n, z_n) X(a, u) q^{L_0}.$$

Finally commuting $X(a, u)$ with q^{L_0} in this last expression we obtain the right hand side of (9.46), proving the proposition. \square

Combining Proposition 9.16 with Proposition 9.17 we obtain a recurrent formula expressing $n+1$ point functions in terms of n -point functions. This is the analog of [2, Prop. 4.3.4] in the degree 0 case.

9.18 Proposition. *With the notation of Proposition 9.16 we have*

$$\begin{aligned}
& F_1^{n+1} \left(a_0, \dots, a_{j-1}, a, a_j, \dots, a_n, F(z_0, \dots, z_{j-1}, w, z_j, \dots, z_n, q) \right) = \\
& \quad \sum_{i=0}^{n-1} \sum_k \sum_{m \geq 0} (-1)^m F_1^n \left(a_0, \dots, a_{[k]} a_i, \dots, a_n, F_{vi, k-m}(z_0, \dots, z_n, q) P_{m+1} \left(\frac{z_n}{z_i}, q \right) \right) \\
& \quad + \sum_k F_1^n \left(a_0, \dots, a_{[k]} a_n, F_{vn, k+1}(z_0, \dots, z_n, q) \right) \\
& \quad - \pi i \sum_k F_1^n \left(a_0, \dots, a_{[k]} a_n, F_{vn, k}(z_0, \dots, z_n, q) \right) \\
& \quad - \sum_k \sum_{j \geq 1} F_1^n \left(a_0, \dots, a_{[k]} a_n, G_{2j}(q) F_{vn, k+1-2j}(z_0, \dots, z_n, q) \right) \\
& + \sum_{i=0}^{j-1} \sum_k \sum_{m \geq 0} F_1^n \left(a_0, \dots, a_{[k]} a_i, \dots, a_n, F_{vi, k-m}(z_0, \dots, z_n, q) \left(P_{m+1} \left(\frac{z_i q}{w}, q \right) - 2\pi i \delta_{m,0} \right) \right) \\
& \quad + \sum_{i=j}^n \sum_k \sum_{m \geq 0} F_1^n \left(a_0, \dots, a_{[k]} a_i, \dots, a_n, F_{vi, k-m}(z_0, \dots, z_n, q) P_{m+1} \left(\frac{z_i}{w}, q \right) \right) \\
& \quad - \sum_{i=1}^n \sum_k F_1^n \left(a, a_1, \dots, a_{0[k]} a_i, \dots, a_n, F_{0ik, z_0, \dots, z_n}(z_1, \dots, z_0, z_j, \dots, z_n, q) \right. \\
& \quad \left. \left(P_1 \left(\frac{z_0 q}{w}, q \right) + P_1 \left(\frac{z_n}{z_0}, q \right) - 2\pi i \right) \right). \tag{9.53}
\end{aligned}$$

10 Convergence

In this section we identify conditions on the vertex algebra V which guarantee that the chiral homology $H_1^{\text{ch}}(V)$ has finite rank as an \mathbb{M}_* -module. This is the content of Proposition 10.6. Under these finiteness hypotheses on V we go on to prove that the chiral homology $H_1^{\text{ch}}(E_\tau, V)$ forms a vector bundle with connection over the disc $\{q \in \mathbb{C} \mid |q| < 1\}$ with regular singularity at $q = 0$. We relate series solutions of flat sections of (the dual of) chiral homology with the Hochschild homology $\text{HH}_1(\text{Zhu}(V))$ of the Zhu

algebra and the Koszul homology $\mathrm{HK}_1(A)$ of the associated graded $A = \mathrm{gr} V$ in 10.15 below, and deduce conditions on V which guarantee vanishing of $H_1^{\mathrm{ch}}(E_\tau, V)$ for all $\tau \in \mathbb{H}$ in Theorem 10.21. Finally, in Theorem 10.23, we deduce convergence of first trace functions under appropriate hypotheses on V .

10.1. In this section we write C_\bullet for the 1-point chiral homology complex $C_\bullet^{n=1}$ of Definition 5.14. According to 5.16 we have $H_i^{\mathrm{ch}}(V)_{\mathrm{tr}} \cong H_i^{\mathrm{ch}}(V) = H_i(C_\bullet^{n=1})$. We regard C_\bullet as a complex of \mathbb{M}_* -modules. As noted in 5.17 the specialization $\mathbb{C} \otimes_{\mathbb{M}_*} H_i^{\mathrm{ch}}(V)$ recovers the chiral homology $H_i^{\mathrm{ch}}(E_\tau, V)$ of the elliptic curve E_τ .

The relations $L_{-1} + \partial/\partial u = 0$ in C_2 and $L_{-1} + \partial/\partial t_0 = 0$ in C_1 allow one to “trade” a partial derivative for a copy of L_{-1} , thus presenting C_k as a quotient of

$$V^{\otimes k+1} \otimes \mathcal{E}_k, \quad \text{where} \quad \mathcal{E}_k = \mathbb{J}_*^k / \left\langle \partial_{t_i} f(t_1, \dots, t_k) \mid f \in \mathbb{J}_*^k \right\rangle_{i=1, \dots, k}.$$

In order to proceed it is convenient to identify a finite set of generators of \mathcal{E}_k . The rank of the \mathbb{M}_* -module \mathcal{E}_k is equal to the k^{th} Betti number of $(X \setminus 0)^k \setminus \Delta$, where $\Delta \subset X^k$ is the diagonal divisor. Explicit generators of \mathcal{E}_k were given in [6] and the differentials d_1 and d_2 written in terms of them [6, Lemma 10.2]. We now recall these formulas for ease of reference. The space \mathcal{E}_0 is generated by the class of the constant function 1. The space \mathcal{E}_1 is generated by the classes of the constant function 1, and of the Weierstrass function $\wp(t_0) = \wp_2(t_0) + g_2(\tau)$. Finally \mathcal{E}_2 is generated by five classes: those of 1, $\wp(u-v)$, $\wp(u)\wp(v)$ as well as

$$\begin{aligned} \underline{\zeta}(u, v) &= \zeta(u-v) + \zeta(v) - \zeta(u) \\ \text{and } \mathfrak{Z}(u, v) &= \wp(u-v)\underline{\zeta}(u, v) + \frac{1}{2}\wp'(u-v). \end{aligned}$$

We now have:

10.2 Lemma [6, Lemma 10.2]. *The following relations hold in C_\bullet :*

$$d_1(a \otimes a_1 \otimes 1) = a_{(0)}a_1 \otimes 1, \quad (10.1)$$

$$d_1(a \otimes a_1 \otimes \wp(t_0)) = a_{(\wp)}a_1 \otimes 1. \quad (10.2)$$

and

$$d_2(a \otimes b \otimes a_1 \otimes 1) = (a \otimes b_{(0)}a_1 - a_{(0)}b \otimes a_1 - b \otimes a_{(0)}a_1) \otimes 1, \quad (10.3)$$

$$d_2(a \otimes b \otimes a_1 \otimes \wp(u-v)) = -a_{(\wp)}b \otimes a_1 \otimes 1 \quad (10.4)$$

$$+ \sum_{j \in \mathbb{Z}_+} \frac{1}{j!} \left(L_{-1}^j a \otimes b(j)a_1 - L_{-1}^j b \otimes a(j)a_1 \right) \otimes \wp(t_0),$$

$$d_2(a \otimes b \otimes a_1 \otimes \wp(u)) = -b \otimes a_{(\wp)}a_1 \otimes \wp(t_0) \quad (10.5)$$

$$+ (a \otimes b_{(0)}a_1 + b_{(0)}a \otimes a_1) \otimes \wp(t_0),$$

$$d_2(-a \otimes b \otimes a_1 \otimes \underline{\zeta}(u, v)) = (a_{(\zeta)}b \otimes a_1 - b \otimes a_{(\zeta)}a_1 - a \otimes b_{(\zeta)}a_1) \otimes 1 \quad (10.6)$$

$$\begin{aligned} &+ \sum_{j \in \mathbb{Z}_+} \frac{(-1)^j}{(j+1)!} L_{-1}^j (a_{(j+1)}b) \otimes a_1 \otimes \wp(t_0) \\ &- \sum_{j \in \mathbb{Z}_+} \frac{1}{(j+1)!} \left(L_{-1}^j a \otimes b(j+1)a_1 + L_{-1}^j b \otimes a(j+1)a_1 \right) \otimes \wp(t_0), \end{aligned}$$

$$d_2(a \otimes b \otimes a_1 \otimes \mathfrak{Z}(u, v)) = - \sum_{j \in \mathbb{Z}_+} \frac{(-1)^j}{(j+1)!} L_{-1}^j (a_{(x^{j+1}\wp(x))}b) \otimes a_1 \otimes \wp(t_0) \quad (10.7)$$

$$+ \sum_{j \in \mathbb{Z}_+} \frac{1}{(j+1)!} \left(L_{-1}^j a \otimes b_{(x^{j+1}\wp(x))}a_1 + {}_{-1}^j b \otimes a_{(x^{j+1}\wp(x))}a_1 \right) \otimes \wp(t_0)$$

$$+ (2\pi i)^2 q \frac{d}{dq} (a_{(\zeta)}b \otimes a_1 - a \otimes b_{(\zeta)}a_1 - b \otimes a_{(\zeta)}a_1) \otimes 1,$$

$$d_2(a \otimes b \otimes a_1 \otimes \wp(u)\wp(v)) = (a \otimes b_{(\wp)}a_1 - b \otimes a_{(\wp)}a_1) \otimes \wp(t_1) \quad (10.8)$$

$$\begin{aligned}
& - \sum_{j \in \mathbb{Z}_+} \frac{(-1)^j}{(j+1)!} L_{-1}^j (a_{(x^{j+1} \wp'(x))} b) \otimes a_1 \otimes \wp(t_0) \\
& - (a_{(\wp)} b \otimes a_1 - b_{(\wp)} a \otimes a_1) \otimes \wp(t_0) - (2\pi i)^2 q \frac{d}{dq} a_{(\wp)} b \otimes a_1 \otimes 1.
\end{aligned}$$

10.3. We now recall the notion of standard filtration [4] on a conformal vertex algebra, and use it to define an increasing filtration on the complex C_\bullet . Let $\{a^i | i \in I\}$ be a strong generating set of the vertex algebra V . We suppose that the strong generators are of homogeneous conformal weight $\Delta(a^i)$. The standard filtration on V , relative to this choice, is the increasing filtration $\{G_p V\}$ in which $G_p V$ is the span of the vectors

$$a_{(-n_1)}^{i_1} \cdots a_{(-n_r)}^{i_r} \mathbf{1}$$

where $r \geq 0$, $a^1, \dots, a^r \in V$ and $n_1, \dots, n_r \in \mathbb{Z}_{>0}$ satisfying $\sum_j \Delta(a^{i_j}) \leq p$. The following properties are easily verified: If $a \in G_p V$ and $b \in G_{p'} V$ then

$$\begin{aligned}
a_{(n)} b & \in G_{p+p'} V & \text{for all } n \in \mathbb{Z}, \\
a_{(n)} b & \in G_{p+p'-1} V & \text{if } n \geq 0, \\
L_{-1} a & \in G_p V.
\end{aligned}$$

For $p \in \mathbb{Z}_+$ we define $G_p C_k$ to be the image in C_k of the subspace of $V^{\otimes k+1} \otimes \mathbb{J}_*^k$ spanned by

$$\bigcup_{\sum p_i \leq p} G_{p_1} V \otimes \cdots \otimes G_{p_{k+1}} V \otimes \mathbb{J}_*^k.$$

This defines an increasing filtration of C_\bullet by \mathbb{M}_* -modules, using here that the filtration G itself is invariant under L_{-1} .

10.4. The associated graded $A = \text{gr}_G V$ carries a commutative algebra structure with product induced by the normally ordered product $a \otimes b \mapsto a_{(-1)} b$, indeed a Poisson vertex algebra structure. We now have

$$\text{gr}_G C_k \cong A^{\otimes k+1} \otimes \mathbb{J}_*^k,$$

with differentials in the associated graded given by

$$d_1 (a \otimes a_1 \otimes 1) = 0, \tag{10.9}$$

$$d_1 (a \otimes a_1 \otimes \wp(t_0)) = a_{(-2)} a_1 \otimes 1, \tag{10.10}$$

and (see Lemma 10.2)

$$d_2 (a \otimes b \otimes a_1 \otimes 1) = 0, \tag{10.11}$$

$$d_2 (a \otimes b \otimes a_1 \otimes \wp(u-v)) = -a_{(-2)} b \otimes a_1 \otimes 1 \tag{10.12}$$

$$d_2 (a \otimes b \otimes a_1 \otimes \wp(u)) = -b \otimes a_{(-2)} a_1 \otimes 1 \tag{10.13}$$

$$d_2 (-a \otimes b \otimes a_1 \otimes \underline{\zeta}(u, v)) = (a_{(-1)} b \otimes a_1 - b \otimes a_{(-1)} a_1 - a \otimes b_{(-1)} a_1) \otimes 1 \tag{10.14}$$

$$d_2 (a \otimes b \otimes a_1 \otimes \mathfrak{Z}(u, v)) = -(a_{(-1)} b \otimes a_1 - a \otimes b_{(-1)} a_1 - b \otimes a_{(-1)} a_1) \otimes \wp(t_0) \tag{10.15}$$

$$d_2 (a \otimes b \otimes a_1 \otimes \wp(u) \wp(v)) = (a \otimes b_{(-2)} a_1 - b \otimes a_{(-2)} a_1) \otimes \wp(t_0). \tag{10.16}$$

To simplify the subsequent discussion we introduce a new complex $\tilde{C}_\bullet = \tilde{C}_\bullet^{(1)} \oplus \tilde{C}_\bullet^{(2)}$, defined as follows:

$$\tilde{C}_0^{(1)} = 0$$

$$\tilde{C}_1^{(1)} = A^{\otimes 2} \otimes \mathbb{C}[1]$$

$$\tilde{C}_2^{(1)} = A^{\otimes 3} \otimes (\mathbb{C}[1] \oplus \mathbb{C}[\wp(u)] \oplus \mathbb{C}[\wp(u-v)] \oplus \mathbb{C}[\underline{\zeta}(u, v)]),$$

with differentials given by the formulas (10.9) and (10.11)-(10.14), and

$$\tilde{C}_0^{(2)} = A \otimes \mathbb{C}[1]$$

$$\begin{aligned}\tilde{C}_1^{(2)} &= A^{\otimes 2} \otimes \mathbb{C}[\wp(u)] \\ \tilde{C}_2^{(2)} &= A^{\otimes 3} \otimes (\mathbb{C}[\mathfrak{Z}(u, v)] \oplus \mathbb{C}[\wp(u)\wp(v)]),\end{aligned}$$

with differentials given by the formulas (10.10), (10.15) and (10.16). In these formulas $\mathbb{C}[\wp(u)]$ is to be understood as the \mathbb{C} -vector space with basis consisting of the formal symbol $[\wp(u)]$, etc. We emphasize that \tilde{C}_\bullet is defined over \mathbb{C} rather than \mathbb{M}_* .

10.5 Lemma. *The differentials defined above make \tilde{C}_\bullet into a complex. There exists a surjection of complexes $\tilde{C}_\bullet \otimes_{\mathbb{C}} \mathbb{M}_* \rightarrow \text{gr}_G C_\bullet$ which induces a surjection of homology groups in degrees 0 and 1.*

Proof. The first claim is essentially a direct consequence of the construction of \tilde{C}_\bullet . However, since it will not take much space, we check that $d^2 = 0$ in \tilde{C}_\bullet directly. For $\tilde{C}_\bullet^{(1)}$ there is nothing to check. For $\tilde{C}_\bullet^{(2)}$ we need to check that

$$(a_{(-1)}b)_{(-2)}c - a_{(-2)}b_{(-1)}c - b_{(-2)}a_{(-1)}c \quad \text{and} \quad a_{(-2)}b_{(-2)}c - b_{(-2)}a_{(-2)}c,$$

vanish in A . For this we use Borcherds identity (2.2) which, in terms of modes, asserts that

$$\sum_{j \in \mathbb{Z}_+} \binom{m}{j} (a_{(n+j)}b)_{(m+k-j)} a_1 = \sum_{j \in \mathbb{Z}_+} (-1)^j \binom{n}{j} (a_{(m+n-j)}b_{(k+j)} - (-1)^n b_{(n+k-j)}a_{(m+j)}) a_1,$$

for all $a, b, a_1 \in V$ and $m, k, n \in \mathbb{Z}$. Putting $k = m = n = -1$ yields

$$\sum_{j \in \mathbb{Z}_+} (-1)^j (a_{(-1+j)}b)_{(-2-j)} a_1 = \sum_{j \in \mathbb{Z}_+} (a_{(-2-j)}b_{(-1+j)} a_1 + b_{(-2-j)}a_{(-1+j)} a_1) = a_{(-2)}b_{(-1)} a_1 + b_{(-2)}a_{(-1)} a_1.$$

In the associated graded we discard terms $a_{(j)}b$ for $j \in \mathbb{Z}_+$, and the identity becomes

$$(a_{(-1)}b)_{(-2)} a_1 = a_{(-2)}b_{(-1)} a_1 + b_{(-2)}a_{(-1)} a_1,$$

proving the first of the two relations. For the other we use the commutator formula

$$[a_{(-2)}, b_{(-2)}] a_1 = \sum_{j \in \mathbb{Z}_+} \binom{-2}{j} (a_{(j)}b)_{(-4-j)} a_1,$$

which, again, vanishes in the associated graded.

The surjection $\tilde{C}_\bullet \otimes_{\mathbb{C}} \mathbb{M}_* \rightarrow \text{gr}_G C_\bullet$ is the obvious one. Since $C_0 = V \otimes \mathbb{J}_*^0 \cong V \otimes \mathbb{M}_*$ the kernel vanishes in degree 0. Passing to the associated long exact sequence, it follows that the induced maps in homology are surjections as claimed. \square

We now consider the commutative algebra $A = \text{gr}_G V$ with derivation induced by L_{-1} . Since in general $Y(L_{-1}a, z) = \partial_z Y(a, z)$ we have $a_{(-2)}b = (L_{-1}a) \cdot b$. First we note that $\tilde{C}_1^{(2)}$ modulo the image of (10.15) coincides with $\text{HH}_1(A) \cong \Omega_{A/\mathbb{C}}$. We now verify that the differential (10.16) coincides with the differential in the Koszul complex (3.2) and conclude that $H_1(\tilde{C}_\bullet^{(2)}) \cong \text{HK}_1(A)$. We remark that the Koszul homology of $A = \text{gr} V$ appeared in a similar way in [6] upon passage to an associated graded.

We now turn to $H_1(\tilde{C}_\bullet^{(1)})$. We recall Zhu's C_2 -algebra R_V , briefly described in Section 2.12. The quotient of $\tilde{C}_1^{(1)} = A \otimes A$ by the images of (10.12) and (10.13) is just $R_V \otimes R_V$. The quotient by the image of the differential (10.14) is precisely the module of Kähler differentials $\text{HH}_1(R_V) \cong \Omega_{R_V/\mathbb{C}}$.

It follows from Lemma 10.5 and the remarks above that $H_1(\text{gr}_G C_\bullet)$ is a quotient of

$$\mathbb{M}_* \otimes_{\mathbb{C}} (\text{HK}_1(A) \oplus \text{HH}_1(R_V)).$$

If we assume finite dimensionality of the term in parentheses then we will obtain a finiteness condition on chiral homology. We now prove a strengthening of this result.

10.6 Proposition. *Let V be a strongly finitely generated conformal vertex algebra, $A = \text{gr} V$ its associated graded and R_V its C_2 -algebra. Suppose that $\dim \text{HK}_1(A) < \infty$. If $\dim \Omega_{R_V} < \infty$, or more generally if $\dim \text{HP}_1(R_V) < \infty$, then the chiral homology group $H_1(C_\bullet)$ is finitely generated as a \mathbb{M}_* -module.*

Proof. The homology groups of C_\bullet are computed by the spectral sequence associated with the filtration G . More precisely $H_\bullet(C_\bullet)$ acquires a filtration, which we also call G , defined by $G_p H_n(C_\bullet) = \text{Im}(H_n(G_p C_\bullet) \rightarrow H_n(C_\bullet))$, and there exist isomorphisms

$$\text{gr}_G H_n(C_\bullet) \cong \bigoplus_{p-q=n} E_{p,q}^\infty. \quad (10.17)$$

Since G is compatible with the \mathbb{M}_* -module structure of C_\bullet , all morphisms in the spectral sequence and in particular the isomorphism (10.17) are morphisms of \mathbb{M}_* -modules. In particular $\text{gr}_G H_1(C_\bullet)$ is a subquotient of $H_1(\text{gr}_G C_\bullet)$ and hence of

$$\mathbb{M}_* \otimes_{\mathbb{C}} (\text{HK}_1(A) \oplus \text{HH}_1(R_V)).$$

Suppose $\dim \text{HH}_1(R_V) < \infty$. Since \mathbb{M}_* is a Noetherian ring, it follows that $\text{gr}_G H_1(C_\bullet)$ is a Noetherian, and hence finitely generated, \mathbb{M}_* -module. In general let R be a ring and M an R -module with increasing filtration F exhaustive and bounded below. If $\text{gr}_F M$ is finitely generated as an R -module then so is M , as can be proved by a straightforward induction. We conclude that $H_1(C_\bullet)$ is a finitely generated \mathbb{M}_* -module.

Now we suppose $\dim \text{HP}_1(R_V) < \infty$ (dropping the hypothesis $\dim \text{HH}_1(R_V) < \infty$). We will show that $\bigoplus_{p-q=1} E_{p,q}^2$ is finitely generated as a \mathbb{M}_* -module.

The total complex of the E^1 page is $H_\bullet(\text{gr}_G C_\bullet)$ with differential which we denote $d^{(1)}$. By Lemma 10.5 the groups $H_\bullet(\text{gr}_G C_\bullet)$ receive surjections from $\mathbb{M}_* \otimes H_\bullet(\tilde{C}_\bullet)$ in degrees 0 and 1. The following diagram presents $\mathbb{M}_* \otimes H_\bullet(\tilde{C}_\bullet)$:

$$\begin{array}{ccc} & \mathbb{M}_* \otimes \Omega_{R_V} & \longleftarrow \mathbb{M}_* \otimes H_2(\tilde{C}_\bullet^{(1)}) \\ & \swarrow \partial_1^P & \searrow \\ \mathbb{M}_* \otimes R_V & \longleftarrow \mathbb{M}_* \otimes \text{HK}_1(A) & \longleftarrow \mathbb{M}_* \otimes H_2(\tilde{C}_\bullet^{(2)}) \end{array} \quad (10.18)$$

the top row being $H_\bullet(\tilde{C}_\bullet^{(1)})$ and the bottom row $H_\bullet(\tilde{C}_\bullet^{(2)})$. The arrows here denote the components of $d^{(1)}$. For example ∂_1^P is induced by (10.1) and thus coincides with (3.5) for the Poisson algebra R_V , while ∂_1 is induced by (10.2) the \wp -product. The kernel of $d_1^{(1)}$ fits into an exact sequence

$$0 \rightarrow \ker(\partial_1^P) \rightarrow \ker(d_1^{(1)}) \rightarrow \text{im}(\partial_1^P) \cap \text{im}(\partial_1) \rightarrow 0, \quad (10.19)$$

the arrows sending x to $(x, 0)$ and (x, y) to $\partial_1(y)$, respectively.

By equation (10.11) terms of the form $a \otimes b \otimes a_1 \otimes 1$ are closed in $\tilde{C}_2^{(1)}$, and so survive in $H_2(\tilde{C}_\bullet^{(1)})$. The differential of such terms is given by (10.3) which we see lies entirely in $\mathbb{M}_* \otimes \Omega_{R_V}$ and is given by

$$a \otimes b_{(0)} a_1 - a_{(0)} b \otimes a_1 - b \otimes a_{(0)} a_1.$$

The image of these terms coincides with the image $\text{im}(\partial_2^P)$ of the differential in the Poisson homology complex of R_V as in (3.6). Therefore we have a surjection

$$\frac{\ker(d_1^{(1)})}{\text{im}(\partial_2^P)} \twoheadrightarrow \frac{\ker(d_1^{(1)})}{\text{im}(d_2^{(1)})} \cong \bigoplus_{p-q=1} E_{p,q}^2.$$

The exact sequence (10.19) induces the following exact sequence

$$0 \rightarrow \text{HP}_1(R_V) \rightarrow \frac{\ker(d_1^{(1)})}{\text{im}(\partial_2^P)} \rightarrow \text{im}(\partial_1^P) \cap \text{im}(\partial_1) \rightarrow 0. \quad (10.20)$$

The rank of the middle term is thus bounded above by $\dim \text{HK}_1(A) + \dim \text{HP}_1(R_V)$. Once again it follows that $H_1(C_\bullet)$ is finitely generated, as required. \square

Proposition 10.6 above is in the same spirit as the following finiteness result on $H_0^{\text{ch}}(V)$. Under the hypothesis $\dim R_V < \infty$ (C_2 -cofiniteness) Proposition 10.7 was proved by Zhu [2] by an induction on conformal weight in $C_0^{n=1} = V$. It was observed by Arakawa and Kawasetsu [3] that Zhu's proof requires only the weaker hypothesis of $\dim \text{HP}_0(R_V) < \infty$.

10.7 Proposition. *Let V be a conformal vertex algebra. If $\dim R_V < \infty$, or more generally if $\dim \text{HP}_0(R_V) < \infty$, then $H_0^{\text{ch}}(V)$ is finitely generated as a \mathbb{M}_* -module.*

Damiolini et. al. [8] extended Zhu's results to to prove finite dimensionality of $H_0^{\text{ch}}(X, \mathcal{A}_V)$ for an arbitrary smooth curve X , where \mathcal{A}_V is the chiral algebra associated to the C_2 -cofinite vertex algebra V . It would be interesting to know if finite dimensionality of $\text{HK}_\bullet(A)$ and $\text{HP}_\bullet(R_V)$ suffice to guarantee finite dimensionality of higher chiral homology in arbitrary genera.

10.8 Lemma. *Let V be a vertex algebra for which $H_i^{\text{ch}}(V)$, for $i = 0$ or $i = 1$, is finitely generated as a \mathbb{M}_* -module, let $\tau \in \mathbb{H}$ and let $E_\tau = \mathbb{C}/\mathbb{Z} + \mathbb{Z}\tau$. Then*

$$\dim_{\mathbb{C}} H_i^{\text{ch}}(E_\tau, V) < \infty.$$

Indeed the dimension is bounded above by the rank of $H_i^{\text{ch}}(V)$ as a \mathbb{M}_* -module.

Proof. The chiral homology of E_τ is obtained specializing q to $e^{2\pi i\tau} \in \mathbb{C}$, i.e., as the tensor product

$$H_i^{\text{ch}}(E_\tau, V) \cong \mathbb{C} \otimes_{\mathbb{M}_*} H_i^{\text{ch}}(V),$$

with \mathbb{M}_* -action on \mathbb{C} given by the evaluation map sending $G_k \in \mathbb{QM}_*$ to $G_k(e^{2\pi i\tau}) \in \mathbb{C}$ for all k . It is immediate that for any fixed $\tau \in \mathbb{H}$ the vector space $H_i^{\text{ch}}(E_\tau, V)$ is finite dimensional over \mathbb{C} , with dimension bounded above by the rank of $H_i^{\text{ch}}(V)$ as a \mathbb{M}_* -module. \square

10.9 Theorem. *Let V be a finitely strongly generated conformal vertex algebra, $A = \text{gr}_F V$ its associated graded and $R_V = A^0$. Let $E_\tau = \mathbb{C}/\mathbb{Z} + \mathbb{Z}\tau$ be a smooth elliptic curve. If V satisfies (1) $\dim \text{HP}_1(R_V) < \infty$ and (2) that the kernel of the canonical surjection $JR_V \rightarrow A$ is finitely generated as a differential ideal, then*

$$\dim H_1^{\text{ch}}(E_\tau, V) < \infty.$$

Proof. By Proposition 3.3 the second of our hypotheses implies that $\dim \text{HK}_1(A) < \infty$. Therefore the hypotheses of Proposition 10.6 are satisfied and hence, by Lemma 10.8, the conclusion follows. \square

10.10. In fact the dimension of $H_i^{\text{ch}}(E_\tau, V)$ is uniformly bounded above by the rank of $H_i^{\text{ch}}(V)$ as a \mathbb{M}_* -module. Below, using techniques from the theory of differential equations and \mathcal{D} -modules, we will obtain stronger results on the dimension of chiral homology. The chiral complexes $\mathbb{QM}_* \otimes_{\mathbb{M}_*} C_\bullet^n$ come equipped with connections defined by equations (5.32) and (5.35). Inspecting these formulas we see that the connections are regular with regular singular point $q = 0$. Indeed, upon restricting to the case of $n = 1$ and passing to homology $H_i^{\text{ch}}(V)_{\text{tr}} \cong H_i^{\text{ch}}(V)$, these connections take the form

$$\nabla_\tau = (2\pi i)^2 q \frac{\partial}{\partial q} - H^{(i)},$$

with $H^{(i)}$ as in equations (5.34) and (5.39) manifestly regular in q . It follows that $H_i^{\text{ch}}(V)$ are regular \mathcal{D} -modules with regular singular point $q = 0$. This \mathcal{D} -module structure was exploited by Zhu in the case $i = 0$ considered in [2] to prove modular invariance of characters of V -modules for V a rational C_2 -cofinite conformal vertex algebra.

10.11 Lemma. *Let $i = 0$ or $i = 1$ and let V be a vertex algebra satisfying the hypotheses of Proposition 10.7 or, respectively, Proposition 10.6. Then the chiral homology $H_i^{\text{ch}}(E_\tau, V)$, for $\tau \in \mathbb{H}$ the upper half complex plane, defines a vector bundle with a flat connection.*

Proof. The \mathbb{QM}_* -module $H_i^{\text{ch}}(V)$ defines a quasicoherent sheaf over \mathbb{H} , after analytification. By [13, Proposition 8.8] an \mathcal{O} -quasicoherent \mathcal{D} -module is locally free. This sheaf is coherent by Lemma 10.8, hence a vector bundle with flat connection. \square

10.12. We now study flat sections of the dual of chiral homology (Definition 5.26) in the case $n = 1$. These are elements of

$$H_i^{\text{ch}}(V)^* = \text{Hom}_{\mathbb{Q}\mathbb{M}_*}(\mathbb{Q}\mathbb{M}_* \otimes_{\mathbb{M}_*} H_i^{\text{ch}}(V), \mathcal{F}),$$

where $\mathcal{F} = \mathcal{F}_0 \cong \mathcal{F}_1$ is the ring of holomorphic functions on \mathbb{H} , which satisfy the differential equation (5.48). We work in the analytic topology. Before proceeding we recall a classical result from the theory of ordinary differential equations (ODE).

10.13 Theorem [14, Chapter III, Theorem 1.3.1]. *Suppose $A(q)$ is an $N \times N$ matrix whose entries are analytic functions in the domain $\{q \in \mathbb{C} \mid 0 < |q| < \epsilon\}$ and has a simple singularity at $q = 0$. Then a fundamental system of solutions of the ODE*

$$w'(q) = \frac{A(q)}{q}w(q), \quad (10.21)$$

is given by

$$W(q) = U(q) \exp(\log(q)M), \quad (10.22)$$

for some constant matrix M , and $U(q)$ an $N \times N$ matrix of functions analytic in $\{q \in \mathbb{C} \mid |q| < \epsilon\}$.

10.14. An equivalent formulation is obtained upon expanding (10.22) relative to a basis of eigenvectors, namely: any solution $w(q)$ of the ODE (10.21) in a connected simply connected domain in $\{q : 0 < |q| < \epsilon\}$ is given by a convergent series of the form

$$w_i(q) = \sum_{p=0}^N \log(q)^p \sum_j q^{\lambda_j} \sum_{n=0}^{\infty} U_{pjn}^{(i)} q^n. \quad (10.23)$$

In general we refer to such an expression as a *Frobenius expansion*. In the present case the complex numbers λ_j are essentially the eigenvalues of the matrix M . It follows, moreover, that a formal series solution $w(q)$ of (10.21) of the form (10.23), is convergent.

10.15. We now suppose, for either $i = 0$ or $i = 1$, that $H_i^{\text{ch}}(V)$ is finitely generated as a \mathbb{M}_* -module, and we let $\{\alpha_1, \dots, \alpha_N\}$ be a set of generators of the finitely generated $\mathbb{Q}\mathbb{M}_*$ -module $\mathbb{Q}\mathbb{M}_* \otimes_{\mathbb{M}_*} H_i^{\text{ch}}(V)$. We denote by $A(q)$ a matrix (with entries in $\mathbb{Q}\mathbb{M}_* \subset \mathbb{C}[[q]]$) of the operator $\frac{1}{(2\pi i)^2} \nabla_\tau$ relative to the generators. We remark that $A(q)$ is not uniquely prescribed in this way since $\{\alpha_1, \dots, \alpha_N\}$ is not necessarily a basis.

Let $S \in H_i^{\text{ch}}(V)^{* \nabla}$ be a flat section of the dual of chiral homology and, for $k = 1, \dots, N$, write $S_k(\tau) = S(\alpha_k, \tau)$. Here, as always, $q = e^{2\pi i \tau}$. We denote by $w(\tau)$ the column vector with entries given by the functions $S_k(\tau)$. Then the differential equation (5.48) implies that $w(\tau)$ is a solution of the ODE

$$\frac{dw}{dq} = \frac{A(q)}{q}w.$$

The entries of $A(q)$ are formal power series with radius of convergence $\epsilon = 1$, we may thus apply Theorem 10.13 to deduce that the functions $S_k(\tau)$ possess convergent Frobenius expansions of the form (10.23).

For any $\alpha \in H_i^{\text{ch}}(V)$ we write $\alpha = \sum_{k=0}^N f_k(q)\alpha_k$ with coefficients $f_k(q) \in \mathbb{M}_*$, so that $S(\alpha, \tau) = \sum_{k=0}^N f_k(q)w_k(\tau)$, and

$$S(\alpha, \tau) = \sum_{p=0}^N \log(q)^p \sum_j q^{\lambda_j} \sum_{n=0}^{\infty} U_{pjn}(\alpha)q^n, \quad (10.24)$$

where now $U_{pjn} : H_i^{\text{ch}}(V) \rightarrow \mathbb{C}$ are \mathbb{C} -linear maps. We shall call a coefficient U_{pjo} a *leading coefficient* if λ_j is minimal in the sense that $\lambda_j - 1$ does not appear in the middle sum in (10.24).

Let U be a leading coefficient of S and let $\alpha \in H_i^{\text{ch}}(V)$ be a class which vanishes at $q = 0$, i.e., which vanishes in $\mathbb{C} \otimes_{\mathbb{Q}\mathbb{M}_*} H_i^{\text{ch}}(V)$ where the $\mathbb{Q}\mathbb{M}_*$ -action on \mathbb{C} is the evaluation map sending $G_k \in \mathbb{Q}\mathbb{M}_*$ to $G_k(0) \in \mathbb{C}$ for all k . Since S is $\mathbb{Q}\mathbb{M}_*$ -linear it is clear that $U(\alpha) = 0$. Thus U descends to a \mathbb{C} -linear map

$$U : H_i^{\text{ch}}(V, q = 0) \rightarrow \mathbb{C}.$$

10.16. It is instructive to recall Zhu's analysis of the connection on chiral homology in degree 0. We first recall the definition and basic properties of the Zhu algebra $\text{Zhu}(V)$ [2]. Let

$$\begin{aligned} f(z) &= 2\pi i \frac{e^{2\pi iz}}{e^{2\pi iz} - 1} = \zeta(z, q=0) + \pi i, \\ g(z) &= (2\pi i)^2 \frac{e^{2\pi iz}}{(e^{2\pi iz} - 1)^2} = \wp(z, q=0), \end{aligned} \tag{10.25}$$

and define, for all $a, b \in V$, the bilinear products

$$\begin{aligned} a \circ b &= \text{res}_w w^{-2} (1+w)^{\Delta(a)} a(w)b dw = a_{[g]}b, \\ \text{and } a * b &= \text{res}_w w^{-1} (1+w)^{\Delta(a)} a(w)b dw = a_{[f]}b. \end{aligned}$$

Then $\text{Zhu}(V)$ is defined to be the quotient vector space $V/V \circ V$ equipped with the associative product $a \otimes b \mapsto a * b$. This is an associative algebra with unit $\mathbf{1}$. The following formula [15, equation (3.15)] for the commutator in $\text{Zhu}(V)$ will be useful¹

$$a * b - b * a = 2\pi i a_{[0]}b \pmod{V_{[g]}V}, \tag{10.26}$$

where in the right hand side we have used the vertex algebra structure $(V, Y[\cdot, z])$ of 2.8. Let $M = \bigoplus_{n=0}^{\infty} M_n$ be a positive energy V -module graded by generalized eigenspaces M_n for L_0 of eigenvalue $\lambda + n$ for some constant $\lambda \in \mathbb{C}$. Then M_0 acquires the structure of a $\text{Zhu}(V)$ -module via the assignment $[a]m = a_0 m$ for all $a \in V$ and $m \in M_0$.

10.17. Now let $S \in H_0^{\text{ch}}(V)^{* \nabla}$ be a flat section of the dual of zeroth chiral homology of the vertex algebra $(V, Y[\cdot, z])$. Concretely this amounts to a map

$$S : \mathbb{Q}\mathbb{M}_* \otimes V \rightarrow \mathcal{F}_0,$$

which annihilates elements of the form

$$a_{[0]}a_1 \quad \text{and} \quad a_{[\wp]}a_1 = a_{[-2]}a_1 + \sum_{k \geq 2} (2k-1)G_{2k}(q)a_{[2k-2]}a_1, \tag{10.27}$$

for all $a, a_1 \in V$, and which satisfies the ODE

$$(2\pi i)^2 q \frac{d}{dq} S(a, \tau) = S(L_{[\zeta]}a, \tau),$$

for all $a \in V$. Let U be a leading coefficient of S . As shown above U descends to an element of the \mathbb{C} -linear dual of

$$H_0^{\text{ch}}(V, q=0) \cong \text{HH}_0(\text{Zhu}(V)).$$

More concretely U annihilates the restrictions of elements (10.27) at $q=0$, namely

$$a_{[0]}b \quad \text{and} \quad a_{[g]}b \quad \text{where} \quad g(t) = \wp(t, q=0) = (2\pi i)^2 \frac{e^{2\pi it}}{(e^{2\pi it} - 1)^2}. \tag{10.28}$$

It follows that U descends to a linear function on $\text{Zhu}(V)/[\text{Zhu}(V), \text{Zhu}(V)] = \text{HH}_0(\text{Zhu}(V))$.

10.18. We now recall some results from [6] on the structure of $H_1^{\text{ch}}(V, q=0)$ and its relation to the Hochschild homology $\text{HH}_1(\text{Zhu}(V))$. In loc. cit. we identified a subcomplex B_{\bullet} of the chiral complex $C_{\bullet}^{m=1}(q=0)$ such that the quotient complex computes Hochschild homology of $\text{Zhu}(V)$ in degrees 0 and 1. The associated graded complex $\text{gr}_G B_{\bullet}$ is closely related to $\tilde{C}_{\bullet}^{(2)}$ introduced above, and its homology is computed by a spectral sequence whose first page is the Koszul homology $\text{HK}_{\bullet}(\text{gr } V)$. Summarizing:

10.19 Proposition [6]. *Let V be a conformal vertex algebra. There exists a surjection*

$$\rho : H_1^{\text{ch}}(V, q=0) \rightarrow \text{HH}_1(\text{Zhu}(V)),$$

and a surjection

$$\text{HK}_1(\text{gr } V) \rightarrow \ker(\rho).$$

In particular if V is classically free then $H_1^{\text{ch}}(V, q=0) \cong \text{HH}_1(\text{Zhu}(V))$.

¹We remark that this equation appears in [2, p. 296] with a typo.

10.20. Dual to the surjection ρ we have an injection

$$\iota : \mathrm{HH}_1(\mathrm{Zhu}(V))^* \rightarrow H_1^{\mathrm{ch}}(V)^*.$$

Since in degree 1 the subcomplex B_\bullet consists of classes of the form $a \otimes a_1 \otimes \varphi(t_0)$, the morphism ι is simply evaluation on homology classes associated with constant function, that is

$$\begin{aligned} \iota(\varphi)[a \otimes a_1 \otimes 1] &= \varphi(a \otimes a_1), \\ \iota(\varphi)[a \otimes a_1 \otimes \varphi(t_0)] &= 0. \end{aligned}$$

Suppose now that V is classically free, i.e., that $\mathrm{HK}_1(\mathrm{gr} V)$, so that ι becomes an isomorphism. It would be very interesting in general to determine the extent to which leading coefficients of flat sections $S \in H_1^{\mathrm{ch}}(V)^{\ast\nabla}$ are controlled by the Hochschild homology $\mathrm{HH}_1(\mathrm{Zhu}(V))$. We hope to study this question in future work.

10.21 Theorem. *Let V be a strongly finitely generated conformal vertex algebra, $A = \mathrm{gr} V$ its associated graded and R_V its C_2 -algebra. Suppose that V is classically free, that $\dim \mathrm{HP}_1(R_V) < \infty$, and that $\mathrm{HH}_1(\mathrm{Zhu}(V)) = 0$. Then the chiral homology $H_1^{\mathrm{ch}}(V)$ vanishes.*

Proof. Since V is classically free we have $\dim \mathrm{HK}_1(A) = 0$, so the conditions of Proposition 10.6 are satisfied, $H_1^{\mathrm{ch}}(V)$ is a finitely generated \mathbb{M}_* -module. Moreover by Proposition 10.19 we have an isomorphism

$$H_1^{\mathrm{ch}}(V, q = 0) \xrightarrow{\sim} \mathrm{HH}_1(\mathrm{Zhu}(V)).$$

Suppose $H_1^{\mathrm{ch}}(V) \neq 0$, and let $S \in H_1^{\mathrm{ch}}(V)^{\ast\nabla}$ be a nonzero flat local section. Then the Frobenius expansion of S possesses some nonzero leading coefficient U which descends to a linear function on $\mathrm{HH}_1(\mathrm{Zhu}(V))$. Since $\mathrm{HH}_1(\mathrm{Zhu}(V)) = 0$ we have an immediate contradiction, and so $H_1^{\mathrm{ch}}(V) = 0$ as required. \square

10.22. We remark on the hypotheses of Theorem 10.21. The hypothesis $\dim \mathrm{HP}_1(R_V) < \infty$ is relatively mild, and is analogous to the condition $\dim \mathrm{HP}_0(R_V) < \infty$ which is weaker than C_2 -cofiniteness. The condition that $\mathrm{HH}_1(\mathrm{Zhu}(V)) = 0$ is satisfied whenever $\mathrm{Zhu}(V)$ is semisimple, for example, and in particular whenever the vertex algebra V is rational. The condition of classical freeness is satisfied for many vertex algebras, and the question of validity or failure seems to be a subtle and interesting question. We discuss this further in Section 12 below on examples.

10.23 Theorem. *Let V be a vertex algebra of central charge c , satisfying the hypotheses of Proposition 10.6. Let M be a V -module and E a self-extension of M , and let*

$$a_0 \otimes a_1 \otimes F(z_0, z_1)$$

be the Fourier series expansion of a 1-cycle $a_0 \otimes a_1 \otimes f(t_0, t_1)$ in the chiral homology complex $C_\bullet^{n=1}$ (5.27). Putting $q = e^{2\pi i\tau}$ where τ lies in the upper half complex plane, the trace function

$$F_1^1(a_0 \otimes a_1 \otimes F(z_0, z_1, q))q^{-\frac{c}{24}} = \mathrm{res}_{z_0} z_1 F(z_0, z_1, q) \mathrm{tr}_M \sigma(a_0, z_0) X(a_1, z_1) q^{L_0 - \frac{c}{24}},$$

converges to a holomorphic function of τ , and is independent of t_1 .

Proof. By Corollary 9.8 the series $F_1^1(a_0 \otimes a_1 \otimes F(z_0, z_1, q))$, which *a priori* depends on z_1 and q , is independent of z_1 . In Theorem 9.9 it is shown that F_1^1 satisfies the differential equation (9.14). This differential equation coincides formally with (5.48) under the identification $z_1 = e^{2\pi i t_1}$ (the extra term involving $\frac{c}{24}$ in (9.14) is cancelled by the factor $q^{-\frac{c}{24}}$), and in particular $q = 0$ is a regular singular point.

The fact that F_1^1 annihilates coboundaries is proved in Proposition 9.7. Since we have restricted F_1^1 to 1-cycles it descends, as in Proposition 10.6, to a morphism from a finitely generated \mathbb{QM}_* -module $\mathbb{QM}_* \otimes_{\mathbb{M}_*} H_1^{\mathrm{ch}}(V)$. Choosing generators produces a finite rank system of ODEs with regular singular point $q = 0$, thus as above the series defining $F_1^1(a_0 \otimes a_1 \otimes F(z_0, z_1, q))$ converges in simply connected subdomains of $0 < |q| < 1$. \square

11 Self-extensions and the first chiral homology

In section 9 we associated to a vertex algebra V , its module M and its self-extension E , a family of linear functionals F_1^n . In section 10 we proved that under certain finiteness conditions on V these linear functionals converge in the case $n = 1$ to holomorphic functions of τ . In this section we prove the convergence of these functionals in the general n case. We show that the limits can be analytically extended to meromorphic elliptic functions and that these extensions, after a suitable normalization factor and exponential change of coordinates, constitute a degree 1 conformal block on the torus.

11.1. Let V be a vertex algebra, M its module and Ψ an $\text{End}(M)$ -valued derivation of V . Let a_0, \dots, a_n in V and $f \in \mathbb{J}_*^{n+1}$ and consider $\alpha = a_0 \otimes \dots \otimes a_n \otimes f \in C_1^n$. Suppose α is a cycle, that is $d_1\alpha = 0$. Let $F(z_0, \dots, z_n, q)$ be the Fourier expansion of f in the domain $|qz_0| < |z_n| < \dots < |z_0|$ as in 4.12. Consider the linear functional $F_1^n(a_0, \dots, a_n, F(z_0, \dots, z_n, q))$ defined by (9.1).

Theorem. *Suppose that the finiteness conditions of Proposition 10.6 are satisfied, then F_1^n is convergent; Moreover, let $\tilde{F}_1^n(a_0, \dots, a_n, F(z_0, \dots, z_n, q))$ be its meromorphic continuation and put*

$$S_1^n(a_0, \dots, a_n, f; t_1, \dots, t_n, \tau) = \tilde{F}_1^n(a_0, \dots, a_n, F(z_0, \dots, e^{2\pi i t_n}, e^{2\pi i \tau})) q^{-\frac{c}{24}}. \quad (11.1)$$

Then the family $S_1 = \{S_1^n\}_{n \geq 1}$ satisfies the genus-one property for the vertex algebra $(V, Y[\cdot, t], \tilde{\omega}, 1)$.

The proof of this Theorem will be given in the next sub-sections. Notice that b) in Definition 6.1 is automatic. We proceed by induction in n . In 11.2 we consider the $n = 1$ case. In 11.3 we prove that F_1^n converges for all n . In 11.3 we prove that S_1^n are well defined on cycles in C_1^n . In 11.5 we prove that S_1^n are elliptic functions and in 11.6 that S_1^n are flat with respect to the connection on the moduli space of marked elliptic curves.

11.2. We want to prove that S_1^n is a well defined flat section of the dual to chiral homology, that is $S_1^n \in H_1^{\text{ch}}(V^{\otimes n})^{*\nabla}$. We will proceed by induction on n . Let us prove first the $n = 1$ case. Notice that c) in definition 6.1 is empty in this case. Let $\alpha \in C_1^1$ be such that $d_1\alpha = 0$. The fact that $S_1^1(\alpha; t_1, \tau) \in \mathcal{F}_1$, that is, it is independent of t_1 , follows immediately from Theorem 10.23. The fact that it vanishes if $\alpha = d_2\beta$ for $\beta \in C_2^1$ follows from Proposition 9.7, hence S_1^1 defines a section of the dual of chiral homology $H_1^{\text{ch}}(V)^*$. The fact that it is a flat section, that is, that it satisfies (5.48), follows at once from Theorem 9.9 after the coordinate change $q = e^{2\pi i \tau}$ and $z_j = e^{2\pi i t_j}$. The extra $\frac{c}{24}$ in the left hand side of (9.14) is cancelled with the factor $q^{-\frac{c}{24}}$ in (11.1).

11.3. Let us now assume that S_1^n is a well defined flat section in $H_1^{\text{ch}}(V^{\otimes n})^{*\nabla}$ and that it satisfies c) in 6.1. We will prove that so is S_1^{n+1} and that it satisfies the insertion formula (6.4). We start by proving that the linear functionals converge and satisfy the insertion formula (6.4).

Assume the notation of Proposition 9.16, so that we have $a_0, \dots, a_n, a \in V$, $1 \leq j \leq n+1$ and $f = f(t_0, \dots, t_{j-1}, s, t_j, \dots, t_n, \tau) \in \mathcal{F}_{n+2}$. Suppose that

$$\alpha = a_0 \otimes \dots \otimes a_{j-1} \otimes a \otimes a_j \otimes \dots \otimes a_n \otimes f \in C_1^{n+1},$$

is a cycle, $d_1\alpha = 0$. In the same way as in 6.2 we notice that the right hand side of (9.53) consists of F_1^n applied to elements in C_1^n that are themselves cycles, therefore they are convergent by induction.

11.4. The meromorphic continuation of the limit in the right hand side of (9.53) is independent of j . Therefore it follows from Proposition 9.18 that for $\sigma \in S_n$ we have

$$\tilde{F}_1^n(a_0, a_{\sigma(1)}, \dots, a_{\sigma(n)}, F(z_0, z_{\sigma(1)}, \dots, z_{\sigma(n)}, q)) = \tilde{F}_1^n(a_0, \dots, a_n, F(z_0, \dots, z_n, q)). \quad (11.2)$$

Thus S_1^{n+1} satisfies (6.2). The fact that S_1^{n+1} vanishes on elements like (5.25) follows from (9.3).

11.5. For $1 \leq j \leq n$ we have from (11.2) that

$$S_1^{n+1}(a_0, \dots, a_{j-1}, a, a_{j+1}, \dots, a_n, f(t_0, \dots, t_{j-1}, t, t_{j+1}, \dots, t_n, \tau); t_1, \dots, t_{j-1}, t, t_{j+1}, \dots, t_n, \tau) = S_1^{n+1}(a_0, a, a_1, \dots, a_n, f(t_0, t, t_1, \dots, t_n, \tau); t, t_1, \dots, t_n, \tau)$$

We proved in 6.2 that the right hand side is an elliptic function in \mathcal{F}_n , it follows that the left hand side is elliptic as well.

11.6. The fact that S_1^{n+1} vanishes on boundaries Follows from Proposition 9.11. The fact that S_1^{n+1} satisfies the differential equations (5.47) follows from (9.2). And finally the fact that S_1^{n+1} satisfies (5.48) follows from Theorem 9.14, thus proving that $S_1^{n+1} \in H_1^{\text{ch}}(V^{\otimes n})^{*\nabla}$ and we have established the induction step.

12 Examples

In this section we work out the constructions and results introduced in this paper in the context of several specific examples. Firstly we derive consequences of the finiteness and vanishing results on chiral homology of Section 10 above for rational vertex algebras. Secondly we work out a simple example of higher trace functions for the Heisenberg vertex algebra.

12.1. The condition that V be classically free is satisfied by universal enveloping vertex algebras such as the Heisenberg vertex algebra, the universal affine vertex algebra $V^k(\mathfrak{g})$ at arbitrary level, their quantum Hamiltonian reductions, that is the universal affine W algebras $W^k(\mathfrak{g}, f)$, and the universal Virasoro vertex algebra Vir^c at arbitrary central charge. The question of classical freeness of rational vertex algebras is quite subtle: the boundary minimal models $\text{Vir}_{2,2k+1}$ are rational and classically free, but other minimal models such as the Ising model $\text{Vir}_{3,4}$ are not. It is also known that the rational affine vertex algebras $V_1(\mathfrak{g})$ are classically free, as are $V_k(\mathfrak{sl}_2)$ for $k \in \mathbb{Z}_+$. This latter statement is believed to generalize from \mathfrak{sl}_2 to other semisimple Lie algebras \mathfrak{g} . We now prove the following corollary of Theorem 10.21.

12.2 Corollary. *The chiral homology $H_1^{\text{ch}}(E_\tau, V)$ of the smooth elliptic curve $E_\tau = \mathbb{C}/\mathbb{Z} + \mathbb{Z}\tau$ with coefficients in the vertex algebra V vanishes in the following cases:*

- (a) $V = \text{Vir}_{2,2s+1}$ is a boundary Virasoro minimal model,
- (b) $V = V_k(\mathfrak{sl}_2)$ is the simple affine vertex algebra at level $k \in \mathbb{Z}_+$,
- (c) $V = V_1(\mathfrak{g})$ is the simple affine vertex algebra at level 1, for \mathfrak{g} a simple Lie algebra.

Proof. For V any of the vertex algebras listed in the statement, $\text{Zhu}(V)$ is semisimple since V is rational, and hence $\text{HH}_1(\text{Zhu}(V)) = 0$. We also have $\dim \text{HP}_1(R_V) < \infty$ as these vertex algebras are all C_2 -cofinite.

For all three parts it suffices, therefore, to verify that V is classically free. Parts (a) and (b) were established in [6] (in the case of (b) relying heavily on results of [16]). Classical freeness of $V_1(\mathfrak{g})$ is proved in [17]. The assertions now follow from Theorem 10.21. \square

12.3 Remark. For \mathfrak{g} of ADE type, item c) above is a particular case of a stronger result of Gaitsgory for lattice vertex algebras [18] (see also [1, §4.9]).

In a similar manner we prove the following.

12.4 Corollary. *The chiral homology $H_1^{\text{ch}}(E_\tau, V)$ of the smooth elliptic curve $E_\tau = \mathbb{C}/\mathbb{Z} + \mathbb{Z}\tau$ with coefficients in the Ising model $\text{Vir}_{3,4}$ is finite dimensional.*

Proof. It was established in [19] that, while $V = \text{Vir}_{3,4}$ is not classically free, the kernel of $JR_V \rightarrow A = \text{gr}_F V$ is generated as a differential ideal by a single element. Hence $\dim \text{HK}_1(A) = 1$. Finite dimensionality now follows from Theorem 10.9. \square

12.5 Remark. The above proof shows that $\dim H_1^{\text{ch}}(E_\tau, \text{Vir}_{3,4}) \leq 1$. In [19] we found an explicit generator for $\text{HK}_1(A)$ in this case, we do not know if the image of this element in $H_1^{\text{ch}}(E_\tau, \text{Vir}_{3,4})$ vanishes or not.

12.6. We describe an example of higher trace function, associated with the Heisenberg vertex algebra. Let $\mathfrak{h} = \mathbb{C}x$ be a one dimensional vector space with a non degenerate bilinear form $(x, x) = 1$, and let $\widehat{\mathfrak{h}} = \mathfrak{h}[t, t^{-1}] \oplus \mathbb{C}K$ be its affinization, i.e., the Lie algebra defined by

$$[xt^m, xt^n] = m\delta_{m,-n}K, \quad [K, \widehat{\mathfrak{h}}] = 0.$$

We write x_m for xt^m . The vacuum module

$$H = U(\widehat{\mathfrak{h}}) \otimes_{U(\mathfrak{h}[t] \oplus \mathbb{C}K)} \mathbb{C}v,$$

in which $Kv = v$ and $\mathfrak{h}[t]v = 0$, carries the natural structure of a conformal vertex algebra with vacuum vector $\mathbf{1} = 1 \otimes v$. This is the rank one Heisenberg vertex algebra. We identify $a_{(-1)}\mathbf{1} = a_{-1}\mathbf{1}$ with $a \in \mathfrak{h}$, the vertex algebra structure is determined by the quantum fields $Y(a, z) = \sum_{n \in \mathbb{Z}} a_n z^{-n-1}$. The conformal vector is given by $\omega = x_{-1}x_{-1}\mathbf{1}$ and has central charge $c = 1$.

12.7. Let D be a finite dimensional $\mathbb{C}[x]$ -module, we consider the induced $\widehat{\mathfrak{h}}$ -module

$$M_D = U(\widehat{\mathfrak{h}}) \otimes_{U(\mathfrak{h}[t] \oplus \mathbb{C}K)} D,$$

in which the action of x_m on D for $m > 0$ is trivial, K acts by 1 and x_0 acts by x . Then M_D is naturally a H -module. The one dimensional $\mathbb{C}[x]$ -module \mathbb{C} corresponds to H itself.

Let us consider $D = \mathbb{C}[x]/(x^2)$ which is a non trivial extension

$$0 \rightarrow \mathbb{C} \rightarrow D \rightarrow \mathbb{C} \rightarrow 0.$$

The H -module M_D is a self-extension of H and possesses a PBW basis

$$x_{-N}^{\lambda_N} \cdots x_{-1}^{\lambda_1} x_0^{\lambda_0} \mathbf{1} \tag{12.1}$$

in which $\lambda_i \in \mathbb{Z}_{\geq 0}$ for all i and $\lambda_0 \leq 1$. The monomials for which $\lambda_0 = 1$ form a basis of the embedded copy $H \subset M_D$.

12.8. The Zhu algebra $\text{Zhu}(H)$ of H is isomorphic to $\mathbb{C}[x]$. Indeed x here is the image of $x_{-1}\mathbf{1}$ in the quotient that defines $\text{Zhu}(H)$.

12.9. We now consider the higher trace function

$$F_1(a \otimes b \otimes f(t_0, t_1)) = \text{res}_{z_0} z_1 \text{tr}_H \sigma(a, z_0) X(b, z_1) F(z_0, z_1) q^{L_0}$$

associated with the self-extension M_D of H described above, and a cycle $a \otimes b \otimes f(t_0, t_1) \in C_1^{n=1}$. In particular if we put $f(t_0, t_1) = 1$ and $b = \mathbf{1}$ then we have

$$d_1(a \otimes b \otimes \mathbf{1}) = a_{(0)}\mathbf{1} = 0.$$

We may thus consider the higher trace functions

$$F_1(a \otimes \mathbf{1} \otimes \mathbf{1}) = \text{tr}_H \sigma(a)_0 \mathbf{1}_0 q^{L_0} = \text{tr}_H \sigma(a)_0 \mathbf{1}_0 q^{L_0}.$$

Since x_0 is central in $\widehat{\mathfrak{h}}$ it follows that x_0 acts relative to the basis of monomials (12.1) of M_D by increasing $\lambda_0 = 0$ to $\lambda_0 = 1$ and annihilating those monomials for which $\lambda_0 = 1$. Therefore $\sigma(x)_0$ is simply the identity on H . Hence

$$F_1(x \otimes \mathbf{1} \otimes \mathbf{1}) = \text{tr}_H q^{L_0} = \prod_{n=1}^{\infty} \frac{1}{1 - q^n}.$$

13 Conclusions

We constructed a complex that computes the chiral homology in degrees 0 and 1 of an elliptic curve E_τ with coefficients in a vertex algebra V supported at the marked point $0 \in E_\tau$. We generalized these complexes to n -insertions of V at the points $t_1, \dots, t_n \in E_\tau$. As the collection $\{t_i\}$ moves in $\text{Sym}^n E_\tau$ and τ moves in \mathbb{H} the homologies of these complexes acquire a flat connection. We give explicit description of this connection in terms of an insertion of the conformal vector and a Weierstrass ζ function, and showed that this connection has a regular singular point as $\tau \rightarrow i\infty$. We found finiteness conditions on the vertex algebra V to guarantee that the first chiral homology of E_τ with coefficients in V is finite dimensional. We defined the degree 1 analog of the space of conformal blocks, or n -point functions of a

vertex algebra, as a compatible collection of flat sections of the dual of the first chiral homology group with coefficients in n insertions of V . We prove the modular invariance of the space of these higher analogs of n -point functions and show that self-extensions of modules produce such compatible systems of flat sections.

As a technical tool to achieve the above, we proved a version of Borcherds formula, relating two different vertex algebra structures on V , as well as elliptic functions and their Fourier expansions. We believe this theorem will be of independent interest.

There are a number of results which we have not achieved. In degree zero, Zhu proved that the space of irreducible modules of V provides a basis for the dual of zeroth chiral homology under certain finiteness conditions of V . We do not have an analogous statement in degree 1. In the classically free case, we have an isomorphism $H_1^{\text{ch}}(V, q = 0) \xrightarrow{\sim} \text{HH}_1(\text{Zhu}(V))$. It would be interesting to find a way of inducing linear functionals on $\text{HH}_1(\text{Zhu}(V))$ to first conformal blocks on the torus. The analogous statement in degree 0 corresponds to inducing irreducible modules of $\text{Zhu}(V)$ to irreducible modules of V .

Our approach follows Zhu's technique of restricting to the point at infinity $q = 0$ in the moduli space of elliptic curves. More generally Miyamoto [20] has studied the zeroth chiral homology of non-rational, C_2 -cofinite vertex algebras, by using *higher Zhu algebras*. These algebras relate to non-reduced neighborhoods $\text{Spec } \mathbb{C}[q]/(q^n)$ of the point at infinity, as the Zhu algebra does in the case $n = 1$. It would be interesting to see if these algebras play a similar role for chiral homology in degree 1.

There are two lines of research that are natural continuations for this work: higher degree chiral homology and higher genera. The first seems to be an immediate extension of the complexes constructed in this article. What remains to be done is to determine the higher analogs of the linear functionals associated to self-extensions of modules. Well known examples of C_2 -cofinite but non-rational vertex algebras, like the triplet vertex algebra, admit higher self-extensions of modules (but not classes in Ext^1). It would be interesting if these classes generate higher chiral homology classes on elliptic curves.

To generalize this work to higher genera a natural approach is to use the techniques of [8]. As mentioned in the introduction, it would be quite interesting to know if the finiteness conditions found in genus 1 in this article suffice to guarantee finite dimensionality at arbitrary genus.

References

- [1] A. Beilinson and V. Drinfeld. *Chiral Algebras*. Number 51 in Colloquium Publications. American Mathematical Society, 2004.
- [2] Y. Zhu. Modular invariance of characters of vertex operator algebras. *J. Amer. Math. Soc.*, 9(1):237–302, 1996.
- [3] T. Arakawa and K. Kawasetsu. Quasi-lisse vertex algebras and modular linear differential equations. In V. Kac and V. Popov, editors, *Lie groups, geometry and representation theory. Progress in mathematics.*, volume 326. Birkäuser, 2018.
- [4] H. Li. Abelianizing vertex algebras. *Comm. Math. Phys.*, 259:391–411, 2005.
- [5] T. Arakawa. A remark on the C_2 -cofiniteness condition on vertex algebras. *Math. Z.*, 270(1-2):559–575, 2012.
- [6] J. Van Ekeren and R. Heluani. Chiral Homology of Elliptic curves and the Zhu Algebra. *To appear in Communications in Mathematical Physics*, <https://arxiv.org/abs/1804.00017>, 2018.
- [7] E. Frenkel and D. Ben-Zvi. *Vertex algebras and algebraic curves*. Number 88 in Mathematical surveys and monographs. AMS, Providence, RI, 2001.
- [8] C. Damiolini, A. Gibney, and N. Tarasca. Conformal blocks from vertex algebras and their connections on $\overline{\mathbb{P}}_{g,n}$, 2020.
- [9] C. Damiolini, A. Gibney, and N. Tarasca. On factorization and vector bundles of conformal blocks from vertex algebras, 2019.

- [10] A. Lichnerowicz et al. Les variétés de poisson et leurs algebres de lie associées. *Journal of differential geometry*, 12(2):253–300, 1977.
- [11] J. Huebschmann. Poisson cohomology and quantization. *J. reine angew. Math*, 408:57–113, 1990.
- [12] A. Libgober. Elliptic genera, real algebraic varieties and quasi-jacobi forms. *Topology of stratified spaces*, 58:95–120, 2009.
- [13] N. M. Katz. Nilpotent connections and the monodromy. *Publications Mathématiques de l’I.H.É.S*, 39:175–232, 1970.
- [14] A. Haefliger. Local theory of meromorphic connections in dimension one (Fuchs theory). In A. Borel, editor, *Algebraic D-modules*. Academic Press, Inc., 1987.
- [15] J. van Ekeren and R. Heluani. A short construction of the Zhu algebra. *J. Algebra*, 528:85–95, 2019.
- [16] A. Meurman and M. Primc. Annihilating fields of standard modules of sl_2^{\sim} and combinatorial identities. *Mem. Amer. Math. Soc.*, 137(652):viii+89, 1999.
- [17] E. Feigin. The PBW filtration, Demazure modules and toroidal current algebras. *SIGMA Symmetry Integrability Geom. Methods Appl.*, 4:Paper 070, 21, 2008.
- [18] D. Gaitsgory. Notes on 2d conformal theories and string theory. In *Quantum fields and strings: a course for mathematicians*, volume 2, pages 1017–1090. AMS, 1999.
- [19] G. Andrews, J. van Ekeren, and R. Heluani. The singular support of the Ising model. *arXiv:2005.10769*, 2020.
- [20] M. Miyamoto. Modular invariance of vertex operator algebras satisfying c_2 -cofiniteness condition. *Duke Mathematical Journal*, 122:51–91, 2004.