

Design and Analysis of Bipartite Experiments under a Linear Exposure-Response Model

Christopher Harshaw¹, Fredrik Sävje¹,
David Eisenstat², Vahab Mirrokni², and Jean Pouget-Abadie²

¹Yale University
²Google Research

July 21, 2022

Abstract

In a bipartite experiment, units that are assigned treatments differ from the units for which we measure outcomes. The two groups of units are connected by a bipartite graph, governing how the treated units can affect the outcome units. Often motivated by experiments in marketplaces, the bipartite experimental framework has been used for example to investigate the causal effects of supply-side changes on demand-side behavior. In this paper, we consider the problem of estimating the average total treatment effect in the bipartite experimental framework under a linear exposure-response model. We introduce the Exposure Reweighted Linear (ERL) Estimator, an unbiased linear estimator of the average treatment effect in this setting. We show that the estimator is consistent and asymptotically normal, provided that the bipartite graph is sufficiently sparse. We derive a variance estimator which facilitates confidence intervals based on a normal approximation. In addition, we introduce EXPOSURE-DESIGN, a cluster-based design which aims to increase the precision of the ERL estimator by realizing desirable exposure distributions. Finally, we demonstrate the effectiveness of the described estimator and design with an application using a publicly available Amazon user-item review graph.

1 Introduction

Two-sided marketplaces are rife with interesting but difficult causal questions. What happens to demand if shipping times or fees are reduced? What happens to people’s willingness to use ride-hailing apps if more drivers are enrolled in specific cities? What happens to long term user behavior if a hotel booking platform changes its recommendation engine? The causal impact of these changes is hard to quantify, even when using randomized experiments, because marketplace dynamics often violate a central tenet of conventional experimentation: the Stable Unit Treatment Value Assumption, abbreviated SUTVA. This assumption stipulates that the treatment assigned to one unit does not affect any other units. Violations of this assumption is a phenomenon known as interference, which is often present in the case of marketplace experiments and complicates causal analysis.

The bipartite experimental framework offers a useful formalism to study two-sided market experiments and other violations of SUTVA that can happen along the edges of a bipartite graph. This stands in contrast with interference that occurs on graphs where all units are of the same type (e.g. users of a social network). In the bipartite experimental framework, we distinguish two types of units: units that can be subject to an intervention and units whose responses are of interest to the experimenter. We assign treatment to the former and measure the outcomes of the latter. The causal impact of treating one group of units is measured on the other group by tracking the *exposure* to treatment that the latter group receives, informed by the knowledge of the bipartite graph between them. We remark that the treatment status of a single unit may affect the measured outcomes of many units and, likewise, a measured outcome may be affected by many treatment units.

For example, consider a marketplace where buyers compete for limited goods, some of which may be perfectly or partially substitutable. Their demand of these goods form a bipartite graph that potentially can be inferred by the marketplace owner. The owner of the marketplace would like to determine the causal effect of discounting prices on buyers’ marketplace behavior through a randomized experiment. Randomly assigning certain buyers to receive a discounted price is often not possible, and might even be prohibited, in which case randomization is only possible at the item-level. At the same time, simply comparing discounted goods with non-discounted goods runs the risk of severe bias: a discounted good may do well against a non-discounted substitutable good, which does not accurately reflect a world where either both or neither are discounted. Instead, the marketplace owner decides to monitor this change at the buyer level, positing that, by tracking both their behavioral changes and their exposure to discounted goods, the causal effect of the discount can be teased out.

As is done in much of the interference literature and other settings where SUTVA is violated, assumptions on potential outcomes are made when the bipartite graph has a many-to-many structure in order to allow for tractable inference. One such assumption is existence of an exposure mapping, which posits that outcomes are some simple function of the treatment assignments of neighboring units in the bipartite graph (Toulis and Kao, 2013; Aronow and Samii, 2017). In this work, we study estimation of an all-or-nothing treatment effect in the bipartite experimental framework under a linear exposure-response model, where exposures are linear functions of assignments and responses are linear functions of the exposures. The main contributions of this work are summarized as follows:

- We describe the Exposure-Rewighted Linear (ERL) estimator, an unbiased linear estimator of the average total treatment effect under the linear exposure-response model. We show that the ERL estimator is consistent and asymptotically normal, provided the graph remains sufficiently sparse.
- We describe a variance estimator, which may be used to construct confidence intervals via a normal approximation. We show that under mild conditions on the exposure distribution, the variance estimator is unbiased. We achieve unbiasedness without assuming constant treatment effects or any other restrictions on the heterogeneity between units’ potential outcomes.
- We describe EXPOSURE-DESIGN, a cluster-based design which aims to increase the precision of the ERL estimator. The design achieves this by increasing the variance of individual exposures while decreasing the covariance between different exposures. This improves precision in several settings of interest.

By constructing the estimator and design together, we can achieve estimates with higher precision.

Organization The remainder of the paper is organized as follows: We review related work in Section 2. In Section 3, we present the bipartite experimental setting and the linear exposure-response model. We present and analyze the ERL estimator in Section 4 and its corresponding confidence intervals in Section 5. We present our cluster-based design EXPOSURE-DESIGN in Section 6. Finally, we apply this methodology on the publicly available Amazon product review graph are presented in Section 7 and we conclude in Section 8.

2 Related Works

Within the wide-ranging causal inference literature, our work falls squarely within the subset relying on the potential outcomes framework (Neyman, 1923; Imbens and Rubin, 2015). The design and analysis of randomized experiments in the presence of interference has garnered plenty of attention, spanning vaccination trials (Struchiner et al., 1990), agricultural studies (Kempton, 1997), voter-mobilization field experiments (Sinclair et al., 2012), and viral marketing campaigns (Aral and Walker, 2011; Eckles et al., 2016b). It is beyond the scope of the current paper to extensively review the literature causal inference with interference. Instead, we direct readers to the review article by Halloran and Hudgens (2016).

Our work is primarily motivated by marketplace experiments. Evidence of interference in marketplaces has been noted across industries for various experimental designs (Gupta et al., 2019). Reiley (2006), Einav et al. (2011) and Holtz et al. (2020) study the interference bias that results from supply-side randomization, while Blake and Coey (2014) and Fradkin (2015) consider this problem in the context of demand/user-side

randomization. Basse et al. (2016) and Liu et al. (2020) compare supply-side randomization to two-sided randomization as well as to budget-split designs, showing bias can be reduced in the context of certain ad auction experiments. More recently, Johari et al. (2020) characterize which randomization scheme (supply-side, demand-side, or two-sided) leads to reduced bias as a function of market balance.

We consider a slightly different experimental setting, introduced by Zigler and Papadogeorgou (2021), characterized by random assignment of treatment on one side of the bipartite graph (demand- or supply-side), while outcomes are measured on the other side. The advantage of this framework is that complex interference relationships can be captured by an exposure function (similar to Aronow and Samii (2017)), which is assumed to solely determine an unit’s outcome. This makes the problem of estimating causal effects tractable despite the complex interference structure. Zigler and Papadogeorgou (2021) study causal estimands which are more closely related to direct effects rather than the all-or-nothing treatment effect considered here. In addition, the analysis of their estimators requires that the bipartite graph be the union of small connected components.

Motivated by marketplace experiments, Pouget-Abadie et al. (2019) introduce a cluster-based design for general bipartite graphs in this framework and consider a similar estimand and exposure-response assumption. Later, Doudchenko et al. (2020) proposed a class of generalized propensity score estimators for this framework, which are unbiased for both experimental and observational settings under standard assumptions and a similar exposure-response assumption.

Our work is the first to propose methods for provably valid inference (e.g., confidence intervals) in the bipartite settings and to jointly consider estimators and designs which improve overall precision of treatment effect estimators. While the cluster design of Pouget-Abadie et al. (2019) is based on the intuition of achieving a large spread of exposures, it disregards the correlation of exposures and is not rigorously tied to the performance of an estimator. Additionally, while the estimators proposed by Doudchenko et al. (2020) are unbiased, they are based on a different approach which requires fitting a generalized propensity score function. Neither of these papers present methods for valid inference.

3 Experimental Setting

In the bipartite experimental framework, the units which receive treatment are distinct from the units on which the outcomes are measured. For example, Zigler and Papadogeorgou (2021) apply the framework to analyze how interventions on power plants’ pollution affect the hospitalization rates among nearby hospitals. We discuss the general bipartite framework in Section 3.1 and the linear exposure response assumption in Section 3.2.

3.1 Bipartite Experiments

In the bipartite experiment setting, there are two groups of units: the *diversion units*, to which treatment is applied, and the *outcome units*, where outcomes are measured. We denote the set of m diversion units by V_d and the set of n outcome units by V_o .

Each of the m diversion units receives a (random) binary treatment $z_i \in \{\pm 1\}$, and we collect these treatments into a treatment vector, $\mathbf{z} = (z_1, z_2, \dots, z_m) \in \{\pm 1\}^m$. The distribution over the random treatment vectors is called the *design* of the experiment and it is chosen by the experimenter. Each of the outcome units $i \in V_o$ is associated with a potential outcome function $Y_i(\mathbf{z})$, which maps the treatment assignments to the observed value, which is a real number. In the bipartite setting, we assume that each potential outcome function depends only on the treatment of a neighborhood set of diversion units. More formally, there exists a *neighborhood mapping* $\mathcal{N} : V_o \rightarrow 2^{V_d}$ such that for all outcome units $i \in V_o$,

$$Y_i(\mathbf{z}) = Y_i(\mathbf{z}') \quad \text{if } z_j = z'_j \text{ for all } j \in \mathcal{N}(i) .$$

Throughout the paper, we assume that the neighborhood mapping is known and correctly specified, so that the above condition holds. We recover the standard Stable Unit Treatment Value Assumption (SUTVA) when the diversion units are identified with the outcome units and the neighborhood mapping is the identity function.

The number of potential outcomes for each outcome unit grows exponentially in the size of its neighborhood. Zigler and Papadogeorgou (2021) avoid this issue by assuming that the bipartite structure is the union of many small connected components. Unfortunately, this is typically not a reasonable assumption in the marketplace settings where we know that more varied interactions occur: buyers may interact with a variety of products. Without further restrictions on the structure of the neighborhoods or the potential outcome functions, inference of any causal estimand is impossible (Basse and Airolidi, 2018; Sävje et al., 2021). Take, for example, an instance where the neighborhood of each outcome unit is all diversion units. For this reason, we introduce a stronger assumption on the potential outcomes.

3.2 Linear Exposure-Response Model

In order to admit tractable inference of causal estimands, we consider a linear exposure-response model, which consists of two underlying assumptions: a linear exposure assumption and a linear response assumption, which we state formally below.

In the linear exposure-response model, we suppose that there is a weighted bipartite graph between diversion units and outcomes units, where the edges have non-negative weights $w_{i,j} \geq 0$, which we arrange into an n -by- m incidence matrix \mathbf{W} . An edge $w_{i,j}$ represents the influence of diversion unit j on the outcome units i . We say that outcome unit i and diversion unit j are *incident* if the weight $w_{i,j}$ is positive. The degree of a diversion unit is the number of outcome units it is incident to, and the largest degree among all diversion units is denoted d_d . The degree of an outcome unit is defined similarly and the largest degree among all outcome units is denoted d_o . For simplicity, we assume that each outcome unit has degree at least 1 and the weights incident to an outcome unit are normalized to sum to one, i.e. the rows of the incidence matrix \mathbf{W} sum to one. We also assume that this weighted bipartite graph is known to the experimenter. In many market experiments, the experimenter may construct an approximation of this graph from historical data.

The *linear exposure assumption* is that the treatment assignments influences the potential outcomes only through a linear combination, which is more structured than arbitrary influence. More formally, for each outcome unit $i \in V_o$, the *exposure* of outcome unit i is

$$x_i(\mathbf{z}) = \sum_{j \in V_d} w_{i,j} z_j ,$$

and for all pairs of assignment vectors \mathbf{z} and \mathbf{z}' with $x_i(\mathbf{z}) = x_i(\mathbf{z}')$, we have that $Y_i(\mathbf{z}) = Y_i(\mathbf{z}')$. This implies that the neighborhood mapping is such that $\mathcal{N}(i) = \{j : w_{i,j} > 0\}$.

We arrange these n exposures into an exposure vector $\mathbf{x}(\mathbf{z}) = (x_1(\mathbf{z}), x_2(\mathbf{z}) \dots x_n(\mathbf{z}))$. Because the exposure is a function of treatment, the experimental design completely determines the exposure distribution. This linear exposure assumption is a generalization of the partial and stratified interference assumptions discussed by Hudgens and Halloran (2008). When the treatment assignment vector \mathbf{z} is clear from context, we write simply x_i and \mathbf{x} for the i th exposure and the exposure vector, respectively. Using matrix-vector notation, we may write the exposure vector as $\mathbf{x}(\mathbf{z}) = \mathbf{W}\mathbf{z}$. Due to the normalization of the weights and the ± 1 values of the treatment assignment, each exposure takes values in the range $[-1, 1]$.

The *linear response assumption* is that for each outcome unit, the potential outcome is a linear function of its exposure. That is, for each outcome unit $i \in V_o$, there exists parameters α_i and β_i such that

$$Y_i(\mathbf{z}) = \alpha_i + \beta_i x_i(\mathbf{z}) .$$

We refer to α_i as the unit-specific intercept and β_i as the unit-specific slope. These terms are unknown to the experimenter, and the experimenter only observes the outcome $Y_i(\mathbf{z})$, along with the sampled assignment vector \mathbf{z} and the resulting exposure vector \mathbf{x} . Similar assumptions have been proposed in the presence of interference by Manski (1993); Bramoullé et al. (2009); Baird et al. (2018); Chin (2019b); Offer-Westort and Dimmery (2021).

We refer to the *linear exposure-response model* as the combination of the linear exposure assumption and the linear response assumption. The linear exposure-response model places certain limits on the potential outcomes, but allows for more complex structure in the bipartite graph. This trade-off is preferable in settings such as marketplace experiments, where we know that a complex bipartite structure exists and

we are more comfortable with making simplifying assumptions about potential outcomes. Note that the bipartite setting with the linear exposure-response assumption recovers the standard SUTVA setting when diversion units are identified with the outcome units and the weight matrix is the identity.

From one perspective, the linear exposure-response model is a strong assumption. It requires that the response for each unit is exactly a linear function in the exposure. This rules out, for example, that different diversion units have different impacts on a single outcome unit. But from another perspective, the model is completely unrestrictive: it does not limit the heterogeneity between units at all. That is, knowing the response function for one unit tells us nothing about the response function of other units. While there are few settings in which the linear exposure-response model will hold exactly, it will often be a useful approximation given its unrestrictiveness with respect to heterogeneity. We leave it to future work to characterize the behavior of estimator when the model holds approximately, and we consider it holding exactly for the rest of the paper.

Although the experimenter is free to choose any treatment design, the exposure distribution is necessarily constrained in certain ways by the bipartite weight matrix, as $\mathbf{x} = \mathbf{W}\mathbf{z}$. For example, if two outcome units $i, j \in V_o$ have exactly the same incident weights, then their exposures x_i and x_j will always be equal in value, regardless of treatment assignment. In this case, the exposures x_i and x_j will be perfectly correlated. Thus, it is challenging or impossible to construct a treatment design which realizes some ideal exposure distribution and this task depends on the bipartite weight matrix itself. This is a central challenge in the bipartite linear exposure-response model, which is both statistical and computational in nature. We address this challenge in Section 6 when we propose our cluster-based treatment design.

3.3 Causal Estimand

We are interested in understanding the contrast between two possible worlds: one where all diversion units receive treatment and one where they all receive control. For an individual outcome unit, this contrast is captured by the individual treatment effect, $\tau_i = Y_i(\mathbf{z} = \mathbf{1}) - Y_i(\mathbf{z} = -\mathbf{1})$ for $i \in V_o$. Just as in the typical SUTVA setting, we cannot hope to estimate the individual treatment effects well because only one potential outcome is observed for any one unit. In light of this, we opt to estimate an aggregated causal quantity. In this paper, we are interested in the Average Total Treatment Effect (ATTE), which is the average contrast between the scenario that all diversion units receive treatment and all diversion units receive control. More precisely, ATTE is defined as

$$\tau = \frac{1}{n} \sum_{i=1}^n \tau_i = \frac{1}{n} \sum_{i=1}^n \left[Y_i(\mathbf{z} = \mathbf{1}) - Y_i(\mathbf{z} = -\mathbf{1}) \right]$$

Under the linear exposure-response model, the ATTE is proportional to the average of the slope terms, as shown in the following proposition.

Proposition 1. *Under the linear exposure-response assumption, the ATTE is $\tau = \frac{2}{n} \sum_{i=1}^n \beta_i$.*

Proof. The individual treatment effect of outcome unit i is proportional to its slope, as

$$\tau_i = Y_i(\mathbf{z} = \mathbf{1}) - Y_i(\mathbf{z} = -\mathbf{1}) = [\beta_i x_i(\mathbf{z} = \mathbf{1}) + \alpha_i] - [\beta_i x_i(\mathbf{z} = -\mathbf{1}) + \alpha_i] = 2\beta_i ,$$

where we have used that $x_i(\mathbf{z} = \mathbf{1}) = 1$ and $x_i(\mathbf{z} = -\mathbf{1}) = -1$. The result follows by taking the average of the individual treatment effects. \square

There are two main challenges in estimating the ATTE in this setting: we want to estimate the average of the slopes of many different linear response functions, but only one point from each of the distinct linear response functions is observed. Although stated in somewhat unfamiliar terms, this is the fundamental problem of causal inference (Holland, 1986). The second challenge is that of constructing a treatment design which realizes a desirable exposure distribution. As previously discussed at the end of Section 3.2, this is a difficult task when the bipartite weight matrix has non-trivial overlapping structures. In the remainder of the paper, we focus on addressing these two challenges by developing an estimator and a class of designs which together accurately estimate the ATTE.

3.4 Cluster Designs

Some of the analysis in this paper assumes that the treatment is assigned according to a *independent cluster designs*, where the diversion units are grouped into clusters and treatment is assigned to an entire cluster. More formally, we say that a partition C_1, C_2, \dots, C_ℓ of the diversion units is a *clustering*, which we denote as $\mathcal{C} = \{C_1, C_2, \dots, C_\ell\}$. That is, all clusters are disjoint and the union of all clusters is set of diversion units V_d . Given a clustering \mathcal{C} , a treatment assignment from the corresponding *independent cluster design* is drawn in the following way: independently for each cluster, we assign all diversion within a cluster to have either treatment $z_i = 1$ with probability p and treatment $z_i = -1$ with probability $1 - p$. For notational simplicity, we consider the treatment probability p to be fixed for all clusters, but our results extend to the setting where each cluster has its own treatment probability. Note that the class of independent cluster designs is completely specified by \mathcal{C} and p .

4 The Exposure Reweighted Linear Estimator

We describe the Exposure Reweighted Linear Estimator, which is an unbiased estimate of the ATTE under the linear exposure-response assumption. The Exposure Reweighted Linear (ERL) estimator is defined below as

$$\hat{\tau} = \frac{2}{n} \sum_{i=1}^n Y_i(\mathbf{z}) \left(\frac{x_i(\mathbf{z}) - \mathbb{E}[x_i(\mathbf{z})]}{\text{Var}(x_i(\mathbf{z}))} \right). \quad (1)$$

The ERL estimator requires knowledge of the mean and variance of each of the marginal exposure distributions under the treatment design. For several commonly used designs such as Bernoulli and independent cluster designs, these exposure characteristics may be computed directly; however, for arbitrary designs, the expectation and variance of the exposures may need to be estimated to high precision using samples drawn from the treatment design. We assume here that these exposure characteristics are known exactly. We emphasize that the ERL estimator may be used under any treatment design and not just the cluster-based treatment design we propose in Section 6.

The ERL estimator belongs to the class of linear estimators, as it is a (random) linear function of the observed outcomes. It shares similarities with the style of Horvitz–Thompson estimators (Narain, 1951; Horvitz and Thompson, 1952), but is not the same. The Horvitz–Thompson estimators re-weights an outcome by the probability of observing that outcome, while the ERL estimator re-weights an outcome by the normalized distance of the exposure from its mean. When there are many exposures, such as under the linear exposure-response model, the type of re-weighting done by the Horvitz–Thompson estimator would lead to excessively large variance.

4.1 Statistical Analysis of the ERL Estimator

In this section, we analyze the behavior of the ERL estimator as a point estimator of the average total treatment effect (ATTE). First, we show that the ERL estimator is unbiased. Then we show consistency and asymptotic normality of the ERL estimator, provided that the bipartite graph is not too dense. Theorem 1 below ensures that under mild conditions on the treatment design, there is no systematic bias in the ERL estimator.

Theorem 1 (Unbiasedness). *Suppose the design is such that each exposure has a positive variance. Under the linear response assumption, the ERL estimator is unbiased for the ATTE: $\mathbb{E}[\hat{\tau}] = \tau$.*

Proof. By linearity, the expectation of the estimator is

$$\mathbb{E}[\hat{\tau}] = \frac{2}{n} \sum_{i=1}^n \mathbb{E} \left[Y_i(\mathbf{z}) \left(\frac{x_i(\mathbf{z}) - \mathbb{E}[x_i(\mathbf{z})]}{\text{Var}(x_i(\mathbf{z}))} \right) \right].$$

By Proposition 1, the ATTE is twice the average of the slope terms β_i . Thus, to complete the proof we show that each expectation terms inside the sum is equal to the corresponding slope β_i . Using the linear response

assumption,

$$\begin{aligned}
\mathbb{E} \left[Y_i(\mathbf{z}) \left(\frac{x_i(\mathbf{z}) - \mathbb{E}[x_i(\mathbf{z})]}{\text{Var}(x_i(\mathbf{z}))} \right) \right] &= \mathbb{E} \left[(\beta_i x_i(\mathbf{z}) + \alpha_i) \left(\frac{x_i(\mathbf{z}) - \mathbb{E}[x_i(\mathbf{z})]}{\text{Var}(x_i(\mathbf{z}))} \right) \right] \\
&= \beta_i \mathbb{E} \left[x_i(\mathbf{z}) \left(\frac{x_i(\mathbf{z}) - \mathbb{E}[x_i(\mathbf{z})]}{\text{Var}(x_i(\mathbf{z}))} \right) \right] + \alpha_i \mathbb{E} \left[\left(\frac{x_i(\mathbf{z}) - \mathbb{E}[x_i(\mathbf{z})]}{\text{Var}(x_i(\mathbf{z}))} \right) \right] \\
&= \beta_i \left(\frac{\mathbb{E}[x_i(\mathbf{z})^2] - \mathbb{E}[x_i(\mathbf{z})]^2}{\text{Var}(x_i(\mathbf{z}))} \right) + \alpha_i \left(\frac{\mathbb{E}[x_i(\mathbf{z})] - \mathbb{E}[x_i(\mathbf{z})]}{\text{Var}(x_i(\mathbf{z}))} \right) \\
&= \beta_i
\end{aligned}$$

□

Next, we analyze the asymptotic behavior of the ERL estimator. In the asymptotic analysis, we suppose that there is a sequence of bipartite experiments, in which the number of units is growing to infinity. Strictly speaking, all quantities of the experiment such as the bipartite graph, the outcomes, the treatment design, etc, should be indexed by an integer N ; however, we drop these subscripts for notational clarity.

We make two additional assumptions about the bipartite experiments in this asymptotic sequence. The first is that the potential outcomes are bounded. The second is that the design has limited dependence between treatment assignments.

Assumption 1 (Bounded Potential Outcomes). *The potential outcomes are bounded in absolute value $|Y_i(\mathbf{z})| \leq M$, where M is a constant.*

Assumption 2 (Design Conditions). *The treatments assignments are distributed according to an independent cluster design, where the probability of treatment p is bounded away from 0 and 1 by a constant in the asymptotic sequence. Additionally, the sizes of clusters are bounded by k , a constant in the asymptotic sequence.*

Assumption 2 rules out certain classes of treatment designs, such as complete randomization (i.e. group balanced designs). While it may be possible to obtain similar asymptotic results under such designs, we limit our consideration to those satisfying Assumption 2. Under these assumptions, we prove that ERL is consistent when the bipartite graph is not too dense.

Theorem 2 (Consistency). *Under Assumptions 1 and 2, and supposing that $d_d d_o^3 = o(n)$ in the asymptotic sequence, the ERL estimator converges in mean square to the ATTE: $\lim_{N \rightarrow \infty} \mathbb{E}[(\hat{\tau} - \tau)^2] = 0$.*

The main technical assumption that we require for consistency is that $d_d d_o^3 = o(n)$ in the asymptotic sequence, where we recall that d_d and d_o are the maximum degrees of the diversion and outcome units, respectively. Informally, this condition ensures that the bipartite graph is not too dense as it grows. While consistency may hold under a weaker condition for a particular design, some density assumption like this must be made. As an example, consider the complete bipartite graph where all outcome units receive the same exposure, in which case consistent estimation is impossible.

We now discuss a setting where this condition $d_d d_o^3 = o(n)$ holds. Suppose that each diversion unit has fixed degree d_d , which is a constant with respect to m and n . The average degree of an outcome unit is then $\bar{d}_o = d_d(m/n)$. Assuming that the maximum outcome degree d_o is within a constant factor of the average, this yields that the term $d_d d_o^3 = \mathcal{O}(d_d^4(m/n)^3)$. Using that the diversion degrees are constant, we get that this term is bounded by $o(n)$ if $m = o(n^{4/3})$. Thus, in graphs with constant diversion degrees where the edges are evenly distributed between outcome units, the hypothesis of Theorem 2 holds when m grows at a rate slower than $n^{4/3}$.

We next describe the asymptotic distribution of the estimator. In particular, we show that the ERL estimator converges in distribution to a normal distribution as the size of the bipartite experiment grows, provided that the graph remains sparse. This result is derived under the same asymptotic regime as above. In order to prove the central limit theorem, we require an additional assumption on the asymptotic sequence of bipartite experiments. Namely, we require that the variance of the ERL estimator decreases no faster than the parametric rate.

Assumption 3 (Not Superefficient). *The normalized variance of the ERL estimator $n \text{Var}(\hat{\tau})$ is bounded away from zero asymptotically.*

Assumption 3 rules out settings in which we can estimate the ATTE at a faster than parametric rate. Such settings are theoretically possible, but not practically relevant. In particular, Assumption 3 rules out two scenarios. The first is when the magnitude of the potential outcomes approaches zero in the asymptotic sequence. Note that this requires that almost all potential outcomes approach zero; the magnitude of the potential outcomes are generally non-zero even when their average is zero. The second scenario is when the design close to perfectly pinpoints the potential outcomes. This can be formalized as the variance of each individual term of the estimator diminishes asymptotically, i.e. $\text{Var}(\hat{\tau}_i) \rightarrow 0$, where $\hat{\tau}_i = 2Y_i(\mathbf{z})(x_i - \mathbb{E}[x_i]) / \text{Var}(x_i)$. Both of these scenarios are knife-edge cases that we have good reason to believe would not materialize in practice. Even if they do, the estimator would still be unbiased and consistent, but its asymptotic distribution might not be normal.

We are now ready to present a central limit theorem which states that under mild regularity conditions, the ERL estimator is asymptotically normal.

Theorem 3 (Asymptotic Normality). *Under Assumptions 1, 2, and 3, and supposing that $d_d^{1.6} d_o^4 = o(n)$, the ERL estimator is asymptotically normal:*

$$\frac{\hat{\tau} - \tau}{\sqrt{\text{Var}(\hat{\tau})}} \xrightarrow{d} \mathcal{N}(0, 1) .$$

The proof relies on Stein’s method for bounding distances between distributions (see, e.g. Ross (2011)). We use Stein’s method because standard techniques for establishing central limit theorems which heavily rely on independence are not applicable in the bipartite experimental framework where exposures are necessarily correlated. We remark that Stein’s method has been recently used for obtaining limiting behavior of other estimators in the interference literature (Aronow and Samii, 2017; Chin, 2019a; Ogburn et al., 2020).

The assumptions on the asymptotic growth of the bipartite graph may be interpreted similarly as those appearing in Theorem 2. Namely, they prevent the bipartite graph from becoming too dense. We remark that the growth assumptions required for asymptotic normality (Theorem 3) are stronger than those required for consistency (Theorem 2). The growth assumptions in Theorem 3 are only sufficient and we conjecture that they are not necessary for asymptotic normality. However, weakening these growth conditions would require a different analysis, either by a more careful application of Stein’s method or by different means all together.

Assumption 2 allows for a broad class of designs. For example, unit-level Bernoulli randomization falls into this class, but this design does not at all consider the structure of the bipartite graph and will generally perform poorly. To derive analytical results for this broad class of designs, the growth conditions on the bipartite graph are quite restrictive, and they may be too restrictive in certain settings where more dense interaction patterns occur. If one restricts focus to a smaller class of designs, these growth conditions could potentially be weakened. The key implication of Assumption 2 together with the growth conditions is that the variance of the exposures is large and the correlation between most pairs of exposures is small. Heuristically, these conditions on the exposure distribution are the main aspects required for consistency and normality. We describe a design in Section 6 that directly targets the exposure distribution to satisfy these conditions, and it will therefore be better behaved than many of the designs allowed by Assumption 2.

5 Variance Estimation

In this section, we present methods for constructing confidence intervals for the ATTE in the bipartite setting under the linear exposure-response assumption. If we knew the variance of the ERL estimator, we could use Theorem 3 directly to construct asymptotically valid confidence intervals. Because the variance of the ERL estimator depends on the unobserved potential outcomes, we must construct an estimator of the variance.

In the typical experimental settings with SUTVA and binary treatments, unbiased variance estimation is not possible without strong assumptions on the heterogeneity between units (Imbens and Rubin, 2015). In light of this negative result, experimenters tend to favor conservative variance estimators that lead to valid but overly wide confidence intervals. In contrast to the typical SUTVA setting with binary treatments,

we show that unbiased variance estimation is possible in the bipartite setting under the linear response assumption when the exposures take many (i.e. non-binary) values.

Our approach to constructing a variance estimator is to decompose the ERL estimator into a weighted average of individual effect estimators, and to decompose the variance of the ERL estimator as the average of the variances and covariances of these individual effect estimators. To this end, define $\hat{\tau}_i \triangleq 2Y_i(\mathbf{z})(x_i - \mathbb{E}[x_i]) / \text{Var}(x_i)$ to be the individual terms in the ERL estimator. We may interpret $\hat{\tau}_i$ as an unbiased, but very imprecise, estimator of the individual treatment effect τ_i . The ERL estimator can be written as the average of these quantities: $\hat{\tau} = (1/n) \sum_{i=1}^n \hat{\tau}_i$. The variance of the ERL estimator may be decomposed as

$$\text{Var}(\hat{\tau}) = \text{Var}\left(\frac{1}{n} \sum_{i=1}^n \hat{\tau}_i\right) = \frac{1}{n^2} \sum_{i=1}^n \left[\text{Var}(\hat{\tau}_i) + \sum_{j \neq i} \text{Cov}(\hat{\tau}_i, \hat{\tau}_j) \right].$$

We will construct unbiased estimators of $\text{Var}(\hat{\tau}_i)$ and $\text{Cov}(\hat{\tau}_i, \hat{\tau}_j)$ and average over these estimators to obtain an estimator of the variance of the ERL estimator. Because we are averaging over many unbiased estimators, the average will often be well-behaved even if the individual estimators perform poorly in mean squared sense.

We begin by deriving an estimator for the individual variance terms, $\text{Var}(\hat{\tau}_i)$. To this end, we define the random variable

$$Q_i = \frac{(x_i - \mathbb{E}[x_i])^2}{\text{Var}(x_i)^2} - \frac{\text{Var}(x_i)(x_i^2 - \mathbb{E}[x_i^2]) - \text{Cov}(x_i, x_i^2)(x_i - \mathbb{E}[x_i])}{\text{Var}(x_i) \text{Var}(x_i^2) - \text{Cov}(x_i, x_i^2)^2},$$

which is a quadratic function of the exposure x_i . Because the exposure distribution is known to the experimenter, Q_i is an observable quantity. The following lemma demonstrates that by re-weighting the observed quantity $Y_i(\mathbf{z})^2$ by Q_i , we obtain an unbiased estimate of the individual variance terms.

Lemma 1. *Fix an outcome unit $i \in V_o$. If the exposure x_i takes at least three values with non-zero probability, then the variance of unit i 's individual treatment effect estimator is equal to*

$$\text{Var}(\hat{\tau}_i) = 4 \cdot \mathbb{E}[Y_i(\mathbf{z})^2 Q_i].$$

We remark that the condition that the support of the exposure contains at least three points is critical for Lemma 1 to go through; in fact, when the exposure takes only binary values, then previously established results in the SUTVA setting show that these individual variance terms cannot generally be estimated without bias (Imbens and Rubin, 2015). Moreover, because the exposure distribution is known before the experiment is run, the experimenter may determine whether this condition holds. At the end of this section, we discuss variance estimation when this support condition on the exposures does not hold.

We estimate the covariance terms $\text{Cov}(\hat{\tau}_i, \hat{\tau}_j)$ in a similar manner. For each pair of outcome units $i, j \in V_o$, we define the random variable

$$R_{i,j} = \frac{(x_i - \mathbb{E}[x_i])(x_j - \mathbb{E}[x_j])}{\text{Var}(x_i) \text{Var}(x_j)} - \frac{a_{i,j}(x_i x_j - \mathbb{E}[x_i x_j]) + b_{i,j}(x_i - \mathbb{E}[x_i]) + c_{i,j}(x_j - \mathbb{E}[x_j])}{\Psi_{i,j}},$$

which is a bivariate degree two polynomial in the exposures x_i, x_j where the coefficients $a_{i,j}, b_{i,j}, c_{i,j}$, and $\Psi_{i,j}$ depend on the joint distribution of the pair of exposures. More precisely, these coefficients are defined as

$$\begin{aligned} a_{i,j} &= \text{Var}(x_i) \text{Var}(x_j) - \text{Cov}(x_i, x_j)^2, \\ b_{i,j} &= \text{Cov}(x_i, x_j) \text{Cov}(x_i x_j, x_j) - \text{Var}(x_j) \text{Cov}(x_i x_j, x_i), \\ c_{i,j} &= \text{Cov}(x_i, x_j) \text{Cov}(x_i x_j, x_i) - \text{Var}(x_i) \text{Cov}(x_i x_j, x_j), \\ \Psi_{i,j} &= \text{Var}(x_i x_j) (\text{Var}(x_i) \text{Var}(x_j) - \text{Cov}(x_i, x_j)^2) - \text{Var}(x_i) \text{Cov}(x_i x_j, x_j)^2 \\ &\quad - \text{Var}(x_j) \text{Cov}(x_i x_j, x_i)^2 + 2 \text{Cov}(x_i, x_j) \text{Cov}(x_i x_j, x_j) \text{Cov}(x_i x_j, x_i). \end{aligned}$$

We remark that these coefficients depend only on the joint distribution of pairs of exposures and so they are known to the experimenter. Thus, just as Q_i is an observable quantity, so too is $R_{i,j}$. The following lemma demonstrates that reweighting the product of two potential outcomes $Y_i(\mathbf{z})Y_j(\mathbf{z})$ by $R_{i,j}$ yields an unbiased estimate of the individual covariance terms.

Lemma 2. Fix a pair of outcome units $i \neq j \in V_o$. If $\Psi_{i,j} \neq 0$, then the covariance between individual treatment effect estimates $\hat{\tau}_i$ and $\hat{\tau}_j$ may be expressed as

$$\text{Cov}(\hat{\tau}_i, \hat{\tau}_j) = 4 \cdot \mathbb{E}[Y_i(\mathbf{z})Y_j(\mathbf{z})R_{i,j}] .$$

The condition that $\Psi_{i,j}$ is nonzero ensures that the re-weighting factor $R_{i,j}$ is well-defined. One situation in which $\Psi_{i,j}$ is zero is when $x_i = x_j$ with probability one: the condition that $\Psi_{i,j}$ is nonzero rules out this scenario. We have not been able to construct a joint distribution for which the denominator $\Psi_{i,j}$ is zero without perfectly correlated exposures. We conjecture that no such distribution exists, but we have not been able to prove this yet. Regardless, the exposure distribution is known, so the experimenter can determine whether $\Psi_{i,j} = 0$ for any pairs of outcome units. At the end of this section, we discuss how one can proceed in the case that $\Psi_{i,j} = 0$ for some pairs of units.

Lemmas 1 and 2 together suggest the following variance estimator:

$$\widehat{\text{Var}}(\hat{\tau}) \triangleq \frac{4}{n^2} \sum_{i=1}^n \left[Y_i(\mathbf{z})^2 Q_i + \sum_{j \neq i} Y_i(\mathbf{z})Y_j(\mathbf{z})R_{i,j} \right] .$$

When the exposure distributions which are symmetric about 0 (e.g. an independent cluster design with treatment probability $p = 1/2$), the random variables Q_i and $R_{i,j}$ simplify considerably as the $\text{Cov}(x_i x_j, x_i)$ and $\text{Cov}(x_i x_j, x_j)$ terms are zero. In this case, these functions may be written simply as

$$Q_i = \frac{x_i^2}{\text{Var}(x_i)^2} - \frac{x_i^2 - \mathbb{E}[x_i^2]}{\text{Var}(x_i^2)} \quad \text{and} \quad R_{i,j} = \frac{x_i x_j}{\text{Var}(x_i) \text{Var}(x_j)} - \frac{(x_i x_j - \mathbb{E}[x_i x_j])}{\text{Var}(x_i x_j)} .$$

For complex designs where closed forms of these quantities are not readily available, these coefficients may be estimated to arbitrary precision via a Monte Carlo procedure (Fattorini, 2006). The following theorem shows that the proposed variance estimator is unbiased.

Theorem 4 (Unbiased Variance Estimator). *Under the conditions in Lemmas 1 and 2, the variance estimator of the ERL point estimator is unbiased, i.e. $\mathbb{E}[\widehat{\text{Var}}(\hat{\tau})] = \text{Var}(\hat{\tau})$.*

When the conditions of Lemmas 1 and 2 do not hold, our proposed variance estimator will be ill-defined or biased. Indeed, it is possible that no unbiased variance estimator exists in such settings (Imbens and Rubin, 2015). In this case, one can replace the problematic $\text{Var}(\hat{\tau}_i)$ and $\text{Cov}(\hat{\tau}_i, \hat{\tau}_j)$ terms, which cannot be estimated directly, with upper bounds that can be estimated. Specifically, if the exposure x_i takes only two values, so Lemma 1 does not hold, then one can replace the variance of the individual treatment effect estimate with the second raw moment: $\text{Var}(\hat{\tau}_i) = \mathbb{E}[\hat{\tau}_i^2] - \mathbb{E}[\hat{\tau}_i]^2 \leq \mathbb{E}[\hat{\tau}_i^2]$, as Aronow and Samii (2013) do when they invoke Young's inequality. We may thus replace the $Y_i(\mathbf{z})Q_i$ terms in our variance estimator with $\hat{\tau}_i^2$. Similarly, if $\Psi_{i,j} = 0$ for some pair of outcome units $i, j \in V_o$ and Lemma 2 does not hold, then one can replace the corresponding covariance term with an upper bound obtained from Cauchy-Schwarz and AM-GM inequalities:

$$\text{Cov}(\hat{\tau}_i, \hat{\tau}_j) \leq \sqrt{\text{Var}(\hat{\tau}_i) \text{Var}(\hat{\tau}_j)} \leq \frac{1}{2} \left(\text{Var}(\hat{\tau}_i) + \text{Var}(\hat{\tau}_j) \right) .$$

Under Assumptions 1 and 2, replacing one of these individual terms in this way leads to a positive bias of the normalized variance estimator on the order $\mathcal{O}(1/n)$. Thus, the variance estimator remains asymptotically unbiased as long as we apply these upper bounds to $o(n)$ terms. Even when the terms can be estimated without bias, it could still be preferable to use the bound here if the denominators of Q_i or $R_{i,j}$ are small, because small denominators will make the estimator imprecise.

We may now use our variance estimator together with the asymptotic normality to construct well-motivated confidence intervals. We may estimate $1 - \alpha$ confidence intervals by

$$\hat{\tau} \pm \Phi^{-1}(1 - \alpha/2) \sqrt{\widehat{\text{Var}}(\hat{\tau})} ,$$

where $\Phi^{-1} : [0, 1] \rightarrow \mathbb{R}$ is the quantile function of the standard normal deviate. It is possible that the proposed variance estimator may take negative values; in particular, this may happen when the true variance

is extremely small (on a normalized scale) or the sample size is small. When the variance estimator takes a negative value, this construction of confidence intervals is not well-defined. We suggest two possible alternatives here: the experimenter may either use the absolute value of the variance estimator under the square root, or the experimenter may use a more conservative variance estimate.

6 A Cluster Design for Targeting Exposure Distribution

In this section, we describe EXPOSURE-DESIGN, an independent cluster design which aims to improve precision of the ERL estimator by constructing a desirable exposure distribution. To this end, we first show in Section 6.1 that increasing the variance of exposures and decreasing the covariance between exposures can lead to improved precision of the ERL estimator in settings of interest. In Section 6.2, we present a clustering objective which aims to achieve such exposure distributions, thereby improving the precision of ERL estimator. Finally, we present a heuristic algorithm for optimizing this clustering objective in Section 6.3.

6.1 An Ideal Exposure Distribution

Like all re-weighted linear estimators, the ERL estimator will incur a large mean squared error when the re-weighting terms are large. In particular, if the variance of an exposure $\text{Var}(x_i(\mathbf{z}))$ is close to zero, the corresponding term of the estimator in (1) will become large, yielding a high mean squared error even though the estimator is unbiased. In general, experimenters should use designs for which the corresponding exposure variances are large.

However, large exposure variances should not be the only property of the exposure distribution that experimenters focus on. Consider a naive design that places equal probability on two treatment vectors: either all diversion units receive treatment ($\mathbf{z} = \mathbf{1}$) or all diversion units receive control ($\mathbf{z} = -\mathbf{1}$). Under this design, all of the exposure variances are 1, which is the largest possible variance. However, we observe either all of the treatment outcomes or all of the control outcomes, but never a mix of the two; in fact, the estimator itself takes only two values. Thus, the ERL estimator will suffer very large MSE under this design, despite the individual exposure variances being as large as possible. This raises the question: how should we construct a design that improve the precision of the ERL estimator?

This is a challenging task, since the precision of the ERL estimator depends on the unobserved outcomes. Indeed, a universally optimal design does not exist (Harshaw et al., 2021). However, we argue that a good heuristic is to construct the design so that the variance of the exposures are large and the covariance between most pairs of exposures are close to zero. As discussed at the end of Section 4.1, a design which directly targets these aspects of the exposure distribution may hope to ensure high precision of the ERL estimator under weaker growth conditions on the bipartite graph than those presented in our analysis.

To further motivate this heuristic, consider the scenario where all of the individual treatment effects are zero, i.e. the response functions are of the form $Y_i(x_i) = \alpha_i$. Studies of these sort are sometimes called uniformity trials or A/A tests. In this scenario, one may derive the MSE of the ERL estimator as

$$\mathbb{E}[(\hat{\tau} - \tau)^2] = \frac{4}{n^2} \left[\sum_{i=1}^m \alpha_i^2 \frac{1}{\text{Var}(x_i)} + 2 \sum_{i < j} \alpha_i \alpha_j \frac{\text{Cov}(x_i, x_j)}{\sqrt{\text{Var}(x_i) \text{Var}(x_j)}} \right].$$

As the individual variance terms increase, the first sum decreases. The effect of the second term depends on the sign of the product of intercepts, $\alpha_i \alpha_j$. Generally speaking, these intercepts are unknown to the experimenter. For the sake of this discussion, consider when the outcomes $Y_i(\mathbf{z})$ are non-negative, in which case all intercepts α_i and their products $\alpha_i \alpha_j$ also are non-negative. In this case, decreasing the correlation between exposures would decrease the second term, leading to an overall decrease in the MSE of the ERL estimator.

6.2 Clustering Objective for Targeting Exposure Distribution

In the previous section, we argued for constructing a design so that the variance of exposures is large and the covariance between most exposures is small. However, as argued in Section 3.2, constructing a treatment distribution which realizes a desired exposure distribution is generally not possible due to overlapping

structures in the bipartite graph. In this section, we present an optimization formulation for an independent cluster design which aims to achieve large exposure variance and small correlations between exposures, to the extent that this is possible given the bipartite graph.

We propose choosing a cluster design which maximizes the following objective function:

$$\max_{\text{clustering } \mathcal{C}} \sum_{i=1}^n \left[\text{Var}(x_i) - \phi \sum_{i \neq j} \text{Cov}(x_i, x_j) \right] \quad (\text{EXPOSURE-DESIGN})$$

The variance and covariance of the exposures above are with respect to the random treatment assignments of the corresponding independent cluster design. The first term in the objective is the sum of the exposure variances, so maximizing this term will encourage large exposure variances. The second term penalizes positive correlation between exposures, and maximizing it encourages small correlation. The correlation penalizing parameter $\phi \geq 0$ controls the relative emphasis between large exposure variances and small exposure correlations. When $\phi = 0$, then the emphasis is placed entirely on increasing individual exposure variance; this is typically undesirable, as the optimal solution is often a single cluster containing all diversion units, which results in the “naive” design where either all diversion units receive treatment or all diversion units receive control. Increasing ϕ places more emphasis on decorrelating exposures.

A key insight to solving the EXPOSURE-DESIGN formulation is that it may be reformulated as a *correlation clustering* problem, which is well-studied in the algorithms literature (Bansal et al., 2002; Swamy, 2004; Charikar et al., 2005). The existing computational understanding of these correlation clustering problems is another reason to use the EXPOSURE-DESIGN objective. The following proposition states the re-formulation of the EXPOSURE-DESIGN objective into the correlation clustering variant, denoted CORR-CLUST.

Proposition 2. *For each pair of diversion units $i, j \in V_d$, define the value $\omega_{i,j} \in \mathbb{R}$ as*

$$\omega_{i,j} = (1 + \phi) \sum_{k=1}^m w_{k,i} w_{k,j} - \phi \left(\sum_{k=1}^m w_{k,i} \right) \left(\sum_{k=1}^m w_{k,j} \right), \quad (2)$$

where $w_{k,i}$ is the weight of the edge between the k th outcome unit and the i th diversion unit. EXPOSURE-DESIGN is equivalent to the following clustering problem:

$$\max_{\text{clusterings } \mathcal{C}} \sum_{C_r \in \mathcal{C}} \sum_{i,j \in C_r} \omega_{i,j}. \quad (\text{CORR-CLUST})$$

Although CORR-CLUST is a variant of the weighted maximization-type correlation clustering problems previously studied in the literature (Charikar et al., 2005; Swamy, 2004), is not equivalent to previously studied formulations in an approximation-preserving sense, as it takes positive and negative values. Given that weighted maximization correlation clustering is NP-Hard (Charikar et al., 2005), it is reasonable to presume that our formulation CORR-CLUST is also computationally hard. However, these computational complexity considerations are beyond the scope of this paper.

We remark that EXPOSURE-DESIGN places no explicit constraint on the number of clusters produced by the clustering algorithm. However, our analysis in Section 4.1 suggests that limiting the cluster sizes, and thereby reducing correlation between exposures, helps to achieve consistency and normality of the ERL estimator. This desirable cluster structure is not captured by the optimization problem itself, but we handle it through our local search heuristic described in Section 6.3. We also remark that the EXPOSURE-DESIGN objective does not directly minimize MSE of the ERL estimator, but should instead be understood as a useful heuristic. Finally, we remark that it is impossible to induce a negative correlation between exposures in the class of independent cluster designs, so the second term of the objective attains its maximum when the exposures are uncorrelated.

The EXPOSURE-DESIGN is conceptually similar to the correlation-clustering based design presented in Pouget-Abadie et al. (2019), but it differs in several key ways. EXPOSURE-DESIGN provides experimenters the flexibility to trade-off larger exposure variances with more de-correlated exposures by setting the parameter ϕ . In contrast, the cluster design of Pouget-Abadie et al. (2019) focuses solely on the exposure variance by maximizing which is referred to as “empirical dose variance” in their paper. As we demonstrate in Appendix C, their objective is equal to ours when the trade-off parameter is set to $\phi = 1/(n - 1)$. In this

sense, their cluster design can be viewed as a specific instance of the more general EXPOSURE-DESIGN, where a greater emphasis is placed on maximizing the exposure variances. More importantly, the EXPOSURE-DESIGN presented in this paper is designed to increase the precision of the ERL estimator, while the correlation-clustering based design of Pouget-Abadie et al. (2019) is motivated by the intuition that extreme exposures are helpful in this setting.

6.3 Local Search Heuristic for Exposure-Design

We now describe a local search heuristic for optimizing EXPOSURE-DESIGN. The local search is initialized with the singleton clustering and iteratively seeks to improve the clustering. In each iteration, the algorithm loops through random pairs of diversion units $i, j \in V_d$ and moves diversion unit j to the cluster currently containing diversion unit i if that change improves the objective value, subject to a user-defined constraint on the clusters. The local search algorithm is presented more formally below as Algorithm 1.

Algorithm 1: Local Search(\mathbf{W}, ϕ, k, T , cluster constraints)

```

1 Initialize singleton clustering  $\mathcal{C} = \{\{1\}, \{2\}, \dots, \{m\}\}$ 
2 for iterations  $t = 1 \dots T$  do
3   Choose a uniformly random permutation  $\pi$  on the diversion units.
4   for diversion units  $i \in \pi$  do
5     Randomly select a diversion unit  $j$  with probability proportional to  $(\mathbf{W}^\top \mathbf{W})_{i,j}$ 
6     Let  $C$  and  $C'$  be the clusters containing diversion units  $i$  and  $j$ , respectively.
7     if moving  $j$  from cluster  $C'$  to  $C$  increases objective value  $\mathcal{E}$  satisfies user-defined constraints
8       then
9         Move diversion unit  $j$  from cluster  $C'$  to  $C$ .
9 return clustering  $\mathcal{C}$ 

```

Given a diversion unit i , we use *wedge sampling* to randomly select another diversion unit j . that is, sampling j proportional to (Cohen and Lewis, 1999). We use the wedge sampling procedure because it encourages picking pairs of units for which $(\mathbf{W}^\top \mathbf{W})_{i,j} = \sum_{k=1}^n w_{k,i} w_{k,j}$ is large, which often results in a large correlation clustering weight $\omega_{i,j}$. Performance improvements are obtained by computing the correlation clustering weights $\omega_{i,j}$ only when they are needed to evaluate changes in the objective. In particular, the first term of (2) is an inner product whose computation scales with the sparsity of the bipartite graph and the second term is the product of sums which may be pre-computed.

Diversion unit j is moved into the cluster containing diversion unit i if two conditions are met: the objective increase and the user-defined cluster constraints are satisfied. We recommend that experimenters choose constraints which limit the cluster sizes in some way. For example, the experimenter may choose to constraint the number of diversion units within a cluster. In our implementation, we constrain the sum of the (unweighted) degrees of diversion units within a cluster to be a fixed fraction of the total number of edges. In this way, no cluster has too many outgoing edges to outcome units. Constraining the clusters in this way implicitly limits the amount of dependence between exposures, which is one of the key aspects underlying the design conditions in Assumption 2 of our analysis.

This local search algorithm is different from the one presented in Pouget-Abadie et al. (2019), which approximates the Gram matrix $\mathbf{W}^\top \mathbf{W}$ offline as the sum of a sparse matrix and a rank-one matrix, so that the algorithm works with an approximation to the objective. In contrast, our algorithm accepts and rejects changes based on the exact value of the objective. Relative to Elsner and Schudy (2009), this local search does not consider moving units to new empty clusters, nor does it consider merging clusters. Moves of the first type seem consistently unprofitable in our setting. As for merges, we find that the algorithm is able to essentially perform them by moving one diversion unit at a time.

7 An Application to Online Marketplace Experiments

In this section, we apply our proposed methodology to a simulated marketplace experiment based on a product review dataset from the Amazon marketplace (McAuley et al., 2015; He and McAuley, 2016). The

Amazon product review dataset contains 83 million reviews made by 121 thousand customers on 9.8 million items. In this application, we imagine running an experiment where we change the pricing mechanism of items in the marketplace, and are interested in how a customer’s reported satisfaction is affected by this change in pricing mechanism. The items sold in the marketplace are the diversion units and the customers in the marketplace are the outcome units. An edge is present in the bipartite graph if a customer reviewed an item and all edges incident to an outcome unit are uniformly weighted. Thus, a customer’s exposure is the unweighted average of the treatment status of the items they have previously reviewed.

In our simulated marketplace experiment, we generate potential outcomes via an exposure-response function. The outcomes themselves are the satisfaction score of a customer given their exposure; the responses in this study are simulated, but we can imagine that they are either reported directly by a customer or constructed based upon text analysis of the customer’s review. In the case of a linear response, a positive slope indicates an increase in customer satisfaction as a result of the new pricing mechanism, while a negative slope indicates a decrease in satisfaction as a result of the new pricing mechanism.

We preprocess the Amazon produce review dataset for computational tractability in the same manner as Pouget-Abadie et al. (2019). We begin by removing customers that have reviewed fewer than 100 items. Next, we execute a balanced partitioning algorithm (Aydin et al., 2019) on the entire bipartite graph to create groups of customers and groups of items. After this preprocessing, we define the diversion units to be the item groups and the outcome units to be the customer groups. The resulting bipartite graph has 1 thousand outcome units, 2.4 million diversion units, and 7.1 million edges. We emphasize that this bipartite graph does not satisfy the growth conditions (specified in Section 4.1) required for consistency and normality under the broad class of designs captured by Assumption 2. In this sense, this application may be considered a test of the efficacy of the proposed EXPOSURE-DESIGN under weaker growth conditions.

We investigate the statistical properties of the ERL estimator, the variance estimator, and the resulting confidence intervals under various treatment designs in this application. In particular, we compare our proposed EXPOSURE-DESIGN to several existing designs: the Bernoulli design, the correlation clustering design of Pouget-Abadie et al. (2019), and the balanced partitioning cluster design of Eckles et al. (2016a), as implemented by Aydin et al. (2019). Although the balanced partitioning design was not developed for the bipartite setting, we may expect it to achieve high precision estimates if the clustering produces decorrelated exposures with large variances.

We generate the potential outcomes in three simulations, where we vary the response functions that are used. The first two simulations feature linear response functions and the third simulation features a non-linear response function. We remark that although the parameters of the response functions are randomly chosen in our simulations, this random parameter draw is made only once and the outcomes themselves are fixed across all sampled assignments of all designs. These simulations are listed below.

- **(Mostly) Positive Treatment Effect.** In this simulation, we set almost all of the individual treatment effects to be positive across units, while varying the responses amongst the units. More precisely, we sample the slope terms as $\beta_i \sim \mathcal{N}(1, 1/4)$ and the intercept terms as $\alpha_i \sim \mathcal{N}(0, 1/8)$.
- **(Nearly) Zero Treatment Effect.** In this simulation, we set all the individual treatment effects close to zero, while varying the baseline outcomes. The outcomes are chosen to be mostly positive. More precisely, we sample the slope terms as $\beta_i \sim \mathcal{N}(0, 1/8)$ and the intercept terms as $\alpha_i \sim \mathcal{N}(2, 1/4)$.
- **Non-Linear Response.** In this simulation, we use a non-linear response function to specify the potential outcomes. In particular, the response of outcome unit i is $Y_i(x_i) = 1 - x_i^2 + \alpha_i$, where $\alpha_i \sim \mathcal{N}(0, 1/8)$. Under this response, all individual treatment effects are 0. Because the linear response assumption is not satisfied, we should not expect our statistical analysis (unbiasedness, consistency, normality, etc) to hold exactly.

When using the EXPOSURE-DESIGN, we set the correlation penalty parameter to $\phi = 0.223$, chosen from a grid of 10 points between $[0, 2]$. The clustering itself is obtained using our local search heuristic presented in Section 6.3. Recall that the correlation clustering objective of Pouget-Abadie et al. (2019) may be obtained by setting $\phi = 1/(n - 1)$. For this reason, we compute the corresponding cluster by running our local search heuristic with $\phi = 0.001 \approx 1/(n - 1)$.

A summary of the main results from these simulations appears in Table 1. For each treatment design and simulation, we sample 20,000 exposure vectors, compute the observed outcomes, and construct the

Design	Simulation 1			Simulation 2			Simulation 3		
	RMSE	Width	Cover.	RMSE	Width	Cover.	RMSE	Width	Cover.
Exposure-Design	0.049	0.219	91.8%	1.81	7.03	94.7%	0.86	3.47	95.6%
Corr Clustering	0.088	0.328	94.1%	2.05	8.00	95.0%	0.90	3.82	96.9%
Balanced Partitioning	0.073	0.283	93.7%	1.85	7.21	94.6%	0.78	3.39	96.9%
Bernoulli	0.659	2.576	95.1%	48.83	190.8	95.1%	24.37	95.15	95.1%

Table 1: Simulation results

corresponding ERL and variance estimators. Given the ERL and variance estimators, we construct the confidence intervals as described in Section 5, with absolute value corrections when the variance estimator takes a negative value. For each simulation and treatment design, we report the root mean square error (RMSE) of the ERL estimator, the average width of the 95% confidence intervals, and the coverage of the 95% confidence intervals.

We draw particular attention to a few trends in these results. EXPOSURE-DESIGN achieves the smallest RMSE in the simulations which satisfy the linear response assumption. All cluster-based designs achieve significantly smaller RMSE than the Bernoulli design, which emphasizes the importance of carefully considering the exposure distribution when the growth conditions (specified in Section 4.1) are not satisfied. The confidence intervals in Simulation 1 under EXPOSURE-DESIGN cover below the nominal 95% level, indicating that either the sampling distribution of the ERL estimator isn't sufficiently approximated by a normal or the variance estimator isn't sufficiently concentrated. The confidence intervals in Simulation 3 cover slightly above the nominal 95% level, which is a result of conservative bias in the variance estimate due to non-linearity of the response.

Figure 1 contains histograms of the ERL estimator for each simulation and design, where the rows correspond to the designs and the columns correspond to the simulations. The dotted vertical line in the plot is the true ATTE. In all simulations, the distribution of the ERL estimator appears unimodal, centered around the ATTE, and (roughly) normal, which is empirical evidence that the normal approximation used to derive confidence intervals may be well-motivated for EXPOSURE-DESIGN and other cluster-based designs. This is to be expected for Simulations 1 and 2 where the linear response assumption holds, but is perhaps surprising in Simulation 3, which features a highly non-linear response. It is possible that the apparent unbiasedness in Simulation 3 is due to symmetry in the response and the exposure distributions, but rigorously investigating this is beyond the scope of the current paper.

Figure 2 contains histograms of the variance estimator for each simulation and treatment design, where the rows correspond to the designs and the columns correspond to the simulations. The dotted vertical line in the plot is the empirical estimate of the variance of the ERL estimator, computed from samples. The variance estimator is unbiased in Simulations 1 and 2, which aligns with Theorem 4; however, because the empirical variance estimate is used, the blue line is close to (but not exactly) the true variance. Increasing the number of sampled exposure vectors decreases this error, but drawing more than 20 thousand samples is prohibitively expensive given the size of the data. In Simulation 1, the mean squared error of all cluster-based designs is so small that the variance estimator takes negative values. In Simulation 3, the response is highly non-linear so that the variance estimator incurs a positive bias, which results in a coverage slightly above the nominal level. Interestingly, the variance estimator under the balanced partitioning design is more concentrated around its mean, which is worth further rigorous investigation.

Figure 3 contains a plot for each simulation, where the mean squared error is plotted against the correlation penalizing parameter ϕ . The mean squared error of the balanced partitioning design appears as a dotted blue line. In Simulations 1 and 2 where the linear response assumption holds, there is a range of values of ϕ where EXPOSURE-DESIGN achieves lower mean squared error than the balanced partitioning design. In our simulations, the choice of $\phi \approx 1/4$ typically achieves lowest mean squared error. However, no design is optimal across all types of potential outcomes (Harshaw et al., 2021) and so we encourage experimenters to select ϕ (and more generally, select designs) by running tests on simulated data.




Figure 1: Histograms of the ERL estimator in simulations.

8 Concluding remarks

In this paper, we have presented methodological contributions towards estimating treatment effects in the linear-exposure response model for bipartite experiments: namely, the Exposure Reweighted Linear (ERL) estimator for consistent and unbiased estimation of average total treatment effect, an unbiased variance estimator which facilitates construction of normal-based confidence intervals, and the EXPOSURE-DESIGN that aims to increase precision of this estimator in various settings of interest.

When employing this design in practice, we recommend that the experimenter choose the value of the trade-off parameter ϕ by running simulations of the experiment using available models of the outcomes when possible. When this is not possible, we find in our simulations that setting $\phi \approx 1/4$ typically yields improvements in the precision of the ERL estimator over the previously correlation clustering design of Pouget-Abadie et al. (2019) where $\phi = 1/(n - 1)$. We suspect that in most settings of interest, the ERL estimator will enjoy increased precision under any treatment design which (either explicitly or implicitly) ensures that exposures have large variance and are decorrelated.

There are several open questions suggested by this work. Given that the design we describe in this paper is heuristically motivated, there is likely room for improved designs. One technical challenge is to construct




Figure 2: Histograms of the variance estimator in simulations.




Figure 3: MSE of the ERL estimator as trade-off parameter ϕ is varied. First values of ϕ are .001 and .01.

a design for which consistency and asymptotic normality of the ERL estimator may be established under weaker growth conditions on the bipartite graph. Another key open question is developing methods for valid inference in bipartite experiments which go beyond the linear exposure-response assumption. Finally, it would be of practical and methodological interest to develop estimation techniques that are robust to

misspecification in the bipartite graph as well as estimation techniques which perform well in the presence of greater structural interference (i.e. when outcomes are influenced by the exposures of other units). The results in Sävje (2021) regarding estimation of treatment effects under a misspecified exposure mapping might extend to this setting, but that remains to be shown. Answering these methodological questions around the bipartite experimental framework will increase its relevance and applicability in practice.

Acknowledgements We would like to thank Peter Aronow, Kay Brodersen, Nick Doudchenko, Ramesh Johari, Khashayar Khosravi, Sebastien Lahaie, Vahan Nanumyan, Lewis Rendell, and Johan Ugander for stimulating discussions which helped shape this work. Christopher Harshaw gratefully acknowledges support from an NSF Graduate Research Fellowship (DGE1122492) as well as support from Google as a Summer Intern and a Student Researcher.

References

- Aral, S. and Walker, D. (2011). Creating social contagion through viral product design: A randomized trial of peer influence in networks. *Management science*, 57(9):1623–1639.
- Aronow, P. M. and Samii, C. (2013). Conservative variance estimation for sampling designs with zero pairwise inclusion probabilities. *Survey Methodology*, 39(1):231–241.
- Aronow, P. M. and Samii, C. (2017). Estimating average causal effects under general interference, with application to a social network experiment. *The Annals of Applied Statistics*, 11(4):1912–1947.
- Aydin, K., Bateni, M., and Mirrokni, V. (2019). Distributed balanced partitioning via linear embedding. *Algorithms*, 12(8):162.
- Baird, S., Bohren, J. A., McIntosh, C., and Özler, B. (2018). Optimal Design of Experiments in the Presence of Interference. *The Review of Economics and Statistics*, 100(5):844–860.
- Bansal, N., Blum, A., and Chawla, S. (2002). Correlation clustering. In *Proceedings of the 43rd Symposium on Foundations of Computer Science*, FOCS '02, page 238. IEEE Computer Society.
- Basse, G. W. and Airoidi, E. M. (2018). Limitations of design-based causal inference and A/B testing under arbitrary and network interference. *Sociological Methodology*, 48(1):136–151.
- Basse, G. W., Soufiani, H. A., and Lambert, D. (2016). Randomization and the pernicious effects of limited budgets on auction experiments. In *Artificial Intelligence and Statistics*, pages 1412–1420. PMLR.
- Blake, T. and Coey, D. (2014). Why marketplace experimentation is harder than it seems: The role of test-control interference. In *Proceedings of the fifteenth ACM conference on Economics and computation*, pages 567–582.
- Bramoullé, Y., Djebbari, H., and Fortin, B. (2009). Identification of peer effects through social networks. *Journal of Econometrics*, 150(1):41–55.
- Charikar, M., Guruswami, V., and Wirth, A. (2005). Clustering with qualitative information. *J. Comput. Syst. Sci.*, 71(3):360–383.
- Chin, A. (2019a). Central limit theorems via Stein’s method for randomized experiments under interference. arXiv:1804.03105.
- Chin, A. (2019b). Regression adjustments for estimating the global treatment effect in experiments with interference. *Journal of Causal Inference*, 7(2).
- Cohen, E. and Lewis, D. D. (1999). Approximating matrix multiplication for pattern recognition tasks. *Journal of Algorithms*, 30(2):211 – 252.
- Doudchenko, N., Zhang, M., Drynkin, E., Airoidi, E., Mirrokni, V., and Pouget-Abadie, J. (2020). Causal inference with bipartite designs. *arXiv preprint arXiv:2010.02108*.

- Eckles, D., Karrer, B., and Ugander, J. (2016a). Design and analysis of experiments in networks: Reducing bias from interference. *Journal of Causal Inference*, 5(1).
- Eckles, D., Kizilcec, R. F., and Bakshy, E. (2016b). Estimating peer effects in networks with peer encouragement designs. *Proceedings of the National Academy of Sciences*, 113(27):7316–7322.
- Einav, L., Kuchler, T., Levin, J. D., and Sundaresan, N. (2011). Learning from seller experiments in online markets. Technical report, National Bureau of Economic Research.
- Elsner, M. and Schudy, W. (2009). Bounding and comparing methods for correlation clustering beyond ilp. In *Proceedings of the Workshop on Integer Linear Programming for Natural Language Processing, ILP '09*, page 19–27. Association for Computational Linguistics.
- Fattorini, L. (2006). Applying the Horvitz-Thompson criterion in complex designs: A computer-intensive perspective for estimating inclusion probabilities. *Biometrika*, 93(2):269–278.
- Fradkin, A. (2015). Search frictions and the design of online marketplaces. *Work. Pap., Mass. Inst. Technol.*
- Gupta, S., Kohavi, R., Tang, D., Xu, Y., Andersen, R., Bakshy, E., Cardin, N., Chandran, S., Chen, N., Coey, D., et al. (2019). Top challenges from the first practical online controlled experiments summit. *ACM SIGKDD Explorations Newsletter*, 21(1):20–35.
- Halloran, M. E. and Hudgens, M. G. (2016). Dependent happenings: A recent methodological review. *Current Epidemiology Reports*, 3(4):297–305.
- Harshaw, C., Sävje, F., Spielman, D., and Zhang, P. (2021). Balancing covariates in randomized experiments with the gram-schmidt walk design. [arXiv:1911.03071](https://arxiv.org/abs/1911.03071).
- He, R. and McAuley, J. (2016). Ups and downs: Modeling the visual evolution of fashion trends with one-class collaborative filtering. In *proceedings of the 25th international conference on world wide web*, pages 507–517.
- Holland, P. W. (1986). Statistics and causal inference. *Journal of the American Statistical Association*, 81(396):945–960.
- Holtz, D., Lobel, R., Liskovich, I., and Aral, S. (2020). Reducing interference bias in online marketplace pricing experiments. *arXiv preprint arXiv:2004.12489*.
- Horvitz, D. G. and Thompson, D. J. (1952). A generalization of sampling without replacement from a finite universe. *Journal of the American Statistical Association*, 47(260):663–685.
- Hudgens, M. G. and Halloran, M. E. (2008). Toward causal inference with interference. *Journal of the American Statistical Association*, 103(482):832–842.
- Imbens, G. W. and Rubin, D. B. (2015). *Causal inference in statistics, social, and biomedical sciences*. Cambridge University Press.
- Johari, R., Li, H., and Weintraub, G. (2020). Experimental design in two-sided platforms: An analysis of bias. In *Proceedings of the 21st ACM Conference on Economics and Computation*, page 851.
- Kempton, R. (1997). Interference between plots. In *Statistical methods for plant variety evaluation*, pages 101–116. Springer.
- Liu, M., Mao, J., and Kang, K. (2020). Trustworthy online marketplace experimentation with budget-split design. *arXiv preprint arXiv:2012.08724*.
- Manski, C. F. (1993). Identification of endogenous social effects: The reflection problem. *The Review of Economic Studies*, 60(3):531–542.

- McAuley, J., Targett, C., Shi, Q., and Van Den Hengel, A. (2015). Image-based recommendations on styles and substitutes. In *Proceedings of the 38th international ACM SIGIR conference on research and development in information retrieval*, pages 43–52.
- Narain, R. (1951). On sampling without replacement with varying probabilities. *Journal of the Indian Society of Agricultural Studies*, (3):169–175.
- Neyman, J. (1923). Sur les applications de la théorie des probabilités aux expériences agricoles: Essai des principes. *Roczniki Nauk Rolniczych*, 10:1–51.
- Offer-Westort, M. and Dimmery, D. (2021). Experimentation for homogenous policy change. arXiv:2101.12318.
- Ogburn, E. L., Sofrygin, O., Diaz, I., and van der Laan, M. J. (2020). Causal inference for social network data. arXiv:1705.08527.
- Pouget-Abadie, J., Aydin, K., Schudy, W., Brodersen, K., and Mirrokni, V. (2019). Variance reduction in bipartite experiments through correlation clustering. In *Advances in Neural Information Processing Systems*, pages 13309–13319.
- Reiley, D. H. (2006). Field experiments on the effects of reserve prices in auctions: More magic on the internet. *The RAND Journal of Economics*, 37(1):195–211.
- Ross, N. (2011). Fundamentals of Stein’s method. *Probability Surveys*, 8:210–293.
- Sävje, F. (2021). Causal inference with misspecified exposure mappings. arXiv:2103.06471.
- Sävje, F., Aronow, P. M., and Hudgens, M. G. (2021). Average treatment effects in the presence of unknown interference. *The Annals of Statistics*, 49(2):673–701.
- Sinclair, B., McConnell, M., and Green, D. P. (2012). Detecting spillover effects: Design and analysis of multilevel experiments. *American Journal of Political Science*, 56(4):1055–1069.
- Struchiner, C. J., Halloran, M. E., Robins, J. M., and Spielman, A. (1990). The behaviour of common measures of association used to assess a vaccination programme under complex disease transmission patterns—a computer simulation study of malaria vaccines. *International journal of epidemiology*, 19(1):187–196.
- Swamy, C. (2004). Correlation clustering: Maximizing agreements via semidefinite programming. SODA ’04, page 526–527. Society for Industrial and Applied Mathematics.
- Toulis, P. and Kao, E. (2013). Estimation of causal peer influence effects. In *Proceedings of the 30th International Conference on Machine Learning*, volume 28, pages 1489–1497.
- Zigler, C. M. and Papadogeorgou, G. (2021). Bipartite causal inference with interference. *Statist. Sci.*, 36(1):109–123.

Appendix

A Analysis of the ERL Estimator

In this section, we present proofs of consistency and asymptotic normality of ERL estimator appearing in Section 4 of the main paper. Before continuing, we introduce some notation used in the proofs. We begin by defining for each outcome unit $i \in V_o$, an estimate of the individual treatment effect τ_i , which is

$$\hat{\tau}_i \triangleq 2Y_i(\mathbf{z}) \left(\frac{x_i(\mathbf{z}) - \mathbb{E}[x_i(\mathbf{z})]}{\text{Var}(x_i(\mathbf{z}))} \right) .$$

Observe that the ERL estimator is the average of these estimates of the individual treatment effects, i.e. $\hat{\tau} = (1/n) \sum_{i=1}^n \hat{\tau}_i$. Throughout the proofs, we will often reason about the behavior of the ERL estimator through the properties of the individual treatment effect estimates.

Next, we introduce the concept of *dependency neighborhoods* (Ross, 2011). Let a_1, a_2, \dots, a_n be random variables indexed by the integers $[n]$ and collect these random variables into the set $\mathcal{A} = \{a_i : i \in [n]\}$. For each variable a_i , we define the *dependency neighborhood* as

$$\mathcal{I}(i) \subset \mathcal{A} \text{ such that } a_i \text{ is jointly independent of the variables } \mathcal{A} \setminus \mathcal{I}(i) .$$

In other words, a random variable a_i is jointly independent of all variables not contained in its dependency neighborhood, but is dependent on variables contained in its dependency neighborhood. We take the convention that $i \in \mathcal{I}(i)$ and so that each dependency neighborhood has cardinality at least 1. A measure of dependence between the random variables is the *maximum dependency degree*, which is $D = \max_{i \in [n]} |\mathcal{I}(i)|$. Note that independent random variables satisfy $D = 1$ and that completely dependent random variables have $D = n$.

For the remainder of the proof, we focus our discussion of dependency neighborhoods and degrees to the collection of errors of the individual treatment effects,

$$a_1 = \tau_1 - \hat{\tau}_1, \quad a_2 = \tau_2 - \hat{\tau}_2, \quad \dots \quad a_n = \tau_n - \hat{\tau}_n .$$

We begin by showing that in this case, the maximum dependency degree may be bounded in terms of the degrees of the bipartite graph and the dependence in the treatment assignments.

Lemma 3. *The dependency degree of the individual treatment effect errors is bounded by $D \leq kd_d d_o$.*

Proof. The first part of this proof is to establish a necessary condition for an individual treatment effect error a_j to be in the dependency neighborhood of a_i , i.e. $a_j \in \mathcal{I}(i)$. We begin by re-writing the exposures under a cluster design. Recall that the exposures are defined as $x_i = \sum_{j=1}^m w_{i,j} z_j$. For each cluster $C \in \mathcal{C}$, define $w_{i,C} = \sum_{j \in C} w_{i,j}$ and define z_C to be the ± 1 cluster treatment assignment variable which is 1 if diversion units in C are treated and -1 otherwise. If $w_{i,C} \neq 0$, then we say that cluster C is *incident* to outcome unit i . Define $S(i) = \{z_C : w_{i,C} \neq 0\}$ to be the cluster treatment assignments which influence the exposure x_i . Under the cluster design, the exposure for outcome unit i may be written as

$$x_i = \sum_{C \in \mathcal{C}} w_{i,C} z_C = \sum_{C \in S(i)} w_{i,C} z_C .$$

By the linear-response assumption, the individual treatment effect error a_i is a function of the exposure x_i . Moreover, a_i is a function of the cluster treatment assignment variables in $S(i)$. Let us denote this relationship by writing $a_i = g_i(S(i))$, where g_i is a function of the cluster treatment variables $z_C \in S(i)$.

Let $B \subset V_o$ be a collection of outcome units. We remark that joint independence of cluster treatment assignments implies joint independence of individual treatment effect errors:

$$S(i) \perp\!\!\!\perp \{S(j) : j \in B\} \Rightarrow a_i \perp\!\!\!\perp \{a_j : j \in B\} .$$

Under an independent cluster design, the cluster treatment assignments $S(i)$ are jointly independent of the cluster treatment assignments $\{S(j) : j \in B\}$ when the corresponding sets of clusters are disjoint, i.e. $S(i) \cap (\cup_{j \in B} S(j)) = \emptyset$. Thus, the individual treatment effect estimate a_i is jointly independent of the collection of individual treatment effect estimates $\{a_j : j \in B\}$ when outcome unit i is not incident to any cluster that is incident to an outcome unit in B . In other words, $a_j \in \mathcal{I}(i)$ only if outcome units i and j are incident to a common cluster.

Fix an outcome unit $i \in V_o$. The remainder of the proof is a simple counting argument which uses this necessary condition to establish that $|\mathcal{I}(i)| \leq kd_d d_o$. In particular, we will count the number of outcome units that are incident to one of the clusters that are incident to i . Because the degree of outcome unit i is at most d_o , it is incident to at most d_o clusters. Each of these clusters has at most k diversion units, by Assumption 2. Because the degree of all diversion units j is at most d_d , the number of outcome units which are incident to at least one of these clusters is at most $kd_d d_o$. Thus, we have established that

$$D = \max_{i \in V_o} |\mathcal{I}(i)| \leq kd_d d_o . \quad \square$$

The following lemma derives a lower bound the exposure variances in terms of the treatment assignment probability and the maximum degree of the outcome units.

Lemma 4. *If each pair of treatment assignments is non-negatively correlated, then each exposure variance is lower bounded as $\text{Var}(x_i) \geq \frac{4p(1-p)}{d_o}$.*

Proof. We begin by expanding the variance of the exposure x_i by

$$\begin{aligned} \text{Var}(x_i) &= \text{Var}\left(\sum_{j=1}^m w_{i,j} z_j\right) && \text{(definition of exposure)} \\ &= \sum_{i=1}^m \left[\text{Var}(w_{i,j} z_j) + \sum_{\ell \neq j} \text{Cov}(w_{i,j} z_j, w_{i,\ell} z_\ell) \right] && \text{(properties of variance)} \\ &= \sum_{i=1}^m \left[w_{i,j}^2 \text{Var}(z_j) + \sum_{\ell \neq j} w_{i,j} w_{i,\ell} \text{Cov}(z_j, z_\ell) \right] && \text{(properties of variance and covariance)} \\ &\geq \sum_{i=1}^m w_{i,j}^2 \text{Var}(z_j) && \text{(non-negative weights and correlations)} \\ &= 4p(1-p) \sum_{i=1}^m w_{i,j}^2 , \end{aligned}$$

where the inequality follows because the weights $w_{i,j}$ are non-negative and the assignments are non-negatively correlated and the last equality follows because z_i are ± 1 random variables with $\Pr(z_i = 1) = p$.

We complete the proof by lower bounding the sum of the squares of the weights. Recall that the sum of the weights is 1 and there are at most d_o non-negative terms in the sum. Using this together with the inequality that relates ℓ_2 to ℓ_1 norms in d -dimensions, $\|\cdot\|_2^2 \geq \frac{1}{d} \|\cdot\|_1^2$, we have that

$$\sum_{i=1}^m w_{i,j}^2 \geq \frac{1}{d_o} \sum_{i=1}^m w_{i,j} = \frac{1}{d_o} . \quad \square$$

The following lemma is a bound on the moments of the errors of the individual treatment effects.

Lemma 5. *The p th moment of the error of the individual treatment effect estimates is bounded by*

$$\mathbb{E}[|\tau_i - \hat{\tau}_i|^p] \leq \left[2M \left(1 + \frac{d_o}{2p(1-p)} \right) \right]^p .$$

Proof. We begin by remarking that $|Y_i(\mathbf{z})| \leq M$ implies that each of the individual slopes are also bounded in absolute value as $|\beta_i| \leq M$. Recall that by the linear response assumption, $Y_i(\mathbf{z}) = \beta_i x_i + \alpha_i$ and by the linear exposure assumption (along with the normalization of the edge weights), setting $\mathbf{z} = \pm \mathbf{1}$ results in an exposure of $\xi = \pm 1$. Thus, when considering $\mathbf{z} = \pm \mathbf{1}$, the bound $|Y_i(\mathbf{z})| \leq M$ implies that $|\pm \beta_i + \alpha_i| \leq M$, which is enough to establish that $|\beta_i| \leq M$.

We now proceed by proving a bound on $|\tau_i - \hat{\tau}_i|$, which holds for any realization of the random variables:

$$\begin{aligned}
|\tau_i - \hat{\tau}_i| &= \left| 2\beta_i - 2Y_i(\mathbf{z}) \left(\frac{x_i - \mathbb{E}[x_i]}{\text{Var}(x_i)} \right) \right| \\
&\leq 2|\beta_i| + 2 \frac{|Y_i(\mathbf{z})| \cdot |x_i - \mathbb{E}[x_i]|}{\text{Var}(x_i)} && \text{(triangle inequality)} \\
&\leq 2M + 2 \frac{M \cdot 2}{\text{Var}(x_i)} && \text{(definition of } M \text{ and above)} \\
&\leq 2M \left(1 + \frac{2}{\text{Var}(x_i)} \right) && \text{(collecting terms)} \\
&\leq 2M \left(1 + \frac{d_o}{2p(1-p)} \right) && \text{(Lemma 4)}
\end{aligned}$$

The moment bound follows by applying the bound above. \square

A.1 Consistency of ERL Estimator (Theorem 2)

We are now ready to establish the consistency of the ERL estimator. Before doing so, we restate the theorem here.

Theorem 2. *Under Assumptions 1 and 2, and supposing that $d_d d_o^3 = o(n)$ in the asymptotic sequence, the ERL estimator converges in mean square to the ATTE: $\lim_{N \rightarrow \infty} \mathbb{E}[(\hat{\tau} - \tau)^2] = 0$.*

Proof. We begin by proving a finite sample bound on the mean squared error of the ERL estimator, and then we finish the proof by taking the limit in the asymptotic sequence. Note that the mean squared error may be broken down into the errors of the individual treatment effect estimates via

$$\mathbb{E}[(\tau - \hat{\tau})^2] = \mathbb{E} \left[\left(\frac{1}{n} \sum_{i=1}^n (\tau_i - \hat{\tau}_i) \right)^2 \right] = \frac{1}{n^2} \sum_{i=1}^n \left(\mathbb{E}[(\tau_i - \hat{\tau}_i)^2] + \sum_{j \neq i} \mathbb{E}[(\tau_i - \hat{\tau}_i)(\tau_j - \hat{\tau}_j)] \right).$$

Note that the term in the inner sum is the covariance of the errors in the individual treatment effect estimators. By definition of the dependency neighborhoods, only terms $j \in \mathcal{I}(i)$ are dependent and so only these terms will have non-zero covariance. Using this and the second moment bound in Lemma 5, we have that

$$\begin{aligned}
\mathbb{E}[(\tau - \hat{\tau})^2] &= \frac{1}{n^2} \sum_{i=1}^n \left(\mathbb{E}[(\tau_i - \hat{\tau}_i)^2] + \sum_{\substack{j \in \mathcal{I}(i) \\ j \neq i}} \mathbb{E}[(\tau_i - \hat{\tau}_i)(\tau_j - \hat{\tau}_j)] \right) && \text{(dependency neighborhoods)} \\
&\leq \frac{1}{n^2} \sum_{i=1}^n \left(\mathbb{E}[(\tau_i - \hat{\tau}_i)^2] + \sum_{\substack{j \in \mathcal{I}(i) \\ j \neq i}} \sqrt{\mathbb{E}[(\tau_i - \hat{\tau}_i)^2] \mathbb{E}[(\tau_j - \hat{\tau}_j)^2]} \right) && \text{(Cauchy-Schwarz)} \\
&\leq \frac{1}{n^2} \sum_{i=1}^n |\mathcal{I}(i)| \cdot \left[2M \left(1 + \frac{d_o}{2p(1-p)} \right) \right]^2 && \text{(Lemma 5)} \\
&\leq \frac{D}{n} \left[2M \left(1 + \frac{d_o}{2p(1-p)} \right) \right]^2. && \text{(max dependency degree)}
\end{aligned}$$

By using the bound $D \leq k d_d d_o$ given in Lemma 3, we have the finite-sample bound on the mean squared error:

$$\mathbb{E}[(\tau - \hat{\tau})^2] \leq \frac{k d_d d_o}{n} \left[2M \left(1 + \frac{d_o}{2p(1-p)} \right) \right]^2$$

We now interpret this finite-sample bound in the context of the asymptotic sequence. By Assumptions 1 and 2, we have that M is a constant, p is bounded away from 0 and 1 by a constant, and k is a constant. It follows that the mean squared error is asymptotically bounded by the rate $\mathbb{E}[(\tau - \hat{\tau})^2] = \mathcal{O}(d_d d_o^3/n)$. By assumption, the asymptotic sequence satisfies $d_d d_o^3 = o(n)$, and thus the mean squared error converges to zero under these conditions. \square

A.2 Asymptotic Normality (Theorem 3)

We establish asymptotic normality of the ERL estimator by using Stein's method. In particular, we use the following result from Ross (2011):

Lemma 6 (Lemma 3.6 of Ross (2011)). *Let a_1, a_2, \dots, a_n be random variables such that $\mathbb{E}[a_i^4] < \infty$, $\mathbb{E}[a_i] = 0$, $\sigma^2 = \text{Var}(\frac{1}{n} \sum_{i=1}^n a_i)$, and define $X = (\frac{1}{n} \sum_{i=1}^n a_i)/\sigma$. Then for a standard normal $Z \sim \mathcal{N}(0, 1)$, we have*

$$d_W(X, Z) \leq \frac{D^2}{\sigma^3 n^3} \sum_{i=1}^n \mathbb{E}[|a_i|^3] + \sqrt{\frac{28}{\pi}} \cdot \frac{D^{3/2}}{n^2 \sigma^2} \sqrt{\sum_{i=1}^n \mathbb{E}[a_i^4]} ,$$

where D is the maximum dependency degree of the random variables and $d_W(\cdot, \cdot)$ is the Wasserstein distance.

We will use Lemma 6 to prove asymptotic normality of the ERL estimator. Before continuing, let us restate the theorem.

Theorem 3. *Under Assumptions 1, 2, and 3, and supposing that $d_d^{1.6} d_o^4 = o(n)$, the ERL estimator is asymptotically normal:*

$$\frac{\hat{\tau} - \tau}{\sqrt{\text{Var}(\hat{\tau})}} \xrightarrow{d} \mathcal{N}(0, 1) .$$

Proof. Our strategy may be described in two main steps: first, we use Lemma 6 to derive a finite-sample bound on the Wasserstein distance between the distribution of $(\tau - \hat{\tau})/\sqrt{\text{Var}(\hat{\tau})}$ and a standard normal. Next, we use this bound to argue that this Wasserstein distance approaches 0 in the limit of the asymptotic sequence under the above conditions.

We seek to apply Lemma 6 where the random variables are the errors of the individual treatment effect estimates; that is,

$$a_1 = \tau_1 - \hat{\tau}_1, a_2 = \tau_2 - \hat{\tau}_2, \dots, a_n = \tau_n - \hat{\tau}_n .$$

Note that $\frac{1}{n} \sum_{i=1}^n a_i = \frac{1}{n} \sum_{i=1}^n \tau_i - \hat{\tau} = \tau - \hat{\tau}$ and $\text{Var}(\frac{1}{n} \sum_{i=1}^n a_i) = \text{Var}(\hat{\tau})$ so that the random variable X in Lemma 6 is equal to $(\tau - \hat{\tau})/\sqrt{\text{Var}(\hat{\tau})}$, which is indeed the random variable we wish to characterize. Let's show that the conditions of Lemma 6 are satisfied: first, recall that $\hat{\tau}_i$ are unbiased estimates of τ_i so that a_i has mean zero. Second, because the potential outcomes are bounded by a constant M , the support of a_i is bounded so the fourth moments are finite. Thus, we may apply Lemma 6 in this setting.

We will use the Lemma 5 to bound the sum of the third and fourth moments. In particular, Lemma 5 implies that

$$\sum_{i=1}^n \mathbb{E}[|a_i|^3] \leq n \cdot \left[2M \left(1 + \frac{d_o}{2p(1-p)} \right) \right]^3 \quad \text{and} \quad \sum_{i=1}^n \mathbb{E}[|a_i|^4] \leq n \cdot \left[2M \left(1 + \frac{d_o}{2p(1-p)} \right) \right]^4 .$$

Using this moment bound together with the bound on the maximum dependence degree D (Lemma 3) on the result of Lemma 6, we obtain that

$$d_W\left(\frac{\tau - \hat{\tau}}{\sqrt{\text{Var}(\hat{\tau})}}, Z\right) \leq \frac{(k d_d d_o)^2}{\sigma^3 n^2} \cdot \left[2M \left(1 + \frac{d_o}{2p(1-p)} \right) \right]^3 + \sqrt{\frac{28}{\pi}} \cdot \frac{(k d_d d_o)^{3/2}}{\sigma^2 n^{3/2}} \cdot \left[2M \left(1 + \frac{d_o}{2p(1-p)} \right) \right]^2$$

We now interpret this finite-sample bound in the context of the asymptotic sequence. By Assumptions 1 and 2, we have that M is a constant, p is bounded away from 0 and 1 by a constant, and k is a constant.

It follows that the Wasserstein distance between $(\tau - \hat{\tau})/\sqrt{\text{Var}(\hat{\tau})}$ and a standard normal is asymptotically bounded as

$$d_W\left(\frac{\tau - \hat{\tau}}{\sqrt{\text{Var}(\hat{\tau})}}, Z\right) = \mathcal{O}\left(\frac{d_d^2 d_o^5}{\sigma^3 n^2} + \frac{d_d^{1.5} d_o^{3.5}}{\sigma^2 n^{3/2}}\right)$$

By Assumption 3, we have that $\text{Var}(\hat{\tau}) = \Omega(n^{-1/2})$, which means that this bound becomes

$$d_W\left(\frac{\tau - \hat{\tau}}{\sqrt{\text{Var}(\hat{\tau})}}, Z\right) = \mathcal{O}\left(\frac{d_d^2 d_o^5}{n^{5/4}} + \frac{d_d^{1.5} d_o^{3.5}}{n}\right).$$

By assumption, the asymptotic sequence satisfies $d_d^{1.6} d_o^4 = o(n)$. Raising both sides to the 5/4 yields that $d_d^2 d_o^5 = o(n^{5/4})$. Additionally, d_d and d_o are positive integers and so they satisfy $d_d^{1.5} d_o^{3.5} \leq d_d^{1.6} d_o^4$, thus $d_d^{1.5} d_o^{3.5} = o(n)$. Thus, the Wasserstein distance between $(\tau - \hat{\tau})/\sqrt{\text{Var}(\hat{\tau})}$ and a standard normal approaches 0 in this asymptotic sequence. \square

B Variance Estimation

In this section, we prove Lemmas 1 and 2, which are then used to prove Theorem 4, which establishes unbiasedness of the proposed variance estimator.

First, we show Lemma 1, which demonstrates how to estimate the variance of an individual treatment effect estimate. To do so, we need to decompose the random variable Q_i . For each outcome unit $i \in V_o$, define the random variable T_i to be the second term in the function Q_i . That is,

$$T_i = \frac{\text{Var}(x_i)(x_i^2 - \mathbb{E}[x_i^2]) - \text{Cov}(x_i, x_i^2)(x_i - \mathbb{E}[x_i])}{\text{Var}(x_i) \text{Var}(x_i^2) - \text{Cov}(x_i, x_i^2)^2}$$

so that $Q_i \triangleq \frac{(x_i - \mathbb{E}[x_i])^2}{\text{Var}(x_i)^2} - T_i$. The following Lemma states the key properties of the random variable T_i , and consequently Q_i , which allow for the proof of Lemma 1.

Lemma 7. *If the exposure x_i takes at least three values with non-zero probability, then the function T_i satisfies the following three properties:*

- $\mathbb{E}[T_i] = 0$
- $\mathbb{E}[x_i T_i] = 0$
- $\mathbb{E}[x_i^2 T_i] = 1$

Proof. For notational simplicity, we drop the subscript as write x_i as x and T_i as T .

First, we will show that the denominator $\text{Var}(x) \text{Var}(x^2) - \text{Cov}(x, x^2)^2$ is positive when the exposure x takes at least three values with non-zero probability. By Cauchy-Schwarz, we have that $\text{Cov}(x, x^2)^2 \leq \text{Var}(x) \text{Var}(x^2)$, which establishes that denominator is non-negative. The Cauchy-Schwarz inequality is strict precisely when x and x^2 are not perfectly correlated, i.e. there does not exist a and b such that $x^2 = ax + b$. Note that this cannot happen when x takes three distinct values. Thus, the Cauchy-Schwarz is strict in this case so that the denominator is positive. For notational convenience, we write the denominator as $\Delta = \text{Var}(x) \text{Var}(x^2) - \text{Cov}(x, x^2)^2$ throughout the remainder of the proof.

We will now show the three properties. The first property follows by linearity of expectation, as

$$\begin{aligned} \Delta \cdot \mathbb{E}[T] &= \mathbb{E}[\text{Var}(x_i)(x^2 - \mathbb{E}[x^2]) - \text{Cov}(x, x^2)(x - \mathbb{E}[x])] \\ &= \text{Var}(x)(\mathbb{E}[x^2] - \mathbb{E}[x^2]) - \text{Cov}(x, x^2)(\mathbb{E}[x] - \mathbb{E}[x]) \\ &= 0, \end{aligned}$$

and the result follows by dividing through by Δ on both sides. Next, we show the second property. Again, by linearity of expectation, we have that

$$\begin{aligned} \Delta \cdot \mathbb{E}[xT] &= \mathbb{E}[\text{Var}(x_i)(x^3 - x\mathbb{E}[x^2]) - \text{Cov}(x, x^2)(x^2 - x\mathbb{E}[x])] \\ &= \text{Var}(x)(\mathbb{E}[x^3] - \mathbb{E}[x]\mathbb{E}[x^2]) - \text{Cov}(x, x^2)(\mathbb{E}[x^2] - \mathbb{E}[x]^2) \\ &= \text{Var}(x) \text{Cov}(x, x^2) - \text{Cov}(x, x^2) \text{Var}(x) \\ &= 0, \end{aligned}$$

and the result follows by dividing through by Δ on both sides. Finally, we show the third property. By linearity of expectation,

$$\begin{aligned}\Delta \cdot \mathbb{E}[x^2 T] &= \mathbb{E}[\text{Var}(x_i)(x^4 - x^2 \mathbb{E}[x^2]) - \text{Cov}(x, x^2)(x^3 - x^2 \mathbb{E}[x])] \\ &= \text{Var}(x)(\mathbb{E}[x^4] - \mathbb{E}[x^2]^2) - \text{Cov}(x, x^2)(\mathbb{E}[x^3] - \mathbb{E}[x^2] \mathbb{E}[x]) \\ &= \text{Var}(x) \text{Var}(x^2) - \text{Cov}(x, x^2)^2 \\ &= \Delta \ ,\end{aligned}$$

and the result follows by dividing both sides by Δ . \square

We are now ready to prove Lemma 1, which we restate here.

Lemma 1. *Fix an outcome unit $i \in V_o$. If the exposure x_i takes at least three values with non-zero probability, then the variance of unit i 's individual treatment effect estimator is equal to*

$$\text{Var}(\hat{\tau}_i) = 4 \cdot \mathbb{E}[Y_i(\mathbf{z})^2 Q_i] \ .$$

Proof. We may use the properties of the random variable T_i proved in Lemma 7 together with the linear response assumption to obtain

$$\begin{aligned}\mathbb{E}[Y_i(\mathbf{z})^2 T_i] &= \mathbb{E}[(\beta_i x_i + \alpha_i)^2 T_i] && \text{(linear response)} \\ &= \beta_i^2 \mathbb{E}[x_i^2 T_i] + 2\beta_i \alpha_i \mathbb{E}[x_i T_i] + \alpha_i^2 \mathbb{E}[T_i] && \text{(expanding terms)} \\ &= \beta_i^2 \ . && \text{(Lemma 7)}\end{aligned}$$

Next, we write the variance of individual treatment effect estimator. Because the individual treatment effect estimator is unbiased,

$$\begin{aligned}\text{Var}(\hat{\tau}_i) &= \mathbb{E}[\hat{\tau}_i^2] - \mathbb{E}[\hat{\tau}_i]^2 \\ &= 4 \cdot \mathbb{E}\left[Y_i(\mathbf{z})^2 \left(\frac{x_i - \mathbb{E}[x_i]}{\text{Var}(x_i)}\right)^2\right] - 4 \cdot \beta_i^2 \\ &= 4 \cdot \mathbb{E}\left[Y_i(\mathbf{z})^2 \left(\frac{x_i - \mathbb{E}[x_i]}{\text{Var}(x_i)}\right)^2\right] - 4 \cdot \mathbb{E}[Y_i(\mathbf{z})^2 T_i] \\ &= 4 \cdot \mathbb{E}\left[Y_i(\mathbf{z})^2 \left[\left(\frac{x_i - \mathbb{E}[x_i]}{\text{Var}(x_i)}\right)^2 - T_i\right]\right] \\ &= 4 \cdot \mathbb{E}[Y_i(\mathbf{z})^2 Q_i] \ .\end{aligned}$$

\square

Next, we will prove Lemma 2 which demonstrates how to estimate the covariance between two individual treatment effects. To do so, we need to decompose the random variable $R_{i,j}$. For each pair of outcome units $i, j \in V_o$, define the random variable $S_{i,j}$ to be the second term in the random variable $R_{i,j}$. That is,

$$S_{i,j} = \frac{a_{i,j}(x_i x_j - \mathbb{E}[x_i x_j]) + b_{i,j}(x_i - \mathbb{E}[x_i]) + c_{i,j}(x_j - \mathbb{E}[x_j])}{\Psi_{i,j}} \ ,$$

where we recall that these coefficients are given as $a_{i,j}, b_{i,j}, c_{i,j}, \Psi_{i,j}$ are defined as

$$\begin{aligned}a_{i,j} &= \text{Var}(x_i) \text{Var}(x_j) - \text{Cov}(x_i, x_j)^2 \\ b_{i,j} &= \text{Cov}(x_i, x_j) \text{Cov}(x_i x_j, x_j) - \text{Var}(x_j) \text{Cov}(x_i x_j, x_i) \\ c_{i,j} &= \text{Cov}(x_i, x_j) \text{Cov}(x_i x_j, x_i) - \text{Var}(x_i) \text{Cov}(x_i x_j, x_j) \\ \Psi_{i,j} &= \text{Var}(x_i x_j)(\text{Var}(x_i) \text{Var}(x_j) - \text{Cov}(x_i, x_j)^2) \\ &\quad - \text{Var}(x_i) \text{Cov}(x_i x_j, x_j)^2 - \text{Var}(x_j) \text{Cov}(x_i x_j, x_i)^2 + 2 \text{Cov}(x_i, x_j) \text{Cov}(x_i x_j, x_j) \text{Cov}(x_i x_j, x_i) \ .\end{aligned}$$

In this way, $R_{i,j} \triangleq \left(\frac{x_i - \mathbb{E}[x_i]}{\text{Var}(x_i)}\right) \left(\frac{x_j - \mathbb{E}[x_j]}{\text{Var}(x_j)}\right) - S_{i,j}$. The following lemma states the key properties of the random variable $S_{i,j}$, and consequently $R_{i,j}$, which allow for the proof of Lemma 2.

Lemma 8. Fix a pair of outcome units $i, j \in V_o$. If $\Psi_{i,j} \neq 0$, then the random variable $S_{i,j}$ satisfies the following properties:

- $\mathbb{E}[S_{i,j}] = 0$
- $\mathbb{E}[x_i S_{i,j}] = \mathbb{E}[x_j S_{i,j}] = 0$
- $\mathbb{E}[x_i x_j S_{i,j}] = 1$

Proof. For notational simplicity, we will drop the subscripts and write x_i as x , x_j as y and $S_{i,j}$ as S . In the same way, we will write $a_{i,j}, b_{i,j}, c_{i,j}$, and $\Psi_{i,j}$ as a, b, c , and Ψ .

Given that $\Psi > 0$, we seek to verify the three properties in the lemma. The first is easy to verify by the linearity of expectation, as

$$\begin{aligned} \Psi \cdot \mathbb{E}[S] &= \mathbb{E}[a(xy - \mathbb{E}[xy]) + b(x - \mathbb{E}[x]) + c(y - \mathbb{E}[y])] \\ &= a(\mathbb{E}[xy] - \mathbb{E}[xy]) + b(\mathbb{E}[x] - \mathbb{E}[x]) + c(\mathbb{E}[y] - \mathbb{E}[y]) \\ &= 0 \end{aligned}$$

so that dividing both sides by Ψ yields the desired result. We now verify the second property. Observe that

$$\begin{aligned} \Psi \cdot \mathbb{E}[xS] &= a \cdot (\mathbb{E}[x^2y] - \mathbb{E}[x]\mathbb{E}[xy]) + b \cdot (\mathbb{E}[x^2] - \mathbb{E}[x]^2) + c \cdot (\mathbb{E}[xy] - \mathbb{E}[x]\mathbb{E}[y]) \\ &= a \cdot \text{Cov}(xy, x) + b \cdot \text{Var}(x) + c \cdot \text{Cov}(x, y) \\ &= \text{Cov}(xy, x)(\text{Var}(x) \text{Var}(y) - \text{Cov}(x, y)^2) \\ &\quad + \text{Var}(x)(\text{Cov}(x, y) \text{Cov}(xy, y) - \text{Var}(y) \text{Cov}(xy, x)) \\ &\quad + \text{Cov}(x, y)(\text{Cov}(x, y) \text{Cov}(xy, x) - \text{Var}(x) \text{Cov}(xy, y)) \\ &= 0 \text{ ,} \end{aligned}$$

and dividing both sides by Ψ yields the desired result. The proof that $\mathbb{E}[yS(x, y)] = 0$ follows in exactly the same fashion. Finally, we verify the third property. Observe that

$$\begin{aligned} \Psi \cdot \mathbb{E}[xyS] &= a \cdot (\mathbb{E}[(xy)^2] - \mathbb{E}[xy]^2) + b \cdot (\mathbb{E}[x^2y] - \mathbb{E}[xy]\mathbb{E}[x]) + c \cdot (\mathbb{E}[xy^2] - \mathbb{E}[xy]\mathbb{E}[y]) \\ &= a \cdot \text{Var}(xy) + b \cdot \text{Cov}(xy, x) + c \cdot \text{Cov}(xy, y) \\ &= \text{Var}(xy)(\text{Var}(x) \text{Var}(y) - \text{Cov}(x, y)^2) \\ &\quad + \text{Cov}(xy, x)(\text{Cov}(x, y) \text{Cov}(xy, y) - \text{Var}(y) \text{Cov}(xy, x)) \\ &\quad + \text{Cov}(xy, y)(\text{Cov}(x, y) \text{Cov}(xy, x) - \text{Var}(x) \text{Cov}(xy, y)) \\ &= \text{Var}(xy)(\text{Var}(x) \text{Var}(y) - \text{Cov}(x, y)^2) \\ &\quad - \text{Var}(x) \text{Cov}(xy, y)^2 - \text{Var}(y) \text{Cov}(xy, x)^2 \\ &\quad + 2 \text{Cov}(x, y) \text{Cov}(xy, y) \text{Cov}(xy, x) \\ &= \Psi \text{ ,} \end{aligned}$$

so that dividing both sides by Ψ yields the desired result. □

Finally, we are ready to prove Lemma 2, which is the last result needed to establish that the variance estimator is unbiased. We restate the lemma below.

Lemma 2. Fix a pair of outcome units $i \neq j \in V_o$. If $\Psi_{i,j} \neq 0$, then the covariance between individual treatment effect estimates $\hat{\tau}_i$ and $\hat{\tau}_j$ may be expressed as

$$\text{Cov}(\hat{\tau}_i, \hat{\tau}_j) = 4 \cdot \mathbb{E}[Y_i(\mathbf{z})Y_j(\mathbf{z})R_{i,j}] \text{ .}$$

Proof. We may use the properties of the random variable $S_{i,j}$ proved in Lemma 8 together with the linear response assumption to obtain

$$\begin{aligned} \mathbb{E}[Y_i(\mathbf{z})Y_j(\mathbf{z})S_{i,j}] &= \mathbb{E}[(\beta_i x_i + \alpha_i)(\beta_j x_j + \alpha_j)S_{i,j}] && \text{(linear response)} \\ &= \beta_i \beta_j \mathbb{E}[x_i x_j S_{i,j}] + \alpha_i \alpha_j \mathbb{E}[S_{i,j}] + \beta_i \alpha_j \mathbb{E}[x_i S_{i,j}] + \beta_j \alpha_i \mathbb{E}[x_j S_{i,j}] && \text{(expanding terms)} \\ &= \beta_i \beta_j \text{ .} && \text{(Lemma 8)} \end{aligned}$$

Next, we write the covariance between the individual treatment effect estimators. Because the individual treatment effect estimators are unbiased,

$$\begin{aligned}
\text{Cov}(\hat{\tau}_i, \hat{\tau}_j) &= \mathbb{E}[\hat{\tau}_i \hat{\tau}_j] - \mathbb{E}[\hat{\tau}_i] \mathbb{E}[\hat{\tau}_j] \\
&= \mathbb{E}[\hat{\tau}_i \hat{\tau}_j] - \tau_i \tau_j \\
&= 4 \cdot \mathbb{E} \left[Y_i(\mathbf{z}) Y_j(\mathbf{z}) \left(\frac{x_i - \mathbb{E}[x_i]}{\text{Var}(x_i)} \right) \left(\frac{x_j - \mathbb{E}[x_j]}{\text{Var}(x_j)} \right) \right] - 4 \cdot \beta_i \beta_j \\
&= 4 \cdot \mathbb{E} \left[Y_i(\mathbf{z}) Y_j(\mathbf{z}) \left(\frac{x_i - \mathbb{E}[x_i]}{\text{Var}(x_i)} \right) \left(\frac{x_j - \mathbb{E}[x_j]}{\text{Var}(x_j)} \right) \right] - 4 \cdot \mathbb{E}[Y_i(\mathbf{z}) Y_j(\mathbf{z}) S_{i,j}] \\
&= 4 \cdot \mathbb{E} \left[Y_i(\mathbf{z}) Y_j(\mathbf{z}) \left[\left(\frac{x_i - \mathbb{E}[x_i]}{\text{Var}(x_i)} \right) \left(\frac{x_j - \mathbb{E}[x_j]}{\text{Var}(x_j)} \right) - S_{i,j} \right] \right] \\
&= 4 \cdot \mathbb{E} \left[Y_i(\mathbf{z}) Y_j(\mathbf{z}) R_{i,j} \right] \quad \square
\end{aligned}$$

We are now ready to prove Theorem 4, which establishes unbiasedness of the variance estimator.

Theorem 4. *Under the conditions in Lemmas 1 and 2, the variance estimator of the ERL point estimator is unbiased, i.e. $\mathbb{E}[\widehat{\text{Var}}(\hat{\tau})] = \text{Var}(\hat{\tau})$.*

Proof. We may calculate the expectation of the variance estimate $\widehat{\text{Var}}(\hat{\tau})$ as

$$\begin{aligned}
\mathbb{E}[\widehat{\text{Var}}(\hat{\tau})] &= \mathbb{E} \left[\frac{4}{n^2} \sum_{i=1}^n \left[Y_i(\mathbf{z})^2 Q_i + \sum_{j \neq i} Y_i(\mathbf{z}) Y_j(\mathbf{z}) R_{i,j} \right] \right] \\
&= \frac{1}{n^2} \sum_{i=1}^n \left[4 \cdot \mathbb{E} \left[Y_i(\mathbf{z})^2 Q_i \right] + \sum_{j \neq i} 4 \cdot \mathbb{E} \left[Y_i(\mathbf{z}) Y_j(\mathbf{z}) R_{i,j} \right] \right] \quad (\text{linearity of } \mathbb{E}) \\
&= \frac{1}{n^2} \sum_{i=1}^n \left[\text{Var}(\hat{\tau}_i) + \sum_{j \neq i} \text{Cov}(\hat{\tau}_i, \hat{\tau}_j) \right] \quad (\text{Lemmas 1 and 2}) \\
&= \text{Var}(\hat{\tau}) \quad . \quad \square
\end{aligned}$$

C Exposure-Design and Correlation Clustering

In this section, we prove the relationship between EXPOSURE-DESIGN, its reformulation CORR-CLUST, the previously proposed correlation clustering design of Pouget-Abadie et al. (2019), and other correlation clustering variants. A summary of the results are:

- In Section C.1, we show that the EXPOSURE-DESIGN may be reformulated as the clustering problem, CORR-CLUST.
- In Section C.2, we compare EXPOSURE-DESIGN to the correlation clustering-based design presented in Pouget-Abadie et al. (2019). In particular, we prove that their design is equivalent to EXPOSURE-DESIGN when the trade-off parameter is set as $\phi = 1/(n-1)$ and no constraint is placed on cluster sizes, i.e. $k = m$.
- In Section C.3, we compare CORR-CLUST to other correlation clustering variants. In particular, we prove that (unconstrained) CORR-CLUST may be viewed as an instance of the weighted maximization correlation clustering considered by Charikar et al. (2005); Swamy (2004) but with a possibly large additive constant which prevents an approximation-preserving reduction.

To begin, we demonstrate how to re-write the CORR-CLUST objective using matrix notation. Let $\omega_{i,j} \in \mathbb{R}$ be the weights for pairs $i, j \in [m]$ and let Ω be the m -by- m matrix whose (i, j) th entry is $\omega_{i,j}$. For a partition

\mathcal{C} of the indices $[m]$, let $Z_{\mathcal{C}}$ be the m -by- m matrix where the (i, j) th entry is 1 if i and j are in the same cluster of \mathcal{C} and 0 otherwise. Then, we may express the CORR-CLUST objective as

$$\sum_{C_r \in \mathcal{C}} \sum_{i, j \in C_r} \omega_{i, j} = \sum_{i=1}^n \sum_{j=1}^n \omega_{i, j} [Z_{\mathcal{C}}]_{(i, j)} = \text{tr}(\Omega Z_{\mathcal{C}}) .$$

Throughout the remainder of the section, it will be useful to write the CORR-CLUST objective using this matrix notation.

C.1 Reformulating Exposure-Design as Corr-Clust

We are now ready to prove Proposition 2, which we restate here for completeness.

Proposition 2. *For each pair of diversion units $i, j \in V_d$, define the value $\omega_{i, j} \in \mathbb{R}$ as*

$$\omega_{i, j} = (1 + \phi) \sum_{k=1}^m w_{k, i} w_{k, j} - \phi \left(\sum_{k=1}^m w_{k, i} \right) \left(\sum_{k=1}^m w_{k, j} \right) , \quad (2)$$

where $w_{k, i}$ is the weight of the edge between the k th outcome unit and the i th diversion unit. EXPOSURE-DESIGN is equivalent to the following clustering problem:

$$\max_{\text{clusterings } \mathcal{C}} \sum_{C_r \in \mathcal{C}} \sum_{i, j \in C_r} \omega_{i, j} . \quad (\text{CORR-CLUST})$$

Proof. Recall that the objective of EXPOSURE-DESIGN is defined as

$$\sum_{i=1}^n \text{Var}(x_i) - \phi \sum_{i \neq j} \text{Cov}(x_i, x_j) ,$$

where the expectation in the variance and covariance terms is taken with respect to the random assignment vector $\mathbf{z} \in \{\pm 1\}^m$, which is drawn from the cluster design given by \mathcal{C} . Recall that the exposures are given by $\mathbf{x} = \mathbf{W}\mathbf{z}$. Using matrix notation, we can more compactly represent this objective as

$$\begin{aligned} \sum_{i=1}^n \text{Var}(x_i) - \phi \sum_{i \neq j} \text{Cov}(x_i, x_j) &= \text{tr} \left((\mathbf{I} - \phi(\mathbf{1}\mathbf{1}^{\top} - \mathbf{I})) \text{Cov}(\mathbf{x}) \right) && \text{(rewriting in terms of tr)} \\ &= \text{tr} \left(((1 + \phi)\mathbf{I} - \phi\mathbf{1}\mathbf{1}^{\top}) \text{Cov}(\mathbf{x}) \right) && \text{(rearranging terms)} \\ &= \text{tr} \left(((1 + \phi)\mathbf{I} - \phi\mathbf{1}\mathbf{1}^{\top}) \text{Cov}(\mathbf{W}\mathbf{z}) \right) && \text{(definition of exposure)} \\ &= \text{tr} \left(((1 + \phi)\mathbf{I} - \phi\mathbf{1}\mathbf{1}^{\top}) \mathbf{W} \text{Cov}(\mathbf{z}) \mathbf{W}^{\top} \right) && \text{(property of covariance)} \\ &= \text{tr} \left(\mathbf{W}^{\top} ((1 + \phi)\mathbf{I} - \phi\mathbf{1}\mathbf{1}^{\top}) \mathbf{W} \text{Cov}(\mathbf{z}) \right) && \text{(cyclic property of trace)} \end{aligned}$$

Because \mathbf{z} is drawn from an independent cluster design, the (i, j) th entry of the covariance matrix $\text{Cov}(\mathbf{z})$ is 1 if diversion units i and j are in the same cluster and 0 otherwise. Thus, by the observation above, this clustering objective is a correlation clustering where the weights are given by the matrix

$$\Omega = \mathbf{W}^{\top} ((1 + \phi)\mathbf{I} - \phi\mathbf{1}\mathbf{1}^{\top}) \mathbf{W} .$$

By inspection, we have that the (i, j) th entry of this matrix Ω is

$$\omega_{i, j} = (1 + \phi) \sum_{k=1}^n w_{k, i} w_{k, j} - \phi \left(\sum_{k=1}^n w_{k, i} \right) \left(\sum_{k=1}^n w_{k, j} \right) ,$$

as desired. □

C.2 An instance of Exposure-Design when $\phi = 1/(n - 1)$

Now we demonstrate that the correlation clustering objective proposed in Pouget-Abadie et al. (2019) is a special case of EXPOSURE-DESIGN when $\phi = 1/(n - 1)$ and no constraint is placed on cluster sizes, i.e. $k = m$. Before giving the formal statement, we re-introduce the clustering objective in that paper; that is,

$$\max_{\text{clusterings } \mathcal{C}} \mathbb{E} \left[\sum_{i=1}^n \left(x_i - \left(\frac{1}{n} \sum_{j=1}^n x_j \right) \right)^2 \right], \quad (\text{EXPOSURE-SPREAD})$$

where the expectation is with respect to the treatment vector $\mathbf{z} \in \{\pm 1\}^m$ drawn according to the independent cluster design given by \mathcal{C} . The quantity in the expectation is a measure of the spread of the exposures. We remark that in Pouget-Abadie et al. (2019), the exposures are called “doses” and the quantity in the expectation is referred to as the “empirical dose variance”.

Proposition 3. *Up to additive and multiplicative constants, EXPOSURE-SPREAD is equivalent to EXPOSURE-DESIGN when the trade-off parameter is set to $\phi = 1/(n - 1)$.*

Proof. Let us denote the exposure spread by

$$Q = \sum_{i=1}^n \left(x_i - \left(\frac{1}{n} \sum_{j=1}^n x_j \right) \right)^2,$$

Note that the exposure spread is equal to the ℓ_2 norm of the *de-meanned* exposure vector $\bar{\mathbf{x}} = (\bar{x}_1, \bar{x}_2, \dots, \bar{x}_n)$, where

$$\bar{x}_i = x_i - \left(\frac{1}{n} \sum_{j=1}^n x_j \right).$$

The entire de-meanned exposure vector may be written as $\bar{\mathbf{x}} = (\mathbf{I} - \frac{1}{n} \mathbf{1}\mathbf{1}^T) \mathbf{x}$. Using the fact that this matrix is a projection and that the exposure vector is $\mathbf{x} = \mathbf{W}\mathbf{z}$, we can write the exposure spread as

$$Q = \|\bar{\mathbf{x}}\|^2 = \|(\mathbf{I} - \frac{1}{n} \mathbf{1}\mathbf{1}^T) \mathbf{x}\|^2 = \mathbf{x}^\top (\mathbf{I} - \frac{1}{n} \mathbf{1}\mathbf{1}^T)^2 \mathbf{x} = \mathbf{x}^\top (\mathbf{I} - \frac{1}{n} \mathbf{1}\mathbf{1}^T) \mathbf{x} = \mathbf{z}^\top \mathbf{W}^\top (\mathbf{I} - \frac{1}{n} \mathbf{1}\mathbf{1}^T) \mathbf{W} \mathbf{z}.$$

Finally, the expectation of the exposure spread may be written as

$$\begin{aligned} \mathbb{E}[Q] &= \mathbb{E} \left[\mathbf{z}^\top \mathbf{W}^\top (\mathbf{I} - \frac{1}{n} \mathbf{1}\mathbf{1}^T) \mathbf{W} \mathbf{z} \right] && \text{(from above)} \\ &= \mathbb{E} \left[\text{tr} \left(\mathbf{z}^\top \mathbf{W}^\top (\mathbf{I} - \frac{1}{n} \mathbf{1}\mathbf{1}^T) \mathbf{W} \mathbf{z} \right) \right] && \text{(trace of a scalar)} \\ &= \mathbb{E} \left[\text{tr} \left(\mathbf{W}^\top (\mathbf{I} - \frac{1}{n} \mathbf{1}\mathbf{1}^T) \mathbf{W} \mathbf{z} \mathbf{z}^\top \right) \right] && \text{(cyclic property of trace)} \\ &= \text{tr} \left(\mathbf{W}^\top (\mathbf{I} - \frac{1}{n} \mathbf{1}\mathbf{1}^T) \mathbf{W} \mathbb{E}[\mathbf{z} \mathbf{z}^\top] \right) && \text{(linearity of trace)} \\ &= \text{tr} \left(\mathbf{W}^\top (\mathbf{I} - \frac{1}{n} \mathbf{1}\mathbf{1}^T) \mathbf{W} \text{Cov}(\mathbf{z}) \right) + c, \end{aligned}$$

where the value c in the last line is $c = \text{tr} \left(\mathbf{W}^\top (\mathbf{I} - \frac{1}{n} \mathbf{1}\mathbf{1}^T) \mathbf{W} \mathbb{E}[\mathbf{z}] \mathbb{E}[\mathbf{z}^\top] \right)$, which follows from $\text{Cov}(\mathbf{z}) = \mathbb{E}[\mathbf{z} \mathbf{z}^\top] - \mathbb{E}[\mathbf{z}] \mathbb{E}[\mathbf{z}^\top]$ and linearity of trace. Moreover, when the probability of treatment assignment p is fixed, this value c is a constant with respect to the clustering being chosen.

Observe that by setting $\phi = 1/(n - 1)$ and multiplying by a factor $(n - 1)/n$, the EXPOSURE-DESIGN objective becomes

$$\frac{n - 1}{n} \text{tr} \left(\mathbf{W}^\top \left(\left(1 + \frac{1}{n - 1} \right) \mathbf{I} - \frac{1}{n - 1} \mathbf{1}\mathbf{1}^\top \right) \mathbf{W} \text{Cov}(\mathbf{z}) \right) = \text{tr} \left(\mathbf{W}^\top (\mathbf{I} - \frac{1}{n} \mathbf{1}\mathbf{1}^T) \mathbf{W} \text{Cov}(\mathbf{z}) \right).$$

Thus, the EXPOSURE-SPREAD objective is equivalent (up to additive and multiplicative constants) to the EXPOSURE-DESIGN objective when $\phi = 1/(n - 1)$. \square

C.3 Comparison to other correlation clustering variants

Recall that we defined the objective of the correlation clustering variant CORR-CLUST as

$$\sum_{C_r \in \mathcal{C}} \sum_{i, j \in C_r} \omega_{i, j} ,$$

where $\omega_{i, j}$ is defined for each pair of diversion units $i, j \in V_d$ as

$$\omega_{i, j} = (1 + \phi) \sum_{k=1}^n w_{k, i} w_{k, j} - \phi \left(\sum_{k=1}^n w_{k, i} \right) \left(\sum_{k=1}^n w_{k, j} \right) ,$$

and $w_{k, i}$ is the weight of the edge between the k th outcome unit and the i th diversion unit. Observe that the term $\omega_{i, j}$ can take positive or negative values.

The maximization weighted correlation clustering variant considered by Charikar et al. (2005); Swamy (2004) is defined as follows. Let $G = (V, E)$ be a graph where each edge $e = (i, j) \in E$ has two *non-negative* weights: $w_{in}(i, j)$ and $w_{out}(i, j)$. Given a clustering \mathcal{C} , an edge $e = (i, j)$ is said to be *in-cluster* if i and j are in the same cluster and *out-cluster* otherwise. The objective function for a given clustering is given by

$$\sum_{\substack{\text{in-cluster} \\ \text{edges } e}} w_{in}(e) + \sum_{\substack{\text{out-cluster} \\ \text{edges } e}} w_{out}(e) \quad (\text{CORR-CLUST-CS})$$

We now show that the CORR-CLUST objective may be written as an instance of the CORR-CLUST-CS objective, but with the addition of a large additive constant. Again, we stress that this reduction is primarily for aesthetic comparison purposes because the appearance of the large additive constant prevents any meaningful approximation-preserving reduction.

Proposition 4. *Our formulation CORR-CLUST may be viewed as an instance of CORR-CLUST-CS with a large additive constant. More precisely, let $w_{in}(i, j) = \max\{0, \omega_{i, j}\}$ and $w_{out}(i, j) = \min\{0, \omega_{i, j}\}$. For a clustering \mathcal{C} , we have that the objectives are related by*

$$\sum_{C_r \in \mathcal{C}} \sum_{i, j \in C_r} \omega_{i, j} - \sum_{i=1}^n \sum_{j=1}^n \min\{0, \omega_{i, j}\} = \sum_{\substack{\text{in-cluster} \\ \text{edges } e}} w_{in}(e) + \sum_{\substack{\text{out-cluster} \\ \text{edges } e}} w_{out}(e)$$

Proof. For each pair of diversion units i, j , define $\omega_{i, j}^+ = \max\{0, \omega_{i, j}\}$ and $\omega_{i, j}^- = -\min\{0, \omega_{i, j}\}$. Observe that $\omega_{i, j} = \omega_{i, j}^+ + \omega_{i, j}^-$ and so we can re-distribute the following sum as

$$\sum_{C_r \in \mathcal{C}} \sum_{i, j \in C_r} \omega_{i, j} = \sum_{C_r \in \mathcal{C}} \sum_{i, j \in C_r} (\omega_{i, j}^+ + \omega_{i, j}^-) = \sum_{C_r \in \mathcal{C}} \sum_{i, j \in C_r} \omega_{i, j}^+ + \sum_{C_r \in \mathcal{C}} \sum_{i, j \in C_r} \omega_{i, j}^- .$$

Subtracting the (instance-dependent) constant $\sum_{i=1}^n \sum_{j=1}^n \min\{0, \omega_{i, j}\}$ from both sides and rearranging yields

$$\begin{aligned} \sum_{C_r \in \mathcal{C}} \sum_{i, j \in C_r} \omega_{i, j} - \sum_{i=1}^n \sum_{j=1}^n \min\{0, \omega_{i, j}\} &= \sum_{C_r \in \mathcal{C}} \sum_{i, j \in C_r} \omega_{i, j}^+ + \sum_{C_r \in \mathcal{C}} \sum_{i, j \in C_r} \omega_{i, j}^- - \sum_{i=1}^n \sum_{j=1}^n \min\{0, \omega_{i, j}\} \\ &= \sum_{C_r \in \mathcal{C}} \sum_{i, j \in C_r} \omega_{i, j}^+ - \sum_{\substack{C_r \neq C'_r \in \mathcal{C} \\ i \in C_r \\ j \in C'_r}} \omega_{i, j}^- \\ &= \sum_{C_r \in \mathcal{C}} \sum_{i, j \in C_r} \omega_{i, j}^+ + \sum_{\substack{C_r \neq C'_r \in \mathcal{C} \\ i \in C_r \\ j \in C'_r}} (-\omega_{i, j}^-) \\ &= \sum_{\substack{\text{in-cluster} \\ \text{edges } e}} w_{in}(e) + \sum_{\substack{\text{out-cluster} \\ \text{edges } e}} w_{out}(e) . \end{aligned}$$

Finally, observe that for each pair (i, j) , the values $w_{in}(i, j)$ and $w_{out}(i, j)$ are non-negative so that the final equation is a valid objective function for the CORR-CLUST-CS formulation. \square