

**FINITE DIMENSIONAL NICHOLS ALGEBRAS OVER SUZUKI
ALGEBRA II: OVER SIMPLE YETTER-DRINFELD MODULES OF**

$$A_{N2n+1}^{\mu\lambda}$$

YUXING SHI

ABSTRACT. In this paper, the author gives a complete set of simple Yetter-Drinfeld modules over Suzuki algebra $A_{N2n+1}^{\mu\lambda}$ [Suz98] and investigates the Nichols algebras over those irreducible Yetter-Drinfeld modules. The finite dimensional Nichols algebras of diagonal type are of Cartan type $A_1, A_1 \times A_1, A_2$, Super type $A_2(q; \mathbb{I}_2)$ and the Nichols algebra $\mathfrak{uf}\mathfrak{o}(8)$. And the involved finite dimensional Nichols algebras of non-diagonal type are 12, $4m$ and m^2 dimensional. The left three unsolved cases are set as open problems.

1. INTRODUCTION

Let \mathbb{k} be an algebraically closed field of characteristic 0. The paper is a sequel of [Shi20] which have started the classification project of finite dimensional Hopf algebras over Suzuki algebra $A_{Nn}^{\mu\lambda}$.

In 2004, Menini et al. studied the quantum lines over \mathcal{A}_{4m} and \mathcal{B}_{4m} , which are isomorphic to A_{1m}^{++} and A_{1m}^{+-} respectively[CDMM04]. In 2019, the author [Shi19] classified finite dimensional Hopf algebras over the Kac-Paljutkin algebra A_{12}^{+-} and Fantino et al. [FGM19] classified finite dimensional Hopf algebras over the dual of dihedral group \widehat{D}_{2m} of order $2m$, with $m = 4a \geq 12$, where \widehat{D}_{2m} is a 2-cocycle deformation of A_{12a}^{++} [Mas00]. In [Shi20], the author gave a complete set of simple Yetter-Drinfeld modules over Suzuki algebra $A_{N2n}^{\mu\lambda}$ and investigated the Nichols algebras over those irreducible Yetter-Drinfeld modules. In this paper, we are going to deal with the case $A_{N2n+1}^{\mu\lambda}$.

Our classification project over Suzuki algebra is based on the *lifting method* which was introduced by Andruskiewitsch and Schneider [AS98]. The lifting method is a general framework to classify finite dimensional Hopf algebras with a fixed sub-Hopf algebra as coradical. Here let us recall the procedure for the

2010 *Mathematics Subject Classification.* 16T05, 16T25, 17B22.

Keywords: Nichols algebra; Hopf algebra; Suzuki algebra.

This work was partially supported by Foundation of Jiangxi Educational Committee (No.12020447) and Natural Science Foundation of Jiangxi Normal University(No. 12018937) .

lifting method briefly. Let H be a Hopf algebra whose coradical H_0 is a Hopf sub-algebra. The associated graded Hopf algebra of H is isomorphic to $R\#H_0$ where $R = \bigoplus_{n \in \mathbb{N}_0} R(n)$ is a braided Hopf algebra in the category ${}^{H_0}\mathcal{YD}$ of Yetter-Drinfeld modules over H_0 , $\#$ stands for the Radford *biproduct* or *bosonization* of R with H_0 . As explained in [AS02], to classify finite-dimensional Hopf algebras H whose coradical is isomorphic to H_0 we have to deal with the following questions:

- (a) Determine all Yetter-Drinfeld modules V over H_0 such that the Nichols algebra $\mathfrak{B}(V)$ has finite dimension; find an efficient set of relations for $\mathfrak{B}(V)$.
- (b) If $R = \bigoplus_{n \in \mathbb{N}_0} R(n)$ is a finite-dimensional Hopf algebra in ${}^{H_0}\mathcal{YD}$ with $V = R(1)$, decide if $R \simeq \mathfrak{B}(V)$. Here $V = R(1)$ is a braided vector space called the *infinitesimal braiding*.
- (c) Given V as in (a), classify all H such that $\text{gr } H \simeq \mathfrak{B}(V)\#H_0$ (lifting).

According to Radford's method [Rad03, Proposition 2], we constructed a complete set of simple Yetter-Drinfeld modules over $A_{N2n+1}^{\mu\lambda}$. There are exactly $8N^2$ one-dimensional, $8N^2n(n+1)$ two-dimensional and $8N^2(2n+1)$ -dimensional non-isomorphic Yetter-Drinfeld modules over $A_{N2n+1}^{\mu\lambda}$, see the Theorem 3.1.

The finite dimensional Nichols algebras of diagonal type over simple Yetter-Drinfeld modules of $A_{N2n+1}^{\mu\lambda}$ can be classified by the following theorem.

Theorem 1.1. *Let M be a simple Yetter-Drinfeld module over $A_{N2n+1}^{\mu\lambda}$. If $\mathfrak{B}(M)$ is of diagonal type and $\dim \mathfrak{B}(M) < \infty$, then $\mathfrak{B}(M)$ can be classified as follows.*

- (1) Cartan type A_1 , see Lemmas 4.1 and 4.3;
- (2) Cartan type $A_1 \times A_1$, see Lemmas 4.5, 4.6, 4.7, 4.8, 4.9, 4.10 and 4.14;
- (3) Cartan type A_2 , see Lemmas 4.5, 4.6, 4.7, 4.8, 4.9, 4.10 and 4.14;
- (4) Super type $\mathbf{A}_2(q; \mathbb{I}_2)$, see Lemmas 4.5 and 4.6;
- (5) The Nichols algebra $\text{uf}\mathfrak{o}(8)$, see Lemmas 4.5 and 4.6.

There are three unsolved cases which are difficult in general in the paper.

Problem 1.2. Determine the dimensions of the following Nichols algebras.

- (1) $\mathfrak{B}(\mathcal{L}_{k,pq}^s)$, $n \geq 2$, see the section 4.3;
- (2) $\mathfrak{B}(V_{abe})$, $b^2 \neq ae \neq 1$, $b^2 \neq (ae)^{-1}$, $b \in \mathbb{G}_m$ for $m \geq 3$;
- (3) $\mathfrak{B}(\mathcal{N}_{k,pq}^s)$, $n \geq 2$, see the section 4.4.

The paper is organized as follows. In section 1, we introduce the background of the paper and summarize our main results. In section 2, we make an introduction for the Suzuki algebra and construct all simple representation of $A_{N2n+1}^{\mu\lambda}$. In section 3, we construct all simple Yetter-Drinfeld modules over $A_{N2n+1}^{\mu\lambda}$ by using Radford's method and we put those Yetter-Drinfeld modules in the appendix. In section 4, we calculate Nichols algebras over simple Yetter-Drinfeld modules of $A_{N2n+1}^{\mu\lambda}$.

2. THE REPRESENTATIONS OF THE HOPF ALGEBRA $A_{N2n+1}^{\mu\lambda}$

Suzuki introduced a family of cosemisimple Hopf algebras $A_{Nn}^{\mu\lambda}$ parametrized by integers $N \geq 1$, $n \geq 2$ and $\mu, \lambda = \pm 1$, and investigated various properties and structures of them [Suz98]. Wakui studied the Suzuki algebra $A_{Nn}^{\mu\lambda}$ in perspectives of polynomial invariant[Wak10], braided Morita invariant[Wak19] and coribbon structures[Wak03]. The Hopf algebra $A_{Nn}^{\mu\lambda}$ is generated by $x_{11}, x_{12}, x_{21}, x_{22}$ subject to the relations:

$$\begin{aligned} x_{11}^2 &= x_{22}^2, & x_{12}^2 &= x_{21}^2, & \chi_{21}^n &= \lambda \chi_{12}^n, & \chi_{11}^n &= \chi_{22}^n, \\ x_{11}^{2N} + \mu x_{12}^{2N} &= 1, & x_{ij}x_{kl} &= 0 \text{ whenever } i + j + k + l \text{ is odd,} \end{aligned}$$

where we use the following notation for $m \geq 1$,

$$\begin{aligned} \chi_{11}^m &:= \overbrace{x_{11}x_{22}x_{11} \dots}^m, & \chi_{22}^m &:= \overbrace{x_{22}x_{11}x_{22} \dots}^m, \\ \chi_{12}^m &:= \overbrace{x_{12}x_{21}x_{12} \dots}^m, & \chi_{21}^m &:= \overbrace{x_{21}x_{12}x_{21} \dots}^m. \end{aligned}$$

The Hopf algebra structure of $A_{Nn}^{\mu\lambda}$ is given by

$$(2.1) \quad \Delta(\chi_{ij}^k) = \chi_{i1}^k \otimes \chi_{1j}^k + \chi_{i2}^k \otimes \chi_{2j}^k, \quad \varepsilon(x_{ij}) = \delta_{ij}, \quad S(x_{ij}) = x_{ji}^{4N-1},$$

for $k \geq 1$, $i, j = 1, 2$.

Let $\overline{i}, \overline{i+j} = \{i, i+1, i+2, \dots, i+j\}$ be an index set. Then the basis of $A_{Nn}^{\mu\lambda}$ can be represented by

$$(2.2) \quad \{x_{11}^s \chi_{22}^t, x_{12}^s \chi_{21}^t \mid s \in \overline{1, 2N}, t \in \overline{0, n-1}\}.$$

Thus for $s, t \geq 0$ with $s+t \geq 1$,

$$\begin{aligned} \Delta(x_{11}^s \chi_{22}^t) &= x_{11}^s \chi_{22}^t \otimes x_{11}^s \chi_{22}^t + x_{12}^s \chi_{21}^t \otimes x_{21}^s \chi_{12}^t, \\ \Delta(x_{12}^s \chi_{21}^t) &= x_{11}^s \chi_{22}^t \otimes x_{12}^s \chi_{21}^t + x_{12}^s \chi_{21}^t \otimes x_{22}^s \chi_{11}^t. \end{aligned}$$

The cosemisimple Hopf algebra $A_{Nn}^{\mu\lambda}$ is decomposed to the direct sum of simple subcoalgebras such as $A_{Nn}^{\mu\lambda} = \bigoplus_{g \in G} \mathbb{k}g \oplus \bigoplus_{\substack{0 \leq s \leq N-1 \\ 1 \leq t \leq n-1}} C_{st}$ [Suz98, Theorem 3.1][Wak10, Proposition 5.5], where

$$\begin{aligned} G &= \{x_{11}^{2s} \pm x_{12}^{2s}, x_{11}^{2s+1} \chi_{22}^{n-1} \pm \sqrt{\lambda} x_{12}^{2s+1} \chi_{21}^{n-1} \mid s \in \overline{1, N}\}, \\ C_{st} &= \mathbb{k}x_{11}^{2s} \chi_{11}^t + \mathbb{k}x_{12}^{2s} \chi_{12}^t + \mathbb{k}x_{11}^{2s} \chi_{22}^t + \mathbb{k}x_{12}^{2s} \chi_{21}^t, \quad s \in \overline{1, N}, t \in \overline{1, n-1}. \end{aligned}$$

The set $\{\mathbb{k}g \mid g \in G\} \cup \{\mathbb{k}x_{11}^{2s} \chi_{11}^t + \mathbb{k}x_{12}^{2s} \chi_{21}^t \mid s \in \overline{1, N}, t \in \overline{1, n-1}\}$ is a full set of non-isomorphic simple left $A_{Nn}^{\mu\lambda}$ -comodules, where the coactions of the comodules listed above are given by the coproduct Δ . Denote the comodule $\mathbb{k}x_{11}^{2s} \chi_{11}^t + \mathbb{k}x_{12}^{2s} \chi_{21}^t$ by Λ_{st} . That is to say the comodule $\Lambda_{st} = \mathbb{k}w_1 + \mathbb{k}w_2$ is defined as

$$\rho(w_1) = x_{11}^{2s} \chi_{11}^t \otimes w_1 + x_{12}^{2s} \chi_{12}^t \otimes w_2, \quad \rho(w_2) = x_{11}^{2s} \chi_{22}^t \otimes w_2 + x_{12}^{2s} \chi_{21}^t \otimes w_1.$$

Proposition 2.1. *Let ω be a primitive $4(2n + 1)N$ -th root of unity, and*

$$\tilde{\mu} = \begin{cases} 1, & \mu = 1, \\ \omega^{2n+1}, & \mu = -1, \end{cases} \quad \bar{\mu} = \begin{cases} 1, & \mu = 1, \\ \omega^{2(2n+1)}, & \mu = -1. \end{cases}$$

Then a full set of non-isomorphic simple left $A_{N2n+1}^{\mu\lambda}$ -modules is given by

(1) V_k , $k \in \overline{0, 2N - 1}$. *The action of $A_{N2n+1}^{\mu\lambda}$ on V_k is given by*

$$x_{12} \mapsto 0, \quad x_{21} \mapsto 0, \quad x_{11} \mapsto \omega^{2k(2n+1)}, \quad x_{22} \mapsto \omega^{2k(2n+1)};$$

(2) V'_k , $k \in \overline{0, 2N - 1}$. *The action of $A_{N2n+1}^{\mu\lambda}$ on V'_k is given by*

$$x_{11} \mapsto 0, \quad x_{22} \mapsto 0, \quad x_{12} \mapsto \omega^{2k(2n+1)}\tilde{\mu}, \quad x_{21} \mapsto \omega^{2k(2n+1)}\tilde{\mu}\lambda;$$

(3) V_{jk} , $k \in \overline{0, N - 1}$, $\frac{j}{2} \in \overline{1, n}$. *The action of $A_{N2n+1}^{\mu\lambda}$ on the row vector (v_1, v_2) for $V_{jk} = \mathbb{k}v_1 \oplus \mathbb{k}v_2$ is given by*

$$x_{11} \mapsto \begin{pmatrix} 0 & \omega^{4k(2n+1)-2jN} \\ \omega^{2jN} & 0 \end{pmatrix}, \quad x_{12}, x_{21} \mapsto 0, \quad x_{22} \mapsto \begin{pmatrix} 0 & \omega^{4k(2n+1)} \\ 1 & 0 \end{pmatrix};$$

(4) V'_{jk} , $k \in \overline{0, N - 1}$, $\frac{j}{2} \in \overline{1, n}$. *The action of $A_{N2n+1}^{\mu\lambda}$ on the row vector (v'_1, v'_2) for $V'_{jk} = \mathbb{k}v'_1 \oplus \mathbb{k}v'_2$ is given by*

$$x_{21} \mapsto \begin{pmatrix} 0 & \lambda\bar{\mu}\omega^{4k(2n+1)-2jN} \\ \lambda\omega^{2jN} & 0 \end{pmatrix}, \quad x_{11}, x_{22} \mapsto 0, \quad x_{12} \mapsto \begin{pmatrix} 0 & \bar{\mu}\omega^{4k(2n+1)} \\ 1 & 0 \end{pmatrix}.$$

Remark 2.2. We left the proof to the readers since it's easy and tedious.

3. YETTER-DRINFELD MODULES OVER $A_{N2n+1}^{\mu\lambda}$

Similarly according to Radford's method [Rad03, Proposition 2], any simple left Yetter-Drinfeld module over a Hopf algebra H could be constructed by the submodule of tensor product of a left module V of H and H itself, where the module and comodule structures are given by :

$$(3.1) \quad h \cdot (\ell \boxtimes g) = (h_{(2)} \cdot \ell) \boxtimes h_{(1)}gS(h_{(3)}),$$

$$(3.2) \quad \rho(\ell \boxtimes h) = h_{(1)} \otimes (\ell \boxtimes h_{(2)}), \forall h, g \in H, \ell \in V.$$

Here we use \boxtimes instead of \otimes to avoid confusion by using too many symbols of the tensor product. we construct all simple left Yetter-Drinfeld modules over $A_{N2n+1}^{\mu\lambda}$ in this way and put them in the appendix without proof since it's tedious verification with the definition of Yetter-Drinfeld modules. Firstly, it's easy to see that there are $8N^2$ pairwise non-isomorphic simple Yetter-Drinfeld modules of one-dimension from the Table 1. Secondly, let us take a closer look at those Yetter-Drinfeld modules which are isomorphic as $A_{N2n+1}^{\mu\lambda}$ -modules and $A_{N2n+1}^{\mu\lambda}$ -comodules, but the modules isomorphism and comodules isomorphism are incompatible. For example:

I Yetter-Drinfeld modules in the part (3), (4) of the list are non-isomorphic since

$$\begin{aligned}\mathcal{C}_{jk,p}^{st} &= \mathbb{k}w_1 \oplus \mathbb{k}w_2 \simeq V_{jk}, & \mathcal{C}_{jk,p}^{st} &= \mathbb{k}w_2 \oplus \mathbb{k}\left((-1)^p \omega^{2k(2n+1)-jN} w_1\right) \simeq \Lambda_{s, 2t+2}, \\ \mathcal{D}_{jk,p}^{st} &= \mathbb{k}w_1 \oplus \mathbb{k}w_2 \simeq V_{jk}, & \mathcal{D}_{jk,p}^{st} &= \mathbb{k}w_1 \oplus \mathbb{k}\left((-1)^p \omega^{jN-2k(2n+1)} w_2\right) \simeq \Lambda_{s, 2t+2}.\end{aligned}$$

II Yetter-Drinfeld modules in the part (8), (9) of the list are non-isomorphic since

$$\begin{aligned}\mathcal{H}_{jk,p}^{st} &= \mathbb{k}w_1 \oplus \mathbb{k}w_2 \simeq V'_{jk}, & \mathcal{H}_{jk,p}^{st} &= \mathbb{k}w_2 \oplus \mathbb{k}\left((-1)^p \sqrt{\lambda\bar{\mu}} \omega^{2k(2n+1)-jN} w_1\right) \simeq \Lambda_{s, 2t+1}, \\ \mathcal{I}_{jk,p}^{st} &= \mathbb{k}w_1 \oplus \mathbb{k}w_2 \simeq V'_{jk}, & \mathcal{I}_{jk,p}^{st} &= \mathbb{k}w_1 \oplus \mathbb{k}\left(\frac{(-1)^p}{\sqrt{\lambda\bar{\mu}}} \omega^{jN-2k(2n+1)} w_2\right) \simeq \Lambda_{s, 2t+1}.\end{aligned}$$

Now we can see that there are $8N^2n(n+1)$ pairwise non-isomorphic simple Yetter-Drinfeld modules of two-dimension from the Table 1. While we break the Yetter-Drinfeld module $M \boxtimes A_{N, 2n+1}^{\mu\lambda}$ into small Yetter-Drinfeld modules for any simple left $A_{N, 2n+1}^{\mu\lambda}$ -module M , there are four class Yetter-Drinfeld modules in total whose dimensions are greater than two. And they have the relations $\mathcal{K}_{k,p}^s \simeq \mathcal{L}_{k,0p}^s$, $\mathcal{M}_{k,p}^s \simeq \mathcal{N}_{k,0p}^s$ as Yetter-Drinfeld modules. Since

$$8N^2 \cdot 1^2 + 8N^2n(n+1) \cdot 2^2 + 8N^2 \cdot (2n+1)^2 = [4N(2n+1)]^2,$$

the list is a complete set of simple Yetter-Drinfeld modules.

Theorem 3.1. *A complete set of simple Yetter-Drinfeld modules over Suzuki algebra $A_{N, 2n+1}^{\mu\lambda}$ is given as follows.*

- (1) *There are $8N^2$ pairwise non-isomorphic Yetter-Drinfeld modules of one dimension:*
 - (a) $\mathcal{A}_{k,p}^s$, $s \in \overline{1, N}$, $k \in \overline{0, 2N-1}$, $p \in \mathbb{Z}_2$;
 - (b) $\mathcal{B}_{k,p}^s$, $s \in \overline{1, N}$, $k \in \overline{0, 2N-1}$, $p \in \mathbb{Z}_2$.
- (2) *There are $8N^2n(n+1)$ pairwise non-isomorphic Yetter-Drinfeld modules of two dimension:*
 - (a) $\mathcal{C}_{jk,p}^{st}$, $s \in \overline{1, N}$, $t \in \overline{0, n-1}$, $\frac{j}{2} \in \overline{1, n}$, $k \in \overline{0, N-1}$, $p \in \mathbb{Z}_2$;
 - (b) $\mathcal{D}_{jk,p}^{st}$, $s \in \overline{1, N}$, $t \in \overline{0, n-1}$, $\frac{j}{2} \in \overline{1, n}$, $k \in \overline{0, N-1}$, $p \in \mathbb{Z}_2$;
 - (c) $\mathcal{E}_{jk,p}^s$, $s \in \overline{1, N}$, $\frac{j}{2} \in \overline{1, n}$, $k \in \overline{0, N-1}$, $p \in \mathbb{Z}_2$;
 - (d) $\mathcal{F}_{k,p}^{st}$, $s \in \overline{1, N}$, $t \in \overline{0, n-1}$, $k \in \overline{0, 2N-1}$, $p \in \mathbb{Z}_2$;
 - (e) $\mathcal{G}_{k,p}^{st}$, $s \in \overline{1, N}$, $t \in \overline{0, n-1}$, $k \in \overline{0, 2N-1}$, $p \in \mathbb{Z}_2$;
 - (f) $\mathcal{H}_{jk,p}^{st}$, $s \in \overline{1, N}$, $t \in \overline{0, n-1}$, $\frac{j}{2} \in \overline{1, n}$, $k \in \overline{0, N-1}$, $p \in \mathbb{Z}_2$;
 - (g) $\mathcal{I}_{jk,p}^{st}$, $s \in \overline{1, N}$, $t \in \overline{0, n-1}$, $\frac{j}{2} \in \overline{1, n}$, $k \in \overline{0, N-1}$, $p \in \mathbb{Z}_2$;
 - (h) $\mathcal{J}_{jk,p}^{st}$, $s \in \overline{1, N}$, $t = n$, $\frac{j}{2} \in \overline{1, n}$, $k \in \overline{0, N-1}$, $p \in \mathbb{Z}_2$.
- (3) *There are $8N^2$ pairwise non-isomorphic Yetter-Drinfeld modules of $2n+1$ dimension:*

- (a) $\mathcal{L}_{k,pq}^s$, $s \in \overline{1, N}$, $k \in \overline{0, N-1}$, $p, q \in \mathbb{Z}_2$;
 (b) $\mathcal{N}_{k,pq}^s$, $s \in \overline{1, N}$, $k \in \overline{0, N-1}$, $p, q \in \mathbb{Z}_2$.

Remark 3.2. As for the description of those simple Yetter-Drinfeld modules, please see the Appendix.

4. NICHOLS ALGEBRAS OVER $A_{N2n+1}^{\mu\lambda}$

In this section, we investigate Nichols algebras over simple Yetter-Drinfeld modules of $A_{N2n+1}^{\mu\lambda}$. So the Yetter-Drinfeld modules discussed in the section are those listed in the Theorem 3.1. For the knowledge about Nichols algebras, please refer to [AS02] [And17] [AA17].

4.1. Nichols algebras of diagonal type. Let $V = \bigoplus_{i \in I} \mathbb{k}v_i$ be a vector space with a braiding $c(v_i \otimes v_j) = q_{ij}v_j \otimes v_i$, then the Nichols algebra $\mathfrak{B}(V)$ is of diagonal type. Our results in this section heavily rely on Heckenberger's classification work [Hec09]. To keep the article concise, we don't repeat this in the following proofs. For more details about Nichols algebras of diagonal type, please consult [AA17].

Lemma 4.1. *Let $s \in \overline{1, N}$, $k \in \overline{0, 2N-1}$, then*

$$\dim \mathfrak{B}(\mathcal{A}_{k,p}^s) = \begin{cases} \infty, & N \mid ks, \\ \frac{N}{(d,N)}, & ks \equiv d \pmod{N}, d \in \overline{1, N-1}. \end{cases}$$

Remark 4.2. When $N = 1$, then $\dim \mathfrak{B}(\mathcal{A}_{k,p}^s) = \infty$.

When $N = 2$, then $\dim \mathfrak{B}(\mathcal{A}_{k,p}^s) = \begin{cases} 2, & s = 1, k = 1 \text{ or } 3, \\ \infty, & \text{otherwise.} \end{cases}$

When N is a prime, then $\dim \mathfrak{B}(\mathcal{A}_{k,p}^s) = \begin{cases} N, & s, k \not\equiv 0 \pmod{N}, \\ \infty, & \text{otherwise.} \end{cases}$

Proof. $c(w \otimes w) = \omega^{4ks(2n+1)}w \otimes w$. □

Lemma 4.3. *Let $s \in \overline{1, N}$, $k \in \overline{0, 2N-1}$, $p \in \mathbb{Z}_2$,*

(1) *when $\lambda = \mu = 1$, then*

$$\dim \mathfrak{B}(\mathcal{B}_{k,p}^s) = \begin{cases} \infty, & 2N \mid [pN + k(2s + 1 + 2n)], \\ \frac{2N}{(2N,d)}, & pN + k(2s + 1 + 2n) \equiv d \pmod{2N}, d \in \overline{1, 2N-1}; \end{cases}$$

(2) *when $\lambda = -1$, $\mu = 1$, then*

$$\dim \mathfrak{B}(\mathcal{B}_{k,p}^s) = \begin{cases} \infty, & 4N \mid [N(2p + 2n + 1) + 2k(2s + 1 + 2n)], \\ \frac{4N}{(4N,d)}, & [N(2p + 2n + 1) + 2k(2s + 1 + 2n)] \equiv d \pmod{4N}, \end{cases}$$

where $d \in \overline{1, 4N-1}$;

	dim	parameters	mod	comod
$\mathcal{A}_{k,p}^s$	1	$s \in \overline{1, N}, p \in \mathbb{Z}_2$ $k \in \overline{0, 2N-1}$	V_k	$\mathbb{k}g_s^+, p = 0$ $\mathbb{k}g_s^-, p = 1$
$\mathcal{B}_{k,p}^s$	1	$s \in \overline{1, N}, p \in \mathbb{Z}_2$ $k \in \overline{0, 2N-1}$	V'_k	$\mathbb{k}h_s^+, p = 0$ $\mathbb{k}h_s^-, p = 1$
$\mathcal{C}_{jk,p}^{st}$	2	$s \in \overline{1, N}, t \in \overline{0, n-1},$ $\frac{j}{2} \in \overline{1, n}, p \in \mathbb{Z}_2,$ $k \in \overline{0, N-1}$	V_{jk}	$\Lambda_{s, 2t+2}$
$\mathcal{D}_{jk,p}^{st}$	2	$s \in \overline{1, N}, t \in \overline{0, n-1},$ $\frac{j}{2} \in \overline{1, n}, p \in \mathbb{Z}_2,$ $k \in \overline{0, N-1}$	V_{jk}	$\Lambda_{s, 2t+2}$
$\mathcal{E}_{jk,p}^s$	2	$s \in \overline{1, N},$ $\frac{j}{2} \in \overline{1, n}, p \in \mathbb{Z}_2,$ $k \in \overline{0, N-1}$	V_{jk}	$\mathbb{k}g_s^+ \oplus \mathbb{k}g_s^-$
$\mathcal{F}_{k,p}^{st}$	2	$s \in \overline{1, N}, t \in \overline{0, n-1},$ $k \in \overline{0, 2N-1}, p \in \mathbb{Z}_2$	$V_k \oplus V_d,$ $k + N \equiv d \pmod{2N}$	$\Lambda_{s, 2t+2}$
$\mathcal{G}_{k,p}^{st}$	2	$s \in \overline{1, N}, t \in \overline{0, n-1},$ $k \in \overline{0, 2N-1}, p \in \mathbb{Z}_2$	$V'_k \oplus V'_d,$ $k + N \equiv d \pmod{2N}$	$\Lambda_{s, 2t+1}$
$\mathcal{H}_{jk,p}^{st}$	2	$s \in \overline{1, N}, t \in \overline{0, n-1},$ $\frac{j}{2} \in \overline{1, n}, p \in \mathbb{Z}_2,$ $k \in \overline{0, N-1}$	V'_{jk}	$\Lambda_{s, 2t+1}$
$\mathcal{I}_{jk,p}^{st}$	2	$s \in \overline{1, N}, t \in \overline{0, n-1},$ $\frac{j}{2} \in \overline{1, n}, p \in \mathbb{Z}_2,$ $k \in \overline{0, N-1}$	V'_{jk}	$\Lambda_{s, 2t+1}$
	2	$s \in \overline{1, N}, t = n,$ $\frac{j}{2} \in \overline{1, n}, p \in \mathbb{Z}_2,$ $k \in \overline{0, N-1}$	V'_{jk}	$\mathbb{k}h_s^+ \oplus \mathbb{k}h_s^-$

TABLE 1. Simple Yetter-Drinfeld modules of one or two dimension. Here $g_s^\pm = x_{11}^{2s} \pm x_{12}^{2s}$, $h_s^\pm = x_{11}^{2s+1} \chi_{22}^{2n} \pm \sqrt{\lambda} x_{12}^{2s+1} \chi_{21}^{2n}$.

(3) when $\lambda = 1, \mu = -1$, then

$$\dim \mathfrak{B}(\mathcal{B}_{k,p}^s) = \begin{cases} \infty, & 4N \mid [2Np + (2s+1+2n)(2k+1)], \\ \frac{4N}{(4N,d)}, & [2Np + (2s+1+2n)(2k+1)] \equiv d \pmod{4N}, d \in \overline{1, 4N-1}; \end{cases}$$

(4) when $\lambda = \mu = -1$, then

$$\dim \mathfrak{B}(\mathcal{B}_{k,p}^s) = \begin{cases} \infty, & 4N \mid [N(2p+2n+1) + (2s+1+2n)(2k+1)], \\ \frac{4N}{(4N,d)}, & [N(2p+2n+1) + (2s+1+2n)(2k+1)] \equiv d \pmod{4N}, \end{cases}$$

where $d \in \overline{1, 4N-1}$.

Remark 4.4. When $N = 1$, then

$$\dim \mathfrak{B}(\mathcal{B}_{k,p}^s) = \begin{cases} 2, & \mu = \lambda = 1, s = 1, (k, p) = (0, 1) \text{ or } (1, 0); \\ 4, & \mu = 1, \lambda = -1, s = 1, k \in \overline{0, 1}, p \in \mathbb{Z}_2; \\ 4, & \mu = -1, \lambda = 1, s = 1, k \in \overline{0, 1}, p \in \mathbb{Z}_2; \\ 2, & \mu = \lambda = -1, s = 1, (k, p) = (0, 1) \text{ or } (1, 0); \\ \infty, & \text{otherwise.} \end{cases}$$

Proof. The braiding of $\mathfrak{B}(\mathcal{B}_{k,p}^s)$ is given by

$$c(w \otimes w) = (-1)^p \sqrt{\lambda} \lambda^n \tilde{\mu}^{2s+1+2n} \omega^{2k(2n+1)(2s+1+2n)} w \otimes w$$

$$= \begin{cases} \omega^{2(2n+1)[pN+k(2s+1+2n)]} w \otimes w, & \lambda = \mu = 1, \\ \omega^{(2n+1)[N(2p+2n+1)+2k(2s+1+2n)]} w \otimes w, & \lambda = -1, \mu = 1, \\ \omega^{(2n+1)[2Np+(2s+1+2n)(2k+1)]} w \otimes w, & \lambda = 1, \mu = -1, \\ \omega^{(2n+1)[N(2p+1+2n)+(2s+1+2n)(2k+1)]} w \otimes w, & \lambda = \mu = -1. \end{cases}$$

□

Lemma 4.5. *Denote*

$$\alpha = 2Nj(t+1) + 4k(2n+1)(t+1+s), \quad \beta = -4Nj(t+1) + 8k(2n+1)(t+1+s),$$

then $\dim \mathfrak{B}(\mathcal{C}_{jk,p}^{st}) < \infty$ iff one of the following conditions holds.

- (1) $4N(2n+1) \nmid \alpha$, $4N(2n+1) \mid \beta$, Cartan type $A_1 \times A_1$;
- (2) $4N(2n+1) \nmid \alpha$, $4N(2n+1) \mid (\alpha + \beta)$, Cartan type A_2 ;
- (3) $\alpha \equiv 2N(2n+1) \pmod{4N(2n+1)}$, $\beta \not\equiv 0$ and $2N(2n+1) \pmod{4N(2n+1)}$,
Super type $\mathbf{A}_2(q; \mathbb{I}_2)$;
- (4) $\alpha - 2\beta \equiv 6\beta \equiv 2N(2n+1) \pmod{4N(2n+1)}$, $4\beta \not\equiv 0 \pmod{4N(2n+1)}$.
The Nichols algebras $\text{uf}\mathfrak{o}(8)$, see [AA17, Page 209].

Proof. The braiding of $\mathfrak{B}(\mathcal{C}_{jk,p}^{st})$ is given by

$$c(w_1 \otimes w_1) = \omega^{[2jN+4k(2n+1)](t+1)+4k(2n+1)s} w_1 \otimes w_1$$

$$c(w_1 \otimes w_2) = \omega^{[4k(2n+1)-2jN](t+1)+4k(2n+1)s} w_2 \otimes w_1$$

$$c(w_2 \otimes w_1) = \omega^{[4k(2n+1)-2jN](t+1)+4k(2n+1)s} w_1 \otimes w_2$$

$$c(w_2 \otimes w_2) = \omega^{[4k(2n+1)+2jN](t+1)+4k(2n+1)s} w_2 \otimes w_2.$$

□

Lemma 4.6. *Denote*

$$\alpha = -2Nj(t+1) + 4k(2n+1)(t+1+s), \quad \beta = 4Nj(t+1) + 8k(2n+1)(t+1+s),$$

then $\dim \mathfrak{B}(\mathcal{D}_{jk,p}^{st}) < \infty$ iff one of the following conditions holds.

- (1) $4N(2n+1) \nmid \alpha$, $4N(2n+1) \mid \beta$, Cartan type $A_1 \times A_1$;
- (2) $4N(2n+1) \nmid \alpha$, $4N(2n+1) \mid (\alpha + \beta)$, Cartan type A_2 ;
- (3) $\alpha \equiv 2N(2n+1) \pmod{4N(2n+1)}$, $\beta \not\equiv 0$ and $2N(2n+1) \pmod{4N(2n+1)}$,
Super type $\mathbf{A}_2(q; \mathbb{I}_2)$;
- (4) $\alpha - 2\beta \equiv 6\beta \equiv 2N(2n+1) \pmod{4N(2n+1)}$, $4\beta \not\equiv 0 \pmod{4N(2n+1)}$.
The Nichols algebras $\text{uf}\mathfrak{b}(8)$, see [AA17, Page 209].

Proof. The braiding of $\mathfrak{B}(\mathcal{D}_{jk,p}^{st})$ is given by

$$\begin{aligned} c(w_1 \otimes w_1) &= \omega^{[4k(2n+1)-2jN](t+1)+4k(2n+1)s} w_1 \otimes w_1, \\ c(w_1 \otimes w_2) &= \omega^{[4k(2n+1)+2jN](t+1)+4k(2n+1)s} w_2 \otimes w_1, \\ c(w_2 \otimes w_1) &= \omega^{[4k(2n+1)+2jN](t+1)+4k(2n+1)s} w_1 \otimes w_2, \\ c(w_2 \otimes w_2) &= \omega^{[4k(2n+1)-2jN](t+1)+4k(2n+1)s} w_2 \otimes w_2. \end{aligned}$$

□

Lemma 4.7. *Denote $q = \omega^{4ks(2n+1)}$, then $\mathfrak{B}(\mathcal{E}_{jk,p}^s)$ is finite dimensional iff one of the following conditions holds.*

- (1) $q = -1$, Cartan type $A_1 \times A_1$;
- (2) $q^3 = 1 \neq q$, Cartan type A_2 .

Proof. The braiding of $\mathfrak{B}(\mathcal{E}_{jk,p}^s)$ is given by

$$\begin{aligned} c(w_1 \otimes w_1) &= x_{11}^{2s} \cdot w_1 \otimes w_1 = \omega^{4k(2n+1)s} w_1 \otimes w_1 \\ c(w_1 \otimes w_2) &= x_{11}^{2s} \cdot w_2 \otimes w_1 = \omega^{4k(2n+1)s} w_2 \otimes w_1 \\ c(w_2 \otimes w_1) &= x_{11}^{2s} \cdot w_1 \otimes w_2 = \omega^{4k(2n+1)s} w_1 \otimes w_2 \\ c(w_2 \otimes w_2) &= x_{11}^{2s} \cdot w_2 \otimes w_2 = \omega^{4k(2n+1)s} w_2 \otimes w_2 \end{aligned}$$

□

Lemma 4.8. *Denote $q = \omega^{2k(2n+1)(2s+2t+1)}$, then*

$$\dim \mathfrak{B}(\mathcal{F}_{k,p}^{st}) = \begin{cases} 4, & q = -1, (\text{Cartan type } A_1 \times A_1), \\ 27, & q^3 = 1 \neq q, (\text{Cartan type } A_2), \\ \infty, & \text{otherwise.} \end{cases}$$

Proof. The braiding of $\mathfrak{B}(\mathcal{F}_{k,p}^{st})$ is given by

$$c(w_1 \otimes w_1) = x_{11}^{2s+1} \chi_{22}^{2t+1} \cdot w_1 \otimes w_1 = \omega^{2k(2n+1)(2s+2t+1)} w_1 \otimes w_1$$

$$\begin{aligned}
c(w_1 \otimes w_2) &= x_{11}^{2s+1} \chi_{22}^{2t+1} \cdot w_2 \otimes w_1 = \omega^{2k(2n+1)(2s+2t+1)} w_2 \otimes w_1 \\
c(w_2 \otimes w_1) &= x_{11}^{2s} \chi_{22}^{2t+2} \cdot w_1 \otimes w_2 = \omega^{2k(2n+1)(2s+2t+1)} w_1 \otimes w_2 \\
c(w_2 \otimes w_2) &= x_{11}^{2s} \chi_{22}^{2t+2} \cdot w_2 \otimes w_2 = \omega^{2k(2n+1)(2s+2t+1)} w_2 \otimes w_2.
\end{aligned}$$

□

4.2. Nichols algebra of type V_{abe} . Let $V_{abe} = \mathbb{k}v_1 \otimes \mathbb{k}v_2$ be a vector space with a braiding given by

$$\begin{aligned}
c(v_1 \otimes v_1) &= av_2 \otimes v_2, & c(v_1 \otimes v_2) &= bv_1 \otimes v_2, \\
c(v_2 \otimes v_1) &= bv_2 \otimes v_1, & c(v_2 \otimes v_2) &= ev_1 \otimes v_1,
\end{aligned}$$

then the braided vector space (V_{abe}, c) is of type V_{abe} . The braided vector space V_{abe} is isomorphic to $V_{ae b 1}$ via $v_1 \mapsto \sqrt{e}v_1, v_2 \mapsto v_2$. When $ae = b^2$, then V_{abe} is of diagonal type and

$$\dim \mathfrak{B}(V_{abe}) = \begin{cases} 4, & b = -1, (\mathfrak{B}(V_{abe}) \text{ is of Cartan type } A_1 \times A_1), \\ 27, & b^3 = 1 \neq b, (\mathfrak{B}(V_{abe}) \text{ is of Cartan type } A_2), \\ \infty, & \text{otherwise.} \end{cases}$$

When $ae \neq b^2$, according to [AG18, section 3.7] and [Shi20], we have

$$(4.1) \quad \dim \mathfrak{B}(V_{abe}) = \begin{cases} 4m, & b = -1, ae \in \mathbb{G}_m, \\ m^2, & ae = 1, b \in \mathbb{G}_m \text{ for } m \geq 2, \\ \infty, & b^2 = (ae)^{-1}, b \in \mathbb{G}_m \text{ for } m \geq 5, \\ \infty, & b \notin \mathbb{G}_m \text{ for } m \geq 2, \\ \text{unknown,} & \text{otherwise,} \end{cases}$$

where \mathbb{G}_m denotes the set of m -th primitive roots of unity.

Lemma 4.9. Denote $q = (-1)^p \lambda^{t+\frac{3}{2}} \tilde{\mu}^{2s+2t+1} \omega^{2k(2n+1)(2s+2t+1)}$, then

$$\dim \mathfrak{B}(\mathcal{G}_{k,p}^{st}) = \begin{cases} 4, & q = -1, (\text{Cartan type } A_1 \times A_1), \\ 27, & q^3 = 1 \neq q, (\text{Cartan type } A_2), \\ \infty, & \text{otherwise.} \end{cases}$$

Proof. $\mathfrak{B}(\mathcal{G}_{k,p}^{st})$ is of type V_{abe} with $a = b = e = (-1)^p \lambda^{t+\frac{3}{2}} \tilde{\mu}^{2s+2t+1} \omega^{2k(2n+1)(2s+2t+1)}$. □

Lemma 4.10. $\mathfrak{B}(\mathcal{H}_{jk,p}^{st})$ is of type V_{abe} with $ae = \lambda \tilde{\mu}^{2s+2t+1} \omega^{2k(2n+1)(4s+4t+2)+jN(4t+2)}$ and $b = (-1)^p \lambda^{t+\frac{1}{2}} \tilde{\mu}^{s+t+\frac{1}{2}} \omega^{2k(2n+1)(2s+2t+1)-jN(2t+1)}$.

Corollary 4.11. Suppose $ae \neq b^2 = (ae)^{-1}$ and $b \in \mathbb{G}_m$ for $m \geq 3$, then

$$\dim \mathfrak{B}(V_{abe}) = \infty.$$

Proof. If $b \in \mathbb{G}_4$, then $b^2 = (ae)^{-1} = -1 = ae$. The case for $m \geq 5$ was proved in [Shi20, Corollary 4.15]. We only need to deal with the case $m = 3$. Let $\mu = \lambda = N = 1$, then $ae = \omega^{2j(2t+1)}$, $b = (-1)^p \omega^{-j(2t+1)}$. And the parameter b covers all 3-th primitive roots of unity. For example, $b = \omega^{-12} \in \mathbb{G}_3$ in case $(n, j, t, p) = (4, 4, 1, 0)$ and $b = \omega^{12} \in \mathbb{G}_3$ in case $(n, j, t, p) = (4, 8, 1, 0)$. In this situation, $\dim \mathfrak{B}(\mathcal{H}_{jk,p}^{st}) = \infty$ since A_{1n}^{++} is isomorphic to a 2-cocycle deformation of \widehat{D}_{4n} [Mas00] and Nichols algebras associated with two dimensional Yetter-Drinfeld modules, over the dihedral group D_{4n} of order $4n$, are either 4-dimension or infinite dimension According to [AF07]. \square

Corollary 4.12. *Let $N = n = 1$, $\mu = 1$, then $\dim \mathfrak{B}(\mathcal{H}_{jk,p}^{st}) = \infty$.*

Proof. In this situation, $ae = \lambda\omega^4$, $b = (-1)^p \lambda^{\frac{1}{2}} \omega^{-2}$. So $ae \neq b^2 = (ae)^{-1}$, $b \in \mathbb{G}_6$ in case $\lambda = 1$ and $b \in \mathbb{G}_{12}$ in case $\lambda = -1$. \square

Remark 4.13. From observation, we have $\frac{ae}{b^2} = \omega^{4jN(2t+1)}$.

- (1) When $ae = 1$, then $b = \omega^{-4N}$ under the case $j = 2$, $t = 0$ and the suitable choice of p . So $\dim \mathfrak{B}(\mathcal{H}_{jk,p}^{st}) = (2n + 1)^2$ for suitable choice of (n, N, s, t, j, k, p) . Denote $\alpha = (n, N, s, t, j, k, p)$. For example, when $\mu = 1$, $\lambda = -1$, $ae = 1$, then

$$\dim \mathfrak{B}(\mathcal{H}_{jk,p}^{st}) = 2^2, \quad \alpha = (4, 6, 1, 1, 6, 3, 0), (4, 6, 2, 1, 6, 3, 1), \dots;$$

$$\dim \mathfrak{B}(\mathcal{H}_{jk,p}^{st}) = 3^2, \quad \alpha = (1, 6, 2, 0, 2, 5, 1), (1, 6, 3, 0, 2, 1, 0), \dots;$$

$$\dim \mathfrak{B}(\mathcal{H}_{jk,p}^{st}) = 5^2, \quad \alpha = (2, 10, 1, 0, 4, 7, 1), (2, 10, 3, 0, 2, 9, 1), \dots;$$

$$\dim \mathfrak{B}(\mathcal{H}_{jk,p}^{st}) = 6^2, \quad \alpha = (1, 6, 2, 0, 2, 5, 0), (1, 6, 5, 0, 2, 5, 1), \dots;$$

$$\dim \mathfrak{B}(\mathcal{H}_{jk,p}^{st}) = 7^2, \quad \alpha = (3, 14, 1, 0, 2, 11, 1), (3, 14, 1, 0, 4, 1, 1), \dots;$$

$$\dim \mathfrak{B}(\mathcal{H}_{jk,p}^{st}) = 9^2, \quad \alpha = (4, 18, 2, 3, 2, 11, 1), (4, 18, 2, 3, 8, 17, 1), \dots;$$

$$\dim \mathfrak{B}(\mathcal{H}_{jk,p}^{st}) = 10^2, \quad \alpha = (2, 10, 3, 0, 4, 3, 0), (2, 10, 7, 1, 2, 7, 1), \dots;$$

$$\dim \mathfrak{B}(\mathcal{H}_{jk,p}^{st}) = 11^2, \quad \alpha = (5, 22, 3, 0, 4, 1, 1), (5, 22, 3, 1, 8, 1, 1), \dots;$$

$$\dim \mathfrak{B}(\mathcal{H}_{jk,p}^{st}) = 13^2, \quad \alpha = (6, 26, 26, 5, 6, 7, 1), (6, 26, 26, 5, 10, 3, 1), \dots.$$

- (2) When $b = -1$, then $ae = \omega^{8N}$ in case $j = 2$ and $t = 0$. So $\dim \mathfrak{B}(\mathcal{H}_{jk,p}^{st}) = 4(2n+1)$ for suitable choice of (n, N, s, t, j, k, p) . Denote $\alpha = (n, N, s, t, j, k, p)$. For example, when $\mu = 1$, $\lambda = -1$, $b = -1$, then

$$\dim \mathfrak{B}(\mathcal{H}_{jk,p}^{st}) = 12, \quad \alpha = (1, 6, 2, 0, 2, 1, 0), (1, 18, 1, 0, 2, 11, 1), \dots;$$

$$\dim \mathfrak{B}(\mathcal{H}_{jk,p}^{st}) = 20, \quad \alpha = (2, 10, 1, 0, 2, 9, 0), (2, 20, 7, 1, 4, 2, 1), \dots;$$

$$\dim \mathfrak{B}(\mathcal{H}_{jk,p}^{st}) = 28, \quad \alpha = (3, 14, 3, 1, 6, 9, 1), (3, 28, 25, 1, 4, 6, 1), \dots;$$

$$\begin{aligned}
\dim \mathfrak{B} \left(\mathcal{H}_{jk,p}^{st} \right) &= 36, & \alpha &= (4, 18, 4, 2, 8, 1, 0), (4, 18, 12, 2, 2, 5, 1), \dots ; \\
\dim \mathfrak{B} \left(\mathcal{H}_{jk,p}^{st} \right) &= 44, & \alpha &= (5, 22, 1, 2, 6, 9, 1), (5, 22, 2, 0, 2, 7, 1), \dots ; \\
\dim \mathfrak{B} \left(\mathcal{H}_{jk,p}^{st} \right) &= 52, & \alpha &= (6, 26, 1, 2, 2, 7, 1), (6, 26, 3, 2, 6, 11, 1), \dots ; \\
\dim \mathfrak{B} \left(\mathcal{H}_{jk,p}^{st} \right) &= 60, & \alpha &= (7, 30, 5, 3, 2, 7, 0), (7, 30, 6, 0, 4, 13, 1), \dots .
\end{aligned}$$

Lemma 4.14. $\mathfrak{B} \left(\mathcal{J}_{jk,p}^{st} \right)$ is of type V_{abe} with $ae = \lambda \bar{\mu}^{2s+2t+1} \omega^{4k(2n+1)(2s+2t+1)-2jN(2t+1)}$ and $b = (-1)^p \lambda^{t-\frac{1}{2}} \bar{\mu}^{s+t+\frac{1}{2}} \omega^{2k(2n+1)(2s+2t+1)+jN(1+2t)}$.

Corollary 4.15. Let $N = n = 1$, $\mu = 1$, then $\mathfrak{B} \left(\mathcal{J}_{jk,p}^{st} \right) = \infty$.

Proof. In this situation, $ae = \lambda \omega^{-4}$, $b = (-1)^p \lambda^{-\frac{1}{2}} \omega^2$. So $b^2 = (ae)^{-1} \neq ae$, $b \in \mathbb{G}_6$ in case $\lambda = 1$ and $b \in \mathbb{G}_{12}$ in case $\lambda = -1$. It's proved by the Corollary 4.11. \square

Remark 4.16. Similar to the remark 4.13, $\mathfrak{B} \left(\mathcal{J}_{jk,p}^{st} \right) = \begin{cases} (2n+1)^2, & ae = 1, \\ 4(2n+1), & b = -1, \end{cases}$ for suitable choice of parameters. Denote $\alpha = (n, N, s, t, j, k, p)$.

(1) When $\mu = 1$, $\lambda = -1$, $ae = 1$, then

$$\begin{aligned}
\dim \mathfrak{B} \left(\mathcal{J}_{jk,p}^{st} \right) &= 2^2, & \alpha &= (4, 6, 6, 3, 6, 5, 1), (4, 10, 1, 1, 6, 5, 1), \dots ; \\
\dim \mathfrak{B} \left(\mathcal{J}_{jk,p}^{st} \right) &= 3^2, & \alpha &= (1, 6, 3, 0, 2, 5, 1), (1, 18, 18, 0, 2, 15, 0), \dots ; \\
\dim \mathfrak{B} \left(\mathcal{J}_{jk,p}^{st} \right) &= 5^2, & \alpha &= (2, 10, 8, 0, 2, 1, 1), (2, 10, 9, 1, 2, 1, 0), \dots ; \\
\dim \mathfrak{B} \left(\mathcal{J}_{jk,p}^{st} \right) &= 6^2, & \alpha &= (1, 12, 3, 0, 2, 10, 0), (1, 24, 23, 0, 2, 4, 0), \dots ; \\
\dim \mathfrak{B} \left(\mathcal{J}_{jk,p}^{st} \right) &= 7^2, & \alpha &= (3, 14, 6, 1, 2, 13, 0), (3, 14, 8, 1, 4, 1, 1), \dots ; \\
\dim \mathfrak{B} \left(\mathcal{J}_{jk,p}^{st} \right) &= 9^2, & \alpha &= (4, 18, 4, 2, 2, 7, 0), (4, 18, 5, 0, 8, 13, 1), \dots ; \\
\dim \mathfrak{B} \left(\mathcal{J}_{jk,p}^{st} \right) &= 10^2, & \alpha &= (2, 20, 11, 0, 2, 18, 1), (2, 20, 12, 1, 4, 2, 1), \dots ; \\
\dim \mathfrak{B} \left(\mathcal{J}_{jk,p}^{st} \right) &= 11^2, & \alpha &= (5, 22, 1, 1, 6, 19, 0), (5, 22, 1, 2, 6, 9, 1), \dots ; \\
\dim \mathfrak{B} \left(\mathcal{J}_{jk,p}^{st} \right) &= 13^2, & \alpha &= (6, 26, 1, 4, 4, 21, 1), (6, 26, 2, 1, 4, 11, 1), \dots .
\end{aligned}$$

(2) When $\mu = 1$, $\lambda = -1$, $b = -1$, then

$$\begin{aligned}
\dim \mathfrak{B} \left(\mathcal{J}_{jk,p}^{st} \right) &= 12, & \alpha &= (1, 6, 2, 0, 2, 5, 1), (1, 18, 7, 0, 2, 5, 1), \dots ; \\
\dim \mathfrak{B} \left(\mathcal{J}_{jk,p}^{st} \right) &= 20, & \alpha &= (2, 10, 2, 1, 4, 9, 1), (2, 10, 4, 0, 4, 9, 1), \dots ; \\
\dim \mathfrak{B} \left(\mathcal{J}_{jk,p}^{st} \right) &= 28, & \alpha &= (3, 14, 4, 1, 4, 11, 1), (3, 14, 5, 2, 6, 5, 0), \dots ; \\
\dim \mathfrak{B} \left(\mathcal{J}_{jk,p}^{st} \right) &= 36, & \alpha &= (4, 18, 5, 3, 2, 5, 1), (4, 18, 5, 3, 4, 1, 0), \dots ; \\
\dim \mathfrak{B} \left(\mathcal{J}_{jk,p}^{st} \right) &= 44, & \alpha &= (5, 22, 2, 0, 6, 1, 1), (5, 22, 2, 0, 8, 5, 0), \dots ;
\end{aligned}$$

$$\begin{aligned} \dim \mathfrak{B}(\mathcal{L}_{jk,p}^{st}) &= 52, & \alpha &= (6, 26, 1, 4, 8, 23, 1), (6, 26, 1, 4, 10, 19, 0), \dots; \\ \dim \mathfrak{B}(\mathcal{L}_{jk,p}^{st}) &= 60, & \alpha &= (7, 30, 30, 3, 2, 13, 1), (7, 30, 30, 3, 14, 1, 1), \dots. \end{aligned}$$

4.3. The Nichols algebras $\mathfrak{B}(\mathcal{L}_{k,pq}^s)$. The braiding of $\mathfrak{B}(\mathcal{L}_{k,pq}^s)$ can be described as follows.

(1) When $a + b \equiv 1 \pmod{2}$, let $b + (2a - 1) = d(2n + 1) + r$, $d \in \mathbb{N}$, $r \in \overline{0, 2n}$.
Then

$$c(m_a \otimes m_b) = \begin{cases} \omega^{4k(2n+1)(s-a)} m_{b+2a-1} \otimes m_a, & d = 0, \\ \omega^{4k(2n+1)(s-a)} m_{2n+1} \otimes m_a, & d = 1, r = 0 \\ (-1)^p \omega^{2k(2n+1)[2(s-a)+2(r-1)+1]} m_{2n+1-(r-1)} \otimes m_a, & d = 1, r \neq 0, \\ (-1)^p \omega^{2k(2n+1)[2(s-a)+4n+1]} m_1 \otimes m_a, & d = 2, r = 0, \\ \omega^{4k(2n+1)[(s-a)+2n+1]} m_r \otimes m_a, & d = 2, r \neq 0. \end{cases}$$

(2) When $a + b \equiv 0 \pmod{2}$, let $2a - 1 = b + 1 + d(2n + 1) + r$, $d \in \mathbb{N}$, $r \in \overline{0, 2n}$.
Then

$$c(m_a \otimes m_b) = \begin{cases} \omega^{4k(2n+1)(s-a+2a-1)} m_{b-(2a-1)} \otimes m_a, & 2a - 1 < b, \\ (-1)^p \omega^{2k(2n+1)[2(s-a)+2b-1]} m_1 \otimes m_a, & 2a - 1 = b, \\ (-1)^p \omega^{2k(2n+1)[2(s-a)+2b-1]} m_{r+1} \otimes m_a, & d = 0, \\ \omega^{4k(2n+1)[s-a+b+r]} m_{2n+1-r} \otimes m_a, & d = 1. \end{cases}$$

Lemma 4.17. *Let (V, c) be a braided vector space such that $c(x \otimes y) \in \mathbb{k}f_x(y) \otimes x$, where the map $f_x : V \rightarrow V$ is bijective for any $x \in V$ under a fixed basis. Then (V, \triangleright) is a rack with $x \triangleright y = f_x(y)$ under the fixed basis.*

Proof. For any $x, y, z \in V$,

$$\begin{aligned} c_1 c_2 c_1(x \otimes y \otimes z) &\in \mathbb{k}[(x \triangleright y) \triangleright (x \triangleright z)] \otimes (x \triangleright y) \otimes x, \\ c_2 c_1 c_2(x \otimes y \otimes z) &\in \mathbb{k}[x \triangleright (y \triangleright z)] \otimes (x \triangleright y) \otimes x. \end{aligned}$$

So $x \triangleright (y \triangleright z) = (x \triangleright y) \triangleright (x \triangleright z)$. □

Corollary 4.18. *The Nichols algebra $\mathfrak{B}(\mathcal{L}_{k,pq}^s)$ is of rack type.*

Lemma 4.19. *When $n = 1$, $p = 1$, $k = 0$, then $\dim \mathfrak{B}(\mathcal{L}_{k,pq}^s) = 12$. It is generated by m_1, m_2, m_3 and with relations*

$$(4.2) \quad m_3 m_2 = m_1 m_3 - m_2 m_1, \quad m_2 m_3 = -m_1 m_2 + m_3 m_1, \quad m_i^2 = 0 \quad \forall i \in \overline{1, 3}.$$

Proof. When $n = 1$, then the braiding of $\mathfrak{B}(\mathcal{L}_{k,pq}^s)$ is given by

$$\begin{aligned} c(m_1 \otimes m_1) &= (-1)^p \omega^{2k(2n+1)(2s-1)} m_1 \otimes m_1, \\ c(m_1 \otimes m_2) &= \omega^{4k(2n+1)(s-1)} m_3 \otimes m_1, \\ c(m_1 \otimes m_3) &= \omega^{4k(2n+1)s} m_2 \otimes m_1, \end{aligned}$$

$$\begin{aligned}
c(m_2 \otimes m_1) &= (-1)^p \omega^{2k(2n+1)[1+2(s-2)]} m_3 \otimes m_2, \\
c(m_2 \otimes m_2) &= (-1)^p \omega^{2k(2n+1)(2s-1)} m_2 \otimes m_2, \\
c(m_2 \otimes m_3) &= (-1)^p \omega^{2k(2n+1)(1+2s)} m_1 \otimes m_2, \\
c(m_3 \otimes m_1) &= \omega^{4k(2n+1)(s-1)} m_2 \otimes m_3, \\
c(m_3 \otimes m_2) &= \omega^{4k(2n+1)s} m_1 \otimes m_3, \\
c(m_3 \otimes m_3) &= (-1)^p \omega^{2k(2n+1)(2s-1)} m_3 \otimes m_3.
\end{aligned}$$

When $p = 1, k = 0$, it's easy to see that the relations (4.2) hold. The Nichols algebra is isomorphic to the 12-dimensional Nichols algebra over the dehdral group D_6 of order 6, see [AG99, Proposition 3.3.9]. \square

Corollary 4.20. *When $N = n = 1$, then $\dim \mathfrak{B}(\mathcal{L}_{k,pq}^s) = \begin{cases} 12, & \text{if } p = 1, \\ \infty, & \text{if } p = 0. \end{cases}$*

Lemma 4.21. *Suppose $n = 2, k = 0, p = 1$, then the Nichols algebra $\mathfrak{B}(\mathcal{L}_{k,pq}^s)$ has the following relations.*

$$\begin{aligned}
m_i^2 &= 0, \quad \forall i \in \overline{1,5}, \\
m_1 m_2 - m_3 m_1 - m_5 m_3 + m_4 m_5 - m_2 m_4 &= 0, \\
m_1 m_3 - m_2 m_1 + m_4 m_2 - m_5 m_4 + m_3 m_5 &= 0, \\
m_1 m_4 - m_5 m_1 + m_2 m_5 + m_3 m_2 + m_4 m_3 &= 0, \\
m_1 m_5 - m_4 m_1 - m_3 m_4 - m_2 m_3 - m_5 m_2 &= 0.
\end{aligned}$$

Proof. When $n = 2$, then the braiding of $\mathfrak{B}(\mathcal{L}_{k,pq}^s)$ is given by

$$\begin{aligned}
c(m_1 \otimes m_1) &= AB^{2s-1} m_1 \otimes m_1, & c(m_1 \otimes m_2) &= B^{2s-2} m_3 \otimes m_1, \\
c(m_1 \otimes m_3) &= B^{2s} m_2 \otimes m_1, & c(m_1 \otimes m_4) &= B^{2s-2} m_5 \otimes m_1, \\
c(m_1 \otimes m_5) &= B^{2s} m_4 \otimes m_1, & c(m_2 \otimes m_1) &= B^{2s-4} m_4 \otimes m_2, \\
c(m_2 \otimes m_2) &= AB^{2s-1} m_2 \otimes m_2, & c(m_2 \otimes m_3) &= AB^{2s-3} m_5 \otimes m_2, \\
c(m_2 \otimes m_4) &= B^{2s+2} m_1 \otimes m_2, & c(m_2 \otimes m_5) &= AB^{2s+1} m_3 \otimes m_2, \\
c(m_3 \otimes m_1) &= AB^{2s-5} m_5 \otimes m_3, & c(m_3 \otimes m_2) &= AB^{2s-3} m_4 \otimes m_3, \\
c(m_3 \otimes m_3) &= AB^{2s-1} m_3 \otimes m_3, & c(m_3 \otimes m_4) &= AB^{2s+1} m_2 \otimes m_3, \\
c(m_3 \otimes m_5) &= AB^{2s+3} m_1 \otimes m_3, & c(m_4 \otimes m_1) &= AB^{2s-3} m_3 \otimes m_4, \\
c(m_4 \otimes m_2) &= B^{2s-4} m_5 \otimes m_4, & c(m_4 \otimes m_3) &= AB^{2s+1} m_1 \otimes m_4, \\
c(m_4 \otimes m_4) &= AB^{2s-1} m_4 \otimes m_4, & c(m_4 \otimes m_5) &= B^{2s+2} m_2 \otimes m_4, \\
c(m_5 \otimes m_1) &= B^{2s-2} m_2 \otimes m_5, & c(m_5 \otimes m_2) &= B^{2s} m_1 \otimes m_5,
\end{aligned}$$

$$\begin{aligned} c(m_5 \otimes m_3) &= B^{2s-2} m_4 \otimes m_5, & c(m_5 \otimes m_4) &= B^{2s} m_3 \otimes m_5, \\ c(m_5 \otimes m_5) &= AB^{2s-1} m_5 \otimes m_5, \end{aligned}$$

where $A = (-1)^p$, $B = \omega^{2k(2n+1)}$. So $A = -1$, $B = 1$ in case $k = 0$, $p = 1$. The relations can be obtained by direct computations. \square

4.4. The Nichols algebra $\mathfrak{B}(\mathcal{N}_{k,pq}^s)$. Define $R^\gamma : \mathcal{N}_{k,pq}^s \otimes \mathcal{N}_{k,pq}^s \longrightarrow \mathcal{N}_{k,pq}^s \otimes \mathcal{N}_{k,pq}^s$ such that $R^\gamma(w_\alpha \otimes w_\beta) =$

$$\left\{ \begin{array}{ll} w_{\beta+\gamma} \otimes w_{2n-\alpha+2}, & \beta + \gamma \leq 2n + 1, \\ (-1)^p \lambda \bar{\mu}^{\frac{1}{2}} \omega^{2k(2n+1)} w_{2n+1} \otimes w_{2n-\alpha+2}, & \beta + \gamma = 2n + 2, \\ (-1)^p \lambda \left[\bar{\mu}^{\frac{1}{2}} \omega^{2k(2n+1)} \right]^{2(\gamma+\beta)-4n-3} w_{4n+3-\gamma-\beta} \otimes w_{2n-\alpha+2}, & \beta + \gamma \in \overline{2n+3, 4n+2}, \\ \lambda \left[\bar{\mu}^{\frac{1}{2}} \omega^{2k(2n+1)} \right]^{4n+2} w_{1+\beta+\gamma-(4n+3)} \otimes w_{2n-\alpha+2}, & \beta + \gamma \geq 4n + 3, \end{array} \right.$$

and $L^\gamma : \mathcal{N}_{k,pq}^s \otimes \mathcal{N}_{k,pq}^s \longrightarrow \mathcal{N}_{k,pq}^s \otimes \mathcal{N}_{k,pq}^s$ such that

$$L^\gamma(w_\alpha \otimes w_\beta) = \left\{ \begin{array}{ll} \left[\bar{\mu}^{\frac{1}{2}} \omega^{2k(2n+1)} \right]^{2\gamma} w_{\beta-\gamma} \otimes w_{2n-\alpha+2}, & \gamma < \beta, \\ (-1)^p \left[\bar{\mu}^{\frac{1}{2}} \omega^{2k(2n+1)} \right]^{2\beta-1} w_{\gamma-\beta+1} \otimes w_{2n-\alpha+2}, & \gamma \in \overline{\beta, \beta+2n}, \\ \lambda \left[\bar{\mu}^{\frac{1}{2}} \omega^{2k(2n+1)} \right]^{2\beta} w_{2n+1} \otimes w_{2n-\alpha+2}, & \gamma = \beta + 2n + 1, \\ \lambda \left[\bar{\mu}^{\frac{1}{2}} \omega^{2k(2n+1)} \right]^{2(\gamma-2n-1)} w_{4n+2+\beta-\gamma} \otimes w_{2n-\alpha+2}, & \gamma > \beta + 2n + 1. \end{array} \right.$$

Then the braiding of $\mathfrak{B}(\mathcal{N}_{k,pq}^s)$ can be described as follows.

(1) When $\alpha = n + 1$, then

$$c(w_\alpha \otimes w_\beta) = (-1)^q \left[\bar{\mu}^{\frac{1}{2}} \omega^{2k(2n+1)} \right]^{2(\alpha+s-1)} w_\beta \otimes w_{2n-\alpha+2};$$

(2) When $\alpha < n + 1$, $\alpha + \beta \equiv 0 \pmod{2}$, then

$$c(w_\alpha \otimes w_\beta) = (-1)^q \left[\bar{\mu}^{\frac{1}{2}} \omega^{2k(2n+1)} \right]^{2(2\alpha+s-n-2)} L^{2(n-\alpha+1)}(w_\alpha \otimes w_\beta);$$

(3) When $\alpha > n + 1$, $\alpha + \beta \equiv 1 \pmod{2}$, then

$$c(w_\alpha \otimes w_\beta) = (-1)^q \left[\bar{\mu}^{\frac{1}{2}} \omega^{2k(2n+1)} \right]^{2(s+n)} L^{2(\alpha-1-n)}(w_\alpha \otimes w_\beta);$$

(4) When $\alpha < n + 1$, $\alpha + \beta \equiv 1 \pmod{2}$, then

$$c(w_\alpha \otimes w_\beta) = (-1)^q \left[\bar{\mu}^{\frac{1}{2}} \omega^{2k(2n+1)} \right]^{2(2\alpha+s-n-2)} R^{2(n-\alpha+1)}(w_\alpha \otimes w_\beta);$$

(5) When $\alpha > n + 1$, $\alpha + \beta \equiv 0 \pmod{2}$, then

$$c(w_\alpha \otimes w_\beta) = (-1)^q \left[\bar{\mu}^{\frac{1}{2}} \omega^{2k(2n+1)} \right]^{2(s+n)} R^{2(\alpha-1-n)}(w_\alpha \otimes w_\beta).$$

Lemma 4.22. *When $n = 1 = \mu$, $k = 0$, $q = 1$, then $\dim \mathfrak{B}(\mathcal{N}_{k,pq}^s) = 12$. It is generated by w_1, w_2, w_3 and with relations:*

$$(4.3) \quad w_1^2 + (-1)^p w_2 w_3 + (-1)^p w_3 w_2 = 0, \quad w_2^2 = 0,$$

$$(4.4) \quad w_1 w_2 + \lambda(-1)^p w_3^2 + w_2 w_1 = 0, \quad w_1 w_3 = 0, \quad w_3 w_1 = 0.$$

Proof. When $n = 1$, the braiding of $\mathfrak{B}(\mathcal{N}_{k,pq}^s)$ is given by

$$\begin{aligned} c(w_1 \otimes w_1) &= (-1)^{p+q} \left[\bar{\mu}^{\frac{1}{2}} \omega^{2k(2n+1)} \right]^{2s-1} w_2 \otimes w_3, \\ c(w_1 \otimes w_2) &= \lambda(-1)^{p+q} \left[\bar{\mu}^{\frac{1}{2}} \omega^{2k(2n+1)} \right]^{2s-1} w_3 \otimes w_3, \\ c(w_1 \otimes w_3) &= (-1)^q \left[\bar{\mu}^{\frac{1}{2}} \omega^{2k(2n+1)} \right]^{2s+2} w_1 \otimes w_3, \\ c(w_2 \otimes w_1) &= (-1)^q \left[\bar{\mu}^{\frac{1}{2}} \omega^{2k(2n+1)} \right]^{2s+2} w_1 \otimes w_2, \\ c(w_2 \otimes w_2) &= (-1)^q \left[\bar{\mu}^{\frac{1}{2}} \omega^{2k(2n+1)} \right]^{2s+2} w_2 \otimes w_2, \\ c(w_2 \otimes w_3) &= (-1)^q \left[\bar{\mu}^{\frac{1}{2}} \omega^{2k(2n+1)} \right]^{2s+2} w_3 \otimes w_2, \\ c(w_3 \otimes w_1) &= (-1)^q \left[\bar{\mu}^{\frac{1}{2}} \omega^{2k(2n+1)} \right]^{2s+2} w_3 \otimes w_1, \\ c(w_3 \otimes w_2) &= (-1)^{p+q} \left[\bar{\mu}^{\frac{1}{2}} \omega^{2k(2n+1)} \right]^{2s+5} w_1 \otimes w_1, \\ c(w_3 \otimes w_3) &= \lambda(-1)^{p+q} \left[\bar{\mu}^{\frac{1}{2}} \omega^{2k(2n+1)} \right]^{2s+5} w_2 \otimes w_1. \end{aligned}$$

It's easy to see that the relations (4.3) and (4.4) hold in case $\mu = 1$, $k = 0$ and $q = 1$. According to direct computation with the relations, we have

$$\begin{aligned} w_1^3 &= (-1)^{p+1} w_1 w_2 w_3 = (-1)^{p+1} w_3 w_2 w_1 = \lambda w_3^3, \\ w_1^2 w_2 &= (-1)^{p+1} w_2 w_3 w_2 = w_2 w_1^2 = -w_1 w_2 w_1, \\ w_2 w_3^2 &= -w_3 w_2 w_3 = w_3^2 w_2 = \lambda(-1)^{p+1} w_2 w_1 w_2, \end{aligned}$$

and the other monomial in degree 3 vanish. This in turn implies

$$\begin{aligned} w_1^4 &= w_1^2 w_2 w_3 = w_3 w_2 w_1^2 = w_3^4 = w_3^2 w_2 w_1 = w_1 w_2 w_3^2 = w_2 w_1^2 w_2 \\ &= w_2 w_1 w_2 w_1 = w_1 w_2 w_1 w_2 = w_3 w_2 w_3 w_2 = w_2 w_3 w_2 w_3 = w_2 w_3^2 w_2 = 0, \\ w_2 w_1^3 &= (-1)^{p+1} w_2 w_1 w_2 w_3 = (-1)^{p+1} w_2 w_3 w_2 w_1 = \lambda w_2 w_3^3 \\ &= -\lambda w_3 w_2 w_3^2 = \lambda w_3^2 w_2 w_3 = -\lambda w_3^3 w_2 = (-1)^p w_3 w_2 w_1 w_2 = w_1^2 w_2 w_1 \\ &= -w_1 w_2 w_1^2 = -w_1^3 w_2 = (-1)^p w_1 w_2 w_3 w_2, \end{aligned}$$

and the other monomials in degree 4 vanish. Now it's easy to see that the Nichols algebra has a basis given by $\{1, w_1, w_2, w_3, w_1^3, w_1^2 w_2, w_2 w_3^2, w_2 w_1^3\}$. \square

Corollary 4.23. *When $N = n = 1 = \mu$, then $\dim \mathfrak{B}(\mathcal{N}_{k,pq}^s) = \begin{cases} 12, & q = 1, \\ \infty, & q = 0. \end{cases}$*

Lemma 4.24. *When $n = 2, q = 1 = \mu, k = 0$, then the Nichols algebra $\mathfrak{B}(\mathcal{N}_{k,pq}^s)$ has the following relations.*

$$\begin{aligned} w_1 w_5 &= 0, & w_2 w_4 &= 0, & w_3^2 &= 0, & w_4 w_2 &= 0, & w_5 w_1 &= 0, \\ (-1)^p w_1^2 + w_3 w_2 + w_2 w_3 + w_4 w_5 + w_5 w_4 &= 0, \\ w_1 w_2 + \lambda(-1)^p w_5^2 + w_2 w_1 + w_3 w_4 + w_4 w_3 &= 0, \\ w_1 w_3 + (-1)^p w_2 w_5 + \lambda w_4^2 + (-1)^p w_5 w_2 + w_3 w_1 &= 0, \\ w_1 w_4 + \lambda(-1)^p w_3 w_5 + \lambda(-1)^p w_5 w_3 + w_4 w_1 + w_2^2 &= 0. \end{aligned}$$

Proof. When $n = 2$, the braiding of $\mathfrak{B}(\mathcal{N}_{k,pq}^s)$ is given by

$$\begin{aligned} c(w_1 \otimes w_1) &= (-1)^{p+q} \left[\bar{\mu}^{\frac{1}{2}} \omega^{2k(2n+1)} \right]^{2s-3} w_4 \otimes w_5, \\ c(w_1 \otimes w_2) &= \lambda(-1)^{p+q} \left[\bar{\mu}^{\frac{1}{2}} \omega^{2k(2n+1)} \right]^{2s-3} w_5 \otimes w_5, \\ c(w_1 \otimes w_3) &= (-1)^{p+q} \left[\bar{\mu}^{\frac{1}{2}} \omega^{2k(2n+1)} \right]^{2s+1} w_2 \otimes w_5, \\ c(w_1 \otimes w_4) &= \lambda(-1)^{p+q} \left[\bar{\mu}^{\frac{1}{2}} \omega^{2k(2n+1)} \right]^{2s+1} w_3 \otimes w_5, \\ c(w_1 \otimes w_5) &= (-1)^q \left[\bar{\mu}^{\frac{1}{2}} \omega^{2k(2n+1)} \right]^{2s+4} w_1 \otimes w_5, \\ c(w_2 \otimes w_1) &= (-1)^q \left[\bar{\mu}^{\frac{1}{2}} \omega^{2k(2n+1)} \right]^{2s} w_3 \otimes w_4, \\ c(w_2 \otimes w_2) &= (-1)^{p+q} \left[\bar{\mu}^{\frac{1}{2}} \omega^{2k(2n+1)} \right]^{2s+3} w_1 \otimes w_4, \\ c(w_2 \otimes w_3) &= (-1)^q \left[\bar{\mu}^{\frac{1}{2}} \omega^{2k(2n+1)} \right]^{2s} w_5 \otimes w_4, \\ c(w_2 \otimes w_4) &= (-1)^q \left[\bar{\mu}^{\frac{1}{2}} \omega^{2k(2n+1)} \right]^{2s+4} w_2 \otimes w_4, \\ c(w_2 \otimes w_5) &= \lambda(-1)^{p+q} \left[\bar{\mu}^{\frac{1}{2}} \omega^{2k(2n+1)} \right]^{2s+3} w_4 \otimes w_4, \\ c(w_3 \otimes w_1) &= (-1)^q \left[\bar{\mu}^{\frac{1}{2}} \omega^{2k(2n+1)} \right]^{2s+4} w_1 \otimes w_3, \end{aligned}$$

$$\begin{aligned}
c(w_3 \otimes w_2) &= (-1)^q \left[\bar{\mu}^{\frac{1}{2}} \omega^{2k(2n+1)} \right]^{2s+4} w_2 \otimes w_3, \\
c(w_3 \otimes w_3) &= (-1)^q \left[\bar{\mu}^{\frac{1}{2}} \omega^{2k(2n+1)} \right]^{2s+4} w_3 \otimes w_3, \\
c(w_3 \otimes w_4) &= (-1)^q \left[\bar{\mu}^{\frac{1}{2}} \omega^{2k(2n+1)} \right]^{2s+4} w_4 \otimes w_3, \\
c(w_3 \otimes w_5) &= (-1)^q \left[\bar{\mu}^{\frac{1}{2}} \omega^{2k(2n+1)} \right]^{2s+4} w_5 \otimes w_3, \\
c(w_4 \otimes w_1) &= (-1)^{p+q} \left[\bar{\mu}^{\frac{1}{2}} \omega^{2k(2n+1)} \right]^{2s+5} w_2 \otimes w_2, \\
c(w_4 \otimes w_2) &= (-1)^q \left[\bar{\mu}^{\frac{1}{2}} \omega^{2k(2n+1)} \right]^{2s+4} w_4 \otimes w_2, \\
c(w_4 \otimes w_3) &= (-1)^q \left[\bar{\mu}^{\frac{1}{2}} \omega^{2k(2n+1)} \right]^{2s+8} w_1 \otimes w_2, \\
c(w_4 \otimes w_4) &= \lambda(-1)^{p+q} \left[\bar{\mu}^{\frac{1}{2}} \omega^{2k(2n+1)} \right]^{2s+5} w_5 \otimes w_2, \\
c(w_4 \otimes w_5) &= (-1)^q \left[\bar{\mu}^{\frac{1}{2}} \omega^{2k(2n+1)} \right]^{2s+8} w_3 \otimes w_2, \\
c(w_5 \otimes w_1) &= (-1)^q \left[\bar{\mu}^{\frac{1}{2}} \omega^{2k(2n+1)} \right]^{2s+4} w_5 \otimes w_1, \\
c(w_5 \otimes w_2) &= (-1)^{p+q} \left[\bar{\mu}^{\frac{1}{2}} \omega^{2k(2n+1)} \right]^{2s+7} w_3 \otimes w_1, \\
c(w_5 \otimes w_3) &= \lambda(-1)^{p+q} \left[\bar{\mu}^{\frac{1}{2}} \omega^{2k(2n+1)} \right]^{2s+7} w_4 \otimes w_1, \\
c(w_5 \otimes w_4) &= (-1)^{p+q} \left[\bar{\mu}^{\frac{1}{2}} \omega^{2k(2n+1)} \right]^{2s+11} w_1 \otimes w_1, \\
c(w_5 \otimes w_5) &= \lambda(-1)^{p+q} \left[\bar{\mu}^{\frac{1}{2}} \omega^{2k(2n+1)} \right]^{2s+11} w_2 \otimes w_1.
\end{aligned}$$

The relations can be checked via direct computations. □

4.5. Finite dimensional Nichols algebras over $A_{13}^{+\lambda}$.

Theorem 4.25. *Finite dimensional Nichols algebras over $A_{13}^{+\lambda}$, associated with simple Yetter-Drinfeld modules, can be classified as follows.*

- (1) $\dim \mathfrak{B}(\mathcal{B}_{k,p}^s) = \begin{cases} 2, & \text{if } \lambda = 1, s = 1, k = 0, p = 1, \\ 4, & \text{if } \lambda = -1, s = 1, k = 0, p \in \mathbb{Z}_2. \end{cases}$
- (2) $\dim \mathfrak{B}(\mathcal{F}_{k,p}^{st}) = 4$ in case $s = 1, t = 0, k = 1, p \in \mathbb{Z}_2$.
- (3) $\dim \mathfrak{B}(\mathcal{G}_{k,p}^{st}) = 4$ in case $\lambda = 1 = s, t = 0, (p, k) = (0, 1)$ or $(1, 0)$.
- (4) $\dim \mathfrak{B}(\mathcal{L}_{k,pq}^s) = 12$ in case $s = p = 1, k = 0, q \in \mathbb{Z}_2$.

(5) $\dim \mathfrak{B}(\mathcal{N}_{k,pq}^s) = 12$ in case $s = q = 1, k = 0, p \in \mathbb{Z}_2$.

Remark 4.26. The proof is just a summation of the section. We will deal with the classification of finite dimensional Hopf algebras over $A_{13}^{+\lambda}$ in a sequel[Shi].

5. APPENDIX

5.1. One dimensional Yetter-Drinfeld modules over $A_{N2n+1}^{\mu\lambda}$.

- (1) $\mathcal{A}_{k,p}^s = \mathbb{k}w$, where $w = v \boxtimes [x_{11}^{2s} + (-1)^p x_{12}^{2s}]$, $s \in \overline{1, N}$, $V_k = \mathbb{k}v$, $p \in \mathbb{Z}_2$.
(2) $\mathcal{B}_{k,p}^s = \mathbb{k}w$, where $w = v \boxtimes [x_{11}^{2s+1} \chi_{22}^{2n} + (-1)^p \sqrt{\lambda} x_{12}^{2s+1} \chi_{21}^{2n}]$, $V'_k = \mathbb{k}v$, $s \in \overline{1, N}$, $p \in \mathbb{Z}_2$.

5.2. Two dimensional Yetter-Drinfeld modules over $A_{N2n+1}^{\mu\lambda}$.

- (1) $\mathcal{C}_{jk,p}^{st} = \mathbb{k}w_1 \oplus \mathbb{k}w_2$, where

$$\begin{aligned} w_1 &= v_1 \boxtimes x_{11}^{2s} \chi_{22}^{2t+2} + (-1)^p \omega^{-2k(2n+1)+jN} v_2 \boxtimes x_{12}^{2s} \chi_{21}^{2t+2} \\ w_2 &= v_2 \boxtimes x_{11}^{2s+1} \chi_{22}^{2t+1} + (-1)^p \omega^{2k(2n+1)-jN} v_1 \boxtimes x_{12}^{2s+1} \chi_{21}^{2t+1}, \end{aligned}$$

$$s \in \overline{1, N}, p \in \mathbb{Z}_2, t \in \overline{0, n-1}, V_{jk} = \mathbb{k}v_1 \oplus \mathbb{k}v_2.$$

- (2) $\mathcal{D}_{jk,p}^{st} = \mathbb{k}w_1 \oplus \mathbb{k}w_2$, where

$$\begin{aligned} w_1 &= v_1 \boxtimes x_{11}^{2s+1} \chi_{22}^{2t+1} + (-1)^p \omega^{jN-2k(2n+1)} v_2 \boxtimes x_{12}^{2s+1} \chi_{21}^{2t+1}, \\ w_2 &= v_2 \boxtimes x_{11}^{2s} \chi_{22}^{2t+2} + (-1)^p \omega^{2k(2n+1)-jN} v_1 \boxtimes x_{12}^{2s} \chi_{21}^{2t+2}, \end{aligned}$$

$$s \in \overline{1, N}, p \in \mathbb{Z}_2, t \in \overline{0, n-1}, V_{jk} = \mathbb{k}v_1 \oplus \mathbb{k}v_2.$$

- (3) $\mathcal{E}_{jk,p}^s = \mathbb{k}w_1 \oplus \mathbb{k}w_2$, where

$$\begin{aligned} w_1 &= v_1 \boxtimes x_{11}^{2s} + (-1)^p \omega^{jN-2k(2n+1)} v_2 \boxtimes x_{12}^{2s}, \\ w_2 &= v_2 \boxtimes x_{11}^{2s} + (-1)^p \omega^{2k(2n+1)-jN} v_1 \boxtimes x_{12}^{2s}, \end{aligned}$$

$$s \in \overline{1, N}, p \in \mathbb{Z}_2, V_{jk} = \mathbb{k}v_1 \oplus \mathbb{k}v_2.$$

- (4) $\mathcal{F}_{k,p}^{st} = \mathbb{k}w_1 \oplus \mathbb{k}w_2$, where

$$\begin{aligned} w_1 &= v \boxtimes [x_{11}^{2s+1} \chi_{22}^{2t+1} + (-1)^p x_{12}^{2s+1} \chi_{21}^{2t+1}], \\ w_2 &= v \boxtimes [x_{11}^{2s} \chi_{22}^{2t+2} + (-1)^p x_{12}^{2s} \chi_{21}^{2t+2}], \end{aligned}$$

$$s \in \overline{1, N}, t \in \overline{0, n-1}, V_k = \mathbb{k}v, p \in \mathbb{Z}_2.$$

- (5) $\mathcal{G}_{k,p}^{st} = \mathbb{k}w_1 \oplus \mathbb{k}w_2$, where

$$\begin{aligned} w_1 &= v \boxtimes [x_{11}^{2s} \chi_{22}^{2t+1} + (-1)^p \sqrt{\lambda} x_{12}^{2s} \chi_{21}^{2t+1}], \\ w_2 &= v \boxtimes [x_{11}^{2s+1} \chi_{22}^{2t} + \frac{(-1)^p}{\sqrt{\lambda}} x_{12}^{2s+1} \chi_{21}^{2t}], \end{aligned}$$

$s \in \overline{1, N}, t \in \overline{0, n-1}, p \in \mathbb{Z}_2, V'_k = \mathbb{k}v.$

(6) $\mathcal{H}_{jk,p}^{st} = \mathbb{k}w_1 \oplus \mathbb{k}w_2$, where

$$w_1 = v'_1 \boxtimes x_{11}^{2s} \chi_{22}^{2t+1} + \frac{(-1)^p}{\sqrt{\lambda\bar{\mu}}} \omega^{jN-2k(2n+1)} v'_2 \boxtimes x_{12}^{2s} \chi_{21}^{2t+1},$$

$$w_2 = v'_2 \boxtimes x_{11}^{2s+1} \chi_{22}^{2t} + (-1)^p \sqrt{\lambda\bar{\mu}} \omega^{2k(2n+1)-jN} v'_1 \boxtimes x_{12}^{2s+1} \chi_{21}^{2t},$$

$V'_{jk} = \mathbb{k}v'_1 \oplus \mathbb{k}v'_2, p \in \mathbb{Z}_2, s \in \overline{1, N}, t \in \overline{0, n-1}.$

(7) $\mathcal{S}_{jk,p}^{st} = \mathbb{k}w_1 \oplus \mathbb{k}w_2$, where

$$w_1 = v'_1 \boxtimes x_{11}^{2s+1} \chi_{22}^{2t} + \frac{(-1)^p}{\sqrt{\lambda\bar{\mu}}} \omega^{jN-2k(2n+1)} v'_2 \boxtimes x_{12}^{2s+1} \chi_{21}^{2t},$$

$$w_2 = v'_2 \boxtimes x_{11}^{2s} \chi_{22}^{2t+1} + (-1)^p \sqrt{\lambda\bar{\mu}} \omega^{2k(2n+1)-jN} v'_1 \boxtimes x_{12}^{2s} \chi_{21}^{2t+1},$$

$V'_{jk} = \mathbb{k}v'_1 \oplus \mathbb{k}v'_2, p \in \mathbb{Z}_2, s \in \overline{1, N}, t \in \overline{0, n-1}.$

5.3. $2n+1$ dimensional Yetter-Drinfeld modules over $A_{N2n+1}^{\mu\lambda}$.

(1) Let $s \in \overline{1, N}, p, q \in \mathbb{Z}_2, V_{jk} = \mathbb{k}v_1 \oplus \mathbb{k}v_2$, and denote

$$a = (-1)^p \omega^{2jN-2k(2n+1)}, \quad b = (-1)^q \sqrt{\lambda} \omega^{-4kn(2n+1)},$$

$$m_1 = (v_1 + av_2) \boxtimes x_{11}^{2s-1} \chi_{22}^{2t} + b \chi_{22}^{2n} \cdot \left[(v_1 + av_2) \boxtimes x_{21}^{2s-1} \right],$$

$$m_i = \begin{cases} x_{22} \cdot m_{i-1} = \chi_{22}^{i-1} \cdot m_1, & i \text{ is even, } 1 < i \leq 2n+1, \\ x_{11} \cdot m_{i-1} = \chi_{11}^{i-1} \cdot m_1, & i \text{ is odd, } 1 < i \leq 2n+1. \end{cases}$$

Then $\mathcal{L}_{k,pq}^s = \bigoplus_{i=1}^{2n+1} \mathbb{k}m_i$ is a $2n+1$ dimensional Yetter-Drinfeld module over $A_{N2n+1}^{\mu\lambda}$ with the module structure given by

$$x_{11} \cdot m_i = \begin{cases} \chi_{11}^i \cdot w_1 = m_{i+1}, & i \text{ is even, } 1 < i \leq 2n+1, \\ \chi_{22}^{i-2} \chi_{11}^2 \cdot m_1 = \omega^{4k(2n+1)} m_{i-1}, & i \text{ is odd, } 1 < i \leq 2n+1, \\ (-1)^p \omega^{2k(2n+1)} m_1, & i = 1, \end{cases}$$

$$x_{22} \cdot m_i = \begin{cases} \chi_{11}^{i-2} \cdot m_1 = \omega^{4k(2n+1)} m_{i-1}, & i \text{ is even, } 1 < i < 2n+1, \\ \chi_{22}^i \cdot m_1 = m_{i+1}, & i \text{ is odd, } 1 \leq i < 2n+1, \\ \chi_{22}^{2n+1} \cdot m_1 = (-1)^p \omega^{2k(2n+1)} m_{2n+1}, & i = 2n+1, \end{cases}$$

$$x_{pq} \cdot m_i = 0, \quad pq = 12 \text{ or } 21, 1 \leq i \leq 2n+1.$$

And the comodule structure is given by

$$\rho(m_i) = \begin{cases} x_{11}^{2(s-i)} \chi_{22}^{2i-1} \otimes m_i + (-1)^q \lambda \sqrt{\lambda} \omega^{4k(2n+1)(i-1-n)} x_{12}^{2(s-i)} \chi_{21}^{2i-1} \otimes m_{2n+2-i}, & i \text{ even,} \\ x_{11}^{2(s-i)} \chi_{11}^{2i-1} \otimes m_i + (-1)^q \lambda \sqrt{\lambda} \omega^{4k(2n+1)(i-1-n)} x_{12}^{2(s-i)} \chi_{12}^{2i-1} \otimes m_{2n+2-i}, & i \text{ odd.} \end{cases}$$

- (2) Let $V'_{jk} = \mathbb{k}v'_1 \oplus \mathbb{k}v'_2$, $a = \frac{(-1)^p}{\sqrt{\mu}} \omega^{-2k(2n+1)}$, $p \in \mathbb{Z}_2$, $q \in \mathbb{Z}_2$, $s \in \overline{1, N}$, $b = (-1)^q (\tilde{\mu} \omega^{4k(2n+1)})^{-n}$, and denote

$$w_1 = (v'_1 + av'_2) \boxtimes x_{11}^{2s} \chi_{22}^{2n} + b \chi_{21}^{2n} \cdot [(v'_1 + av'_2) \boxtimes x_{12}^{2s+1} \chi_{21}^{2n-1}],$$

$$w_i = \begin{cases} x_{21} \cdot w_{i-1} = \chi_{21}^{i-1} \cdot w_1, & i \text{ is even, } 1 < i \leq 2n+1, \\ x_{12} \cdot w_{i-1} = \chi_{12}^{i-1} \cdot w_1, & i \text{ is odd, } 1 < i \leq 2n+1. \end{cases}$$

Then $\mathcal{N}_{k,pq}^s = \bigoplus_{i=1}^{2n+1} \mathbb{k}w_i$ is a $2n+1$ dimensional Yetter-Drinfeld module over $A_{N^{2n+1}}^{\mu\lambda}$ with the module structure given by

$$x_{12} \cdot w_i = \begin{cases} \frac{1}{a} w_1, & i = 1, \\ w_{i+1}, & i \text{ even} \\ \frac{1}{a^2} w_{i-1}, & i \text{ odd and } i > 1 \end{cases}$$

$$x_{21} \cdot w_i = \begin{cases} \frac{1}{a} w_{2n+1}, & i = 2n+1, \\ \frac{1}{a^2} w_{i-1}, & i \text{ even} \\ w_{i+1}, & i \text{ odd and } i < 2n+1, \end{cases}$$

$$x_{\alpha\beta} \cdot w_i = 0, \quad \alpha\beta = 11 \text{ or } 22,$$

and the comodule structure given by

$$\rho(w_i) = \begin{cases} x_{11}^{2s} \chi_{22}^{i-1} \chi_{22}^{2n-i+1} \otimes w_i + \frac{b}{a^{2(i-1)}} x_{12}^{2s} \chi_{21}^{i-1} \chi_{21}^{2n-i+1} \otimes w_{2n-i+2}, & i \text{ even} \\ x_{11}^{2s} \chi_{22}^{2n-i+1} \chi_{11}^{i-1} \otimes w_i + \frac{b}{a^{2(i-1)}} x_{12}^{2s} \chi_{21}^{2n-i+1} \chi_{12}^{i-1} \otimes w_{2n-i+2}, & i \text{ odd.} \end{cases}$$

- (3) $\mathcal{H}_{k,p}^s = \bigoplus_{i=1}^{2n+1} \mathbb{k}w_i$, where

$$w_1 = v \boxtimes x_{11}^{2s-1} + (-1)^p \sqrt{\lambda} \omega^{-4kn(2n+1)} \chi_{22}^{2n} \cdot [v \boxtimes x_{12}^{2s-2} \chi_{21}],$$

$$w_i = \begin{cases} x_{22} \cdot w_{i-1} = \chi_{22}^{i-1} \cdot w_1, & i \text{ is even, } 1 < i \leq 2n+1, \\ x_{11} \cdot w_{i-1} = \chi_{11}^{i-1} \cdot w_1, & i \text{ is odd, } 1 < i \leq 2n+1, \end{cases}$$

$s \in \overline{1, N}$, $p \in \mathbb{Z}_2$, $V_k = \mathbb{k}v$.

- (4) $\mathcal{M}_{k,p}^s = \bigoplus_{i=1}^{2n+1} \mathbb{k}m_i$, where

$$m_1 = v \boxtimes x_{11}^{2s} \chi_{22}^{2n} + a \chi_{21}^{2n} \cdot [v \boxtimes x_{12}^{2s+1} \chi_{21}^{2n-1}],$$

$$m_i = \begin{cases} x_{21} \cdot m_{i-1} = \chi_{21}^{i-1} \cdot m_1, & i \text{ is even, } 1 < i \leq 2n+1, \\ x_{12} \cdot m_{i-1} = \chi_{12}^{i-1} \cdot m_1, & i \text{ is odd, } 1 < i \leq 2n+1, \end{cases}$$

$V'_k = \mathbb{k}v$, $a = (-1)^p \omega^{-4kn(2n+1)} \tilde{\mu}^{-2n}$, $p \in \mathbb{Z}_2$, $s \in \overline{1, N}$.

REFERENCES

- [AA17] N. Andruskiewitsch and I. Angiono, *On finite dimensional Nichols algebras of diagonal type*, Bull. Math. Sci. **7** (2017), no. 3, 353–573. MR 3736568

- [AF07] N. Andruskiewitsch and F. Fantino, *On pointed Hopf algebras associated with alternating and dihedral groups*, Rev. Un. Mat. Argentina **48** (2007), no. 3, 57–71 (2008). MR 2389459
- [AG99] N. Andruskiewitsch and M. Graña, *Braided Hopf algebras over non-abelian finite groups*, Bol. Acad. Nac. Cienc. (Córdoba) **63** (1999), 45–78, Colloquium on Operator Algebras and Quantum Groups (Spanish) (Vaquerías, 1997). MR 1714540 (2001b:16039)
- [AG18] N. Andruskiewitsch and J. Giraldi, *Nichols algebras that are quantum planes*, Linear and Multilinear Algebra **66** (2018), no. 5, 961–991. MR 3775317
- [And17] N. Andruskiewitsch, *An introduction to Nichols algebras*, Quantization, geometry and noncommutative structures in mathematics and physics, Math. Phys. Stud., Springer, Cham, 2017, pp. 135–195. MR 3751453
- [AS98] N. Andruskiewitsch and H.-J. Schneider, *Lifting of quantum linear spaces and pointed Hopf algebras of order p^3* , J. Algebra **209** (1998), no. 2, 658–691. MR 1659895
- [AS02] N. Andruskiewitsch and H.-J. Schneider, *Pointed Hopf algebras*, New directions in Hopf algebras, Math. Sci. Res. Inst. Publ., vol. 43, Cambridge Univ. Press, Cambridge, 2002, pp. 1–68. MR 1913436
- [CDMM04] C. Călinescu, S. Dăscălescu, A. Masuoka, and C. Menini, *Quantum lines over non-cocommutative cosemisimple Hopf algebras*, J. Algebra **273** (2004), no. 2, 753–779. MR 2037722
- [FGM19] F. Fantino, G. García, and M. Mastnak, *On finite-dimensional copointed Hopf algebras over dihedral groups*, J. Pure Appl. Algebra **223** (2019), no. 8, 3611–3634. MR 3926230
- [Hec09] I. Heckenberger, *Classification of arithmetic root systems*, Adv. Math. **220** (2009), no. 1, 59–124. MR 2462836
- [Mas00] A. Masuoka, *Cocycle deformations and Galois objects for some cosemisimple Hopf algebras of finite dimension*, New trends in Hopf algebra theory (La Falda, 1999), Contemp. Math., vol. 267, Amer. Math. Soc., Providence, RI, 2000, pp. 195–214. MR 1800713
- [Rad03] D. E. Radford, *On oriented quantum algebras derived from representations of the quantum double of a finite-dimensional Hopf algebra*, J. Algebra **270** (2003), no. 2, 670–695. MR 2019635
- [Shi] Y.-X. Shi, *Finite dimensional Hopf algebras over Suzuki algebra $A_{13}^{+\lambda}$* , preparing.
- [Shi19] ———, *Finite-dimensional Hopf algebras over the Kac-Paljutkin algebra H_8* , Rev. Un. Mat. Argentina **60** (2019), no. 1, 265–298. MR 3981570
- [Shi20] ———, *Finite dimensional Nichols algebras over Suzuki algebra I: over simple Yetter-Drinfeld modules $A_{N_{2n}}^{\mu,\lambda}$* , arXiv:2011.14274 (2020).
- [Suz98] S. Suzuki, *A family of braided cosemisimple Hopf algebras of finite dimension*, Tsukuba J. Math. **22** (1998), no. 1, 1–29. MR 1637640
- [Wak03] M. Wakui, *The coribbon structures of some finite dimensional braided Hopf algebras generated by 2×2 -matrix coalgebras*, Noncommutative geometry and quantum groups (Warsaw, 2001), Banach Center Publ., vol. 61, Polish Acad. Sci. Inst. Math., Warsaw, 2003, pp. 333–344. MR 2024437
- [Wak10] ———, *Polynomial invariants for a semisimple and cosemisimple Hopf algebra of finite dimension*, J. Pure Appl. Algebra **214** (2010), no. 6, 701–728. MR 2580652
- [Wak19] ———, *Braided Morita equivalence for finite-dimensional semisimple and cosemisimple Hopf algebras*, Proceedings of the Meeting for Study of Number Theory,

Hopf Algebras and Related Topics, Yokohama Publ., Yokohama, 2019, pp. 157–183.
MR 3932380

SCHOOL OF MATHEMATICS AND STATISTICS, JIANGXI NORMAL UNIVERSITY, NANCHANG 330022, P. R.
CHINA
Email address: `yxshi@jxnu.edu.cn`