

# THE NICHOLS ALGEBRA $\mathfrak{B}(V_{abe})$ AND A CLASS OF COMBINATORIAL NUMBERS

YUXING SHI

**ABSTRACT.** We investigate the Nichols algebra  $\mathfrak{B}(V_{abe})$  which are from the Yetter-Drinfeld category of Suzuki algebras. The  $4n$  and  $n^2$  dimensional Nichols algebras, first appeared in [1], are obtained again via a different method. And the connection between the Nichols algebra  $\mathfrak{B}(V_{abe})$  and a class of combinatorial numbers on the subgroups of symmetric groups is established.

## 1. INTRODUCTION

Nichols algebras appeared for the first time in W. Nichols' paper[12], which aimed to construct new examples of Hopf algebras. They also arose independently in Woronowicz, Lusztig, and Rosso's works [18][11][13], via the contexts of non-commutative differential calculus, invariant bilinear form and quantum shuffles respectively.

Nichols algebras over group algebras are of group type. Determining all finite dimensional group type Nichols algebras is a crucial important step for the classification of pointed Hopf algebras[3]. Heckenberger classified all finite dimensional Nichols algebras of diagonal type(Abelian group type)[9], via the Weyl groupoid introduced in [8] and arithmetic root systems. Angiono determined the defining relations of the finite dimensional Nichols algebras of diagonal type in [5][4]. In general, it's difficult to decide whether a given Nichols algebra of non-diagonal type is finite dimensional. Andruskiewitsch and Graña established a bridge between group type Nichols algebras and racks [2], so the study on Nichols algebras of non-Abelian group type(also called rack type) can be carried out in the framework of racks.

When we dealt with Nichols algebras over the Suzuki algebra  $A_{Nr}^{\mu,\lambda}$ [15][16][17], we came across the Nichols algebra  $\mathfrak{B}(V_{abe})$  of non-group type. After we almost had finished the work, we found that our work is overlapped with Andruskiewitsch and Giraldi's work [1, section 3.7]. Since our approach is different and the problem of determining the dimension of  $\mathfrak{B}(V_{abe})$  is not completely solved, our work is still

---

2010 *Mathematics Subject Classification.* 16T05, 16T25, 17B22.

*Keywords:* Nichols algebra; Hopf algebra; Suzuki algebra.

This work was partially supported by Foundation of Jiangxi Educational Committee (No.12020447) and Natural Science Foundation of Jiangxi Normal University(No. 12018937) .

valuable. Besides, we establish a connection between the Nichols algebra  $\mathfrak{B}(V_{abe})$  and a class of combinatorial numbers on the subgroups of the symmetric groups.

## 2. THE NICHOLS ALGEBRA $\mathfrak{B}(V_{abe})$

### 2.1. Nichols algebra.

**Definition 2.1.** [3, Definition 2.1] Let  $H$  be a Hopf algebra and  $V \in {}^H_H\mathcal{YD}$ . A braided  $\mathbb{N}$ -graded Hopf algebra  $R = \bigoplus_{n \geq 0} R(n) \in {}^H_H\mathcal{YD}$  is called the *Nichols algebra* of  $V$  if

- (i)  $\mathbb{k} \simeq R(0)$ ,  $V \simeq R(1) \in {}^H_H\mathcal{YD}$ ,
- (ii)  $R(1) = \mathcal{P}(R) = \{r \in R \mid \Delta_R(r) = r \otimes 1 + 1 \otimes r\}$ .
- (iii)  $R$  is generated as an algebra by  $R(1)$ .

In this case,  $R$  is denoted by  $\mathfrak{B}(V) = \bigoplus_{n \geq 0} \mathfrak{B}^n(V)$ .

*Remark 2.2.* The Nichols algebra  $\mathfrak{B}(V)$  is completely determined by the braiding. More precisely, as proved in [14] and noted in [3],

$$\mathfrak{B}(V) = \mathbb{K} \oplus V \oplus \bigoplus_{n=2}^{\infty} V^{\otimes n} / \ker \mathfrak{S}_n = T(V) / \ker \mathfrak{S},$$

where  $\mathfrak{S}_{n,1} \in \text{End}_{\mathbb{k}}(V^{\otimes(n+1)})$ ,  $\mathfrak{S}_n \in \text{End}_{\mathbb{k}}(V^{\otimes n})$ ,

$$\mathfrak{S}_{n,1} := \text{id} + c_n + c_{n-1}c_n + \cdots + c_1 \cdots c_{n-1}c_n = \text{id} + \mathfrak{S}_{n-1,1}c_n,$$

$$\mathfrak{S}_1 := \text{id}, \quad \mathfrak{S}_2 := \text{id} + c, \quad \mathfrak{S}_n := \mathfrak{S}_{n-1,1}(\mathfrak{S}_{n-1} \otimes \text{id}).$$

### 2.2. The Nichols algebra $\mathfrak{B}(V_{abe})$ .

**Lemma 2.3.** *Let  $V = \mathbb{k}v_1 \oplus \mathbb{k}v_2$  be a vector space,  $c \in \text{End}_{\mathbb{k}}(V \otimes V)$  and suppose  $ab \neq 0$  such that*

$$\begin{aligned} c(v_1 \otimes v_1) &= av_2 \otimes v_2, & c(v_1 \otimes v_2) &= bv_1 \otimes v_2, \\ c(v_2 \otimes v_1) &= \gamma v_2 \otimes v_1, & c(v_2 \otimes v_2) &= ev_1 \otimes v_1. \end{aligned}$$

- (1)  $(V, c)$  is a braided vector space iff  $b = \gamma$ .
- (2) The braided vector space  $(V, c)$  is of diagonal type iff  $b^2 = ae$ .
- (3) Suppose  $b = \gamma$ ,  $b^2 = ae$ , denote

$$w_1 = v_1 + \sqrt{\frac{a}{b}}v_2, \quad w_2 = v_1 - \sqrt{\frac{a}{b}}v_2,$$

then  $(V, c)$  is a braided vector of diagonal type, where the braiding  $c$  on the base  $(w_1, w_2)$  is given by

$$\begin{aligned} c(w_1 \otimes w_1) &= bw_1 \otimes w_1, & c(w_1 \otimes w_2) &= -bw_2 \otimes w_1, \\ c(w_2 \otimes w_1) &= -bw_1 \otimes w_2, & c(w_2 \otimes w_2) &= bw_2 \otimes w_2. \end{aligned}$$

*Remark 2.4.* We denote the braided vector space as  $V_{abe}$  in following. And  $V_{abe}$  is isomorphic to  $V_{aeb1}$  via  $v_1 \mapsto \sqrt{e}v_1, v_2 \mapsto v_2$ .

**Corollary 2.5.** *Let  $b^2 = ae$ , then  $\mathfrak{B}(V_{abe})$  is of diagonal type and*

$$\dim \mathfrak{B}(V_{abe}) = \begin{cases} 4, & b = -1, (\mathfrak{B}(V_{abe}) \text{ is of Cartan type } A_1 \times A_1), \\ 27, & b^3 = 1 \neq b, (\mathfrak{B}(V_{abe}) \text{ is of Cartan type } A_2), \\ \infty, & \text{otherwise.} \end{cases}$$

**2.3. The dimension of  $\mathfrak{B}(V_{abe})$  on special cases.** Let  $N = \{1, 2\}$ , define an action of the symmetric group  $\mathbb{S}_2$  on  $N^2$  as

$$s_1 \cdot 11 = 22, \quad s_1 \cdot 22 = 11, \quad s_1 \cdot 12 = 12, \quad s_1 \cdot 21 = 21.$$

Then there is an action of the symmetric group  $\mathbb{S}_n$  on  $N^n$  induced by the action of  $\mathbb{S}_2$  on  $N^2$ .  $\Phi : \mathbb{S}_n \rightarrow \mathfrak{S}_n$  via  $s_i \mapsto c_i$  is a lifting of  $\mathbb{S}_n$  on the braid group  $\mathfrak{S}_n$ . As for convenience, we denote  $\Phi_\sigma$  as  $\Phi(\sigma)$  for any  $\sigma \in \mathbb{S}_n$ .

Let  $x = i_1 \cdots i_n, y = j_1 \cdots j_n \in N^n$ , define  $v_x = v_{i_1} \cdots v_{i_n}, v_y = v_{j_1} \cdots v_{j_n} \in V_{abe}^n$ ,  $\mathfrak{F}(x|y) = \{\sigma \in \mathbb{S}_n \mid \sigma \cdot x = y\}$  and  $\tilde{\mathfrak{F}}(x|y) \in \mathbb{K}$  such that

$$\sum_{\sigma \in \mathfrak{F}(x|y)} \Phi_\sigma(v_x) = \tilde{\mathfrak{F}}(x|y)v_y.$$

Let  $\bar{i}_k$  equal to 2 in case of  $i_k = 1$  and 1 otherwise for  $1 \leq k \leq n$ , and  $\bar{x} = \bar{i}_1 \cdots \bar{i}_n$ . Denote  $\mathcal{O}(x) = \{y \mid \sigma \cdot x = y, \sigma \in \mathbb{S}_n\}$ , Then it's easy to see that

**Lemma 2.6.** (1)  $\mathfrak{S}_n(v_x) = \sum_{y \in \mathcal{O}(x)} \tilde{\mathfrak{F}}(x|y)v_y$ ,

(2)  $\tilde{\mathfrak{F}}(x|y) = \tilde{\mathfrak{F}}(y|x) \Big|_{a=e}^{a=ae}$ ,

(3)  $\tilde{\mathfrak{F}}(\bar{x}|\bar{y}) = \tilde{\mathfrak{F}}(x|y) \Big|_{a=e}^{a=ae}$ .

(4) Denote  $\vec{x} = i_n \cdots i_1$  if  $x = i_1 \cdots i_n$ , then  $\tilde{\mathfrak{F}}(x|y) = \tilde{\mathfrak{F}}(\vec{x}|\vec{y})$ .

*Proof.* (1) It's easy to see.

(2)  $\tilde{\mathfrak{F}}(y|x) = \{w^{-1} \mid w \in \mathfrak{F}(x|y)\}$ .

(3)  $\tilde{\mathfrak{F}}(\bar{x}|\bar{y}) = \tilde{\mathfrak{F}}(x|y)$ .

(4) There is a bijection between  $\mathfrak{F}(x|y)$  and  $\mathfrak{F}(\vec{x}|\vec{y})$  via a flip of the diagrams of braidings. □

**Lemma 2.7.**

$$\begin{aligned} \mathfrak{F}(2^m(12)^k | 2^m(12)^k) &= (id^{\otimes m} \otimes \mathbb{S}_{2k-1,1}) \mathfrak{F}(2^{m-1}(21)^k | 2^{m-1}(21)^k) \\ &\cup \bigcup_{k_1=0}^{k_1 \leq \frac{m-1}{2}} \prod_{i=m-2k_1}^{m+2k-1} s_i \mathfrak{F}(2^{m-1}(21)^k | 2^{m-2k_1-1} 1^{2k_1} (21)^k), \\ \mathfrak{F}(2^m(21)^k | 2^m(21)^k) &= (id^{\otimes m} \otimes \mathbb{S}_{2k-1,1}) \mathfrak{F}(2^{m+1}(12)^{k-1} | 2^{m+1}(12)^{k-1}) \end{aligned}$$

$$\cup \bigcup_{k_1=0}^{k_1 \leq \frac{m}{2}} \prod_{i=m-2k_1+1}^{m+2k-1} s_i \mathfrak{F} \left( 2^{m+1} (12)^{k-1} | 2^{m-2k_1} 1^{2k_1} 2 (12)^{k-1} \right).$$

*Proof.* We give a proof for the first formula. Since  $\mathbb{S}_{m+2k} = \mathbb{S}_{m+2k-1,1} (\mathbb{S}_{m+2k-1} \otimes 1)$ , we only need to make clear which elements of  $N^{m+2k}$  are sent to  $2^m (12)^k$  under the action of  $\mathbb{S}_{m+2k-1,1}$ . In fact,  $2^m (12)^k$  is stable under the action of  $id^{\otimes m} \otimes \mathbb{S}_{2k-1,1}$ , and the rest elements are shown in Figure 1. The second formula is similar to prove, please refer the Figure 2.  $\square$




FIGURE 1.  $\mathfrak{F} \left( 2^m (12)^k | 2^m (12)^k \right)$




FIGURE 2.  $\mathfrak{F} \left( 2^m (21)^k | 2^m (21)^k \right)$

**Lemma 2.8.**  $\mathfrak{F} (2^n | 2^n)$  is subgroup of  $\mathbb{S}_n$  generated by  $t_i = s_i s_{i+1} s_i$  for  $1 \leq i \leq n-2$ .

*Proof.* When  $n = 2$ ,  $\mathfrak{F} (2^n | 2^n) = \{1\} = \mathbb{S}_1$ . When  $n = 3$ ,  $\mathfrak{F} (2^n | 2^n) = \{1, s_1 s_2 s_1\}$ . According to Lemma 2.7,

$$\begin{aligned} \mathfrak{F} (2^n | 2^n) &= \bigcup_{k_1=0}^{k_1 \leq \frac{n-1}{2}} \prod_{i=n-2k_1}^{n-1} s_i \mathfrak{F} \left( 2^{n-1} | 2^{n-2k_1-1} 1^{2k_1} \right) \\ &= \bigcup_{k_1=0}^{k_1 \leq \frac{n-1}{2}} \prod_{i=n-2k_1}^{n-1} s_i \prod_{k_2=0}^{k_1-1} s_{n-2(k_1-k_2)} \mathfrak{F} \left( 2^{n-1} | 2^{n-1} \right) \end{aligned}$$

$$= \bigcup_{k_1=0}^{k_1 \leq \frac{n-1}{2}} \prod_{k_2=0}^{k_1-1} t_{n-2(k_1-k_2)} \mathfrak{F}(2^{n-1}|2^{n-1}).$$

The Lemma is proved by induction.  $\square$

**Corollary 2.9.**  $|\mathcal{O}(2^{2n})| = \binom{2n}{n}$ ,  $|\mathcal{O}(2^{2n+1})| = \binom{2n+1}{n}$ .

*Proof.* Let  $1 \leq i \leq n-2$ ,  $T_n = \langle t_i \mid i \equiv 1 \pmod{2} \rangle$  and  $T'_n = \langle t_i \mid i \equiv 0 \pmod{2} \rangle$ , then  $\mathfrak{F}(2^{2n}|2^{2n}) = T_n T'_n = T'_n T_n \simeq \mathbb{S}_n \times \mathbb{S}_n$ . So  $|\mathfrak{F}(2^{2n}|2^{2n})| = (n!)^2$ , which implies  $|\mathcal{O}(2^{2n})| = \binom{2n}{n}$ . Similarly,  $\mathfrak{F}(2^{2n+1}|2^{2n+1}) \simeq \mathbb{S}_{n+1} \times \mathbb{S}_n$ , so  $|\mathcal{O}(2^{2n+1})| = \binom{2n+1}{n}$ .  $\square$

**Proposition 2.10.**

$$(2.1) \quad |\mathcal{O}(2^{2(n-k)}(12)^k)| = \binom{2n}{n-k},$$

$$(2.2) \quad |\mathcal{O}(2^{2(n-k)+1}(12)^k)| = \binom{2n+1}{n-k}.$$

*Proof.* Firstly, we observe that  $\mathfrak{F}(2^{m-1}(21)^k|2^{m-1}(21)^k)$  is a subgroup of  $\mathbb{S}_{m+2k}$  and  $\mathfrak{F}(2^{m-1}(21)^k|2^{m-1-2k_1}1^{2k_1}(21)^k)$  is a coset of  $\mathfrak{F}(2^{m-1}(21)^k|2^{m-1}(21)^k)$ . According to Lemma 2.7, we have

$$|\mathfrak{F}(2^{2m}(12)^k|2^{2m}(12)^k)| = (m+2k)|\mathfrak{F}(2^{2m-1}(21)^k|2^{2m-1}(21)^k)|.$$

Similarly, we can prove that

$$\begin{aligned} |\mathfrak{F}(2^{2m-1}(12)^k|2^{2m-1}(12)^k)| &= (m+2k)|\mathfrak{F}(2^{2m-2}(21)^k|2^{2m-2}(21)^k)|, \\ |\mathfrak{F}(2^{2m}(21)^k|2^{2m}(21)^k)| &= (m+2k)|\mathfrak{F}(2^{2m+1}(12)^{k-1}|2^{2m+1}(12)^{k-1})|, \\ |\mathfrak{F}(2^{2m+1}(21)^k|2^{2m+1}(21)^k)| &= (m+2k)|\mathfrak{F}(2^{2m+2}(12)^{k-1}|2^{2m+2}(12)^{k-1})|. \end{aligned}$$

Now we can give a direct computation as follows

$$\begin{aligned} &|\mathfrak{F}(2^{2(n-k)}(12)^k|2^{2(n-k)}(12)^k)| \\ &= (n+k)|\mathfrak{F}(2^{2(n-k)-1}(21)^k|2^{2(n-k)-1}(21)^k)| \\ &= (n+k)(n+k-1)|\mathfrak{F}(2^{2(n-k)}(12)^{k-1}|2^{2(n-k)}(12)^{k-1})| \\ &= (n+k)(n+k-1) \cdots (n-k+1)|\mathfrak{F}(2^{n-k}|2^{n-k})| \\ &= (n+k)!(n-k)! \\ &|\mathfrak{F}(2^{2(n-k)+1}(12)^k|2^{2(n-k)+1}(12)^k)| \\ &= (n-k+1+2k)|\mathfrak{F}(2^{2(n-k)}(21)^k|2^{2(n-k)}(21)^k)| \\ &= (n+k+1)(n-k+2k)|\mathfrak{F}(2^{2(n-k)+1}(12)^{k-1}|2^{2(n-k)+1}(12)^{k-1})| \end{aligned}$$

$$\begin{aligned}
&= (n+k+1)(n+k)\cdots(n-k+2)|\tilde{\mathfrak{F}}(2^{2(n-k)+1}|2^{2(n-k)+1})| \\
&= (n+k+1)!(n-k)!
\end{aligned}$$

□

**Theorem 2.11.**

$$\begin{aligned}
N^{2n} &= \mathcal{O}(2^{2n}) \oplus \bigoplus_{k=1}^n [\mathcal{O}(2^{2(n-k)}(21)^k) \oplus \mathcal{O}(1^{2(n-k)}(12)^k)], \\
N^{2n+1} &= \bigoplus_{k=0}^n [\mathcal{O}(2^{2(n-k)+1}(12)^k) \oplus \mathcal{O}(1^{2(n-k)+1}(21)^k)].
\end{aligned}$$

*Remark 2.12.* The dimensions of the irreducible submodules of  $N^n$  forms exactly the Pascal's triangle according to the Proposition 2.10 and the Theorem 2.11. This is not the first time that Pascal's triangle has been categorified in a module category, please refer to [6], [7], and [10].

*Proof.* According to Proposition 2.10, we only need to check that  $\mathcal{O}(2^{2(n-k)}(21)^k) \neq \mathcal{O}(1^{2(n-k)}(12)^k)$  and  $\mathcal{O}(2^{2(n-k)+1}(12)^k) \neq \mathcal{O}(1^{2(n-k)+1}(21)^k)$ . The latter is obvious since the action of  $\mathbb{S}_{2n+1}$  on  $N^{2n+1}$  always changes even 2's and 1's into each other. Suppose  $\mathcal{O}(2^{2(n-k)}(21)^k) = \mathcal{O}(1^{2(n-k)}(12)^k)$ , then

$$\mathcal{O}(22^{2(n-k)}(21)^k) = \mathcal{O}(21^{2(n-k)}(12)^k).$$

This is a contradiction, since

$$\begin{aligned}
\mathcal{O}(21^{2(n-k)}(12)^k) &= \mathcal{O}(2^{2(n-k)+1}(12)^k), \\
\mathcal{O}(22^{2(n-k)}(21)^k) &= \mathcal{O}(1^{2(n-k)}2(21)^k) = \mathcal{O}(1^{2(n-k+1)+1}(21)^{k-1}).
\end{aligned}$$

□

Denote  $(n)_b = 1 + b + \cdots + b^{n-1} = \frac{b^n-1}{b-1}$ ,  $(n)_b^! = \prod_{k=1}^n (k)_b$ .

**Theorem 2.13.** *Suppose  $b^2 \neq a = 1$ , then  $\dim \mathfrak{B}(V_{1b1}) < \infty$  if and only if  $b$  is a  $n$ -th primitive root of unity for  $n \geq 2$ . In particular,  $\dim \mathfrak{B}(V_{1b1}) = n^2$ .*

*Proof.* When  $a = e = 1$ ,  $\mathfrak{S}_2(v_1^2) = v_1^2 + v_2^2 = \mathfrak{S}_2(v_2^2)$ , so  $\mathfrak{S}_n(v_x) = \mathfrak{S}_n(v_y)$  for any  $y \in \mathcal{O}(x) \subset N^n$ . That is to say,  $\tilde{\mathfrak{F}}(x_1|y_1) = \tilde{\mathfrak{F}}(x_2|y_2)$  for  $x_1, x_2, y_1, y_2 \in \mathcal{O}(x_1) \subset N^n$  and  $\mathfrak{S}_n(v_x) = \tilde{\mathfrak{F}}(x|x) \sum_{y \in \mathcal{O}(x)} v_y$ .

$$\begin{aligned}
&\tilde{\mathfrak{F}}(2^{2(n-k)}(12)^k|2^{2(n-k)}(12)^k) \\
&= (n+k)_b \tilde{\mathfrak{F}}(2^{2(n-k)-1}(21)^k|2^{2(n-k)-1}(21)^k) \\
&= (n+k)_b (n+k-1)_b \tilde{\mathfrak{F}}(2^{2(n-k)}(12)^{k-1}|2^{2(n-k)}(12)^{k-1})
\end{aligned}$$

$$\begin{aligned}
&= (n+k)_b(n+k-1)_b \cdots (n-k+1)_b \tilde{\mathfrak{F}}(2^{n-k}|2^{n-k}) \\
&= (n+k)_b^! (n-k)_b^! \\
&\quad \tilde{\mathfrak{F}}(2^{2(n-k)+1}(12)^k | 2^{2(n-k)+1}(12)^k) \\
&= (n-k+1+2k)_b \tilde{\mathfrak{F}}(2^{2(n-k)}(21)^k | 2^{2(n-k)}(21)^k) \\
&= (n+k+1)_b(n-k+2k)_b \tilde{\mathfrak{F}}(2^{2(n-k)+1}(12)^{k-1} | 2^{2(n-k)+1}(12)^{k-1}) \\
&= (n+k+1)_b(n+k)_b \cdots (n-k+2)_b \tilde{\mathfrak{F}}(2^{2(n-k)+1} | 2^{2(n-k)+1}) \\
&= (n+k+1)_b^! (n-k)_b^!
\end{aligned}$$

Now we can put a basis of  $\mathfrak{B}(V_{1b1})$  on the Pascal's triangle and the coefficients of the symmetrizer's action on the basis is given by the Figure 3. When  $b$  is a




FIGURE 3. coefficients of the braided symmetrizer's action on the basis of  $\mathfrak{B}(V_{1b1})$

primitive  $n$ -th root of unity for  $n \geq 2$ , then all the nonzero coefficients located in a square on the top of the Pascal's triangle, see the case when  $n = 3$  in Figure 3. So  $\dim \mathfrak{B}(V_{1b1}) = n^2$ .  $\square$

**Theorem 2.14.** *Suppose  $b^2 \neq ae$  and  $b = -1$ , then  $\mathfrak{B}(V_{abe}) < \infty$  if  $ae$  is a  $m$ -th primitive root of unity for  $m \geq 1$ . In particular,  $\dim \mathfrak{B}(V_{abe}) = 4m$ .*

*Proof.* Since  $b = -1$ ,  $v_1 v_2 = v_2 v_1 = 0$ . Let  $x = i_1 \cdots i_n \in N^n$  such that  $i_k i_{k+1} = 12$  or  $21$  for some  $k = 0, \dots, n-1$ , then

$$0 = \mathfrak{S}_n(v_x) = \sum_{y \in O(x)} \tilde{\mathfrak{F}}(x, y) v_y \Rightarrow \tilde{\mathfrak{F}}(x, y) = 0.$$

According to Lemma 2.6, we have  $\tilde{\mathfrak{F}}(y, x) = 0$  for any  $y \in O(x)$ . So

$$\mathfrak{S}_n(v_1^n) = \begin{cases} \tilde{\mathfrak{F}}(1^n, 1^n)v_1^n + \tilde{\mathfrak{F}}(1^n, 2^n)v_2^n, & n \text{ is even,} \\ \tilde{\mathfrak{F}}(1^n, 1^n)v_1^n, & n \text{ is odd.} \end{cases}$$

Since  $\mathfrak{S}_n = \mathfrak{S}_{n-1,1}(\mathfrak{S}_{n-1} \otimes \text{id})$ , we have

$$\begin{aligned} \tilde{\mathfrak{F}}(1^{2m}, 2^{2m}) &= a(ab)^{m-1} \tilde{\mathfrak{F}}(1^{2m-1}, 1^{2m-1}) \\ \tilde{\mathfrak{F}}(1^{2m}, 1^{2m}) &= \tilde{\mathfrak{F}}(1^{2m-1}, 1^{2m-1}) \\ &= \tilde{\mathfrak{F}}(1^{2m-2}, 1^{2m-2}) + (eb)^{m-1} \tilde{\mathfrak{F}}(1^{2m-2}, 2^{2m-2}) \\ &= \tilde{\mathfrak{F}}(1^{2m-2}, 1^{2m-2}) + (eb)^{m-1} a(ab)^{m-2} \tilde{\mathfrak{F}}(1^{2m-3}, 1^{2m-3}) \\ &= [1 + (ae)^{m-1} b^{2m-3}] \tilde{\mathfrak{F}}(1^{2m-3}, 1^{2m-3}) \\ &= [1 - (ae)^{m-1}] \tilde{\mathfrak{F}}(1^{2m-3}, 1^{2m-3}) \\ \tilde{\mathfrak{F}}(1^{2m+1}, 1^{2m+1}) &= [1 - (ae)^m] [1 - (ae)^{m-1}] \cdots [1 - (ae)^2] \tilde{\mathfrak{F}}(1^3, 1^3) \\ &= [1 - (ae)^m] [1 - (ae)^{m-1}] \cdots [1 - (ae)^2] (1 - ae) \\ &= (1 - ae)^m (m)_{ae}! \end{aligned}$$

Similarly, we have

$$\begin{aligned} \tilde{\mathfrak{F}}(2^{2m}, 2^{2m}) &= \tilde{\mathfrak{F}}(2^{2m-1}, 2^{2m-1}) = (1 - ae)^{m-1} (m-1)_{ae}! \\ \tilde{\mathfrak{F}}(2^{2m}, 1^{2m}) &= e(eb)^{m-1} \tilde{\mathfrak{F}}(2^{2m-1}, 2^{2m-1}) = e^m b^{m-1} (1 - ae)^{m-1} (m-1)_{ae}! \end{aligned}$$

Since  $ae$  is a  $m$ -th primitive root,  $v_1^{2m+1} = 0$ ,  $v_2^{2m+1} = 0$ , and

$$\begin{aligned} \mathfrak{S}_{2m}(v_1^{2m}) &= (1 - ae)^{m-1} (m-1)_{ae}! [v_1^{2m} + a^m b^{m-1} v_2^{2m}] \\ \mathfrak{S}_{2m}(v_2^{2m}) &= (1 - ae)^{m-1} (m-1)_{ae}! [v_2^{2m} + e^m b^{m-1} v_1^{2m}] \\ &= e^m b^{m-1} \mathfrak{S}_{2m}(v_1^{2m}). \end{aligned}$$

So  $\dim \mathfrak{B}(V_{abe}) = 4m$ . □

### 3. THE POLYNOMIAL $\tilde{\mathfrak{F}}(1^n, 1^n)$ AND A CLASS OF COMBINATORIAL NUMBERS

Since  $V_{abe}$  is isomorphic to  $V_{ae b_1}$  as braided vector spaces and  $\tilde{\mathfrak{F}}(1^n, 1^n)$  is a polynomial of  $b$  and  $ae$ , we set  $e = 1$  for convenience. Let  $x, y \in N^n$ , denote

$$\tilde{\mathfrak{F}}(x, y) = \sum_{k=0} \tilde{\mathfrak{F}}_k(x, y) b^k,$$

where  $\tilde{\mathfrak{F}}_k(x, y)$  is a polynomial of  $a$ . Then

$$\tilde{\mathfrak{F}}(1^n, 1^n) = \sum_{k=0}^{n(n-1)/2} \tilde{\mathfrak{F}}_k(1^n, 1^n) b^k.$$



Let  $w \in \mathfrak{F}(1^n | 1^n)$ , define  $\text{tl}(w)$  as the minimal length of  $w$  in expressions of  $t_i$ 's and  $\text{sl}(w)$  as the minimal length of  $w$  in expressions of  $s_i$ 's. Denote

$$\mathcal{E}_{k,s}^n = \# \left\{ \sigma \in \mathfrak{F}(1^n, 1^n) \mid \text{tl}(\sigma) = k, \text{sl}(\sigma) = s \right\}.$$

**Proposition 3.1.**  $\tilde{\mathfrak{F}}_k(1^n, 1^n) = \sum_{\text{tl}(w)=k} \#\{\text{sl}(w)\} a^{k-[3k-\text{sl}(w)]/2} = \sum_{i=0} \mathcal{E}_{k,2i+k}^n a^i$ .

*Proof.* According to the Lemma 2.8,  $\mathfrak{F}(1^n, 1^n) = \langle t_i = s_i s_{i+1} s_i \mid i = 1, \dots, n-2 \rangle$ . In case  $n = 2m$  is even, then  $\mathfrak{F}(1^{2m}, 1^{2m}) \simeq \mathbb{S}_m \times \mathbb{S}_m$ , where the first  $\mathbb{S}_m$  is the symmetric group on even numbers and the second  $\mathbb{S}_m$  is the symmetric group on odd numbers. If  $a = e = 1$ , then  $\tilde{\mathfrak{F}}(1^{2m}, 1^{2m}) = \binom{m}{b}_a \binom{m}{b}_a$ . So for any  $w \in \mathfrak{F}(1^{2m}, 1^{2m})$ , its contribution to  $\tilde{\mathfrak{F}}(1^{2m}, 1^{2m})$  is  $a^{k-[3k-\text{sl}(w)]/2} b^{\text{tl}(w)}$ . The similar result holds on case that  $n$  is odd.  $\square$

**Lemma 3.2.**  $\tilde{\mathfrak{F}}_0(1^n, 1^n) = 1$ ,  $\tilde{\mathfrak{F}}_1(1^n, 1^n) = (n-2)a$  for  $n \geq 2$ .

**Corollary 3.3.**  $\mathcal{E}_{1,s}^n = \begin{cases} n-2, & s=3, n \geq 2, \\ 0, & \text{otherwise.} \end{cases}$

**Lemma 3.4.**  $\tilde{\mathfrak{F}}_{n(n-1)}(1^{2n}, 1^{2n}) = a^{n(n-1)/2}$  and  $\tilde{\mathfrak{F}}_{n^2}(1^{2n+1}, 1^{2n+1}) = a^{n(n+1)/2}$ .

*Proof.* Let  $w$  be the longest element of  $\mathfrak{F}(1^{2n} | 1^{2n})$  in the sense of  $\text{tl}(w)$ , then

$$\begin{aligned} w &= (t_1 t_3 \cdots t_{2n-3})(t_1 t_3 \cdots t_{2n-5}) \cdots (t_1 t_3) t_1 (t_2 t_4 \cdots t_{2n-2})(t_2 t_4 \cdots t_{2n-4}) \cdots (t_2 t_4) t_2 \\ &= (2n-1, 2n-3, \dots, 3, 1)(2n, 2n-2, \dots, 4, 2) \\ &= (2n-1, 2n, 2n-3, 2n-2, \dots, 3, 4, 1, 2). \end{aligned}$$

So  $\text{sl}(w) = \text{inv}(2n-3, 2n-2, 2n-5, 2n-4, \dots, 3, 4, 1, 2) = 2n(n-1)$ , and

$$\tilde{\mathfrak{F}}_{n(n-1)}(1^{2n}, 1^{2n}) = a^{n(n-1)-[3n(n-1)-\text{sl}(w)]/2} = a^{n(n-1)/2}.$$

Let  $w'$  be the longest element of  $\mathfrak{F}(1^{2n+1} | 1^{2n+1})$  in the sense of  $\text{tl}(w')$ , then

$$\begin{aligned} w' &= (t_1 t_3 \cdots t_{2n-1}) w \\ &= (2n+1, 2n-1, 2n-3, \dots, 3, 1)(2n, 2n-2, \dots, 4, 2) \\ &= (2n+1, 2n, 2n-1, 2n-2, \dots, 4, 3, 2, 1), \end{aligned}$$

which imply that  $\text{tl}(w') = n^2$  and  $\text{sl}(w') = n(2n+1)$ . So

$$\tilde{\mathfrak{F}}_{n^2}(1^{2n+1}, 1^{2n+1}) = a^{n^2-[3n^2-n(2n+1)]/2} = a^{n(n+1)/2}.$$

$\square$

**Lemma 3.5.**  $\tilde{\mathfrak{F}}_2(1^n, 1^n) = \begin{cases} \frac{(n-1)(n-4)}{2} a^2 + (n-3)a, & n \geq 4, \\ 0, & n \leq 3. \end{cases}$

*Proof.* It's easy to see  $\tilde{\mathfrak{F}}_2(1^4, 1^4) = a$ . When  $n \geq 5$ , then

$$\begin{aligned}
\tilde{\mathfrak{F}}_2(1^n, 1^n) &= \tilde{\mathfrak{F}}_2(1^{n-1}, 1^{n-1}) + \tilde{\mathfrak{F}}_1(1^{n-1}, 1^{n-3}2^2) + \tilde{\mathfrak{F}}_0(1^{n-1}, 1^{n-5}2^4) \\
&= \tilde{\mathfrak{F}}_2(1^{n-1}, 1^{n-1}) + [a\tilde{\mathfrak{F}}_1(1^{n-2}, 1^{n-2}) + a\tilde{\mathfrak{F}}_0(1^{n-2}, 1^{n-2})] + a^2 \\
&= \tilde{\mathfrak{F}}_2(1^{n-1}, 1^{n-1}) + (n-3)a^2 + a \\
&= \tilde{\mathfrak{F}}_2(1^4, 1^4) + (2+3+\cdots+n-3)a^2 + (n-4)a \\
&= \frac{(n-1)(n-4)}{2}a^2 + (n-3)a.
\end{aligned}$$

□

**Corollary 3.6.**  $\mathcal{E}_{2,s}^n = \begin{cases} \frac{(n-1)(n-4)}{2}, & s = 6, n \geq 4, \\ n-3, & s = 4, n \geq 4, \\ 0, & \text{otherwise.} \end{cases}$

**Lemma 3.7.**  $\tilde{\mathfrak{F}}_3(1^n, 1^n) = \begin{cases} \frac{(n+1)(n-4)(n-6)}{6}a^3 + (n^2 - 4n - 2)a^2, & n \geq 6, \\ 3a^2, & n = 5, \\ 0, & n \leq 4. \end{cases}$

*Proof.* When  $n \geq 7$ , then

$$\begin{aligned}
\tilde{\mathfrak{F}}_3(1^n, 1^n) &= \tilde{\mathfrak{F}}_3(1^{n-1}, 1^{n-1}) + \tilde{\mathfrak{F}}_0(1^{n-1}, 1^{n-7}2^6) + a[\tilde{\mathfrak{F}}_2(1^{n-2}, 1^{n-2}) \\
&\quad + \tilde{\mathfrak{F}}_1(1^{n-2}, 1^{n-2}) + \tilde{\mathfrak{F}}_0(1^{n-2}, 1^{n-6}2^21^2) + \tilde{\mathfrak{F}}_0(1^{n-3}, 1^{n-7}1^22^2)] \\
&\quad + a^2[\tilde{\mathfrak{F}}_0(1^{n-2}, 1^{n-2}) + \tilde{\mathfrak{F}}_1(1^{n-4}, 1^{n-4}) + \tilde{\mathfrak{F}}_0(1^{n-4}, 1^{n-4})] \\
&= \tilde{\mathfrak{F}}_3(1^{n-1}, 1^{n-1}) + a^3 \\
&\quad + a\left[\frac{(n-3)(n-6)}{2}a^2 + (n-5)a + (n-4)a + a + a\right] \\
&\quad + a^2[1 + (n-6)a + 1] \\
&= \tilde{\mathfrak{F}}_3(1^{n-1}, 1^{n-1}) + \frac{n^2 - 7n + 8}{2}a^3 + (2n-5)a^2 \\
&= \tilde{\mathfrak{F}}_3(1^6, 1^6) + \sum_{k=7}^n \frac{k^2 - 7k + 8}{2}a^3 + \sum_{k=7}^n (2k-5)a^2 \\
&= \frac{(n+1)(n-4)(n-6)}{6}a^3 + (n^2 - 4n - 2)a^2.
\end{aligned}$$

□

**Corollary 3.8.**  $\mathcal{E}_{3,s}^n = \begin{cases} \frac{(n+1)(n-4)(n-6)}{6}, & s = 9, n \geq 6, \\ n^2 - 4n - 2, & s = 7, n \geq 5, \\ 0, & \text{otherwise.} \end{cases}$

**Lemma 3.9.** *When  $n \leq 8$ , then*

$$\tilde{\mathfrak{F}}_4(1^n, 1^n) = \begin{cases} 17a^2 + 52a^3 + 2a^4, & n = 8, \\ 10a^2 + 19a^3 + a^4, & n = 7, \\ 4a^2 + 4a^3, & n = 6, \\ a^3, & n = 5, \\ 0, & n \leq 4. \end{cases}$$

*When  $n > 8$ , then  $\tilde{\mathfrak{F}}_4(1^n, 1^n)$  equals*

$$\frac{(n-7)(n^3 - 7n^2 - 14n + 96)}{24}a^4 + \frac{n^3 - 6n^2 - 13n + 80}{2}a^3 + \frac{n^2 - n - 22}{2}a^2.$$

*Proof.* We only prove the case  $n \geq 9$ .

$$\begin{aligned} \tilde{\mathfrak{F}}_4(1^n, 1^n) &= \tilde{\mathfrak{F}}_4(1^{n-1}, 1^{n-1}) + \tilde{\mathfrak{F}}_0(1^{n-1}, 1^{n-9}2^8) + a\tilde{\mathfrak{F}}_3(1^{n-2}, 1^{n-2}) \\ &\quad + a\tilde{\mathfrak{F}}_2(1^{n-2}, 1^{n-2}) + a\tilde{\mathfrak{F}}_0(1^{n-2}, 1^{n-8}2^41^2) + a\tilde{\mathfrak{F}}_0(1^{n-3}, 1^{n-7}2^4) \\ &\quad + a\tilde{\mathfrak{F}}_0(1^{n-4}, 1^{n-6}2^2) + a^2\tilde{\mathfrak{F}}_1(1^{n-2}, 1^{n-2}) + a^2\tilde{\mathfrak{F}}_0(1^{n-2}, 1^{n-2}) \\ &\quad + a^2\tilde{\mathfrak{F}}_0(1^{n-2}, 1^{n-7}2^21^3) + a^2\tilde{\mathfrak{F}}_0(1^{n-3}, 1^{n-5}2^2) + a^2\tilde{\mathfrak{F}}_0(1^{n-3}, 1^{n-7}21^22) \\ &\quad + a^2\tilde{\mathfrak{F}}_2(1^{n-4}, 1^{n-4}) + a^2\tilde{\mathfrak{F}}_1(1^{n-4}, 1^{n-4}) + a^2\tilde{\mathfrak{F}}_0(1^{n-4}, 1^{n-8}2^21^2) \\ &\quad + a^2\tilde{\mathfrak{F}}_1(1^{n-4}, 1^{n-4}) + (a^2 + a^3)\tilde{\mathfrak{F}}_0(1^{n-4}, 1^{n-4}) + a^2\tilde{\mathfrak{F}}_1(1^{n-5}, 1^{n-5}) \\ &\quad + a^2\tilde{\mathfrak{F}}_0(1^{n-5}, 1^{n-5}) + a^2\tilde{\mathfrak{F}}_0(1^{n-5}, 1^{n-7}2^2) \\ &\quad + a^3\tilde{\mathfrak{F}}_1(1^{n-6}, 1^{n-6}) + a^3\tilde{\mathfrak{F}}_0(1^{n-6}, 1^{n-6}) \\ &= \tilde{\mathfrak{F}}_4(1^{n-1}, 1^{n-1}) + a^4 + a \left[ \frac{(n-1)(n-6)(n-8)}{6}a^3 + (n^2 - 8n + 10)a^2 \right] \\ &\quad + a \left[ \frac{(n-3)(n-6)}{2}a^2 + (n-5)a \right] + a \cdot a^2 + a \cdot a^2 + a \cdot a \\ &\quad + a^2 \cdot (n-4)a + a^2 \cdot 1 + a^2 \cdot a + a^2 \cdot a + a^2 \cdot a^2 \\ &\quad + a^2 \left[ \frac{(n-5)(n-8)}{2}a^2 + (n-7)a \right] + a^2 \cdot (n-6)a + a^2 \cdot a \\ &\quad + a^2 \cdot (n-6)a + (a^2 + a^3) + a^2 \cdot (n-7)a \\ &\quad + a^2 + a^2 \cdot a + a^3 \cdot (n-8)a + a^3 \\ &= \tilde{\mathfrak{F}}_4(1^{n-1}, 1^{n-1}) + \frac{n^3 - 12n^2 + 29n + 36}{6}a^4 + \frac{3n^2 - 15n - 6}{2}a^3 \\ &\quad + (n-1)a^2 \\ &= \tilde{\mathfrak{F}}_4(1^8|1^8) + \frac{a^4}{6} \sum_{k=9}^n k^3 + \left( -2a^4 + \frac{3}{2}a^3 \right) \sum_{k=9}^n k^2 \end{aligned}$$

$$\begin{aligned}
& + \left( \frac{29}{6}a^4 - \frac{15}{2}a^3 + a^2 \right) \sum_{k=9}^n k + (6a^4 - 3a^3 - a^2)(n-9+1) \\
& = \frac{(n-7)(n^3 - 7n^2 - 14n + 96)}{24}a^4 + \frac{n^3 - 6n^2 - 13n + 80}{2}a^3 \\
& \quad + \frac{n^2 - n - 22}{2}a^2.
\end{aligned}$$

□

**Corollary 3.10.**  $\mathcal{E}_{4,s}^n = 0$ , except the following cases.

$$\begin{aligned}
& \mathcal{E}_{4,12}^7 = 1, \quad \mathcal{E}_{4,10}^6 = 4, \quad \mathcal{E}_{4,10}^5 = 1, \\
& \mathcal{E}_{4,s}^n = \begin{cases} \frac{(n-7)(n^3-7n^2-14n+96)}{24}, & s = 12, n \geq 8, \\ \frac{n^3-6n^2-13n+80}{2}, & s = 10, n \geq 7, \\ \frac{n^2-n-22}{2}, & s = 8, n \geq 6. \end{cases}
\end{aligned}$$

**Lemma 3.11.** When  $n < 11$ , then

$$\tilde{\mathfrak{F}}_5(1^n, 1^n) = \begin{cases} 10a^5 + 234a^4 + 226a^3 + 4a^2, & n = 10, \\ 4a^5 + 96a^4 + 131a^3 + 3a^2, & n = 9, \\ 32a^4 + 62a^3 + 2a^2, & n = 8, \\ 10a^4 + 19a^3 + a^2, & n = 7, \\ 4a^3, & n = 6, \\ 0, & n < 6. \end{cases}$$

When  $n \geq 11$ , then

$$\begin{aligned}
\tilde{\mathfrak{F}}_5(1^n, 1^n) & = \frac{n^5 - 20n^4 + 75n^3 + 740n^2 - 5716n + 9360}{120}a^5 + (n-6)a^2 \\
& \quad + \frac{n^4 - 9n^3 - 34n^2 + 474n - 936}{6}a^4 + \frac{n^3 - n^2 - 62n + 172}{2}a^3.
\end{aligned}$$

*Proof.* We only prove the case  $n \geq 11$ .

$$\begin{aligned}
\tilde{\mathfrak{F}}_5(1^n, 1^n) & = \tilde{\mathfrak{F}}_5(1^{n-1}, 1^{n-1}) + a\tilde{\mathfrak{F}}_4(1^{n-2}, 1^{n-2}) + a\tilde{\mathfrak{F}}_3(1^{n-2}, 1^{n-2}) \\
& \quad + a^2\tilde{\mathfrak{F}}_3(1^{n-4}, 1^{n-4}) + a^2\tilde{\mathfrak{F}}_2(1^{n-2}, 1^{n-2}) + 2a^2\tilde{\mathfrak{F}}_2(1^{n-4}, 1^{n-4}) \\
& \quad + a^2\tilde{\mathfrak{F}}_2(1^{n-5}, 1^{n-5}) + a^3\tilde{\mathfrak{F}}_2(1^{n-6}, 1^{n-6}) \\
& \quad + (2n-15)a^5 + (8n-43)a^4 + (4n-5)a^3 + a^2 \\
& = \tilde{\mathfrak{F}}_5(1^{n-1}, 1^{n-1}) + a \left\{ \frac{(n-9)(n^3 - 13n^2 + 26n + 88)}{24}a^4 \right. \\
& \quad \left. + \frac{n^3 - 12n^2 + 23n + 74}{2}a^3 + \frac{n^2 - 5n - 16}{2}a^2 \right\}
\end{aligned}$$

$$\begin{aligned}
& + a \left[ \frac{(n-1)(n-6)(n-8)}{6} a^3 + (n^2 - 8n + 10) a^2 \right] \\
& + a^2 \left[ \frac{(n-3)(n-8)(n-10)}{6} a^3 + (n^2 - 12n + 30) a^2 \right] \\
& + \frac{(n-7)(n-10)}{2} a^5 + (2n^2 - 24n + 67) a^4 + (4n - 27) a^3 \\
& + (2n - 15) a^5 + (8n - 43) a^4 + (4n - 5) a^3 + a^2 \\
& = \tilde{\mathfrak{E}}_5(1^{n-1}, 1^{n-1}) + \frac{n^4 - 18n^3 + 71n^2 + 234n - 1272}{24} a^5 \\
& + \frac{4n^3 - 33n^2 - 37n + 498}{6} a^4 + \frac{3n^2 - 5n - 60}{2} a^3 + a^2 \\
& = \tilde{\mathfrak{E}}_5(1^{10}, 1^{10}) + \sum_{k=11}^n \frac{k^4 - 18k^3 + 71k^2 + 234k - 1272}{24} a^5 \\
& + \sum_{k=11}^n \left[ \frac{4k^3 - 33k^2 - 37k + 498}{6} a^4 + \frac{3k^2 - 5k - 60}{2} a^3 + a^2 \right] \\
& = 10a^5 + 234a^4 + 226a^3 + 4a^2 + (n-10)a^2 \\
& + \frac{(n-10)(n-2)(n^3 - 8n^2 - 41n + 408)}{120} a^5 \\
& + \frac{(n-10)(n^3 + n^2 - 24n + 234)}{6} a^4 + \frac{(n-10)(n^2 + 9n + 28)}{2} a^3 \\
& = \frac{n^5 - 20n^4 + 75n^3 + 740n^2 - 5716n + 9360}{120} a^5 + (n-6)a^2 \\
& + \frac{n^4 - 9n^3 - 34n^2 + 474n - 936}{6} a^4 + \frac{n^3 - n^2 - 62n + 172}{2} a^3.
\end{aligned}$$

□

**Corollary 3.12.**  $\mathcal{E}_{5,s}^n = 0$ , except the following cases.

$$\begin{aligned}
& \mathcal{E}_{5,15}^9 = 4, \quad \mathcal{E}_{5,13}^8 = 32, \quad \mathcal{E}_{5,11}^7 = 19, \quad \mathcal{E}_{5,11}^6 = 4, \\
& \mathcal{E}_{5,s}^n = \begin{cases} \frac{n^5 - 20n^4 + 75n^3 + 740n^2 - 5716n + 9360}{120}, & s = 15, n \geq 10, \\ \frac{n^4 - 9n^3 - 34n^2 + 474n - 936}{6}, & s = 13, n \geq 9, \\ \frac{n^3 - n^2 - 62n + 172}{2}, & s = 11, n \geq 8, \\ n - 6, & s = 9, n \geq 7. \end{cases}
\end{aligned}$$

## REFERENCES

- [1] N. Andruskiewitsch and J. Giraldi. Nichols algebras that are quantum planes. *Linear and Multilinear Algebra*, 66(5):961–991, 2018.

- [2] N. Andruskiewitsch and M. Graña. From racks to pointed Hopf algebras. *Adv. Math.*, 178(2):177–243, 2003.
- [3] N. Andruskiewitsch and H-J. Schneider. Pointed Hopf algebras. In *New directions in Hopf algebras*, volume 43 of *Math. Sci. Res. Inst. Publ.*, pages 1–68. Cambridge Univ. Press, Cambridge, 2002.
- [4] I. Angiono. On Nichols algebras of diagonal type. *J. Reine Angew. Math.*, 683:189–251, 2013.
- [5] I. Angiono. A presentation by generators and relations of Nichols algebras of diagonal type and convex orders on root systems. *J. Eur. Math. Soc. (JEMS)*, 17(10):2643–2671, 2015.
- [6] D. Flath, T. Halverson, and K. Herbig. The planar rook algebra and Pascal’s triangle. *Enseign. Math. (2)*, 55(1-2):77–92, 2009.
- [7] A. M. Gainutdinov, J. L. Jacobsen, H. Saleur, and R. Vasseur. A physical approach to the classification of indecomposable Virasoro representations from the blob algebra. *Nuclear Phys. B*, 873(3):614–681, 2013.
- [8] I. Heckenberger. The Weyl groupoid of a Nichols algebra of diagonal type. *Invent. Math.*, 164(1):175–188, 2006.
- [9] I. Heckenberger. Classification of arithmetic root systems. *Adv. Math.*, 220(1):59–124, 2009.
- [10] M. S. Im and E. Norton. Irreducible calibrated representations of periplectic Brauer algebras and hook representations of the symmetric group. *Journal of Algebra*, 560:442–485, 2020.
- [11] G. Lusztig. *Introduction to quantum groups*, volume 110 of *Progress in Mathematics*. Birkhäuser Boston, Inc., Boston, MA, 1993.
- [12] W.D. Nichols. Bialgebras of type one\*. *Communications in Algebra*, 6(15):1521–1552, 1978.
- [13] M. Rosso. Quantum groups and quantum shuffles. *Invent. Math.*, 133(2):399–416, 1998.
- [14] P. Schauenburg. A characterization of the Borel-like subalgebras of quantum enveloping algebras. *Comm. Algebra*, 24(9):2811–2823, 1996.
- [15] Y.-X. Shi. Finite-dimensional Hopf algebras over the Kac-Paljutkin algebra  $H_8$ . *Rev. Un. Mat. Argentina*, 60(1):265–298, 2019.
- [16] Y.-X. Shi. Finite dimensional Nichols algebras over Suzuki algebra I: over simple Yetter-Drinfeld modules  $A_{N2n}^{\mu\lambda}$ . *arXiv:2011.14274*, 2020.
- [17] Y.-X. Shi. Finite dimensional Nichols algebras over Suzuki algebra II: over simple Yetter-Drinfeld modules of  $A_{N2n+1}^{\mu\lambda}$ . *to appear in arXiv*, 2020.
- [18] S. L. Woronowicz. Differential calculus on compact matrix pseudogroups (quantum groups). *Comm. Math. Phys.*, 122(1):125–170, 1989.

SCHOOL OF MATHEMATICS AND STATISTICS, JIANGXI NORMAL UNIVERSITY, NANCHANG 330022, P. R. CHINA

*Email address:* yxshi@jxnu.edu.cn