

Upward Planar Drawings with Three Slopes

Jonathan Klawitter 

University of Würzburg, Germany

Johannes Zink 

University of Würzburg, Germany

Abstract

We study upward planar straight-line drawings that use only three different slopes. We show that deciding whether a digraph admits such a drawing is NP-hard already for embedded outerplanar digraphs, though linear-time solvable for trees with and without given embedding.

1 Introduction

One of the main goals in graph drawing is to generate clear drawings. To achieve this, we may reduce the visual complexity by using only few different geometric primitives [22] – in our case few slopes. Using only few different slopes is common for schematic drawings; for example, if we allow two different slopes we get orthogonal drawings [9], with three or four slopes we get hexalinear and octilinear drawings [30], respectively. Here, we additionally require our graphs to be drawn upward and planar.

Upward planarity. An *upward planar drawing* of a directed graph (or digraph for short) G is a planar drawing of G where every edge is drawn as a monotonic upward curve. We call G *upward planar* if it admits an upward planar drawing and *upward plane* if it is equipped with an upward planar embedding. Note that an upward planar embedding, given by the cyclic sequences of edges around each vertex, is necessarily bimodal, that is, each cyclic sequence can be split into a contiguous subsequence of incoming edges and a contiguous subsequence of outgoing edges [9].

Upward planarity testing is an NP-complete problem for general digraphs [17], though several fixed-parameter tractable algorithms exist [5, 14, 18]. Furthermore, the problem is polynomial-time solvable, for example for single source digraphs [3], outerplanar digraphs [32], series-parallel digraphs [14], and triconnected digraphs [2]. If the embedding of a digraph is given, upward planarity can be tested in polynomial time [2].


k -slope drawings. A *k -slope drawing* of a graph G is a straight-line drawing of G where every edge is drawn with one of at most k different slopes; see Fig. 1a–b. The (*planar*) *slope number* of G is the smallest k such that G admits a planar k -slope drawing. If only planar drawings are allowed, this is called the *planar slope number* of G . Both numbers have been studied extensively for a variety of classes [4, 11, 13, 15, 16, 20, 21, 24, 26–28, 31, 33]. Determining the planar slope number of a graph is hard in the existential theory of the reals [19].

Di Battista and Tamassia [10] have shown that if a digraph is upward planar, then it even admits an upward planar straight-line drawing. We may therefore ask how many slopes are necessary for a digraph G to admit an upward planar drawing. The answer to this question defines the *upward planar slope number* of G . To the best of our knowledge, this number has so far only been studied by Czyzowicz et al. [6, 7] for lattices and, allowing one bend per edge, by Di Giacomo et al. [12] for series-parallel digraphs and by Bekos et al. [1] for st-graphs.

Most research on slope numbers is concerned with the minimum number of slopes required to draw any graph of a fixed class. We can consider the problem also from a different

2 Upward Planar Drawings with Three Slopes

perspective, namely, given only k slopes, decide whether a given digraph admits an *upward planar k -slope drawing*. For $k = 2$, this question has been investigated by Klawitter and Mchedlidze [23] who show that deciding whether a given upward plane digraph admits an upward planar 2-slope drawing can be done in linear time. For the variable embedding scenario, they give a linear-time algorithm for single-source digraphs, a quartic-time algorithm for series-parallel digraphs, and a fixed-parameter tractable algorithm for general digraphs. Here, we investigate the next natural case $k = 3$. Note that a 2-slope drawing can be stretched in the direction of one slope without affecting the length of edges drawn with the other slope. The fact that this does not hold for three (or more) slopes introduces interesting new geometric aspects for 3-slope drawings.



■ **Figure 1** (a) A digraph G with (b) upward planar 3-slope drawing; (c) drawing rotated by 45° .


Note that the maximum indegree and outdegree of a digraph G are natural lower bounds on its upward planar slope number. Since we have $k = 3$, we assume that our digraphs have maximum indegree and outdegree at most three. Further note that an upward planar 3-slope drawing of a digraph G on any three slopes can be affinely transformed into one with slopes 45° , 90° , and 135° . Hence, we restrict considerations to this slope set. For illustrative purposes however, we rotate drawings by 45° clockwise and thus use the slope set $\{\uparrow, \nearrow, \rightarrow\}$; see Fig. 1c. A *3-slope assignment* of a digraph G assigns each edge of G one of the slopes in $\{\uparrow, \nearrow, \rightarrow\}$. If G is upward plane, we say a 3-slope assignment of G is *consistent* if the assignment complies with the cyclic edge order around each vertex; for example, if a vertex has three incoming edges, then they need to be assigned the slopes \rightarrow , \nearrow , and \uparrow in ccw order. Clearly, if an upward plane embedding does not admit a consistent 3-slope assignment, then it also does not admit an upward plane 3-slope drawing.

Contribution. We initiate research on upward planar 3-slope drawings by examining the complexity of the decision problem. Some of our results are generalizable to $k \geq 3$ slopes. We classify whether an embedded tree T admits an upward planar 3-slope drawing. In the affirmative, we can construct such a drawing. Otherwise, we can change the embedding of T such that it admits a desired drawing. We further show that it is NP-hard to decide whether a given upward outerplanar¹ digraph admits an upward planar 3-slope drawing.

¹ An *upward outerplanar* digraph admits an upward planar drawing with each vertex on the outer face.

2 Trees

We start with the class of directed trees² with indegree and outdegree at most three. While every bimodal embedding of a tree induces an upward planar embedding, it does not necessarily admit an upward planar 3-slope drawing; see tree T in Fig. 2a. There the edge uv of T needs to get slope \uparrow by u but slope \nearrow by v and thus obstructs a consistent 3-slope assignment. However, we can always change the embedding of T such that it admits a consistent 3-slope assignment and drawing; see Fig. 2b. We show that both testing whether an embedding is suitable and finding a suitable embedding can be accomplished in linear time.



■ **Figure 2** (a) The tree T does not admit a consistent 3-slope assignment, (b) unless we change the embedding; (c) an upward planar 3-slope drawing of T with exponentially shrinking edge lengths.

► **Theorem 2.1** (\star). *Under a fixed bimodal embedding, a directed tree T admits an upward planar 3-slope drawing if and only if it admits a consistent 3-slope assignment. Moreover, this can be tested in linear time and, in the affirmative, a drawing can be constructed.*

Proof sketch. While the “only-if”-direction is clear, the converse can be shown by drawing T with edge lengths exponentially shrinking as the distance from a start edge increases; see Fig. 2c. A consistent 3-slope assignment is either directly obstructed by an edge as in Fig. 2a or can be computed straightforwardly. ◀

► **Theorem 2.2** (\star). *Given a directed tree T with indegree and outdegree at most three, we can construct an upward planar 3-slope drawing of T in linear time.*

Proof sketch. To find a suitable embedding, pick a start vertex, order its incident edges bimodally, and continue vertex by vertex to construct a consistent 3-slope assignment. ◀

We conjecture that a similar approach also works for cactus graphs, though constructing an actual drawing is geometrically more involved. We will investigate this further.

3 Outerplanar Graphs


In this section, we show that already for upward outerplanar digraphs, it is NP-hard to decide whether a digraph admits an upward planar 3-slope drawing. It remains open whether the problem is also NP-complete since containment in NP is not immediately clear. More precisely, if the problem was in NP, there would be small proof certificates for yes-instances that a verifier could use to decide the problem in polynomial time. Typically a combinatorial

² A *directed tree* (or *polytree*) is a digraph whose underlying undirected graph is a tree.


4 Upward Planar Drawings with Three Slopes

characterization or a drawing of the input graph could act as such a certificate. However in our case, we do not know whether there are graphs that require irrational (or super-polynomial precise) coordinates and if so, how to treat them implicitly. Yet NP-hardness holds true, even if an upward outerplanar embedding is given. Remember that for an arbitrary number of slopes, upward planarity for outerplanar digraphs can be decided in polynomial time [32] – if an embedding is given, even in linear time. We show NP-hardness by reduction from PLANAR MONOTONE 3-SAT [8], an NP-complete version of 3-SAT, where the three literals of each clause are all either negated or unnegated – from now on called *negative* and *positive* clauses, respectively. Moreover, the incidence graph³ is planar and has a planar drawing where the vertices are rectangles, the edges are vertical straight-line segments, the variables are arranged on a horizontal line, the positive clauses are above, and the negative clauses are below this line. For an example, see Fig. 5a. For a given formula F and a rectangular drawing of its incidence graph, we construct a corresponding upward outerplanar digraph G_F , which can only be drawn upward planar with 3 slopes if F is satisfiable. Our construction follows ideas of Nöllenburg [29] and Kraus [25] and utilizes the following observations.

Up to scaling and mirroring diagonally, G_{\square} in Fig. 3a admits an upward planar 3-slope drawing only as an outerplanar square as in Fig. 3b. We can attach multiple squares (and triangles) to each other as in Fig. 3c–d. The drawing of such a bigger digraph is unique up to scaling and mirroring diagonally. If the squares form a tree, the drawing is outerplanar. We refer to these squares as *unit squares*, since, once set, the side lengths for all attached squares are the same. To allow a certain small degree of freedom, we exploit the following.




■ **Figure 3** (a) The digraph G_{\square} admits only an upward 3-slope drawing as square (b); (c, d) by combining copies of G_{\square} and triangles we can build larger rigid structures.




■ **Figure 4** Upward planar 3-slope drawing of the digraph G_{\leftrightarrow} .

³ In the incidence graph of a SAT formula, there is a vertex for each variable and each clause, and for each occurrence of a variable in a clause, there is an edge between the corresponding vertices.



(a) Rectangular drawing of the incidence graph of a PLANAR MONOTONE 3-SAT formula F .



(b) Outerplanar drawing of the digraph G_F obtained from (a). Chains of unit squares are drawn as straight line segments. The variable/clause/edge gadgets occupy the areas of their corresponding rectangles. Here, x_1 and x_4 are set to false (brush on the left), while x_2 and x_3 are set to true (brush on the right)

■ **Figure 5** Schematic example for our NP-hardness reduction from PLANAR MONOTONE 3-SAT.


► **Lemma 3.1** (\star). *In any upward planar 3-slope drawing of the digraph G_{\leftrightarrow} (see Fig. 4), ...*

- *the edges e_1 and e_2 are parallel and have the same arbitrary length $\ell > 0$,*
- *all other edges are oriented as in Fig. 4 up to mirroring along a diagonal axis, and*
- *all other vertical and horizontal edges have the same lengths, as well as all diagonal edges.*

With this construction kit of useful (sub)graphs in hand, we build a bigger graph whose upward planar drawings represent the satisfying truth assignments for F . The **high-level construction** is depicted in Fig. 5b. We construct, for each variable x_i , an individual digraph – the *variable gadget* for x_i . Similarly, for each clause c_j , there is an individual digraph – the *clause gadget* for c_j . All gadgets mainly consist of chains of G_{\square} s. For a drawing, this enforces a rigid frame structure built from unit squares. We glue all variable gadgets together in a row and connect variable and clause gadgets by *edge gadgets* such that the composite graph remains upward outerplanar (see Fig. 5b) and all G_{\square} s are drawn as unit squares.

A **variable gadget** is depicted in Fig. 6. Its base structure is the (violet) **frame** composed of chains of unit squares. The core element is the (red) **central chain** of unit squares (with a few **side-arms**), which has one degree of flexibility, namely, moving as a whole to the left or to the right without leaving the **frame structure** of the gadget. It looks and behaves a bit like


6 Upward Planar Drawings with Three Slopes



■ **Figure 6** A variable gadget, contained in two positive and one negative clauses, being set to false.

a pipe cleaning brush that is stuck inside the **frame** but can be moved a bit back and forth. Hence, we also call it a *brush*. It is connected via a G_{\leftrightarrow} to the **brush** of the previous variable gadget (see Fig. 6a/d) and the first **brush** is connected to the **frame** via a G_{\leftrightarrow} (see the left of Fig. 5b). This allows only a horizontal shift of the **brushes**, but no vertical movement relative to its anchor point at the **frame structure**. Note that the horizontal position in any variable gadget is independent of those in all other gadgets. If the **brush** is positioned close to the very left (right), the corresponding variable is set to false (true).

For each occurrence of a variable in a positive clause, we have a construction as depicted in Fig. 6b. There, a long chain of (green) G_{\square} s – from now on called *bolt* – is attached to the **frame structure** via two G_{\leftrightarrow} s, which allow only a vertical, but no horizontal shift. The *bolt* has on its left side an **arm**, which can only be placed in one of two **pockets of the frame**. It can always be placed in the **upper pocket**, which pushes the *bolt* outwards with respect to the variable gadget (into an edge and then a clause gadget). It can only be placed in the **lower pocket** if the **brush** is shifted close to the very right (i.e. set to true) – then the *bolt*



■ **Figure 7** Positive clause gadget (negative clause gadget is mirrored vertically) in 8 configurations.

can “fall” into a cove of the **brush**. For each occurrence of a variable in a negative clause, we have the same construction, but upside-down, such that the **bolt** can be pulled into the variable gadget only if the **brush** is shifted close to the very left (i.e. set to false).

Note that, to maintain outerplanarity of the whole construction, the **frame structure** is not contiguous, but connected by G_{\leftrightarrow} s and the **arms** of the **bolts**. Hence, the **frame structure** decomposes into many components that have fixed relative horizontal positions and their unit squares have the same side lengths. However, the components can shift up and down relative to each other. To keep this vertical shift small enough not to affect the correct functioning of our reduction, we use, for each such component, the construction depicted in Fig. 6c. The chain of **brushes** has no flexibility in vertical direction and serves as a base ground for an “anchor” of the **frame structure**. The **frame structure** can move less than one unit up or down unless it violates planarity. If the **frame structure** would be shifted up enough to be completely above the **brush**, it would get in conflict with the adjacent **bolt**.

An **edge gadget** consists of only three straight chains – two are **frame segments**, one is a **bolt** in the middle. Their purpose is to synchronize the distance of the clause gadgets to the variable gadgets and to provide these chains of unit squares for the clause gadgets. Several edge gadgets are depicted on yellow background color in Fig. 5b.

A **clause gadget** for a positive clause is depicted in Fig. 7. Within a **frame**, which is connected at six points to the **frames** of three edge gadgets, there is a horizontal (**orange**) **bar**, which is attached via two G_{\leftrightarrow} s to the **frame** – one G_{\leftrightarrow} allows a horizontal, the other allows a vertical shift. It resembles a crane that can move up and extend its arm, while it holds the horizontal **bar** on a vertical (**orange**) **rope**. The three **bolts** from the corresponding variable gadgets reach into the clause gadget. The lengths of these **bolts** is chosen such that, if they are pushed out of their variable gadget and into the clause gadget, they only

slightly fit inside the gadget. Depending on whether each of the **bolts** is pushed into the clause gadget or pulled out of it, we have eight possible configurations (with sufficiently small vertical slack). They represent the eight possible truth assignments to a clause. In Fig. 7, we illustrate that in each configuration, we can accommodate the horizontal **bar** in an upward planar 3-slope drawing of the clause gadget – except for the case when all three **bolts** push into the clause gadget, which represents the truth assignment **false** to all contained variables. A negative clause gadget uses the same construction, but mirrored vertically. There, three **bolts** pushing into the clause gadget means the contained variables are all set to **true**.

We conclude with our main theorem.

► **Theorem 3.2** (*). *Deciding whether a directed outerplanar graph admits an upward planar 3-slope drawing is NP-hard, even when an upward outerplanar embedding is given.*

In the future, we plan to generalize Theorem 3.2 to $k \geq 3$ slopes. In the fixed embedding setting, the main idea is to simply add $2(k-3)$ dummy leaves to each vertex that automatically occupy the additional slopes.

Acknowledgments. We would like to thank the anonymous reviewers for their helpful comments.

References

- 1 Michael A. Bekos, Emilio Di Giacomo, Walter Didimo, Giuseppe Liotta, and Fabrizio Montecchiani. Universal Slope Sets for Upward Planar Drawings. In Therese Biedl and Andreas Kerren, editors, *Graph Drawing and Network Visualization*, pages 77–91. Springer International Publishing, 2018. doi:10.1007/978-3-030-04414-5_6.
- 2 Paola Bertolazzi, Giuseppe Di Battista, Giuseppe Liotta, and Carlo Mannino. Upward drawings of triconnected digraphs. *Algorithmica*, 12(6):476–497, 1994. doi:10.1007/BF01188716.
- 3 Paola Bertolazzi, Giuseppe Di Battista, Carlo Mannino, and Roberto Tamassia. Optimal Upward Planarity Testing of Single-Source Digraphs. *SIAM Journal on Computing*, 27(1):132–169, 1998. doi:10.1137/S0097539794279626.
- 4 Guido Brückner, Nadine Davina Krisam, and Tamara Mchedlidze. Level-planar drawings with few slopes. In Daniel Archambault and Csaba D. Tóth, editors, *Graph Drawing and Network Visualization*, pages 559–572, 2019. doi:10.1007/978-3-030-35802-0_42.
- 5 Hubert Chan. A Parameterized Algorithm for Upward Planarity Testing. In Susanne Albers and Tomasz Radzik, editors, *Algorithms – ESA 2004*, pages 157–168. Springer, 2004. doi:10.1007/978-3-540-30140-0_16.
- 6 Jurek Czyzowicz. Lattice diagrams with few slopes. *Journal of Combinatorial Theory, Series A*, 56(1):96–108, 1991. doi:10.1016/0097-3165(91)90025-C.
- 7 Jurek Czyzowicz, Andrzej Pelc, and Ivan Rival. Drawing orders with few slopes. *Discrete Mathematics*, 82(3):233–250, 1990. doi:10.1016/0012-365X(90)90201-R.
- 8 Mark de Berg and Amirali Khosravi. Optimal binary space partitions for segments in the plane. *International Journal of Computational Geometry & Applications*, 22(3):187–206, 2012. doi:10.1142/s0218195912500045.
- 9 Giuseppe Di Battista, Peter Eades, Roberto Tamassia, and Ioannis G. Tollis. *Graph Drawing: Algorithms for the Visualization of Graphs*. Prentice-Hall, 1999.
- 10 Giuseppe Di Battista and Roberto Tamassia. Algorithms for plane representations of acyclic digraphs. *Theoretical Computer Science*, 61(2):175–198, 1988. doi:10.1016/0304-3975(88)90123-5.
- 11 Emilio Di Giacomo, Giuseppe Liotta, and Fabrizio Montecchiani. Drawing Outer 1-planar Graphs with Few Slopes. In Christian Duncan and Antonios Symvonis, editors, *Graph Drawing*, pages 174–185. Springer, 2014. doi:10.1007/978-3-662-45803-7_15.


- 12 Emilio Di Giacomo, Giuseppe Liotta, and Fabrizio Montecchiani. 1-Bend Upward Planar Drawings of SP-Digraphs. In Yifan Hu and Martin Nöllenburg, editors, *Graph Drawing and Network Visualization*, pages 123–130. Springer International Publishing, 2016. doi:10.1007/978-3-319-50106-2_10.
- 13 Emilio Di Giacomo, Giuseppe Liotta, and Fabrizio Montecchiani. Drawing subcubic planar graphs with four slopes and optimal angular resolution. *Theoretical Computer Science*, 714:51–73, 2018. doi:10.1016/j.tcs.2017.12.004.
- 14 Walter Didimo, Francesco Giordano, and Giuseppe Liotta. Upward Spirality and Upward Planarity Testing. *SIAM Journal on Discrete Mathematics*, 23(4):1842–1899, 2010. doi:10.1137/070696854.
- 15 Vida Dujmović, David Eppstein, Matthew Suderman, and David R. Wood. Drawings of planar graphs with few slopes and segments. *Computational Geometry*, 38(3):194–212, 2007. doi:10.1016/j.comgeo.2006.09.002.
- 16 Vida Dujmović, Matthew Suderman, and David R. Wood. Graph drawings with few slopes. *Computational Geometry*, 38(3):181–193, 2007. doi:10.1016/j.comgeo.2006.08.002.
- 17 Ashim Garg and Roberto Tamassia. On the Computational Complexity of Upward and Rectilinear Planarity Testing. *SIAM Journal on Computing*, 31(2):601–625, 2001. doi:10.1137/S0097539794277123.
- 18 Patrick Healy and Karol Lynch. Two fixed-parameter tractable algorithms for testing upward planarity. *International Journal of Foundations of Computer Science*, 17(05):1095–1114, 2006. doi:10.1142/S0129054106004285.
- 19 Udo Hoffmann. On the Complexity of the Planar Slope Number Problem. *Journal of Graph Algorithms and Applications*, 21(2):183–193, 2017. doi:10.7155/jgaa.00411.
- 20 Vít Jelínek, Eva Jelínková, Jan Kratochvíl, Bernard Lidický, Marek Tesař, and Tomáš Vyskočil. The Planar Slope Number of Planar Partial 3-Trees of Bounded Degree. *Graphs and Combinatorics*, 29(4):981–1005, 2013. doi:10.1007/s00373-012-1157-z.
- 21 Balázs Keszegh, János Pach, and Dömötör Pálvölgyi. Drawing Planar Graphs of Bounded Degree with Few Slopes. *SIAM Journal on Discrete Mathematics*, 27(2):1171–1183, 2013. doi:10.1137/100815001.
- 22 Philipp Kindermann, Wouter Meulemans, and André Schulz. Experimental analysis of the accessibility of drawings with few segments. *Journal of Graph Algorithms and Applications*, 22(3):501–518, 2018. doi:10.7155/jgaa.00474.
- 23 Jonathan Klawitter and Tamara Mchedlidze. Upward planar drawings with two slopes, 2021. In preparation.
- 24 Kolja Knauer, Piotr Micek, and Bartosz Walczak. Outerplanar graph drawings with few slopes. *Computational Geometry*, 47(5):614–624, 2014. doi:doi.org/10.1016/j.comgeo.2014.01.003.
- 25 Rebecca Kraus. Level-außenplanare Zeichnungen mit wenigen Steigungen, 2020. Bachelor Thesis, University of Würzburg.
- 26 William Lenhart, Giuseppe Liotta, Debajyoti Mondal, and Rahnuma Islam Nishat. Planar and Plane Slope Number of Partial 2-Trees. In Stephen Wismath and Alexander Wolff, editors, *Graph Drawing*, pages 412–423. Springer, 2013. doi:10.1007/978-3-319-03841-4_36.
- 27 Padmini Mukkamala and Dömötör Pálvölgyi. Drawing Cubic Graphs with the Four Basic Slopes. In Marc van Kreveld and Bettina Speckmann, editors, *Graph Drawing*, pages 254–265. Springer, 2012. doi:10.1007/978-3-642-25878-7_25.
- 28 Padmini Mukkamala and Mario Szegedy. Geometric representation of cubic graphs with four directions. *Computational Geometry*, 42(9):842–851, 2009. doi:10.1016/j.comgeo.2009.01.005.
- 29 Martin Nöllenburg. Automated Drawing of Metro Maps, 2010. Diploma Thesis, University of Karlsruhe (TH).

- 30 Martin Nöllenburg and Alexander Wolff. Drawing and labeling high-quality metro maps by mixed-integer programming. *IEEE Transactions on Visualization and Computer Graphics*, 17(5):626–641, 2011. doi:10.1109/TVCG.2010.81.
- 31 János Pach and Dömötör Pálvölgyi. Bounded-degree graphs can have arbitrarily large slope numbers. *Electronic Journal of Combinatorics*, 13(1):N1, 2006.
- 32 Achilleas Papakostas. Upward planarity testing of outerplanar dags. In Roberto Tamassia and Ioannis G. Tollis, editors, *Graph Drawing*, pages 298–306. Springer, 1995. doi:10.1007/3-540-58950-3_385.
- 33 G. A. Wade and J.-H. Chu. Drawability of Complete Graphs Using a Minimal Slope Set. *The Computer Journal*, 37(2):139–142, 1994. doi:10.1093/comjnl/37.2.139.

Appendix

► **Theorem 2.1.** *Under a fixed bimodal embedding, a directed tree T admits an upward planar 3-slope drawing if and only if it admits a consistent 3-slope assignment. Moreover, this can be tested in linear time and, in the affirmative, a drawing can be constructed.*

Proof. As an upward planar 3-slope drawing induces a consistent 3-slope assignment, the “only if”-direction is clear. We show that also the converse direction holds, that is, if the given embedding admits a consistent 3-slope assignment, then we can construct an upward planar 3-slope drawing of T . First, we pick any edge $e = (u, v)$ of T and draw it with its assigned slope and with length 1 if the slope of e is \uparrow or \rightarrow , and with length $\sqrt{2}$ if it is \nearrow . From there we draw any further edge e' with length $(1/4)^{d+1}x$ where d is the distance from e' to e and $x = 1$ or $x = \sqrt{2}$ (again depending on the slope); see Fig. 2c for an example. Note that in the resulting drawing the vertices lie on a grid. The consistent 3-slope assignment ensures that the embedding is maintained. Since $\sum_{i=1}^{\infty} (1/4)^i = 1/3$, the chosen edge lengths ensure that no two edges overlap: let uv be an edge with length c , then any subtree “growing” away from uv at u is contained in a box with side lengths $c \cdot 1/3$ and can thus not overlap with another subtree; see Fig. 8. The argument works recursively. If we only describe the coordinates implicitly, the construction runs in linear time. Since the aspect ratio of the drawing can be exponential, an explicit representation of coordinates may require $\Theta(n)$ bits per coordinate. Therefore, explicit coordinates can be computed in $\mathcal{O}(n^2)$ time.



■ **Figure 8** By shrinking edge lengths by a factor of 0.25, different subtrees cannot overlap.


It remains to show that we can test whether the embedding admits a consistent 3-slope assignment in linear time. For this, we first associate with each edge a *possible slopes set*, which initially contains all three slopes. We then go through the vertices one by one. When visiting a vertex v , we restrict the sets of possible slopes of each incident edge based on the cyclic edge order around v . For example, if v has two incoming edge e and e' in ccw order, then e may have slope \rightarrow or slope \nearrow and e' may have slope \nearrow or slope \uparrow . If this results in an empty set for an edge, then T does not admit an upward plane 3-slope drawing under the given embedding. We claim that if no slope set is empty after we have processed all vertices, then we have a yes-instance. We prove this by constructing a consistent 3-slope assignment based on the possible slopes sets of each edge.

An edge with possible slopes set of size one gets exactly the remaining slope assigned. An edge uv with possible slopes set of size three is the only outgoing edge of u and the only incoming edge of v . It may therefore get any slope and we simply pick slope \nearrow . Suppose now that the edge uv has a possible slope set of size two. We observe that u has outdegree at most two and v has indegree at most two, and for at least one of them it actually is two. Suppose that this is the case for u and let uv' be the second outgoing edge of u . Further

suppose uv' comes ccw after uv . Then uv has possible slope set $\{\nearrow, \rightarrow\}$ and the slope set of uv' is a subset of $\{\nearrow, \uparrow\}$. Hence, it is save to pick slope \rightarrow for uv with regard to uv' . An analogous argument applies to a potential second incoming edge of v . We conclude that if an edge has a potential slope set of size two then we can always pick the slope that is not \nearrow . It is therefore straightforward to go from the potential slope sets to a consistent 3-slope assignment. ◀

► **Theorem 2.2.** *Given a directed tree T with indegree and outdegree at most three, we can construct an upward planar 3-slope drawing of T in linear time.*

Proof. By Theorem 2.1, we only need to find an embedding that admits a consistent 3-slope assignment. For this, we pick any vertex v of T as starting vertex, order the edges incident to v bimodally and assign them slopes accordingly. We then handle the remaining vertices in order of increasing distance to v in the same way as v . The only difference is that, at each vertex u , exactly one edge has already a slope assigned. It is easy to see that this can be done in linear time. ◀



■ **Figure 4** Upward planar 3-slope drawing of the digraph G_{\leftrightarrow} .

► **Lemma 3.1.** *In any upward planar 3-slope drawing of the digraph G_{\leftrightarrow} (see Fig. 4), ...*

- the edges e_1 and e_2 are parallel and have the same arbitrary length $\ell > 0$,
- all other edges are oriented as in Fig. 4 up to mirroring along a diagonal axis, and
- all other vertical and horizontal edges have the same lengths, as well as all diagonal edges.

Proof. As subgraphs, we have twice a pair of attached G_{\square} s. As observed before, each pair can only be drawn as attached unit squares. Without loss of generality, let the first two squares (containing vertices v_2, v_3, v_5) be drawn vertically above each other as in Fig. 4. Since v_2 has outdegree 3 and e_3 has slope \uparrow , we can argue that the edge e_1 gets slope \rightarrow . If e_1 used \nearrow , the edge e_5 would have no slope to close the triangle (e_3, e_4, e_5) because, at v_5 , the rightmost outgoing slope is only \rightarrow . Thus, e_4 gets \nearrow and the only remaining slope for e_1 is \rightarrow . Note that e_3 is a unit square edge and the adjacent edges e_4 and e_5 use the same slopes as their counterparts in a unit square, giving them the corresponding lengths.

The same argument as for the outgoing edges of v_2 applies to the incoming edges of v_4 . However, assume for contradiction, that the whole block on the right side of Fig. 4 is mirrored diagonally, i.e., v_1 and v_4 are on the same horizontal line and e_2 gets slope \uparrow . Then, v_4, v_1 , and v_2 would be on the same horizontal line. This contradicts v_3 being vertically below its neighbors v_2 and v_4 . Hence, the right block has the same orientation as the left one, and e_2 gets slope \rightarrow and is parallel to e_1 . This implies that the distances between v_2 and v_3 on the one hand, and v_1 and v_4 on the other hand are the same and the G_{\square} s (and their attached triangles) have the size and orientation as in Fig. 4. By the orientation of v_1, v_2, v_3, v_4 , the

edges e_1 and e_2 have an identical length ℓ . Only a value $\ell \leq 0$ would cause a non-planar drawing. ◀

► **Theorem 3.2.** *Deciding whether a directed outerplanar graph admits an upward planar 3-slope drawing is NP-hard, even when an upward outerplanar embedding is given.*

Proof. To show NP-hardness, we use the reduction from PLANAR MONOTONE 3-SAT as described above. Let F be a given PLANAR MONOTONE 3-SAT formula with a rectangular drawing of the incidence graph of F and G_F be the digraph obtained by our reduction. Each gadget in G_F has only polynomial size and we can construct it in polynomial time.

If F is satisfiable, then there is a satisfying assignment T^* of truth values to the variables of F . Draw G_F as illustrated in Fig. 5b, Fig. 6, and Fig. 7 such that the **brush** in a variable gadget has a distance of ε (for some sufficiently small $\varepsilon > 0$) to the **frame** on its left if the corresponding variable is set to **false** in T^* and has a distance of ε to the **frame** on its right if the corresponding variable is set to **true**. This drawing uses only three slopes and is upward planar. Observe that it is also outerplanar. On a chain of unit squares, the vertices on both sides are incident to the outer face. The drawings of the variable gadgets are open from one to the other and the drawing of the rightmost variable gadget is open to the outside. Moreover, the drawings of the positive (negative) clause gadgets are open on their top (bottom) right and on their whole bottom (top) sides, which also gives the “covered” clause gadgets and the outer sides of the variable gadgets access to the outer face.

On the other hand, if there is an upward planar 3-slope drawing of G_F and it resembles the structure in Fig. 5b, we can read a satisfying truth assignment depending on the positions of the **brushes** (for the ones in intermediate position where no **bolt** can use the **lower pocket**, we can use any assignment for this variable).

In the remainder of this proof, we argue that any drawing resembles this structure. Consider the G_{\square} of the **frame structure** which is connected via a G_{\leftrightarrow} to the **brush** of the first variable gadget. Clearly, it is drawn as a unit square – it is our reference unit square having side length 1. Observe that all other unit squares are connected via chains of G_{\square} s or G_{\leftrightarrow} s to this reference unit square. By our observation and Lemma 3.1, they have the same side length. When ignoring the central parallel edges in the G_{\leftrightarrow} s, the drawing decomposes to few rigid components. The first rigid **frame component** contains the reference unit square. The central edges of the incident G_{\leftrightarrow} connecting it to the first **brush** can have only relatively small length as they would hit the “back wall” of, again, the first rigid **frame component** otherwise. Hence, the **brush** is indeed drawn inside the **frame** of the first variable gadget. Moreover, the first rigid **frame component** is connected via a G_{\leftrightarrow} to a **bolt**, which in turn is connected to the next rigid **frame components** of the first variable gadget (see Fig. 6b). Since this **bolt** cannot escape the upper boundary of the clause gadget, which is also part of the first rigid **frame component**, the **arm** is in one of the two **pockets**. Then, the start point of the next rigid **frame structure** cannot be above the **arm** of the **brush** in Fig. 6c and the construction depicted there keeps the vertical slack of this rigid **frame component** within the range $(-1, 1)$. This argument inductively propagates for all following rigid components. Hence, we have vertical slack of less than 1 for the **frame structures** and for the **arms** in the **pockets**. In the configuration **false-false-false** of the clause gadget, this would give us $2 - \varepsilon$ vertical gain if two **frames** move away from each other and $1 - \varepsilon$ gain if the **arm** is close to the bottom of its **pocket**. However, we would require a shift of the horizontal **bar** of more than 3 to be below or above the bulge of the left or the right **bolt**. Hence, this configuration is not drawable and all other positions of the horizontal **bar** correspond to one of the 7 other, satisfying configurations of Fig. 7. ◀