

On the Brauer groups of fibrations II

Yanshuai Qin

January 3, 2022

Contents

1	Introduction	1
1.1	Main results	2
1.2	The idea of the proof of Theorem 1.2	3
1.3	Notation and Terminology	4
2	Preliminary	5
3	Proof of Theorem 1.2	9
4	Applications	18
4.1	Reduction of Artin's question	18
4.2	Examples	19

1 Introduction

Let C be the spectrum of the ring of integers in a number field or a smooth projective geometrically connected curve over a finite field with function field K . Let \mathcal{X} be a 2-dimensional regular scheme and $\pi : \mathcal{X} \rightarrow C$ be a proper flat morphism such that the generic fiber X of π is smooth and geometrically connected over K . Artin and Grothendieck (cf. [Gro3, § 4] or [Ulm, Prop. 5.3]) proved that there is an isomorphism

$$\text{III}(\text{Pic}_{X/K}^0) \cong \text{Br}(\mathcal{X})$$

up to finite groups. Assuming the finiteness of $\text{III}(\text{Pic}_{X/K}^0)$ or $\text{Br}(\mathcal{X})$, Milne [Mil4], Gonzales-Aviles [Goa] and Geisser [Gei1] gave a precise relation between their orders. In [Qin1], the author proved a relation between Tate-Shafarevich groups and geometric Brauer groups for fibrations over arbitrary finitely generated fields. The aim of this article is to generalize the original result of Artin and Grothendieck to fibrations of arbitrary relative dimensions over the spectrum of the ring of integers in a number field.

For any noetherian scheme X , the *cohomological Brauer group*

$$\mathrm{Br}(X) := H^2(X, \mathbb{G}_m)_{\mathrm{tor}}$$

is defined to be the torsion part of the étale cohomology group $H^2(X, \mathbb{G}_m)$.

Tate conjecture for divisors

Let us first recall the Tate conjecture for divisors over finitely generated fields.

Conjecture 1.1. (*Conjecture $T^1(X, \ell)$*). *Let X be a projective and smooth variety over a finitely generated field k of characteristic $p \geq 0$, and $\ell \neq p$ be a prime number. Then the cycle class map*

$$\mathrm{Pic}(X) \otimes_{\mathbb{Z}} \mathbb{Q}_{\ell} \longrightarrow H^2(X_{k^s}, \mathbb{Q}_{\ell}(1))^{G_k}$$

is surjective.

It is well-known that $T^1(X, \ell)$ is equivalent to the finiteness of $\mathrm{Br}(X_{k^s})^{G_k}(\ell)$ (cf. Proposition 2.1).

Artin's question

Artin conjectured that $\mathrm{Br}(\mathcal{X})$ is finite for any proper scheme \mathcal{X} over $\mathrm{Spec} \mathbb{Z}$. For a smooth projective variety \mathcal{X} over a finite field, it is well-known that the finiteness of $\mathrm{Br}(\mathcal{X})$ is equivalent to Tate conjecture for divisors on \mathcal{X} (cf. [Tat2]). For a proper flat regular integral scheme \mathcal{X} over $\mathrm{Spec} \mathbb{Z}$, taking $C = \mathrm{Spec} \Gamma(\mathcal{X}, \mathcal{O}_{\mathcal{X}})$, there is a proper flat morphism (the Stein factorization) $\pi : \mathcal{X} \rightarrow C$ with a generic fiber geometrically connected over $K = K(C)$. We will show that our main theorem can imply that the finiteness of $\mathrm{Br}(\mathcal{X})(\ell)$ is equivalent to the finiteness of $\mathrm{III}(\mathrm{Pic}_{X/K}^0)(\ell)$ and $T^1(X, \ell)$ for the generic fiber X of π . In [SZ], Skorobogatov and Zarkhin conjectured that $\mathrm{Br}(X_{k^s})^{G_k}$ is finite for a smooth projective variety X over a finitely generated field k , and they proved the finiteness of $\mathrm{Br}(X_{k^s})^{G_k}$ for abelian varieties and K3 surfaces. Assuming the finiteness of $\mathrm{Br}(X_{K^s})^{G_K}$ and $\mathrm{III}(\mathrm{Pic}_{X/K}^0)$, our main theorem will imply the finiteness of $\mathrm{Br}(\mathcal{X})$.

1.1 Main results

For two abelian groups M and N , we say that they are *almost isomorphic* if there exists a subquotient M_1/M_0 of M (resp. N_1/N_0 of N) such that M_0 (resp. N_0) and M/M_1 (resp. N/N_1) are finite and $M_1/M_0 \cong N_1/N_0$. Let $f : M \rightarrow N$ be a homomorphism with a finite cokernel and a kernel almost isomorphic to an abelian group H , in this situation, we say that the sequence $0 \rightarrow H \rightarrow M \rightarrow N \rightarrow 0$ is exact up to finite groups.

Theorem 1.2. *Let $\pi : \mathcal{X} \rightarrow C$ be a proper flat morphism, where C is $\mathrm{Spec} \mathcal{O}_K$ for some number field K . Assume that \mathcal{X} is regular and the generic fiber X of π is projective and*

geometrically connected over K . Then there is an exact sequence up to finite groups

$$0 \longrightarrow \text{III}(\text{Pic}_{X/K}^0) \longrightarrow \text{Br}(\mathcal{X}) \longrightarrow \text{Br}(X_{\bar{K}})^{G_K} \longrightarrow 0.$$

For arithmetic schemes of dimension ≥ 3 , the above question was first studied by Tankeev (cf. [Tan1, Tan2, Tan3]), and he proved the above result for some special arithmetic schemes. Geisser ([Gei2]) first studied fibrations of arbitrary relative dimensions over finite fields by using étale motivic cohomology theory. In [Qin1], we proved a similar result for fibrations over arbitrary finitely generated fields.

Corollary 1.3. *$\text{Br}(\mathcal{X})(\ell)$ is finite if and only if $\text{III}(\text{Pic}_{X/K}^0)(\ell)$ is finite and $T^1(X, \ell)$ holds.*

The above statement for fibrations over finite fields was proved by Geisser [Gei2]. For a smooth projective geometrically connected variety X over a finitely generated field k of characteristic 0, Orr and Skorobogatov [OS, Thm. 5.1] proved that $\text{Br}(X_{k^s})^{G_k}$ is finite under the assumption of the integral Mumford-Tate conjecture. Thus, Theorem 1.2 implies the following result about Artin's question.

Corollary 1.4. *Assuming the integral Mumford-Tate conjecture and the Tate-Shafarevich conjecture, then $\text{Br}(\mathcal{X})$ is finite for all proper regular schemes over $\text{Spec } \mathbb{Z}$.*

By a theorem of André [And] and a result of Ambrosi [Amb, Cor. 1.6.2.1], the finiteness of $\text{Br}(X_{\bar{k}})^{G_k}$ has been reduced to smooth projective surfaces over k . Thus, Theorem 1.2 can imply the following :

Theorem 1.5. *Assuming that $\text{Br}(\mathcal{X})$ is finite for all 3-dimensional regular proper flat schemes over \mathbb{Z} , then $\text{Br}(\mathcal{X})$ is finite for all regular proper flat schemes over \mathbb{Z} .*

Remark 1.6. The finiteness of Brauer groups for smooth proper varieties over finite fields has been reduced to smooth projective surfaces over finite fields by the work of de Jong [deJ2] and Morrow [Mor, Thm. 4.3].

1.2 The idea of the proof of Theorem 1.2

The left exactness of the sequences can be proved by a combination of Grothendieck's argument in [Gro3, §4] and a pull-back trick developed by Colliot-Thélène and Skorobogatov in [CTS1]. The trick part is the finiteness of the cokernel of the natural map

$$\text{Br}(\mathcal{X}) \longrightarrow \text{Br}(X_{\bar{K}})^{G_K}.$$

Colliot-Thélène [CT, Conj. 1.5(d)] conjectured that the cokernel of the natural map

$$\prod_v A_0(X_{K_v}) \longrightarrow \text{Hom}(\text{Br}(X)/\text{Br}(K), \mathbb{Q}/\mathbb{Z})$$

induced by the Brauer-Manin pairing is a finite group that can be identified with the group

$$\text{Hom}(\text{Br}'(\mathcal{X})/\text{Br}(\mathcal{O}_K), \mathbb{Q}/\mathbb{Z}),$$

where $\text{Br}'(\mathcal{X}) \subset \text{Br}(\mathcal{X})$ is the subgroup of elements vanishing on $X(K_v)$ for all real places of K . The dual statement of the conjecture predicts that the natural map

$$\text{Br}(X)/(\text{Br}(\mathcal{X}) + \text{Br}(K)) \longrightarrow \prod_{v \in C^\circ} \text{Br}(X_{K_v})/(\text{Br}(\mathcal{X}_{\mathcal{O}_v}) + \text{Br}(K_v))$$

has a finite kernel, where C° denotes the set of finite places. In conjunction with a theorem of Colliot-Thélène and Skorobogatov [CTS1] which said that the natural map $\text{Br}(X) \longrightarrow \text{Br}(X_{\bar{K}})^{G_K}$ has a finite cokernel, it can imply that the natural map

$$\text{Br}(X_{\bar{K}})^{G_K}/\text{Br}(\mathcal{X}) \longrightarrow \prod_v \text{Br}(X_{\bar{K}_v})^{G_{K_v}}/\text{Br}(\mathcal{X}_{\mathcal{O}_v})$$

has a finite kernel. By a result of the author [Qin3, Cor. 1.9], the group $\text{Br}(X_{\bar{K}_v})^{G_{K_v}}/\text{Br}(\mathcal{X}_{\mathcal{O}_v})$ is finite for all finite places v . This will imply that the divisible part of $\text{Br}(X_{\bar{K}})^{G_K}/\text{Br}(\mathcal{X})$ is trivial. It follows that $(\text{Br}(X_{\bar{K}})^{G_K}/\text{Br}(\mathcal{X}))(\ell)$ is finite for all prime ℓ . By using the Fontaine-Laffaille-Messing theory (cf. [BM, §3]) for $\mathcal{X}_{\mathcal{O}_v}$ with good reduction, we will show that $(\text{Br}(X_{\bar{K}_v})^{G_{K_v}}/\text{Br}(\mathcal{X}_{\mathcal{O}_v}))(\ell)$ is trivial for all v when ℓ is sufficiently large. This will imply that $(\text{Br}(X_{\bar{K}})^{G_K}/\text{Br}(\mathcal{X}))(\ell)$ is trivial for $\ell \gg 0$.

1.3 Notation and Terminology

Fields

By a *finitely generated field*, we mean a field which is finitely generated over a prime field. For any field k , denote by k^s the separable closure. Denote by $G_k = \text{Gal}(k^s/k)$ the absolute Galois group of k .

Henselization

Let R be a noetherian local ring, denote by R^h (resp. R^{sh}) the henselization (resp. strict henselization) of R at the maximal ideal. If R is a domain, denote by K^h (resp. K^{sh}) the fraction field of R^h (resp. R^{sh}).

Varieties

By a *variety* over a field k , we mean a scheme which is separated and of finite type over k . For a smooth proper geometrically connected variety X over a field k , we use $\text{Pic}_{X/k}^0$ to denote the identity component of the Picard scheme $\text{Pic}_{X/k}$.

Cohomology

The default sheaves and cohomology over schemes are with respect to the small étale site. So H^i is the abbreviation of $H_{\text{ét}}^i$.

Brauer groups

For any noetherian scheme X , denote the *cohomological Brauer group*

$$\mathrm{Br}(X) := H^2(X, \mathbb{G}_m)_{\mathrm{tor}}.$$

Tate-Shafarevich group

For an abelian variety A defined over a global field K , denote the *Tate–Shafarevich group*

$$\mathrm{III}(A) := \mathrm{Ker}(H^1(K, A) \longrightarrow \prod_v H^1(K_v, A)),$$

where K_v denotes the completion of K at a place v . Note that if v is a finite place, the natural map $H^1(K_v^h, A) \rightarrow H^1(K_v, A)$ is an isomorphism (cf. [Mil1, Chap. II, Rem. 3.10]), where K_v^h denotes the fraction field of the henselian local ring at v .

Abelian group

For any abelian group M , integer m and prime ℓ , we set

$$M[m] = \{x \in M \mid mx = 0\}, \quad M_{\mathrm{tor}} = \bigcup_{m \geq 1} M[m], \quad M(\ell) = \bigcup_{n \geq 1} M[\ell^n],$$

$$M(\mathrm{non}\text{-}\ell) = \bigcup_{m \geq 1, \ell \nmid m} M[m], \quad T_\ell M = \mathrm{Hom}_{\mathbb{Z}}(\mathbb{Q}_\ell/\mathbb{Z}_\ell, M) = \varprojlim_n M[\ell^n], \quad V_\ell M = T_\ell(M) \otimes_{\mathbb{Z}_\ell} \mathbb{Q}_\ell.$$

A torsion abelian group M is *of cofinite type* if $M[\ell]$ is finite for all prime ℓ .

For two abelian groups M and N , we say that they are *almost isomorphic* if there exist filtrations $M_0 \subseteq M_1 \subseteq M$ and $N_0 \subseteq N_1 \subseteq N$ such that M_0 (resp. N_0) and M/M_1 (resp. N/N_1) are finite and M_1/M_0 is isomorphic to N_1/N_0 .

Acknowledgements:

I would like to thank my advisor Xinyi Yuan for suggesting this question to me and for telling me the pull-back trick.

2 Preliminary

In this section, we will recall some facts which already appeared in the literature.

Basic exact sequences

Proposition 2.1. *Let X be a smooth projective geometrically connected variety over a field k . Define $\mathrm{NS}(X)$ to be the image of the map $\mathrm{Pic}(X) \rightarrow \mathrm{NS}(X_{\bar{k}})$. Let ℓ be a prime different from $\mathrm{char}(k)$, then we have*

(a) For any $\ell \neq \text{char}(k)$, the exact sequence of G_k -representations

$$0 \longrightarrow \text{NS}(X_{k^s}) \otimes_{\mathbb{Z}} \mathbb{Q}_{\ell} \longrightarrow H^2(X_{k^s}, \mathbb{Q}_{\ell}(1)) \longrightarrow V_{\ell} \text{Br}(X_{k^s}) \longrightarrow 0$$

is split. Taking G_k -invariant, there is an exact sequence

$$0 \longrightarrow \text{NS}(X) \otimes_{\mathbb{Z}} \mathbb{Q}_{\ell} \longrightarrow H^2(X_{k^s}, \mathbb{Q}_{\ell}(1))^{G_k} \longrightarrow V_{\ell} \text{Br}(X_{k^s})^{G_k} \longrightarrow 0.$$

(b) For sufficiently large ℓ , the exact sequence of G_k -modules

$$0 \longrightarrow \text{NS}(X_{k^s})/\ell^n \longrightarrow H^2(X_{k^s}, \mathbb{Z}/\ell^n(1)) \longrightarrow \text{Br}(X_{k^s})[\ell^n] \longrightarrow 0$$

is split for any $n \geq 1$. Moreover, for all but finitely many ℓ , there is an exact sequence

$$0 \longrightarrow \text{NS}(X) \otimes_{\mathbb{Z}} \mathbb{Q}_{\ell}/\mathbb{Z}_{\ell} \longrightarrow H^2(X_{k^s}, \mathbb{Q}_{\ell}/\mathbb{Z}_{\ell}(1))^{G_k} \longrightarrow \text{Br}(X_{k^s})^{G_k}(\ell) \longrightarrow 0.$$

Proof. We refer the proof of (a) and the first claim of (b) to [Qin2, Prop. 2.1]. Taking G_k -invariants, we get an exact sequence for all but finitely many ℓ

$$0 \longrightarrow (\text{NS}(X_{k^s})/\ell^n)^{G_k} \longrightarrow H^2(X_{k^s}, \mathbb{Z}/\ell^n(1))^{G_k} \longrightarrow \text{Br}(X_{k^s})^{G_k}[\ell^n] \longrightarrow 0.$$

Taking direct limit, we get

$$0 \longrightarrow (\text{NS}(X_{k^s}) \otimes_{\mathbb{Z}} \mathbb{Q}_{\ell}/\mathbb{Z}_{\ell})^{G_k} \longrightarrow H^2(X_{k^s}, \mathbb{Q}_{\ell}/\mathbb{Z}_{\ell}(1))^{G_k} \longrightarrow \text{Br}(X_{k^s})^{G_k}(\ell) \longrightarrow 0.$$

To prove the second claim of (b), it suffices to show that the natural map

$$\text{NS}(X) \otimes_{\mathbb{Z}} \mathbb{Q}_{\ell}/\mathbb{Z}_{\ell} \longrightarrow (\text{NS}(X_{k^s}) \otimes_{\mathbb{Z}} \mathbb{Q}_{\ell}/\mathbb{Z}_{\ell})^{G_k}$$

is an isomorphism for all but finitely many ℓ . Consider the exact sequence

$$0 \longrightarrow \text{NS}(X) \otimes_{\mathbb{Z}} \mathbb{Z}_{\ell} \longrightarrow \text{NS}(X_{k^s}) \otimes_{\mathbb{Z}} \mathbb{Z}_{\ell} \longrightarrow \text{NS}(X_{k^s})/\text{NS}(X) \otimes_{\mathbb{Z}} \mathbb{Z}_{\ell} \longrightarrow 0.$$

For ℓ sufficiently large, the last group will be a free \mathbb{Z}_{ℓ} -module. Thus, it is split for all but finitely many ℓ . So tensoring $\mathbb{Q}_{\ell}/\mathbb{Z}_{\ell}$, the sequence is still exact. This proves injectivity. Since the G_k action on $\text{NS}(X_{k^s})$ factors through a finite group G , thus, for ℓ prime to $|G|$, any element in the $(\text{NS}(X_{k^s}) \otimes_{\mathbb{Z}} \mathbb{Q}_{\ell}/\mathbb{Z}_{\ell})^{G_k}$ can be written as $\sum_{g \in G} gx$. This implies that the natural map

$$\text{NS}(X_{k^s})^{G_k} \otimes_{\mathbb{Z}} \mathbb{Q}_{\ell}/\mathbb{Z}_{\ell} \longrightarrow (\text{NS}(X_{k^s}) \otimes_{\mathbb{Z}} \mathbb{Q}_{\ell}/\mathbb{Z}_{\ell})^{G_k}$$

is surjective. Thus, to prove the surjectivity, it suffices to show that

$$\text{NS}(X) \otimes_{\mathbb{Z}} \mathbb{Q}_{\ell} \longrightarrow \text{NS}(X_{k^s})^{G_k} \otimes_{\mathbb{Z}} \mathbb{Q}_{\ell}$$

is surjective. This was proved in [Yua, §2.2]. \square

Cofiniteness

Proposition 2.2. *Let \mathcal{X} be an integral regular scheme flat and of finite type over $\text{Spec } \mathbb{Z}$. Then $\text{Br}(\mathcal{X})$ is of cofinite type.*

Proof. Let ℓ be a prime. By shrinking \mathcal{X} , we may assume that \mathcal{X} is a $\mathbb{Z}[1/\ell]$ -scheme. Let $\pi : \mathcal{X} \rightarrow S = \text{Spec } \mathbb{Z}[1/\ell]$ the structure morphism. By the Kummer exact sequence, there is a surjective map

$$H^2(\mathcal{X}, \mu_\ell) \rightarrow \text{Br}(\mathcal{X})[\ell].$$

Thus, it suffices to show that $H^2(\mathcal{X}, \mu_\ell)$ is finite. By the Leray spectral sequence

$$H^p(S, R^q \pi_* \mu_\ell) \Rightarrow H^{p+q}(\mathcal{X}, \mu_\ell),$$

it suffices to show that $H^p(S, R^q \pi_* \mu_\ell)$ is finite. By [Fu, Thm. 9.5.1], $R^q \pi_* \mu_\ell$ is a constructible ℓ -torsion sheaf. By [Mil1, Chap. II, Thm. 2.13], $H^p(S, R^q \pi_* \mu_\ell)$ is finite. \square

Proposition 2.3. *Let X be a smooth projective geometrically connected variety over a number field K . Define*

$$\text{III}(\text{Pic}_{X/K}) := \text{Ker}(H^1(K, \text{Pic}_{X/K}) \rightarrow \prod_v H^1(K_v^h, \text{Pic}_{X/K})).$$

Then the natural map

$$\text{III}(\text{Pic}_{X/K}^0) \rightarrow \text{III}(\text{Pic}_{X/K})$$

has a finite kernel and a cokernel of finite exponent.

Proof. Since $\text{III}(\text{Pic}_{X/K}^0)$ is of cofinite type (cf. [Mil1, Chap. II, Cor. 5.3 and Lem. 5.5]), it suffices to show that the map has a kernel and a cokernel of finite exponent. By the exact sequence

$$0 \rightarrow \text{Pic}_{X/K}^0 \rightarrow \text{Pic}_{X/K} \rightarrow \text{NS}(X_{\bar{K}}) \rightarrow 0$$

we get a commutative diagram with exact rows

$$\begin{array}{ccccccc} H^1(K, \text{Pic}_{X/K}^0) & \longrightarrow & H^1(K, \text{Pic}_{X/K}) & \longrightarrow & H^1(K, \text{NS}(X_{\bar{K}})) & & \\ & & \downarrow a & & \downarrow b & & \downarrow c \\ \prod_v H^0(K_v^h, \text{NS}(X_{\bar{K}})) & \xrightarrow{\partial} & \prod_v H^1(K_v^h, \text{Pic}_{X/K}^0) & \longrightarrow & \prod_v H^1(K_v^h, \text{Pic}_{X/K}) & \longrightarrow & \prod_v H^1(K_v^h, \text{NS}(X_{\bar{K}})) \end{array}$$

Let \bar{a} denote the map

$$H^1(K, \text{Pic}_{X/K}^0) \rightarrow \prod_v H^1(K_v^h, \text{Pic}_{X/K}^0) / \text{Im}(\partial)$$

By the Snake Lemma, there is an exact sequence

$$\text{Ker}(\bar{a}) \rightarrow \text{Ker}(b) \rightarrow \text{Ker}(c).$$

We also have

$$0 \longrightarrow \text{Ker}(a) \longrightarrow \text{Ker}(\bar{a}) \longrightarrow \text{Im}(\partial).$$

By definition, $\text{Ker}(a) = \text{III}(\text{Pic}_{X/K}^0)$ and $\text{Ker}(b) = \text{III}(\text{Pic}_{X/K})$. Thus, it suffices to show that $\text{Im}(\partial)$ and $\text{Ker}(c)$ are of finite exponent. Since $\text{NS}(X_{\bar{K}})$ is finitely generated, there exists a finite Galois extension L of K such that $\text{Pic}(X_L)$ maps onto $\text{NS}(X_{\bar{K}})$. By the inflation-restriction exact sequence

$$0 \longrightarrow H^1(\text{Gal}(L/K), \text{NS}(X_{\bar{K}})) \longrightarrow H^1(K, \text{NS}(X_{\bar{K}})) \longrightarrow H^1(L, \text{NS}(X_{\bar{K}}))$$

and the fact that $H^1(\text{Gal}(L/K), \text{NS}(X_{\bar{K}}))$ is killed by $[L : K]$ and $H^1(L, \text{NS}(X_{\bar{K}})) = \text{Hom}(G_L, \text{NS}(X_{\bar{K}})_{\text{tor}})$ is killed by $|\text{NS}(X_{\bar{K}})_{\text{tor}}|$, we have that $H^1(K, \text{NS}(X_{\bar{K}}))$ is also of finite exponent. It follows that $\text{Ker}(c)$ is of finite exponent. To show that $\text{Im}(\partial)$ is of finite exponent, it suffices to show that the map

$$H^0(K_v^h, \text{NS}(X_{\bar{K}})) \xrightarrow{\partial_v} H^1(K_v^h, \text{Pic}_{X/K}^0)$$

has an image killed by $[L : K]$. Let w be a place of L over v . Consider the following commutative diagram with exact rows

$$\begin{array}{ccccc} H^0(K_v^h, \text{Pic}_{X/K}) & \longrightarrow & H^0(K_v^h, \text{NS}(X_{\bar{K}})) & \xrightarrow{\partial_v} & H^1(K_v^h, \text{Pic}_{X/K}^0) \\ \downarrow & & \downarrow & & \downarrow g \\ H^0(L_w^h, \text{Pic}_{X/K}) & \xrightarrow{f} & H^0(L_w^h, \text{NS}(X_{\bar{K}})) & \xrightarrow{\partial_w} & H^1(L_w^h, \text{Pic}_{X/K}^0) \end{array}$$

By the choice of L , f is surjective. Thus, $\partial_w = 0$. So $\text{Im}(\partial_v) \subseteq \text{Ker}(g)$. By the inflation-restriction exact sequence, we have that $\text{Ker}(g) = H^1(\text{Gal}(L_w^h/K_v^h), \text{Pic}_{X/K}^0(L_w^h))$ is killed by $[L_w^h : K_v^h]$. Since L/K is finite Galois, so $[L_w^h : K_v^h] | [L : K]$. This proves that $\text{Im}(\partial_v)$ is killed by $[L : K]$. \square

Local results

Proposition 2.4. *Let $\pi : \mathcal{X} \longrightarrow S = \text{Spec } R$ be a proper flat morphism where R is a henselian DVR with a finite residue field k and a quotient field K . Assume that \mathcal{X} is regular and the generic fiber X of π is smooth projective geometrically connected over K . Let $\ell \neq \text{char}(k)$ be a prime. Then the natural map*

$$\text{Br}(\mathcal{X}_{R^{sh}})^{G_k}(\ell) \longrightarrow \text{Br}(X_{K^s})^{G_K}(\ell)$$

has a kernel and a cokernel of finite exponent and is an isomorphism for all but finitely many ℓ . Moreover, if π is smooth, the map is an isomorphism for all $\ell \neq \text{char}(k)$.

Proof. Cf. [Qin3, Lem. 3.9] and [Qin1, Lem. 3.1(b)]. \square

Proposition 2.5. *Let $\pi : \mathcal{X} \rightarrow S = \text{Spec } R$ be a proper flat morphism, where R is a henselian DVR of characteristic 0 with a finite residue field. Assuming that \mathcal{X} is regular and the generic fiber X is geometrically connected over $K = \text{Frac}(R)$, then the natural map*

$$\text{Br}(\mathcal{X}) \rightarrow \text{Br}(X_{\bar{K}})^{G_K}$$

has finite kernel and a finite cokernel.

Proof. Cf. [Qin3, Cor. 1.9]. □

3 Proof of Theorem 1.2

Lemma 3.1. *Let $\pi : \mathcal{X} \rightarrow S = \text{Spec } R$ be a flat morphism with \mathcal{X} integral and regular, where R is a DVR of characteristic 0 with a fraction field K . Let X denote the generic fiber of K . Then there is an exact sequence*

$$0 \rightarrow \text{Br}(\mathcal{X}) \rightarrow \text{Br}(X) \rightarrow \text{Br}(X_{K^h})/\text{Br}(\mathcal{X}_{R^h}).$$

Proof. Let Y denote the special fiber of π . It suffices to show

$$H_Y^3(\mathcal{X}, \mathbb{G}_m) \cong H_Y^3(\mathcal{X}_{R^h}, \mathbb{G}_m).$$

This follows from the Excision theorem (cf. [Mil3, Chap III, Prop. 1.27 and Cor. 1.28]). □

Remark 3.2. In fact, the exact sequence above still holds if we replace R^h by its completion. This actually follows from purity of Brauer groups. We may assume that Y is of codimension 1 with a generic point y . Then there is an injection $\text{Br}(X)/\text{Br}(\mathcal{X}) \hookrightarrow \text{Br}(K(X)_y^h)/\text{Br}(\mathcal{O}_{\mathcal{X},y}^h)$. Since $\text{char}(K(X)) = 0$, we have $\text{Br}(K(X)_y^h) = \text{Br}(\overline{K(X)_y})$ and $\text{Br}(\mathcal{O}_{\mathcal{X},y}^h) = \text{Br}(\overline{\mathcal{O}_{\mathcal{X},y}})$. It is easy to see that the completion does not change if we replace R by \hat{R} .

Lemma 3.3. *Let $\pi : \mathcal{X} \rightarrow C = \text{Spec } \mathcal{O}_K$ be proper flat morphism with \mathcal{X} regular, where K is a number field. Assuming that the generic fiber X of π is geometrically connected over K , then the natural map*

$$\text{Br}(X)/(\text{Br}(\mathcal{X}) + \text{Br}(K)) \rightarrow \prod_{v \in C^\circ} \text{Br}(X_{K_v^h})/(\text{Br}(\mathcal{X}_{\mathcal{O}_v^h}) + \text{Br}(K_v^h))$$

has a finite kernel and the group $\text{Br}(\mathcal{X}) \cap \overline{\text{Br}(K)}$ is finite, where C° denotes the set of closed point of C , \mathcal{O}_v^h denotes the henselian local ring of C at v and $\overline{\text{Br}(K)}$ denotes the image of the pullback map $\text{Br}(K) \rightarrow \text{Br}(X)$.

Proof. Consider the following commutative diagram

$$\begin{array}{ccccccc}
\mathrm{Br}(K) & \longrightarrow & \mathrm{Br}(X) & \longrightarrow & \mathrm{Br}(X)/\mathrm{Br}(K) & \longrightarrow & 0 \\
\downarrow & & \downarrow b & & \downarrow c & & \\
\bigoplus_v \mathrm{Br}(K_v^h) & \xrightarrow{f} & \bigoplus_v \mathrm{Br}(X_{K_v^h})/\mathrm{Br}(\mathcal{X}_{\mathcal{O}_v^h}) & \longrightarrow & \bigoplus_v \mathrm{Br}(X_{K_v^h})/(\mathrm{Br}(\mathcal{X}_{\mathcal{O}_v^h}) + \mathrm{Br}(K_v^h)) & &
\end{array}$$

Let a denote the map

$$\mathrm{Br}(K) \longrightarrow \bigoplus_v \mathrm{Br}(K_v^h)/\mathrm{Ker}(f).$$

By the Snake Lemma, there is a long exact sequence

$$\mathrm{Ker}(a) \longrightarrow \mathrm{Ker}(b) \longrightarrow \mathrm{Ker}(c) \longrightarrow \mathrm{Coker}(a) \xrightarrow{\bar{f}} \mathrm{Coker}(b).$$

By purity of Brauer groups (cf. [Čes]) and Lemma 3.1, $\mathrm{Ker}(b) = \mathrm{Br}(\mathcal{X})$. Thus, it suffices to show that $\mathrm{Ker}(a)$ and $\mathrm{Ker}(f)$ are finite groups. Since

$$\mathrm{Ker}(\mathrm{Ker}(a)) \longrightarrow \bigoplus_{v|\infty} \mathrm{Br}(K_v) \oplus \mathrm{Ker}(f) = \mathrm{Ker}(\mathrm{Br}(K) \longrightarrow \bigoplus_v \mathrm{Br}(K_v)),$$

by the exact sequence coming from the global class field theory (cf. [Mil1, Chap. II, Prop. 2.1])

$$0 \longrightarrow \mathrm{Br}(K) \longrightarrow \bigoplus_v \mathrm{Br}(K_v) \longrightarrow \mathbb{Q}/\mathbb{Z} \longrightarrow 0,$$

we have that the map

$$\mathrm{Ker}(a) \longrightarrow \bigoplus_{v|\infty} \mathrm{Br}(K_v) \oplus \mathrm{Ker}(f)$$

is injective and $\mathrm{Coker}(a)$ is of cofinite type. Thus, it suffices to show that $\mathrm{Ker}(f)$ is finite and $\mathrm{Coker}(\bar{f})$ is of finite exponent. Let L/K be finite Galois extension such that $X(L)$ is not empty. Let $P \in X(L)$. Then P defines a K -morphism $g : Z := \mathrm{Spec} L \rightarrow X$ and the morphism can extend to a C -morphism $g : \mathcal{Z} := \mathrm{Spec} \mathcal{O}_L \rightarrow \mathcal{X}$ since π is proper. Consider the composition

$$\bigoplus_v \mathrm{Br}(K_v^h) \xrightarrow{f} \bigoplus_v \mathrm{Br}(X_{K_v^h})/\mathrm{Br}(\mathcal{X}_{\mathcal{O}_v^h}) \longrightarrow \bigoplus_v \mathrm{Br}(Z_{K_v^h})/\mathrm{Br}(\mathcal{Z}_{\mathcal{O}_v^h}).$$

Obviously, $\mathrm{Ker}(f)$ is a subgroup of the kernel of the composition. We want to show that the kernel of the composition is killed by $[L : K]$. Then, it will imply that $\mathrm{Ker}(f)$ is killed by $[L : K]$. The natural map $\mathcal{Z} \rightarrow C$ is finite flat of degree $[L : K]$. Its base change $Z_{K_v^h} \rightarrow \mathrm{Spec} K_v^h$ is also finite flat of degree $[L : K]$. They induce restriction map $\mathrm{res}_v : \mathrm{Br}(K_v^h) \rightarrow \mathrm{Br}(Z_{K_v^h})$ and corestriction map $\mathrm{cores}_v : \mathrm{Br}(Z_{K_v^h}) \rightarrow \mathrm{Br}(K_v^h)$ (cf. [CTS2, §3.8, p 98]). Since $\mathrm{Br}(\mathcal{O}_v^h) = 0$ for all $v \in C^\circ$, the composition map above can be identified with the restriction map

$$\bigoplus_v \mathrm{Br}(K_v^h)/\mathrm{Br}(\mathcal{O}_v^h) \xrightarrow{\bigoplus_v \mathrm{res}_v} \bigoplus_v \mathrm{Br}(Z_{K_v^h})/\mathrm{Br}(\mathcal{Z}_{\mathcal{O}_v^h}).$$

Since the composition $(\oplus_v \text{cores}_v) \circ (\oplus_v \text{res}_v)$ is equal to the multiplication by $[L : K]$, thus, the kernel of $\oplus_v \text{res}_v$ is killed by $[L : K]$. This proves that $\text{Ker}(f)$ is killed by $[L : K]$. By functoriality, the same argument can imply that $\text{Ker}(\bar{f})$ is also killed by $[L : K]$. To show that $\text{Ker}(f)$ is finite, it suffices to show that for all but finitely many $v \in C^\circ$, the map

$$\text{Br}(K_v^h) \longrightarrow \text{Br}(X_{K_v^h})/\text{Br}(\mathcal{X}_{\mathcal{O}_v^h})$$

is injective. If $X(K_v^h)$ is not empty, the restriction-corestriction argument above will imply that the kernel is killed by 1 and therefore is trivial. Thus, it remains to be shown that $X(K_v^h)$ is not empty for all but finitely many v . Note that if $\mathcal{X}_{\mathcal{O}_v^h} \longrightarrow \text{Spec } \mathcal{O}_v^h$ is smooth and the special fiber over v admits a rational point, by hensel's lemma, $\mathcal{X}_{\mathcal{O}_v^h}$ admits a section. This will imply that $X(K_v^h)$ is not empty. Thus, it suffices to show that for all but finitely many v , the special fiber over v is smooth and has a rational point. For the proof of this, we refer it to [CTSDV, Chap. II, §2]. \square

Remark 3.4. In fact, the above proof shows that the kernel is killed by $[L : K]$ and $\text{Br}(\mathcal{X}) \cap \overline{\text{Br}(K)}$ is killed by $2[L : K]$ for any finite extension L/K such that $X(L) \neq \emptyset$.

Theorem 3.5. *Let $\pi : \mathcal{X} \longrightarrow C$ be a proper flat morphism, where C is $\text{Spec } \mathcal{O}_K$ for some number field K . Assume that \mathcal{X} is regular and the generic fiber X of π is projective and geometrically connected over K . Let ℓ be a prime number. Then there are exact sequences up to finite groups*

$$0 \longrightarrow \text{III}(\text{Pic}_{X/K}^0) \longrightarrow \text{Br}(\mathcal{X}) \longrightarrow \text{Br}(X_{\bar{K}})^{G_K}$$

and

$$0 \longrightarrow \text{III}(\text{Pic}_{X/K}^0)(\ell) \longrightarrow \text{Br}(\mathcal{X})(\ell) \longrightarrow \text{Br}(X_{\bar{K}})^{G_K}(\ell) \longrightarrow 0.$$

Proof. By the Hochschild-Serre spectral sequence

$$E_2^{p,q} = H^p(K, H^q(X_{\bar{K}}, \mathbb{G}_m)) \Rightarrow H^{p+q}(X, \mathbb{G}_m)$$

and the fact $H^3(K, \mathbb{G}_m) = 0$ (cf. [Mil4, Prop. 2.7]), we get a long exact sequence

$$\text{Br}(K) \longrightarrow \text{Ker}(\text{Br}(X)) \longrightarrow \text{Br}(X_{\bar{K}})^{G_K} \longrightarrow H^1(K, \text{Pic}_{X/K}) \longrightarrow 0.$$

Thus, we get a injective natural map

$$H^1(K, \text{Pic}_{X/K}) \longrightarrow \text{Br}(X)/\text{Br}(K).$$

Similarly, we get

$$H^1(K_v^h, \text{Pic}_{X/K}) \longrightarrow \text{Br}(X_{K_v^h})/\text{Br}(K_v^h).$$

Consider the following commutative diagram with exact rows

$$\begin{array}{ccccc} 0 & \longrightarrow & H^1(K, \text{Pic}_{X/K}) & \longrightarrow & \text{Br}(X)/\text{Br}(K) & \xrightarrow{b} & \text{Br}(X_{\bar{K}})^{G_K} \\ & & \downarrow & & \downarrow & & \downarrow \\ & & \prod_v H^1(K_v^h, \text{Pic}_{X/K}) & \xrightarrow{a} & \prod_v \text{Br}(X_{K_v^h})/(\text{Br}(\mathcal{X}_{\mathcal{O}_v^h}) + \text{Br}(K_v^h)) & \longrightarrow & \prod_v \text{Br}(X_{\bar{K}})^{G_v}/\text{Br}(\mathcal{X}_{\mathcal{O}_v^h}) \end{array}$$

where G_v denotes $\text{Gal}(\bar{K}/K^h)$. The above diagram gives the following commutative diagram

$$\begin{array}{ccccccc}
0 & \longrightarrow & H^1(K, \text{Pic}_{X/K}) & \longrightarrow & \text{Br}(X)/\text{Br}(K) & \longrightarrow & \text{Im}(b) \longrightarrow 0 \\
& & \downarrow f & & \downarrow g & & \downarrow h \\
0 & \rightarrow & \prod_v H^1(K_v^h, \text{Pic}_{X/K})/\text{Ker}(a) & \rightarrow & \prod_v \text{Br}(X_{K_v^h})/(\text{Br}(\mathcal{X}_{\mathcal{O}_v^h}) + \text{Br}(K_v^h)) & \rightarrow & \prod_v \text{Br}(X_{\bar{K}})^{G_v}/\text{Br}(\mathcal{X}_{\mathcal{O}_v^h})
\end{array}$$

By the Snake Lemma, we get a long exact sequence

$$0 \longrightarrow \text{Ker}(f) \longrightarrow \text{Ker}(g) \longrightarrow \text{Ker}(h) \longrightarrow \text{Coker}(f).$$

By Lemma 3.3, the natural map

$$\text{Br}(\mathcal{X}) \longrightarrow \text{Ker}(g)$$

has a finite kernel and a finite cokernel. Thus, to prove the theorem, it suffices to show that the natural maps

$$\text{III}(\text{Pic}_{X/K}^0) \longrightarrow \text{Ker}(f)$$

and

$$\text{Ker}(h)(\ell) \longrightarrow \text{Br}(X_{\bar{K}})^{G_K}(\ell)$$

have finite kernels and cokernels and the natural map

$$\text{Ker}(h) \longrightarrow \text{Coker}(f)$$

has a finite image. By Lemma 3.7 below, $\text{Ker}(a)$ is of finite exponent. Since there is an exact sequence

$$0 \longrightarrow \text{III}(\text{Pic}_{X/K}) \longrightarrow \text{Ker}(f) \longrightarrow \text{Ker}(a),$$

and $\text{Ker}(f)$ is of cofinite type (this follows from that $\text{Ker}(g)$ is of cofinite type since $\text{Br}(\mathcal{X})$ is of cofinite type), we have that the injective map

$$\text{III}(\text{Pic}_{X/K}) \longrightarrow \text{Ker}(f)$$

has a finite cokernel. By Lemma 2.3, the natural map

$$\text{III}(\text{Pic}_{X/K}^0) \longrightarrow \text{III}(\text{Pic}_{X/K})$$

has a finite kernel and finite cokernel. Thus, the map

$$\text{III}(\text{Pic}_{X/K}^0) \longrightarrow \text{Ker}(f)$$

also has a finite kernel and finite cokernel. Since $\text{Im}(b)$ is of cofinite type and by Proposition 2.5, the target of h is a product of finite groups, thus h will map the maximal divisible subgroup of $\text{Im}(b)$ to zero. Thus, the divisible part of $\text{Im}(b)$ is contained in $\text{Ker}(h)$. By a theorem of Colliot-Thélène and Skorobogatov [CTS1], $\text{Im}(b)$ has a finite index in $\text{Br}(X_{\bar{K}})^{G_K}$. Thus, they have the same maximal divisible subgroup. So we have

$$\text{Ker}(h)_{\text{div}} = (\text{Br}(X_{\bar{K}})^{G_K})_{\text{div}}.$$

It follows that the inclusion

$$\mathrm{Ker}(h)(\ell) \hookrightarrow \mathrm{Br}(X_{\bar{K}})^{G_K}(\ell)$$

has a finite cokernel for any prime ℓ . It remains to be shown that the natural map

$$\mathrm{Ker}(h) \longrightarrow \mathrm{Coker}(f)$$

has an image of finite exponent. To prove this, we will use a pull-back trick of Colliot-Thélène and Skorobogatov (cf. [CTS1] or [Yua]). The idea is that finding finitely many $\pi_i : \mathcal{Z}_i \rightarrow C$ of relative dimension 1 which satisfies the same condition as π and admits a C -morphism $\mathcal{Z}_i \rightarrow \mathcal{X}$, then use the functoriality of the map $\mathrm{Ker}(h) \rightarrow \mathrm{Coker}(f)$ to get a commutative diagram

$$\begin{array}{ccc} \mathrm{Ker}(h) & \longrightarrow & \mathrm{Coker}(f) \\ \downarrow & & \downarrow \\ \bigoplus_i \mathrm{Ker}(h_i) & \longrightarrow & \bigoplus_i \mathrm{Coker}(f_i) \end{array}$$

Since $\mathrm{Ker}(h_i) = 0$ ($\mathrm{Br}(Z_{\bar{K}})^{G_K} = 0$), to prove the claim, it is enough to find \mathcal{Z}_i such that the second column has a kernel of finite exponent. Since $\mathrm{Ker}(a_i)$ is of finite exponent, by the Snake Lemma, it suffices to find \mathcal{Z}_i such that the natural map

$$\mathrm{Coker}(H^1(K, \mathrm{Pic}_{X/K}) \rightarrow \prod_v H^1(K_v^h, \mathrm{Pic}_{X/K})) \rightarrow \bigoplus_i \mathrm{Coker}(H^1(K, \mathrm{Pic}_{Z_i/K}) \rightarrow \prod_v H^1(K_v^h, \mathrm{Pic}_{Z_i/K}))$$

has a kernel of finite exponent. By Lemma 3.8 below, there exist smooth projective geometrically connected curves $Z_i \subset X$ satisfying the above condition. To get \mathcal{Z}_i , we can take the Zariski closure of Z_i in \mathcal{X} first and then desingularize it. This completes the proof of the theorem. \square

Remark 3.6. The above proof actually shows the cokernel of $\mathrm{Br}(\mathcal{X}) \rightarrow \mathrm{Ker}(h)$ has a cokernel of finite exponent. Since $\mathrm{Im}(b)$ has a finite index in $\mathrm{Br}(X_{\bar{K}})^{G_K}$, thus the natural map

$$\mathrm{Br}(X_{\bar{K}})^{G_K} / \mathrm{Br}(\mathcal{X}) \longrightarrow \prod_v \mathrm{Br}(X_{\bar{K}})^{G_v} / \mathrm{Br}(\mathcal{X}_{\mathcal{O}_v^h})$$

has a finite kernel. To finish the proof of Theorem 1.2, it suffices to show that

$$(\mathrm{Br}(X_{\bar{K}})^{G_v} / \mathrm{Br}(\mathcal{X}_{\mathcal{O}_v^h}))(\ell) = 0$$

for all v when $\ell \gg 0$.

Lemma 3.7. *Let a denote the map*

$$\prod_v H^1(K_v^h, \mathrm{Pic}_{X/K}) \longrightarrow \prod_v \mathrm{Br}(X_{K_v^h}) / (\mathrm{Br}(\mathcal{X}_{\mathcal{O}_v^h}) + \mathrm{Br}(K_v^h)),$$

then $\ker(a)$ is of finite exponent.

Proof. By [Yua, §3.3, p17], there exist smooth projective geometrically connected curves $Z_i \subset X$ over K and an abelian variety A/K with a morphism $A \rightarrow \bigoplus_i \text{Pic}_{Z_i/K}^0$ such that the induced G_K -equivariant map

$$\text{Pic}(X_{\bar{K}}) \times A(\bar{K}) \longrightarrow \bigoplus_i \text{Pic}(Z_i, \bar{K})$$

has a kernel and a cokernel killed by some positive integer N . Taking the Zariski closure of Z_i in \mathcal{X} and then desingularizing it, we get a C -morphism $\mathcal{Z}_i \rightarrow \mathcal{X}$. This gives a commutative diagram

$$\begin{array}{ccc} H^1(K_v^h, \text{Pic}_{X/K}) & \xrightarrow{a_v} & \text{Br}(X_{K_v^h}) / (\text{Br}(\mathcal{X}_{\mathcal{O}_v^h}) + \text{Br}(K_v^h)) \\ \downarrow & & \downarrow \\ \bigoplus_i H^1(K_v^h, \text{Pic}_{Z_i/K}) & \longrightarrow & \bigoplus_v \text{Br}(Z_i, K_v^h) / (\text{Br}(\mathcal{Z}_i, \mathcal{O}_v^h) + \text{Br}(K_v^h)) \end{array}$$

Since $\text{Br}(\mathcal{Z}_i, \mathcal{O}_v^h) = 0$ (cf. [Mil4, Lem. 2.6]), thus, by definition, the second row is injective. It follows that $\ker(a_v)$ is contained in the kernel of the first column, which is a subgroup of the kernel of the map

$$H^1(K_v^h, \text{Pic}_{X/K}) \oplus H^1(K, A) \longrightarrow \bigoplus_i H^1(K_v^h, \text{Pic}_{Z_i/K}).$$

By the long exact sequence of Galois cohomology, it is easy to see the kernel of this map is killed by N^2 . This proves $N^2 \text{Ker}(a) = 0$. \square

Lemma 3.8. *Let X be a smooth projective geometrically connected variety over a number field K . Then, there exist smooth projective geometrically connected curves $Z_i \subset X$ such that the induced map*

$$\text{Coker}(H^1(K, \text{Pic}_{X/K}) \rightarrow \prod_v H^1(K_v^h, \text{Pic}_{X/K})) \rightarrow \bigoplus_i \text{Coker}(H^1(K, \text{Pic}_{Z_i/K}) \rightarrow \prod_v H^1(K_v^h, \text{Pic}_{Z_i/K}))$$

has a kernel of finite exponent.

Proof. We will use the same argument as in the proof of the above lemma. Let Z_i and A as in the above lemma. So we have a G_K -equivariant map

$$\text{Pic}(X_{\bar{K}}) \times A(\bar{K}) \longrightarrow \bigoplus_i \text{Pic}(Z_i, \bar{K})$$

with a kernel and a cokernel killed by some positive integer N . If we think

$$\text{Coker}(H^1(K, -) \rightarrow \prod_v H^1(K_v^h, -))$$

as a functor, obviously, it commutes with direct sum. Thus, it suffices to show that for any morphism of G_K -modules $P_1 \rightarrow P_2$ with a kernel and cokernel killed by N , the induced map

$$\mathrm{Coker}(H^1(K, P_1) \rightarrow \prod_v H^1(K_v^h, P_1)) \rightarrow \mathrm{Coker}(H^1(K, P_2) \rightarrow \prod_v H^1(K_v^h, P_2))$$

has a kernel of finite exponent. Consider the following commutative diagram

$$\begin{array}{ccccccc} H^1(K, P_1) & \xrightarrow{a} & H^1(K, P_2) & \longrightarrow & \mathrm{Coker}(a) & \longrightarrow & 0 \\ \downarrow f & & \downarrow g & & \downarrow \bar{g} & & \\ \mathrm{Ker}(b) & \longrightarrow & \prod_v H^1(K_v^h, P_1) & \xrightarrow{b} & \prod_v H^1(K_v^h, P_2) & \longrightarrow & \mathrm{Coker}(b) \end{array}$$

By the assumption, we have that $\mathrm{Ker}(b)$ and $\mathrm{Coker}(a)$ are killed by N^2 . By the Snake Lemma, there is an exact sequence

$$\mathrm{Ker}(\bar{g}) \rightarrow \mathrm{Coker}(f)/\mathrm{Ker}(b) \rightarrow \mathrm{Coker}(g).$$

It follows that the kernel of $\mathrm{Coker}(f) \rightarrow \mathrm{Coker}(g)$ has a kernel killed by N^4 . \square

By Remark 3.2, we can replace the henselain local ring by the completed local ring in the proof of the above theorem. If we exclude finitely many places, the map $\mathcal{X}_{\mathcal{O}_v} \rightarrow \mathrm{Spec} \mathcal{O}_v$ will be a smooth projective morphism. By the pull-back trick, one can show that

$$\mathrm{Br}(\mathcal{X}_{\mathcal{O}_v}) \rightarrow \mathrm{Br}(\mathcal{X}_{\mathcal{O}_v^h})^{G_{k(v)}}$$

has a cokernel killed by some positive integer independent of v (cf. [Qim3, Prop. 3.6]). In conjunction with Proposition 2.4, this will imply that for all but finitely many v , the natural map

$$\mathrm{Br}(\mathcal{X}_{\mathcal{O}_v})(\mathrm{non}-p_v) \rightarrow \mathrm{Br}(X_{\overline{K}_v})^{G_{K_v}}(\mathrm{non}-p_v)$$

has a cokernel killed by some positive integer independent of v . Thus, to prove Theorem 1.2, by Remark 3.6, it suffices to show that the natural map

$$\mathrm{Br}(\mathcal{X}_{\mathcal{O}_v})(p_v) \rightarrow \mathrm{Br}(X_{\overline{K}_v})^{G_{K_v}}(p_v)$$

is surjective for all but finitely many v , where p_v is the characteristic of the residue field at v . We will prove this in the Corollary 3.10 below.

Proposition 3.9. *Let K/\mathbb{Q}_p be an unramified finite extension and $\pi : \mathcal{X} \rightarrow \mathrm{Spec} \mathcal{O}_K$ be a smooth projective morphism with generic fiber X . Assuming $p \geq 5$, then the natural map*

$$H_{\mathrm{fppf}}^2(\mathcal{X}, \mu_{p^n}) \rightarrow H^2(X_{\overline{K}}, \mu_{p^n})^{G_K}$$

is surjective for any $n \geq 1$

Proof. Let \mathcal{X}_n denote the \mathcal{O}_K/p^n -scheme $\mathcal{X} \otimes_{\mathbb{Z}} \mathbb{Z}/p^n$. Fontaine and Messing [FM, III. 3.1] defined a sheaf $\mathcal{S}_n(r)$ on the small syntomic site $(\mathcal{X}_n)_{syn}$ of \mathcal{X}_n . Following definitions and notations in [EN, §2.1], for $0 \leq r \leq p-1$, define

$$\mathcal{S}_n(r) := \text{Ker}(\mathcal{J}_n^{[r]} \xrightarrow{1-\varphi_r} \mathcal{O}_n^{cr}).$$

In fact, there is an exact sequence of sheaves on $(\mathcal{X}_n)_{syn}$

$$0 \longrightarrow \mathcal{S}_n(r) \longrightarrow \mathcal{J}_n^{[r]} \xrightarrow{1-\varphi_r} \mathcal{O}_n^{cr} \longrightarrow 0.$$

For our purpose, we only consider the case with $r = 1$. The syntomic cohomology $H^i((\mathcal{X}_n)_{syn}, \mathcal{S}_n(1))$ computes the flat cohomology $H_{\text{fppf}}^i(\mathcal{X}, \mu_{p^n})$ (cf. [Kur] and [Sat, §1.4 and §4.5]) and the natural map

$$H_{\text{fppf}}^i(\mathcal{X}, \mu_{p^n}) \longrightarrow H^i(X, \mu_{p^n})$$

is compatible with the Fontaine-Messing map

$$H^i((\mathcal{X}_n)_{syn}, \mathcal{S}_n(1)) \longrightarrow H^i(X, \mu_{p^n}).$$

Thus, it suffices to show that the following map induced by the Fontaine-Messing map

$$H^2((\mathcal{X}_n)_{syn}, \mathcal{S}_n(1)) \longrightarrow H^2(X_{\overline{K}}, \mu_{p^n})^{G_K}$$

is surjective for any $n \geq 1$.

Let Y denote the special fiber \mathcal{X}_1 . Following notations in [Nek, III. 4.10], denote

$$M_n^2 := H^2((\mathcal{X}_n)_{syn}, \mathcal{O}_n^{cr}) = H^2((Y/W_n)_{\text{cris}}, \mathcal{O}_{Y/W_n}) = H_{dR}^2(\mathcal{X}_n/W_n),$$

$$F^r M_n^2 := H^2((\mathcal{X}_n)_{syn}, \mathcal{J}_n^{[r]}) = H^2((\mathcal{X}_n)_{\text{Zar}}, \sigma_{\geq r} \Omega_{\mathcal{X}_n/W_n}^{\bullet}), \quad T_n^2 := H^2(X_{\overline{K}}, \mathbb{Z}/p^n \mathbb{Z}).$$

By [FM, II. 2.7], $(M_n^2, F^r M_n^2, \varphi_r)$ defined an object of $MF_{W, \text{tors}}^{[0,2]} \subseteq MF_{W, \text{tors}}^{[0,p-1]}$ (cf. [Nek, III. Prop. 4.11] or [BM, Thm. 3.2.3]). For $0 \leq r < p-1$, by [Nek, III. 4.8], the functor T (cf. [Nek, III. 4.6] for the definition) induces an isomorphism

$$\alpha_{r, M_n^2} : (F^r M_n^2)^{\varphi_r=1} \longrightarrow (T_n^2(r))^{G_K}.$$

By the exact sequence of sheaves on $(\mathcal{X}_n)_{syn}$

$$0 \longrightarrow \mathcal{S}_n(1) \longrightarrow \mathcal{J}_n^{[1]} \xrightarrow{1-\varphi_1} \mathcal{O}_n^{cr} \longrightarrow 0,$$

taking cohomology, we get a long exact sequence

$$\begin{aligned} H^2((\mathcal{X}_n)_{syn}, \mathcal{S}_n(1)) &\longrightarrow H^2((\mathcal{X}_n)_{syn}, \mathcal{J}_n^{[1]}) \xrightarrow{1-\varphi_1} H^2((\mathcal{X}_n)_{syn}, \mathcal{O}_n^{cr}) \\ &\longrightarrow H^3((\mathcal{X}_n)_{syn}, \mathcal{S}_n(1)). \end{aligned}$$

Thus, there is a surjective natural map

$$H^2((\mathcal{X}_n)_{syn}, \mathcal{S}_n(1)) \longrightarrow H^2((\mathcal{X}_n)_{syn}, \mathcal{J}_n^{[1]})^{\varphi_1=1}.$$

By [Nek, III. Thm. 5.2], there is a commutative diagram

$$\begin{array}{ccc} H^2((\mathcal{X}_n)_{syn}, \mathcal{S}_n(1)) & \longrightarrow & H^2((\mathcal{X}_n)_{syn}, \mathcal{J}_n^{[1]})^{\varphi_1=1} \\ \downarrow \nu & & \downarrow \alpha_{1, M_n^2} \\ H^2(X, \mathbb{Z}/p^n\mathbb{Z}(1)) & \longrightarrow & H^2(X_{\overline{K}}, \mathbb{Z}/p^n\mathbb{Z}(1))^{G_K} \end{array}$$

where ν denotes the Fontaine-Messing map. Since the first row is surjective and α_{1, M_n^2} is an isomorphism, it follows that the natural map

$$H^2((\mathcal{X}_n)_{syn}, \mathcal{S}_n(1)) \longrightarrow H^2(X_{\overline{K}}, \mu_{p^n})^{G_K}$$

is surjective. This completes the proof. \square

Corollary 3.10. *Let $\pi : \mathcal{X} \longrightarrow \text{Spec } \mathcal{O}_K$ be a proper flat morphism, where K is a number field. Assume that \mathcal{X} is regular and the generic fiber X of π is projective and geometrically connected over K . For any finite place v , let p_v denote the characteristic of the residue field of \mathcal{O}_K at v . Then, for all but finitely many places v , the natural map*

$$\text{Br}(\mathcal{X}_{\mathcal{O}_v})(p_v) \longrightarrow \text{Br}(X_{\overline{K}_v})^{G_{K_v}}(p_v)$$

is surjective.

Proof. K/\mathbb{Q} only ramifies at finitely many places. Thus, by the proposition above, for all but finitely many v , the natural map

$$H_{\text{fppf}}^2(\mathcal{X}_{\mathcal{O}_v}, \mu_{p_v^n}) \longrightarrow H^2(X_{\overline{K}_v}, \mu_{p_v^n})^{G_{K_v}}$$

is surjective for any $n \geq 1$. By Proposition 2.1, for all but finitely many v , the natural map

$$H^2(X_{\overline{K}_v}, \mu_{p_v^n})^{G_{K_v}} \longrightarrow \text{Br}(X_{\overline{K}_v})^{G_{K_v}}(p_v)$$

is surjective for any $n \geq 1$. By the following commutative diagram

$$\begin{array}{ccc} H_{\text{fppf}}^2(\mathcal{X}_{\mathcal{O}_v}, \mu_{p_v^n}) & \longrightarrow & \text{Br}(\mathcal{X}_{\mathcal{O}_v})[p_v^n] \\ \downarrow & & \downarrow \\ H^2(X_{\overline{K}_v}, \mu_{p_v^n})^{G_{K_v}} & \longrightarrow & \text{Br}(X_{\overline{K}_v})^{G_{K_v}}[p_v^n], \end{array}$$

for all but finitely many v , the natural map

$$\text{Br}(\mathcal{X}_{\mathcal{O}_v})[p_v^n] \longrightarrow \text{Br}(X_{\overline{K}_v})^{G_{K_v}}[p_v^n]$$

is surjective for any $n \geq 1$. \square

Remark 3.11. By [Qin3, Thm. 1.10], the corollary can imply that for all but finitely many places v , the p_v -primary part of the kernel of the natural map

$$\mathrm{Hom}(\mathrm{Br}(X_{K_v})/\mathrm{Br}(K) + \mathrm{Br}(\mathcal{X}_{\mathcal{O}_v}), \mathbb{Q}/\mathbb{Z}) \longrightarrow \mathrm{Alb}_X(K_v)$$

is trivial.

4 Applications

4.1 Reduction of Artin's question

Lemma 4.1. *Assuming that $\mathrm{Br}(X_{\bar{K}})^{G_K}$ is finite for all smooth projective geometrically connected surfaces over a number field K , then $\mathrm{Br}(X_{\bar{K}})^{G_K}$ is finite for all smooth projective geometrically connected varieties over K .*

Proof. Let X be a smooth projective geometrically connected variety over a number field K . Assuming that $\dim(X) > 2$, by a theorem of Ambrosi [Amb, Cor. 1.6.2.1], there exists a smooth projective geometrically connected hyperplane section D of X such that the induced map

$$\mathrm{NS}(X) \otimes_{\mathbb{Z}} \mathbb{Q} \longrightarrow \mathrm{NS}(D) \otimes_{\mathbb{Z}} \mathbb{Q}$$

is an isomorphism. By Proposition 2.1, for any prime ℓ , there is a commutative diagram with exact rows

$$\begin{array}{ccccccc} 0 & \longrightarrow & \mathrm{NS}(X) \otimes_{\mathbb{Z}} \mathbb{Q}_{\ell} & \longrightarrow & H^2(X_{\bar{K}}, \mathbb{Q}_{\ell}(1))^{G_K} & \longrightarrow & V_{\ell}\mathrm{Br}(X_{\bar{K}})^{G_K} \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & \mathrm{NS}(D) \otimes_{\mathbb{Z}} \mathbb{Q}_{\ell} & \longrightarrow & H^2(D_{\bar{K}}, \mathbb{Q}_{\ell}(1))^{G_K} & \longrightarrow & V_{\ell}\mathrm{Br}(D_{\bar{K}})^{G_K} \longrightarrow 0 \end{array}$$

Since the first column is an isomorphism and the second column is injective (the weak Lefschetz theorem), by the Snake Lemma, the third column is injective. This implies that for any ℓ , the ℓ -primary part of the kernel of the map

$$\mathrm{Br}(X_{\bar{K}})^{G_K} \longrightarrow \mathrm{Br}(D_{\bar{K}})^{G_K}$$

is finite. Since $\mathrm{NS}(X)$ and $\mathrm{NS}(D)$ are finite groups, thus for sufficiently large ℓ , the induced map

$$\mathrm{NS}(X) \otimes_{\mathbb{Z}} \mathbb{Q}_{\ell}/\mathbb{Z}_{\ell} \longrightarrow \mathrm{NS}(D) \otimes_{\mathbb{Z}} \mathbb{Q}_{\ell}/\mathbb{Z}_{\ell}$$

is an isomorphism. By the weak Lefschetz theorem for torsion locally constant sheaves, the induced map

$$H^2(X_{\bar{K}}, \mathbb{Q}_{\ell}/\mathbb{Z}_{\ell}(1)) \longrightarrow H^2(D_{\bar{K}}, \mathbb{Q}_{\ell}/\mathbb{Z}_{\ell}(1))$$

is injective. By Proposition 2.1, we have a similar diagram as before for torsion coefficients when ℓ is sufficiently large. Thus, natural map

$$\mathrm{Br}(X_{\bar{K}})^{G_K}(\ell) \longrightarrow \mathrm{Br}(D_{\bar{K}})^{G_K}(\ell)$$

is injective for all but finitely many ℓ . This proves that the kernel of

$$\mathrm{Br}(X_{\bar{K}})^{G_K} \longrightarrow \mathrm{Br}(D_{\bar{K}})^{G_K}$$

is finite. By induction of the dimension of X , the claim follows. \square

Theorem 4.2. *Assuming that $\mathrm{Br}(X_{\bar{K}})^{G_K}$ is finite for all smooth projective surfaces and $\mathrm{III}(\mathrm{Pic}_{X/K}^0)$ is finite for all smooth projective geometrically connected curves, then $\mathrm{Br}(\mathcal{X})$ is finite for all regular proper flat schemes \mathcal{X} over \mathbb{Z} .*

Proof. Let X be a smooth proper geometrically connected variety over a number field K . By resolution of singularity in characteristic 0, there is a smooth projective variety X' and a birational morphism $X' \rightarrow X$. Since Brauer group is a birational invariants and $\mathrm{Pic}_{X/K}^0$ and $\mathrm{Pic}_{X'/K}^0$ are isogenous to each other, we may assume that X is projective. If $\dim(X) \geq 2$, taking a smooth hyperplane section D in X , then the induced map $\mathrm{Pic}_{X/K}^0 \rightarrow \mathrm{Pic}_{D/K}^0$ has a finite kernel since $H^1(X_{\bar{K}}, \mathbb{Q}_\ell) \rightarrow H^1(D_{\bar{K}}, \mathbb{Q}_\ell)$ is injective. Since $\mathrm{Pic}_{X/K}^0 \rightarrow \mathrm{Pic}_{D/K}^0$ has a finite kernel, there is an abelian variety A/K such that $\mathrm{Pic}_{X/K}^0 \times A$ is isogenous to $\mathrm{Pic}_{D/K}^0$ (cf. [Yua, §3.3, p17]). Thus, the finiteness of $\mathrm{III}(\mathrm{Pic}_{X/K}^0)$ can be reduced to curves. By the above lemma, the finiteness of $\mathrm{Br}(X_{\bar{K}})^{G_K}$ can be reduced to surfaces. Let X denote the generic fiber of \mathcal{X} . By the assumption, $\mathrm{III}(\mathrm{Pic}_{X/K}^0)$ and $\mathrm{Br}(X_{\bar{K}})^{G_K}$ are finite. By Theorem 1.2, $\mathrm{Br}(\mathcal{X})$ is finite. \square

Theorem 4.3. *Assuming that $\mathrm{Br}(\mathcal{X})$ is finite for all 3-dimensional regular proper flat schemes over \mathbb{Z} , then $\mathrm{Br}(\mathcal{X})$ is finite for all regular proper flat schemes over \mathbb{Z} .*

Proof. By the theorem above, it suffices to show that $\mathrm{Br}(X_{\bar{K}})^{G_K}$ and $\mathrm{III}(\mathrm{Pic}_{X/K}^0)$ are finite for all smooth projective geometrically connected surfaces over a number field K under the assumption. By de Jong's theorem [deJ1, Cor. 5.1], there is an alteration $X' \rightarrow X$ such that X' admits a flat proper regular integral model over the ring of integers of a number field. By extending K , we may assume that X' is smooth and geometrically connected over K . Then we have that $\mathrm{Pic}_{X/K}^0 \rightarrow \mathrm{Pic}_{X'/K}^0$ has a finite kernel since $H^1(X_{\bar{K}}, \mathbb{Q}_\ell) \rightarrow H^1(X'_{\bar{K}}, \mathbb{Q}_\ell)$ is injective. Thus, the finiteness of $\mathrm{III}(\mathrm{Pic}_{X'/K}^0)$ can imply the finiteness of $\mathrm{III}(\mathrm{Pic}_{X/K}^0)$. Since $K(X'_{\bar{K}})$ is a finite extension over $K(X_{\bar{K}})$, by the restriction-corestriction argument, the induced map $\mathrm{Br}(K(X_{\bar{K}})) \rightarrow \mathrm{Br}(K(X'_{\bar{K}}))$ has a kernel of finite exponent. It follows that

$$\mathrm{Br}(X_{\bar{K}})^{G_K} \longrightarrow \mathrm{Br}(X'_{\bar{K}})^{G_K}$$

has a finite kernel. By the assumption and Theorem 1.2, $\mathrm{Br}(X'_{\bar{K}})^{G_K}$ and $\mathrm{III}(\mathrm{Pic}_{X'/K}^0)$ are finite. Thus, $\mathrm{Br}(X_{\bar{K}})^{G_K}$ and $\mathrm{III}(\mathrm{Pic}_{X/K}^0)$ are also finite. This completes the proof. \square

4.2 Examples

Proposition 4.4. *Let X be a principal homogeneous space of an abelian variety over a number field K . Assuming that X/K admits a proper regular model $\pi : \mathcal{X} \rightarrow C$ as in*

Theorem 1.2, then there is an isomorphism up to finite groups

$$\text{III}(\text{Pic}_{X/K}^0) \cong \text{Br}(\mathcal{X}).$$

Proof. By [SZ, Thm. 1.1], $\text{Br}(X_{K^s})^{G_K}$ is finite. Then the claim follows directly from Theorem 1.2. \square

Proposition 4.5. (Tankeev) *Let $\pi : \mathcal{X} \rightarrow C$ a proper flat morphism as in Theorem 1.2. Assuming that the generic fiber X is a K3 surface, then $\text{Br}(\mathcal{X})$ is finite.*

Proof. Since $H^1(X, \mathcal{O}_X) = 0$, we have $\text{Pic}_{X/K}^0 = 0$. By [SZ, Thm. 1.2], $\text{Br}(X_{\bar{K}})^{G_K}$ is finite. Then the claim follows directly from Theorem 1.2. \square

References

- [Amb] E. Ambrosi, Specialization of Néron-Severi groups in positive characteristic, arXiv preprint arXiv:1810.06481 .
- [And] Y. André, Pour une théorie inconditionnelle des motifs, Publ. Math. IHES 83, p. 5-49, 1996.
- [BM] C. Breuil, W. Messing. Torsion étale and crystalline cohomologies. Astérisque 279 (2002): 81-124.
- [Čes] K. Česnavičius, Purity for the Brauer group. Duke Mathematical Journal, 168(8)(2019), pp.1461-1486.
- [CT] J.-L. Colliot-Thélène, L'arithmétique du groupe de Chow des zéro-cycles, Journal de théorie des nombres de Bordeaux 7.1 (1995): 51-73.
- [CTS1] J.-L. Colliot-Thélène, A.N. Skorobogatov, Descente galoisienne sur le groupe de Brauer. J. reine angew. Math. 682 (2013) 141-165.
- [CTS2] J.-L. Colliot-Thélène, A.N. Skorobogatov, The Brauer–Grothendieck group, <http://wwwf.imperial.ac.uk/~anskor/brauer.pdf>
- [CTSDV] J.-L. Colliot-Thélène, P. Swinnerton-Dyer, and P. Vojta, Arithmetic Geometry: Lectures Given at the CIME Summer School Held in Cetraro, Italy, September 10-15, 2007. Springer, 2010.
- [Del1] P. Deligne, “SGA 4 1/2.” Lect. Notes Math 569 (1977).
- [Del2] P. Deligne, “Weil’s conjecture: II.” IHÉS Mathematical Publications 52 (1980): 137-252.
- [deJ1] A. J. de Jong, Families of curves and alterations. In Annals of the Fourier Institute (Vol. 47, No. 2, 1997, pp. 599-621).
- [deJ2] A. J. de Jong, Tate conjecture for divisors. Unpublished note.
- [EGA IV] A. Grothendieck, Éléments de géométrie algébrique : IV. Étude locale des schémas et des morphismes de schémas, Troisième partie Publications mathématiques de l’I.H.É.S., tome 28 (1966), p. 5-255.
- [EN] V. Ertl, W. Niziol, Syntomic cohomology and p-adic motivic cohomology. ALGEBRAIC GEOMETRY 6, no. 1 (2019): 100-131.
- [Fal1] G. Faltings, Endlichkeitssätze für abelsche Varietäten über Zahlkörpern. Invent. Math. 73 (1983), no. 3, 349-366.

- [Fal2] G. Faltings, Complements to Mordell. In: G. Faltings, G. Wüstholz et al., Rational points, 2nd edition, F. Viehweg & Sohn, 1986.
- [FM] J.-M. Fontaine, W. Messing, p -adic periods and p -adic étale cohomology, Contemporary Math. 67, 1987, 179-207.
- [Fu] L. Fu, 2011. Etale cohomology theory (Vol. 13). World Scientific.
- [Fuj] K. Fujiwara, A proof of the absolute purity conjecture (after Gabber). In Algebraic geometry 2000, Azumino (pp. 153-183). Mathematical Society of Japan.
- [Gei1] T.H. Geisser, Comparing the Brauer group to the Tate–Shafarevich group. Journal of the Institute of Mathematics of Jussieu, 1-6(2018).
- [Gei2] T.H. Geisser, “TATE’S CONJECTURE AND THE TATE–SHAFAREVICH GROUP OVER GLOBAL FUNCTION FIELDS.” Journal of the Institute of Mathematics of Jussieu (2018): 1-22.
- [Goa] C.D. Gonzalez-Aviles, Brauer Groups and Tate-Shafarevich Groups. J. Math. Sci. Univ. Tokyo, 2003, 10: 391-419.
- [Gro1] A. Grothendieck, Le groupe de Brauer. I. Algèbres d’Azumaya et interprétations diverses, Dix Exposés sur la Cohomologie des Schémas, North-Holland, Amsterdam, 1968, pp. 46–66.
- [Gro2] A. Grothendieck, Le groupe de Brauer. II. Théorie cohomologique, Dix Exposés sur la Cohomologie des Schémas, North-Holland, Amsterdam, 1968, pp. 67–87.
- [Gro3] A. Grothendieck, Le groupe de Brauer. III. Exemples et compléments, Dix Exposés sur la Cohomologie des Schémas, North-Holland, Amsterdam, 1968, pp. 88–188.
- [JS] U. Jannsen, S. Saito, Bertini theorems and Lefschetz pencils over discrete valuation rings, with applications to higher class field theory. Journal of Algebraic Geometry, 21(4)(2012), 683-705.
- [Kur] M. Kurihara, A note on p -adic étale cohomology. Proceedings of the Japan Academy, Series A, Mathematical Sciences 63, no. 7 (1987): 275-278.
- [Mil1] J.S. Milne, Arithmetic duality theorems, Vol. 20. Charleston, SC: BookSurge, 2006.
- [Mil2] J.S. Milne, Abelian Varieties, Chapter V of Arithmetic geometry (Storrs, Conn., 1984), 103–150, Springer, New York, 1986.
- [Mil3] J.S. Milne, Etale cohomology (PMS-33)(1980) (Vol. 5657). Princeton university press.
- [Mil4] J.S. Milne, Comparison of the Brauer Group with the Tate-Safarevic group, J. Fac. Science Univ. Tokyo, Sec. IA 28 (1981), 735–743.

- [MP] K. Madapusi Pera, The Tate conjecture for K3 surfaces in odd characteristic, *Invent. Math.* 201, 625-668 (2015).
- [Mor] M. Morrow, A Variational Tate Conjecture in crystalline cohomology. *Journal of the European Mathematical Society*, 21(11)(2019), 3467-3511.
- [Nek] J. Nekovář, Syntomic cohomology and p -adic regulators, preprint (1998). <https://webusers.imj-prg.fr/~jan.nekovar/pu/syn.pdf>
- [NO] N. Nygaard, A. Ogus, Tate's conjecture for K3 surfaces of finite height, *Ann. of Math.* 122, 461-507 (1985).
- [Nyg] N. O. Nygaard, The Tate conjecture for ordinary K3 surfaces over finite fields, *Invent. Math.* 74, 213-237 (1983).
- [OS] M. Orr, A.N. Skorobogatov, Finiteness theorems for K3 surfaces and abelian varieties of CM type. *Compositio Mathematica*, 2018, 154(8): 1571-1592.
- [Qin1] Y. Qin, On the Brauer groups of fibrations. arXiv preprint arXiv:2012.01324.
- [Qin2] Y. Qin, On geometric Brauer groups and Tate-Shafarevich groups. arXiv:2012.01681.
- [Qin3] Y. Qin, A p -adic local invariant cycle theorem with applications to Brauer groups, arXiv:2103.04945.
- [Sat] K. Sato, p -adic étale Tate twists and arithmetic duality (with an appendix by Hagiwara, K.). *Ann. Sci. École Norm. Sup. (4)* 40, 519–588 (2007)
- [SZ] A.N. Skorobogatov, Y.G. Zarkhin, A finiteness theorem for the Brauer group of Abelian varieties and KS surfaces. *Journal of Algebraic Geometry*, 17(3)(2008), 481-502.
- [Tan1] S.G. Tankeev, On the Brauer group of an arithmetic scheme. *Izvestiya: Mathematics*, 65(2)(2001), 357.
- [Tan2] S.G. Tankeev, On the Brauer group of an arithmetic scheme. II. *Izvestiya: Mathematics*, 67(5)(2003), 1007.
- [Tan3] S.G. Tankeev, On the finiteness of the Brauer group of an arithmetic scheme. *Mathematical Notes*, 95(1-2)(2014), 121-132.
- [Tan4] S. G. Tankeev, Surfaces of K3 type over number fields and the Mumford-Tate conjecture. *Izv. Akad. Nauk SSSR Ser. Mat.* 54 (1990), no. 4, 846-861; translation in *Math. USSR-Izv.* 37 (1991), no. 1, 191- 208.
- [Tan5] S. G. Tankeev, Surfaces of K3 type over number fields and the Mumford-Tate conjecture. II. *Izv. Ross. Akad. Nauk Ser. Mat.* 59 (1995), no. 3, 179-206; translation in *Izv. Math.* 59 (1995), no. 3, 619-646.

- [Tat1] J. Tate, Algebraic cycles and poles of zeta functions. *Arithmetical algebraic geometry*. 1965:93-110.
- [Tat2] J. Tate, Conjectures on algebraic cycles in l -adic cohomology. *Motives* (Seattle, WA, 1991) 55 (1994): 71-83.
- [Tat3] J. Tate, Endomorphisms of abelian varieties over finite fields, *Invent. Math.* 2, 134-144 (1966).
- [Ulm] D. Ulmer, Curves and Jacobians over function fields, *Arithmetic geometry over global function fields*, 2014, pp.281-337.
- [Vid] I. Vidal, Théorie de Brauer et conducteur de Swan. *Journal of Algebraic Geometry* 13.2 (2004): 349-391.
- [Yua] X. Yuan, Comparison of arithmetic Brauer groups with geometric Brauer group
- [Zar1] Y. G. Zarhin, Endomorphisms of Abelian varieties over fields of finite characteristic. *Izv. Akad. Nauk SSSR Ser. Mat.* 39 (1975), 272-277; *Math. USSR Izv.* 9 (1975), 255-260.
- [Zar2] Y. G. Zarhin, Abelian varieties in characteristic p . *Mat. Zametki* 19 (1976), 393-400; *Math. Notes* 19 (1976), 240-244.