

# Conditional strategy equilibrium\*

Lorenzo Bastianello<sup>†</sup>Mehmet S. Ismail<sup>‡</sup>

October 6, 2021

## 1 Definition of pure conditional strategies

Let  $(A_i, u_i)_{i \in N}$  be an  $n$ -person noncooperative game, where  $N = \{1, \dots, n\}$  is the finite set of players,  $A_i = \{a_i^1, a_i^2, \dots, a_i^{m_i}\}$  finite nonempty action set, and  $u_i : A \rightarrow \mathbb{R}_+$  the (von Neumann-Morgenstern) utility function of player  $i \in N$ . A strategy profile is denoted by  $a \in A = \times_{i \in N} A_i$ .<sup>1</sup>

A pure conditional strategy of player  $i$  is a function  $s_i : A_{-i} \rightarrow A_i$ . A pure conditional strategy profile is  $s \in S = \times_{i \in N} S_i$  where  $S_i$  is the set of conditional strategies of player  $i$ . A pure conditional strategy profile  $s \in S$  is an *agreement* if there exists a unique action profile  $a \in A$  such that  $s(a) = a$ —i.e., for all  $i$   $s_i(a_{-i}) = a_i$ —which is the unique fixed point of the conditional strategy profile  $s$ , which always exists because for each  $i$   $A_i$  is nonempty. The intuition is that everybody agrees on the unique strategy profile  $a$ . If  $s \in S$  is not an agreement, then it is called a *disagreement*.

Let  $B = \{s \in S \mid s \text{ is an agreement}\}$  be the set of agreements. The utility function  $U_i : S \rightarrow \mathbb{R}_+$  is defined as follows.

$$U_i(s) = \begin{cases} u_i(a) & \text{if } s \in B \\ 0 & \text{if } s \in S \setminus B, \end{cases}$$

where  $a = s(a)$ . This is a well-defined function since every  $s \in B$  has a unique fixed point, and the disagreement payoff is zero. Note that  $U_i|_B = u_i$ .

---

\*This is a preliminary draft. The prior literature on conditional actions, starting with Nigel (1971), will be reviewed in the next version.

<sup>†</sup>LEMMA, Université Paris 2 Panthéon-Assas, Paris, France. E-mail: lorenzo.bastianello@u-paris2.fr

<sup>‡</sup>Department of Political Economy, King's College London, UK. E-mail: mehmet.s.ismail@gmail.com.

<sup>1</sup>Without loss of generality, we consider nonnegative utility function. The number of pure conditional actions of player  $i$  is given by  $m_i^{\prod_{j \neq i} m_j} = |S_i|$  because  $\prod_{j \neq i} m_j$  gives the number of pure action profiles of everyone but  $i$ .

Let  $(S_i, U_i)_{i \in N}$  denote a game in *conditional extension* of the game  $(A_i, u_i)_{i \in N}$ . We next define conditional strategy equilibrium.

**Definition 1.** A conditional strategy profile  $s^* \in S$  is called a conditional strategy equilibrium if for all  $i$   $s_i^* \in \arg \max U_i(s^*)$  or equivalently

$$U_i(s^*) \geq U_i(s'_i, s_{-i}^*).$$

In other words, a conditional strategy equilibrium (CSE) is a self-enforcing agreement in the space of conditional strategy profiles, just like a Nash equilibrium (Nash, 1950) is a self-enforcing agreement in the space of mixed strategy profiles.

## 2 Conditional mixed extension

Let  $G = (\Delta A_i, u_i)_{i \in N}$  be an  $n$ -person noncooperative game in mixed extension, where  $N = \{1, \dots, n\}$  is the finite set of players,  $\Delta A_i$  the set of all probability distributions over the finite action set  $A_i$ , and  $u_i : \Delta A \rightarrow \mathbb{R}$  the von Neumann-Morgenstern expected utility function of player  $i \in N$  (von Neumann and Morgenstern, 1944). A mixed strategy profile is denoted by  $p \in \Delta A = \times_{i \in N} \Delta A_i$ , where  $\Delta A_i$  is the simplex in  $\mathbb{R}^{m_i-1}$ .

Let  $\mathcal{A}_{-i}$  be the Borel  $\sigma$ -algebra over  $\Delta A_{-i}$ . The set of conditional mixed strategies of player  $i$  is given by  $\Sigma_i = \{ \sigma_i : \Delta A_{-i} \rightarrow \Delta A_i \mid \sigma_i \text{ is a simple and } \mathcal{A}_{-i}\text{-measurable} \}$ . A conditional mixed strategy profile is  $\sigma \in \Sigma = \times_{i \in N} \Sigma_i$  where  $\Sigma_i$  is the set of conditional strategies of player  $i$ . A conditional strategy profile  $\sigma \in \Sigma$  is an *agreement* if there exists a unique  $p \in \Delta A$  such that  $\sigma(p) = p$ —i.e., for all  $i$   $\sigma_i(p_{-i}) = p_i$ —which is the unique fixed point of the conditional strategy profile  $\sigma$ , which always exists because for each  $i$   $\Delta A_i$  is nonempty. The intuition is that everybody agrees on the unique strategy profile  $p$ . If  $\sigma \in \Sigma$  is not an agreement, then it is called a *disagreement*.

Let  $B = \{ \sigma \in \Sigma \mid \sigma \text{ is an agreement} \}$  be the set of agreements. The utility function  $U_i : \Sigma \rightarrow \mathbb{R}_+$  is defined as follows.

$$U_i(\sigma) = \begin{cases} u_i(p) & \text{if } \sigma \in B \\ 0 & \text{if } \sigma \in \Sigma \setminus B, \end{cases}$$

where  $p = \sigma(p)$ . This is a well-defined function since every  $\sigma \in B$  has a unique fixed point. Note that we normalize the disagreement payoff to zero and that  $U_i|_B = u_i$ .

Let  $\Gamma = (\Sigma_i, U_i)_{i \in N}$  denote a game in *conditional mixed extension* of the game  $(A_i, u_i)_{i \in N}$ . We next define conditional strategy equilibrium.

**Definition 2.** A conditional strategy profile  $\sigma^* \in \Sigma$  is called a *conditional strategy equilibrium* if for all  $i$   $\sigma_i^* \in \arg \max U_i(\sigma^*)$  or equivalently

$$U_i(\sigma^*) \geq U_i(\sigma'_i, \sigma_{-i}^*).$$

In other words, a conditional mixed strategy equilibrium (CSE) is a self-enforcing agreement in the space of conditional strategies.

## 2.1 Conditional mixed extension and the mixed extension of the pure conditional strategies

Consider  $\hat{S}_i = \{\hat{s}_i : \Delta A_{-i} \rightarrow A_i \mid \hat{s}_i \text{ is } \mathcal{A}_{-i}\text{-measurable}\}$ , which is the set of pure conditional strategies of player  $i$  against mixed strategies of the others. We denote  $\Delta \hat{S}_i$  the set of all probability measures with finite support over  $\hat{S}_i$ . Every probability measure  $\mu \in \Delta \hat{S}_i$  induces a function  $\sigma_i \in \Sigma_i$ . Define function

$$\begin{aligned} \phi : \Delta \hat{S}_i &\rightarrow \Sigma_i \\ \mu &\mapsto \sigma_i^\mu \end{aligned} \tag{1}$$

where for all  $q_{-i}$

$$\sigma_i^\mu(q_{-i}) = \int_{\hat{S}_i} \hat{s}_i(q_{-i}) d\mu(\hat{s}_i),$$

which is a probability distribution over  $A_i$  given  $q_{-i}$ .

Next, we show that every conditional mixed strategy  $\sigma_i$  can be induced by a probability measure  $\mu$  over conditional (pure) strategies.

**Theorem 1.** For every  $\sigma_i \in \Sigma_i$  there exists a  $\mu \in \Delta \hat{S}_i$  such that  $\phi(\mu) = \sigma_i$  where  $\phi$  is defined in (1).

*Proof.* Fix  $\sigma_i \in \Sigma_i$  and consider the induced partition over  $\Delta A_{-i}$   $\mathcal{P}_{\sigma_i} = \{X_1, \dots, X_{L_{\sigma_i}}\}$ . For ease of notation, we denote  $\mathcal{P} = \mathcal{P}_{\sigma_i}$  and  $L = L_{\sigma_i}$ . Note that  $\sigma_i$  can be written as

$$\begin{aligned} \sigma_i : \mathcal{P} &\rightarrow \Delta A_i \\ l &\mapsto \sigma_i(l) = \mu_l \end{aligned}$$

where  $\mu_l$  is a probability measure over  $A_i$ . Put differently, for  $a_i \in A_i$ ,  $\mu_l(a_i) = \sigma_i(q_{-i})(a_i)$  for all  $q_{-i} \in X_l$ . The finite collection of probability spaces  $(A_i, 2^{A_i}, \mu_l)_{l=1}^L$  induces the product

space  $(\times_{l=1}^L A_i, 2^{\times_{l=1}^L A_i}, \mu)$  where  $\mu$  is the usual product measure over  $2^{\times_{l=1}^L A_i}$  defined by  $\mu((a_i^1, a_i^2, \dots, a_i^L)) = \prod_{l=1}^L \mu_l(a_i^l)$  for all  $(a_i^1, a_i^2, \dots, a_i^L)$  in  $\times_{l=1}^L A_i = \{(a_i^1, a_i^2, \dots, a_i^L) | a_i^l \in A_i\}$ .

Note that every element  $(a_i^1, a_i^2, \dots, a_i^L) \in \times_{l=1}^L A_i$  can be identified with a  $\hat{s}_i \in \hat{S}_i|_{\mathcal{P}}$  where  $\hat{S}_i|_{\mathcal{P}} = \{\hat{s}_i \in \hat{S}_i | \hat{s}_i \text{ is } \mathcal{P}\text{-measurable}\}$ , i.e.,  $\hat{s}_i(q_{-i}) = a_i^l$  for all  $l$  and all  $q_{-i} \in X_l$ . Thus,  $\mu$  is a measure supported by  $\hat{S}_i|_{\mathcal{P}}$  such that for all  $\hat{s}_i \in \hat{S}_i|_{\mathcal{P}}$ ,  $\mu(\hat{s}_i) = \prod_{l=1}^L \mu_l(\hat{s}_i(l))$  where  $\hat{s}_i(l) = \hat{s}_i(q_{-i})$  for all  $q_{-i} \in X_l$ .

Next we show that  $\phi(\mu) = \sigma_i$ . Recall that by (1)  $\phi(\mu) = \sigma_i^\mu$ , which is defined by for all  $q_{-i}$  and all  $a_i$

$$\sigma_i^\mu(q_{-i})(a_i) = \int_{\hat{S}_i} \hat{s}_i(q_{-i})(a_i) d\mu(\hat{s}_i) = \int_{\hat{S}_i|_{\mathcal{P}}} \hat{s}_i(q_{-i})(a_i) d\mu(\hat{s}_i).$$

Note that

$$\hat{s}_i(q_{-i})(a_i) = \begin{cases} 0 & \text{if } \hat{s}_i(q_{-i}) \neq a_i \\ 1 & \text{if } \hat{s}_i(q_{-i}) = a_i, \end{cases}$$

and, therefore,

$$\sigma_i^\mu(q_{-i})(a_i) = \int \mathbb{1}_{\{\hat{s}_i \in \hat{S}_i|_{\mathcal{P}} : \hat{s}_i(q_{-i}) = a_i\}} d\mu(\hat{s}_i) = \mu(\{\hat{s}_i \in \hat{S}_i|_{\mathcal{P}} : \hat{s}_i(q_{-i}) = a_i\}).$$

Fix  $a_i \in A_i$  and  $q_{-i} \in X_{l'}$  for some  $l' \in \{1, \dots, L\}$ . Then,  $\{\hat{s}_i \in \hat{S}_i|_{\mathcal{P}} : \hat{s}_i(q_{-i}) = a_i\} = \{\times_{l=1}^{l'-1} A_i \times \{a_i\} \times \times_{l=l'+1}^L A_i\}$  and since  $\mu$  is the product measure,

$$\mu(\times_{l=1}^{l'-1} A_i \times \{a_i\} \times \times_{l=l'+1}^L A_i) = \mu_{l'}(a_i) = \sigma_i(q_{-i})(a_i).$$

This implies that for all  $q_{-i} \in \Delta A_{-i}$  and all  $a_i \in A_i$ ,  $\sigma_i^\mu(q_{-i})(a_i) = \sigma_i(q_{-i})(a_i)$ , i.e.,  $\phi(\mu) = \sigma_i$ .  $\square$

**Theorem 2.** *In every  $n$ -person game, there exists a pure conditional equilibrium.*

*Proof.* Let  $BR_i(\sigma_{-i}) = \{\sigma_i \in \Sigma_i | \sigma_i \in \arg \max_{\sigma'_i \in \Sigma_i} U_i(\sigma'_i, \sigma_{-i})\}$ . Let  $\bar{S}_i$  denote the set of constant strategies of player  $i$ —i.e.,  $\bar{s}_i \in \bar{S}_i$  if and only if for every  $a_{-i}$   $\bar{s}_i(a_{-i}) = a_i$ .

Player 1 chooses  $\hat{s}_1$  such that  $\hat{s}_1(a_{-1}) \in BR_1(a_{-1})$  for every  $a_{-1} \in A_{-1}$ . Let  $\hat{a}_2(a_3, \dots, a_n) \in \arg \max_{a'_2 \in A_2} u_2(\hat{s}_1(a'_2, a_3, \dots, a_n), a'_2, a_3, \dots, a_n)$ . Then, define the conditional strategy of player 2 as  $\hat{s}_2(a_{-2}) = \hat{a}_2(a_3, \dots, a_n)$  for every  $a_{-2}$ . In words, fixing the action profile  $(a_3, \dots, a_n)$  player 2 chooses an action that maximizes her utility given that player 1 best responds to her action.

Analogously, consider  $\hat{a}_3(a_4, \dots, a_n)$  in

$$\arg \max_{a'_3 \in A_3} u_3(\underbrace{\hat{s}_1(\underbrace{\hat{a}_2(a'_3, \dots, a_n)}_{\hat{a}_2}, a'_3, a_4, \dots, a_n)}_{\hat{a}_1}, \underbrace{\hat{a}_2(a'_3, \dots, a_n)}_{\hat{a}_2}, a'_3, a_4, \dots, a_n).$$

Then, define the conditional strategy of player 3 as  $\hat{s}_3(a_{-3}) = \hat{a}_3(a_4, \dots, a_n)$  for every  $a_{-3}$ . Conditional strategy of player  $k$  is defined as  $\hat{s}_k(a_{-k}) = \hat{a}_k(a_{(k+1)}, \dots, a_n)$  for every  $a_{-k}$  where  $\hat{a}_k$  is defined analogously. For player  $n$ ,  $\hat{a}_n \in \arg \max_{a'_n \in A_n} u_n(\hat{a}_1, \dots, \hat{a}_{n-1}, a'_n)$ , and  $\hat{s}(a_{-n}) = \hat{a}_n$ . Define with a slight abuse of notation  $\hat{a} = (\hat{a}_1, \hat{a}_2, \dots, \hat{a}_n)$  where  $\hat{a}_i = \hat{a}_i(\hat{a}_{i+1}, \dots, \hat{a}_n)$

Next, we show that  $\hat{s} \in B$ , i.e., it is an agreement. We will prove that  $s(\hat{a}) = \hat{a}$  and that for all  $a$  such that  $\hat{s}(a) = a$ , then  $a = \hat{a}$ . This is because  $\hat{s}_n(a_{-n}) = \hat{a}_n$  by definition of  $\hat{s}_n$ , so  $a_n = \hat{a}_n$ . Then, given  $\hat{a}_n$ ,  $\hat{s}_{n-1}(a_{-(n-1)}) = \hat{a}_{n-1}(a_n) = \hat{a}_{n-1}(\hat{a}_n) = \hat{a}_{n-1}$  by definition of  $\hat{s}_{n-1}$ , so  $a_{n-1} = \hat{a}_{n-1}$ . We repeat this process till we obtain  $a_2 = \hat{a}_2$ . Then, notice that  $\hat{s}_1(a_{-1}) = BR_1(a_{-1}) = BR_1(\hat{a}_{-1}) = \hat{a}_1$ .

Now we show that  $\hat{s}$  is a conditional equilibrium. First, we prove that player  $n$  has no unilateral profitable deviation from  $\hat{s}$ . Suppose by way of contradiction that  $\tilde{s}_n$  is such a deviation. Consider  $\tilde{s} = (\tilde{s}_n, \hat{s}_{-n})$ , which must be an agreement in  $B$  because otherwise  $U_n(\tilde{s}) = 0$ . Then, there must be a unique  $\tilde{a} \in A$  such that  $\tilde{s}(\tilde{a}) = \tilde{a}$ . Let  $\tilde{a}_n = \tilde{s}_n(\tilde{a}_{-n})$ . Then,  $u_n(\tilde{a}_n, \hat{a}_{-n}) \leq u_n(\hat{a}) = \max_{a'_n \in A_n} u_n(a'_n, \hat{a}_{-n})$ . Thus,  $\tilde{s}_n$  is not a unilateral profitable deviation. For every player  $k > 1$ , the reasoning is analogous. Finally, we show that player 1 has no unilateral profitable deviation. Fixing  $\hat{s}_{-1}$  induces the action profile  $\hat{a}_{-1}$ . Since  $\hat{s}_1$  is by definition a best reply to  $\hat{a}_{-1}$ , player 1 cannot strictly benefit from a unilateral deviation.  $\square$

## References

- Nash, J. F. (1950). Equilibrium points in n-person games. *Proceedings of the National Academy of Sciences* 36(1), 48–49.
- Nigel, H. (1971). Paradoxes of Rationality: Theory of Metagames and Political Behavior.
- von Neumann, J. and O. Morgenstern (1944). *Theory of Games and Economic Behavior* (1953, 3rd ed.). Princeton: Princeton University Press.